arXiv:2402.15387v1 [q-fin.RM] 23 Feb 2024 [arXiv:2402.15387v1 \[q-fin.RM\] 23 Feb 2024](http://arxiv.org/abs/2402.15387v1)

Higher order measures of risk and stochastic dominance

Alois Pichler[∗]

February 26, 2024

Abstract

Higher order risk measures are stochastic optimization problems by design, and for this reason they enjoy valuable properties in optimization under uncertainties. They nicely integrate with stochastic optimization problems, as has been observed by the intriguing concept of the risk quadrangles, for example.

Stochastic dominance is a binary relation for random variables to compare random outcomes. It is demonstrated that the concepts of higher order risk measures and stochastic dominance are equivalent, they can be employed to characterize the other. The paper explores these relations and connects stochastic orders, higher order risk measures and the risk quadrangle.

Expectiles are employed to exemplify the relations obtained.

Keywords: Higher order risk measure · higher order stochastic dominance · risk quadrangle **Classification:** 62G05, 62G08, 62G20

1 Introduction

Risk measures are considered in various disciplines to assess and quantify risk. Similarly to assigning a premium to an insurance contract with random losses after appraising its risk, risk measures assign a number to a random variable, which itself has stochastic outcomes.

This paper focuses on higher order risk measures, as these risk measures naturally combine with stochastic optimization problems or in 'learning' objectives, as they are given as the result of optimization problems. In addition, these risk measures relate to the risk quadrangle.

The paper derives explicit representations of higher order risk measures for general, elementary risk measures in a first main result. These characterizations are employed to characterize stochastic dominance relations, which are built on general norms. The second main result is a verification theorem. This is a characterization of higher order stochastic dominance relations, which is numerically tractable.

For Hölder norms, stochastic dominance relations have been considered for example in Dupačová and Kopa [\[11\]](#page-22-0), Kopa et al. [\[15](#page-22-1), [16\]](#page-22-2) and Consigli et al. [\[7](#page-22-3)], in portfolio optimization involving commodities (cf. Frydenberg et al. [\[13](#page-22-4)]), and by Dentcheva and Martinez [\[8](#page-22-5)] and Maggioni and Pflug [\[21](#page-23-0), [20\]](#page-23-1) in a multistage setting. The paper employs the characterizations obtained to establish relations for general norms. A comparison of these methods is given in Gutjahr and Pichler [\[14\]](#page-22-6). The paper illustrates these connections for expectiles (Bellini et al. [\[5](#page-22-7)], Bellini and Caperdoni [\[3\]](#page-22-8)) and adds a comparison with other risk measures.

[∗]Technische Universität Chemnitz, Faculty of Mathematics. 90126 Chemnitz, Germany Contact: alois.pichler@math.tu-chemnitz.de © orcid.org/0000-0001-8876-2429

Outline of the paper. The following Section [2](#page-1-0) recalls the mathematical framework for higher order risk measures. Section [3](#page-4-0) addresses the higher order risk measure associated with the spectral risks, as these risk measures constitute an elementary building block for general risk measures. This section develops the first main result, which is an explicit representation of a spectral risk's higher order risk measure. As a special case, the subsequent Section [4](#page-10-0) links and relates stochastic dominance and higher order risk measures. This section presents the second main result, which allows verifying a stochastic dominance relation by involving only finitely many risk levels. The final Section [5](#page-18-0) addresses the expectile and establishes the relations of the preceding sections for this specific risk measure. Section [6](#page-21-0) concludes.

2 Mathematical Framework

Higher order risk measures are a special instance of *risk measures*, often also termed *risk functionals*. To introduce and recall their main properties we consider a space $\mathcal Y$ of R-valued random variables on a probability space with measure P containing at least all bounded random variables, that is, $L^{\infty}(P) \subseteq \mathcal{Y}$. A risk measure then satisfies the following axioms, originally introduced by Artzner et al. [\[1](#page-21-1)].

Definition 2.1 (Risk functional). Let \mathcal{Y} be a space of \mathbb{R} -valued random variables on a probability space (Ω, Σ, P) . A mapping $\mathcal{R}: \mathcal{Y} \to \mathbb{R}$ is

- (i) monotone, if $\mathcal{R}(X) \leq \mathcal{R}(Y)$, provided that $X \leq Y$ almost everywhere;
- (ii) positively homogeneous, if $\mathcal{R}(\lambda Y) = \lambda \mathcal{R}(Y)$ for all $\lambda > 0$;
- (iii) translation equivariant, if $\mathcal{R}(c + Y) = c + \mathcal{R}(Y)$ for all $c \in \mathbb{R}$;
- (iv) subadditive, if $\mathcal{R}(X+Y) \leq \mathcal{R}(X) + \mathcal{R}(Y)$ for all X and $Y \in \mathcal{Y}$.

A mapping satisfying [\(i\)–](#page-1-1)[\(iv\)](#page-1-2) is called a *risk functional*, or a *risk measure*.

The risk quadrangle (cf. Rockafellar and Uryasev [\[34\]](#page-23-2)) interconnects risk measures with the regular measure of deviation, error and regret by

$$
\mathcal{R}(Y) = \inf_{c \in \mathbb{R}} c + \mathcal{V}(Y - c),\tag{2.1}
$$

where V is called *regret function* (or *optimized certainty equivalent*, cf. Ben-Tal and Teboulle [\[6\]](#page-22-9)). It follows from the relation [\(2.1\)](#page-1-3) that \mathcal{R} – if given as in (2.1) – is translation equivariant, i.e, \mathcal{R} satisfies $\mathcal{R}(Y + c) = c + \mathcal{R}(Y)$ for any $c \in \mathbb{R}$ (cf. [\(iii\)](#page-1-4) above). In an economic interpretation, the amount c in [\(2.1\)](#page-1-3) corresponds to an amount of cash spent today, while the remaining quantity $Y - c$ is invested and consumed later, thus subject to V .

The risk functional R is positively homogeneous, if the regret function $\mathcal V$ is positively homogeneous. If $\mathcal V$ is not positively homogeneous, then one may consider the positively homogeneous envelope

$$
\mathcal{V}_{\tilde{\beta}}(Y) = \inf_{t>0} t\left(\tilde{\beta} + \mathcal{V}\left(\frac{Y}{t}\right)\right),\,
$$

where $\tilde{\beta} \ge 0$ is a risk aversion coefficient. The combined functional

$$
\mathcal{R}_{\beta}(Y) = \inf_{c \in \mathbb{R}} c + \mathcal{V}_{\beta}(Y - c)
$$

=
$$
\inf_{\substack{t > 0 \\ q \in \mathbb{R}}} t\left(\tilde{\beta} + q + \mathcal{V}\left(\frac{Y}{t} - q\right)\right)
$$
 (2.2)

is positively homogeneous and translation equivariant (cf. [\(ii\)](#page-1-5) and [\(iii\)\)](#page-1-4). The φ -divergence risk measure is an explicit example of a risk measure, which is defined exactly as [\(2.2\)](#page-1-6), cf. Dommel and Pichler [\[10\]](#page-22-10).

In what follows, we shall address the reverse question first. That is, given the risk functional R , what is the regret functional V so that [\(2.1\)](#page-1-3) holds true? To this end consider a space $\mathcal{Y} \subset L^1(P)$ endowed with norm $\| \cdot \|$. We shall assume the norm to be monotone, that is, $\|X\| \le \|Y\|$ provided that $0 \le X \le Y$ almost everywhere. We associate the following family of risk measure with a given norm.

Definition 2.2 (Higher order risk measure). Let $\|\cdot\|$ be a monotone norm on $\mathcal{Y} \subset L^1(P)$ with $\mathbb{E} 1 = 1$, where $1(\cdot) = 1$ is the identically one function on *Y*. The *higher order risk measure* at risk level $\beta \in [0, 1)$ associated with the norm $\|\cdot\|$ is

$$
\mathcal{R}_{\beta}^{\|\cdot\|}(Y) = \inf_{t \in \mathbb{R}} t + \frac{1}{1 - \beta} \|(Y - t)_{+}\|,\tag{2.3}
$$

where $\beta \in [0, 1)$ is the *risk aversion coefficient* and $x_{+} := \max(0, x)$.

We shall also omit the superscript and write \mathcal{R}_{β} instead of $\mathcal{R}_{\beta}^{\|\cdot\|}$ $^{\parallel}$ $^{\parallel}$ in case the norm is unambiguous given the context. We shall demonstrate first that the higher order risk measure is well-defined for any $\beta \geq 0$.

Proposition 2.3. *Let* $(\mathcal{Y}, \|\cdot\|)$ *be a normed space of random variables. For the functional* \mathcal{R}_{β} *defined in* [\(2.3\)](#page-2-0) *it holds that*

$$
-\|Y\| \le \mathcal{R}_{\beta}(Y) \le \frac{1}{1-\beta} \|Y\|,\tag{2.4}
$$

so that $\mathcal{R}_{\beta}(\cdot)$ *is indeed well-defined on* $(\mathcal{Y}, \| \cdot \|)$ *for every* $\beta \in [0, 1)$ *.*

Proof. The upper bound follows trivially from the definition by choosing $t = 0$ in the defining equation [\(2.3\)](#page-2-0).

For $t \leq 0$, it holds that $-t = -Y + (Y - t) \leq -Y + (Y - t)$. It follows from the triangle inequality that $-t \leq ||Y|| + ||(Y - t)_+||$ and thus

$$
-\|Y\| \le t + \|(Y - t)_+\| \quad \text{for all } t \le 0.
$$

To establish the relation also for $t \geq 0$, we start by observing the following monotonicity property of the objective in [\(2.3\)](#page-2-0) in addition: for $\Delta t \ge 0$, it follows from the reverse triangle inequality that

$$
||Y_{+}|| - ||(Y - \Delta t)_{+}|| \le ||Y_{+} - (Y - \Delta t)_{+}|| \le ||\Delta t \, 1|| = \Delta t,
$$

where we have used that $0 \le Y_+ - (Y - \Delta t)_+ \le \Delta t$ together with monotonicity of the norm. Replacing Y by $Y - t$ in the latter display gives

$$
t + ||(Y - t)_+|| \le t + \Delta t + ||(Y - (t + \Delta t))_+||;
$$

that is, the function $t \mapsto t + ||(Y - t)_+||$ is non-decreasing, which finally establishes that

$$
-\|Y\| \le t + \|(Y - t)_+\| \quad \text{for all } t \in \mathbb{R}.
$$

The lower bound in [\(2.4\)](#page-2-1) thus follows from the latter inequality, as $\mathcal{R}_0(Y) \leq \mathcal{R}_B(Y)$ for any $\beta \geq 0$.

For Hölder spaces (i.e., $L^p(P)$ spaces with $p \ge 1$ and norm $||Y||_p := (||P||^p)^{1/p}$), the higher order risk measure has been introduced in Krokhmal [\[17](#page-22-11)] and studied in Dentcheva et al. [\[9\]](#page-22-12).

Lemma 2.4. $\mathcal{R}_{\beta}(\cdot)$ *is a risk functional, provided that the norm is monotone. Further,* \mathcal{R}_{β} *is Lipschitz continuous with respect to the norm, the Lipschitz constant is* $\frac{1}{1-\beta}$ *.*

Proof. The assertions [\(ii\)](#page-1-5)[–\(iv\)](#page-1-2) in Definition [2.1](#page-1-7) are straight forward to verify; to verify [\(i\)](#page-1-1) it is indispensable to assume that the norm is monotone.

As for continuity, it follows from subadditivity together with [\(2.4\)](#page-2-1) that $\mathcal{R}_{\beta}(Y) - \mathcal{R}_{\beta}(Z) \leq \mathcal{R}_{\beta}(Y - Z) \leq$ $\frac{1}{1-\beta} ||Y - Z||$, and $|\mathcal{R}_{\beta}(Y) - \mathcal{R}_{\beta}(Z)| \leq \frac{1}{1-\beta} ||Y - Z||$ after interchanging the roles of Y and Z. Hence, the \Box assertion.

Note that the higher order risk measure as defined in [\(2.3\)](#page-2-0) defines a risk functional based on a norm. In contrast to this construction, a risk functional R defines a norm via

$$
Y \parallel \coloneqq \mathcal{R}(|Y|) \tag{2.5}
$$

and a Banach space with $\mathcal{Y} = \{Y \in L^1 : \mathcal{R}(|Y|) < \infty\}$ (cf. Pichler [\[30](#page-23-3)]). Its natural dual norm for $Z \in \mathcal{Z} \coloneqq \mathcal{Y}^*$ is

$$
||Z||^* := \sup \{ \mathbb{E} YZ : ||Y|| \le 1 \} = \sup \{ \mathbb{E} YZ : \mathcal{R}(|Y|) \le 1 \} .
$$
 (2.6)

The following relationship allows defining a regret functional to connect a risk functional R with the risk quadrangle.

Proposition 2.5 (Duality). Let R be a risk functional with associated norm $\|\cdot\|$ and dual norm $\|\cdot\|^*$. *For the higher order risk functional it holds that*

$$
\mathcal{R}_{\beta}(Y) = \sup \left\{ \mathbb{E} \, YZ : Z \ge 0, \ \mathbb{E} \, Z = 1 \ and \ ||Z||^* \le \frac{1}{1 - \beta} \right\} \tag{2.7}
$$

$$
= \inf_{t \in \mathbb{R}} t + \frac{1}{1 - \beta} ||(Y - t)_+||, \tag{2.8}
$$

where $\beta \in [0, 1)$ *.*

Remark 2.6. By the interconnecting formula [\(2.1\)](#page-1-3), the higher order risk functional $\mathcal{R}_R^{\|\cdot\|}$ $\frac{N}{\beta}$ associated with the norm $\|\cdot\|$ is the regret function $\mathcal{V}(\cdot) \coloneqq \frac{1}{1-\beta} \|(\cdot)_+\|$.

Proof. It holds by the Hahn–Banach theorem and as $(Y - t)_+ \geq 0$ that

$$
\frac{1}{1-\beta} \cdot \| (Y-t)_+\| = \sup_{\|Z\|^* \le \frac{1}{1-\beta}} \mathbb{E} |Z(Y-t)_+\ge \sup_{\substack{\mathbb{E} |Z=1, \ Z\ge 0, \\ \|Z\|^* \le \frac{1}{1-\beta}}} \mathbb{E} |Z(Y-t)_+.
$$

This establishes the first inequality ' \leq ' in [\(2.8\)](#page-3-0) with $t + (Y - t)_+ \geq Y$, as

$$
t + \frac{1}{1 - \beta} \cdot ||(Y - t)_+|| \ge \sup_{\substack{\mathbb{E} | Z = 1 \\ Z \ge 0, ||Z||^* \le \frac{1}{1 - \beta} \\ \ge \sup_{\substack{\mathbb{E} | Z = 1 \\ \mathbb{E} | Z = 1 \\ Z \ge 0, ||Z||^* \le \frac{1}{1 - \beta} }} \mathbb{E} YZ.
$$

As for the converse inequality assume first that Y is bounded. Note, that

$$
\inf_{t \in \mathbb{R}} t + \mathbb{E}(Y - t)Z = \mathbb{E} YZ + \inf_{t \in \mathbb{R}} t \cdot (1 - \mathbb{E} Z) = \begin{cases} \mathbb{E} YZ & \text{if } \mathbb{E} Z = 1, \\ -\infty & \text{else,} \end{cases}
$$

so that it follows that

$$
\sup_{\substack{\mathbb{E}\,Z=1\\ Z\geq 0,\ \|\mathcal{Z}\|^* \leq \frac{1}{1-\beta}}} \mathbb{E}\, YZ = \sup_{\substack{Z\geq 0,\ \text{if $\mathbb{E}\,(\,Y-\text{f}\,)\,Z$}.\\ \|Z\|^* \leq \frac{1}{1-\beta}}} \inf_{t\in\mathbb{R}}
$$

Further, it holds that $EYZ = t^* + EZ(Y - t^*)_+$ for $t^* \leq Y$ a.s. and thus

$$
\sup_{\substack{\mathbb{E} Z=1, Z\geq 0, \\ \|Z\|^* \leq \frac{1}{1-\beta}}} \mathbb{E} YZ = \sup_{\substack{Z\geq 0, \\ \|Z\|^* \leq \frac{1}{1-\beta}}} t^* + \mathbb{E} Z(Y-t^*)_+ = t^* + \frac{1}{1-\beta} \|(Y-t^*)\| \geq \inf_{t\in\mathbb{R}} t + \frac{1}{1-\beta} \|(Y-t)_+\|,
$$

thus the desired converse inequality, provided that Y is bounded; if Y is not bounded, then there is a bounded Y_{ε} with $Y \leq Y_{\varepsilon}$ ($\varepsilon > 0$) and $||Y_{\varepsilon} - Y|| < \varepsilon$, so that

$$
\mathbb{E} Z(Y_{\varepsilon}-t)_{+}-\varepsilon \mathbb{E} Z \leq \mathbb{E} Z(Y-t)_{+} \leq \mathbb{E} Z(Y_{\varepsilon}-t)_{+},
$$

so that we may conclude that [\(2.8\)](#page-3-0) holds for every $Y \in \mathcal{Y}$.

Example 2.7 (Hölder spaces). The dual norm of the genuine norm $||X||_p \coloneqq (E|X|^p)^{1/p}$ in the Hölder space $L^p(P)$ is $||Z||^* = (\mathbb{E}|Z|^q)^{1/q}$ for the Hölder conjugate exponent with $\frac{1}{p} + \frac{1}{q} = 1$. With Proposition [2.5](#page-3-1) it follows that

$$
\mathcal{R}_{\beta}^{\|\cdot\|_{p}}(Y) = \inf_{t \in \mathbb{R}} t + \frac{1}{1-\beta} \| (Y-t)_{+} \|_{p}
$$

= sup $\left\{ \mathbb{E} YZ : \|Z\|_{q} \le \frac{1}{1-\beta}, Z \ge 0 \text{ and } \mathbb{E} Z = 1 \right\},\$

cf. also Pichler and Shapiro [\[32\]](#page-23-4) and Pichler [\[31](#page-23-5)].

In what follows, we shall elaborate the higher order risk measure and the associated regret function for specific risk measures, specifically the spectral risk measure.

3 Higher order spectral risk

By Kusuoka's theorem (cf. Kusuoka [\[18](#page-22-13)]), every law invariant risk functional can be assembled by elementary risk functionals, each involving the average value-at-risk.

The following section develops the explicit representations of the higher order risk measures associated with spectral risk measures first. The explicit representation then is extended to general risk functionals.

Definition 3.1 (Spectral risk measures). The function σ : $[0, 1) \rightarrow \mathbb{R}$ is called a *spectral function*, if

(i) $\sigma(\cdot) \geq 0$,

- (ii) $\int_0^1 \sigma(u) du = 1$ and
- (iii) $\sigma(\cdot)$ is non-decreasing.

The *spectral risk measure* with spectral function σ is

$$
\mathcal{R}_{\sigma}(Y) \coloneqq \int_0^1 \sigma(u) F_Y^{-1}(u) \, \mathrm{d}u,
$$

where

$$
F_Y^{-1}(u) := \mathsf{V@R}_u(Y) := \inf \{x \in \mathbb{R} \colon P(Y \le x) \ge u\}
$$

is the *value-at-risk*, the *generalized inverse* or *quantile function*.

The higher order risk measure of the spectral risk measure is a spectral risk measure itself. The following theorem presents the corresponding spectral function explicitly and generalizes Pflug [\[28](#page-23-6)]. The result is central towards the main characterization presented in the next sections.

Theorem 3.2 (Higher order spectral risk). Let $\beta \in [0, 1)$ be a risk level. The higher order risk functional *of the risk functional* \mathcal{R}_{σ} *with spectral function* $\sigma(\cdot)$ *has the representation*

$$
\inf_{t \in \mathbb{R}} t + \frac{1}{1 - \beta} \mathcal{R}_{\sigma} \left((Y - t)_{+} \right) = \mathcal{R}_{\sigma_{\beta}}(Y), \tag{3.1}
$$

where σ_B *is the spectral function*

$$
\sigma_{\beta}(u) \coloneqq \begin{cases} 0 & \text{if } u < u_{\beta}, \\ \frac{\sigma(u)}{1-\beta} & \text{else}; \end{cases} \tag{3.2}
$$

here, $u_{\beta} \in \mathbb{R}$ *is the* β *-quantile with respect to the density* σ *, that is, the solution of*

$$
\int_0^{u_\beta} \sigma(u) \, \mathrm{d}u = \beta,\tag{3.3}
$$

which is unique for $\beta > 0$ *.*

Proof. We remark first that σ_{β} indeed is a spectral function, as $\int_0^1 \sigma_{\beta}(u) du = \frac{1}{1-\beta} \int_{u_{\beta}}^1 \sigma(u) du = \frac{1-\beta}{1-\beta}$ $\frac{1-\beta}{1-\beta} = 1$ by the defining property [\(3.3\)](#page-5-0) and [\(ii\)](#page-4-1) in Definition [3.1.](#page-4-2) The quantile u_{β} is uniquely defined for $\beta > 0$, as the function σ is non-decreasing by [\(iii\).](#page-4-3) In what follows we shall demonstrate that the infimum in [\(3.1\)](#page-5-1) is attained at $t^* := F_Y^{-1}(u_\beta)$. Note first that

$$
F_{(Y-t)_+}^{-1}(u) = \begin{cases} 0 & \text{if } u < F_Y(t), \\ F_Y^{-1}(u) - t & \text{else,} \end{cases}
$$

so that

$$
\mathcal{R}_{\sigma}\big((Y-t)_{+}\big) = \int_{0}^{1} \sigma(u) F_{(Y-t)_{+}}^{-1}(u) du = \int_{F_{Y}(t)}^{1} \sigma(u) \big(F_{Y}^{-1}(u) - t\big) du
$$

and

$$
(\mathcal{R}_{\sigma})_{\beta}(Y) = \inf_{t \in \mathbb{R}} t + \frac{1}{1 - \beta} \int_{F_Y(t)}^1 \sigma(u) (F_Y^{-1}(u) - t) \, \mathrm{d}u. \tag{3.4}
$$

.

Assume first that $t \leq t^*$. The inequality $u \leq F_Y(t)$ is equivalent to $F_Y^{-1}(u) \leq t$ (cf. van der Vaart [\[36\]](#page-24-0); this relation of functions F_Y and F_Y^{-1} is occasionally called a *Galois connection*), and thus

$$
\int_{F_Y(t)}^{F_Y(t^*)} \sigma(u) \big(F_Y^{-1}(u) - t \big) \, \mathrm{d}u \le 0,
$$

or equivalently

$$
\int_{F_Y(t)}^1 \sigma(u) \big(F_Y^{-1}(u) - t \big) \, \mathrm{d}u \le \int_{F_Y(t^*)}^1 \sigma(u) \big(F_Y^{-1}(u) - t \big) \, \mathrm{d}u.
$$

Assume next that $u_{\beta} \leq F_Y(t^*)$, then $\int_{F_Y(t^*)}^1 \sigma(u) du \leq 1 - \beta$ so that

$$
\frac{t-t^*}{1-\beta} \int_{F_Y(t^*)}^1 \sigma(u) \, \mathrm{d}u \le t - t^*
$$

Combining the inequalities in the latter displays gives

$$
t^* + \frac{1}{1-\beta} \int_{F_Y(t^*)}^{1} \sigma(u) \left(F_Y^{-1}(u) - t^* \right) \, \mathrm{d}u \le t + \frac{1}{1-\beta} \int_{F_Y(t)}^{1} \sigma(u) \left(F_Y^{-1}(u) - t \right) \, \mathrm{d}u \tag{3.5}
$$

and thus the assertion, provided that $u_{\beta} \leq F_Y(t^*)$ and $t^* \leq t$.

Conversely, assume that $t \leq t^*$. Then the inequality $u \leq F_Y(t^*)$ is equivalent to $F_Y^{-1}(u) \leq t^*$ and thus

$$
\int_{F_Y(t)}^{F_Y(t^*)} \sigma(u) \big(F_Y^{-1}(u) - t^* \big) \, \mathrm{d}u \le 0,
$$

which is equivalent to

$$
\int_{F_Y(t)}^1 \sigma(u) \big(F_Y^{-1}(u) - t^* \big) \, \mathrm{d}u \le \int_{F_Y(t^*)}^1 \sigma(u) \big(F_Y^{-1}(u) - t^* \big) \, \mathrm{d}u.
$$

Assume further that $F_Y(t^*) \le u_\beta$, then $\int_{F_Y(t^*)}^1 \sigma(u) du \ge 1 - \beta$ so that

$$
t^* - t \le \frac{t^* - t}{1 - \beta} \int_{F_Y(t^*)}^{1} \sigma(u) \, \mathrm{d}u.
$$

Combining the latter inequalities gives

$$
t^* + \frac{1}{1-\beta} \int_{F_Y(t)}^1 \sigma(u) \left(F_Y^{-1}(u) - t^* \right) \mathrm{d}u \le t + \frac{1}{1-\beta} \int_{F_Y(t^*)}^1 \sigma(u) \left(F_Y^{-1}(u) - t \right) \mathrm{d}u. \tag{3.6}
$$

It follows from [\(3.5\)](#page-5-2) and [\(3.6\)](#page-6-0) that $t^* := F_Y^{-1}(u_\beta)$ is optimal in [\(3.4\)](#page-5-3). That is,

$$
(\mathcal{R}_{\sigma})_{\beta}(Y) = t^* + \frac{1}{1-\beta} \int_{u_{\beta}}^{1} \sigma(u) (F_Y^{-1}(u) - t^*) du
$$

=
$$
\frac{1}{1-\beta} \int_{u_{\beta}}^{1} \sigma(u) F_Y^{-1}(u) du
$$

=
$$
\int_0^1 \sigma_{\beta}(u) F_Y^{-1}(u) du
$$
 (3.7)

and thus the assertion.

The following statement expresses the higher order risk functional by at the base value u_{β} , and the random variable's aberrations to the right, involving the survival function instead of its inverse distribution function.

Corollary 3.3. *The higher order spectral risk measure is*

$$
(\mathcal{R}_{\sigma})_{\beta}(Y) = \mathsf{V} \circledcirc \mathsf{R}_{u_{\beta}}(Y) + \frac{1}{1-\beta} \int_{\mathsf{V} \circledcirc \mathsf{R}_{u_{\beta}}(Y)}^{\infty} \Sigma(F_Y(y)) \mathrm{d}y \tag{3.8}
$$

(with μ *^{<i>B*} *as in* [\(3.3\)](#page-5-0)*) or, provided that Y is bounded,*

$$
(\mathcal{R}_{\sigma})_{\beta}(Y) = \text{ess inf } Y + \int_{\text{ess inf } Y}^{\infty} \Sigma_{\beta}(F_Y(y)) \, dy,\tag{3.9}
$$

where

$$
\Sigma_{\beta}(u) := \min\left(1, \ \frac{1}{1-\beta} \int_{u}^{1} \sigma(p) \, \mathrm{d}p\right)
$$

is the cumulative spectral function and $\Sigma(u) \coloneqq \sum_0(u) = \int_u^1 \sigma(p) \, dp$.

Proof. Notice first that $\Sigma_{\beta}(u) = 1$ for $u \le u_{\beta}$, where u_{β} is given in [\(3.3\)](#page-5-0). By Theorem [3.2,](#page-5-4) Riemann–Stieltjes integration by parts and changing the variables it holds that

$$
(\mathcal{R}_{\sigma})_{\beta}(Y) = \mathcal{R}_{\sigma_{\beta}}(Y)
$$

\n
$$
= \frac{1}{1-\beta} \int_{u_{\beta}}^{1} \sigma(u) F_{Y}^{-1}(u) du
$$

\n
$$
= -\int_{0}^{1} F_{Y}^{-1}(u) d\Sigma_{\beta}(u)
$$

\n
$$
= -F_{Y}^{-1}(u) \Sigma_{\beta}(u) \Big|_{u=0}^{1} + \int_{0}^{1} \Sigma_{\beta}(u) dF_{Y}^{-1}(u)
$$

\n
$$
= \text{ess inf } Y + \int_{\text{ess inf } Y}^{\infty} \Sigma_{\beta}(F_{Y}(y)) dy,
$$
\n(3.11)

where we have used that $F_Y^{-1}(0) = \text{ess inf } Y$ and $\Sigma_{\beta}(1) = 0$ in [\(3.11\)](#page-7-0). This gives [\(3.9\)](#page-6-1).

The equation [\(3.8\)](#page-6-2) results from sticking to the lower bound u_β (instead of 0) in [\(3.10\)](#page-7-1). That is,

$$
\begin{aligned} \left(\mathcal{R}_{\sigma}\right)_{\beta}(Y) &= -\int_{u_{\beta}}^{1} F_{Y}^{-1}(u) \, \mathrm{d}\Sigma_{\beta}(u) \\ &= -F_{Y}^{-1}(u)\Sigma_{\beta}(u)\Big|_{u=u_{\beta}}^{1} + \int_{u_{\beta}}^{1} \Sigma_{\beta}(u) \, \mathrm{d}F_{Y}^{-1}(u) \\ &= \mathsf{V} \circledast \mathsf{R}_{u_{\beta}}(Y) + \int_{\mathsf{V} \circledast \mathsf{R}_{u_{\beta}}(Y)}^{\infty} \Sigma_{\beta}\big(F_{Y}(y)\big) \, \mathrm{d}y, \end{aligned}
$$

which is assertion [\(3.8\)](#page-6-2). \Box

Corollary 3.4. *The higher order spectral risk measure has the representation*

$$
\mathcal{R}_{\sigma_{\beta}}(Y) = \sup \{ \mathbb{E}[Y \cdot \sigma_{\beta}(U)] : U \in [0,1] \text{ is uniformly distributed} \}.
$$

Proof. Recall first that $Y \sim F_Y^{-1}(U)$ for U uniformly distributed. By the rearrangement inequality, $E Y \sigma_{\beta}(U) \leq E F_Y^{-1}(U) \sigma_{\beta}(U)$, because $F_Y^{-1}(U)$ and $\sigma_{\beta}(U)$ are comonotone and both, $F_Y^{-1}(\cdot)$ and $\sigma_{\beta}(\cdot)$ are non-decreasing functions. The assertion follows with (3.7) .

The celebrated formula (cf. Pflug [\[28\]](#page-23-6), Rockafellar and Uryasev [\[33\]](#page-23-7), Ogryczak and Ruszczyński [\[27](#page-23-8)])

$$
\text{AV@R}_{\alpha}(Y) = \frac{1}{1-\alpha} \int_{\alpha}^{1} \text{V@R}_{u}(Y) \, \mathrm{d}u = \inf_{t \in \mathbb{R}} \ t + \frac{1}{1-\alpha} \, \mathbb{E}(Y - t).
$$

for the average value-at-risk is a special case of preceding Theorem [3.2](#page-5-4) for the spectral function $\sigma(\cdot)$ = $\frac{1}{1-\alpha}1_{[\alpha,1]}(\cdot).$

The following corollary estabilshes this risk functional's higher order variant.

Corollary 3.5 (Average value-at-risk)**.** *The higher order average value-at-risk is*

$$
(\mathsf{AV@R}_{\alpha})_{\beta}(Y) = \mathsf{AV@R}_{1-(1-\alpha)(1-\beta)}(Y),\tag{3.12}
$$

where $Y \in L^1$ *; equivalently,*

$$
\text{AV@R}_{\beta}(Y) = \inf_{t \in \mathbb{R}} t + \frac{1}{1 - \frac{\beta - \alpha}{1 - \alpha}} \text{AV@R}_{\alpha}((Y - t)_{+}),\tag{3.13}
$$

where $\beta \geq \alpha$ *.*

Proof. The spectral function of the average value-at-risk is $\sigma_{\alpha}(\cdot) = \frac{1_{\cdot\geq\alpha}}{1-\alpha}$. It follows from [\(3.3\)](#page-5-0) that $u_{\beta} = \alpha + \beta(1 - \alpha) = 1 - (1 - \alpha)(1 - \beta)$ and $(\sigma_{\alpha})_{\beta} =$ $\int 0$ if $u \leq u_{\beta}$, $\frac{1}{(1-\alpha)(1-\beta)}$ else. function of the average value-at-risk at risk level u_{β} , thus the result.

The assertion [\(3.13\)](#page-7-2) follows by replacing β with $\frac{\beta-\alpha}{1-\alpha}$ in [\(3.12\)](#page-7-3). \Box

$$
\Box
$$

Corollary 3.6 (Kusoka representation spectral risk measures)**.** *Suppose the risk functional is*

$$
\mathcal{R}(Y) = \int_0^1 \mathsf{AV} \mathfrak{S} \mathsf{R}_{\gamma}(Y) \,\mu(\mathrm{d}\gamma),\tag{3.14}
$$

where 𝜇 *is a probability measure on* [0, 1]*. Then the higher order risk measure is*

$$
\mathcal{R}_{\beta}(Y) = \int_0^1 \mathsf{AV} \mathscr{Q} \mathsf{R}_{\gamma}(Y) \,\mu_{\beta}(\mathrm{d}\gamma),
$$

where μ ^{β} (*·*) *is the measure*

$$
\mu_{\beta}(A) := p_0 \cdot \delta_{u_{\beta}}(A) + \frac{1}{1 - \beta} \mu(A \cap (u_{\beta}, 1])
$$
\n(3.15)

and u_{β} and p_0 are determined by the equation and definition

$$
\int_0^{u_\beta} \frac{u_\beta - \alpha}{1 - \alpha} \mu(\mathrm{d}\alpha) = \beta \quad \text{and} \quad p_0 := \frac{1 - u_\beta}{1 - \beta} \int_0^{u_\beta} \frac{\mu(\mathrm{d}\alpha)}{1 - \alpha}.
$$

Proof. Above all, μ_{β} is a probability measure, because $p_0 \ge 0$ and

$$
\mu_{\beta}([0,1]) = p_0 + \frac{1}{1-\beta} \int_{u_{\beta}+}^{1} \mu(\mathrm{d}\alpha)
$$

=
$$
\frac{1}{1-\beta} \int_{0}^{u_{\beta}} \frac{1-\alpha - (u_{\beta}-\alpha)}{1-\alpha} \mu(\mathrm{d}\alpha) + \frac{1}{1-\beta} \int_{u_{\beta}+}^{1} \mu(\mathrm{d}\alpha)
$$

=
$$
\frac{1}{1-\beta} \int_{0}^{1} \mu(\mathrm{d}\alpha) - \frac{\beta}{1-\beta} = 1.
$$

The spectral function of the average value-at-risk at risk level α is $\sigma_{\alpha}(\cdot) = \frac{1_{\cdot\geq\alpha}}{1-\alpha}$. The quantile condition [\(3.3\)](#page-5-0) thus is

$$
\beta = \int_0^1 \frac{\max(0, u_\beta - \alpha)}{1 - \alpha} \mu(\mathrm{d}\alpha)
$$

and thus [\(3.16\)](#page-8-0).

For $u < u_\beta$, the spectral function corresponding to the measure \mathcal{R}_β in [\(3.14\)](#page-8-1) is 0, which coincides with [\(3.2\)](#page-5-5). For $u > u_{\beta}$, the spectral function for \mathcal{R}_{β} is

$$
\frac{p_0}{1 - u_\beta} 1_{u \ge u_\beta} + \int_{u_\beta}^1 \frac{1}{1 - \beta} \frac{1_{u \ge \alpha}}{1 - \alpha} \mu(\mathrm{d}\alpha) = \frac{1}{1 - \beta} \int_0^{u_\beta} \frac{1_{u \ge u_\beta}}{1 - \alpha} \mu(\mathrm{d}\alpha) + \int_{u_\beta}^1 \frac{1}{1 - \beta} \frac{1_{u \ge \alpha}}{1 - \alpha} \mu(\mathrm{d}\alpha) = \frac{1}{1 - \beta} \int_0^1 \frac{\mu(\mathrm{d}\alpha)}{1 - \alpha},
$$

which is the desired result in light of [\(3.2\)](#page-5-5).

In situations of practical interest, the risk measure is often given as finite combination of average values-at-risk at varying levels. The following corollary addresses this situation explicitly.

Corollary 3.7. *Suppose that*

$$
\mathcal{R}(Y) = \sum_{i=1}^{n} p_i \cdot \mathsf{AV} \circ \mathsf{R}_{\alpha_i}(Y) \tag{3.17}
$$

with $p_i \geq 0$, $\sum_{i=1}^n p_i = 1$ *and* $\alpha_i \in [0, 1]$ *for* $i = 1, \ldots, n$ *. Then*

$$
\mathcal{R}_{\beta}(Y) = p_0 \cdot \text{AV@R}_{u_{\beta}}(Y) + \sum_{i:\alpha_i > u_{\beta}} \frac{p_i}{1-\beta} \text{AV@R}_{\alpha_i}(Y),\tag{3.18}
$$

where u_{β} satisfies $\beta = \sum_{i=1}^{n} p_i \frac{\max(0, u_{\beta} - \alpha_i)}{1 - \alpha_i}$ $\frac{(0, u_{\beta} - \alpha_i)}{1 - \alpha_i}$ and $p_0 \coloneqq \sum_{i : \alpha_i \leq u_{\beta}} \frac{p_i}{1 - \alpha_i}$ $1-\alpha_i$ $1 - u_\beta$ $\frac{1-u_{\beta}}{1-\beta}$. *For large risk levels* 𝛽*, specifically if*

$$
\beta \ge 1 - \left(1 - \max_{i=1,\dots,n} \alpha_i\right) \cdot \sum_{i=1}^n \frac{p_i}{1 - \alpha_i},\tag{3.19}
$$

the involved risk measure [\(3.18\)](#page-9-0) *collapses to the average value-at-risk, it holds that*

$$
\mathcal{R}_{\beta}(Y) = \mathsf{AV}@R_{1-(1-\tilde{\alpha})(1-\beta)}(Y),
$$

where $\tilde{\alpha}$ *is the weighed risk quantile* $\tilde{\alpha} := \frac{\sum_{i=1}^{n} \frac{p_i}{1-\alpha_i} \alpha_i}{\sum_{i=1}^{n} p_i}$ $\frac{\sum_{i=1}^{n} \frac{P_i}{1-\alpha_i}}{\sum_{i=1}^{n} \frac{P_i}{1-\alpha_i}}$.

Proof. The result corresponds to the measure $\mu = \sum_{i=1}^{n} p_i \delta_{\alpha_i}$ in [\(3.14\)](#page-8-1), which is a special case in Corollary [3.6.](#page-8-2)

For $u_{\beta} \ge \alpha_i$, $i = 1, ..., n$, it holds that $\beta = \sum_{i=1}^n p_i \frac{u_{\beta} - \alpha_i}{1 - \alpha_i}$ $\frac{a_{\beta}-\alpha_i}{1-\alpha_i} = \sum_{i=1}^n p_i \frac{1-\alpha_i-(1-u_{\beta})}{1-\alpha_i}$ $\frac{i-(1-u_{\beta})}{1-\alpha_i} = 1 - (1$ u_{β}) $\sum_{i=1}^{n} \frac{p_i}{1-\alpha}$ $\frac{p_i}{1-\alpha_i}$, so that u_β ≥ max_{$i=1,...,n$} α_i is equivalent to [\(3.19\)](#page-9-1). It follows that

$$
u_{\beta} = 1 - \frac{1 - \beta}{\sum_{i=1}^{n} \frac{p_i}{1 - \alpha_i}}
$$

= 1 - (1 - \beta) \left(1 - \frac{\sum_{i=1}^{n} \frac{p_i}{1 - \alpha_i} - \sum_{i=1}^{n} \frac{p_i (1 - \alpha_i)}{1 - \alpha_i}}{\sum_{i=1}^{n} \frac{p_i}{1 - \alpha_i}} \right)
= 1 - (1 - \beta) (1 - \tilde{\alpha}), (3.20)

and $p_0 = \sum \frac{p_i}{1 - \alpha_i}$ $1 - u_\beta$ $\frac{1-a_{\beta}}{1-\beta} = 1$, thus the result with [\(3.18\)](#page-9-0). \Box

Remark 3.8*.* Corollary [3.5](#page-7-4) is a special case of [\(3.18\)](#page-9-0) in the preceding corollary, as $\tilde{\alpha} = \alpha$ in this case.

The following statement generalizes the statements from and provides the higher order risk functional for general risk measures.

Theorem 3.9 (Kusuoka representation of higher order risk measures)**.** *Let* R *be a law invariant risk measure with Kusuoka representation*

$$
\mathcal{R}(Y) = \sup_{\mu \in \mathcal{M}} \mathcal{R}_{\mu}(Y). \tag{3.21}
$$

The higher order risk measure is

$$
\mathcal{R}_{\beta}(Y) = \sup_{\mu \in \mathcal{M}} \mathcal{R}_{\mu_{\beta}}(Y),
$$

where the truncated measures μ ^{β *} <i>are given in* [\(3.15\)](#page-8-3)*.*</sup>

Proof. For the risk functional defined in (3.21) it follows from the min-max inequality that

$$
\mathcal{R}_{\beta}(Y) = \inf_{t \in \mathbb{R}} t + \sup_{\mu \in \mathcal{M}} \mathcal{R}_{\mu}((Y - t)_{+})
$$
\n
$$
\geq \sup_{\mu \in \mathcal{M}} \inf_{t \in \mathbb{R}} t + \frac{1}{1 - \beta} \mathcal{R}_{\mu}((Y - t)_{+})
$$
\n
$$
= \sup_{\mu \in \mathcal{M}} (\mathcal{R}_{\mu})_{\beta}(Y) =
$$
\n
$$
= \sup_{\mu \in \mathcal{M}} \mathcal{R}_{\mu_{\beta}}(Y),
$$
\n(3.22)

where we have used Corollary [3.6.](#page-8-2)

For the reverse inequality in [\(3.22\)](#page-10-1) consider the function

$$
(t,\mu) \mapsto t + \mathcal{R}_{\mu}((Y-t)_{+})
$$

on $\mathbb{R}\times\mathcal{M}([0,1])$, where $\mathcal{M}([0,1])$ collects the probability measures on [0, 1] (with its Borel σ -algebra). By its definition [\(3.14\)](#page-8-1), this function is linear in μ , and convex in t, where $t \in \mathbb{R}$ and μ is a measure on [0, 1]. By Prokhorov's theorem, the set $\mathcal{M}([0, 1])$ of probability measures is sequentially compact, as $[0, 1]$ is compact. From Sion's minimax theorem (cf. Sion $[35]$) it follows that equality holds in (3.22) . Thus, the result.

The preceding Theorem [3.9](#page-9-3) provides an explicit characterization for the general higher order risk measure. The following section exploits this representation to characterize general stochastic dominance relations.

4 General stochastic dominance relations

As Section [2](#page-1-0) mentions, the risk measure R defines a norm via the setting $\|\cdot\| := \mathcal{R}(|\cdot|)$ (cf. [\(2.5\)](#page-3-2)), and conversely, the norm $\|\cdot\|$ defines a risk measure via $\mathcal{R}_{\beta}^{\|\cdot\|}$ $\frac{\ln n}{\beta}$, cf. [\(2.3\)](#page-2-0). In what follows we connect a specific stochastic dominance relation with the norm. This stochastic dominance relation can be described by higher order risk measures, developed in the preceding Section [3.](#page-4-0)

We start by defining the stochastic dominance relation based on a monotone norm.

Definition 4.1 (Stochastic dominance). Let $X, Y \in \mathcal{Y}$ be random variables in a Banach space $(\mathcal{Y}, \| \cdot \|)$. The random variable X is *dominated* by Y , denoted

$$
X\leq 0\leq Y,
$$

if

$$
||(t - X)_+|| \ge ||(t - Y)_+|| \text{ for all } t \in \mathbb{R}.
$$
 (4.1)

If the norm is unambiguous from the context, we shall also simply write \leq instead of \leq ^{||.||}.

The cone of random variables triggered by a single variable is convex.

Lemma 4.2 (Convexity of the stochastic dominance cone). *For* $X \in \mathcal{Y}$ *given, the set*

 ${Y: X \leq Y}$

is convex.

Proof. The map $y \mapsto (t - y)_+$ is convex, as follows from reflecting and translating the convex function $x \mapsto x_+$. Suppose that $X \le Y_0$ and $X \le Y_1$. Then it follows for $Y_\lambda := (1 - \lambda)Y_0 + \lambda Y_1$, together with monotonicity of the norm and [\(4.1\)](#page-10-2), that

$$
||(t - Y_{\lambda})_{+}|| \le ||((1 - \lambda)(t - Y_{0}) + \lambda(t - Y_{1}))_{+}||
$$

\n
$$
\le (1 - \lambda) ||(t - Y_{0})_{+}|| + \lambda ||(t - Y_{1})_{+}||
$$

\n
$$
\le (1 - \lambda) ||(t - X)_{+}|| + \lambda ||(t - X)_{+}||
$$

\n
$$
= ||(t - X)_{+}||.
$$

That is, it holds that $X \le Y_\lambda$ and thus the assertion.

4.1 Characterization of stochastic dominance relations

Stochastic dominance relations can be fully characterized by higher order risk measures. The following theorem presents this main result, which integrates the details developed above for these risk functionals and stochastic dominance relations.

Theorem 4.3 (Characterization of stochastic dominance)**.** *The following are equivalent:*

- (i) $X \leq \mathbb{H} \cdot \mathbb{F}$ Y ,
- *(ii)* $\mathcal{R}_{\beta}(-X) \geq \mathcal{R}_{\beta}(-Y)$ *for all* $\beta \in [0, 1)$ *, and*
- *(iii)* $\inf_{Z \in \mathcal{Z}_{\beta}} \mathbb{E} ZX \leq \inf_{Z \in \mathcal{Z}_{\beta}} \mathbb{E} ZY$ *for every* $\beta \in (0, 1)$ *, where*

$$
\mathcal{Z}_{\beta} \coloneqq \left\{ Y \in \mathcal{X}^* \colon ||Z||_* \le \frac{1}{1-\beta}, \ \mathbb{E} |Z = 1, \ Z \ge 0 \right\}
$$

is the positive cone ($Z \ge 0$) *in the dual ball with radius* $\frac{1}{1-\beta}$ ($||Z||_* \le \frac{1}{1-\beta}$), *intersected with the* $simplex$ ($E Z = 1$).

Proof. Suppose that $X \leq \|f\|$ *Y*, then, by definition, $\|(t - X)_+\| \geq \|(t - Y)_+\|$ for every $t \in \mathbb{R}$. It follows that $t + \frac{1}{1-\beta} \|(-X-t)_+\| \geq t + \frac{1}{1-\beta} \|(-Y-t)_+\|$ for all $t \in \mathbb{R}$, and thus assertion [\(ii\)](#page-11-0) after passing to the infimum.

As for the contrary, assume that [\(ii\)](#page-11-0) holds. To demonstrate [\(i\)](#page-11-1) note first that $q \mapsto ||(q - X)_+||$ is convex; indeed, with $q_{\lambda} := (1 - \lambda)q_0 + \lambda q_1$ and $(a + b)_+ \leq a_+ + b_+$ it holds that

$$
(q_{\lambda} - X)_{+} = ((1 - \lambda)(q_0 - X) + \lambda(q_1 - X))_{+} \le (1 - \lambda)(q_0 - X) + \lambda(q_1 - X)_{+}
$$

and thus

$$
||(q_{\lambda} - X)|| \le (1 - \lambda) \cdot ||(q_0 - X)_+|| + \lambda \cdot ||(q_1 - X)_+||
$$

by the triangle inequality of the norm.

For $q \in \mathbb{R}$ fixed, choose

$$
\alpha \in \partial_{\eta} \left\| (\eta - Y)_{+} \right\|_{\eta = q},
$$

that is, the subdifferential (of the convex function $\eta \mapsto ||(\eta - Y)_+||$) evaluated at $\eta = q$, and note that $\alpha \in [0, 1]$. Set $\beta \coloneqq 1 - \alpha$, and observe that

$$
0\in\partial_q\ -q+\frac{1}{1-\beta}\|(q-Y)_+\|
$$

by (2.3) so that

$$
\mathcal{R}_{\beta}(-Y) = -q + \frac{1}{1-\beta} ||(q-Y)_+||.
$$

Employing the definition [\(2.3\)](#page-2-0) again and assumption [\(ii\),](#page-11-0) it follows that

$$
-q + \frac{1}{1-\beta} ||(-X+q)_+|| \ge \mathcal{R}_{\beta}(-X)
$$

\n
$$
\ge \mathcal{R}_{\beta}(-Y)
$$

\n
$$
= -q + \frac{1}{1-\beta} ||(q-Y)_+||,
$$

or equivalently

$$
||(q - X)_+|| \geq ||(q - Y)_+||.
$$

The assertion [\(i\)](#page-11-1) follows, as $q \in \mathbb{R}$ was arbitrary; this establishes equivalence of (i) and [\(ii\).](#page-11-0)

Finally, let $\beta \in (0, 1)$. With [\(ii\)](#page-11-0) and Proposition [2.5](#page-3-1) we have that

$$
\inf_{Z \in \mathcal{Z}_{\beta}} \mathbb{E} \, ZX \leq \inf_{Z \in \mathcal{Z}_{\beta}} \mathbb{E} \, ZY,
$$

where the infimum in both expressions is among $Z \in \mathcal{Z}_{\beta} = \Big\{ Z \in \mathcal{Z} : ||Z||_* \leq \frac{1}{1-\beta} \Big\}$, as the set \mathcal{Z}_{β} collects the constraints in (2.7) . This establishes equivalence between [\(ii\)](#page-11-0) and [\(iii\).](#page-11-2)

Remark 4.4. The quantity $-R(-Y) =: \mathcal{A}(Y)$ arising naturally in Theorem [4.3](#page-11-3) [\(ii\)](#page-11-0) above is often called an *acceptability functional*, cf. Pflug and Römisch [\[29](#page-23-10)].

Corollary 4.5. *Suppose that*

$$
\mathbb{E} \, ZX \leq \mathbb{E} \, ZY \, \text{for all } Z \in \mathcal{Z} \coloneqq \bigcup_{\beta \in (0,1)} \mathcal{Z}_{\beta},\tag{4.2}
$$

then X *is dominated by* Y *,* $X \leq \mathbb{I}^{|X|} Y$ *. Further, the assertion* [\(4.2\)](#page-12-0) *is equivalent to*

$$
\mathcal{R}_{\beta}(X - Y) \le 0 \quad \text{for all } \beta \in (0, 1). \tag{4.3}
$$

Proof. Fix $\beta \in (0, 1)$, then $\inf_{Z \in \mathcal{Z}_{\beta}} \mathbb{E} ZX \leq \inf_{Z \in \mathcal{Z}_{\beta}} \mathbb{E} ZY$ by [\(4.2\)](#page-12-0). With [\(iii\)](#page-11-2) in the preceding Theo-rem [4.3](#page-11-3) it follows that $X \leq Y$.

With [\(2.6\)](#page-3-4), the statement [\(4.3\)](#page-12-1) is equivalent with $\mathbb{E} Z(X-Y) \leq 0$ for $Z \in \mathbb{Z}$ and hence the assertion. \Box

Remark 4.6*.* The assertion [\(4.3\)](#page-12-1), however, is *strictly stronger* than [\(ii\)](#page-11-0) in Theorem [4.3.](#page-11-3) Indeed, it follows with convexity and [\(4.3\)](#page-12-1) that

$$
\mathcal{R}(-Y) \le \mathcal{R}(X - Y) + \mathcal{R}(-X) \le \mathcal{R}(-X),
$$

and hence [\(ii\),](#page-11-0) the assertion, although the reverse implication does not hold true.

4.2 Stochastic dominance in numerical computations

To verify that $X \le Y$ it is necessary to verify the defining condition [\(4.1\)](#page-10-2) for every $t \in \mathbb{R}$. These infinitely many comparisons are intractable for numerical computations. The same holds true for the equivalent characterization [\(ii\)](#page-11-0) in Theorem [4.3,](#page-11-3) as all risk levels $\beta \in (0, 1)$ – again infinitely many – need to be considered. This is difficult, perhaps impossible to ensure in numerical computations.

In what follows we develop an equivalent characterization, which builds on only *finitely many* risk levels. With this, the comparison $X \leq Y$ is numerically tractable.

We start the exposition with the following lemma on convexity (concavity).

Lemma 4.7. *For Y fixed in the domain of* \mathcal{R} *, the mapping*

$$
\beta \mapsto (1 - \beta) \cdot \mathcal{R}_{\beta}(Y)
$$

is concave.

Proof. For $\lambda \in (0, 1)$, define $\beta_{\lambda} := (1 - \lambda)\beta_0 + \lambda \beta_1$. Choose t_{λ} in [\(2.3\)](#page-2-0) minimizing $\mathcal{R}_{\beta_{\lambda}}(Y)$. Then

$$
(1 - \beta_{\lambda})\mathcal{R}_{\beta_{\lambda}}(Y) = (1 - \beta_{\lambda})t_{\lambda} + ||(Y - t_{\lambda})_{+}||
$$

= $(1 - \lambda)((1 - \beta_{0})t_{\lambda} + ||(Y - t_{\lambda})_{+}||) + \lambda((1 - \beta_{1})t_{\lambda} + ||(Y - t_{\lambda})_{+}||)$
 $\geq (1 - \lambda)(1 - \beta_{0})\mathcal{R}_{\beta_{0}}(Y) + \lambda(1 - \beta_{1})\mathcal{R}_{\beta_{1}}(Y)$

by taking the infimum in \mathcal{R}_{β} in the latter expressions.

The latter result on convexity leads to the following result on the derivative of the risk functional with respect to the risk level.

Theorem 4.8. *For* Y in the domain of R and $\beta \in (0, 1)$, the derivative with respect to the risk rate is

$$
\frac{\mathrm{d}}{\mathrm{d}\beta}\mathcal{R}_{\beta}(Y) = \frac{\mathcal{R}_{\beta}(Y) - t_Y(\beta)}{1 - \beta},
$$

where $t_Y(\beta)$ *minimizes the higher order risk measure* $\mathcal{R}_{\beta}(Y)$ *, cf.* [\(2.3\)](#page-2-0)*.*

Proof. As in Lemma [4.7](#page-13-0) above consider the objective

$$
f(\beta, t) := (1 - \beta)t + ||(Y - t)_+||.
$$

For β fixed, the objective is concave, and we may choose $t(\beta)$ in the subgradient, $t(\beta) \in \partial_t f(\beta, t)$. Define the function

$$
f(\beta) := f(\beta, t(\beta)) = \min_{t \in \mathbb{R}} f(\beta, t(\beta)) = (1 - \beta) \mathcal{R}_{\beta}(Y).
$$

It holds that

$$
f'(\beta) = \frac{\partial}{\partial \beta} f(\beta, t(\beta)) + \frac{\partial}{\partial t} f(\beta, t(\beta)) \cdot t'(\beta).
$$

But $0 \in \frac{\partial}{\partial t} f(\beta, t(\beta))$, as $t(\beta)$ is optimal in [\(2.3\)](#page-2-0). With $\frac{\partial}{\partial \beta} f(\beta, t) = -t$, it follows that

$$
f'(\beta) = -t(\beta). \tag{4.4}
$$

It follows that

It follows that

\n
$$
\frac{d}{d\beta}\mathcal{R}_{\beta}(Y) = \frac{d}{d\beta}\frac{f(\beta)}{1-\beta} = \frac{-t(\beta)(1-\beta) + f(\beta)}{(1-\beta)^2} = \frac{\mathcal{R}_{\beta}(Y) - t(\beta)}{1-\beta},
$$
\nthe assertion.

Theorem 4.9 (Verification of stochastic dominance relations). To verify that $X \leq Y$ it is sufficient to verify

$$
\mathcal{R}_{\beta_i}(-X) \ge \mathcal{R}_{\beta_i}(-Y) \tag{4.5}
$$

*for the (*finitely many*) risk levels*

 $\beta_i, \quad i = 1, \ldots, n;$

the risk levels β_i *with* $\beta_i < \gamma_i < \beta_{i+1}$ *are chosen so that*

$$
t_{-X}(\beta) \le t_{-Y}(\beta) \text{ for } \beta \in (\beta_i, \gamma_i) \text{ and } \tag{4.6}
$$

$$
t_{-X}(\beta) \ge t_{-Y}(\beta) \text{ for } \beta \in (\gamma_i, \beta_{i+1})
$$
\n
$$
(4.7)
$$

for $i = 1, 2, ..., n$.

Remark 4.10. For the crucial risk levels (test points) β_i in the preceding Theorem [4.9](#page-13-1) it holds that

$$
t_{-X}(\beta) \ge t_{-Y}(\beta)
$$
 for $\beta \in (\gamma_{i-1}, \beta_i)$ and $t_{-X}(\beta) \le t_{-Y}(\beta)$ for $\beta \in (\beta_i, \gamma_i)$,

so that the curves $t_{-X}(\cdot)$ and $t_{-Y}(\cdot)$ intersect at β_i . For continuous $t_{-X}(\cdot)$ and $t_{-Y}(\cdot)$, the crucial points in the preceding Theorem [4.9](#page-13-1) are among the points, where the minimizers of $\mathcal{R}_{\beta}(-X)$ and $\mathcal{R}_{\beta}(-Y)$ coincide, that is,

$$
t_{-X}(\beta_i)=t_{-Y}(\beta_i), \quad i=1,\ldots,n;
$$

with a similar argument it holds as well that

$$
t_{-X}(\gamma_i)=t_{-Y}(\gamma_i), \quad i=1,\ldots,n.
$$

Proof of Theorem [4.9.](#page-13-1) Recall first from Theorem [4.3](#page-11-3) [\(ii\)](#page-11-0) that $X \leq Y$ is equivalent to

$$
\mathcal{R}_{\beta}(-X) \ge \mathcal{R}_{\beta}(-Y) \text{ for all } \beta \in (0,1).
$$

Assuming [\(4.5\)](#page-13-2), we demonstrate that $\mathcal{R}_{\beta}(-X) \geq \mathcal{R}_{\beta}(-Y)$ for $\beta \in (\beta_i, \beta_{i+1})$ for $i = 1, ..., n$. To this end we distinguish the following two cases, where we reuse the notation introduced in the proof of Theorem [4.8:](#page-13-3)

(i) Suppose that $\beta \in (\beta_i, \gamma_i)$, then, by [\(4.6\)](#page-13-4), $t_{-X}(\beta) \le t_{-Y}(\beta)$. With [\(4.4\)](#page-13-5) we conclude that

$$
f_{-X}(\beta) = f_{-X}(\beta_i) + \int_{\beta_i}^{\beta} f'_{-X}(\gamma) d\gamma
$$

$$
= f_{-X}(\beta_i) - \int_{\beta_i}^{\beta} t_{-X}(\gamma) d\gamma
$$

$$
\geq f_{-Y}(\beta_i) - \int_{\beta_i}^{\beta} t_{-Y}(\gamma) d\gamma
$$

$$
= f_{-Y}(\beta_i) + \int_{\beta_i}^{\beta} f'_{-Y}(\gamma) d\gamma
$$

$$
= f_{-Y}(\beta_i),
$$
 (4.8)

where we have used [\(4.5\)](#page-13-2) in [\(4.8\)](#page-14-0). It follows that $\mathcal{R}_{\beta}(-X) \geq \mathcal{R}_{\beta}(-Y)$ for all $\beta \in (\beta_i, \gamma_i)$.

(ii) If $\beta \in (\gamma_i, \beta_{i+1})$, then $t_{-X}(\beta) \ge t_{-Y}(\beta)$ by assumption [\(4.7\)](#page-13-6). It holds that

$$
f_{-X}(\beta) = f_{-X}(\beta_{i+1}) - \int_{\beta}^{\beta_{i+1}} f'_{-X}(\gamma) d\gamma
$$

= $f_{-X}(\beta_{i+1}) + \int_{\beta}^{\beta_{i+1}} t_{-X}(\gamma) d\gamma$

$$
\geq f_{-Y}(\beta_{i+1}) + \int_{\beta}^{\beta_{i+1}} t_{-Y}(\gamma) d\gamma
$$

= $f_{-Y}(\beta_{i+1}) - \int_{\beta}^{\beta_{i+1}} f'_{-Y}(\gamma) d\gamma$
= $f_{-Y}(\beta),$ (4.9)

where again [\(4.5\)](#page-13-2) was used in [\(4.9\)](#page-14-1). It follows that $\mathcal{R}_{\beta}(-X) \geq \mathcal{R}_{\beta}(-Y)$ for $\beta \in (\gamma_i, \beta_{i+1})$, the remaining case.

Combining the two cases above we find that $\mathcal{R}_{\beta}(-X) \geq \mathcal{R}_{\beta}(-Y)$ for all $\beta \in (\beta_i, \beta_{i+1})$. The assertion for all $\beta \in \mathbb{R}$ thus follows by considering $i = 1, \dots, n$.

Remark 4.11. It is important to note that the critical risk levels β_i , $i = 1, \ldots, n$, in [\(4.5\)](#page-13-2) – in general – depend on *both* random variables, on *X and Y*. The folloiwing remark presents a notable exception to this rule.

Remark 4.12 (Average value-at-risk). For the Hölder norm $\|\cdot\|_1$ with $p = 1$, the optimizers are $t_{-X}(\beta) =$ V@R_{β}(-X), and the function $t_{-X}(\cdot)$ is not continuous for discrete distributions. As $\beta \mapsto t_{-X}(\beta)$ is non-decreasing, the test points are

$$
\beta_i=-x_i, i=1,2,\ldots,n,
$$

where $P(X = x_i) > 0$. In this case, the crucial risk levels β_i , $i = 1, ..., n$, are independent from the variable Y when comparing $X \leq Y$.

4.3 Characterization of stochastic dominance for spectral risk measures

The following builds on the spectral risk measure $\mathcal{R}_{\sigma}(\cdot)$ introduced in Definition [3.1](#page-4-2) and considers the norm

$$
\|\cdot\|_{\sigma} \coloneqq \mathcal{R}_{\sigma}(|\cdot|)
$$

for the spectral function σ . Theorem [4.3](#page-11-3) and the characterization of higher order spectral risk measures (Theorem [3.2\)](#page-5-4) give rise to the following result.

Theorem 4.13. *The stochastic dominance relation*

$$
X\leq \parallel\cdot\parallel_{\sigma} Y
$$

with respect to the norm associated with the spectral risk measure R_{σ} *is equivalent to*

$$
-\sigma_p \mathsf{V@R}_p(Y) + \int_{-\infty}^{\mathsf{V@R}_p(Y)} \Sigma(S_Y(y)) \, dy
$$

$$
\leq -\sigma_p \mathsf{V@R}_p(X) + \int_{-\infty}^{\mathsf{V@R}_p(X)} \Sigma(S_X(x)) \, dx \quad \text{for all } p \in (0,1),
$$

where $\sigma_p := \int_{1-p}^1 \sigma(u) du$ and $S_X(x) := 1 - F_X(x) = P(X > x)$ *is the survival function of the random variable X.*

Proof. We argue with the norm $||Y||_{\sigma} := \mathcal{R}_{\sigma}(|Y|)$. Note, that $(Y-t)_+ \geq 0$, hence the defining equation [\(2.3\)](#page-2-0) is

$$
\mathcal{R}_{\beta}^{\|\cdot\|_{\sigma}}(Y) = \inf_{t \in \mathbb{R}} t + \frac{1}{1-\beta} \|(Y-t)_{+}\|_{\sigma}
$$

$$
= \inf_{t \in \mathbb{R}} t + \frac{1}{1-\beta} \mathcal{R}_{\sigma}((Y-t)_{+})
$$

$$
= \mathcal{R}_{\sigma_{\beta}}(Y), \tag{4.10}
$$

where we have used Theorem [3.2](#page-5-4) in [\(4.10\)](#page-15-0).

From [\(3.8\)](#page-6-2) we have that

$$
\mathcal{R}_{\beta}(-Y) = \mathsf{V} \circledast \mathsf{R}_{u_{\beta}}(-Y) + \frac{1}{1-\beta} \int_{\mathsf{V} \circledast \mathsf{R}_{u_{\beta}}(-Y)}^{\infty} \Sigma(F_{-Y}(y)) \, \mathrm{d}y
$$
\n
$$
= -\mathsf{V} \circledast \mathsf{R}_{1-u_{\beta}}(Y) + \frac{1}{1-\beta} \int_{-\mathsf{V} \circledast \mathsf{R}_{1-u_{\beta}}(Y)}^{\infty} \Sigma(S_Y(-y)) \, \mathrm{d}y
$$
\n
$$
= -\mathsf{V} \circledast \mathsf{R}_{1-u_{\beta}}(Y) + \frac{1}{1-\beta} \int_{-\infty}^{\mathsf{V} \circledast \mathsf{R}_{1-u_{\beta}}(Y)} \Sigma(S_Y(y)) \, \mathrm{d}y,
$$

where we have used that $F_{-Y}(y) = P(-Y \le y) = P(Y \ge -y) = 1 - F_Y(-y) = S_Y(-y)$ and $V \otimes R_\alpha(-Y) =$ $-V@R_{1-\alpha}(Y)$ at points of continuity of $F_Y(\cdot)$.

Now set $1 - u_\beta = p$. Then, by employing the characterizing relation [\(3.3\)](#page-5-0) for the β -quantile of σ , it holds that

$$
1 - \beta = \int_{u_{\beta}}^{1} \sigma(u) du = \int_{1-p}^{1} \sigma(u) du = \sigma_p,
$$

so that

$$
\mathcal{R}_{\beta}(-Y) = -\mathsf{V} \circledast \mathsf{R}_{p}(Y) + \frac{1}{\sigma_{p}} \int_{-\infty}^{\mathsf{V} \circledast \mathsf{R}_{p}(Y)} \Sigma(S_{Y}(y)) \, dy.
$$

By Theorem [4.3,](#page-11-3) the relation $X \leq \| \cdot \|_{\sigma} Y$ is equivalent to $\mathcal{R}_{\beta}^{\|\cdot\|_{\sigma}}(-Y) \leq \mathcal{R}_{\beta}^{\|\cdot\|_{\sigma}}(-X)$ for all $\beta \in (0, 1)$. With that, the assertion follows. \Box

4.4 Higher order stochastic dominance

A traditional way of introducing stochastic dominance relations is by iterating integrals of the cumulative distribution function. This is a special case for the Hölder norm $\|\cdot\|_p$, $p \in [1,\infty)$.

Definition 4.14 (Higher order stochastic dominance, cf. Müller and Stoyan [\[23](#page-23-11)]). The random variable X is dominated by Y in *first order stochastic dominance*, if

$$
F_X(x) \geq F_Y(x) \text{ for all } x \in \mathbb{R},
$$

where $F_X(x) := P(X \le x)$ is the cumulative distribution function. We shall write $X \le (1)$ Y. For $p \in [1, \infty]$, the random variable X is stochastically dominated by Y in p^{th} -stochastic order, if

$$
\mathbb{E}(x - X)_{+}^{p-1} \ge \mathbb{E}(x - Y)_{+}^{p-1} \text{ for all } x \in \mathbb{R};
$$
\n(4.11)

we write $X \leqslant^{(p)} Y$.

Lemma 4.15 (Cf. Ogryczak and Ruszczyński [\[25](#page-23-12), [26](#page-23-13)]). *With* $F_X^{(1)}$ $J_X^{(1)}(\cdot) := F_X(\cdot)$, the k^{th} $(k = 2, 3, ...)$ repeated integral is $F_{\mathbf{x}}^{(k)}$ $J_X^{(k)}(x) := \int_{-\infty}^x F_X^{(k-1)}$ $\chi^{(k-1)}(y)$ dy. The following two points are equivalent, they characterize *stochastic dominance of* integer *orders by repeated integrals:*

- (i) $X \leqslant^{(k)} Y$,
- (iii) $F_Y^{(k)}$ $F_Y^{(k)}(x) \ge F_X^{(k)}$ $J_X^{(k)}(x)$ *for all* $x \in \mathbb{R}$ *.*

Proof. It holds with [Cauchy's formula for repeated integration](https://en.wikipedia.org/wiki/Cauchy_formula_for_repeated_integration) that

$$
F_X^{(k)}(x) = \frac{1}{(k-2)!} \int_{-\infty}^x (x-y)^{k-2} F_X(y) \, dy.
$$

By integration by parts, the latter is

$$
F_X^{(k)}(x) = \frac{1}{(k-1)!} \int_{-\infty}^x (x-y)^{k-1} dF_X(y),
$$

so that

$$
F_X^{(k)}(x) = \frac{1}{(k-1)!} \int_{-\infty}^{\infty} (x - y)_+^{k-1} dF_X(y) = \frac{1}{(k-1)!} \mathbb{E}(x - X)_+^{k-1},
$$

from which the assertion follows from the defining condition [\(4.1\)](#page-10-2) in Definition [4.1.](#page-10-3) \Box

Remark 4.16. It follows from the iterated integral and [\(ii\)](#page-16-0) in Lemma [4.15](#page-16-1) that $X \leq (k) Y \implies X \leq (k+1) Y$ for all *natural* numbers $k = 1, 2, \ldots$. We notice next that

$$
X \preccurlyeq^{(p)} Y \implies X \preccurlyeq^{(p')} Y \text{ for all real numbers } 1 \le p \le p' \in \mathbb{R}.\tag{4.12}
$$

To this end note first that the characterization [\(4.11\)](#page-16-2) is equivalent to

$$
\int_{-\infty}^{x} (x-z)^{p-1} dF_X(z) \ge \int_{-\infty}^{x} (x-z)^{p-1} dF_Y(z) \text{ for all } x \in \mathbb{R}.
$$
 (4.13)

With $\int_{z}^{x} (x - y)^{\alpha-1} (y - z)^{\beta-1} dy = B(\alpha, \beta) (x - z)^{\beta+\alpha-1}$ (*B* is Euler's integral of the first kind) and integration by parts it follows that

$$
\int_{-\infty}^{x} (x - z)^{p'-1} dF_X(z) = \frac{1}{B(p, p' - p)} \int_{-\infty}^{x} \int_{z}^{x} (x - y)^{p'-p-1} (y - z)^{p-1} dy dF_X(z)
$$

\n
$$
= \frac{1}{B(p, p' - p)} \int_{-\infty}^{x} (x - y)^{p'-1-p} \int_{-\infty}^{y} (y - z)^{p-1} dF_X(z) dy
$$

\n
$$
\geq \frac{1}{B(p, p' - p)} \int_{-\infty}^{x} (x - y)^{p'-1-p} \int_{-\infty}^{x} (y - z)^{p-1} dF_Y(z) dy
$$
(4.14)
\n
$$
= \int_{-\infty}^{x} (x - z)^{p'-1} dF_Y(z),
$$

where we have used the characterization [\(4.13\)](#page-17-0) in [\(4.14\)](#page-17-1), as $x - y \ge 0$ and that $B(p, p' - p)$ is well-defined and positive for $p' > p$. The assertion again follows with [\(4.13\)](#page-17-0).

4.5 Comparison of stochastic order relations

Different stochastic dominance relations may vary in strength (the implication [\(4.12\)](#page-17-2) in the preceding Remark [4.16](#page-17-3) is an example). In what follows, we provide an explicit relation to compare stochastic dominance relations, which are built on different spectral functions.

Proposition 4.17 (Comparison of spectral stochastic orders)**.** *Suppose that*

$$
\sigma_{\mu}(u) = \sigma(u) \cdot \int_0^{u_{\beta}} \frac{\mu(\mathrm{d}\beta)}{1-\beta} \tag{4.15}
$$

for some probability measure μ, where u_β is as defined in [\(3.3\)](#page-5-0). Then the stochastic order associated *with* σ_{μ} *is weaker than the genuine stochastic order associated with* σ *. Specifically, for different spectral functions* σ *and* σ_{μ} *, it holds that*

$$
X \leq^{\|\cdot\|_{\sigma}} Y \implies X \leq^{\|\cdot\|_{\sigma_{\mu}}} Y.
$$

Remark 4.18. The function σ_{μ} in [\(4.15\)](#page-17-4) is indeed a spectral function. It is positive, as μ is a positive measure (thus [\(i\)](#page-4-4) in Definition [3.1\)](#page-4-2). The function is non-decreasing, as u_{β} is non-decreasing for β increasing. Finally, the function σ_{μ} is a density: indeed, it holds that

$$
\int_0^1 \sigma_\mu(u) \, \mathrm{d}u = \int_0^1 \sigma(u) \cdot \int_0^{u_\beta} \frac{\mu(\mathrm{d}\beta)}{1-\beta} \mathrm{d}u = \int_0^1 \int_{\beta_u}^1 \sigma(u) \, \mathrm{d}u \, \frac{\mu(\mathrm{d}\beta)}{1-\beta} = \int_0^1 \mu(\mathrm{d}\beta) = 1
$$

by integration by parts, where we have used the definition of u_{β} in [\(3.3\)](#page-5-0).

Proof of Proposition [4.17.](#page-17-5) Since $x \le ||\cdot||_{\sigma}$ *Y*, it holds with Theorem [4.3](#page-11-3) that $\mathcal{R}_{\sigma_{\beta}}(-X) \ge \mathcal{R}_{\sigma_{\beta}}(-Y)$ for all $\beta \in (0, 1)$, where σ_{β} is defined in [\(3.2\)](#page-5-5). By the characterization [\(3.1\)](#page-5-1), this is

$$
\int_{u_{\beta}}^{1} \frac{\sigma(u)}{1 - \beta} F_{-X}^{-1}(u) du \ge \int_{u_{\beta}}^{1} \frac{\sigma(u)}{1 - \beta} F_{-Y}^{-1}(u) du, \qquad \beta \in (0, 1).
$$

Integrating the latter expression with respect to $\mu(d\beta)$ establishes the inequality

$$
\int_{\beta}^{1} \int_{u_{\beta'}}^{1} \frac{\sigma(u)}{1 - \beta'} F_{-X}^{-1}(u) du \, \mu(d\beta') \ge \int_{\beta}^{1} \int_{u_{\beta'}}^{1} \frac{\sigma(u)}{1 - \beta'} F_{-Y}^{-1}(u) du \, \mu(d\beta'), \qquad \beta \in (0, 1).
$$

Interchanging the order of integration together with [\(3.17\)](#page-9-4) gives that

$$
\int_{u_{\beta}}^{1} \int_{\beta}^{\beta_u} \frac{\sigma(u)}{1 - \beta'} \mu(d\beta') F_{-X}^{-1}(u) du \ge \int_{u_{\beta}}^{1} \int_{\beta}^{\beta_u} \frac{\sigma(u)}{1 - \beta'} \mu(d\beta) F_{-Y}^{-1}(u) du, \qquad \beta \in (0, 1),
$$

which in turn is

$$
\int_{u_{\beta}}^{1} \sigma_{\mu}(u) F_{-X}^{-1}(u) du \ge \int_{u_{\beta}}^{1} \sigma_{\mu}(u) F_{-Y}^{-1}(u) du, \qquad \beta \in (0, 1).
$$

This is the assertion.

5 Example: the expectile

The expectile risk measure, originally introduced by Newey and Powell [\[24\]](#page-23-14), has recently gained additional interest (cf. Malandii et al. [\[22\]](#page-23-15), Balbás et al. [\[2\]](#page-21-2) or Farooq and Steinwart [\[12](#page-22-14)] for conditional regressions). A main reason for the additional interest in this risk measure is because it is the only elicitable risk functional (cf. Ziegel [\[37](#page-24-1)]).

As Proposition [2.5](#page-3-1) indicates, the higher order risk measure can be based on the dual norm. For this reason, the following section establishes the dual norm of expectiles first, as it is crucial in understanding its regret function in the risk quadrangle. Next, we provide an explicit characterization of the higher order expectiles, that is, the higher order risk measure based on the expectile risk measure.

The expectile is defined as a *minimizer*. Its Kusuoka representation is central in elaborating the corresponding higher order risk functional.

Definition 5.1. For $\alpha \in (0, 1)$, the expectile is

$$
e_{\alpha}(Y) = \underset{x \in \mathbb{R}}{\arg \min} \mathbb{E} \, \ell_{\alpha}(Y - x),\tag{5.1}
$$

where

$$
\ell_{\alpha}(x) = \begin{cases} \alpha x^2 & \text{if } x \ge 0, \\ (1 - \alpha)x^2 & \text{else} \end{cases}
$$

is the asymmetric loss, or quadratic error function.

The expectile satisfies the first order condition

$$
(1 - \alpha) \mathbb{E}(x - Y)_+ = \alpha \mathbb{E}(Y - x)_+, \tag{5.2}
$$

and $e_{\alpha}(\cdot)$ is a risk measure for $\alpha \in [1/2, 1]$. We mention that condition [\(5.2\)](#page-19-0) provides a definition for $Y \in L^1$, it is thus more general than [\(5.1\)](#page-18-1), which requires $Y \in L^2$. The Kusuoka representation of the expectile (cf. Bellini et al. [\[4](#page-22-15), Proposition 9]) is given by

$$
e_{\alpha}(Y) = \max_{\gamma \in [0,1-\eta]} (1-\gamma) \cdot \mathbb{E}[Y+\gamma \cdot \mathsf{AV@R}_{1-\frac{\gamma}{1-\gamma}\frac{\eta}{1-\eta}}(Y), \tag{5.3}
$$

where $\eta = \frac{1-\alpha}{\alpha}$, so that the risk level in [\(5.3\)](#page-19-1) is $1 - \frac{\gamma}{1-\alpha}$ $1 - \gamma$ η $\frac{\eta}{1-\eta} = \frac{\alpha(2-\gamma)-1}{(2\alpha-1)(1-\gamma)}$ $\frac{\alpha(2-\gamma)-1}{(2\alpha-1)(1-\gamma)}$. Involving spectral risk measures, the expectile can be recast as

$$
e_{\alpha}(Y) = \sup \{ \mathcal{R}_{\sigma_{\gamma}}(Y) : \sigma_{\gamma} \in \mathcal{S} \},
$$

where $S = {\sigma_\gamma : \gamma \in [0, 1 - \eta]}$ collects the spectral functions

$$
s_{\gamma}(u) = \begin{cases} 1 - \gamma & \text{if } u \leq 1 - \frac{\gamma}{1 - \gamma} \frac{\eta}{1 - \eta}, \\ \frac{1 - \gamma}{\eta} & \text{else.} \end{cases}
$$

The higher order expectile can be described by involving its dual norm (cf. (2.8)), as well as its Kusuoka representation (cf. Corollary [3.6\)](#page-8-2). The following two (sub)sections elaborate these possibilities for the expectile.

5.1 The dual norm of expectiles

The higher order expectile can be described with the dual representation [\(2.7\)](#page-3-3), for which the dual norm of the expectile is necessary.

By the characterization of the loss function [\(5.2\)](#page-19-0) it holds that $e_{\alpha}(Y)$ is well-defined for $Y \in L^1(P)$. This is enough to conclude that $\mathbb{E}|Y| \leq C_{\alpha} \cdot e_{\alpha}(|Y|)$ for some constant $C_{\alpha} > 0$ (Lakshmanan and Pichler [\[19,](#page-23-16) Corollary 2.16] elaborate the tight bound $C_{\alpha} = \frac{\alpha}{1-\alpha}$). It follows that $\mathcal{Y}^* = L^{\infty}$, so that $||Z||_{\infty}$ is well-defined for $Z \in \mathcal{Y}^*$.

The following result provides the dual norm of the expectile explicitly.

Proposition 5.2 (Dual norm of the expectile). *For* $\alpha \ge 1/2$ *, the dual norm is*

$$
||Z||_{\alpha}^* := \sup \{ \mathbb{E} YZ : e_{\alpha}(|Y|) \le 1 \}
$$
\n
$$
(5.4)
$$

(cf. [\(2.6\)](#page-3-4)*) . It holds that*

$$
||Z||_{\alpha}^* = \sup_{\beta \in (0,1)} (1 - \beta) \cdot \text{AV@R}_{\beta}(|Z|) + \beta \frac{1 - \alpha}{\alpha} ||Z||_{\infty}.
$$
 (5.5)

Notably, the norm $\|\cdot\|_{\alpha}^{*}$ is *not* a risk measure itself, and [\(5.5\)](#page-19-2) is *not* a Kusuoka representation; indeed, the total weight in the representation (5.5) is

$$
(1 - \beta) + \beta \frac{1 - \alpha}{\alpha} < 1
$$

for $\alpha \in (1/2, 1]$.

Proof of Proposition [5.2.](#page-19-3) We may assume that $Z \ge 0$, as otherwise we may consider sign(Z) · Y instead of Y. For arbitrary sets B and G with $B \subset G$ and $P(G) < 1$ define the random variable

$$
\tilde{Y}_{B,G}(\omega) := \begin{cases}\n0 & \text{if } \omega \in B, \\
1 & \text{if } \omega \in G \setminus B, \text{ and} \\
\frac{1-\alpha}{\alpha} \cdot \frac{P(B)}{1-P(G)} + 1 & \text{else.} \n\end{cases}
$$
\n(5.6)

,

Note, that

$$
(1 - \alpha) \cdot P(B)(1 - 0) = \alpha \cdot (1 - P(G)) \left(\frac{(1 - \alpha)P(B)}{\alpha(1 - P(G))} + 1 - 1 \right)
$$

and hence $e_{\alpha}(\tilde{Y}_{B,G}) = 1$ by the defining equation [\(5.2\)](#page-19-0). It follows with [\(5.4\)](#page-19-4) that

$$
||Z||_{\alpha}^* \geq \mathbb{E} Z Y_{B,G}.
$$

As $B \subset G$ are arbitrary, we conclude in particular that

$$
||Z||_{\alpha}^* \ge \left((1 - P(B)) \cdot \mathsf{AV@R}_{P(B)}(Z) + P(B) \frac{1 - \alpha}{\alpha} \cdot \mathsf{AV@R}_{P(G)}(Z), \right)
$$

because the random variables

$$
\tilde{Y}_{B,G} = \left(1 - P(B)\right) \cdot \frac{1}{1 - P(B)} 1\!\!1_{[P(B),1]}(U) + P(B) \frac{1 - \alpha}{\alpha} \cdot \frac{1}{1 - P(G)} 1\!\!1_{[P(G),1]}(U)
$$

satisfy all conditions from above for any uniform variable U. Now let $P(G) \rightarrow 1$ and by denoting $\beta = P(B)$ it follows that

$$
||Z||_{\alpha}^* \ge \sup_{\beta \in (0,1)} (1 - \beta) \cdot \text{AV@R}_{\beta}(Z) + \beta \frac{1 - \alpha}{\alpha} \operatorname{ess} \sup Z,
$$

as $\mathsf{AV@R}_{\gamma}(Z) \to \mathsf{ess\,sup}\, Z$ for $\gamma \to 1$.

As for the converse observe that we may assume $e_{\alpha}(Y) = 1$ for the optimal random variable in [\(5.4\)](#page-19-4). Consider the Lagrangian

$$
L(Y; \lambda, \mu) := \mathbb{E} ZY - \lambda ((1 - \alpha) \mathbb{E} (1 - Y)_{+} - \alpha \mathbb{E} (Y - 1)_{+}) + \mathbb{E} \mu Y, \tag{5.7}
$$

where the Lagrangian multiplier $\lambda \in \mathbb{R}$ is associated with the equality constraint $e_{\alpha}(Y) = 1$, i.e., [\(5.2\)](#page-19-0), and the measurable variable $\mu \in L^1$, $\mu \ge 0$, is associated with the inequality constraint $Y \ge 0$. Provided That the derivative exists, the first order conditions are

or

$$
0 = \frac{\partial}{\partial Y} L(Y; \lambda, \mu),
$$

\n
$$
Z = \lambda \left(-(1 - \alpha) \mathbb{1}_{\{Y < 1\}} - \alpha \mathbb{1}_{\{Y > 1\}} \right) - \mu \cdot \mathbb{1}_{\{Y = 0\}}.
$$
\n(5.8)

Now note that the left-hand side of [\(5.8\)](#page-20-0) involves the *variable* Z, while the right-hand side only involves *constants*, except on $\{Y = 0\}$, where μ is not necessarily constant. The first order conditions [\(5.8\)](#page-20-0) thus hold true on plateaus of Z, if they coincide with $\{Y < 1\}$ or $\{Y > 1\}$; for $\{Y = 0\}$, equation [\(5.8\)](#page-20-0) is $\mu = -Z - \lambda(1 - \alpha)$; for $\{Y = 1\}$, the derivative of [\(5.7\)](#page-20-1) does not exist or depends on the direction.

It follows, that the optimal Y in [\(5.4\)](#page-19-4) exactly is of form [\(5.6\)](#page-20-2) and hence the assertion. \square

5.2 Higher order expectiles

The Kusuoka representation [\(5.3\)](#page-19-1) is the basis for the expectile's higher order risk measure.

Proposition 5.3. *For* $\beta \in (0, 1)$ *, the higher order expectile is*

$$
(e_{\alpha})_{\beta}(Y) = \max_{\gamma \in [0,1-\eta]} \begin{cases} \left(1 - \frac{\gamma}{1-\beta}\right) \text{AV@R}_{\frac{\beta}{1-\gamma}}(Y) + \frac{\gamma}{1-\beta} \text{AV@R}_{1-\frac{\gamma}{1-\gamma}\frac{\eta}{1-\eta}}(Y) & \text{if } \frac{\gamma}{1-\beta} < 1-\eta, \\ \text{AV@R}_{1-(1-\beta)(1-\tilde{\alpha})}(Y) & \text{else,} \end{cases} \tag{5.9}
$$

where $\eta = \frac{1-\alpha}{\alpha}$ (as above) and $\tilde{\alpha} \coloneqq 1 - \frac{\eta}{1-\alpha}$ $\frac{\eta}{1-\gamma}$.

Proof. The measure in the Kusuoka representation [\(5.3\)](#page-19-1) is $\mu(\cdot) = (1 - \gamma)\delta_0 + \gamma \cdot \delta_{1-\frac{\gamma}{1-\gamma}}$. To apply Corollary [3.7](#page-9-5) we set $p_1 := 1 - \gamma$ and $p_2 = \gamma$, the corresponding risk levels are $\alpha_1 = 0$ and $\alpha_2 = 1 - \frac{\gamma}{1 - \gamma}$ $1 - \gamma$ η $\frac{\eta}{1-\eta}$. The mixed risk level is $\tilde{\alpha} := \frac{\alpha_1 \frac{p_1}{1 - \alpha_1} + \alpha_2 \frac{p_2}{1 - \alpha_2}}{\frac{p_1}{1 - \alpha_1} + \frac{p_2}{1 - \alpha_2}}$ $=\frac{\alpha(2-\gamma)-1}{\gamma}$ $\frac{(2-\gamma)-1}{\alpha(1-\gamma)} = 1 - \frac{\eta}{1-\gamma}$ $\frac{\eta}{1-\gamma}$.

We distinguish the cases $\frac{\gamma}{1-\beta} < 1-\eta$ and $\frac{\gamma}{1-\beta} < 1-\eta$, which are equivalent to $u_{\beta} \le \alpha_2$, i.e., $1 - \frac{1-\beta}{\frac{\gamma}{1-\alpha_1} + \frac{1-\gamma}{1-\alpha_2}}$ $\leq \alpha_2$ in view of [\(3.20\)](#page-9-6). In the first case, the critical equation [\(3.16\)](#page-8-0) is $(1 - \gamma)u_\beta = \beta$, while it is $(1 - \gamma)u_{\beta} + \gamma \frac{u_{\beta} - \alpha_2}{1 - \alpha_2}$ $\frac{1}{1-\alpha_2} = \beta$ in the other case; the solutions thus are $u_{\beta} = \frac{\beta}{1-\beta_2}$ $\frac{\beta}{1-\gamma}$ and $u_{\beta} = \frac{\alpha(2-\beta-\gamma)-1+\beta}{\alpha(1-\gamma)}$ $\frac{-\rho-\gamma-1+\rho}{\alpha(1-\gamma)}$. The corresponding weights p_0 (cf. [\(3.16\)](#page-8-0) again) are $p_0 = \frac{1 - \mu_B}{1 - \beta}$ $\frac{1-u_{\beta}}{1-\beta}(1-\gamma)$, or $p_0 = \frac{1-u_{\beta}}{1-\beta}$ $1-\beta$ $\sqrt{1-\gamma}$ $\frac{1-\gamma}{1-0}$ + $\frac{\gamma}{1-\alpha}$ $1-\alpha_2$ $= 1.$ Finally, note that $u_{\beta} = 1 - (1 - \beta)(1 - \tilde{\alpha})$. The assertion follows with (3.17) and (3.18) in Corollary [3.7.](#page-9-5)

The average value-at-risk is 'closed under higher orders', as its higher order variant is an average value-at-risk as well (cf. (3.12)). This is not the case for the expectile, as the first term in (5.9) is not an expectation as in the genuine Kusuoka representation [\(5.3\)](#page-19-1). Repeating the construction and passing to higher order expectiles leads to more complicated risk measures.

6 Summary

Higher order risk measures naturally integrate with stochastic optimization, as they are stochastic optimization problems themselves. This paper presents and derives explicit forms of higher order risk measures, specifically for spectral risk measures. These risk measures constitute the central building block of general law invariant risk measures.

Extending these results result it is demonstrated that stochastic dominance relations can be characterized by employing higher order risk measures, and vice versa. We provide a verification theorem, which makes higher stochastic dominance relations accisible to numerical compuations.

The results are exemplified for expectiles, a specific risk measure with unique properties.

References

- [1] P. Artzner, F. Delbaen, J.-M. Eber, and D. Heath. Coherent Measures of Risk. *Mathematical Finance*, 9:203–228, 1999. doi[:10.1111/1467-9965.00068.](https://doi.org/10.1111/1467-9965.00068) [2](#page-1-8)
- [2] A. Balbás, B. Balbás, R. Balbás, and J.-P. Charron. Bidual representation of expectiles. *Risks*, 11 (12):220, 2023. ISSN 2227-9091. doi[:10.3390/risks11120220.](https://doi.org/10.3390/risks11120220) [19](#page-18-2)
- [3] F. Bellini and C. Caperdoni. Coherent distortion risk measures and higherorder stochastic dominances. *North American Actuarial Journal*, 11(2):35–42, 2007. doi[:10.1080/10920277.2007.10597446.](https://doi.org/10.1080/10920277.2007.10597446) [1](#page-0-0)
- [4] F. Bellini, B. Klar, A. Müller, and E. Rosazza Gianin. Generalized quantiles as risk measures. *Insurance: Mathematics and Economics*, 54:41–48, 2014. doi[:10.1016/j.insmatheco.2013.10.015.](https://doi.org/10.1016/j.insmatheco.2013.10.015) 20
- [5] F. Bellini, B. Klar, and A. Müller. Expectiles, omega ratios and stochastic ordering. *Methodology and Computing in Applied Probability*, 20(3):855–873, 2016. doi[:10.1007/s11009-016-9527-2.](https://doi.org/10.1007/s11009-016-9527-2) [1](#page-0-0)
- [6] A. Ben-Tal and M. Teboulle. An old-new concept of convex risk measures: The optimized certainty equivalent. *Mathematical Finance*, 17:449–476, 2007. [2](#page-1-8)
- [7] G. Consigli, D. Dentcheva, F. Maggioni, and G. Micheli. Asset liability management under sequential stochastic dominance constraints, 2023. URL <https://optimization-online.org/?p=24837>. [1](#page-0-0)
- [8] D. Dentcheva and G. Martinez. Two-stage stochastic optimization problems with stochastic ordering constraints on the recourse. *European Journal of Operational Research*, 219(1):1–8, 2012. doi[:10.1016/j.ejor.2011.11.044.](https://doi.org/10.1016/j.ejor.2011.11.044) [1](#page-0-0)
- [9] D. Dentcheva, S. Penev, and A. Ruszczyński. Kusuoka representation of higher order dual risk measures. *Annals of Operations Research*, 181:325–335, 2010. doi[:10.1007/s10479-010-0747-5.](https://doi.org/10.1007/s10479-010-0747-5) [3](#page-2-2)
- [10] P. Dommel and A. Pichler. Convex risk measures based on divergence. *Pure and Applied Functional Analysis*, 6(6):1157–1181, 2021. [2](#page-1-8)
- [11] J. Dupačová and M. Kopa. Robustness of optimal portfolios under risk and stochastic dominance constraints. *European Journal of Operational Research*, 234(2):434–441, 2014. doi[:10.1016/j.ejor.2013.06.018.](https://doi.org/10.1016/j.ejor.2013.06.018) [1](#page-0-0)
- [12] M. Farooq and I. Steinwart. Learning rates for kernel-based expectile regression. *Machine Learning*, 108(2):203–227, 2018. doi[:10.1007/s10994-018-5762-9.](https://doi.org/10.1007/s10994-018-5762-9) [19](#page-18-2)
- [13] S. Frydenberg, T. E. Sønsteng Henriksen, A. Pichler, and S. Westgaard. Can commodities dominate stock and bond portfolios? *Annals of Operations Research*, 282(1-2):155–177, 2019. doi[:10.1007/s10479-018-2996-7.](https://doi.org/10.1007/s10479-018-2996-7) [1](#page-0-0)
- [14] W. J. Gutjahr and A. Pichler. Stochastic multi-objective optimization: a survey on non-scalarizing methods. *Annals of Operations Research*, 236(2):1–25, 2013. doi[:10.1007/s10479-013-1369-5.](https://doi.org/10.1007/s10479-013-1369-5) [1](#page-0-0)
- [15] M. Kopa, V. Moriggia, and S. Vitali. Individual optimal pension allocation under stochastic dominance constraints. *Annals of Operations Research*, 260(1–2):255–291, 2016. doi[:10.1007/s10479-016-2387-x.](https://doi.org/10.1007/s10479-016-2387-x) [1](#page-0-0)
- [16] M. Kopa, V. Moriggia, and S. Vitali. Multistage stochastic dominance: an application to pension fund management. *Annals of Operations Research*, 2023. doi[:10.1007/s10479-023-05658-y.](https://doi.org/10.1007/s10479-023-05658-y) [1](#page-0-0)
- [17] P. A. Krokhmal. Higher moment coherent risk measures. *Quantitative Finance*, 7(4):373–387, 2007. doi[:10.1080/14697680701458307.](https://doi.org/10.1080/14697680701458307) [3](#page-2-2)
- [18] S. Kusuoka. On law invariant coherent risk measures. In *Advances in mathematical economics*, volume 3, chapter 4, pages 83–95. Springer, 2001. doi[:10.1007/978-4-431-67891-5.](https://doi.org/10.1007/978-4-431-67891-5) [5](#page-4-5)
- [19] R. Lakshmanan and A. Pichler. Expectiles in risk averse stochastic programming and dynamic optimization. *Pure and Applied Functional Analysis*, 2023. [20](#page-19-5)
- [20] F. Maggioni and G. Ch. Pflug. Bounds and approximations for multistage stochastic programs. *SIAM Journal on Optimization*, 26(1):831–855, 2016. ISSN 1095-7189. doi[:10.1137/140971889.](https://doi.org/10.1137/140971889) [1](#page-0-0)
- [21] F. Maggioni and G. Ch. Pflug. Guaranteed bounds for general non-discrete multistage riskaverse stochastic optimization programs. *SIAM Journal on Optimization*, 29(1):454–483, 2019. doi[:10.1137/17M1140601.](https://doi.org/10.1137/17M1140601) [1](#page-0-0)
- [22] A. Malandii, V. Kuzmenko, and S. Uryasev. Expectile risk quadrangles and applications, 2024. sumbitted for review. [19](#page-18-2)
- [23] A. Müller and D. Stoyan. *Comparison methods for stochastic models and risks*. Wiley series in probability and statistics. Wiley, Chichester, 2002. ISBN 978-0-471-49446-1. URL <https://books.google.com/books?id=a8uPRWteCeUC>. [17](#page-16-3)
- [24] W. K. Newey and J. L. Powell. Asymmetric least squares estimation and testing. *Econometrica*, 55 (4):819–847, 1987. doi[:10.2307/1911031.](https://doi.org/10.2307/1911031) [19](#page-18-2)
- [25] W. Ogryczak and A. Ruszczyński. From stochastic dominance to mean-risk models: Semideviations as risk measures. *European Journal of Operational Research*, 116:33–50, 1999. doi[:10.1016/S0377-2217\(98\)00167-2.](https://doi.org/10.1016/S0377-2217(98)00167-2) [17](#page-16-3)
- [26] W. Ogryczak and A. Ruszczyński. On consistency of stochastic dominance and mean–semideviation models. *Math. Program., Ser. B*, 89:217–232, 2001. doi[:10.1007/s101070000203.](https://doi.org/10.1007/s101070000203) [17](#page-16-3)
- [27] W. Ogryczak and A. Ruszczyński. Dual stochastic dominance and related mean-risk models. *SIAM Journal on Optimization*, 13(1):60–78, 2002. doi[:10.1137/S1052623400375075.](https://doi.org/10.1137/S1052623400375075) [8](#page-7-5)
- [28] G. Ch. Pflug. Some remarks on the Value-at-Risk and the Conditional Value-at-Risk. In S. Uryasev, editor, *Probabilistic Constrained Optimization*, volume 49 of *Nonconvex Optimization and its Application*, chapter 15, pages 272–281. Springer US, 2000. doi[:10.1007/978-1-4757-3150-7.](https://doi.org/10.1007/978-1-4757-3150-7) [5,](#page-4-5) [8](#page-7-5)
- [29] G. Ch. Pflug and W. Römisch. *Modeling, Measuring and Managing Risk*. World Scientific, River Edge, NJ, 2007. doi[:10.1142/9789812708724.](https://doi.org/10.1142/9789812708724) [13](#page-12-2)
- [30] A. Pichler. The natural Banach space for version independent risk measures.*Insurance: Mathematics and Economics*, 53(2):405–415, 2013. doi[:10.1016/j.insmatheco.2013.07.005.](https://doi.org/10.1016/j.insmatheco.2013.07.005) [4](#page-3-5)
- [31] A. Pichler. A quantitative comparison of risk measures. *Annals of Operations Research*, 254(1): 251–275, 2017. doi[:10.1007/s10479-017-2397-3.](https://doi.org/10.1007/s10479-017-2397-3) [5](#page-4-5)
- [32] A. Pichler and A. Shapiro. Minimal representation of insurance prices. *Insurance: Mathematics and Economics*, 62:184–193, 2015. doi[:10.1016/j.insmatheco.2015.03.011.](https://doi.org/10.1016/j.insmatheco.2015.03.011) [5](#page-4-5)
- [33] R. T. Rockafellar and S. Uryasev. Optimization of Conditional Value-at-Risk. *Journal of Risk*, 2(3): 21–41, 2000. doi[:10.21314/JOR.2000.038.](https://doi.org/10.21314/JOR.2000.038) [8](#page-7-5)
- [34] R. T. Rockafellar and S. Uryasev. The fundamental risk quadrangle in risk management, optimization and statistical estimation. *Surveys in Operations Research and Management Science*, 18(1-2):33–53, 2013. doi[:10.1016/j.sorms.2013.03.001.](https://doi.org/10.1016/j.sorms.2013.03.001) [2](#page-1-8)
- [35] M. Sion. On general minimax theorems. *Pacific Journal of Mathematics*, 8(1):171–176, 1958. URL <https://projecteuclid.org/euclid.pjm/1103040253>. [11](#page-10-4)
- [36] A. W. van der Vaart. *Asymptotic Statistics*. Cambridge Uni-versity Press, 1998. doi[:10.1017/CBO9780511802256.](https://doi.org/10.1017/CBO9780511802256) <http://books.google.com/books?id=UEuQEM5RjWgC>. [6](#page-5-6)
- [37] J. F. Ziegel. Coherence and elicitability. *Mathematical Finance*, 26(4):901–918, 2014. doi[:10.1111/mafi.12080.](https://doi.org/10.1111/mafi.12080) [19](#page-18-2)