
SOME RESULTS INVOLVING THE A_α -EIGENVALUES FOR GRAPHS AND LINE GRAPHS *

João Domingos G. da Silva Jr.

Departamento de Engenharia de Produção
Centro Federal de Educação Tecnológica do Rio de Janeiro
Rio de Janeiro, Brazil
joao.dgomes@gmail.com

Carla Silva Oliveira

Departamento de Matemática
Escola Nacional de Ciências Estatísticas
Rio de Janeiro, Brazil
carla.oliveira@ibge.gov.br

Liliana Manuela G. C. da Costa

Departamento de Matemática
Colégio Pedro II
Rio de Janeiro, Brazil
lmgccosta@gmail.com

ABSTRACT

Let G be a simple graph with adjacency matrix $A(G)$, signless Laplacian matrix $Q(G)$, degree diagonal matrix $D(G)$ and let $l(G)$ be the line graph of G . In 2017, Nikiforov defined the A_α -matrix of G , $A_\alpha(G)$, as a linear convex combination of $A(G)$ and $D(G)$, the following way, $A_\alpha(G) := \alpha A(G) + (1 - \alpha)D(G)$, where $\alpha \in [0, 1]$. In this paper, we present some bounds for the eigenvalues of $A_\alpha(G)$ and for the largest and smallest eigenvalues of $A_\alpha(l(G))$. Extremal graphs attaining some of these bounds are characterized.

Keywords Line graphs · Characteristic polynomial · A_α -eigenvalues · A_α -matrix.

1 Introduction

Let $M_{n,m}(\mathbb{R})$ be the set of $n \times m$ real matrices, when $m = n$ we use for short $M_n(\mathbb{R})$. A matrix $M = [m_{ij}]$ is said non-negative ($M \geq 0$) if all its entries, m_{ij} , are non-negative, and M is considered positive ($M > 0$) if all its elements are strictly positive. If $M \in M_n(\mathbb{R})$, the M -characteristic polynomial is defined by $P_M(\lambda) = |\lambda I_n - M|$ and its roots are called M -eigenvalues. We shall index them in non-increasing order and denote by $\lambda_1(M) \geq \dots \geq \lambda_n(M)$. The collection of M -eigenvalues together with their multiplicities is called the M -spectrum, denoted by $\sigma(M)$. The largest M -eigenvalue, $\lambda_1(M)$, is also called spectral radius. The Rayleigh quotient is defined by $R(M, x) = \frac{x^T M x}{x^T x}$, for all nonzero vector $x \in \mathbb{R}^n$.

Let $G = (V, E)$ be a simple graph such that $|V| = n$ and $|E| = m$. For each vertex $v \in V$ the degree of v , denoted by $d(v)$, is defined by the number of edges incident to v . The minimum degree of G , is denoted by $\delta(G) = \min\{d(v) : v \in V\}$ and the maximum degree of G by $\Delta(G) = \max\{d(v) : v \in V\}$. The average degree of the neighbors of $v_i \in V$ is $m_i = \frac{1}{d(v_i)} \sum_{v_j \sim v_i} d(v_j)$. The graph G is called r -regular if each vertex of G has degree r .

The graph G is called non-null if it has at least one edge, and G is connected if every pair of distinct vertices of G is joined by a path in G . The complement of G , denoted by $\overline{G} = (\overline{V}, \overline{E})$, is the graph obtained from G with the same vertex set, $\overline{V} = V$, and $v_i v_j \in \overline{E}$ if and only if $v_i v_j \notin E$. Let $G = (V, E)$ and $H = (W, F)$ be graphs, if $W \subset V$ and $F \subset E$, then H is a subgraph of G . The subgraph denoted by $G - e = (V, E - e)$ is obtained from G by deleting the edge e .

*We would like to thank the National Council for Scientific and Technological Development (CNPq) - Brazil

We denote the path with n vertices by P_n , the complete graph by K_n , the complete bipartite graph with order $n = n_1 + n_2$ by K_{n_1, n_2} and, in particular, the star by $K_{1, n-1}$. A wheel graph of order n , W_n , is a graph that results of connecting all the vertices of a cycle of order $n - 1$ to a single universal vertex known as the hub. The pineapple graph, K_p^q , is obtained by appending q pendant edges to a vertex of a complete graph K_p where $q \geq 1$ and $p \geq 3$. The binomial tree², denoted by BT_k , is a tree defined recursively as follows: BT_0 consists of a single vertex and the binomial tree BT_k consists of two binomial trees BT_{k-1} that are linked together by an edge, as show in Figure 1.




Figure 1: Binomial tree.

The line graph of G , denoted by $l(G)$, is obtained in the following way: each edge in G corresponds to a vertex in $l(G)$, and for two edges in G that share a vertex, make an edge between their corresponding vertices in $l(G)$. From the definition it is important to note that $l(P_n) = P_{n-1}$ and $l(K_{1, n-1}) = K_{n-1}$. An example of a line graph can be seen in Figure 2.




Figure 2: A graph and its line graph.

The first and the second Zagreb indices defined by $Z_1(G) = \sum_{i=1}^n d^2(v_i)$ and $Z_2(G) = \sum_{uv \in E(G)} d(u)d(v)$, respectively,

were introduced by Gutman and Trinajestic, [2]. The first general Zagreb index is defined by $Z^{(p)}(G) = \sum_{i=1}^n d^p(v_i)$,

for $p \in \mathbb{R}$, $p \neq 0$ and $p \neq 1$ and it seems to have been first considered by Li et al. in [3, 4]. For $p = 2$, we have the $Z^{(2)}(G) = Z_1(G)$ and the study of its bounds and properties can be found at [5, 6, 7, 8, 9] and, for $p = 3$, we have $Z^{(3)}(G) = F(G)$, called the forgotten topological index or F-index, whose study appears in [2, 10, 11]. The general

Randić index is defined by $R_a(G) = \sum_{uv \in E(G)} (d(u)d(v))^a$, where $a \in \mathbb{R}^*$ and was introduced by Bollobás and Erdős

in [12]. The study of its bounds can be found at [13] and it is not difficult to see that there are close relations between these topological indices, for example, $R_1(G) = Z_2(G)$.

The adjacency matrix of G , denoted by $A = A(G) = [a_{ij}]$, is a square and symmetric matrix of order n , such that $a_{ij} = 1$ if v_i is adjacent to v_j and $a_{ij} = 0$, otherwise. The incident matrix of G , denoted by $B = B(G) = [b_{ij}]$, is a matrix of order $n \times m$ such that $b_{ij} = 1$ if e_j is an incident edge at v_i and $b_{ij} = 0$, otherwise. The degree matrix of G , denoted by $D(G)$, is the diagonal matrix that has the degree of the vertex v_i , $d(v_i)$, in the i^{th} position. The matrices $L(G) = D(G) - A(G)$ and $Q(G) = D(G) + A(G)$ are called Laplacian matrix and signless Laplacian matrix, respectively. For simplify the notation, we use $\lambda_i(Q(G)) = q_i$ for all $i = 1, \dots, n$.

²For more details we suggest [1].

Some results involving the A_α -eigenvalues for graphs and line graphs

In 2017 Nikiforov, [14], defined for any real $\alpha \in [0, 1]$ the convex linear combination $A_\alpha(G)$ of $A(G)$ and $D(G)$ in the following way:

$$A_\alpha(G) = \alpha D(G) + (1 - \alpha)A(G), \quad (1)$$

which we call the A_α -matrix. From the definition it is easy to see that $A_0(G) = A(G)$, $A_1(G) = D(G)$ and $A_{\frac{1}{2}}(G) = \frac{1}{2}Q(G)$. So, obtaining bounds for A_α -eigenvalues is an interesting problem because it contemplates the study of bounds for the adjacency and signless Laplacian matrices.

In this paper, some bounds for the A_α -eigenvalues are obtained for a simple graph and for its line graph. We present two lower bounds for $\lambda_1(A_\alpha(G))$ and make a comparison between them. Furthermore, we compare the new bounds with those existing in the literature and presented here and to do this, certain criteria needed to be defined. Specifically, we opted to evaluate bounds that pertain to identical extremal graphs.

This paper is organized as follows: in Section 2, we introduce some definitions and results required to prove the main results; in Section 3, we show the main results, together with some comparisons of the obtained bounds.

2 Preliminaries

In this section we present some results that will be useful to prove the main results of the paper. We start with the Theorem of Weyl and So, which inequalities involve eigenvalues of sums of Hermitian matrices.

Theorem 2.1 (Weyl). [15] *Let $A, B \in M_n(\mathbb{R})$ be symmetric and let the spectrum of A , B , and $A + B$ be $\sigma(A) = \{\lambda_1(A), \dots, \lambda_n(A)\}$, $\sigma(B) = \{\lambda_1(B), \dots, \lambda_n(B)\}$ and $\sigma(A + B) = \{\lambda_1(A + B), \dots, \lambda_n(A + B)\}$, respectively. Then,*

$$\lambda_{i+j-1}(A + B) \leq \lambda_i(A) + \lambda_j(B), \quad j = 1, \dots, n - i + 1 \quad (2)$$

for each $i = 1, \dots, n$, with equality for some pair i, j if and only if there is a nonzero vector x such that $Ax = \lambda_i x$, $Bx = \lambda_j x$ and $(A + B)x = \lambda_{i+j-1}x$. Also,

$$\lambda_i(A) + \lambda_j(B) \leq \lambda_{i+j-n}(A + B), \quad j = i, \dots, n \quad (3)$$

for each $i = 1, \dots, n$, with equality for some pair i, j if and only if there is a nonzero vector x such that $Ax = \lambda_i x$, $Bx = \lambda_j x$ e $(A + B)x = \lambda_{i+j-n}x$. If A and B have no common eigenvector, then the inequalities in (2) and (3) are strict.

As consequence of Theorem 2.1, we have Corollary 2.2.

Corollary 2.2. [15] *Let be $A, B \in M_n(\mathbb{R})$ symmetric. Then,*

$$\lambda_i(A) + \lambda_n(B) \leq \lambda_i(A + B) \leq \lambda_i(A) + \lambda_1(B), \quad (4)$$

with $i = 1, \dots, n$. Equality in the upper bound holds if and only if there is a nonzero vector x that is eigenvector of A, B and $A + B$ with corresponding eigenvalues λ_i, λ_1 and λ_i , respectively. Analogously, equality in the lower bound holds if and only if there is nonzero vector x that is eigenvector of A, B and $A + B$ with corresponding eigenvalues λ_i, λ_n and λ_i , respectively.

Theorem 2.3 relates the spectra of a graph and its subgraph, Theorem 2.4 and Proposition 2.5 show the Rayleigh theorem and its adaptation to A_α -matrix.

Theorem 2.3. [16] *Let G be a graph with n vertices and eigenvalues $\lambda_1(A(G)) \geq \dots \geq \lambda_n(A(G))$, and let H an induced subgraph of G with s vertices. If the eigenvalues of H are $\lambda_1(A(H)) \geq \dots \geq \lambda_s(A(H))$ then $\lambda_{n-s+i}(A(G)) \leq \lambda_i(A(H)) \leq \lambda_i(A(G))$, $\forall i = 1, \dots, s$.*

Theorem 2.4. [15] *Let $A \in M_n(\mathbb{R})$ symmetric with eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$. Then,*

$$\lambda_1 = \max_{x \neq 0} R(A, x) \quad \text{and} \quad \lambda_n = \min_{x \neq 0} R(A, x)$$

Proposition 2.5. [14] *If $\alpha \in [0, 1]$ and G is a graph of order n , then*

$$\lambda_1(A_\alpha(G)) = \max_{|x|=1} x^T A_\alpha(G)x \quad \text{and} \quad \lambda_n(A_\alpha(G)) = \min_{|x|=1} x^T A_\alpha(G)x. \quad (5)$$

Furthermore, if x is a unit vector, then $\lambda_1(A_\alpha(G)) = x^T A_\alpha(G)x$ if and only if x is an eigenvector of $\lambda_1(A_\alpha(G))$, and $\lambda_n(A_\alpha(G)) = x^T A_\alpha(G)x$ if and only if x is an eigenvector of $\lambda_n(A_\alpha(G))$.

Some results involving the A_α -eigenvalues for graphs and line graphs

The next result shows a lower bound for the largest eigenvalue of a non-negative matrix.

Lemma 2.6. [17, 18] Let $B = (b_{ij})$ be a non-negative $n \times n$ matrix with $n \geq 2$, $\lambda_1(B)$ be the largest eigenvalue of B , and set $\theta = \min_{1 \leq i \leq n} \{b_{ii}\}$. Then

$$\lambda_1(B) \geq \max_i \left\{ \frac{b_{ii} + \theta}{2} + \sqrt{\frac{(b_{ii} - \theta)^2}{4} + \sum_{i \neq j} b_{ij} b_{ji}} \right\}. \quad (6)$$

Moreover, if B is irreducible with $n \geq 3$, and at least two rows (two columns) of B contain more than one nonzero off-diagonal entry, then inequality is strict.

Lemma 2.7. [19] Let G be a connected graph with n vertices and $A(G) = A$ its adjacency matrix. Let $P(x)$ be any polynomial function and $S_v(P(A))$ be the row sums of $P(A)$ corresponding to each vertex v . Then

$$\min S_v(P(A)) \leq P(\lambda_1(A)) \leq \max S_v(P(A)).$$

Moreover, equality holds if and only if the row sums of $P(A)$ are all equal.

Bounds for the first Zagreb Index, the F-index and the general Randić index are presented in the next results.

Lemma 2.8. [6] Let G be a simple graph with n vertices and m edges. Let δ and Δ be the minimum and the maximum degree of G , respectively. Then, for $n \geq 3$, $Z_1(G) \geq \Delta^2 + \delta^2 + \frac{(2m - \Delta - \delta)^2}{n - 2}$. Furthermore, equality occurs if and only if $d_2 = \dots = d_{n-1}$.

Lemma 2.9. [6] Let G be a connected graph with n vertices and m edges. Let δ be the minimum degree of G . Then, $Z_1(G) \leq 2mn - n(n - 1)\delta + 2m(\delta - 1)$. Moreover, the equality holds if and only if G is a star graph or a regular graph.

Proposition 2.10. [10] Let G be a graph with m edges, whose first Zagreb index is $Z_1(G)$. Then,

$$F(G) \geq \frac{Z_1(G)^2}{2m}$$

Equality is attained in the case of regular graphs.

Theorem 2.11. [13] Let G be a graph with n vertices and $m \geq 1$ edges. Then for $a \geq 1$,

$$R_a \geq 4^a n^{-2a} m^{1+2a}$$

with equality if and only if G is a regular graph.

Theorem 2.12 and Theorem 2.13 show a relation between the vertices degree of the graph and its line graph. Theorem 2.14, Lemma 2.15 and Proposition 2.16, present some results involving line graphs.

Theorem 2.12. [20] Let G be a non-null graph such that $V(G) = \{v_1, \dots, v_n\}$ and m edges. Then

$$(a) \ l(G) \text{ has } m \text{ vertices and } \frac{1}{2} \sum_{i=1}^n d^2(v_i) - m \text{ edges.}$$

$$(b) \ \text{The degree of a vertex } e = v_i v_j \text{ in } l(G) \text{ is } d(e) = d(v_i) + d(v_j) - 2.$$

Theorem 2.13. [20] Let G be a graph with at least one edge. Then,

$$(a) \ \delta(l(G)) \geq 2\delta(G) - 2 \text{ with equality if and only if } G \text{ has two adjacent vertices of degree } \delta(G).$$

$$(b) \ \Delta(l(G)) \leq 2\Delta(G) - 2 \text{ with equality if and only if } G \text{ has two adjacent vertices of degree } \Delta(G).$$

Theorem 2.14. [20] If $\lambda_m(l(G))$ is the smallest eigenvalue of $A(l(G))$, then $-2 \leq \lambda_m(l(G))$.

Lemma 2.15. [21] Let G be a connected graph with n vertices, then $l(G)$ is the complete graph if and only if G is either $K_{1,n-1}$ or K_3 .

Proposition 2.16. [20] If H is a non-null subgraph of G , then $l(H)$ is an induced subgraph of $l(G)$.

As consequence of Proposition 2.16 and Theorem 2.3 we have the Corollary 2.17.

Some results involving the A_α -eigenvalues for graphs and line graphs

Corollary 2.17. *Let G be a graph with n vertices and $m \neq 0$ edges. Then*

$$\lambda_i(A_\alpha(l(G))) \geq \lambda_i(A_\alpha(l(G - e))) \geq \lambda_{i+1}(A_\alpha(l(G))),$$

$\forall i = 0, \dots, m - 1$ and $\alpha \in [0, 1]$.

Some known results in the literature about the incident matrix, Laplacian matrix and signless Laplacian matrix are presented below.

Lemma 2.18. [16] *Let G be a graph with m edges and $B = B(G)$ the incident matrix of G . Then $B^T B = 2I_m + A(l(G))$.*

Remark 2.19. *From Lemma 2.18 follows that*

$$(1 - \alpha)B^T B = (1 - \alpha)A(l(G)) + 2(1 - \alpha)I_m = A_\alpha(l(G)) - \alpha D(l(G)) + 2(1 - \alpha)I_m. \quad (7)$$

Taking $U = -\alpha D(l(G)) + 2(1 - \alpha)I_m$ and substituting in equation (7), we have

$$(1 - \alpha)B^T B = A_\alpha(l(G)) + U, \quad (8)$$

where $U = [u_{ij}]$ is a diagonal matrix of order m . From Theorem 2.12, for all k such that $1 \leq k \leq m$ and $e_k = v_i v_j$ we have that $u_{kk} = -\alpha d(e_k) + 2 - 2\alpha = -\alpha(d(v_i) + d(v_j) - 2) + 2 - 2\alpha = 2 - \alpha(d(v_i) + d(v_j))$.

Lemma 2.20. [16] *Let G be a graph with n vertices and m edges. Consider B and $D(G)$ the incident and the degree matrix of G , respectively. Then $BB^T = D(G) + A(G) = Q(G)$.*

Remark 2.21. *From Lemma 2.20 follows that*

$$(1 - \alpha)BB^T = (1 - \alpha)A(G) + (1 - \alpha)D(G) = A_\alpha(G) - \alpha D(G) + (1 - \alpha)D(G) = A_\alpha(G) + (1 - 2\alpha)D(G) \quad (9)$$

Proposition 2.22. [22] *The least eigenvalue of the signless Laplacian of a connected graph is equal to 0 if and only if the graph is bipartite. In this case 0 is a simple eigenvalue.*

Proposition 2.23. [23] *The matrices $L(G)$ and $Q(G)$ have the same characteristic polynomial if and only if G is a bipartite graph.*

Lemma 2.24 shows a linear correspondence between the eigenvalues of $A_\alpha(G)$ and $A(G)$ and Proposition 2.25 is an adaptation of Perron-Frobenius's Theorem for A_α -matrix.

Lemma 2.24. [14] *If $\alpha \in [0, 1]$ and $k = 1, \dots, n$ and G is a r -regular graph of order n , then there exists a linear correspondence between the eigenvalues of $A_\alpha(G)$ and $A(G)$, the following way*

$$\lambda_k(A_\alpha(G)) = \alpha r + (1 - \alpha)\lambda_k(A(G)). \quad (10)$$

In particular, $\lambda_1(A_\alpha(G)) = r$, $\forall \alpha \in [0, 1]$.

Proposition 2.25. [14] *Let $\alpha \in [0, 1]$, G be a graph and x be a non-negative eigenvector of $\lambda_1(A_\alpha(G))$.*

- (i) *If G is connected, then x is positive and unique minus scalar;*
- (ii) *If G is disconnected and P is the set of vertices with positive entries of x , then the subgraph induced by P is a union of components H of G with $\lambda_1(A_\alpha(H)) = \lambda_1(A_\alpha(G))$;*
- (iii) *If G is connected and μ is an eigenvalue of $A_\alpha(G)$ with a non-negative eigenvector, then $\mu = \lambda_1(A_\alpha(G))$;*
- (iv) *If G is connected, and H is a proper subgraph of G , then $\lambda_1(A_\alpha(H)) < \lambda_1(A_\alpha(G))$.*

Proposition 2.26 and Corollary 2.27 show the A_α -spectrum of K_n and $l(K_{1,n-1})$.

Proposition 2.26. [14] *The eigenvalues of $A_\alpha(K_n)$ are $\lambda_1(A_\alpha(K_n)) = n - 1$ and $\lambda_k(A_\alpha(K_n)) = \alpha n - 1$ for $2 \leq k \leq n$.*

Corollary 2.27. *Let $G \cong K_{1,n-1}$ and $\alpha \in [0, 1]$. Then $\sigma(A_\alpha(l(G))) = \{n - 2^{(1)}, (n - 1)\alpha - 1^{(n-2)}\}$.*

Proof. From Lemma 2.15 we know that $l(G)$ is a complete graph with $n - 1$ vertices and from Proposition 2.26 the result follows. \square

Theorem 2.28 provides relations between $P_{A_\alpha(l(G))}$ and $P_{A_\alpha(G)}$, and between $P_{A_\alpha(l(G))}$ and $P_{A(G)}$. Corollary 2.29 obtains the A_α -spectrum of the $l(K_n)$ and Corollary 2.30 shows a relation between $P_{A_\alpha(l(G))}$ and $P_{Q(G)}$ when G is r -regular.

Some results involving the A_α -eigenvalues for graphs and line graphs

Theorem 2.28. [24] Let G be a r -regular graph with n vertices and m edges such that $r \geq 2$ and $\alpha \in [0, 1]$. Then

$$P_{A_\alpha(l(G))}(\lambda) = (\lambda - 2r\alpha + 2)^{m-n} P_{A_\alpha(G)}(\lambda - r + 2)$$

and

$$P_{A_\alpha(l(G))}(\lambda) = (\lambda - 2r\alpha + 2)^{m-n} (1 - \alpha)^n P_{A(G)}\left(\frac{\lambda - r(\alpha + 1) + 2}{1 - \alpha}\right)$$

Corollary 2.29. Let $\alpha \in [0, 1]$. Then $\sigma(A_\alpha(l(K_n))) = \left\{ 2n - 4, (n(\alpha + 1) - 4)^{(n-1)}, (2\alpha(n-1) - 2)^{\binom{n(n-3)}{2}} \right\}$.

Proof. From Proposition 2.26 and Theorem 2.28 the result follows. \square

Corollary 2.30. Let G be a r -regular graph with n vertices, $m \neq 0$ edges and $\alpha \in [0, 1]$. Then

$$P_{A_\alpha(l(G))}(\lambda) = (\lambda - 2r\alpha + 2)^{m-n} (1 - \alpha)^n P_{Q(G)}\left(\frac{\lambda - 2r\alpha + 2}{1 - \alpha}\right)$$

Proof. From Theorem 2.28 and Lemma 2.20 follows that

$$\begin{aligned} P_{A_\alpha(l(G))}(\lambda) &= (\lambda - 2r\alpha + 2)^{m-n} |(\lambda - 2r\alpha + 2)I_n - (1 - \alpha)BB^T| \\ &= (\lambda - 2r\alpha + 2)^{m-n} (1 - \alpha)^n P_{Q(G)}\left(\frac{\lambda - 2r\alpha + 2}{1 - \alpha}\right). \end{aligned}$$

\square

Remark 2.31. Let G a r -regular graph with $\sigma(Q(G)) = \{q_1, \dots, q_n\}$ and $m > n$. From Corollary 2.30, we have α , $(2r\alpha - 2)$ and $\alpha(2r - q_i) + q_i - 2$ for $i = 1, \dots, n$ belong to $\sigma(A_\alpha(l(G)))$.

Example 1. Consider $G \cong K_n$. We know that $\sigma(Q(K_n)) = \{2n - 2, n - 2^{(n-1)}\}$, see [25, 26]. So, from Corollary 2.30 we have $\sigma(A_\alpha(l(K_n))) = \left\{ 2n - 4, n(\alpha + 1) - 4^{(n-1)}, 2\alpha(n-1) - 2^{\binom{n(n-3)}{2}} \right\}$, which can be seen in [24].

Example 2. Consider $G \cong K_{n,n}$. From Proposition 2.23, $\sigma(L(K_{n,n})) = \sigma(Q(K_{n,n}))$. Moreover, $\sigma(L(K_{n,n})) = \{2n, n^{(2n-2)}, 0\}$, which can be seen in [27, 26]. So from Corollary 2.30, we obtain $\sigma(A_\alpha(l(K_{n,n}))) = \left\{ 2n - 2, n(\alpha + 1) - 2^{(2n-2)}, 2\alpha n - 2^{(n^2-2n+1)} \right\}$.

The following results are bounds for the A_α -eigenvalues. In [14], Proposition 2.33 was introduced without extremal graphs so, we rewrite it and introduce its extremal graph.

Proposition 2.32. [14] If G is a graph with maximum degree Δ , then

$$\lambda_1(A_\alpha(G)) \geq \frac{1}{2} \left(\alpha(\Delta + 1) + \sqrt{\alpha^2(\Delta + 1)^2 + 4\Delta(1 - 2\alpha)} \right) \quad (11)$$

If G is connected, equality holds if and only if $G \cong K_{1,\Delta}$.

Proposition 2.33. [14] Let G be a graph with n vertices and $\alpha \in [0, 1]$. Then,

$$\min_{v_i \in V} \left\{ \sqrt{\alpha d^2(v_i) + (1 - \alpha) \sum_{v_j \sim v_i} d(v_j)} \right\} \leq \lambda_1(A_\alpha(G)) \leq \max_{v_i \in V} \left\{ \sqrt{\alpha d^2(v_i) + (1 - \alpha) \sum_{v_j \sim v_i} d(v_j)} \right\} \quad (12)$$

Moreover, the equalities holds if and only if G is regular.

Proof. We have already seen that $S_{v_i}(A_\alpha(G)) = d(v_i)$ and from [14] we have that $S_{v_i}(A_\alpha^2(G)) = \alpha d^2(v_i) + (1 - \alpha) \sum_{v_j \sim v_i} d(v_j)$.

Some results involving the A_α -eigenvalues for graphs and line graphs

From Lemma 2.7 follows that

$$\min_{v_i \in V} \{S_{v_i}(A_\alpha^2(G))\} \leq \lambda_1^2(A_\alpha(G)) \leq \max_{v_i \in V} \{S_{v_i}(A_\alpha^2(G))\}$$

and then the result follows.

Now, suppose initially that G is a r -regular graph. Hence, $d(v_i) = r, \forall v_i \in V$. Replacing in (12) we obtain $\lambda_1(A_\alpha(G)) = r$. Conversely, if both equalities hold, we have that all row sums are equal and then we can conclude that G is regular. \square

Proposition 2.34. [28] *Let G be a graph with n vertices. If $G \not\cong K_n$, then $\lambda_2(A_\alpha(G)) \geq 0$.*

Proposition 2.35. [29] *The largest eigenvalue of $A_\alpha(P_n)$ satisfies*

$$\lambda_1(A_\alpha(P_n)) \leq \begin{cases} 2\alpha + 2(1 - \alpha) \cos\left(\frac{\pi}{n+1}\right), & \text{if } 0 \leq \alpha < \frac{1}{2}; \\ 2\alpha + 2(1 - \alpha) \cos\left(\frac{\pi}{n}\right), & \text{if } \frac{1}{2} \leq \alpha \leq 1. \end{cases}$$

Equality holds if and only if $\alpha = 0$, $\alpha = \frac{1}{2}$ or $\alpha = 1$.

3 Main Results

In this section we show the main results of this paper that involve bounds for some A_α -eigenvalues of graphs and line graphs and, when is possible, we exhibited extremal graphs. Moreover, comparisons between some bounds are presented.

3.1 Some Bounds for A_α -eigenvalues of Graphs

Theorem 3.1. *Let G be a graph of order $n \geq 2$, Δ and δ be the maximum degree and the minimum degree of G , respectively, and $\alpha \in [0, 1]$. Then*

$$\lambda_1(A_\alpha(G)) \geq \frac{\alpha(\Delta + \delta) + \sqrt{\alpha^2(\Delta - \delta)^2 + 4(1 - \alpha)^2\Delta}}{2} \quad (13)$$

If G is connected, the equality holds if and only if $G \cong K_{1, n-1}$.

Proof. From Lemma 2.6 we have that

$$\begin{aligned} \lambda_1(A_\alpha(G)) &\geq \max_i \left\{ \frac{A_\alpha(G)_{ii} + \alpha\delta}{2} + \sqrt{\frac{(A_\alpha(G)_{ii} - \alpha\delta)^2}{4} + \sum_{i \neq j} A_\alpha(G)_{ij} A_\alpha(G)_{ji}} \right\} \\ &= \frac{\alpha\Delta + \alpha\delta}{2} + \sqrt{\frac{(\alpha\Delta - \alpha\delta)^2}{4} + \max_i \left\{ \sum_{i \neq j} A_\alpha(G)_{ij} A_\alpha(G)_{ji} \right\}} \\ &= \frac{\alpha(\Delta + \delta) + \sqrt{\alpha^2(\Delta - \delta)^2 + 4 \max_i \left\{ \sum_{i \neq j} A_\alpha(G)_{ij} A_\alpha(G)_{ji} \right\}}}{2} \end{aligned}$$

As $A_\alpha(G) = \alpha D(G) + (1 - \alpha)A(G)$, follows that

$$\sum_{i \neq j} A_\alpha(G)_{ij} A_\alpha(G)_{ji} = (1 - \alpha)^2 \sum_{i \neq j} a_{ij} a_{ji}$$

and then

$$\lambda_1(A_\alpha(G)) \geq \frac{\alpha(\Delta + \delta) + \sqrt{\alpha^2(\Delta - \delta)^2 + 4(1 - \alpha)^2\Delta}}{2}.$$

If $G \cong K_{1, n-1}$ we have that $\Delta = n - 1$ and $\delta = 1$, so from [14] we have the equality. Now suppose that equality in (13) holds. Since G is connected, $A_\alpha(G)$ is irreducible. By the equality condition in Lemma 2.6, there exists only one row (column) of $A_\alpha(G)$ containing more than one nonzero off-diagonal entry. Then there exists only a vertex v with $d(v) \geq 2$. So, $G \cong K_{1, n-1}$. \square

Some results involving the A_α -eigenvalues for graphs and line graphs

Corollary 3.2. *Let G be a graph of order $n \geq 2$, Δ the maximum degree, $\delta = 1$ and $\alpha \in [0, 1)$. Then the lower bound presented in (13) and (11) are equal.*

Proof. Taking $\delta = 1$ in (13) we have

$$\begin{aligned}\lambda_1(A_\alpha(G)) &\geq \frac{\alpha(\Delta + 1) + \sqrt{\alpha^2(\Delta - 1)^2 + 4(1 - \alpha)^2\Delta}}{2} \\ &= \frac{\alpha(\Delta + 1) + \sqrt{\alpha^2\Delta^2 - 2\Delta\alpha^2 + \alpha^2 + 4\Delta - 8\alpha\Delta + 4\alpha^2\Delta}}{2} \\ &= \frac{\alpha(\Delta + 1) + \sqrt{\alpha^2(\Delta + 1)^2 + 4\Delta(1 - 2\alpha)}}{2}\end{aligned}$$

□

Theorem 3.3. *Let $\alpha \in [0, 1]$ and G be a graph with $m \neq 0$ edges, $n \geq 3$ vertices, Δ and δ be the maximum and minimum degrees, respectively. Then,*

$$\lambda_1(A_\alpha(G)) \geq \alpha \frac{(\Delta^2 + \delta^2)(n - 2) + (2m - \Delta - \delta)^2}{2m(n - 2)} + (1 - \alpha) \frac{8m^3}{n^2(2\delta m + (n - 1)(2m - n\delta))} \quad (14)$$

The equality occurs if G is a regular graph.

Proof. From Proposition 2.5 we know that there exists an eigenvector $x \in \mathbb{R}^n$ associated with $\lambda_1(A_\alpha(G))$ that satisfies

$\lambda_1(A_\alpha(G)) = \max_{x \in \mathbb{R}^n} \frac{x^T A_\alpha(G) x}{x^T x}$. So for all $y \neq kx$, where $k \in \mathbb{R}$, we have

$$\begin{aligned}\lambda_1(A_\alpha(G)) &\geq \frac{y^T A_\alpha(G) y}{y^T y} = \frac{y^T (\alpha D(G) + (1 - \alpha) A(G)) y}{y^T y} \\ &= \frac{\alpha y^T D(G) y + (1 - \alpha) y^T A(G) y}{y^T y}.\end{aligned}$$

Taking $y = (d(v_1), d(v_2), \dots, d(v_n)) = (d_1, d_2, \dots, d_n)$, we have

$$\begin{aligned}\lambda_1(A_\alpha(G)) &\geq \frac{\alpha \sum_{i=1}^n d_i^3 + (1 - \alpha) y^T A(G) y}{\sum_{i=1}^n d_i^2} = \frac{\alpha \sum_{i=1}^n d_i^3}{\sum_{i=1}^n d_i^2} + \frac{(1 - \alpha) 2 \sum_{v_i \sim v_j} d_i d_j}{\sum_{i=1}^n d_i^2} \\ &= \alpha \frac{\sum_{i=1}^n d_i^3}{\sum_{i=1}^n d_i^2} + (1 - \alpha) \frac{2 \sum_{v_i \sim v_j} d_i d_j}{\sum_{i=1}^n d_i^2} = \alpha \frac{F(G)}{Z_1(G)} + (1 - \alpha) \frac{2R_1(G)}{Z_1(G)}.\end{aligned}$$

From Proposition 2.10 and Theorem 2.11 follows that

$$\lambda_1(A_\alpha(G)) \geq \alpha \frac{Z_1^2(G)}{2m} + (1 - \alpha) \frac{8m^3}{n^2 Z_1(G)} = \alpha \frac{Z_1(G)}{2m} + (1 - \alpha) \frac{8m^3}{n^2 Z_1(G)}.$$

Using Lemmas 2.8 and 2.9 with some algebraic manipulation the result follows.

To prove the equality suppose that G is r -regular graph. From Lemma 2.24 we know that $\lambda_1(A_\alpha(G)) = r$ and moreover that $\Delta = \delta = r$ and $m = \frac{nr}{2}$. Then,

$$\begin{aligned}&\alpha \frac{n((\Delta + \delta)^2 - 4m(\Delta + \delta - m) + (\Delta^2 + \delta^2)(n - 2))}{2m} + (1 - \alpha) \frac{8m^3}{n^2(2\delta m - (n - 1)(2m - n\delta))} = \\ &\frac{n((r + r)^2 - 4 \frac{nr}{2} \left(r + r - \frac{nr}{2}\right) + (r^2 + r^2)(n - 2))}{2 \frac{nr}{2}} + (1 - \alpha) \frac{8 \left(\frac{nr}{2}\right)^3}{n^2 \left(2r \frac{nr}{2} - (n - 1) \left(2 \frac{nr}{2} - nr\right)\right)} = \\ &\alpha r - \frac{2r\alpha}{n} + \frac{2r^2\alpha}{nr} - \alpha r + r = r\end{aligned}$$

□

Theorem 3.4. *Let G be a graph with n vertices, $m \neq 0$ edges, $\Delta(G)$ its maximum degree, $\delta(G)$ its minimum degree and $\alpha \in [0, 1]$. Then,*

$$\lambda_1(A_\alpha(G)) \leq \sqrt{\alpha\Delta^2 + (1-\alpha)(\Delta(\delta-1) - \delta(n-1) + 2m)}. \quad (15)$$

Moreover, equality holds if and only if G is regular.

Proof. Let $S_v(M)$ the row sums of M corresponding to each vertex v . Since $A_\alpha(G) = \alpha D(G) + (1-\alpha)A(G)$, we have that $S_v(A_\alpha(G)) = d(v)$ and $S_v(A(G)D(G)) = S_v(A^2(G)) = \sum_{u \sim v} d(u) = 2m - d(v) - \sum_{\substack{u \sim v \\ u \neq v}} d(u)$. Then,

$$\begin{aligned} S_v(A_\alpha^2(G)) &= S_v((\alpha D(G) + (1-\alpha)A(G))^2) = S_v(\alpha D(G)A_\alpha(G) + \alpha(1-\alpha)A(G)D(G) + (1-\alpha)^2 A^2(G)) \\ &= \alpha S_v(D(G)A_\alpha(G)) + \alpha(1-\alpha)S_v(A(G)D(G)) + (1-\alpha)^2 S_v(A^2(G)) \\ &= \alpha d^2(v) + \alpha(1-\alpha) \left(2m - d(v) - \sum_{\substack{u \sim v \\ u \neq v}} d(u) \right) + (1-\alpha)^2 \left(2m - d(v) - \sum_{\substack{u \sim v \\ u \neq v}} d(u) \right) \\ &= \alpha d^2(v) + (1-\alpha) \left(2m - d(v) - \sum_{\substack{u \sim v \\ u \neq v}} d(u) \right) \\ &\leq \alpha d^2(v) + (1-\alpha)(2m - d(v) - (n - d(v) - 1)\delta) \\ &= \alpha d^2(v) + (1-\alpha)(2m + (\delta-1)d(v) - \delta(n-1)) \\ &\leq \alpha\Delta^2 + (1-\alpha)(2m + (\delta-1)\Delta - \delta(n-1)) \end{aligned}$$

Hence, for every $v \in V$, we have

$$S_v(A_\alpha^2(G)) \leq \alpha\Delta^2 + (1-\alpha)(2m + (\delta-1)\Delta - \delta(n-1)).$$

By Lemma 2.7,

$$\lambda_1^2(A_\alpha(G)) \leq \alpha\Delta^2 + (1-\alpha)(2m + (\delta-1)\Delta - \delta(n-1))$$

Solving the quadratic inequality, we obtain the result.

Suppose initially that G is a r -regular graph, so

$$\begin{aligned} \sqrt{\alpha\Delta^2 + (1-\alpha)(\Delta(\delta-1) - \delta(n-1) + 2m)} &= \sqrt{\alpha r^2 + (1-\alpha) \left(r(r-1) - r(n-1) + 2\frac{nr}{2} \right)} \\ &= \sqrt{\alpha r^2 + (1-\alpha)(r^2 - r - rn + r + nr)} = r \end{aligned}$$

Now, suppose that holds the equality in (15). This implies that all inequalities present in its proof are equalities. So, $d(v) = \Delta$ for every $v \in V$, which implies that G is a regular graph. □

3.2 Bounds Comparison

In this subsection, tables and graphs are presented to compare the proposed bounds by the authors in Subsection 3.1 among themselves and also to compare these bounds with others found in the literature.

3.2.1 Lower bounds proposed

We provide tables illustrating the performance of the two bounds introduced in the preceding section alongside the precise value of $\lambda_1(A_\alpha(G))$. Notice that these bounds have different extremal graphs. Three families of graphs were used, two from special trees, the path and the binomial tree, and the third from the pineapple graph.

Starting with the path P_n , in Tab. 1 we notice that the values obtained using bound (13) are always better than those obtained by bound (14), starting with a notable difference which reduces as α increases, and for values of α close to 1, it ceases to be significant.

Some results involving the A_α -eigenvalues for graphs and line graphs

Lower Bound Comparison for $\mathbf{G} \cong \mathbf{P}_n$											
α		0.0	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9
$n = 100$	$\lambda_1(A_\alpha)$	1.99903	1.99913	1.99922	1.99931	1.99941	1.99951	1.9996	1.9997	1.9998	1.9999
	(13)	1.41421	1.42377	1.43578	1.45125	1.47178	1.5	1.54031	1.6	1.6899	1.8217
	(14)	0.07841	0.26906	0.45971	0.65036	0.841	1.03165	1.2223	1.41295	1.6036	1.79425
$n = 500$	$\lambda_1(A_\alpha)$	1.99996	1.99996	1.99997	1.99997	1.99998	1.99998	1.99998	1.99999	1.99999	2.0
	(13)	1.41421	1.42377	1.43578	1.45125	1.47178	1.5	1.54031	1.6	1.6899	1.8217
	(14)	0.01594	0.21404	0.41215	0.61025	0.80836	1.00647	1.20457	1.40268	1.60078	1.79889
$n = 1000$	$\lambda_1(A_\alpha)$	1.99999	1.99999	1.99999	1.99999	1.99999	2.0	2.0	2.0	2.0	2.0
	(13)	1.41421	1.42377	1.43578	1.45125	1.47178	1.5	1.54031	1.6	1.6899	1.8217
	(14)	0.00798	0.20704	0.40609	0.60514	0.80419	1.00324	1.20229	1.40134	1.6004	1.79945

Table 1: Lower bounds comparison for a path graph increasing the number of nodes.

In Tab.2, the comparative study was carried out using the binomial tree, BT_k . The values obtained by (13) are also better than those obtained by (14), and in this case, regardless of the value of α , the values obtained by the first bound are much better than those obtained by the second.

Lower Bound Comparison for $\mathbf{G} \cong \mathbf{BT}_k$											
α		0.0	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9
$k = 7$ ($n = 128$)	$\lambda_1(A_\alpha)$	3.45291	3.62629	3.82894	4.06529	4.3399	4.65723	5.02142	5.436	5.9036	6.42533
	(13)	2.64575	2.8	3.0	3.25913	3.58997	4.0	4.48806	5.04499	5.65764	6.31293
	(14)	0.06153	0.26415	0.46677	0.66939	0.87201	1.07463	1.27725	1.47988	1.6825	1.88512
$k = 9$ ($n = 512$)	$\lambda_1(A_\alpha)$	3.97329	4.23334	4.53693	4.88921	5.29487	5.75785	6.28124	6.8671	7.51632	8.22825
	(13)	3.0	3.22947	3.52982	3.91868	4.40832	5.0	5.68328	6.44109	7.25576	8.11248
	(14)	0.01556	0.21852	0.42148	0.62443	0.82739	1.03035	1.23331	1.43626	1.63922	1.84218
$k = 10$ ($n = 1024$)	$\lambda_1(A_\alpha)$	4.21077	4.51674	4.87387	5.28746	5.76201	6.30095	6.90661	7.58018	8.32164	9.12947
	(13)	3.16228	3.43141	3.78514	4.24278	4.81534	5.5	6.28161	7.13976	8.05513	9.01233
	(14)	0.0078	0.21	0.41221	0.61441	0.81662	1.01883	1.22103	1.42324	1.62544	1.82765

Table 2: Lower bounds comparison for a binomial tree graph increasing the number of nodes.

The question that arose was: would bound (13) always produce better approximations than bound (14)? To answer this question, we sought to use a family of graphs that were structurally very different from the tree, having chosen the pineapple graph. As we can see in Tab.3 the situation is reversed. Here the values obtained by (13) are always much worse than those provided by (14), which is more evident for values of α close to 0.

Lower Bound Comparison for $\mathbf{G} \cong \mathbf{K}_p^1$											
α		0.0	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9
$p = 99$	$\lambda_1(A_\alpha)$	98.0001	98.00109	98.00209	98.00309	98.00411	98.00513	98.00617	98.00725	98.00842	98.00999
	(13)	9.94987	15.20784	22.62537	31.26653	40.48902	50.0	59.66816	69.42964	79.25048	89.11122
	(14)	95.13821	95.42441	95.71061	95.99681	96.28301	96.56921	96.85541	97.14161	97.42781	97.71401
$p = 499$	$\lambda_1(A_\alpha)$	498.0	498.0002	498.0004	498.0006	498.0008	498.00101	498.00121	498.00141	498.00162	498.00184
	(13)	22.33831	57.00312	102.90936	151.31907	200.49776	250.0	299.66696	349.42878	399.25009	449.11113
	(14)	495.02793	495.32514	495.62234	495.91955	496.21676	496.51397	496.81118	497.10838	497.40559	497.7028
$p = 999$	$\lambda_1(A_\alpha)$	998.0	998.0001	998.0002	998.0003	998.0004	998.0005	998.0006	998.0007	998.0008	998.00091
	(13)	31.60696	107.43866	202.95339	301.32614	400.49888	500.0	599.66681	699.42868	799.25005	899.11112
	(14)	995.01398	995.31258	995.61119	995.90979	996.20839	996.50699	996.80559	997.1042	997.4028	997.7014

Table 3: Lower bounds comparison for a pineapple graph K_p^1 increasing the number of nodes.

Therefore, taking into account the previous observations, we can conclude that the bounds (13) and (14) are incomparable.

3.2.2 Lower bounds proposed and Nikiforov's bounds.

Now, we move on to comparing our bounds with other bounds present in literature. Among the variety of lower bounds that exist, we selected those that are due to Nikiforov and that have the same extremal graphs as ours. The choice of these bounds is due to the fact that they were the first to appear in the literature and also as a simple tribute to the importance of Nikiforov's contributions to Spectral Graph Theory. Thus, having the star as an extremal graph, we compare bound (13) with bound (11) and, having the regular graph as an extremal, we compare bounds (14) and (12).

The results are presented through graphs in the variable α . Each graph shows the difference between the value obtained by each of the bounds and $\lambda_1(A_\alpha(W_n))$, which makes it easier to observe the aspects to be highlighted.

Some results involving the A_α -eigenvalues for graphs and line graphs

Firstly we note that for regular graphs the bound (13) is better than the bound (11), which is proved in the next Proposition.

Proposition 3.5. *If G is a r -regular graph and $\alpha \in [0, 1]$, then the lower bound in (13) is better than the lower bound in (11).*

Proof. Since $r \in \mathbb{N}^*$ and $\alpha \in [0, 1]$ we have

$$\begin{aligned}
& \left(r^{\frac{3}{2}} - r^{\frac{1}{2}} \right) \alpha(1 - \alpha) \geq 0 \\
& -\alpha^2 r^{\frac{3}{2}} + \alpha^2 r^{\frac{1}{2}} + \alpha r^{\frac{3}{2}} - \alpha r^{\frac{1}{2}} \geq 0 \\
& -\alpha^2 r^{\frac{3}{2}} + \alpha^2 r^{\frac{1}{2}} + \alpha r^{\frac{3}{2}} - \alpha r^{\frac{1}{2}} + \frac{\alpha^2 r^2}{4} + \frac{\alpha^2 r}{2} + \frac{\alpha^2}{4} - 2r\alpha + r - \frac{\alpha^2 r^2}{4} - \frac{\alpha^2 r}{2} - \frac{\alpha^2}{4} + 2r\alpha - r \geq 0 \\
& -\alpha^2 r^{\frac{3}{2}} + \alpha^2 r^{\frac{1}{2}} + \alpha r^{\frac{3}{2}} - \alpha r^{\frac{1}{2}} + \frac{\alpha^2 r^2}{4} + \frac{\alpha^2 r}{2} + \frac{\alpha^2}{4} - 2r\alpha + r \geq +\frac{\alpha^2 r^2}{4} + \frac{\alpha^2 r}{2} + \frac{\alpha^2}{4} - 2r\alpha + r \\
& \left(\frac{2r\alpha + 2(1 - \alpha)r^{\frac{1}{2}} - \alpha(r + 1)}{2} \right)^2 \geq \left(\frac{\sqrt{\alpha^2(r + 1)^2 + 4r(1 - 2\alpha)}}{2} \right)^2
\end{aligned}$$

So,

$$\frac{2r\alpha + 2(1 - \alpha)r^{\frac{1}{2}} - \alpha(r + 1)}{2} \geq \frac{\sqrt{\alpha^2(r + 1)^2 + 4r(1 - 2\alpha)}}{2}$$

and then the result follows. \square

For the graph wheel (W_n), which is not a regular graph, Figure 3 shows that the bounds (13) and (11) have a similar behavior, although bound (13) is slightly better. Note that, the greater the number of vertices, the more similar the values provided by the two bounds.




Figure 3: Comparing lower bound for W_n .

From Corollary 3.2, Proposition 3.5 and after some computational tests, the following question arises: is the bound (13) always better than the bound (11)? It is worth noting that even in light of the results, this question is still under discussion.

Now, we see what happens with the bounds (12) and (14). If we consider $G \simeq K_{1,n-1}$ or $G \simeq W_n$, we can observe from Figures 4 and 5, respectively, that for some values of α , Nikiforov's bound is better than our bound, and that for the other values of α , the opposite happens. Therefore, these bounds are incomparable.

Some results involving the A_α -eigenvalues for graphs and line graphs




Figure 4: Comparing lower bound for $K_{1,n-1}$.




Figure 5: Comparing lower bound for W_n .

3.2.3 Upper bounds

Now, utilize the upper bounds, we compare the bounds introduced in (12) and (15). The Proposition 3.6 shows that the bound (12) is always lower or equal than the bound (15).

Proposition 3.6. *Let G be a graph with n vertices, $m \neq 0$ edges, Δ its maximum degree, δ its minimum degree and $\alpha \in [0, 1]$. Then,*

$$\max_{v_i \in V} \left\{ \sqrt{\alpha d^2(v_i) + (1 - \alpha) \sum_{v_j \sim v_i} d(v_j)} \right\} \leq \sqrt{\alpha \Delta^2 + (1 - \alpha)(\Delta(\delta - 1) - \delta(n - 1) + 2m)} \quad (16)$$

Moreover, if equality holds then the maximum in the left side of (16) is achieved when $d(v_i) = \Delta$.

Proof. Suppose that the maximum of the left-hand side of the inequality (16) is reached at vertex v_i and to simplify the notation, we use $d(v_i) = d_i$ for all $i \in \{1, \dots, n\}$. Now,

Some results involving the A_α -eigenvalues for graphs and line graphs

$$\begin{aligned}
& \alpha\Delta^2 + (1-\alpha)(\Delta(\delta-1) - \delta(n-1) + 2m) - \alpha d_i^2 - (1-\alpha) \sum_{v_j \sim v_i} d_j = \\
& \alpha(\Delta^2 - d_i^2) + (1-\alpha) \left(\Delta(\delta-1) - \delta(n-1) + 2m - \sum_{v_j \sim v_i} d_j \right) = \\
& \alpha(\Delta^2 - d_i^2) + (1-\alpha) \left(\Delta(\delta-1) - \delta(n-1) + \sum_{1 \leq s \leq n} d_s - \sum_{v_j \sim v_i} d_j \right) = \\
& \alpha(\Delta^2 - d_i^2) + (1-\alpha) \left(\Delta(\delta-1) - \delta(n-1) + \sum_{v_j \not\sim v_i} d_j + d_i \right) = \\
& \alpha(\Delta^2 - d_i^2) + (1-\alpha) \left(\delta(\Delta - n + 1) + \sum_{v_j \not\sim v_i} d_j + d_i - \Delta \right) \geq \\
& \alpha(\Delta^2 - d_i^2) + (1-\alpha) (\delta(\Delta - n + 1) + \delta(n-1 - d_i) + d_i - \Delta) = \\
& \alpha(\Delta^2 - d_i^2) + (1-\alpha) (\delta(\Delta - d_i) - (\Delta - d_i)) = \\
& \alpha(\Delta^2 - d_i^2) + (1-\alpha)(\delta-1)(\Delta - d_i) \tag{17}
\end{aligned}$$

Here we have two cases to consider: if $d_i \neq \Delta$, follows that (17) is greater or equal to 0; if $d_i = \Delta$, follows that (17) is equal to 0. So, in both cases, the result follows.

Now, suppose that

$$\alpha\Delta^2 + (1-\alpha)(\Delta(\delta-1) - \delta(n-1) + 2m) = \alpha d_i^2 + (1-\alpha) \sum_{v_j \sim v_i} d_j$$

for all $\alpha \in [0, 1]$. Considering both members of the previous equality as polynomials in α and taking into account the equality of polynomials, we have that

$$\Delta^2 - d_i^2 - \sum_{v_j \sim v_i} d_j + \Delta(\delta-1) - \delta(n-1) + 2m = 0 \tag{18}$$

and

$$\Delta(\delta-1) - \delta(n-1) + 2m - \sum_{v_j \sim v_i} d_j = 0. \tag{19}$$

From equality (19) we obtain

$$\Delta(\delta-1) - \delta(n-1) + \sum_{v_j \not\sim v_i} d_j + d_i = 0.$$

As $\sum_{v_j \not\sim v_i} d_j \geq \delta(n-1 - d_i)$ and substituting in the previous equality, we get

$$0 \geq \Delta(\delta-1) - \delta(n-1) + \delta(n-1 - d_i) + d_i$$

and then, after some algebraic manipulation, we get

$$0 \geq (\delta-1)(\Delta - d_i)$$

As $\delta \geq 1$ and $\Delta \geq d_i$ we must have that

$$(\delta-1)(\Delta - d_i) = 0,$$


and then $\delta = 1$ or $d_i = \Delta$.

To satisfy both equations, (18) and (19), we need to have $d_i = \Delta$. □

Some results involving the A_α -eigenvalues for graphs and line graphs


Although the values obtained from bound (12) are less or equal than those obtained from bound (15), in cases of equality, the latter is computed much faster than the former. We reached this conclusion after some computational tests made with, among others, the following families of graphs: $K_{1,n-1}$, W_n , regular graphs, BT_k , Helm graphs³, Windmill graphs⁴. We can observe the elapsed time to compute the aforementioned value in Figure 5.

Elapsed time to compute the upper bound for $K_{1,n-1}$.



(a)

Elapsed time to compute the upper bound for K_n .



(b)

³A Helm graph, H_n , is constructed from a W_n by adding n vertices of degree 1, one adjacent to each terminal vertex. For more details, we suggest [30].

⁴A windmill graph $W(\nu, k)$ consists of ν copies of the complete graph K_k , with every vertices connected to a common vertex.




Figure 5: Average time spent calculating the upper bound 100 times for each number of different vertices in each graph.

3.3 Some results for A_α -spectra of Line Graphs

In this section we present some results involving line graphs. The study of the A_α matrix is very recent and, after a bibliographic research, no results were found, at least so far, involving line graphs and this matrix. Next theorem is based on Theorem 2.14 and presents a lower bound for the smallest eigenvalue of $A_\alpha(l(G))$.

Theorem 3.7. *Let G be a graph with m edges and $\delta(G)$ its minimum degree. Then,*

$$\lambda_m(A_\alpha(l(G))) \geq 2\alpha\delta(G) - 2, \quad (20)$$

for $\alpha \in [0, 1)$.

Proof. Let B the incidence matrix of G . From Remark 2.19 we have

$$(1 - \alpha)B^T B = A_\alpha(l(G)) + 2(1 - \alpha)I_m - \alpha D(l(G)).$$

Let $\{w_1, \dots, w_m\}$ be the orthonormal basis of eigenvectors associated to $\lambda_1(A_\alpha(l(G))), \dots, \lambda_m(A_\alpha(l(G)))$ such that $A_\alpha(l(G))w_i = \lambda_i(A_\alpha(l(G)))w_i$, $\forall i = 1, \dots, m$. Then

$$\begin{aligned} (1 - \alpha)w_m^T B^T B w_m &= w_m^T A_\alpha(l(G)) w_m + 2(1 - \alpha)w_m^T w_m - \alpha w_m^T D(l(G)) w_m \\ &\leq \lambda_m(A_\alpha(l(G))) + 2(1 - \alpha) - \alpha\delta(l(G)). \end{aligned}$$

As $B^T B$ is positive semi-definite, we get

$$0 \leq w_m^T B^T B w_m \leq \frac{\lambda_m(A_\alpha(l(G))) + 2(1 - \alpha) - \alpha\delta(l(G))}{1 - \alpha}.$$

Therefore,

$$0 \leq \lambda_m(A_\alpha(l(G))) + 2(1 - \alpha) - \alpha\delta(l(G))$$

From Theorem 2.13 the result follows

$$\lambda_m(A_\alpha(l(G))) \geq 2\alpha\delta(G) - 2$$

□

Corollary 3.8. *If G is a r -regular graph with n vertices, $m \neq 0$ edges, $r \geq 2$ and $\alpha \in [0, 1)$, then $\lambda_m(A_\alpha(l(G))) = 2r\alpha - 2$ if and only if $m > n$ or G is bipartite.*

Some results involving the A_α -eigenvalues for graphs and line graphs

Proof. Let G a r -regular graph. Firstly suppose that $\lambda_m(A_\alpha(l(G))) = 2r\alpha - 2$. From Corollary 2.30, we have that $m > n$ or 0 is root of $P_{Q(G)}$ and, from Proposition 2.22, it follows that G is bipartite.

If $m > n$, from Theorem 2.28 we have that $2r\alpha - 2$ is an eigenvalue of $A_\alpha(l(G))$ and from Theorem 3.7 it is the smallest one. If G is bipartite, from Proposition 2.22, we have that $0 \in \sigma(Q(G))$ and applying Corollary 2.30, the result follows. \square

Example 3. If $G \cong C_n$ and n is even, from Corollary 3.8 we get that $\lambda_m(A_\alpha(l(G))) = 4\alpha - 2$.

Next propositions present some bounds for the largest eigenvalue of $A_\alpha(l(G))$.

Proposition 3.9. Let G be a connected graph with n vertices and $\alpha \in [0, 1)$. Then, $\lambda_1(A_\alpha(l(G))) < 2$ if and only if $G \cong P_n$.

Proof. Suppose that $G \cong P_n$. From Proposition 2.35,

$$\lambda_1(A_\alpha(l(P_n))) = \lambda_1(A_\alpha(P_{n-1})) < 2.$$

Now, suppose by contradiction that $G \not\cong P_n$. Then $l(G)$ contains at least a cycle C . From Theorem 2.3 and Lemma 2.24 we have

$$\lambda_1(A_\alpha(l(G))) \geq \lambda_1(A_\alpha(C)) = 2,$$

and the result follows. \square

Theorem 3.10. Let G be a graph with maximum degree Δ and minimum degree δ . Then

$$\lambda_1(A_\alpha(G)) + (1 - 2\alpha)\delta - \max_{v_i v_j \in E(G)} \{2 - \alpha(d(v_i) + d(v_j))\} \leq \lambda_1(A_\alpha(l(G))) \quad (21)$$

and

$$\lambda_1(A_\alpha(l(G))) \leq \lambda_1(A_\alpha(G)) + (1 - 2\alpha)\Delta - \min_{v_i v_j \in E(G)} \{2 - \alpha(d(v_i) + d(v_j))\} \quad (22)$$

Proof. Let B be the incidence matrix of G . We know that $\lambda_1(BB^T) = \lambda_1(B^T B)$. Applying Corollary 2.2 in (8) we obtain

$$\lambda_1(A_\alpha(l(G))) + \min_{v_i v_j \in E(G)} \{2 - \alpha(d(v_i) + d(v_j))\} \leq \lambda_1(A_\alpha(l(G)) + U) \leq \lambda_1(A_\alpha(l(G))) + \max_{v_i v_j \in E(G)} \{2 - \alpha(d(v_i) + d(v_j))\} \quad (23)$$

Let be x a unit non-negative eigenvector associated to $\lambda_1(A_\alpha(G))$ and z be a unit non-negative eigenvector associated to $\lambda_1(BB^T)$. From Theorem 2.4, Proposition 2.5 and Remark 2.21 we have

$$\begin{aligned} (1 - \alpha)\lambda_1(BB^T) &= (1 - \alpha) \max_{|z|=1} z^T BB^T z \geq (1 - \alpha)x^T BB^T x = x^T A_\alpha(G)x + (1 - 2\alpha)x^T D(G)x \\ &= \lambda_1(A_\alpha(G)) + (1 - 2\alpha) \sum_{i=1}^n d(v_i)x_i^2 \geq \lambda_1(A_\alpha(G)) + (1 - 2\alpha)\delta. \end{aligned} \quad (24)$$

Thus, from (23)

$$\lambda_1(A_\alpha(l(G))) + \max_{v_i v_j \in E(G)} \{2 - \alpha(d(v_i) + d(v_j))\} \geq \lambda_1(A_\alpha(G)) + (1 - 2\alpha)\delta$$

and, as a consequence, inequality (21) follows.

On the other hand, again from Theorem 2.4, Proposition 2.5 and Remark 2.21 we have

$$\begin{aligned} \lambda_1(A_\alpha(G)) &= \max_{|x|=1} x^T A_\alpha(G)x \geq z^T A_\alpha(G)z = z^T (1 - \alpha)BB^T z - (1 - 2\alpha)z^T D(G)z \\ &\geq \lambda_1(A_\alpha(l(G))) + \min_{v_i v_j \in E(G)} \{2 - \alpha(d(v_i) + d(v_j))\} - (1 - 2\alpha) \sum_{i=1}^n d(v_i)z_i^2 \\ &\geq \lambda_1(A_\alpha(l(G))) + \min_{v_i v_j \in E(G)} \{2 - \alpha(d(v_i) + d(v_j))\} - (1 - 2\alpha)\Delta. \end{aligned} \quad (25)$$

and the inequality (22) follows. \square

Corollary 3.11. Let G be a r -regular graph and $\alpha \in [0, 1]$. Then $\lambda_1(A_\alpha(l(G))) = 2r - 2$.

Some results involving the A_α -eigenvalues for graphs and line graphs

Proof. From Theorem 3.10,

$$\lambda_1(A_\alpha(G)) + (1 - 2\alpha)r - 2 + 2r\alpha \leq \lambda_1(A_\alpha(l(G))) \leq \lambda_1(A_\alpha(G)) + (1 - 2\alpha)r - 2 + 2r\alpha.$$

So, $\lambda_1(A_\alpha(l(G))) = \lambda_1(A_\alpha(G)) + (1 - 2\alpha)r - 2 + 2r\alpha$. As $\lambda_1(A_\alpha(G)) = r$ the result follows. \square

Proposition 3.12. *Let G be a graph with n vertices and $m \neq 0$ edges. Then, $\lambda_1(A_\alpha(l(G))) \leq 2n - 4$ and $\lambda_i(A_\alpha(l(G))) \leq n(\alpha + 1) - 4, \forall i = 2, \dots, m$ and $\alpha \in [0, 1]$.*

Proof. As G is a subgraph of K_n , the result follows from Corollary 2.17 and Example 1. \square

Corollary 3.13. *Let G be a r -regular graph with $n \geq 2$ vertices, $m \neq 0$ edges and $\alpha \in [0, 1]$. Then, $\lambda_1(A_\alpha(l(G))) = 2n - 4$ if and only if $G \cong K_n$.*

Proof. If $G \cong K_n$, from Example 1, $\lambda_1(A_\alpha(l(K_n))) = 2n - 4$. On the other hand, if $\lambda_1(A_\alpha(l(G))) = 2n - 4$ we have that $P_{A_\alpha(l(G))}(2n - 4) = 0$ and, from Theorem 2.28, we obtain that $(2n - 2 - 2r\alpha)^{m-n} P_{A_\alpha(G)}(2n - 2 - r) = 0$, for all $\alpha \in [0, 1]$. Now, consider two cases:

- (i) Suppose $2n - 2 - 2r\alpha = 0$. If $0 < \alpha < 1$, then $r = \frac{n-1}{\alpha} > n - 1$, which is impossible. If $\alpha = 0$, then $n = 1$ which is absurd.
- (ii) Suppose $P_{A_\alpha(G)}(2n - 2 - r) = 0$. As G is a regular graph the multiplicity of $\lambda_1(A_\alpha(G)) = r$ is 1. So we have that $2n - 2 - r = r$. Therefore, $r = n - 1$, which completes the proof.

\square

Finally, next theorem presents bounds for $\lambda_2(A_\alpha(l(G)))$.

Theorem 3.14. *Let G be a connected graph with n vertices, $n \geq 3$. Then*

$$2\alpha - 1 \leq \lambda_2(A_\alpha(l(G))) \leq n(\alpha + 1) - 4.$$

Equality occurs when $G \cong P_3$, for the lower bound, and when $G \cong K_n$, for the upper bound.

Proof. As $n \geq 3$, from Proposition 2.16 we have that $l(P_3)$ is subgraph of $l(G)$. From Theorem 2.3 and Corollary 2.27 we have that $2\alpha - 1 \leq \lambda_2(A_\alpha(l(G)))$ and then we can conclude the lower bound. If $G \cong P_3$, from Corollary 2.27, the equality is achieved. The upper bound and its equality follows straightforwardly from Proposition 3.12. \square

Acknowledgments

The research of C. S. Oliveira is supported by the CNPq Grant 304548/2020-0.

References

- [1] T.H. Cormen, C.E. Leiserson, R.L. Rivest, and C. Stein. Introduction To Algorithms. Mit Electrical Engineering and Computer Science. MIT Press, 2001.
- [2] I. Gutman and N. Trinajstić. Graph theory and molecular orbitals. total ϕ -electron energy of alternant hydrocarbons. Chemical Physics Letters, 17(4):535–538, 1972.
- [3] Xueliang Li and Haixing Zhao. Trees with the first three smallest and largest generalized topological indices. MATCH - Communications in Mathematical and in Computer Chemistry, 50:57–62, 02 2004.
- [4] Xueliang Li and Jie Zheng. A unified approach to the extremal trees for different indices. MATCH - Communications in Mathematical and in Computer Chemistry, 54:195–208, 01 2005.
- [5] Sonja Nikolić, Goran Kovačević, Ante Miličević, and Nenad Trinajstić. The zagreb indices 30 years after. Croatica Chemica Acta, 76:113–124, 06 2003.
- [6] Kinkar Ch. Das. Sharp bounds for the sum of the squares of the degrees of a graph. Kragujevac Journal of Mathematics, 25(25):19–41, 2003.
- [7] Kinkar Ch. Das. Maximizing the sum of the squares of the degrees of a graph. Discrete Mathematics, 285(1):57–66, 2004.

- [8] Sebastian M. Cioabă. Sums of powers of the degrees of a graph. Discrete Mathematics, 306(16):1959–1964, 2006.
- [9] Kinkar Das, Kexiang Xu, and Junki Nam. Zagreb indices of graphs. Frontiers of Mathematics in China, 10:567–582, 03 2015.
- [10] Boris Furtula and Ivan Gutman. A forgotten topological index. Journal of Mathematical Chemistry, 53:1184–1190, 04 2015.
- [11] Muhammad Javaid, Saira Javed, Saima Memon, and Abdulaziz Alanazi. Forgotten index of generalized operations on graphs. Journal of Chemistry, 2021:1–14, 05 2021.
- [12] Béla Bollobás and Paul Erdős. Graphs of extremal weights. Ars Comb., 50, 1998.
- [13] Bo Zhou and Wei Luo. A note on general randić index. MATCH - Communications in Mathematical and in Computer Chemistry, 62:155–162, 01 2009.
- [14] V. Nikiforov. Merging the A- and Q- spectral theories. Applicable Analysis and Discrete Mathematics, 11(1):81–107, 2017.
- [15] R.A. Horn and C.R. Johnson. Matrix Analysis. Matrix Analysis. Cambridge University Press, New York, 2013.
- [16] D. Cvetković, P. Rowlinson, and S. Simić. An Introduction to the Theory of Graph Spectra. London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 2010.
- [17] L.Yu. Kolotilina. Lower bounds for the perron root of a nonnegative matrix. Linear Algebra and its Applications, 180:133–151, 1993.
- [18] Yanqing Chen and Ligong Wang. Sharp bounds for the largest eigenvalue of the signless laplacian of a graph. Linear algebra and its applications, 433(5):908–913, 2010.
- [19] M.N. Ellingham and Xiaoya Zha. The spectral radius of graphs on surfaces. Journal of Combinatorial Theory, Series B, 78(1):45–56, 2000.
- [20] L.W. Beineke and J.S. Bagga. Line Graphs and Line Digraphs. Developments in Mathematics. Springer International Publishing, 2021.
- [21] Chen Yan. Properties of spectra of graphs and line graphs. Applied Mathematics: A Journal of Chinese Universities, 17:371–376, 2002.
- [22] Dragoš Cvetković, Peter Rowlinson, and Slobodan K. Simić. Signless laplacians of finite graphs. Linear Algebra and its Applications, 423(1):155–171, 2007. Special Issue devoted to papers presented at the Aveiro Workshop on Graph Spectra.
- [23] Dragos M. Cvetković, Peter Rowlinson, and Slobodan K. Simic. Eigenvalue bounds for the signless laplacian. Publications De L’institut Mathématique, 81:11–27, 2007.
- [24] João Domingos G. da Silva Junior, Carla Silva Oliveira, and Liliana Manuela G. C. da Costa. On the characteristic polynomial of the A_α -matrix for some operations of graphs. Computational and Applied Mathematics, 42(5), 2023.
- [25] Domingos M. Cardoso, Paula Carvalho, Maria Agueiras A. de Freitas, and Cybele T.M. Vinagre. Spectra, signless laplacian and laplacian spectra of complementary prisms of graphs. Linear Algebra and its Applications, 544:325–338, 2018.
- [26] Dragoš Cvetkovic, Peter Rowlinson, and Slobodan Simic. Spectral Generalizations of Line Graphs: On Graphs with Least Eigenvalue -2. London Mathematical Society Lecture Note Series. Cambridge University Press, 2004.
- [27] Malathy and Kalyani Desikan. Bounds for laplacian spectral radius of the complete bipartite graph. International Journal of Pure and Applied Mathematics, 115(9):343 – 351, 2017.
- [28] Shuang Zhang and Yan Zhu. Some spectral properties of A_α -matrix. Discrete Mathematics, Algorithms and Applications, 11(6):1–7, 2019.
- [29] Vladimir Nikiforov, Germain Pastén, Oscar Rojo, and Ricardo L. Soto. On the A_α -spectra of trees. Linear Algebra and its Applications, 520:286–305, 2017.
- [30] A.M. Marr and W.D. Wallis. Magic Graphs. Springer New York, 2014.