## **Emanuel Herrendorf**

Philipps-Universität Marburg, Fachbereich Mathematik und Informatik, Marburg, Germany

## Christian Komusiewicz 🖂 🗈

Friedrich-Schiller-Universität Jena, Fakultät für Mathematik und Informatik, Jena, Germany

## Nils Morawietz 🖂 🗈

Friedrich-Schiller-Universität Jena, Fakultät für Mathematik und Informatik, Jena, Germany

#### Frank Sommer $\square$

Friedrich-Schiller-Universität Jena, Fakultät für Mathematik und Informatik, Jena, Germany

#### **— Abstract**

Network sparsification is the task of reducing the number of edges of a given graph while preserving some crucial graph property. In community-aware network sparsification, the preserved property concerns the subgraphs that are induced by the communities of the graph which are given as vertex subsets. This is formalized in the II-NETWORK SPARSIFICATION problem: given an edge-weighted graph G, a collection C of c subsets of V(G), called communities, and two numbers  $\ell$  and b, the question is whether there exists a spanning subgraph G' of G with at most  $\ell$  edges of total weight at most b such that G'[C] fulfills  $\Pi$  for each community  $C \in C$ . In this work, we consider two graph properties  $\Pi$ : the connectivity property and the property of having a spanning star. The corresponding problems are called CONNECTIVITY NWS and STARS NWS. Since both problems are NP-hard, we study their parameterized and fine-grained complexity.

First, we provide a tight  $2^{\Omega(n^2+c)} \cdot \operatorname{poly}(n+|\mathcal{C}|)$ -time running time lower bound based on the ETH for both problems, where n is the number of vertices in G. The lower bound holds even in the restricted case when all communities have size at most 4, G is a clique, and every edge has unit weight. For the connectivity property, the unit weight case with G being a clique is the well-studied problem of computing a hypergraph support with a minimum number of edges. We then study the complexity of both problems parameterized by the feedback edge number t of the solution graph G'. For STARS NWS, we present an XP-algorithm for t. This answers an open question by Korach and Stern [Discret. Appl. Math. '08] who asked for the existence of polynomial-time algorithms for t = 0. Our result implies polynomial-time algorithms for all constant values of t. In contrast, we show for CONNECTIVITY NWS that known polynomial-time algorithms for t = 0 [Korach and Stern, Math. Program. '03; Klemz et al., SWAT '14] cannot be extended to larger values of t by showing that CONNECTIVITY NWS is NP-hard for t = 1.

**2012 ACM Subject Classification** Theory of computation  $\rightarrow$  Parameterized complexity and exact algorithms; Theory of computation  $\rightarrow$  Graph algorithms analysis

Keywords and phrases parameterized complexity, hypergraph support, feedback edge number, exponential-time-hypothesis

Acknowledgements Some of the results of this work are also contained in the first author's Master thesis [27].

## 1 Introduction

A common goal in network analysis is to decrease the size of a given network to speed up downstream analysis algorithms or to decrease the memory footprint of the graphs. This leads to the task of network sparsification where one wants to reduce the number of edges of a network while preserving some important property  $\Pi$  [8, 36, 41]. Similarly, in network

design the task is often to construct a minimum-size or minimum-weight network fulfilling a given property, the most famous example being MINIMUM-WEIGHT SPANNING TREE.

In many applications the input contains, in addition to a network, a hypergraph on the same vertex set [20, 28, 37]. The hyperedges of this hypergraph represent, for example, communities that are formed within the network. In presence of such community data, the sparsified network should preserve a property not for the whole network but instead for each community, that is, for each hyperedge of the hypergraph. This leads to the family of *community-aware network sparsification* problems, subsumed by the following problem introduced by Gionis et al. [23].<sup>1</sup>

Π-NETWORK SPARSIFICATION (Π-NWS)

**Input:** A graph G, a collection  $\mathcal{C}$  of c subsets of V(G), called *communities*, an edge-weight function  $\omega : E(G) \to \mathbb{R}_+$ , an integer  $\ell$ , and a positive real number b. **Question:** Is there a graph G' = (V(G), E') with  $E' \subseteq E(G)$ ,  $|E'| \leq \ell$ , and total edge weight at most b such that for all communities  $C_i \in \mathcal{C}$  the subgraph of G induced by  $C_i$  satisfies  $\Pi$ ?

We say that a graph G' fulfilling the requirements is a solution for the instance I. An example instance of  $\Pi$ -NWS and solutions for the two specific properties  $\Pi$  studied in this work is given in Figure 1. A very well-studied property  $\Pi$ , considered by Gionis et al. [23] but also in many previous works [2, 9, 16, 18, 32], in this context is that every community should induce a connected subgraph. When a graph G admits this property for some hypergraph H, then G is called a support for H [6, 7, 32]. We denote the corresponding special case of  $\Pi$ -NWS as CONNECTIVITY NWS. Another variant of  $\Pi$ -NWS, also studied by Gionis et al. [23], is to demand that every community not only induces a connected subgraph but more strongly that it contains a spanning star. In other words, in the solution graph G', every community must be contained in the neighborhood of at least one of its vertices, called a center vertex. We refer to this variant as STARS NWS.

CONNECTIVITY NWS and STARS NWS are both NP-hard [16, 12, 11, 23]. Motivated by this, we study both problems in terms of their parameterized and fine-grained complexity. We also investigate the versions of both problems where each edge has unit weight and refer to them as UNWEIGHTED CONNECTIVITY NWS and UNWEIGHTED STARS NWS.

Our two main results are as follows:

- We show that, based on the Exponential Time Hypothesis (ETH), CONNECTIVITY NWS and STARS NWS do not admit algorithms with running time  $2^{o(n^2)+c} \operatorname{poly}(n+c)$ , even if the input graph is a clique with unit weights and each community has size at most 4. This bound is matched by simple brute-force algorithms.
- We show that STARS NWS admits an XP-algorithm when parameterized by t, the feedback edge number of the solution graph. This positively answers the question of Korach and Stern [34] who asked whether there is a polynomial-time algorithm for finding an optimal solution for STARS NWS that is a tree. In fact, our algorithm extends the polynomial-ime solvable cases to solutions that are tree-like.

We obtain several further results, for example a complexity dichotomy for STARS NWS and UNWEIGHTED STARS NWS parameterized by c, the number of communities.

<sup>&</sup>lt;sup>1</sup> Compared to Gionis et al. [23], we consider the more general scenario where we simultaneously constrain the edge number and total edge weight of the solution graph.



**Figure 1** *a*) The communities (depicted in blue) and the input graph of an instance of II-NWS with unit weights. *b*) and *c*) Optimal solutions (in red) for UNWEIGHTED CONNECTIVITY NWS, and UNWEIGHTED STARS NWS, respectively.

**Known Results.** Already the most basic variant of CONNECTIVITY NWS, where the edges have unit weights and the input graph G is a clique, appears in many applications, ranging from explanation of protein complexes [37] to combinatorial auctions [12] to the construction of P2P overlay networks in publish/subscribe systems [10, 28]. Consequently, the problem has been studied intensively under various names [2, 9, 10, 16, 18, 28] from a parameterized complexity [9, 18, 28] and an approximation algorithms [2, 10, 28] perspective. For example, Du and Miller [16] showed NP-hardness even for instances with maximum community size 3, and Chen et al. [9] presented an FPT-algorithm for the number of communities and an FPT-algorithm for the largest community size plus the feedback edge number t of a solution. From a more practical perspective, (Mixed) Integer Linear Programming formulations were proposed [5, 14]. A particular restriction of the problem is to determine whether there is an acyclic solution. Such solutions are called *tree supports* or *clustered spanning trees* and hypergraphs that have a tree support are also known as *hypertrees*. It can be determined in polynomial-time whether a hypergraph is a hypertree and different polynomial-time algorithms have been described over the years [3, 12, 17, 19, 24, 31, 38, 39].

UNWEIGHTED CONNECTIVITY NWS, with unit weights but with general input graphs G, has applications in the context of placing green bridges [20, 21, 26]. UNWEIGHTED CONNECTIVITY NWS is NP-hard even when the maximum degree of G is 3 [26] and even for seven communities [21]. On the positive side, one can construct in polynomial time a tree support if one exists [25, 32, 33].

For CONNECTIVITY NWS where we may have arbitrary edge-weights, the distinction whether or not G is restricted to be a clique vanishes: any non-clique input graph G may be transformed into a clique by adding the missing edges with a prohibitively large edge weight. The problem of finding a minimum-weight tree support received attention due to its applications in network visualization [32]. As shown by Korach and Stern [33] and Klemz et al. [32], one can compute minimum-weight tree supports in polynomial time. Gioinis et al. [23] provided approximation algorithms for the general problem.

STARS NWS has received less attention than CONNECTIVITY NWS. Gionis et al. [23] showed NP-hardness and provided approximation algorithms. Korach and Stern [34] studied a variant of STARS NWS where the input graph is a clique and the solution is constrained to be a tree T where the closed neighborhood of the center vertex of a community  $C_i$  is exactly the community  $C_i$ . This implies that two different communities need to have different center vertices and thus restricts the allowed set of solution graphs strictly compared to STARS NWS. Korach and Stern [34] showed that this problem is solvable in polynomial time [34]. As an open question, they ask whether this positive result can be lifted to STARS NWS.

Cohen et al. [11] studied the MINIMUM  $\mathcal{F}$ -OVERLAY problem which can be viewed as

the following special case of  $\Pi$ -NWS: The input graph G is a clique and all edges have unit weight;  $\mathcal{F}$  is a family of graphs and the property  $\Pi$  is to have some spanning subgraph which is contained in  $\mathcal{F}$ . It should be noted that CONNECTIVITY NWS and UNWEIGHTED STARS NWS with clique input graphs are special cases of MINIMUM  $\mathcal{F}$ -OVERLAY. Cohen et al. [11] provide a complexity dichotomy with respect to properties of  $\mathcal{F}$ . For most cases of  $\mathcal{F}$ , MINIMUM  $\mathcal{F}$ -OVERLAY is NP-hard. In particular, the dichotomy of Cohen et al. [11] shows that UNWEIGHTED STARS NWS is NP-hard even when G is a clique. Gionis et al. [23] also studied a third property  $\Pi$  where each community needs to induce a subgraph exceeding some prespecified density. They showed NP-hardness and provided some approximation algorithms. Fluschnik and Kellerhals [20] considered several further graph properties  $\Pi$ , for example the property of having small diameter.

Our Results and Organization of the Work. In order to put our main results into context, we first summarize in Section 2 some hardness and tractability results that follow either from simple observations or from previous work. They imply in particular that STARS NWS and CONNECTIVITY NWS have an FPT-algorithm for the parameter solution size  $\ell$  and that they are W[1]-hard with respect to  $k := m - \ell$ , the number of edges not in the solution even in the unit weight case when G is a clique.

Then, in Section 3 we show that UNWEIGHTED CONNECTIVITY NWS and UNWEIGHTED STARS NWS do not admit algorithms with running time  $2^{o(n^2+c)} \cdot \operatorname{poly}(n+c)$  even when G is a clique. This running time lower bound is based on the Exponential Time Hypothesis (ETH) [30]. To show the bound, we develop a compression of 3-SAT instances  $\phi$  into graphs with  $\mathcal{O}(\sqrt{|\phi|})$  vertices.

In Section 4, we then consider parameterization by t, the feedback edge number of the solution graph G'. This is the minimum number of edges that need to be deleted to transform the solution into a forest.<sup>2</sup> The study of this parameter is motivated by the following observation: The solution size parameter  $\ell$  is essentially at least as large as n - 1, and thus neither small in practice nor particularly interesting from an algorithmic point of view. The parameter t can thus essentially be seen as a parameterization above the lower bound n - 1. Our first main result for the parameter t is an XP-algorithm for STARS NWS parameterized by t. Our result positively answers the question of Korach and Stern [34] who asked whether there is a polynomial-time algorithm for t = 0 and substantially extends this tractability further to every constant value of t. We then consider the parameter t for CONNECTIVITY NWS. We show that UNWEIGHTED CONNECTIVITY NWS is NP-hard already if t = 1. Thus, the polynomial-time algorithms of Korach and Stern [33] and Klemz et al. [32] for t = 0 cannot be lifted to larger values of t.

Finally, in Section 5 we study the complexity of STARS NWS with respect to the number c of input communities. The problem is easily seen to be in XP via an  $n^{\mathcal{O}(c)}$ -time algorithm. In light of this, we obtain the following dichotomy: UNWEIGHTED STARS NWS is FPT with respect to c and STARS NWS is W[1]-hard in the most restricted case when G is a clique and all edges have weight 1 or 2.

For an overview of the parameterized complexity results, refer to Table 1.

<sup>&</sup>lt;sup>2</sup> The parameter t can be computed in polynomial time as discussed in Section 2.

**Table 1** An overview of the parameterized complexity results. A ‡ indicates that this result also holds in the unweighted case and a † indicates that this result only holds in the unweighted case.

Parameter	Stars NWS	Connectivity NWS
l	FPT (Proposition 2.2, [20]), no polynomial kernel <sup>‡</sup> (Proposition 2.4, [20])	
$k\coloneqq m-\ell$	W[1]-hard <sup>‡</sup> (Proposition 2.4)	
t	XP (Theorem 4.1)	P for $t = 0$ ([32], Theorem 4.11)
		NP-h for $t = 1^{\ddagger}$ (Theorem 4.13)
с	$FPT^{\dagger}$ (Theorem 5.2)	
	no polynomial kernel <sup>‡</sup> (Theorem $5.3$ )	NP-h for $c = 7^{\ddagger}$ ([21])
	W[1]-h (Theorem 5.1)	
Δ	NP-h for $\Delta = 6^{\ddagger}$ (Corollary 3.4)	NP-h for $\Delta = 3^{\ddagger}$ ([26])

## 2 Preliminaries and Basic Observations

**Preliminaries.** For a set X, we denote by  $\binom{X}{2}$  the collection of all size-two subsets of X. Moreover, for positive integers i and j with  $i \leq j$ , we denote by  $[i, j] := \{k \in \mathbb{N} : i \leq k \leq j\}$ .

An undirected graph G = (V, E) consists of a set of vertices V and a set of edges  $E \subseteq \binom{V}{2}$ . We denote by V(G) and E(G) the vertex and edge set of G, respectively. Furthermore, we let  $n = n(G) \coloneqq |V(G)|$  and  $m \coloneqq |E(G)|$ . For an edge  $e = \{u, v\} \in E$ , we say that uand v are adjacent and that e is incident with u and v. For a vertex set  $V' \subseteq V$ , we denote by  $E_G(V') \coloneqq \{\{u, v\} \in E : u \in V', v \in V'\}$  the edges between the vertices of V'in G. If G is clear from the context, we may omit the subscript. A graph G' is a subgraph of G if  $V(G') \subseteq V(G)$  and  $E(G') \subseteq E(G)$ . Moreover, a subgraph G' of G is spanning if V(G') = V(G). For a vertex set V', we denote by  $G[V'] \coloneqq (V', E_G(V'))$  the subgraph of G induced by V'. For a graph G, a set  $S \subseteq V(G)$  with  $E(S) = \binom{S}{2}$  is called a clique. A size-three clique is also called triangle. A set  $X \subseteq V(G)$  is an independent set of G if the vertices of X are pairwise non-adjacent in G, that is, if  $E_G(X) = \emptyset$ . A graph G is a star of size n-1 with center  $z \in V(G)$  if  $E(G) = \{\{z, v\} : v \in V(G) \setminus \{z\}\}$ . We say that G contains a spanning star if some subgraph G' of G is a star of size n-1. The center of this star is universal for G.

A sequence of distinct vertices  $(v_1, \ldots, v_{\kappa})$  of a graph G is a  $(v_1, v_{\kappa})$ -path, if  $\{v_i, v_{i+1}\} \in$ E(G) for each  $i \in [1, \kappa - 1]$ . Let  $(v_1, \ldots, v_{\kappa})$  be a path with  $\kappa \geq 3$  and  $\{v_1, v_{\kappa}\} \in E(G)$ , then we call  $(v_1, \ldots, v_\kappa)$  a *cycle* in G. A graph without a cycle is called *acyclic*. An edge set  $E' \subseteq E(G)$  is a feedback edge set of G, if the graph  $G' := (V(G), E(G) \setminus E')$  is acyclic. We say that two vertices u and v are *connected* in G if G contains a path between u and v. A graph G is connected if each pair of vertices  $u, v \in V(G)$  is connected. A set  $S \subseteq V(G)$  is a connected component of G, if G[S] is connected and S is inclusion-maximal with this property. Connectivity in hypergraphs is defined similarly: Two vertices u and v are *connected* if there exists a sequence  $C_1, C_2, \ldots, C_p$  of hyperedges, such that  $u \in C_1, v \in C_p$ , and consecutive communities have nonempty intersection. A connected component of this hypergraph is a maximal set of vertices which are connected. We say that the hypergraph is *connected* if it has is exactly one connected component. The number x of connected components of a hypergraph can be computed in polynomial time, for example by BFS. Observe that for a minimal solution graph G' for STARS NWS and CONNECTIVITY NWS, the connected components of G' are exactly the connected components of the community hypergraph. Thus,  $t = \ell - n + x$  and the parameter t can be computed in polynomial time for a given input instance.

For details about parameterized complexity and the exponential-time-hypothesis (ETH), we refer to the standard monographs [15, 13].

**Basic Observations.** To put our main results for CONNECTIVITY NWS and STARS NWS into context, we state some results that either follow easily from previous work or from simple observations.

The naive brute-force approach for each  $\Pi$ -NWS is to perform an exhaustive search over the  $\mathcal{O}(2^m)$  possibilities to select at most  $\ell$  edges from the input graph G. This leads to the following general statement for  $\Pi$ -NWS problems.

▶ **Proposition 2.1.** Let  $\Pi$  be a property which can be decided in poly(n) time. Then,  $\Pi$ -NWS is solvable in  $2^m \cdot c \cdot poly(n)$  time.

For the solution size parameter  $\ell$ , one can obtain the following running time.

▶ **Proposition 2.2.** CONNECTIVITY NWS and STARS NWS can be solved in  $\ell^{\mathcal{O}(\ell)} \cdot \operatorname{poly}(n+c)$  time.

**Proof.** We may use the following branching strategy first observed by Fan et al. [18] for the special case of CONNECTIVITY NWS when G is a clique and the community size is at most d. Later, Cohen et al. [11] described the same branching strategy for some special cases of MINIMUM F OVERLAY including UNWEIGHTED STARS NWS with clique graph G. We recall the strategy for sake of completeness. Start with an initially empty partial solution. If some community is not yet connected by the current partial solution, then branch into the at most  $\binom{d}{2}$  possibilities to add some edge with both endpoints in the community. The depth of the branching is  $\ell$ , the overall search tree size  $d^{\mathcal{O}(\ell)}$ . The branching can be naturally adapted to UNWEIGHTED STARS NWS by branching into the choice of one of d centers for communities without a star. Now observe that instances with a community of size at least  $\ell + 2$  are no-instances for CONNECTIVITY NWS and STARS NWS. Thus, we may safely assume that  $d \leq \ell + 1$ .

The fixed-parameter tractability of UNWEIGHTED CONNECTIVITY NWS with respect to  $\ell$  was also shown by Fluschnik and Kellerhals via a kernelization that gives kernels with at most  $2\ell$  vertices [20].

A further natural parameter that can be considered is  $k := m - \ell$ , the number of edges that are not in the solution. The following proposition summarizes the complexity of the problems with respect to this parameter.

 ▶ Proposition 2.3 (). ■ CONNECTIVITY NWS and STARS NWS are NP-hard for k = 0.
 ■ UNWEIGHTED CONNECTIVITY NWS and UNWEIGHTED STARS NWS can be solved in n<sup>2k</sup> · poly(n) time and are W[1]-hard with respect to k even if G is a clique and if each community has size at most 3.

**Proof.** For CONNECTIVITY NWS and STARS NWS the NP-hardness for k = 0 can be seen as follows: reduce from the unweighted version, set the weight of each edge to one,  $k \coloneqq 0$ , and the weight bound b to the solution size bound of the unweighted instance.

For UNWEIGHTED CONNECTIVITY NWS and UNWEIGHTED STARS NWS the existence of a solution with at most  $\ell$  edges implies the existence of one with exactly  $\ell$  edges. Thus, it is sufficient to consider all  $\binom{m}{k}$  possibilities to remove k edges from G.

The W[1]-hardness for UNWEIGHTED CONNECTIVITY NWS can be seen via an adaption of reduction from Du and Miller [16, Theorem 3.1]. More precisely, we reduce from INDE-PENDENT SET where we are given a graph  $\tilde{G}$  on  $\tilde{n}$  vertices and  $\tilde{m}$  edges and integer  $\tilde{k}$  and

want to decide whether it contains a set of at least  $\tilde{k}$  pairwise nonadjacent vertices. The graph G of the constructed UNWEIGHTED CONNECTIVITY NWS instance is a clique on the vertex set  $V(\tilde{G}) \cup \{x\}$  where x is a new vertex. For each pair u, v of vertices in  $V(\tilde{G})$ , add a community  $\{u, v\}$ . Then, for each edge  $\{u, v\} \in E(\tilde{G})$ , add a community  $\{u, v, x\}$ . Finally, set the budget  $\ell$  to  $\binom{\tilde{n}}{2} + \tilde{n} - \tilde{k}$ . Since the total number of edges in the constructed graph G is  $\binom{\tilde{n}}{2} + \tilde{n}$ , we have  $k = m - \ell = \tilde{k}$ . The equivalence of the instance can be seen as follows: every solution contains all  $\tilde{n}^2$  edges between vertices of  $V(\tilde{G})$ . Thus, every solution misses at least k edges that are incident with x. These edges correspond to a vertex set  $S \subseteq V(G)$  of size at least k. The set S is an independent set in  $\tilde{G}$ : if S contains two adjacent vertices u and v, then the community  $\{u, v, x\}$  does not induce a connected subgraph. The converse direction can be seen analagously.

The results for UNWEIGHTED STARS NWS follow since for communities of size at most 3 the properties of being connected and having a spanning star coincide.

By a reduction from VERTEX COVER, Du and Miller [16] showed that the special case of UNWEIGHTED CONNECTIVITY NWS where G is a clique is NP-hard even if restricted to communities of size at most 3. For UNWEIGHTED CONNECTIVITY NWS, Fluschnik and Kellerhals showed that a polynomial kernel for  $\ell$  and (thus for n) is unlikely, even on planar series-parallel graphs [21]. This result can be also given for the case when the input graph G is a clique.

▶ Proposition 2.4. UNWEIGHTED CONNECTIVITY NWS and UNWEIGHTED STARS NWS do not admit a polynomial kernel for n even if G is a clique, unless  $NP \subseteq coNP/poly$ .

**Proof.** We give a proof for UNWEIGHTED CONNECTIVITY NWS, the result for UNWEIGHTED STARS NWS follows again since for communities of size at most 3 the two properties coincide.

To show the claim we give a polynomial-parameter reduction from HITTING SET parameterized by the size of the universe to UNWEIGHTED CONNECTIVITY NWS parameterized by n. HITTING SET parameterized by the size of the universe is known to not admit a polynomial kernel unless NP  $\subseteq$  coNP/poly [13]. This reduction is again an adaption of the VERTEX COVER reduction of Du and Miller [16]. In HITTING SET we are given a set family  $\mathcal{F}$  over a universe U and an integer k and ask if there is a set  $S \subseteq U$  of size at most k such that for each set  $F \in \mathcal{F}$ , we have  $S \cap F \neq \emptyset$ . For a given HITTING SET instance  $(\mathcal{F}, U, k)$  we construct a graph G with communities as follows. The graph G is a clique with vertex set  $U \cup \{x\}$  for some  $x \notin U$ . For each pair of elements  $u, v \in U$ , add a community  $\{u, v\}$ . Moreover, for each set  $F \in \mathcal{F}$ , add the community  $F \cup \{x\}$ . Finally, set the budget  $\ell$  to  $\binom{|U|}{2} + k$ . Observe that |V(G)| = |U| + 1.

The correctness of the reduction follows from the fact that all edges between vertices of U are fixed, that is, are contained in a community of size 2, and the up to k edges that are added between x and U must have one endpoint in F for each  $F \in \mathcal{F}$ . Thus, these edges correspond to a solution for  $(\mathcal{F}, U, k)$ .

We can also show that the simple brute-force algorithm behind Proposition 2.1 cannot be improved substantially.

▶ **Proposition 2.5.** If the ETH is true, then UNWEIGHTED STARS NWS and UNWEIGHTED CONNECTIVITY NWS cannot be solved in  $2^{o(n+m+c)} \cdot poly(n+c)$  time, even if restricted to instances with community size at most 3.

**Proof.** This can again be shown by adaption of the reduction of Du and Miller [16]. The constructed graph is not a clique; instead it consists of the graph G of the VERTEX COVER



**Figure 2** Sketch of the construction of Theorem 3.1. The communities are blue (solid, dashed and dotted). We only show edges which are contained in at least one community and only some fixed edges (red). a) Part of the variable gadget for  $x_1$  and  $x_2$ . b) The variable communities for a clause  $q = x_1 \vee \overline{x_2} \vee x_3$ . c) The assignment gadget for the first literal  $x_1$  of the clause q. Here, the red edges are the fixed edges with one endpoint in the variable gadget and one in the clause gadget.

instance plus the new universal vertex x. Now for each edge  $\{u, v\} \in E(G)$ , add the two communities  $\{u, v\}$  and  $\{u, v, x\}$ . Finally, we set  $\ell := m + k$ . Since the size of the resulting graph and the number of communities is linear in the size of G, and VERTEX COVER cannot be solved in time  $2^{o(|V(G)|+|E(G)|)}$  if the ETH is true, we obtain the statement.

## 3 A Stronger ETH-Bound

In Proposition 2.5 we observed that algorithms with running time  $2^{o(n+m+c)} \cdot \operatorname{poly}(n+c)$  for UNWEIGHTED CONNECTIVITY NWS and UNWEIGHTED STARS NWS would violate the ETH. We now provide a stronger  $2^{\Omega(n^2+c)} \cdot \operatorname{poly}(n+c)$ -time lower bound for both problems. Notably, this lower bound also applies to the case when all communities have constant size.

## 3.1 ETH Lower Bound for Unweighted Stars NWS

First, we present the lower bound for UNWEIGHTED STARS NWS.

▶ **Theorem 3.1.** If the ETH is true, then UNWEIGHTED STARS NWS cannot be solved in  $2^{o(n^2+c)} \cdot \operatorname{poly}(n+c)$  time, even if G is a clique and if each community has size at most 4.

**Proof.** We present a reduction from 3-SAT to UNWEIGHTED STARS NWS where G is a clique with maximum community size 4 such that the resulting instance has  $\mathcal{O}(\sqrt{|\phi|})$  vertices and  $\mathcal{O}(|\phi|)$  communities, where  $\phi$  denotes the total formula length. Then, the existence of an  $2^{o(n^2+c)} \cdot \text{poly}(n+c)$ -time algorithm for UNWEIGHTED STARS NWS implies the existence of a  $2^{o(|\phi|)}$ -time algorithm for 3-SAT violating the ETH [29, 30]. The input formula  $\phi$  is over the variable set X and each clause  $q \in \Gamma$  contains exactly three literals. For a literal y, we denote by  $\overline{y}$  its complement. A visualization of the construction is given in Figure 2. In all gadgets, we add several communities of size 2. These communities enforce that each solution has to contain the edge of this community. In the following we call such edges *fixed*.

Variable Gadget: We start by describing the construction of the variable gadget  $G_X$ . Recall that  $G_X$  is a clique. The idea is to create for each variable a community C of size 3

with one fixed edge. The two remaining edges of C are referred to as selection edges. The idea is that each solution contains exactly one selection edge of C. One selection edge represents the positive literal, the other one represents the negative literal. The fixed edge of the triangle is used to model that one literal must be set to **true**. The selection edges are arranged compactly, to guarantee that  $|V(G_X)| \in \mathcal{O}(\sqrt{|\phi|})$ . In the following, we describe the graph  $G_X$  together with communities fulfilling the above-described properties. An example of a variable gadget is shown in part a) of Figure 2.

Let  $V(G_X) = U \cup P$  where  $U \coloneqq \{u_1, \ldots, u_{n_x}\}$ ,  $P = P_1 \cup P_2$ , and  $P_i \coloneqq \{p_1^i, \ldots, p_{n_x}^i\}$ for  $i \in [2]$  consist of  $n_x = \lceil \sqrt{|X|} \rceil$  vertices each. It remains to describe the communities: For each variable  $x \in X$ , we add a community  $C_x \coloneqq \{u_j, p_s^1, p_s^2\}$  for  $j, s \in [n_x]$ . This is possible since  $n_x \cdot n_x \ge |X|$ . We refer to these communities as the variable communities  $C^X$ . Afterwards, we set  $\theta(x) \coloneqq \{u_j, p_s^1\}$  and  $\theta(\overline{x}) \coloneqq \{u_j, p_s^2\}$  to assign the positive and negative literal of x to an edge of the variable gadget. Now, we fix the edges of  $G[P] = G[P_1 \cup P_2]$ . Recall that this means that for each edge  $\{p_{j_1}^{i_1}, p_{j_2}^{i_2}\}$  having both endpoints in  $P_1 \cup P_2$ , we add a community  $\{p_{j_1}^{i_1}, p_{j_2}^{i_2}\}$ .

Note that edges with both endpoints in U are not contained in any community. We only add these edges to ensure that G is a clique.

Before describing the other gadgets, let us observe that each selection edge is indeed used in at most one variable gadget.

▶ Claim 1 (\*). Each selection edge of  $E(G_X)$  is contained in only one subgraph induced by a variable community in  $C^X$ .

Proof. Each variable community  $C^x$  consists of one vertex  $u \in U$  and two vertices  $p_i, p_j \in P$ where  $\{u, p_i\}$  and  $\{u, p_j\}$  are the selection edges. Suppose there are two distinct variable communities  $C_1$  and  $C_2$  such that they have at least one selection edge in common. This implies that  $C_1$  and  $C_2$  contain the same vertex of U and have at least one vertex of P in common. By construction, in each variable community  $C_i$  the two vertices of  $P_1 \cup P_2$  are either identical or there is no overlap. Thus,  $C_1$  and  $C_2$  are identical, a contradiction.

Clause Gadget: We continue by describing the construction of the clause gadget  $G_{\Gamma}$ . The idea is that each clause is represented by four vertices of  $V(G_{\Gamma})$  in which a triangle is *fixed*. All three remaining edges of this size-4 clique are referred to as *free*. Note that these free edges form a star with three leaves. Each free edge represents one literal of the clause. For each pair containing two of these three edges, we then create a community containing the three endpoints of these two edges. As in the vertex gadget, these induced subgraphs are arranged compactly, to achieve a clause gadget with  $|V(G_{\Gamma})| \in \mathcal{O}(\sqrt{|\Gamma|})$ . In the following, we describe the graph  $G_{\Gamma}$  together with communities fulfilling these properties. Part b) of Figure 2 shows an example for the representation of a clause.

Let  $V(G_{\Gamma}) = Y \cup Z$  where  $Y = \{y_1, \ldots, y_{n_c}\}, Z = Z_1 \cup Z_2 \cup Z_3$ , and  $Z_i = \{z_1^1, \ldots, z_{n_c}^i\}$ for  $i \in [3]$  consist of  $n_c = \lceil \sqrt{|\Gamma|} \rceil$  vertices each. In the following, we assign each clause to a clique of  $G_{\Gamma}$  having vertex set  $y_j, z_s^1, z_s^2, z_s^3$  for  $j, s \in [n_c]$ . This is possible since  $n_c \cdot n_c \ge |\Gamma|$ . In this clique, we fix the triangle having its endpoints in  $Z_1 \cup Z_2 \cup Z_3$ . Formally, for each clause  $q = \{q_1, q_2, q_3\} \in \Gamma$  we add three communities  $C_q^1 = \{y_j, z_s^2, z_s^3\}, C_q^2 = \{y_j, z_s^1, z_s^3\}$  and  $C_q^3 = \{y_j, z_s^1, z_s^2\}$ . We refer to these communities as the clause communities  $\mathcal{C}^{\Gamma}$ . Afterwards, we set  $\nu(q, q_1) \coloneqq \{y_j, z_s^1\}, \nu(q, q_2) \coloneqq \{y_j, z_s^2\}$ , and  $\nu(q, q_3) \coloneqq \{y_j, z_s^3\}$  to assign each literal in clause q to an edge of the clause gadget. These edges are referred to as free. Second, we fix the edges of the clique  $Z_1 \cup Z_2 \cup Z_3$ .

Note that, similar to the variable gadget, edges with both endpoints in Y are not contained in any community. Again, we only add these edges to ensure that G is a clique.

Observe that the sets of free edges corresponding to two distinct clauses are disjoint:

▶ Claim 2 (\*). Each free edge of  $E(G_{\Gamma})$  is contained in exactly one subgraph induced by a clause community in  $C^{\Gamma}$ .

Proof. Each clause community  $C^q$  consists of one vertex  $y \in Y$  and one vertex  $z_s^i \in Z_i$ . In  $C^q$  the free edges form a star with y as there center and the vertices  $z_s^i$  as leaves. Suppose there are two distinct clause communities  $C_1$  and  $C_2$  such that they have at least one free edge in common. This implies that  $C_1$  and  $C_2$  contain the same vertex of Y and have at least one vertex  $z_s^i$  of  $Z_i$  in common. By construction, in each clause community  $C_i$  the three vertices of  $Z_1 \cup Z_2 \cup Z_3$  are either identical or there is no overlap. Thus,  $C_1$  and  $C_2$  are identical, a contradiction.

**Connecting the Gadgets:** We complete the construction by describing how the variable and clause gadget are connected, using new *assignment* communities. The idea is to add a new community containing the endpoints of a free edge describing a literal in a clause together with the endpoints of the selection edge describing the same opposite literal in the variable gadget. These communities model occurrences of variables in the clauses. Roughly speaking, these communities are satisfied if the selection edge of the variable gadget or the free edge of the clause gadget is part of the solution. To enforce this, we fix further edges of G. An example of an assignment community for one literal is shown in part c) of Figure 2.

We create for each clause  $q = \{q_1, q_2, q_3\} \in \Gamma$  three assignment communities  $C_q^{q_1} = \nu(q, q_1) \cup \theta(\overline{q_1}), C_q^{q_2} = \nu(q, q_2) \cup \theta(\overline{q_2}), \text{ and } C_q^{q_3} = \nu(q, q_3) \cup \theta(\overline{q_3}).$  We denote the assignment communities with  $\mathcal{C}_{\Gamma}^X$ . To enforce that each solution contains the selection edge or the free edge of each assignment community, we fix all edges between the vertex sets U and Z, between the vertex sets P and Y, and between the vertex sets P and Z.

Note that edges between U and Y are not part of any community and only added to ensure that G is a clique.

Finally, we set  $\ell := |X| + 2 \cdot |\Gamma| + {|P| \choose 2} + {|Z| \choose 2} + |U| \cdot |Z| + |P| \cdot |Y| + |P| \cdot |Z|$ . Let *I* denote the constructed instance of UNWEIGHTED STARS NWS. Clearly,  $|V(G)| \in \mathcal{O}(\sqrt{|\phi|})$  and the maximum community size is 4. Instance *I* has at most  $\mathcal{O}(\sqrt{|\phi|} \cdot \sqrt{|\phi|}) = \mathcal{O}(|\phi|)$  communities of size 2,  $|X| + 3 \cdot |\Gamma|$  communities of size 3, and  $3 \cdot |\Gamma|$  communities of size 4. Thus, *I* has  $\mathcal{O}(|\phi|)$  communities.

**Correctness:** We show that the formula  $\phi$  is satisfiable if and only if I is a yes-instance of UNWEIGHTED STARS NWS. Before we prove this statement, we make three observations about the solution.

Since fixed edges are contained in a community of size 2, we have the following

## ▶ Observation 3.2. Each solution of I contains all fixed edges.

Observe that I consists of precisely  $\binom{|P|}{2} + \binom{|Z|}{2} + |U| \cdot |Z| + |P| \cdot |Y| + |P| \cdot |Z| = \ell - (|X| + 2 \cdot |\Gamma|)$  fixed edges. Hence, each solution can contain at most  $|X| + 2 \cdot |\Gamma|$  selection and free edges. Thus, we obtain the following.

**Claim 3** (\*). Each solution of I contains at least |X| selection edges.

Proof. Observe that at least one selection edge of each variable community C in the variable gadget is contained in the solution to satisfy C. By Claim 1, these are at least |X| edges.

<1

▶ Claim 4 (\*). Each solution of I contains at least  $2 \cdot |\Gamma|$  free edges.

Proof. Observe that at least two free edges of each triple of communities  $C_q^1, C_q^2, C_q^3$  of the clause gadget for each clause q are contained in the solution to fulfill the requirements of these communities. By Claim 2, and since no free edge is contained in any community of size 2, these are at least  $2 \cdot |\Gamma|$  edges.

From Observation 3.2 and Claims 3 and 4 we conclude that each solution consists of at least  $\ell$  edges. Now, we show the correctness.

(⇒) Let  $A: X \to \{0, 1\}$  be an assignment satisfying  $\phi$ . We describe how to obtain a solution G' = (V(G), E') with  $|E'| = \ell$  from A. First, by Observation 3.2, the solution contains all  $\ell - |X| - 2 \cdot |\Gamma|$  fixed edges. Second, for each variable, we chose the selection edge of the variable gadget representing the literal that is not satisfied by A. For a formal definition, recall that for a variable  $x \in X$ ,  $\theta(x)$  is the selection edge representing x and  $\theta(\overline{x})$  is the selection edge representing  $\overline{x}$ . We set  $E'_X := \{\theta(x) : x \in X, A(x) = 0\} \cup \{\theta(\overline{x}) : x \in X, A(x) = 1\}$ . Note that  $E'_X$  and the fixed edges fulfill the spanning star property of the variable communities. Third, we construct the set  $E'_{\Gamma}$  of free edges which are contained in the solution. For each clause  $q = \{q_1, q_2, q_3\} \in \Gamma$ , there is at least one variable x such that with the assignment A the literal, say  $q_3$ , corresponding to x satisfies q. We add the edges  $\nu(q, q_1)$  and  $\nu(q, q_2)$  to  $E'_{\Gamma}$ . Recall that in the construction of the clause gadget of q, we created three clause communities  $C^1_q = \nu(q, q_1) \cup \nu(q, q_2)$ ,  $C^2_q = \nu(q, q_1) \cup \nu(q, q_3)$ , and  $C^3_q = \nu(q, q_2) \cup \nu(q, q_3)$ . Observe that the endpoint of  $\nu(q, q_1)$  in Z or the endpoint of  $\nu(q, q_2)$  in Z is a center for each of these three clause communities. Thus, the spanning star property of all clause communities is fulfilled.

Observe that the graph G' has exactly  $\ell$  edges. To verify that I is a yes-instance of UNWEIGHTED STARS NWS, it remains to show that G' also fulfills the spanning star property of the assignment communities. For this, consider the three assignment communities  $C_q^{q_1} = \nu(q, q_1) \cup \theta(\overline{q_1}), C_q^{q_2} = \nu(q, q_2) \cup \theta(\overline{q_2})$ , and  $C_q^{q_3} = \nu(q, q_3) \cup \theta(\overline{q_3})$  corresponding to clause q. Recall that we assume that q is satisfied by literal  $q_3$  and that we defined E' accordingly. Also, recall that all fixed edges are contained in E'. Since  $\nu(q, q_1)$  and  $\nu(q, q_2)$  are edges of E', we conclude that the unique vertex in Z of  $C_q^{q_1}$  and  $C_q^{q_2}$  is the center of a star for these assignment communities. Furthermore, since  $\theta(\overline{q_3}) \in E'$ , we conclude that the unique vertex of the assignment community  $C_q^{q_3}$ . Hence, the requirements of all communities are fulfilled by G'.

 $(\Leftarrow)$  Let G' = (V(G), E') be a solution of I with  $\ell$  edges. From Observation 3.2 and Claims 3 and 4, we conclude that E' contains exactly |X| selection edges, denoted by  $E'_X$ , and exactly  $2 \cdot |\Gamma|$  free edges. Furthermore, note that for each variable community  $C_x$ corresponding to some variable  $x \in X$ , the solution contains at least one of the selection edges  $\theta(x)$  or  $\theta(\overline{x})$ . Recall that, according to Claim 1, the selection edges of each two variable communities are disjoint. Since G' contains exactly |X| selection edges, we conclude that G'contains exactly one selection edge of each variable  $x \in X$ . This, allows us to properly define an assignment A by setting A(x) = 0 if  $\theta(x) \in E'_X$  and A(x) = 1 if  $\theta(\overline{x}) \in E'_X$ .

Similarly, according to Claim 2, the free edges of each two clause communities are disjoint. Since G' contains exactly  $2 \cdot |\Gamma|$  free edges, we conclude that G' contains exactly two free edges per clause  $q \in \Gamma$  (according to Claim 4), that is, we have  $|E' \cap \{\nu(q, q_1), \nu(q, q_2), \nu(q, q_3)\}| = 2$  for each clause  $q \in \Gamma$ .

Since G' contains exactly  $\ell$  edges and |X| of them are selection edges,  $2 \cdot |\Gamma|$  of them are free edges, and  $\ell - |X| - 2 \cdot |\Gamma|$  of them are fixed edges, we conclude that no edge which is not a selection edge, a free edge or a fixed edge can be part of G'. Recall that we created for each clause  $q = \{q_1, q_2, q_3\} \in \Gamma$  three assignment communities  $C_q^{q_1}$ ,  $C_q^{q_2}$ , and  $C_q^{q_3}$  connecting the variable and the clause gadget. Observe that community  $C_q^{q_i}$  contains three fixed edges, the

selection edge  $\nu(q, q_i)$ , the free edge  $\theta(\overline{q_i})$ , and one edge which has none of these 3 types. By the above argumentation on the tightness of the budget  $\ell$  and the fact that G' is a solution, we conclude that in the assignment community  $C_q^{q_i}$  the edge  $\nu(q, q_i)$  or the edge  $\theta(\overline{q_i})$  is contained in E' for each  $i \in \{1, 2, 3\}$  to fulfill the spanning star property of these assignment communities. Now recall that we have  $|E' \cap \{\nu(q, q_1), \nu(q, q_2), \nu(q, q_3)\}| = 2$  for each clause q. Thus, at least one of the edges  $\theta(\overline{q_1}), \theta(\overline{q_2})$ , or  $\theta(\overline{q_3})$  is contained in E'. Without loss of generality, we assume that  $\theta(\overline{q_3}) \in E'$ . This implies that the assignment of the variable xcorresponding to literal  $q_3$  satisfies clause q. Thus, if  $q_3 = x$ , we set A(x) = 1, and otherwise if  $q_3 = \overline{x}$ , we set A(x) = 0. This applies to all clauses, and thus  $\phi$  is satisfied by A.

## 3.2 Adaptions of Theorem 3.1

In the following, we present two further results: First, the same ETH-based bound for UNWEIGHTED CONNECTIVITY NWS (Corollary 3.3). Second, we show that UNWEIGHTED STARS NWS remains NP-hard on graphs with constant maximum degree (Corollary 3.4). The latter result can be shown by reducing from the NP-hard (3,B2)-SAT [4] and by using a similar, but uncompressed, construction to the one introduced in the proof of Theorem 3.1.

Adaption for Unweighted Connectivity NWS. Now, we modify the construction to replace the star requirement for each community with the requirement of being connected. The variable gadget and the clause gadget is constructed exactly as in the construction in Theorem 3.1 for UNWEIGHTED STARS NWS. The assignment gadget, however, is constructed differently: instead of fixing the edges between the vertex sets  $U \cup P$  and Z, and between P and Y, we now only fix the edges between U and Z, and between P and Y. Consequently, all edges with one endpoint in P and one in Z, or one endpoint in U and one in Y are not contained in any community. Then, we set  $\ell := |X| + 2 \cdot |\Gamma| + {|P| \choose 2} + {|Z| \choose 2} + |U| \cdot |Z| + |P| \cdot |Y|$ , that is, compared to UNWEIGHTED STARS NWS, the budget  $\ell$  is decreased by  $|P| \cdot |Z|$ .

Now the proof is similar to the proof of Theorem 3.1: First, the size 2 communities still imply that all fixed edges are part of any solution. Second, since for communities of size 3, the star property and the connectivity property coincide, the argumentation for UNWEIGHTED CONNECTIVITY NWS is analogously to that of UNWEIGHTED STARS NWS. Third, since also in this construction the budget is tight, the solution contains all fixed edges, exactly |X| selection edges, and exactly  $2 \cdot |\Gamma|$  free edges. Now, since the edges of these 3 types in each assignment community form a cycle of length 4 (2 fixed edges, 1 selection edge, and 1 free edge), we conclude that in each assignment community, the selection edge or the free edge (or both) are contained in the solution. From now on, the argumentation is analogous to the argumentation in Theorem 3.1 for UNWEIGHTED STARS NWS. Thus, we obtain the following.

▶ Corollary 3.3. If the ETH is true, then UNWEIGHTED CONNECTIVITY NWS cannot be solved in  $2^{o(n^2)} \cdot \text{poly}(n+c)$  time, even if G is a clique and each community has size at most 4.

Adaption for Unweighted Stars NWS with Constant Maximum Degree. Next, we show that UNWEIGHTED STARS NWS remains NP-hard on graphs with constant maximum degree. We achieve this by three adjustments: first, we do not compress the variable and clause gadget in the construction of Theorem 3.1, second we reduce from the NP-hard (3,B2)-SAT [4] in which each clause consists of three literals and each literal (positive and negative) appears exactly twice, and third, we do not add edges which are not contained in any community.

More precisely, for each variable, we add a triangle with one fixed edge; again this is achieved by adding a community of size 2 containing this edge. Afterwards, for each clause we add a size 4-clique with a fixed triangle. Then, we add the assignment communities and the corresponding fixed edges as described in the construction of Theorem 3.1. Note that each vertex is either contained in a variable triangle or in a size 4-clique corresponding to a clause. Since there are only edges between the variable gadgets and the clause gadgets, each vertex in the variable gadget has exactly two neighbors in the variable gadget and each vertex in the clause gadget has exactly three neighbors in the clause gadget. It remains to show that each vertex in the variable gadget has only a constant number of neighbors in the clause gadget and vice versa.

First, we consider a vertex v in the variable gadget. Since each literal appears exactly twice, there are exactly four assignment communities having vertices of the triangle in the variable gadget corresponding to v. With this observation at hand, one can show that vhas at most four neighbors in the clause gadget. Second, we consider a vertex v in the clause gadget. Since each clause consists of three literals, there are exactly three assignment communities having vertices of the size 4-clique in the clause gadget corresponding to v. With this observation at hand, one can show that v has at most three neighbors in the clause gadget. Hence, we obtain the following.

▶ Corollary 3.4. UNWEIGHTED STARS NWS remains NP-hard and, assuming the ETH, cannot be solved in  $2^{o(n+m+c)} \cdot poly(n+c)$  time on graphs with maximum degree six and community size at most 4.

## 4 Parameterization by the Feedback Edge Number of a Solution

The parameter  $\ell$ , the number of edges in the solution is in most cases not independent from the size of the input instance of STARS NWS or CONNECTIVITY NWS: if the hypergraph  $(V, \mathcal{C})$  is connected, a solution G' has at least n-1 edges. In other words, n-1 is a lower bound for  $\ell$  in this case. In this section, we study STARS NWS and CONNECTIVITY NWS parameterized above this lower bound. Formally, the parameter t is defined as the size of a minimum feedback edge set of the solution of an instance of STARS NWS or CONNECTIVITY NWS. Thus, the parameter t measures how close the solution is to a forest. Formally, the definition is  $t := \ell - n + x$  where x denotes the number of connected components of G'. Recall that t can be computed in polynomial time (see Section 2.).

## 4.1 An XP-Algorithm for Stars NWS

In this subsection, we show that STARS NWS parameterized by t admits an XP-algorithm with the following running time.

▶ Theorem 4.1. STARS NWS can be solved in  $m^{4t} \cdot poly(|I|)$  time.

Korach and Stern [34] asked whether STARS NWS is polynomial-time solvable if t = 0. Theorem 4.1 answers this question positively.

Our XP-algorithm, exploits the fact that there are two different kinds of cycles in G': First, there are *global* cycles. These are the cycles in the solutions that are directly caused by cycles in the input hypergraph. No solution may avoid these cycles. Second, there are *local* cycles. These are cycles which are entirely contained in the subgraph induced by two communities. Since in each solution, each community contains a spanning star, local cycles can only have length 3 or 4. This allows us to bound the number of possible local cycles and



**Figure 3** Examples for solutions with and without local cycles. Red edges indicate the edges of the solution. Part a) shows an example, where both communities induce a local cycle. Part b) shows an example, where the two communities do not induce local cycle. Finally, part c) shows an example, where the solution contains a cycle but no two communities induce a local cycle.

thus to consider all possibilities for the local cycles in XP-time with respect to t. Then, the crux of our algorithm is that after all local cycles have been fixed, all remaining cycles added by our algorithm have to be global and are thus unavoidable. Using this fact, we show that in polynomial time we can compute an optimal solution with feedback edge number at most t that extends a fixed set of local cycles without introducing any further local cycles. To do this, for each community C, we store a set of *potential centers*, that is, vertices of C that may be the center of a spanning star of C in any solution that does not produce new local cycles. We define several *operations* that restrict the potential centers of each community. We show that after all operations have been applied exhaustively, one can greedily pick the best remaining center for each community.

**Algorithm-specific notation.** Next, we present the formal definition of local cycles; an example is shown in Figure 3. For a spanning subgraph H of G and a community  $C \in C$ , let  $\operatorname{univ}_H(C)$  denote the vertices of C that are *universal* for C in H. Recall that a vertex u is universal for some vertex set S in a graph F if  $\{u, w\} \in E(F)$  for each  $w \in S$ . Note that  $\operatorname{univ}_H(C) \subseteq \operatorname{univ}_G(C)$ . In the following, we assume that for each community  $C \in C$ ,  $\operatorname{univ}_G(C) \neq \emptyset$ , as otherwise, there is no solution for the instance I of STARS NWS, and I is a trivial no-instance.

For a solution G', we say that two distinct communities  $C_1$  and  $C_2$  induce a local cycle if for each  $i \in \{1, 2\}$ , there is a vertex  $c_i \in \operatorname{univ}_{G'}(C_i)$  such that the graph  $S_1 \cup S_2$  contains a cycle. Here, for each  $i \in \{1, 2\}$ ,  $S_i$  is the spanning star of  $C_i$  with center  $c_i$  and  $S_1 \cup S_2$  is the union of both these stars defined by  $S_1 \cup S_2 \coloneqq (C_1 \cup C_2, \{\{c_i, w_i\}: w_i \in C_i \setminus \{w_i\}, i \in \{1, 2\}\})$ . Moreover, we say that each cycle of  $S_1 \cup S_2$  is a local cycle in G'. Note that each local cycle has length at most four, and if  $C_1$  and  $C_2$  induce a local cycle, then  $|C_1 \cap C_2| \ge 2$ .

As described above, the first step of the algorithm behind Theorem 4.1 is to test each possibility for the local cycles of the solution. For a fixed guess, we let  $E^*$  denote the set of all edges contained in at least one local cycle and in the following we refer to them as *local edges*. Moreover, we call a minimum solution G' fitting for  $E^*$  if each local cycle of G' uses only edges of  $E^*$  and each edge of  $E^*$  is contained in G'. Hence, to determine whether the choice of local edges  $E^*$  can lead to a solution, we only have to check, whether there is a fitting solution for  $E^*$ . In the following, we show that this can be done in polynomial time.

▶ **Theorem 4.2.** Let  $I = (G = (V, E), C, \omega, \ell, b)$  be an instance of STARS NWS, and let  $E^* \subseteq E$ . In polynomial time, we can

find a solution G' = (V, E') for I with  $E^* \subseteq E'$ ,  $|E'| \le \ell$ , and  $\omega(E') \le b$  or

- correctly output that there is no minimum solution that is fitting for  $E^*$ .

Based on the definition of fitting solutions, we define for each community  $C \in \mathcal{C}$  a set  $\operatorname{fit}_{E^*}(C)$  of possible centers. We initialize  $\operatorname{fit}_{E^*}(C) := \operatorname{univ}_G(C)$  for each community  $C \in \mathcal{C}$ . The goal is to reduce these sets of possible centers of each community as much as possible while preserving the following property, which trivially holds for the initial  $\operatorname{fit}_{E^*}(C)$  for each community  $C \in \mathcal{C}$ .

▶ Property 1. For each minimum solution G' which is fitting for  $E^*$  and each community  $C \in C$ , we have  $\operatorname{univ}_{G'}(C) \subseteq \operatorname{fit}_{E^*}(C)$ .

Note that if Property 1 is fulfilled and if  $\operatorname{fit}_{E^*}(C) = \emptyset$  for some  $C \in \mathcal{C}$ , then we can correctly output that there is no fitting solution for  $E^*$ .

In the following, we define several operations that for some communities  $C \in \mathcal{C}$  remove vertices from fit<sub>E\*</sub>(C) which—when taken as a center vertex for C—would introduce new local cycles, violating the properties of a fitting solution. We show that all of these operations preserve Property 1 and that after all these operations are applied exhaustively, the task of Theorem 4.2 can be performed greedily based on fit<sub>E\*</sub>. Examples for each of our operations are shown in Figure 5.

In the following, we say that a vertex  $v \in V$  is *locally universal* for a vertex set  $A \subseteq V$ , if for each vertex  $w \in A \setminus \{v\}$ , the vertex pair  $\{v, w\}$  is a local edge. Based on this definition, we are now able to present the first operation.

▶ **Operation 1.** Let  $C \in C$  be a community and let  $\{y, z\} \subseteq C$  be a local edge. Remove each vertex v from fit<sub>E\*</sub>(C) which is not locally universal for  $\{y, z\}$ .

The following lemma shows that Operation 1 preserves Property 1.

▶ Lemma 4.3 (\*). Let G' be a minimum solution for I, let C be a community of C and let  $x \in \operatorname{univ}_{G'}(C)$  such that x is not locally universal for some local edge  $\{y, z\} \subseteq C$ . Then, G' is not fitting for  $E^*$ .

**Proof.** If  $\{y, z\}$  is not an edge of G', then G' is not fitting for  $E^*$ , since G' does not contain all edges of  $E^*$ . Hence, in the following we assume that  $\{y, z\}$  is an edge of G'. If the graph G'' obtained by removing the edge  $\{y, z\}$  from G' is a solution for I, then G' is not a minimum solution. Thus, we assume that G'' is not a solution for I. Consequently, there is some community  $D \in \mathcal{C}$  such that  $\operatorname{univ}_{G''}(D) = \emptyset$ . Since G' is a solution for I,  $\operatorname{univ}_{G'}(D) \neq \emptyset$ . Moreover, since G'' is obtained from G' by removing the edge  $\{y, z\}$ , we have  $y \in \operatorname{univ}_{G'}(D)$ or  $z \in \operatorname{univ}_{G'}(D)$ . Hence, C and D induce a local cycle in G' on the vertices of  $\{x, y, z\}$ . Since x is not locally universal for  $\{y, z\}$ , at least one edge of this local cycle is not a local edge. Consequently, G' is not fitting for  $E^*$ .

Note that after the exhaustive application of Operation 1, for each community  $C \in C$  with at least one local edge, the vertices of  $\operatorname{fit}_{E^*}(C)$  induce a clique with only local edges. In the following, we assume that Operation 1 is applied exhaustively.

Next, we define a partition  $\mathfrak{C}$  of the communities of  $\mathcal{C}$ . The idea of this partition is that in each fitting solution for  $E^*$ , all communities of the same part of the partition  $\mathfrak{C}$  have the same unique center. The definition of the partition  $\mathfrak{C}$  is based on the following lemma.

▶ Lemma 4.4 (\*). Let C and D be distinct communities of C with  $|C \cap D| \ge 3$  and where no vertex  $v \in C \cup D$  is locally universal for  $C \cap D$ . Let G' be a solution such that there is no vertex  $w \in C \cap D$  with  $\operatorname{univ}_{G'}(C) = \operatorname{univ}_{G'}(D) = \{w\}$ . Then, C and D induce a local cycle in G' that uses at least one edge which is not a local edge.

**Proof.** Let G' be a solution such that there is no vertex  $w \in C \cap D$  with  $\operatorname{univ}_{G'}(C) = \operatorname{univ}_{G'}(D) = \{w\}$ . Since  $\operatorname{univ}_{G'}(C)$  and  $\operatorname{univ}_{G'}(D)$  are nonempty, there is a vertex  $x \in \operatorname{univ}_{G'}(C)$  and a vertex  $y \in \operatorname{univ}_{G'}(D)$  such that  $x \neq y$ . By the fact that x is not locally universal for  $C \cap D$ , there is some vertex  $x' \in (C \cap D) \setminus \{x\}$ , such that  $\{x, x'\}$  is not a local edge. Similarly, there is some vertex  $y' \in (C \cap D) \setminus \{y\}$ , such that  $\{y, y'\}$  is not a local edge. We distinguish two cases.

**Case 1:** x = y' and y = x'. Since  $|C \cap D| \ge 3$ , there is a vertex  $z \in C \cap D$  distinct from both x and y. Hence, C and D induce a local cycle in G' on the vertices  $\{x, y, z\}$ . This local cycle contains the edge  $\{x, y\}$ , which is not a local edge, since y = x'.

**Case 2:**  $x \neq y'$  or  $y \neq x'$ . Assume without loss of generality that  $x \neq y'$ . Then, if  $x \in C \cap D$  or  $y \in C \cap D$ , then  $\{x, y, y'\}$  is a local cycle (induces by C and D) in G'. Otherwise, if  $C \cap D$  contains neither x nor y, let z be an arbitrary vertex of  $(C \cap D) \setminus \{y'\}$ . Then,  $\{x, y, y', z\}$  is a local cycle (induced by C and D) in G'. Both local cycles contain the edge  $\{y, y'\}$  which is not a local edge.

Consider the auxiliary graph  $G_{\mathfrak{C}}$  with vertex set  $\mathcal{C}$  and where two distinct communities Cand D are adjacent if and only if a)  $|C \cap D| \geq 3$  and b) there is no locally universal vertex for  $C \cap D$  in  $C \cup D$ . The partition  $\mathfrak{C}$  consists of the connected components of  $G_{\mathfrak{C}}$  and for a community  $C \in \mathcal{C}$ , we denote by  $\mathfrak{C}(C)$  the collection of communities in the connected component of C in  $G_{\mathfrak{C}}$ . An example is shown in Figure 4.

By Lemma 4.4 and due to transitivity, we obtain the following.

▶ Corollary 4.5. For each community  $C \in C$  with  $|\mathfrak{C}(C)| \ge 2$  and each fitting solution G' for  $E^*$ , there is a vertex  $v \in \bigcap_{\widetilde{C} \in \mathfrak{C}(C)} \widetilde{C}$  such that  $\operatorname{univ}_{G'}(\widetilde{C}) = \{v\}$  for each  $\widetilde{C} \in \mathfrak{C}(C)$ .

This implies that the following operation preserves Property 1.

▶ **Operation 2.** Let  $C \in C$ . Remove each vertex v from fit<sub>E\*</sub>(C) if v is not contained in  $\bigcap_{\widehat{C} \in \mathfrak{C}(C)}$  fit<sub>E\*</sub>( $\widehat{C}$ ).

After applying Operation 2 exhaustively, for each community  $\widetilde{C} \in \mathfrak{C}(C)$ ,  $\operatorname{fit}_{E^*}(\widetilde{C}) = \operatorname{fit}_{E^*}(C)$ .

Next, we describe an operation for communities that do not contain any local edge. To this end, we observe the following.

▶ Lemma 4.6. Let  $C \in C$  be a community that contains no local edge. Moreover, let  $D \in C$  such that  $|C \cap D| \ge 2$ . Then, for each solution G' where  $\operatorname{univ}_{G'}(C) \not\subseteq C \cap D$ , C and D induce a local cycle in G' that uses at least one edge which is not a local edge.

**Proof.** Let G' be a solution, let x and y be distinct vertices of  $C \cap D$ , and let  $z \in \operatorname{univ}_{G'}(C) \setminus (C \cap D)$ . Since  $z \notin C \cap D$ , G' contains the edges  $\{z, x\}$  and  $\{z, y\}$ . Hence, for each vertex  $w \in \operatorname{univ}_{G'}(D)$ , C and D induce a local cycle in G' on the vertices  $\{x, y, z, w\}$ , since  $z \neq w$ . If  $w \in C \cap D$ , then this cycle has length 3. Otherwise, this cycle has length 4. In both cases, the local cycle contains the edges  $\{z, x\}$  and  $\{z, y\}$ . Since C contains no local edge, neither of these two edges is a local edge.

Note that Lemma 4.6 implies that the following operation preserves Property 1.

▶ **Operation 3.** Let  $C \in C$  such that C contains no local edge. Moreover, let  $D \in C$  such that  $|C \cap D| \ge 2$ . Remove all vertices from fit<sub>E\*</sub>(C) that are not contained in  $C \cap D$ .

Next, we describe two operations for communities that contain at least one local edge.



**Figure 4** Examples for parts of the partition  $\mathfrak{C}$ . Only the local edges are shown. Note that A and C are both contained in  $\mathfrak{C}(B)$ , since A and C share at least three vertices with B and no vertex of  $A \cup B$  or  $C \cup B$  is locally universal for  $A \cap B$  or  $C \cap B$ , respectively. Hence, after exhaustive application of Operation 2, fit<sub>E\*</sub>(A) = fit<sub>E\*</sub>(B) = fit<sub>E\*</sub>(C) =  $\emptyset$ , since A and C share no vertices. Furthermore,  $Y \in \mathfrak{C}(Z)$ , since no vertex of  $Y \cup Z$  is locally universal for  $Y \cap Z$ . Note that  $X \notin \mathfrak{C}(Z)$ , since the black vertex of X is locally universal for  $X \cap Y$  and  $X \cap Z$ . Observe that an exhaustive application of Operation 2 yields fit<sub>E\*</sub>(Z)  $\subseteq Y \cap Z$  and an exhaustive application of Operation 5 yields fit<sub>E\*</sub>(Z)  $\cap (Y \cap Z) = \emptyset$ , since X contains at least one local edge and  $X \cap Z$  contains no local edge. Hence, for both shown hypergraphs, there is no fitting solution for the given set of local edges.

▶ Operation 4. Let C ∈ C such that C contains at least one local edge. Moreover, let D ∉ 𝔅(C) be a community, such that |C ∩ D| = 2 and {x, y} := C ∩ D is not a local edge.
1. If fit<sub>E\*</sub>(C) ∩ {x, y} = Ø, then remove x and y from fit<sub>E\*</sub>(D) or
2. if fit<sub>E\*</sub>(C) ∩ {x, y} = {x}, then set fit<sub>E\*</sub>(C) := {x}.

▶ Lemma 4.7. If Operation 1 is exhaustively applied, then Operation 4 preserves Property 1.

**Proof.** Suppose that Property 1 holds.

First we show that Operation 4 preserves Property 1 if  $\operatorname{fit}_{E^*}(C) \cap \{x, y\} = \emptyset$ . Suppose that  $\operatorname{fit}_{E^*}(C) \cap \{x, y\} = \emptyset$ . Let G' be a solution for I containing all edges of  $E^*$ . We show that G' is not fitting for  $E^*$  if  $\operatorname{univ}_{G'}(D) \cap \{x, y\} \neq \emptyset$ . Suppose that there is some vertex  $z \in \operatorname{univ}_{G'}(D) \cap \{x, y\}$ . Since Property 1 holds and  $\operatorname{fit}_{E^*}(C) \cap \{x, y\} = \emptyset$ , there is some vertex  $w \in \operatorname{univ}_{G'}(C) \setminus \{x, y\}$ . Then, C and D induce a local cycle in G' on the vertices of  $\{w, x, y\}$ . Since  $\{x, y\}$  is not a local edge, G' is not fitting for  $E^*$ .

Next, we show that Operation 4 preserves Property 1 if Operation 1 is exhaustively applied and if  $\operatorname{fit}_{E^*}(C) \cap \{x, y\} = \{x\}$ . Suppose that Operation 1 is exhaustively applied and that  $\operatorname{fit}_{E^*}(C) \cap \{x, y\} = \{x\}$ . Let G' be a solution for I containing all edges of  $E^*$  and let z be an arbitrary vertex of  $\operatorname{univ}_{G'}(D)$ . We show that G' is not fitting for  $E^*$  if there is some vertex  $w \in \operatorname{univ}_{G'}(C)$  distinct from x. Since  $y \notin \operatorname{fit}_{E^*}(C)$ , w is not a vertex of  $C \cap D = \{x, y\}$ .

We show that C and D induce a local cycle in G' containing at least one edge which is not a local edge. If  $z \in \{x, y\}$ , then C and D induce a local cycle in G' on the vertices of  $\{w, x, y\}$ . Since  $\{x, y\}$  is not a local edge, G' is not fitting for  $E^*$ . Otherwise, that is, if  $z \notin \{x, y\}$ , Cand D induce a local cycle in G' on the vertices of  $\{w, x, y, z\}$ . This local cycles contains the edge  $\{w, y\}$ . Hence, to show that G' is not fitting for  $E^*$ , it is sufficient to show that  $\{w, y\}$  is not a local edge. Recall that x is contained in fit<sub> $E^*$ </sub> (C) and that Operation 1 is exhaustively applied. Hence, x is locally universal for each local edge  $\{u, v\}$  in C. By assumption, Ccontains a local edge and  $\{x, y\}$  is not a local edge. Consequently, no local edge in C is incident with y and thus  $\{w, y\}$  is not local edge. Hence, G' is not fitting for  $E^*$ .



**Figure 5** Examples of applications of Operations 1–5. The black edges represent the local edges, the solid (for C) or dashed (for D) red edges show the non-local edges resulting from choosing the respective center for community C or D. For example in 2), z is the center of community C and v is the center of community D, and the edges  $\{z, y\}$  and  $\{v, y\}$  are non-local edges in the solution. For each operation, the violation of the property of being a fitting solution is shown, if a vertex a is selected as a center of a community A where the application of the corresponding operation would remove a from fit<sub>E\*</sub>(A). In 1), 2), 3), and 4.2), the vertex selected as center for community C is removed from fit<sub>E\*</sub>(C) by the respective operation. For example, in 4.2), (assuming fit<sub>E\*</sub>(C)  $\cap$   $\{x, y\} = \{x\}$ ) Operation 4 removes v from fit<sub>E\*</sub>(C), as otherwise selecting v as center of C results in the depicted non-fitting solution. In 4.1) and 5), the vertex selected as center for community C is removed from fit<sub>E\*</sub>(C) by the respective operation. For example in 4.1), (assuming fit<sub>E\*</sub>(C)  $\cap$   $\{x, y\} = \emptyset$ ) Operation 4 removes y from fit<sub>E\*</sub>(D), as otherwise selecting y as center of D results in the depicted non-fitting solution.

▶ **Operation 5.** Let  $C \in C$  be a community containing at least one local edge. Moreover, let  $D \notin \mathfrak{C}(C)$  such that  $|C \cap D| \ge 3$ . For each pair of distinct vertices x and y of  $C \cap D$ , where  $\{x, y\}$  is not a local edge, remove x and y from fit<sub>E\*</sub>(D).

▶ Lemma 4.8. If Operation 1 is exhaustively applied, then Operation 5 preserves Property 1.

**Proof.** Suppose that Property 1 holds.

We show that Operation 5 preserves Property 1 if Operation 1 is exhaustively applied. Hence, assume in the following that Operation 1 is exhaustively applied. Let G' be a solution for I containing all edges of  $E^*$ . We show that G' is not fitting for  $E^*$  if  $\text{univ}_{G'}(D)$  contains xor y.

Since  $|C \cap D| \ge 3$  and  $D \notin \mathfrak{C}(C)$ , there is some vertex  $v \in C \cup D$  which is locally universal for  $C \cap D$ . Note that since  $\{x, y\}$  is not a local edge, v is neither x nor y. We distinguish two cases.

**Case 1:**  $v \in D$ . Since Operation 1 is exhaustively applied and  $\{v, x\}$  and  $\{v, y\}$  are local edges in D, we conclude that each vertex on  $\operatorname{fit}_{E^*}(D)$  is locally universal for x and y. Since  $\{x, y\}$  is not a local edge,  $\operatorname{fit}_{E^*}(D)$  contains neither x nor y.

**Case 2:**  $v \in C \setminus D$ . Since Operation 1 is exhaustively applied, we conclude that each vertex of fit<sub>E\*</sub>(C) is locally universal for  $C \cap D$ . Let w be an arbitrary vertex of  $\operatorname{univ}_{G'}(C)$ . If  $w \notin \operatorname{fit}_{E^*}(C)$ , then G' is not fitting for  $E^*$  since Property 1 holds. Otherwise,  $w \in \operatorname{fit}_{E^*}(C)$ .

**Algorithm 1** Algorithm solving the problem described in Theorem 4.2. **Input** :  $I = (G = (V, E), C, \omega, \ell, b), E^* \subseteq E$ **Output:** A solution G' = (V, E') with at most  $\ell$  edges and total weight at most b, or no, if there is no minimal solution which is fitting for  $E^*$ 1 Compute the partition  $\mathfrak{C}$  of  $\mathcal{C}$ **2** For each  $C \in \mathcal{C}$ , initialize fit<sub>E\*</sub>(C)  $\leftarrow$  univ<sub>G</sub>(C) and apply Operation 1 **3** Apply Operations 1–5 exhaustively 4 if  $\operatorname{fit}_{E^*}(C) = \emptyset$  for some  $C \in \mathcal{C}$  then return no 5  $G_A \leftarrow (V, E^*)$ 6 forall  $\mathcal{L} \in \mathfrak{C}$  do  $C \leftarrow$  some community of  $\mathcal{L}$ 7  $V_{\mathcal{L}} \leftarrow \bigcup_{\widetilde{C} \in \mathcal{L}} \widetilde{C}$ 8  $y \leftarrow \arg\min_{u \in \operatorname{fit}_{E^*}(C)} \omega(\{\{u,v\} : v \in V_{\mathcal{L}} \setminus \{u\}\} \setminus E^*)$ 9 add all edges of  $\{\{y, v\} : v \in V_{\mathcal{L}} \setminus \{y\}\}$  to  $G_A$ 10 11 if  $|E'| \leq \ell$  and  $\omega(E') \leq b$  then return  $G_A$ 12 return no

Consequently, w is distinct from both x and y, since  $\{x, y\}$  is not a local edge. Since  $\{x, y\} \subseteq C \cap D$  and  $w \in \operatorname{univ}_{G'}(C)$ , G' contains the edges  $\{w, x\}$  and  $\{w, y\}$ . If  $\operatorname{univ}_{G'}(D)$  contains x or y, G' contains the edge  $\{x, y\}$ . Hence, C and D induce a local cycle in G' on the vertices  $\{x, y, w\}$ . This local cycle contains the edge  $\{x, y\}$  which is not a local edge. Hence, G' is not fitting for  $E^*$  if  $\operatorname{univ}_{G'}(D)$  contains x or y.

Based on these operations, we are now able to present the algorithm (see Algorithm 1) behind Theorem 4.2. Algorithm 1 works as follows: First, we apply Operations 1–5 exhaustively. Next, if there is a community  $C \in \mathcal{C}$  with  $\operatorname{fit}_{E^*}(C) = \emptyset$ , then we return that there is no fitting solution for  $E^*$ . This is correct, since all defined operations preserve Property 1. Afterwards, we start with an auxiliary graph  $G_A$  with vertex set V and edge set  $E^*$  and we iterate over the partition  $\mathfrak{C}$ . Recall that since Operation 2 is exhaustively applied, for each  $\mathcal{L} \in \mathfrak{C}$ ,  $\operatorname{fit}_{E^*}(C) = \operatorname{fit}_{E^*}(D)$  for any two communities C and D of  $\mathcal{L}$ . For each  $\mathcal{L} \in \mathfrak{C}$ , we find a vertex  $y \in \operatorname{fit}_{E^*}(C)$  that minimizes the total weight of non-local edges required to make y the center of all communities of  $\mathcal{L}$ , where C is an arbitrary community of  $\mathcal{L}$ . Finally, we add all edges between y and each vertex of any community of  $\mathcal{L}$  to  $G_A$ . After the iteration over the partition  $\mathfrak{C}$  is completed, we output  $G_A$  if it contains at most  $\ell$  edges and has total weight at most b. Otherwise, we return that there is no fitting solution for  $E^*$ . It remains to show that this greedy choice for the center vertices is correct.

#### ▶ Lemma 4.9. Algorithm 1 is correct.

**Proof.** If Algorithm 1 reaches Line 4, then  $\operatorname{fit}_{E^*}$  does not fulfill Property 1, and thus Algorithm 1 correctly outputs that there is no fitting solution for  $E^*$ . Otherwise, let  $G_A$ denote the graph constructed by Algorithm 1 and let for each community  $C \in \mathcal{C}$ , center(C)denote the vertex y chosen to be the center of all communities of  $\mathfrak{C}(C)$  in Line 9. By construction,  $G_A$  is a solution since for each community  $C \in \mathcal{C}$ , center(C) is a vertex of  $\operatorname{fit}_{E^*}(C) \subseteq \operatorname{univ}_G(C)$ . If  $G_A$  contains at most  $\ell$  edges and has total weight at most b, then the algorithm correctly outputs the solution  $G_A$  which is fitting for  $E^*$ .

Hence, in the following assume that  $G_A$  contains more than  $\ell$  edges or has weight more than b. Assume towards a contradiction that there is a fitting solution for  $E^*$ . Let  $G_F$  be a

fitting solution for  $E^*$  such that  $\operatorname{Agree}(G_F) := \{C \in \mathcal{C} : \operatorname{center}(C) \in \operatorname{univ}_{G_F}(C)\}$  is as large as possible.

**Case 1:** Agree $(G_F) = \mathcal{C}$ . By construction,  $G_A$  contains all edges of  $E^*$  and only the required edges to achieve that for each community  $C \in \mathcal{C}$ ,  $\operatorname{center}(C) \in \operatorname{univ}_{G_A}(C)$ . Consequently,  $G_A$  is a subgraph of  $G_F$  and thus  $G_F$  contains more than  $\ell$  edges or has weight more than b, a contradiction.

**Case 2:** There is a community  $C \in \mathcal{C} \setminus \operatorname{Agree}(G_F)$ . In the following, we define a fitting solution  $G'_F$  for  $E^*$  with  $\operatorname{Agree}(G'_F) \supseteq \operatorname{Agree}(G_F)$ . By definition,  $\operatorname{center}(C) = \operatorname{center}(\widetilde{C})$  for each community  $\widetilde{C} \in \mathfrak{C}(C)$ . Let  $V_C := \bigcup_{\widetilde{C} \in \mathfrak{C}(C)} \widetilde{C}$  and let  $y := \operatorname{center}(C)$ . Moreover, let x be an arbitrary vertex of  $V_C$  such that  $x \in \operatorname{univ}_{G_F}(\widetilde{C})$  for each community  $\widetilde{C} \in \mathfrak{C}(C)$ . Due to Corollary 4.5 and since  $G_F$  is fitting for  $E^*$ , this vertex exists and is unique if  $|\mathfrak{C}(C)| \ge 2$ . Note that  $C \in \mathcal{C} \setminus \operatorname{Agree}(G_F)$  implies that  $x \neq y$ . This also implies that C has size at least 3, and thus, each community of  $\mathfrak{C}(C)$  has size at least 3. We obtain  $G'_F$  as follows: First, initialize  $G'_F$  as  $G_F$ . Second, for each community  $\widetilde{C} \in \mathfrak{C}(C)$ , remove all edges that are not local edges of  $G_F[\widetilde{C}]$  from  $G'_F$ . Finally, for each community  $\widetilde{C} \in \mathfrak{C}(C)$ , add the minimum number of edges to  $G'_F$  such that  $y \in \operatorname{univ}_{G'_F}(\widetilde{C})$ , that is, the edges  $\{\{y, v\} : v \in V_C \setminus \{y\}\} \setminus E^*$ .

First, we show that  $G'_F$  contains at most as many edges as  $G_F$ . To this end, we first observe the following.

▶ Claim 5 (\*). For each  $z \in V_C \setminus \{x, y\}$ , the edge  $\{x, z\}$  is a local edge if and only if  $\{y, z\}$  is a local edge.

Proof. Assume towards a contradiction that there is some  $z \in V_C \setminus \{x, y\}$  such that exactly one of  $\{y, z\}$  and  $\{x, z\}$  is a local edge. Assume that  $\{y, z\}$  is a local edge and  $\{x, z\}$  is not a local edge. The case if  $\{x, z\}$  is a local edge and  $\{y, z\}$  is not a local edge then follows by similar arguments. Let  $\widetilde{C} \in \mathfrak{C}(C)$  be a community that contains z. Note that  $\widetilde{C}$  also contains the vertices x and y, since they are both contained in fit<sub>E\*</sub> ( $\widetilde{C}$ ). Based on the facts that  $\{y, z\}$  is a local edge and Operation 1 is exhaustively applied, x is locally universal for both y and z, since  $x \in \text{fit}_{E^*}(\widetilde{C})$ . Consequently,  $\{x, z\}$  is a local edge, a contradiction.

Recall that each edge which is in  $G'_F$  and not in  $G_F$  is incident with y and some vertex of  $V_C \setminus \{x, y\}$ . Hence, for each  $z \in V_C \setminus \{x, y\}$  where the edge  $\{y, z\}$  was added to obtain  $G'_F$ , the edge  $\{x, z\}$  was removed to obtain  $G'_F$ . Consequently,  $G'_F$  contains at most as many edges as  $G_F$ . Moreover, this implies that the difference between the total weight of  $G'_F$  and the total weight of  $G_F$  is at most  $\omega(\{\{y, z\} : z \in V_C \setminus \{y\}\} \setminus E^*) - \omega(\{\{x, z\} : z \in V_C \setminus \{x\}\} \setminus E^*)$ . Due to Line 9, this weight difference is not positive. Hence, since  $G_F$  has total weight at most b,  $G'_F$  has total weight at most b.

To show that  $G'_F$  is a solution, it thus remains to show that for each community  $C \in C$ , C has at least one center in  $G'_F$ . To show this, it suffices to show that all communities outside of  $\mathfrak{C}(C)$  have the same centers in  $G_F$  and  $G'_F$ , since y is a center of all communities of  $\mathfrak{C}(C)$ .

▶ Claim 6. For each community  $D \in C \setminus \mathfrak{C}(C)$ ,  $\operatorname{univ}_{G_F}(D) = \operatorname{univ}_{G'_F}(D)$ .

Proof. Due to symmetry, we only show that  $\operatorname{univ}_{G_F}(D) \subseteq \operatorname{univ}_{G'_F}(D)$ . Assume towards a contradiction that there is a vertex  $z \in \operatorname{univ}_{G_F}(D) \setminus \operatorname{univ}_{G'_F}(D)$ . Since  $z \notin \operatorname{univ}_{G'_F}(D)$ , there is an edge  $\{z, w\}$  which is contained in  $G_F$  but not in  $G'_F$ . Moreover,  $\{z, w\}$  is not a local edge, since  $G'_F$  contains all local edges. This further implies that there is a community  $\widetilde{C} \in \mathfrak{C}(C)$  such that  $\{z, w\} \subseteq \widetilde{C}$ . Since  $G_F$  is fitting for  $E^*$ , x is one endpoint of  $\{z, w\}$ , as otherwise,

 $\widetilde{C}$  and D induce a local cycle in  $G_F$  on the vertices  $\{x, z, w\}$  and the edge  $\{z, w\}$  is not a local edge. Next, we distinguish the cases whether  $\widetilde{C}$  contains a local edge.

**Case 1:** there is no local edge in  $\widetilde{C}$ . Since Operation 3 is exhaustively applied,  $\{x, y\} \subseteq$ fit<sub>E\*</sub>( $\widetilde{C}$ )  $\subseteq \widetilde{C} \cap D$ . Hence, if  $|\widetilde{C} \cap D| = 2$ , then x = z and y = w, or vice versa. Consequently, the edge  $\{z, w\}$  is contained in  $G'_F$ , a contradiction. Otherwise, assume  $|\widetilde{C} \cap D| \ge 3$ . We show that in this case, there is no fitting solution for  $E^*$ . Since D is not in  $\mathfrak{C}(C)$ , there is some vertex of  $\widetilde{C} \cup D$  which is locally universal for  $\widetilde{C} \cap D$ . Hence, fit<sub>E\*</sub>( $\widetilde{C}$ )  $\subseteq \widetilde{C} \cap D$ , since  $\widetilde{C}$  contains no local edge and Operation 3 is exhaustively applied. Moreover, since Operation 5 is exhaustively applied and there is no local edge between any two vertices of  $\widetilde{C} \cap D$ , fit<sub>E\*</sub>( $\widetilde{C}$ )  $\cap (\widetilde{C} \cap D) = \emptyset$ . We conclude that fit<sub>E\*</sub>( $\widetilde{C}$ ) =  $\emptyset$ , which implies that there is no fitting solution for  $E^*$ .

**Case 2:** there is some local edge in  $\widetilde{C}$ . Recall that Operation 4 and Operation 5 are applied exhaustively with respect to  $\widetilde{C}$ . If x = z and y = w, or vice versa, then the edge  $\{z, w\}$  is contained in  $G'_F$ , a contradiction. Otherwise, let  $w^*$  be the unique vertex of  $\{z, w\} \setminus \{x\}$ . Since  $\{x, w^*\} = \{z, w\}$  is not a local edge,  $x \in \operatorname{fit}_{E^*}(\widetilde{C})$ , and Operation 1 is exhaustively applied, no vertex of  $\operatorname{fit}_{E^*}(\widetilde{C})$  is locally universal for  $w^*$  and  $w^* \notin \operatorname{fit}_{E^*}(\widetilde{C})$ . Hence, if  $|\widetilde{C} \cap D| = 2$ , then since Operation 4 is exhaustively applied,  $\operatorname{fit}_{E^*}(\widetilde{C})$  has size at most one, a contradiction. Otherwise, if  $|\widetilde{C} \cap D| \geq 3$ , then since Operation 5 is exhaustively applied  $x \notin \operatorname{fit}_{E^*}(D)$  and  $w^* \notin \operatorname{fit}_{E^*}(D)$ . Consequently,  $z \notin \operatorname{fit}_{E^*}(D)$ , a contradiction.

Since  $G_F$  is a solution, for each community  $D \in \mathcal{C} \setminus \mathfrak{C}(C)$ , Claim 6 implies that  $\operatorname{univ}_{G'_F}(D) = \operatorname{univ}_{G_F}(D)$  is nonempty. Hence,  $G'_F$  is a solution. Moreover, since  $G_F$  is a minimum solution, Claim 5 implies that  $G'_F$  is a minimum solution.

Next, we show that  $G'_F$  is fitting for  $E^*$ . To show that  $G'_F$  is a fitting solution for  $E^*$ , it remains to show that each local cycle of  $G'_F$  uses only edges of  $E^*$ .

To show this, we first observe that y is the only center of each community of  $\mathfrak{C}(C)$  in  $G'_F$ .

▶ Claim 7 (\*). For each community  $D \in \mathfrak{C}(C)$ ,  $\operatorname{univ}_{G'_{F}}(D) = \{y\}$ .

Proof. Recall that each community in  $\mathfrak{C}(C)$  has size at least three. This includes the community D. Moreover, y is not a center of D in  $G_F$ , since y is not a center of C in  $G_F$  and x is the unique center of D in  $G_F$ , if  $\mathfrak{C}(C)$  has size at least two. Assume towards a contradiction, that there is a vertex  $z \in \operatorname{univ}_{G'_F}(D)$  distinct from y. By definition of  $G'_F$ , each non-local edge of  $G'_F[D]$  is incident with y. This implies that for each vertex  $w \in D \setminus \{y, z\}$ , the edge  $\{z, w\}$  is a local edge, since z is a center of D in  $G'_F$ . Since D has size at least three, this implies that each vertex of  $D \setminus \{y\}$  is incident with at least one local edge in D. Since  $y \in \operatorname{flt}_{E^*}(D)$  and Operation 1 is exhaustively applied, y is locally universal for D in  $G_F$ . Hence, y is a center of D in  $G_F$ , a contradiction.

We are now ready to show that  $G'_F$  is a fitting solution.

## ▶ Claim 8 (\*). Each local cycle of $G'_F$ uses only edges of $E^*$ .

Proof. Assume towards a contradiction that there is a local cycle in  $G'_F$  with vertex set L and edge set  $E_L$  such that  $E_L \setminus E^* \neq \emptyset$ . Let L be a smallest cycle with this property. Moreover, let Y and D be communities that induce the local cycle on the edges  $E_L$  in  $G'_F$ . First, we argue that we can assume without loss of generality that Y is contained in  $\mathfrak{C}(C)$  and that Dis not contained in  $\mathfrak{C}(C)$ .

On the one hand, due to Claim 6, at least one of Y and D is contained in  $\mathfrak{C}(C)$ , as otherwise, Y and D induce the same local cycle on the edges of  $E_L$  in  $G_F$ . On the other hand, at most one of Y and D is from  $\mathfrak{C}(C)$ , since due to Claim 7,  $\operatorname{univ}_{G_F}(Y) = \operatorname{univ}_{G_F}(D) \subseteq \{y\}$ .

Hence, assume without loss of generality that  $Y \in \mathfrak{C}(C)$  and that  $D \in \mathcal{C} \setminus \mathfrak{C}(C)$ . Let d be an arbitrary vertex of  $\operatorname{univ}_{G'_F}(D) = \operatorname{univ}_{G_F}(D)$  such that the spanning star for D with center d and the spanning star for Y with center y induce the local cycle with the vertex set L and edge set  $E_L$  in  $G'_F$ .

Let  $L' := L \setminus \{y, d\}$ . Note that  $L' \subseteq Y \subseteq V_C$ . Hence, due to Claim 5, for each  $w \in L' \setminus \{x\}$ ,  $\{x, w\}$  is a local edge if and only if  $\{y, w\}$  is a local edge. Moreover, since Operation 1 is exhaustively applied, we observe the following.

**Fact 1.** If there is at least one local edge in Y, then  $\{x, y\}$  is a local edge.

We distinguish three cases.

**Case 1:**  $x \in L \setminus L' = \{y, d\}$ . Note that this implies that x = d, since x is distinct from y. Let e be a non-local edge in  $E_L$ .

If  $e = \{x, y\} = \{d, y\}$ , then |L| = 3, since L is a smallest local cycle in  $G'_F$  containing at least one edge that is not contained in  $E^*$ . Let w be the unique vertex of L'. By construction of  $G'_F$  and since  $w \in Y$ ,  $\{x, w\}$  is a local edge. Hence, there is at least one local edge in Y and thus  $\{x, y\} = e$  is a local edge due to Fact 1, a contradiction.

Otherwise, e is incident with some vertex  $w \in L'$ . Note that the other endpoint of e is either x or y. Hence, due to Claim 5, both  $\{x, w\}$  and  $\{y, w\}$  are not local edges. By construction of  $G'_F$  and since  $w \in Y$ ,  $\{x, w\}$  is not contained in  $G'_F$ , a contradiction.

Case 2:  $x \in L'$ .

**Case 2.1:**  $L' = \{x\}$ . Recall that since Y and D induce the local cycle on the vertices L, there are at least two vertices in  $Y \cap D \cap L$ . Hence  $y \in D$  or  $d \in Y$ . If  $d \notin Y$ , then  $y \in D$  and thus  $G_F$  contains both edges  $\{d, x\}$  and  $\{d, y\}$ . Consequently, since  $\{x, y\}$  is an edge of  $G_F$ , Y and D induce the same local cycle with the edges  $E_L$ , a contradiction. Otherwise, if  $d \in Y$ , then  $\{x, d\}$  is a local edge if and only if  $\{y, d\}$  is a local edge, due to Claim 5. Hence,  $\{x, d\}$  and  $\{y, d\}$  are local edges, as otherwise,  $\{x, d\}$  is not an edge of  $G'_F$ . Consequently  $\{x, y\}$  is the unique edge of  $E_L$  which is not a local edge. Since Y contains at least one local edge, Fact 1 implies that  $\{x, y\}$  is a local edge, a contradiction.

**Case 2.2:** |L'| = 2. Let w be the unique vertex besides x in L'. Recall that since  $w \in Y$ ,  $\{x, w\}$  is an edge of  $G_F$  and that  $\{x, w\}$  is a local edge if and only if  $\{y, w\}$  is a local edge. Hence, Y and D induce a local cycle on the vertices  $\{w, x, d\}$  in  $G_F$ . Since  $G_F$  is fitting for  $E^*$ , each edge of this local cycle is a local edge, that is,  $\{x, w\}$ ,  $\{x, d\}$ , and  $\{y, w\}$  are local edges. By the above, this implies that  $\{y, w\}$  is a local edge and since Y contains at least one local edge,  $\{x, y\}$  is a local edge, due to Fact 1. Hence,  $E_L$  contains only local edges. A contradiction.

Case 3:  $x \notin L$ .

**Case 3.1:** |L'| = 1. Let w be the unique vertex of L'. Recall that since Y and D induce the local cycle on the vertices L, there are at least two vertices in  $Y \cap D \cap L$ . Hence  $y \in D$  or  $d \in Y$ . If  $y \in D$ , then Y and D induce a local cycle in  $G_F$  with the edges  $E'_L := \{\{d, y\}, \{d, w\}, \{x, y\}, \{x, w\}\}$ . Since  $G_F$  is fitting for  $E^*$ , each edge of  $E_L \cap E'_L = \{\{d, y\}, \{d, w\}\}$  is a local edge and  $\{y, w\}$  is the unique edge of  $E_L$  which is not a local edge. Hence, due to Claim 5,  $\{x, w\}$  is not a local edge and thus not all edges of  $E'_L$  are local edges, a contradiction. Otherwise, if  $d \in Y$ , then Y and D induce a local cycle in  $G_F$  with the edges  $E'_L := \{\{d, w\}, \{w, x\}, \{x, d\}\}$ . Since  $G_F$  is fitting for  $E^*$ ,  $\{d, w\} \in E_L \cap E'_L$  is a local edge. Let e be an edge of  $E_L$  which is not a local edge. By the above, e is incident with y. Let w' be the other endpoint of e. Since w' is either d or w, w' is in Y. Hence, due to Claim 5,  $\{x, w'\}$  is not a local edge and thus not all edges of  $E'_L$  are local edge. a contradiction.

**Case 3.2:** |L'| = 2. Let  $w_1$  and  $w_2$  be the two vertices of L'. Recall that since  $L' \subseteq D$ , for each  $i \in \{1, 2\}, \{x, w_i\}$  is an edge of  $G_F$  and  $\{x, w_i\}$  is a local edge if and only if  $\{y, w_i\}$  is a local edge. Hence, Y and D induce the local cycle in  $G_F$  on the edges  $E'_L := \{\{x, w_i\}, \{d, w_i\} : i \in \{1, 2\}\}$ . Since  $G_F$  is fitting for  $E^*$ , each edge of  $E_L \cap E'_L = \{\{d, w_1\}, \{d, w_2\}\}$  is a local edge. Let e be an edge of  $E_L$  which is not a local edge. By the above,  $e = \{y, w_i\}$  for some  $i \in \{1, 2\}$ . Hence,  $\{x, w_i\}$  is not a local edge and thus not all edges of  $E'_L$  are local edges, a contradiction.

Altogether, each local cycle in  $G'_F$  uses only edges of  $E^*$ .

 $\triangleleft$ 

Finally, we show that  $\operatorname{Agree}(G'_F)$  is a proper superset of  $\operatorname{Agree}(G_F)$ . By construction,  $\mathfrak{C}(C) \subseteq \operatorname{Agree}(G'_F)$ , and due to Claim 6, for each community  $D \in \mathcal{C} \setminus \mathfrak{C}(C)$ ,  $\operatorname{univ}_{G'_F}(D) =$   $\operatorname{univ}_{G_F}(D)$ . Hence,  $\operatorname{Agree}(G_F) \subseteq \operatorname{Agree}(G'_F)$ . Moreover, since  $C \notin \operatorname{Agree}(G'_F) \setminus \operatorname{Agree}(G_F)$ we obtain that  $\operatorname{Agree}(G'_F)$  is a proper superset of  $\operatorname{Agree}(G_F)$ . Altogether,  $G'_F$  is a fitting solution for  $E^*$  with  $\operatorname{Agree}(G'_F) \supseteq \operatorname{Agree}(G_F)$ . This contradicts our choice of  $G_F$ .

Hence, if  $G_A$  contains more than  $\ell$  edges or has weight more than b, then the algorithm correctly outputs that there is no fitting solution for  $E^*$ .

Hence, to show Theorem 4.2, it remains to show the running time of Algorithm 1.

**Proof of Theorem 4.2.** Clearly, the partition  $\mathfrak{C}$  of  $\mathcal{C}$  and also the initialization of  $\operatorname{ft}_{E^*}$  in Lines 1 and 2 can be computed in polynomial time. Note that Operations 1–5 can be exhaustively applied in polynomial time by iterating over all local edges and all pairs of communities, since for each community  $C \in \mathcal{C}$ ,  $\operatorname{ft}_{E^*}(C)$  initially has size at most |C| < n and each application of any operation may only remove elements from  $\operatorname{ft}_{E^*}(C)$ . Hence, Lines 3–5 can be performed in polynomial time. Afterwards, Lines 6–10 can be performed in polynomial time since for each partite set of  $\mathfrak{C}$  we compute the vertex y with minimal cost such that y serves as the center of all communities in this partite set. Finally, the check whether the solution has at most  $\ell$  edges and weight at most b can be done in polynomial time. Thus, Algorithm 1 runs in polynomial time.

Finding the correct edge set  $E^*$ : To solve STARS NWS, the main algorithmic difficulty now lies in finding an edge set  $E^*$  that contains all edges of local cycles of any optimal solution of *I*. Hence, to prove Theorem 4.1, it remains to show that such an edge set can be found in  $m^{4t} \cdot \text{poly}(n+c)$  time, if it exists.

▶ Lemma 4.10 (\*). If I is a yes-instance of STARS NWS, then for every optimal solution G' = (V, E'), there is an edge set  $E^* \subseteq E'$  of size at most 4t such that the edge set of each local cycle of G' is a subset of  $E^*$ .

**Proof.** Suppose that I is a yes-instance of STARS NWS and let G' be an optimal solution for I. Consequently, the feedback edge number of G' is at most t. Let  $\mathcal{E}$  denote the collection of edge sets of all local cycles in G' and let  $E^* := \bigcup_{\widetilde{E} \in \mathcal{E}}$ . We show that  $|E^*| \leq 4t$ . To this end, we fix some arbitrary ordering on  $\mathcal{E}$  and let  $\mathcal{E}(i)$  denote the *i*th element of  $\mathcal{E}$  according to this ordering.

For each  $i \in [1, |\mathcal{E}|]$ , we consider the edge set  $E_i$  and the graph  $G_i = (V, E_i)$ , where  $E_i$  is the union of the first i edge sets in  $\mathcal{E}$ . We show that for each  $i \in [1, |\mathcal{E}|]$ , if  $E_i$  is a proper superset of  $E_{i-1}$ , then the feedback edge number of  $G_i$  is larger than the feedback edge number of  $G_{i-1}$ . In other words, while iterating over the order of  $\mathcal{E}$ , at most t different edge sets of  $\mathcal{E}$  can introduce new edges.

Let  $i \in [1, |\mathcal{E}|]$  such that  $E_i$  is a proper superset of  $E_{i-1}$  and let  $\overline{E} := E_i \setminus E_{i-1}$ . Let  $X_i$  denote the endpoints of the edges of  $\mathcal{E}(i)$  and let  $\mathcal{V}$  denote the connected components of  $G_{i-1}$ 

containing at least one vertex of  $X_i$ . Since  $\mathcal{E}(i)$  is the edge set of a cycle,  $\widetilde{V} := \bigcup_{\widetilde{V} \in \mathcal{V}} \widehat{V}$  is a connected component in  $G_i$ , and since  $\overline{E}$  is nonempty,  $|\overline{E}| \ge |\mathcal{V}|$ . Hence, the feedback edge number of  $G_i$  is larger than the feedback edge number of  $G_{i-1}$ .

Since each edge set of  $\mathcal{E}$  has size either 3 or 4, this implies, that  $E^* := E_{|\mathcal{E}|}$  has size at most 4t.

We are finally able to show Theorem 4.1.

**Proof of Theorem 4.1.** Let  $I = (G = (V, E), C, \omega, \ell, b)$  be an instance of STARS NWS. The algorithm works as follows: For each edge set  $E^*$  of size at most 4t, use Algorithm 1 to find a solution for I with at most  $\ell$  edges and total weight at most b that contains  $E^*$  or correctly output that there is no fitting solution for  $E^*$ . If for some  $E^*$ , a solution G' with at most  $\ell$  edges and total weight at most b is found, the algorithm outputs G'. Otherwise, the algorithms outputs "no".

Note that the algorithm runs in the stated running time, since there are at most  $m^{4t}$  edges sets of size at most 4t, all of them can be enumerated in  $m^{4t} \cdot \text{poly}(|I|)$  time, and for each such edge set  $E^*$ , solving the subroutine can be done in polynomial time due to Theorem 4.2.

Finally, we show that the algorithm is correct. If I is a yes-instance of STARS NWS, let G' be an optimal solution for I. Due to Lemma 4.10, G' contains an edges set  $E^*$  of size at most 4t such that each local cycle of G' uses only edges of  $E^*$ . Hence, G' is fitting for  $E^*$ . Consequently for  $E^*$ , a solution for I with at most  $\ell$  edges and total weight at most b is outputted by the algorithm. Otherwise, if I is a no-instance of STARS NWS, then there is no solution for I with at most  $\ell$  edges and total weight at most b. Hence, each subroutine outputs "no" and the whole algorithms correctly outputs "no".

## 4.2 Connectivity NWS

Korach and Stern presented an algorithm for CONNECTIVITY NWS where G is a clique and t = 0 with running time  $\mathcal{O}(c^4 n^2)$  [33]. This result was then improved by Klemz et al. [32] who provided an  $\mathcal{O}(m \cdot (c + \log(n)))$ -time algorithm for CONNECTIVITY NWS with t = 0. Guttmann-Beck et al. [25] presented a similar algorithm for UNWEIGHTED CONNECTIVITY NWS with t = 0.

The algorithms of Klemz et al. [32] and Guttmann-Beck et al. [25] for t = 0 first construct an auxiliary edge-weight function  $\omega$  for the underlying graph G and then use Kruskal's algorithm [35] to find a minimum spanning tree with respect to  $\omega$  which, as they show, corresponds to a solution of the UNWEIGHTED CONNECTIVITY NWS instance. Recall that the communities C define a hypergraph (V(G), C). For the weighted case, Klemz et al. [32] define an isomorphic hypergraph and then use the algorithm for the unweighted case. In the following, we present an, in our opinion, simpler algorithm for CONNECTIVITY NWS with t = 0. Our approach does not need another hypergraph, instead it directly uses Kruskal's algorithm [35] by exploiting a different auxiliary weight function of the underlying graph G.

**A simpler algorithm.** We first show our result for connected hypergraphs. Let  $I = (G = (V, E), \mathcal{C}, \omega, n-1, b)$  be an instance of CONNECTIVITY NWS where the hypergraph  $\mathcal{H} = (V, \mathcal{C})$  is connected. If G is not connected, we output that there is no solution for I. Otherwise, there is at least one spanning tree for G. First, we construct an edge-weight function  $\tilde{\omega}$  for which we then apply Kruskal's algorithm [35]. The weight  $\tilde{\omega}$  of an edge e in G is the number of communities containing edge e plus  $q(e) := \frac{x-\omega(e)}{x}$ . Here, x-1 is the maximum edge weight of G. Note that  $z < \tilde{\omega}(e) < z + 1$  where z is the number of communities

containing edge e. Also observe that for two edges e and e' that are contained in the same number of communities,  $\tilde{\omega}(e) \geq \tilde{\omega}(e')$  if and only if  $\omega(e) \leq \omega(e')$ . Second, we use Kruskal's algorithm [35] to find a maximum-weight spanning tree T on G equipped with  $\tilde{\omega}$ . If this spanning tree T is a solution for I, we output T. Otherwise, we output that there is no solution for I.

▶ **Theorem 4.11** (\*). Let  $I = (G = (V, E), C, \omega, n - 1, b)$  be an instance of CONNECTIVITY NWS where the hypergraph  $\mathcal{H} = (V, C)$  is connected. Then in time  $\mathcal{O}(m \cdot (c + \log(n)))$ , the algorithm described above finds a minimum-weight solution for I or correctly outputs that there is no solution for I.

**Proof.** Observe that the correctness follows, if the following statement is shown: There exists a solution G' for I containing the edges in L of total weight b' if and only if there exists a solution  $G^*$  for I containing the edges in  $L \cup \{\{u, v\}\}$  of weight b'. Here, b' is the minimal weight of any solution and  $\{u, v\}$  is the next edge chosen by Kruskal's algorithm, after exactly the edge of L have been chosen so far.

 $(\Rightarrow)$  Let G' be a solution containing the edges L. We now show that there also exists a solution  $G^*$  containing the edges  $L \cup \{\{u, v\}\}$ . If  $\{u, v\} \in E(G')$ , nothing is to show and thus, in the following we assume that  $\{u, v\} \notin E(G')$ .

Let  $\widetilde{\omega}(\{u,v\}) = y$  and let  $z \coloneqq \lfloor y \rfloor$ . By definition of  $\widetilde{\omega}$ , there exist exactly z pairwise distinct communities  $C_i$  with  $u \in C_i$  and  $v \in C_i$  for each  $i \in [1, z]$ . Since  $\{u, v\} \notin E(G')$ , for each  $i \in [1, z]$  there exists a path  $P_i \coloneqq (u = p_1^i, p_2^i, \dots, p_z^i = v)$  such that each edge of  $P_i$  is contained in E(G'). Since G' is acyclic, we observe that  $P_{i_1} = P_{i_2}$  for each two communities  $C_{i_1}$  and  $C_{i_2}$  containing u and v. Now, let P be the unique path in G' connecting uand v. Observe that  $V(P) \subseteq C_i$  for each  $i \in [1, z]$ . Thus,  $\widetilde{\omega}(e) \ge z$  for each edge e on P.

Recall that  $\{u, v\}$  is an edge chosen by Kruskal's algorithm. Thus,  $L \cup \{\{u, v\}\}$  is acyclic in G. Hence,  $E(P) \not\subseteq L$  and there exists at least one edge  $\{a, p\} \in E(P)$  which is not contained in L. Observe that since  $\{u, v\}$  is chosen by Kruskal's algorithm, each other edge e, such that  $L \cup \{e\}$  is acyclic, has weight at most y. Thus,  $z \leq \tilde{\omega}(\{a, p\}) \leq y$ . Recall that  $\lfloor \tilde{\omega}(\{a, p\}) \rfloor = z = \lfloor \tilde{\omega}(\{u, v\}) \rfloor$  and that we have  $\{a, p, u, v\} \subseteq C_i$  for each  $i \in [1, z]$ . In other words,  $\{u, v\}$  and  $\{a, p\}$  are contained in the same collection of communities.

Now, we can replace the edge  $\{a, p\}$  of the solution G' with the edge  $\{u, v\}$ . Since each community, which contains a and p, contains u and v as well, we obtain a solution  $G^*$  containing the edges  $L \cup \{\{u, v\}\}$ . Since  $\widetilde{\omega}(\{u, v\}) \ge \widetilde{\omega}(\{a, p\})$  and  $\lfloor \widetilde{\omega}(\{a, p\}) \rfloor = z = \lfloor \widetilde{\omega}(\{u, v\}) \rfloor$ , we have  $q(\{u, v\}) \ge q(\{a, p\})$  and thus  $\omega(\{u, v\}) \le \omega(\{a, p\})$ . Since G' has total weight b', we thus conclude that  $G^*$  also has weight b'.

( $\Leftarrow$ ) If there exists a solution  $G^*$  for  $(G, \mathcal{C}, \ell)$  of weight b' containing the edges in  $L \cup \{\{u, v\}\}$ , then  $G^*$  clearly also contains the edges in L.

**Running Time:** Initially, we check whether G is connected in  $\mathcal{O}(n+m)$  time. Afterwards, in  $\mathcal{O}(m \cdot |\mathcal{C}|)$  time we compute the edge-weight function  $\tilde{\omega}$ , followed by applying Kruskal's algorithm in  $\mathcal{O}(m \cdot \log(n))$  time [35], to find a maximum weight spanning tree T for G with respect to  $\tilde{\omega}$ . Finally, we check whether the spanning tree T is a solution in  $\mathcal{O}(m \cdot |\mathcal{C}|)$  time. Hence, we obtain the stated overall running time.

In Theorem 4.11 the instances of CONNECTIVITY NWS are restricted to connected hypergraphs. Next, we generalize the algorithm to hypergraphs with any number of connected components.

▶ Corollary 4.12 (\*). Let  $I = (G = (V, E), C, \omega, n - x, b)$  be an instance of CONNECTIVITY NWS where x is the number of connected components of the hypergraph  $\mathcal{H} = (V, C)$ . Such an instance I is solvable in  $\mathcal{O}(m \cdot (c + \log(n)))$  time.

**Proof.** Initially, we split the hypergraph  $\mathcal{H} = (V, \mathcal{C})$  into its connected components  $\mathcal{H}_1 = (V_1, \mathcal{C}_1), \ldots, \mathcal{H}_x = (V_x, \mathcal{C}_x)$ . For each of these components  $\mathcal{H}_i$  we use the algorithm described in Theorem 4.11 to compute the minimal weight  $b_i$  of a solution for  $\mathcal{H}_i$  which has exactly  $|V_i| - 1$  edges. If any  $\mathcal{H}_i$  does not have a solution which is a tree, we output no. Otherwise, we calculate the sum B of the minimal weights  $b_i$ . If  $B \leq b$ , we output the union of the solutions of all components  $\mathcal{H}_i$ . Otherwise, we output no,

Note that this algorithm is correct. For the overall running time, note that the connected components  $\mathcal{H}_i$  can be determined in linear time. Then, according to Theorem 4.11, the component  $\mathcal{H}_i$  can be solved in  $\mathcal{O}(m_i \cdot (c_i + \log(n_i)))$  time, where  $m_i$  is the number of edges of the subgraph induced by  $V_i$ ,  $c_i = |\mathcal{C}_i|$ , and  $n_i = |V_i|$ . Thus, the instance I of CONNECTIVITY NWS is solvable in  $\mathcal{O}(m \cdot (c + \log(n)))$  time.

Next, we show that the positive result for t = 0 cannot be lifted to t = 1; in this case CONNECTIVITY NWS is NP-hard. We obtain our result by reducing from the NP-hard HAMILTONIAN CYCLE-problem [1, 22], which asks for a given graph G = (V, E) if there is a *Hamiltonian cycle* in G, that is, a cycle containing each vertex of G exactly once.

▶ **Theorem 4.13.** Let  $\Pi$  be a graph class on which HAMILTONIAN CYCLE is NP-hard, then UNWEIGHTED CONNECTIVITY NWS is NP-complete on  $\Pi$  even if t = 1.

**Proof.** Let I := (V, E) be an instance of HAMILTONIAN CYCLE containing at least three vertices. We obtain an equivalent instance  $I' := (G = (V, E), C, \ell)$  of UNWEIGHTED CONNECTIVITY NWS as follows: We start with an empty set C and add for each vertex  $v \in V$  a community  $C_v := V \setminus \{v\}$  to C. Finally, we set  $\ell := |V|$ .

Note that  $t = \ell - n + x$ , where x is the number of connected components of the graph. Thus, t = n - n + 1 = 1.

**Correctness:** We show that I is a yes-instance of HAMILTONIAN CYCLE if and only if I' is a yes-instance of UNWEIGHTED CONNECTIVITY NWS.

(⇒) Suppose that there is a Hamiltonian cycle in I. Let  $E_C$  be the edges of this Hamiltonian cycle and let  $G_C := (V, E_C)$ . We show that  $G_C$  is a solution for I'. Since  $C_v := V \setminus \{v\}$  and  $G_C$  is a cycle the subgraph  $G_C[C_v]$  is connected. Moreover,  $G_C$  contains  $|V| = \ell$  edges. Hence, I' is a yes-instance of UNWEIGHTED CONNECTIVITY NWS.

( $\Leftarrow$ ) Let  $G_C := (V, E_C)$  be a solution of I'. We show that  $G_C$  is a Hamiltonian cycle. To this end, we show that each vertex  $v \in V$  is incident with at least two edges in  $G_C$ . Assume towards a contradiction that there is a vertex  $v \in V$  which is incident with at most one edge in  $G_C$ . Since v is contained in each community of  $C \setminus \{C_v\}$  and each community of Chas size  $|V| - 1 \ge 2$ , v is incident with at least one edge of  $G_C$ . Suppose that w is the unique neighbor of v in  $G_C$ . Hence, v has no neighbor in the community  $C_w = V \setminus \{w\}$  and thus  $G_C[C_w]$  is not connected, a contradiction. Consequently, each vertex of V is incident with at least two edges in  $G_C$ . Since  $G_C$  contains exactly  $\ell = |V|$  edges, this implies that  $G_C$ is a Hamiltonian cycle. Thus, I is a yes-instance of HAMILTONIAN CYCLE.

Since HAMILTONIAN CYCLE is NP-hard on cubic bipartite planar graphs [1], we obtain the following.

▶ Corollary 4.14. UNWEIGHTED CONNECTIVITY NWS is NP-complete even if t = 1 on subcubic bipartite planar graphs.

## 5 Stars NWS Parameterized by the Number of Communities

UNWEIGHTED CONNECTIVITY NWS is NP-hard even for c = 7 [21, Proposition 4]. In contrast, STARS NWS admits an XP-algorithm for c with running time  $n^{\mathcal{O}(c)}$ : For each community C, test each of the at most  $|C| \leq n$  potential center vertices. Then, for each potential solution check whether it consists of at most  $\ell$  edges of total weight at most b. For STARS NWS, we show that it is unlikely that this brute-force algorithm can be improved, by showing W[1]-hardness. For UNWEIGHTED STARS NWS, we obtain an FPT-algorithm for c.

## 5.1 W[1]-hardness for Stars NWS

We first show that the simple  $n^{\mathcal{O}(c)}$ -time algorithm for STARS NWS cannot be lifted to an FPT-algorithm.

▶ Theorem 5.1 (\*). STARS NWS is W[1]-hard when parameterized by c even if G is a clique and each edge weight is 1 or 2.

**Proof.** We provide a parameter-preserving reduction from the W[1]-hard REGULAR MULTI-COLORED CLIQUE problem [13]. The input consists of an *r*-regular graph G, an integer  $\kappa$ , and a partition  $(V_1, \ldots V_{\kappa})$  of V(G). The question is whether there exists a clique of size  $\kappa$ containing exactly one vertex of each partite set  $V_i$ .

We construct an equivalent instance  $(G', \mathcal{C}, \omega, \ell, b)$  of STARS NWS as follows. The vertex set of V(G') consists of a copy of V(G) and  $\kappa$  additional vertex sets  $S_i$ ,  $i \in [\kappa]$ , each of size  $n(G)^3$ . We make G' a clique by adding all edges between vertices of V(G'). To complete the construction, we specify the communities and edge weights. First, for each color class  $i \in$  $[1, \kappa]$ , we add a community  $C_i := V(G) \cup S_i$ . Afterwards, we define the edge weights: For each edge  $\{a, b\} \in E(G')$  such that  $\{a, b\} \in E(G)$ , we set  $\omega(\{a, b\}) := 2$ , for each edge  $\{a, b\}$ with  $a \in S_i$  and  $b \notin V_i$ , we set  $\omega(\{a, b\}) := 2$ , for each edge  $\{a, b\}$  with  $a \in S_i$  and  $b \in S_j$ , we set  $\omega(\{a, b\}) := 2$ , and for each remaining edge  $\{a, b\} \in E(G')$  we set  $\omega(\{a, b\}) := 1$ . Finally, we set  $\ell := \kappa \cdot (n(G)^3 + n(G) - 1) - {\kappa \choose 2}$  and  $b := \kappa \cdot (n(G)^3 + n(G) - 1 + r) - 2{\kappa \choose 2}$ . Note that  $c = \kappa$ , it thus remains to show the equivalence of the two instances.

(⇒) Let  $K := \{v_i : v_i \in V_i\}$  be a multicolored clique in G. We construct a solution of  $(G', \mathcal{C}, \omega, \ell, b)$  as follows. For community  $C_i$  we choose vertex  $v_i$  as its center. Observe that each community has exactly  $n(G)^3 + n(G)$  vertices. Thus, in the solution  $n(G)^3 + n(G) - 1$  edges are contained in each community. Some edges are, however, counted twice: Since the centers of all communities are disjoint and since the center of community  $C_i$  is contained in each other community we count each edge where both endpoints are centers twice. Thus, the solution consists of exactly  $\kappa \cdot (n(G)^3 + n(G) - 1) - {\kappa \choose 2}$  edges. Observe that since the graph G is r-regular and since  $v_i \in V_i$ , the weight of all edges contained in the spanning star of community  $C_i$  is exactly  $n(G)^3 + n(G) - 1 + r$ . The total weight of the solution is thus  $\kappa \cdot (n(G)^3 + n(G) - 1 + r)$  minus the weight of the edges where both endpoints are centers since these edges are counted twice in the sum. Since K is a clique, all these edges have weight 2. Thus, the solution has weight  $\kappa \cdot (n(G)^3 + n(G) - 1 + r) - 2{\kappa \choose 2} = b$ .

 $(\Leftarrow)$  Let G'' be a solution of  $(G', \mathcal{C}, \omega, \ell, b)$ . First, observe that for each  $S_i$ , the solution contains at least  $n(G)^3 - 1$  edges with one endpoint in  $S_i$  and the other endpoint in  $V(G) \cup S_i$ . Let  $E_i$  denote this edge set for each  $S_i$ . Since these edge sets  $E_i$  are pairwise disjoint, the total weight of these edges in  $E_1 \cup \ldots \cup E_{\kappa}$  is at least  $\kappa \cdot (n(G)^3 - 1)$ . Now, if for some  $C_i$ , the center vertex is either contained in  $S_i$  or in  $V(G) \setminus V_i$ , then the set  $E_i$  has weight at least  $2 \cdot (n(G)^3 - 1)$ . Thus, in that case the total edge weight of  $E_1 \cup \ldots \cup E_{\kappa}$  is at least  $(\kappa + 1) \cdot (n(G)^3 - 1)$  which exceeds b since  $n^3(G) > \kappa \cdot (n(G) + r)$ . As a consequence,

the center the center  $c_i$  of each community  $C_i$  is contained in the color class  $V_i$ . We show that  $K := \{c_i : i \in [1, \kappa]\}$  is a clique of size  $\kappa$  in G. First, observe that since each center  $c_i$ has edge weight one to all vertices in  $C_i$  except to its r neighbors in G, the sum of the edge weights in the spanning star of  $C_i$  is exactly  $n(G)^3 + n(G) - 1 + r$ . Hence, the total edge weight of all spanning stars is  $\kappa \cdot (n(G)^3 + n(G) - 1 + r)$  minus the weight of all edges that are contained in multiple spanning stars. Observe that these edges are exactly the  $\binom{\kappa}{2}$  edges between the  $\kappa$  distinct center vertices. Now since  $b = \kappa \cdot (n^3 + n(G) - 1 + r) - 2\binom{\kappa}{2}$ , the sum of all weights of edges which are counted twice is  $2\binom{\kappa}{2}$ . Hence, each edge that is contained in multiple spanning stars has weight 2. Thus, each edge  $\{c_i, c_j\}$  between two centers  $c_i$  and  $c_j$ has weight 2 in G'. Hence,  $\{c_i, c_j\}$  is an edge in G, and we may conclude that K is a clique of size  $\kappa$  in G.

## 5.2 Analysis of Unweighted Stars NWS

Next, we show that, in contrast to the weighted case, UNWEIGHTED STARS NWS admits an FPT-algorithm with respect to c. We also show that a polynomial kernel for c is unlikely.

▶ Theorem 5.2 (\*). UNWEIGHTED STARS NWS is solvable in  $\mathcal{O}(4^{c^2} \cdot (n+m) + n^2 \cdot c)$  time.

**Proof.** Our FPT-algorithm relies on branching on a specific partition of the vertices. We say that two vertices u and w are *center twins* if a) u and w are contained in the same set of communities  $\mathcal{C}' \subseteq \mathcal{C}$  and if b) both vertices are *potential centers* of the same set of communities  $\mathcal{D}' \subseteq \mathcal{C}' \subseteq \mathcal{C}$ . Here, a vertex v is a potential center for a community C, if both  $v \in C$  is a vertex of C and each other vertex of C is adjacent to v. Based on center twins, we can partition the vertex set V(G) into sets  $\mathcal{T} \coloneqq \{T_1, \ldots, T_p\}$ , that is, all vertices in each set  $T_i$  are center twins. Next, we make an observation about yes-instances of UNWEIGHTED STARS NWS regarding center twins.

▶ Claim 9. Let I be an yes-instance of UNWEIGHTED STARS NWS with a minimal solution G' = (V, E') where  $\operatorname{cen}_{G'} : \mathcal{C} \to V(G)$  denotes a mapping of each community  $C \in \mathcal{C}$ to one vertex of C which is universal for C in G'. Let  $u \in V(G)$  be a vertex, let  $T_u$  be the partite set of  $\mathcal{T}$  containing u, and let  $\mathcal{C}_u \subseteq \mathcal{C}$  be the set of communities having its center, with respect to  $\operatorname{cen}_{G'}$ , in  $T_u$ . Then, there exists a solution G'' = (V, E'') with  $|E''| \leq |E'|$ such that u is universal for each community of  $\mathcal{C}_u$  in G'' and for each community  $C \in \mathcal{C} \setminus \mathcal{C}_u$ ,  $\operatorname{cen}_{G'}(C)$  is universal for C in G''.

Proof. We define a new mapping cen" of each community  $C \in \mathcal{C}$  to one vertex of C which serves as a center for C as follows: for each community  $C \in \mathcal{C}_u$ , that is, each community having its center cen<sub>G'</sub>(C) in  $T_u$ , we set cen"(C) := u, and for each remaining community C, that is,  $C \in \mathcal{C} \setminus \mathcal{C}_u$ , we set cen"(C) := cen<sub>G'</sub>(C). Next, we define based on the mapping cen" a new solution G'' := (V, E''). We set  $E'' := \{\{\operatorname{cen}''(D), v\} : D \in \mathcal{C}, v \in D \setminus \{\operatorname{cen}''(D)\}\}$ . Observe that G''[D] contains a spanning star for each community D.

Thus, it remains to show that  $|E''| \leq |E'|$ . To show this statement, we show that for each edge  $\{x, b\} \in E'' \setminus E'$  there exists a corresponding edge in  $E' \setminus E''$  which accounts for  $\{x, b\}$ . Observe that G' and G'' only differ in the centers of all communities in  $\mathcal{C}_u$ . Hence, one of the endpoints of  $\{x, b\}$ , say x, is u. Next, we consider all possibilities for b.

**Case 1:** b is not a center of any community in C with respect to G'. Since  $\{u, b\} \in E''$ , there exists a community  $C \in C$  such that  $\operatorname{cen}''(C) = u, b \in C$ , and  $\operatorname{cen}''(C) = w \neq u$ . Note that  $C \in C_u$ . Since b is no center of any community in G', we observe that  $\{w, b\} \in E' \setminus E''$  and thus  $\{w, b\}$  accounts for  $\{u, b\}$ .

**Algorithm 2** Algorithm for UNWEIGHTED STARS NWS: SolveSNWS Input :  $G = (V, E), C, \ell, E'$ **Output:** A solution G' with at most  $\ell$  edges which contains all edges in E', or no 1 if  $C = \emptyset$  then if  $\ell < |E'|$  then 2 return no 3 return G' = (V, E')4 **5**  $C \leftarrow$  pick community of C6 forall  $T \in \mathcal{T}$  such that  $\operatorname{cand}(T)$  is a potential center for C do  $E'' \leftarrow E' \cup \{ \{ \operatorname{cand}(T), v \} : v \in C \setminus \{ \operatorname{cand}(T) \} \}$ 7 if  $SolveSNWS(G, \mathcal{C} \setminus \{C\}, \ell, E'')$  returns a graph G' then 8 return G'9 10 return no

**Case 2:** *b* is the center of some community  $C_b$  with respect to G'. First, we consider that case that  $C_b \in C_u$ . Since *u* and *b* are center twins, we have  $u \in C_b$ . Thus,  $\{u, b\}$  is contained in E' and in E'' as well. Second, we consider the case that  $C_b \in C \setminus C_u$ . Let  $C_w$  be a community in  $C_u$  whose center is moved from *w* in G' to *u* in G''. If there is no community  $C_b \in C \setminus C_u$  which contains *u* and thus also not *w*, then, similar to Case 1,  $\{w, b\}$  is an edge in  $E' \setminus E''$  and thus  $\{w, b\}$  accounts for  $\{u, b\}$ . This is possible since *u* and *w* are center twins. Hence, in the following we can safely assume that there is a community  $C_b \in C \setminus C_u$  which contains *u* and thus also *w*. Since *b* is the center of  $C_b$  and since  $u, w \in C_b$ , we observe that both  $\{u, b\}$  and  $\{w, b\}$  are contained in  $E' \cap E''$ .

Hence,  $|E''| \leq |E'|$ .

Algorithm: Let  $\mathcal{T}$  be a partitioning into center twins. According to Claim 9, we know that if there is a solution with at most  $\ell$  edges, than there is also a solution with at most  $\ell$  edges such that all communities having its center in some  $T_i \in \mathcal{T}$  have the same center. Thus, we can safely assume that all communities C having their center in  $T_i$  have the same arbitrary but fixed vertex cand $(T_i) \in T_i$  as their center.

Our depth-bounded search tree algorithm SolveSNWS is shown in Algorithm 2. We branch for each community  $C \in \mathcal{C}$  and each partite set T of  $\mathcal{T}$ , whether the center of C is in T, that is, whether cand(T) is the center of T. Note that it is a necessary that cand(T) is a potential center of C. After a center has been selected for each community, it is checked whether the resulting solution has at most  $\ell$  edges. If some branch leads to a solution, then Iis a yes-instance of UNWEIGHTED STARS NWS, and otherwise, if no branch leads to a solution with at most  $\ell$  edges, then I is a no-instance of UNWEIGHTED STARS NWS.

**Correctness:** We show that I is a yes-instance of UNWEIGHTED STARS NWS if and only if the algorithm returns a graph.

 $(\Rightarrow)$  Let *I* be a yes-instance of UNWEIGHTED STARS NWS and let G' = (V, E') be a solution. By applying Claim 9, we are able to obtain a graph G'' = (V, E'') such that  $\operatorname{cen}_{G''}(C_i) = \operatorname{cen}_{G''}(C_j)$  for all pairs of communities  $C_i, C_j \in \mathcal{C}$  where  $c_{G'}(C_i)$  and  $c_{G'}(C_j)$ are center twins. Observe that G'' is found by traversing the search tree built by Algorithm 2 and selecting center cand(*T*) for each center twin class.

 $(\Leftarrow)$  Let G' be the solution returned by Algorithm 2 in Line 4. The conditional statement in Line 2 ensures that G' has at most  $\ell$  edges. The termination condition in Line 1 ensures together with the statement in Line 7 that for each community  $C_i \in \mathcal{C}$  the induced

 $\triangleleft$ 



**Figure 6** Sketch of the construction of Theorem 5.3. The left side shows the HITTING SET instance, and the right side shows the STARS NWS instance. The grey vertices are the new vertices in the set Z. The different line styles show the mapping between the sets in the HITTING SET instance and the communities in the STARS NWS instance.

subgraph  $G'[C_i]$  contains a spanning star. Hence, a solution with at most  $\ell$  edges has been found by the algorithm which implies that I is a yes-instance of UNWEIGHTED STARS NWS.

**Running time:** Let  $C \in \mathcal{C}$  be the community selected for branching in Line 5. The branching vector of the branching in the loop in Lines 6-9 has at most  $|\mathcal{T}|$  entries of value 1, each decreasing the number of communities by 1. We now bound  $|\mathcal{T}|$ : the vertex set V(G) can be partitioned into at most  $2^c$  sets according to part a) of the definition of center twins, that is, vertices belonging to the same set of communities. Then, for each such set there are up to  $2^c$  possibilities on how to partition it according to part b) of the definition of center twins. Thus,  $|\mathcal{T}| \leq 2^c \cdot 2^c = 4^c$ . Since the maximum depth of the search tree is bounded by c, the search tree has size  $\mathcal{O}(4^{c^2})$ . Since the partition  $\mathcal{T}$  is computable in  $\mathcal{O}(n^2 \cdot c)$  time and since the edge set in Line 7 is computable in  $\mathcal{O}(n+m)$  time, the overall running time of  $\mathcal{O}(4^{c^2} \cdot (n+m) + n^2 \cdot c)$  follows.

To complete the parameterized complexity picture, we show that a polynomial kernel for c is unlikely.

▶ **Theorem 5.3** (\*). UNWEIGHTED STARS NWS parameterized by c does not admit a polynomial kernel unless  $NP \subseteq coNP/poly$ .

**Proof.** We give a polynomial parameter transformation from HITTING SET which does not admit a polynomial kernel when parameterized by the number  $|\mathcal{F}|$  of sets, unless NP  $\subseteq$  coNP/poly [13]. Let  $I_{\text{HS}} \coloneqq (U, \mathcal{F}, k)$  be an instance of HITTING SET. We assume that  $|U| \ge 2$ .

Now, we construct an equivalent instance of UNWEIGHTED STARS NWS. An example of the construction is shown in Figure 6. We start by defining the corresponding graph G. First, we add a copy of U to V(G), and second, we add a vertex set Z of size  $|U|^3$  to V(G). Next, we add edges, such that U is a clique and such that each vertex in U is adjacent to each vertex in Z. Now, we define the communities: for each set  $F \in \mathcal{F}$ , we add a community  $C_F := \{F \cup Z\}$ . Finally, we set the parameter  $\ell := k \cdot |U|^3 + |U|^2$ . Let  $I_{\text{SNS}}$  denote the resulting instance of STARS NWS. Note that  $c = |\mathcal{F}|, |E| = {\binom{|U|}{2}} + |U|^4$  and  $|V| = |U| + |U|^3$ .

**Correctness:** We show that  $I_{\text{HS}}$  is a yes-instance of HITTING SET if and only if  $I_{\text{SNS}}$  is a yes-instance of UNWEIGHTED STARS NWS.

(⇒) Let S be a hitting set of size at most k. We show how to obtain a solution G' = (V, E')with  $|E'| \leq \ell$ . We set  $E' := \{\{s, z\} : s \in S, z \in Z\} \cup \{\{u, s\} : s \in S, u \in U \setminus \{s\}\}$ . Observe that  $|E'| \leq k \cdot (|Z| + |U| - 1) \leq k \cdot |U|^3 + |U|^2$ . Recall that for each community  $C_F \in \mathcal{C}$  there exists a set  $F \in \mathcal{F}$  such that  $C_F = F \cup Z$ . Since S is a hitting set, for each community  $C_F$ ,

there exists at least one element  $s \in S$  with  $s \in C_F$ . Such an element s is the center of a spanning star in  $G'[C_F]$  because  $\{\{s,z\} : z \in Z\} \subseteq E'$  and  $\{\{s,y\} : y \in C_F \setminus Z\} \subseteq$  $\{\{s,u\} : u \in U\} \subseteq E'$ . Thus, G' is a solution which implies that  $I_{SNS}$  is a yes-instance of UNWEIGHTED STARS NWS.

( $\Leftarrow$ ) Let  $I_{\text{SNS}}$  be a yes-instance of UNWEIGHTED STARS NWS and let G' = (V, E') be a solution where cen :  $\mathcal{C} \to V$  denotes the mapping of communities to some center vertex in G'. We set  $S \coloneqq \{\text{cen}(C_F) : C_F \in \mathcal{C}\}$  and show that S is a hitting set for  $I_{\text{HS}}$  of size at most k. Since  $Z \subseteq C_F$  and Z is an independent set in G, we obtain that  $\text{cen}(C_F) \notin Z$ for each community  $C_F \in \mathcal{C}$ . This implies that  $S \subseteq U$ . Since  $|E'| \leq \ell = k \cdot |U|^3 + |U|^2$ and  $|U| \geq 2$ , we conclude that  $|S| \leq k$ . Since for each  $F \in \mathcal{F}$ , there exists a community  $C_F$ such that  $F = C_F \setminus Z$ , we conclude that for each  $C_F \in \mathcal{F}$  there exists at least one element u = cen(C) with  $u \in S$  and  $u \in F$ . Thus, X is a hitting set with  $|S| \leq k$  and  $I_{\text{HS}}$  is a yes-instance of HITTING SET.

## 6 Conclusion

Presumably the most interesting open question is whether STARS NWS parameterized by t admits an FPT-algorithm. In Theorem 4.2 we showed that STARS NWS can be solved in polynomial time if for some optimal solution the edge set of all local cycles is known. Hence, to obtain an FPT-algorithm, it would be sufficient to find such an edge set in FPT-time. Moreover, it is open whether UNWEIGHTED CONNECTIVITY NWS can be solved in polynomial time when t is constant and the input graph is a clique. In other words, it is open whether a minimum-edge hypergraph support can be found in polynomial time when it has a constant feedback edge number.

It would also be interesting to close the gap between the running time lower bound of  $2^{\Omega(c)} \cdot \text{poly}(|I|)$  (see Proposition 2.5) and the upper bound of  $2^{\mathcal{O}(c^2)} \cdot \text{poly}(|I|)$  (see Theorem 5.2) for UNWEIGHTED STARS NWS.

Finally, it is interesting to study  $\Pi$ -NWS for other graph properties. For example, Fluschnik and Kellerhals [20] studied a variant of UNWEIGHTED  $\Pi$ -NWS, denoted as *d*-DIAM NWS, where the subgraph induced by each community is required to have diameter at most *d*. They showed that the problem is linear-time solvable for d = 1 and that it is NP-hard even if c = 1 and d = 2. Furthermore, Wallisch [40] showed that 2-DIAM NWS is NP-hard even if  $\Delta = 5$  and each community has size at most 3.

Our XP-algorithm for STARS NWS parameterized by t implies that d-DIAM NWS can be solved in polynomial time when d = 2 and we are searching for a forest, that is, when t = 0because in that case, stars are the only possibility to achieve diameter 2. Can this positive result be extended to an XP-algorithm for t? Is it also possible to achieve a polynomial-time algorithm for t = 0 when  $d \ge 3$ ?

The proofs of our ETH-bounds for UNWEIGHTED CONNECTIVITY NWS and UNWEIGHTED STARS NWS can also be used to exclude  $2^{o(n^2+c)} \cdot \text{poly}(n+c)$  time algorithms for *d*-DIAM NWS: Recall that in Corollary 3.3 we showed this lower bound for UNWEIGHTED CONNECTIVITY NWS even if each community has size at most 4. Since for communities of size at most 4 the properties of being connected and having diameter at most *d* for  $d \ge 3$ coincide, the proof of Corollary 3.3 directly implies the same running time lower bound for *d*-DIAM NWS with  $d \ge 3$ . For the case d = 2, the proof of Theorem 3.1 can be used to obtain the lower bound: For communities of size at most 3, the properties of having a spanning star and having diameter at most 2 coincide. Communities of size 4 with diameter 2 either have a spanning star or contain a cycle of length 4. In the instance constructed

in Theorem 3.1, however, the solution may not contain such cycles since the budget is tight. Hence, we obtain the following.

▶ Corollary 6.1. If the ETH is true, then d-DIAM NWS, for each  $d \ge 2$ , cannot be solved in  $2^{o(n^2+c)} \cdot \operatorname{poly}(n+c)$  time even if G is a clique and each community has size at most 4.

In light of these further hardness results we may ask the following: are there properties  $\Pi$  such that  $\Pi$ -NWS is NP-hard but can be solved in  $2^{\mathcal{O}(n)} \cdot \operatorname{poly}(n+c)$  time?

#### — References –

- 1 Takanori Akiyama, Takao Nishizeki, and Nobuji Saito. NP-completeness of the Hamiltonian cycle problem for bipartite graphs. *Journal of Information Processing*, 3(2):73–76, 1980.
- 2 Dana Angluin, James Aspnes, and Lev Reyzin. Network construction with subgraph connectivity constraints. Journal of Combinatorial Optimization, 29(2):418–432, 2015.
- 3 Catriel Beeri, Ronald Fagin, David Maier, and Mihalis Yannakakis. On the desirability of acyclic database schemes. *Journal of the ACM*, 30(3):479–513, 1983.
- 4 Piotr Berman, Marek Karpinski, and Alex D. Scott. Approximation hardness of short symmetric instances of MAX-3SAT. *Electronic Colloquium on Computational Complexity*, TR03-049, 2003. URL: https://eccc.weizmann.ac.il/eccc-reports/2003/TR03-049/ index.html, arXiv:TR03-049.
- 5 Édouard Bonnet, Diana-Elena Fălămaş, and Rémi Watrigant. Constraint generation algorithm for the minimum connectivity inference problem. In Proceedings of the International Symposium on Experimental Algorithms (SEA<sup>2</sup> '19), volume 11544 of Lecture Notes in Computer Science, pages 167–183. Springer, 2019.
- 6 Ulrik Brandes, Sabine Cornelsen, Barbara Pampel, and Arnaud Sallaberry. Path-based supports for hypergraphs. *Journal of Discrete Algorithms*, 14:248–261, 2012.
- 7 Kevin Buchin, Marc J. van Kreveld, Henk Meijer, Bettina Speckmann, and Kevin Verbeek. On planar supports for hypergraphs. *Journal of Graph Algorithms and Applications*, 15(4):533–549, 2011.
- 8 Chandra Chekuri, Thapanapong Rukkanchanunt, and Chao Xu. On element-connectivity preserving graph simplification. In *Proceedings of the 23rd Annual European Symposium on Algorithms (ESA '15)*, volume 9294 of *Lecture Notes in Computer Science*, pages 313–324. Springer, 2015.
- 9 Jiehua Chen, Christian Komusiewicz, Rolf Niedermeier, Manuel Sorge, Ondrej Suchy, and Mathias Weller. Polynomial-time data reduction for the subset interconnection design problem. SIAM Journal on Discrete Mathematics, 29(1):1–25, 2015.
- 10 Gregory V. Chockler, Roie Melamed, Yoav Tock, and Roman Vitenberg. Constructing scalable overlays for pub-sub with many topics. In Proceedings of the Twenty-Sixth Annual ACM Symposium on Principles of Distributed Computing (PODC '07), pages 109–118. ACM, 2007.
- 11 Nathann Cohen, Frédéric Havet, Dorian Mazauric, Ignasi Sau, and Rémi Watrigant. Complexity dichotomies for the minimum F-overlay problem. Journal of Discrete Algorithms, 52-53:133– 142, 2018.
- 12 Vincent Conitzer, Jonathan Derryberry, and Tuomas Sandholm. Combinatorial auctions with structured item graphs. In Deborah L. McGuinness and George Ferguson, editors, *Proceedings* of the Nineteenth National Conference on Artificial Intelligence (AAAI '04), pages 212–218. AAAI Press / The MIT Press, 2004.
- 13 Marek Cygan, Fedor V. Fomin, Lukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michal Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015.
- 14 Muhammad Abid Dar, Andreas Fischer, John Martinovic, and Guntram Scheithauer. Integer linear programming formulations for the minimum connectivity inference problem and model reduction principles. *Discrete Optimization*, 40:100623, 2021.

- **15** Rodney G. Downey and Michael Ralph Fellows. *Fundamentals of Parameterized Complexity*. Springer Science & Business Media, 2013.
- 16 Ding-Zhu Du and Zevi Miller. Matroids and subset interconnection design. SIAM Journal on Discrete Mathematics, 1(4):416–424, 1988.
- 17 Pierre Duchet. Propriete de Helly et problemes de representation. Colloque International CNRS, Problemes Conbinatoires et Theorie du Graphs, 260:117–118, 1978.
- 18 Hongbing Fan, Christian Hundt, Yu-Liang Wu, and Jason Ernst. Algorithms and implementation for interconnection graph problem. In Proceedings of the Second International Conference on Combinatorial Optimization and Applications (COCOA '08), volume 5165 of Lecture Notes in Computer Science, pages 201–210. Springer, 2008.
- 19 Claude Flament. Hypergraphes arborés. Discrete Mathematics, 21(3):223–227, 1978.
- 20 Till Fluschnik and Leon Kellerhals. Placing green bridges optimally, with a multivariate analysis. In *Proceedings of the 17th Conference on Computability in Europe (CiE '21)*, volume 12813 of *Lecture Notes in Computer Science*, pages 204–216. Springer, 2021.
- 21 Till Fluschnik and Leon Kellerhals. Placing green bridges optimally, with a multivariate analysis. CoRR, abs/2102.04539, 2021. URL: https://arxiv.org/abs/2102.04539, arXiv: 2102.04539.
- 22 M. R. Garey and David S. Johnson. Computers and Intractability: A Guide to the Theory of NP-Completeness. W. H. Freeman, 1979.
- 23 Aristides Gionis, Polina Rozenshtein, Nikolaj Tatti, and Evimaria Terzi. Community-aware network sparsification. In *Proceedings of the 2017 SIAM International Conference on Data Mining (SDM '17)*, pages 426–434. SIAM, 2017.
- 24 Nili Guttmann-Beck, Roni Rozen, and Michal Stern. Vertices removal for feasibility of clustered spanning trees. *Discrete Applied Mathematics*, 296:68–84, 2021.
- 25 Nili Guttmann-Beck, Zeev Sorek, and Michal Stern. Clustered spanning tree conditions for feasibility. Discrete Mathematics & Theoretical Computer Science, 21(1), 2019.
- 26 Maike Herkenrath, Till Fluschnik, Francesco Grothe, and Leon Kellerhals. Placing green bridges optimally, with habitats inducing cycles. In Proceedings of the Thirty-First International Joint Conference on Artificial Intelligence, (IJCAI '22), pages 3825–3831. ijcai.org, 2022.
- 27 Emanuel Herrendorf. On the complexity of community-aware network sparsification. Masterarbeit, Philipps-Universität Marburg, 2022. URL: https://www.fmi.uni-jena.de/fmi\_femedia/fakultaet/institute-und-abteilungen/informatik/algorithm-engineering/master-emanuel-pdf.pdf.
- 28 Jun Hosoda, Juraj Hromkovic, Taisuke Izumi, Hirotaka Ono, Monika Steinová, and Koichi Wada. On the approximability and hardness of minimum topic connected overlay and its special instances. *Theoretical Computer Science*, 429:144–154, 2012.
- 29 Russell Impagliazzo and Ramamohan Paturi. On the complexity of k-SAT. Journal of Computer and System Sciences, 62(2):367–375, 2001.
- **30** Russell Impagliazzo, Ramamohan Paturi, and Francis Zane. Which problems have strongly exponential complexity? *Journal of Computer and System Sciences*, 63(4):512–530, 2001.
- 31 David S. Johnson and Henry O. Pollak. Hypergraph planarity and the complexity of drawing venn diagrams. *Journal of Graph Theory*, 11(3):309–325, 1987.
- 32 Boris Klemz, Tamara Mchedlidze, and Martin Nöllenburg. Minimum tree supports for hypergraphs and low-concurrency Euler diagrams. In Proceedings of the 14th Scandinavian Symposium and Workshops on Algorithm Theory (SWAT '14), volume 8503 of Lecture Notes in Computer Science, pages 265–276. Springer, 2014.
- 33 Ephraim Korach and Michal Stern. The clustering matroid and the optimal clustering tree. Mathematical Programming, 98(1-3):385–414, 2003.
- 34 Ephraim Korach and Michal Stern. The complete optimal stars-clustering-tree problem. Discrete Applied Mathematics, 156(4):444–450, 2008.
- **35** Joseph B. Kruskal. On the shortest spanning subtree of a graph and the traveling salesman problem. *Proceedings of the American Mathematical society*, 7(1):48–50, 1956.

- 36 Gerd Lindner, Christian L. Staudt, Michael Hamann, Henning Meyerhenke, and Dorothea Wagner. Structure-preserving sparsification of social networks. In Proceedings of the 2015 IEEE/ACM International Conference on Advances in Social Networks Analysis and Mining (ASONAM '15), pages 448–454. ACM, 2015.
- 37 Natsu Nakajima, Morihiro Hayashida, Jesper Jansson, Osamu Maruyama, and Tatsuya Akutsu. Determining the minimum number of protein-protein interactions required to support known protein complexes. *PloS one*, 13(4):e0195545, 2018.
- **38** Peter J Slater. A characterization of soft hypergraphs. *Canadian Mathematical Bulletin*, 21(3):335–337, 1978.
- 39 Robert Endre Tarjan and Mihalis Yannakakis. Simple linear-time algorithms to test chordality of graphs, test acyclicity of hypergraphs, and selectively reduce acyclic hypergraphs. SIAM Journal on Computing, 13(3):566–579, 1984.
- 40 Christian Wallisch. Placing green bridges to reconnect habitats densely: Algorithms and complexity. Bachelor's thesis, TU Berlin, 2023. URL: https://fpt.akt.tu-berlin.de/ publications/theses/BA-christian-wallisch.pdf.
- 41 Fang Zhou, Sébastien Mahler, and Hannu Toivonen. Network simplification with minimal loss of connectivity. In *Proceedings of the 10th IEEE International Conference on Data Mining* (*ICDM '10*), pages 659–668. IEEE Computer Society, 2010.