Branch-depth is minor closure of contraction-deletion-depth^{\dagger}

Marcin Briański[‡]

Daniel Kráľ[§]

Kristýna Pekárková[§]

Abstract

The notion of branch-depth for matroids was introduced by DeVos, Kwon and Oum as the matroid analogue of the tree-depth of graphs. The contraction-deletion-depth, another tree-depth like parameter of matroids, is the number of recursive steps needed to decompose a matroid by contractions and deletions to single elements. Any matroid with contraction-deletion-depth at most d has branch-depth at most d. However, the two notions are not functionally equivalent as contraction-deletion-depth of matroids with branch-depth two can be arbitrarily large.

We show that the two notions are functionally equivalent for representable matroids when minor closures are considered. Namely, an \mathbb{F} -representable matroid has small branch-depth if and only if it is a minor of an \mathbb{F} -representable matroid with small contraction-deletion-depth. This implies that any class of \mathbb{F} -representable matroids has bounded branch-depth if and only if it is a subclass of the minor closure of a class of \mathbb{F} -representable matroids with bounded contraction-deletion-depth.

1 Introduction

Graph width parameters play an important role both in structural and algorithmic graph theory. Indeed, graph tree-width, likely the best-known graph width parameter, is one of the core notions in the Graph Minor Project of Robertson and Seymour. The celebrated Courcelle's Theorem [6] asserts that every monadic second order property can be tested in linear times for graphs with bounded tree-width; we refer the reader e.g. to the monograph [7] for further algorithmic applications of tree-width and other graph width parameters. In this paper we study matroid branch-depth, which was introduced by De-Vos, Kwon and Oum [8] as the matroid analogue of tree-depth, a crucial structural graph measure with deep connections to sparsity theory of Nešetril and Ossona de Mendez [22],

 $^{^\}dagger {\rm The}$ second and third authors have been supported by the project 24-11098S of the Czech Science Foundation.

[‡]Theoretical Computer Science Department, Faculty of Mathematics and Computer Science, Jagiellonian University, Kraków, Poland. E-mail: marcin.brianski@doctoral.uj.edu.pl.

[§]Faculty of Informatics, Masaryk University, Botanická 68A, 602 00 Brno, Czech Republic. E-mails: dkral@fi.muni.cz and kristyna.pekarkova@mail.muni.cz.



Figure 1: The relation of classes of representable matroids with bounded depth and width parameters. The statement of Theorem 1 is visualized by the arrow relating classes with bounded contraction-deletion-depth and classes with bounded branch-depth.

graph product structure theorems [10], clustered coloring [23], and many other areas of graph theory. We remark that every graph with small rank-depth is a vertex-minor of a graph with small tree-depth [18], and graph classes with bounded rank-depth coincide with those with bounded shrub-depth [8], which is a graph depth parameter inspired by the classical notion of graph clique-width. Indeed, these two results are counterparts of more classical results concerning clique-width: every graph with small clique-width is a pivot-minor of a graph with small tree-width [21], and graph classes with bounded cliquewidth coincide with those with bounded rank-width [25] (also see [24] for the relation of rank-width, pivot-minors and vertex-minors).

Branch-width is generally considered to be the matroid analogue of graph tree-width, and indeed Hliněný [13–15] proved the matroid analogue of Courcelle's Theorem by showing that every monadic second order property can be tested in polynomial time for matroids represented over a finite field that have bounded branch-width; additional algorithmic results involving branch-width that parallel graph algorithms involving tree-width can be found e.g. [11, 16, 17, 19]. There are two matroid notions that can be considered to be analogues of graph tree-depth. The first is the notion introduced in [20], which is now referred to as contraction^{*}-depth; this parameter turned out to have an unexpected connection to optimal tree-depth of constraint matrices in integer programs [1, 2, 4, 5]. The second is the notion of branch-depth introduced by DeVos, Kwon and Oum in [8] and further studied e.g. in [12]; this notion is defined in a way completely analogous to graph rank-depth, i.e., a matroid has small branch-depth iff it can be represented by a shallow tree with all nodes corresponding to multicuts with small order (a formal definition is given in Section 2).

We now briefly discuss matroid depth and matroid width parameters and their mutual relation, which is visualized in Figure 1, to put our results in the appropriate context. In addition to branch-depth and branch-width, it is possible to define the notion of *contraction-depth*, which is the minimum depth of a procedure recursively splitting a given matroid by contractions to single elements, the notion of *deletion-depth*, which is the minimum depth of a procedure recursively splitting a given matroid by deletions to single elements, and the notion of *contraction-deletion-depth*, which is the minimum depth of a procedure recursively splitting a given matroid by contractions and deletions to single elements; the formal definitions are given in Section 2. These notions were defined by Ding, Oporowski and Oxley [9] in 1995 although the names of contraction-depth, deletion-depth and contraction-depth were coined in [8]; concepts closely related to contractiondepth and deletion-depth were also discussed by Robertson and Seymour in [27]. We remark that, unlike branch-depth and branch-width, none of contraction-depth, deletiondepth, and contraction-deletion-depth is minor-monotone.

Classes of matroids with bounded contraction-depth coincide with classes of matroids with bounded contraction^{*}-depth. In fact, the contraction^{*}-depth of a matroid M is equal to the minimum contraction-depth of a matroid containing M increased by one [3] (except for trivial cases of matroids consisting of loops and bridges). Classes of matroids with bounded contraction-depth do not need to have bounded deletion-depth and vice versa, however, every class of matroids with bounded contraction-depth has bounded contraction-depth, and every class of matroids with bounded deletion-depth has bounded contraction-depth. Furthermore, every class of matroids with bounded contraction-deletion-depth has bounded branch-depth, and every every class of matroids with bounded branch-depth has bounded branch-depth. These relations are depicted in Figure 1 and they all are strict inclusions.

Our main result is the following structural insight in classes of representable matroids with bounded branch-depth, which is depicted by the arrow in Figure 1.

Theorem 1. Let \mathcal{M} be a class of \mathbb{F} -representable matroids. The class has bounded branchdepth if and only if there exists an \mathbb{F} -representable class \mathcal{N} of matroids with bounded contraction-deletion-depth such that \mathcal{M} is a subclass of the minor closure of \mathcal{N} .

In other words, the notion of branch-depth can be thought of as the minor closure of the more restrictive parameter of contraction-deletion-depth. In a certain sense, our result is analogous to the aforementioned result stating that graph classes with bounded rank-depth coincide with those with bounded shrub-depth and are vertex-minor closures of classes with bounded tree-depth. Indeed, rank-depth captures the complexity of cuts in a manner similar to branch-depth, and shrub-depth and tree-depth capture the complexity of decomposing a graph while contraction-deletion-depth attempts to measure similar complexity in the matroid world.

The characterization of classes of representable matroids with bounded branch-depth given in Theorem 1 is implied by the following theorem:

Theorem 2. Every \mathbb{F} -representable matroid M is a minor of an \mathbb{F} -representable matroid N such that

$$cdd(N) \le 2 bd(M) \cdot (4^{bd(M)} - 1) + 1.$$



Figure 2: A fat cycle and a graph that contains the depicted fat cycle as a minor. The branch-depth of the graphic matroid associated with either of the graphs is two, the contraction-deletion-depth of the graphic matroid associated with the former graph is six, and the contraction-deletion-depth of the graphic matroid associated with the latter graph is three.

Since branch-depth is minor-monotone and upper bounded by the contraction-deletiondepth, Theorem 2 asserts that a representable matroid has small branch-depth if and only if it is a minor of a (representable) matroid with small contraction-deletion-depth. We believe that the condition on representability in Theorems 1 and 2 is an artefact of our proof methods, as further discussed in Section 5, and we conjecture that both these results also hold for general matroids.

We now try to gather some intuition why matroids with small branch-depth are minors of matroids with small contraction-deletion-depth. A prototypical example of a matroid with small branch-depth and large contraction-deletion-depth is the graphic matroid of a fat cycle — the graphic matroid associated with the graph obtained from a cycle of length n by replacing each edge by n parallel edges (see Figure 2 for illustration). The contraction-deletion depth of this graphic matroid is n+1 while its branch-depth is only two. However, this graphic matroid is a minor of the graphic matroid associated with the graph obtained from a cycle of length n+1 by replacing all but one edge with n parallel edges, which has contraction-deletion-depth equal to three. Similarly, the proof of Theorem 2 finds an embedding of a given matroid in a larger matroid that fractures into connected components after a small number of contractions and deletions that corresponds to distinct subtrees in a branch-decomposition of the input matroid. On a path towards finding such an embedding, after reviewing the notation used in the paper in Section 2. we prove our key structural lemma (Lemma 4) in Section 3. This lemma asserts that if a collection of subspaces of a vector space has small connectivity, then the subspaces share a low-dimensional subspace and form an almost direct sum. In Section 4, we use the lemma to embed (as minors) matroids with small branch-depth in matroids with small contraction-deletion depth and derive proofs of Theorems 1 and 2. We conclude by stating open problems directly stemming from our results in Section 5.

2 Notation

In this section, we introduce notation used throughout the paper. The set of the first k positive integers is denoted by [k]. We often work with vector spaces and we briefly review notation that we use. If X is a set of vectors, then \overline{X} is the linear hull of the vectors contained in X. If X and Y are subspaces of a vector space, then X + Y is the smallest subspace containing X and Y, and more generally, if X_1, \ldots, X_k are subspaces, then $\sum_{i=1}^k X_i$ is the smallest subspace containing X_1, \ldots, X_k . Finally, if X and Y are subspaces, the quotient space X/Y is the vector space formed by classes of the equivalence relation \sim such that $x \sim x'$ iff $x - x' \in Y$. The quotient space X/Y can be identified with a subspace of X of dimension dim $X - \dim X \cap Y$ (note that we do not require Y to be a subspace of X), which is formed by including a single suitably chosen vector from each class of the equivalence relation \sim .

We assume that the reader is familiar with standard matroid terminology, which can be found e.g. in [26], and we just review notation that may occasionally differ. If M is a matroid, then |M| denotes the number of elements of M. The rank of a set X of elements of M is denoted by rank_M X; we drop the subscript M if the matroid M is clear from context. We slightly abuse the notation and write rank M for the rank of the matroid M. If M is a matroid and e is an element of M, then M/e is the matroid obtained by contracting the element e and $M \setminus e$ is the matroid obtained by deleting the element e. Throughout the paper, all contractions involve non-loop elements of a matroid only. A *minor* of a matroid M is any matroid that can be obtained from M contracting and deleting elements, and the *restriction* of a matroid M to a set X of its elements is the matroid with the ground set X and a subset of X being independent in the restriction if and only if it is independent in M. If \mathcal{M} is a class of matroids, then the *minor closure* of \mathcal{M} is the class containing all matroids that are minors of an element of \mathcal{M} . Finally, a *component* of a matroid is an inclusion-wise maximal subset of elements of M such that any two of them are contained in a common circuit.

A matroid M is \mathbb{F} -representable if it is a vector matroid over a field \mathbb{F} , i.e., its elements can be associated with vectors over \mathbb{F} and a subset of elements of M is independent if and only if the corresponding vectors are linearly independent; such an association is referred to as an \mathbb{F} -representation. If an \mathbb{F} -representation of M is fixed, we say that the matroid M is an \mathbb{F} -represented matroid and identify the elements of M with the vectors over \mathbb{F} . We will always assume that the dimension of the vector space is equal to rank M, that is, the vectors are elements of $\mathbb{F}^{\operatorname{rank} M}$. An \mathbb{F} -represented matroid M can be viewed as a matrix whose columns are the vectors of M. Note that the matroid M is preserved by row operations, i.e., after performing any row operations the obtained matrix is still an \mathbb{F} -representation of the same matroid M. The \mathbb{F} -representation of the matroid M obtained by deleting an element e can be obtained from the \mathbb{F} -representation of M by deleting the column corresponding to e, and the \mathbb{F} -representation of the matroid M obtained by contracting e can be obtained from the \mathbb{F} -representation of M by first performing row operations to make the column corresponding to e to be a unit vector and then by deleting both the only row containing 1 in the column corresponding to e and the column corresponding to e.

We next define the connectivity function for a matroid M and some functions derived from it. If X and Y are disjoint subsets of the elements of a matroid M, we define a two-parameter function $\lambda_M(\cdot, \cdot)$ that

$$\lambda_M(X, Y) = \operatorname{rank}_M X + \operatorname{rank}_M Y - \operatorname{rank}_M (X \cup Y).$$

The connectivity function is the one-parameter function $\lambda_M(\cdot)$ such that $\lambda_M(X) = \lambda_M(X, Y)$ where Y is the complement of X. We also use λ_M for this one-parameter function as it is derived from the more general two-parameter function introduced earlier; we believe that no confusion can arise as the two functions have a different number of parameters. Finally, if X_1, \ldots, X_k are disjoint subsets of the elements of a matroid M, we set

$$\lambda_M^*(X_1,\ldots,X_k) = \max_{I \subseteq [k]} \lambda_M \left(\bigcup_{i \in I} X_i, \bigcup_{i \in [k] \setminus I} X_i \right)$$

In case the matroid M is clear from the context we will drop the subscript, so we will occasionally write $\lambda(\cdot)$, $\lambda(\cdot, \cdot)$ and $\lambda^*(\cdot, \ldots, \cdot)$ instead of $\lambda_M(\cdot)$, $\lambda_M(\cdot, \cdot)$ and $\lambda^*_M(\cdot, \ldots, \cdot)$. We extend the definition of $\lambda^*(\cdot, \ldots, \cdot)$ to vector spaces and set

$$\lambda^*(X_1,\ldots,X_k) = \max_{I \subseteq [k]} \left(\dim \sum_{i \in I} X_i + \dim \sum_{i \in [k] \setminus I} X_i - \dim(X_1 + \ldots + X_k) \right)$$

for vector spaces X_1, \ldots, X_k .

We now define depth parameters used throughout this paper. The *contraction-deletion-depth* a matroid M, which is denoted by cdd(M), is defined recursively as follows:

- If M has a single element, then cdd(M) = 1.
- If M is not connected, then cdd(M) is the maximum contraction-deletion-depth of a component of M.
- Otherwise, $\operatorname{cdd}(M) = 1 + \min_{e \in M} \min\{\operatorname{cdd}(M/e), \operatorname{cdd}(M \setminus e)\}$, i.e., $\operatorname{cdd}(M)$ is one plus the minimum contraction-deletion-depth of a matroid that can be obtained from M by deleting or contracting a single element.

The contraction-depth of a matroid M, denoted by cd(M), is defined analogously but with only contractions of elements permitted in the last case, and the deletion-depth of a matroid M, denoted by dd(M), is defined analogously but with only deletions of elements permitted in the last case.

A (d, r)-decomposition of a matroid M is a tree T such that the radius of T is d, the leaves of T are in a one-to-one correspondence to the elements of M, and the following holds for every inner vertex v of T: $\lambda_M^*(X_1, \ldots, X_k) \leq r$ where k is the number of neighbors of v and X_1, \ldots, X_k are the sets of elements of M assigned to the leaves of the k trees of the forest $T \setminus v$. The *branch-depth* of a matroid M, which is denoted by bd(M), is the minimal k such that M has a (d, r)-decomposition with $d \leq k$ and $r \leq k$. It can be shown [8, Proposition 5.3] that if a matroid M is a minor of a matroid N, then $bd(M) \leq bd(N)$. In other words, branch-depth is a minor-monotone parameter.

For the next definition, recall that the *depth* of a rooted tree T is the maximum number of edges on a path from the root to a leaf of T. A rooted (d, r)-decomposition of a matroid M is a rooted tree T such that the depth of T is d, the leaves of T are in a one-to-one correspondence with the elements of M, and the following holds for every inner vertex vof T: $\lambda_M^*(X_1, \ldots, X_k) \leq r$ where k is the number of children of v and X_1, \ldots, X_k are the sets of elements of M assigned to the leaves of the k subtrees rooted at the children of v.

We conclude this section with the following proposition which relates rooted decompositions and decompositions of matroids.

Proposition 3. If a matroid M has a (d, r)-decomposition, then the matroid M has a rooted (d, r)-decomposition.

Proof. Fix a (d, r)-decomposition T of M, and let w be an inner vertex of T such that any other vertex of T is at distance at most d from w. Since the radius of T is equal to d, there is some vertex of T at distance exactly d from w, and this vertex must be a leaf. Let T' be the rooted tree obtained from T by rooting it at w. Note that the depth of T' is d. Consider now an inner vertex v of T' and let k be the number of children of v. Further, let X_1, \ldots, X_k be the sets of elements of M assigned to the leaves of the k subtrees rooted at the children of v, and let Y be the set of elements of M not assigned to a leaf of the subtree rooted at v, namely, the elements mapped to a leaf not in any of the k subtrees. Since T is a (d, r)-decomposition and it holds that

$$\lambda_M \left(\bigcup_{i \in I} X_i, \bigcup_{i \in [k] \setminus I} X_i \right) \le \lambda_M \left(\bigcup_{i \in I} X_i, Y \cup \bigcup_{i \in [k] \setminus I} X_i \right)$$

for any subset $I \subseteq [k]$, where the latter is at most r, we obtain that $\lambda_M^*(X_1, \ldots, X_k) \leq r$. It follows that T' is a rooted (d, r)-decomposition of the matroid M.

3 Main lemma

The core of our argument is Lemma 4 which we prove in this section. Informally speaking, the lemma asserts that there are only two reasons why a partition E_1, \ldots, E_k of elements of a represented matroid M satisfies that $\lambda_M^*(E_1, \ldots, E_k) \leq r$:

- the subspaces X_1, \ldots, X_k spanned by E_1, \ldots, E_k intersect in a small dimensional subspace, or
- the subspaces X_1, \ldots, X_k form an "almost direct" sum of vector spaces, i.e., they are subspaces in a general position in a space whose dimension is slightly smaller than the sum of the dimensions of the subspaces X_1, \ldots, X_k .

It is not hard to see that either of these two properties implies that $\lambda_M^*(E_1, \ldots, E_k)$ is small. Indeed, if X_1, \ldots, X_k are subspaces of a vector space and there exists a vector space A of dimension a such that

$$\sum_{i=1}^{k} \dim(X_i/A) \le \dim\left(\left(\sum_{i=1}^{k} X_i\right)/A\right) + b,\tag{1}$$

then $\lambda^*(X_1, \ldots, X_k) \leq a + b$. Note that the parameter *a* represents the dimension of the subspace "shared" by X_1, \ldots, X_k , which is the subspace *A*, and the parameter *b* represents how close the subspaces are to the direct sum (after contracting the shared subspace).

We next state the main lemma of this section. Note that, assuming that $A \subseteq X_1 + \cdots + X_k$, it holds that

$$\dim\left(\sum_{i=1}^{k} X_i\right) = \dim\left(\left(\sum_{i=1}^{k} X_i\right)/A\right) + \dim A.$$

Hence, the lemma says that (1) holds with $a = b = \dim A \leq 3r$ for the subspace A from the statement of the lemma.

Lemma 4. Let X_1, \ldots, X_k be vector spaces over a field \mathbb{F} such that $\lambda^*(X_1, \ldots, X_k) \leq r$. There exists a vector space A with dimension at most 3r such that

$$\sum_{i=1}^{k} \dim(X_i/A) \le \dim\left(\sum_{i=1}^{k} X_i\right).$$

Proof. We will construct vector spaces A_m , L_m and R_m for every $m \in [k]$ with the following properties:

- (i) A_m is a subspace of $X_1 + \cdots + X_m$,
- (ii) there exists $I \subseteq [m]$ such that

$$L_m = \sum_{i \in I} X_i$$
 and $R_m = \sum_{i \in [m] \setminus I} X_i$,

- (iii) if m > 1, then $A_{m-1} \subseteq A_m$, $L_{m-1} \subseteq L_m$, and $R_{m-1} \subseteq R_m$,
- (iv) $L_m \cap R_m$ is a subspace of A_m ,
- (v) dim $A_m \leq 3 \cdot \dim L_m \cap R_m$, and
- (vi) it holds that

$$\sum_{i=1}^{m} \dim(X_i/A_m) \le \dim\left(\sum_{i=1}^{m} X_i\right).$$

Once we have constructed A_m , L_m and R_m for every $m \in [k]$, we set the vector space Afrom the statement of the lemma to be the vector space A_k . Note that since the dimension of $L_k \cap R_k$ cannot be larger than $\lambda^*(X_1, \ldots, X_k) \leq r$ by the property (ii), the dimension of $A = A_k$ is at most 3r by the property (v), and the inequality from the statement of the lemma follows from the property (vi) above for m = k. Therefore, it suffices to construct the vector spaces A_m , L_m , and R_m , $m \in [k]$, in order to prove the lemma.

We now iteratively construct the vector spaces A_m , L_m , and R_m , starting with m = 1. We set A_1 and R_1 to be the 0-dimensional vector space and $L_1 = X_1$. It is straightforward to verify that A_1 , L_1 and R_1 satisfy all properties (i)–(vi).

Suppose that we have constructed vector spaces A_{m-1} , L_{m-1} , and R_{m-1} . Consider the vector space A'_m defined as follows:

$$A'_{m} = (X_{m} + A_{m-1}) \cap \sum_{i \in [m-1]} X_{i}.$$
 (2)

Note that A'_m contains A_{m-1} as a subspace, since A_{m-1} is a subspace of $\sum_{i=1}^{m-1} X_i$. We next distinguish two cases: the first is that the spaces A_{m-1} and A'_m are equal, and the second is that A_{m-1} is a proper subspace of A'_m .

If the spaces A_{m-1} and A'_m are equal, we set the subspaces A_m , L_m and R_m as follows:

$$L_m = L_{m-1} + X_m$$
, $R_m = R_{m-1}$ and $A_m = A'_m + (L_m \cap R_m) = A_{m-1} + (L_m \cap R_m)$.

The subspaces A_m , L_m and R_m clearly satisfy the properties (i)–(iv). The property (v), which bounds the dimension of A_m , follows from the next estimates:

$$\dim A_m = \dim(A_{m-1} + (L_m \cap R_m)) = \dim A_{m-1} + \dim(L_m \cap R_m) - \dim(A_{m-1} \cap (L_m \cap R_m)) \leq \dim A_{m-1} + \dim(L_m \cap R_m) - \dim(A_{m-1} \cap (L_{m-1} \cap R_{m-1})) = \dim A_{m-1} + \dim(L_m \cap R_m) - \dim(L_{m-1} \cap R_{m-1}) \leq 3\dim(L_{m-1} \cap R_{m-1}) + \dim(L_m \cap R_m) - \dim(L_{m-1} \cap R_{m-1}) = \dim(L_m \cap R_m) + 2\dim(L_{m-1} \cap R_{m-1}) \leq 3\dim(L_m \cap R_m).$$

We prove the property (vi) jointly for the first and the second cases later.

If the space A_{m-1} is a proper subspace of A'_m , we proceed as follows. Let $d = \dim A'_m - \dim A_{m-1}$; we remark that the previous case $A'_m = A_{m-1}$ can be viewed as the case d = 0 of the arguments that we next present, however, we decided to separate the two cases in the interest of the clarity of presentation. The definition of A'_m as (2) implies that there exist (linearly independent) vectors $x_1, \ldots, x_d \in X_m$ such that any basis of A_{m-1} together with x_1, \ldots, x_d is a basis of A'_m . Note that none of the vectors x_1, \ldots, x_d belongs to A_{m-1} . Since the vector x_i belongs to A'_m , and so $x_i \in X_1 + \cdots + X_{m-1} = L_{m-1} + R_{m-1}$, there exist $\ell_i \in L_{m-1}$ and $r_i \in R_{m-1}$ such that $x_i = \ell_i + r_i$. Let L' be the linear hull of the vectors

 ℓ_1, \ldots, ℓ_d and R' the linear hull of the vectors r_1, \ldots, r_d . Note that the vectors x_1, \ldots, x_d are contained in L' + R', which implies that

$$\dim(L' + R') - \dim((L' + R') \cap A_{m-1}) \ge d.$$
(3)

On the other hand, the space $L' \cap R'$ is a subspace of $L_{m-1} \cap R_{m-1}$ and so of A_{m-1} , which yields that

$$\dim(L' \cap R') - \dim((L' \cap R') \cap A_{m-1}) = 0.$$
(4)

Next observe that

$$\dim L' + \dim R' = \dim(L' + R') + \dim(L' \cap R'), \text{ and}$$

$$\tag{5}$$

$$\dim(L' \cap A_{m-1}) + \dim(R' \cap A_{m-1}) = \dim((L' + R') \cap A_{m-1}) + \dim((L' \cap R') \cap A_{m-1}).$$
(6)

We combine (3), (4), (5) and (6) to obtain that

$$(\dim L' - \dim(L' \cap A_{m-1})) + (\dim R' - \dim(R' \cap A_{m-1})) \ge d.$$

It follows that dim $L' - \dim(L' \cap A_{m-1}) \ge d/2$ or dim $R' - \dim(R' \cap A_{m-1}) \ge d/2$ (or possibly both). The two possibilities are symmetric, so we assume that dim $L' - \dim(L' \cap A_{m-1}) \ge d/2$.

We define the subspaces L_m , R_m , and A_m as

$$L_m = L_{m-1}, \quad R_m = R_{m-1} + X_m \text{ and } A_m = A'_m + (L_m \cap R_m).$$

Note that every $\ell_i = x_i - r_i$ belongs to $L_m \cap R_m$: indeed, $\ell_i \in L_{m-1} \subseteq L_m$ and $x_i - r_i \in X_m + R_{m-1} = R_m$. Hence, the space L', which is the linear hull of the vectors ℓ_1, \ldots, ℓ_d , is a subspace of $L_m \cap R_m$. Since $L_{m-1} \cap R_{m-1}$ is a subspace of A_{m-1} , it follows that

$$\dim(L_m \cap R_m) - \dim(L_{m-1} \cap R_{m-1}) \ge \dim(L' + (L_{m-1} \cap R_{m-1})) - \dim(L_{m-1} \cap R_{m-1})$$
$$\ge \dim(L' + A_{m-1}) - \dim A_{m-1}$$
$$= \dim L' - \dim(L' \cap A_{m-1})$$
$$\ge d/2.$$

Hence, it holds that $d/2 \leq \dim(L_m \cap R_m) - \dim(L_{m-1} \cap R_{m-1})$, which is equivalent to

$$d \le 2 \left(\dim(L_m \cap R_m) - \dim(L_{m-1} \cap R_{m-1}) \right).$$

$$\tag{7}$$

. We can now estimate the dimension of A_m as follows (recall that $L_{m-1} \cap R_{m-1}$ is a subspace of $A_{m-1} \subseteq A'_m$ and also of $L_m \cap R_m$):

$$\dim A_m \leq \dim A'_m + \dim(L_m \cap R_m) - \dim(L_{m-1} \cap R_{m-1}) = \dim A_{m-1} + d + \dim(L_m \cap R_m) - \dim(L_{m-1} \cap R_{m-1}) \leq \dim A_{m-1} + 3 \left(\dim(L_m \cap R_m) - \dim(L_{m-1} \cap R_{m-1})\right) \leq 3 \dim(L_{m-1} \cap R_{m-1}) + 3 \left(\dim(L_m \cap R_m) - \dim(L_{m-1} \cap R_{m-1})\right) = 3 \dim(L_m \cap R_m),$$

where the second inequality follows from (7). We have now verified that the properties (i)-(v) hold in the second case.

We now continue jointly the analysis of the first and second cases and establish the property (vi). We will use the definition of A'_m as (2) and that $A_m = A'_m + (L_m \cap R_m)$. The definition of A'_m yields that

$$\dim(X_m + A_{m-1}) + \dim\left(\sum_{i=1}^{m-1} X_i\right) = \dim\left(\sum_{i=1}^m X_i\right) + \dim A'_m;$$
(8)

indeed, (8) follows from the equality $\dim S + \dim T = \dim S \cup T + \dim S \cap T$ used for $S = X_m + A_{m-1}$ and $T = \sum_{i=1}^{m-1} X_i$. We now establish the property (vi).

$$\sum_{i=1}^{m} \dim(X_i/A_m) \leq \sum_{i=1}^{m} \dim(X_i/A'_m)$$

$$= \sum_{i=1}^{m-1} \dim(X_i/A'_m) + \dim(X_m/A_{m-1}) - (\dim A'_m - \dim A_{m-1})$$

$$= \sum_{i=1}^{m-1} \dim(X_i/A'_m) + \dim(X_m + A_{m-1}) - \dim A_{m-1} - \dim A'_m + \dim A_{m-1}$$

$$= \sum_{i=1}^{m-1} \dim(X_i/A'_m) + \dim(X_m + A_{m-1}) - \dim A'_m$$

$$\leq \sum_{i=1}^{m-1} \dim(X_i/A_{m-1}) + \dim(X_m + A_{m-1}) - \dim A'_m$$

$$\leq \sum_{i=1}^{m-1} \dim X_i + \dim(X_m + A_{m-1}) - \dim A'_m$$

$$= \sum_{i=1}^{m} \dim X_i,$$

where the last equality is a restatement of (8).

4 Main result

In this section, we prove the main result of our paper which is the next theorem. The results announced in Section 1 are its immediate corollaries which we derive just after the proof of the theorem.

Theorem 5. If an \mathbb{F} -representable matroid M has a rooted (d, r)-decomposition, then there exists an \mathbb{F} -representable matroid N such that M is a minor of N and $cdd(N) \leq 2r \cdot (4^d - 1) + 1$. Proof. The proof proceeds by induction on d. We will show the statement for \mathbb{F} -represented matroids and we additionally show that the considered \mathbb{F} -representation of M can be obtained from an \mathbb{F} -representation of N (by deleting rows corresponding to contracted elements and deleting columns corresponding to deleted elements of N). The base of the induction is the case when d = 0. If d = 0, then the matroid M has a single element and we set N = M (note that cdd(N) = 1 as required). We next present the induction step. Fix an \mathbb{F} -represented matroid M and a rooted (d, r)-decomposition T of M. Let k be the number of children of the root of T and let T_1, \ldots, T_k be the (rooted) subtrees of T rooted at the children of T. Further, let E_i , be the elements of M assigned to the leaves of T_i , $i \in [k]$, and let X_i be the linear hull of E_i .

Since T is a (d, r)-decomposition of M, it holds that $\lambda^*(X_1, \ldots, X_k) \leq r$. Hence, Lemma 4 yields that there exists a subspace A of $\mathbb{F}^{\operatorname{rank} M}$ with dimension at most 3r such that

$$\sum_{i=1}^{k} \dim(X_i/A) \le \dim \sum_{i=1}^{k} X_i.$$
(9)

Let d_A be the dimension of the vector space A and let d_i be the dimension of the vector space X_i/A , $i \in [k]$. Note that rank M is equal to $\dim(X_1 + \cdots + X_k)$, which is at most $d_1 + \cdots + d_k + d_A$. On the other hand, the rank of M is at least $d_1 + \cdots + d_k$ by (9).

Fix a basis $B_A = \{a_1, \ldots, a_{d_A}\}$ of the vector space A, and for each $i \in [k]$ let $B_i = \{x_1^i, \ldots, x_{d_i}^i\}$ be vectors of X_i , such that $a_1, \ldots, a_{d_A}, x_1^i, \ldots, x_{d_i}^i$ is a basis of the vector space $X_i + A$. Observe that the vector space $\mathbb{F}^{\operatorname{rank} M} = X_1 + \cdots + X_k$ is spanned by the set of vectors $B_A \cup B_1 \cup \cdots \cup B_k$. It follows that there is a subset of these vectors is a basis of the vector space $X_1 + \ldots + X_k$; we may assume that this subset contains all vectors a_1, \ldots, a_{d_A} (as these vectors are linearly independent). Hence, after possibly permuting the vectors $x_1^i, \ldots, x_{d_i}^i$, we assume that for each $i \in [k]$ there exists $d'_i \in \{0\} \cup [d_i]$, such that the set of vectors $B = B_A \cup \{x_j^i : i \in [k], j \in [d'_i]\}$ form a basis of the vector space $X_1 + \ldots + X_k$. In addition, we assume that the vectors a_1, \ldots, a_{d_A} are the first d_A unit vectors, $x_1^1, \ldots, x_{d'_1}^1$ are the next d'_1 unit vectors, etc. Since the sum $d'_1 + \cdots + d'_k + d_A$ is equal to the rank of M, which is at least $d_1 + \cdots + d_k$, we obtain that

$$\sum_{i=1}^{k} (d_i - d'_i) \leq d_A.$$
(10)

Let d_C be the value of the sum from the left side of (10); note that $d_C \leq d_A \leq 3r$. We now view $X_1 + \cdots + X_k$ as a subspace of $\mathbb{F}^{\operatorname{rank} M + d_C}$, where the vectors in B are the first rank M unit vectors (in the order described above). Next, let $C = \{b_j^i : i \in [k], j \in [d_i] \setminus [d'_i]\}$ be the remaining (last) d_C unit vectors of $\mathbb{F}^{\operatorname{rank} M + d_C}$ (where b_j^1 comes before b_j^2 , which comes before b_j^3 , etc).

We next construct a vector matroid M' of rank equal to rank $M + d_C$ such that M is a minor of M'. The matroid M' contains the following $|M| + d_A + d_C$ vectors:



Figure 3: The F-representation of the matroid M' constructed in the proof of Theorem 5 when k = 3, $d_A = 4$, $d'_1 = 3$, $d_1 = 4$, $d'_2 = 2$, $d_2 = 4$, $d'_3 = 2$ and $d_3 = 3$. Stars depict entries that can be arbitrary (both zero or non-zero), and the entries that are not displayed are zero. The representation of the matroid M'' from the proof is the part of the representation of M' encompassed by the dashed lines.

• For every $i \in [k]$ and every element $x \in E_i$, the matroid M' contains the vector

$$\sum_{j=1}^{d_A} \alpha_j a_j + \sum_{j=1}^{d'_i} \beta_j x_j^i + \sum_{j=d'_i+1}^{d_i} \beta_j b_j^i$$

where $\alpha_j \in \mathbb{F}$, $j \in [d_A]$, and $\beta_j \in \mathbb{F}$, $j \in [d_i]$ are the unique coefficients such that

$$x = \sum_{j=1}^{d_A} \alpha_j a_j + \sum_{j=1}^{d_i} \beta_j x_j^i.$$

In particular, the matroid M' contains the unit vectors x_j^i , $i \in [k]$ and $j \in [d'_i]$ and the unit vectors b_j^i , $i \in [k]$ and $j \in [d_i] \setminus [d'_i]$; the unit vectors b_j^i , $i \in [k]$ and $j \in [d_i] \setminus [d'_i]$ are included because of the elements x_j^i with $j \in [d_i] \setminus [d'_i]$.

- The d_A vectors a_1, \ldots, a_{d_A} .
- For every $i \in [k]$ and $j \in [d_i] \setminus [d'_i]$, the matroid M' contains the vector $z_j^i = b_j^i x_j^i$; note that there are exactly d_B vectors added to M' because of this rule.

The construction of the matroid M' is illustrated in Figure 3.

Observe that the matroid M is a minor of M', specifically, the matroid M can be obtained from M' by deleting the d_A elements a_1, \ldots, a_{d_A} and contracting the d_B elements

 $z_j^i, i \in [k]$ and $j \in [d_i] \setminus [d'_i]$. Let M'' be the matroid obtained from M' by contracting the d_A elements a_1, \ldots, a_{d_A} and deleting the d_B elements $z_j^i, i \in [k]$ and $j \in [d_i] \setminus [d'_i]$. Let M''_i be the restriction of M'' to E_i , and observe M''_i is a union of components of M''.

We aim to apply induction to each of the matroids M''_i , $i \in [k]$. To do so, we first show that T_i is a rooted (d-1, 4r)-decomposition of the matroid M''_i . Observe that the matroid M''_i can be obtained from the matroid M_i by first adding the d_A elements a_1, \ldots, a_{d_A} and then contracting the d_A elements a_1, \ldots, a_{d_A} . It follows that

$$\operatorname{rank}_M X - d_A \le \operatorname{rank}_M X - \dim A \cap \overline{X} = \operatorname{rank}_{M_i''} X \le \operatorname{rank}_M X$$

for every $X \subseteq E_i$. Hence, the following holds for any two disjoint subsets X and Y of E_i :

$$\lambda_{M_i''}(X,Y) = \operatorname{rank}_{M_i''} X + \operatorname{rank}_{M_i''} Y - \operatorname{rank}_{M_i''}(X \cup Y)$$

$$\leq \operatorname{rank}_{M_i''} X + \operatorname{rank}_{M_i''} Y - (\operatorname{rank}_M(X \cup Y) - d_A)$$

$$\leq \operatorname{rank}_M X + \operatorname{rank}_M Y - \operatorname{rank}_M (X \cup Y) + d_A$$

$$= \lambda_M(X,Y) + \dim A \leq \lambda_M(X,Y) + 3r.$$

We conclude that T_i is a rooted (d-1, 4r)-decomposition of the matroid M''_i .

By induction, the $\mathbb F\text{-represented}$ matroid M_i'' is a minor of an $\mathbb F\text{-represented}$ matroid N_i such that

 $\operatorname{cdd}(N_i) \le 8r \cdot (4^{d-1} - 1) + 1 = 2r \cdot (4^d - 1) + 1 - 6r.$

Set $n_i = \operatorname{rank} N_i - \operatorname{rank} M''_i$, $i \in [k]$, and let y_j^i , $i \in [k]$ and $j \in [n_i]$, be the elements of N_i that were contracted. Note that for every $i \in [k]$ the elements x_j^i , $j \in [d_i]$, and y_j^i , $j \in [n_i]$, form a basis of N_i .

We will now construct the sought matroid N; the construction of the matroid N is illustrated in Figure 4. The matroid N is a vector matroid of rank $R = d_1 + \cdots + d_k +$ $n_1 + \cdots + n_k + d_A$ with $|N_1| + \cdots + |N_K| + d_A + d_B$ elements, which one-to-one correspond to the elements of the matroids N_1, \ldots, N_k , and the elements a_1, \ldots, a_{d_A} and z_j^i , $i \in [k]$ and $j \in [d_i] \setminus [d'_i]$, and so we identify them with these elements. The elements a_1, \ldots, a_{d_A} , the elements x_j^i , $i \in [k]$ and $j \in [d_i]$, and the elements y_j^i , $i \in [k]$ and $j \in [n_i]$ will form a basis of N, and so we can assume that these elements are represented by unit vectors in the vector space $\mathbb{F}^{d_1 + \cdots + d_k + n_1 + \cdots + n_k + d_A}$ and the elements a_1, \ldots, a_{d_A} are the first d_A unit vectors. The vector matroid N contains the following vectors:

- each of the unit vectors a_1, \ldots, a_{d_A} ,
- the vector z_j^i , $i \in [k]$ and $j \in [d_i] \setminus [d'_i]$ with all its additional $n_1 + \cdots + n_k$ entries set to zero,
- for every $i \in [k]$ and every vector of N_i contained in M''_i , this vector with its first d_A entries being the same as in M' and all its additional $R d_A d_i n_i$ entries set to zero, and
- for every $i \in [k]$ and every vector of N_i not contained in M''_i , this vector with all its the additional $R d_i n_i$ entries set to zero.



Figure 4: The F-representation of the matroid N constructed in the proof of Theorem 5 when k = 3, $d_A = 4$, $d'_1 = 3$, $d_1 = 4$, $n_1 = 1$, $d'_2 = 2$, $d_2 = 4$, $n_2 = 1$, $d'_3 = 2$, $d_3 = 3$ and $n_3 = 2$. Stars depict entries that can be arbitrary (both zero or non-zero), and the entries that are not displayed are zero. The representations of matroids N_1 , N_2 and N_3 obtained from induction are encompassed by the dashed lines.

Note that the matroid obtained from the matroid N by contracting the d_A elements a_1, \ldots, a_{d_A} and deleting the d_B elements z_j^i , $i \in [k]$ and $j \in [d_i] \setminus [d'_i]$ is the matroid that is the union of the matroids N_1, \ldots, N_k . Since $d_A + d_B \leq 6r$ and $\operatorname{cdd}(N_i) \leq 2r \cdot (4^d - 1) + 1 - 6r$ for every $i \in [k]$, it follows that the contraction-deletion-depth of N is at most $2r \cdot (4^d - 1) + 1$. On the other hand, the matroid obtained from the matroid M by contracting the d_B elements z_j^i , $i \in [k]$ and $j \in [d_i] \setminus [d'_i]$, contracting the $n_1 + \cdots + n_k$ elements y_j^i , $i \in [k]$ and $j \in [n_i]$ and deleting all the remaining elements not contained in $E_1 \cup \cdots \cup E_k$ yields the matroid M. This completes the proof of the induction step and so of the theorem.

We immediately obtain Theorem 2 as a corollary.

Proof of Theorem 2. Let M be an \mathbb{F} -representable matroid with branch-depth k. Since M has a (d, r)-decomposition for some non-negative integers $d \leq k$ and $r \leq k$, Proposition 3 yields that M has a rooted (d, r)-decomposition. The corollary now directly follows from Theorem 5.

We also obtain the characterization of classes of representable matroids with bounded branch-depth given in Theorem 1.

Proof of Theorem 1. Fix a class \mathcal{M} of \mathbb{F} -representable matroids. Suppose that there exists a class \mathcal{N} of matroids with contraction-deletion-depth bounded by d such that \mathcal{M} is a subclass of the minor closure of the class \mathcal{N} . Since the contraction-deletion-depth is an upper bound on the branch-depth [8, Theorem 5.2], the branch-depth of every matroid contained in \mathcal{N} is also bounded by d and so is the branch-depth of every matroid contained in the minor closure of \mathcal{N} as the branch-depth is minor-monotone [8, Proposition 5.3]. Hence, if \mathcal{M} is a subclass of the minor closure of \mathcal{N} , the branch-depth of every matroid contained in \mathcal{M} is bounded by d.

We now prove the other implication. Suppose that the branch-depth of every matroid contained in \mathcal{M} is bounded by d'. For every matroid $M \in \mathcal{M}$, let N_M be an \mathbb{F} -representable matroid with contraction-deletion-depth bounded at most $2d' \cdot (4^{d'} - 1) + 1$ such that Mis a minor of N_M ; such a matroid N_M exists by Theorem 2. The class \mathcal{N} is formed by the matroids N_M for all $M \in \mathcal{M}$. Since the contraction-deletion-depth of every matroid contained in \mathcal{N} is at most $2d' \cdot (4^{d'} - 1) + 1$ and the class M is a subclass of the minor closure of \mathcal{N} , the other implication is now proven.

5 Conclusion

Since our main result, which asserts that every representable matroid with bounded branchdepth is a minor of a (representable) matroid with bounded contraction-deletion-depth, applies to representable matroids it is natural to ask whether it can be extended to all matroids. We believe that indeed the same statement, perhaps with a worse functional dependance on branch-depth, also holds for general matroids. **Conjecture 1.** There exists a function f such that if a matroid M has branch-depth d, then there exists a matroid N with contraction-deletion-depth at most f(d) such that M is a minor of N.

It is possible to state Conjecture 1 purely in terms of relating classes of matroids of bounded branch-depth and the minor closures of matroid classes with bounded contractiondeletion-depth. It is easy to see that Conjecture 1 and Conjecture 2 are equivalent.

Conjecture 2 (Conjecture 1, restated). A class \mathcal{M} of matroids has bounded branch-depth if and only if there exists a class \mathcal{N} of matroids with bounded contraction-deletion-depth such that \mathcal{M} is a subclass of the minor closure of \mathcal{N} .

We have not been able to construct representable matroids M_i that have (d, r)-decompositions and there would not exist matroids N_i with contraction-deletion-depth $\mathcal{O}(dr)$ containing M_i as a minor. So, we pose the following as an open problem.

Problem 3. Is every representable matroid M with branch-depth at most d a minor of a (representable) matroid N with contraction-deletion-depth at most $Kd^2 + L$ for some K and L independent of M?

References

- [1] M. Briański, M. Koutecký, D. Král', K. Pekárková and F. Schröder: *Characterization* of matrices with bounded graver bases and depth parameters and applications to integer programming, to appear in Mathematical Programming.
- [2] M. Briański, M. Koutecký, D. Král', K. Pekárková and F. Schröder: Characterization of Matrices with Bounded Graver Bases and Depth Parameters and Applications to Integer Programming, 49th International Colloquium on Automata, Languages, and Programming (ICALP 2022) (2022), 29:1–29:20.
- [3] M. Briański, D. Král' and A. Lamaison: Closure property of contraction-depth of matroids, preprint arXiv:2311.01945 (2023).
- [4] T. F. N. Chan, J. W. Cooper, M. Koutecký, D. Král' and K. Pekárková: Matrices of Optimal Tree-Depth and Row-Invariant Parameterized Algorithm for Integer Programming, 47th International Colloquium on Automata, Languages, and Programming (ICALP 2020) (2020), 26:1–26:19.
- [5] T. F. N. Chan, J. W. Cooper, M. Koutecký, D. Král' and K. Pekárková: Matrices of optimal tree-depth and row-invariant parameterized algorithm for integer programming, SIAM Journal on Computing 41 (2022), 664–700.
- [6] B. Courcelle: The monadic second-order logic of graphs. I. recognizable sets of finite graphs, Inform. and Comput. 85 (1990), 12–75.

- [7] M. Cygan, F. V. Fomin, L. Kowalik, D. Lokshtanov, D. Marx, M. Pilipczuk, M. Pilipczuk and S. Saurabh: Parameterized Algorithms, Springer, 2015.
- [8] M. DeVos, O. Kwon and S. Oum: *Branch-depth: Generalizing tree-depth of graphs*, European Journal of Combinatorics **90** (2020), 103186, 23pp.
- G. Ding, B. Oporowski and J. Oxley: On infinite antichains of matroids, Journal of Combinatorial Theory Series B 63 (1995), 21–40.
- [10] Z. Dvořák and D. R. Wood: Product structure of graph classes with strongly sublinear separators, arXiv preprint arXiv:2208.10074 (2022).
- [11] T. Gavenčiak, D. Král' and S. Oum: Deciding first order properties of matroids, 39th International Colloquium Automata, Languages, and Programming, ICALP (2012), 239–250.
- [12] J. P. Gollin, K. Hendrey, D. Mayhew and S. Oum: Obstructions for bounded branchdepth in matroids, Advances in Combinatorics (2021), 2021:4, 25pp.
- [13] P. Hliněný: Branch-width, parse trees, and monadic second-order logic for matroids, 20th Annual Symposium on Theoretical Aspects of Computer Science (STACS), LNCS vol. 2607 (2003), 319–330.
- [14] P. Hliněný: On matroid properties definable in the MSO logic, 27th International Symposium on Mathematical Foundations of Computer Science, MFCS (2003), 470– 479.
- [15] P. Hliněný: Branch-width, parse trees, and monadic second-order logic for matroids, Journal of Combinatorial Theory Series B 96 (2006), 325–351.
- [16] P. Hliněný and S. Oum: Finding branch-decompositions and rank-decompositions, 15th Annual European Symposium, ESA (2007), 163–174.
- [17] P. Hliněný and S. Oum: Finding branch-decompositions and rank-decompositions, SIAM Journal on Computing 38 (2008), 1012–1032.
- [18] P. Hliněný, O. Kwon, J. Obdržálek and S. Ordyniak: Tree-depth and vertex-minors, European Journal of Combinatorics 56 (2016), 45–56.
- [19] J. Jeong, E. J. Kim and S. Oum: Finding Branch-Decompositions of Matroids, Hypergraphs, and More, 45th International Colloquium on Automata, Languages, and Programming, ICALP (2018), 80:1–80:14.
- [20] F. Kardoš, D. Král', A. Liebenau and L. Mach: First order convergence of matroids, European Journal of Combinatorics 59 (2017), 150–168.
- [21] O. Kwon and S. Oum: Graphs of small rank-width are pivot-minors of graphs of small tree-width, Discrete Applied Mathematics 168 (2014), 108–118.

- [22] J. Nešetril and P. O. de Mendez: Sparsity: Graphs, Structures, and Algorithms, Algorithms and Combinatorics, volume 28, Springer, 2012.
- [23] S. Norin, A. Scott and D. R. Wood: Clustered colouring of graph classes with bounded treedepth or pathwidth, Combinatorics, Probability and Computing 32 (2023), 122–133.
- [24] S. Oum: Rank-width and vertex-minors, Journal of Combinatorial Theory Series B 95 (2005), 79–100.
- [25] S. Oum and P. Seymour: Approximating clique-width and branch-width, Journal of Combinatorial Theory Series B 96 (2006), 514–528.
- [26] J. Oxley: Matroid Theory, Oxford Graduate Texts in Mathematics, Oxford University Press, 2011.
- [27] N. Robertson and P. Seymour: Graph minors—a survey, Surveys in Combinatorics 1985, London Math. Soc. Lecture Note Ser. vol. 103 (1985), 153–171.