

The Complexity of Diameter on H -free graphs

Jelle J. Oostveen ^{*1}, Daniël Paulusma ², and Erik Jan van Leeuwen ¹

¹Dept. Information and Computing Sciences, Utrecht University, The Netherlands, j.j.oostveen@uu.nl, e.j.vanleeuwen@uu.nl

²Dept. Computer Science, Durham University, United Kingdom, daniel.paulusma@durham.ac.uk

Abstract

The intensively studied DIAMETER problem is to find the diameter of a given connected graph. We investigate, for the first time in a structured manner, the complexity of DIAMETER for H -free graphs, that is, graphs that do not contain a fixed graph H as an induced subgraph. We first show that if H is not a linear forest with small components, then DIAMETER cannot be solved in subquadratic time for H -free graphs under SETH. For some small linear forests, we do show linear-time algorithms for solving DIAMETER. For other linear forests H , we make progress towards linear-time algorithms by considering specific diameter values. If H is a linear forest, the maximum value of the diameter of any graph in a connected H -free graph class is some constant d_{\max} dependent only on H . We give linear-time algorithms for deciding if a connected H -free graph has diameter d_{\max} , for several linear forests H . In contrast, for one such linear forest H , DIAMETER cannot be solved in subquadratic time for H -free graphs under SETH. Moreover, we even show that, for several other linear forests H , one cannot decide in subquadratic time if a connected H -free graph has diameter d_{\max} under SETH.

1 Introduction

The DIAMETER problem asks to find the diameter of an unweighted graph, that is, the longest of the shortest paths between all pairs of nodes. A trivial algorithm executes a Breadth First Search (BFS) from every node in the graph, and has a running time of $O(nm)$. The best known matrix multiplication-based algorithms achieve a running time of $\tilde{O}(n^\omega)$ [16, 45, 48] to find the diameter of a graph, where \tilde{O} hides logarithmic factors and ω is the matrix multiplication constant, with current known value $\omega < 2.371866$ [26]. A search for improvement led to a hardness result; that, under the Strong Exponential Time Hypothesis (SETH), one cannot decide between diameter 2 or 3 on (sparse) split graphs in $O(n^{2-\epsilon})$ time, for any $\epsilon > 0$ [44]. SETH is a hypothesis that states that SATISFIABILITY cannot be solved in $2^{(1-\epsilon)n}$ time, for any $\epsilon > 0$, where n is the

*J.J. Oostveen is supported by the NWO grant OCENW.KLEIN.114 (PACAN).

number of variables [39, 40]. On other simple graph classes like constant degree graphs, truly subquadratic time algorithms for DIAMETER are also ruled out under SETH [33]. Directed versions of the DIAMETER problem admit similar barriers under SETH [4]. No clear bound is known for DIAMETER on dense graphs, but we do know that there is a subcubic equivalence between DIAMETER and computing the reach centrality of a graph, that is, a truly subcubic algorithm for one implies such an algorithm for the other and vice versa [1].

Given that the hardness results are based on long-standing conjectures, it is natural to approach diameter computation and other similar problems on restricted graph classes. Related literature also concerns computation of eccentricities, as computing the diameter of a graph is equivalent to computing the largest eccentricity over all vertices. A conceptually simple algorithm called LexBFS can solve DIAMETER in $O(n + m)$ time for distance-hereditary chordal graphs and interval graphs [25]. Distance-hereditary graphs have been studied separately, and admit linear-time algorithms of all eccentricities [19, 22, 24]. Interval graphs admit computation of the eccentricity of the center of the graph in linear time, next to linear-time diameter computation [43]. Subquadratic algorithms for DIAMETER and computing eccentricities have been studied for more graph classes, including asteroidal triple-free graphs [28], directed path graphs [13], strongly chordal graphs [17], dually chordal graphs [6, 18], Helly graphs and graphs of bounded Helly number [21, 30, 31], α_i -metric graphs [20], retracts [27], δ -hyperbolic graphs [11, 12, 22, 23], planar graphs [2, 8, 35], and outerplanar graphs [34]. DIAMETER was also studied from the parameterized perspective, see e.g. [4, 7, 15, 29, 32], and a large body of work exists on approximation algorithms, see e.g. [4, 5, 10, 13, 14, 20, 44, 46].

For graph classes with forbidden patterns, the result by Ducoffe et al. [32] shows subquadratic time computation of DIAMETER for \mathcal{H} -minor-free graphs, where the precise exponent depends on \mathcal{H} . Johnson et al. [41] showed for \mathcal{H} -subgraph-free graphs that there is a dichotomy for DIAMETER between almost-linear time solvability and quadratic-time conditional lower bounds depending on the family \mathcal{H} . Also note that many of the studied graph classes listed above can be characterized as \mathcal{H} -free graphs for a family of graphs \mathcal{H} , but for each one, \mathcal{H} has size 2 or larger. As far as we are aware, a structured study into forbidden (monogenic) induced patterns is absent in the literature.

Our Contributions. We initiate a structured study into diameter computation on H -free graphs, where H is a single graph. Recall that a graph is H -free if it does not contain H as an induced subgraph. The question we consider is:

For which H -free graph classes \mathcal{G} can the diameter of an n -vertex graph $G \in \mathcal{G}$ be computed in time $O(n^{2-\epsilon})$?

Our first result analyses existing lower bounds to find hardness for H -free graph classes. Recall that a *linear forest* is a disjoint union of one or more paths.

Theorem 1. *Under SETH, if H is not a linear forest, or H contains an induced $2P_2$, computing the diameter of an n -vertex H -free graph cannot be done in time $O(n^{2-\epsilon})$, for all $\epsilon > 0$.*

Theorem 1 shows the most prominent gap to be for graph classes which exclude a small linear forest. To complement the hardness result, we show

linear-time algorithms for several classes of H -free graphs for which H is a small linear forest.

Theorem 2. *Given a graph G that is H -free, we can solve DIAMETER in $O(n + m)$ time, when one of $H \subseteq P_2 + 2P_1$, $H \subseteq P_3 + P_1$ holds.*

We achieve Theorem 2 by careful structural analysis of the graph class and then show that a constant number of Breadth First Searches suffice algorithmically. Note that a running time of $O(n + m)$ clearly beats the naive algorithm of $O(nm)$ time and the matrix multiplication algorithms of $\tilde{O}(n^\omega)$ time, but also rules out any quadratic lower bound in n , as the classes of graphs contain arbitrarily large families of sparse graphs, e.g. stars.

The smallest graph H that is not covered by either Theorem 1 or Theorem 2 is $H = 4P_1$. As a ‘hardness’ result for $4P_1$ -free graphs, one could try to take the split graph construction of Roditty and Williams [44] (see Theorem 20 in Appendix A), and add edges to make the graph consist of three cliques. Conceptually, this would seem to work: the diameter distinction is still 2 or 3 and translates to a SAT positive or negative answer. However, this approach fails due to the quantity of edges one adds to the graph. The lower bound shows that no $O(n^{2-\epsilon})$ time algorithm may exist, which is now a relatively empty lower bound: the graph has a quadratic number of edges, so this lower bound does not even rule out an $O(n + m)$ time algorithm. The density of graphs matters in relation to lower bounds, and seems to provide a barrier to finding a lower bound that rules out a linear-time algorithm.

However, if we adopt the perspective from the other side, a linear-time algorithm for $4P_1$ -free graphs would still be surprising. Indeed, such an algorithm that can decide between diameter 2 or 3 on the three-clique instance described earlier implies an algorithm for ORTHOGONAL VECTORS in time $O(n^2 + d^2)$, where d is the dimension of the vectors and n the size of the vector sets (see a discussion in Appendix A). Although lower bounds do not rule out this possibility, such a result would be highly non-trivial, as the best known algorithms for ORTHOGONAL VECTORS do not achieve this running time for all d [3, 9]. Any linear-time algorithm would even beat the best known matrix-multiplication algorithms of $\tilde{O}(n^\omega)$ time, even if $\omega = 2$. It thus seems we are at an impasse to find or exclude a linear-time algorithm for computing the diameter of $4P_1$ -free graphs.

However, as it turns out, we *can* decide in linear time whether the diameter of a $4P_1$ -free graph is exactly 5. Our approach avoids the above barriers by focusing on specific diameter values instead of deciding on the diameter of a graph completely.

In general, let us call $d_{\max}(\mathcal{G})$ the maximum diameter any graph in the graph class \mathcal{G} can have (see Section 2 for a precise definition). In particular, for $4P_1$ -free graphs, d_{\max} is equal to 5. The research question we further investigate is:

For which H -free graph classes \mathcal{G} can we decide whether the diameter of a graph $G \in \mathcal{G}$ is equal to $d_{\max}(\mathcal{G})$ in linear time?

For some classes \mathcal{G} , it is easy to see $d_{\max}(\mathcal{G})$ is bounded. For instance, the class of cliques has $d_{\max} = 1$. Any graph class that contains paths of arbitrary length has $d_{\max} = \infty$. For deciding whether the diameter of a graph is equal to d_{\max} , only classes with bounded d_{\max} value are interesting to consider. It turns out

that for classes of connected H -free graphs, d_{\max} is bounded exactly when H is a linear forest (see Theorem 5 in the preliminaries).

Our contributions with respect to deciding whether $\text{diam}(G) = d_{\max}(\mathcal{G})$ for a given graph G and graph class \mathcal{G} are twofold.

Firstly, we find several examples of H -free classes \mathcal{G} where H is a linear forest of more than one path where we can decide in linear time whether $\text{diam}(G) = d_{\max}(\mathcal{G})$. Note that d_{\max} can differ vastly for classes where H is a linear forest, depending on H .

Theorem 3. *Given a graph G from a class of graphs \mathcal{G} that is H -free, we can decide whether $\text{diam}(G) = d_{\max}(\mathcal{G})$ in $O(n+m)$ time, when one of $H \subseteq P_2 + 2P_1$, $H \subseteq P_4 + P_1$, $H \subseteq P_3 + 2P_1$, $H \subseteq P_2 + 3P_1$, or $H = 2P_2 + P_1$ holds.*

Secondly, we extend known hardness constructions to hold for $\text{diam}(G) = d_{\max}(\mathcal{G})$ computation for certain H -free graph classes \mathcal{G} .

Theorem 4. *Under SETH, if \mathcal{G} is the class of $2P_2$ -free graphs or P_t -free graphs with $t \geq 5$, t odd, we cannot decide whether the diameter of a graph $G \in \mathcal{G}$ is equal to $d_{\max}(\mathcal{G})$ in time $O(n^{2-\epsilon})$, for all $\epsilon > 0$.*

Theorems 3 and 4 together cover almost all cases where $d_{\max} \leq 4$; $H = 2P_2$ is hard by Theorem 4, $H = P_3 + P_2$ is open, and all other cases with $d_{\max} \leq 4$ are linear-time solvable by Theorem 3. Theorem 3 also gives linear-time algorithms for some cases where $d_{\max} > 4$; $H = P_3 + 2P_1$ and $H = 2P_2 + P_1$ have $d_{\max} = 5$, and $H = P_2 + 3P_1$ has $d_{\max} = 6$. It appears that algorithmically, the presence of a P_1 in H helps out in structural analysis, which may explain the inability to attain a result for $H = P_3 + P_2$. We further discuss particular cases and possible generalizations of our theorems in the conclusion.

Our algorithmic results are mostly attained through careful analysis of the structure of the graph, with respect to the forbidden pattern. This limits the ways in which a shortest path that realizes the diameter can appear in the graph. However, even for small patterns H , such analysis quickly becomes highly technical.

We prove our hardness results in Section 3 and give algorithmic results in Section 4, and prove Theorems 2 and 3 in Section 4.7. We discuss our results, conjectured generalizations, and open questions in the conclusion, see Section 5.

2 Preliminaries

A graph $G = (V, E)$ has a vertex set V and edge set E . Graphs are connected, undirected, and unweighted. For any $v \in V$ denote $N(v)$ as the neighbourhood of v , and $N[v] = N(v) \cup \{v\}$ as the neighbourhood including v . For a set of vertices $S \subseteq V$, let $G[S]$ denote the induced subgraph on the vertices of S , that is, the vertices of S and all edges present in G between those vertices. A vertex $v \in V$ is *complete* to a set $S \subseteq V$ when $S \subseteq N(v)$, and *anti-complete* to a set S when $S \cap N(v) = \emptyset$. A set $A \subseteq V$ is complete to a set $S \subseteq V$ when every vertex in A is complete to S , and A is anti-complete to S when every vertex in A is anti-complete to S . For vertices $v_1, \dots, v_k \in V$, we use $\langle v_1, \dots, v_k \rangle$ to denote an induced path from v_1 to v_k .

A graph is H -free when it does not contain H as an induced subgraph. For graphs G, H let $G + H$ denote the disjoint union. A linear forest is a disjoint

union of one or more paths. We also use a row of integers $(a_i)_{i=1}^k$ for some integer $k \geq 1$, to consider $H = \sum_{i=1}^k P_{a_i}$ -free graphs, which models every possible linear forest for H . $H \subseteq G$ denotes that H is an induced subgraph of G .

Denote for two vertices u, v the distance between u and v as $d(u, v)$. The diameter of a graph G is the length of the longest shortest path, that is, $\text{diam}(G) = \max_{u, v \in V} d(u, v)$. A pair of vertices $u, v \in V$ is called a diametral pair when $d(u, v) = \text{diam}(G)$. A shortest path between a diametral pair u, v of length $d(u, v)$ is called a diametral path. The problem of DIAMETER is to decide on the value of $\text{diam}(G)$ for a given graph G . In the literature, the variant is also considered where we have to report the diametral path. For our algorithms, this makes no difference, as we can always execute a Breadth First Search (BFS) to find diametral paths if we found a diametral pair.

We assume our graphs are connected, and so the diameter is never ∞ . This is not a limiting assumption, as any BFS in time $O(n + m)$ can verify this, and our algorithms run in time $O(n + m)$. Also, for any graph, we can decide in time $O(n + m)$ whether it is a clique, and so whether its diameter is equal to 1.

For a graph class \mathcal{G} (of connected graphs), denote $d_{\max}(\mathcal{G})$ as the supremum of the diameter over all graphs $G \in \mathcal{G}$, that is,

$$d_{\max}(\mathcal{G}) = \sup_{G \in \mathcal{G}} \text{diam}(G).$$

We omit \mathcal{G} when it is clear from context. For instance, for the class of cliques, $d_{\max} = 1$. For many graph classes, d_{\max} is unbounded. However, we can consider H -free graphs where H is some linear forest, as such cases exclude paths of a certain length as induced subgraphs, bounding the diameter. Let us prove this formally.

Theorem 5. *Given a class \mathcal{G} of connected H -free graphs, the following statements hold.*

1. $d_{\max}(\mathcal{G})$ is bounded if and only if H is a linear forest, and
2. if H is a linear forest, say $H = \sum_{i=1}^k P_{a_i}$ for some row of integers $(a_i)_{i=1}^k$ for some integer $k \geq 1$, then $d_{\max}(\mathcal{G}) = k - 3 + \sum_{i=1}^k a_i$.

Proof. Assume for sake of contradiction that \mathcal{G} is a class of connected H -free graphs where H is not a linear forest. We may assume H contains either a cycle or a vertex of degree at least 3. We can conclude that the graphs $G = P_t$ for every integer $t \geq 1$ are contained in \mathcal{G} . But then no bound on the diameter suffices, as we can always pick a $t \geq 1$ larger than the bound to find a graph with larger diameter.

Now assume \mathcal{G} is a class of connected H -free graphs with H a linear forest, say $H = \sum_{i=1}^k P_{a_i}$ for some row of integers $(a_i)_{i=1}^k$, for some integer $k \geq 1$. Note that any $G \in \mathcal{G}$ cannot contain an induced path on $k - 1 + \sum_{i=1}^k a_i$ vertices, as such a path would contain an induced copy of H . We see that the largest induced path in any graph $G \in \mathcal{G}$ has at most $k - 2 + \sum_{i=1}^k a_i$ vertices, and hence the length of any shortest path is at most $k - 3 + \sum_{i=1}^k a_i$. This provides the bound on the $\text{diam}(G)$ for any graph $G \in \mathcal{G}$. \square

Two vertices $u, v \in V$ are twins when $N[u] = N[v]$, also called *true twins*. Two vertices $u, v \in V$ are *false twins* when $N(u) = N(v)$. A *twin class* is a set of

vertices all of which are pairwise twins. With respect to diametral paths, twins are not interesting, as they behave exactly the same in terms of distances. In some procedures, we would like to remove twins from the graph, such that only one each of the vertices in a twin class remains. Habib et al. [36] showed that twins can be identified in time $O(n + m)$, using partition refinement techniques (see also [38]).

Theorem 6 ([36]). *Given a graph G , we can detect true and false twins in G in $O(n + m)$ time.*

Proposition 7. *Given a graph $G = (V, E)$, in $O(n + m)$ time we can detect true twins and remove all-but-one vertex from each twin class resulting in a graph $G' = (V', E')$ with $V' \subseteq V$, $E' \subseteq E$. The following hold*

- (i) *the diameter of G' is equal to the diameter of G , unless G is a clique,*
- (ii) *the distance between two vertices u, v in G' is equal to the distance between u, v in G .*
- (iii) *if G is H -free for some graph H then G' is also H -free.*

Proof. Detect classes of true twins in linear time using Theorem 6. We can remove all-but-one vertex of each twin class, by first marking vertices for removal, and then in linear time enumerating the graph to create a new graph without the marked vertices and their edges. This is the graph G' . Let $\phi : V \rightarrow V'$ map vertices of G to its twin in G' that was not removed; note that this is the identity function for vertices with no twins in G .

If G' has only one vertex, then there are no shortest paths. This can be the case when G is a clique.

Let $u', v' \in V'$. Let $u, v \in V$ be the same vertices in G , and denote the shortest path between u and v with $\langle u = w_0, w_1, \dots, w_k = v \rangle$. Now $\langle u' = \phi(w_0), \phi(w_1), \dots, \phi(w_k) = v' \rangle$ must be a shortest path from u' to v' in G' , as twins have the exact same neighbourhood, and distances cannot be shortened by vertex removal.

Assume G is H -free for some graph H , and assume for sake of contradiction that G' is not H -free. Then, there is some set of vertices $A' \subseteq V'$ such that $G'[A']$ forms an induced copy of H . Take for each $v' \in A'$ some vertex in $v \in V$ such that $\phi(v) = v'$, and call the set of these vertices A . Because v' and v are twins, it must be that $G[A]$ is also an induced copy of H in G , a contradiction. \square

This procedure does not only work for true twins. Assume we have a set $B \subset V(G)$ and a set $V(G) \setminus B$ where we would like to partition B into classes of twins with respect to the edges towards $V(G) \setminus B$ (so irrespective of edges in $G[B]$). We note that we can do this in linear time, by adjusting the algorithm of Theorem 6 by having the base set be B and refining (partitioning the base set) only for every $x \in V(G) \setminus B$ on $N(x) \cap B$. We get the following corollary.

Corollary 8. *Given a graph G and a vertex set $B \subseteq V(G)$, we can partition B into classes of twins with respect to their neighbourhoods towards $V(G) \setminus B$ in time $O(n + m)$.*

3 Proofs of Theorem 1 and 4

In general, showing hardness for DIAMETER for some H also shows hardness for all H' -free graphs where $H' \supseteq H$ is a graph that contains H as an induced subgraph. Showing a linear-time, i.e. $O(n + m)$ time, algorithm for DIAMETER for some H also shows linear-time algorithms for all H' -free graphs, where $H' \subseteq H$ is a graph contained as an induced subgraph in H .

Theorem 1 (Restated). *Under SETH, if H is not a linear forest, or H contains an induced $2P_2$, computing the diameter of an n -vertex H -free graph cannot be done in time $O(n^{2-\epsilon})$, for all $\epsilon > 0$.*

Proof. It is well-known that deciding between diameter 2 or 3 on split graphs is hard [44]; see also Appendix A for an illustration and proof of this result. Split graphs are $(2P_2, C_4, C_5)$ -free [37]. Hence, if H contains a cycle of length at least 4 as an induced subgraph, then computing diameter on H -free graphs is hard by the split graph construction (it is chordal). If H contains a triangle, we can subdivide the split graph construction from [44] to make the graph H -free, as by [41, Lemma 1], hardness of computing the diameter is retained under 2-edge-subdivision. As we 2-subdivide once, the lower bound is preserved (although the distinction values change). So H does not contain a cycle. If H contains a vertex of degree at least three, computing the diameter on H -free graphs is hard, as the construction by Evald and Dahlgaard [33] can be made claw-free by adding chords to all binary trees in the construction (without affecting correctness). Hence, the remaining cases are where H is a linear forest. As the split graph construction shows hardness for $2P_2$ -free graphs, deciding between diameter 2 and 3 is hard for any H -free graph where the linear forest H contains a $2P_2$. \square

The split graph hardness construction by Roditty and Williams [44] shows that deciding between diameter 2 or 3 is hard for all P_t -free graphs with $t \geq 5$. However, many such P_t -free graphs may have diameter much larger than just 2 or 3; in particular, d_{\max} may be much larger. Therefore, this hardness construction does not necessarily rule out being able to compute d_{\max} for many such $H = P_t$. To mitigate this, we extend the hardness construction by Roditty and Williams [44] to rule out d_{\max} computation for all $H = P_t$ -free graphs for odd t , $t \geq 5$.

Theorem 4 (Restated). *Under SETH, if \mathcal{G} is the class of $2P_2$ -free graphs or P_t -free graphs with $t \geq 5$, t odd, we cannot decide whether the diameter of a graph $G \in \mathcal{G}$ is equal to $d_{\max}(\mathcal{G})$ in time $O(n^{2-\epsilon})$, for all $\epsilon > 0$.*

Proof. In this proof we heavily use the split graph hardness construction by Roditty and Williams [44]; see also Appendix A for the construction by Roditty and Williams. The construction shows that we cannot decide between diameter 2 or 3 on m -edge (sparse) split graphs in time $O(m^{2-\epsilon})$ for all $\epsilon > 0$ unless SETH fails. Split graphs are $2P_2$ -free [37]. For $2P_2$ -free graphs $d_{\max} = 3$ by Theorem 5, so hardness for this class immediately follows.

Let $t \geq 5$ be an odd integer. We prove that deciding on $d_{\max}(P_t\text{-free})$ is hard under SETH, that is, we show hardness for deciding between diameter $d_{\max} = t - 2$ and $d_{\max} - 1 = t - 3$, which are the correct values by Theorem 5.

Let G be any split graph hardness instance with n vertices and m edges. As G is a split graph, it is $2P_2$ -free and so P_5 -free. For $t = 5$ this proves




Figure 1: A sketch of a $(P_2 + 2P_1)$ -free graph as seen from some vertex u .

the theorem. Consider the following augmentation: add to every vertex in the independent set a path of $c = \frac{t-5}{2}$ edges. This adds at most cn vertices and edges to the graph. Let G' denote the resulting graph with $O(n + cn) = O(n)$ vertices and $O(m + cn) = O(m)$ edges. The diameter of G' is either $2 + 2c$ or $3 + 2c$, depending on the diameter of G . Deciding on the diameter of G' directly implies deciding on the diameter of G , so the lower bound carries over. Therefore, the hardness of deciding on the diameter of G' holds for deciding between $3 + 2c = t - 2$ and $2 + 2c = t - 3$. Because induced paths have been extended by at most $2c$ vertices and G is P_5 -free, we get that G' is P_{5+2c} -free. The theorem follows. \square

For even t , the construction does not work, as the maximum diameter of the constructed graphs is always odd. We conjecture that Theorem 4 generalizes to all $t \geq 5$, not just odd t ; see the conclusion for a discussion. Note that when $t \leq 4$, the diameter is at most 2, and so it suffices to check whether the graph is a clique, so indeed hardness can only hold for $t \geq 5$.

4 Algorithmic Results

4.1 $(P_2 + 2P_1)$ -free graphs

In this section we prove that the diameter of a $(P_2 + 2P_1)$ -free graph can be computed in linear time. The statement of the theorem is slightly stronger however, as we will need this algorithm as a subroutine in another algorithm later on.

Theorem 9. *Given a graph G , there is an algorithm that in $O(n + m)$ time either (a) correctly decides that G is not $(P_2 + 2P_1)$ -free; or (b) outputs a shortest path, which is diametral if G is $(P_2 + 2P_1)$ -free.*

Proof. Let $G = (V, E)$ be a graph. The diameter of any $(P_2 + 2P_1)$ -free graph is at most 4 by Theorem 5. If the diameter of G is 1, then the graph is a clique, which we can check in $O(n + m)$ time, and return any arbitrary pair of vertices, or a single vertex if $|V| = 1$.

Remove twins from the graph in $O(n + m)$ time. By Proposition 7, distances and the diameter are not affected, and the graph is H -free if it was H -free for some graph H . By abuse of notation, we still call this graph $G = (V, E)$.

Let u be a vertex in G with lowest degree, which can be found in $O(n + m)$ time. Now execute a BFS from u . We distinguish the structure of the graph as seen from u ; see Figure 1. Let $C = V \setminus N[u]$ and let $A \subseteq N(u)$ be the subset of vertices of $N(u)$ with no edges to C . Let $B = N(u) \setminus A$. Note that A, B, C can be identified by the BFS from u . If $C = \emptyset$, then the diameter of G is at most 2, and we are done. Hence, $C \neq \emptyset$ and $B \neq \emptyset$. We note that $A = \emptyset$, which can be seen by the following. For any $a \in A$ it holds that $\deg(a) \leq \deg(u)$ by definition of A . u was picked to be a vertex of lowest degree in G , so for any $a \in A$ we have $\deg(a) = \deg(u)$. But as $N(a) \subseteq \{u\} \cup A \cup B$ for all $a \in A$ and $N(u) = A \cup B$, it follows that every $a \in A$ is a twin of u . But then $A = \emptyset$ as we removed twins from G .

If there is a vertex at distance 5 or more from u , then return that the graph is not $(P_2 + 2P_1)$ -free. If there is a vertex at distance 4 from u , then return this shortest path; it is diametral if G is $(P_2 + 2P_1)$ -free. Both of these cases are identified by the BFS from u . Now observe that any shortest path of length 3 or 4 must have at least one endpoint in C , as the distances between vertices in $A \cup B \cup \{u\}$ are at most 2 by u .

We prove properties of G under the assumption that it is $(P_2 + 2P_1)$ -free.

Claim 9.1. *If G is $(P_2 + 2P_1)$ -free, then (a) $G[C]$ is a complete r -partite graph for some $r \geq 1$, and (b) every $b \in B$ has at most one non-neighbour in every part of $G[C]$.*

Proof. To prove (a), because u is non-adjacent to all of C , $G[C]$ is a $(P_2 + P_1)$ -free graph. The complement of a $(P_2 + P_1)$ -free graph is P_3 -free, which is a disjoint union of cliques, and the complement of a disjoint union of cliques is a complete r -partite graph, for some $r \geq 1$.

To prove (b), for sake of contradiction, assume that a vertex $b \in B$ has two non-neighbours c_1, c_2 in a single part of $G[C]$. Then, (u, b) together with c_1, c_2 form a $P_2 + 2P_1$, a contradiction. \square

Claim 9.2. *In $O(n + m)$ time, we can decide whether $G[C]$ is a complete r -partite graph for some $r \geq 1$ and, if so, return its parts.*

Proof. Detect false twins in $G[C]$ in $O(n + m)$ time using Theorem 6. Let r be the number of false twin classes, note that $r \leq n$. Initialize an array of size r in $O(n)$ time, and count for each class the number of vertices in it in $O(n)$ time total. Now we have to check that each vertex is complete to all vertices except those in its class. To do this, iterate for each vertex v over its adjacency list and count the number of neighbours in C . The degree of a vertex v should come out to be the number of vertices in C minus the size of its class, which are both known.

If this process succeeds, we have r classes, let us call them parts. Every vertex in a part is complete to all other parts, and has no neighbours in its part.

Hence, this is a complete r -partite graph. We can return the parts by returning each false twin class. \square

Claim 9.3. *In $O(n + m)$ time, either (a) we find a length-4 shortest path with both endpoints in C ; (b) we find a length-3 shortest path with both endpoints in C and conclude no such length-4 shortest path exists in G ; (c) we conclude no length-3 or length-4 shortest path with both endpoints in C exists in G ; or (d) we conclude G is not $(P_2 + 2P_1)$ -free.*

Proof. First execute the algorithm of Claim 9.2 to detect if $G[C]$ is a complete r -partite graph for some integer $r \geq 1$. If $G[C]$ is not a complete r -partite graph for some $r \geq 1$, by Claim 9.1, we can safely return option (d). If $G[C]$ is complete r -partite for some $r > 1$, then the distances between vertices of C are at most 2, and a shortest path of length 3 or 4 with both endpoints in C does not exist, and we return option (c).

If $G[C]$ is complete 1-partite, that is, $G[C]$ is an independent set, then look at the neighbourhood of any arbitrary $b \in B$. By Claim 9.1, if G is $(P_2 + 2P_1)$ -free, b is non-adjacent to at most one vertex $c \in C$. If b is adjacent to all of C , then return no shortest path of length 3 or 4 with both endpoints in C exist in G , as all vertices in C have b as a common neighbour, so return option (c). If b has more than one non-adjacency in C , then return option (d). Otherwise, c is the only candidate for an endpoint of a length-3 or length-4 shortest path, as all other pairs $c_1, c_2 \in C$ have b as a common neighbour. Execute a BFS from c and return a length-4 shortest path if found, option (a). If a longer shortest path is found, then return that G is not $(P_2 + 2P_1)$ -free, option (d). If instead only a length-3 shortest path is found, return it as option (b), and by this analysis, there is no shortest path of length 4 with both endpoints in C . \square

Run the algorithm of Claim 9.3. If it returns option (d), then output that G is not $(P_2 + 2P_1)$ -free. If it returns option (a), then output the length-4 shortest path the algorithm gives; it is diametral if G is $(P_2 + 2P_1)$ -free. In both other cases, we argue no length-4 diametral path can exist if G is $(P_2 + 2P_1)$ -free. If G is $(P_2 + 2P_1)$ -free, distances from vertices in B to vertices in C are at most 3, because $G[C]$ is r -partite and every $b \in B$ has at most one non-adjacent vertex per part of $G[C]$ by Claim 9.1. We already found a length-4 shortest path with u as an endpoint, if it exists. But then any length-4 shortest path has both endpoints in C , if it exists, as we already knew at least one endpoint was in C . In both options (b) and (c) we can conclude that no length-4 shortest path with both endpoints in C exists.

We continue as follows. If u has a vertex at distance 3, then we would already know by the BFS from u . Otherwise, every $c \in C$ is at distance 2 from u . If the algorithm of Claim 9.3 returned option (b), we can output a shortest path of length 3 with both endpoints in C ; it is diametral if G is $(P_2 + 2P_1)$ -free. If the algorithm of Claim 9.3 returned option (c), no length-3 shortest path with both endpoints in C exists in G . Hence, the only remaining case for a length-3 shortest path is that there is a vertex in B with distance 3 to a vertex in C . We prove a claim on the structure of $G[C]$ if G is $(P_2 + 2P_1)$ -free and a shortest path of length 3 exists from a vertex in B to a vertex in C .

Claim 9.4. *If G is $(P_2 + 2P_1)$ -free and a length-3 shortest path exists from some $b \in B$ to some $c \in C$, then (a) $G[C]$ has exactly one part with more than one*

vertex, and (b) all vertices in B with a vertex in C at distance 3 are adjacent to only that multi-vertex part and have exactly one non-neighbour in that part.

Proof. Assume G is $(P_2 + 2P_1)$ -free and a length-3 shortest path exists from some vertex in B to some vertex in C . By Claim 9.1, $G[C]$ is a complete r -partite graph for some $r \geq 1$. First, note that any $b \in B$ adjacent to multiple parts has distance at most 2 to all vertices in C , because $G[C]$ is complete r -partite. So, any $b \in B$ with a vertex at distance 3 in C is adjacent to exactly one part of $G[C]$. By assumption, there is such a vertex in B . Moreover, by Claim 9.1, every $b \in B$ has at most one non-neighbour in every part of $G[C]$. Hence, there is at most one part with multiple vertices in $G[C]$. If there is no part with multiple vertices, then every $b \in B$ has distance at most 2 to all $c \in C$. This proves (a). To prove (b), again note that any $b \in B$ adjacent to multiple parts has distance at most 2 to all vertices in C . By Claim 9.1, any $b \in B$ is adjacent to the multi-vertex part and has at most one non-neighbour in that part. Having zero non-neighbours in the multi-vertex part implies the distance to all vertices in C is at most 2. The claim follows. \square

By Claim 9.4, we can look for $G[C]$ to have simple structure. In particular, only one part may have multiple vertices. By Claim 9.2 we can detect if $G[C]$ is complete r -partite in $O(n + m)$ time, and, given the parts, check whether only one part has multiple vertices in $O(n + m)$ time. If this is not the case, then there is no length-3 shortest path from a vertex in B to a vertex in C by Claim 9.4, if G is $(P_2 + 2P_1)$ -free, and we may return a length-2 shortest path with as witness some non-adjacent pair of vertices and a common neighbour.

Otherwise, the structure is as Claim 9.4(a) and (b) suggest. Find all $b \in B$ only adjacent to the multi-vertex part with one non-neighbour in that part in $O(n + m)$ time, by iterating over the adjacency lists of the vertices in B . Let this be the set of vertices B' . Then look for each vertex in B' whether all its neighbours in B have the same non-adjacency in the multi-vertex part in $O(n + m)$ time. If there is a vertex $b \in B'$ that meets this requirement, and G is $(P_2 + 2P_1)$ -free, then this is a witness for diameter-3 shortest path: b is non-adjacent to one vertex $c \in C$ in a multi-vertex part of $G[C]$, which is the only part it is adjacent to, and $N(b) \cap N(c) = \emptyset$. So the distance from b to c is at least (and at most) 3. To verify, execute a BFS from any single one of these vertices. If a shortest path is found of length 4 or longer, then return that G is not $(P_2 + 2P_1)$ -free. If a length-3 shortest path is found, then return it. Otherwise, there is no length-3 shortest path in G from B to C , if G is $(P_2 + 2P_1)$ -free.

If in none of the above cases a confirmation for a shortest path of length 3 or 4 was found, and the graph is $(P_2 + 2P_1)$ -free and not a clique, then the diameter of G must be 2. Return some non-adjacent pair with some common neighbour as a shortest path in $O(n + m)$ time. \square

As an immediate corollary, we get that we can compute the diameter of $3P_1$ -free graphs in linear time, as any $3P_1$ -free graph is $(P_2 + 2P_1)$ -free.

Corollary 10. *Given a $3P_1$ -free graph G , we can compute the diameter of G in $O(n + m)$ time.*

4.2 $(P_3 + P_1)$ -free graphs

We use a characterization by Olariu [42] to argue we can find the diameter of $(P_3 + P_1)$ -free graphs in linear time. As a subroutine, we call on the algorithm of Theorem 9.

Theorem 11. *Given a $(P_3 + P_1)$ -free graph G , we can compute the diameter of G in $O(n + m)$ time.*

Proof. First note that the complement of a $P_3 + P_1$ is a paw. By a result by Olariu [42], a graph is paw-free if and only if each component is triangle-free or complete multi-partite. This characterization must hold for the complement of our $(P_3 + P_1)$ -free input graph G . Hence, our input graph G can be divided into parts, where each part is complete to every other part, and each part is either $3P_1$ -free or a P_3 -free cluster graph (the complements of triangle-free graphs and complete multi-partite graphs).

If G consists of only one part, then it is either cluster with diameter 1, or it is $3P_1$ -free, which we can solve by Corollary 10. When G consists of multiple parts, then its diameter can only be 2 if it is not a clique. However, it is not trivial to detect if we have one or multiple parts in linear time. Instead, if G is not a clique, then run the algorithm of Theorem 9. If G consists of only one part, then the algorithm will return a shortest path that is diametral, because any $3P_1$ -free graph is also $(P_2 + 2P_1)$ -free. If G instead consists of multiple parts, then there are no shortest paths of length more than 2 in G , so the algorithm will either return that G is not $(P_2 + 2P_1)$ -free or return a length-2 shortest path. In either case, we can correctly conclude that the diameter of G is 2. \square

4.3 $(P_4 + P_1)$ -free graphs

We next show that we can decide whether the diameter of a $(P_4 + P_1)$ -free graph is equal to d_{\max} in $O(n + m)$ time. The proof will identify all possibilities of a diameter-4 path occurring in relation to a BFS from an arbitrary vertex. Luckily, most cases reduce to some other case in the proof, and algorithmically speaking, only a few cases require algorithmic computation.

Theorem 12. *Given a $(P_4 + P_1)$ -free graph G , we can decide whether the diameter of G is equal to $d_{\max} = 4$ in $O(n + m)$ time.*

Proof. Let $G = (V, E)$ be a (connected) $(P_4 + P_1)$ -free graph. Indeed, $d_{\max} = 4$ by Theorem 5. We view the structure of G from a BFS from an arbitrary vertex u . Let $C = V \setminus N[u]$ and denote $B = N(u)$. If $C = \emptyset$ or $B = \emptyset$ the diameter of G is at most 2, so assume this is not the case. Note that both sets can be identified during the BFS from u with no overhead. Moreover, $G[C]$ is P_4 -free. We use the convention that $b_i \in B$ and $c_i \in C$.

We first list all possibilities for a diameter-4 shortest path to exist with respect to its structure; see Figure 2 for an illustration. We will call each such possibility a ‘type’. First, observe that at least one of the ends of any distance-4 shortest path must be in C , because distances in $G[B \cup \{u\}]$ are at most 2 via u . Here, we do not write down symmetries, because in an undirected graph the path $\langle v_1, \dots, v_k \rangle$ is equivalent to the path $\langle v_k, \dots, v_1 \rangle$. We now distinguish the following cases:

(1) If the path has u as one endpoint and the other endpoint is in C , then




Figure 2: An illustration of the types of a diameter-4 shortest path appearing in a $(P_4 + P_1)$ -free graph, with respect to the vertex u and its neighbourhood $N(u) = B$. Only the highlighted types **(1a)**, **(2a)**, **(3b)**, **(3c)** require algorithmic computation.

the only possibility is type **(1a)** $\langle u, b, c_1, c_2, c_3 \rangle$, as any vertex in B has u as a neighbour.

(2) If the path has one end in B and the other in C , then we have several options, depending on the number of vertices of B on the path. We have at least one vertex of B on the path because one endpoint is in B . We cannot have only one vertex in B on the path, because then the rest of the path is in C but $G[C]$ is P_4 -free. If we have exactly two vertices of B on the path, they can be adjacent or non-adjacent. Two non-adjacent vertices of B on the path give the possibilities of type **(2a)** $\langle b_2, u, b_1, c_1, c_2 \rangle$ and type **(2b)** $\langle b_1, c_1, b_2, c_2, c_3 \rangle$; the second vertex of B cannot appear later on the path because there would be a shorter path via u . Two adjacent vertices of B on the path give type **(2c)** $\langle b_1, b_2, c_1, c_2, c_3 \rangle$, and no other option exists. If we have three vertices in B on the path the only option is type **(2d)** $\langle b_1, b_2, b_3, c_1, c_2 \rangle$, because if some vertex of B appears later on the path there would be a shorter path via u . There can be at most three vertices in B on the path, because distances between vertices of B are at most 2 via u . Hence, these are all the options for a path with one endpoint in B and one in C .

(3) The last option is for the path to have both endpoints in C . Again, there are multiple cases depending on the number of vertices of B on the path. There cannot be zero vertices of B on the path, because $G[C]$ is P_4 -free. If there is only one vertex of B on the path, we have type **(3a)** $\langle c_1, c_2, b, c_3, c_4 \rangle$ and type **(3b)** $\langle c_1, b, c_2, c_3, c_4 \rangle$, and we cannot pass by u . If we have exactly two vertices

of B on the path and they are non-adjacent, we have type **(3c)** $\langle c_1, b_1, u, b_2, c_2 \rangle$ and type **(3d)** $\langle c_1, b_1, c_2, b_2, c_3 \rangle$. If we have two vertices of B on the path and they are adjacent, we have type **(3e)** $\langle c_1, c_2, b_1, b_2, c_3 \rangle$. If we have exactly three vertices of B on the path we have type **(3f)** $\langle c_1, b_1, b_2, b_3, c_2 \rangle$, which is clearly the only option. There can be at most three vertices in B on the path, because distances between vertices of B are at most 2 via u . Hence, these are all the options for a path with both endpoints in C .

Because we listed all length-4 shortest paths with one endpoint in C and the other endpoint in u, B , or C , and $V = \{u\} \cup B \cup C$, these must be all the options for a length-4 shortest path to appear in G .

To prove the computability of some types, the following structural observation will come in useful.

Claim 12.1. *If a vertex $b \in B$ has a neighbour and a non-neighbour in a single component of $G[C]$, then b is complete to all other components of $G[C]$.*

Proof. Let $c_1, c_2 \in C$ be such that c_1 is a non-neighbour of b in a component of $G[C]$, and $c_2 \in C$ is a neighbour of both c_1 and b (such a pair exists by assumption). Notice that $\langle c_1, c_2, b, u \rangle$ forms an induced P_4 . Consider a vertex $c_3 \in C$ in another component of $G[C]$. It has no edges to either c_1 or c_2 , as it is in another component, and it is non-adjacent to u , as it is in C . But then, as the graph is $(P_4 + P_1)$ -free, (c_3, b) must be an edge of the graph. \square

We show we can decide the diameter of G is 4 in time $O(n + m)$ by resolving each type. Algorithmically speaking, only types **(1a)**, **(2a)**, **(3b)**, and **(3c)** will require computation to find diametral paths corresponding to them. We will show that all other types are either covered by these computations, or are non-existent in G .

(1a) $\langle u, b, c_1, c_2, c_3 \rangle$ This type is identified by the initial BFS from u if and only if it occurs in G .

(2a) $\langle b_2, u, b_1, c_1, c_2 \rangle$ For this type, let us further partition C into sets C' and D , where D consists of the vertices with no neighbours in B (i.e. the vertices at distance 3 from u), and $C' = C \setminus D$. It must be that $c_2 \in D$ for a diametral path of type **(2a)** to exist, otherwise, $\langle c_2, b_i, u, b_2 \rangle$ is a shorter path for some $b_i \in B$. So rename the vertex c_2 as $d = c_2 \in D$ and look for a path $\langle b_2, u, b_1, c_1, d \rangle$. Further partition C' into C_1 and C_2 , where C_1 are the vertices with edges towards D and $C_2 = C' \setminus C_1$. This partitioning can be done during the BFS from u , or alternatively using another linear pass over all vertices and edges. Note that every vertex in D has at least one neighbour in C_1 ; otherwise, a diametral path of type **(1a)** exists in G , and we are done. Let us first describe the algorithmic steps necessary for this type.

(2a.algorithm) Find a vertex $d \in D$ with the smallest degree with respect to C_1 , and execute a BFS from d . If a distance-4 vertex is found, return that the diameter of G is 4. If we do not find a vertex at distance 4, no distance-4 diametral path of type **(2a)** exists in G .

We next prove correctness of **(2a.algorithm)**. We prove correctness when $G[D]$ is not connected in **(2a.1)** and correctness when $G[D]$ is connected in

(2a.2). To do this, we analyse the structure of G under the assumption that a diametral path of type **(2a)** exists, to conclude the structure of G must then be ‘simple’ in some way, to the extent that the above algorithmic steps suffice.

(2a.1) Assume $G[D]$ is not connected. We first prove that every vertex in C_1 is complete to D . Assume for sake of contradiction that there is a $c \in C_1$ which is not complete to D . Let $d' \in D$ be a non-neighbour of c . Let $d \in D$ be some neighbour of c , which exists because $c \in C_1$. Now $\langle d, c, b, u \rangle$ is an induced P_4 for some $b \in B$ which exists by definition of C_1 . But then d' must be a neighbour of d ; otherwise, it would induce a $P_4 + P_1$. We see that every neighbour of c is adjacent to every non-neighbour of c in D . We get a contradiction with the assumption that $G[D]$ is not connected.

So, every vertex in C_1 must be complete to D . But then, from the viewpoint of shortest paths, for any $b \in B$, the distances from all $d \in D$ to b must be equal, as the shortest path to b must go through some vertex of C_1 in the first step, and C_1 is complete to D . Hence, if the BFS that **(2a.algorithm)** executes does not find a diametral path of length 4, no diametral path of type **(2a)** exists in G .

(2a.2) Assume $G[D]$ is connected. Then $G[C_1 \cup D]$ is a connected cograph, so it has diameter at most 2. Let $B_1 \subseteq B$ be the vertices of B with neighbours in C_1 . Vertices in $B_2 = B \setminus B_1$ have no neighbours in C_1 . Every vertex in B_1 has distance at most 3 to any $d \in D$, as the diameter of $G[C_1 \cup D]$ is at most 2. So, for a shortest path of type **(2a)** $\langle b_2, u, b_1, c_1, d \rangle$ we get $b_2 \in B_2$ and $b_1 \in B_1$.

Let us call a pair (d, b_2) with $d \in D, b_2 \in B_2$ ‘good’ if d has distance 4 to b_2 , and ‘bad’ when d has distance at most 3 to b_2 . Assuming a diametral path of form $\langle b_2, u, b_1, c_1, d \rangle$ exists in G , with $b_2 \in B_2, b_1 \in B_1, c_1 \in C_1, d \in D$, it is clear that (d, b_2) is good. Assume that we also have that (d', b_2) is bad for some $d' \neq d \in D$. Then d' has distance exactly 3 to b_2 , as b_2 has no neighbour in C_1 . Let c'_1 be the neighbour of d' on some distance-3 path from d' to b_2 . Then $c_1 \neq c'_1$; otherwise, d has distance 3 to b_2 . Then the shortest path from d' to b_2 is of the form $\langle d', c'_1, c'_2, b_2 \rangle$ with $c'_2 \in C_2$ (case **(2a.2.1)**), or $\langle d', c'_1, b'_1, b_2 \rangle$ with $b'_1 \in B_1, b'_1 \neq b_1$ (case **(2a.2.2)**). See Figure 3 for an illustration of both scenarios.

(2a.2.1) The shortest path from d' to b_2 is of the form $\langle d', c'_1, c'_2, b_2 \rangle$ with $c'_2 \in C_2$. Note that $(c_1, u), (c'_1, u) \notin E$ by definition of C , and $(c_1, b_2), (c'_1, b_2) \notin E$ by definition of B_2 , and $(b_1, b_2), (d, c'_1), (c_1, c'_2), (d, c'_2) \notin E$ as (d, b_2) is good. But then $(c'_1, c_1) \in E$ or $(c'_1, b_1) \in E$; otherwise, $\langle c_1, b_1, u, b_2 \rangle + c'_1$ is an induced $P_4 + P_1$ in the graph.

- If $(c'_1, c_1) \in E$, then $\langle c'_2, c'_1, c_1, d \rangle + u$ is an induced $P_4 + P_1$ in the graph; u is non-adjacent to all of c'_2, c'_1, c_1, d by definition of C and D .
- If $(c'_1, b_1) \in E$ then $\langle b_2, u, b_1, c'_1 \rangle + d$ is an induced $P_4 + P_1$ in the graph; $(c'_1, u) \notin E$ by definition of C , and $(d, b_1), (d, b_2), (d, u) \notin E$ by definition of D .

Hence, we get a contradiction, and this case cannot occur in G .




Figure 3: Structure for type **(2a)** with respect to a $d \in D$ and a $d' \in D$ for which (d, b_2) is good and (d', b_2) is bad for $b_2 \in B_2$. The path from d' to b_2 goes through either some $c'_2 \in C_2$ (left, **(2a.2.1)**) or some $b'_1 \in B_1$ (right, **(2a.2.2)**).

(2a.2.2) The shortest path from d' to b_2 is of the form $\langle d', c'_1, b'_1, b_2 \rangle$ with $b'_1 \in B_1$, $b'_1 \neq b_1$. Note that $(c_1, u) \notin E$ by definition of C , $(d', b_1), (d', b_2), (d', u) \notin E$ by definition of D , and $(b_1, b_2), (c_1, b_2) \notin E$, as (d, b_2) is good. But then $(d', c_1) \in E$: otherwise, $\langle c_1, b_1, u, b_2 \rangle + d'$ is an induced $P_4 + P_1$ in the graph.

Say that d has some other neighbour $c''_1 \in C_1$, so $(c''_1, b'_1) \in E$ for some $b'_1 \in B_1$ (possibly $b'_1 = b_1$), but $(b'_1, b_2) \notin E$ because (d, b_2) is good. Then, c''_1 can fulfil the role of c_1 in the above analysis, so it must be that $(d', c''_1) \in E$. From this analysis we can conclude that, for any $b_2 \in B_2$, if $d \in D$ is such that (d, b_2) is good, and $d' \in D$ is such that (d', b_2) is bad, then it must be that $N(d) \cap C_1 \subset N(d') \cap C_1$. Hence, if some $d \in D$ has minimum degree to C , it may be good.

We now have the tool to prove correctness of **(2a.algorithm)** for this case. If the BFS from the picked $d \in D$ finds a vertex at distance 4, then clearly a length-4 diametral path exists in G . The only risk is that we conclude there is no diametral path corresponding to this case even though it does exist. To this end, let $d \in D$ be the vertex picked by the algorithm, and assume (d, b) is bad for all $b \in B_2$. Assume to the contrary that there exist $d' \in D$, $b_2 \in B_2$ which are at distance 4 in a path of type **(2a)**. Then (d', b_2) is good. So, by the analysis above, it must be that $N(d') \cap C_1 \subset N(d) \cap C_1$, which contradicts the assumption that d was picked to have the minimal size neighbourhood with respect to C_1 . So all pairs (d, b_2) with $d \in D$, $b_2 \in B_2$ must be bad, and we are correct to conclude that a diametral path of type **(2a)** does not exist.

(2b) $\langle b_1, c_1, b_2, c_2, c_3 \rangle$ As b_1 and b_2 are both adjacent to u , if type **(2b)** exists in G , $\langle b_1, u, b_2, c_2, c_3 \rangle$ is also a shortest path of distance 4 from b_1 to c_3 , which is a distance-4 path of type **(2a)**. The algorithm for type **(2a)** finds a diametral path of length 4 or concludes that no diametral path of type **(2a)** exists in G , which also rules out that a diametral path of type **(2b)** exists in G .

(2c) $\langle b_1, b_2, c_1, c_2, c_3 \rangle$ We analyse the structure of the graph if type **(2c)** exists in G . Notice first that c_3 cannot be adjacent to any $b \in B$, as that would make

the distance from c_3 to b_1 at most 3 via u . Now, either the distance from u to c_3 is also 4, which is identified by the algorithm for type **(1a)** if it exists, or c_2 has a neighbour $b_3 \in B$. Note that $b_3 \neq b_1$ and $b_3 \neq b_2$, as the distance from b_1 to c_3 is 4. If $(b_1, b_3) \in E$, then the distance from b_1 to c_3 is not 4. So, $(b_1, b_3) \notin E$. But then $\langle b_1, u, b_3, c_2, c_3 \rangle$ is also a shortest path from b_1 to c_3 of distance 4, which is a length-4 shortest path of type **(2a)**. The algorithm for type **(2a)** finds a diametral path of length 4 or concludes that no diametral path of type **(2a)** exists in G , which also rules out that a diametral path of type **(2c)** exists in G .

(2d) $\langle b_1, b_2, b_3, c_1, c_2 \rangle$ As all $b \in B$ are adjacent to u , if a shortest path of type **(2d)** exists, $\langle b_1, u, b_3, c_1, c_2 \rangle$ is also a shortest path of length 4 from b_1 to c_2 , which is a length-4 shortest path of type **(2a)**. The algorithm for type **(2a)** finds a diametral path of length 4 or concludes that no diametral path of type **(2a)** exists in G , which also rules out that a diametral path of type **(2d)** exists in G .

(3a) $\langle c_1, c_2, b, c_3, c_4 \rangle$ If a diametral path of type **(3a)** exists, notice that there are no edges between c_1, c_2 and c_3, c_4 , and c_1 is not adjacent to b , as otherwise the distance from c_1 to c_4 is not 4. Then, $\langle c_1, c_2, b, u \rangle$ together with c_4 form an induced $P_4 + P_1$ in the graph, a contradiction. So, this type cannot occur in a $(P_4 + P_1)$ -free graph.

(3b) $\langle c_1, b, c_2, c_3, c_4 \rangle$ We first give an algorithm and then prove its correctness.

(3b.algorithm) Check in $O(n+m)$ time that there exists at least one and at most two components $C_1, C_2 \subseteq C$ of $G[C]$ with each at least one vertex $b \in B$ such that b has an edge and a non-edge to that component, by enumerating the adjacencies of vertices $b \in B$. If this is not the case, no diametral path of type **(3b)** exists in G . Let $B_{C_1}, B_{C_2} \subseteq B$ be the set of vertices with an edge and a non-edge to C_1 and C_2 , respectively, which we can find in $O(n+m)$ time. If only one such component exists, execute the following steps only for C_1 and B_{C_1} . For $i \in 1, 2$, mark all vertices in C_i with a neighbour in B_{C_i} , and then mark all vertices in C_i with a marked neighbour. This takes $O(n+m)$ time. Pick an arbitrary unmarked vertex in C_1 and one in C_2 , and execute a BFS from each. Return a distance-4 shortest path if it is found. Otherwise, we conclude that no diametral path of type **(3b)** exists in G .

We next prove the correctness of this algorithm. See Figure 4 for an illustration. If there is no component like C_1 or C_2 , no shortest path of the form $\langle c_1, b, c_2, c_3, c_4 \rangle$ exists, as it must be that $(b, c_2), (c_2, c_3) \in E$ while $(b, c_3) \notin E$ in the diametral path of this type. If there are at least three different components of $G[C]$, $C_1, C_2, C_3 \subseteq C$, which each have a vertex in B with an edge and a non-edge to that component, let these be $b_{C_1}, b_{C_2}, b_{C_3}$ respectively. The distance between any two $c_1, c_2 \in C$ is now at most 2, as by Claim 12.1, $b_{C_1}, b_{C_2}, b_{C_3}$ are complete to all components except C_1, C_2, C_3 , respectively. So, each pair $c_1, c_2 \in C$ has at least one common neighbour among $b_{C_1}, b_{C_2}, b_{C_3}$. So a diametral path of this type cannot occur in G . Hence, we are correct to look for at least one and at most two such components. We also see that one end of the diametral path must be in C_1 or C_2 , if it exists.




Figure 4: Structure for type **(3b)**, where for $i \in \{1, 2\}$, vertices in B_{C_i} are complete to all components of $G[C]$ except for C_i , due to Claim 12.1. C_1 and C_2 have layers corresponding to which vertices are marked by the algorithm, with the bottom layer consisting of unmarked vertices.

For $i \in \{1, 2\}$, all vertices in C_i marked by the algorithm have distance at most 3 to all $c \in C$: vertices in the same component are at distance at most 3, as $G[C]$ is P_4 -free. Next to this, marked vertices are at distance at most 2 from some vertex in B_{C_i} , and by Claim 12.1 vertices in B_{C_i} are complete to all other components of $G[C]$. So, the only candidates in C_1 or C_2 for one end of the distance-4 diametral path are the unmarked vertices. Assume that the algorithm does not find a distance-4 shortest path, but a diametral path of type **(3b)** does exist in G . Assume one end is in C_1 , and let $c_1 \in C_1$ be the vertex the algorithm picked in C_1 . The case where one end is in C_2 is analogous. Note that all vertices not in B_{C_1} are either complete or anti-complete to C_1 . Also, unmarked vertices in C_1 are at distance exactly 2 from all vertices in $C_1 \cap N(B_{C_1})$, as $G[C]$ is P_4 -free. We see that all unmarked vertices in C_1 have equal distance to each $b \in B$, and in particular, to the $b \in B$ the diametral path uses. Also, as the other end of the diametral path is in another component, the distance to that vertex is equal for every unmarked vertex in C_1 . But then the BFS from c_1 would have found a distance-4 shortest path, a contradiction. We can conclude the algorithm is correct, that is, if the algorithm does not find a distance-4 shortest path, no diametral path of type **(3b)** exists in G .

(3c) $\langle c_1, b_1, u, b_2, c_2 \rangle$ First note that $G[C]$ must consist of multiple connected components for a diametral path of type **(3c)** to occur: a connected P_4 -free graph has diameter at most 2, so the shortest path between $c_1, c_2 \in C$ will never go through u .

By Claim 12.1 we have that any vertex in B is either complete or anti-complete to every connected component of $G[C]$, or is not complete nor anti-complete to exactly one connected component of $G[C]$ and complete to all other connected components of $G[C]$. We distinguish cases **(3c.1)** and **(3c.2)**.




Figure 5: An illustration of the conversion of a graph G (left) to a graph G' (right) as in **(3c.1.algorithm)**. Note that for image simplicity, the drawn graph G is not actually a $(P_4 + P_1)$ -free graph.

(3c.1) Every $b \in B$ is complete or anti-complete to every component of $G[C]$. It can be checked in $O(n + m)$ time whether we are in this case. We first give the algorithm to solve this case, and then prove its correctness.

(3c.1.algorithm) For every component of $G[C]$, delete all-but-one vertex in $O(n + m)$ time, each component is a twin class with respect to B . Let the resulting set of vertices be C' . Identify twins within B with respect to their neighbourhood in C' , which can be done in linear time by Corollary 8. We identify every ‘class’ of vertices with the same neighbourhood in C' with single vertices. Let B' be the set of vertices corresponding to classes. We compute the neighbourhoods of the vertices in B' as the union of all neighbourhoods of vertices in the class. In particular, the edges of a $b' \in B'$ towards C' are simply the neighbourhood towards C' of an arbitrary $b \in B$ from that class, as every vertex in a class has an identical neighbourhood to C . To compute the adjacency list of vertices in B' , first initialize an array on the set of vertices of B' , in linear time. We use the array to avoid adding duplicate edges to adjacency lists, but only initialize it once. For each class $b'_1 \in B'$, associate a unique label with it, and enumerate the edge lists of all $b_1 \in B$ in the corresponding class. For an edge (b_1, b_2) , $b_2 \in B$ with class $b'_2 \in B'$, do the following. If the array does not contain the unique label of b'_1 at location b'_2 , overwrite the location b'_2 with the unique label of b'_1 , and add b_2 to the adjacency list of b'_1 . If the array does contain the label of b'_1 at location b'_2 , continue to the next edge. We see that we enumerate all edges in $G[B]$ at most twice, and do constant-time operations for each one. Hence, this procedure takes $O(n + m)$ time. Call the resulting graph $G' = (V', E')$ with $B', C' \subseteq V'$. An illustration of the conversion of a graph G to G' is given in Figure 5. In G' , check that $G'[B']$ consists of two non-empty disjoint cliques B_1, B_2 , where B_1 and B_2 are anti-complete, with possibly a third clique X complete to $B' \setminus X$. This check takes $O(n + m)$ time by inspecting the neighbourhoods of all vertices in B' . If $G'[B']$ does not have this structure, return that there is no diametral path of type **(3c)**. If it does, return that a diametral path of type **(3c)** exists if there are vertices $c_1, c_2 \in C'$ with $N(c_1) = B_1$ and $N(c_2) = B_2$. This can be checked in linear time by inspecting the edge lists of vertices $c \in C'$.

We next prove the correctness of the algorithm. We do this by proving several properties of G' , to then analyse the structure of G' using these properties.

Claim 12.2. *A diametral path of the form $\langle c_1, b_1, u, b_2, c_2 \rangle$ exists in G if and only if a diametral path of the form $\langle c'_1, b'_1, u, b'_2, c'_2 \rangle$ exists in G' , with $c'_1, c'_2 \in C'$ and $b'_1, b'_2 \in B'$.*

Proof. First, note that the process to get G' from G can be modelled as edge contractions and vertex identifications, which means the diameter of G' cannot be larger than that of G , and is at most 4.

Assume a diametral path of the form $\langle c_1, b_1, u, b_2, c_2 \rangle$ exists in G , with $c_1, c_2 \in C$ and $b_1, b_2 \in B$. Then the distance from c_1 to c_2 is 4 in G , and $(b_1, c_2) \notin E$, $(b_2, c_1) \notin E$, $(b_1, b_2) \notin E$, and c_1, c_2 do not have a common neighbour in G . Let $b'_1, b'_2 \in B'$ be the corresponding vertices of b_1, b_2 respectively in $G'[B']$, and let $c'_1, c'_2 \in C'$ be the vertices in the components of c_1, c_2 in $G[C']$ that are in C' . Then $c'_1 \neq c'_2$, as the diameter of a component of $G[C']$ is 2. Also, $b'_1 \neq b'_2$, as b_1 and b_2 do not have identical neighbourhoods w.r.t. C . It also holds that $(b'_1, b'_2) \notin E'$ as the distance between c_1 and c_2 is 4 in G , and so $N(c_1)$ and $N(c_2)$ are disjoint in G . Hence, $\langle c'_1, b'_1, u, b'_2, c'_2 \rangle$ is an induced path in G' , but it remains to show this is the shortest path between c'_1 and c'_2 and the distance between these vertices is 4. Assume to the contrary the distance from c'_1 to c'_2 is at most 3. As $c'_1 \neq c'_2$, the distance is at least 2. The distance between c_1 and c_2 is 4 in G , so $N[c_1]$ and $N[c_2]$ are disjoint, and as every vertex in B is complete or anti-complete to every component of $G[C]$, we see that $N[c'_1]$ and $N[c'_2]$ are disjoint in G' . So the distance between c'_1 and c'_2 is 3. Then, as $G[C']$ is an independent set, there exist $b'_3, b'_4 \in B'$ with $(b'_3, b'_4) \in E'$ and $(b'_3, c_1) \in E'$, $(b'_4, c_2) \in E'$. By the class identification process, we get that c'_1 is adjacent to the whole class corresponding to b'_3 , and similarly, c'_2 is adjacent to the whole class corresponding to b'_4 . But then there must exist $b_3, b_4 \in B$ with $(b_3, b_4) \in E$ and $(b_3, c_1) \in E$, $(b_4, c_2) \in E$, a contradiction. So the distance between c'_1 and c'_2 is also 4 in G' , and hence, c'_1 and c'_2 are a diametral pair in G' with diametral path $\langle c'_1, b'_1, u, b'_2, c'_2 \rangle$.

Assume a diametral path of form $\langle c'_1, b'_1, u, b'_2, c'_2 \rangle$ exists in G' , with $c'_1, c'_2 \in C'$ and $b'_1, b'_2 \in B'$. Then the distance from c'_1 to c'_2 is 4. Let $c_1, c_2 \in C$ be the vertices in G corresponding to c'_1 and c'_2 in G' . Because every vertex in B is complete or anti-complete to every component of $G[C]$, the vertex removal to get to C' did not increase the distance between c'_1 and c'_2 as compared to c_1 and c_2 . As edge/non-edge contraction can never increase the distance between two vertices, the distance between c_1 and c_2 must be at least 4 in G . The distance between c_1 and c_2 cannot be 5, as G is $(P_4 + P_1)$ -free. So the distance between c_1 and c_2 is also 4 in G . Let $b_1, b_2 \in B$ be two arbitrary vertices of the classes corresponding to b'_1, b'_2 , respectively. Then $(c_1, b_1) \in E$ and $(c_2, b_2) \in E$, as the algorithm identified classes with identical neighbourhoods towards C with single vertices and gave those vertices the same neighbourhood. Also, $(b_1, b_2) \notin E$ as $(b'_1, b'_2) \notin E$ and b'_1, b'_2 have as neighbourhoods the union of all neighbourhoods in their respective class. Hence, $\langle c_1, b_1, u, b_2, c_2 \rangle$ is also a diametral path in G . \square

Claim 12.3. *G' does not contain a $P_4 + P_1$ of the form $\langle c'_1, b'_1, u, b'_2 \rangle + c'_2$, with $c'_1, c'_2 \in C'$ and $b'_1, b'_2 \in B'$.*

Proof. Assume to the contrary that there is such a $P_4 + P_1$ in G' . Let it be $\langle c'_1, b'_1, u, b'_2 \rangle + c'_2$, where $b'_1, b'_2 \in B'$ and $c'_1, c'_2 \in C'$. Every component in

$G[C]$ is a twin class with respect to B , and all vertices in B that get identified with the same vertex in B' have identical neighbourhoods with respect to $G[C]$ and $G[C']$. Hence, the identifications from $G[B]$ to $G[B']$ essentially only adjusted adjacencies within $G[B]$ and removed vertices from B . Denote b_1, b_2 two vertices in G that end up identified with b'_1, b'_2 respectively. We immediately have $(c_1, b_1), (b_1, u), (u, b_2) \in E$ and $(c_1, b_2) \notin E$. As $(b'_1, b'_2) \notin E$, we get that $(b_1, b_2) \notin E$, as the identifications take the union of neighbourhoods. So $\langle c_1, b_1, u, b_2 \rangle$ is an induced P_4 in G . We also see that c_2 is disjoint from this P_4 , as $c'_1 \neq c'_2$ and so c_1 and c_2 are in different components, and by the previous arguments, $(c_2, b_1), (c_2, b_2) \notin E$. We get an induced $P_4 + P_1$ in G , a contradiction. \square

Claim 12.4. *For every non-edge $(b'_1, b'_2) \notin E'$ in G' , with $b'_1, b'_2 \in B'$, it holds that b'_1, b'_2 together dominate C' , that is, every $c' \in C'$ is adjacent to b'_1 or b'_2 .*

Proof. Let $(b'_1, b'_2) \notin E'$ be an arbitrary non-edge in G' . As $b'_1 \neq b'_2$, it must be that $N(b'_1) \cap C' \neq N(b'_2) \cap C'$; otherwise, these two vertices would be the same class. C' is an independent set by construction.

Because b'_1 and b'_2 have unequal neighbourhoods towards C' , at least one of the two has a neighbour in C' which is not a neighbour of the other. Without loss of generality, let this be b'_1 . So, b'_1 has a neighbour $c'_1 \in C'$ that is not a neighbour of b'_2 . Because $(b'_1, b'_2) \notin E$ and $(b'_2, c'_1) \notin E$, we have that $\langle c'_1, b'_1, u, b'_2 \rangle$ is an induced P_4 in G' . Hence, every other $c' \in C'$ must be adjacent to some vertex of the P_4 ; otherwise, this c' forms an induced $P_4 + P_1$ which cannot be present in G' by Claim 12.3. The neighbour of any c' cannot be c'_1 , as C' is an independent set. The neighbour of any c' cannot be u , because of the definition of C' . So every other $c' \in C'$ is adjacent to b'_1 or b'_2 . \square

By Claim 12.2 we can look for the diametral path of type **(3c)** in G' to determine its presence in G . To this end, call a non-edge $(b_1, b_2) \notin E'$ in $G'[B']$ ‘good’ if a diametral path of type **(3c)** exists with b_1, b_2 as its vertices in B' . We analyse the structure of $G'[B']$ assuming a good non-edge exists.

Claim 12.5. *In G' , if a good non-edge $(b_1, b_2) \notin E$ exists, then (a) no non-edge triangle is contained in $G'[B']$ as an induced subgraph, (b) every other vertex $b_3 \in B'$ is adjacent to at least one of b_1, b_2 , and (c) if b_1, b_2 have a common neighbour $x \in B'$, then x is complete to $B' \setminus \{x\}$.*

Proof. Let $(b_1, b_2) \notin E'$ be a good non-edge, for $b_1, b_2 \in B'$.

To prove (a), assume there is a non-edge triangle formed by the vertices $a_1, a_2, a_3 \in B'$. Apply Claim 12.4 to all three of the non-edges in this non-edge triangle to see that every vertex in C is adjacent to at least two of a_1, a_2, a_3 . But then (b_1, b_2) is not good, as any two $c_1, c_2 \in C'$ have a common neighbour among a_1, a_2, a_3 . This is a contradiction with the assumption that (b_1, b_2) is good.

To prove (b), assume there is another vertex $b_3 \in B'$. If b_3 is not a neighbour of both b_1 and b_2 , we immediately have a non-edge triangle formed by the non-edges $(b_1, b_2), (b_1, b_3), (b_2, b_3) \notin E'$, a contradiction.

To prove (c), assume b_1, b_2 have a common neighbour $x \in B'$. Assume for sake of contradiction that there is another vertex $b_3 \in B'$ not adjacent to x . By (b), b_3 is adjacent to at least one of b_1, b_2 .

If b_3 is adjacent to only one of b_1, b_2 , say $(b_1, b_3) \in E'$, then every vertex in $C' \setminus N(b_1)$, i.e. the non-neighbours of b_1 in C' , must be adjacent to b_2 by Claim 12.4. Also, by applying Claim 12.4 to $(x, b_3) \notin E'$ we see that every vertex of $C' \setminus N(b_1)$ is also adjacent to either x or b_3 . But then b_1 has distance at most 2 to every $c \in C'$, so (b_1, b_2) is not good, a contradiction.

If instead b_3 is adjacent to b_1, b_2 but not x , then, similar to before, applying Claim 12.4 to the non-edge (x, b_3) gives us that b_1 has distance at most 2 to every $c \in C'$. This is a contradiction with the assumption that (b_1, b_2) is good. We get that b_3 must be adjacent to x . This holds for every $b_3 \in B'$, so x is complete to $B' \setminus \{x\}$. \square

By Claim 12.5, $G'[B']$ admits very strong structural properties if a good non-edge exists. Assume a good non-edge $(b_1, b_2) \notin E'$ exists in $G'[B']$, for $b_1, b_2 \in B'$. Denote by $X \subseteq B'$ the set of vertices of B' complete to B' , and note that this is the same X as the X that appears in the algorithm. We analyse the structure of G' separately for $X \neq \emptyset$ and $X = \emptyset$.

Assume $X \neq \emptyset$. Denote by $B_1 \subseteq B'$ ($B_2 \subseteq B'$) the subset of B' adjacent to b_1, x (b_2, x) but not b_2 (b_1), including b_1 (b_2). Note that, by Claim 12.5(b) and (c), X, B_1, B_2 partition B' . If there is an edge anywhere between B_1 and B_2 , then (b_1, b_2) is not good, as by Claim 12.4 any non-neighbour of b_2 in C' is adjacent to every $b \in B_1$, and any non-neighbour of b_1 in C' is adjacent to every $b \in B_2$. Any non-edge within $G'[B_1]$ or $G'[B_2]$ creates a non-edge triangle with b_2 or b_1 , respectively, so then (b_1, b_2) is not good by Claim 12.5(a). So, if we have a good non-edge (b_1, b_2) , then B' must be a partition of three cliques B_1, B_2, X , where X is complete to $B' \setminus X$ and B_1, B_2 have no edges between them. If this partition exists, then indeed a diametral path of type **(3c)** exists if and only if there exist $c_1, c_2 \in C$ with $N(c_1) = B_1, N(c_2) = B_2$, as the distance between any $b_1 \in B_1$ and $b_2 \in B_2$ is two, and so c_1 to c_2 must have distance 4. The algorithm looks for exactly these sets B_1, B_2, X as described and checks the existence of $c_1, c_2 \in C$ with $N(c_1) = B_1, N(c_2) = B_2$, which is now proven to be correct.

Assume $X = \emptyset$. By Claim 12.5(c), b_1, b_2 have no common neighbour in B' . We get that we can partition B' into $B_1 \cup B_2$, where B_1 consists of the vertices adjacent to b_1 and b_1 itself, and B_2 those adjacent to b_2 and b_2 itself, by Claim 12.5(b). Any non-edge within $G'[B_1]$ or $G'[B_2]$ forms a non-edge triangle with b_2 or b_1 , but then by Claim 12.5(a) (b_1, b_2) is not good. So, B_1 and B_2 must be cliques. If there is an edge anywhere between B_1 and B_2 , then (b_1, b_2) is not good, as by Claim 12.4 any non-neighbour of b_2 in C' is adjacent to every $b \in B_1$, and any non-neighbour of b_1 in C' is adjacent to every $b \in B_2$. We get that B' consists of two disjoint and anti-complete cliques. Now a diametral path of type **(3c)** exists if and only if there exist vertices $c_1, c_2 \in C$ with $N(c_1) = B_1, N(c_2) = B_2$. The algorithm looks for exactly these sets B_1, B_2 as described, if $X = \emptyset$, and checks the existence of $c_1, c_2 \in C$ with $N(c_1) = B_1, N(c_2) = B_2$, which is now proven to be correct.

We conclude that **(3c.1.algorithm)** only returns that there is no diametral path of length 4 when there does not exist a diametral path of type **(3c)** in G .

(3c.2) There exists a $b \in B$ that has an edge and a non-edge to one component of $G[C]$. Let b_1 be such a vertex, and denote C_1 as the component of $G[C]$ to which b_1 has both an edge and a non-edge. Notice that one end of the diametral

path of type **(3c)** must be in C_1 , as the distance between vertices in other connected components of $G[C]$ is at most 2 through b_1 . The other end cannot also be in C_1 , as $G[C_1]$ is a connected P_4 -free graph, and so has diameter at most 2. All vertices in $N(b_1) \cap C_1$ have distance at most 2 to all $c \in C$. If all vertices of C_1 are at distance at most 2 to b_1 , then a diametral path of type **(3c)** does not exist. So we can assume there are vertices $c_3, c_2, c_1 \in C_1$ with $(c_3, c_2), (c_2, c_1), (c_1, b_1) \in E$ and $(c_3, b_1), (c_3, c_1), (c_2, b_1) \notin E$. If a diametral path of type **(3c)** exists, one of its endpoints must be c_3 or another vertex in C_1 at distance 2 to $N(b_1) \cap C_1$. But then a diametral path of the form $\langle c_3, b_3, u, b_2, c_4 \rangle$, for some $b_3, b_2 \in B, c_3 \in C_1, c_4 \in C$ implies that $\langle c_3, c_2, c_1, b_1, c_4 \rangle$ is also a diametral path of type **(3b)**, where $c_2, c_1 \in C_1$ are the vertices on the path from c_3 to b_1 in C_1 . The algorithm for type **(3b)** finds a diametral path of length 4 or concludes that no diametral path of type **(3b)** exists in G , which also rules out that a diametral path of type **(3c)** exists in G .

(3d) $\langle c_1, b_1, c_2, b_2, c_3 \rangle$ As b_1 and b_2 are both adjacent to u , if a diametral path of type **(3d)** exists, $\langle c_1, b_1, u, b_2, c_2 \rangle$ is also a shortest path of distance 4 from c_1 to c_2 which is of type **(3c)**. The algorithm for type **(3c)** finds a diametral path of length 4 or concludes that no diametral path of type **(3c)** exists in G , which also rules out that a diametral path of type **(3d)** exists in G .

(3e) $\langle c_1, c_2, b_1, b_2, c_3 \rangle$ Notice that in a diametral path of type **(3e)**, c_3 has no edges to either c_1, c_2, b_1 , and c_1 is not adjacent to b_1 ; otherwise, the distance from c_1 to c_3 is not 4. Now $\langle c_1, c_2, b_1, u \rangle$ and c_3 form a $P_4 + P_1$. So, this case cannot occur in a $(P_4 + P_1)$ -free graph.

(3f) $\langle c_1, b_1, b_2, b_3, c_2 \rangle$ As b_1 and b_3 are both adjacent to u , if a diametral path of type **(3f)** exists in G , $\langle c_1, b_1, u, b_3, c_2 \rangle$ is also a shortest path of distance 4 from c_1 to c_2 which is of type **(3c)**. The algorithm for type **(3c)** finds a diametral path of length 4 or concludes that no diametral path of type **(3c)** exists in G , which also rules out that a diametral path of type **(3f)** exists in G .

We have now shown that, for each possible structure of the diametral path with respect to u, B, C , we find no diametral path of length 4 only when that structure does not exist in G . So, we have shown that in time $O(n + m)$ we find a distance-4 shortest path if and only if it exists in G . \square

4.4 $(P_3 + 2P_1)$ -free graphs

Theorem 13. *Given a $(P_3 + 2P_1)$ -free graph G , we can decide whether the diameter of G is equal to $d_{\max} = 5$ in $O(n + m)$ time.*

Proof. Let $G = (V, E)$ be a (connected) $(P_3 + 2P_1)$ -free graph. Indeed, $d_{\max} = 5$ by Theorem 5. We start with a BFS from an arbitrary vertex u , and let v be a vertex that is in the second neighbourhood of u . If such a vertex v does not exist, the diameter of G is not 5. We now distinguish the structure of the graph with respect to u and v . This structure can be identified by BFS from both u and v , see also Figure 6 for an illustration. Let $A \subset V$ be the set of all common neighbours of u and v . Note that $A \neq \emptyset$ by the choice of v . Set $B_1 = N(u) \setminus A$ and $B_2 = N(v) \setminus A$. For now, let $C' \subset V$ be the set of all other vertices in the




Figure 6: An illustration of the structure of a $(P_3 + 2P_1)$ -free graph as in the proof of Theorem 13.

graph. The distances between vertices in $V \setminus C'$ are at most 4 due to u and v . So, if C' is empty, we are done. Otherwise, any diametral path of length 5 has an endpoint in C' , so we focus on this. Note that $G[C']$ is P_3 -free, a disjoint union of cliques, as it is non-adjacent to both u and v . For any $a \in A$, the non-neighbours of a in C' must form a complete subgraph, as $\langle u, a, v \rangle$ is a P_3 and the graph is $(P_3 + 2P_1)$ -free. We see that any $a \in A$ is non-adjacent to at most one clique or a part of one clique in C' . If every clique in C' has a vertex with a neighbour in A , then the diameter is not 5: any $c' \in C'$ has distance at most 2 to some $a \in A$, which immediately implies it has distance at most 4 to all vertices of G . So, for the diameter to be 5, there is a clique of vertices $D \subseteq C'$ that is non-adjacent to all of A . Define C as $C = C' \setminus D$. Every vertex $a \in A$ is complete to C by construction. We can identify whether such a clique D exists and find all vertices in D if it does in linear time, by inspection of the adjacency lists of all vertices in A . If the distance from u or v to some vertex is 5, the initial BFS would identify such a diametral path. Vertices in C have distance at most 4 to all vertices in $V \setminus D$.

We can conclude that one end of a diametral path of length 5 has to be in D , if such a path exists. The other end is either a vertex in C or a vertex in $B_1 \cup B_2$.

For a $d \in D$ to have distance 5 to some $c \in C$, it must be that $N(d) \subseteq D$, as u and v have distance 2 to all $c \in C$. Note that all $d \in D$ with $N(d) \subseteq D$ are true twins, as D is a clique. So a single BFS from the lowest degree vertex in D will identify a diametral path of length 5 from D to C , if it exists.

In the following, assume the distance from any $d \in D$ to all $c \in C$ is at most 4. If there are $d_1, d_2 \in D$ such that $N(d_1) \cap B_1 \neq \emptyset$ and $N(d_2) \cap B_2 \neq \emptyset$, then the distance from any $d \in D$ to any other vertex in the graph is at most 4, as the distance from any $d \in D$ is at most 3 to both u and v . We can check in linear time whether D has some edge to B_1 and/or some edge to B_2 , and terminate when there is an edge for both. Without loss of generality, assume that vertices in D are not adjacent to B_1 . Now a diametral path of length 5 can only go from D to B_1 , if the diameter is 5.

Then, we must have $B_1, B_2 \neq \emptyset$. If some $c \in C$ has a neighbour $b_1 \in B_1$, then $\langle u, b_1, c \rangle + v + d$ forms a $P_3 + 2P_1$, where d is any vertex in D . So, either C is empty, or no vertex in C has a neighbour in B_1 . Regardless, we do not




Figure 7: An illustration of the structure of a $(P_2 + 3P_1)$ -free graph as in the proof of Theorem 15.

have to consider finding any shortest path from a vertex in D to any vertex in B_1 that goes through C , as using v instead is equivalent. Let $B'_2 \subseteq B_2$ be the set of vertices in B_2 that each have at least one neighbour in A or B_1 . Let $D' \subseteq D$ be the set of vertices in D with at least one neighbour in B'_2 . Note that we can find B'_2 and D' by inspection of the adjacency lists of all vertices in B_2 and then of all vertices in D . Vertices in D' have distance 3 to u , and so distance at most 4 to all of B_1 . All vertices in $D \setminus D'$ have distance at least 4 to u . If every $b_1 \in B_1$ is complete to A , then any shortest path of length 5 from a vertex in D to a vertex in B_1 would need its endpoint $d \in D$ to have $N(d) \subseteq D$. All such vertices are true twins, and a BFS from a single vertex in D with the lowest degree identifies such a diametral path, if it exists. Otherwise, there exist $b_1 \in B_1$, $a \in A$ with $(b_1, a) \notin E$. But then $G[(B_2 \setminus B'_2) \cup (D \setminus D')]$ must be both P_3 -free and $2P_1$ -free, as b_1 and a form a $2P_1$ but also a P_3 with u . So, $G[(B_2 \setminus B'_2) \cup (D \setminus D')]$ is a clique. But then all $d \in D \setminus D'$ are true twins. Picking one and executing a BFS suffices to find a length-5 diametral path, if it exists. Note that we found the sets B'_2, D' earlier, so the set $D \setminus D'$ is known.

The above analysis shows that we can find a certificate for a shortest path of length 5 if and only if it exists in G . \square

As an immediate corollary, we get that we can decide on d_{\max} for the class of $4P_1$ -free graphs, as any $4P_1$ -free graph is $(P_3 + 2P_1)$ -free, and d_{\max} is equal for both classes.

Corollary 14. *Given a $4P_1$ -free graph G , we can decide whether the diameter of G is equal to $d_{\max} = 5$ in $O(n + m)$ time.*

4.5 $(P_2 + 3P_1)$ -free graphs

Theorem 15. *Given a $(P_2 + 3P_1)$ -free graph G , we can decide whether the diameter of G is equal to $d_{\max} = 6$ in $O(n + m)$ time.*

Proof. Let $G = (V, E)$ be a (connected) $(P_2 + 3P_1)$ -free graph. Indeed, $d_{\max} = 6$ by Theorem 5. We start with a BFS from an arbitrary vertex u , and let v be a vertex that is in the second neighbourhood of u . If such a vertex v does not exist, the diameter of G is not 6. We now distinguish the structure of the graph with respect to u and v . This structure can be identified by BFS from both u and v , see also Figure 7 for an illustration. Let $A \subset V$ be the set of common neighbours of u and v . Note that $A \neq \emptyset$ by the choice of v . Set $B_1 = N(u) \setminus A$ and $B_2 = N(v) \setminus A$. Let $C \subset V$ be the set of all other vertices in the graph. If C is empty, the diameter of G cannot be 6. As $u + v$ is a $2P_1$, $G[C]$ must be $(P_2 + P_1)$ -free, a complete r -partite graph for some integer $r \geq 1$.

Claim 15.1. *The non-neighbours in C of a vertex $b \in B_1 \cup B_2$ form a complete subgraph.*

Proof. Say $b \in B_1$. Then $\langle b, u \rangle + v$ is a $P_2 + P_1$, so the induced subgraph of vertices non-adjacent to all of $\{b, u, v\}$ is $2P_1$ -free. This is the set of vertices in C non-adjacent to b . The case for some $b \in B_2$ is analogous. \square

If the distance from u or v to some vertex is 6, then the initial BFS would identify such a diametral path. Otherwise, at least one endpoint of a length-6 diametral path must be in C . We distinguish algorithmically if $r = 1$ or $r > 1$ by checking whether $G[C]$ contains an edge in $O(n + m)$ time.

If $G[C]$ is 1-partite, an independent set, then every $c \in C$ has a neighbour in $B_1 \cup A \cup B_2$ due to G being a connected graph. Because of this, the distance from any $c \in C$ to any vertex in $B_1 \cup A \cup B_2 \cup \{u, v\}$ is at most 5. So, a diametral path of length 6 then must have both endpoints in C , if it exists. However, by Claim 15.1, the non-neighbours in $G[C]$ of any $b_1 \in B_1$ or $b_2 \in B_2$ form a clique, they must be a single vertex. So, after inspecting the neighbourhood of a single $b_1 \in B_1$ there is at most single vertex $c_1 \in C$ which is a non-neighbour of b_1 . But then the distances between all vertices in $C \setminus \{c_1\}$ are at most 2, and c_1 must be an endpoint of a length-6 diametral path, if it exists. We can find c_1 in time $O(n + m)$ and execute a BFS from it and return a distance-6 shortest path if it is found.

If $G[C]$ is not an independent set, $G[C]$ is a complete r -partite graph for $r > 1$ and the distance between vertices in C is at most 2. The distance from some $a \in A$ to all vertices of G is at most 5, as it has distance 2 to all vertices in $B_1 \cup B_2$. Then, a diametral path of length 6 must have one endpoint in C and one endpoint in either B_1 or B_2 . Without loss of generality, assume that one end of the diametral pair is in B_1 , the other case is symmetric.

Suppose that $G[C]$ is not a clique, and recall that $G[C]$ is a complete r -partite graph. Because the non-neighbours of any $b_1 \in B_1$ in C must form a clique by Claim 15.1, every $b_1 \in B_1$ has a neighbour in C . Hence, the distance from any b_1 to any $c \in C$ is at most 3. Hence, the diameter of G is at most 5. So, if there is a length-6 diametral path from some $c \in C$ to some $b_1 \in B_1$, $G[C]$ is a clique, and there are no edges between C and $B_1 \cup A$. This can be checked in $O(n + m)$ time. Also, the endpoint of the diametral path in C , say $c \in C$, must have $N(c) \subseteq C$, as otherwise it has distance at most 5 to all other vertices. All such vertices are true twins, as $G[C]$ is a clique. So we can execute a BFS from any one of these vertices to find a length-6 shortest path with an endpoint in C and one in B_1 , if it exists. This is simply the lowest degree vertex in C , so finding it and executing a BFS from it takes $O(n + m)$ time.




Figure 8: An illustration of the structure of a $(2P_2 + P_1)$ -free graph as in the proof of Theorem 16.

The above analysis shows that we can find a certificate for a shortest path of length 6 in $O(n + m)$ time if and only if it exists in G . \square

4.6 $(2P_2 + P_1)$ -free graphs

Theorem 16. *Given a $(2P_2 + P_1)$ -free graph G , we can decide whether the diameter of G is equal to $d_{\max} = 5$ in $O(n + m)$ time.*

Proof. Let $G = (V, E)$ be a (connected) $(2P_2 + P_1)$ -free graph. Indeed, $d_{\max} = 5$ by Theorem 5. We start with a BFS from an arbitrary vertex u , and let v be a vertex that is in the second neighbourhood of u , with $N(v) \neq N(u)$. If such a vertex v does not exist, the diameter of G is not 5, but at most 2. Such a vertex v can be identified in time $O(n + m)$. We now distinguish the structure of the graph with respect to u and v . This structure can be identified by BFS from both u and v ; see Figure 8 for an illustration. Let $A \subset V$ be the set of common neighbours of u and v . Note that $A \neq \emptyset$ by the choice of v . Set $B_1 = N(u) \setminus A$ and $B_2 = N(v) \setminus A$. Let $C \subset V$ be the set of all other vertices in the graph.

The algorithm for this problem is given by the following steps. Search for a vertex $c_1 \in C$ with $N(c_1) = B_1$, a vertex $c_2 \in C$ with $N(c_2) = B_2$, and a vertex $c_3 \in C$ with $N(c_3) = A$. For each $i \in \{1, 2, 3\}$, if it exists, execute a BFS from c_i , returning a distance-5 shortest path when found. If none of the options exist or none lead to a distance-5 shortest path, return that the diameter of G is not 5. Notice that this can be executed in $O(n + m)$ time, as we can identify the vertices in $O(n + m)$ time, and we do a constant number of BFS calls. Next, we prove correctness of this algorithm.

We first prove two claims that will be useful in analysing the structure of G .

Claim 16.1. *For all $b_1 \in B_1$, $b_2 \in B_2$, if $(b_1, b_2) \notin E$ then b_1, b_2 together dominate all vertices in C .*

Proof. Let $b_1 \in B_1$, $b_2 \in B_2$ such that $(b_1, b_2) \notin E$. Assume to the contrary there is a vertex $c \in C$ that is not adjacent to either b_1 or b_2 . By construction, c is not adjacent to u or v , and $(u, v) \notin E$. But then we have that $\langle u, b_1 \rangle + \langle v, b_2 \rangle + c$ is an induced $2P_2 + P_1$ in the graph, a contradiction. \square

Claim 16.2. For any edge (c_1, c_2) in $G[C]$, $c_1, c_2 \in C$, for every $b \in B_1 \cup B_2$, it holds that $(b, c_1) \in E$ or $(b, c_2) \in E$.

Proof. Let (c_1, c_2) be an edge in $G[C]$, $c_1, c_2 \in C$. Assume to the contrary that there exist a $b \in B_1 \cup B_2$ with $(b, c_1) \notin E$ and $(b, c_2) \notin E$. Let us assume $b \in B_1$, the other case is symmetric. By construction, c_1, c_2 are not adjacent to u or v , and $(u, v) \notin E$. But then we have that $\langle u, b \rangle + \langle c_1, c_2 \rangle + v$ is an induced $2P_2 + P_1$ in the graph, a contradiction. \square

Since u, v dominate $G[V \setminus C]$ and have a common neighbour, any length-5 shortest path must have at least one endpoint in C , as the distances in $G[V \setminus C]$ are at most 4. If the distance from u or v to some vertex is 5, then we would know by the initial BFS.

Case 1: There exist $b_1 \in B_1, b_2 \in B_2$ with $(b_1, b_2) \notin E$.

Let $C' = N(b_1) \cap N(b_2)$ and let $C_1 = (N(b_1) \cap C) \setminus C'$ and $C_2 = (N(b_2) \cap C) \setminus C'$. Note that C_1, C', C_2 together partition C by Claim 16.1.

We first show that we can find a diametral path of length 5 with one endpoint in B_1 or B_2 and the other in C_1 or C_2 , respectively, if it exists. These two scenarios are symmetric, so assume without loss of generality that a diametral path exists with endpoints $c \in C_1$ and $b_3 \in B_2$. Because the distance from c to b_3 is 5, b_3 is not adjacent to any vertex of B_1 , or any vertex of $C_1 \cup C'$. So, by applying Claim 16.1 with b_1 and b_3 we see that $N(b_3) \cap C = C_2$ as $C_2 \cap N(b_1) = \emptyset$. Furthermore, for any $b'_1 \in B_1$ we get $N(b'_1) \cap C \supseteq C_1 \cup C'$ by applying Claim 16.1 with respect to b'_1 and b_3 . Also, c has no neighbours in A, C', C_2 , or B_2 , as the distance to b_3 is 5. Consider some edge e in $G[C_1]$. By applying Claim 16.2 to e , we get that one of the vertices of e is adjacent to b_3 . We see that c has degree 0 in $G[C_1]$. To sum up, $N(c) = B_1$, and this holds for any $c \in C$ with a vertex in B_2 at distance 5. Hence, all these candidates are twins. We find that detection of a single $c \in C$ with $N(c) = B_1$ and executing a BFS from it indeed suffices to find a length-5 diametral path from a vertex in C_1 to a vertex in B_2 , if it exists. For the symmetric case, this argument shows that it suffices to find a vertex $c \in C$ with $N(c) = B_2$ and execute a BFS from it to find a length-5 diametral path from a vertex in C_2 to a vertex in B_1 , if it exists.

Case 1a: $C' \neq \emptyset$.

In this case, we see that the distances between vertices of C are at most 4. Hence, any diametral path of length 5 has one endpoint in $C_1 \cup C_2$ and one in $B_1 \cup B_2$, if it exists, as the distance from any $c \in C$ to any $a \in A$ is at most 3 via b_1 or b_2 , and C' has distance at most 3 to all vertices. So, no additional algorithmic steps are necessary.

Case 1b: $C' = \emptyset$.

In this case, it may be possible the distance from some $c_1 \in C_1$ to some $c_2 \in C_2$ is 5. Assume the distance from $c_1 \in C_1$ to $c_2 \in C_2$ is 5. Let it be clear that c_1 has no neighbour in $\{u, v\} \cup A \cup B_2 \cup C_2$ and similarly c_2 has no neighbour in $\{u, v\} \cup A \cup B_1 \cup C_1$, as the distance between c_1 and c_2 would not be 5. But then for any $a \in A$ it must be that $(a, b_1), (a, b_2) \in E$, as both $c_1 + \langle u, a \rangle + \langle b_2, c_2 \rangle$ is an induced $2P_2 + P_1$ and $c_2 + \langle v, a \rangle + \langle b_1, c_1 \rangle$ is an induced $2P_2 + P_1$. This is a contradiction with the assumption that the distance from c_1 to c_2 is 5. We conclude that the distance from any $c_1 \in C_1$ to any $c_2 \in C_2$ cannot be 5, and

that the only length-5 diametral path that can exist has one endpoint in B_1 or B_2 and the other in C_1 or C_2 , respectively.

Case 2: There do not exist $b_1 \in B_1, b_2 \in B_2$ with $(b_1, b_2) \notin E$.

We get that B_1 and B_2 are complete to each other, or one of B_1 and B_2 is empty. By choice of v , it cannot be that both B_1 and B_2 are empty sets. We get that vertices in $B_1 \cup B_2$ have distance at most 3 to any other vertex in the graph, as by Claim 16.2 vertices incident to an edge in C are at distance at most 2 from any $b \in B_1 \cup B_2$, and vertices not incident an edge in C are adjacent to at least one vertex in $B_1 \cup A \cup B_2$. Using Claim 16.2 we also see that the distance from any $c \in C$ is at most 4 to anything not in C . Hence, the only option for a distance-5 pair, is starting from a $c_1 \in C$ with an edge in $G[C]$, and ending in a $c_2 \in C$ with $N(c_2) \subseteq A$. Assuming such a pair exists, let the path be $\langle c_1, c'_1, b_1, u, a, c_2 \rangle$ for some $c_1, c'_1, c_2 \in C, b_1 \in B_1, a \in A$. The case for B_2 is symmetric. Say some $a' \in A$ is not a neighbour of c_2 , then $c_2 + \langle v, a' \rangle + \langle c_1, c'_1 \rangle$ is an induced $2P_2 + P_1$, a contradiction. We get that $N(c_2) = A$. Hence, all candidates for c_2 are false twins, and indeed inspecting a single $c \in C$ with $N(c) = A$ and execute a BFS from it suffices to conclude whether a length-5 diametral path of this case exists.

To conclude, we find a length-5 diametral path if and only if it exists in G . \square

4.7 Proofs of Theorem 2 and 3

We now prove our algorithmic theorems, Theorem 2 and Theorem 3.

Theorem 2 (Restated). *Given a graph G that is H -free, we can solve DIAMETER in $O(n + m)$ time, when one of $H \subseteq P_2 + 2P_1, H \subseteq P_3 + P_1$ holds.*

Proof. Theorem 9 shows that we can compute the diameter of an H -free graph in linear time when $H \subseteq P_2 + 2P_1$, and Theorem 11 shows that we can compute the diameter of an H -free graph in linear time when $H \subseteq P_3 + P_1$. \square

Theorem 3 (Restated). *Given a graph G from a class of graphs \mathcal{G} that is H -free, we can decide whether $\text{diam}(G) = d_{\max}(\mathcal{G})$ in $O(n + m)$ time, when one of $H \subseteq P_2 + 2P_1, H \subseteq P_4 + P_1, H \subseteq P_3 + 2P_1, H \subseteq P_2 + 3P_1$, or $H = 2P_2 + P_1$ holds.*

Proof. First note that computing the diameter for any graph class \mathcal{G} with $d_{\max}(\mathcal{G}) \leq 2$ can be done in linear time, as we can check whether the graph is a clique in $O(n + m)$ time. This covers the cases of $H = P_4, P_3, P_2, P_1, 2P_1$, and $P_2 + P_1$. The case of $H = 3P_1$ is shown by Corollary 10 and $H = 4P_1$ is shown by Corollary 14. The case of $H = P_2 + 2P_1$ is given by Theorem 9. $P_2 + 3P_1$ is shown by Theorem 15. The case of $H = 2P_2 + P_1$ is given by Theorem 16. $H = P_3 + P_1$ is given by Theorem 11. $H = P_3 + 2P_1$ is given by Theorem 13. When $H = P_4 + P_1$ we get a linear-time algorithm by Theorem 12. This completes the proof of the theorem. \square

5 Conclusion and Discussion

We analysed the complexity of computing the diameter of H -free graphs. For several H -free graph classes \mathcal{G} where H is a linear forest, we found linear time algorithms for deciding whether $\text{diam}(G) = d_{\max}(\mathcal{G})$ holds. For some such graph classes, hardness results may exist for computing entirely the diameter (note the particular case of $H = 2P_2 + P_1$). However, for $H = 2P_2$ and $H = P_t$ for odd $t \geq 5$, we show that any such a linear-time algorithm would refute SETH. We conjecture that this pattern on P_t -free graphs extends to all $t \geq 5$, as it seems a mere technical limitation that Theorem 4 does not hold for even t , not a barrier for hardness.

Conjecture 17. *Under SETH, if \mathcal{G} is the class of P_t -free graphs with $t \geq 5$, we cannot decide whether the diameter of a graph $G \in \mathcal{G}$ is equal to $d_{\max}(\mathcal{G})$ in time $O(n^{2-\epsilon})$, for all $\epsilon > 0$.*

The hardness implied by Conjecture 17 is in stark contrast to our positive algorithmic results, summarized by Theorem 3. These results suggest that indeed deciding whether the diameter of a graph is equal to $d_{\max}(\mathcal{G})$ may be easier than deciding on the diameter of an H -free graph in general. In an attempt to reveal the underlying pattern, we conjecture our algorithmic results generalize to broad classes of H -free graphs.

Conjecture 18. *For a row of integers $(a_i)_{i=1}^k$, $k \geq 2$, $(a_i)_{i=1}^k \neq (2, 2)$, let \mathcal{G} be the class of $\sum_{i=1}^k P_{a_i}$ -free graphs. Given a graph $G \in \mathcal{G}$, we can decide whether the diameter of G is equal to $d_{\max}(\mathcal{G}) = k - 3 + \sum_{i=1}^k a_i$ in $O(n + m)$ time.*

Conjecture 18 rules out $k = 1$ as these cases are hard under Theorem 4 and Conjecture 17. It also rules out the case of $2P_2$ -free graphs, for which we showed that deciding whether the diameter is equal to $d_{\max} = 3$ is hard under SETH (see Theorem 4).

Because of the generic nature of the statement in Conjecture 18, it may very well turn out the statement is false. We give a more specific version of Conjecture 18, which one may find more plausible. In particular, this version of the conjecture lays emphasis on linear forests H including some set of isolated vertices, as it seems that in our algorithms, the presence of a P_1 in H helps out in structural analysis.

Conjecture 19. *Let \mathcal{G} be the class of $rP_t + sP_1$ -free graphs, where $r, s, t \geq 1$. Given a graph $G \in \mathcal{G}$, we can decide whether the diameter of G is equal to $d_{\max}(\mathcal{G}) = r(t + 1) + 2s - 3$ in $O(n + m)$ time.*

Conjecture 19 demands $s \geq 1$, as $s = 0$ would contradict Theorem 4. $t \geq 1$ must hold for the correctness of the statement of d_{\max} .

Note that none of our results distinguish between Conjecture 18 and Conjecture 19. That is, all our results support the more specific Conjecture 19, which in turn supports the more general conjecture. To the best of our knowledge, there is no evidence against either conjecture. A linear-time algorithm for the only open case with $d_{\max} = 4$, which is for $H = P_3 + P_2$, would support Conjecture 18 but not Conjecture 19, and would suggest Conjecture 19 is not the full truth.

The main open problem posed by our work is then whether the Conjectures 17, 18, and/or 19 hold true. The smallest open cases for algorithmic results to support Conjectures 18 and 19 would be the class of H -free graphs with one of $H = 5P_1$ ($d_{\max} = 7$), $H = 2P_2 + 2P_1$ ($d_{\max} = 7$), $H = P_3 + P_2$ ($d_{\max} = 4$), $H = P_4 + 2P_1$ ($d_{\max} = 6$), or $H = P_5 + P_1$ ($d_{\max} = 5$). The smallest open case for a hardness result to support Conjecture 17 is the class of P_6 -free graphs. Conversely, an algorithmic result for P_6 -free graphs refutes Conjecture 17, and hardness results for the other classes above refute Conjecture 18 or Conjecture 19.

Further open questions are revealed by our results, or in particular, what are results do not show. For instance, for $(2P_2 + P_1)$ -free graphs, we know that we can decide in linear time whether the diameter is equal to 5, and cannot decide in truly subquadratic time whether the diameter is 2 or 3. So, settling the following open question would form a complete dichotomy for $(2P_2 + P_1)$ -free graphs:

Open Question 1. *Given a $(2P_2 + P_1)$ -free graph G , can we decide in linear time whether the diameter of G is equal to 4? Or, conversely, can it be shown that such an algorithm cannot exist (under SETH)?*

Theorem 2 and 3 make progress for DIAMETER computation, but do not settle its complexity completely, so we ask:

Open Question 2. *What is the complexity of the DIAMETER problem on connected H -free graphs, in cases where H is a linear forest that does not contain a $2P_2$?*

References

- [1] Amir Abboud, Fabrizio Grandoni, and Virginia Vassilevska Williams. Subcubic equivalences between graph centrality problems, APSP, and diameter. *ACM Trans. Algorithms*, 19(1):3:1–3:30, 2023.
- [2] Amir Abboud, Shay Mozes, and Oren Weimann. What else can Voronoi diagrams do for diameter in planar graphs? In *Proc. ESA 2023*, volume 274 of *LIPICs*, pages 4:1–4:20. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2023.
- [3] Amir Abboud, Richard Ryan Williams, and Huacheng Yu. More applications of the polynomial method to algorithm design. In *Proc. SODA 2015*, pages 218–230. SIAM, 2015.
- [4] Amir Abboud, Virginia Vassilevska Williams, and Joshua R. Wang. Approximation and fixed parameter subquadratic algorithms for radius and diameter in sparse graphs. In *Proc. SODA 2016*, pages 377–391. SIAM, 2016.
- [5] Arturs Backurs, Liam Roditty, Gilad Segal, Virginia Vassilevska Williams, and Nicole Wein. Towards tight approximation bounds for graph diameter and eccentricities. In *Proc. STOC 2018*, pages 267–280. ACM, 2018.
- [6] Andreas Brandstädt, Victor Chepoi, and Feodor F. Dragan. The algorithmic use of hypertree structure and maximum neighbourhood orderings. *Discret. Appl. Math.*, 82(1-3):43–77, 1998.
- [7] Karl Bringmann, Thore Husfeldt, and Måns Magnusson. Multivariate analysis of orthogonal range searching and graph distances. *Algorithmica*, 82(8):2292–2315, 2020.
- [8] Sergio Cabello. Subquadratic algorithms for the diameter and the sum of pairwise distances in planar graphs. *ACM Trans. Algorithms*, 15(2):21:1–21:38, 2019.
- [9] Timothy M. Chan and R. Ryan Williams. Deterministic APSP, orthogonal vectors, and more: Quickly derandomizing razborov-smolensky. *ACM Trans. Algorithms*, 17(1):2:1–2:14, 2021.
- [10] Shiri Chechik, Daniel H. Larkin, Liam Roditty, Grant Schoenebeck, Robert Endre Tarjan, and Virginia Vassilevska Williams. Better approximation algorithms for the graph diameter. In *Proc. SODA 2014*, pages 1041–1052. SIAM, 2014.
- [11] Victor Chepoi, Feodor F. Dragan, Bertrand Estellon, Michel Habib, and Yann Vaxès. Notes on diameters, centers, and approximating trees of delta-hyperbolic geodesic spaces and graphs. *Electron. Notes Discret. Math.*, 31:231–234, 2008.
- [12] Victor Chepoi, Feodor F. Dragan, Michel Habib, Yann Vaxès, and Hend Alrasheed. Fast approximation of eccentricities and distances in hyperbolic graphs. *JGAA*, 23(2):393–433, 2019.

- [13] Derek G. Corneil, Feodor F. Dragan, Michel Habib, and Christophe Paul. Diameter determination on restricted graph families. *Discret. Appl. Math.*, 113(2-3):143–166, 2001.
- [14] Derek G. Corneil, Feodor F. Dragan, and Ekkehard Köhler. On the power of BFS to determine a graph’s diameter. *Networks*, 42(4):209–222, 2003.
- [15] David Coudert, Guillaume Ducoffe, and Alexandru Popa. Fully polynomial FPT algorithms for some classes of bounded clique-width graphs. *ACM Trans. Algorithms*, 15(3):33:1–33:57, 2019.
- [16] Marek Cygan, Harold N. Gabow, and Piotr Sankowski. Algorithmic applications of baur-strassen’s theorem: Shortest cycles, diameter, and matchings. *J. ACM*, 62(4):28:1–28:30, 2015.
- [17] Feodor F. Dragan. Centers of graphs and the helly property. *Ph. D. Thesis, Moldova State University*, 1989.
- [18] Feodor F. Dragan. HT-graphs: centers, connected r-domination and steiner trees. *Comput. Sci. J. Moldova*, 1(2):64–83, 1993.
- [19] Feodor F. Dragan. Dominating cliques in distance-hereditary graphs. In *Proc. SWAT 1994*, volume 824 of *LNCS*, pages 370–381. Springer, 1994.
- [20] Feodor F. Dragan and Guillaume Ducoffe. α_i -metric graphs: Radius, diameter and all eccentricities. In *Proc. WG 2023*, volume 14093 of *LNCS*, pages 276–290. Springer, 2023.
- [21] Feodor F. Dragan, Guillaume Ducoffe, and Heather M. Guarnera. Fast deterministic algorithms for computing all eccentricities in (hyperbolic) Helly graphs. In *Proc. WADS 2021*, volume 12808 of *LNCS*, pages 300–314. Springer, 2021.
- [22] Feodor F. Dragan and Heather M. Guarnera. Eccentricity function in distance-hereditary graphs. *Theor. Comput. Sci.*, 833:26–40, 2020.
- [23] Feodor F. Dragan, Michel Habib, and Laurent Viennot. Revisiting radius, diameter, and all eccentricity computation in graphs through certificates. *CoRR*, abs/1803.04660, 2018.
- [24] Feodor F. Dragan and Falk Nicolai. LexBFS-orderings of distance-hereditary graphs with application to the diametral pair problem. *Discret. Appl. Math.*, 98(3):191–207, 2000.
- [25] Feodor F. Dragan, Falk Nicolai, and Andreas Brandstädt. Lexbfs-orderings and powers of graphs. In *Proc. WG 1996*, pages 166–180. Springer, 1997.
- [26] Ran Duan, Hongxun Wu, and Renfei Zhou. Faster matrix multiplication via asymmetric hashing. In *Proc. FOCS 2023*, pages 2129–2138. IEEE, 2023.
- [27] Guillaume Ducoffe. Beyond Helly graphs: The diameter problem on absolute retracts. In *Proc. WG 2021*, volume 12911 of *LNCS*, pages 321–335. Springer, 2021.

- [28] Guillaume Ducoffe. The diameter of AT-free graphs. *J. Graph Theory*, 99(4):594–614, 2022.
- [29] Guillaume Ducoffe. Optimal centrality computations within bounded clique-width graphs. *Algorithmica*, 84(11):3192–3222, 2022.
- [30] Guillaume Ducoffe. Distance problems within Helly graphs and k -Helly graphs. *Theor. Comput. Sci.*, 946:113690, 2023.
- [31] Guillaume Ducoffe and Feodor F. Dragan. A story of diameter, radius, and (almost) Helly property. *Networks*, 77(3):435–453, 2021.
- [32] Guillaume Ducoffe, Michel Habib, and Laurent Viennot. Diameter computation on H -minor free graphs and graphs of bounded (distance) VC-dimension. In *Proc. SODA 2020*, pages 1905–1922. SIAM, 2020.
- [33] Jacob Evald and Søren Dahlgaard. Tight hardness results for distance and centrality problems in constant degree graphs. *CoRR*, abs/1609.08403, 2016.
- [34] Arthur M. Farley and Andrzej Proskurowski. Computation of the center and diameter of outerplanar graphs. *Discret. Appl. Math.*, 2(3):185–191, 1980.
- [35] Pawel Gawrychowski, Haim Kaplan, Shay Mozes, Micha Sharir, and Oren Weimann. Voronoi diagrams on planar graphs, and computing the diameter in deterministic $\tilde{o}(n^{5/3})$ time. *SIAM J. Comput.*, 50(2):509–554, 2021.
- [36] Michel Habib, Christophe Paul, and Laurent Viennot. A synthesis on partition refinement: A useful routine for strings, graphs, boolean matrices and automata. In *Proc. STACS 1998*, volume 1373 of *LNCS*, pages 25–38. Springer, 1998.
- [37] Peter L Hammer and Stéphane Földes. Split graphs. *Congressus Numerantium*, 19:311–315, 1977.
- [38] Danny Hermelin, Matthias Mnich, Erik Jan van Leeuwen, and Gerhard J. Woeginger. Domination when the stars are out. *ACM Trans. Algorithms*, 15(2):25:1–25:90, 2019.
- [39] Russell Impagliazzo and Ramamohan Paturi. On the complexity of k -SAT. *J. Comput. Syst. Sci.*, 62(2):367–375, 2001.
- [40] Russell Impagliazzo, Ramamohan Paturi, and Francis Zane. Which problems have strongly exponential complexity? *J. Comput. Syst. Sci.*, 63(4):512–530, 2001.
- [41] Matthew Johnson, Barnaby Martin, Jelle J. Oostveen, Sukanya Pandey, Daniël Paulusma, Siani Smith, and Erik Jan van Leeuwen. Complexity framework for forbidden subgraphs. *CoRR*, abs/2211.12887, 2022.
- [42] Stephan Olariu. Paw-fee graphs. *Information Processing Letters*, 28(1):53–54, 1988.

- [43] Stephan Olariu. A simple linear-time algorithm for computing the center of an interval graph. *Int. J. Comput. Math.*, 34(3-4):121–128, 1990.
- [44] Liam Roditty and Virginia Vassilevska Williams. Fast approximation algorithms for the diameter and radius of sparse graphs. In *Proc. STOC 2013*, pages 515–524. ACM, 2013.
- [45] Avi Shoshan and Uri Zwick. All pairs shortest paths in undirected graphs with integer weights. In *Proc. FOCS 1999*, pages 605–615. IEEE, 1999.
- [46] Oren Weimann and Raphael Yuster. Approximating the diameter of planar graphs in near linear time. *ACM Trans. Algorithms*, 12(1):12:1–12:13, 2016.
- [47] Ryan Williams. A new algorithm for optimal 2-constraint satisfaction and its implications. *Theor. Comput. Sci.*, 348(2-3):357–365, 2005.
- [48] Uri Zwick. All pairs shortest paths using bridging sets and rectangular matrix multiplication. *J. ACM*, 49(3):289–317, 2002.




Figure 9: Reduction of Roditty and Williams [44]. The vertices corresponding to assignments of variables in the sets S_1 and S_2 form an independent set, and the $m + 2$ vertices of C form a clique. Hence, the graph is a split graph.

A Existing Results

The following theorem and proof are due to Roditty and Williams [44]. We provide this result here because we argue about augmented versions of this construction several times.

Theorem 20 ([44]). *Suppose one can distinguish between diameter 2 and 3 in an m -edge undirected unweighted graph in time $O(m^{2-\epsilon})$ for some constant $\epsilon > 0$. Then, CNF-SAT on n variables and m clauses is in $O(2^{n(1-\epsilon/2)})\text{poly}(n, m)$ time, and SETH is false.*

Proof. Given an instance of CNF-SAT on n variables and m clauses, we first partition the variables into two sets S_1 and S_2 on $n/2$ variables each. Create a vertex for every one of the $2^{n/2}$ partial assignments to the variables in S_1 and similarly a vertex for every assignment to the variables in S_2 . Create two nodes t_1 and t_2 and add an edge to t_i from each assignment to the variables of S_i . Create a node for every clause, and connect all clause nodes together with t_1 and t_2 into a clique C of size $m + 2$. Then, connect every assignment node to the clauses that it does not satisfy. See also Figure 9 for an illustration. Now, this graph has diameter 3 if and only if there are two partial assignments, ϕ_1 to S_1 and ϕ_2 to S_2 that together form a satisfying assignment to the CNF formula, i.e. the distance between ϕ_1 and ϕ_2 in the graph is 3 if and only if they form a satisfying assignment, and all other node distances are at most two. The graph has $O(m + 2^{n/2})$ nodes and $O(m2^{n/2})$ edges. Hence, any algorithm for diameter that can distinguish between diameter 2 and 3 in an m' -edge undirected unweighted graph in time $O(m'^{2-\epsilon})$ for some constant $\epsilon > 0$, would imply an algorithm for CNF-SAT in $O\left(m2^{(n/2)(2-\epsilon)}\right) = O\left(2^{n(1-\epsilon/2)}\text{poly}(n, m)\right)$ time, and refute SETH. \square

Related to the above theorem is the ORTHOGONAL VECTORS problem, where, for a value $d = \omega(\log n)$, we are given two sets $A, B \subseteq \{0, 1\}^d$, $|A| = |B| = n$, and are tasked to determine whether there are $a \in A$, $b \in B$ with $a \cdot b = 0$, that is, find two orthogonal vectors. Under SETH, we cannot solve ORTHOGONAL VECTORS in truly subquadratic time [47]. A reduction with the same hardness implication for DIAMETER can be done from ORTHOGONAL VECTORS, and this reduction is essentially equivalent to that of Theorem 20. To see

this, let S_1 have a vertex per vector in A , and S_2 a vertex per vector in B . Let C have $d + 2$ vertices. Again, C is a clique, and has two vertices t_1, t_2 where t_1 is complete to S_1 and t_2 is complete to S_2 . Then, connect a vertex in S_1 or S_2 with a vertex in C if the corresponding vector has a 1 at the corresponding index. Now, the diameter of the graph is 3 if and only if there is an orthogonal pair of vectors.

When discussing $4P_1$ -free graphs in the introduction, we mentioned a possible approach where we make this construction into three cliques. And, a linear-time algorithm for $4P_1$ -free graphs would imply an $O(n^2 + d^2)$ algorithm for ORTHOGONAL VECTORS. To see this, assume we have such an algorithm, and take some input to ORTHOGONAL VECTORS A, B . As above, create the construction that reduces ORTHOGONAL VECTORS to DIAMETER, and also make S_1 and S_2 into a clique. The graph has $O(n + d)$ vertices and $O(n^2 + d^2)$ edges, and is clearly $4P_1$ -free. It takes $O(n^2 + d^2)$ time to construct the graph. But then a linear-time algorithm for DIAMETER on $4P_1$ -free graphs implies an algorithm for ORTHOGONAL VECTORS in $O(n^2 + d^2)$ time.