Wave Propagation in Periodic Plasma Media: A Semiclassical Approach

Reza Torabi

Department of Physics and Astronomy, University of Calgary, Calgary, Alberta T2N 1N4, Canada

e-mail: reza.torabi@ucalgary.ca

Abstract

The propagation of electromagnetic waves in unmagnetized periodic plasma media is studied using the semiclassical wave packet approximation. The formalism gives rise to Berry effect terms in the equation of motion. The Berry effect manifests itself as Rytov polarization rotation law and the polarization-dependent Hall effect. The formalism is also applied to the case of non-periodic inhomogeneous plasma media.

PACS numbers: 03.65.Vf; 03.65.Sq; 52.35.g

1 Introduction

The propagation of electromagnetic waves in inhomogeneous plasmas is an active field of research [1–3] because most of the matter in the universe is in the plasma state [4] and real plasmas are inhomogeneous. A periodic plasma medium is an example of inhomogeneous plasmas. Such a medium can be used to control and confine light because they are analogous to photonic and phononic crystals that permit the passage of waves only at certain frequencies. Ionosphere is another example of inhomogeneous plasma media that the study of its periodicity is of interest in radio physics [5].

Recently, attention has been paid to the phenomena of polarization transport in photonic [6–8] and phononic [9] crystals. These phenomena are related to the topological Berry phase [10] which is a non-integrable phase factor arising from the adiabatic transport of a system around a closed path in the parameter space. In this paper we study the propagation of electromagnetic waves in an unmagnetized slow varying periodic plasma medium via the semiclassical wave packet approximation. The wave packet approximation enables us to solve problems in the presence of periodic structures. It has been used in the study of spin transport in solids [11–13] and also in the photonic [7, 8] and phononic [9] crystals. The formalism yields to the rotation of the polarization plane of a linear polarized wave that is known as Rytov rotation [14, 15]. This is in contrast with the Faraday rotation which is due to the presence of a magnetic field in the medium. In addition, the formalism gives rise to a Berry curvature term in the equation of motion that yields to the spin (polarization) Hall effect [16–18]. According to this effect, waves of left and right circular polarization deflect perpendicular to the direction of propagation oppositely.

The paper is organized as follows. In section II, we derive the wave equation in a periodic plasma medium and introduce Bloch waves as a complete set to construct the wave packet. Section III is devoted to the dynamics of the wave packet. Finally, in section IV we apply the formalism to the case of non-periodic inhomogeneous media. In this case, Berry curvature has the form of opposite signed magnetic monopoles located at the origin of the momentum space. The results for non-periodic inhomogeneous plasma media coincides with the results of [19] obtained from the post geometric approximation.

2 Wave equation and the Bloch waves

An Inhomogeneous plasma medium can be generated, for instance, by applying an external electric field. In this case, the balance equation of the forces in the steady state is

$$\rho_{\alpha 0}(\boldsymbol{x})q_{\alpha}\boldsymbol{E}_{0}(\boldsymbol{x})-m_{\alpha}\nabla p_{\alpha 0}=0,$$

where $\rho_{\alpha 0}$ is unperturbed mass density and $\alpha = e, i$ represent electrons and ions. We are going to derive the equation that determines the wave properties of an inhomogeneous unmagnetized plasma. In this line, similar to the derivation of the wave equation in homogeneous plasmas [20], we consider small harmonic perturbation about the steady state

$$egin{aligned} &
ho_lpha(oldsymbol{x}) =
ho_{lpha 0}(oldsymbol{x}) +
ho_{lpha 1}(oldsymbol{x}) e^{-i\omega t}, \ &oldsymbol{V}_lpha = oldsymbol{K}_0(oldsymbol{x}) + oldsymbol{E}_1(oldsymbol{x}) e^{-i\omega t}, \ &oldsymbol{B} = oldsymbol{B}_1(oldsymbol{x}) e^{-i\omega t}, \end{aligned}$$

but we should notice that the density $\rho_{\alpha 0}(\boldsymbol{x})$ is now a periodic function of \boldsymbol{x} . Putting the above equations into the two fluid momentum equation and keep linear terms, yields to the linearized momentum equation

$$-i\omega\rho_{\alpha 0}(\boldsymbol{x})\boldsymbol{V}_{\alpha 1}(\boldsymbol{x}) = \frac{q_{\alpha}\rho_{\alpha 0}(\boldsymbol{x})}{m_{\alpha}}\boldsymbol{E}_{1}(\boldsymbol{x}) + \rho_{\alpha 1}(\boldsymbol{x})\frac{\nabla p_{\alpha 0}}{n_{\alpha 0}(\boldsymbol{x})} - \gamma\nabla\Big(\frac{\rho_{\alpha 1}(\boldsymbol{x})p_{\alpha 0}}{\rho_{\alpha 0}(\boldsymbol{x})}\Big).$$

In deriving the above, the balance equation and the relation $p_{\alpha 1} = \gamma p_{\alpha 0} \rho_{\alpha 1}(\boldsymbol{x}) e^{-i\omega t} / \rho_{\alpha 0}(\boldsymbol{x})$ have been used. One the other hand, the linear Maxwell equations are

$$\nabla \times B_1 = -\frac{i\omega}{c} E_1 + \frac{4\pi}{c} \sum_{\alpha=e,i} \rho_{\alpha 0}(\boldsymbol{x}) q_{\alpha} V_{\alpha 1}(\boldsymbol{x})$$

and $\nabla \times E_1 = \frac{i\omega}{c} B_1$ which, by taking a curl from both sides, yields to

$$abla imes
abla imes E_1 = \frac{i\omega}{c}
abla imes B_1$$

Combining the linearized momentum equation with linearized Maxwell's equations, results in the wave equation

$$\nabla \times \nabla \times \boldsymbol{E}_{1} - k_{0}^{2} \left(1 - \frac{\omega_{pe}^{2}(\boldsymbol{x})}{\omega^{2}} - \frac{\omega_{pi}^{2}(\boldsymbol{x})}{\omega^{2}} \right) \boldsymbol{E}_{1} = \frac{4\pi}{c^{2}} \sum_{\alpha} \frac{q_{\alpha}}{m_{\alpha}} \left[\gamma \nabla \left(\frac{\rho_{\alpha 1}(\boldsymbol{x})p_{\alpha 0}}{\rho_{\alpha 0}(\boldsymbol{x})} \right) - \frac{\rho_{\alpha 1}(\boldsymbol{x})}{\rho_{\alpha 0}(\boldsymbol{x})} \nabla p_{\alpha 0} \right],$$

where $k_0 = \omega/c,c$ is the speed of light and $\omega_{p\alpha}^2(\boldsymbol{x}) = 4\pi\rho_{\alpha 0}(\boldsymbol{x})q_{\alpha}^2$ is the plasma frequency for particles of species α . The cold plasma approximation and assuming stationary and inhomogeneous density is usually used to describe wave propagation in inhomogeneous media [1, 21]. Under the cold plasma approximation, one can drop the right hand side of the equation and, thus, the wave equation takes the form

$$\nabla \times \nabla \times \boldsymbol{E}_1 = k_0^2 \left(1 - \frac{\omega_{pe}^2(\boldsymbol{x})}{\omega^2} - \frac{\omega_{pi}^2(\boldsymbol{x})}{\omega^2} \right) \boldsymbol{E}_1.$$
(1)

Introducing

$$\epsilon(\boldsymbol{x},\omega) = \left(1 - \frac{\omega_{pe}^2(\boldsymbol{x})}{\omega^2} - \frac{\omega_{pi}^2(\boldsymbol{x})}{\omega^2}\right) \equiv n^2(\boldsymbol{x},\omega)$$

and using, $\nabla \times \nabla \times \boldsymbol{E}_1 = \nabla (\nabla \cdot \mathbf{E}_1) - \nabla^2 \mathbf{E}_1$, to rewrite (1) as

$$\nabla^2 \mathbf{E}_1 - \nabla (\nabla \cdot \mathbf{E}_1) + \frac{\omega^2}{c^2} n^2(\boldsymbol{x}, \omega) \mathbf{E}_1 = 0.$$
⁽²⁾

We decompose \mathbf{E}_1 into a solenoidal part, $\mathbf{E}_{1\perp}$, and an irrotational part, $\mathbf{E}_{1\parallel}$, according to Helmholtz's decomposition, $\mathbf{E}_1 = \mathbf{E}_{1\perp} + \mathbf{E}_{1\parallel}$. The solenoidal and irrotational parts correspond to transverse and longitudinal modes of the wave, respectively. Thence, by taking the curl and the divergence of (2), we obtain two coupled equations

$$\nabla \times \nabla^2 \mathbf{E}_{1\perp} + \frac{\omega^2}{c^2} n^2 \nabla \times \mathbf{E}_{1\perp} = -\frac{\omega^2}{c^2} [(\nabla n^2) \times \mathbf{E}_{1\perp} + (\nabla n^2) \times \mathbf{E}_{1\parallel}], \tag{3}$$

and

$$n^{2}\nabla \cdot \mathbf{E}_{1\parallel} = -(\nabla n^{2}) \cdot \mathbf{E}_{1\parallel} - (\nabla n^{2}) \cdot \mathbf{E}_{1\perp}.$$
(4)

In a smoothly inhomogeneous medium, one can neglect terms on the right hand sides of (3) and (4) since ∇n^2 is very small. So we are left with

$$\nabla \times \left[\nabla^2 \mathbf{E}_{1\perp} + \frac{\omega^2 n^2}{c^2} \mathbf{E}_{1\perp}\right] = 0, \tag{5}$$

and

$$\nabla \cdot \mathbf{E}_{1\parallel} = 0.$$

Therefore (3) and (4) decouple in this approximation, which is called adiabatic approximation. The solenoidal part is divergence free, $\nabla \cdot \mathbf{E}_{1\perp} = 0$, so one can readily show that the divergence of the vector in the bracket of (5) is zero, too. Since, the divergence and the curl of $\mathbf{E}_{1\perp}$ vanish, according to the Helmholtz's theorem, we can conclude that this vector is identically zero, i. e.,

$$\nabla^2 \mathbf{E}_{1\perp} + \frac{\omega^2 n^2}{c^2} \mathbf{E}_{1\perp} = 0.$$
(6)

Similar argument for the irrotational part shows that $\mathbf{E}_{1\parallel} \simeq 0$ in smoothly varying inhomogeneous media. As a result, in the framework of adiabatic approximation the transverse mode decouples and this approximation renders polarization an adiabatic invariant.

Expanding $\mathbf{E}_{1\perp}(\mathbf{x}) = \sum_{\lambda=\pm} \mathbf{e}_{\lambda} E_{1\perp\lambda}(\mathbf{x})$, makes the Eq. (6) an eigenvalue equation

$$-\frac{1}{n^2(\boldsymbol{x},\omega)}\nabla^2 E_{1\perp\lambda} = \frac{\omega^2}{c^2} E_{1\perp\lambda},\tag{7}$$

where \mathbf{e}_{λ} are the orthonormal polarization vectors and $\lambda = +(-)$ stands for right (left)handed polarization. The differential operator in (7) is not Hermitian, However, by introducing the Hermitian operator,

$$\hat{Q} = -\frac{1}{n(\boldsymbol{x})} \nabla^2 \frac{1}{n(\boldsymbol{x})},\tag{8}$$

(7) can be written in the self-adjoint form

$$\hat{Q}\psi_{\lambda} = \frac{\omega^2}{c^2}\psi_{\lambda},\tag{9}$$

where

$$\psi_{\lambda}(\boldsymbol{x}) = n(\boldsymbol{x}) E_{1\perp\lambda}(\boldsymbol{x}).$$

Thus, with the weight function n, eigenfunctions $E_{1\perp\lambda}$ form a complete orthogonal set. We demand a (slow varying) periodic form for $n(\boldsymbol{x})$, so, according to Bloch theorem, the eigenfunctions of (9) are Bloch functions, viz.,

$$\psi_{\lambda \mathsf{n}}(\boldsymbol{x}, \mathbf{q}) = e^{i\mathbf{q}.\boldsymbol{x}} u_{\lambda \mathsf{n}}(\boldsymbol{x}, \mathbf{q}), \qquad (10)$$

where \mathbf{q} is the lattice momentum, \mathbf{n} is the band index and $u_{\lambda \mathbf{n}}(\mathbf{x}, \mathbf{q})$ is periodic in \mathbf{x} with the periodicity imposed on the plasma medium. The Bloch waves (10) form a complete orthonormal set and, thus, can be used as a convenient basis for constructing photons wave packets. In view of our assumption of slow varying inhomogeneity and in the spirit of adiabatic theorem, we drop the band index.

3 The wave packet dynamics

Let us proceed to the quantum level. The Bloch states are $|\psi_{\lambda}(\mathbf{q})\rangle = e^{i\mathbf{q}\cdot\mathbf{x}}|u_{\lambda}(\mathbf{q})\rangle$ and $\psi_{\lambda}(\mathbf{x},\mathbf{q}) = \langle \mathbf{x} | \psi_{\lambda}(\mathbf{q}) \rangle$. We construct the (normalized) wave packet according to

$$|\Psi(t)\rangle = \sum_{\lambda} z_{\lambda}(t) \int d^{3}\mathbf{q} \ a(\mathbf{q}, t) |\psi_{\lambda}(\mathbf{q})\rangle, \qquad (11)$$

where $a(\mathbf{q}, t)$ and $z_{\lambda}(t)$ are amplitudes with the normalizations

$$\int d^{3}\mathbf{q} |a(\mathbf{q},t)|^{2} = 1,$$
$$\sum_{\lambda} |z_{\lambda}(t)|^{2} = 1.$$

We assume that the momentum distribution $|a(\mathbf{q},t)|^2$ is sharply peaked around its mean value \mathbf{q}_c ,

$$\mathbf{q}_c(t) = \int d\mathbf{q} \ |a|^2 \mathbf{q}$$

which is the momentum of the wave packet. Writing the amplitude in the form $a(\mathbf{q}, t) = |a(\mathbf{q}, t)| e^{-i\gamma(\mathbf{q}, t)}$, the wave packet center position, $\langle \Psi | \hat{\boldsymbol{x}} | \Psi \rangle$, will be [9,13]

$$\boldsymbol{x}_{c} = \int d^{3} \mathbf{q} \left| \boldsymbol{a} \right|^{2} \left\{ \nabla_{\mathbf{q}} \gamma(\mathbf{q}, t) + \sum_{\lambda', \lambda} z_{\lambda'}^{*} z_{\lambda} \left\langle u_{\lambda'}(\mathbf{q}) \right| i \nabla_{\mathbf{q}} \left| u_{\lambda}(\mathbf{q}) \right\rangle \right\}.$$
(12)

According to our assumption of a well localized wave packet in the momentum space, (12) reduces to

$$\boldsymbol{x}_{c}(t) \approx \nabla_{\mathbf{q}_{c}} \gamma(\mathbf{q}_{c}, t) + \sum_{\lambda', \lambda} z_{\lambda'}^{*} z_{\lambda} \mathbf{A}_{\lambda' \lambda}(\mathbf{q}_{c}),$$

where $\mathbf{A}_{\lambda'\lambda}(\mathbf{q}) = \langle u_{\lambda'}(\mathbf{q}) | i \nabla_{\mathbf{q}} | u_{\lambda}(\mathbf{q}) \rangle$. By introducing the notation $|z(t)\rangle = \begin{pmatrix} z_{+}(t) \\ z_{-}(t) \end{pmatrix}$, the above equation will be

$$\boldsymbol{x}_{c}(t) \approx \nabla_{\mathbf{q}_{c}} \gamma(\mathbf{q}_{c}, t) + \langle z \mid \hat{\mathbf{A}}(\mathbf{q}_{c}) \mid z \rangle,$$
(13)

where $\hat{\mathbf{A}}$ is a 2 × 2 matrix with vector elements $\mathbf{A}_{\lambda'\lambda}$ and $\mathbf{A}_B = \langle z \mid \hat{\mathbf{A}}(\mathbf{q}_c) \mid z \rangle$ is called Berry connection.

Let us derive equations of motion for such a wave packet. The wave packet dynamics is obtained from the effective Lagrangian ($\hbar = 1$)

$$\mathcal{L} = i \langle \Psi | \dot{\Psi} \rangle - \langle \Psi | \hat{H} | \Psi \rangle, \tag{14}$$

using the Euler-Lagrange equation [22]. The second term in the Lagrangian is $\langle \Psi | \hat{H} | \Psi \rangle = \omega$, and the first one will be

$$i\langle\Psi|\dot{\Psi}\rangle = \frac{\partial\gamma(\mathbf{q}_c,t)}{\partial t} - i\langle z|\dot{z}\rangle.$$
(15)

In deriving the above, we have used (11) and (13). By substituting

$$\frac{\partial \gamma(\mathbf{q}_c, t)}{\partial t} = \frac{d\gamma(\mathbf{q}_c, t)}{dt} - \dot{\mathbf{q}}_c \cdot \nabla_{\mathbf{q}_c} \gamma(\mathbf{q}_c, t),$$

in Eq. (15), the Lagrangian (14) takes the form

$$\mathcal{L} = i \langle z | \dot{z} \rangle + \dot{\mathbf{q}}_c \cdot \mathbf{A}_B + \mathbf{q}_c \cdot \dot{\boldsymbol{x}}_c - \omega.$$
(16)

We have omitted a term of total time derivative $d(\gamma - \boldsymbol{x}_c \cdot \mathbf{q}_c)/dt$ in the Lagrangian, because it does not affect the equations of motion. The equations of motion that follow from the effective Lagrangian (16) are

$$\dot{\boldsymbol{x}}_c = \mathbf{v}_g + \dot{\mathbf{q}}_c \times \boldsymbol{\Omega}(\mathbf{q}_c), \tag{17}$$

$$\dot{\mathbf{q}}_c = \nabla_{\boldsymbol{x}_c} \omega, \tag{18}$$

$$|\dot{z}_c\rangle = -i\dot{\mathbf{q}}_c \cdot \hat{\mathbf{A}}(\mathbf{q}_c)|z_c\rangle,\tag{19}$$

where $\mathbf{v}_g = \nabla_{\mathbf{q}_c} \omega$ is the group velocity and $\mathbf{\Omega} = \nabla \times \mathbf{A}_B$ is called Berry curvature.

The second term on the right hand of Eq. (17), represents a polarization dependent velocity perpendicular to the group velocity. It is spin (polarization) Hall effect of photons. Eq. (19) describes the time evolution of the polarization and has the solution

$$|z(t)\rangle = e^{i\int_C d^3 \mathbf{q}_c \cdot \hat{\mathbf{A}}} |z(0)\rangle, \qquad (20)$$

where C is the wave packet trajectory in the parameter (momentum) space. Eq. (20) can be written in the form $|z(t)\rangle = e^{i\hat{\gamma}}|z(0)\rangle$, where $\hat{\gamma} = \int_C d^3 \mathbf{q}_c \cdot \hat{\mathbf{A}}$ is the Berry-phase matrix. The phase shift is a geometric one which leads to the rotation of the polarization plane for a linear polarized wave known as the Rytov rotation law.

4 Non-periodic inhomogeneous media

The wave packet dynamics formalism is also applicable to the case of slightly inhomogeneous non-periodic media. In this case the basis for the wave packet expansion is $|\psi_{\lambda}(\mathbf{q})\rangle = e^{i\mathbf{q}\cdot\mathbf{x}}|\mathbf{e}_{\lambda}(\mathbf{q})\rangle$ where \mathbf{e}_{λ} are the polarization vectors. Therefore, we can replace $|u_{\lambda}\rangle$ by $|\mathbf{e}_{\lambda}\rangle$ in the aforementioned formalism. We also write \mathbf{e}_{λ} in the helicity basis as $\mathbf{e}_{\lambda} = \frac{1}{\sqrt{2}}(\mathbf{i}_{\theta} \pm i\mathbf{i}_{\phi})$, where \mathbf{i}_{θ} (\mathbf{i}_{ϕ}) is the zenithal (azimuthal) unit vector in the spherical coordinates of the momentum space. Thence, according to $\mathbf{A}_{\lambda'\lambda}(\mathbf{q}) = \langle e_{\lambda'} | i\nabla_{\mathbf{q}} | e_{\lambda} \rangle$, we will have

$$\hat{\mathbf{A}} = \frac{\cot\theta}{q_c} \hat{\sigma}_3 \mathbf{i}_\phi,$$

where $\hat{\sigma}_3$ is the Pauli matrix. Thus, according to (20), right/left circularly polarized waves experience a geometric phase shift of $\pm \int_C \cos\theta \, d\varphi$, respectively. $\theta(\varphi)$ is the zenithal (azimuthal) angle in spherical coordinates. Therefore, if a linear polarized wave propagates in a periodic plasma medium, the polarization plane will rotate which is topological in nature. The Berry curvature will be

$$\mathbf{\Omega}=\mprac{\mathbf{q}_c}{q_c^3}$$

for right/left circularly polarized waves respectively, which, according to (18), propagate along different trajectories and have displacements perpendicular to the wave propagation direction. These results of non-periodic inhomogeneous plasma media coincide with the results of [19] obtained from the post geometric approximation.

5 Conclusion

We studied the propagation of electromagnetic waves in unmagnetized slow varying periodic plasma media via the semiclassical wavepacket approximation . This approximation enables us to solve problems in the presence of periodic structures. The formalism yields to the appearance of an Abelian gauge field (Berry connection) in the effective lagrangian governing the wavepacket dynamics. This results in the rotation of polarization plane, and the polarization-dependent Hall effect for transverse photons in periodic plasma media. The rotation of the polarization plane, of a linear polarized wave, in unmagnetized periodic plasma media is topological in nature and differs from the Faraday rotation which is due to the presence of a magnetic field in the medium. The formalism also applied to the case of non-periodic inhomogeneous plasma media.

References

- S. Hacquin, S. Heuraux, M. Colin and G. Leclert, J. Comp. Phys. 174, 1 (2001).
- [2] S. Cho, Phys. Plasmas **11**, 4399 (2004).
- [3] H. K. Malik, S. Kumer and K.P. Singh, Laser Part. Beams 26, 197 (2008).
- [4] F. F. Chen, Introduction to plasma physics and controlled fusion, 2nd ed., Vol. 1 (Plenum Press, New York, 1984).
- [5] N. V. Bakhmet eva, V. V. Belikovich, M. N. Egereva, A. V. Tolmacheva, Radiophys. Quantum Electron., 53, 69 (2010).
- [6] K. Yu. Bliokh and V. D. Freilikher, Phys. Rev. B 72, 035108 (2005).
- [7] M. Onoda, S. Murakami and N. Nagaosa, Phys. Rev. Lett. **93**, 083901 (2004).
- [8] M. Onoda, S. Murakami and N. Nagaosa, Phys. Rev. E 74, 066610 (2006).
- [9] R. Torabi and M. Mehrafarin, JETP Lett. 88, 590 (2008).
- [10] M. V. Berry, Proc. R. Soc. A **392**, 45 (1984).
- [11] D. Culcer, A. MacDonald and Q. Niu, Phys. Rev. B 68, 045327 (2003).

- [12] M. C. Chang and Q. Niu, Phys. Rev. B 53, 7010 (1996).
- [13] G. Sundaram and Q. Niu, Phys. Rev. B 59, 14915 (1999).
- [14] S. M. Rytov, Dokl. Akad. Nauk SSSR 18, 263 (1938).
- [15] V. V. Vladimirskii, Dokl. Akad. Nauk SSSR 13, 222 (1941).
- [16] R. Torabi, Can. J. Phys. 88, 1 (2010).
- [17] A. V. Dooghin, N. D. Kudnikova, V. S. Liberman and B. Ya. Zeldovich, Phys. Rev. A 45, 8204 (1992).
- [18] V. S. Liberman, B. Ya. Zeldovich, Phys. Rev. A 46, 5199 (1992).
- [19] R. Torabi and M. Mehrafarin, JETP Lett. **95**, 277 (2012).
- [20] N. A. Krall and A. W. Trivelpiece, Principles of Plasma Physics (McGraw-Hill, New York, 1973).
- [21] V. L. Ginzburg, The Propagation of Electromagnetic Waves in Plasmas (Gordon and Breach, NY, 1964).
- [22] R. Jackiw and A. Kerman, Phys. Lett. A **71**, 158 (1979).
- [23] S. P. Kuo, Progress In Electromagnetics Research B, 60, 275 (2014).