A D-module approach to invariant distributions with finitely many orbits

Hiroyuki Ochiai*

July 9, 2024

Abstract

Tauchi provides an example illustrating the action of a real algebraic subgroup H of $GL(2n, \mathbb{R})$ with finitely many orbits on \mathbb{R}^{2n} , while the dimension of the space of relative H-invariant distributions on \mathbb{R}^{2n} is infinite. We offer a perspective on this example from the viewpoint of D-modules, where we explicitly determine the simple quotient regular holonomic D-modules and demonstrate that the distributions exhibit an enlarged symmetry.

1 Introduction

In 2018, Tauchi [10] discovered a representation of a Lie group with finitely many orbits, where the dimension of relatively invariant distributions with a fixed character is infinite. This intriguing representation is motivated by providing an example demonstrating that the smooth regular representation on some homogeneous space of $SL(2n, \mathbb{R})$ with finitely many orbits under a real mirabolic subgroup has infinite multiplicity of a degenerate principal series representation [11], [12]. Distributions of interest are characterized by being annihilated by vector fields corresponding to a Lie algebra. In this paper, we explore the left ideal generated by these vector fields in the ring of differential operators and the corresponding D-modules. This D-module must be non-holonomic due to the infinite dimension of its solutions. We identify countably many quotient D-modules, each turning out to be a simple holonomic D-module. Subsequently, we revisit the Lie algebraic interpretation of these quotient D-modules. It's important to note that the group used here is solvable (upper triangular), potentially resulting in the corresponding D-module having irregular singularities. Nevertheless, these simple holonomic D-modules must exhibit regular singularities.

The interplay between Lie algebra and D-modules for relatively invariant distributions was leveraged by the author [7] to gain insights into the seminal work of van Dijk. In [13], among other contributions, he highlights the non-existence of a spherical distribution — an invariant eigendistribution of the pseudo-Laplacian — on the tangent space of a rank-one semisimple symmetric space $SL(m+1,\mathbb{R})/GL(m,\mathbb{R})$ supported on the singular nilpotent orbits. This marked the beginning of harmonic analysis on this semisimple symmetric space [21, 22, 18, 19] and the exploration of a generalized Gelfand pair [14, 15, 20]. Kowata [5] establishes the non-existence of singular invariant distributions and hyperfunctions [4] without resorting to the pseudo-Laplacian, subject to certain conditions. For instance, when considering the tangent space of $SL(m+1,\mathbb{R})/GL(m,\mathbb{R})$,

^{*}Keywords: D-module, holonomic system, invariant distribution, orbit decomposition, Lie algebra. This work is partially supported by JSPS Grant-in-Aid for Transformative Research Areas (A) (22H05107).

Kowata identifies a larger symmetry, $SO_0(m,m)$, compared to $GL(m,\mathbb{R})$. The author [7] rearticulates this finding in terms of D-modules. Notably, in this tangent space setting, as well as the global setting [8], two Lie algebras, $\mathfrak{gl}(m,\mathbb{R})$ and $\mathfrak{so}(m,m)$, exhibit the same D-modules. As a corollary, we deduce an automatic extension of the symmetry of spherical distribution and the non-existence of singular spherical distribution. In other words, there is no contribution of singular orbits under the smaller Lie group.

Tauchi's example focuses primarily on real projective spaces \mathbb{P}^3 and \mathbb{P}^5 of dimensions three and five, respectively. These spaces correspond to the real vector spaces $\mathbb{R}^4 \setminus \{0\}$ and $\mathbb{R}^6 \setminus \{0\}$ with the origin removed and divided by \mathbb{R}^{\times} . In our analysis, we operate on vector spaces $\mathbb{R}^{2n} = \mathbb{C}^n$ with n=2,3 without dividing by \mathbb{R}^{\times} , and we observe that including the origin poses no issues.

We introduce various real Lie subgroups of $GL(2n,\mathbb{R})$ and their Lie algebras, including their complexifications. Further, we delve into the orbit decomposition of \mathbb{R}^{2n} and \mathbb{C}^{2n} for each case.

In the context of the ring of differential operators with polynomial coefficients, denoted as

$$\mathcal{D} = \mathbb{C}[z_1, \dots, z_{2n}, \frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_{2n}}],$$

we define the left ideal \mathcal{I} of \mathcal{D} and the corresponding D-module \mathcal{D}/\mathcal{I} , along with its hyperfunction or distribution solutions. The reference for D-modules will be [1, 2].

$\mathbf{2}$ n=2

2.1Lie algebra

We define a solvable Lie subalgebra of $\mathfrak{gl}(4,\mathbb{C})$ by

$$\mathfrak{h}_1 := \mathfrak{a} \oplus \mathfrak{n} := (\mathbb{C}E_{11} \oplus \mathbb{C}E_{22} \oplus \mathbb{C}E_{33} \oplus \mathbb{C}E_{44}) \oplus (\mathbb{C}E_{14} \oplus \mathbb{C}E_{24} \oplus \mathbb{C}E_{31} \oplus \mathbb{C}E_{32} \oplus \mathbb{C}E_{34})$$

The Lie algebra \mathfrak{h}_1 is illustrated as

(*	0	0	*)	$\langle z_1 \rangle$		(*	*	*	*)	(z_3)	
0	*	0	*	z_2		0	*	0	*	z_2	
*	*	*	*	z_3	,	0	0	*	*	z_1	,
$\setminus 0$	0	0	*/	$\langle z_4 \rangle$		$\setminus 0$	0	0	*/	$\langle z_4 \rangle$	

where the figure on the left is represented in standard coordinates, whereas the figure on the right

illustrates the upper-triangular realization of the Lie algebra. Put $c = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ and a Lie algebra isomorphism $t : \sigma(A) \to \sigma(A)$ is a finite right of the second seco

algebra isomorphism $\iota : \mathfrak{gl}(4) \to \mathfrak{gl}(4)$ by $\iota(A) = cAc^{-1}$.

We now consider the several subalgebras $\mathfrak{h}_3 \subset \mathfrak{h}_2 \subset \mathfrak{h}_1$ defined by

$$\begin{split} \mathfrak{h}_{2} &:= \mathfrak{a} \oplus (\mathbb{C}E_{14} \oplus \mathbb{C}E_{32}).\\ \mathfrak{h}_{3} &:= (\mathbb{C}(E_{11} + E_{22}) \oplus \mathbb{C}(E_{33} + E_{44})) \oplus (\mathbb{C}E_{14} \oplus \mathbb{C}E_{32}).\\ \mathfrak{h}_{1} &= \begin{pmatrix} * & 0 & 0 & * \\ 0 & * & 0 & * \\ * & * & * & * \\ 0 & 0 & 0 & * \end{pmatrix}, \quad \mathfrak{h}_{2} = \begin{pmatrix} * & 0 & 0 & * \\ 0 & * & 0 & 0 \\ 0 & * & * & 0 \\ 0 & 0 & 0 & * \end{pmatrix}, \quad \mathfrak{h}_{3} = \begin{pmatrix} a_{11} & 0 & 0 & a_{14} \\ 0 & a_{11} & 0 & 0 \\ 0 & a_{32} & a_{44} & 0 \\ 0 & 0 & 0 & a_{44} \end{pmatrix}. \end{split}$$

$$\iota(\mathfrak{h}_1) = \begin{pmatrix} * & * & * & * \\ 0 & * & 0 & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{pmatrix}, \quad \iota(\mathfrak{h}_2) = \begin{pmatrix} * & * & 0 & 0 \\ 0 & * & 0 & 0 \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{pmatrix}, \quad \iota(\mathfrak{h}_3) = \begin{pmatrix} a_{44} & a_{32} & 0 & 0 \\ 0 & a_{11} & 0 & 0 \\ 0 & 0 & a_{11} & a_{14} \\ 0 & 0 & 0 & a_{44} \end{pmatrix}.$$

2.2 Real form

Let $X := \mathbb{C}^4$ be a complex four-dimensional vector space with the standard coordinates $\mathbf{z} = (z_1, z_2, z_3, z_4)^T$. Let $J := \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$. We define a real four-dimensional vector space

$$X_{\mathbb{R}} := \{ \mathbf{z} = (z_1, z_2, z_3, z_4)^T \in \mathbb{C}^4 \mid \mathbf{z} = J\overline{\mathbf{z}} \} = \{ (z_1, z_2, \overline{z_1}, \overline{z_2})^T \mid z_1, z_2 \in \mathbb{C} \}.$$

Then $X_{\mathbb{R}}$ is a real form of the complex vector space $X = \mathbb{C}^4$. The corresponding real form of the complex Lie algebra $\mathfrak{gl}(4,\mathbb{C}) = \operatorname{End}_{\mathbb{C}}(X)$ is $\mathfrak{g}_{\mathbb{R}} := \{X \in \mathfrak{gl}(4,\mathbb{C}) \mid J\overline{X}J = X\}$.

We define the real Lie algebra $\mathfrak{h}_{i\mathbb{R}} := \mathfrak{h}_i \cap \mathfrak{g}_{\mathbb{R}}$ for i = 1, 2, 3.

$$\mathbf{Lemma 1.} \quad \bullet \ \mathfrak{g}_{\mathbb{R}} = \left\{ \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ \frac{a_{13}}{a_{23}} & \frac{a_{14}}{a_{24}} & \frac{a_{11}}{a_{21}} & \frac{a_{12}}{a_{22}} \end{pmatrix} \right\}.$$
$$\bullet \ \mathfrak{h}_{2\mathbb{R}} = \left\{ \begin{pmatrix} a_{11} & 0 & 0 & a_{14} \\ 0 & a_{22} & 0 & 0 \\ 0 & \overline{a_{14}} & \overline{a_{11}} & 0 \\ 0 & 0 & 0 & \overline{a_{22}} \end{pmatrix} \right\} = \left\{ \begin{pmatrix} a_{11} & 0 & 0 & a_{14} \\ 0 & \overline{a_{44}} & 0 & 0 \\ 0 & \overline{a_{14}} & \overline{a_{11}} & 0 \\ 0 & 0 & 0 & a_{44} \end{pmatrix} \right\}, which is a real form of \mathfrak{h}_2.$$

• $\mathfrak{h}_{1\mathbb{R}} = \mathfrak{h}_{2\mathbb{R}}$, which is not a real form of \mathfrak{h}_1 .

•
$$\mathfrak{h}_{3\mathbb{R}} = \left\{ \begin{pmatrix} a_{11} & 0 & 0 & a_{14} \\ 0 & a_{11} & 0 & 0 \\ 0 & \overline{a_{14}} & \overline{a_{11}} & 0 \\ 0 & 0 & 0 & \overline{a_{11}} \end{pmatrix} \right\}, \text{ which is a real form of } \mathfrak{h}_3.$$

The image of each real Lie algebra by ι is given as

$$\iota(\mathfrak{g}_{\mathbb{R}}) = \left\{ \begin{pmatrix} \overline{a_{11}} & \overline{a_{14}} & \overline{a_{13}} & \overline{a_{12}} \\ a_{23} & a_{22} & a_{21} & a_{24} \\ a_{13} & a_{12} & \overline{a_{24}} & \overline{a_{23}} & \overline{a_{22}} \end{pmatrix} \right\},\$$
$$\iota(\mathfrak{h}_{2\mathbb{R}}) = \left\{ \begin{pmatrix} \overline{a_{11}} & \overline{a_{14}} & 0 & 0 \\ 0 & \overline{a_{44}} & 0 & 0 \\ 0 & 0 & a_{11} & a_{14} \\ 0 & 0 & 0 & a_{44} \end{pmatrix} \right\}, \quad \iota(\mathfrak{h}_{3\mathbb{R}}) = \left\{ \begin{pmatrix} \overline{a_{11}} & \overline{a_{14}} & 0 & 0 \\ 0 & a_{11} & 0 & 0 \\ 0 & 0 & a_{11} & a_{14} \\ 0 & 0 & 0 & \overline{a_{41}} \end{pmatrix} \right\}.$$

The dimension of each Lie algebra is

 $\dim_{\mathbb{C}}\mathfrak{h}_1=9,\ \dim_{\mathbb{C}}\mathfrak{h}_2=6,\ \dim_{\mathbb{C}}\mathfrak{h}_3=4,\ \dim_{\mathbb{R}}\mathfrak{h}_{1\mathbb{R}}=\dim_{\mathbb{R}}\mathfrak{h}_{2\mathbb{R}}=6,\ \dim_{\mathbb{R}}\mathfrak{h}_{3\mathbb{R}}=4.$

2.3 Orbit decomposition

For each Lie algebra or its corresponding Lie group, we consider the orbit decomposition by the action of $GL(4, \mathbb{C}) \supset H_1 \supset H_2 \supset H_3$ on $X = \mathbb{C}^4$ and the action of $G_{\mathbb{R}} \supset H_{1\mathbb{R}} = H_{2\mathbb{R}} \supset H_{3\mathbb{R}}$ on $X_{\mathbb{R}}$. The upper triangular realization $\iota(\mathfrak{h}_i)$ is preferred over the standard realization \mathfrak{h}_i to facilitate the observation of the orbit decomposition.

2.3.1 Complex

We define Zariski locally closed subsets O_2, O_1, O_0 of $\mathbb{C}^2 = \{(z_3, z_2)^T \in \mathbb{C}^2\}$ and O'_2, O'_1, O'_0 of $\mathbb{C}^2 = \{(z_1, z_4)^T \in \mathbb{C}^2\}$ by

$$O_{2} := \{ (z_{3}, z_{2})^{T} \in \mathbb{C}^{2} \mid z_{2} \neq 0 \},\$$

$$O_{1} := \{ (z_{3}, z_{2})^{T} \in \mathbb{C}^{2} \mid z_{2} = 0, z_{3} \neq 0 \},\$$

$$O_{0} := \{ (z_{3}, z_{2})^{T} \in \mathbb{C}^{2} \mid z_{2} = z_{3} = 0 \},\$$

$$O'_{2} := \{ (z_{1}, z_{4})^{T} \in \mathbb{C}^{2} \mid z_{4} \neq 0 \},\$$

$$O'_{1} := \{ (z_{1}, z_{4})^{T} \in \mathbb{C}^{2} \mid z_{4} = 0, z_{1} \neq 0 \},\$$

$$O'_{0} := \{ (z_{1}, z_{4})^{T} \in \mathbb{C}^{2} \mid z_{4} = z_{1} = 0 \}.$$

Here, the index denotes the dimension of each orbit. We define $O_{ij} = O_i \times O'_j \subset \mathbb{C}^2 \times \mathbb{C}^2 \subset \mathbb{C}^4$ for i, j = 0, 1, 2. Then

$$\mathbb{C}^2 = \bigsqcup_{i=0,1,2} O_i, \mathbb{C}^2 = \bigsqcup_{j=0,1,2} O'_j, \mathbb{C}^4 = \bigsqcup_{i,j=0,1,2} O_{ij}.$$

For $c \in \mathbb{C}^{\times}$, we define (Tauchi)

$$O_{12c} := \{ (z_3, z_2, z_1, z_4) \in O_{12} \mid z_3 = cz_4 \}, O_{21c} := \{ (z_3, z_2, z_1, z_4) \in O_{21} \mid z_1 = cz_2 \}.$$

Note that $\bigsqcup_{c \in \mathbb{C}^{\times}} O_{12c} = O_{12}$, and $\bigsqcup_{c \in \mathbb{C}^{\times}} O_{21c} = O_{21}$.

Lemma 2. • O_{ij} are $\iota(H_2)$ -orbits.

- O_{ij} remain $\iota(H_3)$ -orbits unless (i, j) = (1, 2), (2, 1).
- O_{12c} and O_{21c} are $\iota(H_3)$ -orbits for $c \in \mathbb{C}^{\times}$.
- $O_{22} \cup O_{12} \cup O_{02}$, O_{21} , O_{20} , $O_{11} \cup O_{01}$, O_{10} , O_{00} are $\iota(H_1)$ -orbits.

2.3.2 Real

The real form $X_{\mathbb{R}} = \{(z_1, z_2, \overline{z_1}, \overline{z_2}) \mid z_1, z_2 \in \mathbb{C}\}$ is a subset of $O_{22} \cup O_{11} \cup O_{00}$.

Lemma 3. • There are three $H_{3\mathbb{R}}$ -orbits

$$O_{22\mathbb{R}} := \{ (z_1, z_2, \overline{z_1}, \overline{z_2}) \in X_{\mathbb{R}} \mid z_2 \neq 0 \},\$$

$$O_{11\mathbb{R}} := \{ (z_1, z_2, \overline{z_1}, \overline{z_2}) \in X_{\mathbb{R}} \mid z_2 = 0, z_1 \neq 0 \},\$$

$$O_{00\mathbb{R}} := \{ (z_1, z_2, \overline{z_1}, \overline{z_2}) \in X_{\mathbb{R}} \mid z_2 = z_1 = 0 \}$$

on $X_{\mathbb{R}}$.

- Each orbit under $H_{3\mathbb{R}}$ is also an orbit under $H_{2\mathbb{R}}$.
- The real manifold $O_{ii\mathbb{R}}$ is a real form of the complex manifold O_{ii} for i = 0, 1, 2.

Remark 4. The groups H_3 and $H_{3\mathbb{R}}$ and their orbit decompositions are given by Tauchi [10]. It is important that H_3 has infinitely many orbits in \mathbb{C}^4 and $H_{3\mathbb{R}}$ has finitely many orbits in the real form $X_{\mathbb{R}}$.

2.4 D-module

In $X = \mathbb{C}^4$, we use coordinates $(z_1, z_2, z_3, z_4) \in X$, and $(z_1, z_2, z_3, z_4, \zeta_1, \zeta_2, \zeta_3, \zeta_4) \in T^*X$ for the cotangent bundle. The ring of differential operators on X is denoted by $\mathcal{D} = \mathcal{D}_X = \mathbb{C}[z_1, z_2, z_3, z_4, \frac{\partial}{\partial z_1}, \frac{\partial}{\partial z_2}, \frac{\partial}{\partial z_3}, \frac{\partial}{\partial z_4}]$. For a complex subspace $Y := \{z_2 = z_4 = 0\}$, which is the closure of O_{11} , we define a D-module

$$\mathcal{B}_{Y|X} = \mathcal{H}^2_{[Y]}(\mathcal{O}_X) = \mathcal{D}/(\mathcal{D}\frac{\partial}{\partial z_1} + \mathcal{D}z_2 + \mathcal{D}\frac{\partial}{\partial z_3} + \mathcal{D}z_4)$$

The characteristic variety of $\mathcal{B}_{Y|X}$ is the conormal bundle $\{\zeta_1 = z_2 = \zeta_3 = z_4 = 0\} = T_Y^* X \subset T^* X$ of Y, and the multiplicity of $T_Y^* X$ is one, that is, $\mathcal{B}_{Y|X}$ is a simple holonomic system. We denote by $\delta(z_2, z_4) \in \mathcal{B}_{Y|X}$ the class represented by $1 \in \mathcal{D}$. For $l \in \mathbb{Z}_{\geq 0}$, we consider the element $T_l := (z_1 \frac{\partial}{\partial z_2})^l \delta(z_2, z_4) \in \mathcal{B}_{Y|X}$. Then

Lemma 5. The annihilator \mathcal{I}_{1l} of T_l is the left ideal of \mathcal{D} generated by

$$z_1 \frac{\partial}{\partial z_1} - l, \quad \frac{\partial^{l+1}}{\partial z_1^{l+1}}, \quad z_2 \frac{\partial}{\partial z_2} + l + 1, \quad z_2^{l+1}, \quad \frac{\partial}{\partial z_3}, \quad z_4. \tag{1}$$

Proof. The partial algebraic Fourier transform

$$z_1 \mapsto z_1, z_2 \mapsto \frac{\partial}{\partial z_2}, z_3 \mapsto z_3, z_4 \mapsto \frac{\partial}{\partial z_4}, \frac{\partial}{\partial z_1} \mapsto \frac{\partial}{\partial z_1}, \frac{\partial}{\partial z_2} \mapsto -z_2, \frac{\partial}{\partial z_3} \mapsto \frac{\partial}{\partial z_3}, \frac{\partial}{\partial z_4} \mapsto -z_4$$

gives an algebra isomorphism from \mathcal{D} to \mathcal{D} . This isomorphism sends $\mathcal{B}_{Y|X}$ to

$$\mathcal{D}/(\mathcal{D}\frac{\partial}{\partial z_1} + \mathcal{D}\frac{\partial}{\partial z_2} + \mathcal{D}\frac{\partial}{\partial z_3} + \mathcal{D}\frac{\partial}{\partial z_4}) \simeq \mathcal{O}_X.$$

The image of $\delta(z_2, z_4)$ is $1 \in \mathcal{O}_X$ and the image of T_l is $(-z_1 z_2)^l \in \mathcal{O}_X$. In the one variable case, the annihilator of z_1^l in $\mathbb{C}[z_1, \frac{\partial}{\partial z_1}]$ is the left ideal of $\mathbb{C}[z_1, \frac{\partial}{\partial z_1}]$ generated by $z_1 \frac{\partial}{\partial z_1} - l$ and $\frac{\partial^{l+1}}{\partial z_1^{l+1}}$. The inverse partial Fourier transform of $z_2 \frac{\partial}{\partial z_2} - l$ is $\frac{\partial}{\partial z_2}(-z_2) - l = -(z_2 \frac{\partial}{\partial z_2} + l + 1)$, which demonstrates the desired lemma.

Let \mathcal{I}_3 be the left ideal generated by

$$z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2} + 1, \ \frac{\partial}{\partial z_3}, \ z_4.$$
 (2)

We define left D-modules $\mathcal{M}_{1l} := \mathcal{D}/\mathcal{I}_{1l}$ and $\mathcal{M}_3 := \mathcal{D}/\mathcal{I}_3$.

Theorem 6. (i) We have a D-module isomorphism $\mathcal{M}_{1l} \ni P \mapsto P \circ (z_1 \frac{\partial}{\partial z_2})^l \delta(z_2, z_4) \in \mathcal{B}_{Y|X}$. Especially, \mathcal{M}_{1l} is a simple holonomic system. (ii) The natural D-module homomorphism $\mathcal{M}_3 \to \mathcal{M}_{1l}$ is surjective.

(iii) For any positive integer L, the natural D-module homomorphism $\mathcal{M}_3 \to \bigoplus_{l=0}^L \mathcal{M}_{1l}$ is surjective.

Proof. (i) follows from Lemma lemma: isom, and (ii) follows from the fact that $\mathcal{I}_3 \subset \mathcal{I}_{1l}$ for all $l \in \mathbb{Z}_{\geq 0}$. We prove (iii). For a given $P_l \in \mathcal{D}$ with $l = 0, 1, \ldots, L$, we put

$$P := \sum_{l=0}^{L} P_l \prod_{0 \le k \le L, k \ne l} \frac{z_1 \frac{\partial}{\partial z_1} - k}{l-k} \in \mathcal{D},$$

then $P \equiv P_l \mod \mathcal{I}_{1l}$ with $l = 0, 1, \dots, L$ follows.

We observe that

$$z_4 \frac{\partial}{\partial z_1}, \ z_2 \frac{\partial}{\partial z_3}, \ z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2} + 1, \ z_3 \frac{\partial}{\partial z_3} + z_4 \frac{\partial}{\partial z_4} + 1 \in \mathcal{I}_3.$$
 (3)

We will interpret these elements as images of a Lie algebra. Let

$$\rho:\mathfrak{gl}(4)=\mathfrak{gl}(4,\mathbb{C})\ni E_{ij}\mapsto -z_j\frac{\partial}{\partial z_i}\in\mathcal{D}$$

be a Lie algebra homomorphism induced from the standard representation of $GL(4, \mathbb{C})$ on $X = \mathbb{C}^4$. We define the character $\chi_l : \mathfrak{h}_1 = \mathfrak{a} \oplus \mathfrak{n} \to \mathbb{C}$ by

$$\chi_l(E_{11}) = -l, \ \chi_l(E_{22}) = l+1, \ \chi_l(E_{33}) = 0, \ \chi_l(E_{44}) = 1$$

and $\chi_l(\mathfrak{n}) = 0$. The restriction of χ_l on \mathfrak{h}_3 is denoted by χ since it does not depend on l. Explicitly,

$$\chi(E_{11} + E_{22}) = 1, \ \chi(E_{33} + E_{44}) = 1.$$

Then Lemma 5 and the equation (3) show that

$$\begin{array}{rcl} (\rho - \chi)(\mathfrak{h}_3) &\subset & \mathcal{I}_3 \\ \cap & & \cap \\ (\rho - \chi_l)(\mathfrak{h}_1) &\subset & \mathcal{I}_{1l} \end{array}$$

$$(4)$$

Remark 7. The Lie algebra homomorphism ρ induces the associative algebra homomorphism

$$\tilde{\rho}: U(\mathfrak{gl}(4)) \to \{ P \in \mathcal{D} \mid [P, z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2} + z_3 \frac{\partial}{\partial z_3} + z_4 \frac{\partial}{\partial z_4}] = 0 \}$$

from the universal enveloping algebra of the Lie algebra $\mathfrak{gl}(4)$ onto the subalgebra of \mathcal{D} of homogeneous degree zero. Note that the map $\tilde{\rho}$ is not injective; for example, a non-zero element $E_{14}E_{32} - E_{12}E_{34} \in U(\mathfrak{gl}(4))$ belongs to the kernel of $\tilde{\rho}$. Therefore, it is possible that two different Lie subalgebras of $\mathfrak{gl}(4)$ generate two distinct left ideals of $U(\mathfrak{gl}(4))$, whose images under $\tilde{\rho}$ may coincide. This is one of the reasons why invariance under a smaller Lie algebra extends to that under a larger Lie algebra.

2.5 Relatively invariant distribution

Let $\mathcal{B}_{X_{\mathbb{R}}}$ be sections of hyperfunctions on $X_{\mathbb{R}}$. From the diagram (4), we conclude the following on solutions of differential equations.

Theorem 8.

Tauchi defines [10] a distribution

$$T_2^l = z_1^l \frac{\partial^l}{\partial z_2^l} \delta(z_2, \overline{z}_2) = (z_1 \frac{\partial}{\partial z_2})^l \delta(z_2, \overline{z}_2)$$
(5)

on the real manifold $X_{\mathbb{R}} = \{(z_1, z_2, \overline{z_1}, \overline{z_2}) \mid z_1, z_2 \in \mathbb{C}\} \simeq \mathbb{R}^4$. Here $\delta(z_2, \overline{z_2})$ denotes the Dirac distribution $\frac{1}{-2\sqrt{-1}}\delta(\Re z_2, \Im z_2)$ on \mathbb{R}^4 . The support of T_2^l is the closure of $O_{11\mathbb{R}}$, that is, $O_{11\mathbb{R}} \sqcup O_{00\mathbb{R}}$.

Proposition 9. $T_2^l \in \{u \in \mathcal{B}_{X_{\mathbb{R}}} \mid Pu = 0, \forall P \in \mathcal{I}_{1l}\} \simeq \operatorname{Hom}_{\mathcal{D}}(\mathcal{M}_{1l}, \mathcal{B}_{X_{\mathbb{R}}})$. Especially, T_2^l is relatively invariant under \mathfrak{h}_1 with the character χ_l , and is relatively invariant under \mathfrak{h}_3 with the character χ .

Note that the relative invariance of T_2^l under \mathfrak{h}_3 is due to Tauchi. We regard Theorem 6 as a D-module counterpart of this result.

3 n = 3

3.1 D-module

In $X = \mathbb{C}^6$, we use coordinates $(z_1, z_2, z_3, z_4, z_5, z_6) \in X$. The ring of differential operators on X is denoted by $\mathcal{D} = \mathcal{D}_X = \mathbb{C}[z_1, z_2, z_3, z_4, z_5, z_6, \frac{\partial}{\partial z_1}, \frac{\partial}{\partial z_2}, \frac{\partial}{\partial z_3}, \frac{\partial}{\partial z_4}, \frac{\partial}{\partial z_5}, \frac{\partial}{\partial z_6}]$. Let $Y = \{z_3 = z_6 = 0\} \subset X = \mathbb{C}^6$, and define a simple holonomic D-module $\mathcal{B}_{Y|X}$ by

$$\mathcal{B}_{Y|X} = \mathcal{H}^2_{[Y]}(\mathcal{O}_X) = \mathcal{D}/(\mathcal{D}\frac{\partial}{\partial z_1} + \mathcal{D}\frac{\partial}{\partial z_2} + \mathcal{D}z_3 + \mathcal{D}\frac{\partial}{\partial z_4} + \mathcal{D}\frac{\partial}{\partial z_5} + \mathcal{D}z_6).$$

The standard generator represented by $1 \in \mathcal{D}$ is denoted by $\delta(z_3, z_6) \in \mathcal{B}_{Y|X}$. We define a section of $\mathcal{B}_{Y|X}$.

$$T_l = (z_2 \frac{\partial}{\partial z_3} + z_4 \frac{\partial}{\partial z_5})^l z_2 z_5 \delta(z_3, z_6)$$
(6)

$$= z_2^l \left(z_2 z_5 \frac{\partial^l}{\partial z_3^l} + l z_4 \frac{\partial^{l-1}}{\partial z_3^{l-1}} \right) \delta(z_3, z_6)$$

$$= (z_2 z_5 - z_3 z_4) (z_2 \frac{\partial}{\partial z_3})^l \delta(z_3, z_6).$$

$$(7)$$

Lemma 10. The annihilator ideal \mathcal{I}_{1l} of T_l is generated by the following elements:

$$\begin{array}{l} \frac{\partial}{\partial z_1}, \ z_6, \ z_2 \frac{\partial}{\partial z_2} + z_3 \frac{\partial}{\partial z_3}, \ z_2 \frac{\partial}{\partial z_2} + z_4 \frac{\partial}{\partial z_4} - 1 - l, \ z_2 \frac{\partial}{\partial z_4} + z_3 \frac{\partial}{\partial z_5}, \\ \\ \frac{\partial^{l+1}}{\partial z_2^{l+1}} \frac{\partial}{\partial z_4}, \ z_3^l \frac{\partial}{\partial z_4}, \ \frac{\partial^{l+2}}{\partial z_2^{l+2}} \frac{\partial}{\partial z_5}, \ z_3^{l+1} \frac{\partial}{\partial z_5}, \\ \\ \frac{\partial^2}{\partial z_4^2}, \ \frac{\partial^2}{\partial z_5^2}, \ \frac{\partial^2}{\partial z_4 \partial z_5}, \ z_4 \frac{\partial}{\partial z_4} + z_5 \frac{\partial}{\partial z_5} - 1. \end{array}$$

Proof. We define a left ideal \mathcal{I}'_l of \mathcal{D} generated by

$$\frac{\partial}{\partial z_1}, \ \frac{\partial}{\partial z_4}, \ \frac{\partial}{\partial z_5}, \ z_6, \ z_2 \frac{\partial}{\partial z_2} - l, \ \frac{\partial^{l+1}}{\partial z_2^{l+1}}, \ z_3 \frac{\partial}{\partial z_3} + l, \ z_3^l.$$

Using a similar argument to Theorem 6(i), we see that $\mathcal{M}'_l := \mathcal{D}/\mathcal{I}'_l$ forms a simple holonomic system. Let \mathcal{I}'' be the left ideal generated by elements in the statement of Lemma other than $z_2 \frac{\partial}{\partial z_4} + z_3 \frac{\partial}{\partial z_5}$. By computing the image of generators of left ideals, we obtain the property:

$$\mathcal{I}''z_4 \subset \mathcal{I}'_l \text{ and } \mathcal{I}''z_5 \subset \mathcal{I}'_{l+1},$$
(8)

$$\mathcal{I}'_l \frac{\partial}{\partial z_4} \subset \mathcal{I}'' \text{ and } \mathcal{I}'_{l+1} \frac{\partial}{\partial z_5} \subset \mathcal{I}''.$$
 (9)

For example, $(z_2 \frac{\partial}{\partial z_2} - l) \frac{\partial}{\partial z_4} = \frac{\partial}{\partial z_4} (z_2 \frac{\partial}{\partial z_2} + z_4 \frac{\partial}{\partial z_4} - l - 1) - z_4 \frac{\partial^2}{\partial z_4^2} \in \mathcal{I}''$. We will see that the map $\mathcal{D} \ni P \mapsto (Pz_4, Pz_5) \in \mathcal{D}^{\oplus 2}$ induces a D-module isomorphism

$$\phi: \mathcal{D}/\mathcal{I}'' \ni P \mapsto (Pz_4, Pz_5) \in \mathcal{M}'_l \oplus \mathcal{M}'_{l+1}.$$

Actually, ϕ is well-defined from the property (8). For a given $P_1, P_2 \in D$, put $P := P_1 \frac{\partial}{\partial z_4} + P_2 \frac{\partial}{\partial z_4} \in \mathcal{D}$, then $Pz_4 - P_1 \in \mathcal{I}'_l$ and $Pz_5 - P_2 \in \mathcal{I}'_{l+1}$. This proves the surjectivity of ϕ . For the injectivity of ϕ , take a $P \in \mathcal{D}$ such that $(P \mod \mathcal{I}'')$ is in the kernel of ϕ . We may write $P = Q_1 \frac{\partial}{\partial z_4} + Q_2 \frac{\partial}{\partial z_5} + Q_3$ with $Q_i \in \mathbb{C}[z_1, z_2, z_3, z_4, z_5, z_6, \frac{\partial}{\partial z_1}, \frac{\partial}{\partial z_2}, \frac{\partial}{\partial z_3}, \frac{\partial}{\partial z_6}]$. Then by assumption, $0 \equiv Pz_4 \equiv Q_1 + Q_3z_4$ mod \mathcal{I}'_l and $0 \equiv Pz_5 \equiv Q_2 + Q_3z_5 \mod \mathcal{I}'_{l+1}$. There exist $Q_4 \in \mathcal{I}'_l, Q_5 \in \mathcal{I}'_{l+1}$ such that $Q_1 = -Q_3z_4 + Q_4, Q_2 = -Q_3z_5 + Q_5$. Then $P = -Q_3(z_4 \frac{\partial}{\partial z_4} + z_5 \frac{\partial}{\partial z_5} - 1) + Q_4 \frac{\partial}{\partial z_4} + Q_5 \frac{\partial}{\partial z_5} \in \mathcal{I}''$ by the property (9).

Note that the image of the map ϕ is the sum of two simple holonomic systems. Therefore, $\mathcal{D}/(\mathcal{I}'' + D(z_2 \frac{\partial}{\partial z_4} + z_3 \frac{\partial}{\partial z_5}))$ is a simple holonomic system or zero. It is easy to see that the annihilator contains $\mathcal{I}'' + D(z_2 \frac{\partial}{\partial z_4} + z_3 \frac{\partial}{\partial z_5})$; then, maximality implies equality.

We define a complex Lie subalgebra $\mathfrak{h}_1 \subset \mathfrak{gl}(6, \mathbb{C})$ by

$$\mathfrak{h}_1 = \bigoplus_{j=1}^6 \mathbb{C}E_{1j} \oplus \bigoplus_{i=2}^6 \mathbb{C}E_{i6} \oplus \mathbb{C}(E_{22} + E_{33}) \oplus \mathbb{C}(E_{44} + E_{55}) \oplus \mathbb{C}(E_{22} + E_{44}) \oplus \mathbb{C}(E_{42} + E_{53}).$$

and its character $\chi_l : \mathfrak{h}_1 \to \mathbb{C}$ by

$$\chi_l(E_{22} + E_{33}) = 0, \ \chi_l(E_{22} + E_{44}) = -1 - l, \ \chi_l(E_{44} + E_{55}) = -1, \ \chi_l(E_{1j}) = \chi_l(E_{i6}) = \chi_l(E_{42} + E_{53}) = 0.$$

Then in terms of a Lie algebra homomorphism

$$\rho:\mathfrak{gl}(6,\mathbb{C})\ni E_{ij}\mapsto -z_j\frac{\partial}{\partial z_i}\in\mathcal{D},$$

Lemma 10 implies $(\rho - \chi_l)(\mathfrak{h}_1) \subset \mathcal{I}_{1l}$.

We define a Lie subalgebra $\mathfrak{h}_3 \subset \mathfrak{h}_1$ by

$$\mathfrak{h}_3 = \mathbb{C}(E_{11} + E_{22} + E_{33}) \oplus \mathbb{C}(E_{44} + E_{55} + E_{66}) \oplus \mathbb{C}(E_{42} + E_{53}) \oplus \mathbb{C}(E_{15} + E_{26}) \oplus \mathbb{C}E_{13} \oplus \mathbb{C}E_{46}.$$

We denote by χ the restriction of χ_l on \mathfrak{h}_3 since the restriction does not depend on l. We define the left ideal $\mathcal{I}_3 \subset \mathcal{D}$ generated by $(\rho - \chi)(\mathfrak{h}_3)$, that is,

$$z_1\frac{\partial}{\partial z_1} + z_2\frac{\partial}{\partial z_2} + z_3\frac{\partial}{\partial z_3}, z_4\frac{\partial}{\partial z_4} + z_5\frac{\partial}{\partial z_5} + z_6\frac{\partial}{\partial z_6} - 1, z_5\frac{\partial}{\partial z_1} + z_6\frac{\partial}{\partial z_2}, z_2\frac{\partial}{\partial z_4} + z_3\frac{\partial}{\partial z_5}, z_3\frac{\partial}{\partial z_1}, z_6\frac{\partial}{\partial z_4} + z_6\frac{\partial}{\partial z_5}, z_6\frac{\partial}{\partial z_6} - 1, z_6\frac{$$

Then $(\rho - \chi)(\mathfrak{h}_3) \subset (\rho - \chi_l)(\mathfrak{h}_{1l})$ for all $l \in \mathbb{Z}_{\geq 0}$, and $(\rho - \chi)(\mathfrak{h}_3) \subset \mathcal{I}_3 \subset \mathcal{I}_{1l}$. We have the same diagram (4).

Now we move on to solutions of these D-modules. We choose a real form $X_{\mathbb{R}} = \{(z_1, z_2, z_3, \overline{z_1}, \overline{z_2}, \overline{z_3}) \mid (z_1, z_2, z_3) \in \mathbb{C}^3\}$ of $X = \mathbb{C}^6$. Denote by $\mathcal{B}_{X_{\mathbb{R}}}$ the set of hyperfunctions on $X_{\mathbb{R}}$.

Lemma 11. Tauchi's distribution $T_0^l = (\overline{z}_1 \frac{\partial}{\partial \overline{z}_2} + z_2 \frac{\partial}{\partial z_3})^l |z_2|^2 \delta(z_3, \overline{z}_3) \in \mathcal{B}_{X_{\mathbb{R}}}$ is a solution of $\mathcal{D}/\mathcal{I}_{1l}$.

This shows T_0^l is relatively invariant under

$$\begin{aligned} \mathfrak{h}_{3\mathbb{R}} &= \mathfrak{h}_3 \cap \mathfrak{g}_{\mathbb{R}} \\ &= \{ a(E_{11} + E_{22} + E_{33}) + \overline{a}(E_{44} + E_{55} + E_{66}) + b(E_{42} + E_{53}) + \overline{b}(E_{15} + E_{26}) + cE_{13} + \overline{c}E_{46} \mid a, b, c \in \mathbb{C} \} \end{aligned}$$

with the character χ . Orbit decomposition is similar to n = 2, so we omit its description.

We would like to express our sincere gratitude to the referee for their valuable comments and suggestions, which have greatly improved the quality and clarity of this paper.

References

- Ryoshi Hotta, Kiyoshi Takeuchi and Toshiyuki Tanisaki, D-modules, perverse sheaves, and representation theory, Progr. Math., 236 Birkhäuser Boston, Inc., Boston, MA, 2008, xii+407 pp.
- [2] Masaki Kashiwara, D-modules and microlocal calculus, translated from the 2000 Japanese original by Mutsumi Saito, Transl. Math. Monogr., 217, Amer. Math. Soc., Providence, RI, 2003.
- [3] M. T. Kosters and G. van Dijk, Spherical distributions on the pseudo-Riemannian space $SL(n,\mathbb{R})/GL(n-1,\mathbb{R})$, J. Funct. Anal. 68 (1986), no. 2, 168–213.
- [4] Atsutaka Kowata, On the construction of spherical hyperfunctions on R^{p+q}, Hiroshima Math. J. 21 (1991), 301–334.

- [5] Atsutaka Kowata, Spherical hyperfunctions on the tangent space of symmetric spaces, Hiroshima Math. J. 21 (1991), 401–418
- [6] Thierry Levasseur and J. Toby Stafford, Invariant differential operators on the tangent space of some symmetric spaces, Annales de l'institut Fourier, tome 49, no 6 (1999), p. 1711–1741.
- [7] Hiroyuki Ochiai, Invariant functions on the tangent space of a rank one semisimple symmetric space, J. Fac. Sci. Univ. Tokyo Sect. IA Math.39(1992), no.1, 17–31.
- [8] Hiroyuki Ochiai, Invariant distributions on a non-isotropic pseudo-Riemannian symmetric space of rank one, Indag. Math. (N.S.) 16 (2005), no. 3-4, 631–638.
- [9] Toshiyuki Tanisaki, D-modules and representation theory. (English summary) Lie theory and representation theory, 177–219. Surv. Mod. Math., 2 International Press, Somerville, MA, 2012.
- [10] Taito Tauchi, Dimension of the space of intertwining operators from degenerate principal series representations, Selecta Math. (N.S.) 24 (2018), no. 4, 3649–3662.
- [11] Taito Tauchi, Relationship between orbit decomposition on the flag varieties and multiplicities of induced representations, Proc. Japan Acad. Ser. A Math. Sci. 95 (2019), no. 7, 75–79.
- [12] Taito Tauchi, Multiplicity of a degenerate principal series for homogeneous spaces with infinite orbits, Proc. Amer. Math. Soc. 150 (2022), no. 2, 849–856.
- [13] Gerrit van Dijk, Invariant eigendistributions on the tangent space of a rank one semisimple symmetric space, Math. Ann. 268 (1984), no. 3, 405–416.
- [14] Gerrit van Dijk, On generalized Gel'fand pairs, Proc. Japan Acad. Ser. A Math. Sci. 60 (1984), no. 1, 30–34.
- [15] Gerrit van Dijk, On a class of generalized Gel'fand pairs, Math. Z. 193 (1986), no. 4, 581–593.
- [16] Gerrit van Dijk, Harmonic analysis on rank one symmetric spaces, Lecture Notes in Phys., 261 Springer-Verlag, Berlin, 1986, 244–252.
- [17] Gerrit van Dijk, Gelfand pairs and beyond, COE Lect. Note, 11, Math-for-Ind. (MI) Lect. Note Ser. Kyushu University, Faculty of Mathematics, Fukuoka, 2008, ii+60 pp.
- [18] Gerrit van Dijk, $(GL(n+1,\mathbb{R}), GL(n,\mathbb{R}))$ is a generalized Gelfand pair, Russ. J. Math. Phys. 15 (2008), no. 4, 548–551.
- [19] Gerrit van Dijk, (U(p,q), U(p-1,q)) is a generalized Gelfand pair, Math. Z. 261 (2009), no. 3, 525–529.
- [20] Gerrit van Dijk, Introduction to Harmonic Analysis and Generalized Gelfand Pairs, Walter de Gruyter, Berlin New York, 2009.
- [21] G. van Dijk and M. Poel, The Plancherel formula for the pseudo-Riemannian space $SL(n,\mathbb{R})/GL(n-1,\mathbb{R})$, Compositio Math. 58 (1986), no. 3, 371–397.
- [22] G. van Dijk and M. Poel, The irreducible unitary $GL(n-1,\mathbb{R})$ -spherical representations of $SL(n,\mathbb{R})$, Compositio Math. 73 (1990), no. 1, 1–30.

Hiroyuki Ochiai Institute of Mathematics for Industry, Kyushu University 744 Motooka Fukuoka 819-0395 Japan ochiai@imi.kyushu-u.ac.jp