Toughness and A_{α} -spectral radius in graphs $*$

Sizhong Zhou¹[†], Yuli Zhang²[†], Tao Zhang³, Hongxia Liu⁴

1. School of Science, Jiangsu University of Science and Technology,

Zhenjiang, Jiangsu 212100, China

2. School of Science, Dalian Jiaotong University,

Dalian, Liaoning 116028, China

3. School of Economics and management, Jiangsu University of Science and Technology,

Zhenjiang, Jiangsu 212100, China

4. School of Mathematics and Information Science, Yantai University,

Yantai, Shandong 264005, China

Abstract

Let $\alpha \in [0, 1)$, and let G be a connected graph of order n with $n \ge f(\alpha)$, where $f(\alpha) = 6$ for $\alpha \in [0, \frac{2}{3}]$ and $f(\alpha) = \frac{4}{1-\alpha}$ for $\alpha \in (\frac{2}{3}, 1)$. A graph G is said to be t-tough if $|S| \geq tc(G-S)$ for each subset S of $V(G)$ with $c(G-S) \geq 2$, where $c(G-S)$ is the number of connected components in $G - S$. The A_{α} -spectral radius of G is denoted by $\rho_{\alpha}(G)$. In this paper, it is verified that G is a 1-tough graph unless $G = K_1 \vee (K_{n-2} \cup K_1)$ if $\rho_\alpha(G) \ge \rho_\alpha(K_1 \vee (K_{n-2} \cup K_1)),$ where $\rho_{\alpha}(K_1 \vee (K_{n-2} \cup K_1))$ equals the largest root of $x^3 - ((\alpha + 1)n + \alpha - 3)x^2 + (\alpha n^2 + (\alpha^2 - \alpha - 3)x^2)$ 1)n – 2 $\alpha + 1$)x – $\alpha^2 n^2 + (3\alpha^2 - \alpha + 1)n - 4\alpha^2 + 5\alpha - 3 = 0$. Further, we present an A_{α} -spectral radius condition for a graph to be a t -tough graph.

Keywords: graph; toughness; A_{α} -spectral radius. (2020) Mathematics Subject Classification: 05C50

1 Introduction

Let G be an undirected simple graph with vertex set $V(G) = \{v_1, v_2, \ldots, v_n\}$ and edge set $E(G)$. The order of G is the number $n = |V(G)|$ of its vertices and its size is the number $m = e(G)$ of its edges. A graph G is called trivial if $|V(G)| = 1$. For arbitrary $v \in V(G)$, the degree of v in G is defined as the number of edges which are adjacent to v and denoted by $d_G(v)$. For a given subset S of $V(G)$, we use G[S] to denote the subgraph of G induced by S, and write $G - S$ for $G[V(G) \setminus S]$. The complement of a graph G is a graph \overline{G} with the same vertex set as G, in which any two distinct vertices are adjacent if and only if they are nonadjacent in G . Let G_1 and G_2 be two vertex-disjoint graphs. We use $G_1 \cup G_2$ to denote the disjoint union of G_1 and G_2 . The join $G_1 \vee G_2$ is the graph

[∗]This work was supported by the Natural Science Foundation of Shandong Province, China (ZR2023MA078)

[†]Corresponding author. E-mail address: zsz cumt@163.com (S. Zhou)

[‡]Corresponding author. E-mail address: zhangyuli djtu@126.com (Y. Zhang)

formed from $G_1 \cup G_2$ by adding all possible edges between $V(G_1)$ and $V(G_2)$. We denote by K_n a complete graph of order n.

Let t be a nonnegative real number. A graph G is said to be t-tough if $|S| \geq tc(G-S)$ for each subset S of $V(G)$ with $c(G-S) \geq 2$, where $c(G-S)$ is the number of connected components in $G-S$. The toughness $t(G)$ of G is the largest real number t for which G is t-tough, or is ∞ if G is complete. This concept was first introduced by Chvátal [\[1\]](#page-8-0) in 1973. Some results on toughness of graphs can be found in $[2-6]$ $[2-6]$.

Given a graph G of order n, the adjacency matrix $A(G)$ of G is the $n \times n$ matrix in which entry a_{ij} is 1 or 0 according to whether v_i and v_j are adjacent or not. The eigenvalues of G are the eigenvalues of its adjacency matrix $A(G)$. Let $\lambda_1(G) \geq \lambda_2(G) \geq \cdots \geq \lambda_n(G)$ be its eigenvalues in nonincreasing order. Note that the adjacency spectral radius of G is equal to $\lambda_1(G)$, written as $\rho(G)$.

Let $D(G)$ denote the diagonal matrix of vertex degree of G. Then $L(G) = D(G) - A(G)$ and $Q(G) = D(G) + A(G)$ are called the Laplacian matrix and signless Laplacian matrix of G, respectively. For any $\alpha \in [0, 1]$, Nikiforov [\[7\]](#page-9-1) introduced the A_{α} -matrix of G as

$$
A_{\alpha}(G) = \alpha D(G) + (1 - \alpha)A(G).
$$

It is easy to see that $A_{\alpha}(G) = A(G)$ if $\alpha = 0$, and $A_{\alpha}(G) = \frac{1}{2}Q(G)$ if $\alpha = \frac{1}{2}$. Note that $A_{\alpha}(G)$ is a real symmetric nonnegative matrix. Hence, the eigenvalues of $A_{\alpha}(G)$ are real, which can be indexed in nonincreasing order as $\lambda_1(A_\alpha(G)) \geq \lambda_2(A_\alpha(G)) \geq \cdots \geq \lambda_n(A_\alpha(G))$. The largest eigenvalue $\lambda_1(A_\alpha(G))$ is called the A_α -spectral radius of G, and denoted by $\rho_\alpha(G)$. Namely, $\rho_\alpha(G) = \lambda_1(A_\alpha(G))$. For some interesting spectral properties of $A_{\alpha}(G)$, we refer the reader to [\[7–](#page-9-1)[14\]](#page-9-2).

Many authors [\[15](#page-9-3)[–20\]](#page-10-0) obtained some results on spectral radius in graphs. Brouwer [\[21\]](#page-10-1) investigated the relationship between toughness and eigenvalues and claimed that for any connected d-regular graph $G, t(G) > \frac{d}{\lambda} - 2$, and he further conjectured that the lower bound $\frac{d}{\lambda} - 2$ can be improved to $\frac{d}{\lambda} - 1$, where $\lambda = \max_{2 \leq i \leq n} |\lambda_i|$. Later, Gu [\[22\]](#page-10-2) strengthened Brouwer's result and claimed the lower bound $\frac{d}{\lambda}$ – 2 can be improved to $\frac{d}{\lambda}$ – $\sqrt{2}$. Very recently, Gu [\[23\]](#page-10-3) confirmed Brouwer's conjecture. Fan, Lin and Lu [\[24\]](#page-10-4) presented two adjacency spectral radius conditions for the existence of a t-tough graph. It is natural and interesting to find a sufficient condition for a graph to be t-tough in view of its A_{α} -spectral radius. In this paper, we first show a A_{α} -spectral radius condition to guarantee that a graph is 1-tough.

Theorem 1.1. Let $\alpha \in [0,1)$, and let G be a connected graph of order n with $n \ge f(\alpha)$, where

$$
f(\alpha) = \begin{cases} 6, & if \alpha \in [0, \frac{2}{3}]; \\ \frac{4}{1-\alpha}, & if \alpha \in (\frac{2}{3}, 1). \end{cases}
$$

If $\rho_{\alpha}(G) \geq \rho_{\alpha}(K_1 \vee (K_{n-2} \cup K_1))$, then G is a 1-tough graph unless $G = K_1 \vee (K_{n-2} \cup K_1)$, where $\rho_{\alpha}(K_1 \vee (K_{n-2} \cup K_1))$ equals the largest root of $x^3 - ((\alpha + 1)n + \alpha - 3)x^2 + (\alpha n^2 + (\alpha^2 - \alpha - 1)n (2\alpha + 1)x - \alpha^2 n^2 + (3\alpha^2 - \alpha + 1)n - 4\alpha^2 + 5\alpha - 3 = 0.$

Further, it is natural and interesting to find an A_{α} -spectral radius condition to ensure that a graph is a t-tough graph. Next, we present an A_{α} -spectral radius condition for a graph to be t-tough.

Theorem 1.2. Let $\alpha \in \left[\frac{1}{2}, \frac{3}{4}\right)$, and let t be a positive integer. If G be a connected graph of order $n \ge \max\{4t^2 + 6t + 1, \frac{12t(1-\alpha)-2\alpha+1}{3-4\alpha}\}$ $\frac{-\alpha-2\alpha+1}{3-4\alpha}$ with $\rho_{\alpha}(G) \geq \rho_{\alpha}(K_{2t-1} \vee (K_{n-2t} \cup K_1)),$ then G is a t-tough graph unless $G = K_{2t-1} \vee (K_{n-2t} \cup K_1).$

2 Preliminaries

In this section, we put forward some necessary lemmas, which play a key role in proving our main results.

Lemma 2.1 (Nikiforov [\[7\]](#page-9-1)). Let K_n be a complete graph of order n. Then

$$
\rho_{\alpha}(K_n)=n-1.
$$

Lemma 2.2 (Nikiforov [\[7\]](#page-9-1)). If G is a connected graph, and H is a proper subgraph of G, then

$$
\rho_{\alpha}(G) > \rho_{\alpha}(H).
$$

Lemma 2.3 (Zhao, Huang and Wang [\[14\]](#page-9-2)). Let $\alpha \in [0,1)$, and let $n_1 \geq n_2 \geq \cdots \geq n_t$ be positive integers with $n = \sum_{i=1}^{t} n_i + s$ and $n_1 \leq n - s - t + 1$. Then

$$
\rho_{\alpha}(K_s \vee (K_{n_1} \cup K_{n_2} \cup \cdots \cup K_{n_t})) \leq \rho_{\alpha}(K_s \vee (K_{n-s-t+1} \cup (t-1)K_1)),
$$

where the equality holds if and only if $(n_1, n_2, \ldots, n_t) = (n - s - t + 1, 1, \ldots, 1)$.

Let M be a real matrix whose rows and columns are indexed by $V = \{1, 2, \ldots, n\}$. Assume that M, with respect to the partition $\pi : V = V_1 \cup V_2 \cup \cdots \cup V_t$, can be written as

$$
M = \begin{pmatrix} M_{11} & M_{12} & \cdots & M_{1t} \\ M_{21} & M_{22} & \cdots & M_{2t} \\ \vdots & \vdots & \ddots & \vdots \\ M_{t1} & M_{t2} & \cdots & M_{tt} \end{pmatrix},
$$

where M_{ij} denotes the submatrix (block) of M formed by rows in V_i and columns in V_j . Let b_{ij} denote the average row sum of M_{ij} , namely, b_{ij} is the sum of all entries in M_{ij} divided by the number of rows. Then matrix $M_{\pi} = (b_{ij})$ is called the *quotient matrix* of M. If the row sum of every block M_{ij} is a constant, then the partition is *equitable*.

Lemma 2.4 (Haemers [\[25\]](#page-10-5), You, Yang, So and Xi [\[26\]](#page-10-6)). Let M be a real matrix with an equitable partition π , and let M_{π} be the corresponding quotient matrix. Then each eigenvalue of M_{π} is an eigenvalue of M. Furthermore, if M is nonnegative, then the spectral radius of M_{π} equals the spectral radius of M.

Lemma 2.5 (Alhevaz, Baghipur, Ganie and Das [\[27\]](#page-10-7)). Let G be a graph of order n with m edges, with no isolated vertices and let $\alpha \in [\frac{1}{2}, 1]$. Then

$$
\rho_{\alpha}(G) \le \frac{2m(1-\alpha)}{n-1} + \alpha n - 1.
$$

If $\alpha \in (\frac{1}{2}, 1)$ and G is connected, equality holds if and only if $G = K_n$.

3 The proof of Theorem 1.1

In this section, we verify Theorem 1.1, which establishes an A_{α} -spectral radius condition for a graph to be 1-tough.

Proof of Theorem 1.1. Suppose to the contrary that G is not a 1-tough graph, there exists some nonempty subset $S \subseteq V(G)$ such that $c(G-S) \geq |S|+1$. Let $|S| = s$. Then G is a spanning subgraph of $G_s^1 = K_s \vee (K_{n_1} \cup K_{n_2} \cup \cdots \cup K_{n_{s+1}})$ for some integers $n_1 \geq n_2 \geq \cdots \geq n_{s+1}$ with $\sum_{i=1}^{s+1}$ $\sum_{i=1}^{\infty} n_i = n - s.$ According to Lemma 2.2, we have

$$
\rho_{\alpha}(G) \le \rho_{\alpha}(G_s^1) \tag{3.1}
$$

with equality if and only if $G = G_s^1$.

Let $G_s^2 = K_s \vee (K_{n-2s} \cup sK_1)$. Using Lemma 2.3, we get

$$
\rho_{\alpha}(G_s^1) \le \rho_{\alpha}(G_s^2),\tag{3.2}
$$

where the equality holds if and only if $(n_1, n_2, \ldots, n_{s+1}) = (n-2s, 1, \ldots, 1)$. Let $\varphi(x) = x^3 - ((\alpha +$ $1)n + \alpha - 3x^2 + (\alpha n^2 + (\alpha^2 - \alpha - 1)n - 2\alpha + 1)x - \alpha^2 n^2 + (3\alpha^2 - \alpha + 1)n - 4\alpha^2 + 5\alpha - 3$. Then $\rho_{\alpha}(K_1 \vee (K_{n-2} \cup K_1))$ equals the largest root of $\varphi(x) = 0$. We are to verify the following claim.

Claim 1. If $\alpha \in [0, 1)$ and $n \ge f(\alpha)$, then we obtain

$$
\rho_{\alpha}(G_s^2) \le \rho_{\alpha}(K_1 \vee (K_{n-2} \cup K_1))
$$

with equality if and only if $G_s^2 = K_1 \vee (K_{n-2} \cup K_1)$. *Proof.* Recall that $G_s^2 = K_s \vee (K_{n-2s} \cup sK_1)$. Obviously, $1 \leq s \leq \frac{n-1}{2}$.

Consider the partition $V(G_s^2) = V(sK_1) \cup V(K_{n-2s}) \cup V(K_s)$. The corresponding quotient matrix of $A_{\alpha}(G_s^2)$ is equal to

$$
B_1 = \begin{pmatrix} \alpha s & 0 & (1 - \alpha)s \\ 0 & n + (\alpha - 2)s - 1 & (1 - \alpha)s \\ (1 - \alpha)s & (1 - \alpha)(n - 2s) & \alpha n - \alpha s + s - 1 \end{pmatrix}.
$$

Then the characteristic polynomial of B_1 is

$$
\varphi_{B_1}(x) = x^3 - ((\alpha + 1)n + (\alpha - 1)s - 2)x^2 + (\alpha n^2 + (\alpha^2 s - \alpha - 1)n - s^2 - (2\alpha - 1)s + 1)x - \alpha^2 sn^2 + (2\alpha^2 - 2\alpha + 1)s^2 n + (\alpha^2 + \alpha)sn - (3\alpha^2 - 5\alpha + 2)s^3 - (\alpha^2 - \alpha + 1)s^2 - \alpha s.
$$
 (3.3)

Since the partition $V(G_s^2) = V(sK_1) \cup V(K_{n-2s}) \cup V(K_s)$ is equitable, it follows from Lemma 2.4 that $\rho_{\alpha}(G_s^2)$ is the largest root of $\varphi_{B_1}(x) = 0$. Namely, $\varphi_{B_1}(\rho_{\alpha}(G_s^2)) = 0$. Let $\eta_1 = \rho_{\alpha}(G_s^2) \ge \eta_2 \ge \eta_3$ be the three roots of $\varphi_{B_1}(x) = 0$ and $Q = \text{diag}(s, n - 2s, s)$. One checks that

$$
Q^{\frac{1}{2}}B_1Q^{-\frac{1}{2}} = \begin{pmatrix} \alpha s & 0 & (1-\alpha)s \\ 0 & n + (\alpha-2)s - 1 & (1-\alpha)s^{\frac{1}{2}}(n-2s)^{\frac{1}{2}} \\ (1-\alpha)s & (1-\alpha)s^{\frac{1}{2}}(n-2s)^{\frac{1}{2}} & \alpha n - \alpha s + s - 1 \end{pmatrix}
$$

is symmetric, and also contains

$$
\left(\begin{array}{cc} \alpha s & 0\\ 0 & n + (\alpha - 2)s - 1 \end{array}\right)
$$

as its submatrix. Since $Q^{\frac{1}{2}}B_1Q^{-\frac{1}{2}}$ and B_1 have the same eigenvalues, the Cauchy interlacing theorem (cf. [\[25\]](#page-10-5)) implies that

$$
\eta_2 \le n + (\alpha - 2)s - 1 < n - 2. \tag{3.4}
$$

If $s = 1$, then $G_s^2 = K_1 \vee (K_{n-2} \cup K_1)$ and $\varphi_{B_1}(x) = \varphi(x)$. And so $\rho_\alpha(G_s^2) = \rho_\alpha(K_1 \vee (K_{n-2} \cup K_1))$. In what follows, we always assume $s \geq 2$.

Note that K_{n-1} is a proper subgraph of $K_1 \vee (K_{n-2} \cup K_1)$. In terms of [\(3.4\)](#page-4-0), Lemmas 2.1 and 2.2, we get

$$
\rho_{\alpha}(K_1 \vee (K_{n-2} \cup K_1)) > \rho_{\alpha}(K_{n-1}) = n-2 > \eta_2.
$$
\n(3.5)

Write $\theta(n) = \rho_{\alpha}(K_1 \vee (K_{n-2} \cup K_1))$. Note that $\varphi(\theta(n)) = 0$. A simple calculation yields that

$$
\varphi_{B_1}(\theta(n)) = \varphi_{B_1}(\theta(n)) - \varphi(\theta(n))
$$

= $(s - 1)((1 - \alpha)(\theta(n))^2 + (\alpha^2 n - 2\alpha - s)\theta(n) - \alpha^2 n^2$
+ $(2\alpha^2 - 2\alpha + 1)sn + (3\alpha^2 - \alpha + 1)n - (3\alpha^2 - 5\alpha + 2)s^2$
- $(4\alpha^2 - 6\alpha + 3)s - 4\alpha^2 + 5\alpha - 3).$ (3.6)

The following proof will be divided into two cases.

Case 1. $0 \leq \alpha \leq \frac{2}{3}$.

Let $h(x) = (1 - \alpha)x^2 + (\alpha^2 n - 2\alpha - s)x - \alpha^2 n^2 + (2\alpha^2 - 2\alpha + 1)s^2 + (3\alpha^2 - \alpha + 1)n - (3\alpha^2 - \alpha + 1)$ $(5\alpha+2)s^2-(4\alpha^2-6\alpha+3)s-4\alpha^2+5\alpha-3$ be a real function in x. Then the symmetry axis of $h(x)$ is $x = \frac{-\alpha^2 n + 2\alpha + s}{2(1-\alpha)}$. Obviously, $h(x)$ is increasing when $x \ge \frac{-\alpha^2 n + 2\alpha + s}{2(1-\alpha)}$. By plugging the value $\theta(n)$ into x of $h(x)$, we obtain

$$
h(\theta(n)) = (1 - \alpha)(\theta(n))^2 + (\alpha^2 n - 2\alpha - s)\theta(n) - \alpha^2 n^2
$$

+ $(2\alpha^2 - 2\alpha + 1)sn + (3\alpha^2 - \alpha + 1)n - (3\alpha^2 - 5\alpha + 2)s^2$
- $(4\alpha^2 - 6\alpha + 3)s - 4\alpha^2 + 5\alpha - 3$

and it follows from [\(3.6\)](#page-4-1) that

$$
\varphi_{B_1}(\theta(n)) = (s-1)h(\theta(n)).\tag{3.7}
$$

Note that $n - 2s \ge 1$. We have $n \ge 2s + 1$. Together with $s \ge 2$ and [\(3.5\)](#page-4-2), we conclude

$$
\frac{-\alpha^2 n + 2\alpha + s}{2(1 - \alpha)} < n - 2 < \rho_\alpha(K_1 \vee (K_{n-2} \cup K_1)) = \theta(n),
$$

and so

$$
h(\theta(n)) > h(n-2)
$$

= $(1 - \alpha)(n-2)^2 + (\alpha^2 n - 2\alpha - s)(n-2) - \alpha^2 n^2$
+ $(2\alpha^2 - 2\alpha + 1)sn + (3\alpha^2 - \alpha + 1)n - (3\alpha^2 - 5\alpha + 2)s^2$
 $- (4\alpha^2 - 6\alpha + 3)s - 4\alpha^2 + 5\alpha - 3$
= $(1 - \alpha)n^2 + ((2\alpha^2 - 2\alpha)s + \alpha^2 + \alpha - 3)n - (3\alpha^2 - 5\alpha + 2)s^2$
 $- (4\alpha^2 - 6\alpha + 1)s - 4\alpha^2 + 5\alpha + 1.$ (3.8)

Let $p(x, s) = (1 - \alpha)x^2 + ((2\alpha^2 - 2\alpha)s + \alpha^2 + \alpha - 3)x - (3\alpha^2 - 5\alpha + 2)s^2 - (4\alpha^2 - 6\alpha + 1)s - 4\alpha^2 + 5\alpha + 1$ be a real function in x . According to (3.8) , we get

$$
h(\theta(n)) > p(n, s). \tag{3.9}
$$

Clearly, $p(x, s)$ is increasing when $x \geq -\frac{(2\alpha^2 - 2\alpha)s + \alpha^2 + \alpha - 3}{2(1 - \alpha)}$ $\frac{2\alpha s + \alpha^2 + \alpha - 3}{2(1-\alpha)}$. If $s \geq 3$, then we can verify that

$$
-\frac{(2\alpha^2 - 2\alpha)s + \alpha^2 + \alpha - 3}{2(1 - \alpha)} < 2s + 1 \le n,
$$

and so

$$
p(n,s) \ge p(2s+1,s)
$$

= $(1 - \alpha)(2s + 1)^2 + ((2\alpha^2 - 2\alpha)s + \alpha^2 + \alpha - 3)(2s + 1)$
 $- (3\alpha^2 - 5\alpha + 2)s^2 - (4\alpha^2 - 6\alpha + 1)s - 4\alpha^2 + 5\alpha + 1$
= $(s^2 - 3)\alpha^2 - (3s^2 - 2s - 5)\alpha + 2s^2 - 3s - 1$
 $\ge \frac{4}{9}(s^2 - 3) - \frac{2}{3}(3s^2 - 2s - 5) + 2s^2 - 3s - 1$
= $\frac{4s^2 - 15s + 9}{9}$
 ≥ 0 ,

where the last two inequalities hold from $\frac{3s^2-2s-5}{2(s^2-3)} > \frac{2}{3} \ge \alpha$ and $s \ge 3$, respectively.

If $s = 2$, then $-\frac{(2\alpha^2 - 2\alpha)s + \alpha^2 + \alpha - 3}{2(1-\alpha)} = -\frac{5\alpha^2 - 3\alpha - 3}{2(1-\alpha)} < 6 = f(\alpha) \le n$ due to $0 \le \alpha \le \frac{2}{3}$. Hence, for $0 \leq \alpha \leq \frac{2}{3}$ and $n \geq f(\alpha) = 6$, we get

$$
p(n, 2) \ge p(6, 2) = 6\alpha^2 - 17\alpha + 9 > 0.
$$

Thus, we obtain $p(n, s) > 0$ for $2 \le s \le \frac{n-1}{2}$. Together with [\(3.7\)](#page-4-4), [\(3.9\)](#page-5-0) and $s \ge 2$, we have

$$
\varphi_{B_1}(\theta(n)) = (s-1)h(\theta(n)) > (s-1)p(n,s) \ge 0.
$$

Recall that $\rho_{\alpha}(G_s^2)$ is the largest root of $\varphi_{B_1}(x) = 0$. As $\eta_2 < n - 2 < \rho_{\alpha}(K_1 \vee (K_{n-2} \cup K_1)) = \theta(n)$ (see [\(3.5\)](#page-4-2)), it follows from [\(3.6\)](#page-4-1), $\frac{-\alpha^2 n+2\alpha+s}{2(1-\alpha)} < n-2 < \theta(n)$ and $\varphi_{B_1}(\theta(n)) > 0$ that

$$
\rho_{\alpha}(G_s^2) < \theta(n) = \rho_{\alpha}(K_1 \vee (K_{n-2} \cup K_1))
$$

for $2 \leq s \leq \frac{n-1}{2}$. **Case 2.** $\frac{2}{3} < \alpha < 1$.

In terms of [\(3.3\)](#page-3-0), we obtain

$$
\varphi_{B_1}(n-2) = (n-2)^3 - ((\alpha + 1)n + (\alpha - 1)s - 2)(n-2)^2
$$

+
$$
(\alpha n^2 + (\alpha^2 s - \alpha - 1)n - s^2 - (2\alpha - 1)s + 1)(n-2)
$$

-
$$
\alpha^2 sn^2 + (2\alpha^2 - 2\alpha + 1)s^2 n + (\alpha^2 + \alpha)sn
$$

-
$$
(3\alpha^2 - 5\alpha + 2)s^3 - (\alpha^2 - \alpha + 1)s^2 - \alpha s
$$

=
$$
(3\alpha - 2)(1 - \alpha)s^3 + ((2\alpha^2 - 2\alpha)n - \alpha^2 + \alpha + 1)s^2
$$

+
$$
((1 - \alpha)n^2 - (\alpha^2 - 3\alpha + 3)n - \alpha + 2)s
$$

$$
+(\alpha - 1)n^2 - (2\alpha - 3)n - 2
$$

$$
:= \Phi(s, n).
$$

Thus, we obtain

$$
\frac{\partial \Phi(s,n)}{\partial s} = 3(3\alpha - 2)(1 - \alpha)s^2 + 2((2\alpha^2 - 2\alpha)n - \alpha^2 + \alpha + 1)s + (1 - \alpha)n^2 - (\alpha^2 - 3\alpha + 3)n - \alpha + 2.
$$

Note that $\frac{2}{3} < \alpha < 1$ and $n \ge f(\alpha) = \frac{4}{1-\alpha}$. By a simple calculation, we have

$$
\frac{\partial \Phi(s, n)}{\partial s}\Big|_{s=2} = (1 - \alpha)n^2 + (7\alpha^2 - 5\alpha - 3)n - 40\alpha^2 + 63\alpha - 18
$$

\n
$$
\geq (1 - \alpha) \left(\frac{4}{1 - \alpha}\right)^2 + (7\alpha^2 - 5\alpha - 3) \left(\frac{4}{1 - \alpha}\right) - 40\alpha^2 + 63\alpha - 18
$$

\n
$$
= \frac{1}{1 - \alpha} (40\alpha^3 - 75\alpha^2 + 61\alpha - 14)
$$

\n
$$
> 0,
$$

and

$$
\frac{\partial \Phi(s,n)}{\partial s}\Big|_{s=\frac{n-1}{2}} = \frac{1}{4}((1-\alpha)(\alpha-2)n^2 + (2\alpha^2 - 6\alpha - 4)n - 5\alpha^2 + 7\alpha + 6)
$$

$$
\leq \frac{1}{4}\left((1-\alpha)(\alpha-2)\left(\frac{4}{1-\alpha}\right)^2 + (2\alpha^2 - 6\alpha - 4)\left(\frac{4}{1-\alpha}\right) - 5\alpha^2 + 7\alpha + 6\right)
$$

$$
= \frac{1}{4(1-\alpha)}(5\alpha^3 - 4\alpha^2 - 7\alpha - 42)
$$

<0.

This implies that $\varphi_{B_1}(n-2) = \Phi(s, n) \ge \min\left\{\Phi(2, n), \Phi\left(\frac{n-1}{2}, n\right)\right\}$ because the leading coefficient of $\Phi(s,n)$ (view as a cubic polynomial of s) is positive, and $2 \leq s \leq \frac{n-1}{2}$. According to $\frac{2}{3} < \alpha < 1$ and $n \ge f(\alpha) = \frac{4}{1-\alpha}$, we obtain

$$
\Phi(2,n) = (1 - \alpha)n^2 + (6\alpha^2 - 4\alpha - 3)n - 28\alpha^2 + 42\alpha - 10
$$

\n
$$
\geq (1 - \alpha) \left(\frac{4}{1 - \alpha}\right)^2 + (6\alpha^2 - 4\alpha - 3) \left(\frac{4}{1 - \alpha}\right) - 28\alpha^2 + 42\alpha - 10
$$

\n
$$
= \frac{2}{1 - \alpha} (14\alpha^3 - 23\alpha^2 + 18\alpha - 3)
$$

\n
$$
> 0,
$$

and

$$
\Phi\left(\frac{n-1}{2},n\right) = \frac{1}{8}((\alpha^2 - 3\alpha + 2)n^3 - (5\alpha^2 - 19\alpha + 16)n^2
$$

+ $(3\alpha^2 - 25\alpha + 34)n + \alpha^2 + \alpha - 20)$

$$
\geq \frac{1}{8}((\alpha^2 - 3\alpha + 2)\left(\frac{4}{1-\alpha}\right)^3 - (5\alpha^2 - 19\alpha + 16)\left(\frac{4}{1-\alpha}\right)^2
$$

+ $(3\alpha^2 - 25\alpha + 34)\left(\frac{4}{1-\alpha}\right) + \alpha^2 + \alpha - 20)$

$$
=\frac{1}{8(1-\alpha)^2}(\alpha^4 - 13\alpha^3 + 11\alpha^2 + 45\alpha - 12)
$$

>0.

Consequently, we conclude $\varphi_{B_1}(n-2) \ge \min\left\{\Phi(2,n), \Phi\left(\frac{n-1}{2}, n\right)\right\} > 0$ for $2 \le s \le \frac{n-1}{2}$. As η_2 $n-2 < \rho_{\alpha}(K_1 \vee (K_{n-2} \cup K_1))$ (see [\(3.5\)](#page-4-2)), we have

$$
\rho_{\alpha}(G_s^2) < n - 2 < \rho_{\alpha}(K_1 \vee (K_{n-2} \cup K_1))
$$

for $2 \leq s \leq \frac{n-1}{2}$. This verifies Claim 1.

In terms of (3.1) , (3.2) and Claim 1, we obtain

$$
\rho_{\alpha}(G) \le \rho_{\alpha}(K_1 \vee (K_{n-2} \cup K_1))
$$

with equality if and only if $G = K_1 \vee (K_{n-2} \cup K_1)$. Which is a contradiction to the A_{α} -spectral condition of Theorem 1.1. This completes the proof of Theorem 1.1. \Box

4 The proof of Theorem 1.2

In this section, we prove Theorem 1.2, which provides an A_{α} -spectral radius condition for a graph to be t-tough.

Proof of Theorem 1.2. Suppose to the contrary that G is not a t-tough graph, there exists some nonempty subset $S \subseteq V(G)$ such that $tc(G - S) - 1 \geq |S|$. Let $|S| = s$ and $c(G - S) = c$. Then G is a spanning subgraph of $G_1 = K_{tc-1} \vee (K_{n_1} \cup K_{n_2} \cup \cdots \cup K_{n_c})$ for some integers $n_1 \geq n_2 \geq \cdots \geq n_c$ with \sum^c $\sum_{i=1} n_i = n - tc + 1$. In terms of Lemma 2.2, we conclude

$$
\rho_{\alpha}(G) \le \rho_{\alpha}(G_1) \tag{4.1}
$$

with equality if and only if $G = G_1$.

Let $G_2 = K_{tc-1} \vee (K_{n-(t+1)c+2} \cup (c-1)K_1)$. By virtue of Lemma 2.3, we obtain

$$
\rho_{\alpha}(G_1) \le \rho_{\alpha}(G_2),\tag{4.2}
$$

where the equality holds if and only if $(n_1, n_2, \ldots, n_c) = (n - (t + 1)c + 2, 1, \ldots, 1).$

If $c = 2$, then we have $G_2 = K_{2t-1} ∨ (K_{n-2t} ∪ K_1)$. According to [\(4.1\)](#page-7-0) and [\(4.2\)](#page-7-1), we get

 $\rho_{\alpha}(G) \leq \rho_{\alpha}(K_{2t-1} \vee (K_{n-2t} \cup K_1)),$

where the equality holds if and only if $G = K_{2t-1} \vee (K_{n-2t} \cup K_1)$. In what follows, we consider $c \geq 3$.

Recall that $G_2 = K_{tc-1} \vee (K_{n-(t+1)c+2} \cup (c-1)K_1)$. According to $\alpha \in [\frac{1}{2}, \frac{3}{4})$ and Lemma 2.5, we obtain

$$
\rho_{\alpha}(G_2) \le \frac{2e(G_2)(1-\alpha)}{n-1} + \alpha n - 1
$$

=
$$
\frac{(1-\alpha)((n-c+1)(n-c)+2(tc-1)(c-1))}{n-1} + \alpha n - 1
$$

=
$$
\frac{1-\alpha}{n-1}((2t+1)c^2 - (2n+2t+3)c + n^2 + n + 2) + \alpha n - 1.
$$
 (4.3)

Let $\varphi(c) = (2t+1)c^2 - (2n+2t+3)c + n^2 + n+2$. Note that $n \ge (t+1)c - 1$. We conclude $3 \le c \le \frac{n+1}{t+1}$. By a direct computation, it follows from $n \ge \max\{4t^2 + 6t + 1, \frac{12t(1-\alpha)-2\alpha+1}{3-4\alpha}\}$ $\frac{-\alpha - 2\alpha + 1}{3 - 4\alpha}$ that

$$
\varphi(3) - \varphi\left(\frac{n+1}{t+1}\right) = \frac{(n-3t-2)(n-4t^2-6t-1)}{(t+1)^2} \ge 0,
$$

which implies that $\varphi(c)$ attains its maximum value at $c = 3$ when $3 \leq c \leq \frac{n+1}{t+1}$. Combining this with $(4.3), \alpha \in \left[\frac{1}{2}, \frac{3}{4}\right)$ $(4.3), \alpha \in \left[\frac{1}{2}, \frac{3}{4}\right)$ and $n \ge \max\{4t^2 + 6t + 1, \frac{12t(1-\alpha)-2\alpha+1}{3-4\alpha}\}$ $\frac{-\alpha - 2\alpha + 1}{3 - 4\alpha}$, we get

$$
\rho_{\alpha}(G_2) \leq \frac{(1-\alpha)\varphi(3)}{n-1} + \alpha n - 1
$$

=
$$
\frac{(1-\alpha)(n^2 - 5n + 12t + 2)}{n-1} + \alpha n - 1
$$

=
$$
n - 2 + \frac{-(3-4\alpha)n + 12t(1-\alpha) - 2\alpha + 1}{n-1}
$$

$$
\leq n - 2 + \frac{-(3-4\alpha) \cdot \frac{12t(1-\alpha) - 2\alpha + 1}{3-4\alpha} + 12t(1-\alpha) - 2\alpha + 1}{n-1}
$$

=
$$
n - 2.
$$
 (4.4)

Since K_{n-1} is a proper subgraph of $K_{2t-1} \vee (K_{n-2t} \cup K_1)$, we conclude

$$
\rho_{\alpha}(K_{2t-1} \vee (K_{n-2t} \cup K_1)) > \rho_{\alpha}(K_{n-1}) = n-2
$$
\n(4.5)

by Lemmas 2.1 and 2.2. It follows from (4.1) , (4.2) , (4.4) and (4.5) that

$$
\rho_{\alpha}(G) \le \rho_{\alpha}(G_1) \le \rho_{\alpha}(G_2) \le n - 2 < \rho_{\alpha}(K_{2t-1} \vee (K_{n-2t} \cup K_1)).
$$

In conclusion, we have

$$
\rho_{\alpha}(G) \le \rho_{\alpha}(K_{2t-1} \vee (K_{n-2t} \cup K_1))
$$

with equality if and only if $G = K_{2t-1} \vee (K_{n-2t} \cup K_1)$. Which is a contradiction to the A_{α} -spectral radius condition of Theorem 1.2. This completes the proof of Theorem 1.2. \Box

Data availability statement

My manuscript has no associated data.

Declaration of competing interest

The authors declare that they have no conflicts of interest to this work.

References

- [1] V. Chvátal, Tough graphs and Hamiltonian circuits, Discrete Mathematics 5 (1973) 215–228.
- [2] H. Enomoto, B. Jackson, P. Katerinis, A. Saito, Toughness and the existence of k-factors, Journal of Graph Theory, 9 (1985) 87–95.
- [3] S. Zhou, Some results on path-factor critical avoidable graphs, Discussiones Mathematicae Graph Theory 43(1) (2023) 233–244.
- [4] S. Zhou, J. Wu, Y. Xu, Toughness, isolated toughness and path factors in graphs, Bulletin of the Australian Mathematical Society 106(2) (2022) 195–202.
- [5] S. Zhou, J. Wu, Q. Bian, On path-factor critical deleted (or covered) graphs, Aequationes Mathematicae 96(4) (2022) 795–802.
- [6] S. Wang, W. Zhang, Isolated toughness for path factors in networks, RAIRO-Operations Research 56(4) (2022) 2613–2619.
- [7] V. Nikiforov, Merging the A- and Q-spectral theories, Applicable Analysis and Discrete Mathematics 11 (2017) 81–107.
- [8] V. Nikiforov, O. Rojo, On the α -index of graphs with pendent paths, Linear Algebra and its Applications 550 (2018) 87–104.
- [9] A. Brondani, F. França, C. Oliveira, Positive semidefiniteness of $A_{\alpha}(G)$ on some families of graphs, Discrete Applied Mathematics 323 (2022) 113–123.
- [10] H. Lin, X. Huang, J. Xue, A note on the A_{α} -spectral radius of graphs, Linear Algebra and its Applications 557 (2018) 430–437.
- [11] H. Lin, X. Liu, J. Xue, Graphs determined by their A_{α} -spectra, Discrete Mathematics 342(2) (2019) 441–450.
- [12] H. Lin, J. Xue, J. Shu, On the A_{α} -spectra of graphs, Linear Algebra and its Applications 556 (2018) 210–219.
- [13] Y. Hao, S. Li, Q. Zhao, On the A_{α} -spectral radius of graphs without large matchings, Bulletin of the Malaysian Mathematical Sciences Society 45 (2022) 3131–3156.
- [14] Y. Zhao, X. Huang, Z. Wang, The A_{α} -spectral radius and perfect matchings of graphs, Linear Algebra and its Applications 631 (2021) 143–155.
- [15] S. O, Spectral radius and matchings in graphs, Linear Algebra and its Applications 614 (2021) 316–324.
- [16] D. Fan, H. Lin, H. Lu, Spectral radius and $[a, b]$ -factors in graphs, Discrete Mathematics 345(7) (2022) 112892.
- [17] Y. Zhang, H. Lin, Q. Liu, J. Zheng, Distance spectrum, 1-factor and vertex-disjoint cycles, Linear Algebra and its Applications 654 (2022) 10–27.
- [18] S. Zhou, Z. Sun, H. Liu, D-index and Q-index for spanning trees with leaf degree at most k in graphs, Discrete Mathematics 347(5) (2024) 113927.
- [19] S. Zhou, H. Liu, Two sufficient conditions for odd [1, b]-factors in graphs, Linear Algebra and its Applications 661 (2023) 149–162.
- [20] S. Zhou, H. Liu, Characterizing an odd [1, b]-factor on the distance signless Laplacian spectral radius, RAIRO-Operations Research 57(3) (2023) 1343–1351.
- [21] A. Brouwer, Toughness and spectrum of a graph, Linear Algebra and its Applications 226-228 (1995) 267–271.
- [22] X. Gu, Toughness in pseudo-random graphs, European Journal of Combinatorics 92 (2021) 103255.
- [23] X. Gu, A proof of Brouwer's toughness conjecture, SIAM Journal of Discrete Mathematics 35 (2021) 948–952.
- [24] D. Fan, H. Lin, H. Lu, Toughness, hamiltonicity and spectral radius in graphs, European Journal of Combinatorics 110 (2023) 103701.
- [25] W. Haemers, Interlacing eigenvalues and graphs, Linear Algebra and its Applications 227 (1995) 593–616.
- [26] L. You, M. Yang, W. So, W. Xi, On the spectrum of an equitable quotient matrix and its application, Linear Algebra and its Applications 577 (2019) 21–40.
- [27] A. Alhevaz, M. Baghipur, H. Ganie, K. Das, On the A_{α} -spectral radius of connected graphs, Ars Mathematica Contemporanea 23 (2023) #P1.06.