

ON OSCILLATIONS OF NONAUTONOMOUS LINEAR IMPULSIVE DIFFERENTIAL EQUATIONS WITH GENERAL PIECEWISE CONSTANT DEVIATING ARGUMENTS

RICARDO TORRES

Dedicated to the memory of Prof. István Györi and Prof. Nicolás Yus Suárez.

ABSTRACT. We give some sufficient conditions that ensure oscillations and nonoscillations for nonautonomous impulsive differential equations with piecewise constant arguments of generalized type. We cover several cases of differential equations with deviated arguments investigated before as particular cases.

1. INTRODUCTION

In many cases, the functions used for modeling the natural phenomena with differential equations may have to be discontinuous, such as a piecewise constant, or have the impulsive effect present to reflect the phenomena's characteristics to be modeled. The reader can find examples of this kind of phenomenon in the works of S. Busenberg and K. Cooke ([7]) and L. Dai and M.C. Singh ([8]). In the first one, the authors modeled diseases of vertical transmission. In the second one, the authors studied the oscillatory motion of spring-mass systems subject to piecewise constant forces of the form $Ax([t])$ or $A\cos([t])$. Indeed, one such system is the famous Geneva wheel, widely used in watches and instruments. The motion of this mechanism is piecewise continuous, and it can be modeled by

$$mx''(t) + kx_1 = A\sin\left(\omega\left[\frac{t}{T}\right]\right),$$

where $[\cdot]$ is the greatest integer function. (See [9]).

As a generalization of the work done by A. Myshkis in [13] (1977) concerning differential equations with deviating arguments, M. Akhmet introduced a very interesting class of differential equations of the form

$$(1.1) \quad z'(t) = f(t, z(t), z(\gamma(t))),$$

where $\gamma(t)$ is a *piecewise constant argument of generalized type*. Such γ is defined as follows: let $(t_n)_{n \in \mathbb{Z}}$ and $(\zeta_n)_{n \in \mathbb{Z}}$ such that $t_n < t_{n+1}, \forall n \in \mathbb{Z}$ with $\lim_{n \rightarrow \infty} t_n = \infty$, $\lim_{n \rightarrow -\infty} t_n = -\infty$ and $\zeta_n \in [t_n, t_{n+1}]$, then $\gamma(t) = \zeta_n$, if $t \in I_n = [t_n, t_{n+1})$. I.e., $\gamma(t)$ is a locally constant function. An elementary example of such functions is $\gamma(t) = [t]$, where $[\cdot]$ is the greatest integer function, which is constant in every interval $[n, n+1[$

2010 *Mathematics Subject Classification.* 34A36, 34K11, 34K45, 39A06, 39A21.

Key words and phrases. Piecewise constant argument, linear functional differential equations, Oscillatory solutions, Impulsive Differential equations, Hybrid dynamics.

with $n \in \mathbb{Z}$.

If a piecewise constant argument is used, the interval I_n is decomposed into an advanced and delayed subintervals $I_n = I_n^+ \cup I_n^-$, where $I_n^+ = [t_n, \zeta_n]$ and $I_n^- = [\zeta_n, t_{n+1}]$. This class of differential equations is known as *Differential Equations with Piecewise Constant Argument of Generalized Type (DEPCAG)*. They have remarkable properties, as the solutions are continuous functions, even when γ is discontinuous. If we assume continuity of the solutions of (1.1), integrating from t_n to t_{n+1} , we define a difference equation, so this type of differential equation has hybrid dynamics (see [2, 14, 21]).

For example, in [18], the author introduced the piecewise constant argument $\gamma(t) = \left[\frac{t}{h}\right]h + \alpha h$ with $h > 0$ and $0 \leq \alpha \leq 1$. Then, we can see that

$$\left[\frac{t}{h}\right]h + \alpha h = (n + \alpha)h, \text{ when } t \in I_n = [nh, (n + 1)h).$$

Its delayed and advanced intervals can be concluded easily: $\gamma(t) - t \geq 0 \Leftrightarrow t \leq (n + \alpha)h$ and $\gamma(t) - t \leq 0 \Leftrightarrow t \geq (n + \alpha)h$. Hence, we have

$$I_n^+ = [nh, (n + \alpha)h), \quad I_n^- = [(n + \alpha)h, (n + 1)h).$$

This piecewise constant argument has been used (case $\alpha = 0$) in the approximation of solutions of ordinary and delayed differential equations (see also [19, 20]).

If an impulsive condition is considered at instants $\{t_n\}_{n \in \mathbb{Z}}$, we define the *Impulsive differential equations with piecewise constant argument of generalized type (IDEPCAG)*,

$$(1.2) \quad \begin{aligned} z'(t) &= f(t, z(t), z(\gamma(t))), & t \neq t_n, \\ \Delta z(t_n) &:= z(t_n) - z(t_n^-) = P_n(z(t_n^-)), & t = t_n, \quad n \in \mathbb{N}, \end{aligned}$$

where $z(t_n^-) = \lim_{t \rightarrow t_n^-} z(t)$, and P_n is the impulsive operator (see [1, 15]).

If the differential equation explicitly shows the piecewise constant argument used, we will call it DEPCA (IDEPCA if it has impulses).

A function $x(t)$ defined on $[\tau, \infty)$ is said to be oscillatory if there exist two real-valued sequences $(a_n), (b_n) \subset [\tau, \infty)$ such that $a_n \rightarrow \infty, b_n \rightarrow \infty$ as $n \rightarrow \infty$ and $x(a_n) \leq 0 \leq x(b_n), \forall n \geq M$, where M is sufficiently large. I.e., if it is neither eventually positive nor eventually negative. Otherwise, it is called nonoscillatory.




FIGURE 1. $f(t) = t - [t]$: an example of an oscillatory piecewise continuous function with infinite zeros ($f(n) = 0, \forall n \in \mathbb{Z}$).

In the piecewise continuous frame, a function $x(t)$ can be oscillatory even if $x(t) \neq 0, \forall t \in [\tau, \infty)$. This is the context of this work, so all the definition of oscillation for the continuous frame are no longer valid (see [12]).




FIGURE 2. $f(t) = (-1)^{[t]}(t + 1 - [t])$. An oscillatory piecewise continuous function with $f(t) \neq 0, \forall t \in \mathbb{R}$.

A nonzero solution $y(n)$ of a difference equation will be *oscillatory* if for every positive integer $M > 0$, there exists $n \geq M$ such that

$$y(n)y(n+1) \leq 0.$$

Otherwise, $y(n)$ will be called *nonoscillatory* (see [10]).

2. AIM OF THE WORK

In [22], J. Wiener and A.R. Aftabzadeh studied the following DEPCA

$$(2.1) \quad x'(t) = a(t)x(t) + a_0(t)x\left(m \left[\frac{t+k}{m}\right]\right), \quad x(0) = C_0,$$

where $a(t), a_0(t)$ are continuous functions on $[0, \infty)$. They conclude the following oscillatory criterion:

Theorem 1. *if either of the conditions*

$$\begin{aligned} \limsup_{n \rightarrow \infty} \int_{mn-k}^{mn} a_0(t) \exp\left(\int_t^{mn} a(s) ds\right) dt &> 1, \\ \liminf_{n \rightarrow \infty} \int_{mn}^{m(n+1)-k} a_0(t) \exp\left(\int_t^{mn} a(s) ds\right) dt &< -1, \end{aligned}$$

holds true, then every solution of (1) is oscillatory.

Inspired in the last work, we will get sufficient conditions for the oscillation of the solutions of the scalar linear nonautonomous IDEPCAG

$$(2.2) \quad \begin{aligned} x'(t) &= a(t)x(t) + b(t)x(\gamma(t)), & t \neq t_k \\ \Delta x|_{t=t_k} &= c_k x(t_k^-), & t = t_k \end{aligned}$$

where $a(t), b(t)$ are continuous real-valued locally integrable functions, $c_k \neq -1, \forall k \in \mathbb{Z}$ and γ is a piecewise constant argument of generalized type.

We will improve some of the results of the last section considering any piecewise constant argument $\gamma(t)$ such that $\gamma(t) = \zeta_k$ if $t \in [t_k, t_{k+1})$ with $t_k \leq \zeta_k \leq t_{k+1}$.

Our work is organized as follows: First, we give some preliminary necessary for a better understanding of the IDEPCAG frame. Next, we study oscillations for the scalar linear IDEPCAG and provide some criteria for oscillatory solutions (extending the classical ones) for the autonomous and nonautonomous cases. Finally, we give some examples showing the effectiveness of our results.

3. SOME RECENT RESULTS OF OSCILLATIONS ON IDEPCAG

In [6] (2013), K-S. Chiu and M. Pinto studied the scalar DEPCAG

$$y'(t) = a(t)y(t) + b(t)y(\gamma(t)), \quad y(\tau) = y_0,$$

where $a(t), b(t)$ are real continuous functions defined on \mathbb{R} . The authors investigated oscillation and periodic solutions. This excellent work seems to be the first to consider oscillations and periodic solutions with a generalized piecewise constant argument. One of the main hypotheses used by the authors is the continuity of the solutions. Still, it has no results for the IDEPCA or IDEPCAG cases.

In [12] (2018), F. Karakoc, A. Unal, and H. Bereketoglu studied the following nonlinear IDEPCA:

$$\begin{cases} x'(t) = -a(t)x(t) - x([t-1])f(y([t]) + h_1(x[t])), & t \neq n \in \mathbb{Z}^+ \\ y'(t) = -b(t)x(t) - y([t-1])g(y([t]) + h_2(y[t])), \\ \\ \begin{cases} x(n) = (1 + c_n)x(n^-), & t = n \in \mathbb{Z}^+ \\ y(n) = (1 + d_n)y(n^-), \end{cases} \end{cases}$$

with initial conditions $x(-1) = x_{-1}$, $x(0) = x_0$, $y(-1) = y_{-1}$, $y(0) = y_0$, where $a, b : [0, \infty) \rightarrow \mathbb{R}$ are continuous functions, $f, g, h_1, h_2 \in C(\mathbb{R}, \mathbb{R})$, c_n and d_n are sequences of real numbers such that $c_n \neq 1$, $d_n \neq 1$, $\forall n \geq 1$. The authors studied the existence and uniqueness of solutions, and they obtained some sufficient conditions for the oscillation of the solution. They stated the following sufficient conditions of oscillation of the solutions:

Theorem 2. Assume that there exist $M_1 > 0$ and $M_2 > 0$ such that $f(t) \geq M_1$ and $g(t) \geq M_2$, $\forall t \in \mathbb{R}$, $t h_1(t), t h_2(t) < 0$ for $t \neq 0$ and $c_n, d_n < 1$ for all $n \in \mathbb{Z}^+$. If the following conditions hold,

$$\begin{aligned} \limsup_{n \rightarrow \infty} (1 - c_n) \int_n^{n+1} \exp \left(\int_{n-1}^s a(u) du \right) ds &> \frac{1}{M_1}, \\ \limsup_{n \rightarrow \infty} (1 - d_n) \int_n^{n+1} \exp \left(\int_{n-1}^s b(u) du \right) ds &> \frac{1}{M_2}. \end{aligned}$$

then all solutions of (3)-(3) are oscillatory.

In [6] (2013), K-S. Chiu and M. Pinto studied the scalar DEPCAG

$$y'(t) = a(t)y(t) + b(t)y(\gamma(t)), \quad y(\tau) = y_0,$$

where $a(t), b(t)$ are real continuous functions defined on \mathbb{R} . The authors investigated oscillation and periodic solutions. This excellent work seems to be the first to consider oscillations and periodic solutions with a generalized piecewise constant argument. One of the main hypotheses used by the authors is the continuity of the solutions. Still, it has no results for the IDEPCA or IDEPCAG cases.

In [5] (2023), K-S. Chiu and I. Berna considered the following impulsive differential equation with a piecewise constant argument

$$(3.1) \quad \begin{aligned} y'(t) &= a(t)y(t) + b(t)y\left(p\left[\frac{t+l}{p}\right]\right), & y(\tau) &= c_0, & t &\neq kp-l \\ \Delta y(kp-l) &= d_k y(kp-l^-), & t &= kp-l, & k &\in \mathbb{Z}, \end{aligned}$$

where $a(t) \neq 0$ and $b(t)$ are real-valued continuous functions, $p < l$ and $d_k \neq -1$, $\forall k \in \mathbb{Z}$. The authors obtained an oscillation criterion for solutions of (3.1).

Finally, in [23] (2023), Z. Yan and J. Gao studied oscillation and non-oscillation of Runge–Kutta methods for the following linear mixed-type impulsive differential equations with piecewise constant arguments

$$(3.2) \quad \begin{aligned} y'(t) &= ay(t) + by([t-1]) + cy([t]) + dy([t+1]), & t &\neq k, \\ \Delta y(k) &= qy(k^-), & t &= k, & k &\in \mathbb{N}, \\ y(0) &= y_0, y(1) = y_1, \end{aligned}$$

where $a, b, c, d, q, y_0, y_1 \in \mathbb{R}$ and $q \neq 1$. They obtained conditions for oscillation and non-oscillation of the numerical solutions. Also, the authors obtained conditions under which Runge–Kutta methods can preserve the oscillation and non-oscillation.

3.1. Why study oscillations on IDEPCAG?.

Example 1. Let the following scalar linear DEPCA

$$(3.3) \quad x'(t) = a(t)(x(t) - x([t+1])), \quad x(\tau) = x_0,$$

and the scalar linear IDEPCA

$$(3.4) \quad \begin{aligned} z'(t) &= a(t)(z(t) - z([t+1])), & t &\neq k \\ z(k) &= c_k z(k^-), & t &= k, & k &\in \mathbb{Z}, \end{aligned}$$

where $a(t)$ is a continuous locally integrable function. As $\gamma(t) = [t+1]$, we have $t_k = k$, $\zeta_k = k+1$ if $t \in [k, k+1]$, $k \in \mathbb{Z}$.

It is very easy to see, by Theorem 5 (considering $c_k = 0$, $\forall k \in \mathbb{Z}$), that the solution of (3.3) is $x(t) = x_0$, $\forall t \geq \tau$. I.e., all the solutions are constant. Thus, solutions are nonoscillatory (we excluded the zero solution).

On the other hand, the solution of (3.4) is

$$z(t) = \left(\prod_{j=k(\tau)+1}^{k(t)} c_j \right) z(\tau), \quad t \geq \tau,$$

where $k(t) = k$ is the only integer such that $t \in [k, k+1]$ (see Example 4). Hence, if $c_k < 0$, $\forall k \in \mathbb{Z}$ all the solutions are oscillatory. This example shows the role of the presence of the impulsive effect. In fact, our main result applies to DEPCAG and IDEPCAG cases.

4. SOME PRELIMINARY ON IDEPCAG THEORY

Let $\mathcal{PC}(X, Y)$ be the set of all functions $r : X \rightarrow Y$ which are continuous for $t \neq t_k$ and continuous from the left with discontinuities of the first kind at $t = t_k$. Similarly, let $\mathcal{PC}^1(X, Y)$ the set of functions $s : X \rightarrow Y$ such that $s' \in \mathcal{PC}(X, Y)$.




FIGURE 3. Solution of (3.4) with $c_k = -\frac{9}{10}$ and $z(0) = -19$.

Definition 1 (DEPCAG solution). A continuous function $x(t)$ is a solution of (1.1) if:

- (i) $x'(t)$ exists at each point $t \in \mathbb{R}$ with the possible exception at the times t_k , $k \in \mathbb{Z}$, where the one side derivative exists.
- (ii) $x(t)$ satisfies (1.1) on the intervals of the form (t_k, t_{k+1}) , and it holds for the right derivative of $x(t)$ at t_k .

Definition 2 (IDEPCAG solution). A piecewise continuous function $y(t)$ is a solution of (1.2) if:

- (i) $y(t)$ is continuous on $I_k = [t_k, t_{k+1})$ with first kind discontinuities at t_k , $k \in \mathbb{Z}$, where $y'(t)$ exists at each $t \in \mathbb{R}$ with the possible exception at the times t_k , where lateral derivatives exist (i.e. $y(t) \in \mathcal{PC}^1([t_k, t_{k+1}), \mathbb{R})$).
- (ii) The ordinary differential equation

$$y'(t) = f(t, y(t), y(\zeta_k))$$

holds on every interval I_k , where $\gamma(t) = \zeta_k$.

- (iii) For $t = t_k$, the impulsive condition

$$\Delta y(t_k) = y(t_k) - y(t_k^-) = P_k(y(t_k^-))$$

holds. I.e., $y(t_k) = y(t_k^-) + P_k(y(t_k^-))$, where $y(t_k^-)$ denotes the left-hand limit of the function y at t_k .

Let the following IDEPCAG:

$$(4.1) \quad \begin{aligned} x'(t) &= f(t, x(t), x(\gamma(t))), & t \neq t_k \\ x(t_k) - x(t_k^-) &= P_k(x(t_k^-)), & t = t_k, \\ x(\tau) &= x_0, \end{aligned}$$

where $f \in C([\tau, \infty) \times \mathbb{R} \times \mathbb{R}, \mathbb{R})$, $P_k \in C(\{t_k\}, \mathbb{R})$ and $(\tau, x_0) \in \mathbb{R}^2$.

Also, let the following hypothesis hold:

- (H1) Let $\eta_1, \eta_2 : \mathbb{R} \rightarrow [0, \infty)$ locally integrable functions and $\omega_k \in \mathbb{R}^+$, $\forall k \in \mathbb{Z}$; such that

$$\begin{aligned} |f(t, x_1, y_1) - f(t, x_2, y_2)| &\leq \eta_1(t) |x_1 - x_2| + \eta_2(t) |y_1 - y_2|, \\ |P_k(x_1(t_k^-)) - P_k(x_2(t_k^-))| &\leq \omega_k |x_1(t_k^-) - x_2(t_k^-)|. \end{aligned}$$

(H2) Assume that

$$\bar{\nu} = \sup_{k \in \mathbb{Z}} \left(\int_{t_k}^{t_{k+1}} (\eta_1(s) + \eta_2(s)) ds \right) < 1.$$

In the following, we mention some valuable results for the rest of the work:

4.1. IDEPCAG Gronwall-Bellman type inequality.

Lemma 1. ([17],[3] Lemma 4.3) Let I an interval and $u, \eta_1, \eta_2 : I \rightarrow [0, \infty)$ such that u is continuous (with possible exception at $\{t_k\}_{k \in \mathbb{N}}$), η_1, η_2 are continuous and locally integrable functions, $\eta : \{t_k\} \rightarrow [0, \infty)$ and $\gamma(t)$ a piecewise constant argument of generalized type such that $\gamma(t) = \zeta_k, \forall t \in I_k = [t_k, t_{k+1})$ with $t_k \leq \zeta_k \leq t_{k+1} \forall k \in \mathbb{N}$. Assume that $\forall t \geq \tau$

$$u(t) \leq u(\tau) + \int_{\tau}^t (\eta_1(s)u(s) + \eta_2(s)u(\gamma(s))) ds + \sum_{\tau \leq t_k < t} \eta(t_k)u(t_k^-),$$

$$\widehat{\vartheta}_k = \int_{t_k}^{\zeta_k} (\eta_1(s) + \eta_2(s)) ds \leq \widehat{\vartheta} := \sup_{k \in \mathbb{N}} \widehat{\vartheta}_k < 1$$

hold. Then, for $t \geq \tau$, we have

$$u(t) \leq \left(\prod_{\tau \leq t_k < t} (1 + \eta(t_k)) \right) \exp \left(\int_{\tau}^t \left(\eta_1(s) + \frac{\eta_2(s)}{1 - \widehat{\vartheta}} \right) ds \right) u(\tau),$$

$$u(\zeta_k) \leq (1 - \vartheta)u(t_k)$$

$$u(\gamma(t)) \leq (1 - \vartheta)^{-1} \left(\prod_{\tau \leq t_k < t} (1 + \eta_3(t_j)) \right) \exp \left(\int_{\tau}^t \left(\eta_1(s) + \frac{\eta_2(s)}{1 - \widehat{\vartheta}} \right) ds \right) u(\tau).$$

4.2. Existence and uniqueness for (4.1).

Theorem 3. (Uniqueness) ([3], Theorem 4.5) Consider the I.V.P for (2.2) with $y(t, \tau, y(\tau))$. Let (H1)-(H2) and Lemma 1 hold. Then, there exists a unique solution y for (2.2) on $[\tau, \infty)$. Moreover, every solution is stable.

Lemma 2. (Existence of solutions in $[\tau, t_k)$ ([3], Lemma 4.6) Consider the I.V.P for (2.2) with $y(t, \tau, y(\tau))$. Let (H1)-(H2) and Lemma 1 hold. Then, for each $y_0 \in \mathbb{R}^n$ and $\zeta_k \in [t_{k-1}, t_k)$ there exists a solution $y(t) = y(t, \tau, y(\tau))$ of (2.2) on $[\tau, t_r)$ such that $y(\tau) = y_0$.

Theorem 4. (Existence of solutions in $[\tau, \infty)$ ([3], Theorem 4.7) Let (H1)-(H2) and Lemma 1 hold. Then, for each $(\tau, y_0) \in \mathbb{R}_0^+ \times \mathbb{R}$, there exists $y(t) = y(t, \tau, y_0)$ for $t \geq \tau$, a unique solution for (2.2) such that $y(\tau) = y_0$.

5. THE NONAUTONOMOUS HOMOGENEOUS LINEAR IDEPCAG

In this section, we will present the nonautonomous homogeneous linear IDEPCAG

$$(5.1) \quad \begin{aligned} z'(t) &= a(t)z(t) + b(t)z(\gamma(t)), & t \neq t_k \\ \Delta z|_{t=t_k} &= c_k z(t_k^-), & t = t_k \end{aligned}$$

where $z \in \mathbb{R}, t \in \mathbb{R}$, $a(t), b(t)$ are real-valued continuous locally integrable functions, $(c_k)_{k \in \mathbb{N}}$ is a real sequence such that $1 + c_k \neq 0 \forall k \in \mathbb{N}$ and $\gamma(t)$ is a generalized

piecewise constant argument.

During the rest of the work, we will assume $\gamma(\tau) := \tau$ if $t_{k(\tau)} \leq \gamma(\tau) < \tau < t_{k(\tau)+1}$, where $k(\tau)$ is the only $k \in \mathbb{Z}$ such that $t_k \leq \tau \leq t_{k+1}$, and we will adopt the following notation for the matricial products and sums:

$$\prod_{j=1}^n A_j = \begin{cases} A_n \cdot A_{n-1} \cdots A_1, & \text{if } n \geq 1, \\ I & \text{if } n < 1. \end{cases} \quad \text{and} \quad \sum_{j=1}^n A_j = \begin{cases} A_1 + \dots + A_n, & \text{if } n \geq 1, \\ 0 & \text{if } n < 1. \end{cases}$$

Let $x(t) = \phi(t, \tau)x(\tau) = \exp\left(\int_{\tau}^t a(u)du\right)x(\tau)$ the solution of the ordinary differential equation

$$\begin{aligned} x'(t) &= a(t)x(t), \\ x_0 &= x(\tau), \quad t, \tau \in [\tau, \infty). \end{aligned}$$

We will assume the following hypothesis:

$$\begin{aligned} (H3) \quad \text{Let } \rho_k^+(a) &= \exp\left(\int_{t_k}^{\zeta_k} |a(u)| du\right), \quad \rho_k^-(a) = \exp\left(\int_{\zeta_k}^{t_{k+1}} |a(u)| du\right), \\ \rho_k(a) &= \rho_k^+(a)\rho_k^-(a) \quad \text{and} \quad \nu_k^{\pm}(b) = \rho_k^{\pm}(a) \ln \rho_k^{\pm}(b), \\ &\text{and assume that} \end{aligned}$$

$$\rho(a) = \sup_{k \in \mathbb{Z}} \rho_k(a) < \infty, \quad \nu^{\pm}(b) = \sup_{k \in \mathbb{Z}} \nu_k^{\pm}(b) < \infty,$$

where

$$(5.2) \quad \nu_k^+(b) < \nu^+(b) < 1, \quad \nu_k^-(b) < \nu^-(b) < 1.$$

Consider the following definitions

$$\begin{aligned} j(t, \zeta_k) &= 1 + \int_{\zeta_k}^t \exp\left(\int_s^{\zeta_k} a(u)du\right) b(s)ds, \\ e(t, \zeta_k) &= \phi(t, \zeta_{k(t)})j(t, \zeta_{k(t)}) \\ &= \left(\exp \int_{\zeta_{k(t)}}^t a(u)du\right) \\ (5.3) \quad &\cdot \left(1 + \int_{\zeta_{k(t)}}^t \exp\left(\int_s^{\zeta_{k(s)}} a(u)du\right) b(s)ds\right). \end{aligned}$$

Remark 1. As a consequence of (H3), it is important to notice the following facts:

(i) Due to condition (5.2), $j^{-1}(t_k, \zeta_k)$ and $j^{-1}(t_{k+1}, \zeta_k)$ are well defined $\forall k \in \mathbb{Z}$, and

$$\begin{aligned} |j^{-1}(t_k, \zeta_k)| &\leq \sum_{k=0}^{\infty} [\nu^+(b)]^k = \frac{1}{1 - \nu^+(b)}, \quad |j(t_{k+1}, \zeta_k)| \leq 1 + \nu^-(b), \\ |j^{-1}(t_{k+1}, \zeta_k)| &\leq \sum_{k=0}^{\infty} [\nu^-(b)]^k = \frac{1}{1 - \nu^-(b)}, \quad |j(t_k, \zeta_k)| \leq 1 + \nu^+(b). \end{aligned}$$

Also, if we set $t_0 = \tau$, we are considering that $j^{-1}(\tau, \gamma(\tau))$ exists.

(ii) On the other hand, if we want a nonzero solution of a linear IDEPCAG, we need $j(t, \zeta_k) \neq 0, \forall k \in \mathbb{Z}$ and $\forall t \in [\tau, \infty)$ (See Remark 2 and [6]).

5.1. The fundamental solution of the homogeneous linear IDEPCAG.

The following results can be found in [17] and [16] in a more general matricial context. They are the IDEPCAG extension of [14]. We will use the next scalar version of them:

Theorem 5. *Let the following linear IDEPCAG system*

$$(5.4) \quad \begin{aligned} z'(t) &= a(t)z(t) + b(t)z(\gamma(t)), & t \neq t_k \\ z(t_k) &= (1 + c_k)z(t_k^-), & t = t_k \\ z_0 &= z(\tau). \end{aligned}$$

If (H3) holds, then the unique solution of (5.4) is

$$z(t) = w(t, \tau)z(\tau), \quad t \in [\tau, \infty),$$

where $w(t, \tau)$ is given by

$$(5.5) \quad w(t, \tau) = w(t, t_{k(t)}) \left(\prod_{r=k(\tau)+2}^{k(t)} (1 + c_r) w(t_r, t_{r-1}) \right) (1 + c_{k(\tau)+1}) w(t_{k(\tau)+1}, \tau)$$

for $t \in I_{k(t)}$ and $\tau \in I_{k(\tau)}$, where $w(t, s)$ is defined as

$$w(t, s) = e(t, \gamma(s))e^{-1}(s, \gamma(s)), \quad \text{if } t, s \in I_k = [t_k, t_{k+1}).$$

In a more explicit form, the unique solution of (5.4) can be written as

$$(5.6) \quad \begin{aligned} z(t) &= w(t, t_{k(t)}) \left(\prod_{r=k(\tau)+2}^{k(t)} (1 + c_r) w(t_r, t_{r-1}) \right) \\ &\cdot (1 + c_{k(\tau)+1}) w(t_{k(\tau)+1}, \tau) z(\tau), \quad \text{for } t \in I_{k(t)}, \tau \in I_{k(\tau)}, \end{aligned}$$

and the discrete solution of (5.4) can be also written by

$$(5.7) \quad z(t_{k(t)}) = \left(\prod_{r=k(\tau)+2}^{k(t)} (1 + c_r) w(t_r, t_{r-1}) \right) (1 + c_{k(\tau)+1}) w(t_{k(\tau)+1}, \tau) z(\tau).$$

The expression given by (5.5) is called the fundamental solution for (5.4).

Proof. Let $t, \tau \in I_k = [t_k, t_{k+1})$ for some $k \in \mathbb{Z}$. In this interval, we are in the presence of the ordinary system

$$z'(t) = a(t)z(t) + b(t)z(\zeta_k).$$

So, the unique solution can be written as

$$(5.8) \quad z(t) = \phi(t, \tau)z(\tau) + \int_{\tau}^t \phi(t, s)b(s)z(\zeta_k)ds.$$

Keeping in mind I_k^+ , evaluating the last expression at $t = \zeta_k$ we have

$$z(\zeta_k) = \phi(\zeta_k, \tau)z(\tau) + \int_{\tau}^{\zeta_k} \phi(\zeta_k, s)b(s)z(\zeta_k)ds,$$

Hence, we get

$$\left(1 + \int_{\zeta_k}^{\tau} \phi(\zeta_k, s)b(s)ds \right) z(\zeta_k) = \phi(\zeta_k, \tau)z(\tau).$$

I.e

$$z(\zeta_k) = j^{-1}(\tau, \zeta_k) \phi(\zeta_k, \tau) z(\tau).$$

Then, by the definition of $e(t, \tau) = \phi(t, \tau) j(t, \tau)$, we have

$$(5.9) \quad z(\zeta_k) = e^{-1}(\tau, \zeta_k) z(\tau).$$

Now, from (5.8) working on I_k^- , considering $\tau = \zeta_k$, we have

$$\begin{aligned} z(t) &= \phi(t, \zeta_k) z(\zeta_k) + \int_{\zeta_k}^t \phi(t, s) b(s) z(\zeta_k) ds \\ &= \phi(t, \zeta_k) \left(1 + \int_{\zeta_k}^t \phi(\zeta_k, s) b(s) ds \right) z(\zeta_k). \end{aligned}$$

I.e.,

$$(5.10) \quad z(t) = e(t, \zeta_k) z(\zeta_k).$$

So, by (5.9), we can rewrite (5.10) as

$$(5.11) \quad z(t) = e(t, \zeta_k) e^{-1}(\tau, \zeta_k) z(\tau).$$

Then, setting

$$w(t, s) = e(t, \gamma(s)) e^{-1}(s, \gamma(s)), \quad \text{if } t, s \in I_k = [t_k, t_{k+1}],$$

we have the solution for (5.4) for $t \in I_k = [t_k, t_{k+1})$,

$$(5.12) \quad z(t) = w(t, \tau) z(\tau).$$

Next, if we consider $\tau = t_k$, and, assuming left side continuity of (5) at $t = t_{k+1}$, we have

$$z(t_{k+1}^-) = w(t_{k+1}, t_k) z(t_k)$$

Then, applying the impulsive condition to the last equation, we get

$$(5.13) \quad z(t_{k+1}) = (1 + c_{k+1}) \cdot w(t_{k+1}, t_k) z(t_k).$$

The last expression defines a finite-difference equation whose solution is (5.7). Now, by (5.12) and the impulsive condition, we have

$$z(t_{k(\tau)+1}) = (1 + c_{k(\tau)+1}) w(t_{k(\tau)+1}, \tau) z(\tau).$$

Hence, considering $\tau = t_k$ in (5) and applying (5.7), we get (5.6). In this way, we have solved (5.4) on $[\tau, t)$. \square

Remark 2.

(i) We used the decomposition of $I_k = I_k^+ \cup I_k^-$ to define W .

In fact, we can rewrite (5.5) in terms of the advanced and delayed parts using (5):

$$\begin{aligned} w(t, \tau) &= e(t, \zeta_{k(t)}) e^{-1}(t_{k(t)}, \zeta_{k(t)}) \left(\prod_{r=k(\tau)+2}^{k(t)} (1 + c_r) e(t_r, \zeta_{r-1}) e^{-1}(t_{r-1}, \zeta_{r-1}) \right) \\ &\quad \cdot (1 + c_{k(\tau)+1}) e(t_{k(\tau)+1}, \gamma(\tau)) e^{-1}(\tau, \gamma(\tau)), \quad \zeta_r = \gamma(t_r), \end{aligned}$$

for $t \in I_{k(t)}$ and $\tau \in I_{k(\tau)}$.

(ii) From (5.6) and (5.7), we can write

$$(5.14) \quad z(t) = w(t, t_{k(t)})z(t_{k(t)}),$$

for $t \in I_k = [t_k, t_{k+1})$, where

$$w(t, t_{k(t)}) = \phi(t, t_{k(t)})j(t, \zeta_{k(t)})j^{-1}(t_{k(t)}, \zeta_{k(t)}),$$

and $z(t_{k(t)})$ is the unique solution of the difference equation

$$(5.15) \quad z(t_{k(t)+1}) = (1 + c_{k(t)+1})w(t_{k(t)+1}, t_{k(t)})z(t_{k(t)}),$$

with $z_0 = z_{k(\tau)}$.

(iii) It is important to notice that a (5.4) has a zero on the interval $I_j = [t_j, t_{j+1})$ at $t = \alpha_j$ if and only if

$$\int_{\zeta_j}^{\alpha_j} \exp\left(\int_s^{\zeta_j} a(u)du\right) b(s)ds = -1,$$

holds, for some $\alpha_j \in I_j$.

Corollary 1. Let the following scalar linear IDEPCAG

$$\begin{aligned} y'(t) &= b(t)y(\gamma(t)), & t &\neq t_k \\ y(t_k) &= (1 + c_k)y(t_k^-), & t &= t_k, \\ y(\tau) &= y_0. \end{aligned}$$

According to Theorem 5, we see that $e(t, t_k) = j(t, t_k)$, where

$$j(t, t_k) = 1 + \int_{t_k}^t b(s)ds \quad \text{and} \quad w(t, s) = j(t, \gamma(s))j^{-1}(s, \gamma(s)).$$

Hence, the solution of the last IDEPCAG is:

$$\begin{aligned} y(t) &= j(t, \zeta_{k(t)})j^{-1}(t_{k(t)}, \zeta_{k(t)}) \left(\prod_{r=k(\tau)+2}^{k(t)} (1 + c_r) j(t_r, \zeta_{r-1})j^{-1}(t_{r-1}, \zeta_{r-1}) \right) \\ &\quad \cdot (1 + c_{k(\tau)+1}) j(t_{k(\tau)+1}, \gamma(\tau))j^{-1}(\tau, \gamma(\tau))y(\tau), \quad \zeta_r = \gamma(t_r). \end{aligned}$$

Remark 3.

- a) Considering $b(t) = 0$, we recover the classical fundamental solution of the impulsive linear differential equation (see [15]).
- b) If $c_k = 0, \forall k \in \mathbb{Z}$, we recover the DEPCAG case studied by M. Pinto in [14].
- c) If we consider $\gamma(t) = p \left[\frac{t+l}{p} \right]$ with $p < l$, we recover the IDEPCA case studied by K-S. Chiu in [5].

6. OSCILLATIONS IN LINEAR IDEPCAG

In this section, we will present sufficient conditions to ensure the oscillatory character of solutions for the scalar and nonscalar version of IDEPCAG (5.4).

6.1. Oscillations in scalar Linear IDEPCAG. *Following the ideas given in [4], we will provide useful results concerning oscillation and nonoscillation conditions of linear IDEPCAG systems, knowing the oscillatory properties of either the discrete or the continuous solution of a linear IDEPCAG.*

Theorem 6. *Let the scalar IDEPCAG:*

$$(6.1) \quad \begin{aligned} z'(t) &= a(t)z(t) + b(t)z(\gamma(t)), & t \neq t_n \\ z(t_n) &= (1 + c_n)z(t_n^-), & t = t_n \\ z_0 &= z(\tau), \end{aligned}$$

with $1 + c_n \neq 0$ for all $n \in \mathbb{N}$. Then:

- a) *If the discrete solution of (6.1) (i.e., the solution of the difference equation (5.13)) is oscillatory, then the solution $z(t)$ of (6.1) is oscillatory.*
- b) *$z(t)$ will be oscillatory if the sequence*

$$\{w_n\} = \left\{ (1 + c_n) \left(\frac{1 + \int_{\zeta_n}^{t_{n+1}} \exp\left(\int_s^{\zeta_n} a(r)dr\right) b(u)du}{1 + \int_{\zeta_n}^{t_n} \exp\left(\int_s^{\zeta_n} a(r)dr\right) b(u)du} \right) \right\}_{n \geq n(\tau)+1}$$

is not eventually positive nor eventually negative.

- c) *If the discrete solution of (6.1) $z(t_n)_{n \geq n(\tau)}$ is nonoscillatory, then $z(t)$ is nonoscillatory if and only if*

$$(6.2) \quad \begin{cases} \int_{\zeta_{k(t)}}^t \exp\left(\int_s^{\zeta_{k(t)}} a(u)du\right) b(s)ds > -1, \\ \text{or} \\ \int_{\zeta_{k(t)}}^t \exp\left(\int_s^{\zeta_{k(t)}} a(u)du\right) b(s)ds < -1, \end{cases}$$

hold for $t \in [t_{k(t)}, t_{k(t)+1})$, $\forall t, t_k \geq M$, where $k = k(t)$ and $M \in \mathbb{N}$ sufficiently large.

Remark 4. *Condition (6.2) is necessary, because $w(t, t_{k(t)})$ involves*

$$j(t, \zeta_{k(t)})j^{-1}(t_{k(t)}, \zeta_{k(t)}) = \left(\frac{1 + \int_{\zeta_{k(t)}}^t \exp\left(\int_s^{\zeta_{k(t)}} a(r)dr\right) b(s)ds}{1 + \int_{\zeta_{k(t)}}^{t_{k(t)}} \exp\left(\int_s^{\zeta_{k(t)}} a(r)dr\right) b(s)ds} \right),$$

$$t, \zeta_{k(t)} \in I_k = [t_k, t_{k+1}), k \geq k(\tau), k \in \mathbb{N}.$$

Proof.

We will proceed as in [4] and [6] with slight changes due to the impulsive effect:

- a) *If the discrete solution $z(t_{k(t)})$ of (6.1) is oscillatory, then (6.1) it will be because the discrete solution is part of the whole solution. Formally, by (5.6) with $t = t_{k(t)}$ for $t \in [t_{k(t)}, t_{k(t)+1})$, we have $z(t) = z(t_{k(t)})$. I.e., (5.7). Then, for the classical definition of difference equations, we have that $z(t)$ is oscillatory because $z(t_{k(t)})$ is oscillatory.*

- b) By (5.3) and (5.7), the solution of (6.1) can be rewritten, in terms of w_n and $w(t_{r+1}, t_r)$ as

$$z(t_{k(t)}) = \exp\left(\int_{t_{k(\tau)+1}}^{t_{k(t)}} a(u)du\right) \left(\prod_{r=k(\tau)+1}^{k(t)} w_r\right) z(t_{k(\tau)+1}),$$

where $w_r = (1 + c_r)w(t_{r+1}, t_r)$ and

$$\begin{aligned} w(t_{r+1}, t_r) &= e(t_{r+1}, \zeta_r)e^{-1}(t_r, \zeta_r) \\ &= \exp\left(\int_{t_r}^{t_{r+1}} a(u)du\right) \left(\frac{1 + \int_{\zeta_r}^{t_{r+1}} \exp\left(\int_u^{\zeta_r} a(r)dr\right) b(u)du}{1 + \int_{\zeta_r}^{t_r} \exp\left(\int_u^{\zeta_r} a(r)dr\right) b(u)du}\right). \end{aligned}$$

It is clear that $z(t_{k(t)})$ is oscillatory if $\{w_n\}$ is not eventually positive or negative. Hence, by part a), $z(t)$ is oscillatory if $\{w_n\}_{n \geq n(\tau)+1}$ is not eventually positive nor eventually negative.

- c) (\Rightarrow) If $z(t_n)_{n \geq n(\tau)}$ is nonoscillatory, we can suppose $z(t_n) > 0$ and $z(t) > 0$ for all $t, n \geq M$ with $M \in \mathbb{N}$ sufficiently large. Also, by (5.12) with $\tau = t_n$ for $t \in [t_n(t), t_{n(t)+1})$ and $t \geq M$, we have

$$z(t) = w(t, t_n)z(t_n),$$

where

$$w(t, t_n) = \exp\left(\int_{t_n}^t a(u)du\right) \left(\frac{1 + \int_{\zeta_n}^t \exp\left(\int_s^{\zeta_n} a(u)du\right) b(s)ds}{1 + \int_{\zeta_n}^{t_n} \exp\left(\int_s^{\zeta_n} a(u)du\right) b(s)ds}\right),$$

for $t \in [t_n, t_{n+1})$.

Then, we have $w(t, t_n) > 0$, i.e. (6.2). The case $z(t_n) < 0$ and $z(t) < 0$ is analog.

(\Leftarrow) Suppose that $z(t_n) > 0$, for all $n \geq M$ sufficiently large and (6.2) holds. We have to prove that $z(t)$ is nonoscillatory. We will proceed by contradiction: assume that $z(t)$ is oscillatory. Then, according to the definition of oscillatory solutions, there must exist $(a_n), (b_n)$ sequences such that $(a_n), (b_n) \rightarrow \infty$ as $t \rightarrow \infty$ and $z(a_n) \leq 0 \leq z(b_n)$. Let $t_n < a_n < t_{n+1}$. From (5.12) with $\tau = t_n$ and $t = a_n$, we get

$$z(a_n) = w(a_n, t_n)z(t_n).$$

Since $z(a_n) \leq 0$ and $z(t_n) > 0$, we conclude that $w(a_n, t_n) \leq 0$, and this fact contradicts to (6.2). The case $z(t_n) < 0$ for $n \geq M$ is analog, taking $z(b_n) \geq 0$ with $t_n < b_n < t_{n+1}$. I.e.,

$$z(b_n) = w(b_n, t_n)z(t_n).$$

Again, since $z(b_n) \geq 0$ and $z(t_n) < 0$, we conclude that $w(b_n, t_n) \leq 0$, and this fact also contradicts to (6.2). \square

Next, inspired by [22], we will state our main result, a version of the Wiener-Aftabizadeh criterion for oscillatory solutions of (6.1):

Theorem 7. (Main Result: Aftabzadeh-Wiener IDEPCAG oscillatory criterion) Let the scalar IDEPCAG (6.1) with $1 + c_k \neq 0$ for all $k \geq M$, $M \in \mathbb{N}$ and M sufficiently large. If either of the conditions:

$$\text{if } 1 + c_{k(t)} > 0 \text{ and } \begin{cases} \limsup_{k \rightarrow \infty} \int_{t_{k(t)}}^{\zeta_{k(t)}} \exp \left(\int_s^{\zeta_{k(s)}} a(u) du \right) b(s) ds > 1, \\ \text{or} \\ \liminf_{k \rightarrow \infty} \int_{\zeta_{k(t)}}^{t_{k(t)+1}} \exp \left(\int_s^{\zeta_{k(s)}} a(u) du \right) b(s) ds < -1, \end{cases}$$

or

$$\text{if } 1 + c_{k(t)} < 0 \text{ and } \begin{cases} \liminf_{k \rightarrow \infty} \int_{t_{k(t)}}^{\zeta_{k(t)}} \exp \left(\int_s^{\zeta_{k(s)}} a(u) du \right) b(s) ds < 1, \\ \text{or} \\ \limsup_{k \rightarrow \infty} \int_{\zeta_{k(t)}}^{t_{k(t)+1}} \exp \left(\int_s^{\zeta_{k(s)}} a(u) du \right) b(s) ds > -1. \end{cases}$$

hold for $t \geq k$, with $k = k(t)$ and $\gamma(t) = \zeta_{k(t)}$ if $t \in I_{k(t)} = [t_{k(t)}, t_{k(t)+1}]$. Then, every solution $z(t)$ of (6.1) oscillates.

Proof. If (6.1) has nonoscillatory solutions, then $z(t) > 0$ or $z(t) < 0$ for $k = k(t)$ sufficiently large, for all $k(t) \geq M$ sufficiently large and $\gamma(t) = \zeta_{k(t)}$ if $t \in I_{k(t)} = [t_{k(t)}, t_{k(t)+1}]$. For the proof, we will consider the advanced and delayed intervals $I_k^+ = [t_k, \zeta_k)$ and $I_k^- = [\zeta_k, t_{k+1})$, respectively. Let's suppose that $z(t) > 0$ for $t \in I_k^-$. Then, it is easy to see that

$$(6.3) \quad \left(z(t) \exp \left(\int_t^{\zeta_k} a(u) du \right) \right)' = b(t) z(\gamma(t)) \exp \left(\int_t^{\zeta_k} a(u) du \right).$$

Integrating the last expression from ζ_k to t_{k+1} we have

$$z(t_{k+1}^-) \exp \left(\int_{t_{k+1}}^{\zeta_k} a(u) du \right) = z(\zeta_k) \left(1 + \int_{\zeta_k}^{t_{k+1}} \exp \left(\int_s^{\zeta_{k(s)}} a(u) du \right) b(s) ds \right).$$

Next, applying the impulsive condition $(1 + c_{k(t)+1})z(t_{k+1}^-) = z(t_{k+1})$, we get

$$z(t_{k+1}) \exp \left(\int_{t_{k+1}}^{\zeta_k} a(u) du \right) = (1 + c_{k+1})z(\zeta_k) \left(1 + \int_{\zeta_k}^{t_{k+1}} \exp \left(\int_s^{\zeta_{k(s)}} a(u) du \right) b(s) ds \right)$$

By hypothesis, $z(t_{k+1}), z(\zeta_k) > 0$. Hence, if $(1 + c_{k+1}) > 0$, then we have

$$\int_{\zeta_k}^{t_{k+1}} \exp \left(\int_s^{\zeta_{k(s)}} a(u) du \right) b(s) ds > -1.$$

I.e.,

$$\liminf_{k \rightarrow \infty} \int_{\zeta_k}^{t_{k+1}} \exp \left(\int_s^{\zeta_{k(s)}} a(u) du \right) b(s) ds \geq -1,$$

Analogously, if $1 + c_{k+1} < 0$, we can conclude

$$\limsup_{k \rightarrow \infty} \int_{\zeta_k}^{t_{k+1}} \exp \left(\int_s^{\zeta_{k(s)}} a(u) du \right) b(s) ds \leq -1,$$

which are a contradiction. Hence, the case $t \in I_k^-$ is proved.

Next, we proceed with the case $t \in I_k^+ = [t_k, \zeta_k]$. In the same way as before, integrating (6.3) from t_k to ζ_k we have

$$z(t_k) \exp \left(\int_{t_k}^{\zeta_k} a(u) du \right) = z(\zeta_k) \left(1 - \int_{t_k}^{\zeta_k} \exp \left(\int_s^{\zeta_k(s)} a(u) du \right) b(s) ds \right).$$

Next, applying the impulsive condition $(1 + c_k)z(t_k^-) = z(t_k)$, we get

$$(1 + c_k)z(t_k^-) \exp \left(\int_{t_k}^{\zeta_k} a(u) du \right) = z(\zeta_k) \left(1 - \int_{t_k}^{\zeta_k} \exp \left(\int_s^{\zeta_k(s)} a(u) du \right) b(s) ds \right)$$

By hypothesis, $z(t_k^-), z(\zeta_k) > 0$. Hence, if $(1 + c_k) > 0$, then we have

$$\int_{t_k}^{\zeta_k} \exp \left(\int_s^{\zeta_k(s)} a(u) du \right) b(s) ds < 1.$$

I.e.,

$$\limsup_{k \rightarrow \infty} \sup_{k \in \mathbb{N}} \int_{\zeta_k}^{t_k} \exp \left(\int_s^{\zeta_k(s)} a(u) du \right) b(s) ds \leq 1,$$

Analogously, if $1 + c_k < 0$, we can conclude

$$\liminf_{k \rightarrow \infty} \inf_{k \in \mathbb{N}} \int_{t_k}^{\zeta_k} \exp \left(\int_s^{\zeta_k(s)} a(u) du \right) b(s) ds \geq 1,$$

which are a contradiction. Hence, the case $t \in I_k^+$ is proved.

The proof is analog for $z(t) < 0$ and t sufficiently large. Hence, (6.1) has oscillatory solutions only. \square

Remark 5.

- For the correspondig DEPCAG result, consider $c_k = 0, \forall k \in \mathbb{Z}$.
- If $c_k = 0, \forall k \in \mathbb{Z}$, we recover the results of K-S. Chiu and M. Pinto given in [6].

Corollary 2. *Let the following IDEPCAG*

$$(6.4) \quad \begin{aligned} y'(t) &= b(t)y(\gamma(t)), & t \neq t_k \\ y(t_k) &= (1 + c_k)y(t_k^-), & t = t_k. \end{aligned}$$

As a consequence of Theorem (7)

$$\text{if } 1 + c_{k(t)} > 0 \text{ and } \begin{cases} \limsup_{k \rightarrow \infty} \sup_{k \in \mathbb{N}} \int_{t_{k(t)}}^{\zeta_{k(t)}} b(s) ds > 1, \\ \text{or} \\ \liminf_{k \rightarrow \infty} \inf_{k \in \mathbb{N}} \int_{\zeta_{k(t)}}^{t_{k(t)+1}} b(s) ds < -1, \end{cases}$$

or

$$\text{if } 1 + c_{k(t)} < 0 \text{ and } \begin{cases} \liminf_{k \rightarrow \infty} \inf_{k \in \mathbb{N}} \int_{t_{k(t)}}^{\zeta_{k(t)}} b(s) ds < 1 \\ \text{or} \\ \limsup_{k \rightarrow \infty} \sup_{k \in \mathbb{N}} \int_{\zeta_{k(t)}}^{t_{k(t)+1}} b(s) ds > -1, \end{cases}$$

hold for $t \geq k$, with $k = k(t)$ and $\gamma(t) = \zeta_{k(t)}$, then every solution $y(t)$ of (6.4) is oscillatory.

In a very similar way to Theorem 7, we can deduce the following result:

Corollary 3. (Aftabizadeh-Wiener IDEPCAG nonoscillatory criterion)

Let the scalar IDEPCAG (6.1) with $1 + c_k \neq 0$ for all $k \geq k(t)$. If the conditions:

$$\text{if } 1 + c_{k(t)} > 0 \text{ and } \begin{cases} \limsup_{k \rightarrow \infty} \sup_{k \in \mathbb{N}} \int_{t_{k(t)}}^{\zeta_{k(t)}} \exp \left(\int_s^{\zeta_{k(s)}} a(u) du \right) b(s) ds \leq 1, \\ \text{and} \\ \liminf_{k \rightarrow \infty} \inf_{k \in \mathbb{N}} \int_{\zeta_{k(t)}}^{t_{k(t)+1}} \exp \left(\int_s^{\zeta_{k(s)}} a(u) du \right) b(s) ds \geq -1, \end{cases}$$

or

$$\text{if } 1 + c_{k(t)} < 0 \text{ and } \begin{cases} \liminf_{k \rightarrow \infty} \inf_{k \in \mathbb{N}} \int_{t_{k(t)}}^{\zeta_{k(t)}} \exp \left(\int_s^{\zeta_{k(s)}} a(u) du \right) b(s) ds \geq 1, \\ \text{and} \\ \limsup_{k \rightarrow \infty} \sup_{k \in \mathbb{N}} \int_{\zeta_{k(t)}}^{t_{k(t)+1}} \exp \left(\int_s^{\zeta_{k(s)}} a(u) du \right) b(s) ds \leq -1. \end{cases}$$

hold for $t \geq k$, with $k = k(t)$ sufficiently large and $\gamma(t) = \zeta_{k(t)}$ if $t \in I_{k(t)} = [t_{k(t)}, t_{k(t)+1})$, then every solution $z(t)$ of (6.1) is nonoscillatory.

7. SOME EXAMPLES OF OSCILLATORY LINEAR IDEPCAG SYSTEMS

Example 2. Let the following IDEPCA:

$$(7.1) \quad x'(t) = (\alpha - 1)x([t]), \quad t \neq n$$

$$(7.2) \quad x(n) = \beta x(n^-), \quad t = n, \quad n \in \mathbb{N},$$

where $t_k = \zeta_k = k$, $k \in \mathbb{N}$, and $\alpha, \beta \neq 1$ (the interesting cases). In virtue of Theorem 7, the solutions of (7.1) are oscillatory if $\alpha > 0$ and $\beta < 0$ or $\alpha < 0$ and $\beta > 0$. I.e. $\alpha\beta < 0$.

Example 3. In [11], I. Györi and G. Ladas studied the following DEPCAs

$$(7.3) \quad x'(t) + px(t) + q_0x([t]) = 0, \quad p, q_0 \in \mathbb{R},$$

$$(7.4) \quad y'(t) + py(t) + q_{-1}y([t-1]) = 0, \quad p, q \in \mathbb{R},$$

and they gave the following theorems, using some tools as the characteristic equation for DDE:

Corollary 4. ([11](Cor. 8.1.1 and 8.1.2))

$$(1) \text{ Every solution of (7.3) oscillates if and only if } q_0 \geq \frac{p}{e^p - 1}.$$

$$(2) \text{ Every solution of (7.4) oscillates if and only if } q_{-1} > \frac{pe^{-p}}{4(e^p - 1)}.$$

For (7.3), applying Theorem 7 with $\gamma(t) = [t]$, $a(t) = -p$, $b(t) = -q_0$ and $c_k = 0$, $\forall k \in \mathbb{Z}$, we obtain the following estimation:

$$\frac{q_0}{p} \left(e^{p((k+1)-k)} - 1 \right) > 1 \Rightarrow q_0 > \frac{p}{e^p - 1}.$$

Finally, for (7.4), also using Theorem 7 and

$$\int_{k-1}^{k+1} e^{-p(\gamma(k(s))-s)} ds = \int_{k-1}^k e^{-p((k-2)-s)} ds + \int_k^{k+1} e^{-p((k-1)-s)} ds,$$

we have $q_{-1} > \frac{pe^{-p}}{2(e^p - 1)}$. The bonds found satisfy the ones given in [11].

Remark 6.

- The estimations obtained by I. Gyori and G. Ladas for (7.3) and (7.4) are sharper than ours. Still, we recover the same results without analyzing the roots of a specific characteristic equation.
- In the light of the previous results, it seems that any $\gamma(t)$ such that $\gamma(t) = \zeta_k \notin I_k = [t_k, t_{k+1})$ for $t \in I_k$ could satisfy Theorem 7. ($\gamma(t) = [t - 1]$ implies $\zeta_k = k - 1$ if $t \in [k, k + 1)$, $k \in \mathbb{Z}$). This fact deserves future research.

Example 4. Let the following scalar linear IDEPCAG

$$(7.5) \quad \begin{aligned} z'(t) &= a(t)(z(t) - z(\gamma(t))), & t \neq k \\ z(k) &= c_k z(k^-), & t = k, \quad k \in \mathbb{Z}, \end{aligned}$$

where $a(t)$ is a continuous locally integrable function, $c_k \neq 0, \forall k \in \mathbb{Z}$, $\gamma(t)$ is any generalized piecewise constant argument, i.e., $\gamma(t) = \zeta_k$, if $t \in I_k = [t_k, t_{k+1})$, $k \in \mathbb{Z}$ and $t_k \leq \zeta_k \leq t_{k+1}$.

As

$$\left(\exp \left(- \int_{\zeta_k}^s a(u) du \right) \right)' = \exp \left(- \int_{\zeta_k}^s a(u) du \right) (-a(s)),$$

by Theorem (7), we have the only possible cases occur when $c_{k(t)} < 0$ and they are

$$\text{when } \liminf_{k \rightarrow \infty} \inf_{k \in \mathbb{N}} \left(1 - \exp \left(- \int_{\zeta_k}^{t_k} a(u) du \right) \right) < 1, \text{ or } \limsup_{k \rightarrow \infty} \sup_{k \in \mathbb{N}} \left(1 - \exp \left(- \int_{t_{k+1}}^{\zeta_k} a(u) du \right) \right) < 1.$$

Hence, if one of the last conditions holds, then the solutions of (3.4) are oscillatory.

As we will see, (7.5) is easily solvable. Let $\phi(t)$ be the fundamental solution of

$$(7.6) \quad x'(t) = a(t)x(t).$$

It is well known that $\phi^{-1}(t)$ is the fundamental solution of the adjoint system associated with (7.6). So, it satisfies

$$(\phi^{-1})'(t) = -\phi^{-1}(t)(a(t)).$$

Therefore, we have

$$\begin{aligned} j(t, t_k) &= 1 - \int_{t_k}^t \phi(t_k, s)a(s)ds \\ &= 1 + \phi(t_k) \left(\int_{t_k}^t -\phi^{-1}(s)a(s)ds \right) \\ &= 1 + \phi(t_k) (\phi^{-1}(t) - \phi^{-1}(t_k)) \\ &= \phi^{-1}(t, t_k). \end{aligned}$$

As $e(t, t_k) = \phi(t, t_k)j(t, t_k)$, we have $e(t, t_k) = 1$ (see Theorem 5). Hence, the solution of (7.5) is

$$z(t) = \left(\prod_{j=k(\tau)+1}^{k(t)} c_j \right) z(\tau), \quad t \geq \tau.$$

(See example 1). The last conclusion applies no matter what $a(t)$ locally integrable function is used.




FIGURE 4. Solution of (7.5) with $c_k = -\frac{60}{67}$ and $z(0) = -11$.

Remark 7. The IDEPCA presented in (3.4) is a particular case of the last IDEPCAG, with $\gamma(t) = [t + 1]$.

Example 5. Let the following scalar linear IDEPCAG

$$(7.7) \quad \begin{aligned} x'(t) &= -ax(t) + \text{sen}(2\pi t)x([t]), & t \neq k, \\ x(k) &= Cx(k^-), & t = k, \quad k \in \mathbb{N}, \\ x(0) &= 1. \end{aligned}$$

We see that $\int_k^{k+1} \exp(-a(k-s)) \sin(2\pi s) ds = \frac{2\pi}{a^2 + 4\pi^2} (1 - e^a)$.

By Theorem 7 and Theorem 3 we have

$$\begin{cases} \text{if } C > 0 \text{ and } \begin{cases} a > 2.07553 \Rightarrow \text{All the solutions are oscillatory.} \\ a \leq 2.07553 \Rightarrow \text{All the solutions are nonoscillatory.} \end{cases} \\ \text{if } C < 0, \text{ all the solutions are oscillatory.} \end{cases}$$

ACKNOWLEDGEMENTS

The author sincerely thanks Prof. Manuel Pinto for the encouragement to work in this subject and for all the support during my career. The author also thanks DESMOS PBC for granting permission to use the images employed in this work. They were created with the DESMOS graphic calculator <https://www.desmos.com/calculator>.




FIGURE 5. Solution of (7.7) with $x(0) = 1, C = -100$ and $a = 1.998$.

REFERENCES

1. Marat Akhmet, *Principles of discontinuous dynamical systems*, Springer, New York, Dordrecht, Heidelberg, London, 2010.
2. ———, *Nonlinear hybrid continuous-discrete-time models*, Atlantis Press, Amsterdam-Paris, 2011.
3. Samuel Castillo, Manuel Pinto, and Ricardo Torres, *Asymptotic formulae for solutions to impulsive differential equations with piecewise constant argument of generalized type*, Electron. J. Differential Equations **2019** (2019), no. 40, 1–22.
4. K.-S. Chiu, *Stability of oscillatory solutions of differential equations with general equations with piecewise constant argument*, Electronic Journal of Qualitative Theory of Differential Equations (2011), no. 88, 1 – 15.
5. K.-S. Chiu and I. Berna, *Nonautonomous impulsive differential equations of alternately advanced and retarded type*, Filomat **37** (2023), 7813–7829.
6. K.-S. Chiu and M. Pinto, *Oscillatory and periodic solutions in alternately advanced and delayed differential equations*, Carpathian Journal of Mathematics **29** (2013), no. 2, 149–158.
7. Kenneth Cooke and Stavros Busenberg, *Models of vertically transmitted diseases with sequential-continuous dynamics*, Nonlinear Phenomena in Mathematical Sciences: Proceedings of an International Conference on Nonlinear Phenomena in Mathematical Sciences, Held at the University of Texas at Arlington, , June 16-20, 1980 (New York), Academic Press, 1982, pp. 179–187.
8. L. Dai and M.C. Singh, *On oscillatory motion of spring-mass systems subjected to piecewise constant forces*, Journal of Sound and Vibration **173** (1994), no. 2, 217–231.
9. Liming Dai, *Nonlinear dynamics of piecewise constant systems and implementation of piecewise constant arguments*, World Scientific Press Publishing Co, New York, 2008.
10. Saber Elaydi, *An introduction to difference equations (undergraduate texts in mathematics) 3rd edition*, Springer, 2005.
11. I. Györi and G. Ladas, *Oscillation theory of delay differential equations: with applications*, Oxford Mathematical Monographs, Oxford University Press, USA, 1992.
12. Fatma Karakoc, Arzu Unal, and Huseyin Bereketoglu, *Oscillation of a nonlinear impulsive differential equation system with piecewise constant argument*, Advances in Difference Equations **2018** (2018), no. 1, 99.
13. Anatoly Myshkis, *On certain problems in the theory of differential equations with deviating argument*, Russian Mathematical Surveys **32** (1977), no. 2, 173–203.
14. Manuel Pinto, *Cauchy and Green matrices type and stability in alternately advanced and delayed differential systems*, Journal of Difference Equations and Applications **17** (2011), no. 2, 235–254.

15. Anatoly Samoilenko, *Impulsive differential equations*, World Scientific Press, Singapur, 1995.
16. R. Torres, *On the linear nonautonomous impulsive differential equations with piecewise constant arguments of generalized type*, (submitted) (2023).
17. Ricardo Torres, *Ecuaciones diferenciales con argumento constante a trozos del tipo generalizado con impulso*, Msc. thesis, Facultad de Ciencias, Universidad de Chile, Santiago, Chile, 2015, Available at <https://repositorio.uchile.cl/handle/2250/188898>.
18. Ricardo Torres, Samuel Castillo, and Manuel Pinto, *How to draw the graphs of the exponential, logistic, and gaussian functions with pencil and ruler in an accurate way*, *Proyecciones, Journal of Mathematics*. (to appear) (2023).
19. Ricardo Torres, Manuel Pinto, Samuel Castillo, and Marco Kostić, *An uniform approximation of an impulsive cnn-hopfield type system by an impulsive differential equation with piecewise constant argument of generalized type on $[\tau, \infty)$* , *Acta Appl. Math.* **8** (2021), no. 171, 1–15.
20. Ricardo Torres, Manuel Pinto, and Ferenc Hartung, *Uniform and exponential approximation error decay of a delayed cnn-hopfield type system with piecewise constant arguments*, (submitted) (2023).
21. J. Wiener, *Generalized Solutions of Functional Differential Equations*, World Scientific, 1993.
22. Joseph Wiener and Abdol Reza Aftabizadeh, *Differential equations alternately of retarded and advanced type*, *Journal of Mathematical Analysis and Applications* **1** (1988), no. 129, 243–255.
23. Zhaolin Yan and Jianfang Gao, *Numerical oscillation and non-oscillation analysis of the mixed type impulsive differential equation with piecewise constant arguments*, *International Journal of Computer Mathematics* (2023), 1–18.

INSTITUTO DE CIENCIAS FÍSICAS Y MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD AUSTRAL DE CHILE, CAMPUS ISLA TEJA S/N, VALDIVIA, CHILE.

INSTITUTO DE CIENCIAS, UNIVERSIDAD NACIONAL DE GENERAL SARMIENTO, LOS POLVORINES, BUENOS AIRES, ARGENTINA

Email address: `ricardo.torres@uach.cl`