THE DISTRIBUTION OF *a*-NUMBERS OF HYPERELLIPTIC CURVES IN CHARACTERISTIC THREE

DEREK GARTON, JEFFREY LIN THUNDER, AND COLIN WEIR

ABSTRACT. In this paper we present a new approach to counting the proportion of hyperelliptic curves of genus g defined over a finite field \mathbb{F}_q with a given a-number. In characteristic three this method gives exact probabilities for curves of the form $Y^2 = f(X)$ with $f(X) \in \mathbb{F}_q[X]$ monic and cubefree, probabilities that match the data presented by Cais et al. in previous work. These results are sufficient to derive precise estimates (in terms of q) for these probabilities when restricting to squarefree f. As a consequence, for positive integers a and g we show that the nonempty strata of the moduli space of hyperelliptic curves of genus g consisting of those curves with a-number a are of codimension 2a - 1. This contrasts with the analogous result for the moduli space of abelian varieties in which the codimensions of the strata are a(a + 1)/2. Finally, our results allow for an alternative heuristic conjecture to that of Cais et al.; one that matches the available data.

Contents

1. Introduction	1
1.1. Main results	2
1.2. Previous work	4
1.3. Organization	5
1.4. Competing interests declaration	5
2. Preliminary results	5
2.1. The Cartier operator and <i>a</i> -numbers	5
2.2. Heights and <i>a</i> -numbers	8
3. Heights of subspaces	11
3.1. Intermediate counting results	11
3.2. Main counting results	19
4. The <i>a</i> -numbers of hyperelliptic curves in characteristic three	22
4.1. The proportion of curves with a given <i>a</i> -number	23
4.2. A heuristic model and conjecture	28
4.3. Results on the moduli space of hyperelliptic curves	29
Acknowledgments	30
References	31

1. INTRODUCTION

Let p be a prime, let q be a power of p, and let \mathbb{F}_q the finite field with q elements. Let X be transcendental over \mathbb{F}_q so that $\mathbb{F}_q(X)$ is a field of rational functions. If C is a smooth

Date: March 4, 2024.

algebraic curve over \mathbb{F}_q , the *a*-number of *C* is an invariant of the *p*-torsion group scheme $\operatorname{Jac}(C)[p]$ which we will denote by a(C); it is defined as

$$a(C) = \dim_{\mathbb{F}_q} (\operatorname{Hom} (\operatorname{Spec}_{\operatorname{\mathsf{Grp}}/\mathbb{F}_q}(\mathbb{F}_q[X]/X^p), \operatorname{Jac} (C)[p])).$$

If C has genus g, then $0 \le a(C) \le g$ (see, e.g., [Oda69]). When g = 1, an elliptic curve is called supersingular if and only if a(C) = 1. An elliptic curve is called ordinary otherwise. Analogously, we say a curve C is ordinary if a(C) = 0, as this is generic. At the other extreme, in [Oor75] Oort shows that a curve C is isomorphic to a product of g supersingular elliptic curves if and only if a(C) = g. The purpose of this paper is to study the distribution of a-numbers of hyperelliptic curves to address such questions as: what is the probability a hyperelliptic curve is ordinary?

For any monic cubefree $f \in \mathbb{F}_q[X]$ let H_f denote the (possibly singular) variety over \mathbb{F}_q with affine equation

$$Y^2 = f(X).$$

If H_f has smooth model H we will write $a(H_f)$ for a(H). In order to study the distribution of $a(H_f)$ as f varies we adopt the following parameterizing sets: for $\epsilon \in \{1, 2\}$ and $g \in \mathbb{Z}_{\geq 0}$ let

$$\mathcal{P}_{\epsilon}(g) = \{ f \in \mathbb{F}_q[X] \mid f \text{ monic, cubefree, } \deg(f) = 2g + \epsilon \}.$$

We often restrict ourselves to squarefree polynomials, so let

$$\mathcal{P}'_{\epsilon}(g) = \{ f \in \mathcal{P}_{\epsilon}(g) \mid f \text{ squarefree} \}.$$

Furthermore, since we are concerned with the proportions of cubefree and squarefree polynomials which yield varieties of given *a*-number, we set

$$\mu_{\epsilon,g}(a) = \frac{\left|\{f \in \mathcal{P}_{\epsilon}(g) \mid a(H_f) = a\}\right|}{\left|\mathcal{P}_{\epsilon}(g)\right|} \quad \text{and} \quad \mu_{\epsilon,g}'(a) = \frac{\left|\{f \in \mathcal{P}_{\epsilon}'(g) \mid a(H_f) = a\}\right|}{\left|\mathcal{P}_{\epsilon}'(g)\right|}$$

for non-negative integers a. We will explicitly compute $\mu_{\epsilon,g}(a)$ in the case p = 3. Specifically, we prove the following. (Here and throughout we use the bracket notation $[\cdot]$ for the greatest integer function.)

1.1. Main results. In Section 4, we prove the following theorems.

Theorem 1.1.1. Suppose p = 3 and g, a are nonnegative integers. If g = 0, then

$$\mu_{2,g}(a) = \mu_{1,g}(a) = \begin{cases} 0 & \text{if } a > 0 \\ 1 & \text{if } a = 0, \end{cases}$$

so suppose $g \geq 1$.

If $g \equiv 0 \pmod{3}$ and $\epsilon = 1, 2$ or if $g \equiv 2 \pmod{3}$ and $\epsilon = 2$, then

$$\mu_{\epsilon,g}(a) = \begin{cases} 1 - q^{-1} & \text{if } a = 0, \\ q^{-2a+1} (1 - q^{-2}) & \text{if } 0 < a < [(g+1)/3], \\ q^{-2a+1} & \text{if } a = [(g+1)/3], \\ 0 & \text{if } a > [(g+1)/3]. \end{cases}$$

On the other hand, if $g \equiv 1 \pmod{3}$ and $\epsilon = 1, 2$ or if $g \equiv 2 \pmod{3}$ and $\epsilon = 1$, then

THE DISTRIBUTION OF a-NUMBERS OF HYPERELLIPTIC CURVES IN CHARACTERISTIC THREE 3

• if $g \leq 2$, then

$$\mu_{\epsilon,g}(a) = \begin{cases} 0 & \text{if } a > 1\\ q^{-1} \left(\frac{1-q^{-1}}{1-q^{-2}}\right) & \text{if } a = 1, \\ \frac{1-q^{-1}}{1-q^{-2}} & \text{if } a = 0 \end{cases}$$

• and if g > 3, then

$$\mu_{\epsilon,g}(a) = \begin{cases} 1 - q^{-1} & \text{if } a = 0, \\ q^{-2a+1} \left(1 - q^{-2}\right) & \text{if } 0 < a < \left[(g-1)/3\right], \\ q^{-2a+1} \left(\frac{1+q^{-1}-q^{-2}}{1+q^{-1}}\right) & \text{if } a = \left[(g-1)/3\right], \\ q^{-2a+1} \left(\frac{1}{1+q^{-1}}\right) & \text{if } a = \left[(g-1)/3\right] + 1, \\ 0 & \text{if } a > \left[(g-1)/3\right] + 1. \end{cases}$$

In some cases Theorem 1.1.1 immediately implies exact probabilities for the *a*-numbers of squarefree polynomials.

Corollary 1.1.2. Suppose p = 3 and $g, a \in \mathbb{Z}_{\geq 0}$. Write $[\cdot]$ for the least integer function. (1) If $a > \lceil \frac{g}{3} \rceil$, then

$$\mu_{1,g}'(a) = \mu_{2,g}'(a) = 0.$$

(2) If $g \equiv 1 \pmod{3}$ and $a = \lceil \frac{g}{3} \rceil$, then

$$\mu'_{1,g}(a) = q^{-2a+1}$$
 and $\mu'_{2,g}(a) = q^{-2a+1} \left(1 + q^{-1}\right)$.

Corollary 1.1.2 has the following application.

Corollary 1.1.3. Suppose p = 3, $\epsilon \in \{1, 2\}$, and g, a, r are nonnegative integers. (1) If $f \in \mathcal{P}'_{q,\epsilon}(g)$ and a(f) = g - r, then

$$g \le \frac{3r}{2} + 1.$$

(2) If $r \equiv 0 \pmod{2}$ and $g \equiv 1 \pmod{3}$, then there exists $f \in \mathcal{P}'_{q,\epsilon}(g)$ with a(f) = g - r.

More generally, Theorem 1.1.1 implies bounds on probabilities for the *a*-numbers of square-free polynomials for all $a \in \mathbb{Z}_{\geq 0}$.

Corollary 1.1.4. Suppose p = 3, $\epsilon \in \{1, 2\}$ and g, a are nonnegative integers.

(1) Then

(2) If
$$a > 0$$
, then

$$\begin{aligned} \mu_{\epsilon,g}(a) \left(1 + q^{-1}\right) - q^{-1} &\leq \mu_{\epsilon,g}'(a) \leq \mu_{\epsilon,g}(a) \left(1 + q^{-1}\right) \\ &\left|\mu_{\epsilon,g}'(a) - q^{-2a+1}\right| < 2q^{-2a}. \end{aligned}$$

Previously the authors of [CEZB13] introduced a heuristic prediction for the values of $\lim_{g\to\infty} \mu'_{1,g}(a)$ via a model of "random *p*-divisible groups". Upon performing numerical computations they made the intriguing remark that the statistics of the *a*-numbers of random *p*-divisible groups appear to differ from those of hyperelliptic curves, especially when p = 3. Specifically they proved that the limiting probability of random 3-divisible groups with *a*-number 0 is $\prod_{i=1}^{\infty} (1+q^{-i})^{-1}$, while on the other hand their computations suggest that when p = 3 the quantities $\mu'_{1,g}(a)$ might tend to $1-q^{-1}$ as *g* increases. Thus, one consequence

of Theorem 1.1.1 is that by broadening the set of hyperelliptic curves considered, the corresponding probabilities suggested by the p = 3 data of [CEZB13] are true not just in the g limit, but for every $g \ge 3$.

In Section 4.2 we introduce an alternative model for studying *a*-numbers of hyperelliptic curves ("random subspaces of fixed height"). Theorem 4.2.1 shows that the statistics of this model match that of the data computed by [CEZB13]. This theorem emboldens us to make Conjecture 4.2.2, which predicts that for p = 3, $\epsilon \in \{1, 2\}$, and $a \in \mathbb{Z}_{\geq 0}$,

$$\lim_{g \to \infty} \mu'_{\epsilon,g}(a) = \begin{cases} 1 - q^{-1} & \text{if } a = 0\\ q^{-2a+1} (1 - q^{-2}) & \text{if } a > 0. \end{cases}$$

This conjecture has the benefit of encompassing all $a \in \mathbb{Z}_{\geq 0}$. (When a > 0, the data of [CEZB13] were not conclusive enough to suggest a limiting probability.)

Finally, in Section 4.3, we apply these estimates of $\mu'_{\epsilon,g}(a)$ to compute the codimensions of the *a*-number strata of the moduli space of smooth hyperelliptic curves of genus g over $\overline{\mathbb{F}_q}$ in characteristic 3. In particular, we prove in Theorem 4.3.1 that when a is positive these codimensions are 2a - 1. By comparison, in [Oda69] the author showed that the codimensions of the *a*-number strata in the full moduli space of abelian varieties are a(a+1)/2. Interestingly, this discrepancy contrasts p-rank strata of hyperelliptic curves; the results of [GP05, Theorem 1] proves that the codimensions of the p-rank strata of hyperelliptic curves and those of abelian varieties are the same.

1.2. **Previous work.** Many previous investigations into heuristics for hyperelliptic curves build upon work of Friedman and Washington [FW89] who compute statistics of invertible matrices over finite fields and compare these statistics to those of the ℓ -parts of Jacobians of hyperelliptic curves, where ℓ is a prime number satisfying $\ell \neq 2p$. This endeavor is analogous to the Cohen-Lenstra heuristics [CL84], where the authors compute statistics of random finite abelian groups and compare them to those of ideal class groups of quadratic number fields. Moreover, the heuristics of Friedman and Washington [FW89] turn out to be provably correct as $q \to \infty$ with $q \not\equiv 0 \pmod{\ell}$, as shown in [EVW16]. In [Gar15] the author addresses the case where $q \not\equiv 1 \pmod{\ell}$, leaving open the case when $p = \ell$. The heuristic investigation of this case begins with [CEZB13], as detailed above.

Other previous results on the *a*-numbers of hyperelliptic curves include the following. Proposition 3.1 of [Re01] is an upper bound on the *a*-numbers of any smooth, complete, irreducible curve in terms its genus. Theorem 1.1 of [Elk11] improves this theorem, restricting to Kummer covers of the projective line. For hyperelliptic curves over finite fields of characteristic three, Corollary 1.1.2 (1) improves upon both these bounds. Moreover, Corollary 1.1.2 (2) implies that Corollary 1.1.2 (1) is sharp when $g \equiv 1 \pmod{3}$ by giving explicit (positive) values of $\mu'_{1,g}(a)$ and $\mu'_{2,g}(a)$ in this case.

Another common means of addressing *a*-numbers is to ask: for which genera g do there exist hyperelliptic curves of *a*-number g-1? We mention the special cases of several results. If a hyperelliptic curve of genus g over a finite field of characteristic three has *a*-number g-1, then:

- [Re01, Proposition 3.1] implies that $q \leq 6$;
- [Elk11, Corollary 1.2] implies that $g \leq 4$; and
- [Fre18, Theorem 1.1] implies that $g \leq 2$.

As mentioned in the previous subsection, we show in Corollary 1.1.3 that if $r \in \mathbb{Z}_{\geq 0}$ and a hyperelliptic curve of genus g over a finite field of characteristic three has a-number g - r, then

$$g \le \frac{3r}{2} + 1.$$

In particular, we recover Frei's result when r = 1. Moreover, when r is even and $q \equiv 1$ (mod 3), we show that this bound cannot be improved.

Finally, [San20] studies the statistics of *a*-numbers in the families of superelliptic curves in characteristic two and Artin-Schreier curves in any positive characteristic. Interestingly, in odd characteristic the limiting probability that an Artin-Schreier curve has a-number 0 (as the genus of the curves grow) is zero, in contrast to what we conjecture to be the case for hyperelliptic curves in Conjecture 4.2.2.

1.3. Organization. This paper is organized as follows. In Section 2 we review a-numbers and the Cartier operator; some of the results in this section hold when p is odd though we restrict to the case p = 3 to simplify the exposition. In Section 3 we use geometry of numbers for function fields to prove our fundamental height results; the results in this section hold for any prime p. In Section 4 we apply these results to deduce our main theorems. As noted, these results are only for the case p = 3.

1.4. Competing interests declaration. The authors declare no competing interests.

2. Preliminary results

In this section we will assume that p = 3 (though much of it is true in the more general case where p is odd). Everything we do will revolve around particular polynomials attached to a given polynomial $f(X) \in \mathbb{F}_q[X]$. Specifically, for any $f \in \mathbb{F}_q[X]$ let $c_0(f), c_1(f)$, and $c_2(f)$ be the unique polynomials in $\mathbb{F}_q[X]$ such that

$$f(X) = c_0(f)^3 + c_1(f)^3 \cdot X + c_2(f)^3 \cdot X^2.$$

In this section we first restrict to the case where f is squarefree. We then recall the definition of the Cartier operator and see how the a-number $a(H_f)$ may be computed via some linear algebra involving the coefficients $c_i(f)$. We then extend this to the case where f is simply cubefree (this of course matches the previous when f happens to be squarefree). Finally, we recall relevant facts from the theory of heights in Diophantine geometry and show how computing the *a*-number may be reformulated in those terms.

In what follows we let K be a fixed algebraic closure of \mathbb{F}_q .

2.1. The Cartier operator and *a*-numbers. Let $f \in \mathbb{F}_q[X]$ be a monic and squarefree polynomial. We denote the K-module of meromorphic differentials on H_f defined over K by Ω_f^1 .

Definition 2.1.1. For any monic squarefree polynomial $f \in \mathbb{F}_q[X]$, the *Cartier operator* of f is the (unique) map

$$\mathcal{C}_f:\Omega^1_f\to\Omega^1_f$$

that satisfies the following properties: for any $\omega, \omega_1, \omega_2 \in \Omega_f^1$ and $z \in \mathcal{O}_{H_f}$,

• $C_f(\omega_1 + \omega_2) = C_f(\omega_1) + C_f(\omega_2);$ • $C_f(z^p\omega) = z \cdot C_f(\omega);$

•
$$C_f(z^p\omega) = z \cdot C_f(\omega)$$

•
$$C_f(\mathrm{d}z) = 0$$
; and

• $C_f\left(\frac{\mathrm{d}z}{z}\right) = \frac{\mathrm{d}z}{z}.$

Set $\omega = \frac{dX}{Y} \in \Omega_f^1$. Then for any $j \in \mathbb{Z}_{\geq 0}$ Elkin [Elk11, Theorem 3.4] proves that

(1)
$$C_f(X^j\omega) = \begin{cases} X^jc_2(f)\omega & \text{if } j \equiv 0 \mod 3, \\ X^{j-1}c_1(f)\omega & \text{if } j \equiv 1 \mod 3, \\ X^{j-2}c_0(f)\omega & \text{if } j \equiv 2 \mod 3. \end{cases}$$

Let g denote the genus of the curve H_f , i.e., $\deg(f) = 2g + \epsilon$ for $\epsilon = 1, 2$. Then by Eq. (1) we see that the Cartier operator yields a linear transformation of the \mathbb{F}_q -subspace of Ω_f^1 spanned by $\{\omega, X\omega, \ldots, X^{g-1}\omega\}$. Section 5 of [Oda69] shows that the a number $a(H_f)$ is the dimension of the kernel of this transformation.

Now let us consider the case where f is possibly just cubefree. In this case we may uniquely write $f = f_1 f_2^2$ where f_1 and f_2 are relatively prime and both are monic and squarefree. (Of course, the case where $f_2 = 1$ is where f is squarefree.) Here $f = f_1 f_2^3 / f_2$. With this in mind, an entirely similar proof as given by Elkin shows that Eq. (1) generalizes to

(2)
$$\mathcal{C}_{f_1}\left(X^j\omega\right) = \begin{cases} X^j c_2(f)\omega & \text{if } j \equiv 0 \mod 3, \\ X^{j-1}c_1(f)\omega & \text{if } j \equiv 1 \mod 3, \\ X^{j-2}c_0(f)\omega & \text{if } j \equiv 2 \mod 3, \end{cases}$$

where now $\omega = \frac{dX}{Yf_2} \in \Omega_{f_1}^1$. Set $d = \deg(f_2)$ and say $f = f_1 f_2^2 \in \mathcal{P}_1(g) \cup \mathcal{P}_2(g)$, so that the genus of H_f , which is the genus of H_{f_1} , is g - d. Of course $a(H_f) = a(H_{f_1})$, too. We will show the following.

Lemma 2.1.2. Let $f = f_1 f_2^2 \in \mathcal{P}_1(g) \cup \mathcal{P}_2(g)$ with f_1 and f_2 monic, squarefree, and relatively prime. Then $a(H_f) = 0$ whenever $\deg(f) < 3$ (i.e., when g = 0). If $\deg f \ge 3$ then Eq. (2) shows that \mathcal{C}_{f_1} restricts to an action on the \mathbb{F}_q -subspace of $\Omega_{f_1}^1$ spanned by $\{\omega, X\omega, \ldots, X^{g-1}\omega\}$. The a-number $a(H_f)$ is the dimension of the kernel of this action.

Example 2.1.3. Let $f = X^3 - X \in \mathcal{P}'_1(1)$. Since

$$f = (X)^3 + (-1)^3 X$$

we see that $c_0(f) = X$, $c_1(f) = -1$, and $c_2(f) = 0$. We set $\omega = dX/Y$ and apply Eq. (1) to compute

$$\mathcal{C}_f(\omega) = c_2(f)\omega = 0.$$

By Lemma 2.1.2 we conclude that the *a*-number $a(H_f) = 1$.

Example 2.1.4. Let

$$f = f_1 f_2^2 = (X^3 - X) (X^2 + X + 2)^2$$

= $X^7 + 2X^6 + X^5 + 2X^4 + 2X^3 + 2X^2 + 2X$
= $(2X^2 + 2X)^3 + (X^2 + 2X + 2)^3 X + (X + 2)X^2$,

so that $f \in \mathcal{P}_1(3)$, and

$$c_0(f) = 2X^2 + 2X,$$

 $c_1(f) = X^2 + 2X + 2,$ and
 $c_2(f) = X + 2.$

As the squarefree part of f is $f_1 = X^3 - X$ we set $\omega = dX/Y(X^2 + X + 2)$ and use Eq. (2) to compute

$$\mathcal{C}_{f_1}(\omega) = (X+2)\omega,$$

$$\mathcal{C}_{f_1}(X\omega) = (X^2+2X+2)\omega, \text{ and}$$

$$\mathcal{C}_{f_1}(X^2\omega) = (2X^2+2X)\omega.$$

Hence the image of $\{\omega, X\omega, X^2\omega\}$ under C_{f_1} has dimension 2, so that $a(H_f) = a(H_{f_1}) = 3-2 = 1$.

Proof. If deg f < 3, i.e., the genus of H_f is zero, then the lemma is trivially true. We thus suppose deg $f \ge 3$ so that $g \ge 1$. If d = 0 we are in the squarefree case, so that the lemma follows from the work of Oda in [Oda69] as remarked above. Hence we may assume for the remainder that $d \ge 1$.

Let $\alpha_1, \ldots, \alpha_d$ be the (distinct) roots of f_2 in K and set $\omega = \frac{dX}{Y} \in \Omega^1_{f_1}$ as above. Via the partial fraction decomposition we see that

$$\operatorname{span}_{K}\left(\left\{\frac{\mathrm{d}X}{Yf_{2}(X)},\ldots,\frac{X^{d-1}\,\mathrm{d}X}{Yf_{2}(X)}\right\}\right) = \operatorname{span}_{K}\left(\left\{\frac{\mathrm{d}(X-\alpha_{1})}{Y(X-\alpha_{1})},\ldots,\frac{\mathrm{d}(X-\alpha_{d})}{Y(X-\alpha_{d})}\right\}\right)$$

For every $i \in \{1, ..., d\}$ choose $\beta_i \in K$ with $\beta_i^2 = f_1(\alpha_i)$, and let P_i be the point (α_i, β_i) on H_{f_1} . With this notation, we see that for any such i, $\operatorname{ord}_{P_i}\left(\frac{\operatorname{d}(X - \alpha_i)}{(X - \alpha_i)}\right) = -1$. Also, if

$$j \in \{1, \dots, d\} \setminus \{i\}$$
 then $\operatorname{ord}_{P_j}\left(\frac{\operatorname{d}(X - \alpha_i)}{(X - \alpha_i)}\right) = 0$. Finally, $\operatorname{ord}_{P_i}\left(\frac{1}{Y}\right) = 0$ since $\operatorname{gcd}\left(f_1, f_2\right) = 1$.

We now recall (see Exercise 4.14 of [Sti09], for example) that if ω is holomorphic, then so is $C_f(\omega)$, and that if P is a point on H_f and $\operatorname{ord}_P(\omega) = -1$, then $\operatorname{ord}_P(\mathcal{C}_f(\omega)) = -1$. Thus \mathcal{C}_{f_1} acts injectively on $\operatorname{span}_K\left(\left\{\frac{\mathrm{d}X}{Y_{f_2}(X)}, \ldots, \frac{X^{d-1}\,\mathrm{d}X}{Y_{f_2}(X)}\right\}\right)$. In particular, if d = g then the genus of H_f is zero and so is the *a*-number $a(H_f)$, which is also the dimension of the kernel of our Cartier operator linear transformation. Thus we now suppose that d < g.

Via long division and the partial fraction decomposition we see that

$$\operatorname{span}_{K}\left(\left\{\frac{\mathrm{d}X}{Yf_{2}(X)}, \dots, \frac{X^{g-1}\,\mathrm{d}X}{Yf_{2}(X)}\right\}\right)$$
$$= \operatorname{span}_{K}\left(\left\{\frac{\mathrm{d}X}{Yf_{2}(X)}, \dots, \frac{X^{d-1}\,\mathrm{d}X}{Yf_{2}(X)}\right\} \bigcup \left\{\frac{\mathrm{d}X}{Y}, \dots, \frac{X^{g-d-1}\,\mathrm{d}X}{Y}\right\}\right)$$
$$= \operatorname{span}_{K}\left(\left\{\frac{\mathrm{d}(X-\alpha_{1})}{Y(X-\alpha_{1})}, \dots, \frac{\mathrm{d}(X-\alpha_{d})}{Y(X-\alpha_{d})}\right\} \bigcup \left\{\omega, X\omega, \dots, X^{g-d-1}\omega\right\}\right).$$

As the genus of H_{f_1} is g - d, we know that $\omega, X\omega, \ldots, X^{g-d-1}\omega$ are holomorphic. Thus, it follows that

$$\{0\} = \operatorname{span}_{K}\left(\left\{\frac{\operatorname{d}(X-\alpha_{1})}{Y(X-\alpha_{1})}, \dots, \frac{\operatorname{d}(X-\alpha_{d})}{Y(X-\alpha_{d})}\right\}\right) \cap \operatorname{span}_{K}\left(\left\{\omega, X\omega, \dots, X^{g-d-1}\omega\right\}\right),$$

finishing the proof.

Lemma 2.1.5. Let $f = f_1 f_2^2 \in \mathbb{F}_q[X]$ be as above and let

$$Q(X)\omega = \sum_{i=0}^{g-1} a_i X^i \omega$$

be an element of the \mathbb{F}_q -subspace of $\Omega^1_{f_1}$ in Lemma 2.1.2. Then $Q(X)\omega$ is in the kernel of C_{f_1} if and only if

(3)
$$c_2(Q)c_0(f) + c_1(Q)c_1(f) + c_0(Q)c_2(f) = 0.$$

Proof. Via Eq. (2) we readily verify the following:

 $C_{f_1}(X^{3j}\omega) = c_2(f)X^j\omega, \qquad C_{f_1}(X^{3j+1}\omega) = c_1(f)X^j\omega, \qquad C_{f_1}(X^{3j+2}\omega) = c_0(f)X^j\omega.$

The lemma follows.

Combining the two lemmas above yields the main result of this subsection. (We adopt the usual convention here that the degree of the zero polynomial is $-\infty$.)

Proposition 2.1.6. Let p = 3 and $g \in \mathbb{Z}_{\geq 0}$. Then for all $f \in \mathcal{P}_1(g) \cup \mathcal{P}_2(g)$ the a-number $a(H_f)$ is the dimension of the \mathbb{F}_q -vector space consisting of those $Q(X)\omega$ where Q satisfies Eq. (3) and

$$\deg\left(c_2(Q)\right) \le \frac{g-3}{3}, \qquad \deg\left(c_1(Q)\right) \le \frac{g-2}{3}, \qquad \deg\left(c_0(Q)\right) \le \frac{g-1}{3}$$

2.2. Heights and *a*-numbers. To ease notation we will write F for the field of rational functions $\mathbb{F}_{q}(X)$ in this subsection. We first recall some more elementary notions for the field F. Of course height machinery encompasses a much wider range of "global fields," but here we will limit ourselves to the needs at hand.

It is well known that the places of F are in one-to-one correspondence with the set of monic irreducible polynomials $P \in \mathbb{F}_{q}[X]$ together with the "place at infinity" v_0 corresponding to the usual degree function. The degree of the place v_0 is one; the degree of the place corresponding to the monic irreducible P is simply the degree of P as a polynomial. For each place v we have the usual notion of the order ord_{v} (the order at v_{0} of a rational function being the negative of its degree, the order at the place corresponding to P being the exact power of P dividing the rational function). We will write Div(F) for the divisor group - the free abelian group generated by the set of places of F. For any non-zero element $\alpha \in F$ we get a well-defined principal divisor $\operatorname{div}(\alpha) = \sum_{v} \operatorname{ord}_{v}(\alpha) \cdot v$. The "product formula" in this particular case states that the degree of any principal divisor is zero:

$$\deg\left(\operatorname{div}(\alpha)\right) = \sum_{v} \operatorname{ord}_{v}(\alpha) \operatorname{deg}(v) = 0.$$

This is readily verified in our case here.

We extend the notion of order to *n*-tuples as follows:

$$\operatorname{ord}_{v}(\alpha_{1},\ldots,\alpha_{n}) = \min\{\operatorname{ord}_{v}(\alpha_{1}),\ldots,\operatorname{ord}_{v}(\alpha_{n})\}.$$

This in turn yields a well-defined divisor (no longer necessarily principal!):

$$\operatorname{div}(\alpha) = \sum_{v} \operatorname{ord}_{v}(\alpha) \cdot v.$$

Via this we have the definition of the usual height of a non-zero vector $\alpha \in F^n$:

 $h(\alpha) = -\deg\big(\operatorname{div}(\alpha)\big).$

Since the degree of a principal divisor is zero, we immediately see that this height function h is really a function on one-dimensional subspaces of F^n .

All the above can be extended to arbitrary subspaces, but for us here we only need the fact that the height h(V) of an (n-1)-dimensional hyperplane $V \subset F^n$ is the height of its orthogonal complement. In other words, if V is the subspace of β such that $\alpha \cdot \beta = 0$, then $h(V) = h(\alpha)$. We also set the height of the full space F^n and its orthogonal complement $\{0\}$ to both be zero.

Example 2.2.1. Via an appropriate scalar multiplication, any non-zero $\alpha \in F^n$ is projectively equivalent to a non-zero *n*-tuple of polynomials (A_1, \ldots, A_n) whose greatest common divisor is 1. For such a tuple, we clearly have $\operatorname{ord}_P(A_1, \ldots, A_n) = 0$ for any place P where P is a monic irreducible polynomial. For the place at infinity v_0 we have

$$\operatorname{ord}_{v_0}(A_1, \ldots, A_n) = \min\{-\deg(A_1), \ldots, -\deg(A_n)\}.$$

Thus

$$h(A_1,\ldots,A_n) = \max\{\deg(A_1),\ldots,\deg(A_n)\}.$$

Moreover, the height of the orthogonal complement $\{\beta : \beta \cdot \alpha = 0\}$ is also the maximum of the degrees of the A_i 's.

Given the example above, the following lemma indicates why one is lead to heights when contemplating the *a*-number $a(H_f)$. (We note for posterity that the obvious generalization of this lemma is valid for all odd primes p, not just p = 3.)

Lemma 2.2.2. For all $f_1, f_2 \in \mathbb{F}_q[X]$, $c_i(g) = f_2c_i(f_1)$ for i = 0, 1, 2 if and only if $g = f_1f_2^3$. In particular, f is cubefree if and only if the coefficients $c_0(f)$, $c_1(f)$, $c_2(f)$ are relatively prime and in that case the height

$$h(c_0(f), c_1(f), c_2(f)) = \max\{\deg(c_0(f)), \deg(c_1(f)), \deg(c_2(f))\}$$

Proof. Clearly it suffices to prove this in the case where f_2 is just a monomial: $f_2 = X^j$ for some j > 0. One readily verifies that for any $f(X) = a_0 + a_1 X + \cdots \in \mathbb{F}_q[X]$

$$c_0(f) = \sum_{i \equiv 0 \mod 3} a_i X^{\frac{i}{3}}$$
$$c_1(f) = \sum_{i \equiv 1 \mod 3} a_i X^{\frac{i-1}{3}}$$
$$c_2(f) = \sum_{i \equiv 2 \mod 3} a_i X^{\frac{i-2}{3}}.$$

The case where $f_2 = X^j$ follows directly from this.

Given any subspace $V \subseteq F^n$ there is the notion of *successive minima*. In the case of a two-dimensional space $V \subset F^3$ there are two minima, $\mu_1(V)$, $\mu_2(V)$, defined as

$$\mu_1(V) = \min\{h(\mathbf{x}_1)\} \mid \mathbf{x}_1 \in V \setminus \{\mathbf{0}\}\} \text{ and}$$
$$\mu_2(V) = \min\{\max\{h(\mathbf{x}_1), h(\mathbf{x}_2)\} \mid \{\mathbf{x}_1, \mathbf{x}_2\} \text{ is a basis for } V\}.$$

We will make frequent use use of Minkowski's Theorem ([Thu08, Theorem 1]), which in this instance reads

(4)
$$\mu_1(V) \le \mu_2(V), \quad \mu_1(V) + \mu_2(V) = h(V).$$

Theorem 2.2.3. Set p = 3, let $g \ge 0$ and $\epsilon \in \{1, 2\}$. Set $m = \lfloor (g - 1)/3 \rfloor$ and suppose $f = f_1 f_2^2 \in \mathcal{P}_{\epsilon}(g)$ with f_1 and f_2 relatively prime. If g = 0 then $a(H_f) = 0$.

Suppose $g \ge 1$ and set $V \subset F^3$ to be the 2-dimensional orthogonal complement of $(c_0(f), c_1(f), c_2(f))$ so that

$$h(V) = \max\{\deg(c_0(f)), \deg(c_1(f)), \deg(c_2(f))\}.$$

Then

$$h(V) = \begin{cases} \frac{2g+\epsilon}{3} = \deg\left(c_0(f)\right) > \max\{\deg\left(c_1(f)\right), \deg\left(c_2(f)\right)\} & \text{if } 2g+\epsilon \equiv 0 \pmod{3}, \\ \frac{2g+\epsilon-1}{3} = \deg\left(c_1(f)\right) > \deg\left(c_2(f)\right) & \text{if } 2g+\epsilon \equiv 1 \pmod{3}, \\ \frac{2g+\epsilon-2}{3} = \deg\left(c_2(f)\right) & \text{if } 2g+\epsilon \equiv 2 \pmod{3}, \end{cases}$$

and $a(H_f) > 0$ only if $\mu_1(V) \leq m$.

Suppose $Q \in \mathbb{F}_q[X]$ is cube-free and satisfies Eq. (3), i.e., $(c_2(Q), c_1(Q), c_0(Q)) \in V$ and $c_0(Q), c_1(Q), c_2(Q)$ are relatively prime. Suppose further that

$$\mu_1(V) = h(c_2(Q), c_1(Q), c_0(Q)) \le m.$$

Unless we have one of the following three situations:

(1) $g \equiv 1 \pmod{3}, \epsilon = 2, \deg(c_2(Q)) = \mu_1(V);$

- (2) $g \equiv 1 \pmod{3}, \ \epsilon = 1, \ \deg(c_1(Q)) = \mu_1(V);$
- (3) $g \equiv 2 \pmod{3}, \ \epsilon = 1, \ \deg(c_2(Q)) = \mu_1(V);$

then the kernel of C_{f_1} is spanned by

$$\{X^{3j}Q\omega: 0 \le j \le m - \mu_1(V)\}$$

so that the a-number

$$a(H_f) = m - \mu_1(V) + 1.$$

In any of (1), (2), or (3) hold then the kernel of C_{f_1} is spanned by

$$\{X^{3j}Q\omega: 0 \le j \le m - \mu_1(V) - 1\}$$

if $m > \mu_1(V)$ or simply $\{0\}$ if $m = \mu_1(V)$, so that the a-number

$$a(H_f) = m - \mu_1(V).$$

Proof. The first part of the theorem follows directly from Lemma 2.2.2 and the definitions.

For the next part, Proposition 2.1.6 and Example 2.2.1 immediately implies that $a(H_f) > 0$ only if $\mu_1(V) \leq \frac{g-1}{3}$.

Now suppose $\mu_1(V) = h(c_2(Q), c_1(Q), c_0(Q)) \leq \frac{g-1}{3}$ where Q satisfies Eq. (3). To ease notation temporarily write l for $\mu_1(V)$. There are only three possible scenarios where $X^{3i}Q$ don't satisfy the degree requirements in Proposition 2.1.6 for all $0 \leq i \leq m-l$: deg $(X^{m-l}c_2(Q)) = \frac{g-1}{3}$ (and so deg $(c_2(Q)) = l$ and $g \equiv 1 \pmod{3}$), deg $(X^{m-l}c_2(Q)) = \frac{g-2}{3}$ (and so deg $(c_2(Q)) = l$ and $g \equiv 1 \pmod{3}$), deg $(X^{m-l}c_2(Q)) = \frac{g-2}{3}$ (and so deg $(c_2(Q)) = l$ and $g \equiv 1 \pmod{3}$) once again). Note that in all three cases here we do have $X^{3i}Q$ satisfies the degree requirements in Proposition 2.1.6 for all $0 \leq i < m-l$ assuming m-l > 0; if m = l then Q does not satisfy the degree requirements in Proposition 2.1.6. In that first scenario, we can't have deg $(c_0(f))$ strictly larger than deg $(c_1(f))$ and deg $(c_2(f))$, since deg $(c_0(Q))$ is at least as large as deg $(c_1(Q))$ and deg $(c_0(Q))$ by hypothesis and Q satisfies Eq. (3). By what we

have already shown this implies that $2g + \epsilon \not\equiv 0 \pmod{3}$, so that $\epsilon \neq 1$. The second and third scenarios are dealt with in an entirely similar manner.

If we don't have one of the situations (1), (2), or (3) and $\mu_1(V) \leq m$, then we have shown that Q satisfies the degree requirements in Proposition 2.1.6 and thus $Q\omega$ is in the kernel of the Cartier operator. Moreover, if $(c_2(P), c_1(P), c_0(P))$ is any other element of V linearly independent from $(c_2(Q), c_1(Q), c_0(Q))$, then its height is at least $\mu_2(V)$. By Eq. (4) and our calculation of h(V), this implies that max{deg $(c_2(P))$, deg $(c_1(P))$, deg $(c_0(P))$ } > $\frac{g-1}{3}$, thus $P\omega$ is not in the kernel. This shows that all $P \in \mathbb{F}_q[X]$ with $P\omega$ in the kernel of the Cartier operator are $\mathbb{F}_q[X]$ -multiples of Q. The theorem now follows from Lemma 2.2.2 and Proposition 2.1.6.

3. Heights of subspaces

The results from this section hold for finite fields of arbitrary characteristic, so suppose that q is any prime power. We continue to write $F = \mathbb{F}_q(X)$, where X is transcendental over \mathbb{F}_q . In this section, we compute statistics of subspaces of F^3 with specified height.

It will prove useful to allow local changes of coordinates. Specifically, if $\mathcal{A} = (\mathcal{A}_v) \in \operatorname{GL}_n(F_{\mathbb{A}})$ (the indexing here is over the places v of F) is an element of the general linear group over the adele ring $F_{\mathbb{A}}$, then for any non-zero $\alpha \in F^n$ we set the "twisted height" $h_{\mathcal{A}}(\alpha)$ to be

$$h_{\mathcal{A}}(\alpha) = -\deg\left(\sum_{v} \operatorname{ord}_{v}(\mathcal{A}_{v}(\alpha)) \cdot v\right).$$

(Since $\mathcal{A} \in \mathrm{GL}_n(F_{\mathbb{A}})$ the local changes of coordinates only change orders at finitely many places, so that this is well-defined.) In this case we also set

$$h_{\mathcal{A}}({\mathbf{0}}) = 0, \qquad h_{\mathcal{A}}(F^n) = -\deg\left(\sum_{v} \det(\mathcal{A}_v) \cdot v\right).$$

Obviously the usual height h without the subscript is simply the case where \mathcal{A} is the identity.

3.1. Intermediate counting results. Our first result uses height machinery to count the number of 2-dimensional subspaces of F^3 containing a given non-zero vector. We begin with some notation. For any $n \in \mathbb{Z}_{\geq 1}$ and $k \in \mathbb{Z}_{\geq 0}$, let $N_{F^n}(k)$ denote the number of one-dimensional subspaces of F^n of (untwisted) height k. Further, if $W \subset F^3$ is a one-dimensional subspace of F^3 , then we let $N_{F^n/W}(k)$ denote the number of 2-dimensional subspaces $V \supset W$ with height h(V) = h(W) + k.

It turns out that

(5)
$$N_{F^n}(k) = \begin{cases} q^{nk-n+1}(q^n-1)(q^{n-1}-1)/(q-1) & \text{if } k \ge 1, \\ (q^n-1)/(q-1) & \text{if } k = 0. \end{cases}$$

Indeed, this follows from the proof of the following proposition.

Proposition 3.1.1. Suppose that $l \in \mathbb{Z}_{\geq 0}$, that $k \in \mathbb{Z}_{\geq 1}$, and that $W \subset F^3$ is a subspace of height l. Then

$$N_{F^3/W}(k) = q^{2k+l-1}(q^2 - 1).$$

Proof. As noted above, our method of proof applies to Eq. (5) as well; we will make appropriate remarks in this regard throughout the proof.

Let $F_{\mathbb{A}}$ be the adele ring of F. Then according to [Thu08, Lemma 8b] there is an $\mathcal{A} \in GL_2(F_{\mathbb{A}})$ giving a "twisted height" $h_{\mathcal{A}}$ on the two-dimensional factor space F^3/W such that

 $h_{\mathcal{A}}(V/W) = h(V) - h(W)$ for all subspaces $V \supseteq W$. Thus $N_{F^3/W}(k)$ is the number of onedimensional subspaces of F^3/W with twisted height k. Similar to Eq. (4) above we have Minkowski's Theorem:

(6)
$$\mu_1(F^3/W) + \mu_2(F^3/W) = h_{\mathcal{A}}(F^3/W) = h(F^3) - h(W) = -l.$$

We see that $\mu_1(F^3/W) \ge 0 - h(W) = -l$ since $h(V) \ge 0$ for all subspaces $V \subseteq F^3$. Hence

(7)
$$-l \le \mu_1(F^3/W) \le -l/2, \qquad \mu_1(F^3/W) \le \mu_2(F^3/W) \le 0.$$

Note that in the case l = 0 we get $\mu_1(F^3/W) = \mu_2(F^3/W) = 0$. We will use Eq. (7) together with techniques from [Thu08] to evaluate the counting function $N_{F^3/W}$.

Our counting function can be expressed in terms of the dimensions of certain Riemann-Roch spaces and divisors. All necessary background on this may be found in [Thu95, Thu08]. For any divisor $\mathfrak{A} \in \operatorname{Div}(F)$, the divisor group of F, and any $\mathcal{A} \in \operatorname{GL}_n(F_{\mathbb{A}})$ set

$$L(\mathfrak{A},\mathcal{A}) = \{ \mathbf{x} \in F^n \mid \operatorname{ord}_v(\mathcal{A}_v \mathbf{x}) \ge -\operatorname{ord}_v(\mathfrak{A}) \text{ for all } v \in M(F) \},\$$

where M(F) denotes the set of places of the field F. As shown in [Thu95, §II], the set $L(\mathfrak{A}, \mathcal{A})$ is a finite-dimensional vector space over \mathbb{F}_q ; denote its dimension by $l(\mathfrak{A}, \mathcal{A})$. Now suppose \mathcal{A} arises from a factor space F^3/W as above. Since $\mu_2(F^3/W) \leq 0$ by Eq. (7), for any divisor \mathfrak{A} with deg $(\mathfrak{A}) \geq -1$ we have

(8)
$$l(\mathfrak{A}, \mathcal{A}) = 2(\deg(\mathfrak{A}) + 1) + \deg \operatorname{div} \det(\mathcal{A}) = 2(\deg(\mathfrak{A}) + 1) + l$$

by [Thu08, Lemma 10]. Note that the height of the full space $h(F^3/W) = -\deg \operatorname{div} \det(\mathcal{A})$ always. Similarly, since $\mu_1(F^n) = \mu_2(F^n) = \cdots = \mu_n(F^n) = 0 = \deg \operatorname{div} \det(I_n)$ for the identity element $I_n \in \operatorname{GL}_n(F_{\mathbb{A}})$,

(8')
$$l(\mathfrak{A}, I_n) = n(\deg(\mathfrak{A}) + 1)$$

for all divisors \mathfrak{A} with deg $(\mathfrak{A}) \geq -1$.

Fix a divisor \mathfrak{A}_0 of degree 1. Then for any $j \geq -1$

(9)
$$l(j\mathfrak{A}_0,\mathcal{A}) = 2(j+1) + l,$$

(9')
$$l(j\mathfrak{A}_0, I_n) = n(j+1)$$

by Eq. (6), Eq. (7), Eq. (8), and Eq. (8').

Next, for any integer $j \ge 0$ set

$$b(j) = \sum_{\substack{\mathfrak{C} \ge 0 \\ \deg(\mathfrak{C}) = j}} \mu(\mathfrak{C}),$$

where here μ denotes the usual Möbius function on effective divisors. By [Thu08, Lemma 12]

(10)
$$b(j) = \begin{cases} 1 & \text{if } j = 0, \\ -(q+1) & \text{if } j = 1, \\ q & \text{if } j = 2, \\ 0 & \text{if } j > 2. \end{cases}$$

With the above notation in mind, [Thu08, (5)] reads

$$(q-1)N_{F^3/W}(k) = \sum_{j=0}^{k-\mu_1(F^3/W)} b(j)(q^{l((k-j)\mathfrak{A}_0,\mathcal{A})}-1).$$

Note by Eq. (6) that $k - \mu_1(F^3/W) \ge 2$ if $l, k \ge 1$ or if l = 0 and $k \ge 2$. Thus the former equation above, in conjunction with Eq. (8), Eq. (9), and Eq. (10), yields

$$\begin{aligned} (q-1)N_{F^3/W}(k) &= \left(q^{l(k\mathfrak{A}_0,\mathcal{A})} - 1\right) - (q+1)\left(q^{l((k-1)\mathfrak{A}_0,\mathcal{A})} - 1\right) + q\left(q^{l((k-2)\mathfrak{A}_0,\mathcal{A})} - 1\right) \\ &= \left(q^{2(k+1)+l} - 1\right) - (q+1)\left(q^{2k+l} - 1\right) + q\left(q^{2(k-1)+l} - 1\right) \\ &= q^{2k+l}\left(q^2 - (q+1) + q^{-1}\right) \\ &= q^{2k+l-1}(q-1)(q^2 - 1) \end{aligned}$$

provided $l, k \ge 1$ or l = 0 and $k \ge 2$. In the case l = 0 we have $\mu_1(F^3/W) = 0$ as noted above, so that when k = 1 we get

$$(q-1)N_{F^3/W}(1) = (q^{l(\mathfrak{A}_0,\mathcal{A})} - 1) - (q+1)(q^{l(0,\mathcal{A})} - 1)$$

= $(q^4 - 1) - (q+1)(q^2 - 1)$
= $q^4 - q^3 - q^2 + q$
= $(q-1)(q^3 - q).$

Similarly, since $\mu_1(F^n) = 0$, we see that [Thu08, (5)] reads

$$(q-1)N_{F^n}(k) = \sum_{j=0}^k b(j) (q^{l((k-j)\mathfrak{A}_0,I_n)} - 1).$$

This together with Eq. (8'), Eq. (9'), and Eq. (10) yield

$$(q-1)N_{F^n}(k) = (q^{n(k+1)} - 1) - (q+1)(q^{nk} - 1) + q(q^{n(k-1)} - 1)$$

= $q^{nk+n} - q^{nk+1} - q^{nk} + q^{nk-n+1}$
= $q^{nk}(q^n - 1) - q^{nk-n+1}(q^n - 1)$
= $q^{nk-n+1}(q^n - 1)(q^{n-1} - 1)$

if $k \ge 2$,

$$(q-1)N_{F^n}(1) = (q^{n^2}-1) - (q+1)(q^n-1)$$

= $q^{n^2} - q^{n+1} - q^n + q$
= $q^n(q^n-1) - q(q^n-1)$
= $q(q^n-1)(q^{n-1}-1),$

and finally

$$(q-1)N_{F^n}(0) = q^n - 1.$$

In what follows it will prove convenient to fix some notation for the various sets we will count.

Definition 3.1.2. For any positive integers m and l let S(m) denote the set of all ordered triples (A_1, A_2, A_3) of relatively prime polynomials in $\mathbb{F}_q[X]$ with maximal degree m and S(m, l) denote the subset of S(m) consisting of those triples such that there is a solution $\mathbf{x} = (x_1, x_2, x_3) \in F^3$ of height l to the homogeneous linear equation

(11)
$$x_1A_1 + x_2A_2 + x_3A_3 = 0.$$

As discussed previously, the set S(m, l) corresponds to the two-dimensional subspaces $V \,\subset F^3$ with h(V) = m that contain a one-dimensional subspace $W \subset V$ with h(W) = l. In fact each such subspace V corresponds to precisely q - 1 triples in S(m, l); this takes into account multiplication by elements of $(\mathbb{F}_q)^{\times}$. The required solution(s) to Eq. (11) of height l are also ordered triples of relatively prime polynomials (P_1, P_2, P_3) after multiplication by a suitable non-zero element of F. We note by Minkowski's Theorem Eq. (4) that this solution (P_1, P_2, P_3) is projectively unique if m > 2l.

Definition 3.1.3. For any positive integers m and l let T(m, l) denote the subset of S(m, l) where there is an ordered triple of relatively prime polynomials (P_1, P_2, P_3) that is a solution to Eq. (11) of height l with the additional stipulation that $\deg(P_2) = l$. Let T'(m, l) denote the subset of S(m, l) where there is a solution to Eq. (11) of height l with $\deg(P_3) = l > \max{\deg(P_1), \deg(P_2)}$.

We will also need to further delineate these ordered triples above according to which entry has the largest degree.

Definition 3.1.4. For any non-empty subset $R \subseteq \{1, 2, 3\}$ let $S_R(m)$, $S_R(m, l)$, $T_R(m, l)$, and T'(m, l) denote the subset of S(m), S(m, l), T(m, l), and T'(m, l), respectively, consisting of those ordered triples (A_1, A_2, A_3) where the degree of A_i is m if and only if $i \in R$.

As a notational convenience we will drop the set brackets in the subscript here, e.g., we will write $S_{1,3}(m,l)$ instead of the more cumbersome $S_{\{1,3\}}(m,l)$.

We have determined the connection between curves with given a-numbers and these sets above in Theorem 2.2.3. Our goal for the remainder of this subsection is to determine the cardinalities of these sets above.

Lemma 3.1.5. The sets S(m), S(m,l), T(m,l), and T'(m,l) are disjoint unions:

$$\begin{split} S(m) &= S_1(m) \cup S_2(m) \cup \dots \cup S_{1,2,3}(m), \\ S(m,l) &= S_1(m,l) \cup S_2(m,l) \cup \dots \cup S_{1,2,3}(m,l) \\ T(m,l) &= T_1(m,l) \cup T_3(m,l) \cup T_{1,3}(m,l) \cup T_{1,2}(m,l) \cup T_{2,3}(m,l) \cup T_{1,2,3}(m,l), \text{ and} \\ T'(m,l) &= T'_1(m,l) \cup T'_2(m,l) \cup T'_{1,2}(m,l). \end{split}$$

Further,

$$|S_{1}(m)| = |S_{2}(m)| = |S_{3}(m)|, \qquad |S_{1,2}(m)| = |S_{1,3}(m)| = |S_{2,3}(m)|,$$

$$|T_{1}(m,l)| = |T_{3}(m,l)|, \qquad |T_{1,2}(m,l)| = |T_{2,3}(m,l)|,$$

$$|T'_{1}(m,l)| = |T'_{2}(m,l)|,$$

and if m > 2l

$$T'_{1,2}(m,l) = S_{1,2}(m,l) \setminus T_{1,2}(m,l).$$

Proof. That S(m) and S(m,l) are the given disjoint unions is obvious from the definitions. The same is true for T(m,l) except here $T_2(m,l)$ is empty. To see why, we note that if $(A_1, A_2, A_3) \in T(m, l)$ with $\deg(A_2) > \max\{\deg(A_1), \deg(A_3\})$ and (P_1, P_2, P_3) is any triple of polynomials with $\deg(P_2) = \max\{\deg(P_1), \deg(P_2), \deg(P_3)\}$, then the degree of $A_1P_1 + A_2P_2 + A_3P_3$ is $\deg(A_2) + \deg(P_2) \ge 0$, so that (P_1, P_2, P_3) is not a solution to Eq. (11). A similar argument shows that $T'_R(m, l)$ is empty whenever $3 \in R$.

Next, any element σ of the symmetric group on $\{1, 2, 3\}$ gives rise to a permutation of S(m)and S(m, l) via $(A_1, A_2, A_3) \mapsto (A_{\sigma(1)}, A_{\sigma(2)}, A_{\sigma(3)})$. One readily checks that this mapping takes $S_i(m)$ to $S_{\sigma(i)}(m)$ and $S_{i,j}(m)$ to $S_{\sigma(i),\sigma(j)}(m)$ for any $i, j \in \{1, 2, 3\}$ in a one-to-one and onto manner. Further, the permutation that permutes 1 with 3 and leaves 2 fixed takes $T_1(m,l)$ to $T_3(m,l)$ and $T_{1,2}(m,l)$ to $T_{2,3}(m,l)$. The permutation that permutes 1 with 2 and leaves 3 fixed takes $T'_1(m,l)$ to $T'_2(m,l)$.

Finally, suppose $(A_1, A_2, A_3) \in S_{1,2}(m, l) \setminus T_{1,2}(m, l)$ and m > 2l. Then by definition $\deg(A_1) = \deg(A_2) = m > \deg(A_3)$ and as noted above the projectively unique solution (P_1, P_2, P_3) to Eq. (11) satisfies $\deg(P_2) < l = \max\{\deg(P_1), \deg(P_3)\}$. This can only be the case if $\deg(P_3) = l > \deg(P_1)$ since otherwise we would have

$$\deg(P_1A_1) = l + m > \max\{\deg(P_2A_2), \deg(P_3A_3)\},\$$

contradicting the assumption that (P_1, P_2, P_3) is a solution to Eq. (11). This completes the proof.

Consider the functions $\phi, \gamma: \mathbb{F}_q[X]^3 \times (\mathbb{F}_q)^{\times} \to \mathbb{F}_q[X]^3$ given by

$$\phi((Q_1, Q_2, Q_3), c) = (Q_1, Q_2, Q_3 + cQ_1)$$

and

$$\gamma((Q_1, Q_2, Q_3), c) = (Q_1 - cQ_3, Q_2, Q_3).$$

Lemma 3.1.6. The function ϕ is one-to-one when restricted to those triples where $Q_1 \neq 0$. Both ϕ and γ take relatively prime triples to relatively prime triples, and neither ϕ nor γ alter the height of (Q_1, Q_2, Q_3) , as long as

 $\deg\left(Q_{1}\right) \geq \max\{\deg\left(Q_{2}\right), \deg\left(Q_{3}\right)\}$

or

$$\deg\left(Q_{1}\right) \leq \max\{\deg\left(Q_{2}\right), \deg\left(Q_{3}\right)\},\$$

respectively.

Proof. Indeed,

$$\phi(\phi((Q_1, Q_2, Q_3), c), -c) = (Q_1, Q_2, Q_3)$$

for all $Q_1, Q_2, Q_3 \in \mathbb{F}_q[X]$, and

$$\phi((Q_1, Q_2, Q_3), c) = \phi((Q_1, Q_2, Q_3), b)$$

with $Q_1 \neq 0$ implies that c = b. An entirely similar argument works for γ . For the second part, the common divisors of two polynomials A and B are the common divisors of A and B + CA for all polynomials C. The statement regarding heights now follows since neither ϕ nor γ alter the maximum degree under the two respective conditions on the degree of Q_1 . \Box

Lemma 3.1.7. We have

$$|S_{1,3}(m)| = (q-1)|S_1(m)|, \qquad |S_{1,2,3}(m)| = (q-1)|S_{1,2}(m)|,$$

$$|T_{1,3}(m,l)| = (q-1)|T_1(m,l)|, \qquad |T_{1,2,3}(m,l)| = (q-1)|T_{1,2}(m,l)|, \qquad and$$

$$|T'_{1,2}(m,l)| = (q-1)|T'_1(m,l)|.$$

Proof. If we fix a $c \in (\mathbb{F}_q)^{\times}$ and let ϕ act on a coefficient triple (A_1, A_2, A_3) defining a twodimensional subspace of F^3 via Eq. (11), then the resulting two-dimensional subspace is given by the image of γ (with the same c) acting on the original solution space. Further, if

 $\deg(A_1) \ge \max\{\deg(A_2), \deg(A_3)\},\$

then any polynomial solution (P_1, P_2, P_3) to Eq. (11) satisfies

 $\deg(P_1) \le \max\{\deg(P_2), \deg(P_3)\}.$

We now see by Lemma 3.1.6 that, when restricted to $S_1(m) \times (\mathbb{F}_q)^{\times}$, $S_{1,2}(m) \times (\mathbb{F}_q)^{\times}$, $T_1(m,l) \times (\mathbb{F}_q)^{\times}$, and $T_{1,2}(m,l) \times (\mathbb{F}_q)^{\times}$, the function ϕ has image $S_{1,3}(m)$, $S_{1,2,3}(m)$, $T_{1,3}(m,l)$, and $T_{1,2,3}(m,l)$, respectively. This together with Lemma 3.1.5 completes the proof of the first four equations.

An entirely similar argument with

$$\phi'((Q_1, Q_2, Q_3), c) = (Q_1, Q_2 + cQ_1, Q_3)$$

and

$$\gamma'((Q_1, Q_2, Q_3), c) = (Q_1 - cQ_2, Q_2, Q_3)$$

proves the last equation and completes the proof.

Proposition 3.1.8. For any integer $m \ge 0$ and all distinct $i, j \in \{1, 2, 3\}$

$$|S(m)| = (q-1)N_{F^3}(m) = \begin{cases} q^{3m-2}(q^3-1)(q^2-1) & \text{if } m > 0, \\ (q^3-1) & \text{if } m = 0, \end{cases}$$
$$|S_i(m)| = \frac{|S(m)|}{1+q+q^2}, \qquad |S_{i,j}(m)| = \frac{(q-1)|S(m)|}{1+q+q^2},$$

and

$$|S_{1,2,3}(m)| = \frac{(q-1)^2 |S(m)|}{1+q+q^2}$$

Fix an integer $l \ge 0$. If m > 2l then

$$|S(m,l)| = q^{2m-l-1}(q-1)(q^2-1)N_{F^3}(l) = \begin{cases} q^{2(m+l)-3}(q^3-1)(q^2-1)^2 & \text{if } l > 0, \\ q^{2m-1}(q^3-1)(q^2-1) & \text{if } l = 0, \end{cases}$$
$$|S_i(m,l)| = \frac{|S(m,l)|}{1+q+q^2}, \qquad |S_{i,j}(m,l)| = \frac{(q-1)|S(m,l)|}{1+q+q^2},$$
$$|S_{i,j}(m,l)| = \frac{(q-1)^2|S(m,l)|}{1+q+q^2},$$

and

$$|S_{1,2,3}(m,l)| = \frac{(q-1)^2 |S(m,l)|}{1+q+q^2}$$

for all distinct $i, j \in \{1, 2, 3\}$. Finally, if m = 2l > 0 then

$$|S_R(m,l)| = \frac{|S_R(m)|(q-1)|}{q}$$

For all subsets $R \subseteq \{1, 2, 3\}$.

Proof. As noted above, there is a (q-1)-to-1 correspondence between the elements of S(m, l) and two-dimensional subspaces $V \subset F^3$ of height m containing a one-dimensional subspace W of height l. Since m > 2l, Minkowski's Theorem Eq. (4) implies that this subspace W is unique to V. Therefore by Proposition 3.1.1 and Eq. (5),

$$|S(m,l)| = (q-1) \sum_{\substack{W \subset F^3 \\ \dim(W)=1 \\ h(W)=l}} N_{F^3/W}(m-l)$$
$$= (q-1)q^{2(m-l)+l-1}(q^2-1)N_{F^3}(l).$$

(Note that Proposition 3.1.1 applies since $m - l \ge 1$ if $l \ge 1$ and $m - l = m \ge 1$ if l = 0.) Next, by Lemma 3.1.5 and Lemma 3.1.7,

$$\begin{split} |S(m,l)| &= |S_1(m,l)| + |S_2(m,l)| + |S_3(m,l)| \\ &+ |S_{1,2}(m,l)| + |S_{1,3}(m,l)| + |S_{2,3}(m,l)| + |S_{1,2,3}(m,l)| \\ &= 3|S_1(m,l)| + 3|S_{1,2}(m,l)| + |S_{1,2,3}(m,l)| \\ &= 3|S_1(m,l)| + (3+q-1)|S_{1,2}(m,l)| \\ &= 3|S_1(m,l)| + (3+q-1)|S_{1,3}(m,l)| \\ &= (3+3(q-1)+(q-1)^2)|S_1(m,l)| \\ &= (1+q+q^2)|S_1(m,l)|. \end{split}$$

This together with Eq. (5), Lemma 3.1.5, and Lemma 3.1.7 once more finishes the proof of the second part of the proposition.

For the first part of the proposition, recall that every one-dimensional subspace of F^3 of height m corresponds to q-1 elements of S(m). The rest of the proof follows exactly as above.

Next, suppose m = 2l > 0. Then by definition and Eq. (4) we have

$$|S(m)| = \sum_{k=0}^{l} |S(m,k)|.$$

By what we have already proven above

$$|S(m,0)| = q^{2(m+0)-3}q^2(q^3-1)(q^2-1) = q^{2(m+0)-3}(q^3-1)(q^2-1)^2 + q^{2m-3}(q^3-1)(q^2-1).$$

Using this together what we have already proven yields

$$|S(m)| = \sum_{k=0}^{l-1} q^{2(m+k)-3} (q^3 - 1)(q^2 - 1)^2 + q^{2m-3}(q^3 - 1)(q^2 - 1) + |S(m,l)|$$

= $q^{2m-3}(q^3 - 1)(q^2 - 1)^2 \frac{q^{2l} - 1}{q^2 - 1} + q^{2m-3}(q^3 - 1)(q^2 - 1) + |S(m,l)|$
= $q^{2m-3}(q^3 - 1)(q^2 - 1)q^{2l} + |S(m,l)|$
= $q^{3m-3}(q^3 - 1)(q^2 - 1) + |S(m,l)|$
= $|S(m)|q^{-1} + |S(m,l)|.$

This shows the case where $R = \emptyset$ of the last part of the proposition. The other cases are entirely similar.

Proposition 3.1.9. Fix an integer $l \ge 0$. If m > 2l then

$$|T'(m,l)| = \frac{|S(m,l)|}{1+q+q^2}, \qquad |T'_1(m,l)| = |T'_2(m,l)| = \frac{|S_1(m,l)|}{q+1}, \qquad and$$
$$|T'_{1,2}(m,l)| = \frac{|S_{1,2}(m,l)|}{q+1}.$$

Proof. We have noted above that any one-dimensional subspace $W \subset F^3$ has a basis element consisting of a relatively prime triple of polynomials (P_1, P_2, P_3) and the height of W is the maximum of the degrees of these polynomials. This triple is unique up to multiplication by

an element of $(\mathbb{F}_q)^{\times}$, thus the indices *i* where deg $(P_i) = h(W)$ are uniquely determined by W.

We may now evaluate |T'(m, l)| in a manner very similar to |S(m, l)| above. Specifically, we have

$$|T'(m,l)| = (q-1) \cdot \sum_{\substack{W \subset F^3 \\ \dim(W)=1 \\ h(W)=l}} N_{F^3/W}(m-l),$$

where the prime indicates that we are to sum only over those W spanned by a relatively prime ordered triple counted in $S_3(l)$. Note that there are q-1 ordered triples in this set for each such W. Via Eq. (5), Proposition 3.1.1, and Proposition 3.1.8

$$|T'(m,l)| = (q-1) \cdot \sum_{\substack{W \in F^3 \\ \dim(W)=l}}^{Y'(m,l)|} N_{F^3/W}(m-l)$$

$$= (q-1)q^{2(m-l)+l-1}(q^2-1)\frac{|S_3(l)|}{q-1}$$

$$= q^{2m-l-1}(q^2-1)\frac{S(l)}{1+q+q^2}$$

$$= q^{2m-l-1}(q^2-1)q^2\frac{(q-1)N_{F^3}(l)}{1+q+q^2}$$

$$= \frac{|S(m,l)|}{1+q+q^2}.$$

Next, by Lemma 3.1.5 and Lemma 3.1.7,

$$|T'(m,l)| = |T'_1(m,l)| + |T'_2(m,l)| + |T'_{1,2}(m,l)|$$

= (q + 1)|T'_1(m,l)|.

The remainder of the Proposition follows from this equation, Eq. (12), Lemma 3.1.5, and Lemma 3.1.7. $\hfill \Box$

Proposition 3.1.10. Fix an integer $l \ge 0$. If m > 2l then

$$|T(m,l)| = \frac{q^2 |S(m,l)|}{1+q+q^2}, \qquad |T_1(m,l)| = |T_3(m,l)| = \frac{q |S_1(m,l)|}{(q+1)},$$

$$T_{1,2}(m,l)| = |T_{2,3}(m,l)| = \frac{q |S_{1,2}(m,l)|}{q+1}, \qquad and \qquad |T_{1,2,3}(m,l)| = \frac{q |S_{1,2,3}(m,l)|}{q+1}.$$

Proof. We argue almost exactly as above in the proof of Proposition 3.1.9, only this time

$$|T(m,l)| = (q-1) \cdot \sum_{\substack{W \subset F^3 \\ \dim(W)=1 \\ h(W)=l}} N_{F^3/W}(m-l),$$

where the prime indicates that we are to sum only over those W spanned by a relatively prime ordered triple counted in the union $S_2(l) \cup S_{1,2}(l) \cup S_{2,3}(l) \cup S_{1,2,3}(l)$. Via Eq. (5), Proposition 3.1.1 and Proposition 3.1.8

(12)

$$|T(m,l)| = (q-1) \cdot \sum_{\substack{W \in F^{3} \\ \dim(W)=1 \\ h(W)=l}} N_{F^{3}/W}(m-l)}$$

$$= (q-1)q^{2(m-l)+l-1}(q^{2}-1)\frac{\left(|S_{2}(l)| + |S_{1,2}(l)| + |S_{1,2,3}(l)| + |S_{1,2,3}(l)|\right)}{q-1}$$

$$= q^{2m-l)-1}(q^{2}-1)\frac{|S(l)|}{1+q+q^{2}}(1+2(q-1)+(q-1)^{2})$$

$$= q^{2m-l-1}(q^{2}-1)q^{2}\frac{(q-1)N_{F^{3}}(l)}{1+q+q^{2}}$$

$$(13) \qquad = \frac{q^{2}}{1+q+q^{2}}|S(m,l)|.$$

Next, by Lemma 3.1.5 and Proposition 3.1.9

$$\frac{(q-1)|S(m,l)|}{(q+1)(1+q+q^2)} = |T'_{1,2}(m,l)|$$
$$= |S_{1,2}(m,l)| - |T_{1,2}(m,l)|$$
$$= \frac{(q-1)|S(m,l)|}{1+q+q^2} - |T_{1,2}(m,l)|,$$

so that

$$|T_{1,2}(m,l)| = \frac{q(q-1)|S(m,l)|}{(q+1)(1+q+q^2)}.$$

Using this together with Lemma 3.1.5 and Lemma 3.1.7 yields

$$\begin{aligned} |T(m,l)| &= |T_1(m,l)| + |T_3(m,l)| + |T_{1,3}(m,l)| + |T_{1,2}(m,l)| + |T_{2,3}(m,l)| + |T_{1,2,3}(m,l)| \\ &= (q+1)|T_1(m,l)| + (q+1)|T_{1,2}(m,l)| \\ &= (q+1)|T_1(m,l)| + \frac{q(q-1)|S(m,l)|}{1+q+q^2}. \end{aligned}$$

The remainder of the Proposition follows from this equation, Eq. (13), Lemma 3.1.5, Lemma 3.1.7, and Proposition 3.1.8.

3.2. Main counting results.

Theorem 3.2.1. Fix integers $m, l \ge 0$. Let $N_1(m, l)$ denote the number of ordered triples (A_1, A_2, A_3) of polynomials such that:

- A_1, A_2, A_3 are relatively prime;
- A_1 is monic of degree m;
- the degrees of both A_2 and A_3 are no greater than m; and
- there is a solution to Eq. (11) of height l.

If m > 2l then

$$N_1(m,l) = \begin{cases} (q^2-1)^2 q^{2(m+l)-1} & if \ l > 0, \\ (q^2-1) q^{2m+1} & if \ l = 0. \end{cases}$$

If m = 2l > 0 then

$$N_1(m,l) = (q-1)(q^2-1)q^{3m-1}.$$

Further, suppose $m > 2l \ge 0$ and let $N'_1(m, l)$ denote the number of those ordered triples as above but where the solution (P_1, P_2, P_3) of height l satisfies $\deg(P_2) < l$. Then

$$N_1'(m,l) = \frac{1}{q+1} N_1(m,l) = \begin{cases} (q^2-1)(q-1)q^{2(m+l)-1} & \text{if } l > 0, \\ (q-1)q^{2m+1} & \text{if } l = 0. \end{cases}$$

Proof. Since we demand that A_1 is monic, in the notation of the previous subsection we have

$$N_1(m,l) = \frac{1}{q-1} (|S_1(m,l)| + |S_{1,2}(m,l)| + |S_{1,3}(m,l)| + |S_{1,2,3}(m,l)|).$$

We now apply Proposition 3.1.8. If m > 2l we have

$$N_{1}(m,l) = \frac{1}{q-1} \left(|S_{1}(m,l)| + |S_{1,2}(m,l)| + |S_{1,3}(m,l)| + |S_{1,2,3}(m,l)| \right)$$

$$= \frac{1+2(q-1)+(q-1)^{2}}{(q-1)(1+q+q^{2})} |S(m,l)|$$

$$= \frac{q^{2}}{q^{3}-1} |S(m,l)|$$

$$= \begin{cases} (q^{2}-1)^{2}q^{2(m+l)-1} & \text{if } l > 0, \\ (q^{2}-1)q^{2m+1} & \text{if } l = 0. \end{cases}$$

The case m = 2l is entirely similar.

The second part we have

$$N_{1}'(m,l) = \frac{1}{q-1} \left(|S_{1}(m,l)| - |T_{1}(m,l)| + |S_{1,2}(m,l)| - |T_{1,2}(m,l)| + |S_{1,3}(m,l)| - |T_{1,3}(m,l)| + |S_{1,2,3}(m,l)| - |T_{1,2,3}(m,l)| \right)$$

= $N_{1}(m,l) \left(1 - \frac{q}{q+1} \right)$
= $\frac{1}{q+1} N_{1}(m,l)$

by Proposition 3.1.10 and our expression for $N_1(m, l)$ above.

Theorem 3.2.2. Fix integers $m, l \ge 0$. Let $N_2(m, l)$ denote the number of ordered triples (A_1, A_2, A_3) of polynomials such that:

- A_1, A_2, A_3 are relatively prime;
- A_1 is monic of degree m;
- the degrees of both A_2 and A_3 are strictly less than m; and
- there is a solution to Eq. (11) of height l.

If m > 2l then

$$N_2(m,l) = \begin{cases} (q^2-1)^2 q^{2(m+l)-3} & \text{if } l > 0, \\ (q^2-1)q^{2m-1} & \text{if } l = 0. \end{cases}$$

If m = 2l > 0 then

$$N_2(m,l) = (q-1)(q^2-1)q^{3m-3}.$$

Further, suppose $m > 2l \ge 0$ and let $N'_2(m, l)$ denote the number of those triples as above but where the solution (P_1, P_2, P_3) satisfies $\deg(P_2) < l$. Then

$$N_{2}'(m,l) = \frac{1}{q+1} N_{2}(m,l) = \begin{cases} (q-1)(q^{2}-1)q^{2(m+l)-3} & \text{if } l > 0, \\ (q-1)q^{2m-1} & \text{if } l = 0. \end{cases}$$

Proof. Since we demand that A_1 is monic, in the notation of the previous section we have

$$N_{2}(m,l) = \frac{1}{q-1} |S_{1}(m,l)|$$

= $\frac{1}{(q-1)(1+q+q^{2})} |S(m,l)|$
= $\begin{cases} \frac{(q^{3}-1)(q^{2}-1)(q+1)}{1+q+q^{2}} q^{2(m+l)-3} & \text{if } l > 0, \\ \frac{(q+1)(q^{3}-1)}{1+q+q^{2}} q^{2m-1} & \text{if } l = 0, \end{cases}$
= $\begin{cases} (q^{2}-1)^{2} q^{2(m+l)-3} & \text{if } l > 0, \\ (q^{2}-1)q^{2m-1} & \text{if } l = 0, \end{cases}$

by Proposition 3.1.8

For the second part, note that since the degree of A_1 is strictly larger than the degrees of both A_2 and A_3 , any solution (P_1, P_2, P_3) of height l to Eq. (11) with deg $(P_2) < l$ necessarily must have deg $(P_3) = l$ strictly larger than the degrees of both P_1 and P_2 . Thus

$$N_{2}'(m,l) = \frac{1}{q-1} |T_{1}'(m,l)|$$
$$= \frac{1}{q^{2}-1} |S_{1}(m,l)|$$

by Proposition 3.1.9.

Theorem 3.2.3. Fix integers $m, l \ge 0$. Let $N_3(m, l)$ denote the number of ordered triples (A_1, A_2, A_3) of polynomials such that:

- A_1, A_2, A_3 are relatively prime;
- A_1 is monic of degree m;
- $\deg(A_2) \leq m;$
- $\deg(A_3) < m$; and
- there is a solution to Eq. (11) of height l.

If m > 2l then

$$N_3(m,l) = \begin{cases} (q^2 - 1)^2 q^{2(m+l)-2} & \text{if } l > 0, \\ (q^2 - 1) q^{2m} & \text{if } l = 0. \end{cases}$$

If m = 2l > 0 then

$$N_3(m,l) = (q-1)(q^2-1)q^{3m-2}.$$

Further, suppose $m > 2l \ge 0$ and let $N'_3(m, l)$ denote the number of those triples above but where the solution (P_1, P_2, P_3) satisfies deg $(P_2) < l$. Then

$$N'_{3}(m,l) = \frac{1}{q+1}N_{3}(m,l) = \begin{cases} (q^{2}-1)(q-1)q^{2(m+l)-1} & \text{if } l > 0, \\ (q-1)q^{2m} & \text{if } l = 0. \end{cases}$$

Proof. Since we demand that A_1 is monic, in the notation of the previous section we have

$$N_{3}(m,l) = \frac{1}{q-1} (|S_{1}(m,l)| + |S_{1,2}(m,l)|)$$
$$= \frac{q}{(q-1)(1+q+q^{2})} |S(m,l)|$$
$$= \begin{cases} (q^{2}-1)^{2}q^{2(m+l)-1} & \text{if } l > 0, \\ (q^{2}-1)q^{2m} & \text{if } l = 0 \end{cases}$$

by Proposition 3.1.8. For the second part, note that since the degree of A_1 is strictly larger than the degree of A_3 and at least as large as the degree of A_2 , any solution (P_1, P_2, P_3) of height l to Eq. (11) with deg $(P_2) < l$ necessarily must have deg $(P_3) = l$ strictly larger than the degrees of both P_1 and P_2 . Thus

$$N'_{3}(m,l) = \frac{1}{q-1} (|T'_{1}(m,l)| + |T'_{1,2}(m,l)|)$$
$$= \frac{1}{q^{2}-1} (|S_{1}(m,l)| + |S_{1,2}(m,l)|)$$

by Proposition 3.1.9.

Proposition 3.2.4. For all g > 0 and $\epsilon \in \{1, 2\}$ we have

$$|\mathcal{P}_{\epsilon}(g)| = q^{2g+\epsilon} (1-q^{-2}).$$

Also,

$$|\mathcal{P}_{\epsilon}(0)| = q^{\epsilon}(1 - q^{-1})$$

for $\epsilon \in \{1, 2\}$. Finally,

$$|\mathcal{P}'_{\epsilon}(g)| = \begin{cases} q^{2g+\epsilon}(1-q^{-1}) & \text{if } g > 0 \text{ or } g = 0 \text{ and } \epsilon = 2, \\ q & \text{if } g = 0 \text{ and } \epsilon = 1. \end{cases}$$

Proof. We will prove the cases for $\mathcal{P}_{\epsilon}(g)$; those for $\mathcal{P}'_{\epsilon}(g)$ are well-known (see Proposition 2.3 of [R02], for example). The case where g = 0 is trivial, so we will assume that g > 0.

Suppose that $g \equiv 0 \pmod{3}$. If $\epsilon = 1$, then by Lemma 2.2.2 and definition $f \in \mathbb{F}_q[X]$ is cube-free of degree 2g+1 if and only if $(c_0(f), c_1(f), c_2(f)) \in S_2(2g/3) \cup S_{1,2}(2g/3)$. The result in this case follows from Proposition 3.1.8 and Eq. (5). If $\epsilon = 2$, then by Lemma 2.2.2 and definition $f \in \mathbb{F}_q[X]$ is cube-free of degree 2g+2 if and only $(c_0(f), c_1(f), c_2(f)) \in S_3(2g/3) \cup S_{1,3}(2g/3) \cup S_{2,3}(2g/3) \cup S_{1,2,3}(2g/3)$, and the result again follows from Proposition 3.1.8 and Eq. (5). The other cases $(g \equiv 1 \pmod{3})$ and $g \equiv 2 \pmod{3}$ are proven in an entirely similar manner.

4. The *a*-numbers of hyperelliptic curves in characteristic three

In this section we restrict to the case p = 3 to deduce the results mentioned in Section 1.

4.1. The proportion of curves with a given *a*-number. We may now prove Theorem 1.1.1 via the counting results in the previous section and Theorem 2.2.3.

Proof of Theorem 1.1.1. Suppose first that $g \equiv 0 \pmod{3}$.

For $f \in \mathcal{P}_{\epsilon}(g)$, if $\epsilon = 1$ then the space V in Theorem 2.2.3 has height

$$h(V) = \frac{2g}{3} = \operatorname{deg}(c_1(f)) > \operatorname{deg}(c_2(f)).$$

We use Theorem 3.2.3 with

 $A_1 = c_1(f), \qquad A_3 = c_2(f), \qquad A_2 = c_0(f), \qquad P_1 = c_2(Q), \qquad P_2 = c_1(Q), \qquad \text{and } P_3 = c_0(Q).$ We set $m = \frac{2g}{3}$. Then for any $0 \le l \le \frac{g}{3}$ the number of $f \in \mathcal{P}_{\epsilon}(g)$ with $a(H_f) = \frac{g}{3} - l = a$ is $N_3(m, l)$ by Theorem 2.2.3. We thus have

$$\mu_{1,g}(a) = \frac{N_3(m, g/3 - a)}{|\mathcal{P}_{\epsilon}(g)|}$$

$$= \begin{cases} \frac{q^{3m-2-2a}(q^2-1)(q-1)}{q^{2g-1}(q^2-1)} & \text{if } a = 0, \\ \frac{q^{2(m+g/3-a)-2}(q^2-1)^2}{q^{2g-1}(q^2-1)} & \text{if } 0 < a < g/3, \\ \frac{q^{2(m+g/3-a)}(q^2-1)}{q^{2g-1}(q^2-1)} & \text{if } a = g/3, \end{cases}$$

$$= \begin{cases} q^{-2a}\frac{q-1}{q} & \text{if } a = 0, \\ q^{-2a}\frac{q^2-1}{q} & \text{if } 0 < a < g/3, \\ q^{-2a}q & \text{if } a = g/3. \end{cases}$$

For $f \in \mathcal{P}_{\epsilon}(g)$, if $\epsilon = 2$ then the space V in Theorem 2.2.3 has height

$$h(V) = \frac{2g}{3} = \deg(c_2(f)).$$

We use Theorem 3.2.1 with

 $A_1 = c_2(f), \qquad A_3 = c_1(f), \qquad A_2 = c_0(f), \qquad P_1 = c_0(Q), \qquad P_2 = c_2(Q), \qquad \text{and } P_3 = c_1(Q).$ We set $m = \frac{2g}{3}$. Then for any $0 \le l \le \frac{g}{3}$ the number of $f \in \mathcal{P}_{\epsilon}(g)$ with $a(H_f) = \frac{g}{3} - l = a$ is $N_1(m, l)$ by Theorem 2.2.3. We thus have

$$\mu_{2,g}(a) = \frac{N_1(m, g/3 - a)}{|\mathcal{P}_{\epsilon}(g)|}$$

$$= \begin{cases} \frac{q^{3m-1}(q^2-1)(q-1)}{q^{2g}(q^2-1)} & \text{if } a = 0, \\ \frac{q^{2(m+g/3-a)-1}(q^2-1)^2}{q^{2g}(q^2-1)} & \text{if } 0 < a < g/3, \\ \frac{q^{2(m+g/3-a)+1}(q^2-1)}{q^{2g}(q^2-1)} & \text{if } a = g/3, \end{cases}$$

$$= \begin{cases} q^{-2a}\frac{q-1}{q} & \text{if } a = 0, \\ q^{-2a}\frac{q^2-1}{q} & \text{if } 0 < a < g/3, \\ q^{-2a}q & \text{if } a = g/3. \end{cases}$$

Next suppose that $q \equiv 1 \pmod{3}$.

For $f \in \mathcal{P}_{\epsilon}(g)$, if $\epsilon = 1$ then the space V in Theorem 2.2.3 has height

$$h(V) = \frac{2g+1}{3} = \deg(c_0(f)) > \max\{\deg(c_1(f)), \deg(c_2(f))\}.$$

We use Theorem 3.2.2 with

 $\begin{array}{ll} A_1 = c_0(f), & A_2 = c_1(f), & A_3 = c_2(f), & P_1 = c_2(Q), & P_2 = c_1(Q), & \text{and } P_3 = c_0(Q). \\ \text{We set } m = \frac{2g+1}{3}. \text{ Note that } m = 2(g-1)/3 + 1, \text{ so that by Eq. (4) the first minima } \mu_1(V) \\ \text{is no greater than } (g-1)/3. \text{ By Theorem 2.2.3 for any } 0 < l \leq \frac{g-1}{3} \text{ the number of } f \in \mathcal{P}_{\epsilon}(g) \\ \text{with } a(H_f) = \frac{g-1}{3} - l + 1 = a \text{ is } N'_2(m,l) + \left(N_2(m,l-1) - N'_2(m,l-1)\right), \text{ the number of } f \in \mathcal{P}_{\epsilon}(g) \\ f \in \mathcal{P}_{\epsilon}(g) \text{ with } a(H_f) = \frac{g-1}{3} + 1 \text{ is } N'_2(m,0), \text{ and the number of } f \in \mathcal{P}_{\epsilon}(g) \text{ with } a(H_f) = 0 \text{ is } \\ N_2(m, \frac{g-1}{3}) - N'_2(m, \frac{g-1}{3}). \end{array}$

Excluding for the moment the case where g = 1 and a = 0, we therefore have

$$\mu_{1,g}(a) = \begin{cases} \frac{N_2\left(m,(g-1)/3\right) - N'_2\left(m,(g-1)/3\right)}{|\mathcal{P}_{\epsilon}(g)|} & \text{if } a = 0, \\ \frac{N'_2\left(m,(g-1)/3 - a + 1\right) + N_2\left(m,(g-1)/3 - a\right) - N'_2\left(m,(g-1)/3 - a\right)}{|\mathcal{P}_{\epsilon}(g)|} & \text{if } 1 \le a \le \frac{g-1}{3}, \\ \frac{N'_2(m,0)}{|\mathcal{P}_{\epsilon}(g)|} & \text{if } a = 0, \\ \frac{q^{2(m+(g-1)/3 - a) - q}(q^2 - 1)^2}{q^{2g-1}(q^2 - 1)} & \text{if } a = 0, \\ \frac{q^{2(m+(g-1)/3 - a) - q}(q^2 - 1)^2}{q^{2g-1}(q^2 - 1)} & \text{if } 0 < a < \frac{g-1}{3}, \\ \frac{q^{2(m+(g-1)/3 - a) - q}(q^2 + q - 1)}{(q^2 + 1)(q^2 - 1)} & \text{if } a = \frac{g-1}{3}, \\ \frac{q^{2(m+(g-1)/3 - a) - q}(q^2 + q - 1)}{(q^2 - 1)q^{2g-1}} & \text{if } a = \frac{g+2}{3}, \end{cases}$$

$$= \begin{cases} q^{-2a} \frac{q-1}{q} & \text{if } a = 0, \\ q^{-2a} \frac{q^2}{q} - 1} & \text{if } a = 0, \\ q^{-2a} \frac{q^2}{q} - 1} & \text{if } a = \frac{g-1}{3}, \\ q^{-2a} \frac{q^2}{q+1} & \text{if } a = \frac{g-1}{3}, \\ q^{-2a} \frac{q^2}{q+1} & \text{if } a = \frac{g-1}{3}, \\ q^{-2a} \frac{q^2}{q+1} & \text{if } a = \frac{g+2}{3}. \end{cases}$$

If g = 1 (so that (g-1)/3 = 0 and m = 1) and a = 0 we have

$$\mu_{1,1}(0) = \frac{N_2(1,0) - N_2'(1,0)}{|\mathcal{P}_{\epsilon}(1)|}$$
$$= \frac{(q^2 - 1)q - (q - 1)q}{q(q^2 - 1)}$$
$$= \frac{q}{q + 1}.$$

For $f \in \mathcal{P}_{\epsilon}(g)$, if $\epsilon = 2$ then the space V in Theorem 2.2.3 has height

$$h(V) = \frac{2g+1}{3} = \deg(c_1) > \deg(c_2)$$

We use Theorem 3.2.3 with

 $\begin{aligned} A_1 &= c_1(f), \quad A_2 &= c_0(f), \quad A_3 &= c_2(f), \quad P_1 &= c_1(Q), \quad P_2 &= c_2(Q), \quad \text{and } P_3 &= c_0(Q). \end{aligned}$ We set $m &= \frac{2g+1}{3}$. Once again the first minima $\mu_1(V)$ is no greater than (g-1)/3. By Theorem 2.2.3 for any $0 < l \leq \frac{g-1}{3}$ the number of $f \in \mathcal{P}_{\epsilon}(g)$ with $a(H_f) &= \frac{g-1}{3} - l + 1 = a$ is $N'_3(m,l) + \left(N_3(m,l-1) - N'_3(m,l-1)\right)$. The number of $f \in \mathcal{P}_{\epsilon}(g)$ with $a(H_f) = \frac{g-1}{3} + 1$ is $N'_3(m,0)$ and the number of $f \in \mathcal{P}_{\epsilon}(g)$ with $a(H_f) = 0$ is $N_3(m, \frac{g-1}{3}) - N'_3(m, \frac{g-1}{3})$. Again temporarily excluding the case where g = 1 and a = 0, we therefore have

$$\begin{split} \mu_{2,g}(a) &= \begin{cases} \frac{N_3\left(m,(g-1)/3\right) - N_3'\left(m,(g-1)/3\right)}{|\mathcal{P}_{\epsilon}(g)|} & \text{if } a = 0, \\ \frac{N_3'\left(m,(g-1)/3 - a + 1\right) + N_3\left(m,(g-1)/3 - a\right) - N_3'\left(m,(g-1)/3 - 1\right)}{|\mathcal{P}_{\epsilon}(g)|} & \text{if } 0 < a \leq \frac{g-1}{3}, \\ \frac{N_3'(m,0)}{|\mathcal{P}_{\epsilon}(g)|} & \text{if } a = 0, \\ \frac{q^{2(m+1)}(q^{2-1})^2}{(q^{2+1})^{2/9}(q^{2-1})} & \text{if } a = 0, \\ \frac{q^{2(m+1)}(q^{2-1})^2}{(q^{2+1})^{2/9}(q^{2-1})} & \text{if } 0 < a < \frac{g-1}{3}, \\ \frac{q^{2(m+(g-1)/3 - a + 1) - 3}(q^2 - 1)^2}{(q^{2+1})^{2/9}(q^{2-1})} & \text{if } a = \frac{g-1}{3}, \\ \frac{q^{2(m+(g-1)/3 - a + 1)}(q-1)}{(q^{2})(q^{2-1})} & \text{if } a = \frac{g+2}{3} + 1, \end{cases} \\ &= \begin{cases} q^{-2a} \frac{q^2}{q} & \text{if } a = 0, \\ q^{-2a} \frac{q^2}{q} - 1 & \text{if } a = 0, \\ q^{-2a} \frac{q^2}{q+1} & \text{if } a = \frac{g-1}{3}, \\ q^{-2a} \frac{q^2}{q+1} & \text{if } a = \frac{g-1}{3}, \\ q^{-2a} \frac{q^2}{q+1} & \text{if } a = \frac{g-1}{3}, \\ q^{-2a} \frac{q^2}{q+1} & \text{if } a = \frac{g+2}{3}. \end{cases} \end{split}$$

If g = 1 and a = 0 we have

$$\mu_{2,1}(0) = \frac{N_3(1,0) - N_3'(1,0)}{|\mathcal{P}_{\epsilon}(1)|}$$
$$= \frac{(q^2 - 1)q^2 - (q - 1)q^2}{q^2(q^2 - 1)}$$
$$= \frac{q}{q + 1}.$$

Finally suppose that $g \equiv 2 \pmod{3}$.

For $f \in \mathcal{P}_{\epsilon}(g)$, if $\epsilon = 1$ then the space V in Theorem 2.2.3 has height

$$h(V) = \frac{2g-1}{3} = \deg(c_2).$$

We use Theorem 3.2.1 with

$$A_1 = c_2(f),$$
 $A_2 = c_0(f),$ $A_3 = c_1(f),$ $P_1 = c_0(Q),$ $P_2 = c_2(Q),$ and $P_3 = c_1(Q).$

We set $m = \frac{2g-1}{3}$. Note that m = 2(g-2)/3 + 1, so that by Eq. (4) the first minima $\mu_1(V)$ is no greater than (g-2)/3. By Theorem 2.2.3 for any $0 < l \leq \frac{g-2}{3}$ the number of $f \in \mathcal{P}_{\epsilon}(g)$ with $a(H_f) = \frac{g-2}{3} - l + 1 = a$ is $N'_1(m, l) + (N_1(m, l-1) - N'_1(m, l-1))$. The number of $f \in \mathcal{P}_{\epsilon}(g)$ with $a(H_f) = \frac{g-2}{3} + 1$ is $N'_1(m, 0)$, and the number of $f \in \mathcal{P}_{\epsilon}(g)$ with $a(H_f) = 0$ is $N_1(m, \frac{g-2}{3}) - N'_1(m, \frac{g-2}{3})$.

Temporarily excluding the case where g = 2 and a = 0, we therefore have

$$\begin{split} \mu_{1,g}(a) &= \begin{cases} \frac{N_1\Big(m,(g-2)/3\Big) - N_1'\Big(m,(g-2)/3\Big)}{|\mathcal{P}_{\epsilon}(g)|} & \text{if } a = 0, \\ \frac{N_1'\Big(m,(g-2)/3 - a + 1\Big) + N_1\Big(m,(g-2)/3 - a\Big) - N_1'\Big(m,(g-2)/3 - a\Big)}{|\mathcal{P}_{\epsilon}(g)|} & \text{if } 0 < a \leq \frac{g-2}{3}, \\ \frac{N_1'(m,0)}{|\mathcal{P}_{\epsilon}(g)|} & \text{if } a = 0, \\ \frac{q^{2(m+4/2)/3 - a}(q^2 - 1)^2}{(q+1)q^{2g-1}(q^2 - 1)} & \text{if } a = 0, \\ \frac{q^{2(m+4/2)/3 - a}(q^2 - 1)^2}{(q+1)q^{2g-1}} & \text{if } 0 < a < \frac{g-2}{3}, \\ \frac{q^{2(m+4/2)/3 - a}(q^2 - 1)^2}{(q+1)q^{2g-1}} & \text{if } a = \frac{g-2}{3}, \\ \frac{q^{2(m+4/2)/3 - a}(q^2 - 1)}{(q+1)q^{2g-1}} & \text{if } a = \frac{g-2}{3}, \\ \frac{q^{2(m+4/2)/3 - a+1) + 1}(q^2 - 1)}{q^{2g-1}(q^2 - 1)} & \text{if } a = \frac{g+1}{3}, \end{cases} \\ &= \begin{cases} q^{-2a}\frac{q-1}{2} & \text{if } a = 0, \\ q^{-2a}\frac{q^2}{2} - 1 & \text{if } 0 < a < \frac{g-2}{3}, \\ q^{-2a}\frac{q^2}{2} - 1 & \text{if } a = \frac{g-2}{3}, \\ q^{-2a}\frac{q^2}{2} + 1 & \text{if } a = \frac{g-2}{3}, \\ q^{-2a}\frac{q^2}{2} + 1 & \text{if } a = \frac{g-2}{3}, \\ q^{-2a}\frac{q^2}{2} + 1 & \text{if } a = \frac{g+1}{3}. \end{cases} \end{split}$$

If g = 2 (so that (g-2)/3 = 0 and m = 1) and a = 0 we have

$$\mu_{1,2}(0) = \frac{N_1(1,0) - N_1'(1,0)}{|\mathcal{P}_{\epsilon}(2)|}$$
$$= \frac{(q^2 - 1)q^3 - (q - 1)q^3}{q^3(q^2 - 1)}$$
$$= \frac{q}{q + 1}.$$

For $f \in \mathcal{P}_{\epsilon}(g)$, if $\epsilon = 2$ then the space V in Theorem 2.2.3 has height

$$h(V) = \frac{2g+2}{3} = \deg(c_0) > \max\{\deg(c_1), \deg(c_2)\}$$

We use Theorem 3.2.2 with

 $A_1 = c_0,$ $A_2 = c_1,$ $A_3 = c_2,$ $P_1 = Q_1,$ $P_2 = Q_2,$ and $P_3 = Q_3.$

We set $m = \frac{2g+2}{3}$. Note that m = 2(g+1)/3, so that by Eq. (4) the first minima $\mu_1(V)$ is no greater than (g+1)/3. By Theorem 2.2.3 $a(H_f) > 0$ only if $\mu_1(V) \le \frac{g-2}{3}$. Moreover, for any $0 \le l \le \frac{g-2}{3}$ the number of $f \in \mathcal{P}_{\epsilon}(g)$ with $a(H_f) = \frac{g-2}{3} - l + 1 = a$ is $N_2(m, l)$. The number of $f \in \mathcal{P}_{\epsilon}(g)$ with $a(H_f) = 0$ is $N_2(m, (g+1)/3)$.

Therefore

$$\mu_{2,g}(a) = \frac{N_2(m, (g-2)/3 - a + 1)}{|\mathcal{P}_{\epsilon}(g)|}$$

$$= \begin{cases} \frac{q^{2g-1}(q^2 - 1)(q - 1)}{q^{2g}(q^2 - 1)} & \text{if } a = 0, \\ \frac{q^{2(m+(g-2)/3 - a + 1) - 3}(q^2 - 1)^2}{q^{2g}(q^2 - 1)} & \text{if } 1 \le a < \frac{g + 1}{3}, \\ \frac{q^{2(m+(g-2)/3 - a + 1) - 1}(q^2 - 1)}{q^{2g}(q^2 - 1)} & \text{if } a = \frac{g + 1}{3}, \end{cases}$$

$$= \begin{cases} q^{-2a} \frac{q - 1}{q} & \text{if } a = 0, \\ q^{-2a} \frac{q^2 - 1}{q} & \text{if } 1 \le a < \frac{g + 1}{3}, \\ q^{-2a} q & \text{if } a = \frac{g + 1}{3}. \end{cases}$$

We now turn to proofs of Corollary 1.1.2 and Corollary 1.1.4.

Proof of Corollary 1.1.2. Part (1) is immediate from Theorem 1.1.1. To prove part (2), let $\epsilon \in \{1,2\}$ and $f \in \mathcal{P}_{\epsilon}(q,g)$, so there are monic squarefree polynomials $f_1, f_2 \in \mathbb{F}_q[X]$ such that $f = f_1(f_2)^2$ and $\gcd(f_1, f_2) = 1$.

Suppose for a moment that we are in the case where f is not squarefree, so that deg $f_2 \ge 1$. Let g_1 be the integer such that $2g_1 + \epsilon = \deg f_1$; since $g \equiv 1 \pmod{3}$ and $g - 1 \ge g_1$, we see that

$$\left\lceil \frac{g}{3} \right\rceil > \frac{g-1}{3} = \left\lceil \frac{g-1}{3} \right\rceil \ge \left\lceil \frac{g_1}{3} \right\rceil.$$

Since $a(H_f) = a(H_{f_1})$ by Lemma 2.1.2, we apply Theorem 1.1.1 to deduce that $a(H_f) < \left\lceil \frac{g}{3} \right\rceil$ in this case. Thus, we see that

$$\mu_{\epsilon,g}'(a) = \mu_{\epsilon,g}(a) \left(\frac{q^{2g+\epsilon} - q^{2g+\epsilon-2}}{q^{2g+\epsilon} - q^{2g+\epsilon-1}} \right) = \mu_{\epsilon,g}(a) \left(1 + q^{-1} \right),$$

so part (2) is true by Theorem 1.1.1.

Proof of Corollary 1.1.4. Let

$$I = \left\{ f \in \mathcal{P}_{q,\epsilon}(g) \setminus \mathcal{P}'_{q,\epsilon}(g) \mid a(H_f) = a \right\}.$$

To prove part (1), we remark that the g = 0 case is immediate by Theorem 1.1.1, so suppose that g > 0. Then

$$\mu_{\epsilon,g}'(a) = \mu_{\epsilon,g}(a) \left(\frac{q^{2g+\epsilon} - q^{2g+\epsilon-2}}{q^{2g+\epsilon} - q^{2g+\epsilon-1}}\right) - \frac{1}{q^{2g+\epsilon} - q^{2g+\epsilon-1}} \sum_{f \in I} 1$$
$$= \mu_{\epsilon,g}(a) \left(1 + q^{-1}\right) - \frac{1}{q^{2g+\epsilon} \left(1 - q^{-1}\right)} \sum_{f \in I} 1,$$

so part (1) follows from the fact that

$$|I| \leq \left| \mathcal{P}_{q,\epsilon}(g) \setminus \mathcal{P}'_{q,\epsilon}(g) \right| = q^{2g+\epsilon-1} \left(1 - q^{-1} \right).$$

Turning to part (2), we see once again that if g = 0, then the result is immediate from Theorem 1.1.1; thus, we suppose g > 0. Since a > 0 we use Theorem 1.1.1 to see that for all $d \in \{0, \ldots, g\}$,

$$\mu_{\epsilon,d}(a) \le q^{-2a+1},$$

so part (1) implies that for all such d,

(14)
$$\mu'_{\epsilon,d}(a) - q^{-2a+1} \le q^{-2a}.$$

Thus, part (2) will follow if we show

$$|I| < 2q^{2g+\epsilon-2a} \left(1-q^{-1}\right).$$

To do so, note that

$$\begin{split} |I| &= \sum_{d=0}^{g-1} \sum_{\substack{f_1 \in \mathcal{P}'_{q,\epsilon}(g) \\ a(H_f) = a}} \sum_{\substack{f_2 \in \mathbb{F}_q[X] \\ \text{squarefree, of degree } g-d, \\ \text{and coprime to } f_1}} 1 & \text{by Lemma 2.1.2} \\ &\leq \sum_{d=0}^{g-1} \sum_{\substack{f_1 \in \mathcal{P}'_{q,\epsilon}(g) \\ a(H_f) = a}} \sum_{\substack{f_2 \in \mathbb{F}_q[X] \\ \text{of degree } g-d}} 1 \\ &= \sum_{d=0}^{g-1} \mu'_{\epsilon,d}(a) \left(q^{2d+\epsilon} - q^{2d+\epsilon-1}\right) q^{g-d} & \text{since } \mu'_{\epsilon,0}(a) = 0 \\ &\leq q^{g+\epsilon-2a+1} \left(1+q^{-1}\right) \sum_{d=0}^{g-1} \left(q^d - q^{d-1}\right) & \text{by Eq. (14)} \\ &= q^{g+\epsilon-2a} \left(\frac{1+q^{-1}}{1-q^{-1}}\right) \left(q^g - q^{g-1} - 1 + q^{-1}\right) \\ &< q^{2g+\epsilon-2a} \left(\frac{1+q^{-1}}{1-q^{-1}}\right) \left(1-q^{-1}\right) \\ &\leq 2q^{2g+\epsilon-2a} \left(1-q^{-1}\right), \end{split}$$

since $q \ge 3$ implies $\left(\frac{1+q^{-1}}{1-q^{-1}}\right) \le 2$.

4.2. A heuristic model and conjecture. We continue to consider the p = 3 case. As in Section 2.2, we write F for $\mathbb{F}_q(X)$. In Theorem 2.2.3 we associated to any monic squarefree polynomial $f \in \mathbb{F}_q[X]$ a particular 2-dimensional subspace V of F^3 . Suppose that g is a positive multiple of 3, say g = 3j for $j \in \mathbb{Z}_{>0}$. In this case for any $f \in \mathcal{P}'_2(g)$:

•
$$h(V) = 2j$$
, and

• $a(H_f) = j - \mu_1(V).$

Thus, we propose to model the *a*-number statistics of hyperelliptic curves of a given genus by random 2-dimensional subspaces of a fixed height. To this end, for any $j, a \in \mathbb{Z}_{\geq 0}$, let

$$\nu_j(a) = \frac{|\{V \subset F^3 \mid \dim_F(V) = 2, \ h(V) = 2j, \ \mu_1(V) = j - a\}|}{|\{V \subset F^3 \mid \dim_F(V) = 2, \ h(V) = 2j\}|}$$

In the notation of Section 3

$$\nu_j(a) = \frac{|S(2j, j - a)|}{|S(2j)|}$$

whenever $0 \le a \le j$, and 0 otherwise, since $0 \le \mu_1(V) \le \frac{h(V)}{2}$ for all 2-dimensional subspaces by Eq. (4). By Proposition 3.1.8 we immediately have the following.

Theorem 4.2.1. For any prime $p, j \in \mathbb{Z}_{>0}$, and $a \in \mathbb{Z}_{\geq 0}$ we have

$$\nu_{j}(a) = \begin{cases} 0 & \text{if } a > j, \\ q^{-2a}q & \text{if } a = j, \\ q^{-2a}(q - q^{-1}) & \text{if } j > a > 0, \\ q^{-2a}(1 - q^{-1}) & \text{if } a = 0. \end{cases}$$

Moreover,

$$\nu_0(a) = \begin{cases} 0 & if \ a > 0, \\ 1 & if \ a = 0. \end{cases}$$

In light of Theorem 4.2.1, we make the following heuristic conjecture.

Conjecture 4.2.2. If p = 3, $\epsilon \in \{1, 2\}$, and $a \in \mathbb{Z}_{>0}$, then

$$\lim_{g \to \infty} \mu_{\epsilon,g}'(a) = \begin{cases} 1 - q^{-1} & \text{if } a = 0\\ q^{-2a+1} \left(1 - q^{-2}\right) & \text{if } a > 0. \end{cases}$$

4.3. Results on the moduli space of hyperelliptic curves. As above, let K be an algebraic closure of \mathbb{F}_q . For any positive integer g, let \mathcal{H}_g denote the moduli space of smooth hyperelliptic curves of genus g over K; for any non-negative integer a, let \mathcal{H}_g^a the open sublocus of \mathcal{H}_g consisting of hyperelliptic curves with a-number a. Using the above counting results we are able to approximate the number of \mathbb{F}_q -rational points on \mathcal{H}_g^a to sufficient accuracy to determine their dimensions. In particular, we can derive the following theorem.

Theorem 4.3.1. Suppose that p = 3, that $g \in \mathbb{Z}_{>0}$, and that $a \in \mathbb{Z}_{\geq 0}$. Write $[\cdot]$ for the least integer function. Then

$$\operatorname{codim}_{\mathcal{H}_g}\left(\mathcal{H}_g^a\right) = \begin{cases} 2g-1 & \text{if } a > \left\lceil \frac{g}{3} \right\rceil, \\ 2a-1 & \text{if } 0 < a \le \left\lceil \frac{g}{3} \right\rceil, \\ 0 & \text{if } a = 0. \end{cases}$$

In fact, if $a > \left\lceil \frac{g}{3} \right\rceil$, then \mathcal{H}_q^a is empty.

Proof. The second statement is simply a rewriting of Corollary 1.1.2 (1). Thus, since \mathcal{H}_g is 2g-1 dimensional, we may assume $a \leq \left\lceil \frac{g}{3} \right\rceil$.

To prove the first statement, we note that the cases of genus 1, 2, 3, and 4 have been verified by direct calculation as carried out in [Igu58], [FvdG04], [EP07], and [Zho20] respectively. For g > 4 we follow the method of proof of Prop 7.1 of [BG01], counting hyperelliptic curves weighted by the size of their automorphism group. Using the terminology of [KS99], let $\mathcal{H}_g^a(\mathbb{F}_q)$ be the set of those curves in \mathcal{H}_g^a that are defined over \mathbb{F}_q , and define the *intrinsic* cardinality of $\mathcal{H}_q^a(\mathbb{F}_q)$ to be

$$\mathrm{IC}(\mathcal{H}_g^a(\mathbb{F}_q)) = \sum_{H \in \mathcal{H}_g^a(\mathbb{F}_q)} \left| \mathrm{Aut}_{\mathbb{F}_q}(H) \right|^{-1}.$$

By [KS99, Theorem 10.7.5], the number of \mathbb{F}_q -points on \mathcal{H}_g^a is IC($\mathcal{H}_g^a(\mathbb{F}_q)$). Thus, if we show that

$$\operatorname{IC}(\mathcal{H}_{g}^{a}(\mathbb{F}_{q})) = \begin{cases} q^{2g-2a} + O\left(q^{2g-2a-1}\right) & \text{if } 0 < a \leq \left\lceil \frac{g}{3} \right\rceil \\ q^{2g-1} + O\left(q^{2g-2}\right) & \text{if } a = 0, \end{cases}$$

we may apply Deligne's proof of the Weil Conjectures [Del74] to conclude that

$$\dim \left(\mathcal{H}_g^a \right) = \begin{cases} 2g - 2a & \text{if } 0 < a \le \left\lceil \frac{g}{3} \right\rceil \\ 2g - 1 & \text{if } a = 0; \end{cases}$$

the first statement will then follow since \mathcal{H}_q is 2g-1 dimensional.

Let S be the set of squarefree polynomials in $\mathbb{F}_q[X]$ of degree 2g + 1 or 2g + 2, so for that every hyperelliptic curve $H \in \mathcal{H}_g^a(\mathbb{F}_q)$ there exists $f \in S$ such that H_f is a model of H. Of course, the set S is acted upon by the group of \mathbb{F}_q -isomorphisms of such curves, all of which are given by a linear fractional map on X and a scalar multiple of Y (see [Nar09] for details). Let G be this group, so that $|G| = (q-1)|\operatorname{PGL}(2,q)| = (q-1)(q^3-q)$.

We claim that for any $\alpha \in (\mathbb{F}_q)^{\times}$ and $\epsilon \in \{1, 2\}$,

$$|\{f \in \alpha \mathcal{P}'_{\epsilon}(g) \mid a(f) = a\}| = (q^{2g+\epsilon} - q^{2g+\epsilon-1})\mu'_{\epsilon,g}(a).$$

Indeed, the element of G mapping $X \mapsto X$ and $Y \mapsto \sqrt{\alpha}$ induces an *a*-number preserving bijection between $\alpha \mathcal{P}'_{\epsilon}(g)$ and $\mathcal{P}'_{\epsilon}(g)$ (in fact, the *a*-number is invariant under any $\overline{\mathbb{F}}_q$ -isomorphism, see Lemma 2.1.2). Thus, as elements of G preserve *a*-numbers, we apply the orbit-stabilizer theorem to see that

$$\begin{split} \mathrm{IC}(\mathcal{H}_{g}^{a}(\mathbb{F}_{q})) &= \frac{1}{|G|} \left| \left\{ f \in S \mid a(f) = a \right\} \right| \\ &= \frac{1}{(q-1)(q^{3}-q)} \left(\sum_{\alpha \in (\mathbb{F}_{q})^{\times}} \left(q^{2g+1} - q^{2g} \right) \mu_{1,g}'(a) + \sum_{\alpha \in (\mathbb{F}_{q})^{\times}} \left(q^{2g+2} - q^{2g+1} \right) \mu_{2,g}'(a) \right) \\ &= \frac{q-1}{(q-1)(q^{3}-q)} \left(\left(q^{2g+1} - q^{2g} \right) \mu_{1,g}'(a) + \left(q^{2g+2} - q^{2g+1} \right) \mu_{2,g}'(a) \right) \\ &= \left(\frac{q^{2g-1}}{1+q^{-1}} \right) \left(q^{-1} \mu_{1,g}'(a) + \mu_{2,g}'(a) \right). \end{split}$$

Next, we apply Corollary 1.1.4(1) and (2) to deduce

$$\mathrm{IC}(\mathcal{H}_{g}^{a}(\mathbb{F}_{q})) = \left(\frac{q^{2g-1}}{1+q^{-1}}\right) \left(q^{-1}\mu_{1,g}'(a) + \mu_{2,g}'(a)\right) = \begin{cases} q^{2g-2a} + O\left(q^{2g-2a-1}\right) & \text{if } 0 < a \le \left[\frac{g}{3}\right] \\ q^{2g-1} + O\left(q^{2g-2}\right) & \text{if } a = 0, \end{cases}$$

as desired.

Corollary 4.3.2. In characteristic 3, if the strata \mathcal{H}_g^a are pure and non-empty, then they are irreducible.

Note that the for $g \leq 4$ the calculations of [Igu58], [FvdG04], [EP07], and [Zho20] show that the non-empty a-number strata in characteristic 3 are indeed irreducible, and so it seems reasonable to conjecture that this continues to be the case for all larger genera.

Acknowledgments

The authors would like to thank Rachel Pries, Jeff Achter, Joseph Gunther, Avinash Kulkarni, and Adam Logan for very helpful conversations.

References

- [BG01] Bradley W. Brock and Andrew Granville, More points than expected on curves over finite field extensions, Finite Fields Appl. 7 (2001), no. 1, 70–91, Dedicated to Professor Chao Ko on the occasion of his 90th birthday. MR 1803936
- [CEZB13] Bryden Cais, Jordan S. Ellenberg, and David Zureick-Brown, Random Dieudonné modules, random p-divisible groups, and random curves over finite fields, J. Inst. Math. Jussieu 12 (2013), no. 3, 651–676. MR 3062874
- [CL84] H. Cohen and H. W. Lenstra, Jr., *Heuristics on class groups of number fields*, Number theory, Noordwijkerhout 1983 (Noordwijkerhout, 1983), Lecture Notes in Math., vol. 1068, Springer, Berlin, 1984, pp. 33–62. MR 756082
- [Del74] Pierre Deligne, *La conjecture de Weil: I*, Inst. Hautes Études Sci. Publ. Math. (1974), no. 43, 273–307. MR 340258
- [Elk11] Arsen Elkin, The rank of the Cartier operator on cyclic covers of the projective line, J. Algebra 327 (2011), 1–12. MR 2746026
- [EP07] Arsen Elkin and Rachel Pries, Hyperelliptic curves with a-number 1 in small characteristic, Albanian J. Math. 1 (2007), no. 4, 245–252. MR 2367217
- [EVW16] Jordan S. Ellenberg, Akshay Venkatesh, and Craig Westerland, Homological stability for Hurwitz spaces and the Cohen-Lenstra conjecture over function fields, Ann. of Math. (2) 183 (2016), no. 3, 729–786. MR 3488737
- [Fre18] Sarah Frei, The a-number of hyperelliptic curves, Women in numbers Europe II, Assoc. Women Math. Ser., vol. 11, Springer, Cham, 2018, pp. 107–116. MR 3882708
- [FvdG04] Carel Faber and Gerard van der Geer, Complete subvarieties of moduli spaces and the Prym map, J. Reine Angew. Math. 573 (2004), 117–137. MR 2084584
- [FW89] Eduardo Friedman and Lawrence C. Washington, On the distribution of divisor class groups of curves over a finite field, Théorie des nombres (Quebec, PQ, 1987), de Gruyter, Berlin, 1989, pp. 227–239. MR 1024565
- [Gar15] Derek Garton, Random matrices, the Cohen-Lenstra heuristics, and roots of unity, Algebra Number Theory 9 (2015), no. 1, 149–171. MR 3317763
- [GP05] Darren Glass and Rachel Pries, *Hyperelliptic curves with prescribed p-torsion*, Manuscripta Math. 117 (2005), no. 3, 299–317. MR 2154252
- [Igu58] Jun-ichi Igusa, Class number of a definite quaternion with prime discriminant, Proc. Nat. Acad. Sci. U.S.A. 44 (1958), 312–314. MR 98728
- [KS99] Nicholas M. Katz and Peter Sarnak, Random matrices, Frobenius eigenvalues, and monodromy, American Mathematical Society Colloquium Publications, vol. 45, American Mathematical Society, Providence, RI, 1999. MR 1659828
- [Nar09] Enric Nart, *Counting hyperelliptic curves*, Adv. Math. **221** (2009), no. 3, 774–787. MR 2511037
- [Oda69] Tadao Oda, The first de Rham cohomology group and Dieudonné modules, Ann. Sci. École Norm. Sup. (4) 2 (1969), 63–135. MR 0241435
- [Oor75] Frans Oort, Which abelian surfaces are products of elliptic curves?, Math. Ann. 214 (1975), 35–47. MR 364264
- [Re01] Riccardo Re, The rank of the Cartier operator and linear systems on curves, J. Algebra 236 (2001), no. 1, 80–92. MR 1808346
- [R02] Rosen, Michael, Number Theory in Function Fields, Graduate Texts in Mathematics, vol 210, Springer-Verlag, Berlin, 2002.
- [San20] Soumya Sankar, Proportion of ordinarity in some families of curves over finite fields, arXiv eprint, v2 (2020).
- [Sti09] Henning Stichtenoth, *Algebraic function fields and codes*, second ed., Graduate Texts in Mathematics, vol. 254, Springer-Verlag, Berlin, 2009. MR 2464941
- [Thu95] Jeffrey Lin Thunder, Siegel's lemma for function fields, Michigan Math. J. 42 (1995), no. 1, 147–162. MR 1322196
- [Thu08] _____, Counting subspaces of given height defined over a function field, J. Number Theory 128 (2008), no. 12, 2973–3004. MR 2464849

[Zho20] Zijian Zhou, *Ekedahl-Oort strata on the moduli space of curves of genus four*, Rocky Mountain J. Math. **50** (2020), no. 2, 747–761. MR 4104409

FARIBORZ MASEEH DEPARTMENT OF MATHEMATICS AND STATISTICS, PORTLAND STATE UNIVERSITY *Email address:* gartondw@pdx.edu

DEPARTMENT OF MATHEMATICAL SCIENCES, NORTHERN ILLINOIS UNIVERSITY *Email address*: jthunder@niu.edu

THE TUTTE INSTITUTE FOR MATHEMATICS AND COMPUTING *Email address*: colinoftheweirs@gmail.com