Estimating the linear relation between variables that are never jointly observed

Polina ARSENTEVA¹, Mohamed Amine BENADJAOUD² and Hervé CARDOT¹

 Institut de Mathématiques de Bourgogne, UMR CNRS 5584, Université de Bourgogne, Dijon, France
 IRSN PSE-SANTE/SERAMED, Fontenay aux roses, France

June 19, 2024

Abstract

In modern experimental science there is a commonly encountered problem of estimating the coefficients of a linear regression in the context where the variables of interest can never be observed simultaneously. Assuming that the global experiment can be decomposed into sub-experiments with distinct first moments, we propose two estimators of the linear regression that take this additional information into account. We consider an estimator based on moments, and an estimator based on optimal transport theory. These estimators are proven to be consistent as well as asymptotically Gaussian under weak hypotheses. The asymptotic variance has no explicit expression, except in some particular cases, for which reason a stratified bootstrap approach is developed to build confidence intervals for the estimated parameters, whose consistency is also shown. A simulation study, assessing and comparing the finite sample performances of these estimators, demonstrated the advantages of the bootstrap approach in multiple realistic scenarios. An application to in vivo experiments, conducted in the context of studying radio-induced adverse effects on mice, revealed important relationships between the biomarkers of interest that could not be identified with the considered naive approach.

1 Introduction

In order to study the effects of a certain treatment on a living organism, in vivo experiments are often conducted. In the context of a complex organism response, scientists may be interested in studying multiple variables describing the effect on different scales. In particular, such variables of interest often include a macroscopic biomarker, only available through in vivo data, and a microscopic biomarker that can also be observed on a cellular level (i.e. in vitro). The interest in this case lies in predicting the former with the latter. For instance, in the context of studying the adverse effects induced by radiotherapy on healthy tissues, the potential outcomes of interest manifest in the form of the severity of macroscopic lesions, and predictors such as gene expression. These measures often require sacrificing the animals that they are taken from. As a result, the quantities of interest cannot be observed on the same animals. Since the goal is to establish relationships between these variables, a problem of statistical data fusion arises.



Figure (1.1) Schematic representation of the design of an in vivo experiment studying the effect of *irradiated volume.*

An illustrative example of an in vivo experiment where the variables of interest are never observed simultaneously is presented in Figure 1.1. In this experiment, mice are irradiated on the lungs with different volume, with a goal of studying the role of irradiated volume in the appearance of radio-induced adverse effects. The latter are assessed by measuring septal thickening, a histological marker of lung injuries. The other variable that is measured with the purpose of predicting the adverse effect-related variable is the expression of multiple pro-inflammatory genes. As shown in Figure 1.1, there are two independent cohorts in the study, one is used to measure gene expression, whereas the other for measuring septal thickness. This is a results of both measures being of destructive nature, which does not allow them to be taken on the same animals.

Comparing distributions of the measurements arising from the two cohorts, such as those presented in Figure 1.2, one may suspect a correlation or even a linear relationship between the variables. In order to assert whether such a relationship exists, one has to connect two variables that are never observed simultaneously, which translates into solving a data fusion problem. In this example, there are four groups indicating time points (1, 3, 6 and 12 months after irradiation) when the corresponding animals are sacrificed and the measurements are taken. Thus, the categorical variable indicating the time point, which is known for every observation, can be used as an additional variable in order to link the predictor and the predicted variables between each other.

The task of linking variables that are never jointly observed cannot be approached as a typical missing values problem, since most methods for inference on incomplete data require a sufficient overlap, which is completely absent in the case treated here. As a result, any approaches applying frameworks such as multiple imputation in the context of data fusion are unsuitable for our application. For example, Carrig et al. (2015) use multiple imputation to integrate disparate datasets allowing for the absence of the overlap, but requiring a calibration



Figure (1.2) Distribution of the data, collected from the irradiated patch under SBRT with 3 mm beam size: the expression of the gene IL6 on the left, and septal thickness on the right. The measurements were made 1, 3, 6 and 12 months after irradiation.

dataset, where all the variables of interest must be jointly observed.

Other approaches to data fusion available in literature include factor analysis (Cudeck, 2000), statistical matching (Mitsuhiro and Hoshino, 2020), Bayesian network inference (Triantafillou et al., 2010; Tsamardinos et al., 2012) and Gaussian Markov combinations (Massa and Riccomagno, 2017). These methods are designed for linking variables that are not observed simultaneously through covariates, present for both variables of interest. This corresponds to the properties of the in vivo data described above. However, the covariates in these approaches are random variables, typically continuous, and often assumed to be Gaussian, which is the case in Cudeck (2000) and Massa and Riccomagno (2017). The grouping variable available through in vivo experiments cannot be represented in a continuous form since categories such as control and sham make it impossible to assume continuity and normality. The Bayesian network approaches introduced by Triantafillou et al. (2010) and Tsamardinos et al. (2012), aimed at inferring binary causal relationships between variables, are more suitable for large datasets with a high number of covariates. Finally, currently available research in statistical matching addresses such aspects as not-at-random missingness (Mitsuhiro and Hoshino, 2021) and high dimensionality (Mitsuhiro and Hoshino, 2020). This approach is based on the idea of comparing distances between the covariates from the datasets of interest, which cannot be done by taking the group variable as the covariate. It can be noted that the goal of the aforementioned examples in statistical matching is to group individuals before imputation, which is not necessary in our case since the groups are already known.

In this work we propose an approach to estimate the relationship between the variables that are not simultaneously observed under the experimental setting described above, based on a conditional model assuming linear relationship within every component, defined by the grouping variable that is assumed to be present. An estimator derived with the method of moments as well as optimal transport solution using Wasserstein distance are considered for the real data problem in question. Both approaches do not require any overlap between the two cohorts of the experiment, and are based on weak assumptions ensuring model identifiability and moments existence. To construct confidence intervals, we propose both asymptotic and bootstrap-based results, the former being often unfeasible in practice, with the latter demonstrating practical advantages in numerous realistic settings.

2 Identification approaches

We consider a real random variable *Y* and a vector of *d* real valued random regressors $\mathbf{X} = (X_1, \ldots, X_d)$ and suppose that the following linear regression hold

$$Y = \beta_0 + \sum_{j=1}^d \beta_j X_j + \epsilon.$$
(2.1)

The residuals ϵ are supposed to be independent of the random covariates X_1, \ldots, X_d , with zero mean and variance σ_{ϵ}^2 . With destructive in vivo experiments, we can never observe simultaneously **X** and *Y*, meaning that we never have at hand the pair (**X**, *Y*) but only (**X**, .) and (., *Y*). This means that only the marginal distributions of **X** and *Y* can be estimated in presence of sampled data.

In the absence of additional information and without any strong additional hypothesis, the parameters $(\beta_0, \beta_1, ..., \beta_d)$ and the variance of the noise σ_{ϵ}^2 cannot be identified. Indeed, if for example X_1 is centered with symmetric distribution, the coefficient β_1 can only be determined up to sign change since $\beta_1 X_1$ and $\beta_1(-X_1)$ have the same distribution.

To deal with this identification issue, we consider that we can perform different experiments in which the mean value of X is allowed to vary. For that, we suppose that there are K groups (corresponding to K different experiments), defined by a discrete variable G taking values in $\{1, \dots, K\}$ observed simultaneously with Y and with X. This means that we have now access to (\mathbf{X}, G) and (Y, G) but not to (\mathbf{X}, Y, G) . We also suppose that ϵ is independent of G.

Given G = k, for k = 1, ..., K, we denote by $\mu_Y^k = \mathbb{E}(Y|G = k)$ and $\mu_{X_j}^k = \mathbb{E}(X_j|G = k)$, j = 1, ..., d, the expected values within each group.

We now present two different approaches developed to identify the vector $\beta = (\beta_0, ..., \beta_d)$ of unknown regression coefficients and the noise variance σ_{ϵ}^2 , taking account of the additional information given by the discrete variable *G*.

2.1 Moment approach

A first and simple approach is based on first moments identification. Taking the conditional expectation, given G = k, in (2.1), we have for k = 1, ..., K,

$$\mu_Y^k = \beta_0 + \sum_{j=1}^d \beta_j \mu_{X_j}^k,$$
(2.2)

since the residual term ϵ is supposed to satisfy $\mathbb{E}(\epsilon | G = k) = 0$ for $k = 1, \dots, K$.

We denote by $\boldsymbol{\mu}_{1,X}$ the $K \times (d+1)$ design matrix, whose *k*th row is equal to $(1, \boldsymbol{\mu}_X^{k\top})$ with $\boldsymbol{\mu}_X^k = (\mu_{X_1}^k, \cdots, \mu_{X_d}^k)^{\top}$, by $\boldsymbol{\mu}_Y$ the *K* dimensional vector with elements $(\mu_Y^1, \dots, \mu_Y^K)$. The *K* linear equations in (2.2) can we written in a matrix way, $\boldsymbol{\mu}_Y = \boldsymbol{\mu}_{1,X}\boldsymbol{\beta}$.

We introduce the following assumption, guaranteeing the identifiability of the model parameters:

$$\mathbf{H}_1 \quad \operatorname{rank}(\boldsymbol{\mu}_{1,X}) = d + 1,$$

meaning that there are at least $K \ge d + 1$ groups and that the d + 1 column vectors of $\mu_{1,X}$ span a vector space of dimension d + 1 in \mathbb{R}^{K} .

Lemma 2.1. If model (2.1) holds and assumption \mathbf{H}_1 is fulfilled, β is uniquely identified in terms of the conditional first order moments of \mathbf{X} and Y given G,

$$\boldsymbol{eta} = \left(\boldsymbol{\mu}_{1,X}^{ op} \boldsymbol{\mu}_{1,X}
ight)^{-1} \boldsymbol{\mu}_{1,X}^{ op} \boldsymbol{\mu}_{Y}$$

Additionally, the noise variance σ_{ϵ}^2 satisfies

$$\sigma_{\epsilon}^2 = \sigma_Y^2 - \boldsymbol{\beta}_{-0}^{\top} \boldsymbol{\Gamma}_X \boldsymbol{\beta}_{-0},$$

where σ_Y^2 is the variance of Y, Γ_X is the covariance matrix of **X** with elements $Cov(X_i, X_j) = \sum_{k=1}^{K} Cov(X_i, X_j | G = k) \mathbb{P}[G = k]$ for i and j in $\{1, \dots, d\}$, and $\beta_{-0} = (\beta_1, \dots, \beta_d)$.

The proof of Lemma 2.1 is direct and thus omitted.

2.2 Optimal transport approach

The second approach is based on optimal transport, in particular on the idea of estimating the linear transformation of the distribution of \mathbf{X} that is the closest to that of Y with respect to the Wasserstein distance (see Panaretos and Zemel (2019) for a general introduction for statisticians).

The optimal transport map T between Gaussian measures on \mathbb{R}^d is linear, and the Wasserstein distance of order 2 between two Gaussian distributions D_1 and D_2 , with $D_1 \sim \mathcal{N}(\boldsymbol{\mu}_1, \boldsymbol{\Gamma}_1)$ and $D_2 \sim \mathcal{N}(\boldsymbol{\mu}_2, \boldsymbol{\Gamma}_2)$, is equal to

$$W_2^2(D_1, D_2) = \|\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2\|^2 + \operatorname{tr}\left(\boldsymbol{\Gamma}_1 + \boldsymbol{\Gamma}_2 - 2\left(\boldsymbol{\Gamma}_2^{1/2}\boldsymbol{\Gamma}_1\boldsymbol{\Gamma}_2^{1/2}\right)^{1/2}\right),$$

where $\|.\|$ denotes the Euclidean norm and tr(A) the trace of matrix A.

If we assume that, given G = k, **X** is a Gaussian random vector and ϵ is Gaussian, we have, if the linear model (2.1) holds, that *Y* is also Gaussian given G = k, with expectation $\mu_Y^k = \beta_0 + \sum_{j=1}^d \beta_j \mu_{X_j}^k$ and variance $\sigma_{\epsilon}^2 + \beta_{-0}^{\top} \Gamma_X^k \beta_{-0}$, where Γ_X^k is the variance matrix of **X** given G = k. Thus, the Wasserstein distance between D_{γ} , the distribution of $\gamma_0 + \gamma_{-0}^{\top} \mathbf{X} + \epsilon$, and D_Y , the distribution of *Y*, is equal to

$$W_2^2(D_\gamma, D_Y) = \mathbb{E}\left[W_2^2(D_\gamma, D_Y)|G\right]$$
$$= \sum_{k=1}^K \pi_k \left[(\mu_Y^k - \alpha_0 - \gamma_{-0}^\top \mu_X^k)^2 + \left(\sigma_{Y,k} - \sqrt{\gamma_{-0}^\top \Gamma_X^k \gamma_{-0} + \sigma_\epsilon^2}\right)^2 \right]$$

where $\pi_k = \mathbb{P}[G = k]$ and $\sigma_{Y,k}^2 = \operatorname{Var}(Y|G = k)$.

We introduce the following loss criterion

$$\varphi(\boldsymbol{\gamma}, \sigma^2) = \sum_{k=1}^{K} \pi_k \left[(\boldsymbol{\mu}_Y^k - \boldsymbol{\gamma}_0 - \boldsymbol{\gamma}_{-0}^\top \boldsymbol{\mu}_X^k)^2 + \left(\sigma_{Y,k} - \sqrt{\boldsymbol{\gamma}_{-0}^\top \boldsymbol{\Gamma}_X^k \boldsymbol{\gamma}_{-0} + \sigma^2} \right)^2 \right].$$
(2.3)

which evaluates the weighted Wasserstein distance between *Y* and a linear combination of the **X** variables, contaminated by a Gaussian noise, with variance σ^2 . We state, without proof, the following Lemma ensuring that the parameters β and σ_{ϵ}^2 can be identified under general conditions.

Lemma 2.2. If model (2.1) holds and if assumption \mathbf{H}_1 is fulfilled, $\varphi(\boldsymbol{\gamma}, \sigma^2)$ has its unique minimum at $\boldsymbol{\gamma} = \boldsymbol{\beta}$ and $\sigma^2 = \sigma_{\epsilon}^2$.

3 Sampled data and estimators

We suppose that experiments are made for $K \ge 2$ different groups and that for each group k, for k = 1, ..., K, we have two independent samples $(Y_1^k, ..., Y_{n_y^k}^k)$ and $(X_{j,1}^k, ..., X_{j,n_x^k}^k)_{j=1,...,d}$, with sizes n_y^k and n_x^k . For each unit $i = 1, ..., n_x^k$ from group k, the vector of covariates is denoted by $\mathbf{X}_i^k = (X_{1,i}, ..., X_{d,i})$. We also define $N_x = \sum_{k=1}^K n_x^k$ and $N_y = \sum_{k=1}^K n_y^k$, the total number of observations of the response Y and the covariates $X_1, ..., X_d$.

As shown in Lemma 2.1 and Lemma 2.2, identification of parameter β relies on the knowledge of the first two conditional moments, given G, of Y and \mathbf{X} . For $k = 1, \ldots, K$, we denote by $\hat{\mu}_Y^k = \frac{1}{n_y^k} \sum_{i=1}^{n_y^k} Y_i^k$ and $\hat{\mu}_{X_j}^k = \frac{1}{n_x^k} \sum_{i=1}^{n_x^k} X_{j,i}^k$ the empirical means within each group k and by $\hat{\sigma}_{Y,k}^2 = \frac{1}{n_y^k} \sum_{i=1}^{n_y^k} (Y_i^k)^2 - (\hat{\mu}_Y^k)^2$, and $\hat{\Gamma}_X^k = \frac{1}{n_x^k} \sum_{i=1}^{n_x^k} \mathbf{X}_i^k (\mathbf{X}_i^k)^\top - \hat{\mu}_X^k (\hat{\mu}_X^k)^\top$, the empirical variances, with $\hat{\mu}_X^k = (\hat{\mu}_{X_1}^k, \ldots, \hat{\mu}_{X_d}^k)$. We also define the overall empirical mean and variance $\hat{\mu}_Y = \frac{1}{N_y} \sum_{k=1}^K n_y^k \hat{\mu}_Y^k$, $\hat{\mu}_X = \frac{1}{N_x} \sum_{k=1}^K n_x^k \hat{\mu}_X^k$, $\hat{\sigma}_Y^2 = \frac{1}{N_y} \sum_{k=1}^K \sum_{i=1}^{n_y^k} (Y_i^k)^2 - (\hat{\mu}_Y)^2$, $\hat{\Gamma}_X = \frac{1}{N_x} \sum_{k=1}^K \sum_{i=1}^{n_x^k} \mathbf{X}_i^k (\mathbf{X}_i^k)^\top - \hat{\mu}_X \hat{\mu}_X^\top$.

We denote by $\hat{\mu}_{1,X}$ the $K \times (d+1)$ matrix, with the first column consisting of ones, and the rest equal to $\hat{\mu}_X$.

3.1 Moment estimators

If $\hat{\mu}_{1,X}$ is full rank, moment estimators of $\beta = (\beta_0, \beta_1, \dots, \beta_d)$ can be built by considering the empirical counterpart of identification equations given in Lemma 2.1,

$$\widehat{\boldsymbol{\beta}}^{M} = \left(\widehat{\boldsymbol{\mu}}_{1,X}^{\top} \widehat{\boldsymbol{\mu}}_{1,X}\right)^{-1} \widehat{\boldsymbol{\mu}}_{1,X}^{\top} \widehat{\boldsymbol{\mu}}_{Y}$$
(3.1)

where $\hat{\mu}_Y = (\hat{\mu}_Y^1, \dots, \hat{\mu}_Y^K)$. We consider in the following a slightly more general moment estimator of β , by introducing a weight w_k given to each group k of observations, with $w_k > 0$ and $\sum_{k=1}^{K} w_k = 1$. The weighted moment estimator β is defined as the minimizer of

$$\psi(\boldsymbol{\gamma}) = \sum_{k=1}^{K} w_k \left[\widehat{\mu}_Y^k - \left(\gamma_0 + \sum_{j=1}^d \gamma_j \widehat{\mu}_{X_j}^k \right) \right]^2,$$

which is unique if $\hat{\mu}_{1,X}$ is full rank and defined by

$$\widehat{\boldsymbol{\beta}}^{M} = \left(\widehat{\boldsymbol{\mu}}_{1,X}^{\top} \mathbf{w} \widehat{\boldsymbol{\mu}}_{1,X}\right)^{-1} \widehat{\boldsymbol{\mu}}_{1,X}^{\top} \mathbf{w} \widehat{\boldsymbol{\mu}}_{Y}$$
(3.2)

with $\mathbf{w} = (w_1, \dots, w_K)$. The first moment estimator considered in (3.1) corresponds to equal weights $w_k = K^{-1}$.

We can then define the following estimator of the noise variance,

$$\widehat{\sigma}_{\epsilon}^{2,M} = \widehat{\sigma}_{Y}^{2} - (\widehat{\boldsymbol{\beta}}_{-0}^{M})^{\top} \widehat{\boldsymbol{\Gamma}}_{X} \widehat{\boldsymbol{\beta}}_{-0}^{M}.$$
(3.3)

3.2 Optimal transport estimators

Estimators of β and σ_{ϵ}^2 based on an optimal transport criterion are derived by minimizing the empirical version $\varphi_n(\gamma, \sigma^2)$ of functional $\varphi(\gamma, \sigma^2)$ defined by

$$\varphi_n(\boldsymbol{\gamma}, \sigma^2) = \sum_{k=1}^K \pi_k \left[(\widehat{\mu}_Y^k - \gamma_0 - \boldsymbol{\gamma}_{-0}^\top \widehat{\boldsymbol{\mu}}_X^k)^2 + \left(\widehat{\sigma}_{Y,k} - \sqrt{\boldsymbol{\gamma}_{-0}^\top \widehat{\boldsymbol{\Gamma}}_X^k \boldsymbol{\gamma}_{-0} + \sigma^2} \right)^2 \right].$$
(3.4)

Note that in absence of a priori information on the probability π_k of observing group k, we can set $\pi_k = K^{-1}$. We denote by $(\hat{\beta}^W, \hat{\sigma}^{2,W})$ minimizers of $\varphi_n(\gamma, \sigma^2)$ which are obtained with iterative optimization algorithms based on gradient descent. The algorithm can be initialized with $(\hat{\beta}^M, \hat{\sigma}^{2,M})$.

4 Consistency and asymptotic distribution

To study the asymptotic behavior of the estimators of β defined in previous section, we suppose that the number K of groups is kept fixed and that for all groups and all variables **X** and Y, the number of observations tends to infinity. This means that $n_{\min} = \min(n_y^1, \dots, n_y^K, n_x^1, \dots, n_x^K)$ the smallest sample size among all experiments is also supposed to tend to infinity.

Under general moment conditions, without any particular assumption on the probability law of Y and \mathbf{X} we get in the following two Lemmas that the moment estimators and the Wasserstein distance based estimators are consistent. The proofs are postponed to the Appendix.

Lemma 4.1. If $\mathbb{E}(Y^2) < +\infty$ and $\mathbb{E}(||\mathbf{X}||^2) < +\infty$, and assumption \mathbf{H}_1 is fulfilled, the sequence of estimators $(\widehat{\boldsymbol{\beta}}^M, \widehat{\sigma}_{\epsilon}^{2,M})$ defined by (3.2) and (3.3) converges in probability to $(\boldsymbol{\beta}, \sigma_{\epsilon}^2)$ when n_{\min} tends to infinity.

For the Wasserstein minimum distance estimator, since there is no explicit expression for the estimators, a compactness assumption is also made to obtain the consistency.

Lemma 4.2. If $\mathbb{E}(Y^2) < +\infty$ and $\mathbb{E}(||\mathbf{X}||^2) < +\infty$, $(\boldsymbol{\beta}, \sigma_{\epsilon}^2) \in \Theta$ and Θ is a compact set that does not contain 0, suppose that model (2.1) holds and hypothesis \mathbf{H}_1 is fulfilled, then the sequence of estimators $(\widehat{\boldsymbol{\beta}}^W, \widehat{\sigma}_{\epsilon}^{2,W})$ that minimize (3.4) converges in probability to $(\boldsymbol{\beta}, \sigma_{\epsilon}^2)$ when n_{\min} tends to infinity.

As far as the asymptotic distribution of the estimators is concerned, and for sake of simplicity and lighter notations, we suppose from now on that the number of experiments is the same for all groups and all variables, that is to say $n = n_y^1 = \cdots n_y^K = n_x^1 \cdots = n_x^K$.

Proposition 4.1. Assume that the assumptions of Lemma 4.1 are fulfilled. Then as n tends to infinity,

$$\sqrt{n}\left(\widehat{\boldsymbol{\beta}}^{M}-\boldsymbol{\beta}\right) \rightsquigarrow \mathcal{N}\left(0,\boldsymbol{\Gamma}_{\beta_{M}}\right)$$

where the expression of the asymptotic covariance matrix Γ_{β_M} is given in the proof.

This result is based on the CLT for empirical means and the application of the delta method, which involves computing the Jacobian of the inverse of matrices, making obtaining the explicit expression for Γ_{β_M} a difficult task when d > 1.

Remark 4.1. The weak convergence toward a Gaussian distribution presented in Proposition 4.1 would remain true, at the expense of heavier notations and a different asymptotic covariance matrix Γ_{β_M} , provided that there exist two constants, $0 < c \leq C$ such that

$$0 < c \le \frac{\max(n_y^1, \cdots, n_y^K, n_x^1, \cdots, n_x^K)}{\min(n_y^1, \cdots, n_y^K, n_x^1, \cdots, n_x^K)} \le C < +\infty,$$
(4.1)

and $\min(n_y^1, \cdots, n_y^K, n_x^1, \cdots, n_x^K) \to \infty$.

The asymptotic normality of $\hat{\beta}^W$ relies on classical results for M-estimators recalled in the Appendix (see Theorem A.4). Note that since we estimate simultaneously β and σ_{ϵ}^2 , an additional condition on the existence of the moments of order four for the covariates is required.

Proposition 4.2. If model (2.1) holds and hypothesis \mathbf{H}_1 is fulfilled, $\mathbb{E}(Y^2) < +\infty$ and $\mathbb{E}(||\mathbf{X}||^4) < +\infty$, $(\boldsymbol{\beta}, \sigma_{\epsilon}^2) \in \Theta$ and Θ is a compact set that does not contain (0, 0), then, as n tends to infinity,

$$\sqrt{n} \left(\begin{pmatrix} \widehat{\boldsymbol{\beta}}^W \\ \widehat{\sigma}_{\epsilon}^{2,W} \end{pmatrix} - \begin{pmatrix} \boldsymbol{\beta} \\ \sigma_{\epsilon}^2 \end{pmatrix} \right) \rightsquigarrow \mathcal{N}(0, \boldsymbol{\Gamma}_W),$$

for some covariance matrix Γ_W .

Similarly to Γ_{β_M} , the expression of the asymptotic covariance matrix Γ_W is almost impossible to explicitly derive manually, with the exception of some particularly simple cases.

5 The particular case of simple linear regression

To show the difficulty, let us examine the case of simple linear regression, that is to say d = 1. The following linear model

$$Y = \beta_0 + \beta_1 X + \epsilon$$

is supposed to hold and if there are two groups k and j such that $\mu_X^k \neq \mu_X^j$, the identification assumption \mathbf{H}_1 is fulfilled and β_0 and β_1 can be uniquely determined. The moment estimators of β_0 and β_1 defined in (3.2) have simple expressions,

$$\widehat{\beta}_0 = \widehat{\mu}_{Y,w} - \widehat{\beta}_1 \widehat{\mu}_{X,w} \tag{5.1}$$

$$\widehat{\beta}_{1} = \frac{\sum_{k=1}^{K} w_{k} \widehat{\mu}_{X}^{k} \widehat{\mu}_{Y}^{k} - \widehat{\mu}_{X,w} \widehat{\mu}_{Y,w}}{\sum_{k=1}^{K} w_{k} \left(\widehat{\mu}_{X}^{k}\right)^{2} - \widehat{\mu}_{X,w}^{2}}$$
(5.2)

where $\hat{\mu}_{Y,w} = \sum_{k=1}^{K} w_k \hat{\mu}_Y^k$ and $\hat{\mu}_{X,w} = \sum_{k=1}^{K} w_k \hat{\mu}_X^k$.

We focus on the asymptotic variance of the estimator $\hat{\beta}_1$ of the slope parameter β_1 , which is often the parameter of interest. We have

Lemma 5.1. Suppose that the model (2.1) is true, with d = 1 and $K \ge 2$. If there are two groups k and j such that $\mu_X^k \neq \mu_X^j$ the vector (β_0, β_1) is identifiable and, as n tends to infinity,

$$\sqrt{n}\left(\widehat{\beta}_1 - \beta_1\right) \rightsquigarrow \mathcal{N}(0, \sigma_{\beta_1}^2)$$

with

$$\sigma_{\beta_1}^2 = \frac{1}{(Var_w(X))^2} \sum_{k=1}^K w_k^2 \left(\beta_1^2 \sigma_{X,k}^2 + \sigma_{Y,k}^2\right) \left(\mu_X^k - \mu_{X,w}\right)^2$$

and $\mu_{X,w} = \sum_{k=1}^{K} w_k \mu_X^k$ and $Var_w(X) = \sum_{k=1}^{K} w_k (\mu_X^k - \mu_{X,w})^2$.

Lemma 5.1 shows us that even in a very simple framework (only one covariate), the asymptotic variance of the estimator of the slope β_1 is quite complicated. It also reveals that minimizing the asymptotic variance $\hat{\beta}_1$ with respect to the weights w_k is not a simple task.

6 Bootstrapping for confidence intervals

Since, as noted in the previous section, it is complicated to compute explicitly the asymptotic variance matrix of $\hat{\beta}^M$ and $\hat{\beta}^W$, we consider stratified bootstrap approaches in order to build confidence sets for β .

Our bootstrap procedure takes account of the independence between the different groups k = 1, ..., K, as well as the independence of inputs $(X_1^k, ..., X_d^k)$ and output Y^k within each group, meaning more formally that, given G = k, the joint probability measure \mathbb{P}^k of Y and \mathbf{X} is a product measure of the marginal measures $\mathbb{P}^k = \mathbb{P}_Y^k \otimes \mathbb{P}_{\mathbf{X}}^k$.

Within each group k, we draw, with equal probability and with replacement, n_y^k observations among $Y_1^k, \ldots, Y_{n_y^k}^k$. We also draw independently, with equal probability and with replacement, n_x^k observations among $\mathbf{X}_1^k, \ldots, \mathbf{X}_{n_x^k}^k$. We denote by μ_Y^{k*} and by μ_X^{k*} the empirical means and by $\sigma_{Y,k}^{2,*}$ and by Γ_X^{k*} the empirical variances of Y and \mathbf{X} in these bootstrap samples. Bootstrapped estimators $\beta^{M,*}$ and $\beta^{W,*}$ of β can now be computed by replacing the empirical moments by the bootstrap moments in (3.2) and (3.4).

To build confidence sets for the components of β based on previous bootstrap procedure, the bootstrap percentile technique, described in Chapter 4 of Shao and Tu (1995), is simple to use.

It can be noted that our estimators are smooth functions of sample means so that classical bootstrap theory applies (see for example Shao and Tu (1995), Chapter 3). For simplicity, as in Proposition 4.1, we suppose that $n = n_y^1 = \ldots = n_y^K = n_x^1 = \ldots = n_x^K$. Because of the considered experimental design, our global "empirical distribution" is made of products of marginal empirical distributions, the bootstrap for means is almost surely consistent for the Kolmogorov metric, and with Theorem 3.1 in Shao and Tu (1995) the same result holds for the estimators of β considered in this work. Then, the application of Theorem 4.1 in Shao and Tu (1995) allows to conclude that bootstrap percentile method gives consistent confidence sets for each component of β .

Proposition 6.1. Suppose that $\mathbb{E}(Y^2) < \infty$ and $\mathbb{E} ||\mathbf{X}||^2 < \infty$ and hypothesis \mathbf{H}_1 is fulfilled. Then as $n \to +\infty$, the bootstrap estimator $\boldsymbol{\beta}^{M,*}$ is strongly consistent for $\boldsymbol{\beta}$ in the Kolmogorov metric.

Consider a risk $\alpha \in (0,1)$, for each component of β , the bootstrap percentile approach provides, for a given nominal level $1 - \alpha$, a consistent confidence set.

We can also state a similar result for the estimators minimizing the Wasserstein distance, under slightly more restrictive moment conditions and compactness assumption.

Proposition 6.2. Suppose that the assumptions of Proposition 4.2 are fulfilled. As $n \to +\infty$, the bootstrap estimator ($\beta^{W,*}, \sigma_{\epsilon}^{2,W,*}$) is strongly consistent for ($\beta, \sigma_{\epsilon}^{2}$) in the Kolmogorov metric. Consider a risk $\alpha \in (0,1)$, for each component of β , the bootstrap percentile approach provides, for a given nominal level $1 - \alpha$, a consistent confidence set.

Propositions 6.1 and 6.2 are very important for practical applications since they ensure that even if we are not able to compute the explicit expression of the asymptotic variance of our estimators, we can still construct asymptotic confidence intervals thanks to simple bootstrap procedures.

7 Simulation study

7.1 Simulation design

We performed a series of simulations in order to evaluate the finite sample performances of the proposed approaches on data that resemble in vivo data originating from real experiments on mice. The number of animals observed per group is chosen for simplicity $n = n_y^1 = \cdots = n_y^K = n_x^1 \cdots = n_x^K$, and thus the weights for every group are also considered to be equal, i.e. $w_1 = \cdots = w_K = \frac{1}{K}$. For every animal $i \in \{1, \ldots, n\}$ and every subpopulation $k \in \{1, \ldots, K\}$, the predictor variable X_i^k is Gaussian univariate: $X_i^k \sim \mathcal{N}(\mu_X^k, \sigma_X^2)$, where $\mu_X^k = 9 + k$. We simulate the predicted variable independently as $Y_i^k = \beta_0 + \beta_1 X_i'^k + \epsilon_i^k$, where $X_i'^k \sim X_i^k$ and $\epsilon_i^k \sim \mathcal{N}(0, \sigma_{\epsilon}^2)$, with regression parameters $\beta_0 = 1$ and $\beta_1 = 2$. It should be noted that X_i^k and $X_i'^k$ are independent, we thus recreate the situation of the predictor and predicted variables not being simultaneously observed.

The variable sets $X = (X_i^k)_{1 \le i \le n, 1 \le k \le K}$ and $Y = (Y_i^k)_{1 \le i \le n, 1 \le k \le K}$ are simulated N_{sim} times. For each simulation, N_{boot} bootstrap samples of the size n are generated from $X^k = (X_i^k)_{1 \le k \le K}$ and $Y^k = (Y_i^k)_{1 \le k \le K}$ for each subpopulation $k \in \{1, \ldots, K\}$ independently, then moment estimators μ_X^{k*} and μ_Y^{k*} are calculated. Finally, the bootstrap sample-based estimators $\beta^{M,*} = (\beta_0^{M,*}, \beta_1^{M,*})$ and $\beta^{W,*} = (\beta_0^{W,*}, \beta_1^{W,*})$ are calculated. Based on N_{boot} estimates, we calculate 95% confidence intervals using the function *quantile* of the Python library NumPy. As a result, we obtain N_{sim} confidence intervals for each regression parameter, that we use to calculate the following quantities of interest: coverage rate of the intervals, their average amplitude, and power, i.e. the proportion of intervals not including 0. Along with the bootstrap estimators, obtained by plugging the estimated values of the moments into the expression for the limit distribution presented in Lemma 5.1. Additionally, we estimate the confidence intervals for the parameter estimators of the regression on the means by group with a naive method, assuming that the deviation of the parameter estimator from the true value divided by the estimator's standard error follows a Student's t-distribution:

$$\frac{\widehat{\beta}_j - \beta_j}{SE(\widehat{\beta}_j)} \rightsquigarrow t_{K-(d+1)} \text{ for } j \in \{0, \dots, d\}.$$



Figure (7.1) The effect of different values of ρ on the data, with K = 4 and $\sigma_X^2 = 0.75$. a) Boxplots constructed from the simulated values of X_i^k . b) Boxplots constructed from the simulated values of Y_i^k with lower relative noise level, i.e. $\rho = 1.1$. c) Boxplots constructed from the simulated values of Y_i^k with higher relative noise level, i.e. $\rho = 1.01$.

Throughout all simulations, we fix the number of simulations $N_{sim} = 500$ and the number of bootstrap samples $N_{boot} = 500$. Multiple parameters are varied to study their effect. We take the number of animals $n \in \{10, 30\}$, in particular to test whether the inference is significantly impaired if the measurements are available only for a small number of animals, which is often the case for real experimental data. We consider the number of groups $K \in \{4, 10\}$, 4 being the number of groups that is often observed in real data, and 10 being a higher number that may produce sufficiently good results with the naive approach to approximating confidence intervals with Student distribution. Additionally, the parameter σ_X^2 can be adjusted to control the extent to which the observations per group can be easily distinguished one from another. We set $\sigma_X^2 \in \{0.75, 2\}$, the first value corresponding to lower overlap between groups, and the second to higher overlap. Finally, we introduce an additional parameter $\rho \in \mathbb{R}^+$ controlling the variance of the response to the variance of noise ratio, i.e. $\rho = \frac{\sigma_Y}{\sigma_c}$, where $\sigma_Y^2 = Var(Y_i^k)$ for all *i* and *k*. The choice of adjusting the noise to signal ratio instead of the quantity of noise itself through σ_{ϵ}^2 is motivated by the fact that σ_Y depends on σ_X , hence the same level of σ_{ϵ} cannot be interpreted the same way for different values of σ_X . The variance of the noise can be expressed as follows: $\sigma_{\epsilon}^2 = \frac{\beta_1^2 \sigma_X^2}{\rho^2 - 1}$. The values of ρ are chosen to correspond to the realistic situation, namely a very noisy case and a slightly less noisy one: $\rho \in \{1.01, 1.1\}$. The effect of different values of ρ on the simulated response variable is illustrated on Figure 7.1.

7.2 Results

The results of the simulation study are presented in Supplementary Figure C.1. Firstly, the results for the moment approach in the asymptotic and bootstrap cases are very similar, thus confirming the effectiveness of the bootstrap approach. Concerning the comparison between the bootstrap procedure and the naive approach, it can be observed that in general the bootstrap estimators produce confidence intervals with smaller amplitudes and with higher power, at the expense of a slightly lower coverage rate. Whereas the empirical coverage rates in all cases are close to the nominal one (95%), the extent to which the amplitudes are smaller and the powers are bigger for the bootstrap estimators is very important in almost all cases. This



Figure (7.2) a) Coverage rates, b) amplitudes, and c) powers of the confidence intervals for the estimators of β_1 obtained from 500 simulations, with number of groups K = 4 and number of animals per group n = 10. The columns of the tables represent combinations of parameters with respect to the group dispersion parameter σ_X^2 and the noise to signal ratio ρ . The lines indicate the method used to estimate the confidence intervals: "mm (asymp)" stands for the method of moments with asymptotic confidence intervals, "mm (boot)" for the method of moments with bootstrap, "ot (boot)" for the optimal transport method with bootstrap, and "mm (student)" for the naive linear regression on means approach based on Student's distribution.

implies that the naive approach based on the Student's distribution is more likely to produce false negatives in terms of significance. This trend is further amplified by the number of groups parameter: whereas the results are overall worsened with the decrease in either the number of animals or groups, it is the case with the small number of groups that demonstrates the biggest difference in the approaches (the case with few groups and animals is presented in Figure 7.2). Indeed, in almost all cases within the tables with K = 4 we observe the amplitudes approximately twice as important for the naive approach, and a similar trend in terms of lower powers. The latter result is important since lower power implies bigger probability of not detecting a significant relationship between the predictor and the predicted variables, when it is actually present. Overall, these results mean that the proposed bootstrap estimators are more effective if the experimental design entails a small number of groups.

Concerning the remaining two parameters, as expected, in general the best results are obtained with lower σ_X^2 and higher ρ . In most cases, the results for the naive and the bootstrap estimators are either both good or both bad in terms of power, with the latter being slightly better. A particularly complicated case can be distinguished, with high dispersion, high noise level, few groups and few animals, where all estimators drastically fail: we observe almost equally bad powers (0.1 for both bootstrap estimators and 0.09 for the naive estimator), despite the significant difference in amplitudes. On the other hand, we can also distinguish two cases where the powers of the bootstrap estimators are more than 90%, whereas those of the naive estimator are under 50%: in both cases there are 4 groups and high noise, in the first case there are only 10 animals but lower dispersion, in the second case high dispersion level is compensated but a higher number of animals. This implies that if the underlying distributions per group are characterized by a reasonable amount of overlap, or a significant overlap is compensated by having more observations, the bootstrap estimators manage to detect the significant relationship in most cases, unlike the naive estimator.

Lastly, it can be observed that the estimator based on optimal transport produces confidence intervals with slightly smaller amplitudes compared to the method of moments estimator. The difference appears to be relatively more important in the cases with higher group overlap $\sigma_X^2 = 2$. However, the powers are not affected by this difference. This may be explained by a more important bias associated with the optimal transport estimator. The estimator will likely produce better results in terms of the power than the method of moments estimator if the bias is corrected.

8 Application to real data

In order to illustrate the proposed estimators on real data, we studied the data previously mentioned in Section 1, obtained from experiments conducted on mice in order to assess the adverse effects induced in the context of different irradiated volume. In the course of these experiments, mice were exposed to either stereotactic body radiation therapy (SBRT) with different beam sizes at 90 Gy on the left lung, or whole-thorax irradiation (WTI) at 19 Gy. For one mouse cohort, the expressions of tho pro-inflammatory genes (IL6 and TNF) were measured, for the other cohort the measurements of the thickness of the alveolar septas were made due to its role as a severity measure of radio-induced pulmonary lesions. In the case of the SBRT, the measurements were taken in multiple locations: the irradiated patch (inside the irradiation field), the remaining part of the left lung referred to as ipsilateral lung, and the right lung (contralateral lung). A control condition is also considered, where the gene expressions are measured without prior radiation exposure. The goal of this statistical analysis is to determine whether there is a statistical association between the gene expression as a predictor and the septal thickening as as outcome. Our approach is applied since the variables are measured on different animals, but within each irradiation condition there are shared groups in terms of the measurement time points.

			М	ETHOD OF MO	MENTS	0	PTIMAL TRANS	SPORT	LIN. REG. ON MEANS			
				(BOOTSTRA	P)		(BOOTSTRA	P)	(Student)			
Loc.	Vol.	Gene	\widehat{eta}_1 95% C.I. SIGN			$\widehat{\beta}_1$	95% C.I.	SIGNIF.	$\widehat{\beta}_1$	95% C.I.	SIGNIF.	
Contro	IL6	2.23	(-1.99, 2.28)	×	0.83	(-0.83, 0.97)	×	2.23	(-2.65, 7.12)	×		
		TNF	2	(-2.0, 2.82)	×	0.88	(-1.0, 1.11)	×	2	(-1.72, 5.72)	×	
	1 mm	IL6	0.43	(-0.23, 1.2)	×	0.2	(-0.2, 0.84)	×	0.43	(-1.32, 2.17)	×	
IPSILATERAL		TNF	0.2	(-0.23, 0.84)	×	0.16	(-0.18, 0.84)	×	0.2	(-1.26, 1.66)	×	
LUNG	3 mm	IL6	0.05	(-0.34, 0.46)	×	0.06	(-0.33, 0.49)	×	0.05	(-1.65, 1.76)	×	
		TNF	0.65	(-0.12, 1.6)	×	0.63	(-0.12, 1.46)	×	0.65	(-1.57, 2.87)	×	
	1 mm	IL6	1.03	(-0.57, 2.19)	×	0.46	(-0.3, 1.0)	×	1.03	(-0.88, 2.94)	×	
RIGHT		TNF	1.05	(-0.47, 2.43)	×	0.66	(-0.31, 1.26)	×	1.05	(-1.01, 3.11)	×	
LUNG	3 mm	IL6	0.3	(-1.45, 1.12)	×	0.37	(-0.74, 0.86)	×	0.3	(-4.94, 5.53)	×	
		TNF	2.02	(0.27, 4.1)	~	1.05	(0.04, 1.44)	~	2.02	(0.03, 4.02)	~	
	1 mm	IL6	0.85	(-0.7, 2.38)	×	0.61	(-0.56, 1.6)	×	0.85	(-3.75, 5.45)	×	
IRRADIATED		TNF	0.85	(-0.6, 2.3)	×	0.69	(-0.54, 1.53)	×	0.85	(-3.96, 5.66)	×	
РАТСН	3 mm	IL6	1.35	(0.22, 2.47)	~	1.3	(0.21, 2.26)	~	1.35	(-1.8, 4.5)	×	
		TNF	3.81	(1.01, 6.37)	~	3.37	(0.99, 5.33)	~	3.81	(-1.86, 9.48)	×	
WHOLE THO	ORAX	IL6	3.7	(1.53, 5.99)	~	2.51	(1.39, 3.43)	~	3.7	(1.11, 6.29)	~	
IRRADIATION		TNF	2.35	(0.57, 4.73)	~	1.67	(0.53, 1.88)	~	2.35	(-3.86, 8.57)	×	

Table (8.1) Results of estimation of the linear regression slope predicting septal thickening with the pro-inflammatory genes expression, with three methods, for control (no irradiation), WTI and SBRT with different beam sizes, with measurements taken in different parts of lungs.



Figure (8.1) Linear model prediction of septal thickness based on IL6 expression, plotted for different locations and beam sizes, with the results from two bootstrap estimators.

In order to ensure the comparability of the results for different treatment conditions and genes, the data were centered and reduced with respect to the global mean and standard deviation prior to estimation. The linear regression parameters were then estimated with three estimators the same way it was done in the simulation study in Section 7. The focus is placed on estimating the slope parameter β_1 in particular. The results are presented in Table 8.1, containing the estimations of β_1 as well as the estimated confidence intervals for the slope estimator, and the corresponding test result on the significance of the estimated relationship.

Major differences between the results for the naive and the bootstrap estimators can be observed in terms of the detected significance. On the one hand, the relationship between the pro-inflammatory genes and septal thickening has been identified by all methods in the case of whole-thorax irradiation (both genes for the bootstrap estimators, and only one for the naive estimator), which is an expected result. The results are also consistent for all methods in the case without radiation exposure (control), where no significant relationship has been identified, as it is expected. On the other hand, we expect to identify a strong correlation in the case of the measurements taken directly from the irradiated patch. This is only the case for the bootstrap estimators, but not for the naive one. This results is in accordance with the results we obtain with simulated data: the confidence intervals are often overestimated with the naive approach, which may result in false negatives in terms of significance.

Among the SBRT irradiation configurations, only the 3mm beam size has shown a significant correlation between inflammatory genes and septal thickening. These results are consistent with literature, indicating it as the beam size starting with which the long-term lesions appear (Bertho et al., 2020). Multiple significant associations have been identified with the bootstrap estimators in the ipsilateral lung and in the right lung for the beam size of 3 mm.

Finally, among the cases where a significant relationship has been detected, the estimated values of the slope are always positive, which indicates a general radio-induced up-regulation trend. These values are in general bigger in case of whole-thorax irradiaion and within the patch than for the ipsilateral or right lung for SBRT, which suggests a stronger correlation between the inflammatory process and lung injuries under high dose/volume irradiation conditions. These results, which are in line with the biological knowledge, could not have been established within the framework of classical statistical regression approaches due to non-simultaneous observations. This effect is illustrated in Figure 8.1 on the example of the linear model prediction made for the gene IL6.

9 Discussion

This work focuses on a statistical framework designed for extracting dependencies from experiments, specifically introducing linear regression estimators in the context where the predictor and the predicted variables are never jointly observed. In this work we chose the basic linear multivariate setting, prioritizing simplicity and computational feasibility. Particularly, the estimator based on the method of moments makes no hypotheses on data distribution and can be calculated explicitly. The estimator based on optimal transport includes a simple optimization problem, and is based on the Gaussian form of the Wasserstein distance but does not technically require the data to be Gaussian, seeking to approximate them with Gaussian variables in whatever case.

However, these approaches are inapplicable in the cases where linear relationship hypothesis cannot be satisfied. For instance, it is the case with predicting survival data with some continuous biomarker, which is of particular interest in research into radio-induced adverse effects. To be able to consider such scenarios, our model can be extended to a more general case, namely with generalized linear model. The optimal transport estimator appears promising in this context given the fact that the Wasserstein distance allows to compare probability distributions of different nature (for example, continuous and discrete).

Another further research direction lies in investigating alternative methods based on integrated likelihood and Bayesian approaches, which are likely to produce better results in many cases but require putting priors on distributions.

Finally, it would be of interest to work on improving theoretical properties of our estimators for finite samples, namely correcting the negative bias that appears for both estimators. The latter is particularly important in the case of the optimal transport estimator, which may be arising naturally with the Wasserstein distance (Wasserman, 2023). Correcting this bias would considerably improve the estimator, making it competitive with the approaches mentioned earlier that make numerous assumptions on the data.

Software availability

Python code and the experimental data are freely available at github.com/parsenteva/vivo_lm.

Acknowledgements

The authors gratefully acknowledge the funding from the European Union through the PO FEDER-FSE Bourgogne 2014/2020 programs as part of the project ModBioCan2020, and from the Institut de Radioprotection et de Sûreté nucléaire as part of the project ROSIRIS. The authors also warmly thank Dr. Agnès François and Dr. Annaïg Bertho for the experimental data, as well as Dr. Patrick Tardivel for his numerous remarks that helped to improve the presentation of the manuscript.

Conflict of Interest: none declared.

References

- Bertho, A., Santos, M. D., Buard, V., Paget, V., Guipaud, O., Tarlet, G., Milliat, F., and François, A. (2020). Preclinical model of stereotactic ablative lung irradiation using arc delivery in the mouse: Effect of beam size changes and dose effect at constant collimation. *International Journal of Radiation Oncology, Biology, Physics*, 107(3):548–562. Publisher: Elsevier.
- Carrig, M. M., Manrique-Vallier, D., Ranby, K. W., Reiter, J. P., and Hoyle, R. H. (2015). A nonparametric, multiple imputation-based method for the retrospective integration of data sets. *Multivariate Behavioral Research*, 50(4):383–397.
- Cudeck, R. (2000). An estimate of the covariance between variables which are not jointly observed. *Psychometrika*, 65(4):539–546.
- Magnus, J. R. and Neudecker, H. (2019). *Matrix differential calculus with applications in statistics and econometrics*. Wiley Ser. Probab. Stat. Hoboken, NJ: John Wiley & Sons, 3rd updated edition edition.
- Massa, M. S. and Riccomagno, E. (2017). Algebraic representations of Gaussian Markov combinations. *Bernoulli*, 23(1):626–644. Publisher: Bernoulli Society for Mathematical Statistics and Probability.
- Mitsuhiro, M. and Hoshino, T. (2020). Kernel canonical correlation analysis for data combination of multiple-source datasets. *Japanese Journal of Statistics and Data Science*, 3(2):651–668.
- Mitsuhiro, M. and Hoshino, T. (2021). Bayesian data combination model with Gaussian process latent variable model for mixed observed variables under NMAR missingness. arXiv:2109.00462 [stat].
- Newey, W. K. and McFadden, D. (1994). Large sample estimation and hypothesis testing. In *Handbook of econometrics, Vol. IV*, volume 2 of *Handbooks in Econom.*, pages 2111–2245. North-Holland, Amsterdam.
- Panaretos, V. M. and Zemel, Y. (2019). Statistical aspects of Wasserstein distances. *Annu. Rev. Stat. Appl.*, 6:405–431.
- Shao, J. and Tu, D. (1995). *The Jackknife and Bootstrap*. Springer Ser. Stat. New York, NY: Springer-Verlag.
- Triantafillou, S., Tsamardinos, I., and Tollis, I. (2010). Learning causal structure from overlapping variable sets. In *Proceedings of the Thirteenth International Conference on Artificial Intelligence and Statistics*, pages 860–867. JMLR Workshop and Conference Proceedings. ISSN: 1938-7228.
- Tsamardinos, I., Triantafillou, S., and Lagani, V. (2012). Towards Integrative Causal Analysis of Heterogeneous Data Sets and Studies. *Journal of Machine Learning Research*, 13(39):1097–1157.
- van der Vaart, A. W. (1998). *Asymptotic statistics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge.
- Wasserman, L. (2023). Optimal transport in statistics. 54th Journées de Statistique of the SFdS, Université libre de Bruxelles.

A Some classical theorems in asymptotic statistics

A proof of the classical continuous mapping theorem can be found in van der Vaart (1998) (Theorem 2.3).

Theorem A.1. (Continuous mapping theorem).

Let $g : \mathbb{R}^d \to \mathbb{R}^m$ be continuous at every point of C such that $\mathbb{P}[X \in C] = 1$. If the sequence of random variables $(X_n)_{n\geq 1}$ converges in distribution (resp. probability, resp. almost surely) to X then $(g(X_n))_{n\geq 1}$ converges in distribution (resp. probability, resp. almost surely) to g(X).

We also recall some well known results that are useful to show the consistency of estimators $\hat{\theta}_n$ defined as the minimizers of functionals $Q_n(\theta)$ which have some regularity properties at the limit.

Theorem A.2. (Lemma 2.9 in Newey and McFadden (1994))

Suppose that $\theta \in \Theta$ and Θ is compact, $Q_0(\theta)$ is continuous and $\forall \theta \in \Theta$, $Q_n(\theta) \to Q_0(\theta)$ in probability as n tends to infinity. If there is $\alpha > 0$ and $B_n = O_p(1)$ such that

$$\forall (\widetilde{\theta}, \theta) \in \Theta \times \Theta, \ |Q_n(\widetilde{\theta}) - Q_n(\theta)| \le B_n \|\widetilde{\theta} - \theta\|^o$$

then

$$\sup_{\theta\in\Theta} |Q_n(\theta) - Q_0(\theta)| \to 0$$
 in probability.

Theorem A.3. (Theorem 2.1 in Newey and McFadden (1994))

Suppose that $\theta \in \Theta$ and Θ is compact, $Q_0(\theta)$ is continuous $\forall \theta \in \Theta$. If $Q_0(\theta)$ is uniquely maximized at θ_0 and, as n tends to infinity, $\sup_{\theta \in \Theta} |Q_n(\theta) - Q_0(\theta)| \to 0$ in probability, then $\widehat{\theta}_n \to \theta_0$ in probability.

Under additional hypotheses, we also get the asymptotic normality of the sequence of estimators $\hat{\theta}_n$ of θ_0 . we denote by $\nabla_{00}Q_n(\theta)$ the Hessian matrix of functional Q_n evaluated at θ .

Theorem A.4. (Theorem 3.1 in Newey and McFadden (1994))

Suppose that $\widehat{\theta}_n \to \theta_0$ in probability, (i) θ_0 is an interior point of Θ , (ii) $Q_n(\theta)$ is twice differentiable in a neighborhood \mathcal{N} of θ_0 , (iii) $\sqrt{n}\nabla_0 Q_n(\theta_0) \rightsquigarrow \mathcal{N}(0, \Sigma)$, (iv) there is $\mathbf{H}(\theta)$ continuous at θ_0 and $\sup_{\theta \in \mathcal{N}} \|\nabla_{00}Q_n(\theta) - \mathbf{H}(\theta)\| \to 0$ in probability (v) $\mathbf{H} = \mathbf{H}(\theta_0)$ is non singular. Then

$$\sqrt{n}\left(\widehat{\theta}_n - \theta_0\right) \rightsquigarrow \mathcal{N}\left(0, \mathbf{H}^{-1} \mathbf{\Sigma} \mathbf{H}^{-1}\right)$$

We also recall the central limit theorem for bootstrap means (see Theorem 23.4 in van der Vaart (1998) for a proof).

Theorem A.5. (*CLT for bootstrap means*)

Let X_1, X_2, \ldots be *i.i.d.* random vectors with mean μ and covariance matrix Γ . Then conditionally on X_1, X_2, \ldots , for almost every sequence X_1, X_2, \ldots

$$\sqrt{n}\left(\overline{X}_{n}^{*}-\overline{X}_{n}\right)\rightsquigarrow\mathcal{N}\left(0,\mathbf{\Gamma}\right)$$

where \overline{X}_n is the empirical mean and \overline{X}_n^* is the empirical mean of *n* independent observations drawn from the empirical distribution.

B Proofs

Proof. of Lemma 4.1

First note that the assumptions $\mathbb{E}(Y^2) < +\infty$ and $\mathbb{E}(\|\mathbf{X}\|^2) < +\infty$ ensure the existence of σ_Y^2 and Γ_X . From the law of large numbers, we have that for all $k \in \{1, \ldots, K\}$, $\hat{\mu}_{1,X}^k \to \mu_X^k$ and $\hat{\mu}_Y^k \to \mu_Y^k$ in probability when n_{\min} tends to infinity.

We deduce from the continuous mapping theorem that $\hat{\mu}_{1,X}^{\top} \hat{\mu}_{1,X} \rightarrow \mu_{1,X}^{\top} \mathbf{w} \mu_{1,X}$ and $\hat{\mu}_{1,X}^{\top} \mathbf{w} \hat{\mu}_{Y} \rightarrow \mu_{1,X}^{\top} \mathbf{w} \mu_{Y}$ in probability. Under hypothesis \mathbf{H}_{1} , the inverse being continuous in a neighborhood of $\mu_{1,X}^{\top} \mathbf{w} \mu_{1,X}$ another application of the continuous mapping theorem gives that $\left(\hat{\mu}_{1,X}^{\top} \mathbf{w} \hat{\mu}_{1,X}\right)^{-1} \rightarrow \left(\mu_{1,X}^{\top} \mathbf{w} \mu_{1,X}\right)^{-1}$ and

$$\widehat{\boldsymbol{\beta}}^{M} = \left(\widehat{\boldsymbol{\mu}}_{1,X}^{\top} \mathbf{w} \widehat{\boldsymbol{\mu}}_{1,X}\right)^{-1} \widehat{\boldsymbol{\mu}}_{1,X}^{\top} \mathbf{w} \widehat{\boldsymbol{\mu}}_{Y} \rightarrow \left(\boldsymbol{\mu}_{1,X}^{\top} \mathbf{w} \boldsymbol{\mu}_{1,X}\right)^{-1} \boldsymbol{\mu}_{1,X}^{\top} \mathbf{w} \boldsymbol{\mu}_{Y} = \boldsymbol{\beta}$$

in probability as n_{\min} tends to infinity.

The law of large numbers gives that $\widehat{\Gamma}_X^2 \to \Gamma_X^2$ and $\widehat{\sigma}_Y^2 \to \sigma_Y^2$ in probability and we deduce, with another application of the continuous mapping theorem, that $\widehat{\sigma}_Y^2 - \widehat{\boldsymbol{\beta}}^\top \widehat{\Gamma}_X^2 \widehat{\boldsymbol{\beta}} \to \sigma_Y^2 - \boldsymbol{\beta}^\top \Gamma_X^2 \boldsymbol{\beta} = \sigma_{\epsilon}^2$ in probability as n_{\min} tends to infinity.

Proof. of Lemma 4.2

The proof is based on Lemma 2.9 and Theorem 2.1 in Newey and McFadden (1994), which are recalled in Appendix A.

The law of large numbers and the continuous mapping theorem give us that for all $(\gamma, \sigma_{\gamma}^2) \in \Theta$, $\varphi_n(\gamma, \sigma_{\gamma}^2) \to \varphi(\gamma, \sigma_{\gamma}^2)$ in probability, when n_{\min} tends to infinity.

Consider now $(\alpha, \sigma_{\alpha}^2) \in \Theta$. We have,

$$\begin{split} (\widehat{\mu}_{Y}^{k} - \gamma_{0} - \boldsymbol{\gamma}_{-0}^{\top} \widehat{\boldsymbol{\mu}}_{X}^{k})^{2} - (\widehat{\mu}_{Y}^{k} - \alpha_{0} - \boldsymbol{\alpha}_{-0}^{\top} \widehat{\boldsymbol{\mu}}_{X}^{k})^{2} \Big| &= \left| (\boldsymbol{\alpha} - \boldsymbol{\gamma})^{T} \begin{pmatrix} 1 \\ \widehat{\boldsymbol{\mu}}_{X}^{k} \end{pmatrix} \left(2\widehat{\mu}_{Y}^{k} - (\boldsymbol{\alpha} + \boldsymbol{\gamma})^{T} \begin{pmatrix} 1 \\ \widehat{\boldsymbol{\mu}}_{X}^{k} \end{pmatrix} \right) \right| \\ &\leq \|\boldsymbol{\alpha} - \boldsymbol{\gamma}\| A_{n}^{k}, \end{split}$$

with Cauchy-Schwarz inequality and $A_{n,k} = \mathcal{O}_p(1)$ because $\|\widehat{\boldsymbol{\mu}}_X^k\| = O_p(1)$, $\widehat{\boldsymbol{\mu}}_Y^k = O_p(1)$ and for some constant C_1 that does not depend on $\boldsymbol{\alpha}$ and $\boldsymbol{\gamma}$, $\|\boldsymbol{\alpha} + \boldsymbol{\gamma}\| \leq C_1 < \infty$ because Θ is supposed to be compact.

On the other hand, we have

$$\begin{vmatrix} \left(\widehat{\sigma}_{Y,k} - \sqrt{\boldsymbol{\gamma}_{-0}^{\top} \widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\gamma}_{-0}} + \sigma_{\gamma}^{2} \right)^{2} - \left(\widehat{\sigma}_{Y,k} - \sqrt{\boldsymbol{\alpha}_{-0}^{\top} \widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\alpha}_{-0}} + \sigma_{\alpha}^{2} \right)^{2} \end{vmatrix} \\ = \left| \sqrt{\boldsymbol{\alpha}_{-0}^{\top} \widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\alpha}_{-0}} + \sigma_{\alpha}^{2} - \sqrt{\boldsymbol{\gamma}_{-0}^{\top} \widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\gamma}_{-0}} + \sigma_{\gamma}^{2}} \right| \left(2\widehat{\sigma}_{Y,k} + \sqrt{\boldsymbol{\alpha}_{-0}^{\top} \widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\alpha}_{-0}} + \sigma_{\alpha}^{2} + \sqrt{\boldsymbol{\gamma}_{-0}^{\top} \widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\gamma}_{-0}} + \sigma_{\gamma}^{2}} \right) \\ = \left| \sqrt{\boldsymbol{\alpha}_{-0}^{\top} \widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\alpha}_{-0}} + \sigma_{\alpha}^{2}} - \sqrt{\boldsymbol{\gamma}_{-0}^{\top} \widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\gamma}_{-0}} + \sigma_{\gamma}^{2}} \right| O_{p}(1)$$

since Θ is compact and $\|\widehat{\Gamma}_X^k\|_{sp} = O_p(1)$, where $\|.\|_{sp}$ denotes the spectral norm. Because $\alpha_{-0}^{\top}\widehat{\Gamma}_X^k\alpha_{-0} - \gamma_{-0}^{\top}\widehat{\Gamma}_X^k\gamma_{-0} = \alpha_{-0}^{\top}\widehat{\Gamma}_X^k(\alpha_{-0} - \gamma_{-0}) + (\alpha_{-0} - \gamma_{-0})^{\top}\widehat{\Gamma}_X^k\gamma_{-0}$ we have, for some constant $C_{2,k} > 0$,

$$\left|\boldsymbol{\alpha}_{-0}^{\top}\widehat{\boldsymbol{\Gamma}}_{X}^{k}\boldsymbol{\alpha}_{-0} - \boldsymbol{\gamma}_{-0}^{\top}\widehat{\boldsymbol{\Gamma}}_{X}^{k}\boldsymbol{\gamma}_{-0}\right| \leq C_{2,k} \left\|\widehat{\boldsymbol{\Gamma}}_{X}^{k}\right\|_{sp} \|\boldsymbol{\alpha} - \boldsymbol{\gamma}\|.$$
(B.1)

Using now the fact that function $x \mapsto \sqrt{x}$ is concave and differentiable, we have for x > 0 and y > 0 that $\sqrt{y} \le \sqrt{x} + \frac{y-x}{2\sqrt{x}}$. Thus, if y > x > 0 then $0 < \sqrt{y} - \sqrt{x} \le \frac{y-x}{2\sqrt{x}}$ and if x > y > 0, then $0 < \sqrt{x} - \sqrt{y} \le \frac{x-y}{2\sqrt{y}}$. Consequently, we have $|\sqrt{y} - \sqrt{x}| \le \frac{|x-y|}{2\min(\sqrt{x},\sqrt{y})}$ and we deduce that,

$$\left|\sqrt{\boldsymbol{\alpha}_{-0}^{\top} \widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\alpha}_{-0} + \sigma_{\alpha}^{2}} - \sqrt{\boldsymbol{\gamma}_{-0}^{\top} \widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\gamma}_{-0} + \sigma_{\gamma}^{2}}\right| \leq B_{n}^{k} \left(\|\boldsymbol{\alpha} - \boldsymbol{\gamma}\| + |\sigma_{\alpha}^{2} - \sigma_{\gamma}^{2}|\right)$$
(B.2)

where $B_n^k = O_p(1)$.

Combining previous inequalities, we get

$$\left|\varphi_{n}(\boldsymbol{\gamma},\sigma_{\gamma}^{2})-\varphi_{n}(\boldsymbol{\alpha},\sigma_{\alpha}^{2})\right| \leq \left(\left\|\boldsymbol{\alpha}-\boldsymbol{\gamma}\right\|+\left|\sigma_{\alpha}^{2}-\sigma_{\gamma}^{2}\right|\right)\sum_{k=1}^{K}\pi_{k}\left(B_{n}^{k}+A_{n}^{k}\right),\tag{B.3}$$

with $\sum_{k=1}^{K} \pi_k (B_n^k + A_n^k) = O_p(1)$. As a result, it can be deduced from Lemma 2.9 in Newey and McFadden (1994) that

$$\sup_{(\boldsymbol{\gamma},\sigma_{\boldsymbol{\gamma}}^2)\in\Theta} \left|\varphi_n(\boldsymbol{\gamma},\sigma_{\boldsymbol{\gamma}}^2) - \varphi(\boldsymbol{\gamma},\sigma_{\boldsymbol{\gamma}}^2)\right| \to 0 \quad \text{ in probability.}$$

We conclude the proof by recalling that $\varphi(\gamma, \sigma_{\gamma}^2)$ attains its unique minimum at $(\beta, \sigma_{\epsilon}^2) \in \Theta$ if assumption \mathbf{H}_1 is fulfilled, so that $(\widehat{\beta}^W, \widehat{\sigma}^{2,W}) \to (\beta, \sigma_{\epsilon}^2)$ in probability in view of Theorem 2.1 in Newey and McFadden (1994).

Proof. of Proposition 4.1

The central limit theorem applies directly to the independent sequences of independent random variables $(\mathbf{X}_1^1, \dots, \mathbf{X}_n^1), \dots, (\mathbf{X}_1^K, \dots, \mathbf{X}_n^K)$ and $(Y_1^1, \dots, Y_n^1), \dots, (Y_1^K, \dots, Y_n^K)$ so that, as *n* tends to infinity

$$\sqrt{n} \begin{pmatrix} \widehat{\boldsymbol{\mu}}_X^1 - \boldsymbol{\mu}_X^1 \\ \vdots \\ \widehat{\boldsymbol{\mu}}_X^K - \boldsymbol{\mu}_X^K \\ \widehat{\boldsymbol{\mu}}_Y^1 - \boldsymbol{\mu}_Y^1 \\ \vdots \\ \widehat{\boldsymbol{\mu}}_Y^K - \boldsymbol{\mu}_Y^K \end{pmatrix} \rightsquigarrow \mathcal{N}(0, \boldsymbol{\Gamma}_{\mu})$$
(B.4)

where Γ_{μ} is a block diagonal matrix, with diagonal elements $(\Gamma_X^1, \ldots, \Gamma_X^K, \sigma_{Y,1}^2, \ldots, \sigma_{Y,K}^2)$, with $\Gamma_X^k = \operatorname{Var}(\mathbf{X}|G = k) = \mathbb{E}\left(\mathbf{X}^k(\mathbf{X}^k)^{\top}\right) - \boldsymbol{\mu}_X^k(\boldsymbol{\mu}_X^k)^{\top}$ and $\sigma_{Y,k}^2 = \operatorname{Var}(Y|G = k)$. Consider the

application $g : \mathbb{R}^{dK+K} \to \mathbb{R}^{d+1}$ defined by

$$g(\boldsymbol{\mu}_X^1,\ldots,\boldsymbol{\mu}_X^K,\boldsymbol{\mu}_Y^1,\ldots,\boldsymbol{\mu}_Y^K) = \left(\boldsymbol{\mu}_{1,X}^\top \mathbf{w} \boldsymbol{\mu}_{1,X}\right)^{-1} \boldsymbol{\mu}_{1,X}^\top \mathbf{w} \boldsymbol{\mu}_Y.$$

Application *g* is differentiable at $\theta = (\mu_X^1, \dots, \mu_X^K, \mu_Y^1, \dots, \mu_Y^K)$, with non null Jacobian matrix denoted by \mathbf{J}_{θ} (see Chapter 8 and more particularly Theorem 8.3 in Magnus and Neudecker (2019)). The application of the Delta method (see Theorem 3.1 in van der Vaart (1998)) permits to get the asymptotic normality convergence result,

$$\sqrt{n}\left(\widehat{\boldsymbol{\beta}}^{M}-\boldsymbol{\beta}\right) \rightsquigarrow \mathcal{N}\left(0,\boldsymbol{\Gamma}_{\boldsymbol{\beta}_{M}}\right),$$

where $\Gamma_{\beta_M} = \mathbf{J}_{\theta} \Gamma_{\mu} \mathbf{J}_{\theta}^{\top}$.

Proof. of Proposition 4.2

The proof consists in checking the different points of Theorem A.4. Point (i) is satisfied by the hypotheses, and the point (ii) follows directly from the fact $\varphi_n(\gamma, \sigma^2)$ is twice-differentiable in a neighborhood of $(\beta, \sigma_{\epsilon}^2)$. To show that (iii) is fulfilled, we consider the following expansion, based on the empirical version of the gradient of φ ,

$$\nabla \varphi_{n} = \begin{pmatrix} -2\sum_{k=1}^{K} \pi_{k} \left(\widehat{\mu}_{Y}^{k} - \beta_{0} - \boldsymbol{\beta}_{-0}^{\top} \widehat{\mu}_{X}^{k} \right) \\ -2\sum_{k=1}^{K} \pi_{k} \left[\left(\widehat{\mu}_{Y}^{k} - \beta_{0} - \boldsymbol{\beta}_{-0}^{\top} \widehat{\mu}_{X}^{k} \right) \widehat{\mu}_{X}^{k} + \left(\frac{\widehat{\sigma}_{Y,k}}{\sqrt{\boldsymbol{\beta}_{-0}^{\top} \widehat{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} + \sigma_{\epsilon}^{2}}} - 1 \right) \widehat{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \right] \\ \sum_{k=1}^{K} \pi_{k} \left(1 - \frac{\widehat{\sigma}_{Y,k}}{\sqrt{\boldsymbol{\beta}_{-0}^{\top} \widehat{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} + \sigma_{\epsilon}^{2}}} \right)$$
(B.5)

Since model (2.1) holds, $\nabla \varphi = 0$ and $\hat{\mu}_Y^k - \beta_0 - \beta_{-0}^\top \hat{\mu}_X^k = (\hat{\mu}_Y^k - \mu_Y^k) - \beta_{-0}^\top (\hat{\mu}_X^k - \mu_X^k)$, we thus deduce with (B.4) the asymptotic normality of the first component of the gradient $\nabla \varphi_n$, that is to say $\sqrt{n} \left(-2\sum_{k=1}^K \pi_k \left(\hat{\mu}_Y^k - \beta_0 - \beta_{-0}^\top \hat{\mu}_X^k\right)\right)$ converges in distribution to a centered Gaussian distribution. As far as the second component is concerned, it can be noted that $\hat{\Gamma}_X^k$ converges in probability to Γ_X^k and by the continuous mapping theorem, $\sqrt{\beta_{-0}^\top \hat{\Gamma}_X^k \beta_{-0} + \sigma_{\epsilon}^2} \rightarrow \sigma_{Y,k}$ in probability. It can also be noted that, under the moment condition $\mathbb{E}\left[\|\mathbf{X}\|^4|G=k\right] < \infty$, the central limit theorem gives that $\sqrt{n} \left(\hat{\Gamma}_X^k - \Gamma_X^k\right)$ converges in distribution to a centered Gaussian multivariate distribution, and we deduce with the Cramer-Wold device, the continuous mapping theorem that the second component of $\nabla \varphi_n$ multiplied by \sqrt{n} also in distribution to a centered Gaussian random vector. It is immediate to deduce that the same convergence result holds for the third component, which is to say that $\sqrt{n} \left(\sum_{k=1}^K \pi_k \left(1 - \frac{\hat{\sigma}_{Y,k}}{\sqrt{\beta_{-0}^\top \hat{\Gamma}_k^T \beta_{-0} + \sigma_{\epsilon}^2}}\right)\right)$ converges in distribution to a centered Gaussian random vector. It is is to say that $\sqrt{n} \left(\sum_{k=1}^K \pi_k \left(1 - \frac{\hat{\sigma}_{Y,k}}{\sqrt{\beta_{-0}^\top \hat{\Gamma}_k^T \beta_{-0} + \sigma_{\epsilon}^2}}\right)\right)$

dom variable. We finally deduce, with the Cramer-Wold device, that (iii) is fulfilled.

To prove that (iv) also holds, consider the Hessian matrix of functional φ_n , evaluated at

$$(\boldsymbol{\beta}, \sigma_{\epsilon}^{2}):$$

$$\nabla_{00}\varphi_{n} = \begin{pmatrix} 2 & 2\left(\sum_{k=1}^{K} \pi_{k} \widehat{\boldsymbol{\mu}}_{X}^{k}\right)^{\top} & 0 \\ 2\sum_{k=1}^{K} \pi_{k} \widehat{\boldsymbol{\mu}}_{X}^{k} & \widehat{\mathbf{H}}(\boldsymbol{\beta}_{-0}) & \sum_{k=1}^{K} \pi_{k} \widehat{\sigma}_{Y,k} \left(\boldsymbol{\beta}_{-0}^{\top} \widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\beta}_{-0} + \sigma_{\epsilon}^{2}\right)^{-3/2} \widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\beta}_{-0} \\ 0 & \sum_{k=1}^{K} \pi_{k} \widehat{\sigma}_{Y,k} \left(\boldsymbol{\beta}_{-0}^{\top} \widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\beta}_{-0} + \sigma_{\epsilon}^{2}\right)^{-3/2} \left(\widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\beta}_{-0}\right)^{\top} & \frac{1}{2} \sum_{k=1}^{K} \pi_{k} \widehat{\sigma}_{Y,k} \left(\boldsymbol{\beta}_{-0}^{\top} \widehat{\boldsymbol{\Gamma}}_{X}^{k} \boldsymbol{\beta}_{-0} + \sigma_{\epsilon}^{2}\right)^{-3/2} \end{pmatrix}$$

where

$$\begin{split} \widehat{\mathbf{H}}(\boldsymbol{\beta}_{-0}) = & 2\sum_{k=1}^{K} \pi_{k} \left[\widehat{\sigma}_{Y,k} \left(\boldsymbol{\beta}_{-0}^{\top} \widehat{\mathbf{\Gamma}}_{X}^{k} \boldsymbol{\beta}_{-0} + \sigma_{\epsilon}^{2} \right)^{-3/2} \left[\left(\widehat{\mathbf{\Gamma}}_{X}^{k} \boldsymbol{\beta}_{-0} \right) \left(\widehat{\mathbf{\Gamma}}_{X}^{k} \boldsymbol{\beta}_{-0} \right)^{\top} - \left(\boldsymbol{\beta}_{-0}^{\top} \widehat{\mathbf{\Gamma}}_{X}^{k} \boldsymbol{\beta}_{-0} + \sigma_{\epsilon}^{2} \right) \widehat{\mathbf{\Gamma}}_{X}^{k} \right] \\ &+ \widehat{\boldsymbol{\mu}}_{X}^{k} \left(\widehat{\boldsymbol{\mu}}_{X}^{k} \right)^{\top} + \widehat{\mathbf{\Gamma}}_{X}^{k} \bigg]. \end{split}$$

By similar arguments as those used to show that $\varphi_n(\beta, \sigma_{\epsilon}^2)$ converges in probability to $\varphi(\beta, \sigma_{\epsilon}^2)$, we deduce that $\nabla_{00}\varphi_n$ converges in probability to some matrix $\mathbf{H}(\beta, \sigma_{\epsilon}^2)$, defined as follows

$$\mathbf{H}(\boldsymbol{\beta}, \sigma_{\epsilon}^{2}) = \begin{pmatrix} 2 & 2\left(\sum_{k=1}^{K} \pi_{k} \boldsymbol{\mu}_{X}^{k}\right)^{\top} & 0 \\ 2\sum_{k=1}^{K} \pi_{k} \boldsymbol{\mu}_{X}^{k} & \mathbf{H}(\boldsymbol{\beta}_{-0}) & \sum_{k=1}^{K} \pi_{k} \sigma_{Y,k} \left(\boldsymbol{\beta}_{-0}^{\top} \boldsymbol{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} + \sigma_{\epsilon}^{2}\right)^{-3/2} \boldsymbol{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \\ 0 & \sum_{k=1}^{K} \pi_{k} \sigma_{Y,k} \left(\boldsymbol{\beta}_{-0}^{\top} \boldsymbol{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} + \sigma_{\epsilon}^{2}\right)^{-3/2} \left(\boldsymbol{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0}\right)^{\top} & \frac{1}{2} \sum_{k=1}^{K} \pi_{k} \sigma_{Y,k} \left(\boldsymbol{\beta}_{-0}^{\top} \boldsymbol{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} + \sigma_{\epsilon}^{2}\right)^{-3/2} \boldsymbol{\rho}_{X}^{k} \boldsymbol{\beta}_{-0} \\ \boldsymbol{\rho}_{X}^{k} \boldsymbol{\rho$$

where

$$\begin{split} \mathbf{H}(\boldsymbol{\beta}_{-0}) = & 2\sum_{k=1}^{K} \pi_{k} \left(\sigma_{Y,k} \left(\boldsymbol{\beta}_{-0}^{\top} \boldsymbol{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} + \sigma_{\epsilon}^{2} \right)^{-3/2} \left[\left(\boldsymbol{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \right) \left(\boldsymbol{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \right)^{\top} - \left(\boldsymbol{\beta}_{-0}^{\top} \boldsymbol{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} + \sigma_{\epsilon}^{2} \right) \boldsymbol{\Gamma}_{X}^{k} \right] \\ & + \boldsymbol{\mu}_{X}^{k} \left(\boldsymbol{\mu}_{X}^{k} \right)^{\top} + \boldsymbol{\Gamma}_{X}^{k} \bigg). \end{split}$$

We now must check that $\mathbf{H}(\boldsymbol{\beta}, \sigma_{\epsilon}^2)$ is a positive definite matrix. For that we show that at the minimizer value $(\boldsymbol{\beta}, \sigma_{\epsilon}^2)$ its determinant is strictly positive. We first note that $\sigma_{Y,k} = (\boldsymbol{\beta}_{-0}^{\top} \boldsymbol{\Gamma}_X^k \boldsymbol{\beta}_{-0} + \sigma_{\epsilon}^2)^{1/2}$ so that $\sigma_{Y,k} (\boldsymbol{\beta}_{-0}^{\top} \boldsymbol{\Gamma}_X^k \boldsymbol{\beta}_{-0} + \sigma_{\epsilon}^2)^{-3/2} = \frac{1}{\sigma_{Y,k}^2}$ and $\mathbf{H}(\boldsymbol{\beta}_{-0})$ can be written in a simpler form,

$$\mathbf{H}(\boldsymbol{\beta}_{-0}) = 2\sum_{k=1}^{K} \pi_{k} \left[\boldsymbol{\mu}_{X}^{k} \left(\boldsymbol{\mu}_{X}^{k} \right)^{\top} + \frac{1}{\sigma_{Y,k}^{2}} \boldsymbol{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \left(\boldsymbol{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \right)^{\top} \right],$$
(B.6)

which is a definite positive matrix under hypothesis H_1 . Using a block matrix determinant formula, we have

$$\left|\mathbf{H}(\boldsymbol{\beta}, \sigma_{\epsilon}^{2})\right| = \begin{vmatrix} 2 & 0 \\ 0 & \frac{1}{2} \sum_{k=1}^{K} \frac{\pi_{k}}{\sigma_{Y,k}^{2}} \end{vmatrix} \left| \mathbf{H}(\boldsymbol{\beta}_{-0}) - \mathbf{C} \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{2}{\sum_{k} \frac{\pi_{k}}{\sigma_{Y,k}^{2}}} \end{pmatrix} \mathbf{C}^{\top} \right|$$
(B.7)

where $\mathbf{C} = \left(2\sum_{k=1}^{K} \pi_k \boldsymbol{\mu}_X^k \quad \sum_{k=1}^{K} \frac{\pi_k}{\sigma_{Y,k}^2} \boldsymbol{\Gamma}_X^k \boldsymbol{\beta}_{-0}\right)$ and it only has to be verified that the second

determinant at the righthand side of (B.7) is strictly positive. We now have to show that

$$\mathbf{H}(\boldsymbol{\beta}_{-0}) - \mathbf{C} \begin{pmatrix} \frac{1}{2} & 0\\ 0 & \frac{2}{\sum_{k} \frac{\pi_{k}}{\sigma_{Y,k}^{2}}} \end{pmatrix} \mathbf{C}^{\top} = 2 \sum_{k=1}^{K} \pi_{k} \boldsymbol{\mu}_{X}^{k} \left(\boldsymbol{\mu}_{X}^{k} \right)^{\top} - 2 \left(\sum_{k=1}^{K} \pi_{k} \boldsymbol{\mu}_{X}^{k} \right) \left(\sum_{k=1}^{K} \pi_{k} \boldsymbol{\mu}_{X}^{k} \right)^{\top} + 2 \sum_{k=1}^{K} \frac{\pi_{k}}{\sigma_{Y,k}^{2}} \mathbf{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \left(\mathbf{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \right)^{\top} - \frac{2}{\sum_{k} \frac{\pi_{k}}{\sigma_{Y,k}^{2}}} \left(\sum_{k=1}^{K} \frac{\pi_{k}}{\sigma_{Y,k}^{2}} \mathbf{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \right) \left(\sum_{k=1}^{K} \frac{\pi_{k}}{\sigma_{Y,k}^{2}} \mathbf{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \right)^{\top}$$
(B.8)

is a positive matrix. We can remark that by Cauchy Schwarz inequality, for $\mathbf{u} \in \mathbb{R}^d$,

$$\mathbf{u}^{\top} \left(\sum_{k=1}^{K} \pi_k \boldsymbol{\mu}_X^k \right) \left(\sum_{k=1}^{K} \pi_k \boldsymbol{\mu}_X^k \right)^{\top} \mathbf{u} = \left(\sum_{k=1}^{K} \pi_k \mathbf{u}^{\top} \boldsymbol{\mu}_X^k \right)^2 \le \sum_{k=1}^{K} \pi_k \left(\mathbf{u}^{\top} \boldsymbol{\mu}_X^k \right)^2$$
$$= \mathbf{u}^{\top} \left(\sum_{k=1}^{K} \pi_k \boldsymbol{\mu}_X^k \left(\boldsymbol{\mu}_X^k \right)^{\top} \right) \mathbf{u}$$

using the fact that $\sum_k (\sqrt{\pi_k})^2 = 1$. It can be noted that if $\mathbf{u} \neq 0$, previous inequality is strict unless $\mathbf{u}^\top \boldsymbol{\mu}_X^1 = \cdots = \mathbf{u}^\top \boldsymbol{\mu}_X^K$, which can not happen under hypothesis \mathbf{H}_1 . The second part at the righthand side of (B.8) is handled the same way. We have

$$\mathbf{u}^{\top} \left(\sum_{k=1}^{K} \frac{\pi_{k}}{\sigma_{Y,k}^{2}} \mathbf{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \right) \left(\sum_{k=1}^{K} \frac{\pi_{k}}{\sigma_{Y,k}^{2}} \mathbf{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \right)^{\top} \mathbf{u} = \left(\mathbf{u}^{\top} \left(\sum_{k=1}^{K} \frac{\pi_{k}}{\sigma_{Y,k}^{2}} \mathbf{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \right) \right)^{2}$$
$$\leq \sum_{k=1}^{K} \sqrt{\frac{\pi_{k}}{\sigma_{Y,k}^{2}}}^{2} \sum_{k=1}^{K} \left(\sqrt{\frac{\pi_{k}}{\sigma_{Y,k}^{2}}}^{2} \mathbf{u}^{\top} \mathbf{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \right)^{2}$$
$$= \sum_{k=1}^{K} \frac{\pi_{k}}{\sigma_{Y,k}^{2}} \sum_{k=1}^{K} \frac{\pi_{k}}{\sigma_{Y,k}^{2}} \sum_{k=1}^{K} \frac{\pi_{k}}{\sigma_{Y,k}^{2}} \mathbf{u}^{\top} \mathbf{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \left(\mathbf{\Gamma}_{X}^{k} \boldsymbol{\beta}_{-0} \right)^{\top} \mathbf{u},$$

and consequently the determinant of $\mathbf{H}(\boldsymbol{\beta}, \sigma_{\epsilon}^2)$ is strictly positive.

To finish the proof, it remains to check that in a neighborhood \mathcal{N} of $(\beta, \sigma_{\epsilon}^2)$, we have

$$\sup_{(\boldsymbol{\gamma},\sigma_{\boldsymbol{\gamma}}^2)\in\mathcal{N}} \|\nabla_{00}\varphi_n(\boldsymbol{\gamma},\sigma_{\boldsymbol{\gamma}}^2) - \mathbf{H}(\boldsymbol{\gamma},\sigma_{\boldsymbol{\gamma}}^2)\| \to 0 \text{ in probability.}$$

This is a direct consequence of the continuous mapping theorem, which gives us that for all $(\gamma, \sigma_{\gamma}^2) \in \mathcal{N}$, $\|\nabla_{00}\varphi_n(\gamma, \sigma_{\gamma}^2) - \mathbf{H}(\gamma, \sigma_{\gamma}^2)\| \to 0$ in probability, and the fact that third order partial derivatives of $\varphi_n(\gamma, \sigma_{\gamma}^2)$ are bounded in probability for $(\gamma, \sigma_{\gamma}^2)$ so that Theorem A.2 can apply.

Proof. of Lemma 5.1 Note that

$$\widehat{\beta}_1 = g(\widehat{\mu}_X^1, \dots, \widehat{\mu}_X^K, \widehat{\mu}_Y^1, \dots, \widehat{\mu}_Y^K),$$

with $g : \mathbb{R}^{K+K} \to \mathbb{R}$ defined as follows,

$$g(\mu_X^1, \dots, \mu_X^K, \mu_Y^1, \dots, \mu_Y^K) = \frac{\sum_{k=1}^K w_k \mu_X^k \mu_Y^k - \mu_{X,w} \mu_{Y,w}}{\sum_{k=1}^K w_k (\mu_X^k)^2 - \left(\sum_{k=1}^K w_k \mu_X^k\right)^2},$$
$$= \frac{\text{Cov}_w(X, Y)}{\text{Var}_w(X)},$$
(B.9)

with the notations $\mu_{X,w} = \sum_{k=1}^{K} w_k \mu_X^k$, $\mu_{Y,w} = \sum_{j=1}^{K} w_j \mu_Y^j$, $\operatorname{Cov}_w(X,Y) = \sum_{k=1}^{K} w_k \mu_X^k \mu_Y^k - \mu_{X,w} \mu_{Y,w}$ and $\operatorname{Var}_w(X) = \sum_{k=1}^{K} w_k (\mu_X^k)^2 - (\mu_{X,w})^2$. The gradient ∇g of g, evaluated at point $(\mu_X^1, \ldots, \mu_X^K, \mu_Y^1, \ldots, \mu_Y^K)$, is equal to

$$\nabla g = \begin{pmatrix} \frac{w_1(\mu_Y^1 - \mu_{Y,w})}{\operatorname{Var}_w(X)} - \frac{2w_1(\mu_X^1 - \mu_{X,w})\operatorname{Cov}_w(X,Y)}{(\operatorname{Var}_w(X))^2} \\ \vdots \\ \frac{w_K(\mu_Y^K - \mu_{Y,w})}{\operatorname{Var}_w(X)} - \frac{2w_K(\mu_X^K - \mu_{X,w})\operatorname{Cov}_w(X,Y)}{(\operatorname{Var}_w(X))^2} \\ \frac{w_1(\mu_X^1 - \mu_{X,w})}{\operatorname{Var}_w(X)} \\ \vdots \\ \frac{w_K(\mu_X^K - \mu_{X,w})}{\operatorname{Var}_w(X)} \end{pmatrix}$$

As in the proof of Proposition 4.1, we get that $\sqrt{n} \left(\widehat{\beta}_1 - \beta_1 \right) \rightsquigarrow \mathcal{N}(0, \sigma_{\beta_1}^2)$ with $\sigma_{\beta_1}^2 = \nabla g^T \Gamma_\mu \nabla g$, so that

$$\sigma_{\beta_{1}}^{2} = \frac{1}{(\operatorname{Var}_{w}(X))^{2}} \sum_{k=1}^{K} w_{k}^{2} \left[\sigma_{X,k}^{2} \left(\mu_{Y}^{k} - \mu_{Y,w} - 2\beta_{1} \left(\mu_{X}^{k} - \mu_{X,w} \right) \right)^{2} + \sigma_{Y,k}^{2} \left(\mu_{X}^{k} - \mu_{X,w} \right)^{2} \right]$$

$$= \frac{1}{(\operatorname{Var}_{w}(X))^{2}} \sum_{k=1}^{K} w_{k}^{2} \left[\sigma_{X,k}^{2} \left(-\beta_{1} \left(\mu_{X}^{k} - \mu_{X,w} \right) \right)^{2} + \sigma_{Y,k}^{2} \left(\mu_{X}^{k} - \mu_{X,w} \right)^{2} \right]$$

$$= \frac{1}{(\operatorname{Var}_{w}(X))^{2}} \sum_{k=1}^{K} w_{k}^{2} \left(\mu_{X}^{k} - \mu_{X,w} \right)^{2} \left(\beta_{1}^{2} \sigma_{X,k}^{2} + \sigma_{Y,k}^{2} \right)$$
(B.10)

remarking that $\beta_1 = \operatorname{Cov}_w(X, Y) / \operatorname{Var}_w(X, Y)$, $\beta_0 = \mu_{Y,w} - \beta_1 \mu_{X,w}$ as well as $\beta_0 = \mu_Y^k - \beta_1 \mu_X^k$.

Proof. of Proposition 6.1

The fact that the bootstrap estimator $\beta^{M,*}$ is strongly consistent for β is a direct consequence of Theorem 3.1 in Shao and Tu (1995), noting that

$$\widehat{\boldsymbol{\beta}}^M = g(\widehat{\boldsymbol{\mu}}_X^1, \dots, \widehat{\boldsymbol{\mu}}_X^K, \widehat{\mu}_Y^1, \dots, \widehat{\mu}_Y^K)$$

is a continuously differentiable function of means at $(\boldsymbol{\mu}_X^1, \dots, \boldsymbol{\mu}_X^K, \boldsymbol{\mu}_Y^1, \dots, \boldsymbol{\mu}_Y^K)$. The fact that confidence sets based on the percentile approach are consistent is proved by checking the assumptions in Theorem 4.1 (iii) Shao and Tu (1995), namely the bootstrap estimator $\boldsymbol{\beta}^{M,*}$ is consistent, $\hat{\boldsymbol{\beta}}^M$ is consistent (Lemma 4.1), with asymptotic Gaussian distribution (Proposition 4.1).

Proof. of Proposition 6.2

We denote by $\theta_0 = (\beta, \sigma_{\epsilon}^{2,W})$ the vector of true parameters, by $\hat{\theta} = (\beta^W, \sigma_{\epsilon}^2)$ the sequence of minimum Wasserstein distance estimators and by $\theta^* = (\beta^{W,*}, \sigma_{\epsilon}^{2,W,*})$ bootstrap estimators of θ_0 . The vector of parameters θ^* is the minimizer of functional φ_n^* defined as follows,

$$\varphi_n^*(\boldsymbol{\gamma}, \sigma^2) = \sum_{k=1}^K \pi_k \left[(\mu_Y^{k,*} - \gamma_0 - \boldsymbol{\gamma}_{-0}^\top \boldsymbol{\mu}_X^{k,*})^2 + \left(\sigma_{Y,k}^* - \sqrt{\boldsymbol{\gamma}_{-0}^\top \boldsymbol{\Gamma}_X^{k,*} \boldsymbol{\gamma}_{-0} + \sigma^2} \right)^2 \right].$$
(B.11)

We first show with arguments similar to those employed in the proof of Lemma 4.2, that θ^* is a consistent estimator for θ_0 , based on the fact that φ_n^* is a smooth function converging to φ and the sample mean theorem for bootstrap (see for example Theorem 23.4 in van der Vaart (1998)). Indeed, we first recall that for all $(\gamma, \sigma_\gamma^2) \in \Theta$, $\varphi_n(\gamma, \sigma_\gamma^2) \to \varphi(\gamma, \sigma_\gamma^2)$ in probability, when n_{\min} tends to infinity and

$$\left|\varphi_{n}^{*}(\boldsymbol{\gamma},\sigma_{\gamma}^{2})-\varphi(\boldsymbol{\gamma},\sigma_{\gamma}^{2})\right| \leq \left|\varphi_{n}^{*}(\boldsymbol{\gamma},\sigma_{\gamma}^{2})-\varphi_{n}(\boldsymbol{\gamma},\sigma_{\gamma}^{2})\right|+\left|\varphi_{n}(\boldsymbol{\gamma},\sigma_{\gamma}^{2})-\varphi(\boldsymbol{\gamma},\sigma_{\gamma}^{2})\right|.$$
(B.12)

Since the bootstrap means converge to the empirical ones we deduce with the continuous mapping theorem that $\varphi_n^*(\gamma, \sigma_\gamma^2) \rightarrow \varphi_n(\gamma, \sigma_\gamma^2)$ in probability, when n_{\min} tends to infinity, so that $\varphi_n^*(\gamma, \sigma_\gamma^2) \rightarrow \varphi(\gamma, \sigma_\gamma^2)$. We also have, as in (B.3), where empirical means are replaced by bootstrap means,

$$\left|\varphi_{n}^{*}(\boldsymbol{\gamma},\sigma_{\gamma}^{2})-\varphi_{n}^{*}(\boldsymbol{\alpha},\sigma_{\alpha}^{2})\right| \leq \left(\left\|\boldsymbol{\alpha}-\boldsymbol{\gamma}\right\|+\left|\sigma_{\alpha}^{2}-\sigma_{\gamma}^{2}\right|\right)\sum_{k=1}^{K}\pi_{k}\left(B_{n}^{k,*}+A_{n}^{k,*}\right),\tag{B.13}$$

for any $(\alpha, \sigma_{\alpha}^2) \in \Theta$, with $\sum_{k=1}^{K} \pi_k \left(B_n^{k,*} + A_n^{k,*} \right) = O_p(1)$. As a result, we deduce from Lemma 4.2, inequality (B.12) and Lemma 2.9 in Newey and McFadden (1994) that

$$\sup_{(\boldsymbol{\gamma},\sigma_{\gamma}^2)\in\Theta}\left|\varphi_n^*(\boldsymbol{\gamma},\sigma_{\gamma}^2)-\varphi(\boldsymbol{\gamma},\sigma_{\gamma}^2)\right|\to 0 \quad \text{ in probability.}$$

We conclude that $\theta^* \to \theta_0$ in probability in view of Theorem 2.1 in Newey and McFadden (1994).

We now prove that $\sqrt{n} \left(\boldsymbol{\theta}^* - \widehat{\boldsymbol{\theta}} \right)$ and $\sqrt{n} \left(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0 \right)$ have the same asymptotic distribution. By definition of $\widehat{\boldsymbol{\theta}}$ and Taylor expansion we have

$$\nabla \varphi_n(\widehat{\boldsymbol{\theta}}) = \nabla \varphi_n(\boldsymbol{\theta}_0) + \nabla_{00} \varphi_n(\overline{\boldsymbol{\theta}}) \left(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0\right) = 0$$
(B.14)

where $\overline{\theta}$ belongs, componentwise, to the segment between θ_0 and $\widehat{\theta}$. We have a similar expansion for boostrap estimators, as well as

$$\nabla \varphi_n^*(\boldsymbol{\theta}^*) = \nabla \varphi_n^*(\boldsymbol{\theta}_0) + \nabla_{00}^* \varphi_n(\overline{\boldsymbol{\theta}}^*) \left(\boldsymbol{\theta}^* - \boldsymbol{\theta}_0\right) = 0$$
(B.15)

where $\overline{\theta}^*$ belongs, componentwise, to the segment between θ_0 and θ^* . Combining (B.14) and

(B.15), we deduce

$$\boldsymbol{\theta}^{*} - \widehat{\boldsymbol{\theta}} = \left(\nabla_{00}^{*}\varphi_{n}(\overline{\boldsymbol{\theta}}^{*})\right)^{-1} \nabla \varphi_{n}^{*}(\boldsymbol{\theta}_{0}) - \left(\nabla_{00}\varphi_{n}(\overline{\boldsymbol{\theta}})\right)^{-1} \nabla \varphi_{n}(\boldsymbol{\theta}_{0}) \\ = \left(\left(\nabla_{00}^{*}\varphi_{n}(\overline{\boldsymbol{\theta}}^{*})\right)^{-1} - \left(\nabla_{00}\varphi_{n}(\overline{\boldsymbol{\theta}})\right)^{-1}\right) \nabla \varphi_{n}^{*}(\boldsymbol{\theta}_{0}) + \left(\nabla_{00}\varphi_{n}(\overline{\boldsymbol{\theta}})\right)^{-1} \left(\nabla \varphi_{n}^{*}(\boldsymbol{\theta}_{0}) - \nabla \varphi_{n}(\boldsymbol{\theta}_{0})\right)$$
(B.16)

Noticing that $\nabla_{00}^* \varphi_n(\overline{\theta}^*)$ and $\nabla_{00} \varphi_n(\overline{\theta})$ both tend in probability to the same limit $\mathbf{H}(\boldsymbol{\beta}, \sigma_{\epsilon}^2)$ and we have, with similar arguments as those used in the proof of Proposition 4.2, that $\nabla \varphi_n^*(\boldsymbol{\theta}_0)$ is $O_p(n^{-1/2})$, it can be deduce that

$$\boldsymbol{\theta}^* - \widehat{\boldsymbol{\theta}} = \left(\nabla_{00}\varphi_n(\overline{\boldsymbol{\theta}})\right)^{-1} \left(\nabla\varphi_n^*(\boldsymbol{\theta}_0) - \nabla\varphi_n(\boldsymbol{\theta}_0)\right) + o_P(n^{-1/2}). \tag{B.17}$$

Using arguments similar to those employed in the expansion of $\nabla \varphi_n$ in the proof of Proposition 4.2, we make appear the difference between bootstrap means and empirical means or a differentiable functional of these quantities:

$$\nabla \varphi_{n}^{*}(\boldsymbol{\theta}_{0}) - \nabla \varphi_{n}(\boldsymbol{\theta}_{0}) = \begin{pmatrix} 2\sum_{k=1}^{K} \pi_{k} \left(\left(\widehat{\mu}_{Y}^{k} - \mu_{Y}^{k,*} \right) - \beta_{0} - \beta_{-0}^{\top} \left(\widehat{\mu}_{X} - \mu_{X}^{k,*} \right) \right) \\ 2\sum_{k=1}^{K} \pi_{k} \left[\left(\widehat{\mu}_{Y}^{k} - \mu_{Y}^{k} - \beta_{0} - \beta_{-0}^{\top} \widehat{\mu}_{X}^{k} \right) \widehat{\mu}_{X}^{k} + \left(\frac{\widehat{\sigma}_{Y,k}}{\sqrt{\beta_{-0}^{\top} \widehat{\Gamma}_{X}^{k} \beta_{-0} + \sigma_{\epsilon}^{2}}} - 1 \right) \widehat{\Gamma}_{X}^{k} \beta_{-0} \right] \\ -2\sum_{k=1}^{K} \pi_{k} \left[\left(\mu_{Y}^{k,*} - \mu_{Y}^{k} - \beta_{0} - \beta_{-0}^{\top} \mu_{X}^{k,*} \right) \mu_{X}^{k,*} + \left(\frac{\sigma_{Y,k}}{\sqrt{\beta_{-0}^{\top} \Gamma_{X}^{k,*} \beta_{-0} + \sigma_{\epsilon}^{2}}} - 1 \right) \Gamma_{X}^{k,*} \beta_{-0} \right] \right) \\ \sum_{k=1}^{K} \left(\frac{\widehat{\sigma}_{Y,k}}{\sqrt{\beta_{-0}^{\top} \Gamma_{X}^{k,*} \beta_{-0} + \sigma_{\epsilon}^{2}}} - \frac{\sigma_{Y,k}^{*}}{\sqrt{\beta_{-0}^{\top} \Gamma_{X}^{k,*} \beta_{-0} + \sigma_{\epsilon}^{2}}} \right) \end{pmatrix}$$
(B.18)

which satisfies the central limit theorem for bootstrap means, or the Delta method for bootstrap estimators (see the Appendix as well as Theorem 23.4 and Theorem 23.5 in van der Vaart (1998)). Consequently, $\nabla \varphi_n^*(\theta_0) - \nabla \varphi_n(\theta_0)$ and $\nabla \varphi_n(\theta_0) - \nabla \varphi(\theta_0)$ have the same asymptotic distribution. By Slustky's theorem, the asymptotic distribution of $\sqrt{n} \left(\theta^* - \hat{\theta} \right)$ is thus the same as the asymptotic distribution of $\mathbf{H}(\beta, \sigma_{\epsilon}^2) \sqrt{n} \nabla \varphi_n(\theta_0)$, and we can conclude that $\sqrt{n} \left(\theta^* - \hat{\theta} \right)$ and $\sqrt{n} \left(\hat{\theta} - \theta_0 \right)$ have also the same asymptotic Gaussian distribution.

figures
nentary
Supplen
U U

ps.=4	0.97	0.97	0.97	0.97
Nb. groul	0.95	0.94	0.94	0.94
nim.=10,	0.97	0.97	0.97	0.97
Nb. a	0.94	0.95	0.95	0.94
	mm (asymp)	mm (boot)	ot (boot)	mm (student)

3.91	3.88	3.45	6.97
10.97	10.83	10.43	18.68
2.40	2.28	2.18	4.34
6.70	6.42	6.36	11.58
mm (asymp)	mm (boot)	ot (boot)	mm (student)
	mm (asymp) 6.70 2.40 10.97 3.91	mm(asymp) 6.70 2.40 10.97 3.91 $mm(boot)$ 6.42 2.28 10.83 3.88	mm (asymp) 6.70 2.40 10.97 3.91 mm (boot) 6.42 2.28 10.83 3.88 ot (boot) 6.36 2.18 10.43 3.45

	_	2	0		=2
ps.=4	0.4]	0.42	0.42	0.2]	sigma2x- rho=1.
Nb. grou	0.08	0.10	0.10	0.08	sigma2x=2 rho=1.01
nim.=10,	0.91	0.94	0.94	0.44	sigma2x=0.75 rho=1.1
Nb. a	0.19	0.21	0.21	0.12	sigma2x=0.75 rho=1.01
	mm (asymp)	mm (boot)	ot (boot)	mm (student)	

parameter combination. The method labels on the left: "mm (asymp)" stands for the method of moments with asymptotic confidence intervals, "mm (boot)" for the method of moments with bootstrap, "ot (boot)" for the optimal transport method with bootstrap, and "mm (student)" for the naive linear regression on means approach based on Student's distribution. Figure (C.1) a) Coverage rates, b) amplitudes, and c) powers of the confidence intervals for the estimators of β_1 obtained from N_{sim} simulations for each 5

	0.96	0.94	06'0	0.88			17.5	15.0 12.5	7.5	2.5			1.0	0.8	0.4	0.2		
.=10	0.89	0.88	0.89	0.94	sigma2x=2 rho=1.1	s.=10	0.58	0.56	0.56	0.64	sigma2x=2 rho=1.1	.=10	1.00	1.00	1.00	1.00	sigma2x=2 rho=1.1	
lb. groups	06.0	0.90	0.89	0.91	sigma2x=2 rho=1.01	lb. groups	1.59	1.56	1.55	1.79	sigma2x=2 rho=1.01	lb. groups	1.00	1.00	1.00	0.98	sigma2x=2 rho=1.01	
m.=30, N	0.91	0.88	0.87	0.94	igma2x=0.75 rho=1.1	m.=30, N	0.35	0.35	0.35	0.39	igma2x=0.75 rho=1.1	m.=30, N	1.00	1.00	1.00	1.00	igma2x=0.75 rho=1.1	
Nb. ani	06.0	0.89	0.89	0.91	sigma2x=0.75 s rho=1.01	Nb. an	0.98	0.95	0.95	1.10	sigma2x=0.75 s rho=1.01	Nb. ani	1.00	1.00	1.00	1.00	sigma2x=0.75 s rho=1.01	
					N						0						0	
os.=4	0.94	0.93	0.93	0.93	sigma2x=2 rho=1.1	os.=4	2.34	2.33	2.23	4.45	sigma2x=2 rho=1.1	os.=4	0.95	0.96	0.96	0.39	sigma2x=2 rho=1.1	
Nb. group	0.93	0.92	0.92	0.91	sigma2x=2 rho=1.01	Nb. grouf	6.53	6.51	6.48	12.01	sigma2x=2 rho=1.01	Nb. grouf	0.24	0.25	0.25	0.14	sigma2x=2 rho=1.01	
nim.=30,	0.93	0.93	0.91	0.91	sigma2x=0.75 rho=1.1	nim.=30,	1.43	1.40	1.38	2.74	sigma2x=0.75 rho=1.1	nim.=30,	1.00	1.00	1.00	0.78	sigma2x=0.75 rho=1.1	
Nb. ai	0.92	0.92	0.92	0.92	sigma2x=0.75 rho=1.01	Nb. aı	3.99	3.90	3.90	7.41	sigma2x=0.75 rho=1.01	Nb. aı	0.46	0.50	0.50	0.20	sigma2x=0.75 rho=1.01	
																		<u>.</u>
_					_	<i>(a</i>					_	9					_	5
s.=10	0.93	0.92	0.90	0.93	sigma2x=2 rho=1.1	(<i>a</i> s.=10	0.98	0.93	0.91	1.11	sigma2x=2 rho=1.1	(<i>b</i> s.=10	1.00	1.00	1.00	1.00	sigma2x=2 rho=1.1	5
Vb. groups.=10	0.92 0.93	0.91 0.92	0.91 0.90	0.93 0.93	sigma2x=2 sigma2x=2 rho=1.01 rho=1.1	(a Vb. groups.=10	2.72 0.98	2.58 0.93	2.57 0.91	3.10 1.11	sigma2x=2 sigma2x=2 rho=1.01 rho=1.1	(b Vb. group <u>s.=</u> 10	0.80 1.00	0.86 1.00	0.86 1.00	0.71 1.00	sigma2x=2 sigma2x=2 rho=1.01 rho=1.1	5
im.=10, Nb. groups.=10	0.92 0.92 0.93	0.91 0.91 0.92	0.91 0.91 0.90	0.93 0.93 0.93	igma2x=0.75 sigma2x=2 sigma2x=2 rho=1.1 rho=1.01 rho=1.1	(<i>a</i> im.=10, Nb. groups.=10	0.61 2.72 0.98	0.57 2.58 0.93	0.57 2.57 0.91	0.69 3.10 1.11	igma2x=0.75 sigma2x=2 sigma2x=2 rho=1.1 rho=1.01 rho=1.1	(<i>b</i> im.=10, Nb. groups.=10	1.00 0.80 1.00	1.00 0.86 1.00	1.00 0.86 1.00	1.00 0.71 1.00	igma2x=0.75 sigma2x=2 sigma2x=2 rho=1.1 rho=1.01 rho=1.1))
Nb. anim.=10, Nb. groups.=10	0.92 0.92 0.92 0.93	0.89 0.91 0.91 0.92	0.89 0.91 0.91 0.90	0.92 0.93 0.93 0.93	igma2x=0.75 sigma2x=0.75 sigma2x=2 sigma2x=2 rho=1.01 rho=1.1 rho=1.01 rho=1.1	(a Nb. anim.=10, Nb. groups.=10	1.68 0.61 2.72 0.98	1.58 0.57 2.58 0.93	1.58 0.57 2.57 0.91	1.91 0.69 3.10 1.11	igma2x=0.75 sigma2x=0.75 sigma2x=2 sigma2x=2 tho=1.01 tho=1.1 tho=1.1	(<i>b</i>) Nb. anim.=10, Nb. groups.=10	1.00 1.00 0.80 1.00	1.00 1.00 0.86 1.00	1.00 1.00 0.86 1.00	0.97 1.00 0.71 1.00	$\label{eq:relation} \begin{array}{llllllllllllllllllllllllllllllllllll$	<i>J</i>)
Nb. anim.=10, Nb. groups.=10	0.92 0.92 0.93 0.93	0.89 0.91 0.91 0.92	0.89 0.91 0.91 0.90	0.92 0.93 0.93 0.93	sigma2x=0.75 sigma2x=0.75 sigma2x=2 sigma2x=2 tho=1.01 rho=1.1 rho=1.01 rho=1.1	(a Nb. anim.=10, Nb. groups.=10	1.68 0.61 2.72 0.98	1.58 0.57 2.58 0.93	1.58 0.57 2.57 0.91	1.91 0.69 3.10 1.11	$\begin{array}{llllllllllllllllllllllllllllllllllll$	(b Nb. anim.=10, <mark>Nb. group</mark> s.=10	1.00 1.00 0.80 1.00	1.00 1.00 0.86 1.00	1.00 1.00 0.86 1.00	0.97 1.00 0.71 1.00	sigma2x=0.75 sigma2x=0.75 sigma2x=2 sigma2x=2 tho=1.01 tho=1.1 tho=1.01 tho=1.1	J)
os.=4 Nb. anim.=10, Nb. groups.=10	0.97 0.92 0.92 0.93 0.93	0.97 0.89 0.91 0.91 0.92	0.97 0.89 0.91 0.91 0.90	0.97 0.92 0.93 0.93 0.93	sigma2x=2 sigma2x=0.75 sigma2x=2.75 sigma2x=2 sigma2x=2 tho=1.1 tho=1.01 tho=1.1 tho=1.01 tho=1.1	(a Nb. anim.=10, Nb. groups.=10	3.91 1.68 0.61 2.72 0.98	3.88 1.58 0.57 2.58 0.93	3.45 1.58 0.57 2.57 0.91	6.97 1.91 0.69 3.10 1.11	sigma2x=2 sigma2x=0.75 sigma2x=0.75 sigma2x=2 sigma2x=2 tho=1.1 tho=1.01 tho=1.1 tho=1.1	(<i>b</i> ss.=4 Nb. anim.=10, Nb. groups.=10	0.41 1.00 1.00 0.80 1.00	0.42 1.00 1.00 0.86 1.00	0.42 1.00 1.00 0.86 1.00	0.21 0.97 1.00 0.71 1.00	sigma2x=2 sigma2x=0.75 sigma2x=0.75 sigma2x=2 sigma2x=2 tho=1.1 tho=1.01 tho=1.1 tho=1.01 tho=1.1	() (
Nb. groups.=4 Nb. anim.=10, Nb. groups.=10	0.95 0.97 0.92 0.92 0.93 0.93	0.94 0.97 0.89 0.91 0.91 0.92	0.94 0.97 0.89 0.91 0.91 0.90	0.94 0.97 0.92 0.93 0.93 0.93	sigma2x=2 sigma2x=0.75 sigma2x=2.5 sigma2x	(a Nb. groups.=4 Nb. anim.=10, Nb. groups.=10	10.97 3.91 1.68 0.61 2.72 0.98	10.83 3.88 1.58 0.57 2.58 0.93	10.43 3.45 1.58 0.57 2.57 0.91	18.68 6.97 1.91 0.69 3.10 1.11	sigma2x=2 sigma2x=0.75 sigma2x=0.75 sigma2x=0.75 sigma2x=2 sigma2x=2 tho=1.01 tho=1.1 tho=1.01 tho=1.1	(b Nb. groups.=4 Nb. anim.=10, Nb. groups.=10	0.08 0.41 1.00 1.00 0.80 1.00	0.10 0.42 1.00 1.00 0.86 1.00	0.10 0.42 1.00 1.00 0.86 1.00	0.08 0.21 0.97 1.00 0.71 1.00	sigma2x=2 sigma2x=0.75 sigma2x=0.75 sigma2x=0.75 sigma2x=2 sigma2x=2 tho=1.01 tho=1.	()
nim.=10, Nb. groups.=4 Nb. anim.=10, Nb. groups.=10	0.97 0.95 0.97 0.92 0.92 0.92 0.93	0.97 0.94 0.97 0.89 0.91 0.91 0.92	0.97 0.94 0.97 0.89 0.91 0.91 0.90	0.97 0.94 0.97 0.92 0.93 0.93 0.93	sigma2x=0.75 sigma2x=2 sigma2x=2.75 sigma2x=0.75 sigma2x=2 sigma2x=2.75 sigmax=2.75 sigmax=2.75 sigmax=2.75 s	(a) nim.=10, Nb. groups.=4 Nb. anim.=10, Nb. groups.=10	2.40 10.97 3.91 1.68 0.61 2.72 0.98	2.28 10.83 3.88 1.58 0.57 2.58 0.93	2.18 10.43 3.45 1.58 0.57 2.57 0.91	4.34 18.68 6.97 1.91 0.69 3.10 1.11	sigma2x=0.75 sigma2x=2 sigma2x=0.75 sigma2x=0.75 sigma2x=0.75 sigma2x=0.75 sigma2x=2 rho=1.1 rho=1.01 rho=1.1 rho=1.1 rho=1.1	(b iim.=10, Nb. groups.=4 Nb. anim.=10, Nb. groups.=10	0.91 0.08 0.41 1.00 1.00 0.80 1.00	0.94 0.10 0.42 1.00 1.00 0.86 1.00	0.94 0.10 0.42 1.00 1.00 0.86 1.00	0.44 0.08 0.21 0.97 1.00 0.71 1.00	sigma2x=0.75 sigma2x=2 sigma2x=0.75 sigma2x=0.75 sigma2x=2 sigma2x=2 mo=1.1 mo=1.01 mo))
Nb. anim.=10, Nb. groups.=4 Nb. anim.=10, Nb. groups.=10	0.94 0.97 0.95 0.97 0.92 0.92 0.93	0.95 0.97 0.94 0.97 0.97 0.91 0.91 0.92	0.95 0.97 0.94 0.97 0.97 0.91 0.91 0.90	0.94 0.97 0.94 0.97 0.92 0.93 0.93 0.93	igma2x=0.75 sigma2x=2.75 sigma2x=2.75 sigma2x=0.75 sigma2x=0.75 sigma2x=2.75 sigmax=2.75	(a Nb. anim.=10, Nb. groups.=4	6.70 2.40 10.97 3.91 1.68 0.61 2.72 0.98	6.42 2.28 10.83 3.88 1.58 0.57 2.58 0.93	6.36 2.18 10.43 3.45 1.58 0.57 2.57 0.91	11.58 4.34 18.68 6.97 1.91 0.69 3.10 1.11	igma2x=0.75 sigma2x=2 sigma2x=2 sigma2x=0.75 sigma2x=0.75 sigma2x=0.75 sigma2x=0.75 sigma2x=2 sigma2x=2 tho=1.01 tho=1.0	(b Nb. anim.=10, Nb. groups.=4 Nb. anim.=10, Nb. groups.=10	0.19 0.91 0.08 0.41 1.00 1.00 0.80 1.00	0.21 0.94 0.10 0.42 1.00 1.00 0.86 1.00	0.21 0.94 0.10 0.42 1.00 1.00 0.86 1.00	0.12 0.44 0.08 0.21 0.97 1.00 0.71 1.00	sigma2x=0.75 sigma2x=2 sigma2x=2 sigma2x=0.75 sigma2x=0.75 sigma2x=2 sigma2x=2 rho=1.01 rho=1	