

TROPICAL TWISTED HURWITZ NUMBERS FOR ELLIPTIC CURVES

MARVIN ANAS HAHN AND HANNAH MARKWIG

ABSTRACT. Hurwitz numbers enumerate branched morphisms between Riemann surfaces. For a fixed elliptic target, Hurwitz numbers are intimately related to mirror symmetry following work of Dijkgraaf. In recent work of Chapuy and Dolega a new variant of Hurwitz numbers with fixed genus 0 target was introduced that includes maps between non-orientable surfaces. These numbers are called b -Hurwitz numbers and are polynomials in a parameter b which measures the non-orientability of the involved maps. An interpretation in terms of factorisations of b -Hurwitz numbers for $b = 1$, so-called twisted Hurwitz numbers, was found in work of Burman and Fesler. In previous work, the authors derived a tropical geometry interpretation of these numbers. In this paper, we introduce a natural generalisation of twisted Hurwitz numbers with elliptic targets within the framework of symmetric groups. We derive a tropical interpretation of these invariants, relate them to Feynman integrals and derive an expression as a matrix element of an operator in the bosonic Fock space.

1. INTRODUCTION

Hurwitz numbers count branched covers of Riemann surfaces with fixed numerical data. They originate from Hurwitz' original work in [Hur92] and have developed to important invariants in enumerative geometry. There are various equivalent definitions of Hurwitz numbers arising from different fields of mathematics. The one most important for this work is its interpretation via monodromy representations as an enumeration of factorisations in the symmetric group. As elliptic curves are the simplest cases of Calabi-Yau varieties, Hurwitz numbers of elliptic curves play a role in mirror symmetry. Dijkgraaf studied the relation between generating functions of Hurwitz numbers of an elliptic curve and Feynman integrals [Dij95].

In recent work of Chapuy and Dolega [CD22] a new class of Hurwitz numbers was introduced, called b -Hurwitz numbers depending on a parameter b . For $b = 0$ one obtains classical Hurwitz numbers, while for $b = 1$ these invariants specialise to an enumeration of covers between possibly non-orientable surface. Following [BF21], this enumeration gives rise to *twisted Hurwitz numbers* which were proved to admit a definition in terms of counting factorisations in the symmetric group in *loc. cit.* that mirrors its classical counterpart. In previous work [HM22], the authors developed a tropical geometry framework for the study of twisted Hurwitz numbers. So far, twisted Hurwitz numbers (and b -Hurwitz numbers) have only been studied for the enumeration of maps with a genus 0 target. In this work, we introduce twisted Hurwitz numbers of an elliptic curve, also in terms of analogous factorisations in the symmetric group. Motivated by the fact that tropical geometry also provides a natural framework for the study of covers of an elliptic curve [BBBM17; HIL22], we also study the tropical geometry of our twisted Hurwitz numbers of an elliptic curve.

1.1. Elliptic Hurwitz numbers. We first introduce the class of Hurwitz numbers of an elliptic curve showing up in the mirror symmetry relation involving Feynman integrals.

Key words and phrases. Tropical geometry, Hurwitz numbers.
2010 Mathematics Subject Classification: 14T15, 14N10, 57M12, 05C30.




FIGURE 1. A sketch of a cut open elliptic curve and the paths in its fundamental group which create the tuples to be counted to obtain a Hurwitz number via monodromy representations.

Definition 1 (Hurwitz numbers of an elliptic curve). Let E be an elliptic curve (i.e. a Riemann surface of genus 1), $g \geq 1$ a non-negative integer, and $d > 0$ a positive integer. Moreover, we fix $p_1, \dots, p_{2g-2} \in E$. Then, we consider covers of degree d , $f: S \rightarrow E$, such that

- S is a Riemann surface of genus g ,
- the ramification profile of p_1, \dots, p_{2g-2} is $(2, 1, \dots, 1)$.

Two covers $f: S \rightarrow E$ and $f': S' \rightarrow E$ are called equivalent if there exists a homeomorphism $h: S \rightarrow S'$, such that $f = f' \circ h$.

Then, we define the **Hurwitz number of the elliptic curve E** as

$$h_{d,g} = \sum_{[f]} \frac{1}{|\text{Aut}(f)|},$$

where the sum runs over all equivalence classes of covers as above.

Such Hurwitz numbers are in fact topological invariants, i.e. they do not depend on the algebraic structure of the Riemann surfaces. Via monodromy representations (see e.g. [CM16]), Hurwitz numbers of an elliptic curve can be computed in terms of factorisations in the symmetric group.

Lemma 2. *The Hurwitz number $h_{d,g}$ equals $\frac{1}{d!}$ times the number of tuples*

$$(\sigma, \tau_1, \dots, \tau_{2g-2}, \alpha) \in (\mathbb{S}_d)^{2g}$$

that satisfy the following:

- (1) each τ_i is a transposition,
- (2) the product of these permutations satisfies the following equation:

$$\tau_{2g-2} \dots \tau_1 \sigma = \alpha \sigma \alpha^{-1},$$

- (3) the subgroup

$$\langle \sigma, \tau_1, \dots, \tau_{2g-2}, \alpha \rangle$$

acts transitively on the set $\{1, \dots, d\}$.

The idea for the proof of Lemma 2 is to lift loops in the fundamental group of the elliptic curve to paths in the covering surface, see Figure 1.

1.2. **Twisted elliptic Hurwitz numbers.** We fix the involution

$$\tau = (1 \ d+1)(2 \ d+2) \dots (d \ 2d) \in \mathbb{S}_{2d}$$

and use the notation

$$B_d = C(\tau) = \{\sigma \in \mathbb{S}_{2d} \mid \sigma\tau\sigma^{-1} = \tau\}, \quad C^\sim(\tau) = \{\sigma \in \mathbb{S}_{2d} \mid \tau\sigma\tau^{-1} = \tau\sigma\tau = \sigma^{-1}\}.$$

We further define the subset $B_d^\sim \subset C^\sim(\tau)$ consisting of those permutations that have no self-symmetric cycles (see [BF21, Lemma 2.1]). We define the **twisted Hurwitz numbers of an elliptic curve** in terms of the symmetric group.

Definition 3 (Twisted Hurwitz numbers of an elliptic curve). Fix a genus g and a degree d . Then the twisted Hurwitz number $\tilde{h}_{d,g}$ of degree d and genus g of an elliptic curve is defined to be $\frac{1}{(2d)!!}$ times the number of tuples

$$(\sigma, \eta_1, \dots, \eta_{g-1}, \alpha) \in (S_{2d})^{g+1}$$

that satisfy the following conditions:

- (1) each η_s is a transposition, $\eta_s = (i_s \ j_s)$ such that $j_s \neq \tau(i_s)$,
- (2) $\sigma \in B_d^\sim$,
- (3) $\alpha \in B_d$,
- (4) the product of these permutations satisfies the following equation:

$$\eta_1 \dots \eta_{g-1} \sigma (\tau \eta_{g-1} \tau) \dots (\tau \eta_1 \tau) = \alpha \sigma \alpha^{-1},$$

- (5) the subgroup

$$\langle \sigma, \eta_1, \dots, \eta_{g-1}, (\tau \eta_{g-1} \tau), \dots, (\tau \eta_1 \tau), \alpha \rangle$$

acts transitively on the set $\{1, \dots, 2d\}$.

The motivation to call these counts of factorisations in the symmetric group twisted Hurwitz numbers of an elliptic curve is coming from Lemma 2. We can also drop the transitivity condition. On the tropical side, this corresponds to allowing disconnected source curves for our covers. We denote these numbers by $\tilde{h}_{d,g}^\bullet$.

Example 4. The twisted Hurwitz number $\tilde{h}_{2,3}$ equals 16.

Remark 5. It is tempting to expect that $\tilde{h}_{d,g}$ admits a geometric interpretation similar to twisted Hurwitz numbers with genus 0 target (see [CD22, Section 2.2] and the discussion after Remark 6 in [HM22]). A reasonable expectation could be that elliptic twisted Hurwitz numbers count maps to an elliptic target E with an orientation reversing involution that respects an orientation reversing involution on E . We leave the question of such a geometric interpretation as an open problem.

1.3. **Main Results.** In Section 2, we develop a tropical approach to elliptic twisted Hurwitz numbers. We introduce tropical elliptic twisted Hurwitz numbers as enumerations of tropical coverings of a tropical elliptic curve, i.e. as certain maps between graphs. As the main result of this section, in Theorem 21 we prove a correspondence theorem stating that elliptic twisted Hurwitz numbers and their tropical counterparts coincide. This tropical interpretation then allows us to derive an expression in Theorem 30 of elliptic twisted Hurwitz numbers in terms of Feynman diagrams in Section 3. We note that the computation of elliptic Hurwitz numbers in terms of Feynman diagrams was first derived in [Dij95] and a new proof employing tropical techniques was given in [BBBM17]. Finally, we follow the slogan *bosonification is tropicalisation* which has now been established in a plethora of works [BG16b; CJMR18a; CJMR21; HL22; HL20] to express elliptic twisted Hurwitz numbers as matrix element on the bosonic Fock space in Section 4.




FIGURE 2. Two abstract tropical curves of genus 3. Edge lengths are not specified in the picture.

Acknowledgements. The authors thanks Raphaël Fesler for many useful discussions. The second author acknowledges support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), Project-ID 286237555, TRR 195. Computations have been made using the Computer Algebra System GAP and OSCAR [15; 22].

2. TWISTED TROPICAL COVERS OF AN ELLIPTIC CURVE

In [HM22], we have introduced twisted versions of tropical Hurwitz numbers. We now generalize to consider twisted versions of tropical covers of an elliptic curve and their counts, building on [CJM10; BBM11; BBBM17]. We start by recalling the basic notions of tropical curves and covers. Then we introduce twisted tropical covers of an elliptic curve, which can roughly be viewed as tropical covers with an involution. By fixing branch points, we produce a finite count of twisted tropical covers of an elliptic curve for which we show in the following that it coincides with the corresponding twisted Hurwitz number of an elliptic curve. Readers with a background in the theory of tropical curves are pointed to the fact that we only consider explicit tropical curves in the following, i.e. there is no genus hidden at vertices.

Definition 6 (Abstract tropical curves). An abstract tropical curve is a graph Γ with the following data:

- (1) The vertex set of Γ is denoted by $V(\Gamma)$ and the edge set of Γ is denoted by $E(\Gamma)$.
- (2) The 1-valent vertices of Γ are called *leaves* and the edges adjacent to leaves are called *ends*.
- (3) The set of edges $E(\Gamma)$ is partitioned into the set of ends $E^\infty(\Gamma)$ and the set of *internal edges* $E^0(\Gamma)$.
- (4) There is a length function

$$\ell: E(\Gamma) \rightarrow \mathbb{R} \cup \{\infty\},$$

such that $\ell^{-1}(\infty) = E^\infty(\Gamma)$.

The genus of an abstract tropical curve Γ is defined as the first Betti number of the underlying graph. An isomorphism of abstract tropical curves is an isomorphism of the underlying graphs that respects the length function. The combinatorial type of an abstract tropical curve is the underlying graph without the length function.

Example 7. A tropical elliptic curve E is a circle of a given length. It may have several two-valent vertices. In the following we will refer to any tropical elliptic curve as E and do not specify the number of vertices when it is clear from the context.

Example 8. Figure 2 shows two abstract tropical curves of genus 3. We have not specified edge lengths in the picture.

Next, we define the notion of tropical covers. We restrict to the case where the target is either a tropical elliptic curve E or a subdivided version of \mathbb{R} , i.e. a line with some 2-valent vertices.

Definition 9 (Tropical covers). Let the target Γ_2 be either a tropical elliptic curve E or a subdivided version of \mathbb{R} . A tropical cover between abstract tropical curves $\pi: \Gamma_1 \rightarrow \Gamma_2$ is a surjective harmonic map, i.e.:




FIGURE 3. Two twisted tropical covers of E of degree 2 and genus 3. The labels on the edges denote the edge weight. Edges which are not labeled have weight one. The involution ι is supposed to exchange respective edges on top resp. bottom of the picture.

- (1) We have $\pi(V(\Gamma_1)) \subset V(\Gamma_2)$.
- (2) Let $e \in E(\Gamma_1)$. Then, we interpret e and $\pi(e)$ as intervals $[0, \ell(e)]$ and $[0, \ell(\pi(e))]$ respectively. We require π restricted to e to be a bijective integer linear function $[0, \ell(e)] \rightarrow [0, \ell(\pi(e))]$ given by $t \mapsto \omega(e) \cdot t$, with $\omega(e) \in \mathbb{Z}$. If $\pi(e) \in V(\Gamma_2)$, we define $\omega(e) = 0$. We call $\omega(e)$ the *weight* of e .
- (3) For a vertex $v \in V(\Gamma_1)$, we denote by $\text{Inc}(v)$ the set of incoming edges at v (edges adjacent to v mapping to the left of $\pi(v)$) and by $\text{Out}(v)$ the set of outgoing edges at v (edges adjacent to v mapping to the right of $\pi(v)$). We then require

$$\sum_{e \in \text{Inc}(v)} \omega(e) = \sum_{e \in \text{Out}(v)} \omega(e).$$

This number is called the local degree of π at v . We call this equality the *harmonicity* or *balancing condition*. For a point v in the interior of an edge e of Γ_1 , the local degree of π at v is defined to be the weight $\omega(e)$.

Moreover, we define the *degree* of π as the sum of local degrees at all vertices of Γ_1 in the pre-image of a given vertex of Γ_2 . By the harmonicity condition, the degree is independent of the choice of vertex of Γ_2 .

For any end e of Γ_2 , we define a partition μ_e as the partition of weights of ends of Γ_1 mapping to e . We call μ_e the *ramification profile* of e .

We call two tropical covers $\pi_1: \Gamma_1 \rightarrow \Gamma_2$ and $\pi_2: \Gamma'_1 \rightarrow \Gamma_2$ equivalent if there exists an isomorphism $g: \Gamma_1 \rightarrow \Gamma'_1$ of metric graphs, such that $\pi_2 \circ g = \pi_1$.

We are now ready to give a definition of twisted tropical covers, which may be viewed as tropical covers admitting an involution with specified locus.

Definition 10 (Twisted tropical covers of E). We define a twisted tropical cover of E to be a tropical cover $\pi: \Gamma_1 \rightarrow E$ with an involution $\iota: \Gamma_1 \rightarrow \Gamma_1$ which respects the cover π , such that:

- we have $g - 1$ branch points $p_1, \dots, p_{g-1} \in E$ which we set as vertices,
- in the preimage of each branch point p_i , there are either two 3-valent vertices or one 4-valent vertex,
- the edges adjacent to a 4-valent vertex all have the same weight,
- the fixed locus of ι is exactly the set of 4-valent vertices.

Example 11. Using the two source curves from Example 8 (see Figure 2), we can build two twisted tropical covers of degree 2, see Figure 3.




FIGURE 4. The quotient covers of the two twisted covers from Example 11, see Figure 3.

As usually for Hurwitz numbers, our enumeration of twisted tropical covers will take automorphisms into account. We give the following definition which specifies automorphisms that take the involution into account.

Definition 12 (Automorphisms). Let $\pi : \Gamma \rightarrow E$ be a twisted tropical cover with involution $\iota : \Gamma \rightarrow \Gamma$. An automorphism of π is a morphism of abstract tropical curves (i.e. a map of metric graphs) $f : \Gamma \rightarrow \Gamma$ respecting the cover and the involution, i.e. $\pi \circ f = \pi$ and $f \circ \iota = \iota \circ f$. We denote the group of automorphisms of π by $\text{Aut}(\pi)$.

Example 13. The twisted tropical cover depicted in Figure 3 on the left has an automorphism group of size 4: we can independently exchange the two pairs of weight 2 edges mapping to the same segment of E . The twisted tropical cover on the right has an automorphism group of size 2, we can exchange the two edges of weight 2, with the two pairs of edges of weight 1 following along.

Definition 14 (Quotient graph Γ/ι , see [HM22]). Let $\pi : \Gamma \rightarrow E$ be a twisted tropical cover with involution $\iota : \Gamma \rightarrow \Gamma$. The involution ι induces a symmetric relation on the vertex and edge sets of Γ : We define for $v, v' \in V(\Gamma)$ (resp. $e, e' \in E(\Gamma)$) that $v \sim v'$ (resp. $e \sim e'$) if and only if $\iota(v) = v'$ (resp. $\iota(e) = e'$). We define Γ/ι as the graph with vertex set $V(\Gamma)/\sim$ and edge set $E(\Gamma)/\sim$ with natural identifications. For $e = [e', e''] \in E(\Gamma/\iota)$ we define the length $\ell(e)$ as $\ell(e') = \ell(e'')$ and its weight $\omega(e)$ with respect to π to be the weight $\omega(e') = \omega(e'')$. In this way, we obtain a tropical cover from the quotient graph Γ/ι to \mathbb{R} , which has 2-valent vertices coming from the 4-valent vertices of Γ , and 3-valent vertices else.

Example 15. The quotient covers of the two twisted covers from Figure 3 are depicted in Figure 4.

Proposition 16. Let $\pi : \Gamma \rightarrow E$ be a twisted tropical cover with involution ι , and $\bar{\pi} : \Gamma/\iota \rightarrow E$ the corresponding tropical cover from the quotient (see Definition 14). Assume Γ has c 4-valent vertices. Then

$$\#\text{Aut}(\pi) = 2^c \cdot \#\text{Aut}(\bar{\pi}).$$

Proof. For every 4-valent vertex, we can exchange the edges of one adjacent twisted pair, leading to an automorphism of π which descends to the identity on the quotient. Also the involution ι descends to the identity on the quotient. As E is a circle, this yields altogether 2^c automorphisms of the twisted cover. Every other automorphism yields a non-trivial automorphism on the quotient. \square

Remark 17. An analogous statement as in Proposition 16 relating automorphisms of a twisted cover and automorphisms of its quotient cover also holds for the case of twisted tropical covers of \mathbb{R} . There we have

$$\#\text{Aut}(\pi) = 2^{c+1} \cdot \#\text{Aut}(\bar{\pi}).$$

The extra factor of 2 arises since \mathbb{R} is not a circle, so we can apply all automorphisms from 4-valent vertices and ι independently.

We are now ready to define twisted tropical Hurwitz numbers of an elliptic curve.

Definition 18 (Twisted tropical Hurwitz number of an elliptic curve). We define the tropical twisted double Hurwitz number of an elliptic curve $\tilde{h}_{d,g}^{\text{trop}}$ to be the weighted enumeration of equivalence classes of twisted tropical covers of degree d and genus g of a tropical elliptic curve E , such that each equivalence class $[\pi: \Gamma \rightarrow \mathbb{R}]$ is counted with multiplicity

$$2^{g-1} \cdot \frac{1}{|\text{Aut}(\pi)|} \cdot \prod_V (\omega_V - 1) \prod_e \omega(e),$$

where the first product goes over all 4-valent vertices and ω_V denotes the weight of the adjacent edges, while the second product is taken over all edges of the quotient graph Γ/ι and $\omega(e)$ denotes their weights.

Example 19. The twisted tropical Hurwitz number $\tilde{h}_{2,3}^{\text{trop}}$ equals 16. There are four twisted tropical covers for $d = 2$ and $g = 3$. We may obtain all four from the sketches in Figure 3. Each depicted map gives rise to two tropical covers depending on the labelling of the branch points, i.e. for the tropical elliptic curve E on the bottom, either the left vertex is labelled p_1 and the right vertex is labelled p_2 or the left vertex is labelled p_2 and the right vertex is labelled p_1 . Each cover coming from the left map in Figure 3 has multiplicity

$$2^2 \cdot \frac{1}{4} \cdot (2-1) \cdot (2-1) \cdot 2 \cdot 2 = 4.$$

Similarly, each cover coming from the right has multiplicity

$$2^2 \cdot \frac{1}{2} \cdot 2 = 4.$$

For the sizes of the automorphism groups, see Example 13. Thus, we obtain $\tilde{h}_{2,3}^{\text{trop}} = 16$ in total. Note that this number coincides with the twisted Hurwitz number we computed in Example 4.

Remark 20. Notice that twisted tropical Hurwitz numbers of an elliptic curve can also be determined by counting quotient covers directly, as quotients are in bijection to twisted covers. By Proposition 16, a quotient cover $\bar{\pi}$ has to be counted with multiplicity

$$2^{g-c-1} \cdot \frac{1}{|\text{Aut}(\bar{\pi})|} \cdot \prod_V (\omega_V - 1) \prod_e \omega(e),$$

where c denotes the number of 2-valent vertices of the source graph Γ , the first product goes over all 2-valent vertices and ω_V denotes the weight of the adjacent edges, while the second product is taken over all edges and $\omega(e)$ denotes their weights.

The equality observed in Example 19 is no coincidence, as shown in the following Theorem:

Theorem 21 (Correspondence Theorem). *The twisted Hurwitz number of E equals its tropical counterpart, i.e.*

$$\tilde{h}_{d,g}^{\text{trop}} = \tilde{h}_{d,g}.$$

Proof. Let $\pi: \Gamma \rightarrow E$ be a twisted tropical cover of degree d . We pick a base point p_0 between p_{g-1} and p_1 and cut the elliptic curve open at p_0 . We also cut the preimages of p_0 under π , thus obtaining a twisted tropical cover $\tilde{\pi}: \tilde{\Gamma} \rightarrow \mathbb{R}$. In the untwisted case, this is explained in detail in Construction 4.4 in [BBBM17].

The twisted Hurwitz number counts tuples

$$\tilde{h}_{d,g} = \frac{1}{(2d)!!} \#\{(\sigma, \eta_1, \dots, \eta_{g-1}, \alpha)\}$$

as in Definition 1. We can split these tuples and first list tuples of the form $\{(\sigma, \eta_1, \dots, \eta_{g-1})\}$, combining each such tuple with a list of possible α . Each tuple $\{(\sigma, \eta_1, \dots, \eta_{g-1})\}$ yields a twisted

tropical cover $\tilde{\pi}$ of \mathbb{R} as in Construction/Theorem 14 [HM22]. This cover has left and right ends of weights given by the cycle lengths of σ . For each twisted tropical cover of \mathbb{R} having ends of the same weights in both directions, the number of tuples of the form above leading to this cover equals its tropical multiplicity by the correspondence theorem (see Proposition 18 and Remark 6 in [HM22]). The tropical multiplicity equals

$$2^{g-1} \prod_V (\omega_V - 1) \prod_e \omega(e) \prod_K \frac{1}{\omega_K} \cdot \frac{1}{|\text{Aut}(\tilde{\pi})|}$$

where the first product goes over all 4-valent vertices V and ω_V denotes the weight of its adjacent edges, the second product goes over all pairs of twisted internal edges of the source $\tilde{\Gamma}$ of the twisted tropical cover $\tilde{\pi}$, and the third over all twisted pairs of components K which consist of a single edge of weight ω_K . Combining tuples of the form $\{(\sigma, \eta_1, \dots, \eta_{g-1})\}$ with possible α amounts to gluing a twisted tropical cover of \mathbb{R} to obtain a twisted tropical cover of E . To determine the number of such gluings, we pass to the quotient covers on each side. Given a twisted tropical cover π of E and its cut cover $\tilde{\pi}$, we consider the quotient cover $\bar{\pi}$ of E and the quotient cut cover $\tilde{\bar{\pi}}$. Note that taking the quotient and cutting the cover commutes.

When gluing a cut quotient cover of \mathbb{R} to a quotient cover of E , we want to pair up left and right ends that should be glued. Each left end of the quotient cover corresponds to a pair of ends of the twisted cover. Assume the pair of cycles $c_1, \tau \circ c_1 \circ \tau$ corresponds to these two ends, and assume that our gluing merges the left end corresponding to c_1 with the right end corresponding to a cycle c_2 of the same length. We want to count the number of α that satisfy $c_2 = \alpha \circ c_1 \circ \alpha^{-1}$. Let $c_1 = (c_{11}, \dots, c_{1\ell(c_1)})$, and $c_2 = (c_{21}, \dots, c_{2\ell(c_2)})$. A choice of α is fixed by setting $\alpha(c_{11}) = c_{2i}$ for any $i = 1, \dots, \ell(c_1)$. As we require that $\alpha\tau = \tau\alpha$ any element in the twisted cycle $\tau \circ c_1 \circ \tau$ of the form $\tau(c_{1j})$ must be mapped to $\tau(\alpha(c_{1j}))$ via α . Thus, choosing a gluing on one of a pair of twisted ends of the cut cover fixes the gluing on the other.

By the same argument as in [BBBM17, Proposition 4.9] the number of such α is given by

$$\prod_{e'} \omega(e')^{c_{e'}} \cdot \frac{|\text{Aut}(\tilde{\pi})|}{|\text{Aut}(\pi)|}, \quad (1)$$

where the product goes over all pairs of twisted edges e' of Γ that contain a preimage of the base point p_0 of E and $c_{e'}$ denotes the number of preimages in e' , $c_{e'} = \#(\pi^{-1}(p_0) \cap e')$.

We can group the tuples in the set according to the twisted tropical cover $\pi : C \rightarrow E$ they provide under the cut-and-join construction, see Construction/Theorem 14 in [HM22]. Thus we can write $\tilde{h}_{d,g}$ as

$$\tilde{h}_{d,g} = \frac{1}{(2d)!!} \cdot \sum_{\pi} \# \{(\sigma, \eta_1, \dots, \eta_{g-1}, \alpha) \text{ yielding the cover } \pi\}.$$

For a fixed cover π , instead of counting tuples yielding π , we can count tuples $(\sigma, \eta_1, \dots, \eta_{g-1})$ yielding the cut twisted tropical cover $\tilde{\pi}$ and then multiply with the number of appropriate α , which we denote by $n_{\tilde{\pi}, \pi}$:

$$\tilde{h}_{d,g} = \frac{1}{(2d)!!} \cdot \sum_{\pi} \# \{(\sigma, \eta_1, \dots, \eta_{g-1}) \text{ that provide the cover } \tilde{\pi}\} \cdot n_{\tilde{\pi}, \pi}.$$

By the above, the count of the tuples yielding a cover $\tilde{\pi}$ divided by $(2d)!!$ equals

$$2^{g-1} \frac{1}{|\text{Aut}(\tilde{\pi})|} \cdot \prod_V (\omega_V - 1) \cdot \prod_{\tilde{e}} \omega(\tilde{e}) \cdot \prod_K \frac{1}{\omega_K}$$

where the first product goes over the 4-valent vertices, the second over all pairs of twisted internal edges \tilde{e} of $\tilde{\Gamma}$ of weight $\omega(\tilde{e})$ and the third over all twisted pairs of components K consisting of a




FIGURE 5. The twisted tropical cover on the right in Figure 3 cut open at the back.

single edge of weight ω_K . From the above, the number $n_{\tilde{\pi}, \pi}$ can be substituted by the expression in Equation (1).

We obtain

$$\tilde{h}_{d,g} = \sum_{\pi} \frac{1}{|\text{Aut}(\tilde{\pi})|} 2^{g-1} \cdot \prod_V (\omega_V - 1) \cdot \prod_{\tilde{e}} \omega(\tilde{e}) \cdot \prod_K \frac{1}{\omega_K} \cdot \prod_{e'} \omega(e')^{c_{e'}} \cdot \frac{|\text{Aut}(\tilde{\pi})|}{|\text{Aut}(\pi)|}.$$

A pair of twisted edges e' of Γ of weight $\omega(e')$ having $c_{e'}$ preimages over the base point provides exactly $c_{e'} - 1$ pairs of single-edge-components of weight $\omega(e')$ in the cut cover $\tilde{\pi}$. Vice versa, each such pair of components comes from a pair of edges with multiple preimages over the base point. Therefore the expression $\prod_K \frac{1}{\omega_K} \cdot \prod_{e'} \omega(e')^{c_{e'}}$ simplifies to $\prod_{e'} \omega(e')$. We obtain

$$\tilde{h}_{d,g} = \sum_{\pi} 2^{g-1} \frac{1}{|\text{Aut}(\pi)|} \cdot \prod_V (\omega_V - 1) \cdot \prod_e \omega(e) = \tilde{h}_{d,g}^{\text{trop}}$$

and the theorem is proved. \square

Example 22. Consider the twisted tropical cover π depicted in Figure 3 on the right with the vertex on the left labelled p_1 and the vertex on the right labelled p_2 . Cutting it open, we obtain the twisted tropical cover $\tilde{\pi}$ of \mathbb{R} depicted in Figure 5.

By the correspondence theorem for twisted tropical covers of \mathbb{R} in [HM22, Theorem 22], it accounts for $(2n)!! = 8$ times its tropical multiplicity many tuples. The cover has an automorphism group of size $2^2 = 4$, as we can independently exchange both pairs of twisted edges of weight 2, with the edges of weight 1 following along. Its tropical multiplicity thus equals

$$2^2 \cdot \frac{1}{4} = 1.$$

This twisted tropical cover of \mathbb{R} thus accounts for 8 tuples of the form $(\sigma_1, \eta_1, \eta_2)$. By Lemma 16 [HM22], there are 2 permutations suitable for σ_1 , (14)(23) and (12)(34). Fix $\sigma_1 = (14)(23)$ momentarily, the other choice is analogous. Then there are two choices for η_1 , (14) or (23). For the next branch point, there are 2 more choices for η_2 , (12) or (34). Altogether, we obtain the 8 tuples as expected.

The extra automorphism that the cut cover obtains (we have $\frac{|\text{Aut}(\tilde{\pi})|}{|\text{Aut}(\pi)|} = \frac{4}{2} = 2$) allows to make an additional choice which left end should be glued to which right end. Let us momentarily fix one of our 8 tuples, $((14)(23), (14), (12))$. If we label all edges with the corresponding permutations, the two right ends are labeled with (12) and (34). Because of the extra automorphism, we can glue the left end labeled (14) either to (12) or to (34). For each choice, we obtain as many α satisfying $\alpha\tau = \tau\alpha$ as the weight of one end, i.e. 2. We thus obtain 4 possible α to add to each of the 8 tuples, yielding 32 tuples of the form $(\sigma_1, \eta_1, \eta_2, \alpha)$. For the tuple fixed above, the 4 possible α we can add are

$$\{(24), (1234), (13), (1432)\}.$$

Dividing the 32 tuples by $(2n)!! = 8$, we expect the tropical multiplicity of the right cover of E in Figure 3 to be 4. Indeed, in Example 19 we already computed its tropical multiplicity to be

$$2^2 \cdot \frac{1}{2} \cdot 2 = 4.$$

3. GENERATING SERIES IN TERMS OF FEYNMAN INTEGRALS

In this section, we express elliptic twisted Hurwitz numbers as Feynman integrals. We assume that $g > 2$ in the following. Consequently, in the quotient covers there cannot be loop edges.

In our context, the following definition of Feynman graph will be needed. These are exactly the graphs that appear as sources for quotients of twisted covers, up to labeling.

Definition 23 (Feynman graph). A Feynman graph is a graph with 2- and 3-valent vertices whose edges are labeled with q_1, \dots, q_r and whose vertices are labeled with x_1, \dots, x_s .

A Feynman integral depends on a Feynman graph and the choice of an order Ω of the vertices.

Definition 24 (Edge Propagator). Let q_k be an edge of a Feynman graph, adjacent to two vertices x_{k_1} and x_{k_2} , where we assume that $x_{k_1} < x_{k_2}$ in the order Ω .

Given $w \in \mathbb{N}$, we define the coefficient c_w of the following propagator function to be

$$c_w := \begin{cases} (w-1) \cdot w & \text{if } x_{k_1} \text{ and } x_{k_2} \text{ are 2-valent} \\ \sqrt{w-1} \cdot w & \text{if } x_{k_1} \text{ or } x_{k_2} \text{ are 2-valent} \\ w & \text{if neither } x_{k_1} \text{ nor } x_{k_2} \text{ are 2-valent.} \end{cases}$$

We then define the propagator function of the edge q_k to be

$$P(q_k) = \sum_{w=1}^{\infty} c_w \left(\frac{x_{k_1}}{x_{k_2}} \right)^w + \sum_{a_k=1}^{\infty} \left(\sum_{w|a_k} c_w \left(\left(\frac{x_{k_1}}{x_{k_2}} \right)^w + \left(\frac{x_{k_2}}{x_{k_1}} \right)^w \right) \right) q_k^{a_k}.$$

Definition 25 (Feynman integral). Let Γ be a Feynman graph and Ω be an order of its vertices. For each edge q_k , we denote its adjacent vertices by x_{k_1} and x_{k_2} , where we assume that $x_{k_1} < x_{k_2}$ in the order Ω . We define the Feynman integral $I_{\Gamma, \Omega}(q_1, \dots, q_r)$ to be

$$I_{\Gamma, \Omega}(q_1, \dots, q_r) = \text{coef}_{[x_1^0 \dots x_s^0]} \prod_{k=1}^r P(q_k).$$

Setting all q_k equal to one variable q , we obtain the Feynman integral $I_{\Gamma, \Omega}(q)$.

Remark 26. Here, we consider Feynman integrals merely as formal power series. In the relation involving (usual) Hurwitz numbers of an elliptic curve, the propagator series can, using a coordinate change, be transformed into a linear combination of the Weierstraß- \wp -function and an Eisenstein series. After this coordinate change, the Feynman integral can be viewed as a complex analytic path integral.

Remark 27. Fix a genus $g > 2$. By an Euler characteristic computation, the source of a twisted tropical cover of E has $2g - 2 - c$ vertices, where c denotes the number of 4-valent vertices. That is, $2g - 2 - 2c$ vertices are 3-valent and c are 4-valent. When passing to the quotient cover, its source graph has $g - 1 - c$ many 3-valent vertices and c many 4-valent vertices. Its total number of vertices is thus $g - 1$, independent of the number of 4-valent vertices in the twisted cover.

Definition 28 (Labeled quotient covers). A *labeled quotient cover* is a quotient of a twisted tropical cover for which the vertices and edges of its source are labeled like a Feynman graph.

Fix a base point b of a tropical elliptic curve E . Fix a genus $g > 2$, and $g-1$ branch points b_1, \dots, b_{g-1} in E .

Given a labeled quotient cover $\bar{\pi}$, we can define its *multidegree* $a \in \mathbb{N}^r$ to be the tuple whose k -th entry equals the sum of the weights of the preimages of the base point b in the edge q_k . Fix an order Ω on $g-1$ elements x_1, \dots, x_{g-1} . Let Γ be a Feynman graph.

We define $\tilde{h}_{\Gamma, \Omega, a}$ to be the weighted number of labeled quotient covers whose source is of combinatorial type Γ , whose multidegree equals a and such that the order given by the preimages of the branch points $\bar{\pi}^{-1}(b_1) < \dots < \bar{\pi}^{-1}(b_{g-1})$ equals Ω . Each such cover is weighted by $2^{g-c-1} \cdot \prod_V (\omega_V - 1) \cdot \omega(e)$. Since there are no nontrivial automorphisms in the presence of labels, this equals the multiplicity given in Remark 20.

Proposition 29. *Let Γ be a Feynman graph of genus $g > 2$ and c be its number of 2-valent vertices. Let a be a multidegree and Ω an order. The count of labeled quotient covers equals a coefficient of a Feynman integral:*

$$\tilde{h}_{\Gamma, \Omega, a} = 2^{g-c-1} \cdot \text{coef}_{[q_1^{a_1} \dots q_r^{a_r}]} I_{\Gamma, \Omega}(q_1, \dots, q_r).$$

The proof follows ideas of [BBBM17; BGM22].

Proof. Expanding the product $\prod_{k=1}^r P(q_k)$, the summands are equal to products of the form

$$\prod_{k=1}^r c_{w_k} \left(\frac{x_i}{x_j} \right)^{w_k} \cdot q_k^{a_k}.$$

If a_k is zero, w_k can be any element in \mathbb{N} , and $i = k_1, j = k_2$. If $a_k > 0$, $w_k | a_k$ and i can be either k_1 or k_2 , and j the remaining. To each such summand, we associate a labeled quotient cover in the following way: We start by fixing as preimages of the branch points the vertices x_i as imposed by the order Ω .

For a factor $c_{w_k} \left(\frac{x_i}{x_j} \right)^{w_k} \cdot q_k^{a_k}$ with $a_k = 0$, we draw an edge labeled q_k which goes from the vertex x_{k_1} to x_{k_2} without crossing over the base point. This is possible since the x_i respect the order Ω . We fix the weight of our edge to be w_k .

For a factor $c_{w_k} \left(\frac{x_i}{x_j} \right)^{w_k} \cdot q_k^{a_k}$ with $a_k > 0$, we draw an edge labeled q_k : If $i = k_1$ we let it start at x_{k_1} and connect with x_{k_2} (where we think of the edges of our cover as oriented in the way imposed by the order Ω). If $i = k_2$, we let it start at x_{k_2} at connect it with x_{k_1} . We "curl" this edge in such a way that it passes $\frac{a_k}{w_k}$ times over the base point b . The weight of the edge in each case is defined to be w_k .

We claim that in this way, we produce a labeled quotient cover contributing to $\tilde{h}_{\Gamma, \Omega, a}$ with $a = (a_1, \dots, a_r)$. Since we used the edge q_k to connect its neighboring vertices in Γ , the source of the covers is of combinatorial type Γ by construction. The multidegree is a , since for each k with $a_k = 0$, we let our edge not pass over the base point, whereas for each k with $a_k > 0$ the edge of weight w_k passes $\frac{a_k}{w_k}$ times over the base point, leading to the entry a_k in the multidegree. The order Ω is also respected by construction.

What remains to be seen is that we obtained indeed a cover, i.e. the balancing condition has to be satisfied. This holds true since a product as above only contributes to the Feynman integral $I_{\Gamma, \Omega}(q_1, \dots, q_r)$ if its constant degree in the x_i vanishes. The total power of x_i equals, by construction, the signed sum of the weights of its adjacent edges. The fact that the degree in x_i is constant is thus equivalent to the balancing condition at vertex x_i .

In this way, we obtain a bijection between summands contributing to the Feynman integral and labeled quotient covers. What about multiplicities? In the Feynman integral, a summand contributes

$\prod_k c_{w_k}$. Thus, the summand contributes $2^{g-c-1} \cdot \prod_k c_{w_k}$ to the right hand side. We have to show that this equals the multiplicity with which the labeled quotient cover is counted in Remark 20, i.e. that $\prod_k c_{w_k} = \prod_V (\omega_V - 1) \cdot \prod_e \omega_e$. Recall that the weight of the edge q_k equals w_k . For a 2-valent vertex V , we have a factor of $\omega_V - 1$, where ω_V denotes the weight of the adjacent edges. We can thus part this contribution into two factors of $\sqrt{\omega_V - 1}$ and shift those towards the adjacent edges.

By definition, if q_k connects two 2-valent vertices, c_{w_k} equals $(w_k - 1) \cdot w_k$ – it obtains two factors of $\sqrt{w_k - 1}$ from both its adjacent vertices, and it also contributes its own weight, as every edge does. If q_k connects a 2-valent with a 3-valent vertex, it obtains only one factor of $\sqrt{w_k - 1}$. If both vertices of q_k are 3-valent, it obtains no such factor. It follows that if we reinterpret the product $\prod_V (\omega_V - 1) \cdot \prod_e \omega_e$ as a product over edges by shifting the vertex contributions as square roots into both adjacent edges, we get exactly the contribution c_{w_k} which is used to define the propagator function for the Feynman integral.

Thus the multiplicity with which a labeled quotient cover contributes to $\tilde{h}_{\Gamma, \Omega, a}$ exactly equals the contribution of its corresponding summand in the Feynman integral (up to the factor of 2^{g-c-1}), and the equality holds. \square

The following is the main theorem of this section and expresses the generation function of elliptic twisted Hurwitz numbers as a finite sum over Feynman integrals.

Theorem 30. *Fix a genus $g > 2$. The generating series of twisted Hurwitz numbers can be expressed in terms of Feynman integrals:*

$$\sum_d \tilde{h}_{d,g} q^d = 2^{g-1} \cdot \sum_{\Gamma} \frac{1}{2^{c_{\Gamma}} \cdot \#\text{Aut}(\Gamma)} \sum_{\Omega} I_{\Gamma, \Omega}(q).$$

Here, the first sum on the right hand side goes over all Feynman graphs of genus g and c_{Γ} denotes their number of 2-valent vertices, while the second sum goes over all order Ω .

Proof. For a fixed graph Γ , let $\tilde{h}_{d, \Gamma}$ be the number of (unlabeled) quotient covers of degree d for which the combinatorial type of the source curve is Γ . As in Remark 20, each cover $\bar{\pi}$ is counted with multiplicity $2^{g-c-1} \cdot \frac{1}{|\text{Aut}(\bar{\pi})|} \prod_V (\omega_V - 1) \prod_e \omega(e)$. There exists a forgetful map ft from the set of labeled quotient covers to the set of unlabeled covers by just forgetting the labels. For an (unlabeled) quotient cover $\bar{\pi}$ whose source is of combinatorial type Γ , the automorphism group of Γ , $\text{Aut}(\Gamma)$, acts transitively on the fiber $\text{ft}^{-1}(\bar{\pi})$ by relabeling vertices and edges. So, to determine the cardinality of the set $\text{ft}^{-1}(\bar{\pi})$, we think of it as the orbit under this action and obtain $\#\text{ft}^{-1}(\bar{\pi}) = \frac{\#\text{Aut}(\Gamma)}{\#\text{Aut}(\bar{\pi})}$, since the stabilizer of the action equals the set of automorphisms of $\bar{\pi}$. Each labeled quotient cover in the set $\text{ft}^{-1}(\bar{\pi})$ is counted with the same multiplicity $2^{g-c-1} \cdot \prod_V (\omega_V - 1) \prod_e \omega(e)$.

The sum $\sum_{a| \sum a_i = d} \sum_{\Omega} \tilde{h}_{\Gamma, \Omega, a}$ can be reorganized as a sum over unlabeled quotient covers, where for each unlabeled cover, we have to sum the multiplicities for each labeled quotient cover in the fiber under ft . As the multiplicity is the same for each element in the fiber, and there are $\frac{\#\text{Aut}(\Gamma)}{\#\text{Aut}(\bar{\pi})}$ elements in the fiber, we can see that this sum equals $\#\text{Aut}(\Gamma) \cdot \tilde{h}_{d, \Gamma}$.

We conclude

$$\sum_d \tilde{h}_{d,g} q^d = \sum_d \sum_{\Gamma} \tilde{h}_{d, \Gamma} q^d = \sum_d \sum_{\Gamma} \frac{1}{\#\text{Aut}(\Gamma)} \sum_{a| \sum a_i = d} \sum_{\Omega} \tilde{h}_{\Gamma, \Omega, a} q^d$$

Now we can replace $\tilde{h}_{\Gamma, \Omega, a}$ by $2^{g-c_{\Gamma}-1}$ times the coefficient of q^a in $I_{\Gamma, \Omega}(q_1, \dots, q_r)$ by Proposition 29. If we insert $q_k = q$ for all k we can conclude that the coefficient of q^d in $I_{\Gamma, \Omega}(q)$ equals $2^{g-c_{\Gamma}-1}$ times $\sum_{a| \sum a_i = d} \tilde{h}_{\Gamma, \Omega, a}$. Thus the generating series above equals

$$2^{g-1} \cdot \sum_{\Gamma} \frac{1}{2^{\text{cr}} \cdot \#\text{Aut}(\Gamma)} \sum_{\Omega} I_{\Gamma, \Omega}(q).$$

□

4. THE FOCK SPACE APPROACH

We shortly review the bosonic Fock space approach for generating series of Hurwitz numbers.

The bosonic Heisenberg algebra \mathcal{H} is the Lie algebra with basis α_n for $n \in \mathbb{Z}$ such that for $n \neq 0$ the following commutator relations are satisfied:

$$[\alpha_n, \alpha_m] = (n \cdot \delta_{n,-m}) \alpha_0,$$

where $\delta_{n,-m}$ is the Kronecker symbol and $[\alpha_n, \alpha_m] := \alpha_n \alpha_m - \alpha_m \alpha_n$. The bosonic Fock space F is a representation of \mathcal{H} . It is generated by a single “vacuum vector” v_\emptyset . The positive generators annihilate v_\emptyset : $\alpha_n \cdot v_\emptyset = 0$ for $n > 0$, α_0 acts as the identity and the negative operators act freely. That is, F has a basis b_μ indexed by partitions, where

$$b_\mu = \alpha_{-\mu_1} \dots \alpha_{-\mu_m} \cdot v_\emptyset.$$

We define an inner product on F by declaring $\langle v_\emptyset | v_\emptyset \rangle = 1$ and α_n to be the adjoint of α_{-n} .

We write $\langle v | A | w \rangle$ for $\langle v | Aw \rangle$, where $v, w \in F$ and the operator A is a product of elements in \mathcal{H} , and $\langle A \rangle$ for $\langle v_\emptyset | A | v_\emptyset \rangle$. The first is called a *matrix element*, the second a *vacuum expectation*.

Definition 31. The vertex operator is defined by:

$$M = 2 \cdot \left(\sum_{k>0} (k-1) \cdot \alpha_{-k} \alpha_k + \frac{1}{2} \sum_{k>0} \sum_{\substack{0 < i, j \\ i+j=k}} \alpha_{-j} \alpha_{-i} \alpha_k + \alpha_{-k} \alpha_i \alpha_j \right)$$

We note that unlike in the Feynman diagram approach here we don't have to shift vertex contributions into neighbouring edges. Moreover, the global factor of 2 is to take the number of branch points into account. We can also view it as vertex contribution.

We obtain the following result.

Proposition 32. *The twisted double Hurwitz number $\tilde{h}_g^\bullet(\mu, \nu)$ (see Definition 4, [HM22]) equals a matrix element on the bosonic Fock space:*

$$\tilde{h}_g^\bullet(\mu, \nu) = \frac{1}{\prod_i \mu_i \cdot \prod_j \nu_j} \langle b_\mu | M^{g-1} | b_\nu \rangle.$$

Proof. This statement follows by combining Wick's Theorem with the Correspondence Theorem for twisted double Hurwitz numbers in [HM22]: Wick's Theorem (Theorem 5.4.3 [CJMR18b], Proposition 5.2 [BG16a], [Wic50]) expresses a matrix element as a weighted count of graphs that are obtained by completing local pictures. It turns out that the graphs in question are exactly the quotient covers we enumerate to obtain $\tilde{h}_g^\bullet(\mu, \nu)$.

Notice that we have to use the disconnected theory here (\bullet), since the matrix element encodes *all* graphs completing the local pictures and cannot distinguish connected and disconnected graphs.

The local pictures are built as follows: we draw one vertex for each vertex operator. For an α_n with $n > 0$, we draw an edge germ of weight n pointing to the right. If $n < 0$, we draw an edge germ of weight n pointing to the left. For the two Fock space elements b_μ and b_ν , we draw germs of ends: of weights μ_i on the left pointing to the right, of weights ν_i on the right pointing to the left. Wick's Theorem states that the matrix element $\langle b_\mu | M^n | b_\nu \rangle$ equals a sum of graphs completing all possible local pictures, where each graph contributes the product of the weights of all its edges (including

the ends) and the vertex contributions arising from the vertex operator. A completion of the local pictures can be interpreted as a quotient cover of \mathbb{R} (with suitable metrization).

The vertex operator sums over all the possibilities of the local pictures for the graphs, i.e. it sums over all possibilities how a vertex of a quotient cover can look like. \square

Combining Proposition 32 with the relation we obtain via cutting in the proof of Theorem 21, we can express twisted Hurwitz numbers of the elliptic curve in terms of matrix elements:

Proposition 33. *A twisted Hurwitz number of the elliptic curve equals a weighted sum of twisted double Hurwitz numbers:*

$$\tilde{h}_{d,g}^\bullet = \sum_{\mu \vdash d} \frac{\prod_i \mu_i}{|\text{Aut}(\mu)|} \tilde{h}_g^\bullet(\mu, \mu).$$

Here, the sum goes over all partitions μ of d .

Proposition 33 is a corollary of the two Correspondence Theorems: given a tropical cover of E , let μ be the partition encoding the weights of the edges mapping to the base point p_0 . We mark the preimages of p_0 , for which we have $|\text{Aut}(\mu)|$ choices. For each choice, we cut off E at p_0 and the covering curve at the preimages of p_0 , obtaining a cover of \mathbb{R} with ramification profiles μ and μ above $\pm\infty$. The cut off tropical cover contributes to $\tilde{h}_g^\bullet(\mu, \nu)$, but its multiplicity differs from the multiplicity of the cover of E by a factor of $\prod \mu_i$, since the edges we cut off are no longer bounded.

Finally, we obtain an expression of elliptic twisted Hurwitz numbers as a matrix element on the bosonic Fock space as a corollary of Propositions 32 and 33.

Corollary 34. *A twisted Hurwitz number of the elliptic curve E equals a sum of matrix elements on the bosonic Fock space:*

$$\tilde{h}_{g,d}^\bullet = \sum_{\mu \vdash d} \frac{1}{|\text{Aut}(\mu)| \prod_i \mu_i} \langle b_\mu | M^n | b_\mu \rangle.$$

REFERENCES

- [BBM11] B. Bertrand, E. Brugallé, and G. Mikhalkin. “Tropical Open Hurwitz numbers”. In: *Rend. Semin. Mat. Univ. Padova* 125 (2011), pp. 157–171.
- [BG16a] F. Block and L. Göttsche. “Fock spaces and refined Severi degrees”. In: *Int. Math. Res. Not. IMRN* 21 (2016), pp. 6553–6580.
- [BG16b] F. Block and L. Göttsche. “Refined curve counting with tropical geometry”. In: *Compositio Mathematica* 152.1 (2016), pp. 115–151.
- [BBBM17] J. Böhm, K. Bringmann, A. Buchholz, and H. Markwig. “Tropical mirror symmetry for elliptic curves”. In: *J. Reine Angew. Math.* 732 (2017), pp. 211–246.
- [BGM22] J. Böhm, C. Goldner, and H. Markwig. “Tropical mirror symmetry in dimension one”. In: *SIGMA* 18 (2022). Preprint, arXiv:1809.10659, p. 046.
- [BF21] Y. Burman and R. Fesler. “Ribbon decomposition and twisted Hurwitz numbers”. Preprint, arXiv:2107.13861. 2021.
- [CJM10] R. Cavalieri, P. Johnson, and H. Markwig. “Tropical Hurwitz numbers”. In: *J. Algebraic Combin.* 32.2 (2010), pp. 241–265.
- [CJMR18a] R. Cavalieri, P. Johnson, H. Markwig, and D. Ranganathan. “A graphical interface for the Gromov-Witten theory of curves”. In: *Algebraic geometry: Salt Lake City 2015*. 2018, pp. 139–167.
- [CJMR18b] R. Cavalieri, P. Johnson, H. Markwig, and D. Ranganathan. “A graphical interface for the Gromov-Witten theory of curves”. In: *Algebraic geometry: Salt Lake City 2015*. Vol. 97. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 2018, pp. 139–167.

- [CJMR21] R. Cavalieri, P. Johnson, H. Markwig, and D. Ranganathan. “Counting curves on Hirzebruch surfaces: tropical geometry and the Fock space”. In: *Mathematical Proceedings of the Cambridge Philosophical Society*. Vol. 171. 1. Cambridge University Press. 2021, pp. 165–205.
- [CM16] R. Cavalieri and E. Miles. *Riemann surfaces and algebraic curves*. Vol. 87. London Mathematical Society Student Texts. A first course in Hurwitz theory. Cambridge University Press, Cambridge, 2016, pp. xii+183.
- [CD22] G. Chapuy and M. Dołęga. “Non-orientable branched coverings, b -Hurwitz numbers, and positivity for multiparametric Jack expansions”. In: *Adv. Math.* 409 (2022), Paper No. 108645.
- [Dij95] R. Dijkgraaf. “The moduli space of curves”. In: vol. 129. *Progr. Math.* Birkhäuser Boston, 1995. Chap. Mirror symmetry and elliptic curves, pp. 149–163.
- [15] *GAP – Groups, Algorithms, and Programming, Version 4.7.8*. The GAP Group. 2015.
- [HL20] M. Hahn and D. Lewański. “Wall-crossing and recursion formulae for tropical Jucys covers”. In: *Transactions of the American Mathematical Society* 373.7 (2020), pp. 4685–4711.
- [HIL22] M. A. Hahn, J.-W. van Ittersum, and F. Leid. “Triply mixed coverings of arbitrary base curves: quasimodularity, quantum curves and a mysterious topological recursion”. In: *Annales de l’Institut Henri Poincaré D* 9.2 (2022), pp. 239–296.
- [HL22] M. A. Hahn and D. Lewanski. “Tropical Jucys covers”. In: *Mathematische Zeitschrift* 301.2 (2022), pp. 1719–1738.
- [HM22] M. A. Hahn and H. Markwig. “Twisted Hurwitz numbers: tropical and polynomial structures”. Preprint, arXiv:2210.00595. 2022.
- [Hur92] A. Hurwitz. “Über algebraische Gebilde mit eindeutigen Transformationen in sich”. In: *Math. Ann.* 41.3 (1892), pp. 403–442.
- [22] *OSCAR – Open Source Computer Algebra Research system, Version 0.8.3-DEV*. The OSCAR Team. 2022.
- [Wic50] G. C. Wick. “The evaluation of the collision matrix”. In: *Physical Rev. (2)* 80 (1950), pp. 268–272.

M. A. HAHN: SCHOOL OF MATHEMATICS 17, WESTLAND ROW, TRINITY COLLEGE DUBLIN, DUBLIN 2, IRELAND
Email address: hahnma@tcd.ie

H. MARKWIG: UNIVERSITÄT TÜBINGEN, FACHBEREICH MATHEMATIK, AUF DER MORGENSTELLE 10, 72076 TÜBINGEN, GERMANY
Email address: hannah@math.uni-tuebingen.de