

Validation of ML-UQ calibration statistics using simulated reference values: a sensitivity analysis

Pascal PERNOT ¹

*Institut de Chimie Physique, UMR8000 CNRS,
Université Paris-Saclay, 91405 Orsay, France^{a)}*

Some popular Machine Learning Uncertainty Quantification (ML-UQ) calibration statistics do not have predefined reference values and are mostly used in comparative studies. In consequence, calibration is almost never validated and the diagnostic is left to the appreciation of the reader. Simulated reference values, based on synthetic calibrated datasets derived from actual uncertainties, have been proposed to palliate this problem. As the generative probability distribution for the simulation of synthetic errors is often not constrained, the sensitivity of simulated reference values to the choice of generative distribution might be problematic, shedding a doubt on the calibration diagnostic. This study explores various facets of this problem, and shows that some statistics are excessively sensitive to the choice of generative distribution to be used for validation when the generative distribution is unknown. This is the case, for instance, of the correlation coefficient between absolute errors and uncertainties (CC) and of the expected normalized calibration error (ENCE). A robust validation workflow to deal with simulated reference values is proposed.

^{a)}Electronic mail: pascal.pernot@cnrs.fr

I. INTRODUCTION

Most calibration statistics are used to compare UQ methods or datasets but are missing a reference value for validation. For instance, the correlation coefficient between absolute errors and uncertainties¹ should be positive, but has no predefined reference value to compare with². This is also the case for some conditional calibration statistics, such as the expected normalized calibration error (ENCE)³, which value depends moreover on the binning scheme⁴.

The use of *probabilistic*⁵ or *simulated*⁶ reference values has been recently proposed to palliate the absence of reference for a calibration statistic. A simulated reference value is estimated by applying the statistic to synthetic datasets containing the actual uncertainties and simulated errors generated from the uncertainties using a model generative distribution D .

The main issue with simulated references is the choice of D . Some calibration statistics depend explicitly on D (typically a normal distribution): calibration curves and calibration error⁷, negative log-likelihood (NLL)... This scenario fixes the choice of D to simulate a reference value. However, this is not the case for many other calibration statistics, such as the calibration error, confidence curves⁵ or reliability diagrams/ENCE⁴. Without constraint on D , the question arises of the sensitivity of such simulated reference values to the choice of generative distribution.

The case of confidence curves⁵ enables to underline the main targets of the present study. A confidence curve is obtained by estimating an error statistic (e.g. the mean absolute error, MAE) for a dataset iteratively pruned from its largest uncertainties. Plotted against the percentage of pruned data, the confidence curve tests how the largest errors are associated with the largest uncertainties. Ideally, a confidence curve should be monotonously decreasing, but there is no predefined reference curve to assess its quality (the so-called *oracle* is unreachable in ML regression tasks⁵). In this context, a confidence curve can inform us on the quality of the link between errors and uncertainties, but not on calibration. Using a simulated reference curve can solve this problem, at the cost of a choice for the generative distribution D , about which Pernot⁵ has raised two main points:

- the sensitivity of the simulated reference curve to D depends on the error statistic used to build the confidence curve: the MAE is very sensitive to D , while the root mean-squared error (RMSE) is not;
- for validation, a confidence band can be estimated from simulated reference curves. For all error statistics, the width of the confidence band depends on D , leading to

ambiguous validation diagnostics when D is not constrained.

The present study considers these two points for other calibration statistics. More specifically, it aims to test (i) how simulated reference values for calibration statistics are sensitive to the choice of a generative distribution, (ii) how this impacts the validation procedure, and (iii) whether the uncertainty on the simulated reference value is a good choice of metric for validation. The focus is limited here to statistics linked to ML regression tasks: the correlation coefficient between absolute errors and uncertainties (CC), an average calibration statistic (the mean squared z -scores, ZMS), and two conditional calibration statistics (the ENCE, its ZMS-derived analog ZMSE).

The next section defines the calibration statistics, the validation approach, and proposes a workflow dealing with the main issues of simulated references. Sect. III presents the application of these methods to an ensemble of nine datasets issued from the ML-UQ literature. The article proceeds with a discussion of the main findings (Sect. IV), and the conclusions are reported in Sect. V.

II. VALIDATION SCORES FOR CALIBRATION AND CORRELATION STATISTICS

This section presents the simulation method for the estimation of calibration statistics references for commonly used correlation, *average* calibration and *conditional* calibration statistics in order to define adequate methods for their validation.

A. Probabilistic generative model

Let us consider a dataset composed of *paired* errors and uncertainties $\{E_i, u_{E_i}\}_{i=1}^M$ to be tested for calibration. A few variance-based UQ validation statistics and methods, based on the correlation between the absolute errors and uncertainties (rank correlation coefficient (Sect. II B) and confidence curve), avoid the need of a probabilistic model linking those variables². In contrast, most of the variance-based UQ calibration statistics are built implicitly on a probabilistic model

$$E_i \sim D(\mu = 0, \sigma = u_{E_i}) \quad (1)$$

or, equivalently,

$$E_i = u_{E_i} \varepsilon_i \quad (2)$$

linking errors to uncertainties, where ε_i is a random number with zero-centered and unit-variance probability density function [$\varepsilon \sim D(\mu = 0, \sigma = 1)$]. This model states that errors

should be unbiased ($\mu = 0$) and that uncertainty quantifies the *dispersion* of errors, according to the metrological definition.⁸

Except in some instances where D is constrained by the method used to generate the dataset (gaussian process, normal likelihood models...), the shape of D is unknown. This is notably the case when uncertainties are obtained *by post-hoc* calibration^{3,9}.

Note that D should not be mistaken for the distribution of errors, which is a *compound distribution* between D and the distribution of uncertainties. Let us assume that the errors are drawn from a distribution D (Eq. 1) with a scale parameter σ , itself distributed according to a distribution G . The distribution of errors, H , is then a *scale mixture distribution* with probability density function

$$p_H(E) = \int_0^\infty p_D(E|\sigma) p_G(\sigma) d\sigma \quad (3)$$

Example - the NIG distribution. For a normal distribution $D = N(0, \sigma)$ and a distribution of variances described by an inverse gamma distribution $\sigma^2 \sim \Gamma^{-1}(\frac{\nu}{2}, \frac{\nu}{2})$, the compound distribution H is a Student's- t distribution with ν degrees of freedom, noted $t(\nu)$. This is a special case of the so-called Normal Inverse Gamma (NIG) distribution used, for instance, in evidential inference¹⁰. □

When D is unknown, there is no evidence that it should be uniform across data space. To avoid unlimited complexity, this hypothesis will be made in the following, without affecting the main conclusions of the study.

B. Calibration statistics derived from the generative model

The generative model described above can be used to derive two families of calibration statistics.

1. The calibration error and related statistics.

The variance of the compound distribution of errors is obtained by the *law of total variance*, i.e.

$$\text{Var}_H(E) = \langle \text{Var}_D(E|\sigma) \rangle_G + \text{Var}_G(\langle E|\sigma \rangle_D) \quad (4)$$

$$= \langle u_E^2 \rangle + \text{Var}_G(\langle E|\sigma \rangle_D) \quad (5)$$

where the first RHS term of Eq. 4 has been identified as the mean squared uncertainty $\langle u_E^2 \rangle$. This expression can be compared to the standard expression for the variance

$$\text{Var}(E) = \langle E^2 \rangle - \langle E \rangle^2 \quad (6)$$

For an unbiased error distribution, one gets $\text{Var}_G(\langle E | \sigma \rangle_D) = 0$ and $\langle E \rangle = 0$, leading to

$$\langle E^2 \rangle = \langle u_E^2 \rangle \quad (7)$$

on which some popular calibration statistics are based.

Example, followed. Considering the NIG model, one can easily verify from the properties of the Student's- t distribution that

$$\langle E^2 \rangle = \sigma^2 \nu / (\nu - 2) \quad (8)$$

(using $\langle E \rangle = 0$), and from the Inverse Gamma distribution¹¹ that

$$\langle u_E^2 \rangle = \sigma^2 \nu / (\nu - 2) \quad (9)$$

so that Eq. 7 is theoretically fulfilled. \square

Based on Eq. 7, the Relative Calibration Error is defined as

$$RCE = \frac{RMV - RMSE}{RMV} \quad (10)$$

where $RMSE$ is the root mean squared error $\sqrt{\langle E^2 \rangle}$ and RMV is the root mean variance ($\sqrt{\langle u_E^2 \rangle}$). The RCE has been shown to be very sensitive to the presence of heavy tails in the uncertainty and error distributions and to be unreliable for a large portion of the studied ML-UQ datasets¹². It is therefore not considered directly here, but it is used in a bin-based statistic of *conditional* calibration, the Expected Normalized Calibration Error³

$$ENCE = \frac{1}{N} \sum_{i=1}^N |RCE_i| \quad (11)$$

where RCE_i is estimated over the data within bin i . According to the binning variable, the ENCE might be used to test *consistency* (binning on u_E) or *adaptivity* (binning on input features).¹³

Pernot⁴ has shown that the ENCE reference value is not zero, and that it depends on the binning scheme. As demonstrated in Appendix B, the ENCE depends on the bins size and does not have a predefined reference value.

In case of heavy-tailed uncertainty distributions, the ENCE is not expected to suffer from the same problem as the RCE when the binning variable is the uncertainty¹². However, it shares the same sensitivity in the case of heavy-tailed error distributions or when the binning variable is not the uncertainty.

2. The ZMS and related statistics.

Another approach to calibration based on Eq. 1 uses scaled errors or z-scores

$$Z_i = \frac{E_i}{u_{E_i}} \sim D(0, 1) \quad (12)$$

with the property

$$\text{Var}(Z) = 1 \quad (13)$$

assessing average calibration for unbiased errors^{2,14}. For biased errors, the calibration equation becomes

$$\text{ZMS} = \langle Z^2 \rangle = 1 \quad (14)$$

which is the preferred form for testing¹³ (ZMS stands for z-score's mean squares). This choice is motivated by the use of the ZMS for binned data, where, even for an unbiased dataset, one should not expect every bin to be unbiased. The ZMS does not depend on a distribution hypothesis and its target value is 1.

The negative log-likelihood (NLL) score can be written as¹⁵

$$\text{NLL} = \frac{1}{2} \left(\langle Z^2 \rangle + \langle \ln u_E^2 \rangle + \ln 2\pi \right) \quad (15)$$

It combines an *average calibration* term, the ZMS,¹⁶ and a *sharpness* term driving the uncertainties towards small values¹⁷ when the NLL is used as a loss function, hence preventing the minimization of $\langle Z^2 \rangle$ by arbitrary large uncertainties. The NLL is the logarithm of a normal probability density function and therefore should be used only when the errors and uncertainties are linked by a standard normal generative distribution ($D = N(0, 1)$). Knowing the reference value of $\langle Z^2 \rangle$, one can assign a reference value to the NLL

$$\text{NLL}_{ref} = \frac{1}{2} \left(1 + \langle \ln u_E^2 \rangle + \ln 2\pi \right) \quad (16)$$

Note that Rasmussen *et al.*⁶ treat the NLL as if it had no predefined reference value and needed a simulated reference. This is not the case when the NLL is defined by Eq. 15, but it might be for other likelihood definitions. In the present case, for a given set of uncertainties, validation of the NLL is equivalent to the validation of the ZMS, with the additional constraint of a normal generative model.

By analogy with the ENCE, one can define a ZMS-based mean calibration error¹⁸

$$\text{ZMSE} = \frac{1}{N} \sum_{i=1}^N |\ln \text{ZMS}_i| \quad (17)$$

where i runs over the N bins and ZMS_i is estimated with the data in bin i . The logarithm accounts for the fact that ZMS is a scale statistic (a ZMS of 2 entails a deviation of the same amplitude as a ZMS of 0.5). As for the ENCE, the ZMSE measures conditional calibration, i.e. consistency if the binning variable is the uncertainty or adaptivity if it is an input feature. Also, the ZMSE depends on the bins size (Appendix B), and does not have a predefined reference value. As observed for the ZVE defined by Pernot⁴, one might expect the ZMSE to be more reliable than the ENCE for heavy-tailed uncertainty distributions, but it shares its sensitivity to heavy-tailed error distributions¹².

C. Correlation

The correlation between absolute errors and uncertainties

$$CC = \text{cor}(|E|, u_E) \quad (18)$$

is used to assess the strength of the link between these variables¹. One expects large absolute errors to be associated with large uncertainties and small uncertainties to be associated with small absolute errors. However, the link is not symmetric, as small absolute errors might be associated with large uncertainties. To account for a possible non linear relation and to reduce the sensitivity to outliers, *Spearman's rank correlation coefficient* is recommended to estimate CC.

A positive value of CC is desirable, but, considering the probabilistic link between errors and uncertainties, its reference value cannot be 1.² In absence of a specified target value, the CC alone should not be used in comparative studies, nor for validation.

D. Validation

The validation protocol has been presented in Ref¹². The main points are summarized here. For a given dataset (E, u_E) and statistic ϑ , one estimates the statistic over the dataset ϑ_{est} and a 95 % confidence interval $I_{BS} = [I_{BS}^-, I_{BS}^+]$ by bootstrapping using the Bias Corrected Accelerated (BC_a) method¹⁹. One can then test that the target value for the statistic, ϑ_{ref} , lies within I_{BS} , i.e.

$$\vartheta_{ref} \in [I_{BS}^-, I_{BS}^+] \quad (19)$$

For a non-binary agreement measure, a standardized score ζ is defined as

$$\zeta(\vartheta_{est}, \vartheta_{ref}, I_{BS}) = \begin{cases} \frac{\vartheta_{est} - \vartheta_{ref}}{I_{BS}^+ - \vartheta_{est}} & \text{if } (\vartheta_{est} - \vartheta_{ref}) \leq 0 \\ \frac{\vartheta_{est} - \vartheta_{ref}}{\vartheta_{est} - I_{BS}^-} & \text{if } (\vartheta_{est} - \vartheta_{ref}) > 0 \end{cases} \quad (20)$$

that can be tested by

$$|\zeta(\vartheta_{est}, \vartheta_{ref}, I_{BS})| \leq 1 \quad (21)$$

which is equivalent to the interval test (Eq. 19). In addition, ζ provides valuable information about the sign and amplitude of the agreement between the statistic and its reference value.

The validation procedure depends then on the availability of ϑ_{ref} , as shown next.

1. *Predefined ϑ_{ref}*

If ϑ_{ref} is known, one can directly use Eq. 20 to estimate

$$\zeta_{BS} = \zeta(\vartheta_{est}, \vartheta_{ref}, I_{BS}) \quad (22)$$

which will be considered below as the benchmark method against which the alternative simulation-based methods will be evaluated.

2. *Estimation of ϑ_{ref} by simulation*

For those statistics without a predefined reference value, one has to make an hypothesis on D , in order to generate ϑ_{ref} from ideally calibrated datasets:

1. Choose a unit-variance generative distribution $D(0, 1)$.
2. Draw a set of pseudo-errors from the actual uncertainties by applying Eq. 1 and estimate the corresponding statistic $\tilde{\vartheta}_D$, where the subscript D denotes the choice of generative distribution.
3. Repeat step 2 N_{MC} times (Monte Carlo sampling).
4. Take the mean value $\tilde{\vartheta}_{D,ref} = \langle \tilde{\vartheta}_D \rangle$ and estimate the *standard error* on $\tilde{\vartheta}_{D,ref}$: $u(\tilde{\vartheta}_{D,ref}) = \text{sd}(\tilde{\vartheta}_D) / \sqrt{N_{MC}}$, where $\text{sd}(x)$ is the standard deviation of variable x .

For N_{MC} on the order of 10^4 , one should have $2u(\tilde{\vartheta}_{D,ref}) \ll U_{BS}$, where U_{BS} is the half range of I_{BS} , and Eq. 20 can be applied without accounting for the uncertainty on $\tilde{\vartheta}_{D,ref}$. One can then estimate a ζ -score

$$\zeta_{SimD} = \zeta(\vartheta_{est}, \tilde{\vartheta}_{D,ref}, I_{BS}) \quad (23)$$

Note that this approach differs from the one proposed previously in the literature^{5,6}, where the value of the statistic was considered as fixed and the reference value as a random variable with uncertainty $\text{sd}(\tilde{\vartheta}_D)$. This scenario is implemented for comparison, with

$$\zeta_{\text{Sim2D}} = \zeta(\vartheta_{\text{est}}, \tilde{\vartheta}_{D,\text{ref}}, I_D) \quad (24)$$

using a confidence interval I_D estimated from the the quantiles of the simulated sample of $\tilde{\vartheta}_D$.

It is important to acknowledge that validation tests based on the latter approach present two drawbacks when compared to the bootstrapping-based procedure: (1) they ignore uncertainty on ϑ_{est} , creating an asymmetric treatment of the statistics, depending on the existence or not of a known reference value; (2) the limits of the confidence interval I_D can be very sensitive to the choice of generative distribution D ,⁵ as will be illustrated in Sect. III.

3. Recommended validation workflow

A recommended validation workflow for a given statistic ϑ is shown in Fig. 1. Note that it is essential that ϑ should be first confirmed as being fit to the purpose of the study, notably for datasets with highly skewed and/or heavy-tailed uncertainty and/or error distributions¹².

For statistics with a known reference value, one can apply directly the benchmark method (Eq. 22). In the absence of a predefined reference value for a statistic, one might generate a simulated one (Sect. IID 2), but a crucial point of this method is the choice of a generative distribution D linking the errors to the uncertainties (Eq. 1).

If D is well constrained, one may proceed to the estimation of the simulated reference value $\tilde{\vartheta}_{D,\text{ref}}$ and use it for validation (Eq. 23). In absence of constraints on D , it is essential to estimate the sensitivity of $\tilde{\vartheta}_{D,\text{ref}}$ to D . For this, at least two alternative shapes for D have to be considered, for instance $D = N(0, 1)$ and $D = t_s(\nu)$, where the unit-variance Student's- t distribution is defined as

$$t_s(\nu) = t(\nu) / \sqrt{\nu/(\nu - 2)} \quad (25)$$

One should use a value of ν small enough to provide a contrast with the normal distribution, but not too small, as it might generate problematic error sets with very heavy tails and many outliers. From numerical experiments in a former study¹², a value of $\nu = 6$ is found to be a good compromise. If the simulated reference values for these choices of D differ more than their statistical uncertainty, the statistic is deemed to be over-sensitive to D and should




Figure 1. Flowchart for the validation of a statistic ϑ . ϑ_{ref} is the reference value used for validation, ϑ_{est} is the actual value of the statistic, I_{BS} is the bootstrapped CI for ϑ_{est} and D is the error generative distribution.

not be used for validation. Otherwise, one might proceed as in the case of a predefined D (Eq. 23).

III. APPLICATIONS

The validation approach presented above is applied to nine datasets extracted from the ML-UQ literature, with a focus on the shape of z-scores distributions and its consequences on the validation process. This is followed by a thorough study of the sensitivity of the validation procedure to various parameters, such as the uncertainty distribution and the

Set #	Name	Size (M)	$\beta_{GM}(u_E^2)$	$\beta_{GM}(E^2)$	$\beta_{GM}(Z^2)$
1	Diffusion_RF ²⁴	2040	0.40	0.82	0.73
2	Perovskite_RF ²⁴	3834	0.72	0.94	0.83
3	Diffusion_LR ²⁴	2040	0.66	0.74	0.69
4	Perovskite_LR ²⁴	3836	0.74	0.82	0.69
5	Diffusion_GPR_Bayesian ²⁴	2040	0.19	0.78	0.79
6	Perovskite_GPR_Bayesian ²⁴	3818	0.50	0.96	0.95
7	QM9_E ⁹	13885	0.93	0.98	0.78
8	logP_10k_a_LS-GCN ⁶	5000	0.30	0.79	0.78
9	logP_150k_LS-GCN ⁶	5000	0.30	0.77	0.75

Table I. The nine datasets used in this study: number, name, size and reference.

shape of the generative distribution.

A. The datasets

Nine test sets with *a priori* calibrated uncertainties have been extracted from the recent ML-UQ literature for the prediction of physico-chemical properties by a diverse panel of ML and UQ methods. These datasets were tested by Pernot¹² for average calibration by the RCE and ZMS statistics. The characterization of the squared uncertainty and squared error distributions by robust skewness and kurtosis statistics (β_{GM} and κ_{CS})²⁰⁻²³ was used to screen heavy-tailed distributions likely to cause reliability problems in the statistical validation of RCE and ZMS. It was found that β_{GM} and κ_{CS} were mostly redundant for these datasets, and only β_{GM} is reported in Table I. β_{GM} is robust to outliers, varies between -1 and 1 and is null for symmetric distributions. For u_E^2 distributions, an upper safety limit of 0.6 for β_{GM} was established based on an Inverse Gamma distribution model, while for E^2 and Z^2 this limit is 0.8, based on a Fisher distribution model¹². Values exceeding these limits are noted in bold face in Table I.

In this previous study¹², the analysis of u_E^2 and E^2 distributions enabled also to show that applying the generative model (Eq. 1) led to reject the normality of D for all the studied datasets. Typically, the errors are much more dispersed than what would be expected from a normal generative distribution. As this is a central point for the estimation of simulated reference values, the distribution of Z , which should reflect the empirical shape of D for calibrated datasets is analyzed next.

Set #	μ_Z	σ_Z	b_Z (%)	ν_Z	ν_{Z^2}
1	-0.027(22)	0.980(30)	3	6.0	7.9
2	-0.018(15)	0.940(26)	2	3.3	4.9
3	0.002(23)	1.058(18)	0	20.1	15.1
4	-0.021(18)	1.107(16)	2	9.1	8.2
5	0.006(20)	0.920(21)	1	3.9	2.7
6	-0.005(16)	0.992(37)	1	1.4	0.9
7	0.0174(84)	0.9858(99)	2	4.4	4.0
8	0.050(14)	0.961(16)	5	3.9	3.7
9	-0.260(13)	0.951(23)	27	3.1	20.2

Table II. Summary statistics for the z-scores (Z): μ_Z is the mean value, σ_Z the standard deviation, $b_Z = 100(\mu_Z - 0)/\sigma_Z$ measures relative bias and ν_Z is the number of degrees of freedom resulting from the fit of the data by a scaled and shifted Student's- t distribution. The last column reports the shape parameter for the fit of Z^2 distributions by a $F(1, \nu_{Z^2})$ distribution¹².

1. Distributions of z-scores

The fit of z-scores distributions by a scaled-and-shifted Student's- t distribution is done by maximum likelihood estimation²⁵. Table II reports the mean values, standard deviations, relative bias and the number of degrees of freedom ν_Z . The smaller ν_Z , the farther one is from a normal distribution. Note that this fit accounts for a possible bias in the dataset, which is non-negligible for Sets 8 and 9 (above 5% of the standard deviation). The values of ν_Z reported here are globally consistent with the ones obtained previously¹² for the fit of the distributions of Z^2 by a Fisher-Snedecor $F(1, \nu_{Z^2})$ distribution, except for Set 9 for which the bias was not taken into account. Fig 2 shows the comparison of the best fits of z-scores by normal and scaled-and-shifted Student's- t distributions.

The only sets for which one gets close to normality are Set 3, with $\nu_Z = 20.1$, and to a lesser degree Set 4, with $\nu_Z = 9.1$. Overall, one has rather small ν_Z values, rejecting unambiguously the normality of z-scores for 7 or 8 out of nine datasets. As a side effect, let us note that the normality of z-scores should not be used as a calibration criterion, unless D is known to be normal.




Figure 2. Z-scores distributions (histograms) with normal (red line) and scaled and shifted Student's- t (blue line) fits. For legibility, the histograms have been truncated to ± 3 standard deviations, hiding a few outlying values.

B. Analysis of the generative model

Considering the generative model, the distribution of z-scores *for calibrated datasets* should be identical to D . To check how close the distribution of z-scores is to the errors generative distribution, simulated error distributions have been generated using the actual uncertainties and two generative distributions: a standard Normal distribution $N(0, 1)$ and a unit-variance Student's- t distribution $t_s(\nu_Z)$ with the degrees of freedom reported in Table II. The results are shown in Fig. 3.

The t_s -based model offers a better fit to the errors histogram for all Sets except 3 and 4,




Figure 3. Recovery of the error distributions with the generative model, using a normal distribution (red line) or a unit-variance Student’s distribution $t_s(\nu_Z)$ (ν_Z from Table II; blue line). For Set 6, the Student’s- t model cannot be generated because of the infinite variance for $\nu_Z < 2$ and it was replaced by $\nu_Z = 2.1$.

for which it is on par with the normal model. Note that for Set 6 the value of ν_Z had to be increased to 2.1 to avoid the infinite variance due to $\nu_Z \leq 2$.

It is thus clear that a normal generative distribution $D = N(0,1)$ is not likely to provide simulated error distributions with properties close to the actual errors, except for a few datasets.

C. Sensitivity analysis

In the previous section, we have seen that if there is a best choice for the generative distribution D , it is rarely the normal distribution. In absence of strong constraints to guide this choice, it is important to assess the sensitivity to D of the simulated reference values for the candidate calibration statistics, and more globally the sensitivity of the validation ζ -scores.

1. Sensitivity of $\tilde{\vartheta}_{D,ref}$ to the uncertainty distribution

A first point is the appreciation of the dependence of the simulated reference values on the uncertainty distribution. For this, a normal generative distribution is chosen [$D = N(0, 1)$], and $\tilde{\vartheta}_{D,ref}$ and $u(\tilde{\vartheta}_{D,ref})$ are estimated for each statistic and the nine example datasets. The sampling size is $N_{MC} = 10^4$ and u_E is sorted and parted into 50 equal-size bins to estimate the ENCE and ZMSE statistics. The results are presented in Fig. 4.

For the ZMS [Fig. 4(a)], the simulation recovers trivially the predefined reference value, as the simulated z-scores are directly samples of the D distribution. For CC [Fig. 4(b)], $\tilde{\vartheta}_{D,ref}$ is seen to depend strongly on the dataset, confirming the absence of a common reference value for these datasets. The ENCE [Fig. 4(c)] and ZMSE [Fig. 4(e)] follow a parallel pattern, which is mostly due to the sensitivity of these variables to the dataset size, M (Appendix B). The correction of this trend by multiplying the statistics by $M^{1/2}$ [Fig. 4(d,f)] shows that the common pattern disappears and that each uncertainty set leads to its own reference value, even if a few CIs overlap. The ENCE has an outstanding value of $\tilde{\vartheta}_{D,ref}$ for Set 7, which is not the case for the ZMSE, and is certainly a symptom of the stratification of the uncertainties resulting from an isotonic regression post-hoc calibration^{13,15,26}.

Globally, apart from the ZMS, one needs therefore to estimate a simulated reference value for each set and statistic.

2. Sensitivity of $\tilde{\vartheta}_{D,ref}$ to D

The simulated reference values are generated for Sets 7 and 8 (as representative of the largest sets with markedly different uncertainty distributions) from their actual uncertainties by using a unit-variance Student's- t generative distribution $D = t_s(\nu)$ for a range of degrees of freedom between 3 and 20, covering the range of values observed for ν_Z (Table II). The sampling size is $N_{MC} = 10^4$ and u_E is sorted and parted into 50 equal-size bins to estimate




Figure 4. Sensitivity of $\tilde{\vartheta}_{D,ref}$ to the dataset for the ZMS (a), CC (b), ENCE (c, d) and ZMSE (e, f) statistics. The generative distribution is standard normal, $D = N(0, 1)$.

the ENCE, ZMSE. The results are presented in Fig. 5.

For ZMS [Fig. 5(a)], one expects to observe the conformity of the simulated reference value with the known reference value (i.e. $\tilde{\vartheta}_{v,ref} \simeq 1$), independently of the chosen generative distribution. This is the case, albeit with a notable increase of Monte Carlo sampling uncertainty as v decreases. For CC [Fig. 5(b)], the sensitivity to v depends on the value of the statistic, and it is less marked for low CC values (Set 8) than for larger ones (Set 7). For the ENCE [Fig. 5(c)] and ZMSE [Fig. 5(d)] statistics, there is no ambiguity, as the simulated ref-




Figure 5. Sensitivity of the simulated reference values to the generative distribution. Calibrated datasets are generated from the actual uncertainties of Sets 7 and 8, using a generative distribution $D = t_s(\nu)$ with ν degrees of freedom.

reference values depend strongly on ν , with a similar behavior for both statistics and datasets.

This sensitivity analysis is based on a worst case scenario where the lower values of ν generate amounts of outliers that can disturb the reliability of calibration statistics. It suggests that using too extreme generative distributions D for the sensitivity analysis of those statistics without reference values might be counterproductive. For instance, a Student's- t distribution with defined skewness and kurtosis requires $\nu \geq 5$, and in the following one will use $t_s(\nu = 6)$ as an alternative to the normal distribution.

Ignoring small values of ν , one still sees that the CC, ENCE and ZMSE statistics can hardly be useful in a validation context when the generative distribution D is unknown.

3. Sensitivity of ζ -scores to D

We now consider how the sensitivity of $\tilde{\vartheta}_{D,ref}$ to D propagates to the ζ -scores. For the simulated reference values, two options for D are considered: a standard normal ($\zeta_{SimN}, \zeta_{Sim2N}$) and unit-variance Student's- t distribution with 6 degrees of freedom $t_s(\nu = 6)$ ($\zeta_{SimT}, \zeta_{Sim2T}$). This value has been chosen to avoid the perturbation of the statistics by extreme values as observed in the previous section.

The results for the ZMS, CC, ENCE and ZMSE statistics and the pertinent simulation scenarios are reported in Figs. 6-7. The intermediate variables for the calculation of these ζ -scores are reported in Appendix A.

Let us first consider the results for the ZMS statistic [Figs. 6(a)]. There is a good agreement between the three bootstrapped CI-based estimations (BS, SimN, SimT). The absence of notable difference between the SimN and SimT results confirms the low sensitivity of this method to the choice of D . It is also clear that the Sim2N and Sim2T scores can differ notably from the other scores and between them, confirming a high sensitivity of the Monte Carlo CI I_D to D . Sets 1-6 have been analyzed in an earlier study¹³, by using the $\text{Var}(Z)$ score and an approximate analytical estimation of confidence intervals²⁷. The validation results differ at the margin, as the approximate method used previously seems to provide slightly wider CIs than the bootstrapped ones used here, and is therefore less stringent.

For the CC statistic [Figs. 6(b)], it appears that the choice of D has always a sizable effect, even for the Sim protocol. In consequence, and unless one trusts more one of the two options for D , the CC statistic cannot be reliably validated for the studied datasets. Note that, by chance and despite the amplitude of the D -induced discrepancies, all the CC validation statistics for a given dataset agree on the binary validation diagnostic. One might be then tempted to conclude that only Sets 3 and 4 have CCs compatible with their simulated reference values. Note that in this case, the Sim and Sim2 ζ -scores for a given choice of D are very similar.

The ζ -scores for the ENCE and ZMSE statistics have been estimated for $N = 20$ bins and present very similar features (Fig.7): a notable sensitivity to D and a large difference between the Sim and Sim2 protocols (except for Set 3). The latter observation confirms that one should not use the Monte Carlo CIs to estimate the ζ -scores. Focusing on the SimN and SimT results, it appears that they globally agree to reject consistency, except for Set 3 where consistency would be validated by SimT, as observed in Appendix C. However, we have shown above that for Sets 3 and 4 the SimN scenario is the most plausible.




Figure 6. Validation of the ZMS and CC statistics with ζ -scores obtained by bootstrapping (BS) and by simulation with two hypotheses on the errors distribution D : Normal (SimN) and $t_s(\nu = 6)$ (SimT), compared to the Monte Carlo approach of Rasmussen *et al.*⁶ (Sim2N and Sim2T). In absence of a predefined reference value, the BS approach is not available for CC.

It is somewhat striking that the validation scores of ENCE and ZMSE are so similar, despite notable differences in the reliability of the RCE and ZMS statistics exposed previously¹². In fact, it was shown that both statistics shared a similar sensitivity to the tailedness of error distributions, unlike their sensitivity to the tailedness of uncertainty distributions, where only the RCE could be notably biased. Binning along u_E is expected to erase the tailedness problem of (binned) uncertainty distributions, placing both ENCE and ZMSE on an equal footing. This would not be the case for another binning variable.




Figure 7. Same as Fig. 6 for the ENCE and ZMSE consistency scores. In absence of a predefined reference value, the BS approach is not available for these statistics.

IV. DISCUSSION

This study focuses on the validation of correlation and calibration statistics by standardized ζ -scores, for which one needs reference values and confidence interval estimates. Reference values are predefined for the ZMS statistic, but not for CC, ENCE and ZMSE. In the latter cases, it is possible to generate reference values by a Monte Carlo approach, which implies to generate simulated error sets from the actual uncertainties, using a generative distribution D . A reference value and its standard error can be obtained by repeating this random sampling. The choice of an interval-based validation method results from the often marked non-normality of the error and uncertainty datasets, which prevents the use of more standard z-scores or t -scores. CIs are therefore estimated by bootstrapping for all statistics,

by taking care that the bootstrap distributions can be non-normal and asymmetric.

In absence of a predefined reference, Pernot⁵ and Rasmussen *et al.*⁶ proposed to use the standard deviation or CI of the simulated reference value as a metric to compare the estimated statistic to its reference value. This study shows unambiguously that several disadvantages come with this approach: (i) it introduces an asymmetry of treatment between the statistics with known and unknown reference values, using bootstrapping-based uncertainties for the former and simulation-based uncertainties for the latter⁶; (ii) the standard deviation (or CI) of the simulated reference values is strongly dependent on the chosen generative distribution D ; and (iii) in such cases, validation becomes dependent on the often *ad hoc* choice of D .

To circumvent these issues, it is suggested here to *use the bootstrapping-based intervals independently of the existence of a reference value, and to characterize the simulated reference by its standard error* (standard deviation of the mean), which is much less sensitive to the choice of D . Moreover, with a moderate effort on the Monte Carlo sampling, it is possible to ensure that this standard error becomes negligible before the bootstrapping-based uncertainty, which solves the problem of the dependence of the ζ -scores on the simulated reference uncertainty. The impact of this approach has been shown by comparing SimN to SimT and Sim2N to Sim2T (Figs.6-7): in all cases the proposed approach reduces the sensitivity to D , except for CC where it is about identical.

Another and less manageable source of sensitivity of the ζ -scores to D comes with the simulated reference value itself. For instance, the mean value of any statistic that involves a sign loss of the simulated errors (e.g. absolute value, squared value...) is likely to be affected by the shape/width of the errors distribution. This is particularly visible for the CC, ENCE and ZMSE statistics, but much less for the ZMS. For confidence curves, Pernot⁵ has shown that the simulated reference based on the RMS of the errors was insensitive to D , which was not the case for the MAE. *The sensitivity of a simulated reference value to the generative model has therefore to be directly assessed before using it for validation.*

V. CONCLUSION

This study sheds a critical look at the validation of an average calibration statistics (ZMS), a correlation statistic (CC) and two conditional calibration statistics (ENCE, ZMSE). A benchmark method was defined for those statistics with a predefined reference value (ZMS) and compared to simulation approaches for statistics with no predefined reference

value (CC, ENCE and ZMSE). A validation workflow was proposed to deal with the main obstacles of the simulation approach (Fig. 1) and notably with the sensitivity of simulated reference values to the generative distribution D used to generate ideal synthetic errors from actual uncertainties.

Important conclusions about the studied statistics have been obtained from a series of ML-UQ datasets from the literature.

- *The generative distribution D is not necessarily normal.* The error distributions obtained by the generative model with a normal distribution D (Eq. 1) rarely agree with the actual error distributions. Better fits are often obtained by using the distribution of scaled errors (z-scores) as a proxy. In some instances, e.g. Bayesian Neural Networks or Gaussian Processes, the generative distribution is prescribed, but this is not generally the case. Thus, before using an hypothetical generative distribution to simulate unknown reference values, one should test its pertinence for the dataset(s) under scrutiny.
- *The use of simulated reference values for the CC, ENCE and ZMSE statistics is not reliable if D is unknown.* It has been shown that the simulated reference values for these statistics are highly sensitive to the choice of generative distribution D . If D is unknown, one cannot estimate a reliable reference value for their validation. For CC, checking that it is positive is the best that can be done and there is not much sense in comparing CC values of different datasets (the largest value is not necessarily the better!). A much more powerful and reliable approach is the plotting of RMSE-based *confidence curves* with bootstrapped CI and simulated reference value. Similarly, the simulated reference values for measures such as the ENCE and ZMSE are very sensitive to the choice of D and therefore cannot generally be used for validation of conditional calibration. When used for comparative studies, they should be limited to datasets of identical sizes and to identical binning schemes. Validation is possible by alternative approaches, such as the “extrapolation to zero bins” method used in Appendix C, or maybe by the estimation of the “fraction of valid bins” proposed by Pernot¹³. However, the reliability of the latter approach still needs to be demonstrated for a diversity of datasets.

Finally, the only case where the use of simulated reference values seems well adapted is for confidence curves, as the RMSE-based reference curve does not depend on D . However, there are probably other calibration statistics for which this concept might be useful, and the proposed workflow should enable to validate them with confidence.

ACKNOWLEDGMENTS

I warmly thank J. Busk for providing the QM9 dataset.

AUTHOR DECLARATIONS

Conflict of Interest

The author has no conflicts to disclose.

CODE AND DATA AVAILABILITY

The code and data to reproduce the results of this article are available at https://github.com/ppernot/2024_SimRef/releases/tag/v1.0 and at Zenodo (<https://doi.org/10.5281/zenodo.10730985>).

REFERENCES

- ¹M. Tynes, W. Gao, D. J. Burrill, E. R. Batista, D. Perez, P. Yang, and N. Lubbers. [Pairwise difference regression: A machine learning meta-algorithm for improved prediction and uncertainty quantification in chemical search](#). *J. Chem. Inf. Model.*, 61:3846–3857, 2021. PMID: 34347460.
- ²P. Pernot. [Prediction uncertainty validation for computational chemists](#). *J. Chem. Phys.*, 157:144103, 2022.
- ³D. Levi, L. Gispan, N. Giladi, and E. Fetaya. [Evaluating and Calibrating Uncertainty Prediction in Regression Tasks](#). *Sensors*, 22:5540, 2022.
- ⁴P. Pernot. [Properties of the ENCE and other MAD-based calibration metrics](#). *arXiv:2305.11905*, May 2023.
- ⁵P. Pernot. [Confidence curves for UQ validation: probabilistic reference vs. oracle](#). *arXiv:2206.15272*, June 2022.
- ⁶M. H. Rasmussen, C. Duan, H. J. Kulik, and J. H. Jensen. [Uncertain of uncertainties? A comparison of uncertainty quantification metrics for chemical data sets](#). *J. Cheminf.*, 15:1–17, December 2023.
- ⁷K. Tran, W. Neiswanger, J. Yoon, Q. Zhang, E. Xing, and Z. W. Ulissi. [Methods for comparing uncertainty quantifications for material property predictions](#). *Mach. Learn.: Sci. Technol.*, 1:025006, 2020.

- ⁸BIPM, IEC, IFCC, ILAC, ISO, IUPAC, IUPAP, and OIML. [Evaluation of measurement data - Guide to the expression of uncertainty in measurement \(GUM\)](#). Technical Report 100:2008, Joint Committee for Guides in Metrology, JCGM, 2008. URL: http://www.bipm.org/utils/common/documents/jcgm/JCGM_100_2008_F.pdf.
- ⁹J. Busk, P. B. Jørgensen, A. Bhowmik, M. N. Schmidt, O. Winther, and T. Vegge. [Calibrated uncertainty for molecular property prediction using ensembles of message passing neural networks](#). *Mach. Learn.: Sci. Technol.*, 3:015012, 2022.
- ¹⁰A. Amini, W. Schwarting, A. Soleimany, and D. Rus. [Deep Evidential Regression](#). *arXiv:1910.02600*, October 2019.
- ¹¹M. Evans, N. Hastings, and B. Peacock. *Statistical Distributions*. Wiley-Interscience, 3rd edition, 2000.
- ¹²P. Pernot. [Negative impact of heavy-tailed uncertainty and error distributions on the reliability of calibration statistics for machine learning regression tasks](#). *arXiv:2402.10043*, February 2024. URL: <https://arxiv.org/abs/2402.10043>.
- ¹³P. Pernot. [Calibration in machine learning uncertainty quantification: Beyond consistency to target adaptivity](#). *APL Mach. Learn.*, 1:046121, 2023.
- ¹⁴P. Pernot. [The long road to calibrated prediction uncertainty in computational chemistry](#). *J. Chem. Phys.*, 156:114109, 2022.
- ¹⁵J. Busk, M. N. Schmidt, O. Winther, T. Vegge, and P. B. Jørgensen. [Graph neural network interatomic potential ensembles with calibrated aleatoric and epistemic uncertainty on energy and forces](#). *Phys. Chem. Chem. Phys.*, 25:25828–25837, 2023.
- ¹⁶W. Zhang, Z. Ma, S. Das, T.-W. Weng, A. Megretski, L. Daniel, and L. M. Nguyen. [One step closer to unbiased aleatoric uncertainty estimation](#). *arXiv:2312.10469*, December 2023.
- ¹⁷T. Gneiting, F. Balabdaoui, and A. E. Raftery. [Probabilistic forecasts, calibration and sharpness](#). *J. R. Statist. Soc. B*, 69:243–268, 2007.
- ¹⁸P. Pernot. [Can bin-wise scaling improve consistency and adaptivity of prediction uncertainty for machine learning regression ?](#) *arXiv:2310.11978*, October 2023.
- ¹⁹T. J. DiCiccio and B. Efron. [Bootstrap confidence intervals](#). *Statist. Sci.*, 11:189–212, 1996. URL: <https://www.jstor.org/stable/2246110>.
- ²⁰E. L. Crow and M. M. Siddiqui. [Robust estimation of location](#). *J. Am. Stat. Assoc.*, 62:353–389, 1967.
- ²¹R. A. Groeneveld and G. Meeden. [Measuring skewness and kurtosis](#). *The Statistician*, 33:391–399, 1984. URL: <http://www.jstor.org/stable/2987742>.

- ²²M. Bonato. Robust estimation of skewness and kurtosis in distributions with infinite higher moments. *Financ Res Lett*, 8:77–87, 2011.
- ²³P. Pernot and A. Savin. Using the Gini coefficient to characterize the shape of computational chemistry error distributions. *Theor. Chem. Acc.*, 140:24, 2021.
- ²⁴G. Palmer, S. Du, A. Politowicz, J. P. Emory, X. Yang, A. Gautam, G. Gupta, Z. Li, R. Jacobs, and D. Morgan. Calibration after bootstrap for accurate uncertainty quantification in regression models. *npj Comput. Mater.*, 8:115, 2022.
- ²⁵M. L. Delignette-Muller and C. Dutang. *fitdistrplus: An R package for fitting distributions*. *J Stat Softw*, 64(4):1–34, 2015.
- ²⁶P. Pernot. Stratification of uncertainties recalibrated by isotonic regression and its impact on calibration error statistics. *arXiv:2306.05180*, June 2023.
- ²⁷E. Cho, M. J. Cho, and J. Eltinge. The variance of sample variance from a finite population. *Int. J. Pure Appl. Math.*, 21:387–394, 2005. URL: <http://www.ijpam.eu/contents/2005-21-3/10/10.pdf>.

APPENDICES

Appendix A: Results tables for standardized scores

The numerical values necessary to build Fig. 6 are reported here (Tables A1-A4), with a summary of the necessary equations. Each table contains the values for one of the ZMS and CC statistics.

All standardized ζ -scores are built from the equation

$$\zeta(\vartheta_{est}, \vartheta_{ref}, I) = \begin{cases} \frac{\vartheta_{est} - \vartheta_{ref}}{I^+ - \vartheta_{est}} & \text{if } (\vartheta_{est} - \vartheta_{ref}) \leq 0 \\ \frac{\vartheta_{est} - \vartheta_{ref}}{\vartheta_{est} - I^-} & \text{if } (\vartheta_{est} - \vartheta_{ref}) > 0 \end{cases} \quad (\text{A1})$$

where ϑ_{est} is the estimated value of the statistic, ϑ_{ref} is its reference value, and $I = [I^-, I^+]$ is a 95% confidence interval. The values of ϑ_{ref} and I are estimated according to three schemes

Scheme	ϑ_{ref}	I
BS	Predefined	Bootstrapped
Sim	Monte Carlo	Bootstrapped
Sim2	Monte Carlo	Monte Carlo

In the absence of a predefined reference value, ζ_{BS} cannot be estimated.

In the Sim and Sim2 schemes, synthetic error sets are generated according to

$$E_i = u_{E,i} \epsilon_i \quad (\text{A2})$$

where $u_{E,i}$ is one of the uncertainties in the original dataset, and ϵ_i is a random value from the generative distribution D ($\epsilon_i \sim D(0, 1)$). Two variants of D are considered in the present study, denoted N and T:

N: a standard normal distribution $D = N(0, 1)$

T: a unit_variance Student's- t distribution $D = t_s(6)$

One has thus five options for the estimation of standardized scores: ζ_{BS} , ζ_{SimN} , ζ_{Sim2N} , ζ_{SimT} , ζ_{Sim2T} , which are reported in the results tables with intermediate results and in Fig. 6.

$\vartheta = \text{ZMS}$		BS				SimN, Sim2N				SimT, Sim2T					
		ϑ	ϑ_{ref}	b_{BS}	I_{BS}	ζ_{BS}	$\tilde{\vartheta}_{ref}$	$u(\tilde{\vartheta}_{ref})$	ζ_{Sim}	I_{Sim2}	ζ_{Sim2}	$\tilde{\vartheta}_{ref}$	$u(\tilde{\vartheta}_{ref})$	ζ_{Sim}	I_{Sim2}
1	0.96	1	5.3e-04	[0.87, 1.12]	-0.25	1.00	3.1e-04	-0.25	[0.94, 1.06]	-0.66	1.00	5.0e-04	-0.26	[0.91, 1.10]	-0.45
2	0.86	1	2.6e-04	[0.80, 0.99]	-1.09	1.00	2.3e-04	-1.09	[0.96, 1.04]	-2.65	1.00	3.7e-04	-1.09	[0.93, 1.07]	-1.72
3	1.12	1	6.5e-04	[1.05, 1.20]	1.66	1.00	3.2e-04	1.66	[0.94, 1.06]	1.89	1.00	4.9e-04	1.67	[0.91, 1.10]	1.17
4	1.23	1	-2.6e-04	[1.16, 1.30]	3.53	1.00	2.3e-04	3.54	[0.96, 1.04]	5.01	1.00	3.6e-04	3.53	[0.94, 1.08]	3.00
5	0.85	1	5.8e-04	[0.78, 0.92]	-1.99	1.00	3.1e-04	-1.99	[0.94, 1.06]	-2.54	1.00	4.9e-04	-1.98	[0.91, 1.10]	-1.73
6	0.98	1	-9.1e-04	[0.86, 1.15]	-0.10	1.00	2.3e-04	-0.09	[0.95, 1.04]	-0.35	1.00	3.6e-04	-0.10	[0.94, 1.07]	-0.25
7	0.97	1	4.9e-04	[0.94, 1.01]	-0.71	1.00	1.2e-04	-0.71	[0.98, 1.02]	-1.23	1.00	1.9e-04	-0.72	[0.96, 1.04]	-0.78
8	0.93	1	-2.9e-04	[0.87, 0.99]	-1.16	1.00	2.0e-04	-1.16	[0.96, 1.04]	-1.92	1.00	3.1e-04	-1.15	[0.94, 1.06]	-1.27
9	0.97	1	1.5e-03	[0.90, 1.08]	-0.27	1.00	2.0e-04	-0.27	[0.96, 1.04]	-0.74	1.00	3.2e-04	-0.27	[0.94, 1.06]	-0.50

Table A1. Intermediate values for the estimation of ζ -scores for the ZMS statistics.

$\vartheta = CC$		BS				SimN, Sim2N				SimT, Sim2T					
Set #	ϑ	ϑ_{ref}	b_{BS}	I_{BS}	ζ_{BS}	$\tilde{\vartheta}_{ref}$	$u(\tilde{\vartheta}_{ref})$	ζ_{Sim}	I_{Sim2}	ζ_{Sim2}	$\tilde{\vartheta}_{ref}$	$u(\tilde{\vartheta}_{ref})$	ζ_{Sim}	I_{Sim2}	ζ_{Sim2}
1	0.50	n/a	-3.1e-05	[0.467, 0.536]	n/a	0.40	1.9e-04	2.76	[0.38, 0.44]	2.67	0.38	1.9e-04	3.39	[0.34, 0.42]	3.35
2	0.62	n/a	-1.2e-04	[0.598, 0.641]	n/a	0.57	1.1e-04	2.40	[0.55, 0.59]	2.38	0.55	1.1e-04	3.44	[0.52, 0.57]	3.37
3	0.26	n/a	-1.8e-04	[0.216, 0.300]	n/a	0.25	2.1e-04	0.21	[0.21, 0.29]	0.21	0.23	2.1e-04	0.61	[0.19, 0.27]	0.60
4	0.40	n/a	6.6e-05	[0.372, 0.428]	n/a	0.42	1.3e-04	-0.77	[0.40, 0.45]	-0.78	0.40	1.4e-04	0.05	[0.37, 0.43]	0.05
5	0.04	n/a	-1.5e-04	[-0.004, 0.081]	n/a	0.11	2.2e-04	-1.63	[0.07, 0.15]	-1.65	0.10	2.2e-04	-1.45	[0.06, 0.14]	-1.47
6	0.40	n/a	-1.4e-04	[0.373, 0.433]	n/a	0.50	1.2e-04	-3.27	[0.48, 0.52]	-4.10	0.48	1.2e-04	-2.57	[0.46, 0.50]	-3.26
7	0.31	n/a	7.9e-06	[0.297, 0.328]	n/a	0.37	7.2e-05	-3.86	[0.36, 0.39]	-4.27	0.35	7.5e-05	-2.54	[0.34, 0.37]	-2.67
8	-0.03	n/a	-1.2e-05	[-0.052, 0.003]	n/a	0.11	1.4e-04	-4.92	[0.08, 0.14]	-5.06	0.10	1.4e-04	-4.61	[0.08, 0.13]	-4.62
9	0.23	n/a	-1.0e-04	[0.207, 0.258]	n/a	0.13	1.4e-04	3.82	[0.10, 0.16]	3.77	0.12	1.4e-04	4.19	[0.09, 0.15]	4.17

Table A2. Same as Table A1 for the CC statistic.

$\vartheta = ENCE$		BS					SimN, Sim2N					SimT, Sim2T				
		Set #	ϑ	ϑ_{ref}	b_{BS}	I_{BS}	ζ_{BS}	$\tilde{\vartheta}_{ref}$	$u(\tilde{\vartheta}_{ref})$	ζ_{Sim}	I_{Sim2}	ζ_{Sim2}	$\tilde{\vartheta}_{ref}$	$u(\tilde{\vartheta}_{ref})$	ζ_{Sim}	I_{Sim2}
1	0.125	n/a	0.016	[0.084, 0.153]	n/a	0.056	9.5e-05	1.66	[0.038, 0.076]	3.52	0.082	1.5e-04	1.04	[0.055, 0.114]	1.36	
2	0.126	n/a	0.033	[0.096, 0.130]	n/a	0.041	7.0e-05	2.78	[0.028, 0.056]	5.98	0.061	1.1e-04	2.13	[0.041, 0.085]	2.80	
3	0.097	n/a	0.024	[0.074, 0.101]	n/a	0.058	9.8e-05	1.70	[0.040, 0.077]	1.98	0.083	1.5e-04	0.59	[0.056, 0.115]	0.44	
4	0.135	n/a	0.007	[0.103, 0.157]	n/a	0.043	7.4e-05	2.94	[0.029, 0.058]	5.88	0.063	1.2e-04	2.30	[0.043, 0.088]	2.88	
5	0.131	n/a	0.023	[0.101, 0.139]	n/a	0.056	9.4e-05	2.52	[0.039, 0.075]	3.88	0.082	1.5e-04	1.66	[0.055, 0.114]	1.52	
6	0.244	n/a	0.044	[0.156, 0.276]	n/a	0.045	8.1e-05	2.26	[0.030, 0.062]	11.7	0.066	1.3e-04	2.02	[0.044, 0.095]	6.21	
7	0.066	n/a	0.006	[0.045, 0.085]	n/a	0.026	5.3e-05	1.93	[0.017, 0.037]	3.58	0.038	8.3e-05	1.33	[0.025, 0.056]	1.55	
8	0.108	n/a	0.015	[0.077, 0.118]	n/a	0.036	6.0e-05	2.28	[0.025, 0.048]	5.77	0.053	9.7e-05	1.72	[0.036, 0.074]	2.65	
9	0.120	n/a	0.013	[0.082, 0.140]	n/a	0.036	6.0e-05	2.21	[0.025, 0.048]	6.95	0.054	9.7e-05	1.74	[0.036, 0.074]	3.25	

Table A3. Same as Table A1 for the ENCE statistic.

$\vartheta = \text{ZMSE}$		BS					SimN, Sim2N					SimT, Sim2T				
		Set #	ϑ	ϑ_{ref}	b_{BS}	I_{BS}	ζ_{BS}	$\tilde{\vartheta}_{ref}$	$u(\tilde{\vartheta}_{ref})$	ζ_{Sim}	I_{Sim2}	ζ_{Sim2}	$\tilde{\vartheta}_{ref}$	$u(\tilde{\vartheta}_{ref})$	ζ_{Sim}	I_{Sim2}
1	0.255	n/a	0.034	[0.172, 0.299]	n/a	n/a	0.112	1.9e-04	1.71	[0.077, 0.152]	3.63	0.164	2.9e-04	1.09	[0.111, 0.223]	1.54
2	0.273	n/a	0.072	[0.207, 0.283]	n/a	n/a	0.082	1.4e-04	2.91	[0.056, 0.111]	6.62	0.121	2.1e-04	2.31	[0.083, 0.165]	3.46
3	0.173	n/a	0.048	[0.136, 0.180]	n/a	n/a	0.112	1.9e-04	1.64	[0.077, 0.151]	1.57	0.163	2.8e-04	0.26	[0.110, 0.223]	0.16
4	0.247	n/a	0.011	[0.191, 0.287]	n/a	n/a	0.082	1.4e-04	2.97	[0.056, 0.110]	5.81	0.121	2.1e-04	2.26	[0.083, 0.165]	2.87
5	0.283	n/a	0.052	[0.221, 0.304]	n/a	n/a	0.112	1.9e-04	2.77	[0.077, 0.151]	4.42	0.163	2.9e-04	1.94	[0.111, 0.224]	1.96
6	0.356	n/a	0.270	[0.240, 0.357]	n/a	n/a	0.082	1.4e-04	2.37	[0.056, 0.111]	9.53	0.121	2.2e-04	2.02	[0.082, 0.168]	5.06
7	0.118	n/a	0.018	[0.078, 0.131]	n/a	n/a	0.043	7.2e-05	1.87	[0.029, 0.058]	5.07	0.066	1.2e-04	1.31	[0.045, 0.089]	2.20
8	0.225	n/a	0.033	[0.162, 0.246]	n/a	n/a	0.071	1.2e-04	2.43	[0.049, 0.097]	6.08	0.107	1.9e-04	1.87	[0.072, 0.147]	2.96
9	0.250	n/a	0.025	[0.171, 0.287]	n/a	n/a	0.071	1.2e-04	2.27	[0.049, 0.096]	7.32	0.107	1.9e-04	1.81	[0.073, 0.147]	3.62

Table A4. Same as Table A1 for the ZMSE statistic.

Appendix B: Reference values for the ENCE and ZMSE statistic

The ENCE and ZMSE statistics have been shown previously to depend on the binning scheme⁴. We consider here other sources of influence, such as the dataset size, and uncertainty distribution shape. This study is performed on synthetic datasets, and the dependence on the generative model is also assessed.

Simulated reference values $\tilde{\vartheta}_{D,ref}$ for the ENCE and ZMSE statistics have been estimated from $N_{MC} = 5000$ synthetic calibrated datasets generated by the NIG model

$$E \sim u_E \times N(0, 1) \quad (B1)$$

$$u_E^2 \sim \Gamma^{-1}(\nu/2, \nu/2) \quad (B2)$$

with three varying parameters: the dataset size $M \in \{2000, 4000, 8000, 12000, 16000\}$, the number of equal-size bins $N \in \{10, 20, 30, 40, 50\}$ and the shape of the uncertainty distribution $\nu \in \{3, 4, 5, 6, 12, 24\}$. The relative uncertainties on the reported mean reference values are around $1.5 \cdot 10^{-3}$. The results are shown in Fig. A1.

The dependence of the ENCE on the number of bins was shown in a previous study⁴ to be in $N^{1/2}$. The log-log plot in Fig. A1(left) reveals that the ENCE depends also on M as a power law. One has thus $ENCE = \alpha M^\beta N^{1/2}$ with $\beta \simeq -1/2$. The same data are then plot on a linear scale as a function of $(N/M)^{1/2}$, from which one can see a perfect linear fit by $\tilde{\vartheta}_{N,ref} = \alpha \times (N/M)^{1/2}$ with $\alpha \simeq 0.56$. The ZMSE presents features very similar to the ENCE, with a different slope, $\tilde{\vartheta}_{N,ref} \simeq 1.14 \times (N/M)^{1/2}$. In both cases, the impact of the uncertainty distribution is minor, albeit larger for ENCE than for ZMSE, and negligible for the validation process.

It seems therefore that, for a normal generative model and a chosen binning scheme, it is possible to define a reference value for the ENCE and ZMSE statistics for each dataset. The availability of a predefined reference value would considerably simplify the validation approach, when compared to the extrapolation-based one proposed by Pernot⁴.

Let us now check the sensitivity of this approach to the generative distribution. The same protocol as above is followed for another choice of generative model (denoted T6IG)

$$E \sim u_E \times t_s(6) \quad (B3)$$

$$u_E^2 \sim \Gamma^{-1}(\nu/2, \nu/2) \quad (B4)$$

and the results are shown in Fig. A2.

The same linear relationship as for the normal generative distribution is observed, but, in each case, with a different slope and a small positive intercept. Besides, a very small




Figure A1. Sensitivity of simulated ENCE and ZMSE values to the dataset size (M), number of bins (N) and shape of the uncertainty distribution (ν), for calibrated datasets generated from the *NIG* model.

deviation from linearity can also be perceived. For ENCE, one gets then $\hat{\theta}_{T,ref} \simeq 0.004 + 0.779 \times (N/M)^{1/2}$ and for ZMSE $\hat{\theta}_{T,ref} \simeq 0.006 + 1.577 \times (N/M)^{1/2}$.

It seems thus impossible to define a reference value for ENCE and ZMSE if the generative distribution is unknown.




Figure A2. Same as Fig. A1 for the T6IG model.

Appendix C: Validation of ZMSE for the application datasets

For each of the nine datasets, one establishes a ZMSE validation diagnostic by a method which is independent of a generative distribution hypothesis. For this, the ZMSE is estimated for a series of bin numbers N between 10 and 150, with the constraint that bins should not contain less than 20 data points. The data with $N > 20$ are fitted by a linear model as a function of $(N/M)^{1/2}$. The intercept of the regression line is then compared to zero.⁴ The reference lines for the NIG and T6IG models defined in Appendix B are also plotted for comparison.

All the datasets are failing this validation test of conditional calibration (consistency), as none of the red intervals encloses the origin. Note however that for some datasets (e.g. Sets 3 and 4), it is possible to choose a number of bins such that the ZMSE is compatible with the T6IG reference (dotted line). This shows the danger of using a reference value from an unconstrained generative model.




Figure A3. Validation of ZMSE by the extrapolation method: ZMSE are estimated for a sequence of bin numbers and plotted as function of $(N/M)^{1/2}$ (blue dots); this sequence (for $N > 20$) is fitted by a linear model (solid red line); reference lines are plotted for the NIG (dashed line) and T6IG (dotted line) models.