

# THE NON-ORIENTABLE 4-GENUS OF 11 CROSSING NON-ALTERNATING KNOTS

MEGAN FAIRCHILD

ABSTRACT. The non-orientable 4-genus of a knot  $K$  in  $S^3$  is defined to be the minimum first Betti number of a non-orientable surface  $F$  in  $B^4$  so that  $K$  bounds  $F$ . We will survey the tools used to compute the non-orientable 4-genus, and use various techniques to calculate this invariant for non-alternating 11 crossing knots. We also will view obstructions to a knot bounding a Möbius band given by the double branched cover of  $S^3$  branched over  $K$ .

## 1 INTRODUCTION

Knots bounding orientable surfaces, both in  $S^3$  and  $B^4$ , has been extensively studied, however much is still to be learned about the non-orientable surfaces in  $B^4$  bounded by knots. Recently, the non-orientable 4-genus of torus knots has been computed for all knots  $T(2, q)$  and  $T(3, q)$  by Allen [1], and most knots  $T(4, q)$  by Binns, Kang, Simone, Truöl, and Sabloff [2, 14]. The non-orientable 4-genus of knots with 10 or fewer crossings has also been computed in detail by Ghanbarian, Jabuka, and Kelly [3, 6], with much focus on alternating knots. This paper aims to shed light on the non-alternating case and strategies to calculate the non-orientable 4-genus. We will explore various techniques in finding this invariant, as well as examining obstructions to knots bounding a Möbius band.

For this paper, a knot  $K$  is in  $S^3$ . The orientable 4-genus of a knot is the minimum genus of an orientable surface in the 4-ball that is bounded by  $K$  and is denoted  $g_4(K)$ , and knots with  $g_4(K) = 0$  are called slice knots. Following Murakami and Yasuhara in [12], the non-orientable 4-genus of a knot  $K$ , denoted  $\gamma_4(K)$ , is defined to be the minimum first Betti number of non-orientable surfaces  $F$  smoothly embedded in  $B^4$  bounded by  $K$ , that is  $\min\{b_1(F) \mid \partial F = K\}$ . Note that the first Betti number is defined to be  $b_1(F) = \dim H_1(F; \mathbb{Z})$ . We have, by definition, for any knot  $K$ ,  $\gamma_4(K) \geq 1$  where equivalence applies when  $K$  bounds a Möbius band. Slice knots that bound a smooth disk embedded in  $B^4$  have non-orientable 4-genus one, as we may attach a non-oriented band to such an embedded disk.

**Theorem 1.1.** *For the 185 non-alternating 11 crossing knots,*

- (a) *121 knots have  $\gamma_4(K) = 1$*
- (b) *58 knots have  $\gamma_4(K) = 2$*

*The remaining 6 knots have  $\gamma_4(K) = 1$  or 2.*

The paper is organized as follows: Section 2 is the background on knot invariants, double branched covers, and useful bounds and obstructions for the non-orientable 4-genus. Section 3 is a survey of the techniques used to solve this problem as well as results.

**Acknowledgements.** I would like to thank Chuck Livingston, Pat Gilmer, and Slaven Jabuka for helpful conversations and comments. Additional thanks to Dror Bar-Natan for

permitting my use of the Knot Atlas figures [8].

## 2 BACKGROUND

We begin by reviewing knot invariants and examining bounds for the non-orientable 4-genus as well as obstructions to a knot bounding a Möbius band. First, the crossing number of a knot is denoted  $n(K)$  and is the crossing number of a diagram of a knot with the fewest crossings that could be drawn on the plane to represent the knot. The unknotting number of a knot  $u(K)$  is the minimum number of crossing changes required to transform  $K$  into the unknot. Similarly,  $u_s(K)$  is the minimum number of crossing changes to change  $K$  into a slice knot. The 4-dimensional clasp number,  $c_4(K)$ , is the minimum number of double points of transversely immersed 2-disks in the 4-ball bounded by  $K$  [12]. We then have the following triple inequality from Jabuka and Kelly [6]:

$$g_4(K) \leq c_4(K) \leq u_s(K) \leq u(K)$$

The orientable genus of a knot also offers an upper bound for the non-orientable 4-genus, respective with smooth and topological for  $i = 4$ , we have [6]:

$$\gamma_i(K) \leq 2g_i(K) + 1 \text{ for } i = 3, 4$$

Similar to the orientable 4-genus, we obtain an upper bound for the non-orientable 4-genus from the non-orientable 3-genus of a knot called the *crosscap number* [9], which is the minimum genus non-orientable surface a knot bounds in  $S^3$ , denoted  $c(K)$ , so we have  $\gamma_4(K) \leq c(K)$ .

Following the notation of Murakami and Yasuhara [11], we define  $\Gamma_4(K) = \min\{b_1(F) | \partial F = K\}$ , or similarly  $\Gamma_4(K) = \min\{2g_4(K), \gamma_4(K)\}$ , and thus  $\Gamma_4(K) \leq \gamma_4(K)$ . Murakami and Yasuhara then give us the following proposition [12]:

**Proposition 2.1** (Proposition 2.3 in [12]). *For any knot  $K$ , the following inequalities hold.*

$$\Gamma_4(K) \leq \begin{cases} c_4(K) & \text{if } c_4(K) \text{ is even} \\ c_4(K) + 1 & \text{otherwise} \end{cases}$$

$$\gamma_4(K) \leq \begin{cases} c_4(K) & \text{if } c_4(K) \text{ is even and } c_4(K) \neq 2 \\ c_4(K) + 1 & \text{otherwise} \end{cases}$$

**Corollary 2.2** (Corollary 2.4 in [12]). *For a knot  $K$ , if  $g_4(K) = c_4(K) \geq 1$ , then  $\Gamma_4(K) = \gamma_4(K)$ .*

The crossing number of a knot offers an upper bound, so we have [11]:

$$\Gamma(K) \leq \left\lfloor \frac{n(K)}{2} \right\rfloor \text{ and } \gamma_4(K) \leq \left\lfloor \frac{n(K)}{2} \right\rfloor$$

The signature of a knot  $\sigma(K)$  is defined to be the signature of the sum of knot's Seifert matrix and it's transpose,  $\sigma(V + V^t)$ . The Arf invariant of a knot is denoted  $\text{Arf}(K)$  and is a concordance invariant in  $\mathbb{Z}_2$  which is calculated using the Seifert form of a knot [9]. These two invariants form a lower bound for the non-oriented 4-genus of a knot, so we have the following proposition.

**Proposition 2.3** (Proposition 2.4 in [3]). *Given a knot  $K$ , if  $\sigma(K) + 4\text{Arf}(K) \equiv 4 \pmod{8}$ , then  $\gamma_4(K) \geq 2$ .*

### Double Branched Cover

Recall the definition of the non-orientable 4-genus is  $\gamma_4(K) = \min\{b_1(F) \mid \partial F = K\}$  and note that  $b_1(F) = \dim H_1(F, \mathbb{Q})$ . Let  $K$  in  $S^3$  bound a connected surface  $F$  in  $B^4$  and denote  $D_F(B^4)$  as the double branched cover of  $B^4$  branched over  $F$ . Gilmer and Livingston proved in [4], Lemma 1, that  $b_2(D_F(B^4)) = b_1(F)$ . The reasoning here is that the double branched cover of  $S^3$  branched over  $K$ , denoted  $D_K(S^3)$ , is a rational homology sphere and  $H_1(D_F(B^4); \mathbb{Q}) = 0$ . We thus may use the linking form of  $D_K(S^3)$  to provide information on the intersection form of  $D_F(B^4)$ .

We also have that the first homology of  $D_K(S^3)$  is finite, so we have a linking form  $\lambda$ , and this is explored in detail by Murakami and Yasuhara in [12]

$$\lambda : H_1(D_K(S^3); \mathbb{Z}) \times H_1(D_K(S^3); \mathbb{Z}) \rightarrow \mathbb{Q}/\mathbb{Z}$$

Given a Goeritz matrix  $G$  for  $K$  (see Section III for details), we have that  $G$  is a relation matrix for  $H_1(D_K(S^3); \mathbb{Z})$  and the linking form  $\lambda$  is given by  $\pm G^{-1}$ , where the sign depends on orientation of  $D_K(S^3)$  [12]. The double branched cover is a useful tool in obstructing knots bounding a Möbius band or a Klein bottle.

**Corollary 2.4** (Corollary 3 in [4]). *Suppose that  $H_1(D_K(S^3)) = \mathbb{Z}_n$  where  $n$  is the product of primes, all with odd exponent. Then if  $K$  bounds a Möbius band in  $B^4$ , there is a generator  $a \in H_1(D_K(S^3))$  such that  $\lambda(a, a) = \pm 1/n$*

**Theorem 2.5** (Theorem 4 in [4]). *Suppose that  $H_1(D_K(S^3)) = \mathbb{Z}_p \oplus \mathbb{Z}_p$  where  $p$  is prime. Then if  $K$  bounds a punctured Klein bottle in  $B^4$ , the discriminant of the linking form is  $\pm 1 \in \mathbb{F}_p^*/(\mathbb{F}_p^*)^2$*

**Theorem 2.6** (Theorem 11 in [4]). *Suppose that  $H_1(D_K(S^3)) = \mathbb{Z}_p \oplus \mathbb{Z}_p \oplus \mathbb{Z}_q$  where  $q \equiv 1 \in \mathbb{F}_p^*/(\mathbb{F}_p^*)^2$ . If  $H_1(D_K(S^3))$  is the boundary of a 4-manifold  $W$  with second Betti number 2 which has an indefinite intersection form, then the linking form restricted to  $\mathbb{Z}_p \oplus \mathbb{Z}_p \subset H_1(D_K(S^3))$  is metabolic.*

## 3 RESULTS AND TECHNIQUES

There are a total of 185 knots that are non-alternating and have 11 crossings, according to the knot info database [9]. Of those knots, there are 16 that are smoothly slice and thus have  $\gamma_4(K) = 1$ .

**Remark 3.1.** There are 16 non-alternating 11 crossing knots that are slice and thus bound a Möbius band:

$$\begin{aligned} &11n_4, 11n_{21}, 11n_{37}, 11n_{39}, 11n_{42}, 11n_{49}, 11n_{50}, 11n_{67}, \\ &11n_{73}, 11n_{74}, 11n_{83}, 11n_{97}, 11n_{116}, 11n_{132}, 11n_{139}, 11n_{172} \end{aligned}$$

**Proposition 3.2.** *The following knots have  $\gamma_4(K) = 1$ :*

$$\begin{aligned} &11n_1, 11n_3, 11n_5, 11n_6, 11n_7, 11n_8, 11n_9, 11n_{11}, 11n_{13}, 11n_{14}, 11n_{15}, \\ &11n_{16}, 11n_{18}, 11n_{19}, 11n_{20}, 11n_{23}, 11n_{24}, 11n_{25}, 11n_{26}, 11n_{27}, 11n_{31}, 11n_{34}, \end{aligned}$$

$11n_{36}, 11n_{41}, 11n_{44}, 11n_{45}, 11n_{46}, 11n_{47}, 11n_{52}, 11n_{54}, 11n_{57}, 11n_{58}, 11n_{59},$   
 $11n_{60}, 11n_{62}, 11n_{64}, 11n_{65}, 11n_{66}, 11n_{68}, 11n_{69}, 11n_{70}, 11n_{71}, 11n_{75}, 11n_{76},$   
 $11n_{77}, 11n_{78}, 11n_{79}, 11n_{80}, 11n_{81}, 11n_{82}, 11n_{86}, 11n_{87}, 11n_{88}, 11n_{89}, 11n_{91},$   
 $11n_{93}, 11n_{94}, 11n_{96}, 11n_{102}, 11n_{104}, 11n_{105}, 11n_{106}, 11n_{107}, 11n_{110}, 11n_{111},$   
 $11n_{113}, 11n_{117}, 11n_{118}, 11n_{120}, 11n_{121}, 11n_{122}, 11n_{123}, 11n_{124}, 11n_{126}, 11n_{127},$   
 $11n_{128}, 11n_{129}, 11n_{134}, 11n_{135}, 11n_{136}, 11n_{142}, 11n_{143}, 11n_{145}, 11n_{146}, 11n_{147},$   
 $11n_{148}, 11n_{150}, 11n_{151}, 11n_{152}, 11n_{153}, 11n_{154}, 11n_{157}, 11n_{158}, 11n_{160}, 11n_{162},$   
 $11n_{163}, 11n_{164}, 11n_{167}, 11n_{168}, 11n_{169}, 11n_{170}, 11n_{173}, 11n_{180}, 11n_{181}, 11n_{183}$

**Proposition 3.3.** *The following knots have  $\gamma_4(K) = 2$ :*

$11n_2, 11n_{10}, 11n_{12}, 11n_{22}, 11n_{28}, 11n_{29}, 11n_{30}, 11n_{32}, 11n_{33}, 11n_{35},$   
 $11n_{38}, 11n_{43}, 11n_{48}, 11n_{51}, 11n_{53}, 11n_{55}, 11n_{56}, 11n_{61}, 11n_{63}, 11n_{72},$   
 $11n_{84}, 11n_{85}, 11n_{90}, 11n_{92}, 11n_{95}, 11n_{98}, 11n_{99}, 11n_{100}, 11n_{101}, 11n_{103},$   
 $11n_{108}, 11n_{109}, 11n_{112}, 11n_{114}, 11n_{115}, 11n_{119}, 11n_{125}, 11n_{130}, 11n_{131}, 11n_{133},$   
 $11n_{137}, 11n_{138}, 11n_{140}, 11n_{141}, 11n_{144}, 11n_{149}, 11n_{155}, 11n_{156}, 11n_{161}, 11n_{165}$   
 $11n_{171}, 11n_{174}, 11n_{175}, 11n_{176}, 11n_{179}, 11n_{182}, 11n_{184}, 11n_{185},$

### Constraints on Invariants

The knot invariant information for this paper was extracted from Knot Info [9].

**Lemma 3.4.** *Given  $K$  is a knot satisfying  $\sigma(K) + 4\text{Arf}(K) \equiv 4 \pmod{8}$ , and  $c_4(K) = 1$ , then  $\gamma_4(K) = 2$ .*

The result is clear from Proposition 2.1 and Corollary 2.3. We now examine knots that have  $g_4(K) = u(K) = 1$  (or  $g_4(K) = u_s(K) = 1$ ) and  $\sigma(K) + 4\text{Arf}(K) \equiv 4 \pmod{8}$  to see the following knots have  $\gamma_4(K) = 2$ :

$11n_{12}, 11n_{28}, 11n_{48}, 11n_{53}, 11n_{55}, 11n_{85}, 11n_{100}$   
 $11n_{114}, 11n_{115}, 11n_{119}, 11n_{130}, 11n_{156}, 11n_{179}, 11n_{182}$

All the above listed knots satisfy  $\sigma(K) + 4\text{Arf}(K) \equiv 4 \pmod{8}$ . Since they satisfy  $g_4(K) = 1 = u(K)$  (or  $g_4(K) = u_s(K) = 1$ ) by the hypothesis, we have  $c_4(K) = 1$ , and thus by Lemma 3.4 we may conclude  $\gamma_4(K) = 2$ .

**Lemma 3.5.** *Given  $K$  is a knot satisfying  $\sigma(K) + 4\text{Arf}(K) \equiv 4 \pmod{8}$ , and  $c_4(K) = 2$ , then  $\gamma_4(K) = 2$ .*

By Corollary 2.2, we have  $\Gamma_4(K) = \gamma_4(K)$ , and thus applying Proposition 2.1 we achieve  $\gamma_4(K) \leq 2$ . Therefore,  $\gamma_4(K) = 2$ . We now observe that the following knots have  $\gamma_4(K) = 2$ :

$11n_2, 11n_{35}, 11n_{95}, 11n_{103}, 11n_{108}, 11n_{109}, 11n_{144}, 11n_{149}, 11n_{174}, 11n_{175}, 11n_{185}$

The above listed knots all satisfy  $\sigma(K) + 4\text{Arf}(K) \equiv 4 \pmod{8}$  and thus  $\gamma_4(K) \geq 2$ . Additionally, these knots all satisfy  $g_4(K) = u(K) = 2$ , and thus  $c_4(K) = 2$ .

### Non-Oriented Band Moves

The primary method used in calculations was via non-oriented band moves. We begin with an oriented knot  $K$  and an oriented band,  $[0, 1] \times [0, 1]$ . Following the conventions of Jabuka and Kelly [6], we attach the band to  $K$  in the sense that the orientation of the band agrees with the orientation of  $K$  on  $[0, 1] \times \{0\}$  but disagrees on  $[0, 1] \times \{1\}$ , or vice versa. One then does surgery along the band. The result of non-orientable band surgery will always be a knot, while the result after *orientable* band surgery is a link. Non-orientable band surgery is explored by Moore and Vazquez in [10] and is called *non-coherent band surgery*.

The notation for a knot  $K$  that has been transformed into a knot  $K'$  by a non-oriented band move is  $K \xrightarrow{h} K'$  where  $h$  is either 0, 1, or -1. These three band moves can be seen in the figure below. From left to right, we have  $\xrightarrow{0}$  is the band move without a twist,  $\xrightarrow{-1}$  is the band move with a left-handed twist, and  $\xrightarrow{1}$  is the band move with a right handed twist.




FIGURE 3.1. Band Moves

**Proposition 3.6** (Proposition 2.4 in [6]). *If the knots  $K$  and  $K'$  are related by a non-oriented band move, then*

$$\gamma_4(K) \leq \gamma_4(K') + 1$$

*If a knot  $K$  is related to a slice knot  $K'$  by a non-oriented band move, then  $\gamma_4(K) = 1$ .*

*Proof of Theorem 1.1 part (a).* Every knot listed in Proposition 3.2 is either a slice knot or one non-oriented band move away from a slice knot. See Figure 4.3 - Figure 4.11 for details.

**Lemma 3.7.** *The following knots have  $\gamma_4(K) = 2$ :*

$$11n_{10}, 11n_{12}, 11n_{30}, 11n_{32}, 11n_{43}, 11n_{48}, 11n_{51}, 11n_{55}, 11n_{61}, 11n_{72} \\ 11n_{85}, 11n_{90}, 11n_{98}, 11n_{103}, 11n_{130}, 11n_{133}$$

We now recall Proposition 2.3 and note the knots listed in the above lemma all satisfy  $\sigma(K) + 4\text{Arf}(K) \equiv 4 \pmod{8}$ . So we know the above knots have  $\gamma_4(K) \geq 2$ . The above listed knots all are one non-oriented band move away from a knot  $K'$  so that  $\gamma_4(K') = 1$  (see Figure 4.12 - Figure 4.15), thus we conclude  $\gamma_4(K) = 2$ .

### Linking Form Calculation

We look for a knot  $K$  so that  $\sigma(K) + 4\text{Arf}(K) \equiv 0, \pm 2 \pmod{8}$ , and thus  $K$  does not meet the obstruction from Proposition 2.3. We calculate the linking form of  $H_1(D_K(S^3))$  to see if  $K$  meets the obstruction from Corollary 2.4. The first thing we do is calculate the Goeritz matrix for  $K$ . We will do an example here, but an interested reader is referred to Gordan and Litherland [5].

To construct the Goeritz matrix, we first make a checkerboard coloring of a knot.




FIGURE 3.2. Checkerboard coloring for  $11n_{155}$

Each white region is labeled  $R_i$  and the unbounded region is  $R_0$ . We then assign a value to each crossing  $C$ ,  $\eta(C) = \pm 1$ , via the figure below, and following the conventions from Gordan and Litherland [5].




FIGURE 3.3. left:  $\eta(C) = 1$ , right:  $\eta(C) = -1$

Next, we construct a matrix  $G'$  with the algorithm:

$$g'(i, j) = \begin{cases} -\sum \eta(C) & \text{where the sum ranges over all crossings } C \text{ incident to } R_i \text{ and } R_j, i \neq j \\ -\sum_{k \neq i} g'(i, k) = g'(i, i) & \text{if } i = j \end{cases}$$

Then, the Goeritz matrix  $G$  is obtained from  $G'$  by deleting the  $0^{\text{th}}$  row and column. The determinant of  $G$  is an invariant of the knot, and  $G$  is a linking matrix for  $H_1(D_K(S^3))$  [5, 12].

Now, we may calculate the linking form. As previously mentioned,  $\pm G^{-1}$  represents the linking form  $\lambda$  where  $\lambda : H_1(D_K(S^3); \mathbb{Z}) \times H_1(D_K(S^3); \mathbb{Z}) \rightarrow \mathbb{Q}/\mathbb{Z}$ . To continue the example, we have  $G$  and  $G^{-1}$  for the knot  $11n_{155}$  as:

$$G = \begin{bmatrix} 3 & -1 & 0 & -1 \\ -1 & 5 & -1 & 0 \\ 0 & -1 & 0 & 2 \\ -1 & 0 & 2 & 0 \end{bmatrix} \quad G^{-1} = \begin{bmatrix} \frac{20}{51} & \frac{2}{17} & \frac{10}{51} & \frac{1}{17} \\ \frac{2}{51} & \frac{1}{17} & \frac{1}{51} & \frac{1}{17} \\ \frac{17}{10} & \frac{1}{17} & \frac{17}{5} & \frac{17}{9} \\ \frac{51}{1} & \frac{17}{2} & \frac{51}{17} & \frac{17}{1} \end{bmatrix}$$

Now we have the linking form  $\lambda(g, g) = \pm 20/51$ . Suppose  $11n_{155}$  bounds a Möbius band. We wish to find an  $n \in \mathbb{Z}$  so that  $\lambda(ng, ng) = \pm 1/51$ . This means  $\pm 20/51 = \lambda(ng, ng) = n^2 \lambda(g, g) = \pm 20n^2/51 = \pm 1/51$ , so  $20n^2 \equiv \pm 1 \pmod{51}$ . A quick calculation shows this is not possible, and thus  $11n_{155}$  does not bound a Möbius band.

## Results

**Theorem 3.8** (Theorem 2 in [4]). *Let  $K$  in  $S^3$  be a knot. The linking form  $(H_1(D_K(S^3)), \lambda)$  splits as a direct sum  $(G_1, \lambda_1) \oplus (G_2, \lambda_2)$  where  $(G_2, \lambda_2)$  is metabolic and  $(G_1, \lambda_1)$  has a presentation of rank  $\lambda_1(F)$ .*

**Lemma 3.9.** *Let  $K$  in  $S^3$  be a knot and suppose that  $H_1(D_K(S^3)) = \mathbb{Z}_{p^2q}$  where  $p$  is prime and  $q$  is a product of primes, all with odd exponent. Then if  $K$  bounds a Möbius band in  $B^4$ , there is a generator  $a \in H_1(D_K(S^3))$  such that either  $\lambda(a, a) = \pm 1/p^2q$  or  $\lambda(a, a) = \pm 1/q$ .*

*Proof.* As we see in Theorem 3.8,  $(H_1(D_K(S^3)), \lambda)$  splits as a direct sum  $(G_1, \lambda_1) \oplus (G_2, \lambda_2)$  where  $(G_2, \lambda_2)$  is metabolic and  $\lambda_1$  is presented by the linking matrix of  $D_K(S^3)$ , which has a presentation of rank one. As  $q$  is square-free, we have that  $\mathbb{Z}_q$  is completely contained in  $G_1$ . Then either  $\mathbb{Z}_{p^2}$  is completely contained in  $G_2$ , which implies it is metabolic, or  $\mathbb{Z}_{p^2}$  is contained in  $G_1$ .

If  $\mathbb{Z}_{p^2}$  is completely contained in  $G_2$ , then there exists a subgroup  $H$  of  $\mathbb{Z}_{p^2}$  so that  $|H|^2 = p^2$  and  $\lambda(g, g') = 0$  for any  $g, g' \in H$ , since  $\lambda_2$  is metabolic. Then, as  $\lambda_1$  must have a presentation of rank one, we have that the presentation matrix must be of the form  $(\pm |G_1|) = (\pm q)$ . Therefore, the linking form  $\lambda_1$  on  $G_1$  is given by  $\pm 1/q$ .

If  $\mathbb{Z}_{p^2}$  is completely contained in  $G_1$ , a similar argument shows  $\lambda_1$  is given by  $\pm 1/q$   $\square$

The following knots:

$$11n_{22}, 11n_{29}, 11n_{33}, 11n_{56}, 11n_{84}, 11n_{92}, 11n_{101}, 11n_{112}, 11n_{125}, 11n_{131}, 11n_{138}, 11n_{155}, \\ 11n_{176}, 11n_{184}$$

have the respective linking forms:

$$\frac{42}{55}, \frac{14}{51}, \frac{22}{51}, \frac{12}{35}, \frac{18}{35}, \frac{2}{15}, \frac{19}{39}, \frac{53}{55}, \frac{61}{63}, \frac{39}{67}, \frac{13}{15}, \frac{20}{51}, \frac{11}{63}, \frac{2}{87}$$

All of which satisfy the obstruction from Corollary 2.4 and Lemma 3.9. Additionally, all of these knots have an non-orientable band move to a knot  $K'$  where  $\gamma_4(K') = 1$  (Figures 4.12 -4.15). Thus, each of these knots has non-orientable 4-genus equal to 2.

### Knot Floer Homology

Ozsváth, Stipsicz, and Szabó explored non-orientable knot floer homology and how the Upsilon invariant can be used as a lower bound for the non-orientable 4-genus [13]. Given  $K$  is a knot, denote  $\Upsilon_K(1)$  as  $v(K)$  (lower case upsilon), and then we have:

$$\left| v(K) - \frac{\sigma(K)}{2} \right| \leq \gamma_4(K)$$

However, if  $K$  is not an L-space knot, this invariant is rather difficult to compute. Additionally, we have from [13] that for an alternating (or quasi-alternating) knot  $K$ ,

$$v(K) = \frac{\sigma(K)}{2}$$

For the 185 non-alternating 11-crossing knots, only 3 are not quasi-alternating. Of those 3, two are slice and one is not. This is thus not a useful lower bound for the knots being considered in this paper. However, this is a useful invariant for torus knots, demonstrated in detail by Binns, Kang, Simone, and Truöl in [2]. Additionally, Allen explored a geography problem where the upsilon invariant was wonderfully utilized in [1].

### 4 SPECIAL CASES

**Lemma 4.1.** *The knot  $11n_{38}$  does not bound a Möbius band.*

The knot  $11n_{38}$  has  $H_1(D_K(S^3)) = \mathbb{Z}_3$  and thus the linking form is represented by the  $1 \times 1$  matrix  $[1/3]$ . This is clear, as the non-zero elements of  $\mathbb{Z}_3$  are 1 and -1. Then, if  $K$  bounds a Möbius band  $F$  in  $B^4$ , we have  $b(F) = b(D_F(B^4)) = 1$  and  $D_F(B^4)$  is negative definite [4]. From Theorem 3 in [5], we have that the intersection form on  $H_2(D_F(B^4))$  is represented by the linking matrix on  $H_1(D_K(S^3))$ , which can be viewed from the entries in the Goeritz matrix. The Goeritz matrix  $G$  is a  $4 \times 4$  matrix that is indefinite, and when diagonalized,  $G = SJS^{-1}$ , the matrix  $J$  is also indefinite. We may suppose that there exists a presentation matrix that represents the linking form, and by checking the diagonal entries on  $-G^{-1}$ , we have that  $1/3$  represents the form. This implies the manifold is positive definite, which is a contradiction. Thus,  $11n_{38}$  does not bound a Möbius band. We then have that there is a non-orientable band move from  $11n_{38}$  to the trefoil knot, which has  $\gamma_4(3_1) = 1$ , therefore we may conclude that  $\gamma_4(11n_{38}) = 2$ . The figure below was obtained from Knot Atlas [8].




FIGURE 4.1. A non-oriented band move from  $11n_{38} \xrightarrow{0} 3_1$



We thus have a combination of Lemmas 3.4, 3.5, and 4.1, Proposition 3.6, and Theorem 3.8 showing Proposition 3.3 is true, thus proving part (b) of Theorem 1.1.

**Lemma 4.2.** *The knots  $11n_{17}$ ,  $11n_{40}$ ,  $11n_{159}$ ,  $11n_{166}$ ,  $11n_{177}$  and  $11n_{178}$  all have  $\gamma_4(K) = 1$  or  $2$ .*

We have the following table:

Knot	linking form	definiteness of $D_F(B^4)$	4-genus
$11n_{17}$	$1/47$	positive	1
$11n_{40}$	$-1/79$	negative	1
$11n_{159}$	$1/71$	positive	1
$11n_{166}$	$1/59$	positive	1
$11n_{177}$	$1/83$	positive	1
$11n_{178}$	$-1/95$	negative	1

*Proof.* Denote  $K$  as a knot listed in Lemma 4.2. We first examine the knot signature and Arf invariant to see  $\sigma(K) + 4\text{Arf}(K) \equiv \pm 2 \pmod{8}$ . Thus, we do not meet the obstruction from Proposition 2.3, so we may only conclude  $\gamma_4(K) \geq 1$ . We then move on to examining the linking form of  $K$ . Note that the determinant of  $K$ ,  $d = \det(K)$ , is either a prime number or a product of exactly 2 prime numbers. As  $d = |H_1(D_K(S^3))|$ , we cannot have a splitting of  $H_1(D_K(S^3))$  into  $G_1 \oplus G_2$  where  $G_2$  is metabolic, since  $d$  is square free. We thus see that the linking form  $\lambda$  for each knot is of the form  $\pm 1/d$ . We also compare the linking form of the knot to the definiteness of  $D_F(B^4)$ . The sign of the 4-manifold  $D_F(B^4)$  corresponds to the sign of the quadratic form [4], thus the linking form, and we see that our signs are corresponding for the linking form and definiteness of  $D_F(B^4)$ . Additionally, each knot is one band move away from a knot  $K'$  so that  $\gamma_4(K') = 1$ , see Figure 4.2, and thus  $\gamma_4(K) \leq 2$ . We thus cannot find an obstruction to these knots bounding a Möbius band, but also cannot find the desired band move to a slice knot. Therefore,  $\gamma_4(K) \leq 2$  for the knots in Lemma 4.2.  $\square$

This concludes the proof for Theorem 1.1.




FIGURE 4.2. Non-oriented band moves from the knots  $11n_{17}$ ,  $11n_{40}$ ,  $11n_{159}$ ,  $11n_{166}$ ,  $11n_{177}$ , and  $11n_{178}$  to knots with non-orientable genus 1.

### Concordance

Knot concordance is a great tool that could be used to solve for the non-orientable 4-genus. For the six remaining knots, their concordance genus is known [9], however the knots to which they are concordant is still unknown. Suppose a given knot  $K$  is concordant to  $K'$ , then it is clear that  $\gamma_4(K) = \gamma_4(K')$ .

**Question 4.3.** Is  $11n_{40}$  concordant to  $10_{57}$ ?

$10_{57}$  is a wonderful candidate for concordance to  $11n_{40}$ , just by a simple analysis of their invariants [9]. If the answer to Question 4.3 is yes, then the  $11n_{40}$  knot has  $\gamma_4(11n_{40}) = 1$ .

**Conjecture 4.4.** The knots  $11n_{17}$ ,  $11n_{159}$ ,  $11n_{166}$ ,  $11n_{177}$ , and  $11n_{178}$  are not concordant to any knot with 11 or fewer crossings. Moreover,  $11n_{17}$ ,  $11n_{159}$ , and  $11n_{166}$  are not concordant to any knot with 12 or fewer crossings.

It should be noted that Kearny has found the concordance genus of 11-crossing knots in [7], as well as specific concordances from 11-crossing knots to knots of lower crossings.




FIGURE 4.3. Non-oriented band moves from the knots  $11n_1, 11n_3, 11n_5, 11n_6, 11n_7, 11n_8, 11n_9, 11n_{11}, 11n_{13}, 11n_{14}, 11n_{15},$  and  $11n_{16}$  to smoothly slice knots.




FIGURE 4.4. Non-oriented band moves from the knots  $11n_{18}$ ,  $11n_{19}$ ,  $11n_{20}$ ,  $11n_{23}$ ,  $11n_{24}$ ,  $11n_{25}$ ,  $11n_{26}$ ,  $11n_{27}$ ,  $11n_{31}$ ,  $11n_{34}$ ,  $11n_{36}$ , and  $11n_{41}$  to smoothly slice knots.




FIGURE 4.5. Non-oriented band moves from the knots  $11n_{44}$ ,  $11n_{45}$ ,  $11n_{46}$ ,  $11n_{47}$ ,  $11n_{52}$ ,  $11n_{54}$ ,  $11n_{57}$ ,  $11n_{58}$ ,  $11n_{59}$ ,  $11n_{60}$ ,  $11n_{62}$ , and  $11n_{64}$  to smoothly slice knots.




FIGURE 4.6. Non-oriented band moves from the knots  $11n_{65}$ ,  $11n_{66}$ ,  $11n_{68}$ ,  $11n_{69}$ ,  $11n_{70}$ ,  $11n_{71}$ ,  $11n_{75}$ ,  $11n_{76}$ ,  $11n_{77}$ ,  $11n_{78}$ ,  $11n_{79}$ , and  $11n_{80}$  to smoothly slice knots.




FIGURE 4.7. Non-oriented band moves from the knots  $11n_{81}$ ,  $11n_{82}$ ,  $11n_{86}$ ,  $11n_{87}$ ,  $11n_{88}$ ,  $11n_{89}$ ,  $11n_{91}$ ,  $11n_{93}$ ,  $11n_{94}$ ,  $11n_{96}$ ,  $11n_{102}$ , and  $11n_{104}$  to smoothly slice knots.




FIGURE 4.8. Non-oriented band moves from the knots  $11n_{105}$ ,  $11n_{106}$ ,  $11n_{107}$ ,  $11n_{110}$ ,  $11n_{111}$ ,  $11n_{113}$ ,  $11n_{117}$ ,  $11n_{118}$ ,  $11n_{120}$ ,  $11n_{121}$ ,  $11n_{122}$ , and  $11n_{123}$  to smoothly slice knots.









FIGURE 4.9. Non-oriented band moves from the knots  $11n_{124}$ ,  $11n_{126}$ ,  $11n_{127}$ ,  $11n_{128}$ ,  $11n_{134}$ ,  $11n_{135}$ ,  $11n_{136}$ ,  $11n_{142}$ ,  $11n_{143}$ ,  $11n_{145}$ ,  $11n_{146}$ , and  $11n_{147}$  to smoothly slice knots.




$$(A) 11n_{148} \xrightarrow{1} 10_{137}$$




$$(B) 11n_{150} \xrightarrow{0} 8_8$$




$$(C) 11n_{151} \xrightarrow{-1} 10_{153}$$




$$(D) 11n_{152} \xrightarrow{0} 10_{153}$$




$$(E) 11n_{153} \xrightarrow{1} 10_{129}$$




$$(F) 11n_{154} \xrightarrow{-1} 12n_{504}$$




$$(G) 11n_{157} \xrightarrow{-1} 9_{27}$$




$$(H) 11n_{158} \xrightarrow{0} 0_1$$




$$(I) 11n_{160} \xrightarrow{1} 12n_{802}$$



$$(J) 11n_{162} \xrightarrow{-1} 10_{140}$$



$$(K) 11n_{163} \xrightarrow{0} 8_8$$



$$(L) 11n_{164} \xrightarrow{0} 8_{20}$$

FIGURE 4.10. Non-oriented band moves from the knots  $11n_{148}$ ,  $11n_{150}$ ,  $11n_{151}$ ,  $11n_{152}$ ,  $11n_{153}$ ,  $11n_{154}$ ,  $11n_{157}$ ,  $11n_{158}$ ,  $11n_{160}$ ,  $11n_{162}$ ,  $11n_{163}$ , and  $11n_{164}$  to smoothly slice knots.




FIGURE 4.11. Non-oriented band moves from the knots  $11n_{167}$ ,  $11n_{168}$ ,  $11n_{169}$ ,  $11n_{170}$ ,  $11n_{173}$ ,  $11n_{180}$ ,  $11n_{181}$ , and  $11n_{183}$  to smoothly slice knots.




FIGURE 4.12. Non-oriented band moves from the knots  $11n_{10}$ ,  $11n_{12}$ ,  $11n_{22}$ ,  $11n_{29}$ ,  $11n_{30}$ ,  $11n_{32}$ ,  $11n_{33}$ ,  $11n_{43}$ ,  $11n_{48}$ ,  $11n_{51}$ ,  $11n_{55}$ , and  $11n_{56}$  to knots with non-orientable genus 1.




FIGURE 4.13. Non-oriented band moves from the knots  $11n_{61}$ ,  $11n_{63}$ ,  $11n_{72}$ ,  $11n_{84}$ ,  $11n_{85}$ ,  $11n_{90}$ ,  $11n_{92}$ ,  $11n_{98}$ ,  $11n_{99}$ ,  $11n_{101}$ ,  $11n_{103}$ , and  $11n_{112}$  to knots with non-orientable genus 1.




FIGURE 4.14. Non-oriented band moves from the knots  $11n_{125}$ ,  $11n_{130}$ ,  $11n_{131}$ ,  $11n_{133}$ ,  $11n_{137}$ ,  $11n_{138}$ ,  $11n_{140}$ ,  $11n_{141}$ ,  $11n_{155}$ ,  $11n_{161}$ ,  $11n_{165}$ , and  $11n_{171}$  to knots with non-orientable genus 1.




FIGURE 4.15. Non-oriented band moves from the knots  $11n_{176}$ ,  $11n_{179}$ , and  $11n_{184}$  to knots with non-orientable genus 1.

## References

- [1] Samantha Allen. Nonorientable surfaces bounded by knots: a geography problem. *New York J. Math.*, 29:1038–1059, 2023.
- [2] Fraser Binns, Sungkyung Kang, Jonathan Simone, and Paula Truöl. On the nonorientable four-ball genus of torus knots. *arXiv preprint arXiv:2109.09187*, 2021.
- [3] Nakisa Ghanbarian. The non-orientable 4-genus for knots with 10 crossings. *J. Knot Theory Ramifications*, 31(5):Paper No. 2250034, 46, 2022.
- [4] Patrick M. Gilmer and Charles Livingston. The nonorientable 4-genus of knots. *J. Lond. Math. Soc.* (2), 84(3):559–577, 2011.
- [5] C. McA. Gordon and R. A. Litherland. On the signature of a link. *Invent. Math.*, 47(1):53–69, 1978.
- [6] Stanislav Jabuka and Tynan Kelly. The nonorientable 4-genus for knots with 8 or 9 crossings. *Algebr. Geom. Topol.*, 18(3):1823–1856, 2018.
- [7] M. Kate Kearney. The concordance genus of 11-crossing knots. *J. Knot Theory Ramifications*, 22(13):1350077, 17, 2013.
- [8] The Knot Atlas. URL: [http://katlas.org/wiki/The\\_Hoste-Thistlethwaite\\_Table\\_of\\_11\\_Crossing\\_Knots](http://katlas.org/wiki/The_Hoste-Thistlethwaite_Table_of_11_Crossing_Knots), February 2024.
- [9] Charles Livingston and Allison H. Moore. Knotinfo: Table of knot invariants. URL: <http://knotinfo.math.indiana.edu>, January 2024.
- [10] Allison H. Moore and Mariel Vazquez. A note on band surgery and the signature of a knot. *Bull. Lond. Math. Soc.*, 52(6):1191–1208, 2020.
- [11] Hitoshi Murakami and Akira Yasuhara. Crosscap number of a knot. *Pacific J. Math.*, 171(1):261–273, 1995.
- [12] Hitoshi Murakami and Akira Yasuhara. Four-genus and four-dimensional clasp number of a knot. *Proc. Amer. Math. Soc.*, 128(12):3693–3699, 2000.
- [13] Peter S. Ozsváth, András I. Stipsicz, and Zoltán Szabó. Unoriented knot Floer homology and the unoriented four-ball genus. *Int. Math. Res. Not. IMRN*, (17):5137–5181, 2017.
- [14] Joshua M. Sabloff. On a refinement of the non-orientable 4-genus of torus knots. *Proc. Amer. Math. Soc. Ser. B*, 10:242–251, 2023.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY

*Email address:* [mfarr17@lsu.edu](mailto:mfarr17@lsu.edu)