# A NOTE ON EXTENDED GENUS FIELDS OF KUMMER EXTENSIONS OF RATIONAL FUNCTION FIELDS

MARTHA RZEDOWSKI-CALDERÓN AND GABRIEL VILLA-SALVADOR

ABSTRACT. We consider the generalization of the extended genus field of a prime degree cyclic Kummer extension of a rational function field obtained by R. Clement in 1992 to general Kummer extensions. We observe that the same approach of Clement works in general.

## 1. INTRODUCTION

The first one to consider genus fields, or more precisely extended genus fields, of function fields was R. Clement in [2]. She considered a cyclic Kummer extension K of  $k := \mathbb{F}_q(T)$ , a global rational function field of prime degree l, where l|q - 1. She used the ideas of H. Hasse developed in [4], where he considered a quadratic extension K of the field of rational numbers  $\mathbb{Q}$ .

In the case of number fields, both, the genus and the extended genus fields of a number field K are canonically defined as the maximal extension of K of the form KF contained in the Hilbert class and in the extended Hilbert class field, respectively, where F is an abelian extension of  $\mathbb{Q}$ . The Hilbert class field (resp. extended Hilbert class field) of K is defined as the maximal unramified (resp. maximal unramified at the finite primes) abelian extension of K.

In the case of global function fields the situation is quite different. There are several reasonable possible definitions of Hilbert class field of a function field K. The main reason is that, in the number field case, the Hilbert class field is a finite extension of K. In the case of a function field K, the extensions of constants of K are unramified so that the maximal unramified abelian of K is of infinite degree.

For a quadratic number field K of  $\mathbb{Q}$ , the extended Hilbert class field  $K_{H^+}$  of K is the abelian extension such that the fully decomposed primes of K in  $K_{H^+}$  are precisely the nonzero principal prime ideals of K generated by a totally positive element. In the case of a quadratic field K of  $\mathbb{Q}$ , an element is totally positive if and only if its norm in  $\mathbb{Q}$  is the square of a real number. In other words, the norm group in the idèle group  $J_K$  of K is  $K^*(\Delta \times \prod_p U_p)$ , where p runs through the prime numbers,  $U_p$  denotes the group of units of  $\mathbb{Q}_p$  and  $\Delta = \{(\alpha_1, \ldots, \alpha_t) \in \prod_{i=1}^t K_{\mathfrak{p}_i} \mid \prod_{i=1}^t N_{K_{\mathfrak{p}_i}/\mathbb{R}}(\alpha_i) \in \mathbb{R}^{*2}\}$  where  $\{\mathfrak{p}_i\}_{i=1}^t$  is the set of infinite primes in K (t = 1 or 2 and  $K_{\mathfrak{p}_i} \in \{\mathbb{R}, \mathbb{C}\}$ ).

For a cyclic Kummer extension K/k of prime degree l, where  $k = \mathbb{F}_q(T)$  and l|q-1, the analogue is to consider as the norm group of  $K_{H^+}$  the subgroup  $K^*(\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}})$  of the idèle group  $J_K$  of K, where  $\Delta = \{(\alpha_1, \ldots, \alpha_t) \in \prod_{i=1}^t K_{\mathfrak{p}_i} \mid$ 

Date: March 1st., 2024.

<sup>2010</sup> Mathematics Subject Classification. Primary 11R58; Secondary 11R29.

Key words and phrases. Global fields, extended genus fields, Kummer extensions.

 $\prod_{i=1}^{t} N_{K_{\mathfrak{p}_{i}}/k_{\infty}}(\alpha_{i}) \in k_{\infty}^{*l}$  and  $U_{\mathfrak{p}}$  denotes the group of units of  $K_{\mathfrak{p}}$ . This is the approach taken by Clement.

In [3], E. O. Curiel-Anaya, M. R. Maldonado-Ramírez and the first author, use this approach to extend the result to general cyclic Kummer extensions.

In this note we will see that this approach still works for general Kummer extensions K of k and obtain a version of extended genus field for this kind of extensions. In the last section we compare three definitions of extended genus fields for this kind of extensions and present some examples.

# 2. General Kummer extensions of k

Let  $k := \mathbb{F}_q(T)$  and let K be a Kummer extension of k of exponent n|q-1. Then

$$\operatorname{Gal}(K/k) \cong C_{n_1} \times \cdots \times C_{n_s}$$

where in general  $C_m$  denotes the cyclic group of order m and  $n_s | \cdots | n_2 | n_1 = n$ . We have

$$K = k \left( \sqrt[n_1]{\gamma_1 D_1}, \dots, \sqrt[n_s]{\gamma_s D_s} \right) = K_1 \cdots K_s,$$
(2.1)

where  $K_i = k( {}^{n_i} \sqrt{\gamma_i D_i}), \gamma_i \in \mathbb{F}_q^*, D_i \in R_T := \mathbb{F}_q[T]$  is a monic polynomial and  $\operatorname{Gal}(K_i/k) \cong C_{n_i}, 1 \le i \le s$ .

Let  $P_1, \ldots, P_r \in R_T^+ := \{P \in R_T \mid P \text{ is a monic irreducible polynomial}\}$  be the primes of k ramified in K with ramification indices  $e_1, \ldots, e_r$  respectively. Let  $\Delta := \{(\alpha_1, \ldots, \alpha_t) \in \prod_{i=1}^t K_{\mathfrak{p}_i} \mid \prod_{i=1}^t N_{K_{\mathfrak{p}_i}/k_{\infty}}(\alpha_i) \in k_{\infty}^{*n}\}$ , where  $\{\mathfrak{p}_i\}_{i=1}^t$  are the infinite primes in K, that is, the primes above  $\mathcal{P}_{\infty} (= \infty)$ , the infinite prime of kand  $k_{\infty} := k_{\mathcal{P}_{\infty}}$ .

Let  $\mathcal{O}_K := \{ \alpha \in K \mid v_{\mathfrak{p}}(\alpha) \ge 0 \text{ for all } \mathfrak{p} \nmid \infty \}$ . Then  $\mathcal{O}_K$  is a Dedekind domain such that the fractional ideals of  $\mathcal{O}_K$  correspond canonically to the finite primes of K. We denote by  $J_K$  the idèle group of K.

**Definition 2.1.** The *extended Hilbert class field*  $K_{H^+}$  of K is the finite abelian extension of K such that the finite primes of K fully decomposed in  $K_{H^+}$  are precisely the principal nonzero ideals generated by an element whose norm from K to k is an n-th power in  $k_{\infty}$ .

The *extended genus field*  $\Gamma$  of K with respect to k is the maximal abelian extension of k contained in  $K_{H^+}$ .

**Proposition 2.2.** We have that  $[J_K : K^*(\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}})]$  is finite.

*Proof.* The same proof in [2] works in this case. We present the proof for the sake of completeness. Let  $A := \Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}}$  and  $B := \prod_{\mathfrak{p} \mid \infty} K_{\mathfrak{p}}^* \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}}$ . We have

$$K^* \left( \Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}} \right) = K^* A \subseteq K^* B = K^* \left( \prod_{\mathfrak{p} \mid \infty} K_{\mathfrak{p}}^* \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}} \right),$$
$$J_K / K^* B = J_K / K^* \left( \prod_{\mathfrak{p} \mid \infty} K_{\mathfrak{p}}^* \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}} \right) \xrightarrow{\cong}_{\psi} Cl(\mathcal{O}_K)$$

where the isomorphism  $\psi$  is the one induced by the natural map  $\theta: J_K \longrightarrow D_K$ given by  $\theta(\vec{\alpha}) = \prod_{\mathfrak{p}} \mathfrak{p}^{\alpha_{\mathfrak{p}}}$ ,  $D_K$  denotes the divisor group of K, and the ideal class group  $Cl(\mathcal{O}_K)$  is isomorphic to  $D_K/D_{\infty}P_K$ , where  $D_{\infty}$  is the free group generated by the infinite primes and  $P_K$  is the group of principal divisors. We have that  $Cl(\mathcal{O}_K)$  is finite.

2

Now  $A \subseteq B$ . Therefore

$$[K^*B:K^*A] = \frac{[B:A]}{[K^*\cap B:K^*\cap A]} = \frac{[\prod_{\mathfrak{p}\mid\infty} K^*_{\mathfrak{p}} \times \prod_{\mathfrak{p}\nmid\infty} U_{\mathfrak{p}}:\Delta \times \prod_{\mathfrak{p}\restriction\infty} U_{\mathfrak{p}}]}{[K^*\cap B:K^*\cap A]}$$
$$= \frac{[\prod_{\mathfrak{p}\mid\infty} K^*_{\mathfrak{p}}:\Delta]}{[U_K:U_K^{(+)}]},$$

where

 $U_K := K^* \cap B = \{ x \in K^* \mid v_p(x) = 0 \text{ for all } p \nmid \infty \} = \mathcal{O}_K^* \quad (\text{units of } \mathcal{O}_K) \text{ and }$ 

$$U_K^{(+)} := \{ x \in U_K \mid \mathcal{N}_{K/k}(x) = \prod_{\mathfrak{p} \mid \infty} \mathcal{N}_{K\mathfrak{p}/k\infty}(x) \in k_\infty^{*n} \}.$$

Now n|q-1 so that gcd(n,p) = 1 where p is the characteristic of k. Therefore  $(U_{\mathfrak{p}}^{(1)})^n = U_{\mathfrak{p}}^{(1)}$ . Since  $k_{\infty}^* = \langle \pi_{\infty} \rangle \times \mathbb{F}_q^* \times U_{\infty}^{(1)}$  where  $\pi_{\infty} = 1/T$  is a uniformizing parameter, we have  $k_{\infty}^{*n} = \langle \pi_{\infty}^n \rangle \times \mathbb{F}_q^{*n} \times (U_{\infty}^{(1)})^n = \langle \pi_{\infty}^n \rangle \times \mathbb{F}_q^{*n} \times U_{\infty}^{(1)}$ . Thus  $\left|\frac{k_{\infty}^*}{k_{\infty}^{*n}}\right| = n^2$ . Consider

$$\varphi \colon \prod_{\mathfrak{p}\mid\infty} K_{\mathfrak{p}}^* \longrightarrow \frac{k_{\infty}^*}{k_{\infty}^{*n}}, \quad \text{given by} \quad \varphi\big((\alpha_{\mathfrak{p}})_{\mathfrak{p}\mid\infty}\big) = \Big(\prod_{\mathfrak{p}\mid\infty} \mathcal{N}_{K_{\mathfrak{p}}/k_{\infty}}(\alpha_{\mathfrak{p}}) \bmod k_{\infty}^{*n}\Big).$$

Then ker  $\varphi = \Delta$  and  $\varphi$  induces an injection

$$\tilde{\varphi} \colon \frac{\left(\prod_{\mathfrak{p}\mid\infty} K_{\mathfrak{p}}^{*}\right)}{\Delta} \hookrightarrow \frac{k_{\infty}^{*}}{k_{\infty}^{*n}}$$

Hence  $[B:A]|n^2$  and finally  $[J_K:K^*A] < \infty$ .

We have that  $K^*(\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_\mathfrak{p})$  is an open subgroup of finite index of  $J_K$ . Let L be the class field corresponding to  $K^*(\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_\mathfrak{p})$ . Then  $N_{L/K} J_L = K^*(\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_\mathfrak{p})$  and  $\operatorname{Gal}(L/K) \cong \frac{J_K}{K^*(\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_\mathfrak{p})}$ .

**Proposition 2.3.** Let  $\mathfrak{q}$  be a finite prime of K, that is,  $\mathfrak{q}$  is a nonzero prime ideal of  $\mathcal{O}_K$ . Then  $\mathfrak{q}$  decomposes fully in L if and only if  $\mathfrak{q}$  is a principal ideal with  $\mathfrak{q} = \langle \delta \rangle$  and  $N_{K/k}(\delta) \in k_{\infty}^{*n}$ .

*Proof.* From class field theory, we have that q decomposes fully in L/K if and only if  $\theta \lceil K_{\mathfrak{q}}^* \rceil_{\mathfrak{q}} \subseteq K^* (\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}})$ , where  $\theta \lceil x \rceil_{\mathfrak{q}} := (\ldots, 1, x, 1, \ldots)$  for  $x \in K_{\mathfrak{q}}^*$  (see [6, Corolario 17.6.196]) if and only if for all  $x \in K_{\mathfrak{q}}^*$  there exist  $\beta_x \in K^*$  and  $\vec{\alpha}_x \in$  $\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}}$  with  $\theta \lceil x \rceil_{\mathfrak{q}} = \beta_x \vec{\alpha}_x$ . Thus  $\vec{\alpha}_x = (\ldots, \beta_x^{-1}, \ldots, \beta_x^{-1}, \beta_x^{-1}x, \beta_x^{-1}, \ldots)$ . This is equivalent to  $v_{\mathfrak{p}}(\beta_x) = 0$  for all  $\mathfrak{p} \neq \mathfrak{q}$  and  $\mathfrak{p} \nmid \infty$  and  $N_{K/k}(\beta_x^{-1}) =$  $\prod_{\mathfrak{p} \mid \infty} N_{K_{\mathfrak{p}}/k_{\infty}}(\beta_x^{-1}) \in k_{\infty}^{*n}$ .

Let  $x \in K_{\mathfrak{q}}^*$ . The principal ideal in  $\mathcal{O}_K$  generated by  $\beta_x$  satisfies  $\langle \beta_x \rangle = \mathfrak{q}^{n_x} = \beta_x \mathcal{O}_K$ , from some  $n_x \in \mathbb{Z}$ . In particular for  $x \in K_{\mathfrak{q}}^*$  with  $v_{\mathfrak{q}}(\beta_x) = 1$  we have  $\langle \beta_x \rangle = \mathfrak{q}, \beta_x \in K^*$ . Hence  $\mathfrak{q}$  is a principal ideal,  $\mathfrak{q} = \langle \beta_x \rangle$  and  $N_{K/k}(\beta_x) \in k_{\infty}^{*n}$ .  $\Box$ 

**Corollary 2.4.** The extended class field  $K_{H^+}$  of K is the class field corresponding to  $K^*(\Delta \times \prod_{\mathfrak{q} \nmid \infty} U_{\mathfrak{q}})$  and a finite prime  $\mathfrak{q}$  of K decomposes fully in  $K_{H^+}$  if and only if  $\mathfrak{q}$  is a principal ideal and the norm from K to k of a generator belongs to  $k_{\infty}^{*n}$ .

**Proposition 2.5.** The field of constants of  $K_{H^+}$  is  $\mathbb{F}_{q^n}$ .

*Proof.* We have (see [6, Teorema 17.6.192]) that the field of constants of  $K_{H^+}$  is  $\mathbb{F}_{q^d}$  where  $d = \min\{ \deg \vec{\alpha} \mid \vec{\alpha} \in \Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}} \text{ and } \deg \vec{\alpha} > 0 \}$ . We will see that d = n.

On the one hand, since  $\prod_{\mathfrak{p}\mid\infty} N_{K_{\mathfrak{p}}/k_{\infty}}^{*} K_{\mathfrak{p}}^{*} = k_{\infty}^{*n}$ , there exists  $\vec{\alpha} \in \Delta \times \prod_{\mathfrak{p}\nmid\infty} U_{\mathfrak{p}}$ with  $v_{\infty} \left( \prod_{\mathfrak{p}\mid\infty} N_{K_{\mathfrak{p}}/k_{\infty}}(\alpha_{\mathfrak{p}}) \right) = v_{\infty}(\pi_{\infty}^{n}) = n$ . On the other hand, let  $\vec{\alpha} \in \Delta \times \prod_{\mathfrak{p}\nmid\infty} U_{\mathfrak{p}}$ . Then  $v_{\mathfrak{p}}(\alpha_{\mathfrak{p}}) = 0$  for all  $\mathfrak{p} \nmid \infty$  and  $\xi = \prod_{\mathfrak{p}\mid\infty} N_{K_{\mathfrak{p}}/k_{\infty}}(\alpha_{\mathfrak{p}}) \in k_{\infty}^{*n}$ . Therefore (see [6, Proposición 17.3.38])

$$\deg \vec{\alpha} = \sum_{\mathfrak{p}} \deg \mathfrak{p} \cdot v_{\mathfrak{p}}(\alpha_{\mathfrak{p}}) = \sum_{\mathfrak{p} \mid \infty} \deg \mathfrak{p} \cdot v_{\mathfrak{p}}(\alpha_{\mathfrak{p}}) = \sum_{\mathfrak{p} \mid \infty} f_{\mathfrak{p}} v_{\mathfrak{p}}(\alpha_{\mathfrak{p}})$$
$$= \sum_{\mathfrak{p} \mid \infty} v_{\infty} \left( N_{K_{\mathfrak{p}}/k_{\infty}}(\alpha_{\mathfrak{p}}) \right) = v_{\infty} \left( \prod_{\mathfrak{p} \mid \infty} N_{K_{\mathfrak{p}}/k_{\infty}}(\alpha_{\mathfrak{p}}) \right) = v_{\infty}(\xi) = na$$

for some  $a \in \mathbb{Z}$ . Hence d = n.

**Proposition 2.6.** Every abelian extension L/k such that  $K \subseteq L$ , L/K is unramified at the finite primes and such that Gal(L/k) is of exponent a divisor of n, is contained in  $K_{H^+}$ .


*Proof.* Since  $K_{H^+}$  is the class field of  $\Delta \times \prod_{\mathfrak{p}\nmid\infty} U_{\mathfrak{p}}$  (Corollary 2.4), we have that  $K_{H^+}/K$  is unramified at the finite primes.

Let L/k be an abelian extension with  $K \subseteq L$ , such that L/K is unramified at the finite primes and such that Gal(L/K) is of exponent a divisor of n.

Let  $N_{L/K} J_L = K^* \mathcal{H} \subseteq J_K$  for some  $\mathcal{H}$ . If  $\mathfrak{q}$  is a finite prime, then  $\mathfrak{q}$  is unramified in L/K so that  $U_{\mathfrak{q}} \subseteq \mathcal{H}$  (see [6, Corolario 17.6.196]). If  $\vec{\alpha} \in \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}}$ , and  $(\underline{\ }, E/F)_{\mathfrak{p}}$ denotes the Artin reciprocity map, we have (see [6, Teorema 17.6.149])

$$(\vec{\alpha},L/K) = \prod_{\mathfrak{p}\mid\infty} (1,L_{\mathfrak{P}}/K_{\mathfrak{p}}) \times \prod_{\mathfrak{p}\nmid\infty} (\alpha_{\mathfrak{p}},L_{\mathfrak{P}}/K_{\mathfrak{p}}) = 1,$$

where  $\mathfrak{P}$  is a prime in *L* above  $\mathfrak{p}$ . It follows that  $\vec{\alpha} \in \mathcal{H}$  and  $\prod_{\mathfrak{p}\nmid\infty} U_{\mathfrak{p}} \subseteq \mathcal{H}$ . If  $\vec{\alpha} \in \Delta$  we have the commutative diagram



where  $\iota$  is the natural embedding. We have that  $N_{K/k}(\Delta) = \{(\delta, 1, ..., 1, ...) \mid \delta \in k_{\infty}^{*n}\}$ . Since  $\operatorname{Gal}(L/k)$  is of exponent n, we have that  $N_{K/k}(\Delta) \subseteq \ker(\underline{., L/k})$  so that  $\Delta \subseteq \ker(\underline{., L/K}) = K^* \mathcal{H}$ . Hence  $\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}} \subseteq K^* \mathcal{H}$  and  $L \subseteq K_{H^+}$ .  $\Box$ 

**Proposition 2.7.** The extension  $K_{H^+}/k$  is a Galois extension.

*Proof.* It is the same as the one in [2]. It follows from the fact that  $\sigma(\Delta \times \prod_{\mathfrak{p}\nmid \infty} U_{\mathfrak{p}}) = \Delta \times \prod_{\mathfrak{p}\nmid \infty} U_{\mathfrak{p}}$  for each *k*-embedding  $\sigma$  of  $K_{H^+}$  in an algebraic closure  $\bar{k}$  of *k*.  $\Box$ 

The following result is the analogue of Lemma 1.5 and Proposition 1.6 in [2]. The same proof works in our more general setting.


**Proposition 2.8.** Let  $I_K$  be the group of fractional ideals of  $\mathcal{O}_K$  and  $P_K$  the subgroup of principal ideals. Let  $J_K^{(+)} = \{\vec{\alpha} \in J_K \mid (\alpha_{\mathfrak{p}})_{\mathfrak{p}|\infty} \in \Delta\}, K^{(+)} = K^* \cap J_K^{(+)}$  and  $P_K^{(+)} = \{\langle \beta \rangle \in P_K \mid \beta \in K^{(+)}\}$ . Then

$$\frac{J_K}{K^*(\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}})} \cong \frac{J_K^{(+)}}{K^{(+)}(\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}})} \cong \frac{I_K}{P_K^{(+)}} =: Cl(\mathcal{O}_K)^{(+)}.$$

Next we recall a result from class field theory.

**Proposition 2.9.** Let E/F be a finite extension of global fields. Let N be an open subgroup of finite index of  $J_E$  such that  $E^* \subseteq N$ . Let  $E_N$  be the class field of N, that is,  $N_{E_N/E}(J_{E_N}) = N$ . Let  $F_0$  be the maximal abelian extension of F contained in  $E_N$ . Then the norm group of  $F_0$  in  $J_F$  is  $F^* N_{E/F}(N)$ .

*Proof.* The norm  $N_{E/F}: J_E \longrightarrow J_F$  is an open map so that  $N_{E/F}(N)$  is an open subgroup of  $J_F$ . Since N is of finite index in  $J_E$ , we have that  $F^* N_{E/F}(N)$  is of finite index in  $J_F$ .



To show that  $F_0$  is the class field of  $F^* N_{E/F}(N)$ , it suffices to show that the group  $N_{E/F}(N)$  is contained in any open subgroup of finite index in  $J_F$  such that its class field F' is contained in  $E_N$ .

Let N' be an open subgroup of finite index in  $J_E$  such that its class field is  $EF' \subseteq E_N$ . Since  $N \subseteq N'$ , it follows that  $N_{E/F}(N) \subseteq N_{E/F}(N')$ . We have the commutative diagram (see [6, Teorema 17.6.137])

$$\begin{array}{c|c} J_E & & \underbrace{(\_, EF'/E)}_{N_{E/F}} & & \operatorname{Gal}(EF'/E) \\ \downarrow & & & \downarrow^{\operatorname{rest}} \\ J_F & & & \underbrace{(\_, F'/F)}_{(\_, F'/F)} & \operatorname{Gal}(F'/F) \end{array}$$

Therefore  $N_{E/F}(N)$  is contained in the subgroup of  $J_F$  corresponding to F'.  $\Box$ 

As a direct consequence of Proposition 2.9 we have

**Corollary 2.10.** The extended genus field  $\Gamma$  of K/k is the class field of  $k^* \operatorname{N}_{K/k} (\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}})$  and

$$[\Gamma:k] = |\operatorname{Gal}(\Gamma/k)| = [J_k:k^* \operatorname{N}_{K/k} (\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}})].$$

To obtain  $\Gamma$ , we have to compute  $N_{K/k} (\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}})$ . First we prove two lemmas.

**Lemma 2.11.** We have  $J_k = k^* (k_\infty^* \times \prod_{P \in R_m^+} U_P)$ .

## M. RZEDOWSKI AND G. VILLA

*Proof.* Let  $\vec{\beta} = (\beta_{\infty}, \beta_P)_{P \in R_T^+} \in J_k$ . Let  $Q_1, \ldots, Q_t \in R_T^+$  be the finite primes such that  $c_i = v_{Q_i}(\beta_{Q_i}) \neq 0$ . We have that  $v_P(\beta_P) = 0$  for all  $P \notin \{Q_1, \ldots, Q_t\}$ . Let  $f \in k^*$  be defined by  $f = \prod_{i=1}^t Q_i^{c_i}$  (f = 1 if t = 0). Then  $f^{-1}\vec{\beta} \in k_{\infty}^* \times \prod_{P \in R_T^+} U_P$ . The result is obtained.

Another proof is as follows, the class field F of  $k^*(k_{\infty}^* \times \prod_{P \in R_T^+} U_P)$  is unramified over k and  $\mathcal{P}_{\infty}$  decomposes fully. Hence F is an extension of constants and the infinite prime (of degree one) decomposes fully in F. Therefore F = k.  $\Box$ 

**Lemma 2.12.** Let  $\mathfrak{p}$  be a finite prime in K. Let P be the irreducible polynomial in  $R_T$  corresponding to the prime in k below  $\mathfrak{p}$ .

- (1) If  $\mathfrak{p}$  is unramified, then  $N_{K_{\mathfrak{p}}/k_P}(U_{\mathfrak{p}}) = U_P$ .
- (2) If  $\mathfrak{p}$  is ramified with ramification index  $e_P$ , then  $N_{K_{\mathfrak{p}}/k_P}(U_{\mathfrak{p}}) = \mathbb{F}_P^{*e_P} \times U_P^{(1)}$ , where  $\mathbb{F}_P$  is the field of constants of  $k_P$ .

*Proof.* For any extension E/F of local fields, we have  $e_{E/F} = [U_F : N_{E/F}(U_E)]$  (see [6, Corolario 17.3.39]). If E/F is tamely ramified,  $N_{E/F}(U_E^{(1)}) = U_F^{(1)}$ . The result follows.

Proposition 2.13. We have

$$[\Gamma:k] = \left[J_k: k^* \operatorname{N}_{K/k} \left(\Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}}\right)\right] = n \prod_{i=1}^r e_i$$

where  $e_i$  is the ramification index of  $P_i$  in K/k,  $1 \le i \le r$ .

*Proof.* From Lemmas 2.11 and 2.12, we obtain

$$\begin{bmatrix} J_k : k^* \operatorname{N}_{K/k} \left( \Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}} \right) \end{bmatrix}$$
  
=  $\begin{bmatrix} k^* \left( \Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}} \right) : k^* \left( k_{\infty}^{*n} \times \prod_{i=1}^r \left( \mathbb{F}_{P_i}^{*e_i} \times U_{P_i}^{(1)} \right) \times \prod_{\substack{P \in R_T^+ \\ P \notin \{P_1, \dots, P_r\}}} U_P \right) \end{bmatrix}$ 

Set  $C := \Delta \times \prod_{\mathfrak{p} \nmid \infty} U_{\mathfrak{p}}$  and  $D := k_{\infty}^{*n} \times \prod_{i=1}^{r} \left( \mathbb{F}_{P_{i}}^{*e_{i}} \times U_{P_{i}}^{(1)} \right) \times \prod_{\substack{P \in R_{T}^{+} \\ P \notin \{P_{1}, \dots, P_{r}\}}} U_{P}.$ 

Then,

$$[k^*C:k^*D] = \frac{[C:D]}{[k^* \cap C:k^* \cap D]}$$

and

$$\frac{C}{D} \cong \frac{k_{\infty}^*}{k_{\infty}^{*n}} \times \prod_{i=1}^r \frac{\mathbb{F}_{P_i}^* \times U_{P_i}^{(1)}}{\mathbb{F}_{P_i}^{*e_i} \times U_{P_i}^{(1)}} \times \prod_{\substack{P \in R_T^+ \\ P \notin \{P_1, \dots, P_r\}}} \frac{U_P}{U_P} \cong C_n^2 \times \prod_{i=1}^r C_{e_i}.$$

Consider now  $\vec{\alpha} \in k^* \cap C$ . Then  $\alpha_P \in U_P$  for all  $P \in R_T^+$ . Thus  $v_P(\alpha_P) = 0$  for all  $P \in R_T^+$ . It follows that  $v_{\infty}(\alpha_{\infty}) = 0$  and that  $\vec{\alpha} = x \in \mathbb{F}_q^*$ . Thus  $k^* \cap C = \mathbb{F}_q^*$ . Similarly  $k^* \cap D = \mathbb{F}_q^{*n}$ . Therefore  $[k^* \cap C : k^* \cap D] = n$ . It follows that  $[k^*C : k^*D] = \frac{n^2 \times \prod_{i=1}^r e_i}{n} = n \cdot \prod_{i=1}^r e_i$ .

We have the main result of this note.

**Theorem 2.14.** Let K be given by (2.1). Then the extended genus field of K/k is

$$\Gamma = \mathbb{F}_{q^n} \left( T, \sqrt[e_1]{P_1}, \sqrt[e_2]{P_2}, \dots, \sqrt[e_r]{P_r} \right).$$

*Proof.* Let  $M = \mathbb{F}_{q^n}(T, \sqrt[e_1]{P_1}, \sqrt[e_2]{P_2}, \dots, \sqrt[e_r]{P_r})$ . Then  $[M : k] = n \cdot \prod_{i=1}^r e_i$ . From Proposition 2.13 we obtain that  $[\Gamma : k] = [M : k]$  and from Proposition 2.6 we obtain that  $M \subseteq K_{H^+}$ . Since M is abelian and  $\Gamma$  is the maximal abelian extension of k contained in  $K_{H^+}$ , it follows that  $M \subseteq \Gamma$ . Hence  $\Gamma = M$ .

**Remark 2.15.** We observe that the extended genus field  $\Gamma$  of a Kummer extension K/k is also a Kummer extension and furthermore that the extended genus field  $\Gamma/k$  is  $\Gamma$  itself.

# 3. COMPARISON AMONG VARIOUS EXTENDED GENUS FIELDS

Let *K* be given by (2.1). We have some different definitions of extended genus fields. We compare three of them.

First we denote by  $K_{\mathfrak{ger}, \text{clement}}$  the extended genus field obtained here, that is,  $K_{\mathfrak{ger}, \text{clement}} = \Gamma = \mathbb{F}_{q^n} (T, \sqrt[e_1]{P_1}, \sqrt[e_2]{P_2}, \dots, \sqrt[e_r]{P_r}).$ 

In [5] the extended genus field of K was defined as  $K_{\mathfrak{ger}, \mathrm{rarzvi}} = E_{\mathfrak{ger}, \mathrm{rarzvi}}K$ , where we have  $E = k \left( \sqrt[n_1]{(-1)^{\deg D_1} D_1}, \sqrt[n_2]{(-1)^{\deg D_2} D_2}, \ldots, \sqrt[n_1]{(-1)^{\deg D_s} D_s} \right)$  and then

$$E_{geg,rarzvi} = k \Big( \sqrt[e_1]{(-1)^{\deg P_1} P_1}, \sqrt[e_2]{(-1)^{\deg P_2} P_2}, \dots, \sqrt[e_r]{(-1)^{\deg P_r} P_r} \Big).$$

Therefore

$$K_{\mathfrak{ge}\mathfrak{g},\mathrm{rarzvi}} = \mathbb{F}_q \big( T, \ {}^{e_1}\sqrt{\epsilon_1}, \ {}^{e_2}\!\sqrt{\epsilon_2}, \ldots, \ {}^{e_r}\!\sqrt{\epsilon_r} \big) \big( \ {}^{e_1}\!\sqrt{P_1}, \ {}^{e_2}\!\sqrt{P_2}, \ldots, \ {}^{e_r}\!\sqrt{P_r} \big),$$

where  $\epsilon_i = (-1)^{\deg P_i} \gamma_i$ ,  $1 \leq i \leq r$ . Since  $e_i | n$  we have  $\mathbb{F}_q(e_i \sqrt{\epsilon_1}, e_i \sqrt{\epsilon_2}, \dots, e_i \sqrt{\epsilon_r}) \subseteq \mathbb{F}_{q^n}$ . Therefore  $K_{\mathfrak{ger}, \operatorname{rarzvi}} \subseteq K_{\mathfrak{ger}, \operatorname{clement}}$ .

Next, using the definition of extended genus field given by Anglès and Jaulent in [1], it was obtained in [7] that  $K_{geg,angjau} = FK$  for some subfield of  $E_{geg,rarzvi}$ . Therefore  $K_{geg,angjau} \subseteq K_{geg,rarzvi}$ .

Finally we have

$$K_{\mathfrak{ger}, \mathrm{angjau}} \subseteq K_{\mathfrak{ger}, \mathrm{rarzvi}} \subseteq K_{\mathfrak{ger}, \mathrm{clement}}.$$

We also remark that while  $K_{ger,clement}$  is defined just for Kummer extensions K/k,  $K_{ger,rarzvi}$  and  $K_{ger,angjau}$  are defined for any separable finite extension K/k.

- **Examples 3.1.** (1) Let  $K := k(\sqrt[l]{\gamma P})$  where l is a prime number such that l|q-1,  $P \in R_T^+$  with deg P < l and  $\gamma \in \mathbb{F}_q^* \setminus (\mathbb{F}_q^*)^l$ . Then  $K_{\mathfrak{geg}, \operatorname{angjau}} = K_{\mathfrak{geg}, \operatorname{rarzvi}}$  (see [7, Example 5.2]).
  - (2) Let l be a prime number such that  $l^2|q-1$  and let  $K = k(\sqrt[l^2]{\gamma D})$  where  $\gamma \in \mathbb{F}_q^*$ ,  $D = P_1^{\alpha_1} \cdots P_r^{\alpha_r} \in R_T$ . Then  $K_{\mathfrak{ger}, \mathrm{angjau}} = K_{\mathfrak{ger}, \mathrm{rarzvi}}$ . (see [7, Example 5.3]).
  - (3) Now, let *l* be a prime number such that *l*|*q* − 1 and let *K* := *k*(<sup>*l*</sup>√(−1)<sup>deg P</sup>P) where *P* ∈ *R*<sup>+</sup><sub>T</sub> and deg *P* denotes the degree of *P*. Then *K*<sub>ger,rarzvi</sub> = *K* ⊆ *K*<sub>*l*</sub> = *K*<sub>ger,clement</sub>, where *K*<sub>*l*</sub> = *K*𝔽<sub>*q*<sup>*l*</sup></sub> denotes the extension of constants of degree *l* of *K*.

**Remark 3.2.** It is very likely that, when K/k is abelian, we have  $K_{geg,angjau} = K_{geg,rarzvi}$ .

#### M. RZEDOWSKI AND G. VILLA

## References

- Anglès, Bruno; Jaulent, Jean–François, *Théorie des genres des corps globaux*, Manuscripta Math. 101, no. 4, 513–532 (2000).
- [2] Clement, Rosario, *The genus field of an algebraic function field*, J. Number Theory 40, no. 3, 359–375, (1992).
- [3] Curiel-Anaya, Edgar Omar, Maldonado-Ramírez, Myriam Rosalía & Rzedowski-Calderón, Martha, Extended genus field of cyclic Kummer extensions of rational function fields, Palestine Journal of Mathematics, 12, No. 4, 440–451, (2023).
- [4] Hasse, Helmut, Zur Geschlechtertheorie in quadratischen Zahlkörpern, J. Math. Soc. Japan 3, 45–51, (1951).
- [5] Ramírez–Ramírez, Elizabeth; Rzedowski-Calderón, Martha & Villa–Salvador, Gabriel, Genus fields of global fields, Palestine Journal of Mathematics 9, no. 2, 999–1019 (2020).
- [6] Rzedowski-Calderón, Martha & Villa-Salvador, Gabriel, Campos ciclotómicos numéricos y de funciones (tercera versión), https://arxiv.org/abs/1407.3238.
- [7] Rzedowski–Calderón, Martha & Villa–Salvador, Gabriel, Class fields, Dirichlet characters, and extended genus fields of global function fields, Math. Nachr. 296, 3606–3618 (2023).

DEPARTAMENTO DE CONTROL AUTOMÁTICO, CENTRO DE INVESTIGACIÓN Y DE ESTUDIOS AVAN-ZADOS DEL I.P.N.

Email address: mrzedowski@ctrl.cinvestav.mx

DEPARTAMENTO DE CONTROL AUTOMÁTICO, CENTRO DE INVESTIGACIÓN Y DE ESTUDIOS AVAN-ZADOS DEL I.P.N.

Email address: gvillasalvador@gmail.com, gvilla@ctrl.cinvestav.mx

#### 8