

FIRST EIGENVALUE CHARACTERIZATION OF CLIFFORD HYPERSURFACES AND VERONESE SURFACES

PEI-YI WU

ABSTRACT. We give an estimate for the first eigenvalue of the Schrödinger operator $L := -\Delta - \sigma$ which defined on the closed minimal submanifold M^n in the unit sphere \mathbb{S}^{n+m} , where σ is the square norm of the second fundamental form.

1. INTRODUCTION

The study of rigidity theorems plays an important role in the theory of minimal submanifolds. Also there has been extensive research in the mathematics literature on rigidity theorems for minimal submanifolds in Spheres since the pioneering results obtained by Simons ([9]), Lawson ([4]) and Chern-Do Carmo-Kobayashi ([3]). To be precise, denote σ as the square norm of second fundamental form, Let M^n be a compact minimal submanifold in a unit sphere \mathbb{S}^{n+m} , they proved that if $0 \leq \sigma \leq \frac{n}{2-\frac{1}{m}}$, then either $\sigma = 0$ or $\sigma = \frac{n}{2-\frac{1}{m}}$ and M is the Clifford hypersurface or the Veronese surface in \mathbb{S}^4 . Later, Li [1] and Xu-Chen [2] improved the pinching number $\frac{n}{2-\frac{1}{m}}$ to $\frac{2n}{3}$ respectively, they proved that if $0 \leq \sigma \leq \frac{2n}{3}$, then either $\sigma = 0$ or $\sigma = \frac{2n}{3}$ and M is the Veronese surface in \mathbb{S}^4 . Recently, Lu generalized their result and got the following rigidity theorem, denote by λ_2 the second largest eigenvalue of the fundamental matrix(see Definition 2.3),

Theorem A (Lu). [5] *Let*

$$0 \leq \sigma + \lambda_2 \leq n.$$

Then either M is totally geodesic, or is one of the Clifford hypersurfaces $M_{r,n-r}$ ($1 \leq r \leq n$) in \mathbb{S}^{n+m} , $m \geq 1$, or is a Veronese surface in \mathbb{S}^{2+m} , $m \geq 2$.

Remark 1. Lu suggests that quantity $\sigma + \lambda_2$ might be the right object to study pinching theorems.

In this paper, using Lu inequality [5][Lemma 2] (see Lemma 2.2), we study the first eigenvalue of the Schrödinger operator $L := -\Delta + q$, where q is some continuous function on M . If there is a nonzero $f \in C^\infty(M)$ satisfying equation $Lf = \mu f$, we call μ the eigenvalue of L . Since Δ is elliptic, so is L . And we have

$$\text{Spec}(L) = \{\mu_i; \mu_1 < \mu_2 \leq \mu_3 \leq \dots\}.$$

we call μ_1 the first eigenvalue of L .

The pinching theorems mentioned above gives a characterization of Clifford hypersurfaces and Veronese Surfaces and their proof shares the same spirit of making good use of the Simons identity. Also mathematicians found that one can use similar argument to give an estimate of the first eigenvalue of the Schrödinger operator, which also can be seen as a new way to characterize the Clifford hypersurfaces and Veronese Surfaces. First, Simons [9] studied the Schrödinger operator $L_I := -\Delta - \sigma$ of minimal hypersurfaces $M^n \rightarrow \mathbb{S}^{n+1}$, proved that its first eigenvalue $\mu_1^I \leq -n$ if M is not totally geodesic. Later, Wu [10] and Perdomo [7] proved that if $\mu_1^I \geq -n$, then M is either totally geodesic or a Clifford hypersurfaces independently. And Wu [10] also got the following results: Define $L_{II} := -\Delta - (2 - \frac{1}{m})\sigma$ of minimal submanifold $M^n \rightarrow \mathbb{S}^{n+m}$ and $L_{III} := -\Delta - \frac{3}{2}\sigma$ of minimal submanifold $M^n \rightarrow \mathbb{S}^{n+m}, m \geq 2$, denote μ_1^{II} and μ_1^{III} by their first eigenvalue respectively. Then, for L_{II} , $\mu_1^{II} \leq -n$ if M is not totally geodesic, and if $\mu_1^{II} \geq -n$, then M is either totally geodesic or $\mu_1^{II} = -n$ and M is either the Clifford hypersurfaces or the Veronese surfaces. For L_{III} , $\mu_1^{III} \leq -n$ if M is not totally geodesic, and if $\mu_1^{III} \geq -n$, then M is either totally geodesic or $\mu_1^{III} = -n$ and M is the Veronese surfaces. Similar result also happens in Legendrian case, by using a pinching rigidity result in [6], Yin-Qi [11] got a sharp estimate for the first eigenvalue of Schrödinger operator defined on minimal Legendrian submanifold $M^3 \rightarrow \mathbb{S}^7$.

Inspired by the correspondence between the pinching theorems and the estimates of the first eigenvalue of certain Schrödinger operators, one expect that these phenomenon would happen to Lu's rigidity theorem A and that leads to our main theorem. Define the Schrödinger operator as

$$L := -\Delta - \sigma$$

and denoted the first eigenvalue of L by μ_1 , our main theorem is

Theorem 1.1 (Main Theorem). *Let M^n be a closed minimal submanifold in $\mathbb{S}^{n+m}(1)$, then*

$$\mu_1 \leq -n + \max_{p \in M} \lambda_2 \tag{1.1}$$

if M is not totally geodesic. Moreover, if $\mu_1 \geq -n + \max_{p \in M} \lambda_2$, then either $\mu_1 = 0$ and M is totally geodesic or $\mu_1 = -n + \max_{p \in M} \lambda_2$ and M is the Clifford hypersurfaces in $\mathbb{S}^{n+m}(1)$ or is the Veronese surface in $\mathbb{S}^{2+m}(1)$.

2. PRELIMINARIES AND LU'S INEQUALITY

Let M^n be a compact minimal submanifold in a unit sphere \mathbb{S}^{n+m} . We shall make use of the following convention on the range of indices:

$$1 \leq A, B, C, \dots \leq n + m; \quad 1 \leq i, j, k, \dots \leq n; \quad n + 1 \leq \alpha, \beta, \gamma, \dots \leq n + m.$$

We choose a local field of orthonormal frames $\{e_1, e_2, \dots, e_{n+m}\}$ in \mathbb{S}^{n+m} such that when restricting on M , the $\{e_1, e_2, \dots, e_n\}$ are tangent to M and $\{e_{n+1}, e_{n+2}, \dots, e_{n+m}\}$ are normal to M . Also $\{\omega_1, \dots, \omega_{n+m}\}$ is the corresponding dual frame. It is well know that

$$\begin{aligned}\omega_{\alpha i} &= \sum_j h_{ij}^\alpha \omega_j, \quad h_{ij}^\alpha = h_{ji}^\alpha, \\ h &= \sum_{\alpha, i, j} h_{ij}^\alpha \omega_i \otimes \omega_j \otimes e_\alpha, \quad H = \frac{1}{n} \sum_{\alpha, i} h_{ii}^\alpha e_\alpha, \\ R_{ijkl} &= \delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk} + \sum_\alpha (h_{ik}^\alpha h_{jl}^\alpha - h_{il}^\alpha h_{jk}^\alpha),\end{aligned}\tag{2.1}$$

$$R_{\alpha\beta kl} = \sum_i (h_{ik}^\alpha h_{il}^\beta - h_{il}^\alpha h_{ik}^\beta),\tag{2.2}$$

$$h_{ijk}^\alpha = h_{ikj}^\alpha,\tag{2.3}$$

where $h, H, R_{ijkl}, R_{\alpha\beta kl}$, are the second fundamental form, the mean curvature vector, the curvature tensor, and the normal curvature tensor of M respectively. We define

$$\sigma = |h|^2, \quad A_\alpha = (h_{ij}^\alpha)_{n \times n}.$$

Denote by h_{ijk}^α the component of the covariant derivative of h_{ij}^α , defined by

$$h_{ijk}^\alpha \omega_k = dh_{ij}^\alpha - \sum_l h_{il}^\alpha \omega_l j - \sum_l h_{lj}^\alpha \omega_l i + \sum_\beta h_{ij}^\beta \omega_{\alpha\beta}.\tag{2.4}$$

From Gauss-Codazzi-Ricci equation above, the following Simons identity is well known,

$$\sum_{i,j} h_{ij}^\alpha \Delta h_{ij}^\alpha = n |A_\alpha|^2 + \sum_t \text{Tr}(A_\alpha A_\beta - A_\beta A_\alpha)^2 - \sum_t (\text{Tr } A_\alpha A_\beta)^2\tag{2.5}$$

Now, we introduce the Lu inequality [5][Lemma 2](see Lemma 2.2), which is the main tool in the proof of Theorem A, and the Theorem 1.1. And the proof of the Lu inequality relies on a algebraic inequality [5][Lemma 1], which we find that there are more cases when the equality holds and use Lagrange Multiplier method to give another proof. To be precise, we restate Lu's lemma [5][Lemma 1] as follows

Lemma 2.1. *Suppose η_1, \dots, η_n are real numbers and*

$$\eta_1 + \dots + \eta_n = 0, \quad \eta_1^2 + \dots + \eta_n^2 = 1.$$

Let $r_{ij} \geq 0$ be nonnegative numbers for $i < j$. Then we have

$$\sum_{i < j} (\eta_i - \eta_j)^2 r_{ij} \leq \sum_{i < j} r_{ij} + \text{Max}(r_{ij}).\tag{2.6}$$

If $\eta_1 \geq \dots \geq \eta_n$, and r_{ij} are not simultaneously zero, then the equality in (2.6) holds in one of the following cases: Fix a integer $k, k \in \{1, \dots, n-1\}$,

(i) $r_{ij} = 0$ if $2 \leq i < j$, $r_{12} = \cdots = r_{1k} = 0, r_{1k+1} = \cdots = r_{1n} > 0$ and

$$\eta_1 = \frac{\sqrt{n-k}}{\sqrt{n-k+1}}, \quad \eta_2 = \cdots = \eta_k = 0, \quad \eta_{k+1} = \cdots = \eta_n = \frac{-1}{\sqrt{(n-k+1)(n-k)}}.$$

(ii) $r_{ij} = 0$ if $i < j < n$, $r_{n-1n} = \cdots = r_{n-k+1n} = 0, r_{n-kn} = \cdots = r_{1n} > 0$

$$\eta_n = \frac{-\sqrt{n-k}}{\sqrt{n-k+1}}, \quad \eta_{n-1} = \eta_{n-2} = \cdots = \eta_{n-k+1} = 0, \quad \eta_{n-k} = \cdots = \eta_1 = \frac{1}{\sqrt{(n-k+1)(n-k)}}.$$

Remark 2. We prove this by two steps. Our first step is the same as Lu's original proof [5][Lemma 1] which reduces the problem to proving the following inequality

$$\sum_{1 < j} (\eta_1 - \eta_j)^2 r_{1j} \leq \sum_{1 < j} r_{1j} + \text{Max}_{1 < j} (r_{1j}).$$

Then, we apply Lagrange Multiplier method to prove this inequality, and for reader's convenience, we will also include the first step in our proof.

Proof. First step. Assume $\eta_1 \geq \cdots \geq \eta_n$. If $\eta_1 - \eta_n \leq 1$ or $n = 2$, then (2.6) is trivial. So we assume $n > 2$, and

$$\eta_1 - \eta_n > 1.$$

Observing that $\eta_i - \eta_j < 1$ for $2 \leq i < j \leq n-1$. Otherwise, one could have

$$1 \geq \eta_1^2 + \eta_n^2 + \eta_i^2 + \eta_j^2 \geq \frac{1}{2}((\eta_1 - \eta_n)^2 + (\eta_i - \eta_j)^2) > 1,$$

which is a contradiction.

Using the same reasoning, if $\eta_1 - \eta_{n-1} > 1$, then we have $\eta_2 - \eta_n \leq 1$; and if $\eta_2 - \eta_n > 1$, then we have $\eta_1 - \eta_{n-1} \leq 1$. Replacing η_1, \cdots, η_n by $-\eta_n, \cdots, -\eta_1$ if necessary, we can always assume that $\eta_2 - \eta_n \leq 1$. Thus $\eta_i - \eta_j \leq 1$ if $2 \leq i < j$, and (2.6) is implied by the following inequality

$$\sum_{1 < j} (\eta_1 - \eta_j)^2 r_{1j} \leq \sum_{1 < j} r_{1j} + \text{Max}_{1 < j} (r_{1j}). \quad (2.7)$$

Before proving the inequality (2.7), we can observe that if the equality in (2.6) holds, we must have $\eta_1 - \eta_n > 1$, otherwise,

$$\sum_{i < j} (\eta_i - \eta_j)^2 r_{ij} \leq \sum_{i < j} r_{ij} < \sum_{i < j} r_{ij} + \text{Max}(r_{ij}),$$

which is a contradiction.

Notice that, when $\eta_1 - \eta_n > 1$, by the discussion above, we have $\eta_i - \eta_j < 1$, for $2 \leq i < j < n$. So, we have $r_{ij} = 0$, for $2 \leq i < j < n$, otherwise, by (2.7) we know that the equality in (2.6) will not hold. Thus, when discussing how the equality holds in (2.6), we only need to analyse the situation of inequality (2.7)(Again we should remind the reader that above analysis comes

from Lu's original proof [5][Lemma 1])

Second step. Let $s_j = r_{1j}$, where $j = 2, \dots, n$, then we write (2.7) as follow

$$\sum_{1 < j} (\eta_1 - \eta_j)^2 s_j \leq \sum_{1 < j} s_j + \text{Max}_{1 < j} (s_j). \quad (2.8)$$

Denote

$$f(\eta_1, \eta_2, \dots, \eta_{n-1}, \eta_n) = \sum_{1 < j} (\eta_1 - \eta_j)^2 s_j$$

we apply Lagrange multiplier method to f with constraints

$$\eta_1 + \dots + \eta_n = 0, \quad \eta_1^2 + \dots + \eta_n^2 - 1 = 0. \quad (2.9)$$

Consider the function

$$\Phi(\eta_1, \eta_2, \dots, \eta_{n-1}, \eta_n) = \sum_{1 < j} (\eta_1 - \eta_j)^2 s_j + \lambda (\eta_1 + \dots + \eta_n) + \mu (\eta_1^2 + \dots + \eta_n^2 - 1)$$

where λ and μ are the Lagrange multipliers.

Partial differentiating with respect to each variable, and let them equal to zero, we obtain the following equations

$$\frac{\partial \Phi}{\partial \eta_1} = \sum_{1 < j} 2(\eta_1 - \eta_j) s_j + \lambda + 2\mu \eta_1 = 0, \quad (2.10)$$

$$\frac{\partial \Phi}{\partial \eta_j} = -2(\eta_1 - \eta_j) s_j + \lambda + 2\mu \eta_j = 0, \quad \text{where } j = 2, \dots, n-1, n. \quad (2.11)$$

by

$$\begin{aligned} \sum_{i=1}^n \frac{\partial \Phi}{\partial \eta_i} &= n\lambda = 0 \\ \sum_{i=1}^n \eta_i \frac{\partial \Phi}{\partial \eta_i} &= 2 \sum_{1 < j} (\eta_1 - \eta_j)^2 s_j + 2\mu = 0 \end{aligned}$$

we can see

$$\begin{aligned} \lambda &= 0 \\ \sum_{1 < j} (\eta_1 - \eta_j)^2 s_j &= -\mu \end{aligned}$$

Hence the critical values of f are given by $-\mu$. Now, we assume $-\mu \neq 0$. Meanwhile, we can also assume that $\mu + \text{Max}_{1 < j} (s_j) < 0$, otherwise, we have

$$-\mu = \sum_{1 < j} (\eta_1 - \eta_j)^2 s_j \leq \text{Max}_{1 < j} (s_j) < \sum_{1 < j} s_j + \text{Max}_{1 < j} (s_j)$$

Then by (2.11), we have

$$\eta_j = \frac{\eta_1 s_j}{\mu + s_j}, \quad j = 2, \dots, n-1, n. \quad (2.12)$$

Substitute (2.12) into $\eta_1 + \dots + \eta_n = 0$, we have

$$1 + \sum_{1 < j} \frac{s_j}{\mu + s_j} = 0$$

Hence

$$0 = 1 + \sum_{1 < j} \frac{s_j}{\mu + s_j} \geq 1 + \sum_{1 < j} \frac{s_j}{\mu + \text{Max}_{1 < j}(s_j)}. \quad (2.13)$$

Multiply both sides of (2.13) by $\mu + \text{Max}_{1 < j}(s_j)$, we get

$$-\mu = \sum_{1 < j} (\eta_1 - \eta_j)^2 s_j \leq \sum_{1 < j} s_j + \text{Max}_{1 < j}(s_j). \quad (2.14)$$

Notice that for any j , if $s_j > 0$, we have $0 > \frac{s_j}{\mu + s_j} \geq \frac{s_j}{\mu + \text{Max}_{1 < j}(s_j)}$, And the equality in (2.14) holds is equivalent to the equality in (2.13) holds. If the equality in (2.13) holds, we have, for each $j > 1$,

$$\frac{s_j}{\mu + s_j} = \frac{s_j}{\mu + \text{Max}_{1 < i}(s_i)}$$

which means that either $s_j = 0$ or the nonzero $s_j = \text{Max}_{1 < i}(s_i)$, that means all nonzero s_j are equal.

Thus, by (2.12) and the assumption above, fix a integer $k, k \in \{1, \dots, n-1\}$, we have $n-1$ cases, for each k ,

$$\eta_1 = \frac{\sqrt{n-k}}{\sqrt{n-k+1}}, \quad \eta_2 = \dots = \eta_k = 0, \quad \eta_{k+1} = \dots = \eta_n = \frac{-1}{\sqrt{(n-k+1)(n-k)}}.$$

Case(ii) is just a permutation of Case(i) under different assumption at the begining, so we finish the proof. \square

Remark 3. (i) When k takes values $n-1$ or 1 in Case(1) of Lemma 2.1, it corresponds to the Case(1) and Case(2) in [5][Lemma 1] respectively.

(ii) This new version would change Lemma 2 in [5], but Lu's rigidity theorem would still hold true, we will discuss this later.

Define the inner product of two $n \times n$ matrices A, B as $\langle A, B \rangle = \text{Tr} AB^\top$ and let $\|A\|^2 = \langle A, A \rangle = \sum_{i,j} a_{ij}^2$, where (a_{ij}) are the entries of A . So the Lu inequality [5][Lemma 2] becomes

Lemma 2.2. *Let A_1 be an $n \times n$ diagonal matrix of norm 1. Let A_2, \dots, A_m be symmetric matrices such that*

(i) $\langle A_\alpha, A_\beta \rangle = 0$ if $\alpha \neq \beta$;

Proposition 2.4 (Lu [5]). *Under notation above, we have*

$$|\nabla f_p|^2 \leq p^2 f_p \sum_{k,\alpha} \lambda_\alpha^{p-2} \left(\nabla_{\frac{\partial}{\partial x_k}} a_{\alpha\alpha} \right)^2. \quad (2.17)$$

$$\begin{aligned} \Delta g_p &= \frac{1}{p} f_p^{\frac{1}{p}-1} \Delta f_p + \frac{1}{p} \left(\frac{1}{p} - 1 \right) f_p^{\frac{1}{p}-2} |\nabla f_p|^2 \\ &\geq 2 f_p^{\frac{1}{p}-1} \sum_{\alpha} \left(\lambda_\alpha^{p-1} \sum_{i,j,k} (h_{ijk}^\alpha)^2 \right) \\ &\quad + 2 f_p^{\frac{1}{p}-1} (r \|A_1\|^{2p} (n - \|A_1\|^2 - \sum_{\alpha=2}^m \|A_\alpha\|^2 - \lambda_2) - 3mn \lambda_{r+1}^p). \end{aligned} \quad (2.18)$$

Then integrating both sides of (2.18), letting $p \rightarrow \infty$, since $\lambda_{r+1}^p / f_p \rightarrow 0$ as p tends to ∞ , Lu derived

$$\int_M \sum_{i,j,k} \sum_{\alpha \leq r} (h_{ijk}^\alpha)^2 + \|A_1\|^2 (n - \|A_1\|^2 - \sum_{\alpha=2}^m \|A_\alpha\|^2 - \lambda_2) \leq 0. \quad (2.19)$$

When the equality holds in (2.19), the equality in (2.15) also holds, A_α takes the form in Lemma 2.2. Then, using the structure equation case by case, Lu proved Theorem A.

Remark 4. Although there are more cases when the equality in (2.15) holds, using similar argument as Lu did in the original proof, we can rule out the new case. To be precise, for each k , let $n > k + 1, j \geq k + 2$, then from $0 = dh_{1j}^{n+1} = h_{11}^{n+1} \omega_{1j}$, we conclude $\omega_{1j} = 0$ if $j \geq k + 2$. Similarly, by computing $dh_{ij}^{n+1}, i \in 2 \dots k + 1$, we also have $\omega_{2j} = \dots = \omega_{k+1j} = 0$ for $j \geq k + 2$. Thus by the structure equations, we have

$$0 = d\omega_{1j} = \omega_1 \wedge \omega_j,$$

a contradiction if $n > k + 1$, thus Theorem A still holds true.

3. PROOF OF THE MAIN THEOREM

Let $g_\epsilon = (g_p + \epsilon)^{\frac{1}{2}}$, where $\epsilon > 0$ is a constant. We first prove the inequality in the main theorem, that is

Proposition 3.1. *Let M^n be a closed nontotally geodesic minimal submanifold in $\mathbb{S}^{n+m}(1)$, then*

$$\mu_1 \leq -n + \max_{p \in M} \lambda_2 - \frac{2}{n+2} \frac{\int_M \left[\frac{1}{r} \sum_{i,j,k} \sum_{\alpha \leq n+r} (h_{ijk}^\alpha)^2 \right]}{\int_M \lambda_1} \quad (3.1)$$

Proof. By direct computation, using (2.18),

$$\begin{aligned}
\Delta g_\epsilon &= \frac{1}{2} (g_p + \epsilon)^{-\frac{1}{2}} \Delta g_p - \frac{1}{4} |\nabla g_p|^2 (g_p + \epsilon)^{-\frac{3}{2}} \\
&= \frac{1}{2} (g_p + \epsilon)^{-\frac{1}{2}} \left(\frac{1}{p} f_p^{\frac{1}{p}-1} \Delta f_p + \frac{1}{p} \left(\frac{1}{p} - 1 \right) f_p^{\frac{1}{p}-2} |\nabla f_p|^2 \right) - \frac{1}{4} |\nabla g_p|^2 (g_p + \epsilon)^{-\frac{3}{2}} \\
&\geq \frac{1}{2} (g_p + \epsilon)^{-\frac{1}{2}} \left(2 f_p^{\frac{1}{p}-1} \sum_\alpha \left(\lambda_\alpha^{p-1} \sum_{i,j,k} (h_{ijk}^\alpha)^2 \right) + 2 f_p^{\frac{1}{p}-1} (r \|A_1\|^{2p} (n+1 - \|A_1\|^2 - \sum_{\alpha=2}^m \|A_\alpha\|^2 - \lambda_2)) \right. \\
&\quad \left. - 6nm f_p^{\frac{1}{p}-1} \lambda_{r+1}^p \right) - \frac{1}{4} |\nabla g_p|^2 (g_p + \epsilon)^{-\frac{3}{2}} \\
&\geq (g_p + \epsilon)^{-\frac{3}{2}} \underbrace{\left[(g_p + \epsilon) f_p^{\frac{1}{p}-1} \sum_\alpha \left(\lambda_\alpha^{p-1} \sum_{i,j,k} (h_{ijk}^\alpha)^2 \right) - \frac{1}{4} |\nabla g_p|^2 \right]}_I \\
&\quad + \underbrace{(g_p + \epsilon)^{-\frac{1}{2}} \left[f_p^{\frac{1}{p}-1} (r \|A_1\|^{2p} (n - \|A_1\|^2 - \sum_{\alpha=2}^m \|A_\alpha\|^2 - \lambda_2)) - 3nm f_p^{\frac{1}{p}-1} \lambda_{r+1}^p \right]}_{II}.
\end{aligned} \tag{3.2}$$

To deal with I , follows from (1.9) and (1.11) in [8][Proposition 1], we have following Lemma **Lemma 3.2** (Shen [8]). *Let M^n be a closed minimal submanifold in $\mathbb{S}^{n+m}(1)$, then we have*

$$|\nabla(|A_\alpha|^2)|^2 \leq \frac{4n}{n+2} |A_\alpha|^2 \left[\sum_{i,j,k} (h_{ijk}^\alpha)^2 \right]. \tag{3.3}$$

Then, applying Lemma 3.2 to (2.17), we have

$$|\nabla g_p|^2 = \frac{1}{p^2} f_p^{\frac{2}{p}-2} |\nabla f_p|^2 \leq f_p^{\frac{2}{p}-1} \sum_\alpha \lambda_\alpha^{p-2} |\nabla \lambda_\alpha|^2 \leq f_p^{\frac{2}{p}-1} \sum_\alpha \frac{4n}{n+2} \left(\lambda_\alpha^{p-1} \sum_{i,j,k} (h_{ijk}^\alpha)^2 \right)$$

Thus, for I , we have

$$\begin{aligned}
I &\geq (g_p + \epsilon)^{-\frac{3}{2}} \left[(g_p + \epsilon) f_p^{\frac{1}{p}-1} \sum_{\alpha} \left(\lambda_{\alpha}^{p-1} \sum_{i,j,k} (h_{ijk}^{\alpha})^2 \right) - \frac{1}{4} \frac{1}{p^2} f_p^{\frac{2}{p}-2} |\nabla f_p|^2 \right] \\
&\geq (g_p + \epsilon)^{-\frac{3}{2}} \left[(g_p + \epsilon) f_p^{\frac{1}{p}-1} \sum_{\alpha} \left(\lambda_{\alpha}^{p-1} \sum_{i,j,k} (h_{ijk}^{\alpha})^2 \right) - \frac{1}{4} f_p^{\frac{2}{p}-1} \frac{4n}{n+2} \sum_{\alpha} \left(\lambda_{\alpha}^{p-1} \sum_{i,j,k} (h_{ijk}^{\alpha})^2 \right) \right] \\
&\geq \frac{2}{n+2} (g_p + \epsilon)^{-\frac{1}{2}} f_p^{\frac{1}{p}-1} \left[\sum_{\alpha} \left(\lambda_{\alpha}^{p-1} \sum_{i,j,k} (h_{ijk}^{\alpha})^2 \right) \right] \\
&\geq 0
\end{aligned}$$

Meanwhile, by definition, put g_{ϵ} into $\mu_1 = \inf_{f \in C^{\infty}(M)} \frac{\int_M L(f) f}{\int_M f^2}$, we get

$$\begin{aligned}
\mu_1 &\leq \frac{\int_M L(g_{\epsilon}) g_{\epsilon}}{\int_M g_{\epsilon}^2} = \frac{\int_M -g_{\epsilon} \Delta g_{\epsilon} - \sigma g_{\epsilon}^2}{\int_M g_{\epsilon}^2} \\
&= \frac{\int_M -g_{\epsilon} (I + II) - \sigma g_{\epsilon}^2}{\int_M g_{\epsilon}^2} \\
&\leq \frac{\int_M -\frac{2}{n+2} f_p^{\frac{1}{p}-1} \left[\sum_{\alpha} \left(\lambda_{\alpha}^{p-1} \sum_{i,j,k} (h_{ijk}^{\alpha})^2 \right) \right]}{\int_M g_{\epsilon}^2} \\
&\quad + \frac{\int_M - \left[f_p^{\frac{1}{p}-1} (r \|A_1\|^{2p} (n - \|A_1\|^2 - \sum_{\alpha=2}^m \|A_{\alpha}\|^2 - \lambda_2)) - 3nm f_p^{\frac{1}{p}-1} \lambda_{r+1}^p \right]}{\int_M g_{\epsilon}^2} \\
&\quad - \frac{\int_M \sigma g_{\epsilon}^2}{\int_M g_{\epsilon}^2}.
\end{aligned}$$

Then, letting $p \rightarrow \infty$ and $\epsilon \rightarrow 0$, using the fact that $\lambda_{r+1}^p / f_p \rightarrow 0$ almost everywhere when $p \rightarrow \infty$, we finish the proof. \square

Now, we prove the main theorem,

Proof. From the proof of Proposition 3.1, we know that if

$$\mu_1 \geq -n + \max_{p \in M} \lambda_2,$$

then either M is totally geodesic, so $\mu_1 = 0$, or $\mu_1 = -n + \max_{p \in M} \lambda_2$, we have

$$\frac{1}{r} \sum_{i,j,k} \sum_{\alpha \leq n+r} (h_{ijk}^{\alpha})^2 = 0,$$

we claim that σ is a constant. By Lemma 2.2, there are following cases:

- [2] Q. Chen and S. Xu, *Rigidity of compact minimal submanifolds in a unit sphere*, Geometriae Dedicata **45** (1993), no. 1, 83–88.
- [3] S.-S. Chern, M D. Carmo, and S. Kobayashi, *Minimal submanifolds of a sphere with second fundamental form of constant length*, Functional analysis and related fields, 1970, pp. 59–75.
- [4] H B. Lawson, *Local rigidity theorems for minimal hypersurfaces*, Annals of Mathematics (1969), 187–197.
- [5] Z. Lu, *Normal scalar curvature conjecture and its applications*, Journal of Functional Analysis **261** (2011), no. 5, 1284–1308.
- [6] Y. Luo, L. Sun, and J. Yin, *An optimal pinching theorem of minimal legendrian submanifolds in the unit sphere*, Calculus of Variations and Partial Differential Equations **61** (2022), no. 5, 1–18.
- [7] O. Perdomo, *First stability eigenvalue characterization of clifford hypersurfaces*, Proceedings of the American Mathematical Society **130** (2002), no. 11, 3379–3384.
- [8] Y. B. Shen, *Curvature and stability for minimal submanifolds*, Sci. Sinica Ser. A **31** (1988), no. 7, 787–797.
- [9] J. Simons, *Minimal varieties in riemannian manifolds*, Annals of Mathematics (1968), 62–105.
- [10] C. Wu, *New characterizations of the clifford tori and the veronese surface*, Archiv der Mathematik **61** (1993), 277–284.
- [11] J. Yin and X. Qi, *Sharp estimates for the first eigenvalue of schrödinger operator in the unit sphere*, Proceedings of the American Mathematical Society **150** (2022), no. 7, 3087–3101.

DEPARTMENT OF MATHEMATICS, FUDAN UNIVERSITY, SHANGHAI 200433, CHINA

Email address: pw17@fudan.edu.cn