

The unitary Cayley graph of upper triangular matrix rings

Waldemar Hołubowski¹, Sergiy Kozerenko^{2,3,4},
Bogdana Oliynyk^{1,2}, Viktoriia Solomko^{1,2}

March 22, 2024

¹Faculty of Applied Mathematics, Silesian University of Technology, Poland

²Department of Mathematics, National University of Kyiv-Mohyla Academy, Ukraine

³Institute of Mathematics of NAS of Ukraine, Ukraine

⁴Kyiv School of Economics, Ukraine

w.holubowski@polsl.pl, kozerenkosergiy@ukr.net, bogdana.oliynyk@polsl.pl,
vsolomko@polsl.pl

Keywords: *unitary Cayley graph, upper triangular matrix ring, antipodal graph, Hamming graph, clique number*

2020 Mathematics Subject Classification: 05C25, 05E16, 15B33, 16S50

Abstract

The unitary Cayley graph C_R of a finite unital ring R is the simple graph with vertex set R in which two elements x and y are connected by an edge if and only if $x - y$ is a unit of R . We characterize the unitary Cayley graph $C_{T_n(\mathbb{F})}$ of the ring of all upper triangular matrices $T_n(\mathbb{F})$ over a finite field \mathbb{F} . We show that $C_{T_n(\mathbb{F})}$ is isomorphic to the semistrong product of the complete graph K_m and the antipodal graph of the Hamming graph $A(H(n, p^k))$, where $m = p^{\frac{kn(n-1)}{2}}$ and $|\mathbb{F}| = p^k$. In particular, if $|\mathbb{F}| = 2$, then the graph $C_{T_n(\mathbb{F})}$ has 2^{n-1} connected components, each component is isomorphic to the complete bipartite graph $K_{m,m}$, where $m = 2^{\frac{n(n-1)}{2}}$. We also compute the diameter, triameter, and clique number of the graph $C_{T_n(\mathbb{F})}$.

Introduction

In algebraic graph theory, various graphs associated with algebraic structures play a prominent role. A notable example is the Cayley graph, denoted as $Cay(G, S)$, of a group G related to some subset S of G . The graph $Cay(G, S)$ is connected if and only if S generates G . Furthermore, $Cay(G, S)$ is regular, and its properties provide significant insights into the structure of finitely generated groups. We refer to books [9] on algebraic graph theory and [11, 20] on geometric group theory for the application of Cayley graphs in groups.

Let R be a finite ring with identity. The *unitary Cayley graph* of a ring R , denoted by C_R , is a simple graph with vertex set R , and two elements x, y are adjacent if and only if $x - y$ is a unit of R . Essentially, it is just a Cayley graph of an additive group of ring R related to the set of units of R .

The unitary Cayley graphs have been studied intensively for the ring Z_n (see, for example [3, 7, 8, 13, 17]). In particular, Dejter and Giudici [7] proved that if n is a prime then the unitary Cayley graph C_{Z_n} is isomorphic to a complete graph with n vertices, and C_{Z_n} is a bipartite graph when n is even, moreover, if $n = 2^s$ then C_{Z_n} is isomorphic to a complete bipartite graph $K_{2^{s-1}, 2^{s-1}}$. In [1] the diameter, girth, eigenvalues, vertex and edge connectivity, and vertex and edge chromatic number were described precisely for a finite commutative ring R .

Let \mathbb{F} be a finite field, n be a positive integer, $n \geq 2$. In their works [15, 16] Kiani, Mollahajiahaei, and Aghaei described the chromatic number, clique number, independence number, diameter of unitary Cayley graph $C_{M_n(\mathbb{F})}$ of matrix ring $M_n(\mathbb{F})$ over a finite field \mathbb{F} and proved that the graph $C_{M_n(\mathbb{F})}$ is regular and strongly regular for $n = 2$. In a related study [23] Rattanakangwanwong and Meemark computed the clique number, chromatic number, and independence number of the subgraph of unitary Cayley graph of a matrix algebra over a finite field induced by the set of idempotent matrices. For some other recent papers on unitary Cayley graphs of matrix ring $M_n(\mathbb{F})$, we refer the reader to [5, 12, 22].

In this paper, we characterize the properties of the unitary Cayley graph $C_{T_n(\mathbb{F})}$ of the ring of all upper triangular $n \times n$ matrices $T_n(\mathbb{F})$ over the field \mathbb{F} . Denote by $(T_n(\mathbb{F}))^*$ the set of all invertible upper triangular $n \times n$ matrices. The vertices of the graph $C_{T_n(\mathbb{F})}$ are upper triangular $n \times n$ matrices $T_n(\mathbb{F})$ over \mathbb{F} and two vertices a and b are adjacent if and only if $a - b$ is an element of $(T_n(\mathbb{F}))^*$.

We will demonstrate that when $|\mathbb{F}| = 2$, the unitary Cayley graph $C_{T_n(\mathbb{F})}$ of the ring of all upper triangular matrices $T_n(\mathbb{F})$ over \mathbb{F} exhibits a fairly simple structure.

Theorem 1. *If $|\mathbb{F}| = 2$, then the graph $C_{T_n(\mathbb{F})}$ has 2^{n-1} connected components, and its each component is isomorphic to the complete bipartite graph $K_{m,m}$, where $m = 2^{\frac{n(n-1)}{2}}$.*

In the case of $|\mathbb{F}| > 2$, the unitary Cayley graph $C_{T_n(\mathbb{F})}$ is connected. Let S be a set of p elements, p and l be positive integers. The *Hamming graph* $H(l, p)$ is defined on the set of ordered l -tuples of elements of S with two vertices being adjacent if they differ in precisely one coordinate. Hamming graphs find active applications in information theory and computer science. Specifically, we will consider Hamming graph $H(n, p^k)$ for $S = \mathbb{F}$, $|\mathbb{F}| = p^k$.

The antipodal graph of a graph G , denoted by $A(G)$, has the same vertex set as G with two vertices u and v being adjacent if the distance between u and v is equal to the diameter of G (see [2, 14]). The antipodal graph $A(H(n, p^k))$ of the Hamming graph $H(n, p^k)$ is defined on the set of all n -tuples of elements from the field \mathbb{F} , and two n -tuples $u = (u_1, \dots, u_n)$ and $v = (v_1, \dots, v_n)$ are adjacent provided $u_i \neq v_i$ for every $1 \leq i \leq n$.

For the case of $|\mathbb{F}| > 2$, we prove the following result that describes the structure of graph $C_{T_n(\mathbb{F})}$ as a semistrong product of two graphs.

Theorem 2. *Let $|\mathbb{F}| > 2$, $|\mathbb{F}| = p^k$ for a prime p and a positive integer k . Then*

$$C_{T_n(\mathbb{F})} \simeq K_m \bullet A(H(n, p^k)), \quad (1)$$

where $m = p^{\frac{kn(n-1)}{2}}$.

Additionally, we also characterize the diameter, triameter, and clique number of the graph $C_{T_n(\mathbb{F})}$.

1 Preliminaries

In this paper, we consider only finite simple graphs. Let $G = (V(G), E(G))$ be a finite connected simple graph. Define a metric d_G on the set of vertices $V(G)$ as follows: for any $u, v \in V(G)$ the distance $d_G(u, v)$ is defined as the length of the shortest path between u and v .

The *diameter* of a connected graph G is the value

$$\text{diam}(G) = \max\{d_G(u, v) : u, v \in V(G)\}.$$

For every triplet of vertices $u, v, w \in V(G)$, we define

$$d_G(u, v, w) = d_G(u, v) + d_G(u, w) + d_G(v, w).$$

The *triameter* of a connected graph G is defined as the value

$$\text{triam}(G) = \max\{d_G(u, v, w) : u, v, w, \in V(G)\}.$$

The triplet of vertices $u, v, w \in V(G)$ is called *triangular* if $d_G(u, v, w) = tr(G)$. The main motivation for studying $tr(G)$ comes from its appearance in lower bounds on the radio k -chromatic number of a graph and the total domination number of a connected graph ([6, 10, 18, 24]).

A *clique* is a subgraph of a graph G that is isomorphic to a complete graph. The *clique number* of G is the size of the largest clique in G , denoted by $\omega(G)$.

Now, consider two finite simple graphs, G and H . The semistrong product $G \bullet H$ (see [19]) is the graph with vertex set $V(G \bullet H) = V(G) \times V(H)$ and edge set

$$E(G \bullet H) = \{(u_1, v_1)(u_2, v_2) \mid u_1 u_2 \in E(G) \text{ and } v_1 v_2 \in E(H) \text{ or } u_1 = u_2 \text{ and } v_1 v_2 \in E(H)\}.$$

2 The properties of the unitary Cayley graph of $T_n(\mathbb{F})$

For any unital finite ring R the unitary Cayley graph C_R is $|R^*|$ -regular [17]. As $|(T_n(\mathbb{F}))^*| = (|\mathbb{F}| - 1)^n \cdot |\mathbb{F}|^{\frac{n^2-n}{2}}$, the next proposition becomes evident.

Proposition 1. *The graph $C_{T_n(\mathbb{F})}$ is $(|\mathbb{F}| - 1)^n |\mathbb{F}|^{\frac{n^2-n}{2}}$ -regular.*

From the structure of upper triangular matrices, we deduce a simple condition for the existence of an edge in the graph $C_{T_n(\mathbb{F})}$.

Proposition 2. *Two matrices $a = \{a_{ij}\}_{1 \leq i, j \leq n}$ and $b = \{b_{ij}\}_{1 \leq i, j \leq n}$ are connected by edge in $C_{T_n(\mathbb{F})}$ if and only if $a_{ii} - b_{ii} \neq 0$ for any $1 \leq i, j \leq n$.*

Proof. Indeed, $a - b$ is invertible if and only if $\det(a - b) \neq 0$. As a and b are upper triangular,

$$\det(a - b) = \prod_{i=1}^n (a_{ii} - b_{ii}).$$

This completes the proof of the proposition. □

Theorem 3. *If $|\mathbb{F}| \neq 2$, then the graph $C_{T_n(\mathbb{F})}$ is connected.*

First, we prove Theorem 1.

Proof of Theorem 1. Let $|\mathbb{F}| = 2$ and let $a = \{a_{ij}\}_{1 \leq i, j \leq n}$ be a matrix from $T_n(\mathbb{F})$. For the vector $(a_{11}, a_{22}, \dots, a_{nn})$ there exists only one vector $(b_{11}, b_{22}, \dots, b_{nn})$ such that

$$a_{ii} - b_{ii} = 1 \pmod{2}, \quad \text{for any } 1 \leq i \leq n.$$

Proposition 2 implies, that the matrix a is connected by an edge with any matrix $b = \{b_{ij}\}_{1 \leq i, j \leq n}$ from $T_n(\mathbb{F})$ with the elements $(b_{11}, b_{22}, \dots, b_{nn})$ on the main diagonal. It is true for all matrices with the elements $(a_{11}, a_{22}, \dots, a_{nn})$ on the main diagonal. However, any matrix with the elements $(b_{11}, b_{22}, \dots, b_{nn})$ on the main diagonal is connected by an edge with any matrix with the elements $(a_{11}, a_{22}, \dots, a_{nn})$ on the main diagonal and is not connected with any another matrix from $T_n(\mathbb{F})$.

Hence, all matrices with the elements $(a_{11}, a_{22}, \dots, a_{nn})$ or $(b_{11}, b_{22}, \dots, b_{nn})$ on the main diagonal form the connected component of the graph $C_{T_n(\mathbb{F})}$. Since we choose matrix a arbitrarily, any connected component of graph $C_{T_n(\mathbb{F})}$ is defined by two vectors $(c_{11}, c_{22}, \dots, c_{nn})$ and $(d_{11}, d_{22}, \dots, d_{nn})$, such that

$$c_{ii} - d_{ii} = 1 \pmod{2}, \text{ for any } 1 \leq i \leq n.$$

Therefore, the graph $C_{T_n(\mathbb{F})}$ has 2^{n-1} connected components and any component is isomorphic to $K_{m,m}$ where m equals the number of upper triangular $n \times n$ matrices over field \mathbb{F}_2 with elements $(c_{11}, c_{22}, \dots, c_{nn})$ on the main diagonal, i.e.

$$m = 2^{\frac{n(n-1)}{2}}.$$

□

Proof of Theorem 3. Let now $|\mathbb{F}| > 2$. Assume, that matrices $a = \{a_{ij}\}_{1 \leq i, j \leq n}$ and $b = \{b_{ij}\}_{1 \leq i, j \leq n}$ are not connected by an edge. Then we can construct a matrix $c = \{c_{ij}\}_{1 \leq i, j \leq n}$ in the following way:

- for all i , $1 \leq i \leq n$, we choose c_{ii} such that $c_{ii} \neq a_{ii}$, $c_{ii} \neq b_{ii}$ (we can choose c_{ii} because $|\mathbb{F}| > 2$);
- for all $1 \leq i, j \leq n$, $i \neq j$ we set $c_{ij} = 0$.

Proposition 2 implies that the matrix c is connected by an edge with the matrices a and b . So, for any matrices a and b that are not connected by an edge, there exists a matrix c which is connected by an edge with both a and b simultaneously. Therefore, the graph $C_{T_n(\mathbb{F})}$ is connected. □

This theorem directly implies the next proposition.

Corollary 1. *If $|\mathbb{F}| > 2$, then*

$$\text{diam}(C_{T_n(\mathbb{F})}) = 2.$$

Proof. The proof of this corollary directly follows from the proof of Theorem 3. □

The next proposition describes the triameter of the graph $C_{T_n(\mathbb{F})}$.

Proposition 3. *If $|\mathbb{F}| > 2$, then*

$$\text{triam}(C_{T_n(\mathbb{F})}) = 6.$$

Proof. It is clear that $\text{triam}(C_{T_n(\mathbb{F})}) \leq 6$, because $\text{triam}(G) = \max\{d_G(u, v, w) : u, v, w, \in V(G)\}$ and $d_G(u, v, w) = d_G(u, v) + d_G(u, w) + d_G(v, w) \leq 3 \cdot \text{diam}(G)$. Since $|\mathbb{F}| > 2$, there exist three pairwise different elements $a, b, c \in \mathbb{F}$. We consider three diagonal matrices $d_1 = \text{diag}(a, a, a, \dots, a)$, $d_2 = \text{diag}(a, b, b, \dots, b)$ and $d_3 = \text{diag}(a, c, c, \dots, c)$. We have $d_{C_{T_n(\mathbb{F})}}(d_1, d_2, d_3) = 2 + 2 + 2 = 6$. This completes the proof of the proposition. \square

The next proposition characterizes the clique number of the graph $C_{T_n(\mathbb{F})}$.

Proposition 4.

$$\omega(C_{T_n(\mathbb{F})}) = |\mathbb{F}|.$$

Proof. Let S be a maximal clique of a graph G_{T_n} . Proposition 2 implies, that

$$a_{ii} \neq b_{ii}$$

for any two different elements $a = \{a_{ij}\}_{1 \leq i, j \leq n}$ and $b = \{b_{ij}\}_{1 \leq i, j \leq n}$ from S and for any i , $1 \leq i \leq n$. But for any i we can choose only $|\mathbb{F}|$ different elements a_{ii} . So, $|S| = |\mathbb{F}|$. \square

3 Connection with Hamming graph

Proof of Theorem 2. We have to show that the unitary Cayley graph $C_{T_n(\mathbb{F})}$ of the ring of all upper triangular matrices $T_n(\mathbb{F})$ over \mathbb{F} is isomorphic to the semistrong product of the complete graph K_m and the antipodal graph of the Hamming graph $A(H(n, p^k))$, where $m = p^{\frac{kn(n-1)}{2}}$.

First, we define a complete graph on the set of all strictly upper triangular matrices. This graph is isomorphic to the graph K_m , where $m = p^{\frac{kn(n-1)}{2}}$.

Now, determine a bijection φ from the set of vertices of the graph $C_{T_n(\mathbb{F})}$ to the Cartesian product of the set of vertices of K_m and the set of vertices of the graph $A(H(n, p^k))$. Let $a = \{a_{ij}\}_{1 \leq i, j \leq n}$ be a matrix from $T_n(\mathbb{F})$. Define

$$\varphi(a) = (\hat{a}, (a_{11}, \dots, a_{nn})),$$

where

$$\hat{a} = \begin{pmatrix} 0 & a_{12} & \dots & a_{1n} \\ 0 & 0 & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & a_{(n-1),n} \\ 0 & 0 & \dots & 0 \end{pmatrix}.$$

We would like to show that two matrices $a = \{a_{ij}\}_{1 \leq i, j \leq n}$ and $b = \{b_{ij}\}_{1 \leq i, j \leq n}$ are connected by an edge in the graph $C_{T_n(\mathbb{F})}$ if and only if they are connected by edge in the semistrong product of graphs K_m and $A(H(n, p^k))$. Assume, that there exist the edge ab in the graph $C_{T_n(\mathbb{F})}$. Proposition 2 implies that $a_{ii} \neq b_{ii}$ for any i , $1 \leq i \leq n$. Then the vectors (a_{11}, \dots, a_{nn}) and (b_{11}, \dots, b_{nn}) are connected by an edge in $A(H(n, p^k))$. If the matrices

$$\hat{a} = \begin{pmatrix} 0 & a_{12} & \dots & a_{1n} \\ 0 & 0 & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & a_{(n-1),n} \\ 0 & 0 & \dots & 0 \end{pmatrix} \quad \text{and} \quad \hat{b} = \begin{pmatrix} 0 & b_{12} & \dots & b_{1n} \\ 0 & 0 & \dots & b_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & b_{(n-1),n} \\ 0 & 0 & \dots & 0 \end{pmatrix}$$

are different, then they are connected by an edge in K_m . Therefore, from the definition of semistrong product of graphs follows, that there exists the edge ab in the graph $K_m \bullet A(H(n, p^k))$.

Assume there is no edge ab in $C_{T_n(\mathbb{F})}$. Proposition 2 implies that there exist i , $1 \leq i \leq n$, such that $a_{ii} = b_{ii}$. Consequently, the vectors (a_{11}, \dots, a_{nn}) and (b_{11}, \dots, b_{nn}) are not connected by edge in $A(H(n, p^k))$. Thus, there is no edge ab in $K_m \bullet A(H(n, p^k))$. □

We can describe the connection between $C_{T_n(\mathbb{F})}$ and $A(H(n, p^k))$ in another way. Define on $T_n(\mathbb{F})$ equivalence relation \equiv :

a matrix $a = \{a_{ij}\}_{1 \leq i, j \leq n}$ is equivalent to a matrix $b = \{b_{ij}\}_{1 \leq i, j \leq n}$ if and only if $a_{ii} = b_{ii}$ for all i , $1 \leq i \leq n$.

That is, each equivalence class is determined by the main diagonal of matrices.

Let \hat{C}_{T_n} be a graph induced by a graph $C_{T_n(\mathbb{F})}$ on the set $T_n(\mathbb{F})|_{\equiv}$. Then we have the next description of \hat{C}_{T_n} .

Proposition 5. $\hat{C}_{T_n} \simeq A(H(n, p^k))$, for all $n \geq 2$.

The proof of this proposition directly follows from the Theorem 2.

4 Acknowledgments

Sergiy Kozerenko is deeply thankful to the Armed Forces of Ukraine for ensuring the safety of Kyiv during the work on this paper. Viktoriia Solomko was supported by a grant from the Institute of International Education's Scholar Rescue Fund.

References

- [1] R. Akhtar, T. Jackson-Henderson, R. Karpman, M. Boggess, I. Jiménez, A. Kinzel and D. Pritikin, On the unitary Cayley graph of a finite ring, *Electron. J. Combin.*, **16(1)** (2009), 13 pages.
- [2] R. Aravamudhan and B. Rajendran, On antipodal graphs, *Discrete Math.* **49(2)** (1984), 193–195.
- [3] P. Berrizbeitia and R. E. Giudici, On cycles in the sequence of unitary Cayley graphs, *Discrete Math.*, **282** (2004), 239–243.
- [4] A. E. Brouwer and W. H. Haemers, Spectra of Graphs, Springer New York (2011), 250 p.
- [5] B. Chen and J. Huang, On unitary Cayley graphs of matrix rings, *Discrete Math.* **345(1)** (2022), article 112671.
- [6] A. Das, Triameter of graphs, *Discuss. Math. Graph Theory* **41** (2021), 601–616.
- [7] I. J. Dejter and R. E. Giudici, On Unitary Cayley Graphs, *J. Combin. Math. Combin. Comput.* **18** (1995), 121–124.
- [8] E. D. Fuchs, Longest induced cycles in circulant graphs, *Electron. J. Comb.* **12(1)** (2005), 12 pages.
- [9] C. D. Godsil and G. Royle, Algebraic graph theory, Springer-Verlag New York Inc. (2001), 443 p.
- [10] A. Hak, S. Kozerenko and B. Oliynyk, A note on the triameter of graphs, *Discrete Appl. Math.*, **309** (2022), 278–284.
- [11] P. de la Harpe, Topics in geometric group theory, Chicago Lectures in Math. University of Chicago Press (2000), 310 p.
- [12] J. Huang and M. Zhang, Spectral properties of Cayley graphs over $M_{m \times n}(F_q)$, *Finite Fields Appl.* **80** (2022), article 102024.

- [13] A. Ilić, The energy of unitary cayley graphs, *Linear Algebra Appl.* **431** (2009), 1881–1889.
- [14] G. Johns and K. Sleno, Antipodal graphs and digraphs, *Int. J. Math. Math. Sci.*, **16** (1993), article 205867.
- [15] D. Kiani and M. M. H. Aghaei, On the unitary Cayley graph of a ring, *Electron. J. Comb.* **19(2)** (2012), P10.
- [16] D. Kiani and M. Mollahajiaghaei, On the unitary Cayley graphs of matrix algebras, *Linear Algebra Appl.* **466** (2015), 421–428.
- [17] W. Klotz and T. Sander, Some properties of unitary Cayley graphs, *Electron. J. Combin.*, **14** (2007), P45.
- [18] S. R. Kola and P. Panigrahi, A lower bound for radio k -chromatic number of an arbitrary graph, *Contrib. Discrete Math.* **10** (2015), 45–56.
- [19] X. Li and X. Zhang, Nowhere-zero 3-flows in semistrong product of graphs, *Discrete Math.*, **338** (2015), 2408–2411.
- [20] C. Loh, Geometric group theory, Universitext Springer (2017), 389 p.
- [21] N. V. Moosavi, K. Khashyarmanesh, S. Mohammadikhah and M. Afkhami, On the Cayley graphs of upper triangular matrix rings, *Quasigroups Related Systems* **26** (2018), 317–326.
- [22] J. Rattanakangwanwong and Y. Meemark, Subconstituents of unitary Cayley graph of matrix algebras, *Finite Fields Appl.*, **80** (2022), article 102004.
- [23] J. Rattanakangwanwong and Y. Meemark, Subgraph of unitary Cayley graph of matrix algebras induced by idempotent matrices, *Discrete Math.*, **347** (2024), article 113739.
- [24] L. Saha and P. Panigrahi, A lower bound for radio k -chromatic number, *Discrete Appl. Math.* **192** (2015), 87–100.