

Rotational surfaces in a normed 3-space whose principal curvatures satisfy a linear relation

Makoto SAKAKI and Kakeru YANASE

Abstract. We classify rotational surfaces in a normed 3-space with rotationally symmetric norm whose principal curvatures satisfy a linear relation.

Mathematics Subject Classification. 53A35, 53A10, 52A15, 52A21, 46B20

Keywords. rotational surface, normed 3-space, principal curvature, linear Weingarten, Birkhoff orthogonal, Birkhoff Gauss map

1 Introduction

It is interesting to generalize differential geometry of surfaces in the Euclidean 3-space to that in normed 3-spaces (cf. [1], [2], [3], [4], [8]). In the Euclidean 3-space, a surface is said to be of linear Weingarten type if its principal curvatures satisfy a linear relation, or if its Gaussian and mean curvatures satisfy a linear relation (cf. [5], [6], [7]).

In this paper, we define surfaces of linear Weingarten type in a normed 3-space as surfaces whose principal curvatures satisfy a linear relation, and classify rotational surfaces of linear Weingarten type in a normed 3-space with rotationally symmetric norm. It is a generalization of [7] in the Euclidean case, and also a generalization of [8] for rotational surfaces with constant mean curvature in a normed 3-space.

This paper is organized as follows. In Section 2, basic facts on surfaces in normed 3-spaces are given (cf. [2]). In Section 3, we give some computations on rotational surfaces in a normed 3-space with rotationally symmetric norm (cf. [8]). In Section 4, we discuss rotational surfaces in the normed 3-space one of whose principal curvatures is constant. In Section 5, we consider rotational surfaces of homogenous linear Weingarten type in the normed 3-space. In Section 6, we study rotational surfaces of inhomogeneous linear Weingarten type in the normed 3-space.

2 Preliminaries

Let $(\mathbb{R}^3, \|\cdot\|)$ be a 3-dimensional normed space. The unit ball B and the unit sphere S are given by

$$B = \{x \in \mathbb{R}^3; \|x\| \leq 1\}, \quad S = \{x \in \mathbb{R}^3; \|x\| = 1\}.$$

We assume that S is smooth and strictly convex, that is, S is a smooth surface and S contains no line segment.

Let v be a non-zero vector in \mathbb{R}^3 and Π be a plane in \mathbb{R}^3 . The vector v is said to be Birkhoff orthogonal to Π , denoted by $v \perp_B \Pi$, if the tangent plane of S at $v/\|v\|$ is parallel to Π .

Let M be a surface immersed in $(\mathbb{R}^3, \|\cdot\|)$, and $T_p M$ be the tangent plane of M at $p \in M$. Then there exists a vector $\eta(p) \in S$ such that $\eta(p) \perp_B T_p M$. This gives a local smooth map $\eta : U \subset M \rightarrow S$, which is called the Birkhoff-Gauss map. It can be global if and only if M is orientable.

Let k_1 and k_2 be the eigenvalues of $d\eta_p$, which are called the principal curvatures. We say that the surface M is of linear Weingarten type if $ak_1 + bk_2 = c$ for some constants a, b and c which are not all zero.

3 Rotational surfaces

Let $(\mathbb{R}^3, \|\cdot\|)$ be a normed 3-space with rotationally symmetric norm

$$\|x\| = ((x_1^2 + x_2^2)^m + x_3^{2m})^{\frac{1}{2m}}, \quad x = (x_1, x_2, x_3),$$

where m is a positive integer. Set

$$\Phi(x_1, x_2, x_3) := (x_1^2 + x_2^2)^m + x_3^{2m}.$$

Then the unit sphere S is given by

$$S = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \mid \Phi(x_1, x_2, x_3) = 1\}.$$

Since the case $m = 1$ is the Euclidean case, we assume that $m \geq 2$ in the following.

Let M be a surface in the above $(\mathbb{R}^3, \|\cdot\|)$ which is rotational around x_3 -axis, and is parametrized by

$$X(u, v) = (\alpha(u) \cos v, \alpha(u) \sin v, \beta(u))$$

where $\alpha > 0$, $\alpha' \neq 0$ and $\beta' \neq 0$. Then

$$X_u = (\alpha' \cos v, \alpha' \sin v, \beta'), \quad X_v = (-\alpha \sin v, \alpha \cos v, 0).$$

The Birkhoff-Gauss map $\eta = \eta(u, v)$ is characterized by the condition

$$(\text{grad}(\Phi))_\eta = \left(\frac{\partial \Phi}{\partial x_1}(\eta), \frac{\partial \Phi}{\partial x_2}(\eta), \frac{\partial \Phi}{\partial x_3}(\eta) \right) = \varphi X_u \times X_v,$$

where φ is a positive function and \times is the standard cross product in \mathbb{R}^3 . Then we have

$$\eta = A^{-\frac{1}{2m}} \left(-(\beta')^{\frac{1}{2m-1}} \cos v, -(\beta')^{\frac{1}{2m-1}} \sin v, (\alpha')^{\frac{1}{2m-1}} \right)$$

where

$$A := (\alpha')^{\frac{2m}{2m-1}} + (\beta')^{\frac{2m}{2m-1}}.$$

We can compute that

$$\eta_u = -\frac{1}{2m-1} A^{-\frac{2m+1}{2m}} (\alpha')^{-\frac{2m-2}{2m-1}} (\beta')^{-\frac{2m-2}{2m-1}} (\alpha' \beta'' - \alpha'' \beta') X_u$$

and

$$\eta_v = -\frac{1}{\alpha} A^{-\frac{1}{2m}} (\beta')^{\frac{1}{2m-1}} X_v,$$

so that

$$k_1 = -\frac{1}{2m-1} A^{-\frac{2m+1}{2m}} (\alpha')^{-\frac{2m-2}{2m-1}} (\beta')^{-\frac{2m-2}{2m-1}} (\alpha' \beta'' - \alpha'' \beta')$$

and

$$k_2 = -\frac{1}{\alpha} A^{-\frac{1}{2m}} (\beta')^{\frac{1}{2m-1}}.$$

In particular, when $\beta(u) = u$, we have

$$k_1 = \frac{1}{2m-1} \left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{2m+1}{2m}} (\alpha')^{-\frac{2m-2}{2m-1}} \alpha'', \quad (3.1)$$

and

$$k_2 = -\frac{1}{\alpha} \left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{1}{2m}}. \quad (3.2)$$

On the other hand, when $\alpha(u) = u$, we have

$$k_1 = -\frac{1}{2m-1} \left(1 + (\beta')^{\frac{2m}{2m-1}} \right)^{-\frac{2m+1}{2m}} (\beta')^{-\frac{2m-2}{2m-1}} \beta'', \quad (3.3)$$

and

$$k_2 = -\frac{1}{u} \left(1 + (\beta')^{\frac{2m}{2m-1}} \right)^{-\frac{1}{2m}} (\beta')^{\frac{1}{2m-1}}. \quad (3.4)$$

4 The case where one of principal curvatures is constant

Let $(\mathbb{R}^3, \|\cdot\|)$ be the normed 3-space as in Section 3. Let M be a rotational surface in $(\mathbb{R}^3, \|\cdot\|)$ parametrized as

$$X(u, v) = (\alpha(u) \cos v, \alpha(u) \sin v, u)$$

where $\alpha > 0$ and $\alpha' \neq 0$.

In this section, we consider the case where k_1 or k_2 is constant.

(i) The case where $k_1 = 0$ identically. In this case, by (3.1), we see that $\alpha(u)$ is a nonconstant linear function, and the surface is a circular cylinder.

(ii) The case where $k_2 = \mu$ for a constant μ . In this case, we have $\mu < 0$ by (3.2). It suffices to consider the case where $\mu = -1$. Then we have

$$-\frac{1}{\alpha} \left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{1}{2m}} = -1,$$

$$\frac{d\alpha}{du} = \pm \frac{(1 - \alpha^{2m})^{\frac{2m-1}{2m}}}{\alpha^{2m-1}},$$

and

$$u(\alpha) = u_{\pm}(\alpha) := \pm \int \frac{\alpha^{2m-1}}{(1 - \alpha^{2m})^{\frac{2m-1}{2m}}} d\alpha = \mp (1 - \alpha^{2m})^{\frac{1}{2m}} + c$$

for a constant c . It satisfies

$$\alpha^{2m} + (u_{\pm}(\alpha) - c)^{2m} = 1,$$

and the resulting surface is a parallel translation of the unit sphere S .

(iii) The case where $k_1 = \mu$ for a non-zero constant μ . Then by (3.1) we have

$$\frac{1}{2m-1} \left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{2m+1}{2m}} (\alpha')^{-\frac{2m-2}{2m-1}} \alpha'' = \mu.$$

Multiplying by $-\alpha'$ we have

$$-\frac{1}{2m-1} \left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{2m+1}{2m}} (\alpha')^{\frac{1}{2m-1}} \alpha'' = -\mu \alpha'.$$

Integrating it we have

$$\left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{1}{2m}} = c_1 - \mu \alpha \ (> 0)$$

for a constant c_1 . Then

$$\frac{d\alpha}{du} = \pm \frac{\{1 - (c_1 - \mu\alpha)^{2m}\}^{\frac{2m-1}{2m}}}{(c_1 - \mu\alpha)^{2m-1}},$$

and we get

$$\begin{aligned} u_{\pm}(\alpha) &= \pm \int \frac{(c_1 - \mu\alpha)^{2m-1}}{\{1 - (c_1 - \mu\alpha)^{2m}\}^{\frac{2m-1}{2m}}} d\alpha \\ &= \pm \frac{1}{\mu} \{1 - (c_1 - \mu\alpha)^{2m}\}^{\frac{1}{2m}} + c_2 \end{aligned}$$

for a constant c_2 . It suffices to consider the case where $\mu = \pm 1$.

(iii-1) In the case where $\mu = 1$, we have

$$u_{\pm}(\alpha) = \pm \{1 - (c_1 - \alpha)^{2m}\}^{\frac{1}{2m}} + c_2,$$

and it satisfies

$$(c_1 - \alpha)^{2m} + (u_{\pm}(\alpha) - c_2)^{2m} = 1, \quad \alpha < c_1. \quad (4.1)$$

(iii-2) In the case where $\mu = -1$, we have

$$u_{\pm}(\alpha) = \mp \{1 - (c_3 + \alpha)^{2m}\}^{\frac{1}{2m}} + c_4$$

for constants c_3 and c_4 . It satisfies

$$(\alpha + c_3)^{2m} + (u_{\pm}(\alpha) - c_4)^{2m} = 1, \quad \alpha > -c_3. \quad (4.2)$$

If we choose $c_1 > 1$, $c_3 = -c_1$ and $c_4 = c_2$, then (4.1) and (4.2) are smoothly connected. Choosing the orientation suitably, we get a rotational torus with $k_1 = 1$.

5 Homogeneous linear Weingarten cases

Let $(\mathbb{R}^3, \|\cdot\|)$ be the normed 3-space as in Section 3. Let M be a rotational surface in $(\mathbb{R}^3, \|\cdot\|)$ parametrized as

$$X(u, v) = (\alpha(u) \cos v, \alpha(u) \sin v, u)$$

where $\alpha > 0$ and $\alpha' \neq 0$.

In this section, we consider the case where $k_1 + \lambda k_2 = 0$ for a non-zero constant λ . Then by (3.1) and (3.2), we have

$$\begin{aligned} \frac{1}{2m-1} \left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{2m+1}{2m}} (\alpha')^{-\frac{2m-2}{2m-1}} \alpha'' \\ - \frac{\lambda}{\alpha} \left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{1}{2m}} = 0, \end{aligned} \quad (5.1)$$

and

$$\frac{\alpha''}{(\alpha')^2 + (\alpha')^{\frac{2m-2}{2m-1}}} = (2m-1) \frac{\lambda}{\alpha}.$$

Multiplying by $2\alpha'$ we have

$$\frac{2\alpha'\alpha''}{(\alpha')^2 + (\alpha')^{\frac{2m-2}{2m-1}}} = 2(2m-1)\lambda \cdot \frac{\alpha'}{\alpha},$$

and

$$2(2m-1)\lambda \log \alpha = \int \frac{((\alpha')^2)'}{(\alpha')^2 + (\alpha')^{\frac{2m-2}{2m-1}}} du.$$

Setting

$$(\alpha')^{\frac{2}{2m-1}} =: Z$$

for the right hand side, we have

$$\begin{aligned} 2\lambda \log \alpha &= \int \frac{Z^{m-1}}{Z^m + 1} dZ = \frac{1}{m} \log(Z^m + 1) + c_1 \\ &= \frac{1}{m} \log((\alpha')^{\frac{2m}{2m-1}} + 1) + c_1 \end{aligned}$$

for a constant c_1 . Then

$$\frac{d\alpha}{du} = \pm \frac{(\alpha^{2m\lambda} - c_2^{2m\lambda})^{\frac{2m-1}{2m}}}{c_2^{(2m-1)\lambda}}$$

for a positive constant c_2 , and

$$u(\alpha) = u_{\pm}(\alpha) := \pm \int \frac{c_2^{(2m-1)\lambda}}{(\alpha^{2m\lambda} - c_2^{2m\lambda})^{\frac{2m-1}{2m}}} d\alpha.$$

Theorem 5.1. *A rotational surface in $(\mathbb{R}^3, \|\cdot\|)$ given by*

$$\bar{X}(\alpha, v) = (\alpha \cos v, \alpha \sin v, u(\alpha))$$

where $\alpha > 0$ and $u' \neq 0$, satisfies $k_1 + \lambda k_2 = 0$ for a non-zero constant λ , if and only if

$$u(\alpha) = \pm \int \frac{c_2^{(2m-1)\lambda}}{(\alpha^{2m\lambda} - c_2^{2m\lambda})^{\frac{2m-1}{2m}}} d\alpha$$

for a positive constant c_2 .

(i) The case where $\lambda > 0$. In this case, we have $\alpha > c_2 (> 0)$ and

$$u_{\pm}(\alpha) = \pm \int_{c_2}^{\alpha} \frac{c_2^{(2m-1)\lambda}}{(\rho^{2m\lambda} - c_2^{2m\lambda})^{\frac{2m-1}{2m}}} d\rho + c_3$$

for a constant c_3 . On the denomiator, we set $f(t) := t^{2m\lambda} - c_2^{2m\lambda}$. Then

$$f'(t) = 2m\lambda t^{2m\lambda-1}, \quad f(c_2) = 0, \quad f'(c_2) \neq 0.$$

So we have $f(t) = (t - c_2)\tilde{f}(t)$ for some smooth function $\tilde{f}(t)$ such that $\tilde{f}(c_2) \neq 0$.

Hence, noting that

$$0 < \frac{2m-1}{2m} < 1,$$

we can see that the above integral converges and

$$\lim_{\alpha \rightarrow c_2} \int_{c_2}^{\alpha} \frac{c_2^{(2m-1)\lambda}}{(\rho^{2m\lambda} - c_2^{2m\lambda})^{\frac{2m-1}{2m}}} d\rho = 0.$$

(i-1) The case where $(2m-1)\lambda > 1$. In this case, we have

$$\lim_{\alpha \rightarrow \infty} \int_{c_2}^{\alpha} \frac{c_2^{(2m-1)\lambda}}{(\rho^{2m\lambda} - c_2^{2m\lambda})^{\frac{2m-1}{2m}}} d\rho = d_1$$

for some positive value d_1 . Then

$$\begin{aligned}\lim_{\alpha \rightarrow c_2} u_{\pm}(\alpha) &= c_3, & \lim_{\alpha \rightarrow \infty} u_{\pm}(\alpha) &= c_3 \pm d_1, \\ \lim_{\alpha \rightarrow c_2} u'_{\pm}(\alpha) &= \pm\infty, & \lim_{\alpha \rightarrow \infty} u'_{\pm}(\alpha) &= 0.\end{aligned}$$

Let $\alpha_+(u)$ be the inverse function of $u_+(\alpha)$, which is increasing on $(c_3, c_3 + d_1)$ and satisfies

$$\lim_{u \rightarrow c_3} \alpha_+(u) = c_2, \quad \lim_{u \rightarrow c_3 + d_1} \alpha_+(u) = \infty, \quad \lim_{u \rightarrow c_3} \alpha'_+(u) = 0.$$

Let $\alpha_-(u)$ be the inverse function of $u_-(\alpha)$, which is decreasing on $(c_3 - d_1, c_3)$ and satisfies

$$\lim_{u \rightarrow c_3} \alpha_-(u) = c_2, \quad \lim_{u \rightarrow c_3 - d_1} \alpha_-(u) = \infty, \quad \lim_{u \rightarrow c_3} \alpha'_-(u) = 0.$$

We define a function $\hat{\alpha}(u)$ on $(c_3 - d_1, c_3 + d_1)$ by

$$\hat{\alpha}(u) = \begin{cases} \alpha_+(u), & c_3 < u < c_3 + d_1 \\ \alpha_-(u), & c_3 - d_1 < u < c_3 \\ c_2, & u = c_3. \end{cases}$$

It is a C^1 -function on $(c_3 - d_1, c_3 + d_1)$ such that

$$\hat{\alpha}'(u) = \begin{cases} \alpha'_+(u), & c_3 < u < c_3 + d_1 \\ \alpha'_-(u), & c_3 - d_1 < u < c_3 \\ 0, & u = c_3. \end{cases}$$

For $u \in (c_3 - d_1, c_3) \cup (c_3, c_3 + d_1)$, $\hat{\alpha}(u)$ satisfies the equation (5.1). Then we can see that

$$\lim_{u \rightarrow c_3} (\hat{\alpha}'(u))^{-\frac{2m-2}{2m-1}} \hat{\alpha}''(u) = \frac{(2m-1)\lambda}{c_2},$$

and since $m \geq 2$,

$$\lim_{u \rightarrow c_3} \hat{\alpha}''(u) = 0.$$

Hence, $\hat{\alpha}(u)$ is a C^2 -function on $(c_3 - d_1, c_3 + d_1)$.

Theorem 5.2. *Under the notation above in (i-1), the rotational surface in $(\mathbb{R}^3, \|\cdot\|)$ given by*

$$\hat{X}(u, v) = (\hat{\alpha}(u) \cos v, \hat{\alpha}(u) \sin v, u), \quad (u, v) \in (c_3 - d_1, c_3 + d_1) \times [0, 2\pi]$$

satisfies $k_1 + \lambda k_2 = 0$ for a constant λ with $(2m-1)\lambda > 1$.

(i-2) The case where $0 < (2m - 1)\lambda \leq 1$. In this case, we have

$$\lim_{\alpha \rightarrow \infty} \int_{c_2}^{\alpha} \frac{c_2^{(2m-1)\lambda}}{(\rho^{2m\lambda} - c_2^{2m\lambda})^{\frac{2m-1}{2m}}} d\rho = \infty.$$

Then

$$\begin{aligned} \lim_{\alpha \rightarrow c_2} u_{\pm}(\alpha) &= c_3, & \lim_{\alpha \rightarrow \infty} u_{\pm}(\alpha) &= \pm\infty, \\ \lim_{\alpha \rightarrow c_2} u'_{\pm}(\alpha) &= \pm\infty, & \lim_{\alpha \rightarrow \infty} u'_{\pm}(\alpha) &= 0. \end{aligned}$$

Let $\alpha_+(u)$ be the inverse function of $u_+(\alpha)$, which is increasing on (c_3, ∞) and satisfies

$$\lim_{u \rightarrow c_3} \alpha_+(u) = c_2, \quad \lim_{u \rightarrow \infty} \alpha_+(u) = \infty, \quad \lim_{u \rightarrow c_3} \alpha'_+(u) = 0.$$

Let $\alpha_-(u)$ be the inverse function of $u_-(\alpha)$, which is decreasing on $(-\infty, c_3)$ and satisfies

$$\lim_{u \rightarrow c_3} \alpha_-(u) = c_2, \quad \lim_{u \rightarrow -\infty} \alpha_-(u) = \infty, \quad \lim_{u \rightarrow c_3} \alpha'_-(u) = 0.$$

We define a function $\hat{\alpha}(u)$ on $(-\infty, \infty)$ by

$$\hat{\alpha}(u) = \begin{cases} \alpha_+(u), & c_3 < u < \infty \\ \alpha_-(u), & -\infty < u < c_3 \\ c_2, & u = c_3. \end{cases}$$

As in the case (i-1), we can see that $\hat{\alpha}(u)$ is a C^2 -function on $(-\infty, \infty)$.

Theorem 5.3. *Under the notation above in (i-2), the rotational surface in $(\mathbb{R}^3, \|\cdot\|)$ given by*

$$\hat{X}(u, v) = (\hat{\alpha}(u) \cos v, \hat{\alpha}(u) \sin v, u), \quad (u, v) \in (-\infty, \infty) \times [0, 2\pi]$$

satisfies $k_1 + \lambda k_2 = 0$ for a constant λ with $0 < (2m - 1)\lambda \leq 1$.

(ii) The case where $\lambda < 0$. In this case, we have $0 < \alpha < c_2$ and

$$\begin{aligned} u_{\pm}(\alpha) &= \pm \int_{c_2}^{\alpha} \frac{c_2^{(2m-1)\lambda}}{(\rho^{2m\lambda} - c_2^{2m\lambda})^{\frac{2m-1}{2m}}} d\rho + c_3 \\ &= \pm \int_{c_2}^{\alpha} \frac{\rho^{(2m-1)(-\lambda)}}{\{c_2^{2m(-\lambda)} - \rho^{2m(-\lambda)}\}^{\frac{2m-1}{2m}}} d\rho + c_3. \end{aligned}$$

As in the case (i), this integral converges and

$$\lim_{\alpha \rightarrow c_2} \int_{c_2}^{\alpha} \frac{\rho^{(2m-1)(-\lambda)}}{\{c_2^{2m(-\lambda)} - \rho^{2m(-\lambda)}\}^{\frac{2m-1}{2m}}} d\rho = 0.$$

Set

$$d_2 := - \lim_{\alpha \rightarrow 0} \int_{c_2}^{\alpha} \frac{\rho^{(2m-1)(-\lambda)}}{\{c_2^{2m(-\lambda)} - \rho^{2m(-\lambda)}\}^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\begin{aligned} \lim_{\alpha \rightarrow c_2} u_{\pm}(\alpha) &= c_3, & \lim_{\alpha \rightarrow 0} u_{\pm}(\alpha) &= c_3 \mp d_2, \\ \lim_{\alpha \rightarrow c_2} u'_{\pm}(\alpha) &= \pm\infty, & \lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha) &= 0. \end{aligned}$$

Correspondingly, we get a closed rotational surface of C^1 -class which is homeomorphic to the 2-sphere.

As in the case (i), we can see that the surface is of C^2 -class at points where $\alpha = c_2$. On the C^2 -smoothness at $\alpha = 0$, we recall that

$$u'_{\pm}(\alpha) = \pm \frac{\alpha^{(2m-1)(-\lambda)}}{\{c_2^{2m(-\lambda)} - \alpha^{2m(-\lambda)}\}^{\frac{2m-1}{2m}}}, \quad \alpha \in (0, c_2).$$

Hence, the limit $\lim_{\alpha \rightarrow 0} u''_{\pm}(\alpha)$ exists if and only if $(2m-1)(-\lambda) \geq 1$.

Moreover, using (3.3) and (3.4), we can see that k_1 and k_2 are extended continuously to $\alpha = 0$ if and only if $(-\lambda) \geq 1$.

Theorem 5.4. *There are closed rotational surface such that $k_1 + \lambda k_2 = 0$ for a constant $\lambda \leq -1$. They are homeomorphic to the 2-sphere.*

6 Inhomogeneous linear Weingarten cases

Let $(\mathbb{R}^3, \|\cdot\|)$ be the normed 3-space as in Section 3. Let M be a rotational surface in $(\mathbb{R}^3, \|\cdot\|)$ parametrized as

$$X(u, v) = (\alpha(u) \cos v, \alpha(u) \sin v, u)$$

where $\alpha > 0$ and $\alpha' \neq 0$.

In this section, we consider the case where $k_1 + \lambda k_2 = \mu$ for non-zero constants λ and μ . The case where $\lambda = 1$ (the constant mean curvature case) was treated in the previous paper [8]. Then by (3.1) and (3.2), we have

$$\begin{aligned} \frac{1}{2m-1} \left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{2m+1}{2m}} (\alpha')^{-\frac{2m-2}{2m-1}} \alpha'' \\ - \frac{\lambda}{\alpha} \left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{1}{2m}} = \mu. \end{aligned} \quad (6.1)$$

Multiplying by $-\alpha^\lambda \alpha'$ we have

$$\begin{aligned} \alpha^\lambda \left\{ -\frac{1}{2m-1} \left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{2m+1}{2m}} (\alpha')^{\frac{1}{2m-1}} \alpha'' \right\} \\ + \lambda \alpha^{\lambda-1} \alpha' \left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{1}{2m}} = -\mu \alpha^\lambda \alpha', \end{aligned}$$

and

$$\alpha^\lambda \left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{1}{2m}} = -\mu \int \alpha^\lambda \alpha' du.$$

6.1 The case where $\lambda = -1$

In this case, we have

$$\frac{1}{\alpha} \left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{1}{2m}} = -\mu \int \frac{\alpha'}{\alpha} du = -\mu \log \alpha + c_1 (> 0)$$

for a constant c_1 . Then

$$\frac{d\alpha}{du} = \pm \frac{\{1 - \alpha^{2m}(c_1 - \mu \log \alpha)^{2m}\}^{\frac{2m-1}{2m}}}{\alpha^{2m-1}(c_1 - \mu \log \alpha)^{2m-1}},$$

and

$$u(\alpha) = u_{\pm}(\alpha) := \pm \int \frac{\alpha^{2m-1}(c_1 - \mu \log \alpha)^{2m-1}}{\{1 - \alpha^{2m}(c_1 - \mu \log \alpha)^{2m}\}^{\frac{2m-1}{2m}}} d\alpha.$$

Theorem 6.1. *A rotational surface in $(\mathbb{R}^3, \|\cdot\|)$ given by*

$$\bar{X}(\alpha, v) = (\alpha \cos v, \alpha \sin v, u(\alpha))$$

where $\alpha > 0$ and $u' \neq 0$, satisfies $k_1 - k_2 = \mu$ for a non-zero constant μ , if and only if

$$u(\alpha) = \pm \int \frac{\alpha^{2m-1}(c_1 - \mu \log \alpha)^{2m-1}}{\{1 - \alpha^{2m}(c_1 - \mu \log \alpha)^{2m}\}^{\frac{2m-1}{2m}}} d\alpha$$

for a constant c_1 . Here we need to have

$$c_1 - \mu \log \alpha > 0, \quad \alpha(c_1 - \mu \log \alpha) < 1.$$

It suffices to consider the case where $\mu = \pm 1$.

(i) The case where $\mu = 1$. We have

$$u_{\pm}(\alpha) = \pm \int \frac{\alpha^{2m-1}(c_1 - \log \alpha)^{2m-1}}{\{1 - \alpha^{2m}(c_1 - \log \alpha)^{2m}\}^{\frac{2m-1}{2m}}} d\alpha$$

where $0 < \alpha < e^{c_1}$ and $\alpha(c_1 - \log \alpha) < 1$. Set

$$g(t) := t(c_1 - \log t), \quad 0 < t < e^{c_1}.$$

Then

$$g'(t) = c_1 - 1 - \log t,$$

and we have the following table:

| | | | | | |
|---------|-------|------------|-------------|------------|-------------|
| t | (0) | \dots | e^{c_1-1} | \dots | (e^{c_1}) |
| $g'(t)$ | | $+$ | 0 | $-$ | |
| $g(t)$ | | \nearrow | e^{c_1-1} | \searrow | |

with

$$\lim_{t \rightarrow 0} g(t) = 0, \quad \lim_{t \rightarrow e^{c_1}} g(t) = 0.$$

(i-1) The case where $c_1 < 1$. In this case, we have $\alpha(c_1 - \log \alpha) < 1$ for $0 < \alpha < e^{c_1}$. So the domain is given by $0 < \alpha < e^{c_1}$, and

$$u_{\pm}(\alpha) = \pm \int_{e^{c_1}}^{\alpha} \frac{\rho^{2m-1}(c_1 - \log \rho)^{2m-1}}{\{1 - \rho^{2m}(c_1 - \log \rho)^{2m}\}^{\frac{2m-1}{2m}}} d\rho + c_2^{\pm}.$$

Set

$$d_1 := - \lim_{\alpha \rightarrow 0} \int_{e^{c_1}}^{\alpha} \frac{\rho^{2m-1}(c_1 - \log \rho)^{2m-1}}{\{1 - \rho^{2m}(c_1 - \log \rho)^{2m}\}^{\frac{2m-1}{2m}}} d\rho.$$

Then we have

$$\lim_{\alpha \rightarrow e^{c_1}} u_{\pm}(\alpha) = c_2^{\pm}, \quad \lim_{\alpha \rightarrow 0} u_{\pm}(\alpha) = c_2^{\pm} \mp d_1,$$

$$\lim_{\alpha \rightarrow e^{c_1}} u'_{\pm}(\alpha) = 0, \quad \lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha) = 0.$$

On the C^2 -smoothness at $\alpha = 0$, we recall that

$$u'_{\pm}(\alpha) = \pm \frac{\{\alpha(c_1 - \log \alpha)\}^{2m-1}}{[1 - \{\alpha(c_1 - \log \alpha)\}^{2m}]^{\frac{2m-1}{2m}}}, \quad \alpha \in (0, e^{c_1}).$$

Since $m \geq 2$, we can see that $\lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha)$ exists. However, by (3.3) and (3.4), we find that k_1 and k_2 are not extended continuously to $\alpha = 0$.

(i-2) The case where $c_1 = 1$. In this case, the domain is given by $0 < \alpha < 1$ or $1 < \alpha < e$.

(i-2-1) On the domain $0 < \alpha < 1$, we have

$$u_{\pm}(\alpha) = \pm \int_0^{\alpha} \frac{\rho^{2m-1}(1 - \log \rho)^{2m-1}}{\{1 - \rho^{2m}(1 - \log \rho)^{2m}\}^{\frac{2m-1}{2m}}} d\rho + c_3^{\pm}.$$

On the demoninator, we set

$$f(t) := 1 - t(1 - \log t).$$

Then

$$f'(t) = \log t, \quad f''(t) = \frac{1}{t}, \quad f(1) = 0, \quad f'(1) = 0, \quad f''(1) = 1 \neq 0.$$

So we have $f(t) = (1-t)^2 \tilde{f}(t)$ for a smooth function $\tilde{f}(t)$ such that $\tilde{f}(1) \neq 0$.

Hence, noting that $m \geq 2$ and

$$1 < 2 \times \frac{2m-1}{2m} < 2,$$

we find that

$$\lim_{\alpha \rightarrow 1} \int_0^{\alpha} \frac{\rho^{2m-1}(1 - \log \rho)^{2m-1}}{\{1 - \rho^{2m}(1 - \log \rho)^{2m}\}^{\frac{2m-1}{2m}}} d\rho = \infty.$$

Thus we have

$$\lim_{\alpha \rightarrow 0} u_{\pm}(\alpha) = c_3^{\pm}, \quad \lim_{\alpha \rightarrow 1} u_{\pm}(\alpha) = \pm\infty, \quad \lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha) = 0.$$

Correspondingly, we get a rotational surface of C^1 -class.

The C^2 -smoothness at $\alpha = 0$ is analogous to the case (i-1). We can see that $\lim_{\alpha \rightarrow 0} u''_{\pm}(\alpha)$ exists, but k_1 and k_2 are not extended continuously to $\alpha = 0$.

(i-2-2) On the domain $1 < \alpha < e$, we have

$$u_{\pm}(\alpha) = \pm \int_e^{\alpha} \frac{\rho^{2m-1}(1 - \log \rho)^{2m-1}}{\{1 - \rho^{2m}(1 - \log \rho)^{2m}\}^{\frac{2m-1}{2m}}} d\rho + c_3^{\pm}.$$

As in the case (i-2-1), we can see that

$$\lim_{\alpha \rightarrow 1} \int_e^{\alpha} \frac{\rho^{2m-1}(1 - \log \rho)^{2m-1}}{\{1 - \rho^{2m}(1 - \log \rho)^{2m}\}^{\frac{2m-1}{2m}}} d\rho = -\infty,$$

and

$$\lim_{\alpha \rightarrow e} u_{\pm}(\alpha) = c_3^{\pm}, \quad \lim_{\alpha \rightarrow 1} u_{\pm}(\alpha) = \mp\infty, \quad \lim_{\alpha \rightarrow e} u'_{\pm}(\alpha) = 0.$$

(i-3) The case where $c_1 > 1$. In this case, there exist two numbers α_1 and α_2 such that

$$\alpha_1(c_1 - \log \alpha_1) = 1, \quad 0 < \alpha_1 < e^{c_1-1},$$

$$\alpha_2(c_1 - \log \alpha_2) = 1, \quad e^{c_1-1} < \alpha_2 < e^{c_1},$$

and the domain is given by $0 < \alpha < \alpha_1$ or $\alpha_2 < \alpha < e^{c_1}$.

(i-3-1) On the domain $0 < \alpha < \alpha_1$, we have

$$u_{\pm}(\alpha) = \pm \int_{\alpha_1}^{\alpha} \frac{\rho^{2m-1}(c_1 - \log \rho)^{2m-1}}{\{1 - \rho^{2m}(c_1 - \log \rho)^{2m}\}^{\frac{2m-1}{2m}}} d\rho + c_4.$$

On the denominator, we set

$$f(t) := 1 - t(c_1 - \log t).$$

Then

$$f'(t) = 1 - c_1 + \log t, \quad f(\alpha_1) = 0,$$

$$f'(\alpha_1) = 1 - c_1 + \log \alpha_1 = 1 - \frac{1}{\alpha_1},$$

where we use that $\alpha_1(c_1 - \log \alpha_1) = 1$ for the last equality. Since $c_1 > 1$, we can see that $\alpha_1 \neq 1$, and $f'(\alpha_1) \neq 0$. Thus we have $f(t) = (\alpha_1 - t)\tilde{f}(t)$ for a smooth function $\tilde{f}(t)$ with $\tilde{f}(\alpha_1) \neq 0$.

Since

$$0 < \frac{2m-1}{2m} < 1,$$

the above integral converges and

$$\lim_{\alpha \rightarrow \alpha_1} \int_{\alpha_1}^{\alpha} \frac{\rho^{2m-1}(c_1 - \log \rho)^{2m-1}}{\{1 - \rho^{2m}(c_1 - \log \rho)^{2m}\}^{\frac{2m-1}{2m}}} d\rho = 0.$$

Set

$$d_2 := - \lim_{\alpha \rightarrow 0} \int_{\alpha_1}^{\alpha} \frac{\rho^{2m-1}(c_1 - \log \rho)^{2m-1}}{\{1 - \rho^{2m}(c_1 - \log \rho)^{2m}\}^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\begin{aligned} \lim_{\alpha \rightarrow \alpha_1} u_{\pm}(\alpha) &= c_4, & \lim_{\alpha \rightarrow 0} u_{\pm}(\alpha) &= c_4 \mp d_2, \\ \lim_{\alpha \rightarrow \alpha_1} u'_{\pm}(\alpha) &= \pm\infty, & \lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha) &= 0. \end{aligned}$$

Correspondingly, we get a rotational surface of C^1 -class which is homeomorphic to the 2-sphere.

Using (6.1), the C^2 -smoothness at $\alpha = \alpha_1$ can be seen as in Section 5. The C^2 -smoothness at $\alpha = 0$ is analogous to the case (i-1). We see that $\lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha)$ exists, but k_1 and k_2 are not extended continuously to $\alpha = 0$.

(i-3-2) On the domain $\alpha_2 < \alpha < e^{c_1}$, we have

$$u_{\pm}(\alpha) = \pm \int_{\alpha_2}^{\alpha} \frac{\rho^{2m-1}(c_1 - \log \rho)^{2m-1}}{\{1 - \rho^{2m}(c_1 - \log \rho)^{2m}\}^{\frac{2m-1}{2m}}} d\rho + c_4.$$

As in the case (i-3-1), it converges. Set

$$d_3 := \lim_{\alpha \rightarrow e^{c_1}} \int_{\alpha_2}^{\alpha} \frac{\rho^{2m-1}(c_1 - \log \rho)^{2m-1}}{\{1 - \rho^{2m}(c_1 - \log \rho)^{2m}\}^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\begin{aligned} \lim_{\alpha \rightarrow \alpha_2} u_{\pm}(\alpha) &= c_4, & \lim_{\alpha \rightarrow e^{c_1}} u_{\pm}(\alpha) &= c_4 \pm d_3, \\ \lim_{\alpha \rightarrow \alpha_2} u'_{\pm}(\alpha) &= \pm\infty, & \lim_{\alpha \rightarrow e^{c_1}} u'_{\pm}(\alpha) &= 0. \end{aligned}$$

The C^2 -smoothness at $\alpha = \alpha_2$ can be seen as before.

(ii) The case where $\mu = -1$. We have

$$u_{\pm}(\alpha) = \pm \int \frac{\alpha^{2m-1}(c_1^* + \log \alpha)^{2m-1}}{\{1 - \alpha^{2m}(c_1^* + \log \alpha)^{2m}\}^{\frac{2m-1}{2m}}} d\alpha$$

where $\alpha > e^{-c_1^*}$ and $\alpha(c_1^* + \log \alpha) < 1$. Set

$$g(t) := t(c_1^* + \log t), \quad t > e^{-c_1^*}.$$

Then, for $t > e^{-c_1^*}$, we have

$$g'(t) = c_1^* + 1 + \log t > 1, \quad \lim_{t \rightarrow e^{-c_1^*}} g(t) = 0, \quad \lim_{t \rightarrow \infty} g(t) = \infty.$$

So there exists a unique number α_3 such that

$$\alpha_3(c_1^* + \log \alpha_3) = 1, \quad \alpha_3 > e^{-c_1^*},$$

and the domain is given by $e^{-c_1^*} < \alpha < \alpha_3$. Thus we have

$$u_{\pm}(\alpha) = \pm \int_{\alpha_3}^{\alpha} \frac{\rho^{2m-1}(c_1^* + \log \rho)^{2m-1}}{\{1 - \rho^{2m}(c_1^* + \log \rho)^{2m}\}^{\frac{2m-1}{2m}}} d\rho + c_5.$$

On the denominator, we set

$$f(t) := 1 - t(c_1^* + \log t).$$

Then

$$\begin{aligned} f'(t) &= -(c_1^* + 1 + \log t), \quad f(\alpha_3) = 0, \\ f'(\alpha_3) &= -(c_1^* + 1 + \log \alpha_3) = -\left(1 + \frac{1}{\alpha_3}\right) < -1, \end{aligned}$$

where we use that $\alpha_3(c_1^* + \log \alpha_3) = 1$ for the last equality. Thus we have $f(t) = (\alpha_3 - t)\tilde{f}(t)$ for a smooth function $\tilde{f}(t)$ with $\tilde{f}(\alpha_3) \neq 0$.

Hence, the above integral converges. Set

$$d_4 := - \lim_{\alpha \rightarrow e^{-c_1^*}} \int_{\alpha_3}^{\alpha} \frac{\rho^{2m-1}(c_1^* + \log \rho)^{2m-1}}{\{1 - \rho^{2m}(c_1^* + \log \rho)^{2m}\}^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\begin{aligned} \lim_{\alpha \rightarrow \alpha_3} u_{\pm}(\alpha) &= c_5, \quad \lim_{\alpha \rightarrow e^{-c_1^*}} u_{\pm}(\alpha) = c_5 \mp d_4, \\ \lim_{\alpha \rightarrow \alpha_3} u'_{\pm}(\alpha) &= \pm\infty, \quad \lim_{\alpha \rightarrow e^{-c_1^*}} u'_{\pm}(\alpha) = 0. \end{aligned}$$

The C^2 -smoothness at $\alpha = \alpha_3$ can be seen as before.

6.2 Connecting argument for $\lambda = -1$

Let $u_{1\pm}$ denote the u_{\pm} in the case (i) of Subsection 6.1, and $u_{2\pm}$ denote the u_{\pm} in the case (ii) of Subsection 6.1.

(C1) Connecting (i-1) and (ii) of Subsection 6.1.

(C1-1) If we choose

$$c_1^* = -c_1, \quad c_2^+ = c_5 - d_4, \quad c_2^- = c_5 + d_4,$$

then by connecting the surfaces in the cases (i-1) and (ii), we get a closed rotational surface of C^1 -class which is homeomorphic to the 2-sphere. Notice that α_3 depends on c_1 .

Noting the orientation, by (3.3) and (3.4) for $\beta = u_{1+}(\alpha)$ and $\beta = u_{2+}(\alpha)$, we have

$$-\frac{1}{2m-1} \left(1 + (u'_{1+})^{\frac{2m}{2m-1}}\right)^{-\frac{2m+1}{2m}} (u'_{1+})^{-\frac{2m-2}{2m-1}} u''_{1+}$$

$$+\frac{1}{\alpha} \left(1 + (u'_{1+})^{\frac{2m}{2m-1}}\right)^{-\frac{1}{2m}} (u'_{1+})^{\frac{1}{2m-1}} = 1, \quad \alpha \in (0, e^{c_1}),$$

and

$$-\frac{1}{2m-1} \left(1 + (u'_{2+})^{\frac{2m}{2m-1}}\right)^{-\frac{2m+1}{2m}} (u'_{2+})^{-\frac{2m-2}{2m-1}} u''_{2+} \\ +\frac{1}{\alpha} \left(1 + (u'_{2+})^{\frac{2m}{2m-1}}\right)^{-\frac{1}{2m}} (u'_{2+})^{\frac{1}{2m-1}} = -1, \quad \alpha \in (e^{c_1}, \alpha_3).$$

Thus we find that

$$\lim_{\alpha \rightarrow e^{c_1}} (u'_{1+}(\alpha))^{-\frac{2m-2}{2m-1}} u''_{1+}(\alpha) = -(2m-1)$$

and

$$\lim_{\alpha \rightarrow e^{c_1}} (u'_{2+}(\alpha))^{-\frac{2m-2}{2m-1}} u''_{2+}(\alpha) = 2m-1.$$

Since $m \geq 2$, we have

$$\lim_{\alpha \rightarrow e^{c_1}} u''_{1+}(\alpha) = \lim_{\alpha \rightarrow e^{c_1}} u''_{2+}(\alpha) = 0.$$

Hence, $u_{1+}(\alpha)$ and $u_{2+}(\alpha)$ are connected C^2 -smoothly at $\alpha = e^{c_1}$, but k_1 is not continuous at $\alpha = e^{c_1}$. The connectedness for $u_{1-}(\alpha)$ and $u_{2-}(\alpha)$ is analogous.

(C1-2) On the other hand, choose

$$c_1^* = -c_1, \quad c_2^+ = c_5 + d_4, \quad c_2^- = c_5 - d_4.$$

If $d_1 \neq d_4$, then we have a rotational surface of C^1 -class.

In this case, noting the difference of the orientation, we have

$$-\frac{1}{2m-1} \left(1 + (u'_{1+})^{\frac{2m}{2m-1}}\right)^{-\frac{2m+1}{2m}} (u'_{1+})^{-\frac{2m-2}{2m-1}} u''_{1+} \\ +\frac{1}{\alpha} \left(1 + (u'_{1+})^{\frac{2m}{2m-1}}\right)^{-\frac{1}{2m}} (u'_{1+})^{\frac{1}{2m-1}} = 1, \quad \alpha \in (0, e^{c_1}),$$

and

$$-\frac{1}{2m-1} \left(1 + (u'_{2-})^{\frac{2m}{2m-1}}\right)^{-\frac{2m+1}{2m}} (u'_{2-})^{-\frac{2m-2}{2m-1}} u''_{2-} \\ +\frac{1}{\alpha} \left(1 + (u'_{2-})^{\frac{2m}{2m-1}}\right)^{-\frac{1}{2m}} (u'_{2-})^{\frac{1}{2m-1}} = 1, \quad \alpha \in (e^{c_1}, \alpha_3).$$

We can see that $u_{1+}(\alpha)$ and $u_{2-}(\alpha)$ are connected C^2 -smoothly at $\alpha = e^{c_1}$, and k_1 is continuous at $\alpha = e^{c_1}$ in this case. The connectedness for $u_{1-}(\alpha)$ and $u_{2+}(\alpha)$ is analogous.

This surface is of C^2 -class, but as mentioned in (i-1) of Subsection 6.1, k_1 and k_2 are not extended continuously to $\alpha = 0$.

Remark. If $d_1 = d_4$, then it has a singularity at a point where $\alpha = 0$.

(C2) Connecting (i-2-2) and (ii) of Subsection 6.1. If we choose

$$c_1^* = -1, \quad c_3^+ = c_5 + d_4, \quad c_3^- = c_5 - d_4,$$

then we have a rotational surface of C^1 -class.

As in the case (C1-2), $u_{1+}(\alpha)$ and $u_{2-}(\alpha)$ are connected C^2 -smoothly at $\alpha = e$, and k_1 is continuous at $\alpha = e$. The connectedness for $u_{1-}(\alpha)$ and $u_{2+}(\alpha)$ is analogous.

(C3) Connecting (i-3-2) and (ii) of Subsection 6.1. If we choose

$$c_1^* = -c_1, \quad c_4 = c_5 + d_4 - d_3,$$

then we have a rotational surface of C^1 -class. Notice that α_2 and α_3 depend on c_1 .

As in the case (C1-2) and (C2), we can see that $u_{1+}(\alpha)$ and $u_{2-}(\alpha)$ are connected C^2 -smoothly at $\alpha = e^{c_1}$, and k_1 is continuous at $\alpha = e^{c_1}$. Analogously, we can see that $u_{1-}(\alpha)$ and $u_{2+}(\alpha)$ have the same derivatives at $\alpha = e^{c_1}$.

Let Γ be the C^2 -curve given by connecting the graphs of u_{1-} , u_{1+} , u_{2-} and u_{2+} . Since Γ has the same derivatives at the end points, it can be extended periodically as a C^2 -curve Γ^* . As a result, we obtain a periodic rotational surface which satisfies $k_1 - k_2 = 1$ for a suitable choice of orientation.

Remark. If $d_3 = d_4$, then we may have a rotational torus such that $k_1 - k_2$ is a non-zero constant.

6.3 The case where $\lambda \neq -1$

In this case, we have

$$\alpha^\lambda \left((\alpha')^{\frac{2m}{2m-1}} + 1 \right)^{-\frac{1}{2m}} = -\mu \int \alpha^\lambda \alpha' du = -\frac{\mu}{\lambda+1} \alpha^{\lambda+1} + c_1 (> 0)$$

for a constant c_1 . Then

$$\frac{d\alpha}{du} = \pm \frac{[(\lambda+1)^{2m} \alpha^{2m\lambda} - \{c_1(\lambda+1) - \mu\alpha^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}}{\{c_1(\lambda+1) - \mu\alpha^{\lambda+1}\}^{2m-1}}$$

and

$$u(\alpha) = \pm \int \frac{\{c_1(\lambda+1) - \mu\alpha^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} \alpha^{2m\lambda} - \{c_1(\lambda+1) - \mu\alpha^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\alpha.$$

Theorem 6.2. A rotational surface in $(\mathbb{R}^3, \|\cdot\|)$ given by

$$\bar{X}(\alpha, v) = (\alpha \cos v, \alpha \sin v, u(\alpha))$$

where $\alpha > 0$ and $u' \neq 0$, satisfies $k_1 + \lambda k_2 = \mu$ for non-zero constants $\lambda \neq -1$ and μ , if and only if

$$u(\alpha) = \pm \int \frac{\{c_1(\lambda+1) - \mu\alpha^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} \alpha^{2m\lambda} - \{c_1(\lambda+1) - \mu\alpha^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\alpha$$

for a constant c_1 . Here we need to have

$$0 < c_1 - \frac{\mu}{\lambda+1} \alpha^{\lambda+1} < \alpha^\lambda.$$

We divide the cases as $\lambda > 0$, $-1 < \lambda < 0$ or $\lambda < -1$. It suffices to consider the case where $\mu = \pm 1$.

(i) The case where $\lambda > 0$ and $\mu = 1$. In this case, we have

$$u_{\pm}(\alpha) = \pm \int \frac{\{c_1(\lambda+1) - \alpha^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} \alpha^{2m\lambda} - \{c_1(\lambda+1) - \alpha^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\alpha$$

and

$$0 < c_1(\lambda+1) - \alpha^{\lambda+1} < (\lambda+1)\alpha^\lambda. \quad (6.2)$$

From the first inequality of (6.2), we need to have $c_1 > 0$, and

$$0 < \alpha < \{c_1(\lambda+1)\}^{\frac{1}{\lambda+1}} =: \alpha_4.$$

For the second inequality of (6.2), set

$$f(t) := (\lambda+1)t^\lambda - c_1(\lambda+1) + t^{\lambda+1}, \quad 0 < t < \alpha_4.$$

Since it is strictly increasing and

$$\lim_{t \rightarrow 0} f(t) = -c_1(\lambda+1) < 0, \quad \lim_{t \rightarrow \alpha_4} f(t) = (\lambda+1)\alpha_4^\lambda > 0,$$

there exists a unique number $\alpha_5 \in (0, \alpha_4)$ such that $f(\alpha_5) = 0$.

Thus the domain is given by $\alpha_5 < \alpha < \alpha_4$, and

$$u_{\pm}(\alpha) = \pm \int_{\alpha_5}^{\alpha} \frac{\{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} \rho^{2m\lambda} - \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho + c_2.$$

On the denominator, since $f(\alpha_5) = 0$ and $f'(\alpha_5) > 0$, we have $f(t) = (t-\alpha_5)\tilde{f}(t)$ for a smooth function $\tilde{f}(t)$ with $\tilde{f}(\alpha_5) \neq 0$. So the above integral converges.

Set

$$d_1 := \lim_{\alpha \rightarrow \alpha_4} \int_{\alpha_5}^{\alpha} \frac{\{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} \rho^{2m\lambda} - \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho.$$

Then we have

$$\lim_{\alpha \rightarrow \alpha_5} u_{\pm}(\alpha) = c_2, \quad \lim_{\alpha \rightarrow \alpha_4} u_{\pm}(\alpha) = c_2 \pm d_1,$$

$$\lim_{\alpha \rightarrow \alpha_5} u'_{\pm}(\alpha) = \pm\infty, \quad \lim_{\alpha \rightarrow \alpha_4} u'_{\pm}(\alpha) = 0.$$

(ii) The case where $\lambda > 0$ and $\mu = -1$. We have

$$u_{\pm}(\alpha) = \pm \int \frac{\{c_1^*(\lambda+1) + \alpha^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} \alpha^{2m\lambda} - \{c_1^*(\lambda+1) + \alpha^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\alpha$$

and

$$0 < c_1^*(\lambda + 1) + \alpha^{\lambda+1} < (\lambda + 1)\alpha^\lambda. \quad (6.3)$$

(ii-1) The case where $c_1^* = 0$. We have

$$\begin{aligned} u_\pm(\alpha) &= \pm \int \frac{\alpha^{(2m-1)(\lambda+1)}}{\{(\lambda+1)^{2m}\alpha^{2m\lambda} - \alpha^{2m(\lambda+1)}\}^{\frac{2m-1}{2m}}} d\alpha \\ &= \pm \int \frac{\alpha^{2m-1}}{\{(\lambda+1)^{2m} - \alpha^{2m}\}^{\frac{2m-1}{2m}}} d\alpha = \mp \{(\lambda+1)^{2m} - \alpha^{2m}\}^{\frac{1}{2m}} + c_3. \end{aligned}$$

It satisfies

$$\alpha^{2m} + (u_\pm(\alpha) - c_3)^{2m} = (\lambda + 1)^{2m},$$

and the surface is homothetic to the unit sphere S .

(ii-2) The case where $c_1^* > 0$. In this case, the first inequality of (6.3) holds obviously. For the second inequality of (6.3), set

$$f(t) := (\lambda + 1)t^\lambda - c_1^*(\lambda + 1) - t^{\lambda+1}, \quad t > 0.$$

Then

$$f'(t) = (\lambda + 1)(\lambda - t)t^{\lambda-1}$$

and

$$\lim_{t \rightarrow 0} f(t) = -c_1^*(\lambda + 1) < 0, \quad \lim_{t \rightarrow \infty} f(t) = -\infty.$$

So we need to have

$$f(\lambda) = \lambda^\lambda - c_1^*(\lambda + 1) > 0,$$

that is,

$$0 < c_1^* < \frac{\lambda^\lambda}{\lambda + 1}.$$

Then there are two numbers α_6 and α_7 such that

$$f(\alpha_6) = f(\alpha_7) = 0, \quad 0 < \alpha_6 < \lambda < \alpha_7.$$

Thus the domain is given by $\alpha_6 < \alpha < \alpha_7$, and

$$u_\pm(\alpha) = \pm \int_{\alpha_6}^{\alpha} \frac{\{c_1^*(\lambda + 1) + \rho^{\lambda+1}\}^{2m-1}}{[(\lambda + 1)^{2m}\rho^{2m\lambda} - \{c_1^*(\lambda + 1) + \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho + c_3.$$

On the denominator, we have

$$f(\alpha_6) = f(\alpha_7) = 0, \quad f'(\alpha_6) \neq 0, \quad f'(\alpha_7) \neq 0.$$

So the above integral converges as α tends to α_6 and α_7 .

Set

$$d_2 := \lim_{\alpha \rightarrow \alpha_7} \int_{\alpha_6}^{\alpha} \frac{\{c_1^*(\lambda + 1) + \rho^{\lambda+1}\}^{2m-1}}{[(\lambda + 1)^{2m}\rho^{2m\lambda} - \{c_1^*(\lambda + 1) + \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\begin{aligned}\lim_{\alpha \rightarrow \alpha_6} u_{\pm}(\alpha) &= c_3, & \lim_{\alpha \rightarrow \alpha_7} u_{\pm}(\alpha) &= c_3 \pm d_2, \\ \lim_{\alpha \rightarrow \alpha_6} u'_{\pm}(\alpha) &= \pm\infty, & \lim_{\alpha \rightarrow \alpha_7} u'_{\pm}(\alpha) &= \pm\infty.\end{aligned}$$

Let Γ be the C^2 -curve given by connecting u_+ and u_- . Since Γ has the same derivatives at the end points, it can be extended periodically as a C^2 -curve Γ^* . Thus we get a periodic rotational surface.

(ii-3) The case where $c_1^* < 0$. From the first inequality of (6.3), we have

$$\alpha > \{(-c_1^*)(\lambda + 1)\}^{\frac{1}{\lambda+1}} =: \alpha_8.$$

For the second inequality of (6.3), set

$$f(t) := (\lambda + 1)t^{\lambda} - c_1^*(\lambda + 1) - t^{\lambda+1}, \quad t > \alpha_8.$$

Then

$$f'(t) = (\lambda + 1)(\lambda - t)t^{\lambda-1}$$

and

$$\lim_{t \rightarrow \alpha_8} f(t) = (\lambda + 1)\alpha_8^{\lambda} > 0, \quad \lim_{t \rightarrow \infty} f(t) = -\infty.$$

So there is a unique number $\alpha_9 \in (\alpha_8, \infty)$ such that $f(\alpha_9) = 0$.

Thus the domain is given by $\alpha_8 < \alpha < \alpha_9$, and

$$u_{\pm}(\alpha) = \pm \int_{\alpha_9}^{\alpha} \frac{\{c_1^*(\lambda + 1) + \rho^{\lambda+1}\}^{2m-1}}{[(\lambda + 1)^{2m} \rho^{2m\lambda} - \{c_1^*(\lambda + 1) + \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho + c_3.$$

On the denominator, we can see that $f(\alpha_9) = 0$ and $f'(\alpha_9) \neq 0$, so that the above integral converges.

Set

$$d_3 := - \lim_{\alpha \rightarrow \alpha_8} \int_{\alpha_9}^{\alpha} \frac{\{c_1^*(\lambda + 1) + \rho^{\lambda+1}\}^{2m-1}}{[(\lambda + 1)^{2m} \rho^{2m\lambda} - \{c_1^*(\lambda + 1) + \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\begin{aligned}\lim_{\alpha \rightarrow \alpha_9} u_{\pm}(\alpha) &= c_3, & \lim_{\alpha \rightarrow \alpha_8} u_{\pm}(\alpha) &= c_3 \mp d_3, \\ \lim_{\alpha \rightarrow \alpha_9} u'_{\pm}(\alpha) &= \pm\infty, & \lim_{\alpha \rightarrow \alpha_8} u'_{\pm}(\alpha) &= 0.\end{aligned}$$

(iii) The case where $-1 < \lambda < 0$ and $\mu = 1$. We have

$$\begin{aligned}u_{\pm}(\alpha) &= \pm \int \frac{\{c_1(\lambda + 1) - \alpha^{\lambda+1}\}^{2m-1}}{[(\lambda + 1)^{2m} \alpha^{2m\lambda} - \{c_1(\lambda + 1) - \alpha^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\alpha \\ &= \pm \int \frac{\alpha^{(2m-1)(-\lambda)} \{c_1(\lambda + 1) - \alpha^{\lambda+1}\}^{2m-1}}{[(\lambda + 1)^{2m} - \alpha^{2m(-\lambda)} \{c_1(\lambda + 1) - \alpha^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\alpha,\end{aligned}$$

and

$$c_1(\lambda + 1) - \alpha^{\lambda+1} > 0, \quad \alpha^{(-\lambda)}\{c_1(\lambda + 1) - \alpha^{\lambda+1}\} < \lambda + 1. \quad (6.4)$$

From the first inequality of (6.4), we have $c_1 > 0$ and

$$0 < \alpha < \{c_1(\lambda + 1)\}^{\frac{1}{\lambda+1}} =: \alpha_{10}.$$

For the second inequality of (6.4), set

$$f(t) := \lambda + 1 - c_1(\lambda + 1)t^{(-\lambda)} + t, \quad 0 < t < \alpha_{10}.$$

Then

$$f'(t) = 1 - c_1(-\lambda)(\lambda + 1)t^{-(\lambda+1)}.$$

Set

$$\alpha_{11} := \{c_1(-\lambda)(\lambda + 1)\}^{\frac{1}{\lambda+1}} < \alpha_{10},$$

where we use that $0 < -\lambda < 1$ for the inequality.

So we have the following table:

| | | | | | |
|---------|-------|------------|---------------|------------|-----------------|
| t | (0) | \cdots | α_{11} | \cdots | (α_{10}) |
| $f'(t)$ | | $-$ | 0 | $+$ | |
| $f(t)$ | | \searrow | | \nearrow | |

with

$$\lim_{t \rightarrow 0} f(t) = \lim_{t \rightarrow \alpha_{10}} f(t) = \lambda + 1 > 0,$$

and

$$f(\alpha_{11}) = (\lambda + 1) \left[1 - (-\lambda)^{\frac{(-\lambda)}{\lambda+1}} \{c_1(\lambda + 1)\}^{\frac{1}{\lambda+1}} \right].$$

(iii-1) The case where $0 < c_1 < \{(\lambda + 1)(-\lambda)^{(-\lambda)}\}^{-1}$. In this case, we have $f(\alpha_{11}) > 0$ and the domain is given by $0 < \alpha < \alpha_{10}$. Then

$$u_{\pm}(\alpha) = \pm \int_0^{\alpha} \frac{\rho^{(2m-1)(-\lambda)}\{c_1(\lambda + 1) - \rho^{\lambda+1}\}^{2m-1}}{[(\lambda + 1)^{2m} - \rho^{2m(-\lambda)}\{c_1(\lambda + 1) - \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho + c_4^{\pm}.$$

Set

$$d_4 := \lim_{\alpha \rightarrow \alpha_{10}} \int_0^{\alpha} \frac{\rho^{(2m-1)(-\lambda)}\{c_1(\lambda + 1) - \rho^{\lambda+1}\}^{2m-1}}{[(\lambda + 1)^{2m} - \rho^{2m(-\lambda)}\{c_1(\lambda + 1) - \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\lim_{\alpha \rightarrow 0} u_{\pm}(\alpha) = c_4^{\pm}, \quad \lim_{\alpha \rightarrow \alpha_{10}} u_{\pm}(\alpha) = c_4^{\pm} \pm d_4,$$

$$\lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha) = 0, \quad \lim_{\alpha \rightarrow \alpha_{10}} u'_{\pm}(\alpha) = 0.$$

On the C^2 -smoothness at $\alpha = 0$, we note that

$$u'_{\pm}(\alpha) = \pm \frac{\{c_1(\lambda + 1)\alpha^{(-\lambda)} - \alpha\}^{2m-1}}{[(\lambda + 1)^{2m} - \{c_1(\lambda + 1)\alpha^{(-\lambda)} - \alpha\}^{2m}]^{\frac{2m-1}{2m}}}.$$

Hence, $\lim_{\alpha \rightarrow 0} u_{\pm}''(\alpha)$ exists if and only if $(2m-1)(-\lambda) \geq 1$.

However, by (3.3), (3.4) and that $-1 < \lambda < 0$, we find that k_1 and k_2 are not extended continuously to $\alpha = 0$.

(iii-2) The case where $c_1 = \{(\lambda+1)(-\lambda)^{(-\lambda)}\}^{-1}$. In this case, we have $f(\alpha_{11}) = 0$ and the domain is given by $0 < \alpha < \alpha_{11}$ or $\alpha_{11} < \alpha < \alpha_{10}$.

(iii-2-1) On the domain $0 < \alpha < \alpha_{11}$, we have

$$u_{\pm}(\alpha) = \pm \int_0^{\alpha} \frac{\rho^{(2m-1)(-\lambda)} \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} - \rho^{2m(-\lambda)} \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho + c_4^{\pm}.$$

On the denominator, since $f(\alpha_{11}) = f'(\alpha_{11}) = 0$ and $f''(\alpha_{11}) \neq 0$, we have $f(t) = (\alpha_{11} - t)^2 \tilde{f}(t)$ for a smooth function $\tilde{f}(t)$ with $\tilde{f}(\alpha_{11}) \neq 0$.

Thus, noting that $m \geq 2$, we have

$$\lim_{\alpha \rightarrow \alpha_{11}} \int_0^{\alpha} \frac{\rho^{(2m-1)(-\lambda)} \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} - \rho^{2m(-\lambda)} \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho = \infty,$$

and

$$\lim_{\alpha \rightarrow 0} u_{\pm}(\alpha) = c_4^{\pm}, \quad \lim_{\alpha \rightarrow \alpha_{11}} u_{\pm}(\alpha) = \pm\infty, \quad \lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha) = 0.$$

Correspondingly, we get a rotational surface of C^1 -class.

As in the case (iii-1), $\lim_{\alpha \rightarrow 0} u_{\pm}''(\alpha)$ exists if and only if $(2m-1)(-\lambda) \geq 1$, but k_1 and k_2 are not extended continuously to $\alpha = 0$.

(iii-2-2) On the domain $\alpha_{11} < \alpha < \alpha_{10}$, we have

$$u_{\pm}(\alpha) = \pm \int_{\alpha_{10}}^{\alpha} \frac{\rho^{(2m-1)(-\lambda)} \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} - \rho^{2m(-\lambda)} \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho + c_4^{\pm}.$$

As in the case (iii-2-1), we can see that

$$\lim_{\alpha \rightarrow \alpha_{10}} u_{\pm}(\alpha) = c_4^{\pm}, \quad \lim_{\alpha \rightarrow \alpha_{11}} u_{\pm}(\alpha) = \mp\infty, \quad \lim_{\alpha \rightarrow \alpha_{10}} u'_{\pm}(\alpha) = 0.$$

(iii-3) The case where $c_1 > \{(\lambda+1)(-\lambda)^{(-\lambda)}\}^{-1}$. In this case, we have $f(\alpha_{11}) < 0$. There are two numbers α_{12} and α_{13} such that

$$f(\alpha_{12}) = f(\alpha_{13}) = 0, \quad 0 < \alpha_{12} < \alpha_{11} < \alpha_{13} < \alpha_{10},$$

and the domain is given by $0 < \alpha < \alpha_{12}$ or $\alpha_{13} < \alpha < \alpha_{10}$.

(iii-3-1) On the domain $0 < \alpha < \alpha_{12}$, we have

$$u_{\pm}(\alpha) = \pm \int_{\alpha_{12}}^{\alpha} \frac{\rho^{(2m-1)(-\lambda)} \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} - \rho^{2m(-\lambda)} \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho + c_5.$$

On the denominator, since $f(\alpha_{12}) = 0$ and $f'(\alpha_{12}) \neq 0$, the above integral converges.

Set

$$d_5 := - \lim_{\alpha \rightarrow 0} \int_{\alpha_{12}}^{\alpha} \frac{\rho^{(2m-1)(-\lambda)} \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} - \rho^{2m(-\lambda)} \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\begin{aligned} \lim_{\alpha \rightarrow \alpha_{12}} u_{\pm}(\alpha) &= c_5, & \lim_{\alpha \rightarrow 0} u_{\pm}(\alpha) &= c_5 \mp d_5, \\ \lim_{\alpha \rightarrow \alpha_{12}} u'_{\pm}(\alpha) &= \pm\infty, & \lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha) &= 0. \end{aligned}$$

Correspondingly, we get a closed rotational surface of C^1 -class which is homeomorphic to the 2-sphere.

As in the case (iii-1), $\lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha)$ exists if and only if $(2m-1)(-\lambda) \geq 1$, but k_1 and k_2 are not extended continuously to $\alpha = 0$.

(iii-3-2) On the domain $\alpha_{13} < \alpha < \alpha_{10}$, we have

$$u_{\pm}(\alpha) = \pm \int_{\alpha_{13}}^{\alpha} \frac{\rho^{(2m-1)(-\lambda)} \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} - \rho^{2m(-\lambda)} \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho + c_5.$$

As in the case (iii-3-1), it converges. Set

$$d_6 := \lim_{\alpha \rightarrow \alpha_{10}} \int_{\alpha_{13}}^{\alpha} \frac{\rho^{(2m-1)(-\lambda)} \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} - \rho^{2m(-\lambda)} \{c_1(\lambda+1) - \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\begin{aligned} \lim_{\alpha \rightarrow \alpha_{13}} u_{\pm}(\alpha) &= c_5, & \lim_{\alpha \rightarrow \alpha_{10}} u_{\pm}(\alpha) &= c_5 \pm d_6, \\ \lim_{\alpha \rightarrow \alpha_{13}} u'_{\pm}(\alpha) &= \pm\infty, & \lim_{\alpha \rightarrow \alpha_{10}} u'_{\pm}(\alpha) &= 0. \end{aligned}$$

(iv) The case where $-1 < \lambda < 0$ and $\mu = -1$. We have

$$\begin{aligned} u_{\pm}(\alpha) &= \pm \int \frac{\{c_1^*(\lambda+1) + \alpha^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} \alpha^{2m\lambda} - \{c_1^*(\lambda+1) + \alpha^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\alpha \\ &= \pm \int \frac{\alpha^{(2m-1)(-\lambda)} \{c_1^*(\lambda+1) + \alpha^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} - \alpha^{2m(-\lambda)} \{c_1^*(\lambda+1) + \alpha^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\alpha, \end{aligned}$$

and

$$c_1^*(\lambda+1) + \alpha^{\lambda+1} > 0, \quad \alpha^{(-\lambda)} \{c_1^*(\lambda+1) + \alpha^{\lambda+1}\} < \lambda+1. \quad (6.5)$$

(iv-1) The case where $c_1^* = 0$. As in the case (ii-1), the surface is homothetic to the unit sphere S .

(iv-2) The case where $c_1^* > 0$. The first inequality of (6.5) holds obviously. For the second inequality of (6.5), set

$$f(t) := \lambda+1 - c_1^*(\lambda+1)t^{(-\lambda)} - t, \quad t > 0.$$

Since it is strictly decreasing and

$$\lim_{t \rightarrow 0} f(t) = \lambda + 1 > 0, \quad \lim_{t \rightarrow \infty} f(t) = -\infty,$$

there is a unique number $\alpha_{14} > 0$ such that $f(\alpha_{14}) = 0$.

The domain is given by $0 < \alpha < \alpha_{14}$, and

$$u_{\pm}(\alpha) = \pm \int_{\alpha_{14}}^{\alpha} \frac{\rho^{(2m-1)(-\lambda)} \{c_1^*(\lambda+1) + \rho^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} - \rho^{2m(-\lambda)} \{c_1^*(\lambda+1) + \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho + c_6.$$

On the denominator, since $f(\alpha_{14}) = 0$ and $f'(\alpha_{14}) < 0$, the above integral converges.

Set

$$d_7 := - \lim_{\alpha \rightarrow 0} \int_{\alpha_{14}}^{\alpha} \frac{\rho^{(2m-1)(-\lambda)} \{c_1^*(\lambda+1) + \rho^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} - \rho^{2m(-\lambda)} \{c_1^*(\lambda+1) + \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\begin{aligned} \lim_{\alpha \rightarrow \alpha_{14}} u_{\pm}(\alpha) &= c_6, & \lim_{\alpha \rightarrow 0} u_{\pm}(\alpha) &= c_6 \mp d_7, \\ \lim_{\alpha \rightarrow \alpha_{14}} u'_{\pm}(\alpha) &= \pm\infty, & \lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha) &= 0. \end{aligned}$$

Thus we have a closed rotational surface of C^1 -class which is homeomorphic to the 2-sphere.

As in the case (iii-1), $\lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha)$ exists if and only if $(2m-1)(-\lambda) \geq 1$, but k_1 and k_2 are not extended continuously to $\alpha = 0$.

(iv-3) The case where $c_1^* < 0$. From the first inequality of (6.5), we have

$$\alpha > \{(-c_1^*)(\lambda+1)\}^{\frac{1}{\lambda+1}} =: \alpha_{15}.$$

For the second inequality of (6.5), set

$$f(t) := \lambda + 1 - c_1^*(\lambda+1)t^{(-\lambda)} - t, \quad t > \alpha_{15}.$$

Since

$$f'(t) = (-c_1^*)(-\lambda)(\lambda+1)t^{-(\lambda+1)} - 1 < -1,$$

and

$$\lim_{t \rightarrow \alpha_{15}} f(t) = \lambda + 1 > 0, \quad \lim_{t \rightarrow \infty} f(t) = -\infty,$$

there is a unique number $\alpha_{16} \in (\alpha_{15}, \infty)$ such that $f(\alpha_{16}) = 0$.

The domain is given by $\alpha_{15} < \alpha < \alpha_{16}$, and

$$u_{\pm}(\alpha) = \pm \int_{\alpha_{16}}^{\alpha} \frac{\rho^{(2m-1)(-\lambda)} \{c_1^*(\lambda+1) + \rho^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} - \rho^{2m(-\lambda)} \{c_1^*(\lambda+1) + \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho + c_6.$$

On the denominator, since $f(\alpha_{16}) = 0$ and $f'(\alpha_{16}) < -1$, the above integral converges.

Set

$$d_8 := - \lim_{\alpha \rightarrow \alpha_{15}} \int_{\alpha_{16}}^{\alpha} \frac{\rho^{(2m-1)(-\lambda)} \{c_1^*(\lambda+1) + \rho^{\lambda+1}\}^{2m-1}}{[(\lambda+1)^{2m} - \rho^{2m(-\lambda)} \{c_1^*(\lambda+1) + \rho^{\lambda+1}\}^{2m}]^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\begin{aligned} \lim_{\alpha \rightarrow \alpha_{16}} u_{\pm}(\alpha) &= c_6, & \lim_{\alpha \rightarrow \alpha_{15}} u_{\pm}(\alpha) &= c_6 \mp d_8, \\ \lim_{\alpha \rightarrow \alpha_{16}} u'_{\pm}(\alpha) &= \pm\infty, & \lim_{\alpha \rightarrow \alpha_{15}} u'_{\pm}(\alpha) &= 0. \end{aligned}$$

(v) The case where $\lambda < -1$ and $\mu = 1$. Set $\omega := -(\lambda+1) > 0$. Then

$$\begin{aligned} u_{\pm}(\alpha) &= \pm \int \frac{(c_1\omega + \alpha^{-\omega})^{2m-1}}{\{\omega^{2m}\alpha^{-2m(\omega+1)} - (c_1\omega + \alpha^{-\omega})^{2m}\}^{\frac{2m-1}{2m}}} d\alpha \\ &= \pm \int \frac{\alpha^{2m-1}(c_1\omega\alpha^{\omega} + 1)^{2m-1}}{\{\omega^{2m} - \alpha^{2m}(c_1\omega\alpha^{\omega} + 1)^{2m}\}^{\frac{2m-1}{2m}}} d\alpha, \end{aligned}$$

where

$$c_1\omega\alpha^{\omega} + 1 > 0, \quad \alpha(c_1\omega\alpha^{\omega} + 1) < \omega. \quad (6.6)$$

(v-1) The case where $c_1 = 0$. The surface is homothetic to the unit sphere S .

(v-2) The case where $c_1 > 0$. The first inequality of (6.6) holds obviously. For the second inequality of (6.6), set

$$f(t) := \omega - t(c_1\omega t^{\omega} + 1), \quad t > 0.$$

Since

$$f'(t) = -c_1\omega(\omega+1)t^{\omega} - 1 < -1 \quad \text{for } t > 0,$$

and

$$\lim_{t \rightarrow 0} f(t) = \omega > 0, \quad \lim_{t \rightarrow \infty} f(t) = -\infty,$$

there is a unique number $\alpha_{17} > 0$ such that $f(\alpha_{17}) = 0$.

The domain is given by $0 < \alpha < \alpha_{17}$, and

$$u_{\pm}(\alpha) = \pm \int_{\alpha_{17}}^{\alpha} \frac{\rho^{2m-1}(c_1\omega\rho^{\omega} + 1)^{2m-1}}{\{\omega^{2m} - \rho^{2m}(c_1\omega\rho^{\omega} + 1)^{2m}\}^{\frac{2m-1}{2m}}} d\rho + c_7.$$

On the denominator, we have $f(\alpha_{17}) = 0$ and $f'(\alpha_{17}) < -1$, so that the above integral converges.

Set

$$d_9 := - \lim_{\alpha \rightarrow 0} \int_{\alpha_{17}}^{\alpha} \frac{\rho^{2m-1}(c_1\omega\rho^{\omega} + 1)^{2m-1}}{\{\omega^{2m} - \rho^{2m}(c_1\omega\rho^{\omega} + 1)^{2m}\}^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\lim_{\alpha \rightarrow \alpha_{17}} u_{\pm}(\alpha) = c_7, \quad \lim_{\alpha \rightarrow 0} u_{\pm}(\alpha) = c_7 \mp d_9,$$

$$\lim_{\alpha \rightarrow \alpha_{17}} u'_{\pm}(\alpha) = \pm\infty, \quad \lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha) = 0.$$

Thus we have a closed rotational surface of C^1 -class which is homeomorphic to the 2-sphere.

On the C^2 -smoothness at $\alpha = 0$, we note that

$$u'_{\pm}(\alpha) = \pm \frac{(c_1\omega\alpha^{\omega+1} + \alpha)^{2m-1}}{\{\omega^{2m} - (c_1\omega\alpha^{\omega+1} + \alpha)^{2m}\}^{\frac{2m-1}{2m}}}.$$

In this case, we find that $\lim_{\alpha \rightarrow 0} u''_{\pm}(\alpha)$ exists, and by (3.3) and (3.4), k_1 and k_2 are extended continuously to $\alpha = 0$.

(v-3) The case where $c_1 < 0$. From the first inequality of (6.6), we have

$$0 < \alpha < \frac{1}{\{(-c_1)\omega\}^{1/\omega}} =: \alpha_{18}.$$

For the second inequality of (6.6), set

$$f(t) := \omega - t(c_1\omega t^{\omega} + 1), \quad 0 < t < \alpha_{18}.$$

Then

$$f'(t) = (-c_1)\omega(\omega + 1)t^{\omega} - 1.$$

Set

$$\alpha_{19} := \frac{1}{\{(-c_1)\omega(\omega + 1)\}^{1/\omega}} < \alpha_{18}.$$

We have the following table:

| | | | | | |
|---------|-----|------------|---------------|------------|-----------------|
| t | (0) | \cdots | α_{19} | \cdots | (α_{18}) |
| $f'(t)$ | | $-$ | 0 | $+$ | |
| $f(t)$ | | \searrow | | \nearrow | |

with

$$\lim_{\alpha \rightarrow 0} f(t) = \lim_{\alpha \rightarrow \alpha_{18}} f(t) = \omega$$

and

$$f(\alpha_{19}) = \omega - \frac{\omega}{\omega + 1} \cdot \frac{1}{\{(-c_1)\omega(\omega + 1)\}^{1/\omega}}.$$

(v-3-1) The case where $c_1 < -\frac{1}{\omega(\omega+1)^{\omega+1}}$. In this case, we have $f(\alpha_{19}) > 0$ and the domain is given by $0 < \alpha < \alpha_{18}$. Then

$$u_{\pm}(\alpha) = \pm \int_0^{\alpha} \frac{\rho^{2m-1}(c_1\omega\rho^{\omega} + 1)^{2m-1}}{\{\omega^{2m} - \rho^{2m}(c_1\omega\rho^{\omega} + 1)^{2m}\}^{\frac{2m-1}{2m}}} d\rho + c_8^{\pm}.$$

Set

$$d_{10} := \lim_{\alpha \rightarrow \alpha_{18}} \int_0^{\alpha} \frac{\rho^{2m-1}(c_1\omega\rho^{\omega} + 1)^{2m-1}}{\{\omega^{2m} - \rho^{2m}(c_1\omega\rho^{\omega} + 1)^{2m}\}^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\begin{aligned}\lim_{\alpha \rightarrow 0} u_{\pm}(\alpha) &= c_8^{\pm}, & \lim_{\alpha \rightarrow \alpha_{18}} u_{\pm}(\alpha) &= c_8^{\pm} \pm d_{10}, \\ \lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha) &= 0, & \lim_{\alpha \rightarrow \alpha_{18}} u'_{\pm}(\alpha) &= 0.\end{aligned}$$

As in the case (v-2), $\lim_{\alpha \rightarrow 0} u''_{\pm}(\alpha)$ exists, and k_1 and k_2 are extended continuously to $\alpha = 0$.

(v-3-2) The case where $c_1 = -\frac{1}{\omega(\omega+1)^{\omega+1}}$. In this case, we have $f(\alpha_{19}) = 0$ and the domain is given by $0 < \alpha < \alpha_{19}$ or $\alpha_{19} < \alpha < \alpha_{18}$.

(v-3-2-1) On the domain $0 < \alpha < \alpha_{19}$, we have

$$u_{\pm}(\alpha) = \pm \int_0^{\alpha} \frac{\rho^{2m-1}(c_1\omega\rho^{\omega} + 1)^{2m-1}}{\{\omega^{2m} - \rho^{2m}(c_1\omega\rho^{\omega} + 1)^{2m}\}^{\frac{2m-1}{2m}}} d\rho + c_8^{\pm}.$$

On the denominator, since $f(\alpha_{19}) = f'(\alpha_{19}) = 0$, $f''(\alpha_{19}) \neq 0$ and $m \geq 2$, we have

$$\lim_{\alpha \rightarrow \alpha_{19}} \int_0^{\alpha} \frac{\rho^{2m-1}(c_1\omega\rho^{\omega} + 1)^{2m-1}}{\{\omega^{2m} - \rho^{2m}(c_1\omega\rho^{\omega} + 1)^{2m}\}^{\frac{2m-1}{2m}}} d\rho = \infty.$$

Then

$$\lim_{\alpha \rightarrow 0} u_{\pm}(\alpha) = c_8^{\pm}, \quad \lim_{\alpha \rightarrow \alpha_{19}} u_{\pm}(\alpha) = \pm\infty, \quad \lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha) = 0.$$

Thus we have a rotational surface of C^1 -class.

As in the case (v-2), $\lim_{\alpha \rightarrow 0} u''_{\pm}(\alpha)$ exists, and k_1 and k_2 are extended continuously to $\alpha = 0$.

(v-3-2-2) On the domain $\alpha_{19} < \alpha < \alpha_{18}$, we have

$$u_{\pm}(\alpha) = \pm \int_{\alpha_{18}}^{\alpha} \frac{\rho^{2m-1}(c_1\omega\rho^{\omega} + 1)^{2m-1}}{\{\omega^{2m} - \rho^{2m}(c_1\omega\rho^{\omega} + 1)^{2m}\}^{\frac{2m-1}{2m}}} d\rho + c_8^{\pm}.$$

As in the case (v-3-2-1), we can see that

$$\lim_{\alpha \rightarrow \alpha_{18}} u_{\pm}(\alpha) = c_8^{\pm}, \quad \lim_{\alpha \rightarrow \alpha_{19}} u_{\pm}(\alpha) = \mp\infty, \quad \lim_{\alpha \rightarrow \alpha_{18}} u'_{\pm}(\alpha) = 0.$$

(v-3-3) The case where $-\frac{1}{\omega(\omega+1)^{\omega+1}} < c_1 < 0$. In this case, we have $f(\alpha_{19}) < 0$. So there are two numbers α_{20} and α_{21} such that

$$f(\alpha_{20}) = f(\alpha_{21}) = 0, \quad 0 < \alpha_{20} < \alpha_{19} < \alpha_{21} < \alpha_{18},$$

and the domain is given by $0 < \alpha < \alpha_{20}$ or $\alpha_{21} < \alpha < \alpha_{18}$.

(v-3-3-1) On the domain $0 < \alpha < \alpha_{20}$, we have

$$u_{\pm}(\alpha) = \pm \int_{\alpha_{20}}^{\alpha} \frac{\rho^{2m-1}(c_1\omega\rho^{\omega} + 1)^{2m-1}}{\{\omega^{2m} - \rho^{2m}(c_1\omega\rho^{\omega} + 1)^{2m}\}^{\frac{2m-1}{2m}}} d\rho + c_9.$$

On the denominator, since $f(\alpha_{20}) = 0$ and $f'(\alpha_{20}) \neq 0$, the above integral converges.

Set

$$d_{11} := - \lim_{\alpha \rightarrow 0} \int_{\alpha_{20}}^{\alpha} \frac{\rho^{2m-1} (c_1 \omega \rho^\omega + 1)^{2m-1}}{\{\omega^{2m} - \rho^{2m} (c_1 \omega \rho^\omega + 1)^{2m}\}^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\begin{aligned} \lim_{\alpha \rightarrow \alpha_{20}} u_{\pm}(\alpha) &= c_9, & \lim_{\alpha \rightarrow 0} u_{\pm}(\alpha) &= c_9 \mp d_{11}, \\ \lim_{\alpha \rightarrow \alpha_{20}} u'_{\pm}(\alpha) &= \pm\infty, & \lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha) &= 0. \end{aligned}$$

Thus we have a closed rotational surface of C^1 -class which is homeomorphic to the 2-sphere.

As in the case (v-2), $\lim_{\alpha \rightarrow 0} u'_{\pm}(\alpha)$ exists, and k_1 and k_2 are extended continuously to $\alpha = 0$.

(v-3-2) On the domain $\alpha_{21} < \alpha < \alpha_{18}$, we have

$$u_{\pm}(\alpha) = \pm \int_{\alpha_{21}}^{\alpha} \frac{\rho^{2m-1} (c_1 \omega \rho^\omega + 1)^{2m-1}}{\{\omega^{2m} - \rho^{2m} (c_1 \omega \rho^\omega + 1)^{2m}\}^{\frac{2m-1}{2m}}} d\rho + c_9.$$

As in the case (v-3-1), it converges. Set

$$d_{12} := \lim_{\alpha \rightarrow \alpha_{18}} \int_{\alpha_{21}}^{\alpha} \frac{\rho^{2m-1} (c_1 \omega \rho^\omega + 1)^{2m-1}}{\{\omega^{2m} - \rho^{2m} (c_1 \omega \rho^\omega + 1)^{2m}\}^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\begin{aligned} \lim_{\alpha \rightarrow \alpha_{21}} u_{\pm}(\alpha) &= c_9, & \lim_{\alpha \rightarrow \alpha_{18}} u_{\pm}(\alpha) &= c_9 \pm d_{12}, \\ \lim_{\alpha \rightarrow \alpha_{21}} u'_{\pm}(\alpha) &= \pm\infty, & \lim_{\alpha \rightarrow \alpha_{18}} u'_{\pm}(\alpha) &= 0. \end{aligned}$$

(vi) The case where $\lambda < -1$ and $\mu = -1$. Setting $\omega := -(\lambda + 1) > 0$, we have

$$\begin{aligned} u_{\pm}(\alpha) &= \pm \int \frac{(c_1^* \omega - \alpha^{-\omega})^{2m-1}}{\{\omega^{2m} \alpha^{-2m(\omega+1)} - (c_1^* \omega - \alpha^{-\omega})^{2m}\}^{\frac{2m-1}{2m}}} d\alpha \\ &= \pm \int \frac{\alpha^{2m-1} (c_1^* \omega \alpha^\omega - 1)^{2m-1}}{\{\omega^{2m} - \alpha^{2m} (c_1^* \omega \alpha^\omega - 1)^{2m}\}^{\frac{2m-1}{2m}}} d\alpha, \end{aligned}$$

where

$$c_1^* \omega \alpha^\omega - 1 > 0, \quad \alpha (c_1^* \omega \alpha^\omega - 1) < \omega. \quad (6.7)$$

From the first inequality of (6.7), we have $c_1^* > 0$ and

$$\alpha > \frac{1}{(c_1^* \omega)^{1/\omega}} =: \alpha_{22}.$$

For the second inequality of (6.7), set

$$f(t) := \omega - t(c_1^* \omega t^\omega - 1), \quad t > \alpha_{22}.$$

Since

$$f'(t) = 1 - c_1^* \omega (\omega + 1) t^\omega < -\omega \quad \text{for } t > \alpha_{22},$$

and

$$\lim_{t \rightarrow \alpha_{22}} f(t) = \omega > 0, \quad \lim_{t \rightarrow \infty} f(t) = -\infty,$$

there is a unique number $\alpha_{23} \in (\alpha_{22}, \infty)$ such that $f(\alpha_{23}) = 0$.

The domain is given by $\alpha_{22} < \alpha < \alpha_{23}$, and

$$u_\pm(\alpha) = \pm \int_{\alpha_{23}}^\alpha \frac{\rho^{2m-1} (c_1^* \omega \rho^\omega - 1)^{2m-1}}{\{\omega^{2m} - \rho^{2m} (c_1^* \omega \rho^\omega - 1)^{2m}\}^{\frac{2m-1}{2m}}} d\rho + c_{10}.$$

Since $f(\alpha_{23}) = 0$ and $f'(\alpha_{23}) \neq 0$, this integral converges.

Set

$$d_{13} := - \lim_{\alpha \rightarrow \alpha_{22}} \int_{\alpha_{23}}^\alpha \frac{\rho^{2m-1} (c_1^* \omega \rho^\omega - 1)^{2m-1}}{\{\omega^{2m} - \rho^{2m} (c_1^* \omega \rho^\omega - 1)^{2m}\}^{\frac{2m-1}{2m}}} d\rho.$$

Then

$$\begin{aligned} \lim_{\alpha \rightarrow \alpha_{23}} u_\pm(\alpha) &= c_{10}, & \lim_{\alpha \rightarrow \alpha_{22}} u_\pm(\alpha) &= c_{10} \mp d_{13}, \\ \lim_{\alpha \rightarrow \alpha_{23}} u'_\pm(\alpha) &= \pm\infty, & \lim_{\alpha \rightarrow \alpha_{22}} u'_\pm(\alpha) &= 0. \end{aligned}$$

6.4 Connecting argument for $\lambda \neq -1$

We use the notation $u_{1\pm}, u_{2\pm}, \dots, u_{6\pm}$ to denote the u_\pm in the case (i), (ii), ..., (vi) of Subsection 6.3, respectively.

(C4) Connecting (i) and (ii-3) of Subsection 6.3. Choose

$$c_1^* = -c_1, \quad c_2 = c_3 + d_3 - d_1.$$

Then u_{1+} and u_{2-} are connected at $\alpha = \alpha_4 = \alpha_8$. As in Subsection 6.2, we can see that they are connected C^2 -smoothly, and also k_1 and k_2 are continuous.

Let Γ be the C^2 -curve given by connecting the graphs of u_{1-}, u_{1+}, u_{2-} and u_{2+} . Since Γ has the same derivatives at the end points, it can be extended periodically as a C^2 -curve Γ^* , and we get a periodic rotational surface. If $d_1 = d_3$, then we may have a rotational torus such that $k_1 + \lambda k_2$ is a non-zero constant, where λ is a positive constant.

(C5) Connecting (iii-1) and (iv-3) of Subsection 6.3. Choose

$$c_1^* = -c_1, \quad c_4^+ = c_6 + d_8 - d_4, \quad c_4^- = c_6 - d_8 + d_4.$$

If $(2m-1)(-\lambda) > 1$ and $d_4 \neq d_8$, we get a rotational surface of C^2 -class, but k_1 and k_2 are not extended continuously to $\alpha = 0$.

(C6) Connecting (iii-2-2) and (iv-3) of Subsection 6.3. Choose

$$c_1^* = -c_1, \quad c_4^+ = c_6 + d_8, \quad c_4^- = c_6 - d_8.$$

Then we have a rotational surface of C^2 -class.

(C7) Connecting (iii-3-2) and (iv-3) of Subsection 6.3. Choose

$$c_1^* = -c_1, \quad c_5 = c_6 + d_8 - d_6.$$

In this case, we can get a periodic rotational surface. If $d_6 = d_8$, then we may have a rotational torus such that $k_1 + \lambda k_2$ is a non-zero constant, where λ is a constant with $-1 < \lambda < 0$.

(C8) Connecting (v-3-1) and (vi) of Subsection 6.3. Choose

$$c_1^* = -c_1, \quad c_8^+ = c_{10} + d_{13} - d_{10}, \quad c_8^- = c_{10} - d_{13} + d_{10}.$$

If $d_{10} \neq d_{13}$, then we have a rotational surface of C^2 -class, and k_1 and k_2 are extended continuously to $\alpha = 0$.

(C9) Connecting (v-3-2-2) and (vi) of Subsection 6.3. Choose

$$c_1^* = -c_1, \quad c_8^+ = c_{10} + d_{13}, \quad c_8^- = c_{10} - d_{13}.$$

Then we have a rotational surface of C^2 -class.

(C10) Connecting (v-3-3-2) and (vi) of Subsection 6.3. Choose

$$c_1^* = -c_1, \quad c_9 = c_{10} + d_{13} - d_{12}.$$

In this case, we have a periodic rotational surface. If $d_{12} = d_{13}$, then we may have a rotational torus such that $k_1 + \lambda k_2$ is a non-zero constant, where λ is a constant with $\lambda < -1$.

References

- [1] V. Balestro, H. Martini and R. Teixeira, Surface immersions in normed spaces from the affine point of view, *Geom. Dedicata* 201 (2019), 21–31.
- [2] V. Balestro, H. Martini and R. Teixeira, Differential geometry of immersed surfaces in three-dimensional normed spaces, *Abh. Math. Semin. Univ. Hambg.* 90 (2020), 111–134.
- [3] V. Balestro, H. Martini and R. Teixeira, On curvature of surfaces immersed in normed spaces, *Monatsh. Math.* 192 (2020), 291–309.
- [4] V. Balestro, H. Martini and R. Teixeira, Some topics in differential geometry of normed spaces, *Adv. Geom.* 21 (2021), 109–118.
- [5] J. A. Galvez, A. Martinez and F. Milan, Linear Weingarten surfaces in \mathbb{R}^3 , *Manatsh. Math.* 138 (2003), 133–144.

- [6] R. Lopez, On linear Weingarten surfaces, *Internat. J. Math.* 19 (2008), 439–448.
- [7] R. Lopez and A. Pampano, Classification of rotational surfaces in Euclidean space satisfying a linear relation between their principal curvatures, *Math. Nachr.* 293 (2020), 735–753.
- [8] M. Sakaki, Rotational surfaces in a 3-dimensional normed space, *Beiträge Alg. Geom.* 65 (2024), 23–41.

Graduate School of Science and Technology
Hirosaki University
Hirosaki 036-8561, Japan
E-mail: sakaki@hirosaki-u.ac.jp