ON THE ANTICYCLOTOMIC MAZUR-TATE CONJECTURE FOR ELLIPTIC CURVES WITH SUPERSINGULAR REDUCTION

CHAN-HO KIM

ABSTRACT. In this short note, we study the anticyclotomic analogue of the "weak" main conjecture of Mazur–Tate on Fitting ideals of Selmer groups for elliptic curves with supersingular reduction.

CONTENTS

1.	Introduction	1
2.	Bertolini–Darmon's theta elements and anticyclotomic <i>p</i> -adic <i>L</i> -functions	2
3.	Anticyclotomic Iwasawa theory for elliptic curves with supersingular reduction	4
4.	The proof of Theorem 1.3	6
Re	ferences	7

1. INTRODUCTION

1.1. The statement of the main result. Let E be an elliptic curve of conductor N over \mathbb{Q} and $p \geq 5$ be a prime of supersingular reduction for E, so $a_p(E) = 0$, such that the mod p representation $\overline{\rho}: G_{\mathbb{Q}} = \operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \to \operatorname{GL}_2(\mathbb{F}_p)$ is surjective.

Let K be an imaginary quadratic field with $(D_K, Np) = 1$ where p splits and and write $p = \mathfrak{p} \cdot \overline{\mathfrak{p}}$ in K. Write

$$N = N^+ \cdot N^-$$

where a prime divisor of N^+ splits in K and a prime divisor of N^- is inert in K.

Assumption 1.1 (odd). Assume that N^- is the square-free product of an odd number of primes.

Let K_{∞} be the anticyclotomic \mathbb{Z}_p -extension of K and K_n be the subextension of K in K_{∞} of degree p^n for $n \ge 0$. Let $\Lambda_n = \mathbb{Z}_p[\operatorname{Gal}(K_n/K)] \simeq \mathbb{Z}_p[X]/((1+X)^{p^n}-1)$ be the finite layer Iwasawa algebra. Under Assumption 1.1, denote by

$$\theta_n(E/K) = \sum_{\sigma \in \operatorname{Gal}(K_n/K)} a_\sigma \cdot \sigma \in \Lambda_n$$

Bertolini–Darmon's theta element of E over K_n reviewed in Section 2. Let ι be the involution on Λ_n defined by inverting group-like elements. Then we have $\iota(\theta_n(E/K)) = \sum_{\sigma \in \text{Gal}(K_n/K)} a_{\sigma}$. σ^{-1} and define

$$L_p(E/K_n) = \theta_n(E/K) \cdot \iota(\theta_n(E/K)).$$

Assumption 1.2. We assume the following hypotheses throughout this article:

(ram) K_{∞}/K is totally ramified at every prime lying above p.

(CR) If a prime ℓ divides N^- and $\ell^2 \equiv 1 \pmod{p}$, then $\overline{\rho}$ is ramified at ℓ .

Date: March 5, 2024.

²⁰¹⁰ Mathematics Subject Classification. 11F67, 11G40, 11R23.

Key words and phrases. refined Iwasawa theory, anticyclotomic Iwasawa theory.

The goal of this article is to prove the following anticyclotomic analogue of the "weak" main conjecture of Mazur and Tate [MT87, Conj. 3].

Theorem 1.3. Under (odd), (ram), and (CR), we have

$$L_p(E/K_n) \in \operatorname{Fitt}_{\Lambda_n} \left(\operatorname{Sel}(K_n, E[p^{\infty}])^{\vee} \right).$$

where $(-)^{\vee}$ means the Pontryagin dual.

Let $\chi : \operatorname{Gal}(K_n/K) \to \overline{\mathbb{Q}}_p^{\times}$ be a character and we extend it to Λ_n linearly. Let $I_{\chi} := \operatorname{ker}(\chi : \Lambda_n \to \overline{\mathbb{Q}}_p) \subseteq \Lambda_n$ be the augmentation ideal at χ and $(E(K_n) \otimes \overline{\mathbb{Q}}_p)^{\chi}$ be the χ -isotypic subspace of $E(K_n) \otimes \overline{\mathbb{Q}}_p$. Thanks to [MT87, Prop. 3], we obtain the "weak" vanishing conjecture.

Corollary 1.4. Under (odd), (ram), and (CR), we have

$$L_p(E/K_n) \in I_{\chi}^{r_{\chi}}/I_{\chi}^{r_{\chi}+1}$$

where $r_{\chi} = \dim_{\overline{\mathbb{Q}}_p} (E(K_n) \otimes \overline{\mathbb{Q}}_p)^{\chi}$.

In particular, $L_p(E/K_n)$ encodes an upper bound of $\operatorname{rk}_{\mathbb{Z}} E(K)$. In the ordinary case, Corollary 1.4 can be deduced from the one-sided divisibility of the anticyclotomic main conjecture more directly.

Remark 1.5. In [MT87], Mazur and Tate formulated "Iwasawa theory for elliptic curves over finite abelian extensions" and several related conjectures. This finite layer setting is much more delicate since the linear algebra techniques for the usual Iwasawa theory (over \mathbb{Z}_p -extensions) are not applicable and we cannot ignore "finite errors" in the computation. The formulation of the "main conjecture" also becomes more subtle since the notion of characteristic ideals is not available. The "weak" main conjecture is formulated in terms of Fitting ideals. See [Kur02, Conj. 0.3] for the strong version.

1.2. Acknowledgement. It is a great pleasure to dedicate this paper to Massimo Bertolini on the occasion of his 60th birthday for his great work and wonderful inspiration. In my graduate years, the celebrated article [BD05] by Bertolini and Darmon was one of the very first serious research articles I read. In particular, my thesis [Kim17] can be explained as "a Greenberg–Vatsal type result for the Bertolini–Darmon setting". It is evident that the content of [BD05] became an important part of my research.

I deeply thank Matteo Longo, Marco Seveso, Stefano Vigni, and Rodolfo Venerucci for inviting me for his birthday conference and for giving me a chance to contribute this volume.

Indeed, the topic of this paper was exactly my original thesis problem suggested by Rob Pollack, but I could not find how to attack it at that time. Later, I found a way to resolve it in the middle of writing the paper with Masato Kurihara [KK21]. I deeply appreciate Rob Pollack and Masato Kurihara. This is the sequel to [Kim], which studies the ordinary case.

This research was partially supported by a KIAS Individual Grant (SP054103) via the Center for Mathematical Challenges at Korea Institute for Advanced Study and by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIT) (No. 2018R1C1B6007009).

2. Bertolini–Darmon's theta elements and anticyclotomic p-adic L-functions

We quickly review the construction of Gross points of conductor p^n and signed anticyclotomic *p*-adic *L*-functions. See [CH15, CH18, Kim19] for details. 2.1. Gross points. Let K be the imaginary quadratic field of discriminant $-D_K < 0$. Define

$$\vartheta := \begin{cases} \frac{D_K - \sqrt{-D_K}}{2} & \text{if } 2 \nmid D_K \\ \frac{D_K - 2\sqrt{-D_K}}{4} & \text{if } 2 \mid D_K \end{cases}$$

so that $\mathcal{O}_K = \mathbb{Z} + \mathbb{Z}\vartheta$. Let B_{N^-} be the definite quaternion algebra over \mathbb{Q} of discriminant N^- . Then there exists an embedding of K into B_{N^-} [Vig80]. More explicitly, we choose a K-basis (1, J) of B_{N^-} so that $B_{N^-} = K \oplus K \cdot J$ such that $\beta := J^2 \in \mathbb{Q}^{\times}$ with $\beta < 0, J \cdot t = \overline{t} \cdot J$ for all $t \in K, \beta \in (\mathbb{Z}_q^{\times})^2$ for all $q \mid pN^+$, and $\beta \in \mathbb{Z}_q^{\times}$ for all $q \mid D_K$. Fix a square root $\sqrt{\beta} \in \overline{\mathbb{Q}}$ of β . For a \mathbb{Z} -module A, write $\widehat{A} = A \otimes \widehat{\mathbb{Z}}$. Fix an isomorphism

$$i := \prod i_q : \widehat{B}_{N^-}^{(N^-)} \simeq \mathcal{M}_2(\mathbb{A}^{(N^-\infty)})$$

as follows:

• For each finite place $q \mid N^+ p$, the isomorphism $i_q : B_{N^-,q} \simeq M_2(\mathbb{Q}_q)$ is defined by

$$i_q(\vartheta) = \begin{pmatrix} \operatorname{trd}(\vartheta) & -\operatorname{nrd}(\vartheta) \\ 1 & 0 \end{pmatrix}, \qquad i_q(J) = \sqrt{\beta} \cdot \begin{pmatrix} -1 & \operatorname{trd}(\vartheta) \\ 0 & 1 \end{pmatrix}$$

where trd and nrd are the reduced trace and the reduced norm on B, respectively.

• For each finite place $q \nmid pN^+$, the isomorphism $i_q : B_{N^-,q} \simeq M_2(\mathbb{Q}_q)$ is chosen so that $i_q (\mathcal{O}_K \otimes \mathbb{Z}_q) \subseteq M_2(\mathbb{Z}_q).$

Under the fixed isomorphism *i*, for any rational prime *q*, the local Gross point $\varsigma_q \in B_{N^-,q}^{\times}$ is defined as follows:

•
$$\varsigma_q := 1$$
 in $B_{nN^-,q}^{\times}$ for $q \nmid pN^+$.
• $\varsigma_q := \frac{1}{\sqrt{D_K}} \cdot \begin{pmatrix} \vartheta & \overline{\vartheta} \\ 1 & 1 \end{pmatrix} \in \operatorname{GL}_2(K_{\mathfrak{q}}) = \operatorname{GL}_2(\mathbb{Q}_q)$ for $q \mid N^+$ with $q = \mathfrak{q}\overline{\mathfrak{q}}$ in \mathcal{O}_K .
• $\varsigma_p^{(n)} = \begin{pmatrix} \vartheta & -1 \\ 1 & 0 \end{pmatrix} \cdot \begin{pmatrix} p^n & 0 \\ 0 & 1 \end{pmatrix} \in \operatorname{GL}_2(K_{\mathfrak{p}}) = \operatorname{GL}_2(\mathbb{Q}_p)$ where $p = \mathfrak{p}\overline{\mathfrak{p}}$ splits in K .

By using the fixed embedding of K into B_{N^-} , we define $x_n : \widehat{K}^{\times} \to \widehat{B}_{N^-}^{\times}$ by $x_n(a) = a \cdot \varsigma^{(n)} := a \cdot \left(\varsigma_p^{(n)} \times \prod_{q \neq p} \varsigma_q\right)$. The collection $\left\{x_n(a) : a \in \widehat{K}^{\times}\right\}$ of points is called the **Gross points of conductor** p^n on $\widehat{B}_{N^-}^{\times}$. The fixed embedding $K \hookrightarrow B_{N^-}$ also induces an optimal embedding of $\mathcal{O}_n = \mathbb{Z} + p^n \mathcal{O}_K$ into the Eichler order $B_{N^-} \cap \varsigma^{(n)} \widehat{R}_{N^+}(\varsigma^{(n)})^{-1}$ where R_{N^+} is the Eichler order of level N^+ under the fixed isomorphism i.

2.2. Theta elements and *p*-adic *L*-functions.

2.2.1. Let $\phi_f: B_{N^-}^{\times} \setminus \widehat{B}_{N^+}^{\times} \to \mathbb{C}$ be the Jacquet–Langlands transfer of f. Since $B_{N^-}^{\times} \setminus \widehat{B}_{N^-}^{\times} / \widehat{R}_{N^+}^{\times}$ is a finite set and f is a Hecke eigenform, we are able to and do normalize

$$\phi_f: B_{N^-}^{\times} \backslash \widehat{B}_{N^-}^{\times} / \widehat{R}_{N^+}^{\times} \to \mathbb{Z}_p$$

such that the image of ϕ_f does not lie in $p\mathbb{Z}_p$. This normalization is closely related to the N^- -new congruence ideal [PW11, Kim17, KO23]. Let

$$\widetilde{\theta}_n(E/K) = \sum_{[a]\in\mathcal{G}_n} \phi_f(x_n(a)) \cdot [a] \in \mathbb{Z}_p[\mathcal{G}_n]$$

where $\mathcal{G}_n = K^{\times} \setminus \widehat{\mathcal{K}}^{\times} / \widehat{\mathcal{O}}_n^{\times}$ and [a] is the image of $a \in \widehat{K}^{\times}$ in \mathcal{G}_n . Then **Bertolini–Darmon's** theta element $\theta_n(E/K)$ of E over K_n is defined by the image of $\widetilde{\theta}_n(E/K)$ in Λ_n

$$\mathbb{Z}_p[\mathcal{G}_n] \xrightarrow{} \Lambda_n$$

$$\widetilde{\theta}_n(E/K) \xrightarrow{} \theta_n(E/K)$$

It is known that $\theta_n(E/K)$ interpolates "an half of" $L(E, \chi, 1)$ where χ runs over characters on $\text{Gal}(K_n/K)$ [CH18, Kim19].

2.2.2. Let $\omega_n = \omega_n(X) = (1+X)^{p^n} - 1$ and $\Phi_n(1+X) = \omega_n(X)/\omega_{n-1}(X)$ where Φ_n is the p^n -th cyclotomic polynomial. Fix a generator γ of $\operatorname{Gal}(K_{\infty}/K)$ and take a generator γ_n of $\operatorname{Gal}(K_n/K)$ as the image of γ . Then we have isomorphisms $\Lambda_n \simeq \mathbb{Z}_p[X]/(\omega_n(X))$ and $\Lambda \simeq \mathbb{Z}_p[X]$ by sending the generators to 1+X. Via the latter isomorphism, we also regard $\omega_n \in \Lambda$. Let $\omega_0^{\pm}(X) := X$, $\widetilde{\omega}_0^{\pm}(X) := 1$, and

$$\omega_n^+ = \omega_n^+(X) := X \cdot \prod_{\substack{2 \le m \le n,m: \text{ even}}} \Phi_m(1+X), \qquad \omega_n^- = \omega_n^-(X) := X \cdot \prod_{\substack{1 \le m \le n,m: \text{ odd}}} \Phi_m(1+X),$$
$$\widetilde{\omega}_n^+ = \widetilde{\omega}_n^+(X) := \prod_{\substack{2 \le m \le n,m: \text{ even}}} \Phi_m(1+X), \qquad \widetilde{\omega}_n^- = \widetilde{\omega}_n^-(X) := \prod_{\substack{1 \le m \le n,m: \text{ odd}}} \Phi_m(1+X).$$

Then we have $\omega_n(X) = \omega_n^{\pm}(X) \cdot \widetilde{\omega}_n^{\mp}(X)$, respectively. We also regard ω_n^{\pm} , $\widetilde{\omega}_n^{\pm}$ as elements in Λ_n or Λ .

Proposition 2.1. Let ϵ be the sign of $(-1)^n$. Then:

(1) $\omega_n^{\epsilon} \theta_n(E/K) = 0.$ (2) There exists a unique element $\theta_n^{\epsilon}(E/K)$ in $\Lambda/\omega_n^{\epsilon}\Lambda$ such that

(2.1)
$$\theta_n(E/K) = \widetilde{\omega}_n^{-\epsilon} \cdot \theta_n^{\epsilon}(E/K)$$

Proof. See [DI08, Prop. 2.8].

Thanks to [DI08, Lem. 2.9], $\{(-1)^{n/2} \cdot \theta_n^+(E/K) : n \text{ is even}\}$ and $\{(-1)^{(n+1)/2} \cdot \theta_n^-(E/K) : n \text{ is odd}\}$ form compatible sequences with respect to $\Lambda/\omega_n^+\Lambda$ with even n and $\Lambda/\omega_n^-\Lambda$ with odd n, respectively. The signed *p*-adic *L*-functions for *E* and *K* are defined by

(2.2)

$$L_{p}^{+}(E/K_{\infty}) := \lim_{\substack{n, \text{ even}}} \left((-1)^{n/2} \cdot \theta_{n}^{+}(E/K) \cdot (-1)^{n/2} \cdot \iota(\theta_{n}^{+}(E/K)) \right)$$

$$= \lim_{\substack{n, \text{ even}}} \left(\theta_{n}^{+}(E/K) \cdot \iota(\theta_{n}^{+}(E/K)) \right) \in \Lambda,$$

$$L_{p}^{-}(E/K_{\infty}) := \lim_{\substack{n, \text{ odd}}} \left((-1)^{(n+1)/2} \cdot \theta_{n}^{-}(E/K) \cdot (-1)^{(n+1)/2} \cdot \iota(\theta_{n}^{-}(E/K)) \right)$$

$$= \lim_{\substack{n, \text{ odd}}} \left(\theta_{n}^{-}(E/K) \cdot \iota(\theta_{n}^{-}(E/K)) \right) \in \Lambda.$$

3. Anticyclotomic Iwasawa theory for elliptic curves with supersingular reduction

3.1. Basic objects of \pm -Iwasawa theory. We quickly recall the basic objects of \pm -Iwasawa theory, which is initiated by S. Kobayashi [Kob03] and R. Pollack [Pol03]. See also [IP06].

3.1.1. Local conditions at p. Let E be an elliptic curve over \mathbb{Q} with $a_p(E) = 0$. Since $p = \mathfrak{p} \cdot \overline{\mathfrak{p}}$ in K and the (ram) condition, we are able to write

(3.1)
$$K_{n,p} = K_{n,\mathfrak{p}} \oplus K_{n,\overline{\mathfrak{p}}} \simeq \mathbb{Q}_{n,p} \oplus \mathbb{Q}_{n,p}$$

where $K_{n,p} = K_n \otimes \mathbb{Q}_p$ and $K_{n,v}$ is the completion of K_n at $v \in \{\mathfrak{p}, \overline{\mathfrak{p}}\}$. Then we define

$$E^{+}(K_{n,p}) := \{ P \in E(K_{n,p}) : \operatorname{Tr}_{n/m+1}(P) \in E(K_{m,p}) \text{ for even } m \ (0 \le m < n) \}$$
$$E^{-}(K_{n,p}) := \{ P \in E(K_{n,p}) : \operatorname{Tr}_{n/m+1}(P) \in E(K_{m,p}) \text{ for odd } m \ (0 \le m < n) \}$$

where $\operatorname{Tr}_{n/m+1} : E(K_{n,p}) \to E(K_{m+1,p})$ is the trace map.

3.1.2. The norm subgroups. Let \widehat{E} be the formal group associated to E and \mathfrak{m}_n be the maximal ideal of $\mathbb{Q}_{n,p}$. We define

$$\widehat{E}^{+}(\mathfrak{m}_{n}) := \{ P \in \widehat{E}(\mathfrak{m}_{n}) : \operatorname{Tr}_{n/m+1}(P) \in \widehat{E}(\mathfrak{m}_{m}) \text{ for even } m \ (0 \le m < n) \}$$
$$\widehat{E}^{-}(\mathfrak{m}_{n}) := \{ P \in \widehat{E}(\mathfrak{m}_{n}) : \operatorname{Tr}_{n/m+1}(P) \in \widehat{E}(\mathfrak{m}_{m}) \text{ for odd } m \ (0 \le m < n) \}$$

where $\operatorname{Tr}_{n/m+1}: \widehat{E}(\mathfrak{m}_n) \to \widehat{E}(\mathfrak{m}_{m+1})$ is the trace map.

3.1.3. \pm -Selmer groups. Following [Kim13], we define the \pm -Selmer groups of $E[p^{\infty}]$ over K_n by

$$\operatorname{Sel}^{\pm}(K_n, E[p^{\infty}]) := \ker \left(\operatorname{Sel}(K_n, E[p^{\infty}]) \to \frac{\operatorname{H}^1(K_{n,p}, E[p^{\infty}])}{E^{\pm}(K_{n,p}) \otimes \mathbb{Q}_p / \mathbb{Z}_p} \right)$$

where $Sel(K_n, E[p^{\infty}])$ is the standard Selmer group of $E[p^{\infty}]$ over K_n . The \pm -Selmer groups of $E[p^{\infty}]$ over K_{∞} is defined by

$$\operatorname{Sel}^{\pm}(K_{\infty}, E[p^{\infty}]) := \varinjlim_{n} \operatorname{Sel}^{\pm}(K_{n}, E[p^{\infty}]),$$

respectively.

3.2. Signed Iwasawa theory.

3.2.1. Signed main conjectures. We recall the Euler system divisibility of the signed main conjectures.

Theorem 3.1. Under (odd), (ram), and (CR), we have the following statements:

- (1) $\operatorname{Sel}^{\pm}(K_{\infty}, E[p^{\infty}])$ is Λ -cotorsion. (2) $\left(L_{p}^{\pm}(E/K_{\infty})\right) \subseteq \operatorname{char}_{\Lambda}\left(\operatorname{Sel}^{\pm}(K_{\infty}, E[p^{\infty}])^{\vee}\right).$

Proof. See [Vat02, DI08, PW11, KPW17, BBL].

The signed main conjecture claims that

(3.2)
$$\left(L_p^{\pm}(E/K_{\infty})\right) = \operatorname{char}_{\Lambda}\left(\operatorname{Sel}^{\pm}(K_{\infty}, E[p^{\infty}])^{\vee}\right)$$

3.2.2. Non-existence of proper Λ -submodules of finite index. We recall B.D.Kim's result on the non-existence of proper Λ -submodules of finite index. See [Shi] for the *p*-inert case.

Theorem 3.2. If $\operatorname{Sel}^{\pm}(K_{\infty}, E[p^{\infty}])$ is Λ -cotorsion, then $\operatorname{Sel}^{\pm}(K_{\infty}, E[p^{\infty}])$ has no proper Λ submodule of finite index, respectively.

Proof. See [Kim13, Thm. 1.1].

Corollary 3.3. Under (odd), (ram), and (CR), $\operatorname{Sel}^{\pm}(K_{\infty}, E[p^{\infty}])$ has no proper Λ -submodule of finite index; thus,

$$\operatorname{char}_{\Lambda}\operatorname{Sel}^{\pm}(K_{\infty}, E[p^{\infty}]) = \operatorname{Fitt}_{\Lambda}\operatorname{Sel}^{\pm}(K_{\infty}, E[p^{\infty}]).$$

Proof. It follows from Theorems 3.1 and 3.2.

3.2.3. Signed control theorems. We recall the signed version of the control theorem.

Theorem 3.4. The restriction map

$$\operatorname{Sel}^{\pm}(K_n, E[p^{\infty}])[\omega_n^{\pm}] \to \operatorname{Sel}^{\pm}(K_{\infty}, E[p^{\infty}])[\omega_n^{\pm}]$$

is injective with the finite cokernel whose size is bounded independently of n. If we further assume that $p \nmid \operatorname{Tam}(E)$, then the restriction map is an isomorphism.

Proof. The $a_p(E) = 0$ condition ensures that E(K)[p] is trivial, and it implies that the restriction map is injective as in [Kob03, Lem. 9.1]. The failure of the surjectivity comes only from prime-to-p local conditions; thus, the situation coincides with the ordinary case. The $p \nmid \operatorname{Tam}(E)$ condition ensures that the failure vanishes. See [Kob03, Thm. 9.3], [Gre99, Prop. 3.8], and [IP06, Thm. 6.8] for detail.

3.3. The consequence.

Corollary 3.5. Under (odd), (ram), and (CR), we have

$$\left(\widetilde{\omega}_n^{\mp} \cdot L_p^{\pm}(E/K_{\infty}) \pmod{\omega_n}\right) \subseteq \left(\widetilde{\omega}_n^{\mp}\right) \cdot \operatorname{Fitt}_{\Lambda_n}\left(\operatorname{Sel}^{\pm}(K_n, E[p^{\infty}])^{\vee}\right)$$

in Λ_n , respectively.

Proof. By applying Theorem 3.1 and Corollary 3.3, we have

$$\left(L_p^{\pm}(E/K_{\infty})\right) \subseteq \operatorname{Fitt}_{\Lambda}\left(\operatorname{Sel}^{\pm}(K_{\infty}, E[p^{\infty}])^{\vee}\right).$$

Taking the quotient by ω_n^{\pm} , we obtain

$$\left(L_p^{\pm}(E/K_{\infty}) \pmod{\omega_n^{\pm}}\right) \subseteq \operatorname{Fitt}_{\Lambda_n/\omega_n^{\pm}}\left(\left(\operatorname{Sel}^{\pm}(K_{\infty}, E[p^{\infty}])[\omega_n^{\pm}]\right)^{\vee}\right)$$

in Λ_n/ω_n^{\pm} , respectively. By the signed control theorem (Theorem 3.4), we obtain

$$\left(L_p^{\pm}(E/K_{\infty}) \pmod{\omega_n^{\pm}}\right) \subseteq \operatorname{Fitt}_{\Lambda_n/\omega_n^{\pm}}\left(\left(\operatorname{Sel}^{\pm}(K_n, E[p^{\infty}])[\omega_n^{\pm}]\right)^{\vee}\right)$$

in Λ_n/ω_n^{\pm} , respectively. Since Fitting ideals are compatible with base change, we have the equality

$$\operatorname{Fitt}_{\Lambda_n/\omega_n^{\pm}}\left(\left(\operatorname{Sel}^{\pm}(K_n, E[p^{\infty}])[\omega_n^{\pm}]\right)^{\vee}\right) = \frac{\operatorname{Fitt}_{\Lambda_n}\left(\operatorname{Sel}^{\pm}(K_n, E[p^{\infty}])^{\vee}\right) + (\omega_n^{\pm})}{(\omega_n^{\pm})}$$

in Λ_n/ω_n^{\pm} . Thus, we have inclusions

$$\left(L_p^{\pm}(E/K_{\infty}) \pmod{\omega_n}\right) + \left(\omega_n^{\pm}\right) \subseteq \operatorname{Fitt}_{\Lambda_n}\left(\operatorname{Sel}^{\pm}(K_n, E[p^{\infty}])^{\vee}\right) + \left(\omega_n^{\pm}\right)$$

in Λ_n , respectively. Multiplying $\widetilde{\omega}_n^{\mp}$, the conclusion immediately follows.

Remark 3.6. If we further assume $p \nmid \text{Tam}(E)$ and the \pm -main conjectures (3.2), then the inclusion in Corollary 3.5 becomes an equality.

4. The proof of Theorem 1.3

By using the global Poitou–Tate duality, consider the exact sequence of Λ_n -modules (cf. [KO18, (4.2)])

$$\left(\frac{E(K_{n,p})\otimes\mathbb{Q}_p/\mathbb{Z}_p}{E^{\pm}(K_{n,p})\otimes\mathbb{Q}_p/\mathbb{Z}_p}\right)^{\vee} \xrightarrow{\iota^{\pm}} \operatorname{Sel}(K_n, E[p^{\infty}])^{\vee} \longrightarrow \operatorname{Sel}^{\pm}(K_n, E[p^{\infty}])^{\vee} \longrightarrow 0.$$

Then we have

$$\operatorname{Fitt}_{\Lambda_n}\left(\left(\frac{E(K_{n,p})\otimes\mathbb{Q}_p/\mathbb{Z}_p}{E^{\pm}(K_{n,p})\otimes\mathbb{Q}_p/\mathbb{Z}_p}\right)^{\vee}/\operatorname{ker}(\iota^{\pm})\right)\cdot\operatorname{Fitt}_{\Lambda_n}\left(\operatorname{Sel}^{\pm}(K_n, E[p^{\infty}])^{\vee}\right)\subseteq\operatorname{Fitt}_{\Lambda_n}\left(\operatorname{Sel}(K_n, E[p^{\infty}])^{\vee}\right)$$

thanks to the behavior of Fitting ideals in the short exact sequence. Due to the behavior of Fitting ideals under the quotient map, we also have

$$\operatorname{Fitt}_{\Lambda_n}\left(\left(\frac{E(K_{n,p})\otimes\mathbb{Q}_p/\mathbb{Z}_p}{E^{\pm}(K_{n,p})\otimes\mathbb{Q}_p/\mathbb{Z}_p}\right)^{\vee}\right)\subseteq\operatorname{Fitt}_{\Lambda_n}\left(\left(\frac{E(K_{n,p})\otimes\mathbb{Q}_p/\mathbb{Z}_p}{E^{\pm}(K_{n,p})\otimes\mathbb{Q}_p/\mathbb{Z}_p}\right)^{\vee}/\operatorname{ker}(\iota^{\pm})\right).$$

Applying [KO18, Lem. 3.14] with (3.1), we observe that

$$\left(\frac{E(K_{n,p}) \otimes \mathbb{Q}_p / \mathbb{Z}_p}{E^{\pm}(K_{n,p}) \otimes \mathbb{Q}_p / \mathbb{Z}_p} \right)^{\vee} \simeq \left(\frac{\widehat{E}(\mathfrak{m}_n) \otimes \mathbb{Q}_p / \mathbb{Z}_p}{\widehat{E}^{\pm}(\mathfrak{m}_n) \otimes \mathbb{Q}_p / \mathbb{Z}_p} \right)^{\vee} \oplus \left(\frac{\widehat{E}(\mathfrak{m}_n) \otimes \mathbb{Q}_p / \mathbb{Z}_p}{\widehat{E}^{\pm}(\mathfrak{m}_n) \otimes \mathbb{Q}_p / \mathbb{Z}_p} \right)^{\vee} \\ \simeq \left(\frac{\widehat{E}(\mathfrak{m}_n)}{\widehat{E}^{\pm}(\mathfrak{m}_n)} \otimes \mathbb{Q}_p / \mathbb{Z}_p \right)^{\vee} \oplus \left(\frac{\widehat{E}(\mathfrak{m}_n)}{\widehat{E}^{\pm}(\mathfrak{m}_n)} \otimes \mathbb{Q}_p / \mathbb{Z}_p \right)^{\vee}.$$

Due to [IP06, Prop. 4.11], we have the following exact sequence

where f is the diagonal embedding and $g: (a,b) \mapsto a-b$. Note that $\widetilde{\omega}_n^+ \widetilde{\omega}_n^- \Lambda_n \simeq \Lambda_n / X \Lambda_n \simeq \mathbb{Z}_p$. Since $\widehat{E}(\mathfrak{m}_n) / \widehat{E}^{\pm}(\mathfrak{m}_n) \simeq (\widetilde{\omega}_n^+, \widetilde{\omega}_n^-) \Lambda_n / \widetilde{\omega}_n^{\mp} \Lambda_n$, we have

$$\operatorname{Fitt}_{\Lambda_n}\left(\left(\frac{(\widetilde{\omega}_n^+,\widetilde{\omega}_n^-)\Lambda_n}{\widetilde{\omega}_n^+\Lambda_n}\otimes \mathbb{Q}_p/\mathbb{Z}_p\right)^\vee\right) = \operatorname{Fitt}_{\Lambda_n}\left(\left(\frac{\widehat{E}(\mathfrak{m}_n)}{\widehat{E}^{\pm}(\mathfrak{m}_n)}\otimes \mathbb{Q}_p/\mathbb{Z}_p\right)^\vee\right).$$

Proposition 4.1.

$$\operatorname{Fitt}_{\Lambda_n}\left(\left(\frac{(\widetilde{\omega}_n^+,\widetilde{\omega}_n^-)\Lambda_n}{\widetilde{\omega}_n^{\mp}\Lambda_n}\otimes \mathbb{Q}_p/\mathbb{Z}_p\right)^{\vee}\right)=\widetilde{\omega}_n^{\mp}\Lambda_n,$$

respectively.

Proof. See [KK21, Prop. 4.1].

To sum up, we have

$$\begin{aligned} & \left(\widetilde{\omega}_{n}^{\mp}\right)^{2} \cdot \operatorname{Fitt}_{\Lambda_{n}} \left(\operatorname{Sel}^{\pm}(K_{n}, E[p^{\infty}])^{\vee}\right) \\ &= \operatorname{Fitt}_{\Lambda_{n}} \left(\left(\frac{\left(\widetilde{\omega}_{n}^{+}, \widetilde{\omega}_{n}^{-}\right) \Lambda_{n}}{\widetilde{\omega}_{n}^{\mp} \Lambda_{n}} \otimes \mathbb{Q}_{p} / \mathbb{Z}_{p} \right)^{\vee} \right)^{2} \cdot \operatorname{Fitt}_{\Lambda_{n}} \left(\operatorname{Sel}^{\pm}(K_{n}, E[p^{\infty}])^{\vee} \right) \\ &= \operatorname{Fitt}_{\Lambda_{n}} \left(\left(\widehat{E}(\mathfrak{m}_{n}) / \widehat{E}^{\pm}(\mathfrak{m}_{n}) \otimes \mathbb{Q}_{p} / \mathbb{Z}_{p} \right)^{\vee} \right)^{2} \cdot \operatorname{Fitt}_{\Lambda_{n}} \left(\operatorname{Sel}^{\pm}(K_{n}, E[p^{\infty}])^{\vee} \right) \\ &\subseteq \operatorname{Fitt}_{\Lambda_{n}} \left(\left(\frac{E(K_{n,p}) \otimes \mathbb{Q}_{p} / \mathbb{Z}_{p}}{E^{\pm}(K_{n,p}) \otimes \mathbb{Q}_{p} / \mathbb{Z}_{p}} \right)^{\vee} / \operatorname{ker}(\iota^{\pm}) \right) \cdot \operatorname{Fitt}_{\Lambda_{n}} \left(\operatorname{Sel}^{\pm}(K_{n}, E[p^{\infty}])^{\vee} \right) \\ &\subseteq \operatorname{Fitt}_{\Lambda_{n}} \left(\operatorname{Sel}(K_{n}, E[p^{\infty}])^{\vee} \right). \end{aligned}$$

By Corollary 3.5 with the multiplication by $\widetilde{\omega}_n^{\mp},$ we have

$$\left(\left(\widetilde{\omega}_n^{\mp}\right)^2 \cdot L_p^{\pm}(E/K_{\infty}) \pmod{\omega_n}\right) \subseteq \operatorname{Fitt}_{\Lambda_n}\left(\operatorname{Sel}(K_n, E[p^{\infty}])^{\vee}\right).$$

By using (2.1), (2.2), and the functional equation for Bertolini–Darmon's theta elements (see [BD96, Prop. 2.13] and [BD05, Lem. 1.5]), we have

$$\left(\left(\widetilde{\omega}_{n}^{\mp}\right)^{2} \cdot L_{p}^{\pm}(E/K_{\infty}) \pmod{\omega_{n}}\right) = \left(\left(\widetilde{\omega}_{n}^{\mp}\right)^{2} \cdot \theta^{\pm}(E/K_{n}) \cdot \iota(\theta^{\pm}(E/K_{n}))\right)$$
$$= \left(\left(\widetilde{\omega}_{n}^{\mp}\right)^{2} \cdot \theta^{\pm}(E/K_{n}) \cdot \theta^{\pm}(E/K_{n})\right)$$
$$= \left(\theta(E/K_{n}) \cdot \theta(E/K_{n})\right)$$
$$= \left(\theta(E/K_{n}) \cdot \iota(\theta(E/K_{n}))\right)$$
$$= \left(L_{p}(E/K_{n})\right).$$

Thus, Theorem 1.3 follows.

References

- [BBL] A. A. Burungale, K. Büyükboduk, and A. Lei, Anticyclotomic Iwasawa theory of abelian arieties of GL₂-type at non-ordinary primes, preprint, arXiv:2211.03722.
- [BD96] M. Bertolini and H. Darmon, Heegner points on Mumford-Tate curves, Invent. Math. 126 (1996), no. 3, 413–456.

- [BD05] _____, Iwasawa's main conjectures for elliptic curves over anticyclotomic \mathbb{Z}_p -extensions, Ann. of Math. (2) **162** (2005), no. 1, 1–64.
- [CH15] M. Chida and M.-L. Hsieh, On the anticyclotomic Iwasawa main conjecture for modular forms, Compos. Math. 151 (2015), no. 5, 863–897.
- [CH18] _____, Special values of anticyclotomic L-functions for modular forms, J. Reine Angew. Math. 741 (2018), 87–131.
- [DI08] H. Darmon and A. Iovita, The anticyclotomic main conjecture for elliptic curves at supersingular primes, J. Inst. Math. Jussieu 7 (2008), no. 2, 291–325.
- [Gre99] R. Greenberg, Iwasawa theory for elliptic curves, Arithmetic theory of elliptic curves (Cetraro, 1997) (Berlin) (C. Viola, ed.), Lecture Notes in Math., vol. 1716, Centro Internazionale Matematico Estivo (C.I.M.E.), Florence, Springer-Verlag, 1999, Lectures from the 3rd C.I.M.E. Session held in Cetraro, July 12--19, 1997, pp. 51—-144.
- [IP06] A. Iovita and R. Pollack, Iwasawa theory of elliptic curves at supersingular primes over \mathbb{Z}_p -extensions of number fields, J. Reine Angew. Math. **598** (2006), 71–103.
- [Kim] C.-H. Kim, An anticyclotomic Mazur–Tate conjecture for modular forms, submitted, arXiv:1612.03743.
- [Kim13] B. D. Kim, The plus/minus Selmer groups for supersingular primes, J. Aust. Math. Soc. 95 (2013), no. 2, 189–200.
- [Kim17] C.-H. Kim, Anticyclotomic Iwasawa invariants and congruences of modular forms, Asian J. Math. 21 (2017), no. 3, 499–530.
- [Kim19] _____, Overconvergent quaternionic forms and anticyclotomic p-adic L-functions, Publ. Mat. 63 (2019), no. 2, 727–767.
- [KK21] C.-H. Kim and M. Kurihara, On the refined conjectures on Fitting ideals of Selmer groups of elliptic curves with supersingular reduction, Int. Math. Res. Not. IMRN 2021 (2021), no. 14, 10559–10599.
- [KO18] T. Kitajima and R. Otsuki, On the plus and the minus Selmer groups for elliptic curves at supersingular primes, Tokyo J. Math. 41 (2018), no. 1, 273–303.
- [KO23] C.-H. Kim and K. Ota, On the quantitative variation of congruence ideals and integral periods of modular forms, Res. Math. Sci. 10 (2023), no. 22, 34 pages.
- [Kob03] S. Kobayashi, Iwasawa theory for elliptic curves at supersingular primes, Invent. Math. 152 (2003), no. 1, 1–36.
- [KPW17] C.-H. Kim, R. Pollack, and T. Weston, On the freeness of anticyclotomic Selmer groups of modular forms, Int. J. Number Theory 13 (2017), no. 6, 1443–1455.
- [Kur02] M. Kurihara, On the Tate Shafarevich groups over cyclotomic fields of an elliptic curve with supersingular reduction I, Invent. Math. 149 (2002), 195–224.
- [MT87] B. Mazur and J. Tate, Refined conjectures of the "Birch and Swinnerton-Dyer type", Duke Math. J. 54 (1987), no. 2, 711–750.
- [Pol03] R. Pollack, On the p-adic L-function of a modular form at a supersingular prime, Duke Math. J. 118 (2003), no. 3, 523–558.
- [PW11] R. Pollack and T. Weston, On anticyclotomic μ-invariants of modular forms, Compos. Math. 147 (2011), 1353–1381.
- [Shi] R. Shii, On non-trivial Λ -submodules with finite index of the plus/minus Selmer group over anticyclotomic \mathbb{Z}_p -extension at inert primes, preprint, arXiv:2308.16384.
- [Vat02] V. Vatsal, Uniform distribution of Heegner points, Invent. Math. 148 (2002), no. 1, 1–48.
- [Vig80] M.-F. Vignéras, Arithmétique des algèbres de quaternions, Lecture Notes in Math., vol. 800, Springer, 1980.

JUNE E HUH CENTER FOR MATHEMATICAL CHALLENGES, KOREA INSTITUTE FOR ADVANCED STUDY, 85 HOEGIRO, DONGDAEMUN-GU, SEOUL 02455, REPUBLIC OF KOREA

Email address: chanho.math@gmail.com