# **REVERSIBILITY IN THE SEIFERT-FIBERED SPACES**

ANUSHREE DAS AND DEBATTAM DAS

ABSTRACT. An element a in a group  $\Gamma$  is called *reversible* if there exists  $g \in \Gamma$  such that  $gag^{-1} = a^{-1}$ . The reversible elements are also known as 'real elements' or 'reciprocal elements' in literature. In this paper, we classify the reversible elements in Fuchsian groups, and use this classification to find all reversible elements in a Seifert-fibered group. In the last section we apply this classification to the braid groups, particularly to the braid group on 3 strands.

### 1. INTRODUCTION

An element a in a group  $\Gamma$  is said to be 'reversible' if it is conjugate to its own inverse, and the conjugating elements are called 'reverser' or 'reversing symmetry'. If the conjugating element is also an involution, then it is called 'strongly reversible' element. It has also been called by other terms like 'reciprocals' or 'real elements' in other literature, e.g., [BR], [La] etc. In this paper our discussion is limited to some properties of reversible elements, and a discussion of reversible elements in some discrete geometric groups. Reversibility has its own geometric motivations and has been discussed widely in literature. For instance, in a Fuchsian group, the collection of conjugacy classes of reversible elements have a 1-1 correspondence to the conjugacy classes of infinite dihedral groups contained in that Fuchsian group. But in general, supposing G is a 3manifold group (the fundamental group of some 3-manifold), if G is torsion free then a reversible element q in G is represented as a loop which is homotopic to its own inverse (i.e., it will form an embedded Klein bottle with the reversing symmetry). So, it can be considered that finding a reversible element with a reverser in a 3-manifold is equivalent to finding an embedded Klein bottle up to homotopy inside the manifold. In this paper we are specially interested in Seifert fibered manifolds since the fundamental group of a Seifert fibered manifold is a universal extension of Fuchsian groups or non-orientable 2-orbifold groups.

A Seifert fibered manifold is a compact 3-manifold which can be seen as a  $S^1$  bundle over a 2-dimensional orbifold as the base space, such that every fiber has a solid fibered torus neighbourhood. These spaces were first studied by Herbert Seifert in his doctoral thesis [ST], and they form an important example of 3-manifolds. In fact, all the the compact orientable 3-manifolds in 6 of the 8 possible 3-manifold geometric structures can be seen as Seifert fibered spaces. If we consider the space of fibers of M with a quotient topology, we get a resulting surface called the orbit surface of M. We denote the orbit surface by G. The orbit surface can also be seen as a quotient of the base space of M. There is a map  $\phi : \pi_1(G) \to \{-1, 1\}$  called the classifying

<sup>2020</sup> Mathematics Subject Classification. Primary 57K; Secondary 20H10, 20H05.

Key words and phrases. Reversible elements, Fuchsian groups, surface groups, Seifert fibered space, Braid groups.

homomorphism. A loop  $\alpha$  on G has image  $\phi(\alpha) = 1$  if the fiber above  $\alpha$  is mapped to itself in an orientation preserving way as we follow the loop  $\alpha$ , and has image  $\phi(\alpha) = -1$  otherwise. The classifying homomorphism plays a crucial role in determining the fiber type of a Seifert fibered space.

Following Seifert's work and Brin's note in [Br], the closed Seifert fibered spaces can be completely classified based on certain symbols. If M is a closed and oriented Seifert fibered space, then M is uniquely determined by a symbol

$$(O, o, g|b, \mu_1, \beta_1, \ldots, \mu_n, \beta_n)$$

or

$$(O, n, k | b, \mu_1, \beta_1, \ldots, \mu_n, \beta_n).$$

Here O refers to the orientability of M, o and n refer to the orientability or non-orientability of the base space respectively, g is the genus of the orientable base space, and k is the number of crosscaps of the non-orientable base space. b is an integer depending on the fibration of the space, and the  $\mu_i$  and  $\beta_i$  depend on the fiber type of the  $i^{th}$  essential fiber. If M is non-orientable, then there are 4 possible symbols associated:

$$(N, o, g|b, \mu_1, \beta_1, \dots, \mu_n, \beta_n)$$

if the base space is orientable, and otherwise, one of the following 3 types based on the classifying homomorphism of the base space:

$$(N, n, I, k | b, \mu_1, \beta_1, \dots, \mu_n, \beta_n);$$
  
 $(O, n, II, k | b, \mu_1, \beta_1, \dots, \mu_n, \beta_n);$   
 $(O, n, III, k | b, \mu_1, \beta_1, \dots, \mu_n, \beta_n).$ 

Given any such Seifert fibered space, this paper is concerned with checking the reversibility of the elements of  $\pi_1(M)$ , the fundamental group of M. The fundamental group of M depends on the orientability of the orbit surface G. If G is orientable, it can be shown that the fundamental group of M is of the form

(1.1) 
$$\pi_1(M) = \langle a_i, b_i, c_i, d_i, h | a_i h a_i^{-1} = h^{\phi(a_i)},$$

(1.2) 
$$b_i h b_i^{-1} = h^{\phi(b_i)}, c_i h c_i^{-1} = h, d_i h d_i^{-1} = h^{\phi(d_i)}, c_i h c_i^{-1} =$$

(1.3) 
$$c_i^{\mu_i} = h_i^\beta, \Pi[a_i, b_i] \Pi c_i \Pi d_i h^b = 1 \rangle$$

If G is not orientable,  $\pi_1(M)$  has the presentation

(1.4) 
$$\pi_1(M) = \langle x_i, c_i, d_i, h | x_i h x_i^{-1} = h^{\phi(x_i)}, c_i h c_i^{-1} = h, d_i h d_i^{-1} = h^{\phi(d_i)},$$

(1.5) 
$$c_i^{\mu_i} = h^{\beta_i}, \Pi x_i^2 \Pi c_i \Pi d_i h^b = 1 \rangle.$$

In both cases the  $\mu_i$ ,  $\beta_i$  and b are the same as used in the representation of M, and h represents an ordinary fiber. The  $a_i$  and  $b_i$  are the generators of the handlebodies of the orbit surface, the  $d_i$ represent the boundaries of the orbit surface, the  $x_i$  generate the crosscaps in the orbit surface, and the  $c_i$  are the generators of the essential fibers.  $\phi$  is the classifying homomorphism.  $\langle h \rangle$  is a normal cyclic subgroup of  $\pi_1(M)$ , and so we can consider the quotient group  $\pi_1(M)/\langle h \rangle$ . If the orbit surface G is orientable, this quotient is a Fuchsian group which can be written as

$$\pi_1(M)/\langle h\rangle = \langle a_i, b_i, c_i, d_i | c_i^{\mu_i} = 1, \Pi[a_i, b_i] \Pi c_i \Pi d_i = 1 \rangle.$$

If G is non-orientable, then

$$\pi_1(M)/\langle h \rangle = \langle x_i, c_i, d_i | c_i^{\mu_i} = 1, \Pi x_i^2 \Pi c_i \Pi d_i = 1 \rangle$$

From the representation of  $\pi_1(M)$ , we can see that the reversibility of the element h is dependent on the classifying homomorphism  $\phi$ . If there is any generator z of the form  $a_i, b_i, x_i$ , or  $d_i$  such that  $\phi(z) = -1$ , then the relations in the definition of  $\pi_1(M)$  give that h is reversible, i.e.,  $zhz^{-1} = h^{\phi(z)} = h^{-1}$ . Our aim is to classify all the reversible elements of  $\pi_1(M)$ . The main result of our paper is:

**Theorem 1.1.** If M has an orientable orbit surface and  $\phi$  is nontrivial, the reversible elements of  $\pi_1(M)$  are the elements conjugate to h and  $c_i^{\mu_i}$  for all i and the conjugates of the elements of either of the form  $c_i^{\frac{\mu_i}{2}}kc_j^{\frac{\mu_j}{2}}k^{-1}$  with  $\phi(k) = -1$  or of the form  $c_i^{\frac{\mu_i}{2}}kc_j^{-\frac{\mu_j}{2}}k^{-1}$  with  $\phi(k) = 1$ , where  $\mu_i$  and  $\mu_j$  are even, and k is any element in  $\pi_1(M)$ ,  $\beta_i = \beta_j$ .  $\pi_1(M)$  has reversible elements as the elements conjugate to  $c_i^{\frac{\mu_i}{2}}kc_j^{-\frac{\mu_j}{2}}k^{-1}$  if  $\phi$  is trivial.

We also similarly classify the reversible elements in case of Seifert fibered spaces with nonorientable orbit surfaces.

In Section 2, we discuss about the reversible elements of fundamental groups of two-dimensional hyperbolic orbifolds, and we classify the reversible elements of such groups. This classification is then applied to the fundamental groups of any Seifert-fibered space in Section 3. Section 4 then discusses an application of the results of Section 3 on the fundamental group of  $B_3$ , the braid group on 3 strands. The possibility of using the same methods for any  $B_n/\langle h \rangle$  is also discussed there.

After completion of the first draft, the authors were informed by Prof. Masakazu Teragaito regarding the article [HMT] written by Himeno, Motegi, and Teragaito, where they classify 3manifolds whose fundamental groups admit generalised torsion elements of order two, and further classify such elements. Their proof is topological and completely different from the proof in this paper. The two articles were written completely independently and at around the same time. The proof in [HMT] crucially uses important results by Jaco-Shalen and Hass, while the techniques in this article are simpler by comparison.

Acknowledgment. The authors would like to acknowledge and thank Prof. Masakazu Teragaito for his helpful comments and for pointing out an error in the first draft. The authors further thank Prof. Krishnendu Gongopadhyay and Dr. Soma Maity for helpful feedback. A. Das acknowledges full support from an NBHM research fellowship during this work. D. Das acknowledges full support from a UGC-NFSC research fellowship during this work.

#### 2. Reversiblity in Fundamental groups of the surfaces

2.1. Reversible classes of Fuchsian group. In [DG2, Remark 1], we have seen the classifications of the z-classes for hyperbolic elements. In the same way, we can see the classifications of reciprocal elements for any finite type Fuchsian groups. We have discussed the reciprocal elements of Hecke groups in [DG1], which is a particular class of Fuchsian groups.

From now on, the hyperbolic axis for an hyperbolic element h is written as  $I_h$ .

**Theorem 2.1.** Let  $\Gamma$  be an arbitrary finitely generated Fuchsian group. Let  $g \in \Gamma$ . Then the following are equivalent.

- (1) g is reciprocal.
- (2) g is either an involution or conjugate to a hyperbolic element h such that the hyperbolic axis of h passes through some of the even order vertices of the fundamental domain of the group  $\Gamma$ .

To prove this theorem, we first need to prove the following lemmas.

**Lemma 2.2.** If a hyperbolic element h in the Fuchsian group  $\Gamma$  is reciprocal, then the fixed point of the reciprocator of the reciprocal elements lies on the hyperbolic axis of h, i.e., on  $I_h$ .

**Proof.**  $\Gamma$  is in PSL(2,  $\mathbb{R}$ ), and so we will conjugate  $\Gamma$  by a Möbius map f such that in the new group after conjugation, h will be a hyperbolic element in the Poincaré disk where the hyperbolic axis of h passes through (0,0). Now, the reciprocator for h is g, which is an involution having fixed point z. So, let us suppose z does not lie on  $I_h$  and  $I_h$  is a straight line passing through the origin. Then, z lies one side of the axis. In our earlier work [DG1, Lemma 2.2] we have shown that in the Fuchsian group, the reciprocators of a reciprocal element fix the hyperbolic axis of that particular hyperbolic element and interchange the fixed points of hyperbolic axis. Now, we apply the map g to the disk. This map is a  $\pi$ -rotation along  $I_h$  of the disk, and therefore the point z must change its position. This is contradiction. So, the fixed point of g must lie on  $I_h$ .

**Lemma 2.3.** A hyperbolic element g in the Fuchsian group  $\Gamma$  is reciprocal if and only if it passes through some of the even-order vertices of the tessellation which is done by the fundamental domain of  $\Gamma$ .

**Proof.** We first show that if g is reciprocal, then  $I_g$  passes through at least one even order vertex in the fundamental domain. First, the fixed points of the reciprocator of g will always be a vertex of the domain. From [DG2, Theorem 3], we obtain that  $I_g$  passing through a fixed domain represents a z-class of that Fuchsian group. And if it is reciprocal then there is a fixed point of one of the reciprocator of g that lies on  $I_g$  by 2.2, and it is an even ordered-vertex of the fundamental domain (from the definition of fundamental domain). This proves one side of the statement.

Conversely, if the axis of the primitive hyperbolic element g passes through an even-order vertex z of the tiling which is done by the fundamental domain, then let k be the elliptic finite order element in the Fuchsian group such that, k(z) = z and  $k^{2m} = Id$ . So,  $k^m$  is an involution and it

is a is  $\pi$ -rotation. Then, it will interchange the fixed points of g, and so  $k^m g k^m = g^{-1}$  since g is primitive. This shows that any non-trivial power of g is reciprocal. This proves the lemma.

2.2. Proof of Theorem 2.1. Combining Lemma 2.2 and Lemma 2.3, we get the proof of the theorem.  $\hfill \Box$ 

2.3. Reversible classes of non-orientable surface fundamental groups. Let S be a compact connected non-orientable surface. Due to the classification of surfaces, the fundamental group of S can be presented in the form

(2.1) 
$$\langle x_1, x_2, \dots, x_k, d_1, d_2, \dots, d_m | \Pi x_i^2 \Pi d_i = 1 \rangle$$

where k represents number of cross-caps and the  $d_i$  represent the boundary components. In such cases, we have the following lemma.

**Lemma 2.4.** If S is a compact, connected non-orientable surface having the fundamental group 2.1, then there are no reversible elements in  $\pi_1(S)$  except when S is the Klein-bottle or  $\mathbb{RP}^2$ .

**Proof.** By the classification of non-orientable surfaces, we know that S is homeomorphic to  $\mathbb{RP}^2 \# \mathbb{RP}^2 \# \dots \# \mathbb{RP}^2.$ We also know that a loop l is reversible iff it will freely homotope to its  $\underbrace{\mathbb{RP}^2 \# \mathbb{RP}^2 \# \dots \# \mathbb{RP}^2}_{k \text{ times}}.$ 

own inverse, that is, they will form a Klein bottle. So, if S is itself a Klein bottle, then any power of the generating loop is reversible. If S is  $\mathbb{RP}^2$ , then the fundamental group will be  $\mathbb{Z}_2$ , and the generator is reversible.

If k > 2, there is no finite order element in the fundamental group. Then our claim is that there are no more reversible elements in S. If on the contrary there exists any loop l that homotopes to its own inverse, then that element along with the reverser will form a Klein bottle which is a closed surface (if the reverser is infinite order). That implies, there is an embedding of the Klein bottle in S. Since both S and the Klein bottle have same dimension and are both complete surfaces, therefore by the Hopf–Rinow theorem, they must be homeomorphic. This contradicts our hypothesis and hence proves the lemma.

Now we classify the reversible elements in the fundamental group of a non-orientable twodimensional orbifold  $\Gamma'$  which is neither  $\mathbb{RP}^2$  nor Klein bottle.

**Lemma 2.5.** Infinite order reversible elements in  $\Gamma'$  are strongly reversible elements.

The proof is similar to the proof of [DG1, Lemma 3.4].

**Proof.** The group  $\Gamma'$  is a discrete subgroup of  $PGL(2, \mathbb{R})$ . Let r in  $\mathbf{H}^2/\Gamma'$  be a reversible element in  $\Gamma'$ . Then there is an element b in  $\Gamma'$  such that  $b^{-1}rb = r^{-1}$ . At first we consider that r is orientation preserving isometry in  $\mathbf{H}^2$  (which implies that will not be a parabolic element). Then, b either fixes the fixed points of r or reflects the fixed points to each other. If b is fixes the fixed points of r then it will either commute with r or reverse r. In the case where it commutes with r it leads us to a contradiction that b is not a reverser, and in the case where it is reversing, b will be an involution. If b interchanges the fixed points, then b must be an involution.

In r is a orientation reversing element, then  $r^2$  will be orientation preserving element with the same reverser b. From the previous case,  $b^2 = 1$ . Hence, in any case, r will be strongly reversible.

**Theorem 2.6.** The infinite ordered reversible elements in  $\Gamma'$  are conjugate to  $c_i^{\frac{\mu_i}{2}} k c_j^{\frac{\mu_j}{2}} k^{-1}$  where  $\mu_i$  and  $\mu_j$  are even and  $k \in \Gamma'$ .

**Proof.** Since there is no finite order element in  $\Gamma'$  other than the  $c_i$ , this theorem follows from the previous lemma.

#### 3. Seifert fibered space

To find the reversible elements of the fundamental group of a Seifert fibered space, we first focus on the quotient group  $\pi_1(M)/\langle h \rangle$  and discuss its reversibility. We first deal with those spaces which have orientable orbit surfaces.

**Corollary 3.1.** Any element conjugate to either  $c_i^{\frac{\mu_i}{2}}$  or  $c_i^{\frac{\mu_i}{2}} k c_j^{\frac{\mu_j}{2}} k^{-1}$  where  $k \in \pi_1(M)/\langle h \rangle$  is reversible in  $\pi_1(M)/\langle h \rangle$  iff  $\mu_i$  and  $\mu_j$  are even. No other elements in  $\pi_1(M)/\langle h \rangle$  are reversible.

**Proof.** Since  $\pi_1(M)/\langle h \rangle$  is a Fuchsian group  $\Gamma$ , all reversible elements in  $\Gamma$  are strongly reversible. By [DG1, Theorem 1.1], it is known that an element in a Fuchsian group is reversible if and only if it is an involution or a product of 2 involutions. By the representation of the group, the involutions are the elements conjugate to any  $c_i^{\frac{\mu_i}{2}}$  where  $\mu_i$  is even. Hence, the claim follows.  $\Box$ 

Now we can discuss the reversibility of the fundamental group of M. Since there is no finite order element in  $\pi_1(M)$ , we cannot talk about strongly reversible elements here. But there may be reversible elements, as stated earlier. Not that if an element is reversible, then so are its conjugates.

**Lemma 3.2.** Elements conjugate to the power of the fiber generator h is a reversible element iff the classifying homomorphism on  $\pi_1(G)$  is non-trivial.

**Proof.** If an element in  $\pi_1(M)$  is reversible then it must be free homotopic to its inverse. This free homotopy forms an embedded Klein bottle in M. Let  $p: \pi_1(M) \to \pi_1(G)$  be the projection map from M to the orbit space. If there is a homotopy class of a loop z that is generated by the generating elements of  $\pi_1(M)$  and has a non-trivial part which cuts the fiber h transitively, then p(z) is a loop in the orbit space. So h is reversible iff the image of p(z) under  $\phi$  is non-trivial. Since the image of  $\phi$  of z depends on image of  $\phi$  of generating set elements, this implies the lemma.

Alternatively, since  $\phi$  is a homomorphism, it is determined by its actions on the generators of  $\pi_1(G)$ . If  $\phi$  is non-trivial, there is some generator x of the form  $a_i, b_i, x_i$  or  $d_i$  such that  $\phi(x) = -1$ . Then by the presentation of  $\pi_1(M), xhx^{-1} = h^{\phi(x)} = h^{-1}$ , and so h is reversible.

**Corollary 3.3.** If h is reversible in  $\pi_1(M)$ , the  $c_i^{\mu_i}$  are reversible elements although each  $c_i$  is an infinite order element in  $\pi_1(M)$ .

**Proof.** Let h be reversible. Then so is  $h^{\beta_i}$ . The relation  $c_i^{\mu_i} = h^{\beta_i}$  now implies that  $c_i^{\mu_i}$  is also reversible.

The two following lemmas investigate the reversibility of the other elements of  $\pi_1(M)$ .

**Lemma 3.4.** The powers of  $c_i^{\mu_i}$  are reversible, but any other multiple of  $c_i$  is not reversible for any  $\pi_1(M)$ .

**Proof.** On the contrary, let a multiple of  $c_i$  be a reversible element in  $\pi_1(M)$  with reverser  $\rho$ . We can consider that  $\rho$  is not a power of h, since h is contained in the centralizer of  $c_i$ . Then, the reverser  $\rho$  is transverse to h, and this implies any power of  $\rho$  will always be transverse to h. Hence  $\rho$  has a non-trivial image in the group  $\pi_1(M)/\langle h \rangle$ , denoted by  $[\rho]$ . Since, from [DG1], every reversible element in a Fuchsian group is strongly reversible,  $[\rho]$  must be of order 2. This implies,  $\rho^2 = h^k$  for some  $k \in \mathbb{Z}$ , which contradicts our assumption. So, there is no such reverser that reverses any other power of  $c_i$  except for the powers of the  $c_i^{\mu_i}$ .

**Lemma 3.5.** If *M* has an orientable orbit surface, the only reversible elements in  $\pi_1(M)$  can be the conjugates of the powers of *h* and the conjugates of the elements either of the form  $c_i^{\frac{\mu_i}{2}}kc_j^{\frac{\mu_j}{2}}k^{-1}$ with  $\phi(k) = -1$  or of the form  $c_i^{\frac{\mu_i}{2}}kc_j^{\frac{-\mu_j}{2}}k^{-1}$  with  $\phi(k) = 1$ , where  $\mu_i$  and  $\mu_j$  are even, and *k* is any element in  $\pi_1(M)$ ,  $\beta_i = \beta_j$ .

**Proof.** Let x be an element in  $\pi_1(M)$ , other than any power of h. Let x be reversible. Then x is freely homotopic to  $x^{-1}$ . Consider the projection of this free homotopy to the Fuchsian group  $\pi_1(M)/\langle h \rangle$ . This would be a free homotopy of a loop in this Fuchsian complex to its inverse. But this would imply that this is a reversible element in  $\pi_1(G)/\langle h \rangle$ . But as we have shown earlier, the reversible elements in this group are of the form of conjugates of  $c_i^{\frac{\mu_i}{2}} k c_j^{\frac{\mu_j}{2}} k^{-1}$  or  $c_i^{\frac{\mu_i}{2}}$  for even  $\mu_i$  and  $\mu_j$ . But, for elements that are conjugates of  $c_i^{\frac{\mu_i}{2}}$ , i.e.,  $\alpha c^{\frac{\mu_i}{2}} \alpha^{-1}$ , the reverser of x must be x itself. Then the element in  $x \in \pi_1(M)$  will satisfy,

$$\alpha c^{\frac{\mu_i}{2}} \alpha^{-1} \alpha c^{\frac{\mu_i}{2}} \alpha^{-1} \alpha c^{-\frac{\mu_i}{2}} \alpha^{-1} = \alpha c^{-\frac{\mu_i}{2}} \alpha^{-1}$$
$$\Rightarrow h^{\beta_i} = 1.$$

Since h is an infinite order element, this is a contradiction. Hence, the lifts of the elements conjugate to  $c^{\frac{\mu_i}{2}}$  are not reversible in  $\pi_1(M)$ . So, the only other candidates from  $\pi_1(M)/\langle h \rangle$  that may be reversible in  $\pi_1(M)$ , are the products of the involutions.

Let the element 
$$x = c_i^{\frac{\mu_i}{2}} k c_j^{\frac{\gamma_j}{2}} k^{-1}$$
 be reversible in  $\pi_1(M)/\langle h \rangle$  with reverser  $c^{\frac{\mu_i}{2}}$ . Then in  $\pi_1(M)$ ,  
 $c_i^{\frac{\mu_i}{2}} c_i^{\frac{\mu_j}{2}} k c_j^{\frac{\mu_j}{2}} k^{-1} c_i^{-\frac{\mu_i}{2}} c_i^{\frac{\mu_j}{2}} k^{-1} = c_i^{\mu_i} k c_j^{\frac{\mu_j}{2}} k^{-1} k c_j^{\frac{\mu_j}{2}} k^{-1} = h^{\beta_i} k h^{\beta_j} k^{-1}.$ 

Then,  $h^{\beta_i}kh^{\beta_j}k^{-1}$  will be identity only if  $\beta_i = \beta_j$  and  $\phi(k) = -1$ . It will be same if we choose any other reverser for that particular reversible element since all the reversers are in the form of  $x^l c^{\frac{\mu_i}{2}}$ . In the similar way, we can show that  $c_i^{\frac{\mu_i}{2}}kc_j^{-\frac{\mu_j}{2}}k^{-1}$  is reversible if  $\phi(k) = 1$ . If x is not reversible in  $\pi_1(M)/\langle h \rangle$ , a reverser of x in  $\pi_1(M)$ , say  $\rho$ , must be a power of h, since the projection to  $\pi_1(M)/\langle h \rangle$  must be constant. Then in  $\pi_1(M)/\langle h \rangle$ ,  $[x] = [x^{-1}]$ . This implies that  $[x^2] = 1$ . Then,  $x^2$  is a power of h, and  $x^2$  is also reversible with same reverser  $\rho$  that x is. Then,

$$h^{k}x^{2}h^{-k} = x^{-2}$$
$$\Rightarrow x^{2} = x^{-2}$$
$$\Rightarrow x^{4} = Id.$$

This is contradiction since  $\pi_1(M)$  is torsion free. This proves our lemma.

3.1. **Proof of Theorem 1.1.** The previous lemma shows that there are no reversible elements in  $\pi_1(M)$  other than those conjugate to h. As shown earlier, the reversibility of h and its conjugates depends on the classifying homomorphism. This is summarised in the statement of Theorem 1.1.

In the case of non-orientable surfaces, we have certain exceptions to this situation, as shown in the previous section.

The reversible elements of  $\pi_1(M)$  in case of a non-orientable orbit surface can be summarised as follows, by the same proofs as above.

**Corollary 3.6.** If M has an non-orientable orbit surface other than the Klein bottle or  $\mathbb{RP}^2$ , the reversible elements of  $\pi_1(M)$  are either conjugate to any power of h or either of the form  $c_i^{\frac{\mu_i}{2}}kc_j^{\frac{\mu_j}{2}}k^{-1}$  with  $\phi(k) = -1$  or of the form  $c_i^{\frac{\mu_i}{2}}kc_j^{\frac{-\mu_j}{2}}k^{-1}$  with  $\phi(k) = 1$ , where  $\mu_i$  and  $\mu_j$  are even, and k is any element in  $\pi_1(M)$ ,  $\beta_i = \beta_j$ .

When the orbit surface is the Klein bottle or  $\mathbb{RP}^2$ , we get additional reversible elements which are the generators of the orbit surface. Thus, in the case of a given orbit surface, the reversibility of the elements of the fundamental group of a Seifert fibered space over that orbit surface is solely determined by  $\phi$ .

**Remark 3.7.** The classification of the reversible elements in a given Seifert-fibered space depends only on the classifying homomorphism.

#### 4. Application to universal extension of Fuchsian groups

In the previous section, we have classified the reversible elements of the fundamental group of a Seifert-fibered space. We can see the fundamental group in the form of an exact sequence:

$$(4.1) 1 \longrightarrow \mathbb{Z} \longrightarrow \pi_1(M) \longrightarrow F \longrightarrow 1$$

Here,  $\pi_1(M)$  is the fundamental group of M, and F is a Fuchsian group or discrete isometry group of PGL(2,  $\mathbb{R}$ ). So,  $\pi_1(M)$  is then a universal extension of F, which can now be seen as  $\pi_1(M)/\langle h \rangle$ . This gives us a new view-point to consider certain groups as fundamental groups of a Seifert-fibered space. Let us consider  $F = \text{PSL}(2, \mathbb{Z})$  which is presented by  $\langle a, b | a^2, b^3 \rangle$ . We have already mentioned the presentation of  $\pi_1(M)$  for a given M in Equation (1) and Equation (1).

We can choose an orientable connected 3-manifold M in such a way that the  $\pi_1(M) \cong B_3$ .  $B_3$  has a presentation as  $\langle \sigma_1 \sigma_2 | \sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2 \rangle$ . The normal subgroup of  $B_3$  is  $\mathbb{Z}$  and it is generated by  $(\sigma_1 \sigma_2)^3$ . Thus,  $B_3/\langle \sigma_1 \sigma_2 \rangle = \langle \sigma_1 \sigma_2 \sigma_1, \sigma_1 \sigma_2 | (\sigma_1 \sigma_2)^3, (\sigma_1 \sigma_2 \sigma_1)^2 \rangle = \text{PSL}(2, \mathbb{Z}) \cong \Delta(2, 3, \infty)$ . Choosing the  $c_i$  and  $\beta_i$  suitably, we can consider the Seifert-fibered manifold M determined by (O, o, 0|1, (2, 1), (3, 1)). The presentation of the fundamental group of M is then

(4.2) 
$$\langle c_1, c_2, d, h | c_i h c_i^{-1} = h, dh d^{-1} = h^{\phi(d)}, c_i^{\mu_i} = h_i^{\beta}, c_1 c_2 dh = 1 \rangle.$$

Now, consider the generating elements  $c_1 = \sigma_1 \sigma_2 \sigma_1, c_2 = \sigma_1 \sigma_2$ , which implies that  $h = (\sigma_1 \sigma_2)^3$ and  $d^{-1} = \sigma_1 \sigma_2 \sigma_1^2 \sigma_2$ .

This determines the Fuchsian complex  $\pi_1(M)/\langle h \rangle$ , and we obtain the resultant orbit surface G as the open disk D by further quotienting it by the  $c_i$ . Then,  $\phi : \pi_1(D) \longrightarrow \mathbb{Z}_2$  is the trivial homomorphim. Then, the group 4.2 can be presented as

$$\langle \sigma_1 \sigma_2 \sigma_1, \sigma_1 \sigma_2, \sigma_1 \sigma_2 \sigma_1^2 \sigma_2, (\sigma_1 \sigma_2)^3 | (\sigma_1 \sigma_2)^3 \sigma_1 \sigma_2 \sigma_1 = \sigma_1 \sigma_2 \sigma_1 (\sigma_1 \sigma_2)^3, (\sigma_1 \sigma_2 \sigma_1)^2 = (\sigma_1 \sigma_2)^3 \rangle$$
  
=  $\langle \sigma_1, \sigma_2 | \sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2 \rangle = B_3.$ 

So, we can deduce the exact sequence

$$(4.3) 1 \longrightarrow \mathbb{Z} \longrightarrow B_3 \longrightarrow \mathrm{PSL}(2,\mathbb{Z}) \longrightarrow 1$$

**Corollary 4.1.** The only reversible elements of  $B_3$  are conjugates to  $[\sigma_1 \sigma_2 \sigma_1, k]$  for any element k in  $B_3$ .

**Proof.** From the above discussion,  $\phi$  is trivial, and applying Theorem 3.5 we know that reversible elements for the orientable case depend only on  $\phi$ . The claim thus follows from this theorem.

Note that we cannot use the same technique to find the reversible elements of  $B_n$  since  $B_n$  is not a Seifert-fibered group for any n > 3. This follows from the theorem below.

## **Theorem 4.2.** For any n > 3 the braid group $B_n$ is not isomorphic to any Seifert-fibered group.

**Proof.** On the contrary let us assume that braid groups are Seifert-fibered groups. So, let there be a Seifert-fibered manifold M such that  $B_n \cong \pi_1(M)$ , with maximal normal subgroup  $\langle h \rangle$ where h represents an ordinary fiber. It is well known that the braid group  $B_n$  where n > 2 is not virtually abelian. Then, from [Ma, Corollary 10.4.6], the center of  $\pi_1(M)$  is either  $\langle h \rangle$  or trivial. Since the center of  $B_n$  is not trivial, therefore  $Z(B_n)$  is  $\langle h \rangle$ . Let us consider the braid group  $B_n$ 

$$B_n = \langle \sigma_1 \sigma_2 \dots \sigma_{n-1} | \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \sigma_i \sigma_j = \sigma_j \sigma_i \text{ for } |i-j| > 2 \rangle$$

In another presentation we can rewrite the group as mentioned in [MK, Chapter 2, Section 2, Exercise 2.4]:

(4.4) 
$$B_n = \langle x, y | x^n = (xy)^{n-1}, x^i y x^{-i} y = y x^i y x^{-i} \text{ where } i = 2, 3, \dots, m \rangle$$

where,  $m = \lfloor \frac{n}{2} \rfloor$ . Then, the fiber  $h = (\sigma_1 \sigma_2 \dots \sigma_{n-1})^n = x^n$  is the generator of  $Z(\pi_1(M))$ . Then,

$$B_n/\langle h \rangle = \langle x, y | x^n = (xy)^{n-1} = 1, x^i y x^{-i} y = y x^i y x^{-i} \text{ where } i = 2, 3, \dots, m \rangle$$

is a Fuchsian group or a fundamental group of non-orientable orbifold surface. So, it acts as an isometry group in  $\mathbf{H}^2$ . If  $B_n/\langle h \rangle$  acts in  $\mathbf{H}^2$ , then  $x^iyx^{-i}$  and y are powers of the same element  $\psi \in B_n/\langle h \rangle$ , that is,  $x^iyx^{-i} = \psi^u$  and  $y = \psi^v$  for some integers u, v. But in [DG1], we noticed that  $x^i$  sends fixed points of y to the fixed points of  $x^iyx^{-i}$ . That means either  $x^i$  fixes the fixed points of y, or interchanges them for every i. If it fixes the fixed points then that implies xy = yx. Then x is an element in the center of  $B_n/\langle h \rangle$ , so it must be trivial. This is contradiction in our assumption.

So,  $x^i$  must interchange the fixed points of y for every i. Then  $x^2$  must be an involution (orientation preserving or orientation reversing). This implies n = 4, and thus  $x^2yx^{-2} = y^{-1}$ .  $y^{-1}$  is  $\sigma_3$  in  $B_4$ , and then  $\sigma_1\sigma_3$  is an element in the center which contradicts our assumption. So,  $B_n/\langle h \rangle$  will not be the fundamental group of a Seifert-fibered space for any n > 3.

**Remark 4.3.** From Corollary 4.1, there are some reversible elements in  $B_n$  which are conjugate to the elements of the form [a, b], where the image of a is an involution in the quotient space  $B_n/\langle x^n \rangle$  and b is any other element in  $B_n$ . This is a sufficient condition to be a reversible element in  $B_n$ .

**Remark 4.4.** From the above theorem, we obtain that the same idea that we have used for Seifertfibered spaces will not work for  $B_n$ . Furthermore, by [HMT, Theorem 1.3] and Theorem 4.2,  $B_n$ cannot be a fundamental group of any 3-manifold. This has been proven earlier using different techniques, for example in [Go] where they use the theory of coherent groups, however this gives a different perspective using reversible elements.

### References

[Br]	M. G. Brin. Seifert Fibered Spaces: Notes for a course given in the Spring of 1993. arXiv preprint arXiv:0711.1346 1
[BR]	M. Baake and J. A. G. Roberts, The structure of reversing symmetry groups. Bull. Austral. Math.
[DG1]	<ul> <li>Soc. 73 (2006), no. 3, 445–459. 1</li> <li>D. Das, K. Gongopadhyay. <i>Reciprocity in the Hecke Groups.</i> arXiv preprint arXiv:2307.00496.(2023) 2.1, 2.1, 2.3, 3, 3, 4</li> </ul>
[DG2]	D. Das, K. Gongopadhyay. On The z-classes in Fuchsian group. (In preparation) 2.1, 2.1
[Go]	C. M. Gordon. Artin groups, 3-manifolds and coherence. Boletín de la Sociedad Matemática Mexi- cana, Vol. 10, (2004), 193-198. https://web.ma.utexas.edu/users/gordon/artin-dedicated.pdf 4.4
[HMT]	K. Himeno, K. Motegi, and M. Teragaito. Classification of generalized torsion elements of order two in 3-manifold groups. https://sites.google.com/izt.uam.mx/citaandppicta2023/home 1, 4.4
[La]	J. S. W. Lamb, Reversing symmetries in dynamical systems. J. Phys. A 25 (1992), no. 4, 925–937. 1
[Ma]	B. Martelli. Introduction to Geometric Topology arXiv preprint arXiv:1610.02592 4
[MK]	K. Murasugi and B. I. Kurpita, A study of braids, Mathematics and its Applications 484, Kluwer Academic Publishers, Dordrecht, 1999. 4
[ST]	H. Seifert, W. Threlfall. Seifert and Threlfall: a textbook of topology Pure Appl. Math., 89 Academic

Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1980. xvi+437 pp. 1

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH (IISER) MOHALI, KNOWLEDGE CITY, SECTOR 81, S.A.S. NAGAR 140306, PUNJAB, INDIA

Email address: nshrdas@gmail.com Email address: debattam123@gmail.com