CALIBRATING DOUBLY-ROBUST ESTIMATORS WITH UNBALANCED TREATMENT ASSIGNMENT

WORKING PAPER

Daniele Ballinari^{*†} Swiss National Bank Börsenstrasse 15 8001 Zurich, Switzerland daniele.ballinari@snb.ch

May 20, 2024

ABSTRACT

Machine learning methods, particularly the double machine learning (DML) estimator (Chernozhukov et al., 2018), are increasingly popular for the estimation of the average treatment effect (ATE). However, datasets often exhibit unbalanced treatment assignments where only a few observations are treated, leading to unstable propensity score estimations. We propose a simple extension of the DML estimator which undersamples data for propensity score modeling and calibrates scores to match the original distribution. The paper provides theoretical results showing that the estimator retains the DML estimator's asymptotic properties. A simulation study illustrates the finite sample performance of the estimator.

JEL classification C14 · C21 · C52 · C55

Keywords Causal machine learning \cdot Double machine learning \cdot Average treatment effect \cdot Unbalanced treatment assignment \cdot Undersampling

1 Introduction

Estimation of the average treatment effect (ATE) is of central importance in empirical research. The interest generally lies in the effect that a binary treatment has on an outcome variable. For example, we might be interested in the effect that a training program has on the unemployment duration. With the increasing availability of large datasets, the use of machine learning (ML) methods to estimate the ATE has become popular (Athey and Imbens, 2019; Yao et al., 2021). The double machine learning (DML) estimator (Chernozhukov et al., 2018) is a widely adopted ATE estimator that relies on ML-estimated nuisance functions. The DML estimator has been shown to be consistent and asymptotically normal, even when using ML methods that converge at a slower rate than the parametric rate (Chernozhukov et al., 2018).

While dataset sizes have increased, the number of treated units often remains small. The treatment is often costly and/or time-consuming. For example, the submission of a new drug to a patient might take several months, or a training program for the unemployed is very costly. Control outcomes and covariates, on the other hand, are more easily collected from, e.g., administrative agencies, medical records, or financial markets (Künzel et al., 2019; Bouchaud, 2022; Hujer et al., 2006). The dataset might then consist of only a few treated, but many control observations. This unbalancedness can lead to unstable propensity score estimations (Huber et al., 2013). Even though the DML estimator

^{*}The views, opinions, findings, and conclusions or recommendations expressed in this paper are strictly those of the author(s). They do not necessarily reflect the views of the Swiss National Bank (SNB). The SNB takes no responsibility for any errors or omissions in, or for the correctness of, the information contained in this paper.

[†]For helpful comments and suggestions, we thank Nora Bearth, Michael Lechner, Thomas Maag, Alexander Wehrli, Michael Zimmert, and two anonymous reviewers.

relies on a doubly robust approach that combines the conditional outcome expectations with the propensity score, instability in the propensity score estimation can lead to high variability in the ATE estimate.

In this paper, we propose a simple extension of the DML estimator that addresses the issue of unbalanced treatment assignment. Inspired by the ML classification literature (Japkowicz and Stephen, 2002), the proposed approach undersamples the data used for fitting the ML model for the propensity score. Using the relation between the true and the undersampled propensity score, we calibrate the propensity scores to match the original data distribution. We show that the proposed estimator has the same asymptotic distribution as the DML estimator, attains the parametric rate of convergence \sqrt{N} and its variance achieves the semi-parametric efficiency bound (Hahn, 1998). We illustrate the finite sample performance of the estimator in a simulation study. While we present the results in the context of the DML estimator, the proposed approach applies to any ATE estimator that relies on the efficient score function.

2 Calibration estimator

2.1 Notation and causal identification

We define causal effects using Rubin's 1972 potential outcome framework. The interest lies in the effect of a binary treatment variable³ D on an outcome variable Y. We denote the potential outcome for treatment D = d, that is the outcome one would observe if the treatment was d, as Y^d . The effect of interest is the average treatment effect (ATE) defined as:

$$\theta = \mathbb{E}[Y^1 - Y^0] \tag{1}$$

The potential outcomes are not directly observed and the parameter of interest has to be identified from observational data. The researcher observes a sample of i.i.d. random variables $\{Z_1, \ldots, Z_N\}$ where $Z_i := (X_i, D_i, Y_i)$, with X_i being a *p*-dimensional vector of exogenous control variables with support \mathcal{X} , D_i the binary treatment random variable, and Y_i the observed outcome. Define the conditional outcome expectations as $\mu_d(X) = \mathbb{E}[Y|D = d, X]$ and the propensity score as $p(X) = \mathbb{E}[D|X] = \mathbb{P}[D = 1|X]$. The respective estimated quantities are denoted as $\hat{\mu}_d(X)$ and $\hat{p}(X)$. Finally, we denote by $\tau(Z)$ the efficient score function:

$$\tau(Z) = \mu_1(X) - \mu_0(X) + \frac{D}{p(X)}(Y - \mu_1(X)) - \frac{1 - D}{1 - p(X)}(Y - \mu_0(X)).$$

Identification of the ATE from observable outcomes is achieved under the following assumption. Assumption 1 (Identification). For observation Z = (X, D, Y) assume that (Rosenbaum and Rubin, 1983):

- (i) $Y^0, Y^1 \perp D | X = x$ for any $x \in \mathcal{X}$ (conditional independence assumption).
- (ii) The observed outcome is $Y = DY^1 + (1 D)Y^0$ (stable unit treatment value assumption).
- (iii) For any $x \in \mathcal{X}$ it holds that $\eta < p(x) < 1 \eta$ for some $\eta > 0$ (common support).
- (iv) $X = X^1 = X^0$ where X^d denotes the random covariate vector under treatment d (exogenity of covariates).

Under Assumption 1 the ATE can be characterized as a functional of the joint distribution of the observed data (X, D, Y) (Athey and Imbens, 2019):

$$\theta = \mathbb{E}[Y^1 - Y^0] = \mathbb{E}[\mu_1(X) - \mu_0(X)] = \mathbb{E}[\tau(Z)].$$
(2)

2.2 Double machine learning estimator

The estimator proposed in this study extends the popular ATE estimator developed by Chernozhukov et al. (2018) generally referred to as the double machine learning (DML) estimator. DML builds on two key ingredients. First, it uses the efficient score function $\tau(Z)$ to construct an estimator of the ATE. Second, it uses cross-fitting to estimate the nuisance functions $\mu_d(X)$ and p(X). In more detail, the DML estimator for the ATE is defined as follows:

STEP 1 For some fixed $K \in \{2, ..., N\}$, randomly partition the observation indices into K sets $\mathcal{I}_1, ..., \mathcal{I}_K$ of equal size. Denote the complement of \mathcal{I}_k by $\mathcal{I}_{-k} = \{1, ..., N\} \setminus \mathcal{I}_k$. Denote the cardinality of each set of indices by $|\mathcal{I}_k|$.

³The estimator presented in this paper can be directly generalized to multivalued treatments (see, among others, Farrell, 2015; Knaus, 2020).

STEP 2 for k = 1 to K do:

Estimate the nuisance functions $\mu_d(x)$ and p(x) on the sample defined by indices \mathcal{I}_{-k} and denote the estimated functions by $\widehat{\mu}_d^{\mathcal{I}_{-k}}(x)$ and $\widehat{p}^{\mathcal{I}_{-k}}(x)$.

end for

STEP 3 Estimate the ATE using the estimator:

$$\widehat{\theta}^{\text{DML}} = \frac{1}{N} \sum_{k=1}^{K} \sum_{i \in \mathcal{I}_k} \widehat{\tau}^{\mathcal{I}_{-k}}(Z_i)$$
(3)

where:

$$\hat{\tau}^{\mathcal{I}_{-k}}(Z_i) = \hat{\mu}_1^{\mathcal{I}_{-k}}(X_i) - \hat{\mu}_0^{\mathcal{I}_{-k}}(X_i) + \frac{D_i}{\hat{p}^{\mathcal{I}_{-k}}(X_i)}(Y_i - \hat{\mu}_1^{\mathcal{I}_{-k}}(X_i)) - \frac{1 - D_i}{1 - \hat{p}^{\mathcal{I}_{-k}}(X_i)}(Y_i - \hat{\mu}_0^{\mathcal{I}_{-k}}(X_i))$$

Chernozhukov et al. (2018) provide an asymptotic theory for the DML estimator. In particular, they show that the estimator is consistent and asymptotically normal, even when using machine learning (ML) methods that converge at a slower rate than the parametric rate.

2.3 Calibration estimator

A shortcoming of the DML estimator is its poor finite sample performance when the treatment assignment is unbalanced, i.e. when either very few or very many observations are treated. In the following, without loss of generality, we consider only the case where very few are treated. ML models perform poorly when the data is unbalanced (Japkowicz and Stephen, 2002) and predict propensity scores that are potentially close to zero or one. A common approach to improving the performance of ML models is to undersample the observations that are overrepresented, i.e. the observations that are not treated (He and Garcia, 2009).⁴

Undersampling the observations changes the underlying distribution of the data and the predicted propensity scores will be biased for the true propensity scores from the unbalanced distribution (Pozzolo et al., 2015). To formalize this concept, let S_i be a random variable equal to 1 if observation *i* is sampled and 0 otherwise. It then follows that $\mathbb{P}[S_i = 1|D_i = 1] = 1$ since all treated observations are kept in the sample. For the untreated observations we have that $\mathbb{P}[S_i = 1|D_i = 0] = \mathbb{E}[D_i]/(1 - \mathbb{E}[D_i]) =: \gamma$. Moreover, since the undersampling strategy is not dependent on the control variables *X*, we have that $\mathbb{P}[S_i = 1|D_i = d, X_i] = \mathbb{P}[S_i = 1|D_i = d]$. Using Bayes' rule it can be shown that the propensity score for the undersampled data $p_S(X_i)$ is given by (Pozzolo et al., 2015):

$$p_S(X_i) = \frac{p(X_i)}{p(X_i) + \gamma \cdot (1 - p(X_i))}$$
(4)

from where it follows that $p_S(X_i) \neq p(X_i)$ for $\gamma < 1$.

A naïve solution to address this issue would be to not only undersample the data used for estimating the propensity score but also the data on which the efficient score function is computed. In other words, a valid strategy would be to undersample the entire dataset and apply DML to the undersampled data (hereafter referred to as U-DML). However, this approach reduces the number of observations from N to $2 \cdot \mathbb{E}[D_i] \cdot N$. In cases where only 5% of the observations are treated, we would discard 90% of the observations.

We propose a calibration estimator that uses the entire sample and corrects for the bias in the propensity score. The main idea is to only undersample the data used for fitting the ML model for the propensity score. Using the relation between the true and the undersampled propensity score in Equation (4), we calibrate the propensity scores to match the original data distribution. In more detail, the calibrated-undersampled DML (CU-DML) estimator is defined as follows:

STEP 1 Estimate γ as:

$$\widehat{\gamma} = \frac{\sum_{i=1}^{N} D_i}{\sum_{i=1}^{N} (1 - D_i)}$$
(5)

⁴Another common technique used in the ML literature is to oversample the minority class (Japkowicz and Stephen, 2002). This approach might be problematic in the context of DML since it introduces dependence in the data, complicating the asymptotic theory of the proposed estimator. The convergence rates of common ML methods needed for the asymptotic results presented in Section 2.4 have been proven for the case of independent and identically distributed data; see, e.g. Belloni and Chernozhukov (2013) for the Lasso, Luo et al. (2016) for L_2 boosting, Wager and Walther (2016) for random forests, and Chen and White (1999) for neural networks.

STEP 2 For some fixed $K \in \{2, ..., N\}$, randomly partition the observation indices into K sets $\mathcal{I}_1, ..., \mathcal{I}_K$ of equal size. Denote the complement of \mathcal{I}_k by $\mathcal{I}_{-k} = \{1, ..., N\} \setminus \mathcal{I}_k$. Denote the cardinality of each set of indices by $|\mathcal{I}_k|$.

STEP 3 for k = 1 to K do:

- STEP 3.1 Estimate the nuisance functions $\mu_d(X)$ on the sample defined by indices \mathcal{I}_{-k} and denote the estimated functions by $\hat{\mu}_d^{\mathcal{I}_{-k}}(X)$.
- STEP 3.2 Draw random variables S_i for $i \in \mathcal{I}_{-k}$ from a Bernoulli distribution with probability $\mathbb{P}[S_i = 1 | D_i = 1] = 1$ and $\mathbb{P}[S_i = 1 | D_i = 0] = \hat{\gamma}$. Define the undersampled indices as:

$$\mathcal{I}_{-k}^S = \{i \in \mathcal{I}_{-k} | S_i = 1\}$$

Estimate the nuisance function $p_S(X)$ on the sample defined by indices \mathcal{I}_{-k}^S and denote the estimated function by $\hat{p}_S^{\mathcal{I}_{-k}^S}(X)$. Calibrate the propensity score estimated on the undersampled data to match the original data distribution:

$$\hat{p}^{\mathcal{I}_{-k}}(X) = \frac{\hat{\gamma} \cdot \hat{p}_S^{\mathcal{I}_{-k}^s}(X)}{\hat{\gamma} \cdot \hat{p}_S^{\mathcal{I}_{-k}^s}(X) + \left(1 - \hat{p}_S^{\mathcal{I}_{-k}^s}(X)\right)}$$
(6)

end for

STEP 4 Estimate the ATE using the estimator:

$$\widehat{\theta}^{\text{CU-DML}} = \frac{1}{N} \sum_{k=1}^{K} \sum_{i \in \mathcal{I}_k} \widehat{\tau}^{\mathcal{I}_{-k}}(Z_i)$$
(7)

where:

$$\hat{\tau}^{\mathcal{I}_{-k}}(Z_i) = \hat{\mu}_1^{\mathcal{I}_{-k}}(X_i) - \hat{\mu}_0^{\mathcal{I}_{-k}}(X_i) + \frac{D_i}{\hat{p}^{\mathcal{I}_{-k}}(X_i)}(Y_i - \hat{\mu}_1^{\mathcal{I}_{-k}}(X_i)) - \frac{1 - D_i}{1 - \hat{p}^{\mathcal{I}_{-k}}(X_i)}(Y_i - \hat{\mu}_0^{\mathcal{I}_{-k}}(X_i))$$

Notice that, in contrast to the nuisance functions, γ can be estimated from the entire sample. For $\hat{\gamma} = 1$ the CU-DML estimator reduces to the classical DML estimator. Furthermore, the results presented in the next section show that asymptotically $\hat{\theta}^{\text{CU-DML}}$ converges in probability to $\hat{\theta}^{\text{DML}}$, also for $\hat{\gamma} < 1$.

2.4 Asymptotic results

We now present the asymptotic results for the CU-DML estimator. We start by introducing the following assumptions. Assumption 2 (Boundedness of conditional variances). *For the conditional variance of the outcome it holds that:*

$$\sup_{x \in \mathcal{X}} \operatorname{Var}\left[Y|D = d, X = x\right] < \zeta$$

for some $\zeta < \infty$.

Assumption 3 (Boundedness of the propensity score).

(i) For the unconditional propensity score $\lambda := \mathbb{E}[p(X)] = \mathbb{E}[D]$ and its estimator $\widehat{\lambda} = N^{-1} \sum_{i=1}^{N} D_i$ it holds that:

$$\epsilon < \lambda < 1 - \epsilon \qquad \epsilon < \lambda < 1 - \epsilon$$

for some $\epsilon > 0$.

(ii) For all $x \in \mathcal{X}$ it holds that:

$$\eta < p_S(x) < 1 - \eta$$

for some $\eta > 0$.

Assumption 4 (Convergence of the ML estimators).

(i) The ML methods are sup-norm consistent:

$$\sup_{x \in \mathcal{X}} |\widehat{\mu}_d(x) - \mu_d(x)| \stackrel{p}{\to} 0 \qquad \sup_{x \in \mathcal{X}} |\widehat{p}_S(x) - p_S(x)| \stackrel{p}{\to} 0$$

(ii) The ML methods have risk-decay rates that satisfy (risk-decay assumption):

$$\mathbb{E}\left[\left(\widehat{\mu}_d(x) - \mu_d(x)\right)^2\right] \mathbb{E}\left[\left(\widehat{p}_S(x) - p_S(x)\right)^2\right] = o(N^{-1})$$

These assumptions closely resemble the ones required for the asymptotic results of the DML estimator (Chernozhukov et al., 2018; Wager, 2022). In addition to the boundedness of the propensity score, Assumption 3 (i) requires the unconditional propensity score to be bounded away from zero and one. This assumption is natural, as in a situation where the expected propensity score equals zero (one), there are no (only) treated. Importantly, Assumption 3 (ii) relaxes the usual bounds imposed on the propensity score, since $\tilde{\eta} < p(X) < 1 - \tilde{\eta}$ with $\tilde{\eta} < \eta$ whenever $\gamma < 1.^5$ Assumption 4 states the convergence rate requirements of the ML estimators in terms of the undersampled propensity score $p_S(X)$. The theoretical result is then given by the following theorem.

Theorem 1. Under Assumptions 2, 3 and 4 it holds that:

$$\sqrt{N}\left(\widehat{\theta}^{CU\text{-}DML} - \theta\right) \xrightarrow{d} \mathcal{N}\left(0, V^*\right)$$

where:

$$V^* = \operatorname{Var}[\mu_1(X_i) - \mu_0(X_i)] + \mathbb{E}\left[\frac{\sigma_1^2(X_i)}{p(X_i)}\right] + \mathbb{E}\left[\frac{\sigma_0^2(X_i)}{1 - p(X_i)}\right]$$

with $\sigma_d^2(X_i) = \operatorname{Var}[Y_i^d | X_i].$

The proof of Theorem 1 is relegated to Appendix A. Theorem 1 shows that the CU-DML estimator has the same asymptotic distribution as the DML estimator. In particular, the estimator attains the parametric rate of convergence \sqrt{N} and its variance achieves the semi-parametric efficiency bound (Hahn, 1998).

3 Simulation study

In this section, we investigate the finite sample performance of the CU-DML estimator in a simulation study. We consider two simulation strategies. In the first strategy, we generate data from a synthetic data generating process (DGP). In more detail, we generate data from the DGP used by Nie and Wager (2020):

$$X_{i} \sim \text{Unif}[0, 1]^{20}, \quad D_{i} | X_{i} \sim \text{Bernoulli} (p(X_{i})), \quad \epsilon_{i} \stackrel{\text{nu}}{\sim} \mathcal{N}(0, 1),$$

$$Y_{i} = b(X_{i}) + (D_{i} - 0.5) (X_{i,1} + X_{i,2}) + \sigma \epsilon_{i}.$$
(8)

The baseline main effect is the scaled Friedman (1991) function $b(X_i) = \sin(\pi X_{i,1}X_{i,2}) + 2(X_{i,3} - 0.5)^2 + X_{i,4} + 0.5X_{i,5}$. For the propensity score we follow Künzel et al. (2019) and set $p(X_i) = \alpha (1 + \beta_{2,4} (\min(X_{i,1}, X_{i,2})))$ where $\beta_{2,4}(\cdot)$ is the beta cumulative distribution function with shape parameters 2 and 4. The share of treated is $\mathbb{E}[D_i] = (31/21)\alpha$ and the true ATE is 1. The innovation standard deviation is set to $\sigma = 1$, and results for $\sigma = 5$ are relegated to Appendix B.

In the second strategy, we analyze the CU-DML estimator's performance using an Empirical Monte Carlo Study (EMCS) approach (Huber et al., 2013; Lechner and Wunsch, 2013). Our EMCS, following Knaus et al. (2022), uses a dataset from Lechner et al. (2020) of Swiss unemployed individuals in 2003. The dataset includes data on the effect of a job search program (treatment) on the number of months employed in the first six months after the start of the program (outcome), and 49 covariates providing information on individual characteristics of the unemployed, the regional employment agency, and regional labour market characteristics. For more data details, see Knaus et al. (2022). The EMCS proceeds as follows. We estimate the propensity score using a logistic regression on the entire sample of 91'339 unemployed individuals. Then, we define a new sample consisting of only the non-treated individuals whose fitted propensity score \hat{p}_i lies between 0.05 and 0.95. In each simulation, we draw a random sample with replacement of size N from this new sample and randomly assign treatments as $D_i = \mathbbm 1\{V_i < \hat{p}_i/\lambda\}$, where \hat{p}_i is the fitted propensity score from the logistic regression, $V_i \sim \text{Unif}[0, 1]$ and λ controls the share of treated $\mathbb{E}[D_i]$. The true ATE is 0 by construction.

⁵From Equation (4) it follows that $\tilde{\eta} < p(X) < 1 - \tilde{\eta}$ with $\tilde{\eta} = \gamma \eta / (1 - \eta + \gamma \eta)$.

We compare the CU-DML estimator with the DML estimator (Chernozhukov et al., 2018), and three popular adjustments of the DML estimator that are designed to improve its finite sample performance when the treatment assignment is unbalanced.⁶ The W-DML estimator uses winsorized propensity scores at 0.01 and 0.99 (Imbens, 2004). In the N-DML estimator we normalize the weights $w_i^{\mathcal{I}-k} := D_i/\hat{p}^{\mathcal{I}-k}(X_i)$ to sum to unity (e.g. Słoczyński and Wooldridge, 2018). Following Huber et al. (2013), the T-DML estimator normalizes the weights $w_i^{\mathcal{I}-k}$ and then truncates them to not exceed 0.04, before normalizing them again. Finally, we also consider the U-DML estimator, where the entire sample is undersampled and the DML estimator is applied to the undersampled data. All estimators use 5-fold crossfitting. The nuisance functions are estimated using random forests (Breiman, 2001) with 500 trees. The maximal depth of the trees and the minimal number of observations in the leaf nodes are determined by 5-fold cross-validation over 20 simulation replications and set to the most frequently selected values.⁷ Details on the software used for the simulations are provided in Appendix B.1.

The results are presented in Table 1. The table reports the root mean squared error (RMSE), the absolute value of the average bias, the standard deviation, and the coverage of the 95% confidence interval for each ATE estimator. The simulation is repeated 1'000 times for each of the following tuples of sample size and share of treated: $(N, \mathbb{E}[D_i]) \in \{(2'000, 0.05), (2'000, 0.1), (4'000, 0.025), (4'000, 0.05), (4'000, 0.025), (8'000, 0.05), (8'000, 0.01)\}$. The case where N = 2'000 and $\mathbb{E}[D_i] = 0.025$ is excluded since we would only have 50 treated observations. The best performance in terms of RMSE in each simulation is highlighted in bold. For the synthetic DGP (Panel A), the lowest RMSE is achieved by the CU-DML estimator in 5 out of 8 settings. Only when at least 400 observations are treated, the CU-DML estimator is outperformed in terms of RMSE by the normalized and trimmed DML estimators. However, the RMSE differences between N-DML, T-DML, and CU-DML estimators are considerably small in these cases. These results are unaffected by a higher innovation standard deviation $\sigma = 5$, see Table B1. For the EMCS (Panel B), CU-DML achieves the smallest RMSE for all combinations of $(N, \mathbb{E}[D_i])$. Also for the EMCS, we observe that the outperformance of the CU-DML estimator decreases for larger samples and/or less imbalanced samples. These patterns are in line with the findings from the machine learning classification literature, showing that class imbalanced-ness is a relative problem, related to both the degree of imbalancedness and the sample size (Japkowicz and Stephen, 2002).

The bias and standard deviation of the estimators unveil that, while the analyzed estimators do not differ substantially in terms of their bias, the CU-DML estimator has the lowest standard deviation across all simulation strategies and settings. The simulation study also confirms the theoretical result presented in the previous section. First, the coverage of the proposed estimator is in all strategies and settings close to 95%. Second, its RMSE decreases at rate \sqrt{N} .

4 Conclusion

This paper addresses a common finite sample problem of the double-machine learning ATE estimator (DML) (Chernozhukov et al., 2018): in settings with unbalanced treatment assignment estimations of the propensity scores become either too close to zero or one. This causes the DML estimator to become unstable, especially in small samples. We propose a simple yet effective adjustment of the DML estimator (CU-DML) where the machine learning models for the propensity scores are estimated over an undersampled dataset. The resulting propensity score predictions are then calibrated to adjust for the undersampling.

We provide theoretical results for the CU-DML estimator, showing that it has the same asymptotic distribution as the DML estimator. In particular, the estimator attains the parametric rate of convergence \sqrt{N} and its variance achieves the semi-parametric efficiency bound (Hahn, 1998). Furthermore, a small simulation study provides evidence for the finite sample performance of the proposed estimator, showing that it is of particular use in settings with highly unbalanced treatment assignments or small samples.

Future research could adapt the proposed approach to other estimators that rely on the estimation of the propensity score, such as the inverse probability-weighted estimator. Furthermore, CU-DML could be extended to estimators of the conditional average treatment effect (Fan et al., 2022; Zimmert and Lechner, 2019).

⁶Over the past two decades, these adjustments have been primarily employed to improve the finite sample performance of the inverse-probability-weighted estimator when the treatment assignment is unbalanced; see, among others, the extensive simulation study of Huber et al. (2013). More recently, these approaches have also been employed to improve the performance of the DML estimator.

⁷Properely tuning the hyperparameters of the random forests is crucial, especially for DML when the treatment assignment is unbalanced. The popular Python implementation of random forest, scikit-learn (Pedregosa et al., 2011), sets the default minimal number of observations in the leaf nodes to 1. This may lead to highly unstable propensity scores when the treatment assignment is unbalanced. See Bach et al. (2024) for a recent simulation study on the importance of hyperparameter-tuning for the DML estimator.

Panel A: Synthetic DGP													
	N =	2000, E[$D_i] = 2.5\%$	$\delta, \sigma = 1$	N =	4000, E[$[D_i] = 2.5\%$	$\delta, \sigma = 1$	N =	8000, E	$[D_i] = 2.5\%$	$\delta, \sigma = 1$	
	RMSE	Bias	Std. dev.	Coverage	RMSE	Bias	Std. dev.	Coverage	RMSE	Bias	Std. dev.	Coverage	
DML	-	-	-	-	0.188	0.008	0.188	0.984	0.094	0.005	0.094	0.979	
U-DML	-	-	-	-	0.173	0.070	0.158	0.928	0.122	0.057	0.108	0.924	
CU-DML	-	-	-	-	0.129	0.038	0.123	0.938	0.086	0.018	0.084	0.953	
W-DML	-	-	-	-	0.145	0.031	0.142	0.971	0.091	0.010	0.091	0.978	
N-DML	-	-	-	-	0.149	0.028	0.146	0.937	0.089	0.012	0.088	0.943	
T-DML	-	-	-	-	0.145	0.035	0.141	0.942	0.089	0.012	0.088	0.943	
	N =	$D_i] = 5.0\%$	$\delta, \sigma = 1$	$N = 4000, \mathbb{E}[D_i] = 5.0\%, \sigma = 1$				$N = 8000, \mathbb{E}[D_i] = 5.0\%, \sigma = 1$					
	RMSE	Bias	Std. dev.	Coverage	RMSE	Bias	Std. dev.	Coverage	RMSE	Bias	Std. dev.	Coverage	
DML	0.160	0.022	0.158	0.985	0.089	0.009	0.089	0.976	0.060	0.002	0.060	0.962	
U-DML	0.176	0.071	0.161	0.929	0.118	0.051	0.107	0.937	0.091	0.047	0.077	0.910	
CU-DML	0.132	0.039	0.126	0.946	0.086	0.017	0.084	0.960	0.059	0.008	0.059	0.950	
W-DML	0.156	0.023	0.155	0.985	0.089	0.009	0.089	0.976	0.060	0.002	0.060	0.962	
N-DML	0.143	0.035	0.138	0.934	0.086	0.014	0.085	0.957	0.059	0.004	0.059	0.940	
T-DML	0.143	0.036	0.138	0.935	0.086	0.014	0.085	0.956	0.059	0.004	0.059	0.940	
	$N = 2000, \mathbb{E}[D_i] = 10.0\%, \sigma = 1$					$N = 4000, \mathbb{E}[D_i] = 10.0\%, \sigma = 1$				$N = 8000, \mathbb{E}[D_i] = 10.0\%, \sigma = 1$			
	RNDE	Dias	Stu. ucv.	Coverage	KNISL	Dias	Siu. ucv.	Coverage	RNDL	Dias	Siu. ucv.	Coverage	
DML	0.097	0.016	0.095	0.952	0.060	0.009	0.060	0.961	0.043	0.002	0.043	0.943	
U-DML	0.126	0.057	0.113	0.924	0.088	0.046	0.075	0.915	0.069	0.042	0.055	0.867	
CU-DML	0.094	0.022	0.091	0.947	0.062	0.013	0.060	0.954	0.044	0.005	0.043	0.928	
W-DML	0.097	0.016	0.095	0.952	0.060	0.009	0.060	0.961	0.043	0.002	0.043	0.943	
N-DML	0.095	0.020	0.093	0.924	0.060	0.010	0.059	0.950	0.043	0.003	0.043	0.932	
T-DML	0.095	0.020	0.093	0.924	0.060	0.010	0.059	0.950	0.043	0.003	0.043	0.932	
Panel B: Empirical Monte Carlo Study													
	N	7 <u>- 2000</u>						-	$N = 8000 \mathbb{E}[D_i] = 2.5\%$				
				5%	Λ	I = 4000	$\mathbb{E}[D_i] = 2$	5%	Δ	I = 8000) $\mathbb{E}[D_i] = 2$	2.5%	
	RMSE	Bias	$\mathbb{E}[D_i] = 2$ Std. dev.	2.5% Coverage	N RMSE	I = 4000 Bias	$\mathbb{E}[D_i] = 2$ Std. dev.	2.5% Coverage	۸ RMSE	V = 8000 Bias	$\mathbb{E}[D_i] = 2$ Std. dev.	2.5% Coverage	
	RMSE	= 2000 Bias	$\mathbb{E}[D_i] = 2$ Std. dev.	2.5% Coverage	RMSE	V = 4000 Bias	$\mathbb{E}[D_i] = 2$ Std. dev.	2.5% Coverage	RMSE	V = 8000 Bias	$\mathbb{E}[D_i] = 2$ Std. dev.	2.5% Coverage	
DML	RMSE -	Bias	$[D_i] = 2$ Std. dev.	Coverage	RMSE	V = 4000 Bias 0.051 0.024	$\mathbb{E}[D_i] = 2$ Std. dev. 0.309 0.260	0.984	0.194	V = 8000 Bias 0.041 0.012	$\mathbb{E}[D_i] = 2$ Std. dev. 0.189 0.102	2.5% Coverage 0.975	
DML U-DML	RMSE	Bias	$\mathbb{E}[D_i] = 2$ Std. dev.	2.5% Coverage - -	RMSE 0.314 0.261 0.200	V = 4000 Bias 0.051 0.024	$\mathbb{E}[D_i] = 2$ Std. dev. 0.309 0.260 0.200	2.5% Coverage 0.984 0.950	0.194 0.183	V = 8000 Bias 0.041 0.013 0.022	$\mathbb{E}[D_i] = 2$ Std. dev. 0.189 0.183 0.146	2.5% Coverage 0.975 0.962	
DML U-DML CU-DML	RMSE - -	Bias	$\mathbb{E}[D_i] = 2$ Std. dev.	Coverage - -	N RMSE 0.314 0.261 0.208	V = 4000 Bias 0.051 0.024 0.016 0.014	$D_{i} \mathbb{E}[D_{i}] = 2$ Std. dev. 0.309 0.260 0.208 0.208	2.5% Coverage 0.984 0.950 0.946 0.970	A RMSE 0.194 0.183 0.148 0.148	V = 8000 Bias 0.041 0.013 0.023 0.051	$D_{i} \mathbb{E}[D_{i}] = 2$ Std. dev. 0.189 0.183 0.146 0.154	2.5% Coverage 0.975 0.962 0.944 0.941	
DML U-DML CU-DML W-DML	RMSE - - -	Bias - - - -	$[E[D_i] = 2$ Std. dev.	2.5% Coverage - - - -	N RMSE 0.314 0.261 0.208 0.233 0.252	V = 4000 Bias 0.051 0.024 0.016 0.044 0.057	$\begin{array}{c} \text{D}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \end{array}$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948	N RMSE 0.194 0.183 0.148 0.162 0.174	V = 8000 Bias 0.041 0.013 0.023 0.051 0.050	$D_{i} \mathbb{E}[D_{i}] = 2$ Std. dev. 0.189 0.183 0.146 0.154 0.166	2.5% Coverage 0.975 0.962 0.944 0.961	
DML U-DML CU-DML W-DML N-DML	RMSE - - - -	Bias	$\int_{i} \mathbb{E}[D_i] = 2$ Std. dev.	2.5% Coverage - - - - -	N RMSE 0.314 0.261 0.208 0.233 0.253	V = 4000 Bias 0.051 0.024 0.016 0.044 0.057 0.051	$\begin{array}{c} 0, \mathbb{E}[D_i] = 2\\ \hline 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.221 \end{array}$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.927	N RMSE 0.194 0.183 0.148 0.162 0.174 0.167	V = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052	$\begin{array}{c} \text{0.189} \\ \hline 0.189 \\ \hline 0.183 \\ 0.146 \\ 0.154 \\ 0.166 \\ 0.150 \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.928	
DML U-DML CU-DML W-DML N-DML T-DML	RMSE	Bias	$\mathbb{E}[D_i] = 2$ Std. dev.	2.5% Coverage - - - - - - -	N RMSE 0.314 0.261 0.208 0.233 0.253 0.237	V = 4000 Bias 0.051 0.024 0.016 0.044 0.057 0.051 $V = 4000$	$D_{i} \mathbb{E}[D_{i}] = 2$ Std. dev. 0.309 0.260 0.208 0.228 0.246 0.231 $\mathbb{E}[D_{i}] = 5$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937	A RMSE 0.194 0.183 0.148 0.162 0.174 0.167	V = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052 $V = 8000$	$\mathbb{E}[D_i] = 2$ Std. dev. 0.189 0.183 0.146 0.154 0.166 0.159 $\mathbb{E}[D_i] = 5$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938	
DML U-DML CU-DML W-DML N-DML T-DML	RMSE - - - - - N RMSE	$\frac{1}{2000}$ Bias	$\mathbb{E}[D_i] = 2$ Std. dev.	2.5% Coverage - - - - - - - - - - - - - - - - - - -	N RMSE 0.314 0.261 0.208 0.233 0.253 0.237	V = 4000 Bias 0.051 0.024 0.016 0.044 0.057 0.051 $V = 4000$ Bias	$\begin{array}{l} (b) & \mathbb{E}[D_i] = 2\\ & \text{Std. dev.} \end{array} \\ \hline \\ \hline \\ 0.309 \\ 0.260 \\ 0.208 \\ 0.228 \\ 0.246 \\ 0.231 \\ 0. \mathbb{E}[D_i] = 5\\ & \text{Std. dev.} \end{array}$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937 5.0% Coverage	A RMSE 0.194 0.183 0.148 0.162 0.174 0.167 A RMSE	7 = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052 $7 = 8000$ Bias	$\begin{array}{l} \text{(b)} \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.189\\ 0.183\\ 0.146\\ 0.154\\ 0.166\\ 0.159 \\ \hline \\ \text{(b)} \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938 5.0% Coverage	
DML U-DML CU-DML W-DML N-DML T-DML	RMSE - - - - - - - - - - - - - - - - - - -	$\frac{1}{2000}$ Bias $\frac{1}{1000}$ $\frac{1}{1000}$ Bias $\frac{1}{1000}$ Bias	$\mathbb{E}[D_i] = 2$ Std. dev. $\mathbb{E}[D_i] = 5$ Std. dev. 0.252	2.5% Coverage - - - - - - - - - - - - - - - - - - -	N RMSE 0.314 0.261 0.208 0.233 0.253 0.237 N RMSE 0.172	V = 4000 Bias 0.051 0.024 0.016 0.044 0.057 0.051 $V = 4000$ Bias 0.043	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.231 \\ \hline \\ \text{0}, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.167 \\ \hline \end{array}$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937 5.0% Coverage 0.962	A RMSE 0.194 0.183 0.148 0.162 0.174 0.167 RMSE 0.111	7 = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052 7 = 8000 Bias 0.032	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.189\\ 0.183\\ 0.146\\ 0.154\\ 0.166\\ 0.159 \\ \hline \\ \text{0}, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.107 \\ \hline \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938 5.0% Coverage 0.959	
DML U-DML CU-DML W-DML N-DML T-DML DML	RMSE - - - - - - - - - - - - - - - - - - -	V = 2000 Bias V = 2000 Bias 0.049 0.012	$\mathbb{E}[D_i] = 2$ Std. dev. $\frac{1}{2}$ $\mathbb{E}[D_i] = 5$ Std. dev. $\frac{1}{2}$ $\mathbb{E}[D_i] = 5$ Std. dev. $\frac{1}{2}$ $\mathbb{E}[D_i] = 5$ $\mathbb{E}[D_i] = 5$ $\mathbb{E}[D_i] = 5$ $\mathbb{E}[D_i] = 5$	2.5% Coverage - - - - - - - - - - - - - - - - - - -	N RMSE 0.314 0.261 0.208 0.233 0.253 0.237 N RMSE 0.172 0.180	V = 4000 Bias 0.051 0.024 0.016 0.044 0.057 0.051 $V = 4000$ Bias 0.043 0.012	$\begin{array}{c} (i), \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.231\\ 0, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline 0.167\\ 0.179\\ \end{array}$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937 6.0% Coverage 0.962 0.962	A RMSE 0.194 0.183 0.148 0.162 0.174 0.167 A RMSE 0.111 0.129	7 = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052 7 = 8000 Bias 0.032 0.022	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.189\\ 0.183\\ 0.146\\ 0.154\\ 0.166\\ 0.159 \\ \hline \\ \text{0}, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.107\\ 0.127 \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.940 0.940 0.938 5.0% Coverage 0.959 0.948	
DML U-DML CU-DML W-DML N-DML T-DML DML U-DML CU-DML	RMSE - - - - - - - - - - - - - - - - - - -	$\frac{1}{10000} = \frac{10000}{10000}$	$\mathbb{E}[D_i] = 2$ Std. dev. $\mathbb{E}[D_i] = 5$ Std. dev. 0.252 0.261 0.213	2.5% Coverage - - - - - - - - - - - - - - - - - - -	A RMSE 0.314 0.261 0.208 0.233 0.253 0.237 A RMSE 0.172 0.180 0.143	$\begin{array}{c} I = 4000\\ \text{Bias} \\ \hline 0.051\\ 0.024\\ 0.016\\ 0.044\\ 0.057\\ 0.051 \\ \hline I = 4000\\ \text{Bias} \\ \hline 0.043\\ 0.012\\ 0.023 \\ \hline \end{array}$	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.231\\ 0, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.167\\ 0.179\\ 0.141\\ \end{array}$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937 5.0% Coverage 0.962 0.953 0.943	A RMSE 0.194 0.183 0.148 0.162 0.174 0.167 A RMSE 0.111 0.129 0.101	V = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052 V = 8000 Bias 0.032 0.022 0.022	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.189\\ 0.183\\ 0.146\\ 0.154\\ 0.166\\ 0.159 \\ \hline \\ \text{Std. dev.} \\ \hline \\ 0.107\\ 0.127\\ 0.099 \\ \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938 5.0% Coverage 0.959 0.948 0.941	
DML U-DML CU-DML W-DML N-DML T-DML DML U-DML U-DML W-DML	RMSE - - - - - - - - - - - - - - - - - - -	V = 2000 Bias 	$\mathbb{E}[D_i] = 2$ Std. dev. $[\mathbb{E}[D_i] = 5$ Std. dev. 0.252 0.261 0.213 0.239	2.5% Coverage - - - - - - - - - - - - - - - - - - -	N RMSE 0.314 0.261 0.208 0.233 0.253 0.237 N RMSE 0.172 0.180 0.143	7 = 4000Bias 0.051 0.024 0.016 0.044 0.057 0.051 7 = 4000 Bias 0.043 0.012 0.023 0.043	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.231\\ 0.228\\ 0.246\\ 0.231\\ 0.228\\ 0.246\\ 0.231\\ 0.216\\ 0.171\\ 0.158\\ \end{array}$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937 6.0% Coverage 0.962 0.953 0.943 0.943	A RMSE 0.194 0.183 0.162 0.174 0.167 A RMSE 0.111 0.129 0.101	7 = 8000Bias 0.041 0.013 0.023 0.051 0.050 0.052 7 = 8000 Bias 0.032 0.022 0.023 0.023	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.189\\ 0.183\\ 0.146\\ 0.154\\ 0.166\\ 0.159 \\ \text{0}, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.107\\ 0.127\\ 0.099\\ 0.103 \\ \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938 5.0% Coverage 0.959 0.948 0.941 0.957	
DML U-DML CU-DML W-DML N-DML T-DML U-DML CU-DML CU-DML W-DML N-DMI	RMSE 	V = 2000 Bias 	$\mathbb{E}[D_i] = 2$ Std. dev. $\mathbb{E}[D_i] = 5$ Std. dev. 0.252 0.261 0.213 0.239 0.226	2.5% Coverage - - - - - - - - - - - - - - - - - - -	N RMSE 0.314 0.261 0.208 0.233 0.253 0.237 N RMSE 0.172 0.180 0.143 0.164 0.164	7 = 4000 Bias 0.051 0.024 0.016 0.044 0.057 0.051 7 = 4000 Bias 0.043 0.012 0.023 0.044 0.048	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.231\\ 0, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.167\\ 0.179\\ 0.141\\ 0.158\\ 0.156\\ \end{array}$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937 6.0% Coverage 0.962 0.953 0.943 0.962 0.962	∧ RMSE 0.194 0.183 0.148 0.162 0.174 0.167 ∧ RMSE 0.111 0.129 0.101 0.108 0.108	7 = 8000Bias 0.041 0.013 0.023 0.051 0.050 0.052 7 = 8000 Bias 0.032 0.022 0.022 0.023 0.033 0.033	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.189\\ 0.183\\ 0.146\\ 0.154\\ 0.166\\ 0.159 \\ \hline \\ \text{0}, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.107\\ 0.127\\ 0.099\\ 0.103\\ 0.104 \\ \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938 5.0% Coverage 0.959 0.948 0.957 0.941	
DML U-DML CU-DML W-DML N-DML T-DML U-DML U-DML W-DML N-DML T-DML	RMSE - - - - - - - - - - - - - - - - - - -	V = 2000 Bias $V = 2000$ Bias 0.049 0.012 0.015 0.056 0.056	$\mathbb{E}[D_i] = 2$ Std. dev. $\mathbb{E}[D_i] = 5$ Std. dev. 0.252 0.261 0.213 0.226 0.219	2.5% Coverage - - - - - - - - - - - - - - - - - - -	N RMSE 0.314 0.261 0.208 0.233 0.237 N RMSE 0.172 0.180 0.143 0.164 0.163	7 = 4000 Bias 0.051 0.024 0.016 0.044 0.057 0.051 7 = 4000 Bias 0.043 0.012 0.023 0.044 0.048 0.048	$\begin{array}{l} \text{(b)} \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.231\\ 0.228\\ \text{Std. dev.} \\ \hline \\ \text{Std. dev.} \\ \hline \\ 0.167\\ 0.179\\ 0.141\\ 0.158\\ 0.156\\ 0.154\\ \end{array}$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937 6.0% Coverage 0.962 0.953 0.943 0.962 0.941	∧ RMSE 0.194 0.183 0.148 0.162 0.174 0.167 ∧ RMSE 0.111 0.129 0.101 0.108 0.110	7 = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052 7 = 8000 Bias 0.032 0.022 0.023 0.033 0.036	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.189\\ 0.183\\ 0.146\\ 0.154\\ 0.166\\ 0.159 \\ \hline \\ \text{0}, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.107\\ 0.127\\ 0.099\\ 0.103\\ 0.104\\ 0.104 \\ \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938 5.0% Coverage 0.959 0.948 0.957 0.941 0.957	
DML U-DML CU-DML N-DML T-DML U-DML U-DML CU-DML W-DML N-DML T-DML	RMSE - - - - - - - - - - - - - - - - - - -	V = 2000 Bias $V = 2000$ Bias 0.049 0.012 0.015 0.051 0.056 0.056 0.056	$\mathbb{E}[D_i] = 2$ Std. dev. $\mathbb{E}[D_i] = 5$ Std. dev. $\mathbb{E}[D_i] = 5$ Std. dev. 0.252 0.261 0.213 0.239 0.226 0.219	2.5% Coverage - - - - - - - - - - - - - - - - - - -	N RMSE 0.314 0.261 0.208 0.233 0.237 N RMSE 0.172 0.180 0.143 0.164 0.163 0.161	7 = 4000 Bias 0.051 0.024 0.016 0.044 0.057 0.051 7 = 4000 Bias 0.043 0.012 0.023 0.044 0.048 0.048 0.048 0.048	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.231\\ 0, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.167\\ 0.179\\ 0.141\\ 0.158\\ 0.156\\ 0.154\\ \hline \end{array}$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937 Coverage 0.962 0.953 0.943 0.962 0.953 0.943 0.962	∧ RMSE 0.194 0.183 0.148 0.162 0.174 0.167 ∧ RMSE 0.111 0.129 0.101 0.108 0.110 0.110	7 = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052 7 = 8000 Bias 0.032 0.022 0.023 0.033 0.036 0.036	$\begin{array}{c} \text{(b)} \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.189\\ 0.183\\ 0.146\\ 0.154\\ 0.166\\ 0.159 \\ \hline \\ \text{Std. dev.} \\ \hline \\ 0.107\\ 0.127\\ 0.099\\ 0.103\\ 0.104\\ 0.104 \\ \hline \\ \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938 5.0% Coverage 0.959 0.948 0.941 0.957 0.941 0.941	
DML U-DML CU-DML W-DML T-DML T-DML U-DML CU-DML N-DML N-DML T-DML	RMSE - - - - - - - - - - - - - - - - - - -	$\frac{1}{10000} = \frac{10000}{10000}$	$\mathbb{E}[D_i] = 1$ Std. dev. $\mathbb{E}[D_i] = 5$ Std. dev. $\mathbb{E}[D_i] = 5$ Std. dev. $\mathbb{E}[D_i] = 1$ Std. dev.	2.5% Coverage - - - - - - - - - - - - - - - - - - -	N RMSE 0.314 0.261 0.208 0.233 0.237 N RMSE 0.172 0.180 0.143 0.164 0.164 0.161 N RMSE	7 = 4000 Bias 0.051 0.024 0.044 0.057 0.051 7 = 4000 Bias 0.043 0.012 0.023 0.044 0.048 0.048 = 4000 Bias	$\begin{array}{l} (\mathcal{B}[D_i]=2\\ \mathrm{Std.~dev.}\\ \hline 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.231\\ 0.\mathbb{E}[D_i]=5\\ \mathrm{Std.~dev.}\\ \hline 0.167\\ 0.179\\ 0.141\\ 0.158\\ 0.156\\ 0.154\\ 0.154\\ \mathbf{E}[D_i]=1\\ \mathrm{Std.~dev.}\\ \end{array}$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937 6.0% Coverage 0.962 0.953 0.943 0.962 0.941 0.941 0.941	∧ RMSE 0.194 0.183 0.148 0.162 0.174 0.167 ∧ RMSE 0.111 0.129 0.101 0.108 0.110 ∧	7 = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052 7 = 8000 Bias 0.032 0.022 0.023 0.036 0.036 = 8000 Bias	$\begin{array}{c} \text{(b)} & \mathbb{E}[D_i] = 2\\ & \text{Std. dev.} \end{array} \\ \hline \\ & 0.189\\ & 0.183\\ & 0.146\\ & 0.154\\ & 0.166\\ & 0.159 \end{array} \\ \text{(b)} & \mathbb{E}[D_i] = 5\\ & \text{Std. dev.} \end{array} \\ \hline \\ & 0.107\\ & 0.107\\ & 0.107\\ & 0.109\\ & 0.104\\ & 0.104\\ & 0.104 \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938 5.0% Coverage 0.959 0.948 0.941 0.957 0.941 0.941 0.941	
DML U-DML CU-DML W-DML T-DML T-DML U-DML CU-DML N-DML N-DML T-DML	RMSE 	V = 2000 Bias 0.049 0.012 0.015 0.056 0.056 0.056 0.056 0.056 0.033	$\mathbb{E}[D_i] = 2$ Std. dev. $\mathbb{E}[D_i] = 5$ Std. dev. 0.252 0.261 0.213 0.239 0.226 $0.219\mathbb{E}[D_i] = 1 Std. dev.0.153$	2.5% Coverage - - - - - - - - - - - - - - - - - - -	N RMSE 0.314 0.261 0.208 0.233 0.237 N RMSE 0.172 0.180 0.143 0.164 0.164 0.161 N RMSE	7 = 4000 Bias 0.051 0.024 0.044 0.057 0.051 7 = 4000 Bias 0.043 0.012 0.023 0.044 0.048 = 4000 Bias	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.231\\ 0.228\\ 0.246\\ 0.231\\ 0.231\\ 0.228\\ 0.246\\ 0.231\\ 0.246\\ 0.231\\ 0.167\\ 0.179\\ 0.141\\ 0.158\\ 0.156\\ 0.154\\ 0.154\\ 0.154\\ 0.154\\ 0.154\\ 0.154\\ 0.155\\ 0.154\\ 0.155\\ 0.154\\ 0.155\\ 0.154\\ 0.155\\ 0.154\\ 0.155\\ 0.154\\ 0.155\\ 0.154\\ 0.155\\ 0.154\\ 0.155\\ 0.154\\ 0.155\\ 0.155\\ 0.154\\ 0.155\\ 0.15\\$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937 6.0% Coverage 0.962 0.953 0.943 0.962 0.941 0.941 0.0% Coverage	N RMSE 0.194 0.183 0.148 0.162 0.174 0.167 N RMSE 0.111 0.129 0.101 0.108 0.110 N RMSE	7 = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052 7 = 8000 Bias 0.032 0.022 0.023 0.036 0.036 = 8000 Bias 0.014	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.189\\ 0.183\\ 0.146\\ 0.154\\ 0.166\\ 0.159 \\ \hline \\ \text{0}, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.107\\ 0.127\\ 0.099\\ 0.103\\ 0.104\\ 0.104 \\ \hline \\ \text{Std. dev.} \\ \hline \\ \mathbb{E}[D_i] = 1\\ \text{Std. dev.} \\ \hline \\ 0.071 \\ \hline \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938 5.0% Coverage 0.959 0.948 0.941 0.957 0.941 0.954 0.941 0.954 0.941	
DML U-DML CU-DML W-DML N-DML T-DML U-DML CU-DML W-DML T-DML T-DML T-DML	RMSE - - - - - - - - - - - - - - - - - - -	7 = 2000 Bias $$	$\mathbb{E}[D_i] = 1$ Std. dev. $\mathbb{E}[D_i] = 5$ Std. dev. $\mathbb{E}[D_i] = 5$ Std. dev. 0.252 0.261 0.213 0.239 0.226 0.213 0.239 0.226 $0.219\mathbb{E}[D_i] = 1 Std. dev.0.153$ 0.183	2.5% Coverage 	N RMSE 0.314 0.261 0.208 0.233 0.237 N RMSE 0.172 0.180 0.143 0.164 0.163 0.164 0.163 0.164 0.163 0.164 0.163 0.164 0.163 0.164 0.163 0.164 0.163 0.164 0.163 0.164 0.163 0.164 0.163 0.164 0.164 0.163 0.164 0.167	7 = 4000 Bias 0.051 0.024 0.016 0.044 0.057 0.051 7 = 4000 Bias 0.043 0.012 0.023 0.044 0.048 0.048 0.048 0.048 0.048 0.024 0.017	$\begin{array}{l} \text{D}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.231\\ 0.228\\ 0.246\\ 0.231\\ 0.231\\ 0.246\\ 0.231\\ 0.246\\ 0.231\\ 0.246\\ 0.231\\ 0.246\\ 0.231\\ 0.246\\ 0.151\\ 0.154\\ 0.156\\ 0.154\\ 0.155\\ 0.154\\ 0.105\\ 0.127\\$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937 6.0% Coverage 0.962 0.953 0.943 0.962 0.941 0.941 0.941 0.941 0.945	N RMSE 0.194 0.183 0.148 0.162 0.174 0.167 N RMSE 0.111 0.129 0.101 0.108 0.110 0.110 0.110 0.110 0.110 0.102 0.103	7 = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052 7 = 8000 Bias 0.032 0.033 0.036 0.036 0.036 0.036 0.036 0.036 0.036 0.034 0.014	$\begin{array}{l} \text{D}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.189\\ 0.183\\ 0.146\\ 0.154\\ 0.166\\ 0.159 \\ \hline \\ \text{D}, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.107\\ 0.127\\ 0.099\\ 0.103\\ 0.104\\ 0.104 \\ \hline \\ \text{Std. dev.} \\ \hline \\ \mathbb{E}[D_i] = 1\\ \text{Std. dev.} \\ \hline \\ 0.071\\ 0.087 \\ \hline \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938 5.0% Coverage 0.959 0.948 0.941 0.957 0.941 0.941 0.941 0.941 0.941 0.941 0.941	
DML U-DML CU-DML W-DML T-DML T-DML U-DML CU-DML W-DML T-DML T-DML U-DML CU-DML CU-DML	RMSE - - - - - - - - - - - - - - - - - - -	$\begin{array}{c} -2000\\ \text{Bias} \\ \hline \\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -$	$\mathbb{E}[D_i] = 1$ Std. dev. $\mathbb{E}[D_i] = 5$ Std. dev. 0.252 0.261 0.213 0.239 0.226 $0.219\mathbb{E}[D_i] = 1 Std. dev.0.153$ 0.183 0.183	2.5% Coverage - - - - - - - - - - - - - - - - - - -	N RMSE 0.314 0.261 0.203 0.233 0.233 0.233 0.233 0.233 0.233 0.233 0.233 0.233 0.233 0.233 0.233 0.233 0.172 0.180 0.161 N RMSE 0.107 0.128 0.007	7 = 4000 Bias 0.051 0.024 0.016 0.044 0.057 0.051 7 = 4000 Bias 0.043 0.012 0.023 0.044 0.048 0.048 0.048 = 4000 Bias 0.024 0.017 0.016	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.231\\ 0.228\\ 0.246\\ 0.231\\ 0.228\\ 0.246\\ 0.231\\ 0.218\\ 0.167\\ 0.179\\ 0.141\\ 0.158\\ 0.167\\ 0.158\\ 0.156\\ 0.154\\ 0.154\\ 0.165\\ 0.127\\ 0.098\\ 0.098\\ 0.098\\ 0.098\\ 0.008\\$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937 6.0% Coverage 0.962 0.953 0.943 0.943 0.941 0.941 0.941 0.955 0.950	N RMSE 0.194 0.183 0.162 0.174 0.167 N RMSE 0.111 0.129 0.101 0.108 0.110 N RMSE 0.0101 0.0102 0.0101 0.0101 0.0102 0.0072 0.0088 0.0628	7 = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052 7 = 8000 Bias 0.032 0.022 0.023 0.036 0.036 = 8000 Bias 0.014 0.014 0.014 0.014	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.189\\ 0.183\\ 0.146\\ 0.154\\ 0.166\\ 0.159 \\ \hline \\ \text{0}, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.107\\ 0.127\\ 0.099\\ 0.103\\ 0.104\\ 0.104\\ \hline \\ \text{0}, \mathbb{E}[D_i] = 1\\ \text{Std. dev.} \\ \hline \\ 0.071\\ 0.087\\ 0.068 \\ \hline \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938 5.0% Coverage 0.959 0.948 0.941 0.957 0.941 0.941 0.941 0.941 0.941	
DML U-DML CU-DML W-DML T-DML T-DML U-DML U-DML W-DML T-DML U-DML U-DML U-DML U-DML U-DML W-DMI	RMSE - - - - - - - - - - - - - - - - - - -	7 = 2000 Bias $7 = 2000$ Bias 0.049 0.012 0.056 0.056 0.056 0.056 0.056 0.033 0.016 0.023 0.033	$\mathbb{E}[D_i] = 1$ Std. dev. $[D_i] = 1$ Std. dev. $[D_i] = 1$ Std. dev. [0.252] $[0.261]$ $[0.213]$ $[0.239]$ $[0.226]$ $[0.219]\mathbb{E}[D_i] = 1 Std. dev.[0.153]$ $[0.153]$ $[0.144]$ $[0.152]$	2.5% Coverage 	N RMSE 0.314 0.261 0.208 0.233 0.233 0.237 N RMSE 0.172 0.180 0.143 0.164 0.163 0.161 N RMSE 0.107 0.128 0.099 0.109	$\begin{array}{c} & I = 4000\\ \text{Bias} \\ 0.051\\ 0.024\\ 0.016\\ 0.044\\ 0.057\\ 0.051 \\ \hline I = 4000\\ \text{Bias} \\ 0.043\\ 0.012\\ 0.023\\ 0.044\\ 0.048\\ 0.048\\ \hline I = 4000\\ \text{Bias} \\ \hline 0.024\\ 0.017\\ 0.016\\ 0.023 \\ \hline \end{array}$	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.231\\ 0.228\\ 0.246\\ 0.231\\ 0.228\\ 0.246\\ 0.231\\ 0.228\\ 0.167\\ 0.179\\ 0.141\\ 0.158\\ 0.166\\ 0.154\\ 0.156\\ 0.154\\ \text{Std. dev.} \\ \hline 0.105\\ 0.127\\ 0.098\\ 0.104\\ \end{array}$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937 6.0% Coverage 0.962 0.953 0.943 0.962 0.941 0.941 0.941 0.941 0.941	N RMSE 0.194 0.183 0.162 0.174 0.167 N RMSE 0.111 0.129 0.101 0.100 0.101 0.110 0.101 0.102 0.002 0.0052 0.0052	7 = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052 7 = 8000 Bias 0.032 0.032 0.033 0.036 0.036 0.036 = 80000 Bias 0.014 0.014 0.014 0.014 0.014	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.189\\ 0.183\\ 0.146\\ 0.154\\ 0.166\\ 0.159 \\ \hline \\ \text{0}, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.107\\ 0.127\\ 0.099\\ 0.103\\ 0.104\\ 0.104\\ \hline \\ \text{Std. dev.} \\ \hline \\ 0.071\\ 0.087\\ 0.068\\ 0.071 \\ \hline \\ \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938 5.0% Coverage 0.959 0.948 0.941 0.957 0.941 0.941 0.941 0.941 0.941	
DML U-DML CU-DML W-DML T-DML T-DML U-DML CU-DML N-DML T-DML U-DML CU-DML CU-DML CU-DML CU-DML CU-DML CU-DML N-DMI N-DMI	RMSE 	$\begin{array}{c} -2000\\ \text{Bias} \\ \hline \\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -$	$\mathbb{E}[D_i] = 1$ Std. dev. $[D_i] = 1$ Std. dev.	2.5% Coverage 	N RMSE 0.314 0.261 0.208 0.233 0.237 N RMSE 0.172 0.180 0.143 0.163 0.161 N RMSE 0.107 0.128 0.099 0.106 0.106	7 = 4000 Bias 0.051 0.024 0.016 0.044 0.057 0.051 $7 = 4000$ Bias 0.043 0.012 0.023 0.044 0.048 0.048 0.048 0.048 0.048 0.024 0.017 0.016 0.023 0.024 0.023 0.024	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.231\\ 0, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline 0.167\\ 0.179\\ 0.141\\ 0.158\\ 0.156\\ 0.154\\ 0.156\\ 0.154\\ \text{Std. dev.} \\ \hline 0.105\\ 0.127\\ 0.098\\ 0.104\\ 0.103\\ \end{array}$	2.5% Coverage 0.984 0.950 0.948 0.979 0.948 0.937 6.0% Coverage 0.962 0.953 0.943 0.962 0.941 0.941 0.941 0.941 0.941 0.955 0.950 0.951 0.955	N RMSE 0.194 0.183 0.148 0.162 0.174 0.167 N RMSE 0.111 0.129 0.101 0.108 0.110 N RMSE 0.0072 0.0088 0.0072 0.072 0.072	7 = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052 7 = 8000 Bias 0.032 0.022 0.023 0.033 0.036 0.036 = 8000 Bias 0.014 0.014 0.012 0.014 0.016	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.189\\ 0.183\\ 0.146\\ 0.154\\ 0.166\\ 0.159 \\ \hline \\ 0, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.107\\ 0.127\\ 0.099\\ 0.103\\ 0.104\\ 0.104\\ \hline \\ \text{Std. dev.} \\ \hline \\ 0.071\\ 0.087\\ 0.068\\ 0.071\\ 0.078\\ 0.068\\ 0.071\\ 0.070 \\ \hline \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938 5.0% Coverage 0.959 0.948 0.941 0.957 0.941 0.941 0.941 0.941 0.941 0.941	
DML U-DML CU-DML W-DML T-DML U-DML CU-DML CU-DML N-DML T-DML U-DML CU-DML W-DML CU-DML W-DML CU-DML W-DML T-DMI	RMSE - - - - - - - - - - - - -	V = 2000 Bias $-$ $-$ $-$ Bias 0.049 0.012 0.015 0.051 0.056 0.056 0.056 0.056 0.056 0.033 0.016 0.020 0.033 0.037 0.03 0.03	$\mathbb{E}[D_i] = 2$ Std. dev. Std. dev. $\mathbb{E}[D_i] = 5$ Std. dev. 0.252 0.261 0.213 0.239 0.226 0.219 $\mathbb{E}[D_i] = 1$ Std. dev. $\mathbb{E}[D_i] = 1$ Std. dev. 0.153 0.149 0.153 0.149 0.149 0.153 0.149 0.149 0.149 0.149 0.153 0.149 0.153 0.149 0.153 0.153 0.153 0.149 0.153 0.153 0.149 0.153 0.149 0.153 0.153 0.149 0.153 0.149 0.153 0.153 0.149 0.153 0.153 0.154 0.155 0.149 0.153 0.154 0.155 0.155 0.155 0.155 0.155 0.155 0.155 0.155 0.155 0.155 0.149 0.155 0	2.5% Coverage - - - - - - - - - - - - - - - - - - -	N RMSE 0.314 0.261 0.208 0.233 0.237 N RMSE 0.172 0.180 0.143 0.164 0.164 0.161 N RMSE 0.107 0.128 0.0099 0.106 0.107 0.106	7 = 4000 Bias 0.051 0.024 0.016 0.044 0.057 0.051 7 = 4000 Bias 0.043 0.012 0.023 0.044 0.048 = 4000 Bias 0.024 0.017 0.016 0.023 0.024 0.025 0.025	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.309\\ 0.260\\ 0.208\\ 0.228\\ 0.246\\ 0.231\\ 0.228\\ 0.246\\ 0.231\\ 0.228\\ 0.246\\ 0.231\\ 0.231\\ 0.246\\ 0.231\\ 0.167\\ 0.179\\ 0.141\\ 0.158\\ 0.167\\ 0.179\\ 0.141\\ 0.158\\ 0.156\\ 0.154\\ 0.154\\ 0.154\\ 0.154\\ 0.103\\ 0.10$	2.5% Coverage 0.984 0.950 0.946 0.979 0.948 0.937 6.0% Coverage 0.962 0.953 0.943 0.962 0.941 0.941 0.941 0.0% Coverage 0.955 0.950 0.951	N RMSE 0.194 0.183 0.148 0.162 0.174 0.167 N RMSE 0.111 0.129 0.101 0.108 0.110 N RMSE 0.072 0.088 0.069 0.072 0.072 0.072 0.072	7 = 8000 Bias 0.041 0.013 0.023 0.051 0.050 0.052 7 = 8000 Bias 0.032 0.022 0.023 0.036 0.036 = 8000 Bias 0.014 0.014 0.012 0.014 0.016 0.016	$\begin{array}{c} \text{0}, \mathbb{E}[D_i] = 2\\ \text{Std. dev.} \\ \hline \\ 0.189\\ 0.183\\ 0.146\\ 0.154\\ 0.166\\ 0.159 \\ \hline \\ 0.159\\ 0, \mathbb{E}[D_i] = 5\\ \text{Std. dev.} \\ \hline \\ 0.107\\ 0.127\\ 0.099\\ 0.103\\ 0.104\\ 0.104\\ \hline \\ \text{Std. dev.} \\ \hline \\ 0.071\\ 0.087\\ 0.068\\ 0.071\\ 0.070\\ 0.070 \\ \hline \end{array}$	2.5% Coverage 0.975 0.962 0.944 0.961 0.940 0.938 5.0% Coverage 0.959 0.948 0.941 0.957 0.941 0.951 0.960 0.952 0.949 0.960 0.951	

Table 1: Statistics of the two simulation strategies

NOTE: The table reports the root mean squared error (RMSE), the absolute value of the average bias, the standard deviation, and the coverage of the 95% confidence interval for the ATE estimators across 1'000 simulations. Panel A reports the results for the synthetic DGP where data is generated according to Equation (8), while Panel B reports the results for the Empirical Monte Carlo Study. For each of the two simulation strategies, the table reports the results for 8 different simulation settings, where the sample size N and the share of treated $\mathbb{E}[D_i]$ are varied. For the synthetic DGP, the innovation standard deviation σ is set to 1, which corresponds to a signal-to-noise ratio of $Var[y]/\sigma^2 = 1.3$. The best performance in terms of RMSE is highlighted in bold.

References

ATHEY, S. AND G. W. IMBENS (2019): "Machine Learning Methods That Economists Should Know About," *Annual Review of Economics*, 11, 685–725.

BACH, P., O. SCHACHT, V. CHERNOZHUKOV, S. KLAASSEN, AND M. SPINDLER (2024): "Hyperparameter Tuning for Causal Inference with Double Machine Learning: A Simulation Study," Preprint (arXiv:2402.04674).

- BELLONI, A. AND V. CHERNOZHUKOV (2013): "Least squares after model selection in high-dimensional sparse models," *Bernoulli*, 19, 521–547.
- BOUCHAUD, J.-P. (2022): "The inelastic market hypothesis: a microstructural interpretation," *Quantitative Finance*, 22, 1785–1795.
- BREIMAN, L. (2001): "Random forests," *Machine learning*, 45, 5–32.
- CHEN, X. AND H. WHITE (1999): "Improved rates and asymptotic normality for nonparametric neural network estimators," *IEEE Transactions on Information Theory*, 45, 682–691.
- CHERNOZHUKOV, V., D. CHETVERIKOV, M. DEMIRER, E. DUFLO, C. HANSEN, W. NEWEY, AND J. ROBINS (2018): "Double/debiased machine learning for treatment and structural parameters," *The Econometrics Journal*, 21, C1–C68.
- FAN, Q., Y.-C. HSU, R. P. LIELI, AND Y. ZHANG (2022): "Estimation of conditional average treatment effects with high-dimensional data," *Journal of Business & Economic Statistics*, 40, 313–327.
- FARRELL, M. H. (2015): "Robust inference on average treatment effects with possibly more covariates than observations," *Journal of Econometrics*, 189, 1–23.
- FRIEDMAN, J. H. (1991): "Multivariate Adaptive Regression Splines," The Annals of Statistics, 19, 1-67.
- HAHN, J. (1998): "On the Role of the Propensity Score in Efficient Semiparametric Estimation of Average Treatment Effects," *Econometrica*, 66, 315–331.
- HE, H. AND E. A. GARCIA (2009): "Learning from Imbalanced Data," *IEEE Transactions on Knowledge and Data Engineering*, 21, 1263–1284.
- HUBER, M., M. LECHNER, AND C. WUNSCH (2013): "The performance of estimators based on the propensity score," *Journal of Econometrics*, 175, 1–21.
- HUJER, R., S. L. THOMSEN, AND C. ZEISS (2006): "The effects of vocational training programmes on the duration of unemployment in Eastern Germany," AStA Advances in Statistical Analysis, 90, 299–321.
- IMBENS, G. W. (2004): "Nonparametric Estimation of Average Treatment Effects Under Exogeneity: A Review," *The Review of Economics and Statistics*, 86, 4–29.
- JAPKOWICZ, N. AND S. STEPHEN (2002): "The class imbalance problem: A systematic study," *Intelligent Data Analysis*, 6, 429–449.
- KNAUS, M. C. (2020): "A Double Machine Learning Approach to Estimate the Effects of Musical Practice on Student's Skills," *Journal of the Royal Statistical Society Series A: Statistics in Society*, 184, 282–300.
- KNAUS, M. C., M. LECHNER, AND A. STRITTMATTER (2022): "Heterogeneous employment effects of job search programs: A machine learning approach," *Journal of Human Resources*, 57, 597–636.
- KÜNZEL, S. R., J. S. SEKHON, P. J. BICKEL, AND B. YU (2019): "Metalearners for estimating heterogeneous treatment effects using machine learning," *Proceedings of the National Academy of Sciences*, 116, 4156–4165.
- LECHNER, M., M. KNAUS, M. HUBER, M. FRÖLICH, S. BEHNCKE, G. MELLACE, AND A. STRITTMATTER (2020): "Swiss Active Labor Market Policy Evaluation [Dataset]," https://doi.org/10.23662/FORS-DS-1203-1.
- LECHNER, M. AND C. WUNSCH (2013): "Sensitivity of matching-based program evaluations to the availability of control variables," *Labour Economics*, 21, 111–121.
- LUO, Y., M. SPINDLER, AND J. KÜCK (2016): "High-Dimensional L_2 Boosting: Rate of Convergence," Preprint (arXiv:1602.08927).
- NIE, X. AND S. WAGER (2020): "Quasi-oracle estimation of heterogeneous treatment effects," *Biometrika*, 108, 299–319.
- PEDREGOSA, F., G. VAROQUAUX, A. GRAMFORT, V. MICHEL, B. THIRION, O. GRISEL, M. BLONDEL, P. PRET-TENHOFER, R. WEISS, V. DUBOURG, J. VANDERPLAS, A. PASSOS, D. COURNAPEAU, M. BRUCHER, M. PER-ROT, AND E. DUCHESNAY (2011): "Scikit-learn: Machine Learning in Python," *Journal of Machine Learning Research*, 12, 2825–2830.
- POZZOLO, A. D., O. CAELEN, R. A. JOHNSON, AND G. BONTEMPI (2015): "Calibrating Probability with Undersampling for Unbalanced Classification," in 2015 IEEE Symposium Series on Computational Intelligence, 159–166.
- ROSENBAUM, P. R. AND D. B. RUBIN (1983): "The central role of the propensity score in observational studies for causal effects," *Biometrika*, 70, 41–55.

- RUBIN, D. (1972): "Estimating causal effects of treatments in randomized and nonrandomized studies," *Journal of Educational Psychology*, 66, 688–701.
- SŁOCZYŃSKI, T. AND J. M. WOOLDRIDGE (2018): "A general double robustness result for estimating average treatment effects," *Econometric Theory*, 34, 112–133.

WAGER, S. (2022): "STATS 361: Causal Inference," Lecture Notes.

- WAGER, S. AND G. WALTHER (2016): "Adaptive concentration of regression trees, with application to random forests," Preprint (arXiv:1503.06388).
- YAO, L., Z. CHU, S. LI, Y. LI, J. GAO, AND A. ZHANG (2021): "A Survey on Causal Inference," ACM Trans. Knowl. Discov. Data, 15.
- ZIMMERT, M. AND M. LECHNER (2019): "Nonparametric estimation of causal heterogeneity under high-dimensional confounding," *Preprint (arXiv:1908.08779)*.

A Appendix: Proofs

Lemma A1. Let $\gamma = \mathbb{E}[D]/(1 - \mathbb{E}[D])$ and $\widehat{\gamma} = \left(\sum_{i=1}^{N} D_i\right) / \left(1 - \sum_{i=1}^{N} D_i\right)$. Under Assumption 3 it holds that: $|\widehat{\gamma} - \gamma| = o_p(N^{-1/2})$

Proof of Lemma A1. Let $\hat{p} = N^{-1} \sum_{i=1}^{N} D_i$ and consequently $\hat{\gamma} = \hat{p}/(1-\hat{p})$. Since $\mathbb{E}[\hat{p}] = \mathbb{E}[D] = p$, we have that: $\mathbb{E}[|\hat{p} - p|^2] = \operatorname{Var}[\hat{p}] = \frac{p(1-p)}{N} = o(N^{-1})$

and therefore $|\widehat{p}-p|=o_p(N^{-1/2}).$ For $\widehat{\gamma}$ we have:

$$|\widehat{\gamma} - \gamma| = \left|\frac{\widehat{p}}{1 - \widehat{p}} - \frac{p}{1 - p}\right| = \left|\frac{\widehat{p}(1 - p) - p(1 - \widehat{p})}{(1 - \widehat{p})(1 - p)}\right| \le \frac{1}{\epsilon^2}|\widehat{p} - p| = o_p(N^{-1/2})$$

Proof of Theorem 1. The proof follows the approach of Wager (2022). If the true nuisance functions $\mu_d(X)$ and p(X) are known, the oracle estimator for θ given by:

$$\widetilde{\theta} = \frac{1}{N} \sum_{i=1}^{N} \tau(Z_i)$$

$$= \frac{1}{N} \sum_{i=1}^{N} \left(\mu_1(X_i) - \mu_0(X_i) + \frac{D_i}{p(X_i)} (Y_i - \mu_1(X_i)) - \frac{1 - D_i}{1 - p(X_i)} (Y_i - \mu_0(X_i)) \right)$$

is a sample average of i.i.d. random variables and by the central limit theorem we have that:

$$\sqrt{N}\left(\widetilde{\theta}-\theta\right) \stackrel{d}{\longrightarrow} \mathcal{N}\left(0,V^*\right).$$

It therefore sufficies to show that $\sqrt{N}\left(\tilde{\theta} - \hat{\theta}^{\text{CU-DML}}\right) = o_p(1)$. Notice that $\theta = \theta_1 - \theta_0$ where $\theta_d = \mathbb{E}[Y^d]$. Moreover, $\tilde{\theta} = \tilde{\theta}_1 - \tilde{\theta}_0$ and $\hat{\theta}^{\text{CU-DML}} = \hat{\theta}_1^{\text{CU-DML}} - \hat{\theta}_0^{\text{CU-DML}}$. Therefore, it sufficies to show that $\sqrt{N}\left(\tilde{\theta}_1 - \hat{\theta}_1^{\text{CU-DML}}\right) = o_p(1)$, the result for $\tilde{\theta}_0$ follows analogously. The estimator can be further re-written as:

$$\widehat{\theta}_{1}^{\text{CU-DML}} = \frac{1}{N} \sum_{k=1}^{K} \sum_{i \in \mathcal{I}_{k}} \widehat{\mu}_{1}^{\mathcal{I}_{-k}}(X_{i}) + \frac{D_{i}}{\widehat{p}^{\mathcal{I}_{-k}}(X_{i})} (Y_{i} - \widehat{\mu}_{1}^{\mathcal{I}_{-k}}(X_{i}))$$
$$= \sum_{k=1}^{K} \frac{|\mathcal{I}_{k}|}{N} \widehat{\theta}_{1,\mathcal{I}_{k}}^{\text{CU-DML}}$$

where:

$$\widehat{\theta}_{1,\mathcal{I}_{k}}^{\text{CU-DML}} = \frac{1}{|\mathcal{I}_{k}|} \sum_{i \in \mathcal{I}_{k}} \widehat{\mu}_{1}^{\mathcal{I}_{-k}}(X_{i}) + \frac{D_{i}}{\widehat{p}^{\mathcal{I}_{-k}}(X_{i})} (Y_{i} - \widehat{\mu}_{1}^{\mathcal{I}_{-k}}(X_{i})).$$

Analogously, we can defined the oracle estimator $\tilde{\theta}_{1,\mathcal{I}_k}$. It therefore sufficies to show that $\sqrt{N}\left(\tilde{\theta}_{1,\mathcal{I}_k} - \hat{\theta}_{1,\mathcal{I}_k}^{\text{CU-DML}}\right) = o_p(1)$ for all $k = 1, \ldots, K$. We can decompose the difference between the oracle and the CU-DML estimator as:

$$\widetilde{\theta}_{1,\mathcal{I}_{k}} - \widehat{\theta}_{1,\mathcal{I}_{k}}^{\text{CU-DML}} = \underbrace{\frac{1}{|\mathcal{I}_{k}|} \sum_{i \in \mathcal{I}_{k}} \left(\widehat{\mu}_{1}^{\mathcal{I}_{-k}}(X_{i}) - \mu_{1}(X_{i}) \right) \left(1 - \frac{D_{i}}{p(X_{i})} \right)}_{\text{(A)}} + \underbrace{\frac{1}{|\mathcal{I}_{k}|} \sum_{i \in \mathcal{I}_{k}} D_{i} \left((Y_{i} - \mu_{1}(X_{i})) \left(\frac{1}{\widehat{p}^{\mathcal{I}_{-k}}(X_{i})} - \frac{1}{p(X_{i})} \right) \right)}_{\text{(B)}} - \underbrace{\frac{1}{|\mathcal{I}_{k}|} \sum_{i \in \mathcal{I}_{k}} D_{i} \left(\left(\widehat{\mu}_{1}^{\mathcal{I}_{-k}}(X_{i}) - \mu_{1}(X_{i}) \right) \left(\frac{1}{\widehat{p}^{\mathcal{I}_{-k}}(X_{i})} - \frac{1}{p(X_{i})} \right) \right)}_{\text{(C)}}$$

We will show that each of the three terms converges to zero in probability at rate $N^{-1/2}$. Term (A) is not dependent on the estimated propensity score and is not affected by undersampling. As such, the usual arguments apply for its convergence and we refer to Wager (2022) for the details. For term (B) we can show that its squared L_2 -norm is $o(N^{-1/2})$:

$$\begin{split} & \left\| \frac{1}{|\mathcal{I}_{k}|} \sum_{i \in \mathcal{I}_{k}} D_{i} \left((Y_{i} - \mu_{1}(X_{i})) \left(\frac{1}{\hat{p}^{\mathcal{I}_{-k}}(X_{i})} - \frac{1}{p(X_{i})} \right) \right) \right\|_{2}^{2} \\ &= \mathbb{E} \left[\left[\left(\frac{1}{|\mathcal{I}_{k}|} \sum_{i \in \mathcal{I}_{k}} D_{i} \left((Y_{i} - \mu_{1}(X_{i})) \left(\frac{1}{\hat{p}^{\mathcal{I}_{-k}}(X_{i})} - \frac{1}{p(X_{i})} \right) \right) \right)^{2} \right] \\ &= \mathbb{E} \left[\mathbb{E} \left[\left(\frac{1}{|\mathcal{I}_{k}|} \sum_{i \in \mathcal{I}_{k}} D_{i} \left((Y_{i} - \mu_{1}(X_{i})) \left(\frac{1}{\hat{p}^{\mathcal{I}_{-k}}(X_{i})} - \frac{1}{p(X_{i})} \right) \right) \right)^{2} \right] \{Z_{i}\}_{i \in \mathcal{I}_{-k}}, \{D_{i}\}_{i=1}^{N} \right] \right] \\ &= \mathbb{E} \left[\mathbb{Var} \left[\frac{1}{|\mathcal{I}_{k}|} \sum_{i \in \mathcal{I}_{k}} D_{i} \left((Y_{i} - \mu_{1}(X_{i})) \left(\frac{1}{\hat{p}^{\mathcal{I}_{-k}}(X_{i})} - \frac{1}{p(X_{i})} \right) \right) \right] \{Z_{i}\}_{i \in \mathcal{I}_{-k}}, \{D_{i}\}_{i=1}^{N} \right] \right] \\ &= \frac{1}{|\mathcal{I}_{k}|^{2}} \mathbb{E} \left[\sum_{i \in \mathcal{I}_{k}} \mathbb{Var} \left[D_{i} \left((Y_{i} - \mu_{1}(X_{i})) \left(\frac{1}{\hat{p}^{\mathcal{I}_{-k}}(X_{i})} - \frac{1}{p(X_{i})} \right) \right) \right] \{Z_{i}\}_{i \in \mathcal{I}_{-k}}, \{D_{i}\}_{i=1}^{N} \right] \right] \\ &= \frac{1}{|\mathcal{I}_{k}|} \mathbb{E} \left[\mathbb{Var} \left[D_{i} \left((Y_{i} - \mu_{1}(X_{i})) \left(\frac{1}{\hat{p}^{\mathcal{I}_{-k}}(X_{i})} - \frac{1}{p(X_{i})} \right) \right) \right] \{Z_{i}\}_{i \in \mathcal{I}_{-k}}, \{D_{i}\}_{i=1}^{N} \right] \right] \\ &= \frac{1}{|\mathcal{I}_{k}|} \mathbb{E} \left[D_{i} \left(Y_{i} - \mu_{1}(X_{i} \right) \right]^{2} \left(\frac{1}{\hat{p}^{\mathcal{I}_{-k}}(X_{i})} - \frac{1}{p(X_{i})} \right) \right] \\ &\leq \frac{1}{|\mathcal{I}_{k}|} \zeta \mathbb{E} \left[\left(\frac{1}{\hat{p}^{\mathcal{I}_{-k}}(X_{i})} - \frac{1}{p(X_{i})} \right)^{2} \right] = \frac{o(1)}{N}. \end{split}$$

The fourth equality in Equation (A1) follows from the fact that for any $i, j \in \mathcal{I}_k$ with $i \neq j$ we have:

$$\begin{split} &\operatorname{Cov}\left[D_{i}\left(\left(Y_{i}-\mu_{1}(X_{i})\right)\left(\frac{1}{\hat{p}^{\mathcal{I}_{-k}}(X_{i})}-\frac{1}{p(X_{i})}\right)\right),\\ &D_{j}\left(\left(Y_{j}-\mu_{1}(X_{j})\right)\left(\frac{1}{\hat{p}^{\mathcal{I}_{-k}}(X_{j})}-\frac{1}{p(X_{j})}\right)\right)\right)\left|\{Z_{i}\}_{i\in\mathcal{I}_{-k}},\{D_{i}\}_{i=1}^{N}\right]\\ &=\mathbb{E}\left[D_{i}\left(\left(Y_{i}-\mu_{1}(X_{i})\right)\left(\frac{1}{\hat{p}^{\mathcal{I}_{-k}}(X_{j})}-\frac{1}{p(X_{j})}\right)\right)\right)\\ &D_{j}\left(\left(Y_{j}-\mu_{1}(X_{j})\right)\left(\frac{1}{\hat{p}^{\mathcal{I}_{-k}}(X_{j})}-\frac{1}{p(X_{j})}\right)\right)\left|\{Z_{i}\}_{i\in\mathcal{I}_{-k}},\{D_{i}\}_{i=1}^{N}\right]\\ &=\mathbb{E}\left[\left(\frac{1}{\hat{p}^{\mathcal{I}_{-k}}(X_{i})}-\frac{1}{p(X_{i})}\right)\left(\frac{1}{\hat{p}^{\mathcal{I}_{-k}}(X_{j})}-\frac{1}{p(X_{j})}\right)\\ &\mathbb{E}[D_{i}(Y_{i}-\mu_{1}(X_{i}))|X_{i},D_{i}]\mathbb{E}[D_{j}(Y_{j}-\mu_{1}(X_{j}))|X_{j},D_{j}]\right|\{Z_{i}\}_{i\in\mathcal{I}_{-k}},\{D_{i}\}_{i=1}^{N}\right]=0 \end{split}$$

by the law of iterated expectations and independence of the observations. The last equality in Equation (A1) follows from Assumption 2 the fact that, under Assumptions 3 and 4, the calibrated propensity score is sup-norm consistent since:

$$\begin{aligned} |\hat{p}_{S}^{\mathcal{I}_{-k}}(x)\widehat{\gamma} - p_{S}(x)\gamma| &= |\hat{p}_{S}^{\mathcal{I}_{-k}}(x)\widehat{\gamma} - p_{S}(x)\widehat{\gamma} + p_{S}(x)\widehat{\gamma} - p_{S}(x)\gamma| \\ &\leq |\hat{p}_{S}^{\mathcal{I}_{-k}}(x) - p_{S}(x)| \cdot |\widehat{\gamma}| + |\widehat{\gamma} - \gamma| \cdot |p_{S}(x)| \\ &\leq |\hat{p}_{S}^{\mathcal{I}_{-k}}(x) - p_{S}(x)| + |\widehat{\gamma} - \gamma| \end{aligned}$$

and $|\hat{\gamma} - \gamma| = o_p(N^{-1/2})$ as shown in Lemma A1. We conclude that (B) is $o_p(N^{-1/2})$. Lastly, we bound the term (C) by showing that its expectation converges to zero at rate $N^{-1/2}$:

$$\begin{split} & \mathbb{E}\left[D_{i}\left(\left(\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right)\left(\frac{1}{\hat{p}^{\mathcal{I}-k}(X_{i})}-\frac{1}{p(X_{i})}\right)\right)\right] \\ &\leq \mathbb{E}\left[\left(\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right)\left(\frac{1}{\hat{p}^{\mathcal{I}-k}(X_{i})}-\frac{1}{p(X_{i})}\right)\right] \\ &\leq \mathbb{E}\left[\left|\left(\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right)\left(\frac{1}{\hat{p}^{\mathcal{I}-k}(X_{i})}-\frac{1}{p(X_{i})}\right)\right|\right] \\ &= \mathbb{E}\left[\left|\left(\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right)\left(\frac{1}{\gamma p_{S}(X_{i})}-\frac{1}{\hat{\gamma} \hat{p}_{S}^{\mathcal{I}-k}(X_{i})}-\frac{1}{\gamma}+\frac{1}{\hat{\gamma}}\right)\right|\right] \\ &\leq \mathbb{E}\left[\left|\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right|\left|\left(\frac{1}{\gamma p_{S}(X_{i})}-\frac{1}{\hat{\gamma} \hat{p}_{S}^{\mathcal{I}-k}(X_{i})}\right)+\left(\frac{1}{\hat{\gamma}}-\frac{1}{\gamma}\right)\right|\right] \\ &\leq \mathbb{E}\left[\left|\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right|\left|\left|\hat{\gamma} \hat{p}_{S}^{\mathcal{I}-k}(X_{i})-\gamma p_{S}(X_{i})\right|\right]+\mathbb{E}\left[\left|\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right|\left|\gamma-\hat{\gamma}\right|\right] \\ &\leq c_{1}\mathbb{E}\left[\left|\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right|\left|\hat{\gamma} \hat{p}_{S}^{\mathcal{I}-k}(X_{i})-p_{S}(X_{i})\right|\right] \\ &+ c_{2}\mathbb{E}\left[\left|\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right|\left|\hat{\gamma}-\gamma|p_{S}(X_{i})\right] \\ &+ c_{2}\mathbb{E}\left[\left|\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right|\left|\hat{p}_{S}^{\mathcal{I}-k}(X_{i})-p_{S}(X_{i})\right|\right] \\ &+ c_{1}\mathbb{E}\left[\left|\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right|\left|\hat{p}_{S}^{\mathcal{I}-k}(X_{i})-p_{S}(X_{i})\right|\right] \\ &+ c_{2}\mathbb{E}\left[\left|\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right|\left|\hat{p}_{S}^{\mathcal{I}-k}(X_{i})-p_{S}(X_{i})\right|\right] \\ &+ c_{2}\mathbb{E}\left[\left|\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right|\left|\hat{p}_{S}^{\mathcal{I}-k}(X_{i})-p_{S}(X_{i})\right|\right] \\ &+ c_{2}\mathbb{E}\left[\left|\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right|\left|\hat{\gamma}-\gamma\right|\right] \\ &+ c_{2}\mathbb{E}\left[\left|\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right|\left|\hat{\gamma}-\gamma\right|\right] \\ &= c_{1}\|\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right|\left|\hat{\gamma}-\gamma\right|\right] \\ &= c_{1}\|\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right|_{2}\|\hat{p}_{S}^{\mathcal{I}-k}(X_{i})-p_{S}(X_{i})\right\|_{2} + (c_{1}+c_{2})\left\|\hat{\mu}_{1}^{\mathcal{I}-k}(X_{i})-\mu_{1}(X_{i})\right\|_{2}\|\hat{\gamma}-\gamma\right\|_{2} \end{aligned}$$

where the first term in the last step is $o(N^{-1/2})$ by Assumption 4 (ii) and the second term is $o(N^{-1/2})$ by Lemma A1 and Assumption 4 (i). The positive constants c_1 and c_2 come from the boundness of $p_S(x)$ and γ (and that of their estimators). From the law of large numbers, it follows that (C) is $o_p(N^{-1/2})$.

B Appendix: Details on the simulation study

B.1 Implementation details

The simulation study is implemented in Python 3.12. The data generating process is implemented using the numpy 1.26.4 package. The machine learning models are estimated using the scikit-learn 1.4.0 package, using the acceleration extension scikit-learn-intelex 2024.4.0. The empirical data is processed using pandas 2.2.2. The code for the simulation study is availabile at https://github.com/dballinari/Calibrating-doubly-robust-estimators-with-unbalanced-treatment-assignment

B.2 Additional simulation results

	N =	2000. E[$D_i = 2.5\%$	$\sigma = 5$	$N = 4000, \mathbb{E}[D_i] = 2.5\%, \sigma = 5$				$N = 8000, \mathbb{E}[D_i] = 2.5\%, \sigma = 5$			
	RMSE	Bias	Std. dev.	Coverage	RMSE	Bias	Std. dev.	Coverage	RMSE	Bias	Std. dev.	Coverage
DML	-	-	-	-	0.885	0.025	0.884	0.983	0.448	-0.001	0.448	0.980
U-DML	-	-	-	-	0.761	0.097	0.755	0.948	0.523	0.065	0.519	0.954
CU-DML	-	-	-	-	0.588	0.056	0.585	0.944	0.402	0.019	0.401	0.963
W-DML	-	-	-	-	0.675	0.053	0.673	0.974	0.429	0.005	0.428	0.979
N-DML	-	-	-	-	0.698	0.046	0.696	0.932	0.420	0.008	0.420	0.948
T-DML	-	-	-	-	0.671	0.055	0.669	0.934	0.420	0.008	0.420	0.948
	$N = 2000, \mathbb{E}[D_i] = 5.0\%, \sigma = 5$				$N = 4000, \mathbb{E}[D_i] = 5.0\%, \sigma = 5$				$N = 8000, \mathbb{E}[D_i] = 5.0\%, \sigma = 5$			
	RMSE	Bias	Std. dev.	Coverage	RMSE	Bias	Std. dev.	Coverage	RMSE	Bias	Std. dev.	Coverage
DML	0.742	0.038	0.741	0.982	0.421	0.020	0.420	0.973	0.287	0.008	0.287	0.961
U-DML	0.773	0.100	0.767	0.949	0.510	0.045	0.508	0.961	0.375	0.055	0.372	0.951
CU-DML	0.600	0.056	0.597	0.950	0.395	0.025	0.395	0.958	0.284	0.017	0.283	0.951
W-DML	0.727	0.034	0.726	0.982	0.420	0.021	0.420	0.973	0.287	0.008	0.287	0.961
N-DML	0.656	0.054	0.654	0.934	0.402	0.028	0.401	0.950	0.283	0.012	0.283	0.947
T-DML	0.651	0.045	0.649	0.936	0.402	0.028	0.401	0.950	0.283	0.012	0.283	0.947
	$N = 2000, \mathbb{E}[D_i] = 10.0\%, \sigma = 5$			$N = 4000, \mathbb{E}[D_i] = 10.0\%, \sigma = 5$				$N = 8000, \mathbb{E}[D_i] = 10.0\%, \sigma = 5$				
	RMSE	Bias	Std. dev.	Coverage	RMSE	Bias	Std. dev.	Coverage	RMSE	Bias	Std. dev.	Coverage
DML	0.456	0.039	0.454	0.953	0.288	0.028	0.287	0.965	0.208	0.010	0.208	0.949
U-DML	0.550	0.080	0.544	0.944	0.362	0.047	0.359	0.951	0.272	0.047	0.268	0.932
CU-DML	0.438	0.045	0.436	0.942	0.292	0.029	0.290	0.958	0.210	0.015	0.209	0.936
W-DML	0.456	0.039	0.454	0.953	0.288	0.028	0.287	0.965	0.208	0.010	0.208	0.949
N-DML	0.446	0.043	0.444	0.924	0.286	0.031	0.284	0.949	0.207	0.012	0.207	0.939
T-DML	0.446	0.043	0.444	0.924	0.286	0.031	0.284	0.949	0.207	0.012	0.207	0.939

Table B1: Statistics of the synthetic DGP with $\sigma = 5$

NOTE: The table reports the root mean squared error (RMSE), the absolute value of the average bias, the standard deviation, and the coverage of the 95% confidence interval for the ATE estimators across 1'000 simulations. The data is generated from the synthetic DGP according to Equation (8), with innovation standard deviation $\sigma = 5$, which corresponds to a signal-to-noise ratio of $\operatorname{Var}[y]/\sigma^2 = 1.0$. The table reports the results for 8 different simulation settings, where the sample size N and the share of treated $\mathbb{E}[D_i]$ are varied. The best performance in terms of RMSE is highlighted in bold.