

NOISE SENSITIVITY AND STABILITY ON GROUPS

RYOKICHI TANAKA

ABSTRACT. We provide finitely generated infinite groups on which natural random walks are noise sensitive in total variation as well as ones on which natural random walks are noise stable in total variation.

1. INTRODUCTION

Let Γ be a countable group and μ be a probability measure on it. A μ -**random walk** $\{w_n\}_{n=0}^\infty$ starting from the identity id is defined by $w_n := x_1 \cdots x_n$ and $w_0 := \text{id}$ for an independent, identically distributed sequence x_1, x_2, \dots with the common law μ . The distribution of w_n is the n -fold convolution $\mu_n := \mu^{*n}$. For a real $\rho \in [0, 1]$, let

$$\pi^\rho := \rho\mu \times \mu + (1 - \rho)\mu_{\text{diag}} \quad \text{on } \Gamma \times \Gamma,$$

where $\mu \times \mu$ denotes the product measure and $\mu_{\text{diag}}((x, y)) := \mu(x)$ if $x = y$ and 0 otherwise. Let us consider a π^ρ -random walk $\{\mathbf{w}_n\}_{n \in \mathbb{Z}_+}$ starting from the identity on $\Gamma \times \Gamma$. On the one hand, we say that the μ -random walk is **noise sensitive** in total variation if

$$\lim_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} = 0 \quad \text{for all } \rho \in (0, 1],$$

where $\|\cdot\|_{\text{TV}}$ stands for the total variation distance that coincides with the half of ℓ^1 -norm. On the other hand, we say that the μ -random walk is **noise stable** in total variation if

$$\lim_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} = 1 \quad \text{for all } \rho \in [0, 1).$$

This latter notion is a strong negation of the former. Since we use only total variation distance in the definitions, let us simply say noise sensitive or noise stable respectively if there is no danger of confusion.

If a μ -random walk on Γ is noise sensitive, then informally speaking, the situation is as in the following: For each fixed $\rho \in (0, 1]$ even though it is close to 0, independent refreshes of ρ -portions in the increments produce an asymptotically independent copy of the original μ -random walk. If a μ -random walk on Γ is noise stable, then the situation is completely opposite in the following sense: For each fixed $\rho \in [0, 1)$ even though it is close to 1, independent refreshes of ρ -portions in the increments only produce a pair which is statistically distinguishable from the pair of two independent μ -random walks.

The noise sensitivity for random walks on groups was introduced by Benjamini and Brioussell in [BB23, Definition 2.1]. In their paper, they discuss the notion and the variants not only in total variation (there it is called ℓ^1 -noise sensitivity) but also in other distances or in terms of entropy. See also a related discussion in [Kal18, Section 3.3.4]. It has been

observed that on finite groups random walks are noise sensitive in all natural notions [BB23, Proposition 5.1]. This raises a challenge to find finitely generated *infinite* groups on which random walks are noise sensitive. Further the problem becomes more restrictive by measuring the distance in total variation. A simple observation on the local central limit theorem shows that standard random walks on finite rank free abelian groups \mathbb{Z}^m are *neither* noise sensitive *nor* noise stable (see Appendix A). Benjamini and Brioussell have shown that on the infinite dihedral group some lazy simple random walk is noise sensitive [ibid. Theorem 1.4]. So far, this has been the only known random walk which is noise sensitive in total variation on a finitely generated infinite group. We provide a class of such groups on which natural random walks are noise sensitive.

Theorem 1.1. *Let (Γ, S) be an affine Weyl group, and μ be a probability measure on Γ such that the support of μ equals $S \cup \{\text{id}\}$. For all $\rho \in (0, 1]$,*

$$\lim_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} = 0,$$

i.e., the μ -random walk on Γ is noise sensitive in total variation.

Theorem 1.1 is shown in Theorem 3.10. The infinite dihedral group

$$D_\infty := \langle s_1, s_2 \mid s_1^2 = s_2^2 = \text{id} \rangle \quad \text{with } S = \{s_1, s_2\}$$

is a special case of affine Weyl group called type \tilde{A}_1 . See more on explicit examples of affine Weyl groups in Section 3.8. In fact, there exist a constant $C > 0$ and an integer $m > 0$ such that for all large enough integer n ,

$$\|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} \leq \frac{C(\log n)^m}{n}.$$

In Theorem 1.1, the laziness (i.e., $\mu(\text{id}) > 0$) is crucial since otherwise random walk on that group is not necessarily noise sensitive. This in particular shows that the noise sensitivity (in total variation) is a property of the random walk rather than the group as it has been pointed out in [BB23]. Let us note that the μ -random walk on the infinite dihedral group considered there has a particular form: $\mu(s_1) = \mu(s_2) = \mu(\text{id}) = 1/3$. It is not already clear from their proof that changing the laziness (i.e., the measure on the identity element) would still provide a noise sensitive random walk. We show that this is indeed the case, furthermore, μ is allowed to be a non-uniform distribution on $S \cup \{\text{id}\}$.

Let us discuss the noise stability. The following is known for a finitely supported probability measure μ : Either if the group Γ admits a surjective homomorphism onto \mathbb{Z} , or if (Γ, μ) is **non-Liouville**, i.e., there exists a non-constant bounded μ -harmonic function on Γ , then a μ -random walk on Γ is *not* noise sensitive [BB23, Theorem 1.1]. In a more specific class of (Γ, μ) (where possibly Γ does or does not admit a surjective homomorphism onto \mathbb{Z}), a strong negation of noise sensitivity has been shown for non-elementary word hyperbolic groups, e.g., free groups of rank at least 2. In which case, if μ has the support generating Γ and a finite first moment, then there exists a $\rho_0 \in (0, 1]$ such that $\|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} \rightarrow 1$ as $n \rightarrow \infty$ for all $\rho \in [0, \rho_0)$ [Tan22, Theorem 1.3]. In this generality, it is not known as to whether it holds that $\rho_0 = 1$. (Note that it is rather straightforward to check this for simple random walks on free *semi*-groups of rank at least

2, cf. [ibid, Introduction].) We show that in a more restricted class of groups and random walks it is indeed the case.

Theorem 1.2. *Let Γ be a non-elementary word hyperbolic group which admits a surjective homomorphism onto \mathbb{Z} and μ be a probability measure on Γ . If the support of μ generates Γ as a semigroup and μ has a finite second moment with respect to a word metric, then*

$$\lim_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} = 1 \quad \text{for all } \rho \in [0, 1),$$

i.e., the μ -random walk is noise stable in total variation.

Theorem 1.2 is shown in Theorem 2.1, where, in fact, it is shown that

$$\lim_{n \rightarrow \infty} \|\pi_n^\rho - \pi_n^{\rho_\star}\|_{\text{TV}} = 1 \quad \text{for all } \rho, \rho_\star \in [0, 1] \text{ with } \rho \neq \rho_\star. \quad (1.1)$$

In particular, Theorem 1.2 includes the case when Γ is a free group of rank at least 2 with μ supported on the standard set of generators. This special case has an implication on a family of simple random walks on high girth finite graphs (Corollary 2.7).

The following result provides further examples of groups on which random walks can be noise stable. Let $G \wr \Lambda := (\bigoplus_\Lambda G) \rtimes \Lambda$ be the lamplighter group with the lamp group G and the base group Λ (see Section 2.2). A probability measure μ on $G \wr \Lambda$ is called **switch-or-walk** if μ has the support in the set $\{(\delta^s, \text{id}), (\mathbf{id}, x) : s \in G, x \in \Lambda\}$ where $\delta^s(x) = s$ if $x = \text{id}$ and id else, and $\mathbf{id}(x) = \text{id}$ for all $x \in \Lambda$.

Theorem 1.3. *Let G be a nontrivial finite group, m be a positive integer, and μ be a finitely supported switch-or-walk probability measure on $G \wr \mathbb{Z}^m$ such that the support of μ generates $G \wr \mathbb{Z}^m$ as a semigroup. If the projection μ_{base} of μ on \mathbb{Z}^m defines a transient random walk, then*

$$\lim_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} = 1 \quad \text{for all } \rho \in [0, 1),$$

i.e., the μ -random walk on $G \wr \mathbb{Z}^m$ is noise stable in total variation.

Theorem 1.3 is shown in a slightly more general form in Theorem 2.3. Furthermore, (1.1) holds also in this setting. The result applies, for example, to the case when $(\mathbb{Z}/2) \wr \mathbb{Z}^m$ for $m \geq 3$ and μ is the uniform distribution on the set of generators $(\delta^s, 0)$, $(\mathbf{id}, \pm \mathbf{e}_i)$ for $s \in \mathbb{Z}/2$, $i = 1, \dots, m$, where $\{\mathbf{e}_1, \dots, \mathbf{e}_m\}$ denotes the standard basis in \mathbb{Z}^m .

Outlines of proofs. First let us discuss the noise sensitivity result in Theorem 1.1. An affine Weyl group (Γ, S) with the standard set of generators S is associated with some Euclidean space \mathbb{R}^m . The group Γ acts on \mathbb{R}^m isometrically and generators act as reflections relative to hyperplanes. The Γ has the form $\Lambda \rtimes W$ where Λ is identified with a lattice in \mathbb{R}^m and W is a finite group (called a spherical Weyl group). There is a point o in \mathbb{R}^m such that the orbit map $\Phi : x \mapsto x.o$ is injective. Taking a conjugate by a translation if necessary, we assume that o is the origin in \mathbb{R}^m (see Section 3.1 for the precise discussion). A main ingredient is to establish a local central limit theorem (Theorem 3.6). We define a discrete normal distribution $\mathcal{N}_{n\Sigma}^\Phi$ on Γ induced from the normalized restriction on $\Phi(\Gamma)$ of the m -dimensional Gaussian density function with a covariance matrix $n\Sigma$. Further we show the following: The distribution μ_n is approximated by $\mathcal{N}_{n\Sigma}^\Phi$ uniformly on Γ within an

error of order $n^{-\frac{m}{2}-1}$ as n tends to infinity. The local central limit theorem itself follows from a classical argument based on characteristic functions. Some more general results (other than the Cayley graphs of affine Weyl groups) have been known, e.g., in [KS83], [PS94], [KSS98] and [Sun13]. We provide the proof of the local central limit theorem in our setting with an error estimate. Furthermore, we use an explicit form of matrix Σ in terms of harmonic 1-forms on a finite quotient graph of the Cayley graph by the lattice Λ . The matrix Σ is obtained as some limiting form, which is known in the literature of symbolic dynamics, however, the explicit form in Theorem 3.6 plays an important role.

We apply this discussion to $(\Gamma \times \Gamma, \pi^\rho)$ for $\rho \in (0, 1]$. The local central limit theorem enables us to show that there exists a constant $C > 0$ such that for all large enough n ,

$$\|\pi_n^\rho - \mathcal{N}_{n\Sigma^\rho}^\Phi\|_{\text{TV}} \leq \frac{C(\log n)^m}{n}$$

(Theorem 3.8). This bound is sharp up to the factor $O((\log n)^m)$ (Remark 3.9). If μ has the support precisely $S \cup \{\text{id}\}$, then π^ρ has the support precisely $(S \cup \{\text{id}\})^2$ consisting of element of order at most 2. The explicit formula of the covariance matrix implies that Σ^ρ has a block diagonal form and $\Sigma^\rho = \Sigma^1$ for all $\rho \in (0, 1]$. (This is the only part where we use the particular structure of the generating sets.) Thus by the triangle inequality we conclude Theorem 1.1 (in Theorem 3.10). Let us note that if the support of μ does not contain id , then the support of π^ρ does not generate the group $\Gamma \times \Gamma$ (cf. Section 3.7). Furthermore, there is an example of noise sensitive random walk on the infinite dihedral group D_∞ not covered by this general statement; see Section 3.8.1.

Next let us discuss the noise stability results in Theorems 1.2 and 1.3. The proof uses a resolution of the identification problem on Poisson boundary. This is known in [Kai00] for the case of (products of) word hyperbolic groups, and in [LP21] for the case of lamplighter groups, respectively (see Sections 2.1 and 2.2). In the case of a product of word hyperbolic groups $\Gamma \times \Gamma$, the π^ρ -random walk $\{\mathbf{w}_n\}_{n \in \mathbb{Z}_+}$ converges to a point \mathbf{w}_∞ in the product of geometric (Gromov) boundaries $\partial\Gamma \times \partial\Gamma$ almost surely in $(\Gamma \cup \partial\Gamma)^2$. The distribution of \mathbf{w}_∞ is the harmonic (or, stationary) measure ν_{π^ρ} on $\partial\Gamma \times \partial\Gamma$. A result by Kaimanovich [Kai00] implies that the tail (and invariant) σ -field of the π^ρ -random walk is obtained by the σ -algebra $\sigma(\mathbf{w}_\infty)$ generated by \mathbf{w}_∞ modulo null measure sets.

If Γ admits a surjective homomorphism φ onto \mathbb{Z} , then $\{\varphi(w_n^{(1)}) - \varphi(w_n^{(2)})\}_{n \in \mathbb{Z}_+}$ for $\mathbf{w}_n = (w_n^{(1)}, w_n^{(2)})$ satisfies the Hartman-Wintner law of the iterated logarithm with the variance proportional to ρ . This produces a tail (in fact, invariant) event which has a full measure for the one ρ but null for all the others. Since the tail (and invariant) events are obtained by $\sigma(\mathbf{w}_\infty)$ -measurable sets up to ν_{π^ρ} -measure null in $\partial\Gamma \times \partial\Gamma$, the harmonic measures ν_{π^ρ} are mutually singular for different parameters $\rho \in [0, 1]$. Noting that the distribution π_n^ρ weakly converges to ν_{π^ρ} in the compact metrizable space $(\Gamma \cup \partial\Gamma)^2$ as $n \rightarrow \infty$, we conclude that the μ -random walk on Γ is noise stable in total variation. The argument works if μ has a finite second moment and the support generates Γ as a semigroup. In the case of lamplighter groups, the proof follows the same line, relying on the result by Lyons and Peres [LP21].

Organization. In Section 2, we show the noise stability for word hyperbolic groups (Theorem 2.1), and for lamplighter groups (Theorem 2.3). A result on families of high girth finite graphs is obtained in Corollary 2.7. In Section 3, we show the local central limit theorem in a slightly extended setting (Theorem 3.6) and deduce the noise sensitivity for affine Weyl groups (Theorem 3.10); explicit examples are presented in Section 3.8. In Appendix A, we include a result (Theorem A.1) on \mathbb{Z}^m for the sake of convenience.

Notations. For a constant C , we write $C = C_\Sigma$ to indicate its dependence on Σ . For non-negative real valued functions f and g on a common (sub-)set of non-negative integers \mathbb{Z}_+ , we write $f(n) = O(g(n))$ or $f \ll g$ if there exists a constant C such that $f(n) \leq Cg(n)$ for all large enough n . Also, $f(n) = O_\Sigma(g(n))$ if $C = C_\Sigma$ in the above notation. Further we write $f(n) = \Omega(g(n))$ if there exists a constant $c > 0$ such that $f(n) \geq cg(n)$ for all large enough n , and $f(n) = \Theta(g(n))$ if $f \ll g$ and $g \ll f$. For a set A , we denote by $\#A$ the cardinality. For a real $a \in \mathbb{R}$, we denote by $[a]$ the integer part of a .

2. NOISE STABILITY

2.1. Noise stability on word hyperbolic groups. For background on word hyperbolic groups, we refer to the original paper [Gro87]. See also a useful exposition [Cal13].

Let Γ be a non-elementary word hyperbolic group which admits a surjective homomorphism φ onto the infinite cyclic group \mathbb{Z} . Let μ be a probability measure on Γ where the support generates the whole group as a semigroup. For example, Γ is a free group of finite rank greater than one, or a fundamental group of closed orientable surface of genus greater than one. In those cases, such a φ is obtained through the abelianization $\Gamma/[\Gamma, \Gamma]$, which is a free abelian group.

Theorem 2.1. *Let Γ be a non-elementary word hyperbolic group which admits a surjective homomorphism onto \mathbb{Z} , and μ be a probability measure on Γ . If the support of μ generates the whole group as a semigroup and μ has a finite second moment with respect to a word metric, then*

$$\lim_{n \rightarrow \infty} \|\pi_n^\rho - \pi_n^{\rho_\star}\|_{\text{TV}} = 1 \quad \text{for all } \rho, \rho_\star \in [0, 1] \text{ with } \rho \neq \rho_\star.$$

In particular,

$$\lim_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} = 1 \quad \text{for all } \rho \in [0, 1),$$

i.e., the μ -random walk is noise stable in total variation.

For a countable group Γ and a probability measure on it, let $\Omega = \Gamma^{\mathbb{Z}_+}$ and \mathcal{F} be the σ -algebra generated by cylinder sets. Let \mathbf{P}_μ be the distribution of the μ -random walk starting from id on Ω , and we consider the probability space $(\Omega, \mathcal{F}, \mathbf{P}_\mu)$. We define a Γ -action on Ω by the diagonal action: $x \cdot \{w_n\}_{n \in \mathbb{Z}_+} := \{xw_n\}_{n \in \mathbb{Z}_+}$ for $x \in \Gamma$. Let us denote by \mathcal{I} the **invariant σ -field** on Ω , i.e., $\mathcal{I} = \sigma\{A \in \mathcal{F} : \vartheta^{-1}(A) = A\}$ where ϑ is the shift $\vartheta(\{w_n\}_{n \in \mathbb{Z}_+}) := \{w_{n+1}\}_{n \in \mathbb{Z}_+}$ on Ω .

In the case when Γ is a word hyperbolic group, if the support of μ generates a non-elementary subgroup as a group, then a μ -random walk $\{w_n\}_{n \in \mathbb{Z}_+}$ converges to a limit w_∞ in the Gromov boundary $\partial\Gamma$ in the natural compact metrizable space $\Gamma \cup \partial\Gamma$ almost

surely in \mathbf{P}_μ . The distribution ν_μ of w_∞ on $\partial\Gamma$ is called the **harmonic measure**. This is non-atomic and is the unique probability measure which is μ -stationary on $\partial\Gamma$, i.e., $\mu * \nu_\mu = \nu_\mu$ where $\mu * \nu_\mu := \sum_{x \in \Gamma} \mu(x) x_* \nu_\mu$ and $x_* \nu_\mu := \nu_\mu \circ x^{-1}$. We consider the natural Borel measurable structure $\mathcal{B}(\partial\Gamma)$ on $\partial\Gamma$. Note that w_∞ is defined \mathbf{P}_μ -almost everywhere on Ω . Fix $\xi \in \partial\Gamma$. Letting Ω_{conv} be the set of $\mathbf{w} = \{w_n\}_{n \in \mathbb{Z}_+}$ in Ω having a unique limit in $\partial\Gamma$. The set Ω_{conv} is \mathcal{F} -measurable and $\mathbf{P}_\mu(\Omega_{\text{conv}}) = 1$. Let us define

$$\mathbf{bnd} : \Omega \rightarrow \partial\Gamma,$$

by $\mathbf{bnd}(\mathbf{w}) := w_\infty(\mathbf{w})$ if $\mathbf{w} \in \Omega_{\text{conv}}$, and ξ otherwise. Note that Ω_{conv} is invariant under the Γ -action; the map \mathbf{bnd} is Γ -equivariant on Ω_{conv} , i.e., $\mathbf{bnd}(x \cdot \mathbf{w}) = x \cdot \mathbf{bnd}(\mathbf{w})$ for $x \in \Gamma$ and $\mathbf{w} \in \Omega_{\text{conv}}$. Moreover, since $\mathbf{P}_\mu(\Omega_{\text{conv}}) = 1$, the harmonic measure ν_μ is obtained as the pushforward of \mathbf{P}_μ by \mathbf{bnd} , i.e.,

$$\nu_\mu = \mathbf{bnd}_* \mathbf{P}_\mu \quad \text{on } \partial\Gamma.$$

Let $\sigma(\mathbf{bnd})$ be the σ -field generated by \mathbf{bnd} ; we have

$$\sigma(\mathbf{bnd}) = \{\mathbf{bnd}^{-1}(B) : B \in \mathcal{B}(\partial\Gamma)\}.$$

In the case when μ has a finite first moment (or more generally finite logarithmic moment with finite entropy), it is known that $(\partial\Gamma, \nu_\mu)$ is a Poisson boundary [Kai00]. Namely, the invariant σ -field \mathcal{I} and $\sigma(\mathbf{bnd})$ coincide modulo \mathbf{P}_μ -null sets. In other words, two σ -fields agree after taking completions under \mathbf{P}_μ . In particular, for every $A \in \mathcal{I}$ there exist $B_0, B_1 \in \mathcal{B}(\partial\Gamma)$ such that $\mathbf{bnd}^{-1}(B_0) \subset A \subset \mathbf{bnd}^{-1}(B_1)$ and $\nu_\mu(B_1 \setminus B_0) = 0$. The result of [Kai00] is generalized in a wider setting. The case which we are focusing on is a product of word hyperbolic group $\Gamma \times \Gamma$ endowed with a probability measure π . If both the pushforwards of π under projections from $\Gamma \times \Gamma$ onto factors satisfy the condition as above, then the harmonic measure ν_π is defined on $\partial\Gamma \times \partial\Gamma$ as the distribution of \mathbf{P}_π -almost every limiting point

$$\mathbf{w}_n = (w_n^{(1)}, w_n^{(2)}) \rightarrow (w_\infty^{(1)}, w_\infty^{(2)}) \quad \text{as } n \rightarrow \infty \text{ in } (\Gamma \cup \partial\Gamma)^2.$$

This in particular shows that π_n^ρ weakly converges to ν_{π^ρ} as $n \rightarrow \infty$ in $(\Gamma \cup \partial\Gamma)^2$ for each $\rho \in [0, 1]$. Let us define

$$\mathbf{bnd} := \mathbf{bnd}^{(1)} \times \mathbf{bnd}^{(2)} : \Omega \times \Omega \rightarrow \partial\Gamma \times \partial\Gamma,$$

where each $\mathbf{bnd}^{(i)} : \Omega \rightarrow \partial\Gamma$ for $i = 1, 2$ is the map constructed for a single Γ . We have $\nu_\pi = \mathbf{bnd}_* \mathbf{P}_\pi$. If in addition π has a finite first moment, then $(\partial\Gamma \times \partial\Gamma, \nu_\pi)$ is a Poisson boundary for $(\Gamma \times \Gamma, \pi)$.

Theorem 2.2. *If Γ is a non-elementary word hyperbolic group and π is a probability measure on $\Gamma \times \Gamma$ with finite first moment such that both pushforwards under the projections onto factors are μ whose support generates Γ as a semigroup, then $(\partial\Gamma \times \partial\Gamma, \nu_\pi)$ is a Poisson boundary for $(\Gamma \times \Gamma, \pi)$.*

This is a direct consequence of the ray (or strip) criterion due to [Kai00, Theorem 5.5]. We use the consequence of this fact: The invariant σ -field \mathcal{I} and $\sigma(\mathbf{bnd})$ coincide after taking completions with respect to \mathbf{P}_π .

Proof of Theorem 2.1. Let $\varphi(\mathbf{x}) := \varphi(x^{(1)}) - \varphi(x^{(2)})$ for $\mathbf{x} = (x^{(1)}, x^{(2)}) \in \Gamma \times \Gamma$. For a π^ρ -random walk $\{\mathbf{w}_n\}_{n \in \mathbb{Z}_+}$ starting from id on $\Gamma \times \Gamma$, the process $\{\varphi(\mathbf{w}_n)\}_{n \in \mathbb{Z}_+}$ defines a $\varphi_*\pi^\rho$ -random walk on \mathbb{Z} . Note that the pushforward $\varphi_*\pi^\rho$ has a finite second moment since φ is a homomorphism. It holds that $\mathbf{E}_{\pi^\rho}\varphi(\mathbf{x}) = 0$ and the variance

$$\mathrm{Var}_{\pi^\rho}(\varphi) = \mathbf{E}_{\pi^\rho}(\varphi(x^{(1)}) - \varphi(x^{(2)}))^2 = 2\rho v(\mu, \varphi),$$

where $v(\mu, \varphi) := \mathrm{Var}_\mu(\varphi)$. We have that $v(\mu, \varphi) > 0$ since the homomorphism φ is surjective. The Hartman-Wintner law of the iterated logarithm (cf. [MP10, Theorem 5.17]) reads the following: Letting

$$\mathrm{LIL}(\rho) := \left\{ \{\mathbf{w}_n\}_{n \in \mathbb{Z}_+} \in \Omega \times \Omega : \limsup_{n \rightarrow \infty} \frac{\varphi(\mathbf{w}_n)}{\sqrt{n \log \log n}} = \sqrt{2\rho v(\mu, \varphi)} \right\},$$

we have $\mathbf{P}_{\pi^\rho}(\mathrm{LIL}(\rho)) = 1$ for all $\rho \in [0, 1]$. Note that the event $\mathrm{LIL}(\rho)$ belongs to the invariant σ -field \mathcal{I} .

Fix $\rho, \rho_* \in [0, 1]$ with $\rho \neq \rho_*$. Since $(\partial\Gamma \times \partial\Gamma, \nu_{\pi^\rho})$ is a Poisson boundary for $(\Gamma \times \Gamma, \pi^\rho)$ by Theorem 2.2, the completions of the invariant σ -field \mathcal{I} and $\sigma(\mathbf{bnd})$ under \mathbf{P}_{π^ρ} coincide. In particular, since $\mathrm{LIL}(\rho) \in \mathcal{I}$, there exist Borel sets $B_0, B_1 \in \mathcal{B}(\partial\Gamma \times \partial\Gamma)$ such that $\mathbf{bnd}^{-1}(B_0) \subset \mathrm{LIL}(\rho) \subset \mathbf{bnd}^{-1}(B_1)$ and $\nu_{\pi^\rho}(B_1 \setminus B_0) = 0$. It holds that

$$\nu_{\pi^\rho}(B_0) = \mathbf{P}_{\pi^\rho}(\mathbf{bnd}^{-1}(B_0)) = \mathbf{P}_{\pi^\rho}(\mathrm{LIL}(\rho)) = 1,$$

and since $\rho \neq \rho_*$,

$$\nu_{\pi^{\rho_*}}(B_0) = \mathbf{P}_{\pi^{\rho_*}}(\mathbf{bnd}^{-1}(B_0)) \leq \mathbf{P}_{\pi^{\rho_*}}(\mathrm{LIL}(\rho)) = 0.$$

Therefore we have

$$\begin{aligned} \|\nu_{\pi^\rho} - \nu_{\pi^{\rho_*}}\|_{\mathrm{TV}} &= \sup\{|\nu_{\pi^\rho}(B) - \nu_{\pi^{\rho_*}}(B)| : B \in \mathcal{B}(\partial\Gamma \times \partial\Gamma)\} \\ &\geq |\nu_{\pi^\rho}(B_0) - \nu_{\pi^{\rho_*}}(B_0)| = 1, \end{aligned}$$

whence $\|\nu_{\pi^\rho} - \nu_{\pi^{\rho_*}}\|_{\mathrm{TV}} = 1$. Recall that π_n^ρ converges weakly to ν_{π^ρ} as $n \rightarrow \infty$ in $(\Gamma \cup \partial\Gamma)^2$, which is a compact metrizable space. Hence it follows that for $\rho \neq \rho_*$,

$$\liminf_{n \rightarrow \infty} \|\pi_n^\rho - \pi_n^{\rho_*}\|_{\mathrm{TV}} \geq \|\nu_{\pi^\rho} - \nu_{\pi^{\rho_*}}\|_{\mathrm{TV}} = 1.$$

This concludes the first claim by noting that the total variation distance between two probability measures is at most 1. The second claim follows since π^{ρ_*} equals $\mu \times \mu$ for $\rho_* = 1$. \square

2.2. Noise stability on lamplighter groups. For (countable) groups G and Λ , let $G \wr \Lambda := (\bigoplus_\Lambda G) \rtimes \Lambda$ be the (restricted) wreath product of G and Λ , where the product is defined by

$$((\Psi(x))_{x \in \Lambda}, z) \cdot ((\Psi'(x))_{x \in \Lambda}, z') = ((\Psi(x) \cdot \Psi'(z^{-1}x))_{x \in \Lambda}, z \cdot z')$$

for $((\Psi(x))_{x \in \Lambda}, z)$ and $((\Psi'(x))_{x \in \Lambda}, z')$ in $G \wr \Lambda$. The group $G \wr \Lambda$ is called the **lamplighter group** with the lamp group G over the base group Λ . If G and Λ are finitely generated, then $G \wr \Lambda$ is finitely generated, in which case $|\cdot|$ denotes a word metric. For a probability measure μ on a countable group Γ , the **entropy** of μ is defined by $H(\mu) := -\sum_{x \in \Gamma} \mu(x) \log \mu(x)$, where $0 \log 0 := 0$.

Theorem 2.3. *Let G be a nontrivial finite group, m be a positive integer, and μ be a switch-or-walk probability measure on $G \wr \mathbb{Z}^m$ such that $H(\mu) < \infty$ and the support of μ generates the group as a semigroup. If the projection μ_{base} of μ on \mathbb{Z}^m defines a transient random walk on \mathbb{Z}^m and μ_{base} has a finite second moment, then*

$$\lim_{n \rightarrow \infty} \|\pi_n^\rho - \pi_n^{\rho_*}\|_{\text{TV}} = 1 \quad \text{for all } \rho, \rho_* \in [0, 1] \text{ with } \rho \neq \rho_*.$$

In particular,

$$\lim_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} = 1 \quad \text{for all } \rho \in [0, 1),$$

i.e., the μ -random walk on $G \wr \mathbb{Z}^m$ is noise stable in total variation.

For a probability measure μ on $G \wr \Lambda$, let $\{w_n\}_{n \in \mathbb{Z}_+}$ be the μ -random walk on $G \wr \Lambda$ such that $w_0 = \text{id}$. The probability space is denoted by $(\Omega, \mathcal{F}, \mathbf{P}_\mu)$ where $\Omega = \Gamma^{\mathbb{Z}_+}$. For $\{w_n\}_{n \in \mathbb{Z}_+} \in \Omega$, letting $w_n = (\Psi_n, z_n)$, we define

$$\Omega_{\text{conv}} := \left\{ \{w_n\}_{n \in \mathbb{Z}_+} \in \Omega : \text{the pointwise limit of } \{\Psi_n\}_{n \in \mathbb{Z}_+} \text{ exists} \right\},$$

and $\Psi_\infty := \lim_{n \rightarrow \infty} \Psi_n$ for $\{w_n\}_{n \in \mathbb{Z}_+} \in \Omega_{\text{conv}}$. Let G^Λ be the product space endowed with the natural product topology (where G is equipped with the discrete topology) and the associated Borel σ -field $\mathcal{B}(G^\Lambda)$. If $\mathbf{P}_\mu(\Omega_{\text{conv}}) = 1$, i.e., Ψ_∞ is defined almost surely in \mathbf{P}_μ , then the distribution of Ψ_∞ on G^Λ is denoted by ν_μ . For various classes of $G \wr \Lambda$ and μ , it is shown that $\mathbf{P}_\mu(\Omega_{\text{conv}}) = 1$ and (G^Λ, ν_μ) is a Poisson boundary for $(G \wr \Lambda, \mu)$ in [LP21]. Among other results, one reads the following:

Theorem 2.4 (Theorem 4.7 in [LP21]). *Let G be a nontrivial finite group and Λ be a finitely generated, infinite group with an abelian subgroup of finite index. Let μ be a switch-or-walk probability measure of finite entropy on $G \wr \Lambda$ whose support generates $G \wr \Lambda$ as a semigroup. If the projection μ_{base} of μ on Λ defines a transient random walk, then G^Λ endowed with the law ν_μ of Ψ_∞ is a Poisson boundary for $(G \wr \Lambda, \mu)$.*

The case which we focus on is when $\Lambda = \mathbb{Z}^m$ for $m \geq 1$ (see also [LP21, Theorem 4.6] and a special case in [LP16, Theorem 14.39]). The proof uses the following criterion developed upon the one due to Kaimanovich [Kai00].

Corollary 2.5 (Corollary 2.2 in [LP21]). *Let $\{w_n\}_{n \in \mathbb{Z}_+}$ be a μ -random walk on Γ with $H(\mu) < \infty$. Let \mathcal{I} be the associated invariant σ -field and $\mathcal{J} \subseteq \mathcal{I}$ be a sub- σ -field such that $x.A \in \mathcal{J}$ for all $x \in \Gamma$ and all $A \in \mathcal{J}$. If for each $\varepsilon > 0$, there exists a random sequence $\{Q_{n,\varepsilon}\}_{n \in \mathbb{Z}_+}$ of finite subsets of Γ such that*

- (i) $Q_{n,\varepsilon}$ is \mathcal{J} -measurable,
- (ii) $\limsup_{n \rightarrow \infty} (1/n) \log \#Q_{n,\varepsilon} < \varepsilon$ almost everywhere in \mathbf{P}_μ , and
- (iii) $\limsup_{n \rightarrow \infty} \mathbf{P}_\mu(w_n \in Q_{n,\varepsilon}) > 0$,

then $\mathcal{J} = \mathcal{I}$ modulo \mathbf{P}_μ -measure null sets.

We use the corresponding result on a π^ρ -random walk on $(G \wr \mathbb{Z}^m) \times (G \wr \mathbb{Z}^m)$. Note that in this case each marginal on each factor is a μ -random walk on $G \wr \mathbb{Z}^m$. If a μ_{base} -random walk on \mathbb{Z}^m is transient, then $\Psi_\infty := (\Psi_\infty^{(1)}, \Psi_\infty^{(2)})$ is defined almost surely as the pointwise

limit of $\{(\Psi_n^{(1)}, \Psi_n^{(2)})\}_{n \in \mathbb{Z}_+}$ in $G^{\mathbb{Z}^m} \times G^{\mathbb{Z}^m}$. The analogous claim to Theorem 2.4 holds for $((G \wr \mathbb{Z}^m) \times (G \wr \mathbb{Z}^m), \pi^\rho)$. This follows from the proof of Theorem 2.4 and Corollary 2.5. We provide only a sketch of the proof for the sake of convenience.

Theorem 2.6. *Let G be a nontrivial finite group and m be a positive integer. Let μ be a switch-or-walk probability measure of finite entropy on $G \wr \mathbb{Z}^m$ and the support generates $G \wr \mathbb{Z}^m$ as a semigroup. If the projection μ_{base} of μ on \mathbb{Z}^m defines a transient random walk, then for each $\rho \in [0, 1]$ the π^ρ -random walk on $(G \wr \mathbb{Z}^m) \times (G \wr \mathbb{Z}^m)$, the space $G^{\mathbb{Z}^m} \times G^{\mathbb{Z}^m}$ endowed with the law ν_{π^ρ} of $(\Psi_\infty^{(1)}, \Psi_\infty^{(2)})$ is the Poisson boundary for $((G \wr \mathbb{Z}^m) \times (G \wr \mathbb{Z}^m), \pi^\rho)$.*

A sketch of proof. The proof of Theorem 2.4 (Theorem 4.7 in [LP21]) shows the following: For each $\varepsilon > 0$, there exists a sequence $\{Q_{n,\varepsilon}\}_{n \in \mathbb{Z}_+}$ of finite subsets in $G \wr \mathbb{Z}^m$ such that

- (i) $Q_{n,\varepsilon}$ is $\sigma(\Psi_\infty)$ -measurable,
- (ii) $\limsup_{n \rightarrow \infty} (1/n) \log \#Q_{n,\varepsilon} < \varepsilon$ almost surely in \mathbf{P}_μ , and
- (iii) $\lim_{n \rightarrow \infty} \mathbf{P}_\mu(w_{t_0 \cdot n} \in Q_{t_0 \cdot n, \varepsilon}) = 1$ for a positive integer t_0 .

This in particular implies that $\sigma(\Psi_\infty) = \mathcal{I}$ modulo \mathbf{P}_μ -measure null sets by Corollary 2.5. Let $\rho \in [0, 1]$. For a π^ρ -random walk $\{\mathbf{w}_n\}_{n \in \mathbb{Z}_+}$ on $(G \wr \mathbb{Z}^m) \times (G \wr \mathbb{Z}^m)$, each $\{w_n^{(i)}\}_{n \in \mathbb{Z}_+}$ for $i = 1, 2$ where $\mathbf{w}_n = (w_n^{(1)}, w_n^{(2)})$ is a μ -random walk on $G \wr \mathbb{Z}^m$ satisfying the assumption in Theorem 2.4. Therefore $\Psi_\infty^{(i)}$ is defined almost surely as a pointwise limit of $\{\Psi_n^{(i)}\}_{n \in \mathbb{Z}_+}$ for $w_n^{(i)} = (\Psi_n^{(i)}, z_n^{(i)})$ on \mathbb{Z}^m for $i = 1, 2$ since $\{z_n^{(i)}\}_{n \in \mathbb{Z}_+}$ is a μ_{base} -random walk which is transient. For each $\varepsilon > 0$, let $Q_{n,\varepsilon}^{(i)}$ denote $\sigma(\Psi_\infty^{(i)})$ -measurable sets satisfying the above (i), (ii) and (iii). Letting $Q_{n,\varepsilon} := Q_{n,\varepsilon}^{(1)} \times Q_{n,\varepsilon}^{(2)}$, we have

- (i) $Q_{n,\varepsilon}$ is $\sigma(\Psi_\infty^{(1)}, \Psi_\infty^{(2)})$ -measurable,
- (ii) $\limsup_{n \rightarrow \infty} (1/n) \log \#Q_{n,\varepsilon} < 2\varepsilon$ almost surely in \mathbf{P}_{π^ρ} , and
- (iii) $\lim_{n \rightarrow \infty} \mathbf{P}_{\pi^\rho}(\mathbf{w}_{t_0 \cdot n} \in Q_{t_0 \cdot n, \varepsilon}) = 1$,

where t_0 is the same positive integer as above. To obtain (iii), we have used by (iii),

$$\mathbf{P}_{\pi^\rho}(\mathbf{w}_{t_0 \cdot n} \notin Q_{t_0 \cdot n, \varepsilon}) \leq \mathbf{P}_\mu(w_{t_0 \cdot n}^{(1)} \notin Q_{t_0 \cdot n, \varepsilon}^{(1)}) + \mathbf{P}_\mu(w_{t_0 \cdot n}^{(2)} \notin Q_{t_0 \cdot n, \varepsilon}^{(2)}) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Therefore Corollary 2.5 (Corollary 2.2 in [LP21]) concludes the claim. \square

Let $\Omega := ((G \wr \mathbb{Z}^m) \times (G \wr \mathbb{Z}^m))^{\mathbb{Z}_+}$ and

$$\Omega_{\text{conv}} := \left\{ \{\mathbf{w}_n\}_{n \in \mathbb{Z}_+} \in \Omega : \text{the pointwise limit of } \{\Psi_n^{(i)}\}_{n \in \mathbb{Z}_+} \text{ exists for } i = 1, 2 \right\}.$$

Let us define

$$\mathbf{bnd} : \Omega \rightarrow G^{\mathbb{Z}^m} \times G^{\mathbb{Z}^m}, \quad \mathbf{bnd}(\mathbf{w}) := \Psi_\infty = (\Psi_\infty^{(1)}, \Psi_\infty^{(2)}),$$

for $\mathbf{w} = \{(\Psi_n, \mathbf{z}_n)\}_{n \in \mathbb{Z}_+} \in \Omega_{\text{conv}}$ and $\mathbf{bnd}(\mathbf{w}) = \xi$ for $\mathbf{w} \in \Omega \setminus \Omega_{\text{conv}}$ where ξ is an arbitrary fixed element in $G^{\mathbb{Z}^m} \times G^{\mathbb{Z}^m}$ (e.g., $\xi = (\mathbf{id}, \mathbf{id})$).

Proof of Theorem 2.3. Let $\Gamma := G \wr \mathbb{Z}^m$ where $m \geq 1$. For the π^ρ -random walk $\mathbf{w}_n = (\Psi_n, \mathbf{z}_n)$ and for $\rho, \rho_* \in [0, 1]$, we have by the definition of the total variation distance,

$$\|\pi_n^\rho - \pi_n^{\rho_*}\|_{\text{TV}} \geq \|\mathbf{P}_{\pi^\rho}(\Psi_n \in \cdot) - \mathbf{P}_{\pi^{\rho_*}}(\Psi_n \in \cdot)\|_{\text{TV}} \quad \text{for all } n \in \mathbb{Z}_+.$$

Therefore,

$$\liminf_{n \rightarrow \infty} \|\pi_n^\rho - \pi_n^{\rho_*}\|_{\text{TV}} \geq \|\mathbf{P}_{\pi^\rho}(\Psi_\infty \in \cdot) - \mathbf{P}_{\pi^{\rho_*}}(\Psi_\infty \in \cdot)\|_{\text{TV}} = \|\nu_{\pi^\rho} - \nu_{\pi^{\rho_*}}\|_{\text{TV}}. \quad (2.1)$$

This follows since $\Psi_n \rightarrow \Psi_\infty$ as $n \rightarrow \infty$ almost surely in \mathbf{P}_{π^ρ} , and the distributions of Ψ_n weakly converges to the one of Ψ_∞ on the compact metrizable space $G^{\mathbb{Z}^m} \times G^{\mathbb{Z}^m}$ for ρ and ρ_* , respectively. Let us fix a surjective homomorphism $\varphi : \mathbb{Z}^m \rightarrow \mathbb{Z}$ (e.g., the projection onto the first coordinate). For $\mathbf{x} = (x^{(1)}, x^{(2)}) \in \mathbb{Z}^m \times \mathbb{Z}^m$, we write $\varphi(\mathbf{x}) = \varphi(x^{(1)}) - \varphi(x^{(2)})$. For each $\rho \in [0, 1]$, let

$$\text{LIL}(\rho) := \left\{ \{\mathbf{w}_n\}_{n \in \mathbb{Z}_+} \in (\Gamma \times \Gamma)^{\mathbb{Z}_+} : \limsup_{n \rightarrow \infty} \frac{\varphi(\mathbf{w}_n)}{\sqrt{n \log \log n}} = \sqrt{2\rho v(\mu, \varphi)} \right\},$$

where $v(\mu, \varphi) := \text{Var}_\mu(\varphi)$ and $\text{Var}_{\pi^\rho}(\varphi) = 2\rho v(\mu, \varphi)$. We have $v(\mu, \varphi) > 0$ by the assumption that the support of μ generates Γ as a semigroup. Note that $\mathbf{P}_{\pi^\rho}(\text{LIL}(\rho)) = 1$ for all $\rho \in [0, 1]$ by the Hartman-Wintner law of the iterated logarithm (cf. [MP10, Theorem 5.17]). Furthermore, $\text{LIL}(\rho)$ is in the invariant σ -field for every $\rho \in [0, 1]$. By Theorem 2.6, $(G^{\mathbb{Z}^m} \times G^{\mathbb{Z}^m}, \nu_{\pi^\rho})$ is the Poisson boundary for $(\Gamma \times \Gamma, \pi^\rho)$ for each $\rho \in [0, 1]$. This implies that there exist B_0 and B_1 in $\mathcal{B}((G \times G)^{\mathbb{Z}^m \times \mathbb{Z}^m})$ such that

$$\mathbf{bnd}^{-1}(B_0) \subset \text{LIL}(\rho) \subset \mathbf{bnd}^{-1}(B_1) \quad \text{and} \quad \mathbf{P}_{\pi^\rho}(\mathbf{bnd}^{-1}(B_1) \setminus \mathbf{bnd}^{-1}(B_0)) = 0.$$

Therefore for $\rho, \rho_* \in [0, 1]$ with $\rho \neq \rho_*$,

$$\mathbf{P}_{\pi^\rho}(\mathbf{bnd}^{-1}(B_0)) = \mathbf{P}_{\pi^\rho}(\text{LIL}(\rho)) = 1 \quad \text{and} \quad \mathbf{P}_{\pi^{\rho_*}}(\mathbf{bnd}^{-1}(B_0)) \leq \mathbf{P}_{\pi^{\rho_*}}(\text{LIL}(\rho)) = 0.$$

This shows that

$$\|\nu_{\pi^\rho} - \nu_{\pi^{\rho_*}}\|_{\text{TV}} = \sup \{ |\nu_{\pi^\rho}(B) - \nu_{\pi^{\rho_*}}(B)| : B \in \mathcal{B}(G^{\mathbb{Z}^m} \times G^{\mathbb{Z}^m}) \} = 1,$$

concluding the first claim by (2.1), and the second claim by setting $\rho_* = 1$. \square

2.3. Finite Cayley graphs. Let $G = (V(G), E(G))$ be a finite graph. All graphs we discuss in this subsection are unoriented, simple and with no loops. For a sequence of vertices $\{x_i\}_{i=0}^n$, we say that it is a path if $\{x_i, x_{i+1}\} \in E(G)$ for all $0 \leq i < n$, and that it is a cycle if in addition to that $x_0 = x_n$. The length of a path or a cycle $\{x_i\}_{i=0}^n$ is n . The **girth** of G denoted by $\mathbf{girth}(G)$ is the minimum length of cycles $\{x_i\}_{i=0}^n$ such that $x_i \neq x_j$ for $i \neq j$ and $0 \leq i, j < n$ with $n > 2$. It is defined as ∞ if there is no such cycle. A family \mathcal{G} of finite (connected) graphs is said to have **logarithmically large girths** if there exists a constant $c > 0$ such that $\mathbf{girth}(G) \geq c \log \#V(G)$ for all $G \in \mathcal{G}$.

For a finitely generated group Γ with a finite set of generators S invariant under involution $s \mapsto s^{-1}$, let $\text{Cay}(\Gamma, S)$ be the corresponding (right) Cayley graph (see the beginning of Section 3). The simple random walk $\{w_n\}_{n \in \mathbb{Z}_+}$ with $w_0 = \text{id}$ on a Cayley graph $\text{Cay}(\Gamma, S)$ is the μ -random walk on Γ with the uniform distribution μ on S starting from id .

For a finite index normal subgroup N , let $S_{N \setminus \Gamma}$ denote the image of S under the canonical surjection $\Pi_N : \Gamma \rightarrow N \setminus \Gamma$, and let us consider the associated Cayley graph on the finite group,

$$G_S(N \setminus \Gamma) := \text{Cay}(N \setminus \Gamma, S_{N \setminus \Gamma}).$$

The injectivity radius $\mathbf{inj}(G_S(N\backslash\Gamma))$ is the maximum $r \in \mathbb{Z}_+$ such that Π_N restricted on the ball of radius r (centered at id in the graph distance) in $\text{Cay}(\Gamma, S)$ is injective. Let us say that a family of graphs \mathcal{G} of the form $G_S(N\backslash\Gamma)$ has **large injectivity radii** if there exists a function $R : \mathbb{N} \rightarrow \mathbb{N}$ such that $R(n) \rightarrow \infty$ as $n \rightarrow \infty$ and $\mathbf{inj}(G) \geq R(\#V)$ for all $G \in \mathcal{G}$. If in particular $R(n) \geq c \log n$ for some fixed constant $c > 0$, then we say that \mathcal{G} has **logarithmically large injectivity radii**. Since the Cayley graph of a free group Γ with a free generator S is a tree, every family of the form $G_S(N\backslash\Gamma)$ of logarithmically large injectivity radii has logarithmically large girths. Explicit examples of such families have been constructed in [LPS88] and [AB22].

Corollary 2.7. *Let Γ be a word hyperbolic group as in Theorem 2.1 and \mathcal{G} be a family of finite Cayley graphs of the form $G_S(N\backslash\Gamma)$ with large injectivity radii. For the μ -random walk on $N\backslash\Gamma$ starting from the identity such that the support of μ coincides with $S_{N\backslash\Gamma}$, for all $\rho, \rho_* \in [0, 1]$ with $\rho \neq \rho_*$,*

$$\lim_{n \rightarrow \infty} \|\pi_{R(n)}^\rho - \pi_{R(n)}^{\rho_*}\|_{\text{TV}} = 1,$$

where $n := \#(N\backslash\Gamma)$ for some $R(n)$ such that $R(n) \rightarrow \infty$ as $n \rightarrow \infty$. In particular, if \mathcal{G} has logarithmically large injectivity radii, then there exists a constant $c > 0$ such that for all $\rho \in [0, 1]$,

$$\lim_{n \rightarrow \infty} \|\pi_{\lfloor c \log n \rfloor}^\rho - \mu_{\lfloor c \log n \rfloor} \times \mu_{\lfloor c \log n \rfloor}\|_{\text{TV}} = 1.$$

Proof. Let us define a lift of π^ρ -random walk for μ on $N\backslash\Gamma \times N\backslash\Gamma$ to $\Gamma \times \Gamma$: If $\mathbf{inj}(G_S(N\backslash\Gamma))$ is positive, then there exists a probability measure $\tilde{\mu}$ on S in Γ such that μ with the support $S_{N\backslash\Gamma}$ is obtained as the pushforward of $\tilde{\mu}$ under the map $\Pi_N : \Gamma \rightarrow N\backslash\Gamma$. Letting $\tilde{\pi}^\rho$ be the one defined for $\tilde{\mu}$ for $\rho \in [0, 1]$, we have a $\tilde{\pi}^\rho$ -random walk $\{\tilde{\mathbf{w}}_n\}_{n=0}^\infty$ on $\Gamma \times \Gamma$ starting from id and a π^ρ -random walk $\{\mathbf{w}_n\}_{n=0}^\infty$ starting from id on $N\backslash\Gamma \times N\backslash\Gamma$ is obtained by $\mathbf{w}_n = (\Pi_N \times \Pi_N)(\tilde{\mathbf{w}}_n)$. On the Cayley graph on $\Gamma \times \Gamma$ with respect to $S \times S$, the $\tilde{\pi}^\rho$ -random walk changes graph distance by one at each step and similarly for the π^ρ -random walk on $N\backslash\Gamma \times N\backslash\Gamma$. Let $r \leq \mathbf{inj}(G_S(N\backslash\Gamma))$. Since $\Pi_N \times \Pi_N : \Gamma \times \Gamma \rightarrow N\backslash\Gamma \times N\backslash\Gamma$ is injective on the ball of radius r centered at id in $\text{Cay}(\Gamma \times \Gamma, S \times S)$, it holds that for $m = R(\#(N\backslash\Gamma))$,

$$\|\pi_m^\rho - \pi_m^{\rho_*}\|_{\text{TV}} = \|\tilde{\pi}_m^\rho - \tilde{\pi}_m^{\rho_*}\|_{\text{TV}} \quad \text{for all } \rho, \rho_* \in [0, 1].$$

Therefore since $\tilde{\mu}$ has a finite support which generates Γ as a semigroup, Theorem 2.1 on the $\tilde{\pi}^\rho$ -random walk yields the first claim for a large injectivity radii family. The second claim follows since $R(n) \geq c \log n$ for a fixed constant $c > 0$ for a family of logarithmically large injectivity radii. \square

3. NOISE SENSITIVITY

For a group Γ and for a subset A in Γ , we write $\Gamma = \langle A \rangle$ if Γ is generated by A as a *semigroup*, i.e., every element in Γ is obtained as a product of some finite sequence of elements from A . Let Γ be a finitely generated group with a finite symmetric set of generators S , i.e., $\Gamma = \langle S \rangle$ and S is invariant under the map $s \mapsto s^{-1}$. It holds that, in fact, $s = s^{-1}$ for all $s \in S$ if S consists of involutions. (This is the case of an (affine)

Weyl group (Γ, S) in the following discussion.) Let $\text{Cay}(\Gamma, S)$ be the (right) Cayley graph of Γ with respect to S , i.e., the set of vertices is Γ and an edge $\{x, y\}$ is defined if and only if $x^{-1}y \in S$. Since S is invariant under $s \mapsto s^{-1}$, the Cayley graph is defined as an undirected graph. For $x \in \Gamma$, let $|x|_S$ denote the word norm with respect to S , i.e., the graph distance between id and x in $\text{Cay}(\Gamma, S)$.

3.1. Affine Weyl groups. Let (Γ, S) be an **affine Weyl group** where S is a canonical finite set of generators, consisting of involutions, i.e., $s^2 = \text{id}$ for every $s \in S$. The group Γ admits a semi-direct product structure $\Gamma = \Lambda \rtimes W$ where the subgroup W called the **spherical Weyl group** is finite and the normal subgroup Λ is isomorphic to a free abelian group of finite rank. For a thorough background on the subject, we refer to [AB08].

The group Γ is equipped with an isometric action on the standard Euclidean space \mathbb{R}^m for some $m \geq 1$, where each generator $s \in S$ acts as a reflection with respect to an affine hyperplane. The action is properly discontinuous and admits a relatively compact convex fundamental domain with nonempty interior C_0 called a **chamber**. The group Γ acts on the set of chambers $\mathcal{C} := \{xC_0\}_{x \in \Gamma}$ simply transitively, i.e., for all $C_1, C_2 \in \mathcal{C}$ there exists $x \in \Gamma$ such that $C_1 = xC_2$, and if $xC_0 = C_0$, then $x = \text{id}$. The normal subgroup Λ acts freely (i.e., without fixed points) as translations on \mathbb{R}^m . The Λ -orbit of the origin is a lattice:

$$\{a_1v_1 + \cdots + a_mv_m : a_1, \dots, a_m \in \mathbb{Z}\},$$

where v_1, \dots, v_m form a basis in \mathbb{R}^m . We identify Λ with the lattice in \mathbb{R}^m . Note that the action of W preserves Λ .

The affine Weyl group (Γ, S) is called **reducible** if there exist nontrivial affine Weyl groups generated by S_1 and S_2 respectively with $S = S_1 \times \{\text{id}\} \cup \{\text{id}\} \times S_2$ for which $\Gamma = \langle S_1 \rangle \times \langle S_2 \rangle$, and **irreducible** otherwise. The group Γ we consider is possibly (and basically) reducible. Irreducible ones are completely classified in terms of root systems. For example, the affine Weyl group of type \tilde{A}_1 is the infinite dihedral group

$$\langle s_1, s_2 \mid s_1^2 = s_2^2 = \text{id} \rangle,$$

where s_1 and s_2 act as reflections with respect to 0 (the origin) and 1 respectively in \mathbb{R} . A chamber has the form of interval $[0, 1)$.

Let us fix a point o in the interior of a chamber and define

$$\Phi : \Gamma \rightarrow \mathbb{R}^m, \quad x \mapsto x.o.$$

The map Φ is injective since Γ acts on the set of chambers simply transitively, and is Γ -equivariant, i.e., $\Phi(xy) = x.\Phi(y)$ for all $x, y \in \Gamma$. Let us call $\Phi : \Gamma \rightarrow \mathbb{R}^m$ an associated **equivariant embedding**. Since the generators act as reflections with respect to affine hyperplanes, it is illustrative to consider that $\text{Cay}(\Gamma, S)$ is realized in \mathbb{R}^m via the map Φ . Namely, the vertices are placed insides of the chambers as the orbit $x.o$ for $x \in \Gamma$ and an edge is a line segment connecting two vertices for which one is obtained from the other by a reflection of the form xsx^{-1} for $s \in S$ and $x \in \Gamma$.

The group Λ itself acts on $\text{Cay}(\Gamma, S)$ from left freely as automorphisms of the graph. Let us consider the quotient graph $G = \Lambda \backslash \text{Cay}(\Gamma, S)$. The graph $G = (V(G), E^{un}(G))$ is

finite, the set of vertices $V(G)$ is W and the set of edges $E^{un}(G)$ consists of *undirected* edges. Note, however, that G is *not* the right Cayley graph of W with respect to the image \overline{S} of S under the quotient map $\Lambda \rtimes W \rightarrow W$. This is because the quotient map restricted on S is not bijective onto \overline{S} . The graph G has possibly multiple edges.

3.2. Pointed finite networks as quotients. The main interest is on an affine Weyl group Γ and the canonical set of generators S . It is, however, useful to discuss a slightly more general setting. Let Γ be a virtually finite rank free abelian group, i.e., Γ admits a finite rank free abelian group Λ as a finite index subgroup. We assume that Γ acts on an \mathbb{R}^m isometrically with a relatively compact fundamental domain with nonempty interior, and that Λ acts as translations and is identified with a lattice in \mathbb{R}^m . Let us fix a point o in the interior of such a fundamental domain of Γ and define $\Phi : \Gamma \rightarrow \mathbb{R}^m$ by $x \mapsto x.o$. The map Φ is equivariant with Γ -actions and injective. For a finite symmetric set of generators S in Γ , let

$$G := \Lambda \backslash \text{Cay}(\Gamma, S).$$

The quotient $G = (V(G), E^{un}(G))$ is a finite (undirected) graph possibly with multiple edges (whence a multi-graph) and with loops. It holds that $V(G) = \Lambda \backslash \Gamma$ and

$$E^{un}(G) = \{\{x, x.s\} : x \in \Lambda \backslash \Gamma, s \in S\},$$

where $\{x, x.s\}$ and $\{x.s, x\}$ are identified.

Let μ be a probability measure on Γ such that the support $\text{supp } \mu$ of μ is finite, that $\Gamma = \langle \text{supp } \mu \rangle$, and that μ is **symmetric**, i.e., $\mu(s) = \mu(s^{-1})$ for every $s \in \text{supp } \mu$. If we define $S = \text{supp } \mu$, then S is a finite symmetric set of generators, and μ defines a Markov chain on $G = \Lambda \backslash \text{Cay}(\Gamma, S)$. Namely, the transition probability is defined by

$$p(x, x.s) := \mu(s) \quad \text{for } x \in \Lambda \backslash \Gamma \text{ and } s \in S.$$

Note that this Markov chain is irreducible, i.e., it visits every vertex from every other vertex after some time since $\Gamma = \langle \text{supp } \mu \rangle$. Furthermore, it is reversible with respect to the uniform distribution π on the set of vertices $V(G) = \Lambda \backslash \Gamma$. Indeed, since μ is symmetric, it holds that

$$\pi(x)p(x, x.s) = \pi(x.s)p(x.s, x) \quad \text{for } x \in \Lambda \backslash \Gamma \text{ and } s \in S, \quad (3.1)$$

where $\pi(x) = 1/\#V(G)$ for $x \in V(G)$. Let $\overline{\mu}$ denote the pushforward under the map $\Gamma \rightarrow \Lambda \backslash \Gamma$, and $x_0 \in \Lambda \backslash \Gamma$ denote the coset containing id . Note that if $\overline{\mu}(x_0) > 0$, then

$$p(x_0, x_0) = \sum_{s \in S, x_0 = x_0.s} \mu(s) = \overline{\mu}(x_0) > 0.$$

If $\mu(\text{id}) > 0$, then in fact $p(x, x) \geq \mu(\text{id}) > 0$ for every $x \in V(G)$. For each $\{x, y\} \in E^{un}(G)$, let us define the **conductance** by

$$c(x, y) := \pi(x)p(x, y).$$

This is well-defined since $c(x, y) = c(y, x)$ by (3.1). Note that $c(x, y) > 0$ for all $\{x, y\} \in E^{un}(G)$. Let us call (G, c, x_0) the pointed (finite) **network** as the finite multi-graph G

equipped with the conductance $c : E^{un}(G) \rightarrow (0, \infty)$ and the point x_0 (corresponding to the coset containing id).

3.3. Harmonic 1-forms on finite graphs. Let (G, c, x_0) be the pointed finite network. Henceforth it is convenient to consider G as a **graph with orientations** where each edge (and loop) has both possible orientations. Let

$$E(G) := \{(x, y), (y, x) : \{x, y\} \in E^{un}(G)\}.$$

For $e = (x, y)$, we write $\bar{e} = (y, x)$. The “reversing direction” operation $\bar{\cdot} : E(G) \rightarrow E(G)$, $e \mapsto \bar{e}$, defines a bijection and $\bar{\bar{e}} = e$ for $e \in E(G)$. For $e = (x, y) \in E(G)$, let us denote by $oe := x$ the **origin** and by $te := y$ the **terminus** of e respectively. We have that $o\bar{e} = te$ for $e \in E(G)$. Let us also consider $\text{Cay}(\Gamma, S)$ as a graph with orientations, defining both possible orientations for each edge and loop. Letting $c(e) := c(x, y)$ and $p(e) := p(x, y)$ for $e = (x, y)$, we have that $c(e) = c(\bar{e})$ and $c(e) = \pi(oe)p(e)$ for $e \in E(G)$. It holds that by the definition of conductance,

$$\pi(x) = \sum_{e \in E_x} c(e), \quad \text{where } E_x := \{e \in E(G) : oe = x\} \text{ for } x \in V(G).$$

Let us define the \mathbb{C} -linear space of complex-valued functions on $V(G)$ by

$$C^0(G, \mathbb{C}) := \{f : V(G) \rightarrow \mathbb{C}\}$$

equipped with the inner product $\langle f_1, f_2 \rangle_\pi := \sum_{x \in V(G)} f_1(x) \overline{f_2(x)} \pi(x)$, where \bar{a} stands for the complex-conjugate of $a \in \mathbb{C}$. Similarly, let $C^0(G, \mathbb{R})$ be the \mathbb{R} -linear space of real-valued functions on $V(G)$ endowed with the inner product as the restriction of $\langle \cdot, \cdot \rangle_\pi$. Further let us define the \mathbb{R} -linear space of real-valued **1-forms** on $E(G)$ by

$$C^1(G, \mathbb{R}) := \{\omega : E(G) \rightarrow \mathbb{R} : \omega(\bar{e}) = -\omega(e) \text{ for } e \in E(G)\}$$

equipped with the inner product $\langle \omega_1, \omega_2 \rangle_c := (1/2) \sum_{e \in E(G)} \omega_1(e) \omega_2(e) c(e)$. The **differential** $d : C^0(G, \mathbb{R}) \rightarrow C^1(G, \mathbb{R})$ is the \mathbb{R} -linear map defined by

$$df(e) := f(te) - f(oe) \quad \text{for } e \in E(G),$$

and the **adjoint** $d^* : C^1(G, \mathbb{R}) \rightarrow C^0(G, \mathbb{R})$ with respect to the inner products is obtained by

$$d^* \omega(x) := - \sum_{e \in E_x} \frac{1}{\pi(x)} c(e) \omega(e) \quad \text{for } x \in V(G).$$

It holds that for $f \in C^0(G, \mathbb{R})$ and $\omega \in C^1(G, \mathbb{R})$,

$$\langle df, \omega \rangle_c = \langle f, d^* \omega \rangle_\pi. \quad (3.2)$$

Note that if we define the transition operator P on $C^0(G, \mathbb{R})$ to itself by

$$Pf(x) := \sum_{e \in E_x} \frac{1}{\pi(x)} c(e) f(te) \quad \text{for } x \in V(G),$$

then $d^*d = I - P$ where I is the identity operator. Let us define the space of **harmonic 1-forms** by

$$H^1 := \{\omega \in C^1(G, \mathbb{R}) : d^* \omega = 0\}.$$

Note that $H^1 = (\text{Im } d)^\perp$ the orthogonal complement of the image $\text{Im } d$ by (3.2). The fact that we use in the sequel is that for every 1-form $\omega \in C^1(G, \mathbb{R})$ there exists a *unique* harmonic 1-form $u \in H^1$ and some $f \in C^0(G, \mathbb{R})$ such that

$$u + df = \omega.$$

The u is obtained as the H^1 -part in the orthogonal decomposition $C^0(G, \mathbb{R}) = H^1 \oplus \text{Im } d$. Note that f is not unique as every f added a constant function satisfies the relation.

Remark 3.1. If we endow G with a structure of 1-dimensional CW complex, then the 1-cohomology group with real coefficient is defined as $H^1(G, \mathbb{R}) := C^1(G, \mathbb{R})/\text{Im } d$. The fact mentioned above means that every 1-cohomology class is represented by a unique harmonic 1-form. Although all these notions are not needed in our discussion, it might be useful to grasp an idea behind some of our computations.

3.4. Perturbations of transfer operators. For each 1-form $\omega \in C^1(G, \mathbb{R})$, let us define the **transfer operator** on $C^0(G, \mathbb{C})$ by

$$\mathcal{L}_\omega f(x) := \sum_{e \in E_x} p(e) e^{2\pi i \omega(e)} f(te) \quad \text{for } x \in V(G).$$

Here $i = \sqrt{-1}$. In this particular setting where μ is symmetric, the transfer operator is self-adjoint on $(C^0(G, \mathbb{C}), \langle \cdot, \cdot \rangle_\pi)$. Hence it has real eigenvalues. Let $\lambda(\omega)$ be the largest eigenvalue of \mathcal{L}_ω . If $\omega = 0$, then $\lambda(0) = 1$ and this is a simple eigenvalue by the Perron-Frobenius theorem since $\mathcal{L}_0 = P$ and P is irreducible. We apply to an analytic perturbation in ω : For a small enough neighborhood U of 0 in $C^1(G, \mathbb{R})$, the function $U \rightarrow \mathbb{R}$, $\omega \mapsto \lambda(\omega)$ is real analytic. Moreover, corresponding eigenvectors f_ω depend analytically in $\omega \in U$ with $f_0 = \mathbf{1}$ (the constant vector with all 1's). This follows from the Hamilton-Cayley theorem and the implicit function theorem in this finite graph setting. Note that if we consider $\omega + d\varphi$ for $\varphi \in C^0(G, \mathbb{R})$ in place of ω , then $\mathcal{L}_{\omega+d\varphi} = e^{-\varphi} \mathcal{L}_\omega e^{\varphi}$, where $(e^\varphi f)(x) := e^{\varphi(x)} f(x)$ for $x \in V(G)$, and thus

$$\lambda(\omega + d\varphi) = \lambda(\omega).$$

This shows that $\lambda(\omega)$ depends only on the harmonic H^1 -part of ω . Let

$$\beta(\omega) := \log \lambda(\omega).$$

Since $\mathcal{L}_0 = P$ which has a simple eigenvalue $\lambda(0) = 1$, there exists a small enough open neighborhood U of 0 in $C^1(G, \mathbb{R})$ such that $\lambda(\omega)$ is a simple eigenvalue of \mathcal{L}_ω and $\beta(\omega) = \log \lambda(\omega)$ is well-defined for all $\omega \in U$. It holds that $\beta(0) = 0$ since $\lambda(0) = 1$.

Lemma 3.2. *Let $\lambda(\omega) = e^{\beta(\omega)}$ and f_ω be the eigenvalue and the corresponding eigenvector of \mathcal{L}_ω such that $f_0 = \mathbf{1}$ and $\langle f_\omega, \mathbf{1} \rangle_\pi = 1$ for $\omega \in U$ where U is a neighborhood of 0 in $C^1(G, \mathbb{R})$. For all harmonic 1-forms $u, u_i \in H^1$ on G and real parameters r, r_i for $i = 1, 2, 3$, the following holds:*

$$\left. \frac{d}{dr} \right|_{r=0} \beta(ru) = 0, \tag{3.3}$$

$$\left. \frac{d}{dr} \right|_{r=0} f_{ru}(x) = 0 \quad \text{for all } x \in V(G), \tag{3.4}$$

$$\frac{\partial^2}{\partial r_1 \partial r_2} \Big|_{(r_1, r_2) = (0, 0)} \beta(r_1 u_1 + r_2 u_2) = -4\pi^2 \sum_{e \in E(G)} u_1(e) u_2(e) c(e), \quad (3.5)$$

and

$$\frac{\partial^3}{\partial r_1 \partial r_2 \partial r_3} \Big|_{(r_1, r_2, r_3) = (0, 0, 0)} \beta(r_1 u_1 + r_2 u_2 + r_3 u_3) = 0. \quad (3.6)$$

Proof. Let f_ω be the eigenvector normalized as stated. Since $\overline{\mathcal{L}_\omega f_\omega} = \mathcal{L}_{-\omega} \overline{f_\omega}$ holds for all ω and $\lambda(\omega)$ is real for all ω in a small enough neighborhood of 0 in $C^1(G, \mathbb{R})$, it holds that $\lambda(-\omega) = \overline{\lambda(\omega)} = \lambda(\omega)$. Thus $\beta(-\omega) = \beta(\omega)$ for all $\omega \in U$ and all odd time derivatives of β at 0 vanish. This in particular implies (3.3) and (3.6).

For every 1-form u , it holds that $ru \in U$ for all small enough real r and

$$\left\langle \frac{d}{dr} \Big|_{r=0} f_{ru}, \mathbf{1} \right\rangle_\pi = 0. \quad (3.7)$$

Moreover, since P is self-adjoint with respect to $\langle \cdot, \cdot \rangle_\pi$ and $P\mathbf{1} = \mathbf{1}$, by (3.7) it holds that

$$\left\langle P \left(\frac{d}{dr} \Big|_{r=0} f_{ru} \right), \mathbf{1} \right\rangle_\pi = \left\langle \frac{d}{dr} \Big|_{r=0} f_{ru}, P\mathbf{1} \right\rangle_\pi = \left\langle \frac{d}{dr} \Big|_{r=0} f_{ru}, \mathbf{1} \right\rangle_\pi = 0. \quad (3.8)$$

We will also use the analogous identities to (3.7) and (3.8) for the second and third derivatives of the normalization: $\langle f_\omega, \mathbf{1} \rangle_\pi = 1$ for $\omega \in U$.

First differentiating $\mathcal{L}_{ru} f_{ru} = e^{\beta(ru)} f_{ru}$ at $r = 0$ yields for each $x \in V(G)$,

$$\sum_{e \in E_x} \left(p(e)(2\pi i u(e)) + p(e) \frac{d}{dr} \Big|_{r=0} f_{ru}(te) \right) = \frac{d}{dr} \Big|_{r=0} \beta(ru) + \frac{d}{dr} \Big|_{r=0} f_{ru}(x), \quad (3.9)$$

where we have used $f_0 = \mathbf{1}$. Let us note that (3.9) above yields by (3.3) which we have just shown and by that $d^*u = 0$,

$$P \left(\frac{d}{dr} \Big|_{r=0} f_{ru} \right) (x) = \frac{d}{dr} \Big|_{r=0} f_{ru}(x) \quad \text{for each } x \in V(G).$$

Since P has the simple eigenvalue 1, this implies that $(d/dr)|_{r=0} f_{ru}$ is constant. By (3.7), for every harmonic 1-form u , it holds that $(d/dr)|_{r=0} f_{ru}(x) = 0$ for all $x \in V(G)$, showing (3.4).

For all 1-forms u_1 and u_2 and for all small enough reals r_1 and r_2 , it holds that

$$\mathcal{L}_{r_1 u_1 + r_2 u_2} f_{r_1, r_2} = e^{\beta_{r_1, r_2}} f_{r_1, r_2} \quad \text{where } \beta_{r_1, r_2} := \beta(r_1 u_1 + r_2 u_2) \text{ and } f_{r_1, r_2} = f_{r_1 u_1 + r_2 u_2}.$$

Taking the second derivatives at $(r_1, r_2) = (0, 0)$ of both terms yields by (3.3), (3.4) and that $f_0 = \mathbf{1}$, for each $x \in V(G)$,

$$\begin{aligned} \sum_{e \in E_x} \left(p(e)(-4\pi^2 u_1(e) u_2(e)) + p(e) \frac{\partial^2}{\partial r_1 \partial r_2} \Big|_{(r_1, r_2) = (0, 0)} f_{r_1, r_2}(te) \right) \\ = \frac{\partial^2}{\partial r_1 \partial r_2} \Big|_{(r_1, r_2) = (0, 0)} \beta_{r_1, r_2} + \frac{\partial^2}{\partial r_1 \partial r_2} \Big|_{(r_1, r_2) = (0, 0)} f_{r_1, r_2}(x). \end{aligned}$$

Evaluating the inner products of the above terms with $\mathbf{1}$ leads

$$\begin{aligned} -4\pi^2 \sum_{e \in E(G)} c(e)u_1(e)u_2(e) + \left\langle P \left(\frac{\partial^2}{\partial r_1 \partial r_2} \Big|_{(r_1, r_2) = (0, 0)} f_{r_1, r_2} \right), \mathbf{1} \right\rangle_{\pi} \\ = \frac{\partial^2}{\partial r_1 \partial r_2} \Big|_{(r_1, r_2) = (0, 0)} \beta_{r_1, r_2} + \left\langle \frac{\partial^2}{\partial r_1 \partial r_2} \Big|_{(r_1, r_2) = (0, 0)} f_{r_1, r_2}, \mathbf{1} \right\rangle_{\pi}. \end{aligned}$$

The second terms in the left hand side and in the right hand side respectively are 0 since they are the second derivatives on the normalization (cf. (3.7) and (3.8)). This proves (3.5). \square

3.5. An explicit Hessian formula in terms of harmonic 1-forms. Recall that $\Phi : \Gamma \rightarrow \mathbb{R}^m$, $x \mapsto x.o$. Taking a conjugate to the action of Γ by a translation in \mathbb{R}^m , we assume that o is the origin, whence $\Phi(\text{id}) = 0$. The function $(x, y) \mapsto \Phi(y) - \Phi(x)$ for (oriented) edges (x, y) in $\text{Cay}(\Gamma, S)$ is invariant under the action by Λ . Indeed, this follows since Λ is identified with a lattice and acts as translations in \mathbb{R}^m . Therefore this descends to an \mathbb{R}^m -valued function on $E(G)$, which we denote by $e \mapsto \Phi_e$ for $e \in E(G)$. Note that $\Phi_{\bar{e}} = -\Phi_e$ for each $e \in E(G)$. Let $\langle \cdot, \cdot \rangle$ be the standard inner product in \mathbb{R}^m . For each $v \in \mathbb{R}^m$, let

$$\widehat{v}(e) := \langle v, \Phi_e \rangle \quad \text{for } e \in E(G),$$

which defines a 1-form on G . For $v \in \mathbb{R}^m$ near 0, let $\beta(v) := \beta(\widehat{v})$ where $e^{\beta(v)}$ is the largest eigenvalue of the transfer operator $\mathcal{L}_{\widehat{v}}$. Let us define the Hessian of β at 0 in \mathbb{R}^m with the standard coordinate (r_1, \dots, r_m) by

$$\text{Hess}_0 \beta := \left(\frac{\partial^2}{\partial r_k \partial r_l} \Big|_{(r_1, \dots, r_m) = (0, \dots, 0)} \beta(r_1, \dots, r_m) \right)_{k, l = 1, \dots, m}.$$

For the pointed finite network (G, c, x_0) and $\Phi : \Gamma \rightarrow \mathbb{R}^m$, we compute $\text{Hess}_0 \beta$.

Lemma 3.3. *The Hessian $\text{Hess}_0 \beta$ of β at 0 in \mathbb{R}^m is non-degenerate and negative definite. Moreover, it holds that*

$$\langle v_1, \text{Hess}_0 \beta v_2 \rangle = -4\pi^2 \sum_{e \in E(G)} u_1(e)u_2(e)c(e), \quad (3.10)$$

where u_i is defined as the harmonic part of \widehat{v}_i for $v_i \in \mathbb{R}^m$ for $i = 1, 2$.

Proof. For every $v \in \mathbb{R}^m$, we have $\widehat{v}(e) = \langle v, \Phi_e \rangle$ for $e \in E(G)$, and u is the harmonic part of \widehat{v} , i.e., the unique $u \in H^1$ such that $u + df = \widehat{v}$ for some $f \in C^0(G, \mathbb{R})$. Note that the resulting map $v \mapsto u$ is \mathbb{R} -linear. Since $\beta(v)$ depends only on the harmonic part of v , Lemma 3.2 (3.5) implies that

$$\begin{aligned} \langle v_1, \text{Hess}_0 \beta v_2 \rangle &= \frac{\partial^2}{\partial r_1 \partial r_2} \Big|_{(r_1, r_2) = (0, 0)} \beta(r_1 v_1 + r_2 v_2) \\ &= \frac{\partial^2}{\partial r_1 \partial r_2} \Big|_{(r_1, r_2) = (0, 0)} \beta(r_1 u_1 + r_2 u_2) = -4\pi^2 \sum_{e \in E(G)} u_1(e)u_2(e)c(e). \end{aligned}$$

This shows (3.10).

Let v_1, \dots, v_m be a basis of the lattice in \mathbb{R}^m : $\Lambda = \{a_1 v_1 + \dots + a_m v_m : a_1, \dots, a_m \in \mathbb{Z}\}$. For each $v_k = \Phi(v_k) \in \Lambda$ under the identification of Λ with the lattice, there exists a path $(\tilde{e}_1, \dots, \tilde{e}_n)$ from id to v_k in $\text{Cay}(\Gamma, S)$ since the Cayley graph is connected. Let (e_1, \dots, e_n) be the image in G of that path under the covering map from $\text{Cay}(\Gamma, S)$. Note that the image is a cycle: $x_0 = oe_1$, $te_i = oe_{i+1}$ for $i = 1, \dots, n-1$ and $te_n = x_0$. Thus $\sum_{l=1}^n df(e_l) = 0$ and $\sum_{l=1}^n u(e_l) = \sum_{l=1}^n (u(e_l) + df(e_l)) = \sum_{l=1}^n \hat{v}(e_l)$. Furthermore,

$$\sum_{l=1}^n \hat{v}(e_l) = \sum_{l=1}^n \langle v, \Phi_{e_l} \rangle = \sum_{l=1}^n \langle v, \Phi(te_l) - \Phi(oe_l) \rangle = \langle v, \sum_{l=1}^n (\Phi(te_l) - \Phi(oe_l)) \rangle.$$

The last term equals $\langle v, v_k \rangle$ since $\sum_{l=1}^n (\Phi(te_l) - \Phi(oe_l)) = \Phi(te_n) - \Phi(oe_1) = v_k$. Therefore for each $k = 1, \dots, m$ there exists a cycle (e_1, \dots, e_n) in G with $x_0 = oe_1$ and $te_n = x_0$ such that

$$\sum_{l=1}^n u(e_l) = \langle v, v_k \rangle. \quad (3.11)$$

For $v \in \mathbb{R}^m$, let us assume that $\langle v, \text{Hess}_0 \beta v \rangle = 0$. It holds that $u = 0$ by (3.10), and thus $\langle v, v_k \rangle = 0$ for every $k = 1, \dots, m$ by (3.11). Hence $v = 0$ since v_1, \dots, v_m form a basis of a lattice in \mathbb{R}^m . This shows that $\text{Hess}_0 \beta$ is non-degenerate. Furthermore $\text{Hess}_0 \beta$ is negative definite by (3.10). \square

Remark 3.4. Let us consider the Hessian $\text{Hess}_{H^1} \beta$ of β at 0 on H^1 , i.e.,

$$\langle u_1, \text{Hess}_{H^1} \beta u_2 \rangle_\pi = \left. \frac{\partial^2}{\partial r_1 \partial r_2} \right|_{(r_1, r_2) = (0, 0)} \beta(r_1 u_1 + r_2 u_2) \quad \text{for } u_1, u_2 \in H^1,$$

where $H^1 \rightarrow H^1 : u \mapsto \text{Hess}_{H^1} \beta u$ defines an \mathbb{R} -linear map. Lemma 3.2 (3.5) implies that

$$\langle u_1, \text{Hess}_{H^1} \beta u_2 \rangle_\pi = -4\pi^2 \sum_{e \in E(G)} u_1(e) u_2(e) c(e),$$

which shows that $\text{Hess}_{H^1} \beta$ is non-degenerate and negative definite on H^1 . There exists a natural inclusion $\mathbb{R}^m = H^1(\mathbb{R}^m/\Lambda, \mathbb{R}) \rightarrow H^1(G, \mathbb{R})$, represented by $\mathbb{R}^m \rightarrow H^1 : v \mapsto u$ in Lemma 3.3. In this identification, $\text{Hess}_0 \beta$ is the restriction of $\text{Hess}_{H^1} \beta$ to \mathbb{R}^m , and this implies that $\text{Hess}_0 \beta$ is non-degenerate and negative definite. A thorough framework is found in [Sun13]. We use the explicit form of $\text{Hess}_0 \beta$ later in our discussion.

3.6. Local central limit theorems. For every positive integer $n \in \mathbb{Z}_{>0}$, it holds that

$$\mathcal{L}_{\hat{v}}^n \mathbf{1}(x_0) = \sum_{(e_1, \dots, e_n)} p(e_1) \cdots p(e_n) e^{2\pi i (\hat{v}(e_1) + \dots + \hat{v}(e_n))}. \quad (3.12)$$

In the above the summation runs over all directed paths (e_1, \dots, e_n) starting from x_0 in G , i.e., $oe_1 = x_0$ and $te_k = oe_{k+1}$ for each $k = 1, \dots, n-1$. For each such path (e_1, \dots, e_n) , there exists a unique path $(\tilde{e}_1, \dots, \tilde{e}_n)$ which is a lift of the path, starting from id in $\text{Cay}(\Gamma, S)$. By a lift we mean that the path (e_1, \dots, e_n) is the image of $(\tilde{e}_1, \dots, \tilde{e}_n)$ under the covering map $\text{Cay}(\Gamma, S) \rightarrow G = \Lambda \backslash \text{Cay}(\Gamma, S)$. The definition of the 1-form \hat{v} on G

implies that

$$\begin{aligned}\widehat{v}(e_1) + \cdots + \widehat{v}(e_n) &= \langle v, \Phi_{e_1} \rangle + \cdots + \langle v, \Phi_{e_n} \rangle \\ &= \langle v, \Phi(t\tilde{e}_1) - \Phi(o\tilde{e}_1) \rangle + \cdots + \langle v, \Phi(t\tilde{e}_n) - \Phi(o\tilde{e}_n) \rangle = \langle v, \Phi(t\tilde{e}_n) \rangle,\end{aligned}$$

where $\Phi(o\tilde{e}_1) = \Phi(\text{id}) = 0$. Recall that $p(e) = \mu(s)$ for $e = (x, x.s) \in E(G)$ and $s \in S$ in the pointed finite network (G, c, x_0) . By (3.12), it holds that

$$\mathcal{L}_{\widehat{v}}^n \mathbf{1}(x_0) = \sum_{(\tilde{e}_1, \dots, \tilde{e}_n)} \mu(s_1) \cdots \mu(s_n) e^{2\pi i \langle v, \Phi(t\tilde{e}_n) \rangle} = \sum_{x \in \Gamma} \mu_n(x) e^{2\pi i \langle v, \Phi(x) \rangle},$$

where the (edge) path $(\tilde{e}_1, \dots, \tilde{e}_n)$ is represented as the (vertex) path $(\text{id}, s_1, s_1 s_2, \dots, s_1 \cdots s_n)$ in $\text{Cay}(\Gamma, S)$. Therefore letting

$$\varphi_{\mu_n}(v) := \sum_{x \in \Gamma} \mu_n(x) e^{2\pi i \langle v, \Phi(x) \rangle} \quad \text{for } v \in \mathbb{R}^m,$$

we have that for all $v \in \mathbb{R}^m$ and all $n \in \mathbb{Z}_{>0}$,

$$\mathcal{L}_{\widehat{v}}^n \mathbf{1}(x_0) = \varphi_{\mu_n}(v). \quad (3.13)$$

Let Λ^* be the dual lattice of Λ , i.e.,

$$\Lambda^* := \left\{ a_1 v_1^* + \cdots + a_m v_m^* : a_1, \dots, a_m \in \mathbb{Z} \right\},$$

where v_1^*, \dots, v_m^* form the dual basis of v_1, \dots, v_m in \mathbb{R}^m : $\langle v_k^*, v_l \rangle = 1$ if $k = l$ and 0 else. The fundamental parallelepiped of Λ^* in \mathbb{R}^m is denoted by

$$D := \left\{ r_1 v_1^* + \cdots + r_m v_m^* \in \mathbb{R}^m : |r_i| \leq 1/2, i = 1, \dots, m \right\}.$$

The volume of D is assumed to be 1 up to a homothety in \mathbb{R}^m . The Fourier inversion formula shows that for all $n \in \mathbb{Z}_{>0}$,

$$\mu_n(x) = \int_D \varphi_{\mu_n}(v) e^{-2\pi i \langle v, \Phi(x) \rangle} dv \quad \text{for } x \in \Gamma. \quad (3.14)$$

For $\delta > 0$, let

$$D_\delta := \left\{ r_1 v_1^* + \cdots + r_m v_m^* \in \mathbb{R}^m : |r_i| < \delta, i = 1, \dots, m \right\}.$$

Lemma 3.5. *If $p(x_0, x_0) > 0$ in the pointed finite network (G, c, x_0) , then for all small enough $\delta > 0$, there exists a constant $c_\delta > 0$ such that for all $n \in \mathbb{Z}_{>0}$,*

$$|\varphi_{\mu_n}(v)| \leq \sqrt{\#V(G)} \cdot e^{-c_\delta n} \quad \text{for all } v \in D \setminus D_\delta.$$

Proof. This uses a standard perturbation argument; we provide a proof for the sake of completeness. For $v \in \mathbb{R}^m$, let $\|\mathcal{L}_{\widehat{v}}\| := \max_{\|f\|=1} \|\mathcal{L}_{\widehat{v}} f\|$ where $\|\cdot\|$ is the associated norm in $C^0(G, \mathbb{C})$. Since $\mathcal{L}_{\widehat{v}}$ is self-adjoint, the eigenvalues are real and the operator norm $\|\mathcal{L}_{\widehat{v}}\|$ is the spectral radius $|\lambda(v)|$, i.e., the largest eigenvalue in absolute value. Note that $|\lambda(v)| \leq 1$. For $v \in D$, if $|\lambda(v)| = 1$, then the condition $\mu(\text{id}) > 0$ implies that $v = 0$ (in which case in fact $\lambda(0) \neq -1$). Indeed, for the maximal eigenvalue $\lambda(v)$ in absolute value and a corresponding eigenvector f_v , we have $\mathcal{L}_{\widehat{v}} f_v = \lambda(v) f_v$. Taking absolute values shows that $|f_v| \leq P|f_v|$, implying that $|f_v|$ is a non-zero constant since P is irreducible. Further since f_v is an eigenvector with the eigenvalue 1 in absolute value, $\lambda(v) f_v(x)$ and

$e^{2\pi i \langle v, \Phi_e \rangle} f_v(te)$ for $e \in E_x$ are on a common circle in the complex plane for each $x \in V(G)$. Since $\mathcal{L}_{\widehat{v}} f_v = \lambda(v) f_v$, it holds that for all $x \in V(G)$ and for all $e \in E_x$,

$$\lambda(v) f_v(x) = e^{2\pi i \langle v, \Phi_e \rangle} f_v(te). \quad (3.15)$$

If $p(x_0, x_0) > 0$, then for each $v_k \in \Lambda$ there exist a $v' \in \Lambda$ and an edge path (as a lift) from id to $v' + v_k$ in $\text{Cay}(\Gamma, S)$ of length with a given (in particular, even) parity. Applying to (3.15) along the image of the path in G successively yields $\langle v, v' + v_k \rangle \in \mathbb{Z}$, and thus $\langle v, v_k \rangle \in \mathbb{Z}$. This holds for a basis v_1, \dots, v_k of Λ , implying that $v \in \Lambda^*$. Therefore if $v \in D$, then $v = 0$.

We have shown that $|\lambda(v)| < 1$ for all $v \in D \setminus \{0\}$, and in this finite dimensional setting, $v \mapsto \|\mathcal{L}_{\widehat{v}}\| = |\lambda(v)|$ is continuous. Thus for a small enough $\delta > 0$, there exists a constant $c_\delta > 0$ such that $|\lambda(v)| \leq e^{-c_\delta}$ on a compact set $D \setminus D_\delta$. Since

$$\|\mathcal{L}_{\widehat{v}}^n \mathbf{1}(x_0)\| \sqrt{\pi(x_0)} \leq \|\mathcal{L}_{\widehat{v}}^n \mathbf{1}\|_\pi \leq |\lambda(v)|^n \|\mathbf{1}\|_\pi,$$

$\|\mathbf{1}\|_\pi = 1$ and $\pi(x_0) = 1/\#V(G)$, by (3.13), we conclude the claim. \square

For an associated Γ -equivariant embedding $\Phi : \Gamma \rightarrow \mathbb{R}^m$, $x \mapsto x.o$ and a non-degenerate positive definite matrix Σ , let

$$\xi_\Sigma(x) := \frac{1}{(2\pi)^{\frac{m}{2}} \sqrt{\det \Sigma}} e^{-\frac{1}{2} \langle \Phi(x), \Sigma^{-1}(\Phi(x)) \rangle} \quad \text{for } x \in \Gamma.$$

Theorem 3.6 (Local central limit theorem). *Let Γ be a virtually finite rank abelian group acting on \mathbb{R}^m isometrically with a relatively compact fundamental domain which contains the origin o in the interior. Let μ be a probability measure on Γ such that the support $\text{supp } \mu$ is finite, $\Gamma = \langle \text{supp } \mu \rangle$ and μ is symmetric. If the corresponding pointed finite network (G, c, x_0) satisfies that $p(x_0, x_0) > 0$, then the following holds: There exist a non-degenerate positive definite matrix Σ and a constant $C > 0$ such that*

$$\sup_{x \in \Gamma} |\mu_n(x) - \xi_{n\Sigma}(x)| \leq \frac{C}{n^{\frac{m}{2}+1}} \quad \text{for all } n \in \mathbb{Z}_{>0}.$$

Moreover, the matrix Σ is obtained by

$$\langle v_1, \Sigma v_2 \rangle = \sum_{e \in E(G)} u_1(e) u_2(e) c(e), \quad (3.16)$$

where u_i is the harmonic part of \widehat{v}_i for $v_i \in \mathbb{R}^m$ for $i = 1, 2$.

Proof. For all $\delta > 0$, the Fourier inversion formula (3.14) and the change of variable yield for $n \in \mathbb{Z}_{>0}$ and for $x \in \Gamma$,

$$\mu_n(x) = \frac{1}{n^{\frac{m}{2}}} \int_{D_{\delta, \sqrt{n}}} \varphi_{\mu_n} \left(\frac{v}{\sqrt{n}} \right) e^{-2\pi i \langle v, \Phi(x) \rangle / \sqrt{n}} dv + \int_{D \setminus D_\delta} \varphi_{\mu_n}(v) e^{-2\pi i \langle v, \Phi(x) \rangle} dv.$$

Since $p(x_0, x_0) > 0$ in the pointed finite network (G, c, x_0) , Lemma 3.5 shows that for all small enough $\delta > 0$, there exists a constant $c_\delta > 0$ such that for all $n \in \mathbb{Z}_{>0}$,

$$\sup_{v \in D \setminus D_\delta} |\varphi_{\mu_n}(v)| = O(e^{-c_\delta n}).$$

Further by (3.13), up to replacing c_δ by a smaller positive value if necessary,

$$\varphi_{\mu_n}(v) = e^{n\beta(v)} \langle \mathbf{1}, f_v \rangle_\pi f_v(x_0) + O(e^{-c_\delta n}) \quad \text{for } v \in D_\delta,$$

where $f_v := f_{\hat{v}}$ is an eigenvector of $\mathcal{L}_{\hat{v}}$ corresponding to $e^{\beta(v)}$. By Lemma 3.3, the Hessian of β at 0 on \mathbb{R}^m is obtained by $\text{Hess}_0 \beta = -4\pi^2 \Sigma$. Lemma 3.2 implies that by the Taylor theorem,

$$\beta(v) = -2\pi^2 \langle v, \Sigma v \rangle + O(\|v\|^4) \quad \text{for } v \in D_\delta.$$

Therefore for all $n \in \mathbb{Z}_{>0}$ and for all $v \in D_{\delta\sqrt{n}}$,

$$\beta\left(\frac{v}{\sqrt{n}}\right) = -\frac{2\pi^2}{n} \langle v, \Sigma v \rangle + O\left(\frac{\|v\|^4}{n^2}\right).$$

Furthermore for all $n \in \mathbb{Z}_{>0}$ and for all $v \in D_{\delta\sqrt{n}}$,

$$\varphi_{\mu_n}\left(\frac{v}{\sqrt{n}}\right) = e^{-2\pi^2 \langle v, \Sigma v \rangle} (1 + R_n(v)) \cdot \langle \mathbf{1}, f_{v/\sqrt{n}} \rangle_\pi f_{v/\sqrt{n}}(x_0) + O(e^{-c_\delta n}),$$

where $R_n(v) = O(\|v\|^4/n)$. For the (normalized) eigenvector f_v of $e^{\beta(v)}$ for v near 0, since $f_0 = \mathbf{1}$, by Lemma 3.2 (3.4) the Taylor theorem shows the following: For each $x \in V(G)$,

$$f_{v/\sqrt{n}}(x) = 1 + O\left(\frac{\|v\|^2}{n}\right) \quad \text{and} \quad \langle \mathbf{1}, f_{v/\sqrt{n}} \rangle_\pi f_{v/\sqrt{n}}(x_0) = 1 + O\left(\frac{\|v\|^2}{n}\right) \quad \text{for } v \in D_{\delta\sqrt{n}}.$$

Note that for every real $\alpha \geq 0$,

$$\int_{D_{\delta\sqrt{n}}} e^{-2\pi^2 \langle v, \Sigma v \rangle} \frac{\|v\|^\alpha}{n} dv \leq \frac{1}{n} \int_{\mathbb{R}^m} \|v\|^\alpha e^{-2\pi^2 \langle v, \Sigma v \rangle} dv = O\left(\frac{1}{n}\right).$$

Summarizing the above estimates with $e^{-c_\delta n} \ll n^{-\frac{m}{2}-1}$ yields for $x \in \Gamma$ and for $n \in \mathbb{Z}_{>0}$,

$$\mu_n(x) = \frac{1}{n^{\frac{m}{2}}} \int_{D_{\delta\sqrt{n}}} e^{-2\pi^2 \langle v, \Sigma v \rangle} e^{-2\pi i \langle v, \Phi(x) \rangle / \sqrt{n}} dv + O\left(\frac{1}{n^{\frac{m}{2}+1}}\right). \quad (3.17)$$

A direct computation on the Fourier transform yields for all $n \in \mathbb{Z}_{>0}$ and for all $x \in \Gamma$,

$$\xi_{n\Sigma}(x) = \frac{1}{n^{\frac{m}{2}}} \int_{\mathbb{R}^m} e^{-2\pi i \langle v, \Phi(x) \rangle / \sqrt{n}} e^{-2\pi^2 \langle v, \Sigma v \rangle} dv.$$

Abusing notations, we have a (possibly different) constant $c_\delta > 0$ such that for all $n \in \mathbb{Z}_{>0}$ and uniformly in $x \in \Gamma$,

$$\xi_{n\Sigma}(x) = \frac{1}{n^{\frac{m}{2}}} \int_{D_{\delta\sqrt{n}}} e^{-2\pi i \langle v, \Phi(x) \rangle / \sqrt{n}} e^{-2\pi^2 \langle v, \Sigma v \rangle} dv + O(e^{-c_\delta n}). \quad (3.18)$$

Therefore by (3.17) and (3.18), for all $n \in \mathbb{Z}_{>0}$ and uniformly in $x \in \Gamma$,

$$\mu_n(x) = \xi_{n\Sigma}(x) + O\left(\frac{1}{n^{\frac{m}{2}+1}}\right).$$

Furthermore the explicit form of Σ is obtained by Lemma 3.3, as claimed. \square

Lemma 3.7. *Let μ be a probability measure on Γ such that $\text{supp } \mu$ is finite and $\Gamma = \langle \text{supp } \mu \rangle$, and $\{w_n\}_{n \in \mathbb{Z}_+}$ be a μ -random walk with $w_0 = \text{id}$. There exists a constant $C > 0$ such that for all $n \in \mathbb{Z}_{>0}$ and for all integer $t > C$,*

$$\mathbf{P}(|w_n|_S \geq t) \leq C \exp\left(-\frac{t^2}{Cn}\right).$$

Proof. Note that if $\{\Phi(w_n)\}_{n \in \mathbb{Z}_+}$ is a martingale with respect to the filtration associated with $\{w_n\}_{n \in \mathbb{Z}_+}$, then the proof follows from a concentration inequality. In general, $\{\Phi(w_n)\}_{n \in \mathbb{Z}_+}$ is not a martingale. What we do below is to replace Φ by another Λ -equivariant map which makes the images of w_n form a martingale.

First we claim that there exists a Λ -equivariant harmonic map $\Phi_H : \Gamma \rightarrow \mathbb{R}^m$. This is a map satisfying that $\Phi_H(gx) = \Phi_H(x) + g$ for all $x \in \Gamma$ and for all $g \in \Lambda$ under the identification between Λ and a lattice in \mathbb{R}^m , and

$$\sum_{s \in \text{supp } \mu} (\Phi_H(xs) - \Phi_H(x)) \mu(s) = 0 \quad \text{for each } x \in \Gamma.$$

This map is obtained from a Λ -equivariant lift of a Dirichlet energy minimizing map from $G = (V(G), E(G))$ with weights on edges $c(e)$ for $e \in E(G)$ into the flat torus \mathbb{R}^m/Λ equipped with metric as a quotient of the standard Euclidean space. The existence of such a map is shown by a simple variational calculus [KS01, Theorem 2.3] (see also [Sun13, Chapter 7]). (In general, Φ_H is not necessarily injective, but this does not affect the following discussion.) For a Λ -equivariant harmonic map Φ_H , we have a martingale $\{\Phi_H(w_n)\}_{n \in \mathbb{Z}_+}$ with respect to the natural filtration.

Next note that the map Φ_H yields a quasi-isometry between $\text{Cay}(\Gamma, S)$ and \mathbb{R}^m . In particular, there exist constants $c_0, c_1 > 0$ such that

$$\|\Phi_H(x)\| \geq c_0|x|_S - c_1 \quad \text{for all } x \in \Gamma. \quad (3.19)$$

Finally, each component of $\Phi_H(w_n)$ in the coordinate of \mathbb{R}^m is a martingale with a uniformly bounded difference B for some $B > 0$. Hence a union bound and the Azuma-Hoeffding inequality show that by (3.19), for all $r \in \mathbb{Z}_+$ and for all $n \in \mathbb{Z}_{>0}$,

$$\mathbf{P}(|w_n|_S \geq (r + c_1)/c_0) \leq \mathbf{P}(\|\Phi_H(w_n)\| \geq r) \leq 2m \exp\left(-\frac{r^2}{2B^2n}\right).$$

Therefore taking a large enough constant $C > 0$ concludes the inequality as claimed. \square

For the ξ_Σ in Theorem 3.6, let us define a **discrete normal distribution** \mathcal{N}_Σ^Φ on Γ by

$$\mathcal{N}_\Sigma^\Phi(x) := \frac{1}{Z} \xi_\Sigma(x) \quad \text{where } Z := \sum_{x \in \Gamma} \xi_\Sigma(x) \text{ for } x \in \Gamma.$$

Theorem 3.8. *In the same setting and assumption as in Theorem 3.6, there exists a constant $C > 0$ such that for all large enough integers n ,*

$$\|\mu_n - \mathcal{N}_{n\Sigma}^\Phi\|_{\text{TV}} \leq \frac{C(\log n)^{\frac{m}{2}}}{n}.$$

Proof. The local central limit theorem (Theorem 3.6) implies that there exists a constant $C > 0$ such that for all $n \in \mathbb{Z}_{>0}$,

$$\sup_{x \in \Gamma} |\mu_n(x) - \xi_{n\Sigma}(x)| \leq \frac{C}{n^{\frac{m}{2}+1}}. \quad (3.20)$$

For a μ -random walk $\{w_n\}_{n \in \mathbb{Z}_+}$ with $w_0 = \text{id}$ on Γ , Lemma 3.7, there exists a constant $C > 0$ such that for all $n \in \mathbb{Z}_{>0}$ and for all $r > C$,

$$\mathbf{P}(|w_n|_S \geq r) \leq C \exp\left(-\frac{r^2}{Cn}\right). \quad (3.21)$$

A direct computation on $\xi_{n\Sigma}$ yields for a (possibly different) constant $C > 0$, for all $n > C$ and for all $r > C\sqrt{n}$,

$$\sum_{|x|_S \geq r} \xi_{n\Sigma}(x) \leq C \exp\left(-\frac{r^2}{Cn}\right). \quad (3.22)$$

Indeed, this follows from an approximation by a Gaussian f_Σ on \mathbb{R}^m for which $\xi_\Sigma = f_\Sigma \circ \Phi$, and that Φ yields a quasi-isometry between $\text{Cay}(\Gamma, S)$ and \mathbb{R}^m . Note that $\mathbb{R}^m = \bigcup_{x \in \Gamma} x\overline{C}_0$ for the closure \overline{C}_0 of a relatively compact fundamental domain C_0 of Γ . Estimating the gradient of $f_{n\Sigma}(v)$ in $v \in \mathbb{R}^m$ shows the following: There exists a constant $c > 0$ such that for all $x \in \Gamma$, all $v_1, v_2 \in x\overline{C}_0$ and all $n \in \mathbb{Z}_{>0}$,

$$f_{n\Sigma}(v_1) \leq c \left(1 + \frac{\|v_2\|}{n}\right) f_{n\Sigma}(v_2).$$

Further note that $\|\Phi(x)\| \geq c_0|x|_S - c_1$ for all $x \in \Gamma$. Let $\|\Sigma^{\frac{1}{2}}\|$ denote the matrix norm of $\Sigma^{\frac{1}{2}}$. If $(c_0r - c_1)/(\|\Sigma^{\frac{1}{2}}\|\sqrt{n}) \geq 1$, then by the change of variables $v \mapsto \sqrt{n}\Sigma^{\frac{1}{2}}v$,

$$\begin{aligned} \sum_{|x|_S \geq r} \xi_{n\Sigma}(x) &\leq c \int_{\|v\| \geq c_0r - c_1} \left(1 + \frac{\|v\|}{n}\right) \frac{1}{(2\pi n)^{\frac{m}{2}} \sqrt{\det \Sigma}} \exp\left(-\frac{1}{2n} \langle v, \Sigma^{-1}v \rangle\right) dv \\ &\leq \frac{c}{(2\pi)^{\frac{m}{2}}} \left(1 + \frac{\|\Sigma^{\frac{1}{2}}\|}{\sqrt{n}}\right) \int_{\|v\| \geq (c_0r - c_1)/(\|\Sigma^{\frac{1}{2}}\|\sqrt{n})} \|v\| e^{-\frac{1}{2}\|v\|^2} dv \leq C \exp\left(-\frac{(c_0r - c_1)^2}{\|\Sigma^{\frac{1}{2}}\|^2 n}\right). \end{aligned}$$

This shows (3.22).

Combining (3.20), (3.21) and (3.22) yields for all $n > C$ and for all $r > C\sqrt{n}$,

$$\begin{aligned} \|\mu_n - \xi_{n\Sigma}\|_1 &= \sum_{|x|_S \leq r} |\mu_n(x) - \xi_{n\Sigma}(x)| + \sum_{|x|_S > r} |\mu_n(x) - \xi_{n\Sigma}(x)| \\ &\leq \#B_S(r) \frac{C}{n^{\frac{m}{2}+1}} + 2C \exp\left(-\frac{r^2}{Cn}\right), \end{aligned}$$

where $B_S(r) := \{x \in \Gamma : |x|_S \leq r\}$. Note that $\#B_S(r) = \Theta(r^m)$, in particular, $\#B_S(r) \leq Cr^m$ for all $r \in \mathbb{Z}_+$ for a constant $C > 0$ independent of r since Φ is a Λ -equivariant injective map from Γ into \mathbb{R}^m . Letting $r = A\sqrt{n \log n}$ for a constant $A > 0$, we have $r > C\sqrt{n}$ for all large enough $n \in \mathbb{Z}_{>0}$, and

$$\|\mu_n - \xi_{n\Sigma}\|_1 \leq \frac{C^2 A^m (\log n)^{\frac{m}{2}}}{n} + \frac{2C}{n^{A^2/C}}.$$

Fixing a large enough constant A such that $A^2/C > 1$ shows that there exists a constant C_1 such that for all large enough n ,

$$\|\mu_n - \xi_{n\Sigma}\|_1 \leq \frac{C_1(\log n)^{\frac{m}{2}}}{n}. \quad (3.23)$$

Note that there exists a constant $c > 0$ such that for all $n \in \mathbb{Z}_{>0}$,

$$\sum_{x \in \Gamma} \xi_{n\Sigma}(x) = 1 + \sum_{x \in \Gamma \setminus \{\text{id}\}} \xi_{n\Sigma}(x) = 1 + O(e^{-cn}). \quad (3.24)$$

Indeed, since \mathbb{Z}^m is a finite index subgroup of Γ and Φ is a \mathbb{Z}^m -equivariant embedding with a discrete image in \mathbb{R}^m , the Poisson summation formula (cf. (A.1) in Appendix A) on finitely many orbits of \mathbb{Z}^m shows (3.24). This implies that for all $n \in \mathbb{Z}_{>0}$,

$$\|\xi_{n\Sigma} - \mathcal{N}_{n\Sigma}^\Phi\|_1 = \sum_{x \in \Gamma} \left| \xi_{n\Sigma}(x) - \frac{1}{\sum_{z \in \Gamma} \xi_{n\Sigma}(z)} \xi_{n\Sigma}(x) \right| = O(e^{-cn}).$$

Therefore this together with (3.23), adjusting a constant factor C yields the claim. \square

Remark 3.9. The inequality (3.23) suffices for our purpose in Theorem 3.10 below. Note that in this generality the estimate in Theorem 3.8 is sharp up to the factor $O((\log n)^{\frac{m}{2}})$. Indeed, the local central limit theorem provides an example satisfying that

$$\|\mu_n - \mathcal{N}_{n\Sigma}^\Phi\|_{\text{TV}} = \Omega(1/n),$$

e.g., a lazy simple random walk on \mathbb{Z}^m .

3.7. Noise sensitivity on affine Weyl groups. Let (Γ, S) be an affine Weyl group where S is a canonical set of generators consisting of involutions. We have $\Gamma = \Lambda \rtimes W$ as in Section 3.1. For a probability measure μ on Γ and for all $\rho \in [0, 1]$, we recall that $\pi^\rho = \rho\mu \times \mu + (1-\rho)\mu_{\text{diag}}$, where $\mu \times \mu$ denote the product measure and $\mu_{\text{diag}}((x, y)) = \mu(x)$ if $x = y$ and 0 otherwise.

For each $\rho \in [0, 1]$, the measure π^ρ is defined on $\Gamma \times \Gamma = (\Lambda \times \Lambda) \rtimes (W \times W)$, for which $S_* := (S \cup \{\text{id}\})^2$ is a generating set of order at most 2. On the one hand, if $\text{supp } \mu = S \cup \{\text{id}\}$, then $\text{supp } \pi^\rho = S_*$ for all $\rho \in (0, 1]$. On the other hand, however, if $\rho = 0$ or $\text{supp } \mu = S$ (in which case, the support does not contain id), then $\text{supp } \pi^\rho$ never generate $\Gamma \times \Gamma$. Indeed, every affine Weyl group Γ admits a surjective homomorphism onto $\{\pm 1\}$ through the determinant of the isometry part in the natural affine representation. The product group $\Gamma \times \Gamma$ admits a surjective homomorphism onto $\{\pm 1\}^2$. If $\text{supp } \mu$ does not contain id , then $\text{supp } \pi^\rho$ for $\rho \in (0, 1]$ only generates a proper subgroup of $\Gamma \times \Gamma$ (of index 2). Furthermore if $\rho = 0$, then $\text{supp } \pi^\rho$ generates the diagonal subgroup isomorphic to Γ in $\Gamma \times \Gamma$.

Theorem 3.10. *Let (Γ, S) be an affine Weyl group, and μ be a probability measure on Γ such that the support of μ equals $S \cup \{\text{id}\}$. For all $\rho \in (0, 1]$, the π^ρ -random walk on $\Gamma \times \Gamma$ starting from the identity satisfies the following: There exist a constant $C > 0$ and an integer $m > 0$ such that for all large enough integer n ,*

$$\|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} \leq \frac{C(\log n)^m}{n}.$$

In particular, for all $\rho \in (0, 1]$,

$$\lim_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} = 0,$$

i.e., the μ -random walk on Γ is noise sensitive in total variation.

Proof. Let us apply to $\Gamma \times \Gamma$ and π^ρ the discussion we have made so far. Fix a point o in the interior of a chamber for Γ in the associated Euclidean space \mathbb{R}^m . We assume that o is the origin after taking a conjugate by a translation for the action of Γ if necessary. Let $\Phi := (\Phi^{(1)}, \Phi^{(2)}) : \Gamma \times \Gamma \rightarrow \mathbb{R}^m \times \mathbb{R}^m$, $(x_1, x_2) \mapsto (x_1.o, x_2.o)$, where $\Phi^{(i)}(x) = x.o$ for $x \in \Gamma$ and for $i = 1, 2$. For all $\rho \in (0, 1]$, the support of π^ρ is finite, $\Gamma \times \Gamma = \langle \text{supp } \pi^\rho \rangle$ and π^ρ is symmetric since every element in $\text{supp } \pi^\rho = S_* = (S \cup \{\text{id}\})^2$ has the order at most 2. Further let $G := (\Lambda \times \Lambda) \setminus \text{Cay}(\Gamma \times \Gamma, S_*)$ equipped with the conductance $c(\mathbf{e})$ for $\mathbf{e} \in E(G)$ induced from π^ρ . We consider the corresponding pointed finite network (G, c, x_0) where x_0 is the identity element in $W \times W = (\Lambda \times \Lambda) \setminus (\Gamma \times \Gamma)$. For all $\rho \in (0, 1]$, the corresponding Markov chain on G is irreducible and satisfies that $p(x_0, x_0) > 0$, in fact, $p(x, x) > 0$ for every $x \in V(G)$ since $\pi^\rho(\text{id}) > 0$.

Let Σ^ρ be the matrix for π^ρ in the local central limit theorem (Theorem 3.6), let us show that $\Sigma^\rho = \Sigma^1$ for all $\rho \in (0, 1]$. In fact, in the block diagonal form along the decomposition $\mathbb{R}^m \times \mathbb{R}^m$, we prove that for all $\rho \in (0, 1]$,

$$\Sigma^\rho = \begin{pmatrix} \Sigma^\mu & 0 \\ 0 & \Sigma^\mu \end{pmatrix},$$

where Σ^μ is the matrix Σ^μ for μ in the local central limit theorem. Theorem 3.6 shows that Σ^ρ is obtained by

$$\langle \mathbf{v}_1, \Sigma^\rho \mathbf{v}_2 \rangle = \sum_{\mathbf{e} \in E(G)} \mathbf{u}_1(\mathbf{e}) \mathbf{u}_2(\mathbf{e}) c(\mathbf{e}), \quad (3.25)$$

for $\mathbf{v}_1, \mathbf{v}_2 \in \mathbb{R}^m \times \mathbb{R}^m$ and \mathbf{u}_i is the harmonic part of the 1-form $\widehat{\mathbf{v}}_i = (\langle \mathbf{v}_i, \Phi_{\mathbf{e}} \rangle)_{\mathbf{e} \in E(G)}$ for $i = 1, 2$.

First let us show that $\langle \mathbf{v}_1, \Sigma^\rho \mathbf{v}_2 \rangle = 0$ for all $\mathbf{v}_1 = (v_1, 0), \mathbf{v}_2 = (0, v_2) \in \mathbb{R}^m \times \mathbb{R}^m$. For each edge $\mathbf{e} = ((x_1, x_2), (x_1.s_1, x_2.s_2)) \in E(G)$, we write $e_i = (x_i, x_i.s_i)$ for $x_i \in W$ and $s_i \in S \cup \{\text{id}\}$ for $i = 1, 2$. For such $\mathbf{v}_1, \mathbf{v}_2$, we have that

$$\widehat{\mathbf{v}}_i(\mathbf{e}) = \langle \mathbf{v}_i, \Phi_{\mathbf{e}} \rangle = \langle \mathbf{v}_i, \Phi_{e_i}^{(i)} \rangle = \widehat{\mathbf{v}}_i(e_i) \quad \text{for } i = 1, 2. \quad (3.26)$$

Let \mathbf{u}_i be the harmonic part of $\widehat{\mathbf{v}}_i$, and u_i be the harmonic part of $\widehat{\mathbf{v}}_i$ for each $i = 1, 2$. It holds that

$$\mathbf{u}_i(\mathbf{e}) = u_i(e_i) \quad \text{for } i = 1, 2. \quad (3.27)$$

Indeed, let $\widetilde{u}_i(\mathbf{e}) := u_i(e_i)$ for $i = 1, 2$. These define harmonic 1-forms on G : This follows since $S_* = (S \cup \{\text{id}\})^2$ and each marginal of π^ρ is μ . Furthermore by the definition of u_i , it holds that for some $f_i : W \rightarrow \mathbb{R}$,

$$\widehat{\mathbf{v}}_i(x, x.s) = u_i(x, x.s) + df_i(x, x.s) \quad \text{for } x \in W \text{ and } s \in S. \quad (3.28)$$

Hence by (3.26) and (3.28), letting $\tilde{f}_i : V(G) \rightarrow \mathbb{R}$ by $\tilde{f}_i((x_1, x_2)) := f_i(x_i)$ for $(x_1, x_2) \in V(G) = W \times W$ for $i = 1, 2$ yields

$$\widehat{v}_i(\mathbf{e}) = \tilde{u}_i(\mathbf{e}) + d\tilde{f}_i(\mathbf{e}) \quad \text{for } \mathbf{e} \in E(G).$$

The uniqueness of harmonic part concludes (3.27).

The right hand side in (3.25) is computed as in the following: By (3.27), it holds that

$$\begin{aligned} \sum_{\mathbf{e} \in E(G)} \mathbf{u}_1(\mathbf{e})\mathbf{u}_2(\mathbf{e})c(\mathbf{e}) &= \sum_{\mathbf{e} \in E(G)} u_1(e_1)u_2(e_2)c(\mathbf{e}) \\ &= \sum_{(x_1, x_2) \in W \times W} \sum_{(s_1, s_2) \in S_*} u_1(x_1, x_1 \cdot s_1)u_2(x_2, x_2 \cdot s_2)\pi((x_1, x_2))\pi^\rho((s_1, s_2)). \end{aligned}$$

Note that the summation over $(s_1, s_2) \in S_*$ is restricted to $S \times S$ since $u_i(x, x) = 0$ for $x \in W$ and for each $i = 1, 2$ (in fact, all 1-form has value 0 on loops). Furthermore $\pi((x, y)) = 1/(\#W)^2$ for all $(x, y) \in W \times W$. Hence the last term times the factor $(\#W)^2$ equals

$$\begin{aligned} &\sum_{x_1 \in W, s_1 \in S} u_1(x_1, x_1 \cdot s_1) \sum_{x_2 \in W, s_2 \in S} u_2(x_2, x_2 \cdot s_2)\pi^\rho((s_1, s_2)) \\ &= \sum_{x_1 \in W, s_1 \in S} u_1(x_1, x_1 \cdot s_1) \frac{1}{2} \sum_{x_2 \in W, s_2 \in S} (u_2(x_2, x_2 \cdot s_2) + u_2(x_2 \cdot s_2, x_2))\pi^\rho((s_1, s_2)) = 0. \end{aligned}$$

In the above we have used that $s_2 = s_2^{-1}$ and $\pi^\rho((s_1, s_2)) = \pi^\rho((s_1, s_2^{-1}))$ for each $(s_1, s_2) \in S \times S$ in the first equality, and that $u_2(x_2, x_2 \cdot s_2) = -u_2(x_2 \cdot s_2, x_2)$ for all $x_2 \in W$ and all $s_2 \in S$ in the last equality. Therefore for all $\mathbf{v}_1 = (v_1, 0), \mathbf{v}_2 = (0, v_2) \in \mathbb{R}^m \times \mathbb{R}^m$, it holds that $\langle \mathbf{v}_1, \Sigma^\rho \mathbf{v}_2 \rangle = 0$.

Next since each marginal of π^ρ is μ , for $\mathbf{v}_1 = (v_1, 0), \mathbf{v}_2 = (0, v_2) \in \mathbb{R}^m \times \mathbb{R}^m$,

$$\langle \mathbf{v}_1, \Sigma^\rho \mathbf{v}_2 \rangle = \langle v_1, \Sigma^\mu v_2 \rangle.$$

The same equality holds for $\mathbf{v}_1 = (0, v_1), \mathbf{v}_2 = (0, v_2) \in \mathbb{R}^m \times \mathbb{R}^m$. Summarizing all the above discussion, we obtain $\Sigma^\rho = \Sigma^1$ for all $\rho \in (0, 1]$.

Finally, Theorem 3.8 shows that for all $\rho \in (0, 1]$, there exists a constant $C > 0$ such that for all large enough n , by the triangle inequality,

$$\|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} \leq \|\pi_n^\rho - \mathcal{N}_{n\Sigma^\rho}^\Phi\|_{\text{TV}} + \|\mu_n \times \mu_n - \mathcal{N}_{n\Sigma^1}^\Phi\|_{\text{TV}} \leq \frac{2C(\log n)^m}{n}.$$

This concludes the first claim. The second claim follows from the first claim. \square

3.8. Examples. Let us provide explicit examples of random walks on the affine Weyl groups of type $\widetilde{A}_1 \times \widetilde{A}_1, \widetilde{A}_2$ and \widetilde{C}_2 . Figures 1, 4 and 5 respectively describe the Cayley graphs of the groups with the corresponding sets of generators (the solid lines with dots). The associated action of each group on \mathbb{R}^2 consists of reflections with respect to lines (indicated as dotted lines) with a fundamental domain (colored in dark gray). The lattice has a larger fundamental domain (colored in light gray).

3.8.1. *Type $\tilde{A}_1 \times \tilde{A}_1$.* Let us consider the infinite dihedral group:

$$D_\infty = \langle s_1, s_2 \mid s_1^2 = s_2^2 = \text{id} \rangle.$$

Let $S := \{s_1, s_2\}$. The pair (D_∞, S) is the affine Weyl group of type \tilde{A}_1 , and the product group $D_\infty \times D_\infty$ with the standard set of generators $S \times \{\text{id}\} \cup \{\text{id}\} \times S$ is the affine Weyl group of type $\tilde{A}_1 \times \tilde{A}_1$. The group $D_\infty \times D_\infty$ is isomorphic to $\mathbb{Z}^2 \rtimes (\mathbb{Z}/2)^2$. We define $\Phi : D_\infty \times D_\infty \rightarrow \mathbb{R}^2$ in a way that the origin is the barycenter of a fundamental chamber (which is a square of side length $1/2$). The lattice is identified with the standard integer lattice (Figure 1).




FIGURE 1. The Cayley graph of $D_\infty \times D_\infty$ with the set of generators S_* (where loops corresponding to id are omitted) and the group action on \mathbb{R}^2 .

Let μ be a probability measure on D_∞ such that $\text{supp } \mu = S \cup \{\text{id}\}$, and π^ρ be the associated probability measure on $D_\infty \times D_\infty$ for $\rho \in (0, 1]$. The measure π^ρ has the support $S_* = (S \cup \{\text{id}\})^2$. The quotient graph $G = \mathbb{Z}^2 \setminus \text{Cay}(D_\infty \times D_\infty, S_*)$ is described in Figure 2.

First we consider the case when $\mu(s_1)$ and $\mu(s_2)$ are equal, i.e.,

$$\mu(s_1) = \mu(s_2) = \frac{1}{2}(1 - \mu(\text{id})) \quad \text{and} \quad 0 < \mu(\text{id}) < 1.$$

In this case, for $v_1 = (1, 0), v_2 = (0, 1) \in \mathbb{R}^2$, the 1-forms $(\langle v_i, \Phi_e \rangle)_{e \in E(G)}$ for $i = 1, 2$ are harmonic. A direct computation yields

$$\Sigma^\rho = \begin{pmatrix} \frac{1}{4}(1 - \mu(\text{id})) & 0 \\ 0 & \frac{1}{4}(1 - \mu(\text{id})) \end{pmatrix} \quad \text{for } \rho \in (0, 1].$$

Next in the case when $\mu(s_1)$ and $\mu(s_2)$ are not necessarily equal, the harmonic 1-forms

$$u_i(\mathbf{e}) = \langle v_i, \Phi_{\mathbf{e}} \rangle - df_i(\mathbf{e}) \quad \text{for } \mathbf{e} \in E(G) \text{ and } i = 1, 2,$$

are obtained by (possibly non-constant) functions $f_i : V(G) \rightarrow \mathbb{R}$. For $i = 1$, let

$$f_1((0, 0)) = f_1((0, 1)) = \frac{\mu(s_2)}{2(1 - \mu(\text{id}))} \quad \text{and} \quad f_1((1, 0)) = f_1((1, 1)) = \frac{\mu(s_1)}{2(1 - \mu(\text{id}))}.$$

The values of the harmonic 1-form u_1 on the two oriented edges from $(0, 0)$ to $(1, 0)$ satisfy that

$$u_1((0, 0), (0.s_1, 0)) + f_1((1, 0)) - f_1((0, 0)) = \Phi((s_1.o, o)) - \Phi((o, o)) = \frac{1}{2},$$

$$u_1((0, 0), (0.s_2, 0)) + f_1((1, 0)) - f_1((0, 0)) = \Phi((s_2.o, o)) - \Phi((o, o)) = -\frac{1}{2},$$

and similarly, on the four oriented edges from $(0, 0)$ to $(1, 1)$. The values of u_1 on the two oriented edges from $(0, 0)$ to $(0, 1)$, and from $(1, 0)$ to $(1, 1)$, respectively, are 0. The harmonic 1-form u_2 is obtained similarly. A direct computation yields

$$\Sigma^\rho = \begin{pmatrix} \frac{\mu(s_1)\mu(s_2)}{1-\mu(\text{id})} & 0 \\ 0 & \frac{\mu(s_1)\mu(s_2)}{1-\mu(\text{id})} \end{pmatrix} \quad \text{for } \rho \in (0, 1].$$




FIGURE 2. The quotient graph $G = \mathbb{Z}^2 \backslash \text{Cay}(D_\infty \times D_\infty, S_*)$ where $S_* = (S \cup \{\text{id}\})^2$ and $S = \{s_1, s_2\}$.

The following example is not covered by Theorem 3.10: Let μ be the probability measure on D_∞ defined by

$$\mu(s_1) = \mu(s_1 s_2) = \mu(s_2 s_1) = \frac{1}{3}.$$

Let $S' := \{s_1, s_1 s_2, s_2 s_1\}$. We have $\text{supp } \mu = S'$ but $\text{id} \notin S'$. The associated probability measure π^ρ for $\rho \in (0, 1]$ on $D_\infty \times D_\infty$ has the support $S_* = (S')^2$ which does not satisfy

the condition in Theorem 3.10. The quotient graph of the corresponding Cayley graph for $D_\infty \times D_\infty$ is described in Figure 3. The matrix is explicitly computed as

$$\Sigma^\rho = \begin{pmatrix} \frac{1}{12} & 0 \\ 0 & \frac{1}{12} \end{pmatrix} \quad \text{for } \rho \in (0, 1].$$

The proof of Theorem 3.10 is directly applied to this case and the μ -random walk on D_∞ is noise sensitive in total variation.




FIGURE 3. The quotient graph $G = \mathbb{Z}^2 \setminus \text{Cay}(D_\infty \times D_\infty, S'_*)$ where $S'_* = (S')^2$ and $S' = \{s_1, s_1 s_2, s_2 s_1\}$.

3.8.2. *Type \tilde{A}_2 .* Let us consider the affine Weyl group of type \tilde{A}_2 :

$$\Gamma = \langle s_1, s_2, s_3 \mid s_1^2 = s_2^2 = s_3^2 = (s_1 s_2)^3 = (s_2 s_3)^3 = (s_3 s_1)^3 = 1 \rangle$$

with $S = \{s_1, s_2, s_3\}$. The Cayley graph $\text{Cay}(\Gamma, S)$ and the group action on \mathbb{R}^2 is described in Figure 4 (left). The group Γ is isomorphic to $\mathbb{Z}^2 \rtimes \mathfrak{S}_3$ where \mathfrak{S}_3 is the symmetric group on the set $\{1, 2, 3\}$. The quotient graph $G = \mathbb{Z}^2 \setminus \text{Cay}(\Gamma, S)$ is described in Figure 4 (right).

Let us consider $\Phi : \Gamma \rightarrow \mathbb{R}^2$ such that the origin is the barycenter of a fundamental chamber (which is an equilateral triangle of side length $(2\sqrt{3})^{1/2}/3$). A fundamental




FIGURE 4. The Cayley graph of the affine Weyl group of type \tilde{A}_2 and the group action on \mathbb{R}^2 (left). The quotient graph $G = \mathbb{Z}^2 \setminus \text{Cay}(\Gamma, S)$ for the affine Weyl group (Γ, S) of type \tilde{A}_2 (right).

domain for the \mathbb{Z}^2 -action is a hexagon of unit area. For a probability measure μ on Γ such that

$$\mu(s_1) = \mu(s_2) = \mu(s_3) = \frac{1}{3}(1 - \mu(\text{id})) \quad \text{and} \quad 0 < \mu(\text{id}) < 1,$$

the matrix for μ in the local central limit theorem (Theorem 3.6) is computed as

$$\Sigma^\mu = \begin{pmatrix} \frac{4\sqrt{3}}{9}(1 - \mu(\text{id})) & 0 \\ 0 & \frac{4\sqrt{3}}{9}(1 - \mu(\text{id})) \end{pmatrix}.$$

3.8.3. *Type \tilde{C}_2 .* Let us consider the affine Weyl group of type \tilde{C}_2 :

$$\Gamma = \langle s_1, s_2, s_3 \mid s_1^2 = s_2^2 = s_3^2 = (s_1 s_2)^4 = (s_2 s_3)^4 = (s_1 s_3)^2 = 1 \rangle$$

with $S = \{s_1, s_2, s_3\}$. The Cayley graph $\text{Cay}(\Gamma, S)$ and the group action on \mathbb{R}^2 is described in Figure 5 (left). The group Γ is isomorphic to $\mathbb{Z}^2 \rtimes ((\mathbb{Z}/2)^2 \rtimes \mathfrak{S}_2)$ where $(\mathbb{Z}/2)^2 \rtimes \mathfrak{S}_2$ is the signed permutations on the set $\{1, 2\}$. The quotient graph G of the Cayley graph with the set of generators S by the lattice \mathbb{Z}^2 is described in Figure 5 (right).

Let us consider $\Phi : \Gamma \rightarrow \mathbb{R}^2$ such that the origin is the barycenter of a fundamental chamber (which is an isosceles right triangle of equal side length $1/2$). A fundamental domain for the \mathbb{Z}^2 -action has unit area. For a probability measure μ on Γ such that

$$\mu(s_1) = \mu(s_2) = \mu(s_3) = \frac{1}{3}(1 - \mu(\text{id})) \quad \text{and} \quad 0 < \mu(\text{id}) < 1,$$

the matrix for μ in the local central limit theorem (Theorem 3.6) is computed as

$$\Sigma^\mu = \begin{pmatrix} \frac{1}{512}(1 - \mu(\text{id})) & 0 \\ 0 & \frac{1}{512}(1 - \mu(\text{id})) \end{pmatrix}.$$




FIGURE 5. The Cayley graph of the affine Weyl group of type \tilde{C}_2 and the group action on \mathbb{R}^2 (left). The quotient graph $G = \mathbb{Z}^2 \setminus \text{Cay}(\Gamma, S)$ for the affine Weyl group (Γ, S) of type \tilde{C}_2 (right).

APPENDIX A. NOISE SENSITIVITY PROBLEM ON \mathbb{Z}^m

Fix an integer $m \geq 1$. We say that a μ -random walk (or μ) on \mathbb{Z}^m is **irreducible** if $\bigcup_{n \in \mathbb{Z}_+} \text{supp } \mu_n = \mathbb{Z}^m$, **aperiodic** if for each $x \in \mathbb{Z}^m$ for all large enough n one has $\mu_n(x) > 0$. For μ of a finite first moment, we say that μ has **mean zero** if $\sum_{x \in \mathbb{Z}^m} x \mu(x) = 0$.

Theorem A.1. *Let m be a positive integer. If μ is an irreducible probability measure with finite second moment and mean zero on \mathbb{Z}^m , then*

$$\lim_{\rho \rightarrow 1} \limsup_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} = 0 \quad \text{and} \quad \lim_{\rho \rightarrow 0} \liminf_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} = 1.$$

Let Σ be a non-degenerate symmetric positive definite matrix of size m and

$$f_\Sigma(v) := \frac{1}{\sqrt{(2\pi)^m \det \Sigma}} e^{-\frac{1}{2}\langle v, \Sigma^{-1}v \rangle} \quad \text{for } v \in \mathbb{R}^m.$$

Let us define the function $F(v) := \sum_{x \in \mathbb{Z}^m} f_\Sigma(x+v)$ for $v \in \mathbb{R}^m$. Note that since F is \mathbb{Z}^m -periodic on \mathbb{R}^m , it is regarded as a function on $[0, 1)^m$. Let

$$\widehat{F}(x) := \int_{[0, 1)^m} F(v) e^{-2\pi i \langle v, x \rangle} dv \quad \text{and} \quad \widehat{f}_\Sigma(x) := \int_{\mathbb{R}^m} f_\Sigma(v) e^{-2\pi i \langle v, x \rangle} dv \quad \text{for } x \in \mathbb{Z}^m.$$

Since f_Σ is in the Schwartz class on \mathbb{R}^m , we have $\widehat{F}(x) = \widehat{f}_\Sigma(x)$ for $x \in \mathbb{Z}^m$ and

$$F(v) = \sum_{x \in \mathbb{Z}^m} \widehat{f}_\Sigma(x) e^{2\pi i \langle x, v \rangle},$$

where the right hand side is absolutely convergent. In the case when $v = 0$, the Poisson summation formula is obtained by a direct computation,

$$\sum_{x \in \mathbb{Z}^m} f_\Sigma(x) = \sum_{x \in \mathbb{Z}^m} \widehat{f}_\Sigma(x) = \sum_{x \in \mathbb{Z}^m} e^{-2\pi^2 \langle x, \Sigma x \rangle}.$$

Thus there exists a constant $c_\Sigma > 0$ such that for all $n \in \mathbb{Z}_+$,

$$\sum_{x \in \mathbb{Z}^m} f_{n\Sigma}(x) = \sum_{x \in \mathbb{Z}^m} e^{-2\pi^2 n \langle x, \Sigma x \rangle} = 1 + O_\Sigma(e^{-c_\Sigma n}). \quad (\text{A.1})$$

Let Σ_1 and Σ_2 be non-degenerate symmetric positive definite matrices of size m . First we consider the upper bound part. For every real $\lambda > 0$ and for every $n \in \mathbb{Z}_{>0}$,

$$\begin{aligned} & \sum_{\|x\| \leq \lambda n^{1/2}} |f_{n\Sigma_1}(x) - f_{n\Sigma_2}(x)| \\ &= \sum_{\|x\| \leq \lambda n^{1/2}} \left| f_{n\Sigma_1}(x) - \sqrt{\frac{\det \Sigma_2}{\det \Sigma_1}} f_{n\Sigma_2}(x) \right| + \sum_{\|x\| \leq \lambda n^{1/2}} \left| 1 - \sqrt{\frac{\det \Sigma_2}{\det \Sigma_1}} \right| f_{n\Sigma_2}(x). \end{aligned} \quad (\text{A.2})$$

The first sum in (A.2) is estimated as follows: For $x \in \mathbb{Z}^m$,

$$\left| 1 - e^{-\frac{1}{2n} \langle x, (\Sigma_2^{-1} - \Sigma_1^{-1})x \rangle} \right| \leq e^{\frac{1}{2n} |\langle x, (\Sigma_2^{-1} - \Sigma_1^{-1})x \rangle|} - e^{-\frac{1}{2n} |\langle x, (\Sigma_2^{-1} - \Sigma_1^{-1})x \rangle|}.$$

Since $|\langle x, (\Sigma_2^{-1} - \Sigma_1^{-1})x \rangle| \leq \|\Sigma_2^{-1} - \Sigma_1^{-1}\| \|x\|^2$, for $x \in \mathbb{Z}^m$ with $\|x\| \leq \lambda n^{1/2}$,

$$\left| 1 - e^{-\frac{1}{2n} \langle x, (\Sigma_2^{-1} - \Sigma_1^{-1})x \rangle} \right| \leq e^{\frac{\lambda^2}{2} \|\Sigma_2^{-1} - \Sigma_1^{-1}\|} - e^{-\frac{\lambda^2}{2} \|\Sigma_2^{-1} - \Sigma_1^{-1}\|} = 2 \sinh \left(\frac{\lambda^2}{2} \|\Sigma_2^{-1} - \Sigma_1^{-1}\| \right).$$

Therefore the first sum in (A.2) is at most by (A.1), for all $n \in \mathbb{Z}_+$ and for all $\lambda > 0$, if $\|\Sigma_2^{-1} - \Sigma_1^{-1}\| \leq 2/\lambda^2$, then

$$\sum_{\|x\| \leq \lambda n^{1/2}} \left| 1 - e^{-\frac{1}{2n} \langle x, (\Sigma_2^{-1} - \Sigma_1^{-1})x \rangle} \right| f_{n\Sigma_1}(x) \leq 3\lambda^2 \|\Sigma_2^{-1} - \Sigma_1^{-1}\| (1 + O_{\Sigma_1}(e^{-c_{\Sigma_1} n})).$$

The second sum in (A.2) equals by (A.1), for all $n \in \mathbb{Z}_+$,

$$\left| 1 - \sqrt{\frac{\det \Sigma_2}{\det \Sigma_1}} \right| \sum_{x \in \mathbb{Z}^m} f_{n\Sigma_2}(x) = \left| 1 - \sqrt{\frac{\det \Sigma_2}{\det \Sigma_1}} \right| (1 + O_{\Sigma_2}(e^{-c_{\Sigma_2} n}))$$

Since the right hand side in (A.1) tends to 1 as $n \rightarrow \infty$, summarizing the above estimates yields for every $\lambda > 0$, there exist constants $C_{1,2,\lambda} = C_{\Sigma_1, \Sigma_2, \lambda} > 0$ and $c_{1,2} = c_{\Sigma_1, \Sigma_2} > 0$ such that for all $n \in \mathbb{Z}^+$,

$$\sum_{\|x\| \leq \lambda n^{1/2}} |f_{n\Sigma_1}(x) - f_{n\Sigma_2}(x)| \leq 3\lambda^2 \|\Sigma_2^{-1} - \Sigma_1^{-1}\| + \left| 1 - \sqrt{\frac{\det \Sigma_2}{\det \Sigma_1}} \right| + C_{1,2,\lambda} e^{-c_{1,2} n}. \quad (\text{A.3})$$

Next let us consider the lower bound part. Noting that $\sqrt{f_1 - f_2} \geq \sqrt{f_1} - \sqrt{f_2}$ for $f_1 \geq f_2 \geq 0$, we have by squaring both sides and summing over \mathbb{Z}^m ,

$$\|f_{n\Sigma_1} - f_{n\Sigma_2}\|_1 \geq \sum_{x \in \mathbb{Z}^m} f_{n\Sigma_1}(x) + \sum_{x \in \mathbb{Z}^m} f_{n\Sigma_2}(x) - 2 \sum_{x \in \mathbb{Z}^m} \sqrt{f_{n\Sigma_1}(x) f_{n\Sigma_2}(x)}.$$

In the following, we assume that $\Sigma_1^{-1} + \Sigma_2^{-1}$ is invertible. In that case, by (A.1),

$$\begin{aligned} \sum_{x \in \mathbb{Z}^m} \sqrt{f_{n\Sigma_1}(x)f_{n\Sigma_2}(x)} &= \sum_{x \in \mathbb{Z}^m} \frac{1}{(2\pi)^{\frac{m}{2}} (\det(n\Sigma_1) \det(n\Sigma_2))^{\frac{1}{4}}} e^{-\frac{1}{4n} \langle x, (\Sigma_1^{-1} + \Sigma_2^{-1})x \rangle} \\ &= \frac{2^{\frac{m}{2}} (\det((\Sigma_1^{-1} + \Sigma_2^{-1})^{-1}))^{\frac{1}{2}}}{(\det \Sigma_1 \det \Sigma_2)^{\frac{1}{4}}} \sum_{x \in \mathbb{Z}^m} e^{-4\pi^2 n \langle x, (\Sigma_1^{-1} + \Sigma_2^{-1})^{-1}x \rangle}. \end{aligned}$$

Let us focus on the special case when $m = 2$, and

$$\Sigma^1 = \sigma^2 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad \Sigma^\rho = \sigma^2 \begin{pmatrix} 1 & 1-\rho \\ 1-\rho & 1 \end{pmatrix} \quad \text{for } \sigma > 0 \text{ and } 0 < \rho \leq 1.$$

In this case,

$$(\Sigma^1)^{-1} + (\Sigma^\rho)^{-1} = \frac{1}{\sigma^2} \begin{pmatrix} 1 + \frac{1}{\rho(2-\rho)} & -\frac{1-\rho}{\rho(2-\rho)} \\ -\frac{1-\rho}{\rho(2-\rho)} & 1 + \frac{1}{\rho(2-\rho)} \end{pmatrix}.$$

Furthermore, $\det \Sigma^1 = \sigma^4$, $\det \Sigma^\rho = \sigma^4 \rho(2-\rho)$ and

$$\det((\Sigma^1)^{-1} + (\Sigma^\rho)^{-1}) = \frac{1}{\sigma^4} \left(\left(1 + \frac{1}{\rho(2-\rho)} \right)^2 - \left(\frac{1-\rho}{\rho(2-\rho)} \right)^2 \right).$$

Hence one computes for all small enough $\rho > 0$,

$$\frac{2}{(\rho(2-\rho))^{\frac{1}{4}} \left(\left(1 + \frac{1}{\rho(2-\rho)} \right)^2 - \left(\frac{1-\rho}{\rho(2-\rho)} \right)^2 \right)^{\frac{1}{2}}} = \frac{2(\rho(2-\rho))^{\frac{3}{4}}}{\sqrt{(\rho(2-\rho) + 1)^2 - (1-\rho)^2}} < 2\rho^{\frac{1}{4}}.$$

Summarizing the above computations and letting $n \rightarrow \infty$ yield

$$\liminf_{n \rightarrow \infty} \|f_{n\Sigma^\rho} - f_{n\Sigma^1}\|_1 \geq 2 - 2\rho^{\frac{1}{4}}. \quad (\text{A.4})$$

Proof of Theorem A.1. Let μ be an irreducible probability measure with finite second moment and mean zero on \mathbb{Z}^m . We assume that μ is aperiodic and the general case is reduced to this case. For $0 < \rho \leq 1$, we note that π^ρ is aperiodic since π^ρ and $\mu \times \mu$ have the same support in \mathbb{Z}^{2m} . Let Σ^ρ and Σ^1 denote the covariance matrices of size $2m$ for π^ρ and $\mu \times \mu$ respectively.

First let us show the upper bound in the claim. For all real $\lambda > 0$ and integers $n > 0$, by the Chebyshev inequality,

$$\sum_{\|\mathbf{x}\| > \lambda n^{1/2}} \pi_n^\rho(\mathbf{x}) = \mathbf{P}_{\pi^\rho}(\|\mathbf{w}_n\| > \lambda n^{1/2}) \leq \frac{1}{\lambda^2 n} \mathbf{E}_{\pi_n^\rho} \|\mathbf{w}_n\|^2 = \frac{2m}{\lambda^2} \mathbf{E}_\mu |x|^2.$$

This shows that (recalling that $\pi^1 = \mu \times \mu$)

$$\sum_{\|\mathbf{x}\| > \lambda n^{1/2}} |\pi_n^\rho(\mathbf{x}) - \mu_n \times \mu_n(\mathbf{x})| \leq \frac{4m}{\lambda^2} \mathbf{E}_\mu |x|^2. \quad (\text{A.5})$$

Since π^ρ is aperiodic for $\rho \in (0, 1]$, then the local central limit theorem [LL10, Theorem 2.3.9] shows that there exists a sequence $\delta_n > 0$ such that $\delta_n \rightarrow 0$ as $n \rightarrow \infty$, for all $n > 0$ and for all $\mathbf{x} \in \mathbb{Z}^{2m}$,

$$|\pi_n^\rho(\mathbf{x}) - f_{n\Sigma^\rho}(\mathbf{x})| \leq \frac{\delta_n}{n^m} \quad \text{and} \quad |\mu_n \times \mu_n(\mathbf{x}) - f_{n\Sigma^1}(\mathbf{x})| \leq \frac{\delta_n}{n^m}. \quad (\text{A.6})$$

Note that there exists a constant $C_m > 0$ such that for all real $\lambda > 0$ and all $n \in \mathbb{Z}_{>0}$, the number of $\mathbf{x} \in \mathbb{Z}^{2m}$ with $\|\mathbf{x}\| \leq \lambda n^{1/2}$ is at most $C_m \lambda^{2m} n^m$. Therefore it holds that

$$\sum_{\|\mathbf{x}\| \leq \lambda n^{1/2}} |\pi_n^\rho(\mathbf{x}) - \mu_n \times \mu_n(\mathbf{x})| \leq C \lambda^{2m} n^m \cdot \frac{2\delta_n}{n^m} + \sum_{\|\mathbf{x}\| \leq \lambda n^{1/2}} |f_{n\Sigma^\rho}(\mathbf{x}) - f_{n\Sigma^1}(\mathbf{x})|. \quad (\text{A.7})$$

Hence by (A.3), (A.5) and (A.7), for every $\lambda > 0$,

$$\limsup_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_1 \leq 3\lambda^2 \|(\Sigma^\rho)^{-1} - (\Sigma^1)^{-1}\| + \left| 1 - \sqrt{\frac{\det \Sigma^\rho}{\det \Sigma^1}} \right| + \frac{4m}{\lambda^2} \mathbf{E}_\mu |x|^2.$$

Since $\Sigma^\rho \rightarrow \Sigma^1$ as $\rho \rightarrow 1$, letting $\rho \rightarrow 1$ yields

$$\limsup_{\rho \rightarrow 1} \limsup_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_1 \leq \frac{4m}{\lambda^2} \mathbf{E}_\mu |x|^2.$$

This holds for all $\lambda > 0$, and thus we obtain in the total variation distance

$$\lim_{\rho \rightarrow 1} \limsup_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} = 0.$$

This shows the upper bound in the claim.

Next let us show the lower bound in the claim. The general case reduces to the case when $m = 1$ since a projection $\mathbb{Z}^m \rightarrow \mathbb{Z}$ (whence $\mathbb{Z}^{2m} \rightarrow \mathbb{Z}^2$) to a coordinate only decreases the total variation distance. Note that a direct computation shows that there exist constants $c_\rho, C_\rho > 0$ depending only on ρ such that for all $\lambda > 0$ and all $n > 0$ with $\lambda n^{1/2} - \sqrt{2} \geq 1$,

$$\sum_{\|\mathbf{x}\| > \lambda n^{1/2}} f_{n\Sigma^\rho}(\mathbf{x}) \leq \int_{\|\mathbf{x}\| \geq \lambda n^{1/2} - \sqrt{2}} \frac{1}{\sqrt{(2\pi)^2 \det(n\Sigma^\rho)}} e^{-\frac{1}{2n} \langle \mathbf{x}, (\Sigma^\rho)^{-1} \mathbf{x} \rangle} d\mathbf{x} \leq C_\rho e^{-c_\rho \lambda^2}.$$

Therefore together with (A.5) and (A.6), for (possibly different) constants $c_\rho, C_\rho > 0$ depending only on ρ and a constant $c > 0$, for all $\lambda > 2$ and all $n > 0$,

$$\|\pi_n^\rho - \mu_n \times \mu_n\|_1 \geq \|f_{n\Sigma^\rho} - f_{n\Sigma^1}\|_1 - C_\rho e^{-c_\rho \lambda^2} - 2C \lambda^{2m} \delta_n - \frac{4m}{\lambda^2} \mathbf{E}_\mu |x|^2.$$

Thus by (A.4), letting $n \rightarrow \infty$, we obtain

$$\liminf_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_1 \geq 2 - 2\rho^{\frac{1}{4}} - C_\rho e^{-c_\rho \lambda^2} - \frac{4m}{\lambda^2} \mathbf{E}_\mu |x|^2.$$

In total variation distance, letting $\lambda \rightarrow \infty$ and then $\rho \rightarrow 0$ yields

$$\lim_{\rho \rightarrow 0} \liminf_{n \rightarrow \infty} \|\pi_n^\rho - \mu_n \times \mu_n\|_{\text{TV}} = 1.$$

In the case when μ is not aperiodic, then we apply the above discussion to the even times and the odd times respectively for those aperiodic random walks, and obtain the same results. \square

Acknowledgment. The author is partially supported by JSPS Grant-in-Aid for Scientific Research JP20K03602.

REFERENCES

- [AB08] Peter Abramenko and Kenneth S. Brown. *Buildings*, volume 248 of *Graduate Texts in Mathematics*. Springer, New York, 2008. Theory and applications.
- [AB22] Goulmira Arzhantseva and Arindam Biswas. Logarithmic girth expander graphs of $SL_n(\mathbb{F}_p)$. *J. Algebraic Combin.*, 56(3):691–723, 2022.
- [BB23] Itai Benjamini and Jérémie Brieussel. Noise sensitivity of random walks on groups. *ALEA Lat. Am. J. Probab. Math. Stat.*, 20(2):1139–1164, 2023.
- [Cal13] Danny Calegari. The ergodic theory of hyperbolic groups. In *Geometry and topology down under*, volume 597 of *Contemp. Math.*, pages 15–52. Amer. Math. Soc., Providence, RI, 2013.
- [Gro87] M. Gromov. Hyperbolic groups. In S. M. Gersten, editor, *Essays in group theory*, volume 8 of *Mathematical Sciences Research Institute Publications*, pages 75–263, New York, 1987. Springer-Verlag.
- [Kai00] Vadim A. Kaimanovich. The Poisson formula for groups with hyperbolic properties. *Ann. of Math. (2)*, 152(3):659–692, 2000.
- [Kal18] Gil Kalai. Three puzzles on mathematics, computation, and games. In *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. I. Plenary lectures*, pages 551–606. World Sci. Publ., Hackensack, NJ, 2018.
- [KS83] András Krámli and Domokos Szász. Random walks with internal degrees of freedom. I. Local limit theorems. *Z. Wahrsch. Verw. Gebiete*, 63(1):85–95, 1983.
- [KS01] Motoko Kotani and Toshikazu Sunada. Standard realizations of crystal lattices via harmonic maps. *Trans. Amer. Math. Soc.*, 353(1):1–20, 2001.
- [KSS98] Motoko Kotani, Tomoyuki Shirai, and Toshikazu Sunada. Asymptotic behavior of the transition probability of a random walk on an infinite graph. *J. Funct. Anal.*, 159(2):664–689, 1998.
- [LL10] Gregory F. Lawler and Vlada Limic. *Random Walk: A Modern Introduction*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2010.
- [LP16] Russell Lyons and Yuval Peres. *Probability on Trees and Networks*, volume 42 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, New York, 2016.
- [LP21] Russell Lyons and Yuval Peres. Poisson boundaries of lamplighter groups: proof of the Kaimanovich-Vershik conjecture. *J. Eur. Math. Soc. (JEMS)*, 23(4):1133–1160, 2021.
- [LPS88] A. Lubotzky, R. Phillips, and P. Sarnak. Ramanujan graphs. *Combinatorica*, 8(3):261–277, 1988.
- [MP10] Peter Mörters and Yuval Peres. *Brownian Motion*, volume 30 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, 2010. With an appendix by Oded Schramm and Wendelin Werner.
- [PS94] Mark Pollicott and Richard Sharp. Rates of recurrence for \mathbf{Z}^q and \mathbf{R}^q extensions of subshifts of finite type. *J. London Math. Soc. (2)*, 49(2):401–416, 1994.
- [Sun13] Toshikazu Sunada. *Topological Crystallography: With a View Towards Discrete Geometric Analysis*, volume 6 of *Surveys and Tutorials in the Applied Mathematical Sciences*. Springer, Tokyo, 2013.
- [Tan22] Ryokichi Tanaka. Non-noise sensitivity for word hyperbolic groups. *Ann. Fac. Sci. Toulouse Math.*, to appear., arXiv:2211.03951, 2022.

DEPARTMENT OF MATHEMATICS, KYOTO UNIVERSITY, KYOTO 606-8502 JAPAN

Email address: rtanaka@math.kyoto-u.ac.jp