### Weakly modular graphs with diamond condition, the interval function and axiomatic characterizations

Lekshmi Kamal K. Sheela<sup>10</sup><sup>†</sup>, Jeny Jacob<sup>10</sup><sup>‡</sup>, and Manoj Changat<sup>10</sup><sup>§</sup>

Department of Futures Studies, University of Kerala, Trivandrum 695 581, India lekshmisanthoshgr@gmail.com, jenyjacobktr@gmail.com, mchangat@keralauniversity.ac.in,

Abstract. Weakly modular graphs are defined as the class of graphs that satisfy the *triangle condition* (TC) and the *quadrangle condition* (QC). We study an interesting subclass of weakly modular graphs that satisfies a stronger version of the triangle condition, known as the *triangle diamond condition* (TDC). and term this subclass of weakly modular graphs as the *diamond-weakly modular graphs*. It is observed that this class contains the class of bridged graphs and the class of weakly bridged graphs.

The interval function  $I_G$  of a connected graph G with vertex set V is an important concept in metric graph theory and is one of the prime example of a transit function; a set function defined on the Cartesian product  $V \times V$  to the power set of V satisfying the expansive, symmetric and idempotent axioms.

In this paper, we derive an interesting axiom denoted as (J0'), obtained from a well-known axiom introduced by Marlow Sholander in 1952, denoted as (J0). It is proved that the axiom (J0') is a characterizing axiom of the diamond-weakly modular graphs. We propose certain types of independent first-order betweenness axioms on an arbitrary transit function R and prove that an arbitrary transit function becomes the interval function of a diamond-weakly modular graph if and only if R satisfies these betweenness axioms. Similar characterizations are obtained for the interval function of bridged graphs and weakly bridged graphs.

#### 1 Introduction

In this paper, we consider only finite, simple, and connected graphs. Weakly modular graphs form an interesting class of graphs and are introduced by Chepoi in [10]. Weakly modular graphs generalize a variety of classes of graphs in *metric* graph theory such as modular graphs, Helly graphs, bridged graphs, and dual polar graphs. Weakly modular graphs have applications beyond metric graph

<sup>† 🖸 0000-0002-8527-3280</sup> 

<sup>&</sup>lt;sup>‡</sup> **()**0000-0001-6477-154X

<sup>§</sup> D0000-0001-7257-6031

theory and studied in other fields of mathematics like geometric group theory, incidence geometries and buildings, theoretical computer science, and combinatorial optimization, see, Chalopine et al. in [5], also, see the survey due to Bandelt and Chepoi [4].

The main goal of the studies of graph classes in metric graph theory involves the investigation that how far the properties of the standard path metric d in these graph classes can be approximated with the main properties of classical metric geometries like Euclidean  $\ell_2$ -geometry (and more generally, the  $\ell_1$ - and  $\ell_{\infty}$ -geometries), hyperbolic spaces, hypercubes, and trees. Several interesting classes of graphs are identified to have such properties.

Weakly modular graphs and their subclasses are one of the central classes of metric graph theory. Bridged graphs form an interesting subclass of weakly modular graphs. A graph G = (V, E) is a *bridged graph* if G has no isometric cycles of length greater than 3. It is trivial to note that the family of bridged graphs contain the family of chordal graphs. In [10], bridged graphs are characterized as the so called  $(C_4, C_5)$ -free weakly modular graphs and in [11], Farber proved that a graph G is a bridged graph if and only if all neighborhoods of convex sets are convex. Another interesting subclass of weakly modular graphs, defined in [5]. A graph G is called weakly bridged if it is weakly modular without induced cycle of length four.

In this paper, we introduce another subclass of weakly modular graphs, which we name as *diamond-weakly modular graphs*. We prove that these graphs form a super-class of weakly bridged graphs and bridged graphs. We characterize these graphs using properties of the *interval function* of the graph. The interval function  $I_G$  of a graph G can be related to *metric betweenness* of G.

Recently, a new property in terms of the 'natural betweenness', namely the metric betweenness in graphs is observed on the main classes of graphs in metric graph theory in [6]. That is, the central classes of graphs in metric graph theory are capable of a characterization in first order logic (definable in first order logic) with the "betweenness predicate" induced by the natural betweenness on graphs. More precisely, the metric betweenness (or shortest path betweenness) resulting from the graph metric d of graphs G = (V, E) are defined using the ternary relation B(abc) on the vertex set V in such a way that "the vertex b lies on some shortest path of G between the vertices a and c." It is established in [6] that the first order logic with betweenness, denoted in short as **FOLB**, is a powerful logic as far as the first order axiomatization of graph properties are concerned.

Metric betweenness can be expressed more conveniently by the *interval func*tion, defined for a connected graph G as the function  $I_G: V \times V \longrightarrow 2^V$  with

$$I_G(u,v) = \{ w \in V : d(u,w) + d(w,v) = d(u,v) \}$$
  
=  $\{ w \in V : w \text{ lies on some shortest } u, v - \text{ path in } G \}.$ 

In this paper, we avoid the language of logic and use the interval function  $I_G$  rather than the betweenness relation B due to the simplicity of  $I_G$ . The function

 $I_G$  is a well-known tool in metric graph theory and several authors have studied the function  $I_G$ , in particular, Mulder has given a systematic study of  $I_G$  in the axiomatic setting in [14]. The interval function of a weakly modular graph has very special properties and a nice axiomatic characterization of I exists for G in terms of an arbitrary set function known as a transit function in [3]. The term transit function is due to Mulder [15], and it is introduced to generalize the three classical notions in mathematics, namely, convexity, interval and betweenness in an axiomatic approach. The concept of transit function is already known as interval operator and is used by Van de vel in [24] in the context of convexity, see also, [1,13]. Bandelt and Chepoi in [3] discussed the same concept for studying metric properties of discrete metric spaces and graphs. In this paper, we follow Mulder and use the term transit function.

Given a non-empty set V, a *transit function* is defined as a function  $R : V \times V \longrightarrow 2^V$  satisfying the following three axioms:

- (t1)  $u \in R(u, v)$ , for all  $u, v \in V$ ,
- (t2) R(u,v) = R(v,u), for all  $u, v \in V$ ,
- (t3)  $R(u, u) = \{u\}$ , for all  $u \in V$ .

We may refer R as a transit function on V. If V is the vertex set of a graph G, then we say that R is a transit function on G.

Given a transit function R on V, one can define the underlying graph  $G_R$  of a transit function R on V as the graph with vertex set V, where two distinct vertices u and v are joined by an edge if and only if  $R(u, v) = \{u, v\}$ .

Nebeský initiated an interesting problem of characterizing the interval function I of a connected graph G = (V, E) using a set of simple first-order axioms defined on an arbitrary transit function R on V during the 1990s. Nebeský [17,18] proved that there exists such a characterization for the interval function I(u, v) in terms of an arbitrary transit function R. More such characterizations are described in [19,20,21,22,16].

In the rest of this section, we fix some of the graph theoretical notations and terminology used in this paper. Let G be a graph and H a subgraph of G. The subgraph H is called an *isometric* subgraph of G if the distance  $d_H(u, v)$  between any pair of vertices, u, v in H coincides with that of the distance  $d_G(u, v)$ . H is called an *induced* subgraph if u, v are vertices in H such that uv is an edge in G, then uv must be an edge in H also. A graph G is said to be H-free, if G has no induced subgraph isomorphic to H. A k-wheel is a graph consisting of a cycle on k-vertices,  $C_k$  ( $k \ge 4$ ), and a central vertex adjacent to all the vertices of the cycle. A wheel graph is a k-wheel for some  $k \ge 4$ .

We consider the following metric conditions on a graph G:

- Triangle condition (TC): for any three vertices u, v, w with 1 = d(v, w) < d(v, u) = d(u, w), there exists a common neighbor z of v and w such that d(u, z) = d(u, v) 1.
- Quadrangle condition (QC): for any four vertices u, v, w, y with d(v, y) = d(w, y) = 1 and  $2 = d(v, w) \le d(u, v) = d(u, w) = d(u, y) 1$ , there exists a

common neighbor z of v and w such that d(u, z) = d(u, v) - 1.

It can be easily verified that the conditions (TC) and (QC) can also be defined in terms of the interval  $I_G(u, v)$  instead of the distance d in G. Also, it follows that graphs satisfying (QC) are precisely the class of modular graphs [5].

A graph G is *weakly modular* if its distance function d satisfies the triangle and quadrangle conditions.

We organize the paper as follows. In Section 2, we define diamond-weakly modular graphs and prove that the family of diamond-weakly modular graphs contain the class of bridged graphs and weakly bridged graphs. Further, we prove that the family of diamond-weakly modular graphs are closed under the operation of gated amalgamations. Section ??, we provide the implications of the axioms that we consider in this paper for a general transit function R. In Section 4, we characterize the diamond-weakly modular graph G using the axiom (J0') on its interval function  $I_G$  and discuss the axiomatic characterization of the interval function R. In Section 5, we provide the axiomatic characterizations of bridged graphs and weakly bridged graphs, respectively, using a set of first-order axioms on an arbitrary transit function.

#### 2 Diamond-weakly modular graphs

We first define the diamond-weakly modular graph, for that we introduce a stronger version of triangle condition known as triangle diamond condition (TDC), defined as follows.

- Triangle diamond condition (TDC): For any three vertices u, v, w with 1 = d(v, w) < d(u, v) = d(u, w), there exists a common neighbor z of v and w such that d(u, z) = d(u, v) - 1 and the vertices v, w, z should form diamonds with vertices x and y where d(x, v) = d(y, w) = 1 and d(u, x) = d(u, v) - 1, and d(u, y) = d(u, w) - 1.

A graph G is called *diamond-weakly modular* if its distance function d satisfies (QC) and (TDC). It can be easily verified that the wheel graphs satisfy the (TDC) and hence belong to the class of diamond-weakly modular graphs. Furthermore, diamond-weakly modular graphs contain the class of modular graphs by definition as it satisfies the quadrangle condition.

Now, we prove the interesting subclasses of weakly modular graphs such as bridged graphs and weakly bridged graphs are diamond-weakly modular. For that, we define some more graph theoretic terms defined in [12]. A cycle C is *well-bridged* if and only if, for each v in C, either the neighbors of v in C are adjacent, or  $d_G(v, x) < d_C(v, x)$  for some antipode x of v in C. An *antipode* of a node v of C is a node of C at maximum distance from v along C.

**Theorem 1.** [12] If G is a bridged graph, then every cycle in G is well-bridged.

Using this result, we have the following Lemma that bridged graphs are diamondweakly modular graphs.

## **Lemma 1.** Bridged graphs satisfy (TDC) and hence are diamond-weakly modular graphs.

*Proof.* Suppose G is a bridged graph. Since bridged graphs are weakly modular G satisfies (TC). Let u, y, v be the vertices such that 1 = d(v, y) < d(u, v) = d(u, y) = m and let z be the vertex adjacent to both v and y with d(u, z) = m-1. Suppose x be the vertex adjacent to y on I(u, y) and x' be the vertex adjacent to v on I(u, v). Consider the cycle C:  $xyz \to u \to x$  and has length at least four. Since G is bridged, the cycle C is well bridged by Theorem 1. Consider the vertex y, then either the neighbors of y in C are adjacent, or  $d_G(y, a) < d_C(y, a)$  for some antipode a of y in C. But we have antipode of y in C is u and by assumption  $d_G(u, y) = d_C(u, y) = m$ . Then the only possibility is that the neighbors of y in C are adjacent. That is, x and z are adjacent. Similarly, we can prove that x' and z are adjacent. Hence G satisfies (TDC) and hence G is a diamond-weakly modular graph.

Clearly, the class of bridged graphs form a strict subclass of diamond-weakly modular graphs. For example,  $W_4$  and  $W_5$  are diamond-weakly modular, but they are not bridged graphs.

Weakly bridged graphs are superclass of bridged graphs. In [5], weakly bridged graphs are defined as weakly modular graphs without an induced cycle of length four. Next Lemma shows that weakly bridged graphs are diamond-weakly modular graphs.

#### Lemma 2. Weakly bridged graphs are diamond-weakly modular graphs.

*Proof.* Suppose G is a weakly bridged graph which is not a diamond-weakly modular graph. Since G is weakly modular, it satisfies (TC), but it does not satisfy (TDC). Let u, y, v be the vertices such that 1 = d(v, y) < d(u, v) = d(u, y) = m and let z be the vertex adjacent to both v and y with d(u, z) = m-1. Suppose x be the vertex adjacent to y on I(u, y) and x' be the vertex adjacent to v on I(u, v). We may suppose that z is not adjacent to x. Consider the cycle C:  $xyz \to u \to x$ , which has a length of at least four. Now consider the vertices x, z and u, 2 = d(x, z) and d(u, x) = d(u, z) = m - 1. Apply (QC) on x, z and u, we have a w with w adjacent to both x and z with d(u, w) = m - 2. (Note that w may be equal to u if cycle C has length four). Then the vertices y, x, z, w will induce a cycle of length four, a contradiction to the assumption that G is weakly bridged.

So far, we have modular graphs, bridged graphs, weakly bridged graphs, the graphs  $W_4$  and  $W_5$  are diamond-weakly modular graphs. Note that there are other diamond-weakly modular graphs which do not belong to any of these graphs and some of them are given in figures 1.

Besides these graphs, the graphs obtained by identifying an outer edge of  $W_5$  with an edge in  $K_n$  are diamond-weakly modular graphs. The graphs obtained



Fig. 1. diamond-weakly modular graphs

by identifying an outer edge of  $W_4$  with an edge uv in  $K_n$  and making the central vertex of  $W_4$  adjacent to all the vertices in  $K_n$  other than the vertices u and v are also diamond-weakly modular. Clearly, the graphs formed from any vertex identification of any of the different combination of  $W_4$  and  $W_5$  with  $W_4$ ,  $W_5$  and bridged graph are diamond-weakly modular.

Let  $G_1$  be  $W_4$  or  $W_5$  and  $G_2$  be  $W_4$ ,  $W_5$  or bridged graph. Let  $G_0$  be any isomorphic subgraph of both  $G_1$  and  $G_2$ . Now consider the graph G obtained by identifying  $G_0$  of  $G_1$  and  $G_2$  and making all the vertices in  $V(G_1) \setminus V(G_2)$  adjacent to all the vertices in  $V(G_2) \setminus V(G_1)$ . Then G is diamond-weakly modular.

Let  $y \in V(G)$ . Then a vertex x in the set  $S \subset V$  is a gate in S for y, if  $x \in I(y, w)$  for each  $w \in S$ . The set S is said to be gated if every vertex in G has a gate in S. It is clear that for a vertex  $s \in S$ , s itself is the gate of S. That is, S is gated if and only if every vertex  $y \notin S$  has a gate in S. From the definition of a gated set S, it follows easily that the intersection of two gated sets is gated. The subgraph induced by a gated set S is usually called a gated subgraph of G. Gated amalgamation is an operation that can be performed on gated subsets (subgraphs) to form new gated sets (subgraphs). Gated amalgamation is well known operation in metric graph theory [2]. It is defined for two gated sets  $G_1$  and  $G_2$  of G as follows. Let  $G_1$  and  $G_2$  be gated subgraphs of a graph G such that  $G_1 \cup G_2 = G$  and  $G_1 \cap G_2 \neq \emptyset$ . If in addition there are no edges between  $G_1 \setminus G_2$  and  $G_2 \setminus G_1$  then G is a gated amalgam of  $G_1$  and  $G_2$ , where  $H_1$  and  $H_2$  are isomorphic graphs. Then the gated amalgam of  $G_1$  and  $G_2$  is obtained from  $G_1$  and  $G_2$  by identifying their subgraphs  $H_1$  and  $H_2$ .

**Theorem 2.** Diamond-weakly modular graphs are closed under gated amalgamations.

Proof. Let  $G_1$  and  $G_2$  be two diamond-weakly modular graphs and G be the graph obtained by the gated amalgamation of  $G_1$  and  $G_2$ . Let  $H_1 \sim H_2 = G_0$  be the isomorphic gated subgraphs of  $G_1$  and  $G_2$  which are identified to obtain G. Now consider the vertex u and the edge vy in G with d(u, v) = d(u, y) = m. Then the edge vy lies either in  $G_1$  or in  $G_2$ , since there are no edges between  $G_1 \setminus G_2$  and  $G_2 \setminus G_1$ . Clearly, these three vertices satisfies (TDC) if they all lie either in  $G_1$  or in  $G_2$ . Suppose the vertex u lies in  $G_1 \setminus G_2$  and the edge vy lies

7

in  $G_2 \setminus G_1$ . Let g(u) be the gate of u in  $G_2$ . Then g(u) will be the unique vertex in  $G_2$  closest to u and  $g(u) \in I(u, v) \cap I(u, y)$ . Since d(u, v) = d(u, y) = m and g(u) is the gate of u in  $G_2$ , we get, d(g(u), v) = d(g(u), y). Then the vertices v, y, g(u), with d(v, y) = 1 and d(g(u), v) = d(g(u), y) satisfies (TDC) on G, since, they lie in the diamond-weakly modular graph  $G_2$ . Hence the vertices u, v, ysatisfy (TDC) on G. Similarly, we can prove that G satisfies (QC).

Note that, diamond-weakly modular graphs are not closed under Cartesian products. For example, consider the graph  $K_3 \Box K_2$ . It follows that the graph  $K_3 \Box K_2$  (the prism graph) is a weakly modular graph that doesn't satisfy *TDC*. We also have the following proposition.

**Proposition 1.** A graph G is diamond-weakly modular then G is weakly modular and if G contain  $C_5, W_4^-$  or house as induced subgraphs then there exists a vertex adjacent to all the five vertices of these subgraphs.

*Proof.* Suppose G is a diamond-weakly modular graph. Then clearly, G is a weakly modular graph. Assume G contains  $C_5$  as an induced subgraph with u, x, y, v, x' as its adjacent vertices. For the vertex u and the edge yv with d(u, y) = d(u, v) = 2, there is a vertex z adjacent to all the y, v, x, x' and u, since G is diamond-weakly modular. So there exists a vertex  $z \in G$  such that z is adjacent to all the vertices of  $C_5$ . Similar situation hold in the case of induced  $W_4^-$  and house.

The converse of above Proposition 1 need not be true and is illustrated in Figure  $\,2$ 



**Fig. 2.** The graph in the figure is weakly modular and for the induced subgraphs,  $C_5$ , house, and  $W_4^-$ , there exists a vertex adjacent to all the five vertices of these subgraphs. But it is not diamond-weakly modular, since the vertex u and the edge yv satisfies (TC) but not (TDC). Here  $z_1$  is not adjacent to x and  $z_2$  is not adjacent to x'. Also, the vertices  $u, x, y, z_1$  and v induce a house and  $z_2$  is a vertex adjacent to all the five vertices of the house. But still, the graph do not satisfy (TDC).

#### 3 Betweenness axioms for arbitrary transit function and their implications

In this section, we consider a transit function R defined on a non-empty finite set V and discuss the axioms and their relationships for R. These axioms will be used for the characterization of diamond-weakly modular graphs using the interval function as well as the axiomatic characterization of the interval function of diamond-weakly modular graphs, bridged graphs and weakly bridged graphs using an arbitrary transit function R.

The following three axioms denoted as (b2), (b3), and (b4) together with the defining transit axioms (t1), (t2) are essential and considered by Nebeský and Mulder in the characterizations of the interval function I of a graph.

(b2) if  $x \in R(u, v)$  and  $y \in R(u, x)$ , then  $y \in R(u, v)$ , (b3) if  $x \in R(u, v)$  and  $y \in R(u, x)$ , then  $x \in R(y, v)$ , (b4) if  $x \in R(u, v)$ , then  $R(u, x) \cap R(x, v) = \{x\}$ .

A transit function R satisfying axioms (b2) and (b3) is known as a geometric transit function. Bandelt and Chepoi et al. in [3] introduced the axiom known as (ta) for proving a necessary condition for a geometric transit function to be the interval function of a connected graph. Further in [3], it is proved that the axiom (ta) is satisfied by the interval function of a weakly modular graph.

 $\begin{array}{l} (ta): \text{ if } R(u,v) \cap R(u,w) = \{u\}, R(u,v) \cap R(v,w) = \{v\}, R(u,w) \cap R(v,w) = \{w\} \\ \text{ and } R(u,v) = \{u,v\}, \text{ then } R(u,w) = \{u,w\} \text{ and } R(v,w) = \{v,w\}, \text{ for all } \\ u,v,w \in V. \end{array}$ 

The following axiom (J0) is due to Sholander in his attempt to prove the interval function of trees in [23] during 1952. In [8], this axiom is shown to be a characterizing axiom for the interval function of a Ptolemaic graph, see also [9].

(J0): if 
$$x \in R(u, y), y \in R(x, v)$$
, then  $x \in R(u, v)$ , for all distinct  $u, v, x, y \in V$ .

We consider a slightly modified version of the axiom (J0), defined as follows;

(J0'): if  $x \in R(u, y), y \in R(x, v), R(u, y) \cap R(x, v) \subset \{u, x, y, v\}$ , then  $x \in R(u, v)$ , for all distinct  $u, x, y, v \in V$ .

We prove that this axiom is a characterizing axiom of the diamond-weakly modular graphs in Section 4, where we give a characterization of diamond-weakly modular graphs G using its interval function  $I_G$  and also provide an axiomatic characterization of  $I_G$  in terms of a set of first order axioms on an arbitrary transit function R.

From the definitions of the axioms (J0) and (J0'), we have that the axiom (J0) implies axiom (J0'), while the reverse implication is not true. In other words, axiom (J0') is a weaker axiom than (J0). Example 1 shows that  $(J0') \neq (J0)$ .

Example 1. For a transit function R on V,  $(J0') \neq (J0)$ Let  $V = \{a, b, c, d, e\}$ . Let  $R : V \times V \to 2^V$  defined as follows.  $R(a, e) = \{a, e\}, R(a, b) = \{a, b\}, R(b, e) = \{b, e\}, R(b, c) = \{b, c\}, R(c, e) = \{c, e\}, R(c, d) = \{c, d\}, R(d, e) = \{d, e\}, R(a, c) = \{a, b, c, e\}, R(a, d) = \{a, e, d\}, R(b, d) = \{c, d\}, R(b, d) = \{c, d\}, R(c, d) = \{c, d$   $\{b, c, d, e\}, R(x, y) = R(y, x) \text{ and } R(x, x) = \{x\}, \text{ for all } x, y \text{ in } V.$  We can see that  $b \in R(a, c)$  and  $c \in R(b, d)$  but  $b \notin R(a, d)$ , so that R does not satisfy (J0) axiom. We can see that there exists no u, v, x, y and z satisfying the assumptions of the axiom (J0') and hence the axiom (J0') follows trivially.

In addition to the axiom (J0'), we consider the axioms (br) and (br') for a transit function R. These axioms are essential for the characterizations of bridged graphs and weakly bridged graphs, which we prove in Section 5.

- (br): if  $R(x, y) = \{x, y\}, R(x, u) = \{x, u\}, R(v, y) = \{v, y\}$  and  $z \in R(u, v)$ , then  $R(x, z) = \{x, z\}$  or  $R(y, z) = \{y, z\}$ , for any u, v, x, y, z.
- (br'): if  $R(u, x) = \{u, x\}, R(x, v) = \{x, v\}$  and  $z \in R(u, v)$ , then  $R(x, z) = \{x, z\}$ , for any u, v, x, z.

We now state the Mulder- Nebeský Theorem stated in [16] characterizing the interval function of an arbitrary connected graph using axioms on an arbitrary transit function. In addition to the transit axioms (t1), (t2) and the betweenness axioms (b2), (b3), (b4), the following axioms (s1) and (s2) are required in the theorem.

- $(s1): R(u,\bar{u}) = \{u,\bar{u}\}, R(v,\bar{v}) = \{v,\bar{v}\}, u \in R(\bar{u},\bar{v}) \text{ and } \bar{u}, \bar{v} \in R(u,v), \text{ then } v \in R(\bar{u},\bar{v}).$
- $(s2) : R(u,\bar{u}) = \{u,\bar{u}\}, R(v,\bar{v}) = \{v,\bar{v}\}, \bar{u} \in R(u,v), v \notin R(\bar{u},\bar{v}), \bar{v} \notin R(u,v),$ then  $\bar{u} \in R(u,\bar{v}).$

**Theorem 3.** [16] Let  $R: V \times V \longrightarrow 2^V$  be a function on V, satisfying the axioms (t1), (t2), (b2), (b3), (b4) with the underlying graph  $G_R$  and let I be the interval function of  $G_R$ . The following statements are equivalent.

- (a) R = I
- (b) R satisfies axioms (s1) and (s2).

The axiom (ta) defined by Chepoi et al. in [3] is needed for the characterization of  $I_G$  of these graph classes (diamond-weakly modular, bridged graphs and weakly bridged graphs). Also, it is proved in [3] that discrete geometric interval space X satisfying the (ta) axiom is graphic. In our terminology, it means that if R satisfies axioms (b2), (b3) and (ta) then R will coincide with the interval function of the underlying graph  $G_R$ .

Though, we give another proof establishing the dependency of the axioms (b2), (b3), (ta), (s1) and (s2).

**Lemma 3.** Let R be a transit function on a nonempty set V satisfies the axioms (b2) and (b3) then for any u, v and w, there exists  $x \in R(u, v) \cap R(u, w)$  such that  $R(x, v) \cap R(x, w) = \{x\}$ .

*Proof.* We have  $u \in R(u, v) \cap R(u, w)$ . If  $R(u, v) \cap R(u, w) = \{u\}$ , then u is the required x and completes the proof. If not, there exist a  $x_1$  such that  $x_1 \in R(u, v) \cap R(u, w)$ . Then  $x_1 \in R(u, v)$  and  $x_1 \in R(u, w)$  and by (b1) we have  $u \notin R(x_1, v)$  and  $u \notin R(x_1, w)$  imply that  $R(x_1, v) \subset R(u, v)$  and  $R(x_1, w) \subset R(u, v)$  and  $R(x_1, w)$  and  $R(x_1, w) \subset R(u, v)$  and  $R(x_1, w)$  and  $R(x_1, w)$  and  $R(x_1, w)$  and  $R(x_1, w) \subset R(u, v)$  and  $R(x_1, w) \subset R(u, v)$  and  $R(x_1, w)$  an

 $\begin{array}{l} R(u,w) \text{ by } (b2). \text{ Hence } R(x_1,v) \cap R(x_1,w) \subset R(u,v) \cap R(u,w) \text{ and } \mid R(x_1,v) \cap R(x_1,w) \mid < \mid R(u,v) \cap R(u,w) \mid . \text{ If } R(x_1,v) \cap R(x_1,w) = \{x_1\}, \text{ complete the proof with } x = x_1. \text{ Otherwise, there exist } x_2 \text{ such that } x_2 \in R(x_1,v) \cap R(x_1,w) \text{ and by the above argument we have } R(x_2,v) \cap R(x_2,w) \subset R(x_1,v) \cap R(x_1,w) \subset R(u,v) \cap R(u,w) \text{ and } \mid R(x_2,v) \cap R(x_2,w) \mid < \mid R(x_1,v) \cap R(x_1,w) \mid < \mid R(u,v) \cap R(u,w) \mid. \text{ If } R(x_2,v) \cap R(x_2,w) = \{x_2\}, \text{ complete the proof with } x = x_2. \text{ Otherwise, since } V \text{ is finite, we can find } x_n \text{ with } x_n \in R(x_{n-1},v) \cap R(x_{n-1},w) \text{ such that } R(x_n,v) \cap R(x_n,w) \subset R(x_{n-1},v) \cap R(x_{n-1},w) \subset \dots \subset R(u,v) \cap R(u,w) \text{ and } \mid R(x_n,v) \cap R(x_n,w) \mid < \mid R(x_{n-1},v) \cap R(x_{n-1},w) \mid < \dots < \mid R(u,v) \cap R(u,w) \mid \\ \text{ and } R(x_n,v) \cap R(x_n,w) = \{x_n\}. \end{array}$ 

**Proposition 2.** A transit function R defined on a non-empty set V satisfies axioms (b2), (b3) and (ta) then R satisfies axioms (s1) and (s2).

*Proof.* Let R be a transit function satisfying (b2), (b3) and (ta). We have to show that R satisfies (s1). Suppose not. That is  $R(u, \bar{u}) = \{u, \bar{u}\}, R(v, \bar{v}) = \{v, \bar{v}\}, u \in R(\bar{u}, \bar{v})$  and  $\bar{u}, \bar{v} \in R(u, v)$ , and  $v \notin R(\bar{u}, \bar{v})$ . Consider the vertices  $\bar{u}, v, \bar{v}$ . claim 1:  $\bar{v} \notin R(\bar{u}, v)$  and  $u \notin R(\bar{u}, v)$ .

If  $\bar{v} \in R(\bar{u}, v)$ , we have  $\bar{u} \in R(u, v)$  and  $\bar{v} \in R(\bar{u}, v) \Rightarrow \bar{u} \in R(u, \bar{v})$  by (b3) and then  $u \in R(\bar{u}, \bar{v}), \bar{u} \in R(u, \bar{v}) \Rightarrow u \in R(\bar{u}, \bar{u})$ , a contradiction. Therefore  $\bar{v} \notin R(\bar{u}, v)$ . Also since  $\bar{u} \in R(u, v)$ , by (b1) we have  $u \notin R(\bar{u}, v)$ . Hence the claim.

Claim 2:  $R(\bar{u}, \bar{v}) \cap R(\bar{u}, v) \setminus \{\bar{u}\} \neq \phi$ .

If  $R(\bar{u}, \bar{v}) \cap R(\bar{u}, v) \setminus \{\bar{u}\} = \phi$ , then  $R(\bar{u}, \bar{v}) \cap R(\bar{u}, v) = \{\bar{u}\}, R(\bar{u}, \bar{v}) \cap R(\bar{v}, v) = \{\bar{u}\}, R(\bar{v}, v) \in \{\bar{v}\}, R(\bar{v},$  $\{\bar{v}\}$  and  $R(\bar{u},v) \cap R(\bar{v},v) = \{v\}$  and  $R(v,\bar{v}) = \{v,\bar{v}\}$  then by (ta) we can say that  $R(\bar{u}, \bar{v}) = \{\bar{u}, \bar{v}\}$ , a contradiction since  $u \in R(\bar{u}, \bar{v})$ . Therefore there exist a  $w_1$  with  $w_1 \neq \bar{u}$  such that  $w_1 \in R(\bar{u}, \bar{v}) \cap R(\bar{u}, v)$ . Hence the claim. Since  $w_1 \in R(\bar{u}, \bar{v}) \cap R(\bar{u}, v)$  by lemma 3, we can find  $y_1$  such that  $R(y_1, \bar{v}) \cap R(y_1, v) =$  $\{y_1\}$ , and by (ta) on  $y_1, v, \bar{v}$ , we have  $R(y_1, \bar{v}) = \{y_1, \bar{v}\}$  and  $R(y_1, v) = \{y_1, v\}$ . Since  $\bar{u} \in R(u, v), y_1 \in R(\bar{u}, v)$  by (b3) we have  $\bar{u} \in R(u, y_1)$ . Now consider the vertices  $u, \bar{v}, y_1$ , then analogous to claim1, here we can prove that  $\bar{v} \notin R(u, y_1)$ and  $\bar{u}, y_1 \notin R(u, \bar{v})$  by (b3). Also we can prove  $R(u, \bar{v}) \cap R(u, y_1) \setminus \{u\} \neq \phi$  by applying (ta) on  $u, \bar{v}, y_1$ , therefore there exist a  $y_2$  with  $y_2 \neq u$  such that  $y_2 \in$  $R(u, \bar{v}) \cap R(u, y_1)$  and  $R(y_2, \bar{v}) \cap R(y_2, y_1) = \{y_2\}$ . Then by (ta) on  $y_2, y_1, \bar{v}$ , we have  $R(y_2, \bar{v}) = \{y_2, \bar{v}\}$  and  $R(y_2, y_1) = \{y_2, y_1\}$ . Since  $u \in R(\bar{u}, \bar{v}), y_2 \in R(u, \bar{v})$ , we have  $u \in R(\bar{u}, y_2)$  by (b3). Now consider  $\bar{u}, y_1, y_2$ , and apply the same as above, we can find a  $y_3$  such that  $y_3 \in R(\bar{u}, y_1) \cap R(\bar{u}, y_2)$  and  $R(y_3, y_1) = \{y_3, y_1\}$ and  $R(y_3, y_2) = \{y_3, y_2\}$ . Also  $\bar{u} \in R(u, y_3)$ . Continuing like this we can find  $y_4, y_5, \dots$  such that  $R(y_k, y_{k-1}) = \{y_k, y_{k-1}\}$  and  $R(y_k, y_{k-2}) = \{y_k, y_{k-2}\}$  and for some k = 2m,  $u \in R(\bar{u}, y_k)$  and  $\bar{u} \in R(u, y_{k-1})$ . Since V is finite, we can find some n such that either  $y_n = u$  or  $y_n = \bar{u}$ . If  $y_n = u$ , we have  $R(y_n, y_{n-1}) = \{y_n, y_{n-1}\}$  and  $R(y_n, y_{n-2}) = \{y_n, y_{n-2}\}$  and  $u \in R(\bar{u}, y_n)$ and also  $\bar{u} \in R(u, y_{n-1})$ , a contradiction since  $R(u, y_{n-1}) = R(y_n, y_{n-1}) =$  $\{y_n, y_{n-1}\}$ . If  $y_n = \bar{u}$ , then  $\bar{u} \in R(u, y_n)$  and also  $u \in R(\bar{u}, y_{n-1})$ , a contradiction since  $R(\bar{u}, y_{n-1}) = R(y_n, y_{n-1}) = \{y_n, y_{n-1}\}$ . In both case we have a contradiction. Therefore  $v \in R(\bar{u}, \bar{v})$ .

Now we have to show that R satisfies axiom (s2). Suppose not. That is  $R(u,\bar{u}) = \{u,\bar{u}\}, R(v,\bar{v}) = \{v,\bar{v}\}, \bar{u} \in R(u,v), v \notin R(\bar{u},\bar{v}), \bar{v} \notin R(u,v)$  and  $\bar{u} \notin R(u,\bar{v})$ .

Claim 3:  $u \notin R(\bar{u}, \bar{v})$ . Suppose  $u \in R(\bar{u}, \bar{v})$ . Consider the vertices  $\bar{u}, \bar{v}, v$ . Since  $\bar{u} \in R(u, v)$  by (b1) we have  $u \notin R(\bar{u}, v)$ . Also  $\bar{v} \notin R(\bar{u}, v)$  since  $\bar{v} \notin R(u, v)$  and  $v \notin R(\bar{u}, \bar{v})$ . From these we have  $R(\bar{u}, v) \cap R(\bar{v}, v) = \{v\}$ ,  $R(\bar{u}, \bar{v}) \cap R(\bar{v}, v) = \{\bar{v}\}$  and  $R(\bar{v}, v) = \{\bar{v}, v\}$  also  $u \in R(\bar{u}, \bar{v})$ . Then by (ta),  $R(\bar{u}, v) \cap R(\bar{u}, \bar{v}) \setminus \{\bar{u}\} \neq \phi$ . That is there exist  $w_1$  such that  $w_1 \in R(\bar{u}, v) \cap R(\bar{u}, \bar{v})$ . Hence by lemma 3, there exist a  $y_1$  such that  $R(y_1, v) \cap R(y_1, \bar{v}) = \{y_1\}$ . Then by (ta) on  $y_1, \bar{v}, v$ , we have  $R(y_1, \bar{v}) = \{y_1, \bar{v}\}$  and  $R(y_1, v) = \{y_1, v\}$ . Also  $\bar{u} \in R(u, y_1)$ . That is we have  $R(u, \bar{u}) = \{u, \bar{u}\}, R(\bar{v}, y_1) = \{\bar{v}, y_1\}, u, y_1 \in R(\bar{u}, \bar{v})$ , and  $\bar{u} \in R(u, y_1)$ , then by (s1)  $\bar{v} \in R(u, y_1) \subseteq R(u, v)$ . That is  $\bar{v} \in R(u, v)$ , a contradiction. Therefore  $u \notin R(\bar{u}, \bar{v})$  and hence claim 3.

Now consider  $u, \bar{u}, \bar{v}$ . We have  $u \notin R(\bar{u}, \bar{v}), \bar{u} \notin R(u, \bar{v})$  and  $v \notin R(\bar{u}, \bar{v})$ . Also  $v \notin R(u, \bar{v})$ , since  $\bar{u} \notin R(u, \bar{v})$ . Apply (ta) on  $u, \bar{u}, \bar{v}$ , we have two cases

**case 1**  $R(u, \bar{v}) \cap R(\bar{u}, \bar{v}) = \{\bar{v}\}$ . Which implies that  $R(\bar{u}, \bar{v}) = \{\bar{u}, \bar{v}\}$  and  $R(u, \bar{v}) = \{u, \bar{v}\}$  by (ta). Then consider  $v, \bar{u}, \bar{v}$ , also it is clear that  $R(\bar{u}, \bar{v}) = \{\bar{u}, \bar{v}\}$  and  $R(v, \bar{v}) = \{v, \bar{v}\}$ , then by (ta) on  $v, \bar{u}, \bar{v}$ , we have  $R(\bar{u}, v) = \{\bar{u}, v\}$ . Consider  $u, v, \bar{v}$ . We have  $\bar{u}, v \notin R(u, \bar{v})$  and  $\bar{v} \notin R(u, v)$ . Then  $R(u, \bar{v}) \cap R(v, \bar{v}) = \{\bar{v}\}$ ,  $R(u, v) \cap R(v, \bar{v}) = \{v\}$  and  $R(u, v) \cap R(u, \bar{v}) = \{u\}$  and  $R(v, \bar{v}) = \{v, \bar{v}\}$ , by (ta) we have  $R(u, v) = \{u, v\}$ , contradiction since  $\bar{u} \in R(u, v)$ .

**case 2**  $R(u, \bar{v}) \cap R(\bar{u}, \bar{v}) \neq \{\bar{v}\}$ . That is there exist a  $x_1$  such that  $x_1 \in R(u, \bar{v}) \cap R(\bar{u}, \bar{v})$ , and by lemma 3, we may assume that  $R(x_1, u) \cap R(x_1, \bar{u}) = \{x_1\}$  and by (ta) on  $u, x_1, \bar{u}$ , we have  $R(u, x_1) = \{u, x_1\}$  and  $R(\bar{u}, x_1) = \{\bar{u}, x_1\}$ . Then  $x_1 \in R(u, \bar{v})$ , but  $x_1 \notin R(\bar{u}, v)$ . Now consider  $\bar{u}, x_1, v$ .

Claim 4:  $\bar{u} \notin R(x_1, v)$ . Suppose  $\bar{u} \in R(x_1, v)$ . Consider the vertices  $\bar{u}, \bar{v}, v$ , we have  $R(\bar{u}, \bar{v}) \cap R(v, \bar{v}) = \{\bar{v}\}$ ,  $R(\bar{u}, v) \cap R(v, \bar{v}) = \{v\}$ ,  $R(v, \bar{v}) = \{v, \bar{v}\}$ . If  $R(\bar{u}, \bar{v}) \cap R(\bar{u}, v) = \{\bar{u}\}$ , then by (ta), we get  $R(\bar{u}, \bar{v}) = \{\bar{u}, \bar{v}\}$ , a contradiction as  $x_1 \in R(\bar{u}, \bar{v})$ . So there exist a  $x_2$  such that  $x_2 \in R(\bar{u}, \bar{v}) \cap R(\bar{u}, v)$ .  $\bar{u} \in R(x_1, v)$ and  $x_2 \in R(\bar{u}, \bar{v})$  implies  $\bar{u} \in R(x_1, x_2)$  by (b3). Now applay (s1) on the vertices  $\bar{u}, x_1, \bar{v}, x_2$ , we get  $\bar{v} \in R(x_1, x_2)$ . Also  $\bar{u} \in R(x_1, v)$ ,  $x_2 \in R(\bar{u}, v)$  implies  $x_2 \in R(x_1, v)$  by (b2) and  $\bar{v} \in R(x_1, x_2)$ ,  $x_2 \in R(x_1, v)$  implies  $\bar{v} \in R(x_1, v)$ by (b2). By (s1) on  $\bar{u}, x_1, \bar{v}, v$ , we get  $v \in R(\bar{u}, \bar{v})$ , a contradiction. Therefore  $\bar{u} \notin R(x_1, v)$ , hence the claim. Now we have  $x_1, \bar{v} \notin R(\bar{u}, v)$  and  $\bar{u} \notin R(x_1, v)$ . Then  $R(\bar{u}, x_1) \cap R(\bar{u}, v) = \{\bar{u}\}$ ,  $R(\bar{u}, x_1) \cap R(x_1, v) = \{x_1\}$ ,  $R(\bar{u}, x_1) = \{\bar{u}, x_1\}$ , then we have to consider two cases

**case 2.1** If  $R(\bar{u}, v) \cap R(x_1, v) = \{v\}$ , then by (ta) on  $\bar{u}, x_1, v$ , we get  $R(\bar{u}, v) = \{\bar{u}, v\}$  and  $R(x_1, v) = \{x_1, v\}$ . Now consider  $\bar{u}, v, \bar{v}$ , we have  $R(\bar{u}, v) = \{\bar{u}, v\}$ ,  $R(v, \bar{v}) = \{v, \bar{v}\}$  and  $x_1 \notin R(\bar{u}, v)$ . Apply (ta) on  $\bar{u}, \bar{v}, v$ , implies that  $R(\bar{u}, \bar{v}) = \{\bar{u}, \bar{v}\}$ , contradiction to  $x_1 \in R(\bar{u}, \bar{v})$ .

**case 2.2**  $R(\bar{u}, v) \cap R(x_1, v) \neq \{v\}$ . Then there exist  $x_2$  such that  $x_2 \in R(\bar{u}, v) \cap R(x_1, v)$ , and also by the lemma 3 we may assume that  $R(\bar{u}, x_2) \cap R(x_1, x_2) = \{x_2\}$  and by (ta) on  $x_2, x_1, \bar{u}$  we have  $R(\bar{u}, x_2) = \{\bar{u}, x_2\}$  and  $R(x_1, x_2) = \{x_1, x_2\}$ . Also  $x_2 \in R(\bar{u}, v) \subseteq R(u, v)$ . That is  $x_2 \in R(u, v)$ , but  $x_2 \notin R(u, \bar{v})$ . Now consider  $x_1, x_2, \bar{v}$ , by (ta) we get a  $x_3$  such that  $x_3 \notin R(\bar{u}, v)$ , but  $x_3 \in R(\bar{u}, v)$ .  $R(u, \bar{v})$ . Continuing like this we get  $x_4, x_5, ..., x_n$  such that  $x_n = v$  or  $x_n = \bar{v}$ . If  $x_n = v$ , then  $x_n \in R(u, v)$ , but  $x_n \notin R(u, \bar{v})$  and if  $x_n = \bar{v}$ , then  $x_n \notin R(u, v)$ , but  $x_n \in R(u, \bar{v})$ . Therefore in all the cases we have  $R(u, \bar{v}) \cap R(v, \bar{v}) = \{\bar{v}\}, R(u, v) \cap R(v, \bar{v}) = \{v\}$  and  $R(u, v) \cap R(u, \bar{v}) = \{u\}$  and  $R(v, \bar{v}) = \{v, \bar{v}\}$ , and by (ta) on  $v, u, \bar{v}$ , we get  $R(u, v) = \{u, v\}$ , which is the final contradiction, since  $\bar{u} \in R(u, v)$ . Therefore  $\bar{u} \in R(u, \bar{v})$ .

The following straightforward Lemma for the connectedness of the underlying graph  $G_R$  of a transit function R is proved in [7].

**Lemma 4.** [7] If the transit function R on a non-empty set V satisfies axioms (b1) and (b2), then the underlying graph  $G_R$  of R is connected.

We have the following Theorem.

**Theorem 4.** If R is a transit function on V satisfying the axioms (J0') and (b3), then R satisfies axiom (b2) and  $G_R$  is connected.

*Proof.* Let R satisfies axioms (J0') and (b3). To prove R satisfies (b2). For  $u, x, y, v \in V$ , let  $x \in R(u, v)$ , and  $y \in R(u, x)$ . Since R satisfies (b3), we have  $x \in R(u, v), y \in R(u, x) \implies x \in R(y, v)$ . In axiom (J0') the four elements are distinct, so the required minimum cardinality of the set V is four and for  $|V| \leq 3$ , (b2) holds trivially. We use induction on  $|V| \geq 4$  and we explicitly prove cases where |V| = 4, 5, 6, 7. We also use the fact that axiom (b3) implies (b1).

**Case 1:** |V| = 4.

Let  $V = \{u, x, y, v\}$ . Here  $y \in R(u, x), x \in R(y, v)$ . Since  $x \in R(u, v)$ , we have  $v \notin R(u, x)$  by (b1). Then  $R(u, x) \cap R(y, v) \subset \{u, x, y, v\}$  implies that  $y \in R(u, v)$  by (J0').

**Case 2:** |V| = 5.

Let  $V = \{u, x, w_1, y, v\}$ . We have  $y \in R(u, x), x \in R(y, v)$  and if  $R(u, x) \cap R(y, v) \subset \{u, x, y, v\}$  implies that  $y \in R(u, v)$  by (J0') and we are done. Assume  $w_1 \in R(u, x) \cap R(y, v)$  with  $w_1 \notin \{u, x, y, v\}$ . Now apply (b3) we get the following.  $x \in R(u, v), w_1 \in R(u, x) \implies x \in R(w_1, v), w_1 \in R(y, v), x \in R(w_1, v) \implies w_1 \in R(y, x)$  and  $y \in R(u, x), w_1 \in R(y, x) \implies y \in R(u, w_1)$ .

**Claim**  $x, v \notin R(u, w_1)$ . Since  $w_1 \in R(u, x)$ ,  $x \notin R(u, w_1)$  by (b1). Suppose  $v \in R(u, w_1)$ , also we have  $x \in R(v, w_1) \implies v \in R(u, x)$  by (b3), a contradiction to (b1) as  $x \in R(u, v)$ . That is  $x, v \notin R(u, w_1)$ . Hence the claim. Now we have  $y \in R(u, w_1), w_1 \in R(y, v)$  and  $R(u, w_1) \cap R(y, v) \subset \{u, w_1, y, v\} \implies y \in R(u, v)$  by (J0').

**Case 3:** |V| = 6.

Let  $V = \{u, x, w_1, w_2, y, v\}$ . Here also  $y \in R(u, x), x \in R(y, v)$  and if  $R(u, x) \cap R(y, v) \subset \{u, x, y, v\}$  implies that  $y \in R(u, v)$ . If  $R(u, x) \cap R(y, v) \not\subset \{u, x, y, v\}$ . That is either  $w_1$  or  $w_2$  or both belongs to  $R(u, x) \cap R(y, v)$ . Let us assume  $w_1 \in R(u, x) \cap R(y, v)$  with  $w_1 \notin \{u, x, y, v\}$ . Then similar to case 2, we get either  $y \in R(u, v)$ , if  $R(u, w_1) \cap R(y, v) \subset \{u, w_1, y, v\}$  or  $w_2 \in R(u, w_1) \cap R(y, v)$  with  $w_2 \notin \{u, x, w_1, y, v\}$ . Then by (b3), we get  $w_1 \in R(w_2, x)$ , since  $w_1 \in R(w_2, x)$ .  $R(u, x), w_2 \in R(u, w_1).$ 

**Claim**  $u, y \notin R(w_1, v)$ . Since  $w_1 \in R(y, v), y \notin R(w_1, v)$  by (b1). Suppose  $u \in R(w_1, v)$  and we have  $w_1 \in R(y, v), u \in R(w_1, v) \Longrightarrow w_1 \in R(y, u)$  by (b3), a contradiction to (b1) as  $y \in R(u, w_1)$ . Hence the claim. Now  $x \in R(v, w_1), w_1 \in R(x, w_2)$  and  $R(v, w_1) \cap R(x, w_2) \subset \{v, w_1, x, w_2\}$  implies  $x, w_1 \in R(v, w_2)$  by (J0'). Using (b3) we have the following.  $w_2 \in R(y, v), x \in R(w_2, v) \Longrightarrow w_2 \in R(y, x)$  and  $y \in R(u, x), w_2 \in R(y, x) \Longrightarrow y \in R(u, w_2)$ .

**Claim**  $x, v, w_1 \notin R(u, w_2)$ . Since  $w_2 \in R(u, w_1)$  by (b1) we get  $w_1 \notin R(u, w_2)$ . Suppose  $x \in R(u, w_2)$  and we have  $w_1 \in R(x, w_2) \implies x \in R(u, w_1)$  by (b3), a contradiction to (b1) as  $w_1 \in R(u, x)$ . Suppose  $v \in R(u, w_2)$  and we have  $x \in R(v, w_2)$  then by  $(b3), v \in R(u, x)$ , a contradiction to (b1) as  $x \in R(u, v)$ . Hence the claim. Thus  $R(u, w_2) \cap R(y, v) \subset \{u, w_2, y, v\}$ . That is  $y \in R(u, w_2), w_2 \in R(y, v)$  and  $R(u, w_2) \cap R(y, v) \subset \{u, w_2, y, v\}$  implies that  $y \in R(u, v)$  by (J0'). **Case 4:** |V| = 7.

Let  $V = \{u, x, w_1, w_2, w_3, y, v\}$ . Then similar to case 3, we can prove the following. Either  $y \in R(u, v)$ , or  $w_3 \in R(u, w_2) \cap R(y, v)$  with  $w_3 \notin \{u, x, w_1, w_2, y, v\}$  and by (b3) we get  $w_2 \in R(w_3, w_1)$ , since  $w_2 \in R(u, w_1), w_3 \in R(u, w_2)$ . Also it is easy to get  $u, y, v \notin R(w_2, x)$ . Then by (J0)' on  $x, w_1, w_2, w_3$ , we get  $w_1, w_2 \in R(x, w_3)$ . Using (b1) and (b3) we can easily prove the Claims:  $u, y, \notin R(w_2, v)$  and  $x \notin R(w_3, w_1)$ , since  $w_1, w_2 \in R(x, w_3)$ . Then  $w_1 \in R(v, w_2), w_2 \in R(w_1, w_3)$  and  $R(v, w_2) \cap R(w_1, w_3) \subset \{v, w_1, w_2, w_3\}$  implies  $w_1, w_2 \in R(u, w_3)$  by (J0'). Then by continuous application of axiom (b3), we get  $y \in R(u, w_3)$  and we can prove the claim  $x, v, w_1, w_2 \notin R(u, w_3)$  easily as in case 3. Then  $y \in R(u, w_3), w_3 \in R(y, v)$  and  $R(u, w_3) \cap R(y, v) \subset \{u, w_3, y, v\}$  implies that  $y \in R(u, v)$  by axiom (J0').

**Case** n: |V| = n, n > 7.

Let  $V = \{u, x, w_1, w_2, w_3, \dots, w_n, y, v\}$ . Like the above cases, we have either  $y \in R(u, v)$  or  $w_n \in R(u, w_{n-1}) \cap R(y, v)$  with  $w_n \notin \{u, x, w_1, w_2, \dots, w_{n-1}, y, v\}$ . From the above cases we get the following:  $w_i \in R(u, w_{i-1}), w_{i-2}, w_{i-1} \in$  $R(w_{i-3}, w_i), w_{i-2}, w_{i-1} \in R(v, w_i), y \in R(u, w_i) \text{ and } w_{i-1} \in R(w_i, w_{i-2}) \text{ for } 4 \leq 0$  $i \leq n-1$ . Using these we can easily prove the case |V| = n like the above cases. Now  $w_{n-1} \in R(u, w_{n-2}), w_n \in R(u, w_{n-1}) \implies w_{n-1} \in R(w_n, w_{n-2})$  by (b3). Also it is easy to get the claim  $u, y, v, x, w_1, \ldots, w_{n-4} \notin R(w_{n-1}, w_{n-3})$ . Then by (J0)' on  $w_{n-3}, w_{n-2}, w_{n-1}, w_n$ , we get  $w_{n-2}, w_{n-1} \in R(w_{n-3}, w_n)$ . Also we can prove the Claim:  $u, y \notin R(w_{n-1}, v)$  and  $x, w_1, \ldots, w_{n-3} \notin R(w_n, w_{n-2})$  using axiom (b1) and (b3) easily. That is  $w_{n-2} \in R(v, w_{n-1}), w_{n-1} \in R(w_n, w_{n-2})$  and  $R(v, w_{n-1}) \cap R(w_n, w_{n-2}) \subset \{v, w_{n-2}, w_{n-1}, w_n\} \implies w_{n-2}, w_{n-1} \in R(v, w_n)$ by (J0'). Then by (b3) we have the following.  $w_n \in R(y, v), w_{n-1} \in R(w_n, v) \implies$  $w_n \in R(y, w_{n-1})$  and  $y \in R(u, w_{n-1}), w_n \in R(y, w_{n-1}) \implies y \in R(u, w_n)$ . We can prove the claim  $x, v, w_1, w_2, \ldots, w_{n-1} \notin R(u, w_n)$  easily as in the above cases. Then  $y \in R(u, w_n), w_n \in R(y, v)$  and  $R(u, w_n) \cap R(y, v) \subset \{u, w_n, y, v\}$ implies that  $y \in R(u, v)$  by axiom (J0'), which completes the proof that R satisfies axiom (b2). Since for a transit function R, axiom (b3) implies axiom (b1)and by Lemma 4, it follows that  $G_R$  is connected.

14 M. Changat, L.K. Sheela

#### 4 Interval function of diamond-weakly modular graphs

In this section, we prove the main Theorem characterizing a diamond-weakly modular graph G using the (J0') axiom on the interval function  $I_G$  of G, which in turn provide us an axiomatic characterization of the interval function  $I_G$  of G.

**Theorem 5.** The interval function  $I_G$  of a connected graph G satisfies the axiom (J0') if and only if G is a diamond-weakly modular graph.

*Proof.* We prove the contra-positive of the statement of the theorem. That is, we prove that the interval function  $I_G$  of a connected graph G doesn't satisfy the axiom (J0') if and only if G is not a diamond-weakly modular graph. First assume G is not a diamond-weakly modular graph. Then the distance function ddoesn't satisfy either (TDC) or (QC) or both on G. Assume that d doesn't satisfy the (TDC). That is, for any three vertices u, v, y with 1 = d(v, y) < d(u, v) =d(u, y) = k and there does not exists a common neighbor z of v, x, x' and y such that d(u, z) = d(u, v) - 1 where x is the neighbor of y in the u, y-shortest path and x' is the neighbor of v in the u, v-shortest path. Then clearly,  $x \in I(u, y)$ ,  $y \in I(x, v)$  and  $I(u, y) \cap I(x, v) \subset \{u, x, y, v\}$ , but  $x \notin I(u, v)$ . So I do not satisfy axiom (J0') on G. Now assume that d doesn't satisfy the (QC). That is, for any four vertices u, v, w, y with d(v, w) = d(w, y) = 1 and  $2 = d(v, y) \leq d(u, v) = 0$ d(u, y) = d(u, w) - 1 and there does not exists a common neighbor z of v and y such that d(u, z) = d(u, v) - 1. Let x be the neighbor of y in the u, y-shortest path. Then clearly  $x \in I(u, y), y \in I(x, v)$  and  $I(u, y) \cap I(x, v) \subset \{u, x, y, v\}$ , but  $x \notin I(u, v)$ . So I does not satisfy axiom (J0') on G.

To prove the converse part: Assume that the interval function I of a connected graph G doesn't satisfy the axiom (J0'). That is, there exists vertices u, x, y, v in G such that  $x \in I(u, y), y \in I(x, v), I(u, y) \cap I(x, v) \subset \{u, x, y, v\}$  and  $x \notin I(u, v)$ . That is, there exist a u, y-shortest path containing x and a x, v-shortest path containing y such that x doesn't belong to a u, v-shortest path in G. From this, we infer that x and y are adjacent and that  $d(u, y), d(x, v), d(u, v) \ge 2$ . Without loss of generality, we can take the vertices u, x, y, v such that y is at a maximum distance say m from u and let d(y, v) = n.

Consider the vertex  $y_1 \in I(y, v)$  which is adjacent to y. We may choose  $y_1$  such that the distance  $d(u, y_1)$  is minimum. Then the possibilities of  $d(u, y_1)$  are m-1, m or m+1. If  $d(u, y_1) = m-1$ , then  $y_1 \in I(u, y) \cap I(x, v)$ , which is a contradiction to the assumption that u, x, y, v does not satisfy axiom (J0'), so that  $d(u, y_1) = m-1$  is not possible.

When  $d(u, y_1) = m$ : consider the vertices  $u, x, y, y_1$ . Clearly, these vertices do not satisfy the axiom (J0'). Let  $y' \in I(u, y_1)$  which is adjacent to  $y_1$ . Then,  $d(u, y) = d(u, y_1) = m$  and  $d(y, y_1) = 1$ . If possible, let the vertices  $u, y, y_1$ satisfies (TDC). Then there exists a vertex z adjacent to both y and  $y_1$  with d(u, z) = m - 1 and z is adjacent to both x and y'. Then,  $z \in I(u, y) \cap I(x, v)$ and so  $I(u, y) \cap I(x, v) \not\subset \{u, x, y, v\}$  and hence the vertices u, x, y, v satisfies the axiom (J0'). Hence the vertices  $u, y, y_1$  will not satisfy (TDC) and so G is not diamond-weakly modular graph.

When  $d(u, y_1) = m + 1$ : consider the vertices  $u, y, y_1, v$ . Clearly,  $y \in I(u, y_1)$ and  $y_1 \in I(y, v)$ . If  $I(u, y_1) \cap I(y, v) \subset \{u, y_1, y, v\}$ , then either  $y \in I(u, v)$  or  $y \notin I(u, v)$ . But both of these will yield a contradiction. For, if  $y \in I(u, v)$  then  $x \in I(u, v)$  (by axiom (b2)); if  $y \notin I(u, v)$ , then we get four vertices  $u, y, y_1, v$ which does not satisfy (J0') and  $d(u, y_1) > m$ , a contradiction to the maximality of d(u, y). So this implies that  $I(u, y_1) \cap I(y, v) \not\subset \{u, y_1, y, v\}$ . Then, there exist a  $w \neq u, y_1, y, v$  such that  $w \in I(u, y_1) \cap I(y, v)$ . Let P be the shortest  $uy_1$ path containing w and Q be the shortest y, v-path containing w. Also let t be the length of shortest u, w- subpath of P and  $\ell$  be the length of shortest w, ysubpath of Q .

**Claim**:  $\ell + t = m + 2$ . We have d(u, y) = m and  $d(u, y_1) = m + 1$ . So possibilities of  $\ell + t$  are m, m + 1 and m + 2. Let  $w_1$  be the neighbor of y in the shortest w, ysubpath of Q. Also  $w_1 \in I(y, v)$ . So  $d(u, w_1) = m - 1$  and  $d(u, w_1) = m$  are not possible, since it is a contradiction to the choice of  $y_1$  that  $d(u, y_1)$  minimum. Therefore  $\ell + t = m + 2$ . Hence the claim. That is  $w \to y_1 \to y$  is a shortest w, y- path. So can replace  $w_1$  with  $y_1$ . Consider the vertices  $u, y, y_1 y'$ , it is clear that d(u, y) = d(u, y') = m and d(y, y') = 2. If d satisfies (QC), then there exist a z which is adjacent to y and y' with d(u, z) = m - 1. Then  $z \in I(y, w)$ and since  $I(y, w) \subseteq I(y, v)$  implies that  $z \in I(y, v)$ . Also  $z \in I(u, y)$  and hence  $z \in I(u, y) \cap I(x, v)$ . So the vertices u, x, y, satisfies (J0'), contradiction. Hence d does not satisfy (QC) on u, y, y' and hence G is not diamond-weakly modular.

That is, we have proved that if I doesn't satisfy axiom (J0'), then G doesn't satisfy (TDC) or (QC) and hence G is not a diamond-weakly modular graph.

From Proposition 2, Theorem 3, Theorem 4, Theorem 5 and we have the following theorem.

**Theorem 6.** Let R be a function from  $V \times V$  to  $2^V$ , where V is a non empty set. Then R satisfies the axioms (t1), (t2), (t3), (b3), (J0') and (ta) if and only if  $G_R$ is a diamond-weakly modular graph and R coincides with the interval function  $I_{G_R}$  of  $G_R$ .

The examples below establishes the independence of the axioms (t1), (t2), (t3),(b3), (J0') and (ta).

Example 2 ((t2), (t3), (b3), (J0') and (ta) but not (t1)).

Let  $V = \{u, v, x, y\}$  and define a transit function R on V as R(u, v) = R(v, u) = $\{u\}, R(u,x) = \{u,x\} R(u,y) = \{u,x,y\}, R(v,x) = \{v,x\}, R(v,y) = \{v,x,y\}$  $R(x,y) = \{x,y\}, R(a,a) = \{a\}$  and R(a,b) = R(b,a) for all  $a, b \in V$ . We can see that R satisfies  $(t_2)$ ,  $(t_3)$ ,  $(b_3)$ ,  $(J_0)$  and  $(t_a)$ . But R does not satisfy axiom (t1), since  $v \notin R(u, v)$  and  $v \notin R(v, u)$ .

Example 3 ((t1), (t3), (b3), (J0') and (ta) but not (t2)).Let  $V = \{u, v, x, y\}$  and define a transit function R on V as follows: R(u, v) = $\{u, v\} = R(v, u), R(u, x) = \{u, v, x\}, R(x, u) = \{x, u\}, R(u, y) = \{u, v, x, y\} = \{u, v, x, y\}$  $R(y,u), R(v,x) = \{v,x\} = R(x,v), R(v,y) = \{v,x,y\}, R(y,v) = \{y,v\}, R(x,y) = \{v,v\}, R(x,y) = \{v,v\}, R(x,y) = \{v,v\}, R(y,v) =$  $\{x,y\} = R(y,x), R(a,a) = \{a\}$  for all  $a \in V$ . We can see that R satisfies (t1),(t3), (b3), (J0') and (ta). But  $R(u, v) \neq R(v, u)$ . Therefore R does not satisfy the (t2) axiom.

15

Example 4 ((t1), (t2), (b3), (J0') and (ta) but not (t3)). Let  $V = \{u, v, x, y\}$  and define a transit function R on V as follows:  $R(u, u) = \{u, v\}, R(u, v) = \{u, v\}, R(u, x) = \{u, x\}, R(u, y) = \{u, y\}, R(v, x) = \{v, u, x\}, R(v, y) = \{v, u, y\} R(x, y) = \{x, y\}, R(a, a) = \{a\}$  and R(a, b) = R(b, a) for all  $a, b \in V$ . We can see that R satisfies (t1), (t2), (b3), (J0') and (ta). But  $v \in R(u, u)$ . Therefore R does not satisfy the (t3) axiom.

Example 5 ((t1), (t2), (t3), (b3), (ta) but not (J0')).

Let  $V = \{u, v, w, x, y, z\}$  and define a transit function R on V as follows:  $R(u, v) = \{u, w, v, z\}, R(x, v) = \{x, w, y, v\}, R(x, z) = \{x, y, u, z\} R(w, z) = \{w, u, v, z\}, R(w, y) = \{w, x, v, y\}, \text{ and } R(a, a) = \{a\} \text{ and } R(a, b) = R(b, a)$ for all  $a, b \in V$ . We can see that R satisfies (t1), (t2), (t3), (b3), (ta). But  $x \in R(u, y), y \in R(x, v), R(u, y) \cap R(x, v) \subset \{u, x, y, v\}$  and  $x \notin R(u, v)$  Therefore R does not satisfy the (J0') axiom.

Example 6 ((t1), (t2), (t3), (b3), (J0') but not (ta)). Let  $V = \{u, v, w, x, y\}$  and define a transit function R on V as follows:  $R(u, w) = \{u, x, y, w\}$ ,  $R(v, x) = \{v, y, w, x\}$  and  $R(a, a) = \{a\}$  and R(a, b) = R(b, a) for all  $a, b \in V$ . We can see that R satisfies (t1), (t2), (t3), (b3). Also  $w \in R(v, x), x \in R(w, u)$  and  $R(v, x) \cap R(w, u) = \{y, x, w\} \not\subset \{v, x, w, u\}$ . That is R satisfy (J0'). But  $R(u, v) \cap R(u, w) = \{u\}$ ,  $R(u, v) \cap R(v, w) = \{v\}$ ,  $R(u, w) \cap R(v, w) = \{w\}$  and  $R(u, v) = \{u, v\}$ ,  $R(v, w) = \{v, w\}$  and  $R(u, w) = \{u, x, y, w\}$ . Therefore R does not satisfy the (ta) axiom.

Example 7 ((t1), (t2), (t3), (ta), (J0') but not (b3)). Let  $V = \{u, v, w, x, y\}$  and define a transit function R on V as follows:  $R(u, y) = \{u, x, y, w\}, R(x, v) = \{x, y, v, w\}, R(u, v) = V$  and  $R(a, a) = \{a\}$  and R(a, b) = R(b, a) for all  $a, b \in V$ . We can see that R satisfies (t1), (t2), (t3), (J0') and (ta). But  $y \in R(u, v), w \in R(u, y)$ , and  $y \notin R(w, v)$ . Therefore R does not satisfy the (b3) axiom.

# 5 Interval function of bridged graphs and weakly bridged graphs

In this section, we characterize the interval function of bridged graphs using axioms (J0') and (br) and weakly bridged graphs by (J0') and (br') on an arbitrary transit function. First, we prove the following Theorem characterizing the interval function I of the bridged graph.

**Theorem 7.** Let G be a graph. The interval function  $I_G$  satisfies the axioms (J0') and (br) if and only if G is a bridged graph.

*Proof.* First assume that I satisfy axioms (J0') and (br) on G. We have to show that G is a bridged graph. By Theorem 5, the interval function I on a graph G satisfies (J0') if and only if G is a diamond-weakly modular graph. That is, if I satisfy axiom (J0'), then G is a diamond-weakly modular graph which is a

subclass of weakly modular graphs. According to the result by Chepoi et al. [10], bridged graphs are exactly weakly modular graphs that do not contain induced  $C_4$  and  $C_5$ . Now it is enough to show that I satisfy (br), then G is  $(C_4, C_5)$ free. If G contains an induced  $C_5$  with consecutive vertices as u, x, y, v, z, it is clear that I does not satisfy (br) on  $C_5$ , since  $I(x, y) = \{x, y\}, I(x, u) =$  $\{x, u\}, I(v, y) = \{v, y\}$  and  $z \in I(u, v)$  and  $I(x, z) \neq \{x, z\}$  and  $I(y, z) \neq \{y, z\}$ . Similarly, if G contains an induced  $C_4$  with consecutive vertices as u, x = y, v, z, it is clear that I does not satisfy (br) on  $C_4$ .

Conversely, suppose that G is a bridged graph. We have to show that Isatisfy axioms (J0') and (br). Since bridged graphs are diamond-weakly modular graphs, I satisfy axiom (J0') on G. It remains to show that I satisfy axiom (br). Suppose not. That is  $I(x,y) = \{x,y\}, I(x,u) = \{x,u\}, I(v,y) = \{v,y\}$  and  $z \in I(u, v)$  and  $I(x, z) \neq \{x, z\}$  and  $I(y, z) \neq \{y, z\}$ . Let P: uxyv be the u, vpath containing x and y. Since  $z \in I(u, v)$ ,  $I(x, z) \neq \{x, z\}$  and  $I(y, z) \neq \{y, z\}$ , there exist a u, v- shortest path containing z which is different from the path P. Then possibility of d(u, v) are 3 and 2. When d(u, v) = 3, then z is either adjacent to u or v. Without loss of generality, we may assume z is adjacent to u. Let z' be the neighbor of z in the u, v- shortest path. Then u, x, y, v, z', z will form a cycle of length 6. If  $z/y \in E(G)$  and  $z/x \notin E(G)$ , the vertices u, x, y, z/zwill induce a cycle of length 5. If both  $z'y, z'x \in E(G)$ , then the vertices u, x, z', zwill induce a  $C_4$ . When d(u, v) = 2, then z is adjacent to both u and v. Then the vertices u, x, y, z, v will induce a cycle of length 5. If  $uy \in E(G)$ , the vertices u, y, z, v will induce a cycle of length 4 and if  $xv \in E(G)$ , the vertices u, x, v, zwill induce a cycle of length 4. When x = y, the vertices u, x = y, z, v will induce a 4-cycle. In all cases, we get a contradiction to the assumption that G is a bridged graph.

From Proposition 2, Theorem 3, Theorem 4, and Theorem 7, we have the following theorem.

**Theorem 8.** Let R be a function from  $V \times V$  to  $2^V$ , where V is a non empty set. Then R satisfies the axioms (t1), (t2), (t3), (J0'), (b3), (ta) and (br) if and only if  $G_R$  is a bridged graph and R coincides with the interval function  $I_{G_R}$ .

The examples below establish the independence of the axioms (t1), (t2), (t3), (b3), (br), (J0') and (ta). The function defined in Example 2 satisfy axioms (t2), (t3), (b3), (br), (J0') and (ta) but does not satisfy axiom (t1). The function defined in Example 3 satisfy axioms (t1), (t3), (b3), (br), (J0') and (ta) but does not satisfy axiom (t2) and (ta) but does not satisfy axiom (t2). The function defined in Example 4 satisfy axioms (t1), (t2), (b3), (br), (J0') and (ta) but does not satisfy axiom (t3).

Example 8 ((t1), (t2), (t3), (ta), (br), (b3) but not (J0')).

Let  $V = V(C_8)$  and R = I on  $C_8$ . The consecutive vertices of  $C_8$  be  $v_1, v_2, ..., v_8$ . Then R satisfies axioms (t1), (t2), (t3), (br), (ta) and (b3). Assume  $u = v_1, x = v_3, y = v_4$  and  $v = v_6$ . Then  $x \in R(u, y), y \in R(x, v), R(u, y) \cap R(x, v) \subset \{u, x, y, v\}$  and  $x \notin R(u, v)$ . Therefore R does not satisfy the (J0') axiom.

Example 9 ((t1), (t2), (t3), (br), (b3), (J0') but not (ta)). Let  $V = \{u, v, w, x, y\}$  and define a transit function R on V as follows: R(u, w) =  $\{u, y, w\}, R(v, x) = \{v, y, w, x\}, R(u, x) = \{u, y, x\} \text{ and } R(a, a) = \{a\} \text{ and } R(a, b) = R(b, a) \text{ for all } a, b \in V.$  We can see that R satisfies (t1), (t2), (t3), (b3), (br) and (J0'). But  $R(u, v) \cap R(u, w) = \{u\}, R(u, v) \cap R(v, w) = \{v\}, R(u, w) \cap R(v, w) = \{w\} \text{ and } R(u, v) = \{u, v\}, R(v, w) = \{v, w\} \text{ and } R(u, w) = \{u, y, w\}.$  Therefore R does not satisfy the (ta) axiom.

Example 10 ((t1), (t2), (t3), (ta), (br), (J0') but not (b3)).

Let  $V = \{u, v, w, x, y\}$  and define a transit function R on V as follows:  $R(u, y) = \{u, x, y, w\}, R(x, v) = \{x, y, v, w\}, R(u, v) = V$  and  $R(a, a) = \{a\}$  and R(a, b) = R(b, a) for all  $a, b \in V$ . We can see that R satisfies (t1), (t2), (t3), (J0'), (br) and (ta). But  $y \in R(u, v), w \in R(u, y)$ , and  $y \notin R(w, v)$ . Therefore R does not satisfy the (b3) axiom.

Example 11 ((t1), (t2), (t3), (ta), (b3), (J0') but not (br)). Let  $V = \{u, v, x, y, z, w\}$  and define a transit function R on V as follows:  $R(u, y) = \{u, x, z, y\}, R(u, v) = \{u, v, z, w\}R(x, v) = \{x, y, z, v\}, R(x, w) = \{x, w, z, u\}, R(w, y) = \{w, y, z, v\}$  and  $R(a, a) = \{a\}$  and R(a, b) = R(b, a) for all  $a, b \in V$ . We can see that R satisfies (t1), (t2), (t3), (J0'), (ta) and (b3). Rdoes not satisfy the (br), since  $I(x, y) = \{x, y\}, I(x, u) = \{x, u\}, I(v, y) = \{v, y\},$  $z \in I(u, v)$  and  $I(x, z) \neq \{x, z\}$  and  $I(y, z) \neq \{y, z\}$ .

For the characterization of the interval function of weakly bridged graphs, we need the Lemma 2. From the definition of axioms (br) and (br'), we can see that axiom (br') is easily followed from axiom (br) by taking x = y.

The axioms (br') and (J0') characterize the interval function of weakly bridged graphs. We have the following theorems in which the proof is similar to that of theorems, Theorem 7 and 8.

**Theorem 9.** Let G be a graph. The interval function  $I_G$  satisfies the axioms (J0') and (br') if and only if G is a weakly bridged graph.

**Theorem 10.** Let R be a function from  $V \times V$  to  $2^V$ , where V is a non empty set. Then R satisfies the axioms (t1), (t2), (t3), (J0'), (b3), (ta) and (br') if and only if  $G_R$  is a weakly bridged graph and R coincides with the interval function  $I_{G_R}$ .

The independence of the axioms (t1), (t2), (t3), (b3), (br'), (J0') and (ta) will easily follow from Example 2, Example 3, Example 4, Example 8, Example 9, and Example 10 except the following one.

Example 12 ((t1), (t2), (t3), (ta), (b3), (J0') but not (br')). Let  $V = \{u, v, x, z\}$ and define a transit function R on V as follows: R(u, v) = R(x, z) = V and  $R(a, a) = \{a\}$  and R(a, b) = R(b, a) for all  $a, b \in V$ . We can see that R satisfies (t1), (t2), (t3), (J0'), (ta) and (b3). R does not satisfy the (br'), since I(x, u) = $\{x, u\}, I(v, x) = \{v, x\}, z \in I(u, v)$  and  $I(x, z) \neq \{x, z\}$ .

**Concluding Remarks:** In this paper we introduced a subclass of weakly modular graphs which satisfy a stronger version (diamond condition) of the triangle condition of the weakly modular graphs and name it as the diamond-weakly modular graphs. We have proved that the class of diamond-weakly modular graphs contain the family of bridged graphs and weakly bridged graphs. Further we have characterized this class of graphs in terms of the axioms of its interval function. Nevertheless, a purely graph theoretic characterization for diamondweakly modular graphs remains as an open problem.

As a byproduct, we have obtained that the diamond-weakly modular graphs, bridged graphs and weakly bridged graphs are definable in FOLB.

From the Mulder- Nebeský Theorem (Theorem 3), the following definition of *graphic interval structure* is derived by Chalopine at al. in [6]. We quote the following definitions and conventions regarding the FOLB definability from [6].

A graphic interval structure is a  $\sigma$ -structure (V, B) where V is a finite set and B is a ternary predicate ( a ternary relation) on V satisfying the following axioms:

- (IB1)  $\forall u \forall v B(u, u, v)$
- (IB2)  $\forall u \forall v \forall x B(u, x, v) \implies B(v, x, u)$
- (IB3)  $\forall u \forall x B(u, x, u) \implies x = u$
- (IB4)  $\forall u \forall v \forall w \forall x B(u, w, v) \land B(u, x, w) \implies B(u, x, v)$
- (IB5)  $\forall u \forall v \forall w \forall x B(u, v, x) \land B(u, w, x) \land B(u, v, w) \implies B(v, w, x)$
- $(\text{IB6}) \quad \forall u \forall u' \forall v \forall v' B(u, u', u') \land B(v, v', v') \land B(u', u, v') \land B(u, u', v) \land B(u, v', v) \implies B(u', v, v')$
- (IB7)  $\forall u \forall u' \forall v \forall v' B(u, u', u') \land B(v, v', v') \land B(u, u', v) \land \neg B(u, v', v) \land \neg B(u', v, v') \implies B(u, u', v').$

In the class C of  $\sigma$ -structures (V, B) satisfying axioms (IB1)–(IB7), a query Q on C is definable in first order logic with betweeness (*FOLB-definable*), if it can be defined by a first order formula F over (V, B).

From the Theorems 5, 7 and 9, we can easily derive the following FOLB queries for diamond-weakly modular graphs, bridged graphs and weakly bridged graphs.

 $\begin{array}{l} diamond-weakly\ modular \equiv \forall u, x, y, v,\ B(u, x, y) \land B(x, y, v) \land [(B(u, z, y) \land B(x, z, v) \implies (z = u \lor z = x \lor z = y) \land z \neq v) \lor ((B(u, z, y) \land B(x, z, v) \implies (z = u \lor z = x \lor z = v) \land z \neq y) \lor ((B(u, z, y) \land B(x, z, v) \implies (z = u \lor z = v \lor z = y) \land z \neq x) \lor ((B(u, z, y) \land B(x, z, v) \implies (z = v \lor z = y) \land z \neq u)] \implies B(u, x, v). \end{array}$ 

 $\begin{array}{l} bridged \equiv diamond-weakly \ modular \ \land \forall u, x, y, v, z, w[((B(x, w, y) \implies w = x \lor w = y) \land (B(x, w, u) \implies w = x \lor w = u) \land (B(v, w, y) \implies w = v \lor w = y) \land B(u, z, v)) \implies ((B(x, w, z) \implies w = x \lor w = z) \lor (B(z, w, y) \implies w = z \lor w = y))] \end{array}$ 

 $\begin{array}{l} weakly \ bridged \equiv diamond-weakly \ modular \ \land \forall u, x, v, z, w \ [((B(u, w, x) \implies w = u \lor w = x) \land (B(x, w, v) \implies w = x \lor w = v) \land B(u, z, v)) \implies \\ ((B(x, w, z) \implies w = x \lor w = z))] \end{array}$ 

#### References

- H.-J. Bandelt, M. van de Vel, and E. Verheul, Modular interval spaces, Math. Nachr. 163 (1993), 177–201.
- H.-J.Bandelt, H. M. Mulder, and E.Wilkeit, Quasi-median graphs and algebras, J. Graph Theory 18 (1994), 681–703.
- H.J. Bandelt, V. Chepoi, A Helly theorem in weakly modular space, Discrete Mathematics, (1996),160, 25-39 Pages 25-39,.
- H.-J. Bandelt and V. Chepoi, Metric graph theory and geometry: a survey, Surveys on Discrete and Computational Geometry: Twenty Years Later, J.E. Goodman, J. Pach, and R. Pollack (eds), Contemp. Math., 453 (2008), pp. 49–86.
- 5. J. Chalopine, V. Chepoi, and H. Hirai, and D. Osajda, Weakly modular graphs and nonpositive curvature, Memoirs of AMS, 268 (2020), no. 1309.
- J. Chalopin, M. Changat, V. Chepoi, and J. Jacob, First-Order Logic Axiomatization of Metric Graph Theory, Theoretical Computer Science. 2024 Feb 21:114460.
- M. Changat, J. Mathew, H.M. Mulder The induced path function, monotonicity and betweenness Disc. Appl. Math. 158(5) 2010 426–433
- M. Changat, A.K. lakshmikuttyamma, J. Mathew, I. Peterin, P.G. Narasimha-Shenoi, G. Seethakuttyamma, S. Špacapan A forbidden subgraph characterization of some graph classes using betweenness axioms Disc. Math. 313 2013 951–958
- M. Changat, L.K.Sheela, P.G.Narasimha-Shenoi The Interval function, Induced path function, axiomatic characterizations, Ptolemaic and chordal graphs, 2022 (submitted)
- V. Chepoi, Classifying graphs by metric triangles, Metody Diskretnogo Analiza 49 (1989), 75–93 (Russian).
- M. Farber, Bridged graphs and geodesic convexity, Discrete Mathematics, 1987, Volume 66, Issue 3, 249-257, ISSN 0012-365X
- M.Farber, R.E Jamison, On local convexity in graphs, Discrete Mathematics, (1987), 66, 3, 231-247.
- J. Hedlíková, Ternary spaces, media, and Chebyshev sets, Czechoslovak Math. J. 33 (1983), 373–389.
- 14. H.M. Mulder The Interval function of a GraphMC Tract 132 Mathematisch Centrum, Amsterdam 1980
- H.M. Mulder, Transit functions on graphs (and posets) Convexity in Discrete Structures M. Changat, S. Klavžar, H.M. Mulder, A. Vijayakumar, eds. Lecture Notes Ser. 5 Ramanujan Math. Soc. 2008 117–130
- H.M. Mulder, L. Nebeský Axiomatic characterization of the interval function of a graph European J. Combin 30 2009 1172–1185
- 17. L. Nebeský A characterization of the interval function of a connected graph Czech. Math. J. 44 1994 173–178
- L. Nebeský A characterization of the set of all shortest paths in a connected graph Mathematica Bohemica 119.1 1994 15–20
- 19. L. Nebeský A characterization of geodetic graphs Czech. Math. J. 45.3 1995 491–493
- L. Nebeský Characterizing the interval function of a connected graph Math. Bohem. 123.2 1998 137–144
- 21. L. Nebeský A new proof of a characterization of the set of all geodesics in a connected graphCzech. Math. J.48.41998809–813
- L. Nebeský Characterization of the interval function of a (finite or infinite) connected graphCzech. Math. J. 51 2001 635–642

- 23. M. Sholander Trees, lattices, order, and betweenness<br/>Proc. Amer. Math. Soc. 31952  $369{-}381$
- 24. M. van de Vel, Theory of Convex Structures, Elsevier, Amsterdam, 1993.