# On 3-dimensional MRD codes of type $\langle x^{q^t}, x + \delta x^{q^{2t}}, G(x) \rangle$

Daniele Bartoli<sup>\*</sup>, and Francesco Ghiandoni<sup>†</sup>

March 5, 2024

#### Abstract

In this work we present results on the classification of  $\mathbb{F}_{q^n}$ -linear MRD codes of dimension three. In particular, using connections with certain algebraic varieties over finite fields, we provide non-existence results for MRD codes  $\mathcal{C} = \langle x^{q^t}, F(x), G(x) \rangle \subseteq \mathcal{L}_{n,q}$  of exceptional type, i.e. such that  $\mathcal{C}$  is MRD over infinite many extensions of the field  $\mathbb{F}_{q^n}$ . These results partially address a conjecture of Bartoli, Zini and Zullo in 2023.

### 1 Introduction

Let q be a prime power, n be a positive integer, and denote by  $\mathbb{F}_{q^n}$  the finite field with  $q^n$  elements and by  $\mathbb{P}^N(\mathbb{K})$  (resp.  $\mathbb{A}^N(\mathbb{K})$ ) the N-dimensional projective (resp. affine) space over the field  $\mathbb{K}$ . Let  $\mathcal{L}_{n,q} = \{\sum_{i=0}^{n-1} a_i X^{q^i} : a_i \in \mathbb{F}_{q^n}\}$  denote the  $\mathbb{F}_q$ -algebra of the  $\mathbb{F}_q$ -linearized polynomials (or q-

Let  $\mathcal{L}_{n,q} = \{\sum_{i=0}^{n-1} a_i X^q : a_i \in \mathbb{F}_{q^n}\}$  denote the  $\mathbb{F}_q$ -algebra of the  $\mathbb{F}_q$ -linearized polynomials (or q-polynomials) of q-degree smaller than n. For any  $f(x) = \sum_{i=0}^{n-1} a_i X^{q^i}$ , we define  $\deg_q(f(X)) = \max\{i : a_i \neq 0\}$  and  $\min \deg_q(f(X)) = \min\{i : a_i \neq 0\}$ . We identify a polynomial  $g(X) \in \mathcal{L}_{n,q}$  with the  $\mathbb{F}_q$ -linear map  $x \mapsto g(x)$  over  $\mathbb{F}_{q^n}$ ; in this way,  $\mathbb{F}_q$ -linearized polynomials over  $\mathbb{F}_{q^n}$  are in one-to-one correspondence with  $\mathbb{F}_q$ -linear maps over  $\mathbb{F}_{q^n}$ .

The rank metric on the  $\mathbb{F}_q$ -vector space  $\mathbb{F}_q^{m \times n}$  is defined by

$$d(A,B) := \operatorname{rank}(A-B) \text{ for } A, B \in \mathbb{F}_{a}^{m \times n}$$

We call a subset of  $\mathbb{F}_q^{m \times n}$  equipped with the rank metric a *rank-metric code*. For a rank-metric code  $\mathcal{C}$  containing at last two elements, its *minimum distance* is given by

$$d(\mathcal{C}) := \min_{A, B \in \mathcal{C}, A \neq B} d(A, B).$$

When  $\mathcal{C}$  is an  $\mathbb{F}_q$ -subspace of  $\mathbb{F}_q^{m \times n}$ , we say that  $\mathcal{C}$  is an  $\mathbb{F}_q$ -linear code of dimension  $\dim_{\mathbb{F}_q}(\mathcal{C})$ . Under the assumption that  $m \leq n$ , it is well known (and easily verified) that every rank-metric code  $\mathcal{C}$  in  $\mathbb{F}_q^{m \times n}$  with minimum distance d satisfies the Singleton-like bound

$$|\mathcal{C}| < q^{n(m-d+1)}$$

<sup>\*</sup>Dipartimento di Matematica e Informatica, Università degli Studi di Perugia, Perugia, Italy. daniele.bartoli@unipg.it

<sup>&</sup>lt;sup>†</sup>Dipartimento di Matematica e Informatica "Ulisse Dini", Università degli studi di Firenze, Firenze, Italy, francesco.ghiandoni@unifi.it

In case of equality, C is called a *maximum rank-metric* code, or MRD code for short. MRD codes have been studied since the 1970s by Delsarte [16] and Gabidulin [17] and have seen much interest in recent years due to an important application in network coding and cryptography [26].

From a different perspective, rank-metric codes can also be seen as sets of (restrictions of)  $\mathbb{F}_q$ -linear homomorphisms from  $(\mathbb{F}_{q^n})^m$  to  $\mathbb{F}_{q^n}$  equipped with the rank metric; see [4, Sections 2.2 and 2.3]. In this case, it is evident that multivariate linearized polynomials can be seen as the natural algebraic counterpart of rank-metric codes. In particular, when m = n, a rank metric code  $\mathcal{C}$  can be seen as set of  $\mathbb{F}_q$ -linear endomorphisms of  $\mathbb{F}_{q^n}$ , i.e.  $\mathcal{C} \subseteq \mathcal{L}_{n,q}$ . From now on we will consider m = n and d < n. In the case of univariate linearized polynomials, Sheekey pointed out in [30] the following connection between  $\mathbb{F}_{q^n}$ -linear MRD codes and the so called *scattered* polynomials:  $\mathcal{C}_f = \langle x, f(x) \rangle_{\mathbb{F}_{q^n}}$  is an MRD code with  $\dim_{\mathbb{F}_{q^n}}(\mathcal{C}) = 2$ if and only if

$$\dim_{\mathbb{F}_a} \ker(f(x) - mx) \le 1$$

for every  $m \in \mathbb{F}_{q^n}$ . The concept of scattered polynomial introduced in [30] has been slightly generalized in [7].

**Definition 1.1.** [7, 30] An  $\mathbb{F}_q$ -linearized polynomial  $f(X) \in \mathbb{F}_{q^n}[X]$  is called a scattered polynomial of index  $t \in \{0, \ldots, n-1\}$  if

$$\lim_{\mathbb{F}_q} \ker(f(x) - mx^{q^{\iota}}) \le 1,$$

for every  $m \in \mathbb{F}_{q^n}$ . Also, a scattered polynomial of index t is **exceptional** if it is scattered of index t over infinitely many extensions  $\mathbb{F}_{q^{nm}}$  of  $\mathbb{F}_{q^n}$ .

While several families of scattered polynomials have been constructed in recent years [6, 10, 12, 15, 18, 20, 27, 30, 32], only two families of exceptional ones are known:

- (Ps)  $f(x) = x^{q^t}$  of index 0, with gcd(t, n) = 1 (polynomials of so-called pseudoregulus type);
- (LP)  $f(x) = x + \delta x^{q^{2t}}$  of index t, with gcd(t, n) = 1 and  $N_{q^n/q}(\delta) = \delta^{(q^n-1)/(q-1)} \neq 1$  (so called LP polynomials).

From a coding theory point of view, if f is exceptional scattered of index t, the corresponding rank distance code  $C_{f,t}^m = \langle x^{q^t}, f(x) \rangle_{\mathbb{F}_{q^{mn}}} \subseteq \mathcal{L}_{nm,q}$  turns out to be an MRD code for infinitely many m; codes of this kind are called exceptional  $\mathbb{F}_{q^n}$ -linear MRD codes (see [9]). Moreover, in [4] the authors introduce the notions of h-scattered sequences and exceptional h-scattered sequences which parameterize exceptional MRD codes. Only two families of exceptional  $\mathbb{F}_{q^n}$ -linear MRD codes are known so far:

(G) 
$$\mathcal{G}_{r,s} = \langle x, x^{q^s}, \dots, x^{q^{s(r-1)}} \rangle_{\mathbb{F}_{q^n}}$$
, with  $gcd(s, n) = 1$ , see [16, 17];

(T) 
$$\mathcal{H}_{r,s}(\delta) = \langle x^{q^s}, \dots, x^{q^{s(r-1)}}, x + \delta x^{q^{sr}} \rangle_{\mathbb{F}_{q^n}}$$
, with  $gcd(s, n) = 1$  and  $N_{q^n/q}(\delta) \neq (-1)^{nr}$  see [20, 30].

The first family is known as generalized Gabidulin codes and the second one as generalized twisted Gabidulin codes.

In [8] it has been shown that the only exceptional  $\mathbb{F}_{q^n}$ -linear MRD codes spanned by monomials are the codes (G), in connection with so-called Moore exponent sets, while in [9] the authors investigated exceptional  $\mathbb{F}_{q^n}$ -linear MRD codes not generated by monomials and proved that an exceptional *r*-dimensional

 $\mathbb{F}_{q^n}$ -linear MRD code contains an exceptional scattered polynomial (see Theorem 4.2).

Motivated by this last necessary condition on MRD codes of exceptional type, we considered codes of type  $\mathcal{C} = \langle x^{q^t}, F(x), G(x) \rangle_{\mathbb{F}_{q^n}}$  and we address a conjecture in [9] for r = 3 and F(x) a LP polynomial. The techniques that we use to prove our results are from algebraic geometry over finite fields. This approach has already proved successful in the investigation of families of functions with many applications in coding theory and cryptography; see for instance [1–3, 5, 22, 24, 25, 31]. Our main result can be summarized as follows (see Theorem 6.4).

**Main Theorem.** If  $(t,q) \notin \{(1,3); (1,4); (1,5); (2,3); (2,4); (2,5), (4,3)\}$ , then there are no exceptional 3-dimensional  $\mathbb{F}_{q^n}$ -linear MRD codes of type  $\mathcal{C} = \langle x^{q^t}, x + \delta x^{q^{2t}}, G(x) \rangle \subseteq \mathcal{L}_{n,q}$ , with  $\deg_q(G(x)) > 2t$ .

### 2 Preliminaries on algebraic curves and varieties

Let  $F(X,Y) \in \mathbb{K}[X,Y]$ ,  $\mathbb{K}$  a field, be a polynomial defining an affine plane curve  $\mathcal{C} : F(X,Y) = 0$ . A plane curve is absolutely irreducible if there are no non-trivial factorizations of its defining polynomial F(X,Y)in  $\overline{\mathbb{K}}[X,Y]$ , where  $\overline{\mathbb{K}}$  is the algebraic closure of  $\mathbb{K}$ . If  $F(X,Y) = \prod_i F^{(i)}(X,Y)$ , with  $F^{(i)}(X,Y) \in \overline{\mathbb{K}}[X,Y]$ of positive degree, then  $\mathcal{C}_i : F^{(i)}(X,Y) = 0$  are called components of  $\mathcal{C}$ . A component is  $\mathbb{F}_q$ -rational if it is fixed by the Frobenius morphism  $\varphi$  or equivalently  $\lambda F^{(i)}(X,Y) \in \mathbb{K}[X,Y]$  for some  $\lambda \in \overline{\mathbb{K}}$ .

Let  $P = (u, v) \in \mathbb{A}^2(\mathbb{K})$  be a point in the plane, and write

$$F(X + u, Y + v) = F_0(X, Y) + F_1(X, Y) + F_2(X, Y) + \cdots$$

where  $F_i$  is either zero or homogeneous of degree *i*. The multiplicity of  $P \in C$ , written as  $m_P(\mathcal{C})$  or  $m_P(F)$ , is the smallest integer *m* such that  $F_m \neq 0$  and  $F_i = 0$  for i < m;  $F_m = 0$  is the tangent cone of  $\mathcal{C}$  at *P*. A linear component of the tangent cone is called a tangent of  $\mathcal{C}$  at *P*. The point *P* is on the curve  $\mathcal{C}$  if and only if  $m_P(\mathcal{C}) \geq 1$ . If *P* is on  $\mathcal{C}$ , then *P* is a simple point of  $\mathcal{C}$  if  $m_P(\mathcal{C}) = 1$ , otherwise *P* is a singular point of  $\mathcal{C}$ . It is possible to define in a similar way the multiplicity of an ideal point of  $\mathcal{C}$ , that is a point of the curve lying on the line at infinity. We denote by  $Sing(\mathcal{C})$  the set of singular points of the curve  $\mathcal{C}$ .

Given two plane curves  $\mathcal{A}$  and  $\mathcal{B}$  and a point P on the plane, the *intersection number* (or *intersection multiplicity*)  $I(P, \mathcal{A} \cap \mathcal{B})$  of  $\mathcal{A}$  and  $\mathcal{B}$  at the point P can be defined by seven axioms. We do not include its precise and long definition here. For more details, we refer to [21] and [23] where the intersection number is defined equivalently in terms of local rings and in terms of resultants, respectively.

For a given plane curve  $\mathcal{C}$  and a point  $P \in \mathcal{C}$ , we denote by  $I_{P,max}(\mathcal{C})$  the maximum possible intersection multiplicity of two components of  $\mathcal{C}$  at  $P \in Sing(\mathcal{C})$ . We list here two useful results in this direction.

**Lemma 2.1.** [21, Section 3.3] [29, Lemma 4.3] [7, Lemma 2.5] Let q be a prime power and  $F(X,Y) \in \mathbb{F}_q[X,Y]$ . Let  $P = (\alpha, \beta) \in \mathbb{F}_q^2$  and write

$$F(X + \alpha, Y + \beta) = F_m(X, Y) + F_{m+1}(X, Y) + \dots,$$

where  $F_i \in \mathbb{F}_q[X,Y]$  is zero or homogeneous of degree *i* and  $F_m \neq 0$ . The following properties hold.

- (i) If  $F_m(X,Y)$  is separable, then  $I_{P,max}(\mathcal{C}) \leq |m^2/2|$ .
- (ii) Suppose that  $F_m = L^m$  with L a linear form;

- if  $L \nmid F_{m+1}$  then  $I_{P,max}(\mathcal{C}) = 0$ , - if  $L^2 \nmid F_{m+1}$  then  $I_{P,max}(\mathcal{C}) \leq m$ .

**Criterion 2.2.** [25, Lemma 11] Let C: h(X,Y) = 0 be a curve of degree n defined over  $\mathbb{F}_q$ . If  $\sum_{P \in Sing(h)} I_{P,max}(C) < \frac{2}{9} \deg^2(h)$  then C possesses at least one absolutely irreducible component

defined over  $\mathbb{F}_q$ 

Consider the set  $\overline{\mathbb{F}_q}[[t]]$  of the formal power series on t. Let  $(x_0, y_0) \in \overline{\mathbb{F}_q}^2$  be an affine point of  $\mathcal{C}$ : F(X,Y) = 0. A branch of center  $(x_0, y_0)$  of  $\mathcal{C}$  is a point  $(x(t), y(t)) \in (\overline{\mathbb{F}_q}[[t]])^2$  such that F(x(t), y(t)) = 0, where

$$\begin{aligned} x(t) &= x_0 + u_1 t + u_2 t^2 + \dots, \\ y(t) &= y_0 + v_1 t + v_2 t^2 + \dots. \end{aligned}$$

See [23, Chapter 4] for more details on branches. There exists a unique branch centered at a simple point of  $\mathcal{C}$ . If there exists only a branch centered in a point  $P \in \mathcal{C}$  then  $I_{P,max}(\mathcal{C}) = 0$ .

**Remark 2.3.** In order to determine branches centered at singular points of a curve  $\mathcal{C}$  we make use of quadratic transformations; see [23, Section 4]. Consider a curve C defined by

$$F(X,Y) = F_r(X,Y) + F_{r+1}(X,Y) + \dots = 0,$$

where each  $F_i(X,Y)$  is homogeneous in X and Y and of degree i and  $F_r \neq 0$ . First, we can suppose that the singular point under examination is the origin O = (0,0) and that X = 0 is not a tangent line at O. Let r be its multiplicity. The geometric transform of a curve C is the curve C' given by F'(X,Y) = $F(X, XY)/X^r$ . (If Y = 0 is not a tangent line at O then we can also consider  $\mathcal{C}'$  defined by F'(X, Y) = $F(XY,Y)/Y^r$ ). By [23, Theorem 4.44], there exists a bijection between the branches of C centered at the origin and the branches of  $\mathcal{C}'$  centered at an affine point on X = 0. In our proofs we will perform chains of local quadratic transformations until the total number of branches is determined. In particular, if the tangent cone  $F_r(X,Y)$  at O splits into non-repeated linear factors (over the algebraic closure), then there are precisely r distinct branches centered at O. In fact, distinct linear factors of  $F_r(X,Y)$  correspond to distinct affine points of  $\mathcal{C}'$  on X = 0.

An algebraic hypersurface is an algebraic variety that may be defined by a single polynomial equation. An algebraic hypersurface defined over a field  $\mathbb{K}$  is *absolutely irreducible* if the associated polynomial is irreducible over every algebraic extension of K. An absolutely irreducible K-rational component of a hypersurface  $\mathcal{V}$ , defined by the polynomial F, is simply an absolutely irreducible hypersurface which is associated to a non-costant factor of F defined over  $\mathbb{K}$ .

**Lemma 2.4.** [1, Lemma 2.1] Let  $X, H \subseteq \mathbb{P}^{N}(\mathbb{F}_{q})$  be projective hypersurfaces. If  $X \cap H$  has a non-repeated absolutely irreducible component defined over  $\mathbb{F}_q$ , then X has an absolutely irreducible component defined over  $\mathbb{F}_q$ .

In our investigation we will need bounds on the number of  $\mathbb{F}_q$ -rational points of algebraic varieties and we will make use of the following result a number of times.

**Theorem 2.5.** [11, Theorem 7.1] Let W be an absolutely irreducible variety defined over  $\mathbb{F}_q$  of dimension n and degree d. If  $q > 2(n+1)d^2$ , then

$$\#(\mathcal{W} \cap \mathbb{A}^N(\mathbb{F}_q)) \ge q^n - (d-1)(d-2)q^{n-1/2} + 5d^{13/3}q^{n-1}.$$

#### 3 Scattered sequences of order 1 and MRD codes

In this section we recall the notion of scatteredness for subspaces and sequences in  $\mathbb{F}_{q^n}^r$ , and how they are related to rank-metric codes.

**Definition 3.1.** [14] Let h, r, n be positive integers, such that h < r. An  $\mathbb{F}_q$ -subspace  $U \subseteq \mathbb{F}_{q^n}^r$  is said to be h-scattered if for every h-dimensional  $\mathbb{F}_{q^n}$ -subspace  $H \subseteq \mathbb{F}_{q^n}^r$ , it holds  $\dim_{\mathbb{F}_q}(U \cap H) \leq h$ . When h = 1, a 1-scattered subspace is simply called scattered.

For what concerns h-scattered subspaces, there is a well-known bound on their  $\mathbb{F}_q$ -dimension. Namely, an h-scattered subspace  $U \subseteq \mathbb{F}_{q^n}^r$ , if U does not define a subgeometry, satisfies

$$\dim_{\mathbb{F}_q}(U) \le \frac{rn}{h+1};\tag{1}$$

see [14]. An h-scattered subspace meeting (1) with equality is called a maximum h-scattered subspace.

Let 
$$\mathcal{G} = \{g_1, \dots, g_k\} \subseteq \mathcal{L}_{n,q}[X_1, \dots, X_m]$$
, where  $\mathcal{L}_{n,q}[X_1, \dots, X_m] := \left\{ \sum_{i=1}^m \sum_{j=0}^{n-1} \gamma_{i,j} X_i^{q^j} : \gamma_{i,j} \in \mathbb{F}_{q^n} \right\}$ , and consider the  $\mathbb{F}_{i}$ -space

consider the  $\mathbb{F}_q$ -space

$$U_{\mathcal{G}} := \{ (g_1(x_1, \dots, x_m), \dots, g_r(x_1, \dots, x_m)) : x_1, \dots, x_m \in \mathbb{F}_{q^n} \} \subseteq \mathbb{F}_{q^n}^r.$$

$$\tag{2}$$

**Definition 3.2.** [4] Let  $\mathcal{I} := (i_1, i_2, \dots, i_m) \in (\mathbb{Z}/n\mathbb{Z})^m$  and consider  $f_1, \dots, f_r \in \mathcal{L}_{n,q}[X_1, \dots, X_m]$ . Let  $U_{\mathcal{I},\mathcal{F}} := U_{\mathcal{F}'}$ , where  $\mathcal{F}' = (X_1^{q^{i_1}}, \dots, X_m^{q^{i_m}}, f_1, \dots, f_s) \subseteq \mathcal{L}_{n,q}[X_1, \dots, X_m]$ . The s-tuple  $\mathcal{F} := (f_1, \dots, f_s)$  is said to be an  $(\mathcal{I}; h)_{q^n}$ -scattered sequence of order m if  $U_{\mathcal{I},\mathcal{F}}$  is maximum

h-scattered in  $\mathbb{F}_{a^n}^{m+s}$ .

An  $(\mathcal{I};h)_{q^n}$ -scattered sequence  $\mathcal{F} := (f_1,\ldots,f_s)$  of order m is said to be **exceptional** if it is h-scattered over infinitely many extensions  $\mathbb{F}_{q^n}^{\ell}$  of  $\mathbb{F}_{q^n}$ .

As the following remark shows,  $(\mathcal{I}; h)_{q^n}$ -scattered sequences, with  $|\mathcal{I}| = 1$ , have been considered also in [9], though with slightly different terminology.

**Remark 3.3.** It is not difficult to see that, for m = 1 and  $\mathcal{I} = \{t\}$ , an (r-1)-tuple  $(f_2, \ldots, f_r) \subseteq \mathcal{L}_{n,q}$ is a  $(\mathcal{I}; r-1)_{q^n}$ -scattered (or simply  $(t, r-1)_{q^n}$ -scattered) sequence of order 1 if and only if, for any  $\alpha_1,\ldots,\alpha_r\in\mathbb{F}_{q^n},$ 

$$\det \begin{pmatrix} \alpha_1^{q^t} & f_2(\alpha_1) & \cdots & f_r(\alpha_1) \\ \alpha_2^{q^t} & f_2(\alpha_2) & \cdots & f_r(\alpha_2) \\ \vdots & \vdots & \cdots & \vdots \\ \alpha_r^{q^t} & f_2(\alpha_r) & \cdots & f_r(\alpha_r) \end{pmatrix} = 0 \implies \dim_{\mathbb{F}_q} \langle \alpha_1, \dots, \alpha_r \rangle_{\mathbb{F}_q} < r;$$

see for istance [8, 9, 32] for an explicit link between scattered spaces and Moore matrices.

If the previous property holds,  $\underline{f} = (x^{q^t}, f_2, \ldots, f_r)$  is said to be a **Moore polynomial set** for q and n of index t (see [9, Definition 9]).

Moore polynomial sets can be characterized in terms of MRD codes as follows.

**Theorem 3.4.** [9] Let r and n be positive integers with  $r \leq n+1$ , and let  $\underline{f} = (x^{q^t}, f_2(x), \ldots, f_r(x))$ , where  $x^{q^t}, f_2(x), \ldots, f_r(x) \in \mathcal{L}_{n,q}$  are  $\mathbb{F}_{q^n}$ -linearly independent. The  $\mathbb{F}_{q^n}$ -linear rank metric code

$$\mathcal{C}_{\underline{f}} = \langle x^{q^t}, f_2(x), \dots, f_r(x) \rangle_{\mathbb{F}_{q^n}}$$

is an MRD code if and only if f is a Moore polynomial set for q and n.

Now we focus on the exceptionality of  $\mathbb{F}_{q^n}$ -linear MRD codes  $\mathcal{C} \subseteq \mathcal{L}_{n,q}$  of dimension r, or equivalently by Theorem 3.4, on the exceptionality of scattered sequences of order 1. We can assume infact, without restrictions, the following properties on the polynomials generating a non-degenerate  $\mathbb{F}_{q^n}$ -linear code  $\mathcal{C}$  (for details see [9] and [4, Definition 2.8]). In particular, from [4, Proposition 2.11] and [28, Corollary IV.10] we can assume that  $\mathcal{C}$  contains a monomial.

**Remark 3.5.** [9, Properties 13] Given a non-degenerate  $\mathbb{F}_{q^n}$ -linear code  $\mathcal{C}$  of dimension r, there exist  $f_1(x), \ldots, f_r(x) \in \mathcal{C}$  such that the following properties hold:

- (1)  $f_1(x) = x^{q^t};$
- (2)  $f_1(x), \ldots, f_r(x)$  are  $\mathbb{F}_{q^n}$ -linearly independent;
- (3)  $M_1 := \deg_a(f_1(x)), \ldots, M_r := \deg_a(f_r(x))$  are all distinct;
- (4)  $m_1 := \min \deg_q(f_1(x)), \ldots, m_r := \min \deg_q(f_r(x))$  are all distinct, and  $m_i = 0$  for some i;
- (5)  $f_1(x), \ldots, f_r(x)$  are monic;
- (6) for any *i*, if  $f_i(x)$  is a monomial then  $m_i = M_i \ge t$ .

A Moore polynomial set  $\underline{f} = (f_1(x), \ldots, f_r(x)) \subseteq \mathcal{L}_{n,q}$  satisfying the previous six properties is said to be a Moore polynomial set for q and n of index t.

#### 4 Scattered sequences and algebraic varieties

In this section, we consider varieties introduced in [9] to traslate the determination of scattered sequences of order 1 into an algebraic geometry problem.

• 
$$\mathcal{U} := \mathcal{U}_{\underline{f}} \subset \mathbb{P}^r(\overline{\mathbb{F}_{q^n}}), \qquad \mathcal{U} : F_{\underline{f}}(X_1, \dots, X_r) := \det(M_{\underline{f}}(X_1, \dots, X_r)) = 0,$$
  
where  
$$M_{\underline{f}}(X_1, \dots, X_r) = \begin{pmatrix} f_1(X_1) & f_2(X_1) & \cdots & f_r(X_1) \\ f_1(X_2) & f_2(X_2) & \cdots & f_r(X_2) \\ \vdots & \vdots & \cdots & \vdots \\ f_1(X_r) & f_2(X_r) & \cdots & f_r(X_r) \end{pmatrix};$$

- $\mathcal{V} := \mathcal{U}_{(x,x^q,\dots,x^{q^{r-1}})} \subset \mathbb{P}^r(\overline{\mathbb{F}_{q^n}}), \qquad \mathcal{V} : F_{(x,x^q,\dots,x^{q^{r-1}})}(X_1,\dots,X_r) = 0$ where  $F_{(x,x^q,\dots,x^{q^{r-1}})}(X_1,\dots,X_r) = \prod_{(a_1,\dots,a_r)\in\mathbb{P}^{r-1}(\mathbb{F}_q)} (a_1X_1 + \dots + a_rX_r); \qquad (3)$
- $\mathcal{W} \subset \mathbb{P}^r(\overline{\mathbb{F}_{q^n}})$ , with affine equation

$$\mathcal{W}: \frac{F_{\underline{f}}(X_1, \dots, X_r)}{F_{(x, x^q, \dots, x^{q^{r-1}})}(X_1, \dots, X_r)} = 0.$$
(4)

The link between scattered sequence of order 1 and algebraic hypersurfaces is straightforward.

**Proposition 4.1.** [9] The (r-1)-tuple  $(f_2, \ldots, f_r)$  is a  $(\{t\}, r-1)_{q^n}$ -scattered sequence of order 1 if and only if all the affine  $\mathbb{F}_{q^n}$ -rational points of  $\mathcal{W}$  lie on  $\mathcal{V}$ .

**Theorem 4.2.** [9, Main Theorem] Let  $C \subseteq \mathcal{L}_{n,q}$  be an exceptional r-dimensional  $\mathbb{F}_{q^n}$ -linear MRD code containing at least a separable polynomial f(x) and a monomial. If r > 3, assume also that q > 5. Let t be the minimum integer such that  $x^{q^t} \in C$ .

If t > 0 and  $C = \langle x^{q^t}, f(x), g_3(x), \dots, g_r(x) \rangle_{\mathbb{F}_{q^n}}$ , with  $\deg(g_i(x)) > \max\{q^t, \deg(f(x))\}$  for each  $i = 3, \dots, r$ , then f(x) is exceptional scattered of index t.

**Remark 4.3.** Note that without loss of generality, we can always assume that a code C contains a separable polynomial ([9, Remark 12]), and if C is non-degenerate it also contains a monomial (see [19, Lemma 2.1] and [28, Corollary IV.10]).

As previously stated in the introduction, until today, the only known non-monomial example of exceptional scattered polynomials, for arbitrary t, is given by the LP polynomials; therefore it is natural to search for exceptional MRD codes where f(x) is of such type. In the following we will focus on RD codes of dimension 3.

# 5 Moore polynomial sets of type $\underline{f} = (x^{q^t}, x + \delta x^{q^{2t}}, G(x))$

In this section we investigate curves arising from Moore polynomial sets for q and n, of index t, of type  $\underline{f} = (x^{q^t}, x + \delta x^{q^{2t}}, G(x))$ . They correspond to  $(\{t\}, 2)_{q^n}$ -scattered sequences of order 1 as we have seen in Remark 3.3. Let q be a prime power and t, n positive integers.

**Proposition 5.1.** [9, Proposition 23] If the pairs  $(x + \delta x^{q^{2t}}, G(x)) \subseteq \mathcal{L}_{n,q}$  is a  $(\{t\}, 2)_{q^n}$ -scattered sequence of order 1 and  $n > 4 \deg_q(G) + 2$ , then  $\min \deg_q(G) = 2t$  or  $\min \deg_q(G) = t/2$ .

Let consider first the case  $\min \deg_q(G) = 2t$ . Set

$$F := X + \delta X^{q^{2t}}, \qquad N_{q^n|q}(\delta) \neq 1, \tag{5}$$

$$G := X^{q^{2t}} + \dots + CX^{q^k}, \qquad C \neq 0,$$
(6)

where  $F, G \in \mathbb{F}_{q^n}[X]$ , and 0 < 2t < k < n. Fix an element  $\lambda \in \mathbb{F}_{q^n} \setminus \mathbb{F}_q$  such that  $F(\lambda) \neq 0 \neq G(\lambda)$  and

$$L_{\xi}(\lambda) := (\xi - \xi^{q^{t}})^{q^{k+t}} (\xi^{q^{k-t}} - \xi)^{q^{t}} F(\lambda)^{q^{t}+1} + \delta(\xi^{q^{k}} - \xi)^{q^{t}(q^{t}+1)} \lambda^{q^{t}(q^{t}+1)} \neq 0,$$
(7)

for each  $\xi \in \mathbb{F}_{q^{k-2t}} \setminus \mathbb{F}_{q^{k-t}}$ . Such an element exists in any field  $\mathbb{F}_{q^n}$ , with  $n \ge k+t+1$ . Indeed the polynomial  $L_{\xi}$  is not zero and of degree  $q^{3t} + q^{2t}$ , for each  $\xi \in \mathbb{F}_{q^{k-2t}} \setminus \mathbb{F}_{q^{k-t}}$ , so

$$\#\left(\bigcup_{\xi \in \mathbb{F}_{q^{k-2t}}} \{\eta : L_{\xi}(\eta) = 0\}\right) \le (q^{3t} + q^{2t})q^{k-2t} = q^{k+t} + q^k < q^n$$

for  $n \ge k + t + 1$ . Consider the curves

$$C: \begin{vmatrix} X^{q^t} & F(X) & G(X) \\ Y^{q^t} & F(Y) & G(Y) \\ \lambda^{q^t} & F(\lambda) & G(\lambda) \end{vmatrix} = 0,$$
(8)  
$$\mathcal{A}: \frac{\begin{vmatrix} X^{q^t} & F(X) & G(X) \\ Y^{q^t} & F(Y) & G(Y) \\ \lambda^{q^t} & F(\lambda) & G(\lambda) \end{vmatrix}}{\begin{vmatrix} X & X^q & X^{q^2} \\ Y & Y^q & Y^{q^2} \\ \lambda & \lambda^q & \lambda^{q^2} \end{vmatrix}} = 0.$$
(9)

**Lemma 5.2.** [9, Lemma 18] Let  $\mathcal{V}: F_{(x,x^q,x^{q^2})}(X,Y,Z) = 0$  and  $\mathcal{W}: \frac{F_{(x^{q^t},x+\delta x^{q^{2t}},G(x))}(X,Y,Z)}{F_{(x,x^q,x^{q^2})}(X,Y,Z)} = 0$  as in Equations 3 and 4. If  $\mathcal{A}$  has a non-repeated  $\mathbb{F}_{q^n}$ -rational absolutely irreducible component not contained in

the curve defined by  $F_{(x,x^q,x^{q^2})}(X,Y,\lambda) = 0$ , then  $\mathcal{W}$  has a non-repeated  $\mathbb{F}_{q^n}$ -rational absolutely irreducible component not contained in  $\mathcal{V}$ .

In the following we exploit a method introduced in [24] via the investigation of singular points of the curves  $\mathcal{A}$  and  $\mathcal{C}$ , see Criterion 2.2.

In particular, our aim is to prove the existence of an absolutely irreducible  $\mathbb{F}_{q^n}$ -rational component of the curve  $\mathcal{A}$ . We proceed as follows.

- 1. We determine the set  $Sing(\mathcal{A})$  of singular points of  $\mathcal{A}$ ; see below.
- 2. For each point  $P \in Sing(\mathcal{A})$  we provide upperbounds on  $I_{P,max}(\mathcal{A})$ ; see Propositions 5.3, 5.6, Lemma 5.7 and Proposition 6.2.
- 3. We compute an upper bound on  $\sum_{P \in Sing(\mathcal{A})} I_{P,max}(\mathcal{A})$  and obtain the desired result via Criterion 2.2; see Theorems 6.1 and 6.3.

We will investigate the singular points of C; in fact, as it can be easily seen, the set of its singular points contains also the singular points of A.

Note that an affine point  $P = (\overline{x}, \overline{y}) \in \mathcal{C}$  is singular if and only if

$$\begin{vmatrix} \overline{x}^{q^t} & G(\overline{x}) \\ \lambda^{q^t} & G(\lambda) \end{vmatrix} = \begin{vmatrix} \overline{y}^{q^t} & G(\overline{y}) \\ \lambda^{q^t} & G(\lambda) \end{vmatrix} = 0.$$

One can see immediately that

$$\begin{vmatrix} (X+\overline{x})^{q^t} & F(X+\overline{x}) & G(X+\overline{x}) \\ (Y+\overline{y})^{q^t} & F(Y+\overline{y}) & G(Y+\overline{y}) \\ \lambda^{q^t} & F(\lambda) & G(\lambda) \end{vmatrix} = H_{q^t} + H_{q^t+1} + \dots,$$

where

$$H_{q^{t}} = \begin{vmatrix} F(\overline{y}) & G(\overline{y}) \\ F(\lambda) & G(\lambda) \end{vmatrix} X^{q^{t}} - \begin{vmatrix} F(\overline{x}) & G(\overline{x}) \\ F(\lambda) & G(\lambda) \end{vmatrix} Y^{q^{t}};$$
(10)

$$H_{q^{t}+1} = G(\lambda)(X^{q^{t}}Y - XY^{q^{t}}).$$
(11)

As a direct consequence of Lemma 2.1 we have the following.

**Proposition 5.3.** Let  $C: F_{\underline{f}}(X, Y, \lambda) = 0$  and  $P = (\overline{x}, \overline{y}) \in C$  such that

$$\begin{vmatrix} \overline{x}^{q^t} & G(\overline{x}) \\ \lambda^{q^t} & G(\lambda) \end{vmatrix} = \begin{vmatrix} \overline{y}^{q^t} & G(\overline{y}) \\ \lambda^{q^t} & G(\lambda) \end{vmatrix} = 0.$$

Then

• 
$$I_{P,max}(\mathcal{C}) \leq \frac{(q^t+1)^2}{4}$$
 if  $\begin{vmatrix} F(\overline{y}) & G(\overline{y}) \\ F(\lambda) & G(\lambda) \end{vmatrix} = \begin{vmatrix} F(\overline{x}) & G(\overline{x}) \\ F(\lambda) & G(\lambda) \end{vmatrix} = 0;$ 

•  $I_{P,max}(\mathcal{C}) \leq q^t$  otherwise.

Remark 5.4. Observe that

$$\begin{vmatrix} \overline{x}^{q^t} & G(\overline{x}) \\ \lambda^{q^t} & G(\lambda) \end{vmatrix} = [\overline{x}G(\lambda)^{1/q^t} - \lambda(\overline{x}^{q^t} + \dots + C^{1/q^t}\overline{x}^{q^{k-t}})]^{q^t} = g(\overline{x})^{q^t},$$

where g is a separable polynomial of degree  $q^{k-t}$ . Thus the number of affine singular points of C is at most  $q^{2(k-t)}$ . Moreover, if  $m_P(C) = q^t + 1$ , that is

$$\begin{vmatrix} F(\overline{y}) & G(\overline{y}) \\ F(\lambda) & G(\lambda) \end{vmatrix} = \begin{vmatrix} F(\overline{x}) & G(\overline{x}) \\ F(\lambda) & G(\lambda) \end{vmatrix} = 0,$$
(12)

then we obtain

$$\frac{\overline{x}^{q^t}}{\lambda^{q^t}} = \frac{G(\overline{x})}{G(\lambda)} = \frac{F(\overline{x})}{F(\lambda)},$$

i.e.

$$F(\overline{x})\lambda^{q^t} - F(\lambda)\overline{x}^{q^t} = \delta\lambda^{q^t}\overline{x}^{q^{2t}} - (\lambda + \delta\lambda^{q^{2t}})\overline{x}^{q^t} + \lambda^{q^t}\overline{x} = 0.$$
(13)

By combining Equation 13 with  $g(\overline{x}) = \overline{x}G(\lambda)^{1/q^t} - \lambda(\overline{x}^{q^t} + \dots + C^{1/q^t}\overline{x}^{q^{k-t}}) = 0$ , we get that  $\overline{x}$  (or  $\overline{y}$  equivalently) must be a root of a polynomial

$$h(X)^{q^{t}} := [\alpha_{0}X + \dots + \alpha_{k-2t}X^{q^{k-2t}} + \tilde{\alpha}X^{q^{t}}]^{q^{t}}$$
(14)

for suitable  $\alpha_0, \ldots, \alpha_{k-2t}, \tilde{\alpha} \in \mathbb{F}_{q^n}$ , where  $\deg(h) \leq q^{\max\{k-2t,t\}}$ . Thus

$$\#\{P \in \mathcal{C} : m_P(\mathcal{C}) = q^t + 1\} \le q^{\max\{2k - 4t, 2t\}}$$

Consider now the line at infinity  $\ell_{\infty}: Z = 0$ . A homogeneous equation of C is given by

$$\begin{vmatrix} X^{q^{t}} & XZ^{q^{2t}-1} + \delta X^{q^{2t}} & X^{q^{2t}}Z^{q^{k}-q^{2t}} + \dots + CX^{q^{k}} \\ Y^{q^{t}} & YZ^{q^{2t}-1} + \delta Y^{q^{2t}} & Y^{q^{2t}}Z^{q^{k}-q^{2t}} + \dots + CY^{q^{k}} \\ \lambda^{q^{t}} & F(\lambda)Z^{q^{2t}-q^{t}} & G(\lambda)Z^{q^{k}-q^{t}} \end{vmatrix} = 0.$$
(15)

Thus  $\mathcal{C} \cap \ell_{\infty}$  is defined by

$$\begin{vmatrix} \delta X^{q^{2t}} & C X^{q^k} \\ \delta Y^{q^{2t}} & C Y^{q^k} \end{vmatrix} = \delta C (X^{q^{2t}} Y^{q^k} - X^{q^k} Y^{q^{2t}}) = \delta C X^{q^{2t}} Y^{q^{2t}} \prod_{\xi \in \mathbb{F}_{q^{k-2t}} \setminus \{0\}} (Y - \xi X)^{q^{2t}} = 0, \quad (16)$$

so points at infinity of C are of type  $P = (1:\xi:0), \xi \in \mathbb{F}_{q^{k-2t}}$ , or P = (0:1:0).

Let consider the projectivity  $\Psi : (x : y : z) \mapsto (x : y - \xi x : z)$  that maps  $(1 : \xi : 0)$  into (1 : 0 : 0). An affine equation of  $\Psi(\mathcal{C})$  (dehomogenizing with respect to X) is given by

$$f(Y,Z) := \begin{vmatrix} 1 & Z^{q^{2t}-1} + \delta & Z^{q^k-q^{2t}} + \dots + C \\ (Y+\xi)^{q^t} & F^*(Y+\xi,Z) & G^*(Y+\xi,Z) \\ \lambda^{q^t} & F(\lambda)Z^{q^{2t}-q^t} & G(\lambda)Z^{q^k-q^t} \end{vmatrix} = 0,$$
(17)

where

$$F^*(Y+\xi,Z) = YZ^{q^{2t}-1} + \delta Y^{q^{2t}} + \xi Z^{q^{2t}-1} + \delta \xi^{q^{2t}};$$
(18)

$$G^*(Y+\xi,Z) = Y^{q^{2t}}Z^{q^k-q^{2t}} + \dots + CY^{q^k} + \xi^{q^{2t}}Z^{q^k-q^{2t}} + \dots + C\xi^{q^k}.$$
(19)

It is not hard to see that

$$\begin{vmatrix} 1 & Z^{q^{2t}-1} + \delta & Z^{q^k-q^{2t}} + \dots + C \\ (Y+\xi)^{q^t} & F^*(Y+\xi,Z) & G^*(Y+\xi,Z) \\ \lambda^{q^t} & F(\lambda)Z^{q^{2t}-q^t} & G(\lambda)Z^{q^k-q^t} \end{vmatrix} = B_{q^{2t}-q^t} + B_{q^{2t}-1} + B_{q^{2t}} + L(Y,Z),$$

where

$$B_{q^{2t}-q^t} = -CF(\lambda)(\xi^{q^{k-t}} - \xi)^{q^t} Z^{q^{2t}-q^t},$$
(20)

$$B_{q^{2t}-1} = C\lambda^{q^{t}}(\xi^{q^{k}} - \xi)Z^{q^{2t}-1}, \qquad (21)$$

$$B_{q^{2t}} = C[F(\lambda)Y^{q^{t}}Z^{q^{2t}-q^{t}} - \lambda^{q^{t}}(YZ^{q^{2t}-1} + \delta Y^{q^{2t}})], \qquad (22)$$

and

$$L(Y,Z) = \sum_{i,j} \alpha^{(i,j)} Y^i Z^j, \qquad i+j \ge q^k - q^{k-1} = q^{k-1}(q-1) \ge q^{2t}(q-1).$$
(23)

**Proposition 5.5.** Let  $P_{\xi} = (1 : \xi : 0), \ \xi \in \mathbb{F}_{q^{k-2t}}$ . Then  $m_{P_{\xi}}(\mathcal{C}) = q^{2t} - q^t$  or  $m_{P_{\xi}}(\mathcal{C}) = q^{2t}$ . Also  $m_{P_{\xi}}(\mathcal{C}) = q^{2t}$  if and only if  $\xi \in \mathbb{F}_{q^{\operatorname{gcd}(k,t)}}$ .

*Proof.* The statement follows immediately from

$$\begin{split} \mathbb{F}_{q^{k-2t}} \cap \mathbb{F}_{q^{k-t}} &= \mathbb{F}_{q^{\gcd(k-2t,k-t)}}, \\ \mathbb{F}_{q^{k-t}} \cap \mathbb{F}_{q^{k}} &= \mathbb{F}_{q^{\gcd(k-t,k)}}, \\ \gcd(k-2t,k-t) &= \gcd(k-t,t) = \gcd(k,t) = \gcd(k,k-t) \end{split}$$

**Proposition 5.6.** Let  $P = (1:\xi:0)$ , where  $\xi \in \mathbb{F}_{q^{\text{gcd}(k,t)}}$ , or P = (0:1:0). Then  $I_{P,max}(\mathcal{C}) \leq \frac{q^{4t}}{4}$ .

 $Proof. \quad \bullet \ P = (1:\xi:0), \text{ with } \xi \in \mathbb{F}_{q^{\gcd(k,t)}}.$ 

The statement follows from Lemma 2.1 and Proposition 5.5.

• P = (0:1:0). Since C is fixed by the projectivity  $(x:y:z) \mapsto (y:x:z)$ , we get that  $I_{(0:1:0),max} = I_{(1:0:0),max} \leq \frac{q^{4t}}{4}$ .

# 5.1 Study of the intersection multiplicity of branches at the singular points $P_{\xi} = (1 : \xi : 0)$ , with $\xi \notin \mathbb{F}_{q^{\text{gcd}(k,t)}}$

In this subsection we consider singular points of type  $P_{\xi} = (1 : \xi : 0)$ , where  $\xi \notin \mathbb{F}_{q^{\text{gcd}(k,t)}}$ . In particular we study branches centred at those points an we determine upper bounds on the multiplicity of intersection of putative components of  $\mathcal{C}$  (and therefore  $\mathcal{A}$ ).

**Lemma 5.7.** Let  $P_{\xi} = (1 : \xi : 0)$ , with  $\xi \notin \mathbb{F}_{q^{gcd}(k,t)}$ . Then there is a unique branch centered at  $P_{\xi}$ . Thus, the multiplicity of two putative components of C (and therefore  $\mathcal{A}$ ) in  $P_{\xi}$  is 0.

*Proof.* Consider the local quadratic transformation  $f^{(1)}(Y, Z) = f(Y, YZ)/Y^{q^{2t}-q^{t}}$ , where f(Y, Z) is defined as in (17). We have that

$$f^{(1)}(Y,Z) = \alpha^{(0,q^{2t}-q^t)} Z^{q^{2t}-q^t} + \alpha^{(0,q^{2t}-1)} Y^{q^t-1} Z^{q^{2t}-1} + \alpha^{(q^t,q^{2t}-q^t)} Y^{q^t} Z^{q^{2t}-q^t}$$
  
+  $\alpha^{(1,q^{2t}-1)} Y^{q^t} Z^{q^{2t}-1} + \alpha^{(q^{2t},0)} Y^{q^t} + L^{(1)}(Y,Z),$ 

where

$$\alpha^{(0,q^{2t}-q^t)} = -CF(\lambda)(\xi^{q^{k-t}}-\xi)^{q^t};$$
(24)

$$\alpha^{(0,q^{2t}-1)} = C\lambda^{q^t}(\xi^{q^k} - \xi); \tag{25}$$

$$\alpha^{(1,q^{2t}-1)} = -C\lambda^{q^t}; \tag{26}$$

$$\alpha^{(q^{2t},0)} = -C\lambda^{q^t}\delta; \tag{27}$$

$$\alpha^{(q^t, q^{2t} - q^t)} = CF(\lambda); \tag{28}$$

$$L^{(1)}(Y,Z) = \sum_{i,j} \alpha^{(i,j)} Y^{i+j-q^{2t}+q^t} Z^j.$$
(29)

Since  $f^{(1)}(0, Z) = \alpha^{(0,q^{2t}-q^t)} Z^{q^{2t}-q^t}$ , the branches centered at (0,0) belonging to f = 0 correspond to the branches centered at (0,0) belonging to  $f^{(1)} = 0$ . Now the origin is a non-ordinary  $q^t$ -point of the curve defined by  $f^{(1)}$  and the line Z = 0 is not the tangent line at the origin; so we can apply the local quadratic transformation  $f^{(2)}(Y,Z) = f^{(1)}(YZ,Z)/Z^{q^t}$ . After applying the same transformation  $q^t - 2$  times, we obtain

$$\begin{split} f^{(q^{t}-1)}(Y,Z) &= \alpha^{(0,q^{2t}-q^{t})}Z^{q^{t}} + \alpha^{(0,q^{2t}-1)}Y^{q^{t}-1}Z^{q^{2t}-q^{t}+1} + \alpha^{(q^{t},q^{2t}-q^{t})}Y^{q^{t}}Z^{q^{2t}-q^{t}} \\ &+ \alpha^{(1,q^{2t}-1)}Y^{q^{t}}Z^{q^{2t}-1} + \alpha^{(q^{2t},0)}Y^{q^{t}} + L^{(q^{t}-1)}(Y,Z), \end{split}$$

where  $L^{(q^t-1)}(Y,Z) = \sum_{i,j} \alpha^{(i,j)} Y^{i+j-q^{2t}+q^t} Z^{j+(q^t-2)(i+j-q^{2t})}.$ Note that, for each  $2 \le \ell \le q^t - 1$ ,  $L^{(\ell)}(Y,Z) = \sum_{i,j} \alpha^{(i,j)} Y^{i+j-q^{2t}+q^t} Z^{j+(\ell-1)(i+j-q^{2t})}.$ 

Also,  $j + (\ell - 1)(i + j - q^{2t}) \ge i + j - q^{2t} > 0$ , so  $L^{(\ell)}(Y, 0) = 0$  and  $f^{(\ell)}(Y, 0) = \alpha^{(q^{2t}, 0)}Y^{q^t}$ . Therefore, the branches centered at (0, 0) belonging to f = 0 are in one-to-one correspondence with the branches centered at (0, 0) belonging to  $f^{(\ell)} = 0$ , for each  $2 \le \ell \le q^t - 1$ .

Now consider the transformation  $(Y, Z) \mapsto (Y - \beta_1 Z, Z)$ , where  $\alpha^{(q^{2t}, 0)} \beta_1^{q^t} + \alpha^{(0, q^{2t} - q^t)} = 0$ , i.e.

$$\beta_1^{q^t} = -\frac{F(\lambda)(\xi^{q^{k-t}} - \xi)^{q^t}}{\lambda^{q^t}\delta}.$$
(30)

We get

$$f^{(q^{t})}(Y,Z) = f^{(q^{t}-1)}(Y + \beta_{1}Z,Z) = \alpha^{(0,q^{2t}-1)} \sum_{\mu=0}^{q^{t}-1} \beta_{1}^{q^{t}-1-\mu} Y^{\mu} Z^{q^{2t}-\mu} + \alpha^{(q^{t},q^{2t}-q^{t})}(Y^{q^{t}} Z^{q^{2t}-q^{t}} + \beta_{1}^{q^{t}} Z^{q^{2t}}) + \alpha^{(1,q^{2t}-1)}(Y^{q^{t}} Z^{q^{2t}-1} + \beta_{1}^{q^{t}} Z^{q^{2t}+q^{t}-1}) + \alpha^{(q^{2t},0)} Y^{q^{t}} + L^{(q^{t})}(Y,Z),$$

where  $L^{(q^t)}(Y,Z) = \sum_{i+j \ge q^{2t}(q-1)} \alpha^{(i,j)} \sum_{\nu=0}^{i+j-q^{2t}+q^t} {i+j-q^{2t}+q^t \choose \nu} \beta_1^{i+j-q^{2t}+q^t-\nu} Y^{\nu} Z^{j+(q^t-1)(i+j-q^{2t})+q^t-\nu}.$ Since the origin is still a non-ordinary  $q^t$ -point of  $f^{(q^t)} = 0$  and the line Z = 0 is not the tangent line at the origin, we can apply the local quadratic transformation  $f^{(q^t+1)}(Y,Z) = f^{(q^t)}(YZ,Z)/Z^{q^t}$ . After applying the same transformation  $q^t - 1$  times, we obtain

$$f^{(2q^{t}-1)}(Y,Z) = \alpha^{(0,q^{2t}-1)} \sum_{\mu=0}^{q^{t}-1} (-1)^{\mu} \beta_{1}^{q^{t}-1-\mu} Y^{\mu} Z^{q^{t}+(q^{t}-2)\mu} + \alpha^{(q^{t},q^{2t}-q^{t})} (Y^{q^{t}} Z^{q^{2t}-q^{t}} + \beta_{1}^{q^{t}} Z^{q^{t}})$$
  
+  $\alpha^{(1,q^{2t}-1)} (Y^{q^{t}} Z^{q^{2t}-1} + \beta_{1}^{q^{t}} Z^{2q^{t}-1}) + \alpha^{(q^{2t},0)} Y^{q^{t}} + L^{(2q^{t}-1)}(Y,Z),$ 

where  $L^{(2q^t-1)}(Y,Z) = \sum_{i+j \ge q^{2t}(q-1)} \alpha^{(i,j)} \sum_{\nu=0}^{i+j-q^{2t}+q^t} {i+j-q^{2t}+q^t \choose \nu} \beta_1^{i+j-q^{2t}+q^t-\nu} Y^{\nu} Z^{j+(q^t-1)(i+j-q^{2t})+(q^t-2)(\nu-q^t)}.$ Note that, for each  $1 \le \ell \le q^t - 2$ ,

$$j + (q^{t} - 1)(i + j - q^{2t}) + \ell(\nu - q^{t}) \ge j + \ell(i + j - q^{2t} - q^{t} + \nu) \ge i + j - q^{2t}(1 + \frac{1}{q^{t}}) > 0,$$

so  $L^{(\ell)}(Y,0) = 0$  and  $f^{(\ell)}(Y,0) = \alpha^{(q^{2t},0)}Y^{q^t}$ . Therefore the branches centered at (0,0) for f = 0 are in one-to-one correspondence with the branches centered at (0,0) for  $f^{(2q^t-1)} = 0$ . Consider the map  $(Y,Z) \mapsto (Y - \beta_2 Z, Z)$ , with  $\alpha^{(0,q^{2t}-1)}\beta_1^{q^t-1} + \alpha^{(q^t,q^{2t}-q^t)}\beta_1^{q^t} + \alpha^{(q^{2t},0)}\beta_2^{q^t} = 0$ , i.e.

$$\beta_2^{q^{2t}} = \frac{\beta_1^{q^t(q^t-1)} [\lambda^{q^t}(\xi^{q^k} - \xi) + F(\lambda)\beta_1]^{q^t}}{\lambda^{q^{2t}} \delta^{q^t}} \\ = \frac{F(\lambda)^{q^t-1} (\xi^{q^{k-t}} - \xi)^{q^t(q^t-1)} [\lambda^{q^t(q^t+1)} \delta(\xi^{q^k} - \xi)^{q^t} - F(\lambda)^{q^t+1} (\xi^{q^{k-t}} - \xi)^{q^t}]}{\lambda^{2q^{2t}} \delta^{2q^t}}.$$

We get

$$\begin{split} f^{(2q^{t})}(Y,Z) &= \alpha^{(0,q^{2t}-1)} \sum_{\mu=1}^{q^{t}-1} \sum_{\tilde{\mu}=0}^{\mu} (-1)^{\mu} \binom{\mu}{\tilde{\mu}} \beta_{1}^{q^{t}-1-\mu} \beta_{2}^{\mu-\tilde{\mu}} Y^{\tilde{\mu}} Z^{q^{t}+(q^{t}-1)\mu-\tilde{\mu}} + \alpha^{(q^{t},q^{2t}-q^{t})} (Y^{q^{t}} Z^{q^{2t}-q^{t}} + \beta_{2}^{q^{t}} Z^{q^{2t}}) \\ &+ \alpha^{(1,q^{2t}-1)} (Y^{q^{t}} Z^{q^{2t}-1} + \beta_{2}^{q^{t}} Z^{q^{2t}+q^{t}-1} + \beta_{1}^{q^{t}} Z^{2q^{t}-1}) + \alpha^{(q^{2t},0)} Y^{q^{t}} + L^{(2q^{t})}(Y,Z), \\ \text{where } L^{(2q^{t})}(Y,Z) &= \sum_{i+j \ge q^{2t}(q-1)} \sum_{\nu=0}^{i+j-q^{2t}+q^{t}} \sum_{\tilde{\nu}=0}^{\nu} \tilde{\alpha}_{\nu\tilde{\nu}}^{(i,j)} Y^{\tilde{\nu}} Z^{j+(q^{t}-1)(i+j-q^{2t})+(q^{t}-2)(\nu-q^{t})+\nu-\tilde{\nu}}, \end{split}$$

for suitable  $\tilde{\alpha}_{\nu\tilde{\nu}}^{(i,j)} \in \mathbb{F}_{q^n}$ . Let apply again the quadratic transformation  $f^{(2q^t+1)}(Y,Z) = f^{(2q^t)}(YZ,Z)/Z^{q^t}$ . It follows that

$$f^{(2q^{t}+1)}(Y,Z) = \alpha^{(0,q^{2t}-1)} \sum_{\mu=1}^{q^{t}-1} \sum_{\tilde{\mu}=0}^{\mu} (-1)^{\mu} {\binom{\mu}{\tilde{\mu}}} \beta_{1}^{q^{t}-1-\mu} \beta_{2}^{\mu-\tilde{\mu}} Y^{\tilde{\mu}} Z^{(q^{t}-1)\mu} + \alpha^{(q^{t},q^{2t}-q^{t})} (Y^{q^{t}} Z^{q^{2t}-q^{t}} + \beta_{2}^{q^{t}} Z^{q^{2t}-q^{t}}) + \alpha^{(1,q^{2t}-1)} (Y^{q^{t}} Z^{q^{2t}-1} + \beta_{2}^{q^{t}} Z^{q^{2t}-1} + \beta_{1}^{q^{t}} Z^{q^{t}-1}) + \alpha^{(q^{2t},0)} Y^{q^{t}} + L^{(2q^{t}+1)}(Y,Z),$$

where  $L^{(2q^t+1)}(Y,Z) = \sum_{i+j \ge q^{2t}(q-1)} \sum_{\nu=0}^{i+j-q^{2t}+q^t} \sum_{\tilde{\nu}=0}^{\nu} \tilde{\alpha}_{\nu\tilde{\nu}}^{(i,j)} Y^{\tilde{\nu}} Z^{j+(q^t-1)(i+j-q^{2t}+\nu-q^t)}.$ 

Thus the smallest homogeneous part of  $f^{(2q^t+1)}(Y,Z)$  is given by  $(\alpha^{(1,q^{2t}-1)}\beta_1^{q^t} - \alpha^{(0,q^{2t}-1)}\beta_1^{q^t-2}\beta_2)Z^{q^t-1}$ ,

where

$$(\alpha^{(1,q^{2t}-1)}\beta_1^{q^t} - \alpha^{(0,q^{2t}-1)}\beta_1^{q^t-2}\beta_2)^{q^{2t}} = -C^{q^{2t}}\lambda^{q^{3t}}\beta_1^{q^{2t}(q^t-2)}[\beta_1^{2q^{2t}} + (\xi^{q^k} - \xi)^{q^t}\beta_2^{q^{2t}}]$$

Therefore from (30) it follows that  $m_{(0,0)}(f^{(2q^t+1)}) = q^t - 1$  if and only

$$\beta_1^{2q^{2t}} + (\xi^{q^k} - \xi)^{q^t} \beta_2^{q^{2t}} \neq 0,$$

i.e., after some computations,

$$\frac{F(\lambda)^{q^t-1}}{\lambda^{2q^{2t}}\delta^{2q^t}} (\xi^{q^{k-t}} - \xi)^{q^{2t}-q^t} [(\xi - \xi^{q^t})^{q^{k+t}} (\xi^{q^{k-t}} - \xi)^{q^t} F(\lambda)^{q^t+1} + \delta(\xi^{q^k} - \xi)^{q^t(q^t+1)} \lambda^{q^t(q^t+1)}] \neq 0$$
(31)

(recall Equation 7).

Also, the homogeneous part of degree  $q^t$  in  $f^{(2q^t+1)}(Y,Z)$  is not zero and contains the term  $\alpha^{(q^{2t},0)}Y^{q^t}$ . By Lemma 2.1(i) it follows that  $I_{P_{\xi},max}(\mathcal{C}) = 0$ , i.e. there is a unique branch of  $\mathcal{C}$  centered at  $P_{\xi}$ .

#### 6 Main result

#### 6.1 Case min deg<sub>q</sub>(G(X)) = 2t

Thanks to the results of the previous section, we now are able to prove the following theorem.

**Theorem 6.1.** Let q, t, k be intergers, with q > 2, t > 0 and k > 2t. If  $(t, q) \notin \{(1,3); (1,4); (1,5); (2,3)\}$ and  $n \ge k + t + 1$ , then  $\mathcal{A}$  has an absolutely irreducible component defined over  $\mathbb{F}_{q^n}$  and not contained in the curve defined by  $F_{(x,x^q,x^{q^2})}(X,Y,\lambda) = 0$ .

*Proof.* Since C has a finite number of singularities, the curve defined by  $F_{(x,x^q,x^{q^2})}(X,Y,\lambda) = 0$  is not a component of A. Moreover, any upper bound on  $I_{P,max}(C)$  is also an upper bound for  $I_{P,max}(A)$ , since the curve A is a component of the curve C.

We want to prove that  $\sum_{P \in Sing(\mathcal{A})} I_{P,max}(\mathcal{A}) < \frac{2}{9} \deg^2(\mathcal{A}) = \frac{2}{9} (q^k + q^{2t} - (q^2 + q + 1))^2; \text{ the statement will then follow by Criterion 2.2.}$ 

Let

$$\begin{split} \Omega &:= \{(1:\xi:0):\xi\in\mathbb{F}_{q^{k-2t}}\}\cup\{(0:1:0)\},\\ \Pi &:= \{(1:\xi:0):\xi\in\mathbb{F}_{q^{\gcd(k,t)}}\},\\ \Theta &:= \{(\overline{x},\overline{y}):\overline{x}^{q^{t}}G(\lambda)-\lambda^{q^{t}}G(\overline{x})=\overline{y}^{q^{t}}G(\lambda)-\lambda^{q^{t}}G(\overline{y})=0\},\\ \Sigma &:= \{(\overline{x},\overline{y})\in\Theta:F(\overline{x})\lambda^{q^{t}}-F(\lambda)\overline{x}^{q^{t}}=F(\overline{y})\lambda^{q^{t}}-F(\lambda)\overline{y}^{q^{t}}=0\}. \end{split}$$

We have that  $Sing(\mathcal{C}) = \Omega \cup \Theta$ ,  $\#\Omega = q^{k-2t}+1$ ,  $\#\Pi = q^{\operatorname{gcd}(k,t)}+1$ ,  $\#\Theta \leq q^{2(k-t)}$ ,  $\#\Sigma \leq q^{\min\{\max\{2(k-2t),2t\},4t\}}$  (see Remark 5.4) and by Proposition 5.3, Proposition 5.6, and Lemma 5.7

$$I_{P,\max}(\mathcal{A}) \leq \begin{cases} 0 & P \in \Omega \setminus \Pi; \\ \frac{q^{4t}}{4} & P \in \Pi; \\ q^t & \Theta \setminus \Sigma; \\ \frac{(q^t+1)^2}{4} & P \in \Sigma. \end{cases}$$

Observe that  $Sing(\mathcal{C})$  is a finite set, so the curve defined by  $F_{(x,x^q,x^{q^2})}(X,Y,\lambda) = 0$  is not a component of  $\mathcal{A}$ . We distinguish the following cases.

• Case k > 3t

It follows that

$$\sum_{P \in Sing(\mathcal{A})} I_{P,max}(\mathcal{A}) \leq (q^{2(k-t)} - q^{4t})q^t + q^{4t} \cdot \frac{(q^t + 1)^2}{4} + (q^{\gcd(k,t)} + 1)\frac{q^{4t}}{4}$$

$$\leq q^{2k-t} - q^{5t} + \frac{q^{6t}}{4} + \frac{q^{5t}}{2} + \frac{q^{4t}}{4} + \frac{q^{5t}}{4} + \frac{q^{4t}}{4} = q^{2k-t} + \frac{q^{6t}}{4} - \frac{q^{5t}}{4} + \frac{q^{4t}}{2}$$

$$\leq q^{2k-t} + \frac{q^{6t}}{4} \leq \frac{5}{4}q^{\max\{2k-t,6t\}} \leq \frac{2}{9}q^{2k} < \frac{2}{9}(q^k + q^{2t} - (q^2 + q + 1))^2,$$

for  $t \ge 2$  and  $q \ge 3$ .

Finally, for t = 1, the inequality

$$(q^{2(k-t)} - q^{4t})q^t + q^{4t} \cdot \frac{(q^t + 1)^2}{4} + (q^{\gcd(k,t)} + 1)\frac{q^{4t}}{4} < \frac{2}{9}(q^k + q^{2t} - (q^2 + q + 1))^2$$
(32)

is satisfied when even

$$\frac{2}{9} - \frac{1}{q} - \frac{4(q+1)}{9q^k} - \frac{1}{4q^2} > 0,$$

that holds for  $q \geq 5$ .

• Case k = 3tWe have

$$\sum_{P \in Sing(\mathcal{A})} I_{P,max}(\mathcal{A}) \leq (q^{4t} - q^{2t})q^t + q^{2t} \cdot \frac{(q^t + 1)^2}{4} + (q^t + 1)\frac{q^{4t}}{4}$$

$$\leq \frac{5}{4}q^{5t} + \frac{q^{4t}}{2} - \frac{q^{3t}}{2} + \frac{q^{2t}}{4} \leq \frac{5}{4}q^{5t} + \frac{q^{4t}}{2}$$

$$\leq \frac{2}{9}q^{6t} < \frac{2}{9}(q^{3t} + q^{2t} - (q^2 + q + 1))^2,$$
(33)

for  $q^t \ge 6$ , so q > 2 and  $t \ge 2$ . For t = 1, the inequality  $\sum_{P \in Sing(\mathcal{A})} I_{P,max}(\mathcal{A}) < \frac{2}{9}q^{2k}$  is satisfied when

$$\frac{2}{9} - \frac{5}{4q} - \frac{4(q+1)}{9q^3} - \frac{1}{2q^2} > 0,$$

that holds for  $q \geq 7$ .

#### • Case 2t < k < 3tIt results

$$\sum_{P \in Sing(\mathcal{A})} I_{P,max}(\mathcal{A}) \le (q^{2(k-t)} - q^{2t})q^t + q^{2t} \cdot \frac{(q^t + 1)^2}{4} + (q^s + 1)\frac{q^{4t}}{4},$$
(34)

where  $s := \gcd(k, t), s \le t/2, k = \tilde{k}s, t = \tilde{t}s, \gcd(\tilde{k}, \tilde{t}) = 1$ . Note that k > 2t implies  $\tilde{k} \ge 2\tilde{t} + 1$ , so  $s + 4t = (1 + 4\tilde{t})s < (1 + 2\tilde{t})2s \le 2\tilde{k}s = 2k$ . It follows that

$$\sum_{P \in Sing(\mathcal{A})} I_{P,max}(\mathcal{A}) \leq q^{2k-t} + \frac{q^{4t}}{2} - \frac{q^{3t}}{2} + \frac{q^{2t}}{4} + \frac{q^{4t+s}}{4}$$

$$\leq q^{2k-t} + \frac{q^{4t+s}}{2} \leq \begin{cases} \frac{3}{2}q^{2k-1} & \text{for } t \geq 1\\ (1+\frac{q}{2})q^{2k-2} & \text{for } t \geq 2\\ (1+\frac{q^2}{2})q^{2k-3} & \text{for } t \geq 3 \end{cases}$$

$$\leq \frac{2}{9}q^{2k} < \frac{2}{9}(q^k + q^{2t} - (q^2 + q + 1))^2,$$

for  $t \ge 3$  and  $q \ge 3$ ,  $t \ge 2$  and  $q \ge 4$ , or t = 1 and  $q \ge 7$ .

# **6.2** Case min deg<sub>q</sub>(G(X)) = t/2

The procedure is exactly the same as for  $\min \deg_q(G(X)) = 2t$ . Consider

$$G := X^{q^{t/2}} + \dots + CX^{q^k} \qquad C \neq 0, \quad 0 < 2t < k < n,$$
(35)

and  $F, \lambda, C, A$  as in Equations 5, 7, 8, 9. Actually, all the results in Section 6 apply here. In particular, by repeating all the computations in Section 6 we get the analogous of Propositions 5.5, 5.6 and Lemma 5.7.

**Proposition 6.2.** Let  $n \ge k + t + 1$  and  $F, G, \lambda, C, A$  as in Equations 5, 7, 8, 9, 35. Then

•  $Sing(\mathcal{C}) \cap \{Z = 0\} = \{(1:\xi:0): \xi \in \mathbb{F}_{q^{k-2t}}\} \cup \{(0:1:0)\},\$ 

• 
$$m_{(1:\xi:0)}(\mathcal{C}) = q^{2t} - q^t$$
 or  $m_{(1:\xi:0)}(\mathcal{C}) = q^{2t}$ 

•  $m_{(1:\xi:0)}(\mathcal{C}) = q^{2t}$  if and only if  $\xi \in \mathbb{F}_{q^{\text{gcd}(k,t)}}$ ,

• 
$$I_{(1:\xi:0),\max}(\mathcal{C}) \leq \begin{cases} \frac{q^{4t}}{4} & \text{if } \xi \in \mathbb{F}_{q^{\gcd(k,t)}};\\ 0 & \text{if } \xi \in \mathbb{F}_{q^{k-2t}} \setminus \mathbb{F}_{q^{\gcd(k,t)}}. \end{cases}$$

For affine singularities of C, we proceed exactly as in Section 5. A point  $P = (\overline{x}, \overline{y}) \in C$  is singular if and only if

$$\begin{vmatrix} \overline{x}^{q^t} & G(\overline{x}) \\ \lambda^{q^t} & G(\lambda) \end{vmatrix} = \begin{vmatrix} \overline{y}^{q^t} & G(\overline{y}) \\ \lambda^{q^t} & G(\lambda) \end{vmatrix} = 0.$$

By direct computations one can observe that

$$\begin{vmatrix} (X+\overline{x})^{q^t} & F(X+\overline{x}) & G(X+\overline{x}) \\ (Y+\overline{y})^{q^t} & F(Y+\overline{y}) & G(Y+\overline{y}) \\ \lambda^{q^t} & F(\lambda) & G(\lambda) \end{vmatrix} = H_{q^{t/2}} + H_{q^{t/2}+1} + \dots,$$

where

$$H_{q^{t/2}} = \begin{vmatrix} \overline{y}^{q^t} & F(\overline{y}) \\ \lambda^{q^t} & F(\lambda) \end{vmatrix} X^{q^{t/2}} - \begin{vmatrix} \overline{x}^{q^t} & F(\overline{x}) \\ \lambda^{q^t} & F(\lambda) \end{vmatrix} Y^{q^{t/2}};$$
(36)

$$H_{q^{t/2}+1} = \lambda^{q^t} (X^{q^{t/2}}Y - XY^{q^{t/2}}).$$
(37)

Thus  $I_{P,max}(\mathcal{C}) \leq q^{t/2}$  or  $I_{P,max}(\mathcal{C}) \leq \frac{(q^{t/2}+1)^2}{4}$  by Lemma 2.1. Also, observe that

$$\frac{\overline{x}^{q^{t}}}{\lambda^{q^{t}}} \frac{G(\overline{x})}{G(\lambda)} \bigg| = \left[ \overline{x}^{q^{t/2}} G(\lambda)^{1/q^{t/2}} - \lambda^{q^{t/2}} (\overline{x} + \dots + C^{1/q^{t/2}} \overline{x}^{q^{k-t/2}}) \right]^{q^{t/2}} = g(\overline{x})^{q^{t/2}},$$

where g is a separable polynomial of degree  $q^{k-t/2}$ . Thus the number of affine singular points of C is at most  $q^{2k-t}$ . Finally, if  $m_P(\mathcal{C}) = q^{t/2} + 1$ , that is

$$\begin{vmatrix} \overline{y}^{q^t} & F(\overline{y}) \\ \lambda^{q^t} & F(\lambda) \end{vmatrix} = \begin{vmatrix} \overline{x}^{q^t} & F(\overline{x}) \\ \lambda^{q^t} & F(\lambda) \end{vmatrix} = 0,$$
(38)

we obtain

$$F(\overline{x})\lambda^{q^{t}} - F(\lambda)\overline{x}^{q^{t}} = \delta\lambda^{q^{t}}\overline{x}^{q^{2t}} - (\lambda + \delta\lambda^{q^{2t}})\overline{x}^{q^{t}} + \lambda^{q^{t}}\overline{x} = 0.$$
(39)

By combining last equation with  $g(\overline{x}) = \overline{x}^{q^{t/2}} G(\lambda)^{1/q^{t/2}} - \lambda^{q^{t/2}} (\overline{x} + \dots + C^{1/q^{t/2}} \overline{x}^{q^{k-t/2}}) = 0$ , we get that  $\overline{x}$  (or  $\overline{y}$  equivalently) must be a root of a *q*-polynomial

$$h(X)^{q^{t/2}} := [\alpha_0 X + \dots + \alpha_{k-2t} X^{q^{k-t}} + \tilde{\alpha} X^{q^{3t/2}}]^{q^{t/2}},$$
(40)

for suitable  $\alpha_0, \ldots, \alpha_{k-2t}, \tilde{\alpha} \in \mathbb{F}_{q^n}$ , where  $\deg(h) \leq q^{\max\{k-t, 3t/2\}}$ .

Now we are able to prove the analogous of Theorem 6.1 for  $\operatorname{min} \operatorname{deg}_{a}(G) = t/2$ .

**Theorem 6.3.** Let q, t, k be integers, with q > 2, t > 0 even and k > 2t. Let  $n \ge k+t+1$  and  $F, G, \lambda, C, \mathcal{A}$  as in Equations 5, 7, 8, 9, 35. If  $(t,q) \notin \{(2,3); (2,4); (2,5), (4,3)\}$ , then  $\mathcal{A}$  has an absolutely irreducible component defined over  $\mathbb{F}_{q^n}$  and not contained in the curve defined by  $F_{(x,x^q,x^{q^2})}(X,Y,\lambda) = 0$ .

Proof. Since  $|Sing(\mathcal{C})| < \infty$ , the curve defined by  $F_{(x,x^q,x^{q^2})}(X,Y,\lambda) = 0$  is not a component of  $\mathcal{A}$ ; thus by Criterion 2.2 it is enough to prove that  $\sum_{P \in Sing(\mathcal{A})} I_{P,max}(\mathcal{A}) < \frac{2}{9} \deg^2(\mathcal{A}) = \frac{2}{9}(q^k + q^{2t} - (q^2 + q + 1))^2$ . Let

$$\begin{split} \Omega &:= \{(1:\xi:0):\xi\in\mathbb{F}_{q^{k-2t}}\}\cup\{(0:1:0)\},\\ \Pi &:= \{(1:\xi:0):\xi\in\mathbb{F}_{q^{\gcd(k,t)}}\},\\ \Theta &:= \{(\overline{x},\overline{y}):\overline{x}^{q^{t}}G(\lambda)-\lambda^{q^{t}}G(\overline{x})=\overline{y}^{q^{t}}G(\lambda)-\lambda^{q^{t}}G(\overline{y})=0\},\\ \Sigma &:= \{(\overline{x},\overline{y})\in\Theta:F(\overline{x})\lambda^{q^{t}}-F(\lambda)\overline{x}^{q^{t}}=F(\overline{y})\lambda^{q^{t}}-F(\lambda)\overline{y}^{q^{t}}=0\}. \end{split}$$

We have that  $Sing(\mathcal{C}) = \Omega \cup \Theta$ ,  $\#\Omega = q^{k-2t} + 1$ ,  $\#\Pi = q^{\operatorname{gcd}(k,t)} + 1$ ,  $\#\Theta \le q^{2k-t}$ ,  $\#\Sigma \le q^{\min\{\max\{2(k-t),3t\},4t\}}$ , and by Equation 36 and Proposition 6.2

$$I_{P,\max}(\mathcal{A}) \leq \begin{cases} 0 & P \in \Omega \setminus \Pi; \\ \frac{q^{4t}}{4} & P \in \Pi; \\ q^{t/2} & \Theta \setminus \Sigma; \\ \frac{(q^{t/2}+1)^2}{4} & P \in \Sigma. \end{cases}$$

We distinguish the following cases.

• Case k > 5t/2One gets

$$\sum_{P \in Sing(\mathcal{A})} I_{P,max}(\mathcal{A}) \leq (q^{2k-t} - q^{4t})q^{t/2} + q^{4t} \cdot \frac{(q^{t/2} + 1)^2}{4} + (q^{\gcd(k,t)} + 1)\frac{q^{4t}}{4}$$

$$\leq q^{2k-t/2} - q^{9t/2} + \frac{q^{5t}}{4} + \frac{q^{9t/2}}{2} + \frac{q^{4t}}{4} + \frac{q^{5t}}{4} + \frac{q^{4t}}{4}$$

$$= q^{2k-t/2} - \frac{q^{9t/2}}{2} + \frac{q^{5t}}{2} + \frac{q^{4t}}{2}$$

$$\leq q^{2k-t/2} + \frac{q^{5t}}{2} \leq \frac{3}{2}q^{\max\{2k-t/2,5t\}} \leq \frac{2}{9}q^{2k} < \frac{2}{9}(q^k + q^{2t} - (q^2 + q + 1))^2,$$

for  $t \ge 4$  and  $q \ge 3$ , or t = 2 and  $q \ge 7$ .

• Case k = 5t/2We have

$$\sum_{P \in Sing(\mathcal{A})} I_{P,max}(\mathcal{A}) \leq (q^{4t} - q^{3t})q^{t/2} + q^{3t} \cdot \frac{(q^{t/2} + 1)^2}{4} + (q^t + 1)\frac{q^{4t}}{4}$$
$$\leq \frac{q^{5t}}{4} + q^{9t/2} - \frac{q^{7t/2}}{2} + \frac{q^{4t}}{2} + \frac{q^{3t}}{4} \leq \frac{q^{5t}}{4} + q^{9t/2} + \frac{q^{4t}}{2}$$
$$\leq \frac{3}{4}q^{5t} + \frac{q^{4t}}{2} \leq \frac{2}{9}q^{6t} < \frac{2}{9}(q^{3t} + q^{2t} - (q^2 + q + 1))^2,$$

for  $q^t \ge 4$ , so for all q, t.

• Case 2t < k < 5t/2We have

$$(q^{2k-t} - q^{3t})q^{t/2} + q^{3t} \cdot \frac{(q^{t/2} + 1)^2}{4} + (q^s + 1)\frac{q^{4t}}{4} < \frac{2}{9}(q^k + q^{2t} - (q^2 + q + 1))^2$$

where  $s := \gcd(k, t), s \le t/2, k = \tilde{k}s, t = \tilde{t}s, \gcd(\tilde{k}, \tilde{t}) = 1$ . Of course k > 2t implies  $\tilde{k} \ge 2\tilde{t} + 1$ , so  $s + 4t = (1 + 4\tilde{t})s < (1 + 2\tilde{t})2s \le 2\tilde{k}s = 2k$ . It follows that

$$\sum_{P \in Sing(\mathcal{A})} I_{P,max}(\mathcal{A}) \leq (q^{2k-t} - q^{3t})q^{t/2} + q^{3t} \cdot \frac{(q^{t/2} + 1)^2}{4} + (q^s + 1)\frac{q^{4t}}{4}$$
$$= q^{2k-t/2} + \frac{q^{4t}}{2} - \frac{q^{7t/2}}{2} + \frac{q^{3t}}{4} + \frac{q^{4t+s}}{4}$$
$$\leq q^{2k-t/2} + \frac{q^{4t+s}}{2} \leq \begin{cases} \frac{3}{2}q^{2k-1} & \text{for } t \ge 2\\ (1+q/2)q^{2k-2} & \text{for } t \ge 4\\ (1+q^2/2)q^{2k-2} & \text{for } t \ge 6\end{cases}$$
$$\leq \frac{2}{9}q^{2k} < \frac{2}{9}(q^k + q^{2t} - (q^2 + q + 1))^2,$$

for  $t \ge 6$  and  $q \ge 3$ ,  $t \ge 4$  and  $q \ge 4$  or t = 2 and  $q \ge 7$ .

#### 6.3 Main theorem

Now we are in position to prove our main result.

**Theorem 6.4.** Let  $C = \langle x^{q^t}, x + \delta x^{q^{2t}}, G(x) \rangle \subseteq \mathcal{L}_{n,q}$  be an exceptional 3-dimensional  $\mathbb{F}_{q^n}$ -linear MRD code, where t is the minimum integer such that  $x^{q^t} \in C$ . If  $(t,q) \notin \{(1,3); (1,4); (1,5); (2,3); (2,4); (2,5), (4,3)\}$ , then  $\deg_q(G(x)) < 2t$ .

*Proof.* By Theorem 3.4 and Proposition 5.1 it follows that  $\min \deg_q(G(X)) = 2t$  or t/2.

Suppose by way of contradiction that  $\deg_q(G(X)) > 2t$ ; then G(X) contains at least two terms (see Equations 6, 35). Let us consider  $\mathcal{W}, \lambda, \mathcal{A}$  as in Equations 4, 7, 9. Theorems 6.1 and 6.3 ensure the existence of an absolutely irreducible component defined over  $\mathbb{F}_{q^n}$  of  $\mathcal{A}$  not contained in the curve defined by  $F_{(x,x^q,x^{q^2})}(X,Y,\lambda) = 0$ . By Lemma 5.2,  $\mathcal{W}$  has an absolutely irreducible  $\mathbb{F}_{q^n}$ -rational component not contained in  $\mathcal{V}$ . Therefore Theorem 2.5 guarantees the existence of  $\mathbb{F}_{q^n}$ -rational points in  $\mathcal{W} \setminus \mathcal{V}$  for n large enough, and a contradiction arises from Theorem 3.4 and Proposition 4.1.

### Acknowledgments

This research was supported by the Italian National Group for Algebraic and Geometric Structures and their Applications (GNSAGA - INdAM).

## References

- Y. Aubry, G. McGuire and F. Rodier. A few more functions that are not APN infinitely often, finite fields theory and applications. *Contemporary Math.*, 518:23–31, 2010.
- [2] D. Bartoli, G. Fatabbi and F. Ghiandoni. On the exceptionality of rational APN functions. Des. Codes Cryptogr., 91(10):3167–3186, 2023.
- [3] D. Bartoli, G. Longobardi, G. Marino and M. Timpanella. Scattered trinomials of  $\mathbb{F}_{q^6}[X]$  in even characteristic. arXiv:2307.12829, 2023.
- [4] D. Bartoli, G. Marino, A. Neri, and L. Vicino. Exceptional scattered sequences. arXiv:2211.11477, 2022.
- [5] D. Bartoli and M. Timpanella. On a conjecture on APN permutations. Cryptogr. Commun., 14(4):925–931, 2022.
- [6] D. Bartoli, C. Zanella and F. Zullo. A new family of maximum scattered linear sets in  $PG(1, q^6)$ . Ars Math. Contemporanea, 19(1):125-145, 2020
- [7] D. Bartoli and Y. Zhou. Exceptional scattered polynomials. J. Algebra, 509:507–534, 2018.
- [8] D. Bartoli and Y. Zhou Asymptotics of Moore exponent sets. J. Comb. Theory Ser. A, 175:105281, 2020.
- D. Bartoli, G. Zini and F. Zullo. Linear Maximum Rank Distance Codes of Exceptional Type IEEE Tran. Inf. Theory, 69(6):3627–3636, 2023.
- [10] A. Blokhuis and M. Lavrauw. Scattered spaces with respect to a spread in PG(n,q). Geometriae Dedicata, 81(1):231-243, 2000.
- [11] A. Cafure and G. Matera. Improved explicit estimates on the number of solutions of equations over a finite field. *Finite Fields Appl.*, 12(2):155–185, 2006.
- [12] B. Csajbók, G. Marino, O. Polverino and C. Zanella. A new family of MRD-codes. *Linear Algebra Appl.*, 548:203–220, 2018.
- [13] B. Csajbók, G. Marino, O. Polverino and F. Zullo. Maximum scattered linear sets and MRD codes. J. Algebraic Comb., 46(3-4):517–531, 2017.
- [14] B. Csajbók, G. Marino, O. Polverino and F. Zullo. Generalising the scattered property of subspaces. *Combinatorica*, 41(2):237–262, 2021.
- [15] B. Csajbók, G. Marino and F. Zullo, New maximum scattered linear sets of the projective line. *Finite Fields Their Appl.*, 54:133–150, 2018.
- [16] P. Delsarte. Bilinear forms over a finite field, with applications to coding theory. J. Comb. Theory Ser. A, 25(3):226–241, 1978.

- [17] E.M. Gabidulin. Theory of codes with maximum rank distance. Problemy Peredachi Informatsii, 21(1):3–16, 1985.
- [18] G. Longobardi and C. Zanella, Linear sets and MRD-codes arising from a class of scattered linearized polynomials. J. Algebr. Combinatorics, 53:639–661, 2021.
- [19] G. Lunardon, R. Trombetti and Y. Zhou, On kernels and nuclei of rank metric codes. J. Algebr. Combinat., 46(2):313–340, 2017.
- [20] G. Lunardon, R. Trombetti and Y. Zhou, Generalized twisted Gabidulin codes. J. Comb. Theory Ser. A, 159:79–106, 2018.
- [21] W. Fulton, Algebraic curves, Advanced Book Classics, Addison-Wesley Publishing Company Advanced Book Program, Redwood City, CA, 1989.
- [22] F. Hernando and G. McGuire. Proof of a conjecture on the sequence of exceptional numbers classifying cyclic codes and APN functions. J. Algebra, 343:78–92, 2011.
- [23] J. W. P. Hirschfeld, G. Korchmáros and F. Torres. Algebraic Curves over a Finite Field. Princeton Series in Applied Mathematics, Princeton. 2008.
- [24] H. Janwa, G. McGuire and R. Wilson. Double-error correcting cyclic codes and absolutely irreducible polynomials over GF(2). J. Algebra, 178:665–676, 1995.
- [25] D. Jedlicka. APN monomials over  $GF(2^n)$  for infinitely many n. Finite Fields Appl., 13:1006–1028, 2007.
- [26] R. Koetter and F. Kschischang. Coding for errors and erasure in random network coding. IEEE Trans. Inform. Theory, 54(8):3579-3591, 2008.
- [27] A. Neri, P. Santonastaso and F. Zullo. Extending two families of maximum rank distance codes *Finite Fields Their Appl.*, 81:102045, 2022.
- [28] O. Polverino, P. Santonastaso, J. Sheekey and F. Zullo. Divisible Linear Rank Metric Codes IEEE Trans. Inf. Theory, 69(7):4528-4536, 2023.
- [29] K.-U. Schmidt and Y. Zhou. Planar functions over fields of characteristic two. J. Algebraic Comb., 40:503–526, 2014.
- [30] J. Sheekey. A new family of linear maximum rank distance codes. Adv. Math. Commun., 10(3):475-488, 2016.
- [31] M. Timpanella and G. Zini On a family of linear MRD codes with parameters  $[8 \times 8, 16, 7]_q$ , Des. Codes Cryptogr. DOI:10.1007/s10623-022-01179-0, 2023.
- [32] C. Zanella, A condition for scattered linearized polynomials involving Dickson matrices. J. Geometry, 110(3):1–9, 2019.