

Cosmological evolution of matter with interacting dust fluids

Geoffrey Okeng'o¹, Nceba Mhlahlo^{2*}, Roy Maartens^{3*}

¹Physics Department, University of Nairobi, P. O. Box 30197-00100
GPO, Nairobi, Kenya.

^{2*}Physics Department, University of the Witwatersrand, 1 Jan Smuts
Avenue, Johannesburg, 2000, South Africa.

^{3*}Department of Physics & Astronomy, University of the Western Cape,
Cape Town, 7535, South Africa.

*Corresponding author(s). E-mail(s): nmgaxamba@gmail.com;
roy.maartens@gmail.com;

Abstract

We split the total matter fluid into a bound (halo) component and an unbound (free particles) fluid component that is accreted by the halos. We adopt a different framework that treats the structure formation problem as a gravitational interaction between these virialised cold dark matter halos and the unbound inter-halo cold dark matter (and cold baryon) particles. This interaction involves in general an exchange of energy and momentum during the accretion process. We then explore the evolution of the average matter density and of large-scale structure formation, using a simplified phenomenological model that is based on results from extended Press-Schechter and N-body simulations. At high redshifts most matter is in diffuse form and is not part of the haloes. As particles are accreted by the virialised halos, the particle number density decreases and that of the bound matter increases. We also present a general analysis of the background and linear perturbations for the interacting fluids, showing in detail the energy and momentum exchange terms.

1 Introduction

In the standard Λ CDM cosmological model, at late times when radiation may be neglected, the observed large-scale structure in the universe is understood to be based

on cold dark matter (CDM) haloes which act as gravitational sinks for baryonic matter, leading to the formation of stars, galaxies and higher-level clustering of luminous matter. Here we consider a simplified matter model in which the CDM is an elementary particle (e.g. the neutralino) and the baryons are cold. We neglect the complexities associated with non-cold baryonic matter by confining our analysis to linear scales. In this scenario, the first collapsed bound (virialised) objects are CDM haloes of mass determined by the mass of the CDM particle – e.g., approximately Earth mass for a neutralino particle [1]. Subsequently, haloes grow by accretion of the unbound inter-halo particles. Moving forward in time, an increasing fraction of the total matter is captured in bound structures, with a spectrum of masses [1–6].

Analytical models of structure formation are based on a Friedmann-Lemâitre-Robertson-Walker (FLRW) background with perturbations that grow under gravitational instability. The implicit assumption involved here is that the evolving mixture of particles of different (and changing) masses may be treated as a single pressure-free ‘dust’ model, all of whose particles are considered as test particles – i.e. effectively massless particles. Standard perturbation theory does not specify the masses of the dust particles and their changing mass spectrum. The agreement between the results of the perturbative analysis and those of N-body simulations on large enough scales suggests that this assumption is reasonable. Nevertheless, it is interesting to ask what the nature of the evolving spectrum of masses is and whether and how it affects cosmic evolution and structure formation. This question can be answered by N-body simulations [1–6] but an analytical approach is also of interest. In this paper we adopt a simplified analytical approach to tackle this question (see e.g. [7–9] for more sophisticated approaches).

Studies of interacting fluids (see e.g [10]) have so far focussed on the “dark sector” interaction between dark matter and dark energy and have provided useful insights into the coincidence problem and structure formation. In this paper we explore a different framework that treats the structure formation problem as a gravitational interaction between virialised CDM halos and the unbound inter-halo CDM (and cold baryon) particles. This interaction is an exchange of energy and momentum during the accretion process. We use the results from [1], based on extended Press-Schechter formalism and Monte Carlo simulations and tested with N-body simulations. These results indicate that by redshift zero, $\sim 80 - 95\%$ of matter is within halos of all possible masses – with $\sim 60 - 70\%$ of matter in halos of mass $\gtrsim 3 \times 10^9 M_\odot$ (see their Fig. 3). The percentages depend on the details assumed in their models. These details are not important for our purposes, since our focus is on a simple analytical model that reflects the key qualitative features.

2 The 2-fluid model

The two fluids are the halo fluid $A = h$ and the free (unbound) particle fluid $A = f$, each with a dust equation of state, $w_A = 0$, and vanishing speed of sound $c_{sA} = 0$. The total matter density is

$$\rho_m = \rho_h + \rho_f, \tag{1}$$

which satisfies the background and perturbation equations of the standard Λ CDM model. For example,

$$\mathcal{H}^2 = \frac{8\pi G}{3} \bar{\rho}_m a^2 + \frac{\Lambda}{3} a^2, \quad \bar{\rho}'_m + 3\mathcal{H}\bar{\rho}_m = 0, \quad (2)$$

where the prime denotes a conformal time derivative and $\mathcal{H} = a'/a$ is the conformal Hubble rate.

We assume a simplified model for the density transfer from free particles to halos:

$$\frac{\rho_h}{\rho_m} = \mathcal{F}(\rho_m). \quad (3)$$

In other words, the fraction of mass in halos is given by a function of the total density.

2.1 Background equations

In the background,

$$\bar{\rho}'_h + 3\mathcal{H}\bar{\rho}_h = \bar{\rho}_m \bar{\mathcal{F}}', \quad (4)$$

$$\bar{\rho}'_f + 3\mathcal{H}\bar{\rho}_f = -\bar{\rho}_m \bar{\mathcal{F}}', \quad (5)$$

where we used (1) and (2). Note that $\bar{\mathcal{F}}' = -3\mathcal{H}\bar{\rho}_m \partial \bar{\mathcal{F}} / \partial \bar{\rho}_m$.

The general background equations for two interacting fluids (see e.g. the extensive review in [10]) reduce in our case to

$$\bar{\rho}'_h + 3\mathcal{H}\bar{\rho}_h = a\bar{Q}_h, \quad (6)$$

$$\bar{\rho}'_f + 3\mathcal{H}\bar{\rho}_f = a\bar{Q}_f = -a\bar{Q}_h. \quad (7)$$

Here \bar{Q}_A are the rates of energy density transfer:

$$a\bar{Q}_h = \bar{\rho}_m \bar{\mathcal{F}}' = -a\bar{Q}_f. \quad (8)$$

Clearly \bar{Q}_h is positive since the halo fluid is accreting the free-particle fluid. We can also rewrite these equations in terms of the dimensionless density parameters:

$$\Omega'_h = \frac{\bar{\mathcal{F}}'}{\bar{\mathcal{F}}} \Omega_h + 3\mathcal{H}(\Omega_h + \Omega_f - 1)\Omega_h, \quad (9)$$

$$\Omega'_f = -\frac{\bar{\mathcal{F}}'}{\bar{\mathcal{F}}} \Omega_h + 3\mathcal{H}(\Omega_h + \Omega_f - 1)\Omega_f. \quad (10)$$

2.2 Linear perturbations

Linear perturbation of (3) gives

$$\delta_h - \delta_m = \frac{\delta \mathcal{F}}{\bar{\mathcal{F}}} = \frac{\partial \ln \bar{\mathcal{F}}}{\partial \ln \bar{\rho}_m} \delta_m = \bar{\mathcal{G}} \delta_m \quad \text{where} \quad \bar{\mathcal{G}} \equiv \frac{\partial \ln \bar{\mathcal{F}}}{\partial \ln \bar{\rho}_m}. \quad (11)$$

From this we obtain

$$\delta_h = (1 + \bar{\mathcal{G}}) \delta_m, \quad (12)$$

$$\delta_f = \left(1 - \frac{\Omega_h}{\Omega_f} \bar{\mathcal{G}}\right) \delta_m. \quad (13)$$

It follows from these equations that the power spectra of the halo fluid and the free-particle fluid are

$$P_h(a, k) = [1 + \bar{\mathcal{G}}(a)]^2 P_m(a, k), \quad (14)$$

$$P_f(a, k) = \left[1 - \frac{\Omega_h(a)}{\Omega_f(a)} \bar{\mathcal{G}}(a)\right]^2 P_m(a, k), \quad (15)$$

where P_m is the matter power spectrum.

Equations (9)–(15) allow us to evade the solving of the perturbed energy-momentum conservation equations. Nevertheless, it is instructive to see how our equations correspond to the general perturbation equations for coupled fluids. In order to make this comparison, we choose the Newtonian gauge. The perturbed line element at late times is then

$$ds^2 = a^2 [-(1 + 2\Phi)d\eta^2 + (1 - 2\Phi)d\mathbf{x}^2], \quad (16)$$

where Φ is the gravitational potential. The four velocity of fluid A is

$$u_A^\mu = \frac{1}{a} (1 - \Phi, \partial^i v_A), \quad (17)$$

where v_A is the peculiar velocity potential, which vanishes in the background. The general perturbation equations for two interacting fluids are based on [10]

$$\nabla_\nu T_A^{\mu\nu} = Q_A^\mu, \quad (18)$$

where Q_A^μ is the energy-momentum transfer four-vector, which satisfies $\sum_A Q_A^\mu = 0$ from conservation of total energy-momentum. In general the four-vector Q_A^μ has the form

$$Q_A^\mu = Q_A u^\mu + F_A^\mu \quad \text{where} \quad Q_A = \bar{Q}_A + \delta Q_A \quad \text{and} \quad u_\mu F_A^\mu = 0. \quad (19)$$

Here F_A^μ is the momentum density transfer rate relative to the total four-velocity $u^\mu = a^{-1}(1 - \Phi, \partial^i v)$ and f_A is the intrinsic momentum transfer potential. It follows that

$$Q_0^A = -a [\bar{Q}_A(1 + \Phi) + \delta Q_A], \quad (20)$$

$$Q_i^A = a \partial_i [f_A + \bar{Q}_A v]. \quad (21)$$

In the case of 2 dust fluids, it is natural to choose the total 4-velocity as that of matter: $v = v_m$. Then the equations reduce to

$$\delta'_A - k^2 v_A - 3\mathcal{H}\Phi' = \frac{a\bar{Q}_A}{\bar{\rho}_A}(\Phi - \delta_A) + \frac{a}{\bar{\rho}_A} \delta Q_A, \quad (22)$$

$$v'_A + \mathcal{H}v_A + \Phi = \frac{a\bar{Q}_A}{\bar{\rho}_A}(v_m - v_A) + \frac{af_A}{\bar{\rho}_A}. \quad (23)$$

The total matter density contrast evolves according to the equation

$$\delta'_m - k^2 v_m - 3\Phi' = 0. \quad (24)$$

Our simplified model does not include the small effect of the gravitational potential on the density contrast. Since $|\Phi| \ll |\delta_A|$ on the scales of interest, we can neglect the gravitational potential in (22) and (24), so that

$$\delta'_A - k^2 v_A = \frac{a}{\bar{\rho}_A}(\delta Q_A - \delta_A), \quad (25)$$

$$\delta'_m - k^2 v_m = 0. \quad (26)$$

Since our model is confined to scales where linear perturbation theory is valid, we can ignore the velocity bias between the halos and the unbound particles. This bias arises on small nonlinear scales because the halos represent peaks in the matter distribution, with a threshold for formation [7, 9, 11]. Therefore it is reasonable to assume

$$v_A = v_m. \quad (27)$$

Consequently, there is no intrinsic momentum transfer in the matter frame:

$$f_A = 0. \quad (28)$$

With these additional simplifications, the perturbed conservation equations reduce to

$$\delta'_A - k^2 v_m = \frac{a}{\bar{\rho}_A}(\delta Q_A - \delta_A), \quad (29)$$

$$\delta'_m - k^2 v_m = 0, \quad (30)$$

$$v'_m + \mathcal{H}v_m + \Phi = 0. \quad (31)$$

By comparing (11)–(13) with (29)–(30), we find that

$$\begin{aligned} \frac{a}{\bar{\rho}_h} \delta Q_h &= \left[\bar{\mathcal{G}}' + (f - 3)\mathcal{H}\bar{\mathcal{G}} - 3\mathcal{H}\bar{\mathcal{G}}^2 \right] \delta_m \\ &= -\frac{a}{\bar{\rho}_f} \frac{\Omega_f}{\Omega_h} \delta Q_f. \end{aligned} \quad (32)$$

Here we used $\delta Q_f = -\delta Q_h$, and the linear growth rate is defined as

$$f = \frac{d \ln \delta_m}{d \ln a}, \quad (33)$$

so that $\delta'_m = f \mathcal{H} \delta_m$.




Fig. 1 The energy density parameters for the bound (halo) fluid h and the unbound (free-particle) fluid f , shown together with the total matter energy density Ω_m for the model (34)

3 A simple model

A simple power-law fit to the results of [1] (their Fig. 3, ellipsoidal collapse model) is

$$\mathcal{F} = \mathcal{F}_0 \left(\frac{\rho_m}{\rho_{m0}} \right)^{-\alpha [(\rho_m/\rho_{m0})^{1/3} - 1]}, \quad (34)$$

where the constants are $\mathcal{F}_0 = 0.9$ and $\alpha = 0.07$. Note that in the background, $\bar{\mathcal{F}} = \mathcal{F}_0(1+z)^{-3\alpha z}$, where z is the cosmological redshift. The background equations from 9 and 10 now become

$$\frac{d\Omega_h}{da} = \frac{3}{a^2} \alpha (1-a-\ln a) \Omega_h + \frac{3}{a} (\Omega_m - 1) \Omega_h \quad (35)$$

$$\frac{d\Omega_f}{da} = -\frac{3}{a^2} \alpha (1-a-\ln a) \Omega_f + \frac{3}{a} (\Omega_m - 1) \Omega_f \quad (36)$$




Fig. 2 Plots for the functions $\mathcal{F}(z)$ (top) and $\mathcal{F}(a)$ (bottom) for the power law. Here $\mathcal{F}_0 = \rho_{h0}/\rho_{m0} = 0.9$ from our fitting

The numerical integration of these equations leads to the background energy density parameters for this model, which are shown in Figure 1. The plot shows how the matter density of the bound structures grows while that of the unclustered matter decreases over cosmic time. This growth and decay in matter densities happens until $\Omega_h - \Omega_f$ equality is reached at $a \sim 0.07$, when $\Omega_h = \Omega_f \sim 0.5$, which is when the clustered matter existed in the same quantity as the unclustered matter. Past this time, Ω_h continues to grow while Ω_f continues with its decline. The total matter energy density, Ω_m , remains nearly constant, up to $a \sim 0.15$, where it starts to decay due to the effect of Dark Energy, before reaching its current value of ~ 0.25 at $a \sim 1$. Bound matter reaches its highest density ($\Omega = 0.8$) when the redshift is ~ 3 .

Plots of the function 34 are displayed in Figure 2 where the behaviour of the mass halo fraction with respect to the redshift and the scale factor over cosmic time can be observed. The fraction of all matter predicted to be in haloes at $z = 0$, i.e. to be part of clustered matter, is close to hundred per cent. However, at high redshifts or early

times a much higher fraction of Dark Matter particles is in diffuse form and is not part of the haloes.

4 Discussion

We have presented a new formalism to study structure formation by treating the total matter fluid as composed of two components. We have derived the background and perturbation equations for the case of two interacting fluids and solved them when the ratio between the bound and unbound matter is modeled as power law, approximating the results from the extended Press-Schechter formalism and N-body simulations [1]. We have modelled the energy transfer rate using the power law model of the halo mass fraction \mathcal{F} .

Our model results, on the one hand, show that clustered matter grows due to accretion of DM particles, while the unclustered matter density decays over cosmic time. On the other hand, the total matter energy density, Ω_m , remains nearly constant for a large part of the cosmic evolution, and decays at late times, due to the effect of Dark Energy.

The halo mass fraction, \mathcal{F} , grows over cosmic time. This means that the fraction of all matter predicted to be in diffuse form is much larger at early time while that predicted to be in gravitationally bound structures is much larger at late times, close to hundred per cent at the current epoch. At low values of a a large fraction of DM particles is free, up to Ω_h - Ω_f equality, beyond which a much higher fraction of DM particles has become part of haloes. These results show that in the framework of hierarchical structure formation, gravitationally bound structures can form not only from the assembly of smaller haloes, but directly from free diffuse DM particles (see also [1]).

Acknowledgments RM was supported by the South African Radio Astronomy Observatory and National Research Foundation (grant 75415). NM acknowledges support by the NRF (grant 111735) and the Anderson-Capelli fund administered by the Wits DVC office.

References

- [1] Angulo, R.E., White, S.D.M.: The birth and growth of neutralino haloes. *Mon. Not. Roy. Astron. Soc.* **401**(3), 1796–1803 (2010) <https://doi.org/10.1111/j.1365-2966.2009.15742.x> [arXiv:0906.1730](https://arxiv.org/abs/0906.1730) [astro-ph.CO]
- [2] Chan, K.C., Scoccimarro, R., Sheth, R.K.: Gravity and large-scale nonlocal bias. *Phys. Rev. D* **85**(8), 083509 (2012) <https://doi.org/10.1103/PhysRevD.85.083509> [arXiv:1201.3614](https://arxiv.org/abs/1201.3614) [astro-ph.CO]

- [3] van Daalen, M.P., Schaye, J.: The contributions of matter inside and outside of haloes to the matter power spectrum. *Mon. Not. Roy. Astron. Soc.* **452**(3), 2247–2257 (2015) <https://doi.org/10.1093/mnras/stv1456> arXiv:1501.05950 [astro-ph.CO]
- [4] Shattow, G.M., Croton, D.J.: Beyond the halo: redefining environment with unbound matter in N-body simulations. *Mon. Not. Roy. Astron. Soc.* **452**(2), 1779–1791 (2015) <https://doi.org/10.1093/mnras/stv1464> arXiv:1507.05656 [astro-ph.CO]
- [5] Angulo, R.E., Hahn, O., Ludlow, A.D., Bonoli, S.: Earth-mass haloes and the emergence of NFW density profiles. *Mon. Not. Roy. Astron. Soc.* **471**(4), 4687–4701 (2017) <https://doi.org/10.1093/mnras/stx1658> arXiv:1604.03131 [astro-ph.CO]
- [6] Zavala, J., Frenk, C.S.: Dark Matter Haloes and Subhaloes. *Galaxies* **7**(4), 81 (2019) <https://doi.org/10.3390/galaxies7040081> arXiv:1907.11775 [astro-ph.CO]
- [7] Biagetti, M., Desjacques, V., Kehagias, A., Riotto, A.: Halo velocity bias. *Phys. Rev. D* **90**(10), 103529 (2014) <https://doi.org/10.1103/PhysRevD.90.103529> arXiv:1408.0293 [astro-ph.CO]
- [8] Biagetti, M., Desjacques, V., Kehagias, A., Racco, D., Riotto, A.: The halo Boltzmann equation. *JCAP* **2016**(4), 040 (2016) <https://doi.org/10.1088/1475-7516/2016/04/040> arXiv:1508.07330 [astro-ph.CO]
- [9] Matsubara, T.: Velocity bias and the nonlinear perturbation theory of peaks. *Phys. Rev. D* **100**(8), 083504 (2019) <https://doi.org/10.1103/PhysRevD.100.083504> arXiv:1907.13251 [astro-ph.CO]
- [10] Välviviita, J., Majerotto, E., Maartens, R.: Large-scale instability in interacting dark energy and dark matter fluids. *JCAP* **2008**(7), 020 (2008) <https://doi.org/10.1088/1475-7516/2008/07/020> arXiv:0804.0232 [astro-ph]
- [11] Baldauf, T., Desjacques, V., Seljak, U.: Velocity bias in the distribution of dark matter halos. *Phys. Rev. D* **92**(12), 123507 (2015) <https://doi.org/10.1103/PhysRevD.92.123507> arXiv:1405.5885 [astro-ph.CO]