

Minimum scaling model and exact exponents for the Nambu-Goldstone modes in the Vicsek Model

Harukuni Ikeda*

Department of Physics, Gakushuin University, 1-5-1 Mejiro, Toshima-ku, Tokyo 171-8588, Japan
(Dated: March 5, 2024)

We investigate the scaling behavior of Nambu-Goldstone (NG) modes in the ordered phase of the Vicsek model, introducing a phenomenological equation of motion (EOM) incorporating a previously overlooked non-linear term. This term arises from the interaction between velocity fields and density fluctuations, leading to new scaling behaviors. We derive exact scaling exponents in two dimensions, which reproduce the isotropic scaling behavior reported in a prior numerical simulation.

Introduction.— When a continuous symmetry is broken spontaneously, there emerge soft modes known as Nambu-Goldstone (NG) modes [1, 2]. It is known that continuous symmetry is not broken in the equilibrium state of two-dimensional models with short-range interactions since strong fluctuation along the NG mode destroys the long-range order. This principle is widely recognized as the Hohenberg-Mermin-Wagner (HMW) theorem [3, 4].

The HMW theorem does not hold out of equilibrium. Several out-of-equilibrium systems show continuous symmetry breaking even in $d = 2$ [5–13]. A prototypical example is the Vicsek model [5]. The model consists of XY spins moving at a constant speed in their orientation direction, which mimics the flocking behavior of active matter such as birds and bacteria [5, 14]. Numerical simulations in $d = 2$ revealed that the model undergoes a continuous symmetry breaking from the disordered phase, where the mean velocity of the spins vanishes, $\langle \mathbf{v} \rangle = 0$, to the ordered phase, where $\langle \mathbf{v} \rangle \neq 0$ [5].

In 1995, Toner and Tu proposed a hydrodynamic theory to model the flocking behavior of active matter, such as the Vicsek model [15]. Their theory (TT95) successfully explained the existence of the ordered phase of the Vicsek model in $d = 2$. Furthermore, they derived the exact scaling behavior of the NG modes in $d = 2$, using an elegant argument combining a pseudo-Galilean transformation and perturbative renormalization group methods [15]. Moreover, a recent more detailed investigation has shown that their results also become exact in $d > 2$, if the system can be considered as *incompressible* fluids [16]. However, for compressible fluids, subsequent re-analysis of the hydrodynamic theory by Toner, one of the authors of TT95, pointed out that several additional terms were missing in the original theory [17]. These terms hinder the exact calculations of the scaling exponents even in $d = 2$. Consequently, the applicability of TT95 to compressible fluids, like the Vicsek model, remains uncertain [17].

Numerical studies of the Vicsek model for moderate system size $N \sim 10^6$ reported results consistent with TT95 [18, 19]. However, a recent extensive numerical simulation for larger system size $N \sim 10^9$ uncovered

scaling behaviors that is distinct from the predictions of TT95. Notably, the simulation indicates almost isotropic scaling in the ordered phase [20], while TT95 predicts anisotropic scaling [15]. Similar isotropic scaling was observed in a recent experiment of Janus colloidal particles in an AC electric field [21].

Here, we reconcile the above discrepancy between the theory and numerical simulation. We introduce a simple phenomenological equation of motion (EOM) for the NG modes that dominate the scaling behavior in the ordered phase [1, 2]. A critical insight is that the symmetry of NG modes differs from the velocity field due to the hybridization of the velocity field and density fluctuations. Consequently, our EOM incorporates an additional non-linear term that was overlooked in previous studies. We calculate the exact scaling exponents of the EOM in $d = 2$. The new scaling exponents obtained by this work confirm the isotropic scaling observed in the recent numerical simulation of the Vicsek model [20].

Previous works.— Here, we briefly summarize previous theoretical and numerical studies. We first discuss the *minimal* hydrodynamic theory introduced and investigated in Ref. [22]:

$$\frac{\partial \mathbf{v}}{\partial t} + \lambda(\mathbf{v} \cdot \nabla)\mathbf{v} = \nabla^2 \mathbf{v} - \frac{\delta F[\mathbf{v}]}{\delta \mathbf{v}} + \boldsymbol{\xi}, \quad (1)$$

where $\boldsymbol{\xi}$ denotes the thermal white noise, and F denotes the standard quartic potential $F[\mathbf{v}] = a(\mathbf{v} \cdot \mathbf{v})/2 + b(\mathbf{v} \cdot \mathbf{v})^2/4$ with $b > 0$. Here, $a > 0$ and $a < 0$ represent the disordered and ordered phases, respectively [23]. When the advection λ is zero and in $d = 2$, Eq. (1) reduces to the standard Langevin equation for the XY model and only shows a quasi-long-range order, which disappears in the thermodynamic limit [24, 25]. With advection $\lambda \neq 0$, on the contrary, the model exhibits the true long-range order [22] [26]. A significant benefit of the minimal model (1) is that the analysis in TT95 becomes exact for this model. Numerical results of (1) in the ordered phase indeed well agree with TT95 [22].

Compared to the full hydrodynamic theory by Toner and Tu [15], the minimal model (1) neglects the effects of the density fluctuations $\delta\rho$, which appear through the

pressure-gradient forces such as ∇P and $\mathbf{v}(\mathbf{v} \cdot \nabla)P$. Here, the pressure P depends on both $\delta\rho$ and \mathbf{v} . Due to this complexity, the renormalization group calculations of the full hydrodynamic theory are extremely challenging, and exact scaling exponents are yet to be known even in $d = 2$ [17]. In this work, instead, we propose a simple phenomenological EOM for the NG modes, which can be treated exactly in $d = 2$, as we will see below.

New phenomenological EOM.— When the rotational symmetry is broken in a d -dimensional model, the number of the corresponding NG modes is $d - 1$ [1, 2]. Let $\{u_\mu(\mathbf{x}, t)\}_{\mu=1, \dots, d-1}$ represent magnitudes of the NG modes. We shall assume that the dynamics of u_μ is described by the following phenomenological EOM:

$$\dot{u}_\mu = \nabla^2 u_\mu + f_{\text{nl}}[u_\mu] + \xi_\mu, \quad (2)$$

where f_{nl} denotes (unknown) non-linear terms, and ξ_μ denotes the white noise of zero mean and variance

$$\langle \xi_\mu(\mathbf{x}, t) \xi_\nu(\mathbf{x}', t') \rangle = 2T \delta_{\mu\nu} \delta(\mathbf{x} - \mathbf{x}') \delta(t - t'). \quad (3)$$

Let us first consider the linear model with $f_{\text{nl}} = 0$. To investigate the large spatio-temporal scaling behavior, we apply the following scaling transformations [15, 27]:

$$\mathbf{x}_\perp \rightarrow l\mathbf{x}_\perp, \quad x_\parallel \rightarrow l^\zeta x_\parallel, \quad t \rightarrow l^z t, \quad u_\mu \rightarrow l^\chi u_\mu, \quad (4)$$

where and hereafter the subscripts \parallel and \perp denote components of any vector parallel and perpendicular to the mean velocity $\langle \mathbf{v} \rangle$, respectively. Using (3), we find the scaling dimension of the noise as $\xi_\mu \rightarrow l^{-(z+\zeta+d-1)/2} \xi_\mu$ [27]. Then, the EOM reduces to

$$l^{\chi-z} \dot{u}_\mu = l^{\chi-2} \nabla_\perp^2 u_\mu + l^{\chi-2\zeta} \partial_\parallel^2 u_\mu + l^{-(z+\zeta+d-1)/2} \xi_\mu. \quad (5)$$

Requiring that the all terms have the same scaling dimension, we get the following scaling relations:

$$\chi - z = \chi - 2 \quad (6)$$

$$\chi - z = \chi - 2\zeta \quad (7)$$

$$\chi - z = -\frac{z + \zeta + d - 1}{2}, \quad (8)$$

leading to [15, 27]

$$z = 2, \quad \zeta = 1, \quad \chi = \frac{2-d}{2}. \quad (9)$$

For $d < 2$, the fluctuation $\langle \delta u^2 \rangle \sim l^{2\chi}$ diverges in the limit $l \rightarrow \infty$, implying that the fluctuations of the NG modes destroy the long-range order. Therefore, the lower critical dimension for continuous symmetry breaking is $d_{\text{low}} = 2$. The result is consistent with the HMW theorem [3, 4, 28].

Next, we investigate the effects of the non-linear terms. The analysis can be simplified in $d = 2$ because there is

only one NG mode $u_\mu = u$. Above the lower critical dimension, $u \sim l^\chi$, $\partial_\perp \sim l^{-1}$, and $\partial_\parallel \sim l^{-\zeta}$ vanish in the thermodynamic limit $l \rightarrow \infty$, enabling the expansion of the non-linear term f_{nl} by u and its spatial derivatives, $\partial_{\perp\text{or}\parallel} u$, $\partial_{\perp\text{or}\parallel}^2 u$, \dots . Since the restoring force of the NG mode should vanish for spatially uniform deformations, each term in f_{nl} should involve at least one spatial derivative $\partial_{\perp\text{or}\parallel}$ [29]. Therefore, f_{nl} can be expanded as

$$f_{\text{nl}}[u, \partial_{\perp\text{or}\parallel} u, \partial_{\perp\text{or}\parallel}^2 u, \dots] = c_1 u \partial_\perp u + c_2 u \partial_\parallel u + O\left(u^2 \partial_{\perp\text{or}\parallel} u, u (\partial_{\perp\text{or}\parallel} u)^2, u \partial_{\perp\text{or}\parallel}^2 u\right), \quad (10)$$

where c_1 and c_2 are constants. The higher-order terms have smaller scaling dimensions and are irrelevant for $l \gg 1$. In summary, the EOM in $d = 2$ reduces to

$$\dot{u} \approx (\partial_\perp^2 + \partial_\parallel^2) u + c_1 u \partial_\perp u + c_2 u \partial_\parallel u + \xi. \quad (11)$$

Below, we shall argue that the non-linear terms stabilize the long-range order and enable the continuous symmetry breaking in $d = 2$.

Exact exponents in two dimensions.— For the minimal hydrodynamic theory (1), the NG mode is identified with the velocity component orthogonal to the order parameter $u = v_\perp$. In this case, the EOM must remain invariant under the transformation $x_\perp \rightarrow -x_\perp$ and $v_\perp \rightarrow -v_\perp$, necessitating $c_2 = 0$ [30]. This specific case, (11) with $c_2 = 0$, has been well investigated in previous studies [15, 30]. For completeness, we will briefly outline the derivation of the scaling exponents. First, note that the non-linear term can be rewritten as a total derivative with respect to ∂_\perp as $c_1 u \partial_\perp u = \partial_\perp (c_1 u^2/2)$. A perturbative calculation of this term generates terms such as $\partial_\perp(\text{some function})$, which only affect the scaling dimensions of terms involving ∂_\perp . Therefore, the scaling dimensions $\dot{u} \sim l^{\chi-z}$, $\partial_\parallel^2 u \sim l^{\chi-2\zeta}$, and $\xi \sim l^{-(z+\zeta+d-1)/2}$ remain unchanged. Requiring that those three terms have the same scaling dimension, we get the scaling relations (7), and (8) for $d = 2$. Next, note that (11) with $c_2 = 0$ is invariant under the pseudo-Galilean transformation: $u \rightarrow u + U$, $x_\perp \rightarrow x_\perp + c_1 U t$, which implies the following scaling relation [15, 31]:

$$1 = \chi + z. \quad (12)$$

Using (7), (8), and (12), we get [30]

$$z = \frac{2(1+d)}{5}, \quad \zeta = \frac{d+1}{5}, \quad \chi = \frac{3-2d}{5}, \quad (13)$$

which are consistent with TT95 [15]. Note that the scaling relations have been proven only in $d = 2$, and it is not clear if the exponents (13) become exact in $d > 2$. Remarkably, a recent detailed investigation has established that the exponents (13) become exact for incompressible fluids $\nabla \cdot \mathbf{v} = 0$ in $d > 2$ [16]. The scaling behaviors (13)

TABLE I. Scaling exponents. Data are taken from Ref [20].

	$d = 2$			$d = 3$		
	Vicsek	TT95	This work	Vicsek	TT95	This work
χ	-0.31(2)	-0.2	-0.33	-0.62	-0.6	-0.67
ζ	0.95(2)	0.6	1	1	0.8	1
z	1.33(2)	1.2	1.33	1.77	1.6	1.67
α	0.84(1)	0.8	0.83	0.79(2)	0.77	0.78

imply $\langle \delta v_{\perp}^2 \rangle \sim l^{2\chi} \rightarrow 0$ in $d = 2$, meaning that the non-linear term stabilizes the long-range order in $d = 2$ [15]. The exponents (13) well agree with a numerical simulation of the minimal model (1) in $d = 2$ [22].

Let us turn to the Vicsek model. It is crucial to note that the velocity field generally couples to the density fluctuations [17], and hence, the NG mode is not identical to v_{\perp} . In other words, there is no reason to assume $c_2 = 0$. Below, we calculate the scaling exponents for the case $c_2 \neq 0$. In (11), the non-linear terms proportional to c_i can be rewritten as the total derivatives with respect to ∂_{\parallel} and ∂_{\perp} , meaning that the perturbative renormalization group calculations yield no corrections to the scaling of \dot{u} and ξ [15, 20, 30]. Requiring $\dot{u} \sim \xi$, we get (8) for $d = 2$ [20]. This scaling relation is often referred to as the hyperscaling and has been confirmed in the recent numerical simulation of the Vicsek model [20]. Note also that (11) is invariant under the pseudo-Galilean transformation: $u \rightarrow u + U$, $x_{\perp} \rightarrow x_{\perp} + c_1 U t$, and $x_{\parallel} \rightarrow x_{\parallel} + c_2 U t$, which implies the following scaling relations [15, 31]

$$1 = \chi + z, \quad (14)$$

$$\zeta = \chi + z. \quad (15)$$

Using (8), (14), and (15), we can determine the scaling exponents:

$$z = \frac{d+2}{3}, \quad \zeta = 1, \quad \chi = \frac{1-d}{3}. \quad (16)$$

The results are consistent with our previous heuristic scaling argument [32]. Given that (11) is the most general form of EOM taking into account the lowest order non-linear terms, we conclude that (16) are the exact scaling exponents of the Vicsek model in $d = 2$. In $d > 2$, it is unclear if the EOM satisfies the scaling relations (8), (14), and (15), and we can not rule out the possibility that some corrections appear to the exponents (16).

Comparison with numerical simulation.— In Table I, we compare the numerical results of the Vicsek model [20], TT95 [15], and our theoretical predictions (16). Overall, our results show better agreement with numerical results than that of TT95. In particular, we get an almost perfect agreement for $d = 2$. For $d = 3$, there are small discrepancies between the numerical and theoretical results. We expect that further theoretical [15, 30, 31, 33–35], and numerical studies [20, 36] will




FIG. 1. Correlation functions. Markers denote the numerical results of the Vicsek model in $d = 2$ for $L = 8000$, taken from Fig.1 (a) in Ref. [20]. The dashed and dotted lines represent theoretical predictions.

lead to more accurate estimates of the scaling exponents in $d = 3$.

Another important physical quantity is the correlation function $C(\mathbf{x}) = \langle \mathbf{v}_{\perp}(\mathbf{x}) \cdot \mathbf{v}_{\perp}(0) \rangle$. The scaling behaviors $\mathbf{x}_{\perp} \sim l$, $x_{\parallel} \sim l^{\zeta}$, and $\mathbf{v}_{\perp} \sim u \sim l^{\chi}$ lead to

$$C(\mathbf{x}) \sim l^{2\chi} C(l^{-1} \mathbf{x}_{\perp}, l^{-\zeta} x_{\parallel}) \sim |\mathbf{x}_{\perp}|^{2\chi} \sim x_{\parallel}^{2\chi/\zeta}. \quad (17)$$

In the Fourier space, it leads to [15, 31, 37]:

$$\begin{aligned} \tilde{C}(\mathbf{q}) &= \int d\mathbf{x} e^{i\mathbf{q} \cdot \mathbf{x}} C(\mathbf{x}) \\ &\sim l^{2\chi + \zeta + d - 1} \tilde{C}(l\mathbf{q}_{\perp}, l^{\zeta} q_{\parallel}) \sim |\mathbf{q}_{\perp}|^{-z} \sim q_{\parallel}^{-z/\zeta}. \end{aligned} \quad (18)$$

To precisely calculate $\tilde{C}(\mathbf{q})$, one should perform a numerical simulation for a sufficiently large linear system size L because the minimal wave vector q_* scales as $q_* \sim 2\pi/L$. The best numerical result for the Vicsek model has been obtained in $d = 2$ for $L = 8000$ in Ref. [20]. In Fig. 1, we compare the numerical result, our theoretical prediction, and that of TT95 [15, 37]. Our theory shows a better agreement with the numerical results than TT95.

In the ordered phase of the Vicsek model, the density fluctuations are significantly enhanced due to the NG modes. This phenomenon is referred to as the giant number fluctuations (GNF) [38]. The GNF is popularly studied in numerical and experimental studies [20, 21, 38]. So, we here derive the scaling exponent α that characterizes the strength of the GNF. To be more precise, we shall observe the number fluctuations in a box of linear size l :

$$\begin{aligned} \langle \delta N^2 \rangle &= \left\langle \left(\int_{[0,l]^d} d\mathbf{x} \delta \rho(\mathbf{x}) \right)^2 \right\rangle \\ &= l^d \int_{[0,l]^d} d\mathbf{x} \langle \delta \rho(\mathbf{x}) \delta \rho(0) \rangle. \end{aligned} \quad (19)$$

We assume that the density fluctuations are dominated by the NG modes, $\delta\rho \sim u$, and the correlation function $C_\rho(\mathbf{x}) = \langle \delta\rho(\mathbf{x})\delta\rho(0) \rangle$ satisfies $C_\rho(\mathbf{x}_\perp, x_\parallel) = l^{2\chi} C_\rho(l^{-1}\mathbf{x}_\perp, l^{-\zeta}x_\parallel)$. Then, the scaling analysis leads to $\int d\mathbf{x} C_\rho(\mathbf{x}) \sim l^{2\chi+\zeta+d-1} \sim l^z$. Substituting this into (19), we get

$$\langle \delta N^2 \rangle \sim l^{d+z} \sim (N/\rho)^{2\alpha}, \quad (20)$$

where $\rho = N/l^d$ denotes the mean-density, and we have introduced the exponent [15, 37]

$$\alpha = \frac{1}{2} + \frac{z}{2d}. \quad (21)$$

The central limit theorem predicts $\alpha = 1/2$. In the ordered phase of the Vicsek model, one has $\alpha > 1/2$: the fluctuations grow much faster than the naive expectation of the central limit theorem [38]. This anomalous enhancement of the density fluctuations is the signature of the GNF [38]. In Table I, we compare the numerical result [20, 38], our theoretical prediction, and that of TT95 [15]. In $d = 2$, our theory shows a better agreement with the numerical result than that of TT95. In $d = 3$, the predictions of the two theories are so close that we can not judge which theory is more precise from the current numerical data. Extensive numerical simulations for larger system sizes are required to elucidate this point.

Summary.— In summary, we have investigated the NG modes of the Vicsek model in the ordered phase. We derived a phenomenological EOM of the NG modes and investigated its scaling behavior. Our essential observation is that the NG modes should not be simply identified with the vertical components of the velocity field but should depend both on the velocity field and the density fluctuations. As a consequence, the EOM has an additional non-linear term overlooked in the previous studies, which leads to a new scaling behavior. We calculated the exponents that characterize the scaling of the NG modes. In particular, the pseudo-Galilean invariance allows us to calculate the exact scaling exponents in $d = 2$. We found that our results agree well with a recent numerical simulation of the Vicsek model in $d = 2$.

Although we derived the EOM by a phenomenological argument, the resultant EOM incorporates all the relevant non-linear terms, making it quite universal. Therefore, we expect that the EOM can be applied to other two-dimensional systems that exhibit long-range order, as well as the Vicsek model. For instance, two-dimensional long-range orders have been reported by recent experiments with bacteria and Janus particles [14, 21]. It is an important future work to check if these systems follow the scaling laws found in this work.

After we completed the previous version of the manuscript (arXiv:2401.01603), P. Jentsch and C. F. Lee submitted a preprint addressing the same problem of the scaling exponents of the Vicsek model

(arXiv:2402.01316). They introduced and investigated a simplified version of the Toner-Tu hydrodynamic theory [15, 17, 37]. Their renormalization group calculations for the simplified hydrodynamic theory predict weak anisotropy: $\zeta = 0.975$ in $d = 2$. It remains an intriguing question whether similar calculations for the full hydrodynamic theory can reproduce the isotropic scaling behavior $\zeta = 1$.

We thank H. Tasaki for his careful reading of the manuscript and constructive comments. We also thank H. Nakano, Y. Kuroda, and D. Nishiguchi for helpful discussions. This work was supported by KAKENHI 23K13031.

* harukuni.iked@gakushuin.ac.jp

- [1] Y. Nambu, Phys. Rev. **117**, 648 (1960).
- [2] J. Goldstone, A. Salam, and S. Weinberg, Phys. Rev. **127**, 965 (1962).
- [3] P. C. Hohenberg, Phys. Rev. **158**, 383 (1967).
- [4] N. D. Mermin and H. Wagner, Phys. Rev. Lett. **17**, 1133 (1966).
- [5] T. Vicsek, A. Czirók, E. Ben-Jacob, I. Cohen, and O. Shochet, Phys. Rev. Lett. **75**, 1226 (1995).
- [6] K. E. Bassler and Z. Rácz, Phys. Rev. E **52**, R9 (1995).
- [7] M. D. Reichl, C. I. Del Genio, and K. E. Bassler, Phys. Rev. E **82**, 040102 (2010).
- [8] S. A. M. Loos, S. H. L. Klapp, and T. Martynec, Phys. Rev. Lett. **130**, 198301 (2023).
- [9] L. P. Dadhichi, J. Kethapelli, R. Chajwa, S. Ramaswamy, and A. Maitra, Phys. Rev. E **101**, 052601 (2020).
- [10] L. Galliano, M. E. Cates, and L. Berthier, Phys. Rev. Lett. **131**, 047101 (2023).
- [11] H. Ikeda, Phys. Rev. E **108**, 064119 (2023).
- [12] H. Ikeda and Y. Kuroda, arXiv preprint arXiv:2304.14235 (2023).
- [13] H. Ikeda, arXiv preprint arXiv:2309.03155 (2023).
- [14] D. Nishiguchi, K. H. Nagai, H. Chaté, and M. Sano, Phys. Rev. E **95**, 020601 (2017).
- [15] J. Toner and Y. Tu, Phys. Rev. Lett. **75**, 4326 (1995).
- [16] L. Chen, C. F. Lee, and J. Toner, New Journal of Physics **20**, 113035 (2018).
- [17] J. Toner, Phys. Rev. E **86**, 031918 (2012).
- [18] Y. Tu, J. Toner, and M. Ulm, Phys. Rev. Lett. **80**, 4819 (1998).
- [19] F. Giavazzi, C. Malinverno, S. Corallino, F. Ginelli, G. Scita, and R. Cerbino, Journal of Physics D: Applied Physics **50**, 384003 (2017).
- [20] B. Mahault, F. Ginelli, and H. Chaté, Phys. Rev. Lett. **123**, 218001 (2019).
- [21] J. Iwasawa, D. Nishiguchi, and M. Sano, Phys. Rev. Res. **3**, 043104 (2021).
- [22] M. Besse, H. Chaté, and A. Solon, Phys. Rev. Lett. **129**, 268003 (2022).
- [23] H. Nishimori and G. Ortiz, *Elements of phase transitions and critical phenomena* (Oxford university press, 2011).
- [24] J. M. Kosterlitz and D. Thouless, Journal of Physics C: Solid State Physics **5**, L124 (1972).
- [25] J. M. Kosterlitz and D. J. Thouless, in *Basic Notions Of Condensed Matter Physics* (CRC Press, 1994) pp. 493–

- 515.
- [26] The stability of the ordered phase of the minimal model remains under debate. A numerical simulation reported nucleation that destabilizes the ordered phase for moderate a [22]. However, the characteristic timescale of the nucleation rapidly increases as a decreases, making it difficult to discuss whether similar nucleation occurs for small a .
- [27] E. Medina, T. Hwa, M. Kardar, and Y.-C. Zhang, *Phys. Rev. A* **39**, 3053 (1989).
- [28] H. Tasaki, *Phys. Rev. Lett.* **125**, 220601 (2020).
- [29] A. Altland and B. D. Simons, *Condensed matter field theory* (Cambridge university press, 2010).
- [30] P. Sartori and C. F. Lee, *New Journal of Physics* **21**, 073064 (2019).
- [31] J. Toner, Y. Tu, and S. Ramaswamy, *Annals of Physics* **318**, 170–244 (2005).
- [32] H. Ikeda, arXiv preprint arXiv:2401.01603 (2024).
- [33] L. Chen, C. F. Lee, and J. Toner, *Nature communications* **7**, 12215 (2016).
- [34] L. Di Carlo and M. Scandolo, *New Journal of Physics* **24**, 123032 (2022).
- [35] P. Jentsch and C. F. Lee, *Phys. Rev. Res.* **5**, 023061 (2023).
- [36] G. Grégoire and H. Chaté, *Phys. Rev. Lett.* **92**, 025702 (2004).
- [37] J. Toner and Y. Tu, *Phys. Rev. E* **58**, 4828 (1998).
- [38] D. Nishiguchi, *Journal of the Physical Society of Japan* **92** (2023), 10.7566/jpsj.92.121007.