

The spectrum of the Corona of Hypergraphs

Liya Jess Kurian^{1*}, Chithra A. V^{1†}

1 Department of Mathematics, National Institute of Technology Calicut,
Calicut-673 601, Kerala, India

Abstract

The corona of hypergraphs is an extension of the corona operation applied to graphs. The corona $G_0^* \odot_1^n G_1^*$ of two hypergraphs is obtained by taking n copies of G_1^* (where n is the order of G_0^*) and by joining the i -th vertex of G_0^* with the i -th copy of G_1^* . In this paper, we estimate the complete spectrum (adjacency and Seidel) and eigenvectors of the corona $G_0^* \odot_1^n G_1^*$ of two hypergraphs when G_1^* is regular. Additionally, we define the corona hypergraph $G_0^{*(m)} = G_0^{*(m-1)} \odot_1^n G_0^*$ and determined its adjacency spectrum. Also, we extend the definition coronal of the adjacency matrix. Moreover, we estimate the characteristic polynomial of Seidel matrix of the generalised corona of hypergraphs. Applying these results, we obtain infinitely many non-regular non-isomorphic adjacency and Seidel cospectral hypergraphs.

Keywords: Seidel matrix, adjacency matrix, hypergraph, (k, r) -regular hypergraph, corona.
MSC 05C65 05C50 15A18

1 Introduction

Hypergraph is the generalisation of a graph in which hyperedge can connect more than 2 vertices at a time. In this article, we study hypergraphs that are simple and finite. More precisely, a hypergraph $G^* = (V, E)$ of order n with a vertex set $V(G^*) = \{v_1, v_2, v_3, \dots, v_n\}$ and a collection $E(G^*) = \{e_1, e_2, e_3, \dots, e_m\}$ of hyperedges such that each hyperedge is a non-empty subset ($|e_i| \geq 2$) of the vertex set V . The degree $d(v)$ of a vertex $v \in V$ is defined as the number of hyperedges that include the vertex v . A hypergraph G^* is said to be a k -uniform hypergraph [5, 10] if each of its hyperedges contains exactly k vertices, where $k \geq 2$. A hypergraph with $d(v_i) = r$ for all $v_i \in V$ is called a r -regular hypergraph. A hypergraph is said to be a (k, r) -regular hypergraph if it is both k -uniform and r -regular. Hypergraphs that are 2-uniform and 2-regular are referred to as cycles in graph theory. The properties of (k, r) regular hypergraph are studied in [10]. A k -uniform hypergraph of order n is said to be a complete k -uniform hypergraph K_n^k if every possible k subset of the vertex set forms a hyperedge [4]. Let G_1^* and G_2^* be two k -uniform hypergraphs with vertex sets $V_1(G_1^*)$ and $V_2(G_2^*)$ and edge sets $E_1(G_1^*)$ and $E_2(G_2^*)$, respectively. Then the join of two k -uniform hypergraphs [13] G_1^* and G_2^* is also a k -uniform hypergraph $G^* = G_1^* \oplus G_2^*$ with vertex set $V(G^*) = V_1(G_1^*) \cup V_2(G_2^*)$ and edge set $E(G^*) = E_0 \cup \bigcup_{i=1}^2 E_i(G_i^*)$, where E_0 contains all possible k -subsets of V satisfying $e \cap V_i \neq \phi, \forall i = 1, 2, e \in E_0$. Let $G_1^*[V_1']$ be the subhypergraph induced by the vertex set $V_1' \subset V_1(G_1^*)$. Then $G_1^*[V_1'] \otimes G_2^*$ is a hypergraph with vertex set $V_1(G_1^*) \cup V_2(G_2^*)$ and edge set contains the hyperedges of G_1^* together with hyperedges of $G_1^*[V_1'] \oplus G_2^*$.

*liyajess@gmail.com

†chithra@nitc.ac.in

Research on the spectrum of hypergraphs has gained significant attention. In 2012, Cooper and Dutle [5] suggested tensors (hypermatrices) to analyse the spectral characteristics of hypergraphs, which gives analogous results to spectral graph theory. Here, we consider adjacency and Seidel matrices associated with the hypergraph. The adjacency matrix $A(G^*) = (a_{ij})$ of G^* [11] is an $n \times n$ matrix whose rows and columns are indexed by the vertices of G^* and for all $v_i, v_j \in V$,

$$a_{ij} = \begin{cases} |\{e_k \in E(G^*) : \{v_i, v_j\} \subset e_k\}| & , v_i \neq v_j, k = 1, 2, 3, \dots, m \\ 0 & , v_i = v_j \end{cases}.$$

Let J_n and I_n denote the all one and identity matrix of order n and $J_{k,n}$ denote all one matrix of order $k \times n$, respectively. Then, the Seidel matrix $S(G^*)$ of a hypergraph G^* is defined as $S(G^*) = J_n - I_n - 2A(G^*)$ [15]. For any square matrix M we can find a scalar λ such that $M\mathbf{x} = \lambda\mathbf{x}$ where \mathbf{x} is the non-zero eigenvector corresponding to the eigenvalue λ . The collection of all the eigenvalues of $A(G^*)$ and $S(G^*)$ together with their multiplicities is known as the adjacency spectrum and the Seidel spectrum of G^* , respectively. Let $\lambda_1, \lambda_2, \lambda_3, \dots, \lambda_d$, be the eigenvalues of an adjacency matrix $A(G^*)$ of hypergraph G^* with multiplicities $m_1, m_2, m_3, \dots, m_d$. Then, the adjacency spectrum $\sigma_A(G^*)$ of G^* is denoted as,

$$\sigma_A(G^*) = \begin{pmatrix} \lambda_1 & \lambda_2 & \lambda_3 & \cdots & \lambda_d \\ m_1 & m_2 & m_3 & \cdots & m_d \end{pmatrix}.$$

The adjacency spectral radius (Seidel spectral radius) of G^* , denoted by $\rho(A(G^*))$ ($\rho(S(G^*))$) is the largest eigenvalue of $A(G^*)$ ($S(G^*)$). Two hypergraphs G^* and H^* are Seidel-cospectral (adjacency-cospectral) if they have the same Seidel spectrum (adjacency spectrum). For a k -uniform hypergraph $G^* = (V, E)$, $v \in V$ is said to be the neighbour of some $(k-1)$ subset of V , say U if $\{v\} \cup U$ is a hyperedge of G^* . The neighbourhood of U is denoted by $N(U)$.

In [8], Frucht and Harary proposed the construction of corona graphs. Subsequently, the analysis of the spectra of the corona product of two graphs has received huge attention. In [2], Barik et al., have provided the complete spectra of the corona product of two graphs as a means of determining the spectrum of larger graphs in terms of simple graphs. By introducing the concept of coronal McLeman [12] formulated a method to establish the spectrum of the corona product of two hypergraphs. Later, Sharma [14] determined the structural properties of the corona graph and its spectrum. The study on the corona product of graphs was recently extended to hypergraphs [13]. In connection with the spectrum of hypergraphs, it is of interest to identify a family of non-isomorphic cospectral hypergraphs. Obviously, isomorphic graphs are cospectral. There are several methods to find a cospectral family of hypergraphs. In this paper, we use signed M-GM switching to construct a family of non-isomorphic Seidel cospectral hypergraphs. Throughout this paper, we use the notation $j \in \mathbb{Z}^{[a,b]}$ to denote j that takes all the integer values satisfying the condition $a \leq j \leq b$.

This paper focuses on the study of the corona product of k -uniform hypergraphs. In Section 2, we give fundamental definitions and results that will be used later. In Section 3, we determine the adjacency and seidel spectrum of generalised corona of k -uniform hypergraphs. In Section 4, we estimate the adjacency and Seidel spectrum of corona product of two k -uniform hypergraphs. Also, we calculate the coronal of (k, r) -regular hypergraph and find an infinite pair of adjacency cospectral hypergraphs. In Section 5, we compute the adjacency spectrum of the corona hypergraph. In Section 6, we present a new method for the construction of Seidel-cospectral hypergraphs. Using this we get infinitely many non-isomorphic Seidel-cospectral k -uniform hypergraphs.

2 Preliminaries

Lemma 2.1. [6] *Let M_{11}, M_{12}, M_{21} , and M_{22} be matrices with M_{22} invertible. Let*

$$M = \begin{bmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \end{bmatrix}$$

Then, $\det(M) = \det(M_{22})\det(M_{11} - M_{12}M_{22}^{-1}M_{21})$.

Lemma 2.2. [9] For any two real numbers r and s ,

$$(rI_n - sJ_n)^{-1} = \frac{1}{r}I_n + \frac{s}{r(r - ns)}J_n.$$

Lemma 2.3. [7] Let $M \in \mathbb{R}^{n \times n}$ be an invertible matrix, U and W are $n \times 1$ matrices. Then

$$\det(UW^T + M) = (1 + W^T M^{-1}U) \det(M).$$

Lemma 2.4. [14] Let $n, m \in \mathbb{R}$. Then

$$\sum_{j=0}^{m-1} 2^j n(n-1)(n+1)^{m-j-1} + 2^m n = n(n+1)^m.$$

Definition 2.5. [13](Generalized corona of hypergraphs) The corona $G^* = G_0^* \odot_p^t G_i^*$, $i \in \mathbb{Z}^{[1,t]}$ of k -uniform hypergraphs G_0^* and G_i^* (where G_0^* and G_i^* has n and n_i vertices respectively) with partition $P = \{U_1, U_2, \dots, U_t\}$ (with $|U_i| = p$, $p \in \mathbb{N}$) of the vertex set V_0 of hypergraph G_0^* is defined as a k -uniform hypergraph $\bigcup_{i=1}^t \left(\bigcup_{j=1}^p G_0^*[U_i] \otimes G_i^{*(j)} \right)$ obtained by taking p copies of G_i^* say $G_i^{*(j)}$, where $j \in \mathbb{Z}^{[1,p]}$.

3 Generalised corona of hypergraphs

In this section, we determine the spectral properties of the generalised corona of k -uniform hypergraphs. In [13], Sarkar and Banerjee provided the spectrum of the generalised corona of k -uniform hypergraphs by considering another definition of the adjacency matrix. We modify the spectrum by taking into account the definition stated in [11].

3.1 Adjacency spectrum of corona of hypergraphs

Let G_0^* be a k -uniform hypergraph with a vertex set V_0 and partition $P = \{U_1, U_2, \dots, U_t\}$ such that $|U_i| = p$, $p \in \mathbb{N}$. Also, G_i^* be the (k, r) -regular hypergraphs with vertex set V_i and edge set E_i , where $i \in \mathbb{Z}^{[1,t]}$. Let $|V_i| = m$, then the corresponding adjacency matrix of G^* given by

$$A(G^*) = \begin{bmatrix} X & H \\ H^T & Y \end{bmatrix},$$

where X , H and Y denotes the adjacency relation between the vertices of V_0 , vertices of V_0 and V_i , and vertices of V_i respectively.

$$\begin{aligned} X &= A(G_0^*) + aI_t \otimes (J_p - I_p), \\ H &= bI_t \otimes J_{p,pm}, \\ Y &= \text{diag}(I_p \otimes Y_1, I_p \otimes Y_2, I_p \otimes Y_3, \dots, I_p \otimes Y_t), \end{aligned}$$

where $a = \begin{cases} p \left(\binom{p+m-2}{k-2} - \binom{p-2}{k-2} \right) & \text{if } p \in \mathbb{Z}^{[2,n]} \\ 0 & \text{if } p = 1 \end{cases}$, $b = \binom{p+m-2}{k-2}$, $Y_i = A(G_i^*) + c(J_m - I_m)$ and $c = \binom{p+m-2}{k-2} - \binom{m-2}{k-2}$.

Theorem 3.1.1. *The characteristic polynomial of $G^* = G_0^* \odot_p^t G_i^*$, $i \in \mathbb{Z}^{[1,t]}$ is given by,*

$$P_{A(G^*)}(\lambda) = \left(\prod_{i=1}^t \det(Y_i - \lambda I_m) \right)^p \det(A(G_0^*) + I_t \otimes \left((a - \frac{b^2 pm}{r(k-1) + c(m-1) - \lambda}) J_p - (a + \lambda) I_p \right)),$$

$$\text{where } a = \begin{cases} p \binom{p+m-2}{k-2} - \binom{p-2}{k-2} & \text{if } p \in \mathbb{Z}^{[2,n]} \\ 0 & \text{if } p = 1 \end{cases}, \quad b = \binom{p+m-2}{k-2}, \quad Y_i = A(G_i^*) + c(J_m - I_m) \text{ and } c = \binom{p+m-2}{k-2} - \binom{m-2}{k-2}.$$

Proof. The characteristic polynomial of G^* is given by,

$$P_{A(G^*)}(\lambda) = \det \begin{pmatrix} A(G_0^*) + aI_t \otimes (J_p - I_p) - \lambda I_n & bI_t \otimes J_{p,pm} \\ bI_t \otimes J_{pm,p} & Y - \lambda I_{ptk} \end{pmatrix}.$$

From Lemma 2.1, we get

$$P_{A(G^*)}(\lambda) = \det(Y - \lambda I_{ptk}) \det(A(G_0^*) + aI_t \otimes (J_p - I_p) - \lambda I_n - b^2 I_t \otimes J_{p,pm} (Y - \lambda I_{ptk})^{-1} I_t \otimes J_{pm,p}). \quad (1)$$

Since row sum of the adjacency matrix of a (k, r) -regular hypergraph is $r(k-1)$, we get

$$(Y - \lambda I_{ptk})(I_t \otimes J_{pm,p}) = (r(k-1) + c(m-1) - \lambda)(I_t \otimes J_{pm,p}).$$

Then,

$$(I_t \otimes J_{pm,p}) = \frac{1}{(r(k-1) + c(m-1) - \lambda)} (Y - \lambda I_{ptk})(I_t \otimes J_{pm,p}). \quad (2)$$

Also,

$$(I_t \otimes J_{pm,p})(I_t \otimes J_{pm,p}) = pm I_t \otimes J_p. \quad (3)$$

Hence the theorem follows from (1),(2) and (3). \square

Using Theorem 3.1.1, we obtain the characteristic polynomial of G^* in terms of the eigenvalues of the factor hypergraphs G_0^* and G_i^* , $i \in \mathbb{Z}^{[1,n]}$ when $p = 1$ and $t = n$.

Corollary 3.1.2. *Let G_0^* be a k -uniform hypergraph with eigenvalues $\lambda'_1 \geq \lambda'_2 \geq \lambda'_3 \geq \dots \geq \lambda'_n$ and G_i^* , $i \in \mathbb{Z}^{[1,n]}$ be (k, r) -regular hypergraphs with eigenvalues $\lambda_1^{(i)} = r(k-1) \geq \lambda_2^{(i)} \geq \lambda_3^{(i)} \geq \dots \geq \lambda_m^{(i)}$. Then the characteristic polynomial of $G^* = G_0^* \odot_1^n G_i^*$ is given by,*

$$P_{A(G^*)}(\lambda) = (r(k-1) - c(m-1) - \lambda)^n \prod_{i=1}^n \prod_{j=2}^m (\lambda_j^{(i)} - c - \lambda) \prod_{i=1}^n \left(\lambda'_i + \frac{\lambda^2 - (r(k-1) + c(m-1))\lambda - b^2 m}{r(k-1) + c(m-1) - \lambda} \right).$$

Proof. By substituting $t = n$ and $p = 1$ in Theorem 3.1.1, we have

$$P_{A(G^*)}(\lambda) = \left(\prod_{i=1}^n \det(Y_i - \lambda I_m) \right) \det \left(A(G_0^*) + \left(\frac{\lambda^2 - (r(k-1) + c(m-1))\lambda - b^2 m}{r(k-1) + c(m-1) - \lambda} \right) I_n \right).$$

Hence the result. \square

3.2 Seidel Spectrum of the corona of hypergraphs

Let G_0^* be a k -uniform hypergraph of order n with the partition $P = \{U_1, U_2, U_3, \dots, U_t\}$, where $|U_i| = p$, $p \in \mathbb{N}$ and G_i^* , $i \in \mathbb{Z}^{[1,t]}$ be a k -uniform hypergraph of order n_i . Then the Seidel matrix of $G^* = G_0^* \odot_p^t G_i^*$ is given by,

$$S(G^*) = \begin{bmatrix} X_S & H_S \\ H_S^T & Y_S \end{bmatrix}$$

such that,

$$\begin{aligned} X_S &= S(G_0^*) - 2aI_t \otimes (J_p - I_p), \\ H_S &= (J_t - 2bI_t) \otimes J_{p,pm_i}, \\ Y_S &= J_{pm_i t} + \text{diag}(I_p \otimes (Y_{S_1} - J_{n_i}), \dots, I_t \otimes (Y_{S_t} - J_{n_i})), \end{aligned}$$

where $Y_{S_i} = S(G_i^*) - 2c(J_{n_i} - I_{n_i})$.

Theorem 3.2.1. *Let G_0^* be a k -uniform hypergraph of order n with the partition $P = \{U_1, U_2, U_3, \dots, U_t\}$, where $|U_i| = p$, $p \in \mathbb{N}$ and G_i^* , $i \in \mathbb{Z}^{[1,t]}$ be a (k, r) -regular hypergraph of order m . Then the characteristic polynomial of Seidel matrix of $G^* = G_0^* \odot_p^t G_i^*$ is*

$$\begin{aligned} P_{S(G^*)}(\mu) &= \left(1 + \frac{pmt}{h}\right) \left(\prod_{i=1}^t \det(Y_{S_i} - J_m - \mu I_m)\right)^p \det\left(S(G_0^*) + I_t \otimes \left(-2a + \frac{4pmb^2}{h}\right)J_p - I_p\right) \\ &\quad + (2a - \mu)I_p - \frac{pm}{h} \left((t - 4b) - \frac{pm(t - 2b)^2}{h + pmt}\right)J_n, \end{aligned}$$

where $a = \begin{cases} p \binom{p+m-2}{k-2} - \binom{p-2}{k-2} & \text{if } p \in \mathbb{Z}^{[2,n]} \\ 0 & \text{if } p = 1 \end{cases}$, $b = \binom{p+m-2}{k-2}$, $Y_{S_i} = S(G_i^*) - 2c(J_m - I_m)$, $c = \binom{p+m-2}{k-2} - \binom{m-2}{k-2}$ and $h = -(1 + 2r(k-1) + 2c(m-1) + \mu)$.

Proof. The characteristic polynomial of the Seidel matrix of G^* is given by

$$P_{S(G^*)}(\mu) = \det \begin{pmatrix} S(G_0^*) - 2aI_t \otimes (J_p - I_p) - \mu I_n & H_S \\ H_S^T & Y_S - \mu I_{pmt} \end{pmatrix},$$

where $Y_S - \mu I_{pmt} = J_{pmt} + \text{diag}(I_p \otimes (Y_{S_1} - J_m - \mu I_m), \dots, I_p \otimes (Y_{S_t} - J_m - \mu I_m))$ and $H_S = (J_t - 2bI_t) \otimes J_{p,pm}$.

From Lemma 2.1, we get

$$P_{S(G^*)}(\mu) = \det(Y_S - \mu I) \det(S(G_0^*) - 2aI_t \otimes (J_p - I_p) - \mu I_n - H_S(Y_S - \mu I_{pmt})^{-1} H_S^T). \quad (4)$$

Since G_i^* 's are r -regular m -uniform hypergraphs, we have

$$S(G_i^*)J_{m,p} = (m - 1 - 2r(k-1))J_{m,p}.$$

On simplification, we get

$$\left. \begin{aligned} &\text{diag}(I_p \otimes (Y_{S_1} - J_m - \mu I_m), \dots, I_p \otimes (Y_{S_t} - J_m - \mu I_m))(J_t \otimes J_{m,p}) \\ &= -(1 + 2r(k-1) + 2c(m-1) + \mu)(J_t \otimes J_{m,p}), \\ &\text{diag}(I_p \otimes (Y_{S_1} - J_m - \mu I_m), \dots, I_p \otimes (Y_{S_t} - J_m - \mu I_m))(I_t \otimes J_{m,p}) \\ &= -(1 + 2r(k-1) + 2c(m-1) + \mu)(I_t \otimes J_{m,p}). \end{aligned} \right\} \quad (5)$$

Now consider,

$$(Y_S - \mu I_{pmt})H_S^T = (J_{pmt} + \text{diag}(I_p \otimes (Y_{S_1} - J_m - \mu I_m), \dots, I_p \otimes (Y_{S_t} - J_m - \mu I_m)))((J_t - 2bI_t) \otimes J_{pm,p}).$$

From (5), we get

$$\begin{aligned} (Y_S - \mu I_{pmt})H_S^T &= J_{pmt}((J_t - 2bI_t) \otimes J_{pm,p}) - (1 + 2r(k-1) + 2c(m-1) + \mu)((J_t - 2bI_t) \otimes J_{pm,p}) \\ &= (J_{pmt} + hI_{pmt})H_S^T, \end{aligned}$$

where $h = -(1 + 2r(k-1) + 2c(m-1) + \mu)$. Therefore,

$$H_S^T = (J_{pmt} + hI_{pmt})^{-1}(Y_S - \mu I_{pmt})H_S^T.$$

By Lemma 2.2, we get $(hI_{pmt} + J_{pmt})^{-1} = \frac{1}{h}I_{pmt} - \frac{1}{h(h+pmt)}J_{pmt}$. Hence,

$$H_S^T = \left(\frac{1}{h}I_{pmt} - \frac{1}{h(h+pmt)}J_{pmt}\right)(Y_S - \mu I_{pmt})H_S^T. \quad (6)$$

Again,

$$\begin{aligned} H_S H_S^T &= \left((J_t - 2bI_t) \otimes J_{p,pm}\right) \left((J_t - 2bI_t) \otimes J_{pm,p}\right) \\ &= ((t-4b)J_t + 4b^2I_t) \otimes pmJ_p. \end{aligned} \quad (7)$$

Also,

$$\begin{aligned} H_S J_{pmt} H_S^T &= \left((J_t - 2bI_t) \otimes J_{p,pm}\right) J_{pmt} \left((J_t - 2bI_t) \otimes J_{pm,p}\right) \\ &= \left(J_{pt,pmt} J_{pmt} - 2b(I_t \otimes J_{p,pm}) J_{pmt}\right) \left((J_t - 2bI_t) \otimes J_{pm,p}\right) \\ &= (pmt - 2bpm) J_{pt,pmt} \left((J_t - 2bI_t) \otimes J_{pm,p}\right) \\ &= (pmt - 2bpm)^2 J_{pt} = p^2 m^2 (t - 2b)^2 J_{pt}. \end{aligned} \quad (8)$$

From (6), we have

$$\begin{aligned} H_S (Y_S - \mu I_{pmt})^{-1} H_S^T &= H_S (Y_S - \mu I_{pmt})^{-1} \left(\frac{1}{h}I_{pmt} - \frac{1}{h(h+pmt)}J_{pmt}\right) (Y_S - \mu I_{pmt}) H_S^T \\ &= \frac{1}{h} H_S H_S^T - \frac{1}{h(h+pmt)} H_S (Y_S - \mu I_{pmt})^{-1} J_{pmt} (Y_S - \mu I_{pmt}) H_S^T. \end{aligned}$$

Since $Y_S - \mu I_{pmt}$ commute with J_{pmt} , we get

$$H_S (Y_S - \mu I_{pmt})^{-1} H_S^T = \frac{1}{h} H_S H_S^T - \frac{1}{h(h+pmt)} H_S J_{pmt} H_S^T.$$

From (7) and (8), we obtain

$$H_S (Y_S - \mu I_{pmt})^{-1} H_S^T = \frac{pm}{h} \left(\left((t-4b) - \frac{pm(t-2b)^2}{h+pmt} \right) J_{pt} + 4b^2 (I_t \otimes J_p) \right). \quad (9)$$

Take $D = \text{diag}(I_p \otimes (Y_{S_1} - J_m - \mu I_m), \dots, I_p \otimes (Y_{S_t} - J_m - \mu I_m))$, Then $DJ_{pmt,1} = hJ_{pmt,1}$. Also,

$$J_{pmt,1} = \frac{1}{h} D J_{pmt,1}. \quad (10)$$

Now consider,

$$\det(Y_S - \mu I) = \det(J_{pmt} + D) = \det(J_{pmt,1} J_{1,pmt} + D).$$

By Lemma 2.3, we have

$$\det(Y_S - \mu I) = (1 + J_{1,pmt} D^{-1} J_{pmt,1}) \det(D).$$

Using (9) and (10) in (4), we get

$$\begin{aligned} P_{S(G^*)}(\mu) &= \left(1 + \frac{pmt}{h}\right) \left(\prod_{i=1}^t \det(Y_{S_i} - J_m - \mu I_m)\right)^p \det\left(S(G_0^*) + I_t \otimes \left(-\left(2a + \frac{4pmb^2}{h}\right)J_p - I_p\right)\right. \\ &\quad \left.+ (2a - \mu)I_p\right) - \frac{pm}{h} \left(\left(t - 4b\right) - \frac{pm(t - 2b)^2}{h + pmt}\right) J_n. \end{aligned}$$

□

Next, we determine the characteristic polynomial of G^* in terms of the factor hypergraphs when $p = 1$ and $t = n$.

Theorem 3.2.2. *Let G_0^* be a (k, r_0) -regular hypergraph of order n with partition $P = \{U_1, U_2, U_3, \dots, U_t\}$, where $|U_i| = p, p \in \mathbb{N}$ and $G_i^*, i \in \mathbb{Z}^{[1,t]}$ be a (k, r) -regular hypergraph of order m . Let $\mu_1^{(0)} = n - 1 - 2r_0(k - 1)$, $\mu_2^{(0)} \geq \mu_3^{(0)} \geq \dots \geq \mu_n^{(0)}$ be the Seidel eigenvalues of G_0^* and $\mu_1^{(i)} = m - 1 - 2r(k - 1)$, $\mu_2^{(i)} \geq \mu_3^{(i)} \geq \dots \geq \mu_m^{(i)}$ be the Seidel eigenvalues of G_i^* . Then the characteristic polynomial of the Seidel matrix of $G^* = G_0^* \odot_1^n G_i^*$ is*

$$P_{S(G^*)}(\mu) = ((\mu_1^{(0)} - \mu)(h + mn) + 4bm(n - b) - mn^2) \left(\prod_{i=2}^n (\mu_i^{(0)} h - 4mb^2 - \mu h)\right) \left(\prod_{j=1}^n \prod_{i=2}^m (\mu_i^{(j)} + 2c - \mu)\right)$$

where $b = \binom{p+m-2}{k-2}$, $c = \binom{p+m-2}{k-2} - \binom{m-2}{k-2}$ and $h = -(1 + 2r(k - 1) + 2c(m - 1) + \mu)$.

Proof. By substituting $t = n$ and $p = 1$ in Theorem 3.2.1, we obtain

$$P_{S(G^*)}(\mu) = \left(\frac{h + mn}{h}\right) \left(\prod_{i=1}^n \det(Y_{S_i} - J_m - \mu I_m)\right) \det\left(S(G_0^*) + \left(\frac{-4mb^2 - \mu h}{h}\right)I_n - \frac{m}{h} \left(\frac{nh - 4bh - 4b^2m}{h + mn}\right)J_n\right). \quad (11)$$

Consider,

$$\begin{aligned} &\det\left(S(G_0^*) + \left(\frac{-4mb^2 - \mu h}{h}\right)I_n - \frac{m}{h} \left(\frac{nh - 4bh - 4b^2m}{h + mn}\right)J_n\right) \\ &= \left(\mu_1^{(0)} - \frac{mn(nh - 4bh - 4b^2m)}{h(h + mn)} - \frac{4mb^2 + \mu h}{h}\right) \left(\prod_{i=2}^n \left(\mu_i^{(0)} - \frac{4mb^2 + \mu h}{h}\right)\right) \\ &= \left(\frac{\mu_1^{(0)} h(h + mn) - mn(nh - 4bh - 4b^2m) - (4mb^2 + \mu h)(h + mn)}{h^n(h + mn)}\right) \left(\prod_{i=2}^n (\mu_i^{(0)} h - 4mb^2 + \mu h)\right) \\ &= \left(\frac{(\mu_1^{(0)} - \mu)(h + mn) + 4bm(n - b) - mn^2}{h^n(h + mn)}\right) \left(\prod_{i=2}^n (\mu_i^{(0)} h - 4mb^2 + \mu h)\right). \quad (12) \end{aligned}$$

Also, we have

$$\prod_{i=1}^n \det(Y_{S_i} - J_m - \mu I_m) = \prod_{i=1}^n \det(S(G_i^*) - (2c + 1)J_m + (2c - \mu)I_m)$$

$$\begin{aligned}
&= (m-1-2r(k-1) - (2c+1)m + 2c - \mu)^n \prod_{j=1}^n \prod_{i=2}^m (\mu_i^{(j)} + 2c - \mu) \\
&= h^n \prod_{i=2}^m (\mu_i^{(j)} + 2c - \mu). \tag{13}
\end{aligned}$$

Substituting (12),(13) in (11), we get the desired result. \square

4 Spectrum of the corona of two hypergraphs

The corona product of two k -uniform hypergraphs $G^* = G_0^* \odot_1^n G_1^*$, is the k -uniform hypergraph obtained by taking $p = 1$, $k = n$, and $G_i^* = G_1^*$ for $i \in \mathbb{Z}^{[1,n]}$ in Definition 2.5. By proper labelling of vertices, we obtain the adjacency matrix and Seidel matrix of G^* as follows:

$$\begin{aligned}
A(G^*) &= \begin{bmatrix} A(G_0^*) & \binom{m-1}{k-2} J_{1,m} \otimes I_n \\ \binom{m-1}{k-2} J_{m,1} \otimes I_n & A(G_1^*) \otimes I_n \end{bmatrix}, \\
S(G^*) &= \begin{bmatrix} S(G_0^*) & J_{1,m} \otimes \left(J_n - 2 \binom{m-1}{k-2} I_n \right) \\ J_{m,1} \otimes \left(J_n - 2 \binom{m-1}{k-2} I_n \right) & J_m \otimes (J_n - I_n) + S(G_1^*) \otimes I_n \end{bmatrix}.
\end{aligned}$$

The following theorem describes the adjacency spectrum and spectral radius of corona product of two k -uniform hypergraphs G_0^* and G_1^* , where G_1^* is r -regular hypergraph.

Theorem 4.1. *Let G_0^* be any k -uniform hypergraph of order n with eigenvalues $\lambda_1^{(0)} \geq \lambda_2^{(0)} \geq \lambda_3^{(0)} \geq \dots \geq \lambda_n^{(0)}$ and G_1^* be a (k, r) -regular hypergraph of order m with eigenvalues $\lambda_1^{(1)} = r(k-1) \geq \lambda_2^{(1)} \geq \lambda_3^{(1)} \geq \dots \geq \lambda_m^{(1)}$, then for $i \in \mathbb{Z}^{[1,n]}$, $j \in \mathbb{Z}^{[2,m]}$ the spectrum of $G^* = G_0^* \odot_1^n G_1^*$ is given by*

$$\sigma_A(G^*) = \left(\begin{array}{c} \frac{\lambda_i^{(0)} + r(k-1) \pm \sqrt{(\lambda_i^{(0)} - r(k-1))^2 + 4m \binom{m-1}{k-2}^2}}{2} \\ 1 \end{array} \quad \begin{array}{c} \lambda_j^{(1)} \\ n \end{array} \right)$$

$$\text{and } \rho_A(G^*) = \frac{\rho(A(G_0^*)) + r(k-1) + \sqrt{(\rho(A(G_0^*)) - r(k-1))^2 + 4m \binom{m-1}{k-2}^2}}{2}.$$

Proof. For $i \in \mathbb{Z}^{[1,n]}$, let \mathbf{x}_i be an eigenvector of $A(G_0^*)$ corresponding to the eigenvalue $\lambda_i^{(0)}$. We have to determine λ_i 's and t_i 's such that $\mathbf{x}_i = [\mathbf{x}_i \quad t_i J_{m,1} \otimes \mathbf{x}_i]^T$ is an eigenvector corresponding to the eigenvalue λ_i of $A(G^*)$. Since $A(G^*)\mathbf{x}_i = \lambda_i \mathbf{x}_i$, we have the following equations

$$\begin{aligned}
\lambda_i^{(0)} \mathbf{x}_i + t_i \binom{m-1}{k-2} (J_{1,m} \otimes I_n)(J_{m,1} \otimes \mathbf{x}_i) &= \lambda_i \mathbf{x}_i, \\
\binom{m-1}{k-2} (J_{m,1} \otimes I_n) \mathbf{x}_i + t_i (A(G_1^*) \otimes I_n)(J_{m,1} \otimes \mathbf{x}_i) &= \lambda_i t_i (J_{m,1} \otimes \mathbf{x}_i).
\end{aligned}$$

On simplification, we get

$$\lambda_i^{(0)} + t_i m \binom{m-1}{k-2} = \lambda_i, \quad \binom{m-1}{k-2} + t_i r(k-1) = \lambda_i t_i.$$

After solving these equations, we obtain

$$\lambda_i = \frac{\lambda_i^{(0)} + r(k-1) \pm \sqrt{(\lambda_i^{(0)} - r(k-1))^2 + 4m \binom{m-1}{k-2}^2}}{2},$$

and

$$t_i = \frac{-(\lambda_i^{(0)} - r(k-1)) \pm \sqrt{(\lambda_i^{(0)} - r(k-1))^2 + 4m \binom{m-1}{k-2}^2}}{2m \binom{m-1}{k-2}} = \frac{\binom{m-1}{k-2}}{\lambda_i - r(k-1)}.$$

Hence, for $i \in \mathbb{Z}^{[1,n]}$

$$\tilde{\lambda}_i = \frac{\lambda_i^{(0)} + r(k-1) + \sqrt{(\lambda_i^{(0)} - r(k-1))^2 + 4m \binom{m-1}{k-2}^2}}{2}, \quad \tilde{\tilde{\lambda}}_i = \frac{\lambda_i^{(0)} + r(k-1) - \sqrt{(\lambda_i^{(0)} - r(k-1))^2 + 4m \binom{m-1}{k-2}^2}}{2}$$

are the eigenvalues of $A(G^*)$ corresponding to the eigenvectors

$$\begin{bmatrix} \mathbf{x}_i \\ \frac{\binom{m-1}{k-2}}{\tilde{\lambda}_i - r(k-1)} J_{m,1} \otimes \mathbf{x}_i \end{bmatrix}, \quad \begin{bmatrix} \mathbf{x}_i \\ \frac{\binom{m-1}{k-2}}{\tilde{\tilde{\lambda}}_i - r(k-1)} J_{m,1} \otimes \mathbf{x}_i \end{bmatrix}$$

respectively. Let \mathbf{y}_j be an eigenvector corresponding to the eigenvalue $\lambda_j^{(1)}$ of $A(G_1^*)$ and let $\mathbf{e}_i \in \mathbb{R}^n$ be a vector with 1 in the i -th coordinate and 0 elsewhere. Since G_1^* is a (k, r) -regular hypergraph, for $j \in \mathbb{Z}^{[2,m]}$, \mathbf{y}_j 's are orthogonal to \mathbf{j} . It is clear that, for $i \in \mathbb{Z}^{[1,n]}$

$$A(G^*) \begin{bmatrix} \mathbf{0}_{n,1} \\ \mathbf{y}_j \otimes \mathbf{e}_i \end{bmatrix} = \lambda_j^{(1)} \begin{bmatrix} \mathbf{0}_{n,1} \\ \mathbf{y}_j \otimes \mathbf{e}_i \end{bmatrix}.$$

For $2 \leq j \leq m$, $\lambda_j^{(1)}$ is an eigenvalue of $A(G^*)$ with multiplicity n . Note that $\lambda_j^{(1)} \leq r(k-1)$, hence $\rho(A(G^*)) = \tilde{\lambda}_1$. This completes the proof. \square

The following corollary gives the adjacency spectrum and spectral radius of G^* when $G_1^* = K_m^m$.

Corollary 4.2. *Let G_0^* be any m -uniform hypergraph of order n with eigenvalues $\lambda_1^{(0)} \geq \lambda_2^{(0)} \geq \lambda_3^{(0)} \geq \dots \geq \lambda_n^{(0)}$ and $G_1^* = K_m^m$ be an m -uniform complete hypergraph of order m , then for $i \in \mathbb{Z}^{[1,n]}$, $j \in \mathbb{Z}^{[2,m]}$ the spectrum of $G^* = G_0^* \odot_1^n G_1^*$ is given by*

$$\sigma_A(G^*) = \begin{pmatrix} \frac{\lambda_i^{(0)} + m - 1 \pm \sqrt{(\lambda_i^{(0)} - (m-1))^2 + 4m(m-1)^2}}{2} & & -1 \\ & 1 & n(m-1) \end{pmatrix}$$

$$\text{and } \rho(A(G^*)) = \frac{\rho(A(G_0^*)) + (m-1) + \sqrt{(\rho(A(G_0^*)) - (m-1))^2 + 4m(m-1)}}{2}.$$

Example 4.3. *Let $G_0^* = G_1^* = K_m^m$ be the complete m uniform hypergraph of order m . Then spectrum of $G^* = K_m^m \odot_1^n K_m^m$ is*

$$\sigma_A(G^*) = \begin{pmatrix} (m-1)(1 \pm \sqrt{m}) & \frac{m-2 \pm \sqrt{m^2 + 4m(m-1)^2}}{2} & & -1 \\ & 1 & m-1 & m(m-1) \end{pmatrix}$$

$$\text{and } \rho(A(G^*)) = (m-1)(1 + \sqrt{m}).$$

The following theorem gives the Seidel spectrum of corona of two k -uniform hypergraphs.

Theorem 4.4. *Let G_0^* be an n order (k, r_0) -regular hypergraph with Seidel eigenvalues $\mu_1^{(0)} \geq \mu_2^{(0)} \geq \mu_3^{(0)} \geq \dots \geq \mu_n^{(0)}$ and G_1^* be m -order (k, r_1) -regular hypergraph with Seidel eigenvalues $\mu_1^{(1)} \geq \mu_2^{(1)} \geq \mu_3^{(1)} \geq \dots \geq \mu_m^{(1)}$, then for $i \in \mathbb{Z}^{[1, n]}$, $j \in \mathbb{Z}^{[1, m]}$, $\mu_i^{(0)} \neq n - 1 - 2r_0(k - 1)$ and $\mu_j^{(1)} \neq m - 1 - 2r_1(k - 1)$ the Seidel spectrum of $G^* = G_0^* \odot_1^n G_1^*$ is given by*

$$\sigma_S(G^*) = \begin{pmatrix} \frac{\mu_i^{(0)} - 1 - 2r_1(k-1) \pm \sqrt{(\mu_i^{(0)} + 1 + 2r_1(k-1))^2 + 16m \binom{m-1}{k-2}^2}}{2} & \mu_j^{(1)} & \alpha_1 & \alpha_2 \\ 1 & n & 1 & 1 \end{pmatrix},$$

where α_1 and α_2 are the roots of the equation $\mu^2 + (2 - n(m + 1) + 2(r_0 + r_1)(k - 1))\mu + (n - 1 - 2r_0(k - 1))(mn - 1 - 2r_1(k - 1)) - m(n - 2 \binom{m-1}{k-2})^2 = 0$.

Proof. Let \mathbf{x}_i be an eigenvector associated with an eigenvalue $\mu_i^{(0)}$ ($\mu_i^{(0)} \neq n - 1 - 2r_0(k - 1)$) and $i \in \mathbb{Z}^{[1, n]}$ of $S_{G_0^*}^*$. We have to determine μ_i 's and t_i 's such that $\mathbf{x}_i = [\mathbf{x}_i \ t_i J_{m,1} \otimes \mathbf{x}_i]^T$ is an eigenvector associated with the eigenvalue μ_i of $S_{G^*}^*$. Since $S(G^*)\mu_i = \mu_i \mathbf{x}_i$, we have the following equations,

$$\mu_i^{(0)} \mathbf{x}_i + t_i m \left(J_n - 2 \binom{m-1}{k-2} I_n \right) \mathbf{x}_i = \mu_i \mathbf{x}_i,$$

$$\left(J_{m,1} \otimes (J_n - 2 \binom{m-1}{k-2} I_n) \right) \mathbf{x}_i + t_i \left(m J_{m,1} \otimes (J_n - I_n) \mathbf{x}_i + S(G_1^*) J_{m,1} \otimes \mathbf{x}_i \right) = \mu_i t_i J_{m,1} \otimes \mathbf{x}_i.$$

Also, row sum of $S(G_1^*)$ is $m - 1 - 2r_1(k - 1)$ and \mathbf{x}_i is orthogonal to \mathbf{j} . Therefore,

$$\begin{aligned} \mu_i &= \mu_i^{(0)} - 2t_i m \binom{m-1}{k-2}, \\ \mu_i t_i &= -2 \binom{m-1}{k-2} + t_i (-1 - 2r_1(k - 1)). \end{aligned}$$

On solving the system, we get

$$t_i = \frac{1 + 2r_1(k - 1) + \mu_i^{(0)} \pm \sqrt{(1 + 2r_1(k - 1) + \mu_i^{(0)})^2 + 16m \binom{m-1}{k-2}^2}}{4m \binom{m-1}{k-2}} = \frac{-2 \binom{m-1}{k-2}}{1 + 2r_1(k - 1) + \mu_i}$$

and

$$\mu_i = \frac{\mu_i^{(0)} - 1 - 2r_1(k - 1) \pm \sqrt{(1 + 2r_1(k - 1) + \mu_i^{(0)})^2 + 16m \binom{m-1}{k-2}^2}}{2}, \mu_i^{(0)} \neq n - 1 - 2r_0(k - 1).$$

Now we can see that, for $i \in \mathbb{Z}^{[1, n]}$

$$S(G^*) \begin{bmatrix} \mathbf{0}_{1, n} \\ \mathbf{y}_j \otimes e_i \end{bmatrix} = \mu_j^{(1)} \begin{bmatrix} \mathbf{0}_{1, n} \\ \mathbf{y}_j \otimes e_i \end{bmatrix},$$

where \mathbf{y}_j 's are the eigenvectors corresponding to the eigenvalues $\mu_j^{(1)}$'s ($\mu_j^{(1)} \neq m - 1 - 2r_1(k - 1)$) and $j \in \mathbb{Z}^{[1, m]}$. Thus $\mu_j^{(1)}$'s are eigenvalues of $S(G^*)$ with multiplicity n .

We obtain the remaining eigenvalues of $S(G^*)$ from the eigenvalues of the quotient matrix Q of $S(G^*)$,

$$Q = \begin{bmatrix} n-1-2r_0(k-1) & m(n-2\binom{m-1}{k-2}) \\ n-2\binom{m-1}{k-2} & mn-1-2r_1(k-1) \end{bmatrix}.$$

Thus the remaining two eigenvalues of $S(G^*)$ can be obtained from the characteristic polynomial of Q ($= \mu^2 + (2-n(m+1)+2(r_0+r_1)(k-1))\mu + (n-1-2r_0(k-1))(mn-1-2r_1(k-1)) - m(n-2\binom{m-1}{k-2}) = 0$). Hence the theorem. \square

4.1 Coronal of the hypergraph

Now we introduce a new invariant called coronal of a matrix, which is helpful for determining the spectrum of the corona of two hypergraphs.

Definition 4.1.1. Let G^* be a k -uniform hypergraph on n vertices with adjacency matrix A . If $A - \lambda I$ is invertible, then the coronal $\chi_{A(G^*)}(\lambda)$ is defined to be the sum of entries of $(A - \lambda I)^{-1}$. This can be written as

$$\chi_{A(G^*)}(\lambda) = J_{1,n}(A - \lambda I)^{-1}J_{n,1}.$$

By applying Definition 4.1.1, we determine the characteristic polynomial of the corona product of two k -uniform hypergraphs in a more computationally easier manner.

Theorem 4.1.2. Let G_0^* and G_1^* be k -uniform hypergraphs of order n and m , respectively. Then, the characteristic polynomial of $G^* = G_0^* \odot_1^n G_1^*$ is

$$P_{A(G^*)}(\lambda) = \left(P_{A(G_1^*)}(\lambda)\right)^n P_{A(G_0^*)}\left(\lambda + \binom{m-1}{k-2} \chi_{A(G_1^*)}(\lambda)\right).$$

Proof. The characteristic polynomial of G^* is,

$$P_{A(G^*)}(\lambda) = \det \begin{pmatrix} A(G_0^*) - \lambda I_n & \binom{m-1}{k-2} J_{1,m} \otimes I_n \\ \binom{m-1}{k-2} J_{m,1} \otimes I_n & (A(G_1^*) - \lambda I_m) \otimes I_n \end{pmatrix}.$$

From Lemma 2.1, we get

$$\begin{aligned} P_{A(G^*)}(\lambda) &= \det \left(A(G_0^*) - \lambda I_n - \binom{m-1}{k-2} (J_{1,m} \otimes I_n) ((A(G_1^*) - \lambda I_m) \otimes I_n)^{-1} (J_{m,1} \otimes I_n) \right) \\ &\quad \det \left((A(G_1^*) - \lambda I_m) \otimes I_n \right) \\ &= \det \left(A(G_0^*) - \left(\binom{m-1}{k-2} \chi_{A(G_1^*)}(\lambda) + \lambda \right) I_n \right) \det \left(A(G_1^*) - \lambda I_m \right)^n \\ &= \left(P_{A(G_0^*)} \left(\binom{m-1}{k-2} \chi_{A(G_1^*)}(\lambda) + \lambda \right) \right) \left(P_{A(G_1^*)}(\lambda) \right)^n. \end{aligned}$$

\square

Corollary 4.1.3. Let G_0^* and H_0^* are non-isomorphic cospectral k -uniform hypergraph, and G_1^* be any k -uniform hypergraph. Then $G_0^* \odot_1^n G_1^*$ and $H_0^* \odot_1^n G_1^*$ are non-isomorphic cospectral k -uniform hypergraphs.

By finding the corona product with G_1^* repeatedly, we get infinite pair of adjacency-cospectral hypergraphs. Clearly $G_1^* \odot_1^n G_0^*$ and $G_1^* \odot_1^n H_0^*$ need not be cospectral k -uniform hypergraph. In addition, if $\chi_{A(G_0^*)}(\lambda) = \chi_{A(H_0^*)}(\lambda)$, then $G_1^* \odot_1^n G_0^*$ and $G_1^* \odot_1^n H_0^*$ are non-isomorphic cospectral hypergraphs.

Next, we determine the coronal of (k, r) -regular hypergraph, which can be used to find the spectrum of the corona of two hypergraphs.

Theorem 4.1.4. Let G_1^* be a (k, r_1) -regular hypergraph on m vertices. Then coronal of $A(G_1^*)$ is given by,

$$\chi_{A(G_1^*)}(\lambda) = \frac{m}{r_1(k-1) - \lambda}.$$

Proof. As a consequence of regularity of $A(G_1^*)$, it follows that $(A(G_1^*) - \lambda I_m)J_{m,1} = (r(m-1) - \lambda)J_{m,1}$ and also $J_{m,1} = \frac{(A(G_1^*) - \lambda I_m)J_{m,1}}{(r(m-1) - \lambda)}$. Hence the coronal of $A(G_1^*)$ is,

$$\chi_{A(G_1^*)}(\lambda) = J_{1,m}(A(G_1^*) - \lambda I_m)^{-1}J_{m,1} = \frac{J_{1,m}J_{m,1}}{r_1(k-1) - \lambda} = \frac{m}{r_1(k-1) - \lambda}.$$

□

Using the coronal of (k, r) -regular hypergraph G_1^* , we get the characteristic polynomial of $G^* = G_0^* \odot_1^n G_1^*$ as follows

$$P_{A(G^*)}(\lambda) = \left(P_{A(G_1^*)}(\lambda)\right)^n P_{A(G_0^*)} \left(\lambda + \frac{m}{r(k-1) - \lambda} \left(\frac{m-1}{k-2}\right)^2\right). \quad (14)$$

We can easily find the spectrum of G^* from (14), if the spectrum of G_0^* and G_1^* are known. The following example gives the spectrum of G^* if $G_1^* = K_3^3$ and G_0^* as in Figure 1(b).

Example 4.1.5. Let G_0^* and K_3^3 be 3-uniform hypergraphs as given in Figure 1 and $G^* = G_0^* \odot_1^n K_3^3$. The characteristic polynomial of $A(G_0^*)$ is $P_{A(G_0^*)}(\lambda) = \lambda^4 - 8\lambda^2 - 8\lambda$, $P_{A(K_3^3)}(\lambda) = -\lambda^3 + 3\lambda + 2$ and the coronal of K_3^3 is $\chi_{K_3^3}(\lambda) = \frac{-3}{\lambda-2}$. Here $n = 4$, $m = 3$ and $k = 3$, we have

$$\begin{aligned} P_{A(G^*)}(\lambda) &= (-\lambda^3 + 3\lambda + 2)^4 \left(\left(\lambda - \frac{12}{\lambda-2}\right)^4 - 8 \left(\lambda - \frac{12}{\lambda-2}\right)^2 - 8 \left(\lambda - \frac{12}{\lambda-2}\right) \right) \\ &= (\lambda + 1)^8 (\lambda^2 - 16)(\lambda^2 - 2\lambda - 12)(\lambda^4 - 6\lambda^3 - 16\lambda^2 + 80\lambda + 80). \end{aligned}$$

Then the spectrum of G^* is,

$$\sigma_{A(G^*)} = \left(\begin{array}{cccccccccc} -1 & 16 & -16 & \frac{829}{180} & \frac{-3567}{1369} & \frac{10046}{1637} & \frac{1106}{263} & \frac{-2877}{836} & \frac{-1525}{1693} \\ 8 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{array} \right)$$




Figure 1: Example of Corona product of two hypergraphs

5 Spectrum of the corona hypergraph $G_0^{*(m)}$

In this section, we define the corona hypergraph based on the definition of the corona product of two hypergraphs. In addition, we determine the spectrum of the corona hypergraph.

Definition 5.1. Let G_0^* be a k -uniform hypergraph on n vertices. Then the corona hypergraph $G_0^{*(m)}$ with respect to the base hypergraph $G_0^* = G_0^{*(1)}$ is obtained by,

$$G_0^{*(m)} = G_0^{*(m-1)} \odot_1^n G_0^{*(1)}.$$




Figure 2: $K_3^{3(3)}$

Figure 2 illustrates the corona hypergraph $G_0^{*(m)}$, when $G_0^* = K_m^m$. Following are some results on the corona hypergraphs.

1. The order of $G_0^{*(m)}$ is $n(n+1)^{m-1}$.
2. The size of $G_0^{*(m)}$ is $\tilde{e}(n+1)^{(m-1)} + \binom{n}{k-1}((n+1)^{m-2} - 1)$, where \tilde{e} is the size of G_0^* .
3. If G_0^* is connected, then the corona hypergraph $G_0^{*(m)}$ is also connected.

Now we obtain the adjacency spectrum of the corona hypergraph $G_0^{*(m)}$ in terms of the eigenvalues of G_0^* by applying Theorem 4.1.

For any (k, r) -regular hypergraph of order n , we denote

$$\phi(\lambda) = \frac{\lambda + r(k-1) \pm \sqrt{(\lambda - r(k-1))^2 + 4n \binom{n-1}{k-2}^2}}{2},$$

$$\phi^0(\lambda) = \lambda \text{ and } \phi^m(\lambda) = \phi(\phi^{m-1}(\lambda)).$$

Theorem 5.2. Let G_0^* be a (k, r) regular hypergraph and its spectrum be

$$\sigma_A(G_0^*) = \begin{pmatrix} \lambda_1 = r(k-1) & \lambda_2 & \lambda_3 & \cdots & \lambda_n \\ 1 & 1 & 1 & \cdots & 1 \end{pmatrix}.$$

Then the spectrum of corona hypergraph $G_0^{*(m)}$ is,

$$\sigma_A(G_0^{*(m)}) = \begin{pmatrix} \phi^{m-1}(\lambda_i)_{i \in \mathbb{Z}[1, n]} & \phi^{m-2}(\lambda_i)_{i \in \mathbb{Z}[2, n]} & \cdots & \phi^j(\lambda_i)_{i \in \mathbb{Z}[2, n]} & \cdots & \phi^0(\lambda_i)_{i \in \mathbb{Z}[2, n]} \\ 1 & n & \cdots & n(n+1)^{m-j-2} & \cdots & n(n+1)^{m-2} \end{pmatrix}.$$

Proof. To prove this we use mathematical induction on m . From Theorem 4.1, for $i \in \mathbb{Z}^{[1,n]}, j \in \mathbb{Z}^{[2,n]}$ the spectrum of $G_0^{*(2)} = G_0^* \odot G_0^*$ is given by

$$\sigma_A(G_0^{*(2)}) = \left(\begin{array}{c} \frac{\lambda_i + r(k-1) \pm \sqrt{(\lambda_i - r(k-1))^2 + 4n \binom{n-1}{k-2}}}{2} \\ 1 \end{array} \lambda_j \right) = \left(\begin{array}{cc} \phi^1(\lambda_i)_{i \in \mathbb{Z}^{[1,n]}} & \phi^0(\lambda_i)_{i \in \mathbb{Z}^{[2,n]}} \\ 1 & n \end{array} \right).$$

Since $\phi^1(\lambda_i)_{i \in \mathbb{Z}^{[1,n]}}$ gives $2n$ eigenvalues and $\phi^0(\lambda_i)_{i \in \mathbb{Z}^{[2,n]}}$ gives the rest of the $n(n-1)$ eigenvalues of $G_0^{*(2)}$, we get all the $n(n+1)$ eigenvalues of $G_0^{*(2)}$.

Let the result holds for $G_0^{*(m)} (m > 2)$. Next we prove for $G_0^{*(m+1)} = G_0^{*(m)} \odot_1^n G_0^*$. By Theorem 4.1, we have

$$\begin{aligned} \sigma_A(G_0^{*(m+1)}) &= \left(\begin{array}{cc} \phi(\sigma_A(G_0^{*(m)})) & \phi^0(\lambda_i)_{i \in \mathbb{Z}^{[2,n]}} \\ 1 & n \end{array} \right) \\ &= \left(\begin{array}{cccc} \phi(\phi^{(m-1)}(\lambda_i)_{i \in \mathbb{Z}^{[1,n]}}) & \phi(\phi^{(m-2)}(\lambda_i)_{i \in \mathbb{Z}^{[2,n]}}) & \cdots & \phi(\phi^0(\lambda_i)_{i \in \mathbb{Z}^{[2,n]}}) \\ 1 & n & \cdots & n(n+1)^{m-2} \\ & & & n(n+1)^{m-1} \end{array} \right). \end{aligned} \quad (15)$$

It is clear that $\phi^m(\lambda_i)_{i \in \mathbb{Z}^{[1,n]}}$ gives $2^m n$ eigenvalues and $\phi^j(\lambda_i)_{i \in \mathbb{Z}^{[2,n]}}$ gives $2^j n(n-1)(n+1)^{m-j-1}$ eigenvalues of $G_0^{*(m+1)}$. Therefore, from (15) and Lemma 2.4 we obtain all the $n(n+1)^m$ eigenvalues of $G_0^{*(m+1)}$. Hence the theorem. \square

6 Construction of Seidel-cospectral hypergraphs

Sarkar and Banerjee [13] introduced a method for constructing cospectral k -uniform hypergraphs based on the vertex corona of hypergraphs. Later, in [1], various techniques for generating non-isomorphic adjacency-cospectral hypergraphs were discussed.

In this section, we focus on constructing non-isomorphic Seidel-cospectral k -uniform hypergraphs by extending the method of signed M-GM switching [3] to find a family of switching equivalent hypergraphs.

Let G^* be a k -uniform hypergraph with vertex set V and hyperedge set E , and $\{U_1, U_2, U_3, \dots, U_{2t}, U\}$, $t \in \mathbb{N}$ be the partition of V . Then, proper labelling of the vertices, enables us to express the Seidel matrix of G^* as follows:

$$S(G^*) = \begin{bmatrix} S_{1,1} & S_{1,2} & S_{1,3} & \cdots & S_{1,2t} & S_1 \\ S_{2,1} & S_{2,2} & S_{2,3} & \cdots & S_{2,2t} & S_2 \\ S_{3,1} & S_{3,2} & S_{3,3} & \cdots & S_{3,2t} & S_3 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ S_{2t,1} & S_{2t,2} & S_{2t,3} & \cdots & S_{2t,2t} & S_{2t} \\ S_1^T & S_2^T & S_3^T & \cdots & S_{2t}^T & S_0 \end{bmatrix}.$$

Theorem 6.1. *Let $G^* = (V, E)$ be a k -uniform hypergraph and $\{U_1, U_2, U_3, \dots, U_{2t}, U\}$ be the partition of V that satisfies the following conditions:*

- $|U_i| = m$, $m \in \mathbb{Z}^+$, $0 < i \leq 2t$ and $|U| = k - 1$.
- For any odd i , $N(U) \cap (U_i \cup U_{i+1}) = U_i$, where $N(U) = \{v \in V : v \cup U \in E\}$.

- For any odd i, j ($1 \leq i, j \leq 2t$),

$$\text{for } 1 \leq s \leq m, \sum_{r=1}^m ((S_{i,j})_{rs} - (S_{i,j+1})_{rs}) = \sum_{r=1}^m ((S_{i+1,j+1})_{rs} - (S_{i+1,j})_{rs}) = l,$$

$$\text{for } 1 \leq r \leq m, \sum_{s=1}^m ((S_{i,j})_{rs} - (S_{i+1,j})_{rs}) = \sum_{s=1}^m ((S_{i+1,j+1})_{rs} - (S_{i,j+1})_{rs}) = l.$$

Then a non-isomorphic Seidel cospectral mate k -uniform hypergraph H^* of G^* forms by replacing the edge $\{u_i\} \cup U$, $\forall u_i \in U_i$ with the new edges $\{u_{i+1}\} \cup U$, $u_{i+1} \in U_{i+1}$.

Proof. Consider an orthogonal block matrix P

$$P = \begin{bmatrix} P_1 & P_2 & \mathbf{O} & \mathbf{O} & \cdots & \mathbf{O} \\ P_2 & P_1 & \mathbf{O} & \mathbf{O} & \cdots & \mathbf{O} \\ \mathbf{O} & \mathbf{O} & P_1 & P_2 & \cdots & \mathbf{O} \\ \mathbf{O} & \mathbf{O} & P_2 & P_1 & \cdots & \mathbf{O} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \mathbf{O} & \mathbf{O} & \mathbf{O} & \mathbf{O} & \cdots & I_{k-1} \end{bmatrix} \text{ and } \tilde{S} = \begin{bmatrix} \tilde{S}_{1,1} & \tilde{S}_{1,2} & \cdots & \tilde{S}_{1,2t} & \tilde{S}_1 \\ \tilde{S}_{2,1} & \tilde{S}_{2,2} & \cdots & \tilde{S}_{2,2t} & \tilde{S}_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \tilde{S}_{2t,1} & \tilde{S}_{2t,2} & \cdots & \tilde{S}_{2t,2t} & \tilde{S}_{2t} \\ \tilde{S}_1 & \tilde{S}_2 & \cdots & \tilde{S}_{2t} & \tilde{S}_0 \end{bmatrix}$$

such that $PSP = \tilde{S}$, where $P_1 = I_m - \frac{1}{m}J_m$ and $P_2 = \frac{1}{m}J_m$. Therefore, for any odd i, j

$$\begin{bmatrix} P_1 & P_2 \\ P_2 & P_1 \end{bmatrix} \begin{bmatrix} S_{i,j} & S_{i,j+1} \\ S_{i+1,j} & S_{i+1,j+1} \end{bmatrix} \begin{bmatrix} P_1 & P_2 \\ P_2 & P_1 \end{bmatrix} = \begin{bmatrix} \tilde{S}_{i,j} & \tilde{S}_{i,j+1} \\ \tilde{S}_{i+1,j} & \tilde{S}_{i+1,j+1} \end{bmatrix}.$$

Now we prove that $\tilde{S}_{i,j} = S_{i,j}$

$$\begin{aligned} \tilde{S}_{i,j} &= (P_1 S_{i,j} + P_2 S_{i+1,j})P_1 + (P_1 S_{i,j+1} + P_2 S_{i+1,j+1})P_2 \\ &= S_{i,j} + \frac{1}{m} \left(-J_m(S_{i,j} - S_{i+1,j}) - (S_{i,j} - S_{i,j+1})J_m \right) + \frac{1}{m^2} \left(J_m(S_{i,j} - S_{i,j+1})J_m \right. \\ &\quad \left. + J_m(S_{i+1,j+1} - S_{i+1,j})J_m \right) \\ &= S_{i,j} + \frac{1}{m} \left(-lJ_m - lJ_m \right) + \frac{1}{m^2} \left(lmJ_m + lmJ_m \right) \\ &= S_{i,j}. \end{aligned}$$

From a similar argument we obtain,

$$\begin{bmatrix} \tilde{S}_{i,j} & \tilde{S}_{i,j+1} \\ \tilde{S}_{i+1,j} & \tilde{S}_{i+1,j+1} \end{bmatrix} = \begin{bmatrix} S_{i,j} & S_{i,j+1} \\ S_{i+1,j} & S_{i+1,j+1} \end{bmatrix}.$$

Next, we will show that for an odd i ($0 \leq i \leq 2t$), $\tilde{S}_i = S_{i+1}$ and $\tilde{S}_{i+1} = S_i$ according to which all the values of Seidel matrix corresponding to $k-1$ entries of U and U_i is switched with U_{i+1} . Since $N(U) \cap (U_i \cup U_{i+1}) = U_i$, we can observe that S_i and S_{i+1} are block matrix with all entries equal to -1 and 1 respectively. Then, $J_m S_i = -m J_{m,k-1}$ and $J_m S_{i+1} = m J_{m,k-1}$. Therefore,

$$\tilde{S}_i = P_1 S_i + P_2 S_{i+1} = S_i + 2J_{m,k-1} = S_{i+1}$$

and

$$\tilde{S}_{i+1} = P_2 S_i + P_1 S_{i+1} = S_{i+1} - 2J_{m,k-1} = S_i$$

Hence, we can conclude that S and \tilde{S} are similar matrices corresponding to Seidel-cospectral hypergraphs G^* and H^* , respectively. \square

Example 6.2. Let G^* be a hypergraph with vertex set $V(G^*) = \{v_0, v_1, u_1, u_2, u_3, u_4, u_5, u_6\}$ and edge set $E = \{\{u_1, u_2, u_3\}, \{u_1, u_4, u_5\}, \{u_2, u_5, u_6\}, \{u_3, u_4, u_6\}, \{v_0, v_1, u_1\}, \{v_0, v_1, u_2\}, \{v_0, v_1, u_3\}\}$. Consider the partition $\{U_1 = \{u_1, u_2, u_3\}, U_2 = \{u_4, u_5, u_6\}, U = \{v_0, v_1, \}\}$ of V . Then $N(U) = \{u_1, u_2, u_3\} = U_1$. Now, we get a non-regular non-isomorphic Seidel-cospectral hypergraph H^* with vertex set V and edge set $E/\{\{v_0, v_1, u_1\}, \{v_0, v_1, u_2\}, \{v_0, v_1, u_3\}\} \cup \{\{v_0, v_1, u_4\}, \{v_0, v_1, u_5\}, \{v_0, v_1, u_6\}\}$.

Next, we prove the existence of infinitely many non-isomorphic Seidel-cospectral hypergraphs for any two hypergraphs that are not isomorphic.

Theorem 6.3. Let G_1^* and H_1^* be non-isomorphic cospectral (k, r) -regular hypergraphs of order m . Then $G_0^* \odot_1^n G_1^*$ and $G_0^* \odot_1^n H_1^*$ are non-regular Seidel-cospectral hypergraphs.

Proof. The proof follows from Theorem 3.2.2 and Theorem 4.4. □

7 Conclusion

In this paper, the characteristic polynomial of generalised adjacency and generalised Seidel matrix of the corona of hypergraphs are determined. Also, proposed a method to generate infinitely many non-regular and non-isomorphic Seidel-cospectral hypergraphs. In addition, estimated the adjacency spectrum, Seidel spectrum and adjacency eigenvectors of the corona of two hypergraphs. Corona hypergraphs are constructed by iteratively applying the corona product of the base hypergraph. When the base hypergraph is regular, we determined the adjacency spectrum of corona hypergraphs. Moreover, a new invariant the coronal of the adjacency matrix of a hypergraph, has been introduced to find the spectrum of the corona of hypergraphs in a more computably easier way.

8 Declarations

On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

- [1] Aida Abiad and Antonina P Khramova. Constructing cospectral hypergraphs. *arXiv preprint arXiv:2211.06087*, 2022.
- [2] S. Barik, S. Pati, and B. K. Sarma. The spectrum of the corona of two graphs. *SIAM J. Discrete Math.*, 21(1):47–56, 2007.
- [3] Francesco Belardo, Maurizio Brunetti, Matteo Cavaleri, and Alfredo Donno. Constructing cospectral signed graphs. *Linear Multilinear Algebra*, 69(14):2717–2732, 2021.
- [4] Claude Berge. *Graphs and hypergraphs*. vol. 6, North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York, 1973.
- [5] Joshua Cooper and Aaron Dutle. Spectra of uniform hypergraphs. *Linear Algebra Appl.*, 436(9):3268–3292, 2012.
- [6] Arpita Das and Pratima Panigrahi. Spectra of R -vertex join and R -edge join of two graphs. *Discuss. Math. Gen. Algebra Appl.*, 38(1):19–31, 2018.
- [7] Jiu Ding and Aihui Zhou. Eigenvalues of rank-one updated matrices with some applications. *Appl. Math. Lett.*, 20(12):1223–1226, 2007.

- [8] Roberto W. Frucht and Frank Harary. The corona of two graphs. *Scientia (Valparaíso)*, 139:42–49, 1970.
- [9] TK Jahfar and AV Chithra. Central vertex join and central edge join of two graphs. *AIMS Math.*, 5(6):7214–7234, 2020.
- [10] Reji Kumar and Renny P Varghese. Spectrum of (k, r) -regular hypergraphs. *International J.Math. Combin*, 2:52–59, 2017.
- [11] Hongying Lin and Bo Zhou. Spectral radius of uniform hypergraphs. *Linear Algebra Appl.*, 527:32–52, 2017.
- [12] Cam McLeman and Erin McNicholas. Spectra of coronae. *Linear Algebra Appl.*, 435(5):998–1007, 2011.
- [13] Amitesh Sarkar and Anirban Banerjee. Joins of hypergraphs and their spectra. *Linear Algebra Appl.*, 603:101–129, 2020.
- [14] Rohan Sharma, Bibhas Adhikari, and Abhishek Mishra. Structural and spectral properties of corona graphs. *Discrete Appl. Math.*, 228:14–31, 2017.
- [15] A Y Zakiyyah. Some result on integrality of several matrix representation of complete r -uniform hypergraph. *Journal of Physics: Conference Series*, 2157:012006, 1 2022.