COMMON NEIGHBORHOOD (SIGNLESS) LAPLACIAN SPECTRUM AND ENERGY OF CCC-GRAPH

FIRDOUS EE JANNAT AND RAJAT KANTI NATH*

ABSTRACT. In this paper, we consider commuting conjugacy class graph (abbreviated as CCC-graph) of a finite group G which is a graph with vertex set $\{x^G : x \in G \setminus Z(G)\}$ (where x^G denotes the conjugacy class containing x) and two distinct vertices x^G and y^G are joined by an edge if there exist some elements $x' \in x^G$ and $y' \in y^G$ such that they commute. We compute common neighborhood (signless) Laplacian spectrum and energy of CCC-graph of finite non-abelian groups whose central quotient is isomorphic to either $\mathbb{Z}_p \times \mathbb{Z}_p$ (where p is any prime) or the dihedral group D_{2n} $(n \geq 3)$; and determine whether CCC-graphs of these groups are common neighborhood (signless) Laplacian hyperenergetic/borderenergetic. As a consequence, we characterize certain finite non-abelian groups viz. D_{2n} , T_{4n} , U_{6n} , $U_{(n,m)}$, SD_{8n} and V_{8n} such that their CCC-graphs are common neighborhood (signless) Laplacian hyperenergetic/borderenergetic.

1. INTRODUCTION

Characterizing finite groups through various graphs defined on them have been an active area of research over the last 50 years. A number of graphs have been defined on groups [5]. Among those, in our paper, we consider commuting conjugacy class graph (abbreviated as CCC-graph) of a finite non-abelian group G. For any element $x \in G$, we write x^G to denote the conjugacy class of G containing x. The CCC-graph of G, denoted by Γ_G , is defined as a simple undirected graph whose vertex set is the set of conjugacy classes of non-central elements of G and two vertices x^G and y^G are adjacent if there exists some elements $x' \in x^G$ and $y' \in y^G$ such that x'y' = y'x'. In 2009, Herzog et al. [25] introduced the concept of CCC-graph of a group. In 2016, Mohammadian et al. [30] have characterized finite groups such that their CCC-graph is triangle-free. Later on Salahshour and Ashrafi [32, 33], obtained structures of CCC-graph of several families of finite CA-groups. Salahshour [34] also described Γ_G for the groups whose central quotient is isomorphic to a dihedral group. Characterizations of various classes of finite non-abelian groups through energy, (signless) Laplacian

²⁰²⁰ Mathematics Subject Classification. 05C25, 05C50, 15A18.

Key words and phrases. Common Neighborhood; Spectrum; Energy; Commuting Conjugacy Class Graph.

Corresponding author.

energy, common neighborhood energy (abbreviated as CN-energy) and genus of their CCC-graphs can be found in [3, 4, 26].

The energy of a graph \mathcal{G} (denoted by $E(\mathcal{G})$) is the sum of absolute values of all the eigenvalues of the adjacency matrix of \mathcal{G} . This notion was also used in obtaining π -electron energy of a conjugated carbon molecule in theoretical chemistry. The study of energy of a graph was initiated by Gutman [18] in 1978. After a long time, in 2006 and 2008, Gutman et al. introduced two more graph-energy-like quantity, known as Laplacian energy [23] (denoted by $LE(\mathcal{G})$) and signless Laplacian energy [20] (denoted by $LE^+(\mathcal{G})$), using Laplacian and signless Laplacian eigenvalues of a graph respectively. These energies are used to study various properties of graphs (see [6, 15]). Later on, mathematicians have introduced several kinds of graph energies (see [21, 22]) and studied graph properties. In 2011, Alwardi et al. [2] have introduced CN-energy of a graph. Let \mathcal{G} be a graph with vertex set $V(\mathcal{G}) = \{v_1, v_2, v_3, \dots, v_n\}$. Let $C(v_i, v_j)$ be the set of vertices of a graph \mathcal{G} other than v_i and v_j which are adjacent to both v_i and v_j . Then the common neighborhood matrix (CN-matrix) of \mathcal{G} , denoted by $CN(\mathcal{G})$, is a matrix of size n whose (i, j)-th entry is given by

$$CN(\mathcal{G})_{i,j} = \begin{cases} |C(v_i, v_j)|, & \text{if } i \neq j \\ 0, & \text{otherwise.} \end{cases}$$

The CN-energy of \mathcal{G} is the sum of absolute values of all the eigenvalues of $CN(\mathcal{G})$. Motivated by the study of (signless) Laplacian energy, Jannat et al. [28] have introduced the notions of common neighborhood Laplacian energy (CNL-energy) and common neighborhood signless Laplacian energy (CNSL-energy) of a graph.

The common neighborhood Laplacian matrix (CNL-matrix) and the common neighborhood signless Laplacian matrix (CNSL-matrix) of \mathcal{G} , denoted by $\text{CNL}(\mathcal{G})$ and $\text{CNSL}(\mathcal{G})$, respectively, are given by

 $\operatorname{CNL}(\mathcal{G}) := \operatorname{CNRS}(\mathcal{G}) - \operatorname{CN}(\mathcal{G}) \text{ and } \operatorname{CNSL}(\mathcal{G}) := \operatorname{CNRS}(\mathcal{G}) + \operatorname{CN}(\mathcal{G}),$

where $\text{CNRS}(\mathcal{G})$ is a matrix of size $|V(\mathcal{G})| = n$ whose (i, j)-th entry is given by

$$\operatorname{CNRS}(\mathcal{G})_{i,j} = \begin{cases} \sum_{k=1}^{n} \operatorname{CN}(\mathcal{G})_{i,k}, & \text{if } i = j \text{ and } i = 1, 2, \dots, n \\ 0, & \text{if } i \neq j. \end{cases}$$

The common neighborhood Laplacian spectrum of \mathcal{G} (abbreviated as CNL-spectrum and denoted by CNL-spec(\mathcal{G})) is the set of eigenvalues of CNL(\mathcal{G}) with multiplicities. We write CNL-spec(\mathcal{G}) = { $(\alpha_1)^{a_1}, (\alpha_2)^{a_2}, \ldots, (\alpha_k)^{a_k}$ }, where $\alpha_1, \alpha_2, \ldots, \alpha_k$ are the distinct eigenvalues of CNL(\mathcal{G}) with corresponding multiplicities a_1, a_2, \ldots, a_k . Similarly, common neighborhood signless Laplacian spectrum of \mathcal{G} (abbreviated as CNSL-spectrum and denoted by CNSL-spec(\mathcal{G})) is the set of eigenvalues of CNSL(\mathcal{G}) with multiplicities. We write CNSL-spec(\mathcal{G}) = { $(\beta_1)^{b_1}, (\beta_2)^{b_2}, \ldots, (\beta_\ell)^{b_\ell}$ }, where $\beta_1, \beta_2, \ldots, \beta_\ell$ are the distinct eigenvalues of $\text{CNSL}(\mathcal{G})$ with corresponding multiplicities b_1, b_2, \ldots, b_ℓ . A graph \mathcal{G} is called CNL-integral (CNSL-integral) if CNL-spectrum (CNSL-spectrum) contains only integers. The notions of CNL-integral and CNSLintegral graphs were introduced in [28] motivated by the notions of integral (introduced by Harary and Schwenk [24]), L-integral (introduced by Grone and Merris [17]), Q-integral (introduced by Simic and Stanic [35]) and CN-integral (introduced by Alwardi et al. [2]) graphs. A finite graph is called super integral if it is integral, L-integral and Q-integral (see [3]). Integral graphs have some interests for designing the network topology of perfect state transfer networks (see [1] and the references there in).

The CNL-energy and CNSL-energy of \mathcal{G} , denoted by $LE_{CN}(\mathcal{G})$ and $LE_{CN}^+(\mathcal{G})$ respectively, are defined as

(1)
$$LE_{CN}(\mathcal{G}) := \sum_{i=1}^{k} a_i |\alpha_i - \Delta(\mathcal{G})|$$

and

(2)
$$LE_{CN}^{+}(\mathcal{G}) := \sum_{i=1}^{\ell} b_i \left| \beta_i - \Delta(\mathcal{G}) \right|,$$

where $\Delta(\mathcal{G}) = \frac{\operatorname{tr}(\operatorname{CNRS}(\mathcal{G}))}{|V(\mathcal{G})|}$ and $\operatorname{tr}(\operatorname{CNRS}(\mathcal{G}))$ is the trace of $\operatorname{CNRS}(\mathcal{G})$. In [28], various facets of the CNL-spectrum, CNL-energy, CNSL-spectrum and CNSL-energy of graphs were discussed; their connections with other well-known graph energies and Zagreb indices were also established. It was observed that

(3)
$$LE_{CN}(K_n) = LE_{CN}^+(K_n) = 2(n-1)(n-2),$$

where K_n is the complete graph of order n. A graph \mathcal{G} of order n is called CNLhyperenergetic or CNSL-hyperenergetic according as $LE_{CN}(\mathcal{G}) > 2(n-1)(n-2)$ or $LE_{CN}^+(\mathcal{G}) > 2(n-1)(n-2)$. Further, it is called CNL-borderenergetic (CNSLborderenergetic) if CNL-energy (CNSL-energy) of \mathcal{G} is equal to 2(n-1)(n-2). These classes of graphs were introduced in [27, 28] motivated by the notions of various types of hyperenergetic graphs (see [2, 13, 16, 19, 37, 38, 39]).

In Section 2, we compute CNL-spectrum, CNSL-spectrum, CNL-energy and CNSLenergy of CCC-graphs of finite non-abelian groups whose central quotient is isomorphic to either $\mathbb{Z}_p \times \mathbb{Z}_p$ (where p is any prime) or the dihedral group D_{2n} $(n \geq 3)$. In Section 3, we determine whether CCC-graphs of these groups are CNL-integral, CNSL-integral, CNL-hyperenergetic, CNSL-hyperenergetic, CNL-borderenergetic and CNSL-borderenergetic. As a consequence, we characterize the groups viz. D_{2n} , T_{4n} , U_{6n} , $U_{(n,m)}$, SD_{8n} and V_{8n} such that their CCC-graphs have above mentioned properties. In Subsection 3.1, we compare various CN-energies of CCC-graphs of the groups

F. E. JANNAT AND R. K. NATH

G considered in Section 2 and show that $E_{CN}(\Gamma_G) = LE_{CN}(\Gamma_G) = LE_{CN}^+(\Gamma_G)$ or $E_{CN}(\Gamma_G) < LE_{CN}^+(\Gamma_G) < LE_{CN}(\Gamma_G)$. We also characterize the groups viz. D_{2n} , T_{4n} , U_{6n} , $U_{(n,m)}$, SD_{8n} and V_{8n} such that their CCC-graphs satisfy above mentioned equality/inequality. Finally, we conclude the paper in Section 4 by listing certain problems that arise naturally after our investigation.

2. Computations of spectrum and energies

In this section, we compute CNL-spectrum, CNSL-spectrum and their respective energies of CCC-graphs of various families of non-abelian finite groups. In particular, we consider finite non-abelian groups whose central quotients are isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$ (where p is any prime) or $D_{2n} = \langle x, y : x^n = y^2 = 1, yxy^{-1} = x^{-1} \rangle$ (for $n \ge 3$) in the following subsections. We shall also consider a generalization of dihedral groups, namely $U_{(n,m)} = \langle x, y : x^{2n} = y^m = 1, x^{-1}yx = y^{-1} \rangle$ (for $n \ge 2, m \ge 2$) along with the dicyclic group $T_{4n} = \langle x, y : x^{2n} = 1, x^n = y^2, y^{-1}xy = x^{-1} \rangle$ (for $n \ge 2$), the semidihedral groups $SD_{8n} = \langle x, y : x^{4n} = y^2 = 1, yxy = x^{2n-1} \rangle$ (for $n \ge 2$), the groups $U_{6n} = \langle x, y : x^{2n} = y^3 = 1, x^{-1}yx = y^{-1} \rangle$ (for $n \ge 2$) and $V_{8n} = \langle x, y : x^{2n} = y^4 = 1, yx = x^{-1}y^{-1}, y^{-1}x = x^{-1}y \rangle$ (for $n \ge 2$). The following result is useful in our computation.

Theorem 2.1. [28] Let $\mathcal{G} = l_1 K_{m_1} \cup l_2 K_{m_2} \cup l_3 K_{m_3}$, where $l_i K_{m_i}$ denotes the disjoint union of l_i copies of K_{m_i} for i = 1, 2, 3. Then

CNL-spec(
$$\mathcal{G}$$
)
= { $(0)^{l_1+l_2+l_3}, (m_1(m_1-2))^{l_1(m_1-1)}, (m_2(m_2-2))^{l_2(m_2-1)}, (m_3(m_3-2))^{l_3(m_3-1)}$ }

and

CNSL-spec(
$$\mathcal{G}$$
) = { $(2(m_1 - 1)(m_1 - 2))^{l_1}, ((m_1 - 2)^2)^{l_1(m_1 - 1)}, (2(m_2 - 1)(m_2 - 2))^{l_2}, ((m_2 - 2)^2)^{l_2(m_2 - 1)}, (2(m_3 - 1)(m_3 - 2))^{l_3}, ((m_3 - 2)^2)^{l_3(m_3 - 1)}$ }.

2.1. Groups whose central quotient is isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$. This class of groups have been considered by Salahshour and Ashrafi [32, Theorem 3.1] and showed that

(4)
$$\Gamma_G = (p+1)K_{\underline{(p-1)|Z(G)|}}.$$

CN-energy of CCC-graphs of this class of groups have been studied in [26]. It is worth mentioning that commuting and non-commuting graphs of this class of groups are also studied in [7, 8, 9, 10, 11, 14, 12, 31]. In the following theorem, we derive CNL-spectrum, CNSL-spectrum, CNL-energy and CNSL-energy of CCC-graphs of this class of groups. **Theorem 2.2.** Let G be a finite non-abelian group with $|Z(G)| = z \ge 2$ and $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$, where p is a prime. Then CNL-spectrum, CNSL-spectrum, CNL-energy and CNSL-energy of CCC-graph of G are given by

CNL-spec(
$$\Gamma_G$$
) = $\left\{ (0)^{p+1}, \left(\frac{1}{p^2} (pz - z)(pz - z - 2p) \right)^{\frac{(p+1)}{p}(pz - z - p)} \right\},$

 $\text{CNSL-spec}(\Gamma_G) =$

$$\left\{ \left(\frac{2}{p^2}(pz-z-p)(pz-z-2p)\right)^{p+1}, \left(\frac{1}{p^2}(pz-z-2p)^2\right)^{\frac{(p+1)}{p}(pz-z-p)} \right\}$$

and

$$LE_{CN}(\Gamma_G) = LE_{CN}^+(\Gamma_G) = \begin{cases} \frac{3}{2}, & \text{for } p = 2 \& z = 3\\ \frac{4(p-2)(p+1)}{p^2}, & \text{for } p \ge 2 \& z = 2\\ \frac{2(p+1)(p(z-2)-z)(p(z-1)-z)}{p^2}, & \text{otherwise.} \end{cases}$$

Proof. From (4), we have $\Gamma_G = (p+1)K_n$, where $n = \frac{(p-1)z}{p}$. Therefore, by Theorem 2.1, we get

CNL-spec(Γ_G) = {(0)^{p+1}, ($\frac{1}{p^2}(pz-z)(pz-z-2p)$)^{$\frac{(p+1)}{p}(pz-z-p)$}} and CNSL-spec(Γ_G) = {($\frac{2}{p^2}(pz-z-p)(pz-z-2p)$)^{p+1}, ($\frac{1}{p^2}(pz-z-2p)^2$)^{$(p+1)\frac{1}{p}(pz-z-p)$}}. Here $|V(\Gamma_G)| = \frac{(p^2-1)z}{p}$ and tr(CNRS(Γ_G)) = $\frac{(p-1)(p+1)z(p(z-2)-z)(p(z-1)-z)}{p^3}$. Therefore, $\Delta(\Gamma_G) = \frac{(p(z-2)-z)(p(z-1)-z)}{p^2}$.

Now we calculate CNL-energy of Γ_G . We have

$$L_1 := |0 - \Delta(\Gamma_G)| = \left| -\frac{(p(z-2) - z)(p(z-1) - z)}{p^2} \right|.$$

Let $\alpha_1(p, z) = -(p(z-2)-z)(p(z-1)-z)$. Then $\alpha_1(p, z) = -2p^2 - 3pz - z^2 + \frac{1}{2}p^2 z(6-z) + \frac{1}{2}pz^2(4-p) < 0$ for $p \ge 4$ and $z \ge 6$. It can be seen that $\alpha_1(2, z) = -z(z-6)-8 = 1$ or ≤ 0 according as z = 3 or $z \ne 3$; $\alpha_1(3, z) = -2(z-3)(2z-3) = 2$ or ≤ 0 according as z = 2 or $z \ne 2$; $\alpha_1(p, 2) = 2p - 4 \ge 0$; $\alpha_1(p, 3) = -2p^2 + 9p - 9 = 1$ or ≤ 0 according as p = 2 or $p \ne 2$; $\alpha_1(p, 4) = -6p^2 + 20p - 16 \le 0$ and $\alpha_1(p, 5) = -12p^2 + 35p - 25 \le 0$. Therefore

$$L_1 = \begin{cases} -\frac{(p(z-2)-z)(p(z-1)-z)}{p^2}, & \text{for } p = 2 \& z = 3; p \ge 2 \& z = 2\\ \frac{(p(z-2)-z)(p(z-1)-z)}{p^2}, & \text{otherwise.} \end{cases}$$

Also

$$L_2 := \left| \frac{(pz-z)(pz-2p-z)}{p^2} - \Delta(\Gamma_G) \right| = \left| \frac{pz-2p-z}{p} \right|.$$

Let $\alpha_2(p, z) = pz - 2p - z$. Then $\alpha_2(p, z) = (z - 2)p - z \ge z - 4 \ge 0$ for all $z \ge 4$ since $p \ge 2$. It can be seen that $\alpha_2(p, 2) = -2 < 0$ and $\alpha_2(p, 3) = p - 3 \ge 0$ or < 0 according as $p \ge 3$ or p = 2. Therefore

$$L_{2} = \begin{cases} -\frac{pz-2p-z}{p}, & \text{for } p = 2 \& z = 3; p \ge 2 \& z = 2\\ \frac{pz-2p-z}{p}, & \text{otherwise.} \end{cases}$$

Hence, by (1), we get

$$LE_{CN}(\Gamma_G) = (p+1) \times L_1 + \frac{p+1}{p}(pz - z - p) \times L_2$$
$$= \begin{cases} \frac{3}{2}, & \text{for } p = 2 \& z = 3\\ \frac{4(p-2)(p+1)}{p^2}, & \text{for } p \ge 2 \& z = 2\\ \frac{2(p+1)(p(z-2)-z)(p(z-1)-z)}{p^2}, & \text{otherwise.} \end{cases}$$

For CNSL-energy of Γ_G we have

$$B_{1} := \left| \frac{2(pz - p - z)(pz - 2p - z)}{p^{2}} - \Delta(\Gamma_{G}) \right| = \left| \frac{(pz - 2p - z)(pz - p - z)}{p^{2}} \right|$$
$$= -L_{1} = \begin{cases} -\frac{(pz - 2p - z)(pz - p - z)}{p^{2}}, & \text{for } p = 2 \& z = 3; p \ge 2 \& z = 2\\ \frac{(pz - 2p - z)(pz - p - z)}{p^{2}}, & \text{otherwise.} \end{cases}$$

Also

$$B_{2} := \left| \frac{(pz - 2p - z)^{2}}{p^{2}} - \Delta(\Gamma_{G}) \right| = \left| \frac{-pz + 2p + z}{p} \right| = -L_{2}$$
$$= \begin{cases} \frac{2p + z - pz}{p}, & \text{for } p = 2 \& z = 3; p \ge 2 \& z = 2\\ -\frac{2p + z - pz}{p}, & \text{otherwise.} \end{cases}$$

Hence, by (2), we get

$$LE_{CN}^{+}(\Gamma_{G}) = (p+1) \times B_{1} + \frac{p+1}{p}(pz-z-p) \times B_{2}$$
$$= \begin{cases} \frac{3}{2}, & \text{for } p = 2 \& z = 3\\ \frac{4(p-2)(p+1)}{p^{2}}, & \text{for } p \ge 2 \& z = 2\\ \frac{2(p+1)(p(z-2)-z)(p(z-1)-z)}{p^{2}}, & \text{otherwise.} \end{cases}$$

Hence the result follows.

As a corollary of the above theorem we get the following result.

Corollary 2.3. Let G be a non-abelian group of order p^n with $|Z(G)| = p^{n-2}$. Then CNL-spectrum, CNSL-spectrum, CNL-energy and CNSL-energy of CCC-graph of G are given by

$$CNL-spec(\Gamma_G) = \left\{ (0)^{p+1}, \left(\frac{1}{p^2} (p^{n-1} - p^{n-2})(p^{n-1} - p^{n-2} - 2p) \right)^{\frac{(p+1)}{p} \left(p^{n-1} - p^{n-2} - p \right)} \right\},$$

$$CNSL-spec(\Gamma_G) = \left\{ \left(\frac{2}{p^2} (p^{n-1} - p^{n-2} - p)(p^{n-1} - p^{n-2} - 2p) \right)^{p+1}, \left(\frac{1}{p^2} (p^{n-1} - p^{n-2} - 2p)^2 \right)^{\frac{(p+1)}{p} \left(p^{n-1} - p^{n-2} - p \right)} \right\}$$

and

$$LE_{CN}(\Gamma_G) = LE_{CN}^+(\Gamma_G) = \begin{cases} 0, & \text{for } p = 2 \& n = 3\\ \frac{2(p+1)\left(p\left(p^{n-2}-2\right)-p^{n-2}\right)\left(p\left(p^{n-2}-1\right)-p^{n-2}\right)}{p^2}, & \text{otherwise.} \end{cases}$$

Proof. Here $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$. Hence the result follows from Theorem 2.2.

2.2. Groups whose central quotient is isomorphic to a dihedral group. The CCC-graph of this class of group was first studied by Salahshour [34] in 2020. Salahshour [34, Theorem 1.2] obtained the following structures of Γ_G (where $\frac{G}{Z(G)} \cong D_{2n}$)

(5)
$$\Gamma_G = \begin{cases} K_{\frac{(n-1)|Z(G)|}{2}} \cup 2K_{\frac{|Z(G)|}{2}}, & \text{if } 2 \mid n \\ K_{\frac{(n-1)|Z(G)|}{2}} \cup K_{|Z(G)|}, & \text{if } 2 \nmid n. \end{cases}$$

The CN-energy of such Γ_G was studied in [26]. Also spectrum, L-spectrum and Q-spectrum of commuting and non-commuting graphs of this class of groups was studied in [7, 8, 9, 10, 11, 12] along with their respective energies.

Theorem 2.4. Let G be a finite group such that |Z(G)| = z and $\frac{G}{Z(G)} \cong D_{2n}$ (where $n \ge 3$). Then CNL-spectrum, CNSL-spectrum, CNL-energy and CNSL-energy of CCC-graph of G are as given below:

(a) If n is even then

(i) CNL-spec(
$$\Gamma_G$$
) = $\left\{ (0)^3, (\frac{1}{4}(nz-z)(nz-z-4))^{\frac{1}{2}(nz-z-2)}, (\frac{1}{4}z(z-4))^{z-2} \right\}$ and
 $LE_{CN}(\Gamma_G) = \frac{((n-1)z-2)(n(z+1)((n-2)z-4)+11z-4)}{2(n+1)}.$

(ii) CNSL-spec $(\Gamma_G) = \left\{ \left(\frac{1}{2}(nz-z-2)(nz-z-4)\right)^1, \left(\frac{1}{4}(nz-z-4)^2\right)^{\frac{1}{2}(nz-z-2)}, \right\}$

 $\overline{7}$

 $\left(\frac{1}{2}(z-2)(z-4)\right)^2, \left(\frac{1}{4}(z-4)^2\right)^{z-2}$

and

$$LE_{CN}^{+}(\Gamma_G) = \begin{cases} \frac{28}{5}, & \text{for } n = 4 \& z = 2\\ \frac{3}{5}z^2(4z-6), & \text{for } n = 4 \& z \ge 3\\ \frac{(n-2)(n-1)z^2(nz-6)}{2(n+1)}, & \text{otherwise.} \end{cases}$$

(b) If n is odd then

$$\begin{array}{ll} \text{(i) CNL-spec}(\Gamma_G) = \left\{ (0)^2, (\frac{1}{4}(nz-z)(nz-z-4))^{\frac{1}{2}(nz-z-2)}, (z(z-2))^{z-1} \right\} \\ and \ LE_{CN}(\Gamma_G) = \left\{ \begin{array}{ll} 0, & \text{for } n = 3 \ \& z = 1 \\ 4(z-1)(z-2), & \text{for } n = 3 \ \& z \ge 2 \\ \frac{((n-1)z-2)((n-3)(n+1)z^2+((n-6)n+17)z-4(n+1))}{2(n+1)}, & \text{otherwise.} \end{array} \right. \\ (\text{ii) CNSL-spec}(\Gamma_G) = \left\{ (\frac{1}{2}(nz-z-2)(nz-z-4))^1, (\frac{1}{4}(nz-z-4)^2)^{\frac{1}{2}(nz-z-2)}, \\ & (2(z-1)(z-2))^1, ((z-2)^2)^{z-1} \right\} \\ and \ LE_{CN}^+(\Gamma_G) = \left\{ \begin{array}{ll} 0, & \text{for } n = 3 \ \& z = 1; \\ 4(z-1)(z-2), & \text{for } n = 3 \ \& z \ge 2 \\ \frac{(n-5)(n-3)(n+3)}{2(n+1)}, & \text{for } n \ge 7 \ \& z = 1 \\ \frac{(n-3)(n-1)z^2(nz+z-6)}{2(n+1)}, & \text{otherwise.} \end{array} \right. \end{array} \right.$$

Proof. From (5), we have $\Gamma_G = K_{\frac{(n-1)z}{2}} \cup 2K_{\frac{z}{2}}$ or $K_{\frac{(n-1)z}{2}} \cup K_z$ according as n is even or odd.

(a)(i) If n is even, then by Theorem 2.1

CNL-spec(
$$\Gamma_G$$
) = $\left\{ (0)^1, \left(\frac{(n-1)z}{2} \left(\frac{(n-1)z}{2} - 2 \right) \right)^{\frac{(n-1)z}{2} - 1}, (0)^2, \left(\frac{z}{2} \left(\frac{z}{2} - 2 \right) \right)^{2(\frac{z}{2} - 1)} \right\}.$

Here $|V(\Gamma_G)| = \frac{1}{2}(n+1)z$ and $\operatorname{tr}(\operatorname{CNRS}(\Gamma_G)) = \frac{1}{8}z((n((n-3)n+3)+1)z^2 - 6((n-2)n+3)z + 8(n+1)))$. So, $\Delta(\Gamma_G) = \frac{(n((n-3)n+3)+1)z^2 - 6((n-2)n+3)z + 8(n+1))}{4(n+1)}$. Note that $z \ge 2$. We have

$$L_1 := |0 - \Delta(\Gamma_G)| = \left| -\frac{(n((n-3)n+3)+1)z^2 - 6((n-2)n+3)z + 8(n+1))}{4(n+1)} \right|.$$

Let $\alpha_1(n,z) = (n((n-3)n+3)+1)z^2 - 6((n-2)n+3)z + 8(n+1)$. Then $\alpha_1(n,z) = 8+8n+6z(2n-3)+z^2+3nz^2+\frac{n^2z}{2}(nz-12)+\frac{n^2z^2}{2}(n-6) > 0$ for $n \ge 12$. Also, $\alpha_1(4,z) = 29z^2 - 66z + 40 \ge 0$, $\alpha_1(6,z) = 127z^2 - 162z + 56 \ge 0$, $\alpha_1(8,z) = 345z^2 - 306z + 72 \ge 0$ and $\alpha_1(10,z) = 731z^2 - 498z + 88 \ge 0$. Therefore

$$L_1 = \frac{(n((n-3)n+3)+1)z^2 - 6((n-2)n+3)z + 8(n+1)}{4(n+1)}$$

We have

$$L_2 := \left| \frac{1}{4} (nz - z)(nz - z - 4) - \Delta(\Gamma_G) \right| = \left| \frac{n(z+1)((n-2)z - 4) + 11z - 4}{2(n+1)} \right|.$$

Let $\alpha_2(n, z) = \{n(z+1)((n-2)z-4)+11z-4\}$. Then $\alpha_2(n, z) > 0$ for $n \ge 6$, since $n-2 \ge 4 \implies z(n-2)-4 \ge 0 \implies n(z+1)(z(n-2)-4) \ge 0$. Also, $\alpha_2(4, z) = 8z^2 + 3z - 20 \ge 0$. Therefore

$$L_2 = \frac{n(z+1)((n-2)z-4) + 11z - 4}{2(n+1)}$$

We have

$$L_3 := \left| \frac{1}{4} z(z-4) - \Delta(\Gamma_G) \right| = \left| \frac{14z - 8 - 8n - nz(2(z+8) + n(-6 + (n-3)z))}{4(n+1)} \right|.$$

Let $\alpha_3(n, z) = 14z - 8 - 8n - nz(2(z+8) + n(-6 + (n-3)z))$. For $n \ge 10$, n((n-3)z-6) > 0and 2(z+8) > 0. So, $\alpha_3(n, z) < 0$ for all $n \ge 10$. Also, $\alpha_3(4, z) = -24z^2 + 46z - 40 \le 0$, $\alpha_3(6, z) = -120z^2 + 134z - 56 \le 0$ and $\alpha_3(8, z) = -336z^2 + 270z - 72 \le 0$. Therefore

$$L_3 = -\frac{14z - 8 - 8n - nz(2(z+8) + n(-6 + (n-3)z))}{4(n+1)}.$$

Hence, by (1), we get

$$LE_{CN}(\Gamma_G) = 3 \times L_1 + \frac{1}{2}(nz - z - 2) \times L_2 + (z - 2) \times L_3$$
$$= \frac{((n - 1)z - 2)(n(z + 1)((n - 2)z - 4) + 11z - 4)}{2(n + 1)}.$$

(a)(ii) If n is even, then by Theorem 2.1

CNSL-spec
$$(\Gamma_G) = \left\{ \left(2\left(\frac{(n-1)z}{2} - 1\right) \left(\frac{(n-1)z}{2} - 2\right) \right)^1, \left(\left(\frac{(n-1)z}{2} - 2\right)^2 \right)^{\frac{(n-1)z}{2} - 1}, \left(\frac{1}{2}(z-2)(z-4)\right)^2, \left(\frac{1}{4}(z-4)^2\right)^{z-2} \right\}. \right\}$$

Here $|V(\Gamma_G)| = \frac{1}{2}(n+1)z$ and $\operatorname{tr}(\operatorname{CNRS}(\Gamma_G) = \frac{1}{8}z((n((n-3)n+3)+1)z^2 - 6((n-2)n+3)z + 8(n+1)))$. So, $\Delta(\Gamma_G) = \frac{(n((n-3)n+3)+1)z^2 - 6((n-2)n+3)z + 8(n+1))}{4(n+1)}$. Note that $z \ge 2$. We have

$$B_1 := \left| \frac{1}{2} (nz - z - 2)(nz - z - 4) - \Delta(\Gamma_G) \right|$$
$$= \left| \frac{(n (n^2 + n - 5) + 1) z^2 - 6n(n+2)z + 8n + 30z + 8}{4(n+1)} \right|$$

Let $\beta_1(n,z) = (n(n^2 + n - 5) + 1)z^2 - 6n(n + 2)z + 8n + 30z + 8$. Then $\beta_1(n,z) = 8 + 8n + 30z + z^2 + nz^2(n-5) + \frac{nz}{2}(n^2z - 24) + \frac{n^2z}{2}(nz - 12)$. Clearly for $n \ge 12$, $\beta_1(n,z) > 0$,

as $n^2 z - 24 \ge 0$ and $nz - 12 \ge 0$. It can be seen that $\beta_1(4, z) = 61z^2 - 114z + 40 \ge 0$, $\beta_1(6, z) = 223z^2 - 258z + 56 \ge 0$, $\beta_1(8, z) = 537z^2 - 450z + 72 \ge 0$ and $\beta_1(10, z) = 1051z^2 - 690z + 88 \ge 0$. Therefore

$$B_1 = \frac{\left(n\left(n^2 + n - 5\right) + 1\right)z^2 - 6n(n+2)z + 8n + 30z + 8}{4(n+1)}.$$

We have

$$B_2 := \left| \frac{1}{4} (nz - z - 4)^2 - \Delta(\Gamma_G) \right| = \left| \frac{n((n-2)z^2 - (n+6)z + 4) + 13z + 4}{2(n+1)} \right|$$

Let $\beta_2(n, z) = n((n-2)z^2 - (n+6)z + 4) + 13z + 4$. Then $\beta_2(n, z) = 4 + 4n + 13z + \frac{nz}{3}(nz - 18) + \frac{n^2z}{3}(z-3) + \frac{nz^2}{3}(n-6) > 0$ for $n \ge 6$ and $z \ge 3$. It can be seen that $\beta_2(4, z) = 8z^2 - 27z + 20 = -2$ or ≥ 0 according as z = 2 or $z \ge 3$ and $\beta_2(n, 2) = 2n(n-8) + 30 = -2$ or ≥ 0 according as n = 4 or $n \ge 6$. Therefore

$$B_2 = \begin{cases} \frac{1}{5}, & \text{for } n = 4 \& z = 2\\ \frac{n((n-2)z^2 - (n+6)z + 4) + 13z + 4}{2(n+1)}, & \text{otherwise.} \end{cases}$$

We have

$$B_3 := \left| \frac{1}{2} (z-2)(z-4) - \Delta(\Gamma_G) \right|$$
$$= \left| \frac{-\left(\left(n^3 - 3n^2 + n - 1 \right) z^2 \right) + 6(n-4)nz + 8n + 6z + 8}{4(n+1)} \right|.$$

Let $\beta_3(n,z) = -((n^3 - 3n^2 + n - 1)z^2) + 6(n - 4)nz + 8n + 6z + 8$. Then $\beta_3(n,z) = 8(1 - nz) + 8n(1 - z) + 2z(3 - 4n) + z^2(1 - n) + \frac{n^2z}{2}(12 - nz) + \frac{n^2z^2}{2}(6 - n) < 0$ for $n \ge 12$. It can be seen that $\beta_3(4,z) = -19z^2 + 6z + 40 \le 0$, $\beta_3(6,z) = -113z^2 + 78z + 56 \le 0$, $\beta_3(8,z) = -327z^2 + 198z + 72 \le 0$ and $\beta_3(10,z) = -709z^2 + 366z + 88 \le 0$. Therefore

$$B_3 = -\frac{-\left(\left(n^3 - 3n^2 + n - 1\right)z^2\right) + 6(n-4)nz + 8n + 6z + 8}{4(n+1)}$$

We have

$$B_4 := \left| \frac{1}{4} (z-4)^2 - \Delta(\Gamma_G) \right| = \left| \frac{n(8 - z(n((n-3)z-6) + 2(z+10))) + 10z + 8)}{4(n+1)} \right|$$

Let $\beta_4(n,z) = n(8 - z(n((n-3)z-6) + 2(z+10))) + 10z + 8$. Then $\beta_4(n,z) = 8n - 10nz + 10z - 10nz + 8 - 2nz^2 + \frac{n^2z}{2}(12 - nz) + \frac{n^2z^2}{2}(6 - n) < 0$ for $n \ge 12$. It can be seen that $\beta_4(4,z) = -24z^2 + 26z + 40 \le 0, \ \beta_4(6,z) = -120z^2 + 106z + 56 \le 0, \ \beta_4(8,z) = -336z^2 + 234z + 72 \le 0$ and $\beta_4(10,z) = -720z^2 + 410z + 88 \le 0$. Therefore

$$B_4 = -\frac{n(8 - z(n((n-3)z - 6) + 2(z+10))) + 10z + 8}{4(n+1)}$$

Hence, by (2), we get

$$LE_{CN}^{+}(\Gamma_{G}) = 1 \times B_{1} + \frac{1}{2}(nz - z - 2) \times B_{2} + 2 \times B_{3} + (z - 2) \times B_{4}$$
$$= \begin{cases} \frac{28}{5}, & \text{for } n = 4 \& z = 2\\ \frac{3}{5}z^{2}(4z - 6), & \text{for } n = 4 \& z \ge 3\\ \frac{(n-2)(n-1)z^{2}(nz - 6)}{2(n+1)}, & \text{otherwise.} \end{cases}$$

(b)(i) If n is odd, then by Theorem 2.1

CNL-spec
$$(\Gamma_G) = \left\{ (0)^1, \left(\frac{(n-1)z}{2} \left(\frac{(n-1)z}{2} - 2 \right) \right)^{\frac{(n-1)z}{2} - 1}, (0)^1, (z(z-2))^{z-1} \right\}.$$

Here $|V(\Gamma_G)| = \frac{1}{2}(n+1)z$ and $\operatorname{tr}(\operatorname{CNRS}(\Gamma_G) = \frac{1}{8}z(nz+z-4)(((n-4)n+7)z-2(n+1)).$ So, $\Delta(\Gamma_G) = \frac{(nz+z-4)(((n-4)n+7)z-2(n+1))}{4(n+1)}$. We have

$$L'_{1} := |0 - \Delta(\Gamma_{G})| = \left| -\frac{(nz + z - 4)(((n-4)n + 7)z - 2(n+1)))}{4(n+1)} \right|$$

Let $\alpha'_1(n,z) = (nz+z-4)(((n-4)n+7)z-2(n+1))$. Then $\alpha'_1(n,z) = (nz+z-4)(7z-2+\frac{nz}{2}(n-8)+\frac{n}{2}(nz-4)) > 0$ for $n \ge 8$, since $z \ge 1$. Again $\alpha'_1(3,z) = 16z^2-48z+32 \ge 0$, $\alpha'_1(5,z) = 72z^2-120z+48 \ge 0$ and $\alpha'_1(7,z) = 224z^2-240z+64 \ge 0$, as $z \ge 1$. Therefore (nz+z-4)(((n-4)n+7)z-2(n+1))

$$L'_{1} = \frac{(nz+z-4)(((n-4)n+7)z-2(n+1))}{4(n+1)}$$

We have

$$L'_{2} := \left| \frac{1}{4} (nz - z)(nz - z - 4) - \Delta(\Gamma_{G}) \right| = \left| \frac{n^{2}z^{2} + n^{2}z - 2nz^{2} - 6nz - 4n - 3z^{2} + 17z - 4}{2(n+1)} \right|$$

Let $\alpha'_2(n,z) = n^2 z^2 + n^2 z - 2nz^2 - 6nz - 4n - 3z^2 + 17z - 4$. Then $\alpha'_2(n,z) = 17z - 4 + \frac{n}{2}(nz-8) + \frac{nz}{2}(n-12) + \frac{z^2}{2}(n^2-6) + \frac{nz^2}{2}(n-4) > 0$ for $n \ge 8$. It can be seen that $\alpha'_2(3,z) = 8z - 16 = -8$ or ≥ 0 according as z = 1 or $z \ge 2$; $\alpha'_2(5,z) = 12z^2 + 12z - 24 \ge 0$ and $\alpha'_2(7,z) = 32z^2 + 24z - 32 \ge 0$, as $z \ge 1$. Therefore

$$L'_{2} = \begin{cases} 1, & \text{for } n = 3 \& z = \\ \frac{n^{2}z^{2} + n^{2}z - 2nz^{2} - 6nz - 4n - 3z^{2} + 17z - 4}{2(n+1)}, & \text{otherwise.} \end{cases}$$

1

We have

$$L'_{3} := |z(z-2) - \Delta(\Gamma_{G})| = \left| \frac{-n^{3}z^{2} + 3n^{2}z^{2} + 6n^{2}z + nz^{2} - 20nz - 8n - 3z^{2} + 22z - 8}{4(n+1)} \right|.$$

Let $\alpha'_3(n,z) = -n^3 z^2 + 3n^2 z^2 + 6n^2 z + nz^2 - 20nz - 8n - 3z^2 + 22z - 8$. Then $\alpha'_3(n,z) = -8 - 8n - 2z(10n - 11) - 3z^2 - \frac{n^2 z}{3}(nz - 18) - \frac{nz^2}{3}(n^2 - 3) - \frac{n^2 z^2}{3}(n - 9) < 0$ for $n \ge 19$. It can be seen that $\alpha'_3(3,z) = 16z - 32 = -16$ or ≥ 0 according as z = 1 or $z \ge 2$, $\alpha'_3(5,z) = -48z^2 + 72z - 48 \le 0$, $\alpha'_3(7,z) = -192z^2 + 176z - 64 \le 0$, $\alpha'_3(9,z) = -480z^2 + 176z - 64 \le 0$.

 $\begin{array}{l} 328z-80\leq 0,\,\alpha_3'(11,z)=-960z^2+528z-96\leq 0,\,\alpha_3'(13,z)=-1680z^2+776z-112\leq 0,\\ \alpha_3'(15,z)=-2688z^2+1072z-128\,\leq\, 0 \ \text{and} \ \alpha_3'(17,z)=-4032z^2+1416z-144\,\leq\, 0.\\ \text{Therefore} \end{array}$

$$L'_{3} = \begin{cases} z - 2, & \text{for } n = 3 \& z \ge 2\\ -\frac{-n^{3}z^{2} + 3n^{2}z^{2} + 6n^{2}z + nz^{2} - 20nz - 8n - 3z^{2} + 22z - 8}{4(n+1)}, & \text{otherwise.} \end{cases}$$

Hence, by (1), we get

$$LE_{CN}(\Gamma_G) = 2 \times L_1' + \frac{1}{2}(nz - z - 2) \times L_2' + (z - 1) \times L_3'$$

$$= \begin{cases} 0, & \text{for } n = 3 \& z = 1 \\ 4(z - 1)(z - 2), & \text{for } n = 3 \& z \ge 2 \\ \frac{((n - 1)z - 2)((n - 3)(n + 1)z^2 + ((n - 6)n + 17)z - 4(n + 1))}{2(n + 1)}, & \text{otherwise.} \end{cases}$$

(b)(ii) If n is odd, then by Theorem 2.1

CNSL-spec
$$(\Gamma_G) = \left\{ \left(2\left(\frac{(n-1)z}{2} - 1\right) \left(\frac{(n-1)z}{2} - 2\right) \right)^1, \\ \left(\left(\frac{(n-1)z}{2} - 2\right)^2 \right)^{\frac{(n-1)z}{2} - 1}, (2(z-1)(z-2))^1, ((z-2)^2)^{z-1} \right\}.$$

Here $|V(\Gamma_G)| = \frac{1}{2}(n+1)z$ and $\operatorname{tr}(\operatorname{CNRS}(\Gamma_G) = \frac{1}{8}z(nz+z-4)(((n-4)n+7)z-2(n+1))$. So, $\Delta(\Gamma_G) = \frac{(nz+z-4)(((n-4)n+7)z-2(n+1))}{4(n+1)}$. We have

$$B'_{1} := \left| \frac{1}{2} (nz - z - 2)(nz - z - 4) - \Delta(\Gamma_{G}) \right|$$
$$= \left| \frac{n^{3}z^{2} + n^{2}z^{2} - 6n^{2}z - 5nz^{2} - 12nz + 8n - 5z^{2} + 42z + 8}{4(n+1)} \right|$$

Let $\beta'_1(n, z) = n^3 z^2 + n^2 z^2 - 6n^2 z - 5nz^2 - 12nz + 8n - 5z^2 + 42z + 8$. Then $\beta'_1(n, z) = 8 + 8n + 42z + z^2(n(n-5)-5) + nz(n(nz-6)-12)$. For $n \ge 9$ we have $nz - 6 \ge 3$ which gives n(nz-6) - 12 > 0 and n(n-5) - 5 > 0. Thus, $\beta'_1(n, z) > 0$. Again $\beta'_1(3, z) = 16z^2 - 48z + 32 \ge 0$, $\beta'_1(5, z) = 120z^2 - 168z + 48 \ge 0$ and $\beta'_1(7, z) = 352z^2 - 336z + 64 \ge 0$, as $z \ge 1$. Therefore

$$B_1' = \frac{n^3 z^2 + n^2 z^2 - 6n^2 z - 5n z^2 - 12n z + 8n - 5z^2 + 42z + 8}{4(n+1)}$$

We have

$$B'_{2} := \left| \frac{1}{4} (nz - z - 4)^{2} - \Delta(\Gamma_{G}) \right| = \left| \frac{n^{2}z^{2} - n^{2}z - 2nz^{2} - 6nz + 4n - 3z^{2} + 19z + 4}{2(n+1)} \right|.$$

Let $\beta_2'(n,z) = n^2 z^2 - n^2 z - 2n z^2 - 6n z + 4n - 3z^2 + 19z + 4$. Then $\beta_2'(n,z) = 4 + 4n + 19z + \frac{nz}{4}(nz-24) + \frac{n^2 z}{4}(z-4) + \frac{z^2}{4}(n^2-12) + \frac{nz^2}{4}(n-8) > 0$ for $n \ge 9$ and $z \ge 5$. It can be seen that

 $\beta'_2(3,z) = 16 - 8z = 8 \text{ or } \le 0 \text{ according as } z = 1 \text{ or } z \ge 2; \ \beta'_2(5,z) = 12z^2 - 36z + 24 \ge 0; \ \beta'_2(7,z) = 32z^2 - 72z + 32 = -8 \text{ or } \ge 0 \text{ according as } z = 1 \text{ or } z \ge 2; \ \beta'_2(n,1) = 20 - 4n \ge 0 \text{ or } < 0 \text{ according as } n = 1,3,5 \text{ or } n \ge 7; \ \beta'_2(n,2) = 2n(n-8) + 30 \ge 0 \text{ for all } n \text{ and } \beta'_2(n,3) = 6n^2 - 32n + 34 = -8 \text{ or } \ge 0 \text{ according as } n = 3 \text{ or } n \ne 3. \text{ Therefore}$

$$B_2' = \begin{cases} z - 2, & \text{for } n = 3 \& z \ge 2\\ \frac{n^2 z^2 - n^2 z - 2n z^2 - 6n z + 4n - 3z^2 + 19z + 4}{2(n+1)}, & \text{otherwise.} \end{cases}$$

We have

$$B'_3 := |2(z-1)(z-2) - \Delta(\Gamma_G)| = \left| \frac{-n^3 z^2 + 3n^2 z^2 + 6n^2 z + 5n z^2 - 36n z + 8n + z^2 + 6z + 8}{4(n+1)} \right|$$

Let $\beta'_3(n,z) = -n^3 z^2 + 3n^2 z^2 + 6n^2 z + 5nz^2 - 36nz + 8n + z^2 + 6z + 8$. Then $\beta'_3(n,z) = -8nz + 8 - 8nz + 8n - 20nz + 6z - \frac{n^2 z}{4}(nz - 24) - \frac{z^2}{4}(n^3 - 4) - \frac{nz^2}{4}(n^2 - 5) - \frac{n^2 z^2}{4}(n - 12) < 0$ for $n \ge 25$. Again, $\beta'_3(3,z) = 16z(z-3) + 32 \ge 0$, $\beta'_3(5,z) = -24z^2 - 24z + 48 \le 0$, $\beta'_3(7,z) = -160z^2 + 48z + 64 \le 0$, $\beta'_3(9,z) = -440z^2 + 168z + 80 \le 0$, $\beta'_3(11,z) = -912z^2 + 336z + 96 \le 0$, $\beta'_3(13,z) = -1624z^2 + 552z + 112 \le 0$, $\beta'_3(15,z) = -2624z^2 + 816z + 128 \le 0$, $\beta'_3(17,z) = -3960z^2 + 1128z + 144 \le 0$, $\beta'_3(19,z) = -5680z^2 + 1488z + 160 \le 0$, $\beta'_3(21,z) = -7832z^2 + 1896z + 176 \le 0$ and $\beta'_3(23,z) = -10464z^2 + 2352z + 192 \le 0$. Therefore,

$$B'_{3} = \begin{cases} z(z-3)+2, & \text{for } n = 3 \& z \ge 1 \\ -\frac{n^{3}(-z^{2})+3n^{2}z^{2}+6n^{2}z+5nz^{2}-36nz+8n+z^{2}+6z+8}{4(n+1)}, & \text{for } n \ge 5 \& z \ge 1. \end{cases}$$

We have

$$B'_4 := \left| (z-2)^2 - \Delta(\Gamma_G) \right| = \left| \frac{-n^3 z^2 + 3n^2 z^2 + 6n^2 z + nz^2 - 28nz + 8n - 3z^2 + 14z + 8}{4(n+1)} \right|.$$

Let $\beta'_4(n,z) = -n^3 z^2 + 3n^2 z^2 + 6n^2 z + nz^2 - 28nz + 8n - 3z^2 + 14z + 8$. Then $\beta'_4(n,z) = -8n(z-1) - 14z(n-1) + (8 - 6nz - 3z^2) - \frac{n^2 z}{3}(nz - 18) - \frac{nz^2}{3}(n^2 - 3) - \frac{n^2 z^2}{3}(n-9) < 0$ for $n \ge 19$. It can be seen that $\beta'_4(3,z) = 32 - 16z = 16$ or ≤ 0 according as z = 1 or $z \ge 2$; $\beta'_4(5,z) = 24z(1-2z) + 48 = 24$ or ≤ 0 according as z = 1 or $z \ge 2$; $\beta'_4(5,z) = 24z(1-2z) + 48 = 24$ or ≤ 0 according as z = 1 or $z \ge 2$; $\beta'_4(7,z) = -192z^2 + 112z + 64 \le 0$; $\beta'_4(9,z) = -480z^2 + 248z + 80 \le 0$; $\beta'_4(11,z) = -960z^2 + 432z + 96 \le 0$; $\beta'_4(13,z) = -1680z^2 + 664z + 112 \le 0$; $\beta'_4(15,z) = -2688z^2 + 944z + 128 \le 0$ and $\beta'_4(17,z) = -4032z^2 + 1272z + 144 \le 0$, as $z \ge 1$. Therefore,

$$B'_{4} = \begin{cases} \frac{-n^{3}z^{2} + 3n^{2}z^{2} + 6n^{2}z + nz^{2} - 28nz + 8n - 3z^{2} + 14z + 8}{4(n+1)}, & \text{for } n = 3, 5 \& z = 1\\ -\frac{-n^{3}z^{2} + 3n^{2}z^{2} + 6n^{2}z + nz^{2} - 28nz + 8n - 3z^{2} + 14z + 8}{4(n+1)}, & \text{otherwise.} \end{cases}$$

Hence, by (2), we get

$$LE_{CN}^{+}(\Gamma_{G}) = 1 \times B_{1}' + \frac{1}{2}(nz - z - 2) \times B_{2}' + 1 \times B_{3}' + (z - 1) \times B_{4}'$$

$$= \begin{cases} 0, & \text{for } n = 3 \& z = 1; \ n = 5 \& z = 1 \\ 4(z - 1)(z - 2), & \text{for } n = 3 \& z \ge 2 \\ \frac{(n - 5)(n - 3)(n + 3)}{2(n + 1)}, & \text{for } n \ge 7 \& z = 1 \\ \frac{(n - 3)(n - 1)z^{2}(nz + z - 6)}{2(n + 1)}, & \text{otherwise.} \end{cases}$$

Hence the result follows.

As a corollary of the above Theorem 2.4, we get the following results.

Corollary 2.5. The CNL-spectrum, CNSL-spectrum, CNL-energy and CNSL-energy of CCC-graph of the dihedral group D_{2n} (where $n \ge 3$) are as given below: (a) If n is odd then (i) CNL-spec $(\Gamma_{D_{2n}}) = \{(0)^2, (\frac{1}{4}(n-1)(n-5))^{\frac{1}{2}(n-3)}\}$ and $LE_{CN}(\Gamma_{D_{2n}}) = \frac{(n-5)(n-3)(n-1)}{n+1}.$ (ii) CNSL-spec $(\Gamma_{D_{2n}}) = \{(0)^1, (\frac{1}{2}(n-3)(n-5))^1, (\frac{1}{4}(n-5)^2)^{\frac{1}{2}(n-3)}\}$ and $LE_{CN}^+(\Gamma_{D_{2n}}) = \frac{(n-5)(n-3)(n+3)}{2(n+1)}.$

(b) If n is even then

(i) CNL-spec
$$(\Gamma_{D_{2n}}) = \{(0)^3, (\frac{1}{4}(n-2)(n-6))^{\frac{1}{2}(n-4)}\}$$
 and

$$LE_{CN}(\Gamma_{D_{2n}}) = \frac{3(n-6)(n-4)(n-2)}{2(n+2)}.$$

(ii) CNSL-spec $(\Gamma_{D_{2n}}) = \{(0)^2, (\frac{1}{2}(n-4)(n-6))^1, (\frac{1}{4}(n-6)^2)^{\frac{1}{4}(n-4)}\} and$ $LE_{CN}^+(\Gamma_{D_{2n}}) = \begin{cases} \frac{28}{5}, & \text{for } n = 8\\ \frac{(n-6)(n-4)(n-2)}{n+2}, & \text{for } n \neq 8. \end{cases}$

Proof. We know that $\frac{D_{2n}}{Z(D_{2n})} \cong D_{2 \times \frac{n}{2}}$ or D_{2n} according as n is even or odd. Therefore, by Theorem 2.4, we get the required result.

Corollary 2.6. The CNL-spectrum, CNSL-spectrum, CNL-energy and CNSL-energy of CCC-graph of the dicyclic group T_{4n} (where $n \ge 2$) are as given below:

(a) CNL-spec $(\Gamma_{T_{4n}}) = \{(0)^3, ((n-1)(n-3))^{n-2}\}$ and

$$LE_{CN}(\Gamma_{T_{4n}}) = \frac{6(n-3)(n-2)(n-1)}{n+1}.$$

(b) CNSL-spec
$$(\Gamma_{T_{4n}}) = \{(0)^2, (2(n-2)(n-3))^1, ((n-3)^2)^{n-2}\}$$
 and
 $LE_{CN}^+(\Gamma_{T_{4n}}) = \begin{cases} \frac{28}{5}, & \text{for } n = 4\\ \frac{4(n-3)(n-2)(n-1)}{n+1}, & \text{for } n \neq 4. \end{cases}$

Proof. We know that $\frac{T_{4n}}{Z(T_{4n})} \cong D_{2n}$. Therefore, by Theorem 2.4, we get the required result.

Corollary 2.7. The CNL-spectrum, CNSL-spectrum, CNL-energy and CNSL-energy of CCC-graph of the group $U_{6n} = \langle x, y : x^{2n} = y^3 = 1, x^{-1}yx = y^{-1} \rangle$ (where $n \ge 2$) are as given below:

(a) CNL-spec
$$(\Gamma_{U_{6n}}) = \{(0)^2, (n(n-2))^{2(n-1)}\}\ and\ LE_{CN}(\Gamma_{U_{6n}}) = 4(n-2)(n-1).$$

(b) CNSL-spec
$$(\Gamma_{U_{6n}}) = \{(2(n-1)(n-2))^2, ((n-2)^2)^{2(n-1)}\}\ and$$

 $LE_{CN}^+(\Gamma_{U_{6n}}) = 4(n-2)(n-1).$

Proof. We know that $\frac{U_{6n}}{Z(U_{6n})} = D_{2\times 3}$. Therefore, by Theorem 2.4, we get the required result.

Corollary 2.8. The CNL-spectrum, CNSL-spectrum, CNL-energy and CNSL-energy of CCC-graph of the group $U_{(n,m)}$ (where $m \ge 2$ and $n \ge 2$) are as given below: (a) If m is even then

(i) CNL-spec
$$(\Gamma_{U_{(n,m)}}) =$$

$$\left\{ (0)^3, (n(n-2))^{2(n-1)}, \left(\frac{1}{4}(nm-2n)(nm-2n-4)\right)^{\frac{1}{2}(nm-2n-2)} \right\}$$

and

$$LE_{CN}(\Gamma_{U_{(n,m)}}) = \begin{cases} 4(n-1)(2n-1), & \text{for } m = 2 \& n \ge 2\\ 6(n-2)(n-1), & \text{for } m = 4 \& n \ge 2\\ 8(n-1)(2n-1), & \text{for } m = 6 \& n \ge 2\\ \frac{((m-2)n-2)(m(2n+1)((m-4)n-4)+44n-8)}{2(m+2)}, & \text{otherwise.} \end{cases}$$

(ii) CNSL-spec
$$(\Gamma_{U_{(n,m)}}) = \left\{ (2(n-1)(n-2))^2, ((n-2)^2)^{2(n-1)}, \\ \left(\frac{1}{2}(nm-2n-2)(nm-2n-4)\right)^1, \left(\frac{1}{4}(nm-2n-4)^2\right)^{\frac{1}{2}(nm-2n-2)} \right\}$$

and

$$LE_{CN}^{+}(\Gamma_{U_{(n,m)}}) = \begin{cases} 4(n-1)(2n-1), & \text{for } m = 2 \& n \ge 2\\ 6(n-2)(n-1), & \text{for } m = 4 \& n \ge 2\\ 8n(2n-3) + 8, & \text{for } m = 6 \& n \ge 2\\ \frac{24}{5}n^2(4n-3), & \text{for } m = 8 \& n \ge 2\\ \frac{(m-4)(m-2)n^2(mn-6)}{m+2}, & \text{otherwise.} \end{cases}$$

(b) If m is odd then

(i) CNL-spec
$$(\Gamma_{U_{(n,m)}}) = \left\{ (0)^2, (n(n-2))^{n-1}, \left(\frac{1}{4}(nm-n)(nm-n-4)\right)^{\frac{1}{2}(nm-n-2)} \right\}$$

and

$$LE_{CN}(\Gamma_{U_{(n,m)}}) = \begin{cases} 4(n-1)(n-2), & \text{for } m = 3 \& n \ge 2\\ \frac{((m-1)n-2)\left((m-3)(m+1)n^2 + ((m-6)m+17)n - 4(m+1)\right)}{2(m+1)}, & \text{otherwise.} \end{cases}$$
(ii) CNSL-spec $(\Gamma_{U_{(n,m)}}) = \{(2(n-1)(n-2))^1, ((n-2)^2)^{n-1}, \\ \end{cases}$

$$\left(\frac{1}{2}(nm-n-2)(nm-n-4)\right)^{1}, \left(\frac{1}{4}(nm-n-4)^{2}\right)^{\frac{1}{2}(nm-n-2)}\right\}$$

and

$$LE_{CN}^{+}(\Gamma_{U_{(n,m)}}) = \begin{cases} 4(n-1)(n-2), & \text{for } m = 3 \& n \ge 2\\ \frac{(m-3)(m-1)n^2(mn+n-6)}{2(m+1)}, & \text{otherwise.} \end{cases}$$

Proof. We know that $\frac{U_{(n,m)}}{Z(U_{(n,m)})}$ is isomorphic to $D_{2\times\frac{m}{2}}$ or D_{2m} according as m is even or odd. Therefore, by Theorem 2.4, we get the required result.

Corollary 2.9. The CNL-spectrum, CNSL-spectrum, CNL-energy and CNSL-energy of CCC-graph of the group SD_{8n} (where $n \ge 2$) are as given below:

(a) If n is even then
(b) CNL-spec(
$$\Gamma_{SD_{8n}}$$
) = {(0)³, ((2n - 1)(2n - 3))²ⁿ⁻²} and
 $LE_{CN}(\Gamma_{SD_{8n}}) = \frac{12(n - 1)(4(n - 2)n + 3)}{2n + 1}$.
(ii) CNSL-spec($\Gamma_{SD_{8n}}$) = {(0)², (2(2n - 2)(2n - 3))¹, ((2n - 3)²)²ⁿ⁻²} and
 $LE_{CN}^{+}(\Gamma_{SD_{8n}}) = \begin{cases} \frac{28}{5}, & \text{for } n = 2\\ \frac{8(n - 1)(2n - 3)(2n - 1)}{2n + 1}, & \text{for } n \ge 4. \end{cases}$

(b) If n is odd then
(i) CNL-spec(
$$\Gamma_{SD_{8n}}$$
) = {(0)², (8)³, ((2n - 2)(2n - 4))²ⁿ⁻³} and
 $LE_{CN}(\Gamma_{SD_{8n}}) = \begin{cases} 24, & \text{for } n = 3\\ \frac{4(2n-3)(5(n-3)n+4)}{n+1}, & \text{for } n \ge 5. \end{cases}$
(ii) CNSL-spec($\Gamma_{SD_{8n}}$) = {(2(2n - 3)(2n - 4))¹, ((2n - 4)²)²ⁿ⁻³, 12¹, 4³} and
 $LE_{CN}^{+}(\Gamma_{SD_{8n}}) = \begin{cases} 24, & \text{for } n = 3\\ \frac{16(n-3)(n-1)(2n-1)}{n+1}, & \text{for } n \ge 5. \end{cases}$

Proof. We know that $\frac{SD_{8n}}{Z(SD_{8n})}$ is isomorphic to $D_{2\times 2n}$ or D_{2n} according as n is even or odd. Therefore, by Theorem 2.4, we get the required result.

Corollary 2.10. The CNL-spectrum, CNSL-spectrum, CNL-energy and CNSL-energy of CCC-graph of the group V_{8n} (where $n \geq 2$) are as given below:

- (a) If n is even then
- (i) CNL-spec($\Gamma_{V_{8n}}$) = {(0)⁵, ((2n 2)(2n 4))²ⁿ⁻³} and $LE_{CN}(\Gamma_{V_{8n}}) = \frac{20(n - 2)(n - 1)(2n - 3)}{n + 1}.$ (ii) CNCL and ($\Gamma_{V_{8n}}$) = {(0)⁴/₂/₂(2n - 4))¹/₂((2n - 4))²/₂(2n - 4))²/₂(2n - 4))²/₂(2n - 4))²/₂(2n - 4)²/₂(2n - 4))²/₂(2n - 4))²/₂(2n - 4)²/₂(2n - 4))²/₂(2n - 4)²/₂(2n - 4))²/₂(2n - 4)²/₂(2n - 4))²/₂(2n - 4)²/₂(2n - 4)²/₂

(ii) CNSL-spec
$$(\Gamma_{V_{8n}}) = \{(0)^4, (2(2n-3)(2n-4))^1, ((2n-4)^2)^{2n-3}\}$$
 and
 $LE_{CN}^+(\Gamma_{V_{8n}}) = \frac{16(n-2)(n-1)(2n-3)}{n+1}.$

(b) If n is odd then

(i) CNL-spec
$$(\Gamma_{V_{8n}}) = \{(0)^3, ((2n-1)(2n-3))^{2n-2}\}$$
 and
 $LE_{CN}(\Gamma_{V_{8n}}) = \frac{12(n-1)(4(n-2)n+3)}{2n+1}.$

(ii) CNSL-spec $(\Gamma_{V_{8n}}) = \{(0)^2, (2(2n-2)(2n-3))^1, ((2n-3)^2)^{2n-2}\}$ and $LE_{CN}^+(\Gamma_{V_{8n}}) = \frac{8(n-1)(2n-3)(2n-1)}{2n+1}.$

Proof. (a) If n is even then, by [33, Proposition 2.4], we have $\Gamma_{V_{8n}} = K_{2n-2} \cup 2K_2$. (i) By Theorem 2.1, we get

CNL-spec(
$$\Gamma_{V_{8n}}$$
) = {(0)⁵, ((2n-2)(2n-4))²ⁿ⁻³}.

Here $|V(\Gamma_{V_{8n}})| = 2(n+1)$ and $\operatorname{tr}(\operatorname{CNRS}(\Gamma_{V_{8n}})) = 4(n-2)(n-1)(2n-3)$. So, $\Delta(\Gamma_{V_{8n}}) = \frac{2(n-2)(n-1)(2n-3)}{n+1}$. We have

$$L_1 := |0 - \Delta(\Gamma_{V_{8n}})| = \left| -\frac{2(n-2)(n-1)(2n-3)}{n+1} \right| = \frac{2(n-2)(n-1)(2n-3)}{n+1},$$

since -2(n-2)(n-1)(2n-3) < 0, as $n \ge 2$, so 2n-3 > 0, $n-2 \ge 0$ and n-1 > 0. Also

$$L_2 := |(2n-2)(2n-4) - \Delta(\Gamma_{V_{8n}})| = \left|\frac{10(n-2)(n-1)}{n+1}\right|$$
$$= \frac{10(n-2)(n-1)}{n+1}, \text{ as } n \ge 2$$

Therefore, by (1), we get

$$LE_{CN}(\Gamma_{V_{8n}}) = 5 \times L_1 + (2n-3) \times L_2 = \frac{20(n-2)(n-1)(2n-3)}{n+1}.$$

(ii) By Theorem 2.1, we get

CNSL-spec(
$$\Gamma_{V_{8n}}$$
) = {(0)⁴, (2(2n-3)(2n-4))¹, ((2n-4)²)²ⁿ⁻³}

1.1

We have

$$B_1 := |0 - \Delta(\Gamma_{V_{8n}})| = L_1,$$

$$B_2 := |2(2n-3)(2n-4) - \Delta(\Gamma_{V_{8n}})| = \left|\frac{2(n-2)(n+3)(2n-3)}{n+1}\right|$$

$$= \frac{2(n-2)(n+3)(2n-3)}{n+1}, \text{ as } n \ge 2,$$

and

$$B_3 := \left| (2n-4)^2 - \Delta(\Gamma_{V_{8n}}) \right| = \left| \frac{2(n-2)(3n-7)}{n+1} \right| = \frac{2(n-2)(3n-7)}{n+1}$$

as $n \ge 2$, so $2(n-2)(3n-7) \ge 0$. Therefore, by (2), we get

$$LE_{CN}^{+}(\Gamma_{V_{8n}}) = 4 \times B_1 + 1 \times B_2 + (2n-3) \times B_3 = \frac{16(n-2)(n-1)(2n-3)}{n+1}.$$

(b) If n is odd then, [33, Proposition 2.4], we have $\Gamma_{V_{8n}} = K_{2n-1} \cup 2K_1 = \Gamma_{D_{2 \times 4n}}$. Hence, the result follows from Corollary 2.5.

3. Some consequences

In this section, we discuss some consequences of the results obtained in Section 2. Looking at the CNL-spectrum and CNSL-spectrum of CCC-graphs of the groups considered in Section 2, we get the following result.

Theorem 3.1. Let G be a finite non-abelian group with center Z(G). Then the CCCgraph of G is (CNSL) CNL-integral if

(a)
$$\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$$
.

(b) $\frac{G}{Z(G)} \cong D_{2n}$. (c) G is isomorphic to D_{2n} , T_{4n} , U_{6n} , $U_{(n,m)}$, SD_{8n} and V_{8n} .

Now we shall determine whether CCC-graphs of these groups are (CNSL) CNL-hyperenergetic.

Theorem 3.2. Let G be a finite non-abelian group and $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$. Then the CCC-graph of G is not (CNSL) CNL-hyperenergetic.

Proof. Let
$$|Z(G)| = z$$
. Then $z \ge 2$ and $|V(\Gamma_G)| = \frac{(p^2 - 1)z}{p}$. By Theorem 2.2 and (3)

$$LE_{CN}(K_{|V(\Gamma_G)|}) - LE_{CN}(\Gamma_G) = \begin{cases} \frac{4((p-2)p(2p^2+p-2)+4)}{p^2}, & \text{for } p \ge 2 \& z = 2\\ 16, & \text{for } p = 2 \& z = 3\\ \frac{2p^3z^2 - 2p^2z^2 - 4p^2 - 2pz^2 + 2z^2}{p}, & \text{otherwise.} \end{cases}$$

Let $f_1(p) = 4 ((p-2)p (2p^2 + p - 2) + 4)$ and $f_2(p, z) = 2p^3 z^2 - 2p^2 z^2 - 4p^2 - 2pz^2 + 2z^2$, where $z \ge 3$. Then $f_1(p) > 0$. Also, $f_2(p, z) = \frac{2}{3}(p-3)p^2 z^2 + \frac{2}{3}p^2 (pz^2 - 6) + \frac{2}{3}(p^2 - 3)pz^2 + 2z^2 > 0$ for $p \ge 3$. For p = 2 we have $f_2(p, z) = 6z^2 - 16 > 0$. Hence, $LE_{CN}(K_{|V(\Gamma_G)|}) - LE_{CN}(\Gamma_G) > 0$. By Theorem 2.2, we also have $LE_{CN}(\Gamma_G) = LE_{CN}^+(\Gamma_G)$. Therefore, $LE_{CN}^+(\Gamma_G) - LE_{CN}^+(K_{|V(\Gamma_G)|}) > 0$. Hence the result follows.

An immediate corollary of the above theorem is given below.

Corollary 3.3. Let G be a non-abelian group of order p^n and center $|Z(G)| = p^{n-2}$. Then CCC-graph of G is not (CNSL) CNL-hyperenergetic.

Theorem 3.4. Let G be a finite non-abelian group and $\frac{G}{Z(G)} \cong D_{2n}$ (where $n \ge 3$). Then the CCC-graph of G is

- (a) CNL-borderenergetic if n = 3, 11 & z = 1.
- (b) CNL-hyperenergetic except for $n = 4, 6 \& z = 2; n = 4 \& z = 3; n = 4 \& z = 4; n = 3 \& z \ge 2; n = 5, 7, 9 \& z = 1 and n = 5 \& z = 2, 3.$
- (c) CNSL-borderenergetic for n = 3 & z = 1.
- (d) CNSL-hyperenergetic except for n = 4 & z = 2, 3, 4, 5; n = 6, 8 & z = 2; n = 6 & z = 3; n = 5 & z = 1; n = 3 & $z \ge 2; n \ge 7$ (n is odd) & z = 1; n = 5, 7, 9 & z = 2 and n = 5 & z = 3, 4.

Proof. We have

$$V(\Gamma_G)| = \begin{cases} \frac{1}{2}(n+1)z, & \text{for } n \text{ is even} \\ \frac{1}{2}(n+1)z, & \text{for } n \text{ is odd.} \end{cases}$$

Case 1: n is even

In this case $z \ge 2$. By Theorem 2.9 and (3), we have

$$LE_{CN}(K_{|V(\Gamma_G)|}) - LE_{CN}(\Gamma_G) = \frac{-n^3 z^3 + 3n^2 z^3 + 12n^2 z^2 - 2nz^3 - 18nz^2 - 24nz + 12z^2 + 12z}{2(n+1)}$$

Let $f_1(n,z) = -n^3 z^3 + 3n^2 z^3 + 12n^2 z^2 - 2nz^3 - 18nz^2 - 24nz + 12z^2 + 12z$. Then $f_1(n,z) = \frac{1}{2}n^2 z^3(6-n) + \frac{1}{2}n^2 z^2(24-nz) - 2nz^3 + 6z^2(2-3n) + 12z(1-2n) < 0$ for $n \ge 6$ and $z \ge 4$. We have $f_1(n,2) = 8n^2(9-n) + 72 - 136n < 0$ for $n \ge 10$. Also $f_1(4,2) = 168, f_1(6,2) = 120$ and $f_1(8,2) = -504$. Therefore, $f_1(n,2) > 0$ or < 0according as n = 4, 6 or $n \ge 8$. We have $f_1(n, 3) = 27n^2(7 - n) + 144 - 288n < 0$ for $n \geq 8.$

Also $f_1(4,3) = 288$ and $f_1(6,3) = -612$. Therefore, $f_1(4,3) > 0$ or < 0 according as n = 4 or $n \ge 6$.

Now we need to check for n = 4 and $z \ge 4$. We have $f_1(4, z) = 12z^2(11-2z)-84z < 12z^2(11-2z)-82z^2(11$ 0 for $z \ge 6$. Also $f_1(4,4) = 240$ and $f_1(4,5) = -120$. Therefore, $f_1(4,z) > 0$ or < 0 according as z = 4 or $z \geq 5$. Hence, $LE_{CN}(K_{|V(\Gamma_G)|}) - LE_{CN}(\Gamma_G) > 0$ for n = 4, 6 & z = 2 and n = 4 & z = 3, 4. Otherwise, $LE_{CN}(K_{|V(\Gamma_G)|}) - LE_{CN}(\Gamma_G) < 0.$

By Theorem 2.9 and (3), we also have

$$\begin{split} LE_{CN}^+(K_{|V(\Gamma_G)|}) - LE_{CN}^+(\Gamma_G) \\ &= \begin{cases} \frac{92}{5}, & \text{for } n = 4 \& z = 2\\ \frac{1}{10}z \left(-24z^2 + 161z - 150\right) + 4, & \text{for } n = 4 \& z \geq 3\\ \frac{-n^3z^3 + n^3z^2 + 3n^2z^3 + 9n^2z^2 - 6n^2z - 2nz^3 - 15nz^2 - 12nz + 8n + 13z^2 - 6z + 8}{2(n+1)}, & \text{otherwise.} \end{cases} \end{split}$$

Let $f_2(z) = \frac{1}{10}z(-24z^2 + 161z - 150) + 4$ and $f_3(n, z) = -n^3z^3 + n^3z^2 + 3n^2z^3 + 9n^2z^2 - 16n^2z^2 + 161z - 150) + 4$ $6n^2z - 2nz^3 - 15nz^2 - 12nz + 8n + 13z^2 - 6z + 8$. Then $f_2(z) = \frac{1}{10}z(z(161 - 24z) - 150) + 6n^2z - 2nz^3 - 15nz^2 - 12nz + 8n + 13z^2 - 6z + 8$. 4 > 0 or < 0 according as z = 3, 4, 5 or $z \ge 6$. Also, $f_3(n, z) = \frac{1}{3}n^3(3-z)z^2 + \frac{1}{3}(9-z)z^2 + \frac{1}{$ $n)n^{2}z^{3} + \frac{1}{3}n^{2}z^{2}(27 - nz) - 6n^{2}z + z^{2}(13 - 2nz) + (8n - 15nz^{2}) + (8 - 12nz) - 6z < 0$ for $n \ge 10$ and $z \ge 3$.

We have $f_3(n,2) = 4n^2(12-n) + 48 - 92n < 0$ for $n \ge 12$. Also, $f_3(6,2) = 360$, $f_3(8,2) = 336$ and $f_3(10,2) = -72$. Therefore, $f_3(n,2) > 0$ or < 0 according as n = 6, 8 or n > 10.

We have $f_3(6, z) = z^2(463 - 120z) - 294z + 56 < 0$ for $z \ge 4$. Also, $f_3(6, 3) = 101$. Therefore, $f_3(6,z) > 0$ or < 0 according as z = 3 or $z \ge 4$. We have $f_3(8,z) =$ $z^{2}(981 - 336z) - 486z + 72 < 0$ for $z \ge 3$. Therefore, $f_{3}(n, z) > 0$ if n = 6, 8 & z = 2and n = 6 & z = 3. Otherwise, $f_3(n, z) < 0$. Hence, $LE_{CN}^+(K_{|V(\Gamma_G)|}) - LE_{CN}^+(\Gamma_G) > 0$ if n = 4 & z = 2, 3, 4, 5; n = 6, 8 & z = 2 and n = 6 & z = 3. Otherwise, $LE_{CN}^+(K_{|V(\Gamma_G)|}) - LE_{CN}^+(\Gamma_G) < 0.$

Case 2: n is odd

By Theorem 2.9 and (3), we have

$$LE_{CN}(K_{|V(\Gamma_G)|}) - LE_{CN}(\Gamma_G)$$

$$= \begin{cases} 0, & \text{for } n = 3 \& z = 1 \\ 4z^2 - 4, & \text{for } n = 3 \& z \ge 2 \\ \frac{24z - 24nz + 12z^2 - 24nz^2 + 12n^2z^2 - 3z^3 + nz^3 + 3n^2z^3 - n^3z^3}{2(n+1)}, & \text{otherwise.} \end{cases}$$

Clearly $4z^2 - 4 > 0$ for $z \ge 2$. Let $f_4(n, z) = 24z - 24nz + 12z^2 - 24nz^2 + 12n^2z^2 - 3z^3 + nz^3 + 3n^2z^3 - n^3z^3$. Then $f_4(n, z) = \frac{1}{3}n^2z^3(9-n) + \frac{1}{3}nz^3(3-n^2) + \frac{1}{3}n^2z^2(36-nz) - 3z^3 + 12z^2(1-2n) + 24z(1-n) < 0$ for $n \ge 9$ and $z \ge 4$. We have $f_4(n, 1) = n^2(15-n) + 33 - 47n < 0$ for $n \ge 15$. Also, $f_4(5, 1) = 48$, $f_4(7, 1) = 96$, $f_4(9, 1) = 96$, $f_4(11, 1) = 0$ and $f_4(13, 1) = -240$. Therefore

$$f_4(n,1) \begin{cases} = 0, & \text{for } n = 11 \\ > 0, & \text{for } n = 5,7,9 \\ < 0, & \text{for } n \ge 13. \end{cases}$$

We have $f_4(n,2) = 8n^2(9-n) + 72 - 136n < 0$ for $n \ge 9$, $f_4(5,2) = 192$ and $f_4(7,2) = -96$. Therefore, $f_4(n,2) > 0$ or < 0 according as n = 5 or $n \ge 7$.

We have $f_4(n,3) = 27n^2(7-n) + 99 - 261n < 0$ for $n \ge 7$ and $f_4(5,3) = 144$. Therefore, $f_4(n,3) > 0$ or < 0 according as n = 5 or $n \ge 7$.

Again, we have $f_4(5, z) = 48z^2(4-z) - 96z < 0$ and $f_4(7, z) = 48z^2(9-4z) - 144z < 0$ for $z \ge 4$. Therefore

$$f_4(n,z) \begin{cases} = 0, & \text{for } n = 11 \& z = 1 \\ > 0, & \text{for } n = 5, 7, 9 \& z = 1; n = 5 \& z = 3 \\ < 0, & \text{otherwise.} \end{cases}$$

Hence

$$LE_{CN}(K_{|V(\Gamma_G)|}) - LE_{CN}(\Gamma_G)$$

$$\begin{cases} = 0, & \text{for } n = 3, 11 \& z = 1 \\ > 0, & \text{for } n = 3 \& z \ge 2; \\ & n = 5, 7, 9 \& z = 1; \ n = 5 \& z = 2, 3 \\ < 0, & \text{otherwise.} \end{cases}$$

By Theorem 2.9 and (3), we also have

$$\begin{split} LE_{CN}^+(K_{|V(\Gamma_G)|}) - LE_{CN}^+(\Gamma_G) \\ &= \begin{cases} 0, & \text{for } n = 3 \,\&\, z = 1 \\ 4, & \text{for } n = 5 \,\&\, z = 1 \\ 4(z^2 - 1), & \text{for } n = 3 \,\&\, z \geq 2 \\ \frac{n^2 + 4n - 21}{n + 1}, & \text{for } n \geq 7 \,\&\, z = 1 \\ \frac{-n^3 z^3 + n^3 z^2 + 3n^2 z^3 + 9n^2 z^2 - 6n^2 z + nz^3 - 21nz^2 - 12nz + 8n - 3z^3 + 19z^2 - 6z + 8}{2(n + 1)}, \text{ otherwise.} \end{split}$$

Clearly $4(z^2 - 1) > 0$ for $z \ge 2$ and $n^2 + 4n - 21 > 0$ for $n \ge 7$. Let $f_5(n, z) = -n^3 z^3 + n^3 z^2 + 3n^2 z^3 + 9n^2 z^2 - 6n^2 z + nz^3 - 21nz^2 - 12nz + 8n - 3z^3 + 19z^2 - 6z + 8$. Then $f_5(n, z) = \frac{1}{4}n^3 z^2(4 - z) + \frac{1}{4}n^2 z^3(12 - n) + \frac{1}{4}n^2 z^2(36 - nz) + \frac{1}{4}nz^3(4 - n^2) - 21nz^2 + (8n - 12nz) - 6n^2 z + z^2(19 - 3z) + (8 - 6z) < 0$ for $n \ge 13$ and $z \ge 7$. We have $f_5(n, 2) = 4n^2(12 - n) + 48 - 92n < 0$ for $n \ge 13$. Again $f_5(5, 2) = 288$, $f_5(7, 2) = 384$, $f_5(9, 2) = 192$ and $f_5(11, 2) = -480$. Therefore, $f_5(n, 2) > 0$ or < 0 according as n = 5, 7, 9 or $n \ge 11$.

We have $f_5(n,3) = 18n^2(8-n) + 80 - 190n < 0$ for $n \ge 9$. Again, $f_5(5,3) = 480$ and $f_5(7,3) = -368$. Therefore, $f_5(n,3) > 0$ or < 0 according as n = 5 or $n \ge 7$.

We have $f_5(n,4) = 24n^2(13-2n) + 96 - 312n < 0$ for $n \ge 7$. Also, $f_5(5,4) = 336$. Therefore, $f_5(n,4) > 0$ or < 0 according as n = 5 or $n \ge 7$.

We have $f_5(n,5) = 10n^2(57-10n) + 78 - 452n < 0$ for $n \ge 7$. Also, $f_5(5,5) = -432$. Therefore, $f_5(n,5) < 0$.

We have $f_5(n, 6) = 36n^2(26 - 5n) + 8 - 604n < 0$ for $n \ge 7$. Also, $f_5(5, 6) = -2112$. Therefore, $f_5(n, 6) < 0$.

Now we shall check for n = 5, 7, 9, 11 and $z \ge 7$. We have $f_5(5, z) = 24z^2(11 - 2z) + 48 - 216z < 0$, $f_5(7, z) = 16z^2(41 - 12z) + 64 - 384z < 0$, $f_5(9, z) = 8z^2(161 - 60z) + 80 - 600z < 0$ and $f_5(11, z) = 96z^2(23 - 10z) + 96 - 864z < 0$, as $z \ge 7$. Thus, $f_5(n, z) > 0$ if n = 5, 7, 9 & z = 2 and n = 5 & z = 3, 4. Otherwise, $f_5(n, z) < 0$. Hence

$$LE_{CN}^{+}(K_{|V(\Gamma_{G})|}) - LE_{CN}^{+}(\Gamma_{G})$$

$$= \begin{cases} = 0, & \text{for } n = 3 \& z = 1 \\ > 0, & \text{for } n = 5 \& z = 1; \ n = 3 \& z \ge 2; \\ n \ge 7 \& z = 1; \ n = 5, 7, 9 \& z = 2; \ n = 5 \& z = 3, 4 \\ < 0, & \text{otherwise.} \end{cases}$$

Hence the result follows.

As a corollary of the above theorem we get the following results.

Corollary 3.5. The CCC-graph of U_{6n} $(n \ge 2)$ is not (CNSL) CNL-hyperenergetic.

Corollary 3.6. Let $G = D_{2n}, T_{4n}, U_{6n}, SD_{8n}$ or $U_{(n,m)}$. Then

- (a) Γ_G is CNL-borderenergetic if and only if $G = D_6$, D_{22} and $U_{(n,2)}$ for $n \ge 2$.
- (b) Γ_G is CNL-hyperenergetic if and only if $G = D_{2n}$ for $n \ge 13$; T_{4n} for $n \ge 7$; SD_{8n} for $n \ge 4$ and $U_{(n,m)}$ except for $m = 3 \& n \ge 2$, m = 5 & n = 2, 3, $m = 4 \& n \ge 2$, m = 8 & n = 2 and $m = 6 \& n \ge 2$.
- (c) Γ_G is CNSL-borderenergetic if and only if $G = D_6$ and $U_{(n,2)}$ for $n \ge 2$.
- (d) Γ_G is CNSL-hyperenergetic if and only if $G = D_{2n}$ for n is even and $n \ge 20$; T_{4n} for $n \ge 10$; SD_{8n} for $n \ge 6$ and $U_{(n,m)}$ except for $m = 3 \& n \ge 2$, $m = 5, 7, 9 \& n = 2, m = 5 \& n = 3, 4, m = 4 \& n \ge 2, m = 8 \& n = 2$, $m = 6 \& n \ge 2$ and m = 10 & n = 2.

Corollary 3.7. The CCC-graph of V_{8n} $(n \ge 2)$ is CNL-hyperenergetic for $n \ge 6$ and CNSL-hyperenergetic for $n \ge 4$.

Proof. Case 1: n is even

We have $|V(\Gamma_{V_{8n}})| = (2n+2)$. By Theorem 2.10 and (3), we get

$$LE_{CN}(K_{|V(\Gamma_{V_{8n}})|}) - LE_{CN}(\Gamma_{V_{8n}}) = \frac{120 - 32(n-4)(n-2)n}{n+1} \begin{cases} > 0, & \text{for } 2 \le n \le 4 \\ < 0, & \text{for } n \ge 6. \end{cases}$$

$$LE_{CN}^{+}(K_{|V(\Gamma_{V_{8n}})|}) - LE_{CN}^{+}(\Gamma_{V_{8n}}) = -\frac{6(n-1)(n(5n-19)+16)}{n+1} \begin{cases} > 0, & \text{for } n=2\\ < 0, & \text{for } n\geq 4. \end{cases}$$

Thus, $\Gamma_{V_{8n}}$ is not CNL-hyperenergetic if n = 2, 4 and $\Gamma_{V_{8n}}$ is CNL-hyperenergetic if $n \ge 6$. Also, it is not CNSL-hyperenergetic if n = 2 and $\Gamma_{V_{8n}}$ is CNSL-hyperenergetic if $n \ge 4$.

Case 2: n is odd

We have $\Gamma_{V_{8n}} = K_{2n-1} \cup 2K_1 = \Gamma_{D_{2\times 4n}}$. Then, by Corollary 3.6, we have that $\Gamma_{V_{8n}}$ is CNL-hyperenergetic if $n \ge 4$ and CNSL-hyperenergetic if $n \ge 5$. Hence, the result follows.

3.1. Comparing various CN-energies. In this subsection, we compare various CN-energies of CCC-graphs of the groups considered in Section 2.

Theorem 3.8. Let G be a finite group such that $|Z(G)| = z \ge 2$ and $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$. If p = 2 & z = 3 or $p \ge 3 \& z = 2$ then

$$E_{CN}(\Gamma_G) < LE_{CN}(\Gamma_G) = LE_{CN}^+(\Gamma_G).$$

For all other cases, $E_{CN}(\Gamma_G) = LE_{CN}(\Gamma_G) = LE_{CN}^+(\Gamma_G)$.

Proof. In view of Theorem 2.2, it is sufficient to compare $E_{CN}(\Gamma_G)$ and $LE_{CN}(\Gamma_G)$. By Theorem 2.2 and [26, Theorem 2.9], we have

$$LE_{CN}(\Gamma_G) - E_{CN}(\Gamma_G) = \begin{cases} 3, & \text{for } p = 2 \& z = 3\\ \frac{8(p-2)(p+1)}{p^2}, & \text{for } p \ge 2 \& z = 2\\ 0, & \text{otherwise.} \end{cases}$$

Clearly, 8(p-2)(p+1) = 0 or > 0 according as p = 2 or p > 2. Hence, the result follows.

As a corollary to Theorem 3.8 we have the following result.

Corollary 3.9. Let G be a non-abelian p-group of order p^n and $|Z(G)| = p^{n-2}$, where p is a prime and $n \ge 3$. Then $E_{CN}(\Gamma_G) = LE_{CN}(\Gamma_G) = LE_{CN}^+(\Gamma_G)$.

Theorem 3.10. Let G be a finite group and $\frac{G}{Z(G)} \cong D_{2n}$ $(n \ge 3)$. If $n = 3 \& z \ge 1$ or n = 5 & z = 1 (where |Z(G)| = z) then

$$E_{CN}(\Gamma_G) = LE_{CN}(\Gamma_G) = LE_{CN}^+(\Gamma_G).$$

For all other cases, $E_{CN}(\Gamma_G) < LE_{CN}^+(\Gamma_G) < LE_{CN}(\Gamma_G)$.

Proof. Case 1: n is even

In this case $z \ge 2$. By Theorem 2.4 and [26, Theorem 2.14], we have

$$LE_{CN}^{+}(\Gamma_{G}) - E_{CN}(\Gamma_{G})$$

$$= \begin{cases} \frac{8}{5}, & \text{for } n = 4 \& z = 2\\ \frac{z^{2}(24z-91)+150z-120}{10}, & \text{for } n = 4 \& z \ge 3\\ \frac{(n-2)(n-1)nz^{3}-(n(n(n+5)-17)+15)z^{2}+6(n+1)^{2}z-24(n+1))}{2(n+1)}, & \text{otherwise.} \end{cases}$$

Let $f_1(z) = z^2(24z - 91) + 150z - 120$ and $f_2(n, z) = (n-2)(n-1)nz^3 - (n(n(n+5)-17)+15)z^2 + 6(n+1)^2z - 24(n+1)$. Then $f_1(z) > 0$ for $z \ge 4$. Also, $f_1(3) = \frac{159}{10}$. Therefore, $f_1(z) > 0$ for $z \ge 3$. It can be seen that $f_2(n, z) = \frac{1}{3}n^3(z-3)z^2 + \frac{1}{3}(n-9)n^2z^3 + \frac{1}{3}n^2z^2(nz-15) + 6(n^2z-4) + 2nz^3 + (17n-15)z^2 + 12n(z-2) + 6z > 0$ for $n \ge 10$ and $z \ge 3$. Also, $f_2(n, 2) = 4(n-2)(n-3)^2 > 0$ for $n \ge 6$. We have $f_2(6, z) = z^2(120z-309)+294z-168 > 0$ and $f_2(8, z) = z^2(336z-711)+486z-216 > 0$ for $z \ge 3$. Therefore, $f_2(n, z) > 0$ for $n \ge 6$ and $z \ge 2$. Hence

$$LE_{CN}^+(\Gamma_G) - E_{CN}(\Gamma_G) > 0.$$

Again

$$LE_{CN}^{+}(\Gamma_{G}) - LE_{CN}(\Gamma_{G})$$

$$= \begin{cases} -\frac{8}{5}, & \text{for } n = 4 \& z = 2\\ \frac{-z(29z - 66) - 40}{10}, & \text{for } n = 4 \& z \ge 3\\ -\frac{(n((n-3)n+3)+1)z^{2} - 6((n-2)n+3)z + 8(n+1))}{2(n+1)}, & \text{otherwise.} \end{cases}$$

Clearly, -z(29z-66) - 40 < 0 for $z \ge 3$. Let $f_3(n,z) = -((n((n-3)n+3)+1)z^2 - 6((n-2)n+3)z + 8(n+1))$. Then $f_3(n,z) = -\frac{1}{2}(n-6)n^2z^2 - \frac{1}{2}n^2z(nz-12) - 3nz^2 - 6(2n-3)z - 8n - z^2 - 8 < 0$ for $n \ge 6$ and $z \ge 2$. Therefore

$$LE_{CN}^+(\Gamma_G) - LE_{CN}(\Gamma_G) < 0.$$

Hence, $E_{CN}(\Gamma_G) < LE_{CN}^+(\Gamma_G) < LE_{CN}(\Gamma_G)$. Case 2: *n* is odd

By Theorem 2.4 and [26, Theorem 2.14], we have

$$LE_{CN}^{+}(\Gamma_{G}) - E_{CN}(\Gamma_{G})$$

$$= \begin{cases} 0, & \text{for } n = 3, 5 \& z = 1 \\ 0, & \text{for } n = 3 \& z \ge 2 \\ \frac{(n-5)(n-3)}{n+1}, & \text{for } n \ge 7 \& z = 1 \\ \frac{z((n-3)(n-1)(n+1)z^{2} - (n(n(n+5)-21)+23)z+6(n+1)^{2}) - 16(n+1)}{2(n+1)}, & \text{otherwise.} \end{cases}$$

Clearly (n-5)(n-3) > 0 for $n \ge 7$. Let $f_4(n,z) = z((n-3)(n-1)(n+1)z^2 - (n(n(n+5)-21)+23)z+6(n+1)^2) - 16(n+1)$, where $n \ge 5$ and $z \ge 2$. Then $f_4(n,z) = \frac{1}{4}n^3(z-4)z^2 + \frac{1}{4}(n-12)n^2z^3 + \frac{1}{4}(n^2-4)nz^3 + \frac{1}{4}n^2z^2(nz-20)+2(3n^2z-8) + (21n-23)z^2 + 4n(3z-4) + 3z^3 + 6z > 0$ for $n \ge 13$ and $z \ge 4$. We have $f_4(n,2) = 4(n-2)(n-3)^2 > 0$ as $n \ge 5$; $f_4(n,3) = 18n^2(n-6) + 182n - 124 > 0$ for $n \ge 7$; and $f_4(5,3) = 336$. Therefore, $f_4(n,3) > 0$. Further, for $z \ge 4$ we have $f_4(5,z) = 24z^2(2z-7) + 216z - 96 > 0$; $f_4(7,z) = 16z^2(12z-29) + 384z - 128 > 0$; $f_4(9,z) = 8z^2(60z-121) + 600z - 160 > 0$ and $f_4(11,z) = 192z^2(5z-9) + 864z - 192 > 0$. Therefore, $f_4(n,z) > 0$. Thus

$$LE_{CN}^{+}(\Gamma_{G}) - E_{CN}(\Gamma_{G}) \begin{cases} = 0, & \text{for } n = 3 \& z \ge 1; n = 5 \& z = 1 \\ > 0, & \text{otherwise.} \end{cases}$$

Again

$$LE_{CN}^{+}(\Gamma_G) - LE_{CN}(\Gamma_G) = \begin{cases} 0, & \text{for } n = 3 \& z = 1\\ 0, & \text{for } n = 3 \& z \ge 2\\ 0, & \text{for } n = 5 \& z = 1\\ -\frac{(n-5)^2(n-3)}{2(n+1)}, & \text{for } n \ge 7 \& z = 1\\ -\frac{(nz+z-4)(((n-4)n+7)z-2(n+1))}{2(n+1)}, & \text{otherwise.} \end{cases}$$

Clearly, $-(n-5)^2(n-3) < 0$ for $n \ge 7$. Let $f_5(n, z) = -(nz+z-4)(((n-4)n+7)z-2(n+1))$. Then $f_5(n, z) = -\frac{1}{2}(n-6)n^2z^2 - \frac{1}{2}n^2z(nz-12) - 3nz^2 - 6(2n-5)z - 8n - 7z^2 - 8 < 0$ for $n \ge 7$ and $z \ge 2$. For $z \ge 2$ we have $f_5(5, z) = -24(z-1)(3z-2) < 0$. Therefore, $f_5(n, z) < 0$. Thus

$$LE_{CN}^{+}(\Gamma_{G}) - LE_{CN}(\Gamma_{G}) \begin{cases} = 0, & \text{for } n = 3 \& z \ge 1; \ n = 5 \& z = 1 \\ < 0, & \text{otherwise.} \end{cases}$$


Hence, $E_{CN}(\Gamma_G) = LE_{CN}^+(\Gamma_G) = LE_{CN}(\Gamma_G)$, if $n = 3 \& z \ge 1$ or n = 5 & z = 1. For all other cases, $E_{CN}(\Gamma_G) < LE_{CN}^+(\Gamma_G) < LE_{CN}(\Gamma_G)$. This completes the proof. \Box

We conclude this section with the following corollary.

Corollary 3.11. Let $G = D_{2n}, T_{4n}, U_{6n}, SD_{8n}, V_{8n}$ or $U_{(n,m)}$. Then

- (a) $E_{CN}(\Gamma_G) = LE_{CN}^+(\Gamma_G) = LE_{CN}(\Gamma_G)$ if and only if $G = D_6, D_8, D_{10}, D_{12}, T_8, T_{12}, SD_{28}, V_{16}, U_{6n} \text{ for } n \ge 2 \text{ and } U_{(n,m)} \text{ for } m = 3, 4, 6 \text{ and } n \ge 2.$
- (b) $E_{CN}(\Gamma_G) < LE_{CN}^+(\Gamma_G) < LE_{CN}(\Gamma_G)$ if and only G is not among the groups listed in (a).

In the following figures, closeness of various CN-energies of CCC-graphs of $D_{2n}, T_{4n}, U_{6n}, SD_{8n}, V_{8n}$ and $U_{(n,m)}$ are depicted.






FIGURE 1. CN-energies of $\Gamma_{D_{2n}}$, n is odd

FIGURE 2. CN-energies of $\Gamma_{D_{2n}}$, n is even




FIGURE 3. CN-energies of $\Gamma_{T_{4n}}$




FIGURE 7. CN-energies of $\Gamma_{V_{8n}}$, n is even





FIGURE 4. CN-energies of $\Gamma_{U_{(4,m)}}, m$ is odd






FIGURE 8. CN-energies of $\Gamma_{V_{8n}}$, n is odd

4. CONCLUSION

In this paper, we compute common neighborhood (signless) Laplacian spectrum and energy of CCC-graphs of certain finite non-abelian groups. We show that CCCgraphs of all the groups considered in this paper are CNL-integral and CNSL-integral. The common neighborhood spectrum and energy of CCC-graphs of theses groups are already computed in [26]. Analogous to the notion of super integral graph, we call a finite graph *super CN-integral* if it is CN-integral, CNL-integral and CNSL-integral. Thus, CCC-graphs of the groups considered in this paper are super CN-integral. It may be interesting to consider the following problem.

Problem 4.1. Characterize all finite non-abelian groups G such that Γ_G is super CN-integral.

The existence of finite non-abelian groups G such that Γ_G is CN-hyperenergetic is not clear (see [26]). However, there are finite non-abelian groups G such that Γ_G is CNborderenergetic (See [26, Theorem 3.6]), CNL-hyperenergetic/CNL-borderenergetic and CNSL-hyperenergetic/CSNL-borderenergetic (See Corollary 3.6). Thus the following problem is worth considering.

Problem 4.2. Characterize all finite non-abelian groups G such that Γ_G is CNborderenergetic/ CNL-hyperenergetic/ CNL-borderenergetic/ CNSL-hyperenergetic/ CNL-borderenergetic.

We have found several classes of finite non-abelian groups G such that $E_{CN}(\Gamma_G) = LE_{CN}(\Gamma_G) = LE_{CN}^+(\Gamma_G)$ in Subsection 3.1. Thus, we pose the following problem.

Problem 4.3. Characterize all finite non-abelian groups G such that

$$E_{CN}(\Gamma_G) = LE_{CN}(\Gamma_G) = LE_{CN}^+(\Gamma_G).$$

In Subsection 3.1, we have also found several classes of finite non-abelian groups G such that $E_{CN}(\Gamma_G) < LE_{CN}^+(\Gamma_G) < LE_{CN}(\Gamma_G)$. In [3, Theorem 4.6], it was observed that there are several classes of finite non-abelian groups G such that $E(\Gamma_G) < LE^+(\Gamma_G) < LE(\Gamma_G)$. It follows that there exist finite non-abelian groups such that $E(\Gamma_G)$, $LE^+(\Gamma_G)$, $LE(\Gamma_G)$ and $E_{CN}(\Gamma_G)$, $LE_{CN}^+(\Gamma_G)$, $LE_{CN}(\Gamma_G)$ behave similarly. Thus, the following problem arises naturally.

Problem 4.4. Determine all the finite non-abelian groups G such that $E(\Gamma_G)$, $LE^+(\Gamma_G)$, $LE(\Gamma_G)$ and $E_{CN}(\Gamma_G)$, $LE_{CN}^+(\Gamma_G)$, $LE_{CN}(\Gamma_G)$ behave similarly.

It is worth noting that problem similar to Problem 4.4 can also be asked for any finite graph.

In [20], Gutman et al. conjectured that $E(\mathcal{G}) \leq LE(\mathcal{G})$ for any finite graph \mathcal{G} but soon after the announcement, this conjecture was refuted [29, 36]. For the groups G, we consider in this paper, we have

(6) $E_{CN}(\Gamma_G) \leq L E_{CN}(\Gamma_G).$

In view of this it is too early to conjecture that the inequality (6) holds for CCCgraphs for any finite non-abelian group. However, one may consider the following problem.

Problem 4.5. Determine all the finite non-abelian groups such that the inequality (6) does not hold. In general, determine all the finite graphs \mathcal{G} such that the inequality $E_{CN}(\mathcal{G}) \leq LE_{CN}(\mathcal{G})$ does not hold.

Acknowledgments

The first author would like to thank DST for the INSPIRE Fellowship (IF200226).

References

- O. Ahmadi, N. Alon, I. F. Blake and I. E. Shaparlinski, Graphs with integral spectrum, *Linear Algebra and its Application* 430(1), 547-552, 2009.
- [2] A. Alwardi, N. D. Soner and I. Gutman, On the common-neighborhood energy of a graph, Bulletin. Classe des Sciences Mathématiques et Naturelles. Sciences Mathématiques 36, 49-59, 2011.
- [3] P. Bhowal and R. K. Nath, Spectral aspects of commuting conjugacy class graph of finite groups, Algebraic Structures and Their Applications 8(2), 67-118, 2021.
- [4] P. Bhowal and R. K. Nath, Genus of commuting conjugacy class graph of certain finite groups, Algebraic Structures and Their Applications 9(1), 93-108, 2022.
- [5] P. J. Cameron, Graphs defined on groups, International Journal of Group Theory 11(2), 53-107, 2022.
- [6] K. C. Das and S. A. Mojallal, On Laplacian energy of graphs, *Discrete Mathematics* 325, 52-64, 2014.
- [7] P. Dutta, B. Bagchi and R. K. Nath, Various energies of commuting graphs of finite non-abelian groups, *Khayyam Journal of Mathematics* 6(1), 27–45, 2020.
- [8] P. Dutta, J. Dutta and R. K. Nath, Laplacian spectrum of non-commuting graphs of finite groups, *Indian journal of pure and applied mathematics* 49(2), 205-216, 2018.
- [9] J. Dutta and R. K. Nath, Finite groups whose commuting graphs are integral, Matematički Vesnik 69(3), 226–230, 2017.
- [10] J. Dutta and R. K. Nath, Laplacian and signless Laplacian spectrum of commuting graphs of finite groups, *Khayyam Journal of Mathematics* 4(1), 77–87, 2018.
- [11] P. Dutta and R. K. Nath, On Laplacian energy of non-commuting graphs of finite groups, Journal of Linear and Topological Algebra 7(2), 121–132, 2018.
- [12] W. N. T. Fasfous and R. K. Nath, Inequalities involving energy and Laplacian energy of noncommuting graphs of finite groups, *Indian Journal of Pure and Applied Mathematics*, 2023 https://doi.org/10.1007/s13226-023-00519-7.
- [13] W. N. T. Fasfous, R. K. Nath and R. Sharafdini, Various spectra and energies of commuting graphs of finite rings, *Hacettepe Journal of Mathematics and Statistics* 49(6), 1915-1925, 2020.

- [14] W. N. T. Fasfous, R. Sharafdini, and R. K. Nath, Common neighborhood spectrum of commuting graphs of finite groups, Algebra and Discrete Mathematics 32(1), 33-48, 2021.
- [15] H. A. Ganie and S. Pirzada, On the bounds for signless Laplacian energy of a graph, Discrete Applied Mathematics 228(10), 3–13, 2017.
- [16] S. Gong, X. Li, G. Xu, I. Gutman, and B. Furtula, Borderenergetic Graphs, MATCH Communications in Mathematical and in Computer Chemistry 74, 321-332, 2015.
- [17] R. Grone and R. Merris, The Laplacian spectrum of a graph II, SIAM Journal of Discrete Mathematics 7(2), 221-299, 1994.
- [18] I. Gutman, The energy of a graph, Berichte der Mathematisch-Statistischen Sektion im Forschungszentrum Graz 103, 1-22, 1978.
- [19] I. Gutman, Hyperenergetic molecular graphs, Journal of the Serbian Chemical Society 64, 199-205, 1999.
- [20] I. Gutman, N. M. M. Abreu, C. T. M. Vinagre, A. S. Bonifacioa and S. Radenkovic, Relation between energy and Laplacian energy, *MATCH Communications in Mathematical and in Computer Chemistry* 59, 343-354, 2008.
- [21] I. Gutman and B. Furtula, Survey of Graph Energies, Mathematics Interdisciplinary Research 2, 85-129, 2017.
- [22] I. Gutman and B. Furtula, Graph energies and their applications, Bulletin. Classe des Sciences Mathématiques et Naturelles. Sciences Mathématiques 44, 29-45, 2019.
- [23] I. Gutman and B. Zhou, Laplacian energy of a graph, *Linear Algebra and its Applications* 414, 29-37, 2006.
- [24] F. Harary and A. J. Schwenk, Which graphs have integral spectra?, Graphs and Combinatorics, Lect. Notes Maths. 406, 45-51, 1974.
- [25] M. Herzog, M. Longobardi and M. Maj, On a commuting graph on conjugacy classes of groups, *Communications in Algebra* 37(10), 3369-3387, 2009.
- [26] F. E. Jannat and R. K. Nath, Common neighbourhood spectrum and energy of commuting conjugacy class graph, *Journal of Algebraic Systems* 12(2), 301-326, 2025.
- [27] F. E. Jannat and R. K. Nath, Common neighbourhood Laplacian and signless Laplacian spectra and energies of commuting graph, https://arxiv.org/pdf/2403.01082.pdf.
- [28] F. E. Jannat, R. K. Nath and K. C. Das, Common neighborhood energies and their relations with Zagreb index, https://arxiv.org/abs/2402.15416.
- [29] J. Liu, and B. Liu, On the relation between energy and Laplacian energy, MATCH Commun. Math. Comput. Chem. 61, 403–406, 2009.
- [30] A. Mohammadian, A. Erfanian, D. G. M. Farrokhi and B. Wilkens, Triangle-free commuting conjugacy class graphs, *Journal of Group Theory* 19, 1049-1061, 2016.
- [31] R. K. Nath, W. N. T. Fasfous, K. C. Das and Y. Shang, Common neighbourhood energy of commuting graphs of finite groups, *Symmetry* 13(9), 1651-1662, 2021.
- [32] M. A. Salahshour and A. R. Ashrafi, Commuting conjugacy class graphs of finite groups, Algebraic Structures and Their Applications 7(2), 135-145, 2020.
- [33] M. A. Salahshour and A. R. Ashrafi, Commuting conjugacy class graph of finite CA-groups, *Khayyam Journal of Mathematics* 6(1), 108-118, 2020.
- [34] M. A. Salahshour, Commuting conjugacy class graph of G when $\frac{G}{Z(G)} \cong D_{2n}$, Mathematics Interdisciplinary Research 1, 379-385, 2020.
- [35] S. K. Simic and Z. Stanic, Q-integral graphs with edge-degrees at most five, Discrete Mathematics 308, 4625-4634, 2008.

- [36] D. Stevanovic, I. Stankovic and M. Milosevic, On the relation between energy and Laplacian energy of graphs, MATCH Commun. Math. Comput. Chem. 61, 395–401, 2009.
- [37] Q. Tao, and Y. Hou, Q-borderenergetic graphs, AKCE International Journal of Graphs and Combinatorics 17(1), 38-44, 2020.
- [38] F. Tura, L-borderenergetic graphs, MATCH Communications in Mathematical and in Computer Chemistry 77, 37-44, 2017.
- [39] H. B. Walikar, H. S. Ramane, and P. R. Hampiholi, On the energy of a graph, *Graph connections*, Eds. R. Balakrishnan, H. M. Mulder and A. Vijayakumar, Allied publishers, New Delhi, 120-123, 1999.

FIRDOUS EE JANNAT, DEPARTMENT OF MATHEMATICAL SCIENCE, TEZPUR UNIVERSITY, NA-PAAM -784028, SONITPUR, ASSAM, INDIA.

Email address: firdusej@gmail.com

RAJAT KANTI NATH, DEPARTMENT OF MATHEMATICAL SCIENCE, TEZPUR UNIVERSITY, NA-PAAM -784028, SONITPUR, ASSAM, INDIA.

 $Email \ address: \verb"rajatkantinath@yahoo.com"$