

Tsirelson bounds for quantum correlations with indefinite causal order

Zixuan Liu^{1,2} and Giulio Chiribella^{1,2,3,4,*}

¹*QICI Quantum Information and Computation Initiative, Department of Computer Science, The University of Hong Kong, Pokfulam Road, Hong Kong*

²*HKU-Oxford Joint Laboratory for Quantum Information and Computation*

³*Department of Computer Science, University of Oxford, Wolfson Building, Parks Road, Oxford, UK*

⁴*Perimeter Institute for Theoretical Physics, 31 Caroline Street North, Waterloo, Ontario, Canada*

Quantum theory is in principle compatible with processes that violate causal inequalities, an analogue of Bell inequalities that constrain the correlations observed by a set of parties operating in a definite causal order. Since the introduction of causal inequalities, determining their maximum quantum violation, analogue to Tsirelson’s bound for Bell inequalities, has remained an open problem. Here we provide a general method for bounding the violation of arbitrary causal inequalities, establishing limits on the correlations achievable by arbitrary local experiments and by arbitrary quantum processes with indefinite causal order. We prove that the maximum violation is generally smaller than the algebraic maximum, and determine Tsirelson-like bounds for a class of causal inequalities including some of the most paradigmatic examples. Surprisingly, we find that the algebraic maximum of arbitrary causal inequalities can be achieved by a new type of processes that allow for information to flow in an indefinite temporal direction within the parties’ laboratories.

I. INTRODUCTION

Traditional formulations of physics generally assume that events take place in a well-defined causal order. On the other hand, it has been observed quantum theory is in principle compatible with scenarios where the order of events is indefinite [1–4], a phenomenon that has potential applications to quantum information [5–10] and is expected to play an integral role in a future theory of quantum gravity [11]. The possibility of indefinite causal order gives rise to an analogue of quantum nonlocality, originally discovered by Oreshkov, Costa, and Brukner (OCB) [2], who developed a framework for studying the most general correlations arising from quantum experiments performed in a set of local laboratories. In this context, the assumption of definite causal order among the experiments implies an analogue of Bell inequalities, called causal inequalities. Strikingly, OCB found that the validity of quantum theory in the local laboratories is in principle compatible with correlations that violate a causal inequality, hereafter referred to as the OCB inequality.

Over the past decade, quantum violations of causal inequalities were found in a variety of scenarios [12–17]. These findings raise a fundamental question: what is the maximum violation of causal inequalities allowed by quantum mechanics? Can it reach the maximum algebraic value of the corresponding correlations? The analogues of these questions have been extensively studied in the context of Bell inequalities, where they led to the problem of bounding the maximum Bell violations allowed by quantum mechanics [18–27]. In the paradigmatic case of the Clauser-Horne-Shimony-Holt (CHSH) inequality [28], the problem was famously solved by Tsirelson [18], who derived the maximum quantum violation, now known as the Tsirelson bound.

In stark contrast with the Bell scenario, little is known about the boundary of the set of quantum correlations

with indefinite causal order. For the OCB inequality, an upper bound on the violations achieved by a restricted class of local experiments was derived in Ref. [29], but whether larger violations could be achieved by more general experiments remained as an open question. For another causal inequality, associated to the Guess-Your-Neighbor’s Input game [30], it was recently proven that the maximum violation has to be strictly smaller than the algebraic maximum [31]. In general, however, no upper bound other than the algebraic maximum has been found for any causal inequality so far. In addition, new questions have recently arisen from the introduction of a new class of scenarios where not only the causal order of the experiments, but also the temporal direction of the information flow within the local laboratories can be indefinite [32]. Can these scenarios lead to even larger violations? And in the affirmative case, where does the boundary lie between the correlations achievable with indefinite causal order alone and those achievable when indefinite causal order is combined with indefinite temporal direction?

Here we develop a general method for bounding the maximum violation of arbitrary causal inequalities by quantum processes with indefinite causal order. As an application of the general method, we establish the analogue of Tsirelson bound for the OCB inequality and for a class of causal inequalities, which we name single-trigger inequalities. We then show that allowing information to flow in an indefinite temporal direction within the local laboratories leads to a violation of all causal inequalities to their algebraic maximum for arbitrary numbers of parties, outcomes, and settings. Remarkably, this extreme violation of causal inequalities can be achieved even if all local laboratories are restricted to classical operations. Overall, our findings open up a search for physical principles determining the boundaries of the set of quantum correlations with indefinite causal order, and a search for applications of causal inequalities to quantum cryptography and other quantum technologies.

II. RESULTS

Single-trigger causal inequalities. Here we introduce a special class of causal inequalities that provide the foundation of our method. For the inequalities in this class, the maximum violation can be determined explicitly by a semidefinite program, which in turn can be used to provide upper bounds on the violation of arbitrary causal inequalities.

In the framework of causal inequalities [2, 12, 16], a set of parties operate in different regions of spacetime, performing local operations in their laboratories. The interaction between the parties' laboratories and the outside world takes place only at specific moments: in the simplest presentation of the framework, the i -th laboratory is assumed to be shielded from the outside world at all times, except for two moments t_i and $t'_i \geq t_i$ when a shutter is opened, allowing physical systems to enter and exit the laboratory, respectively [2].

In the time between t_1 and t'_i the i -th party performs an experiment, obtaining an outcome. We denote by x_i (a_i) the setting (outcome) of the experiment performed by the i -th party, and by $\vec{x} = (x_1, \dots, x_N)$ ($\vec{a} = (a_1, \dots, a_N)$) the vector of all parties' settings (outcomes). An N -partite correlation function is an expression of the form

$$\mathcal{I} = \sum_{\vec{x}, \vec{a}} \alpha_{\vec{x}, \vec{a}} p(\vec{a} | \vec{x}), \quad (1)$$

where each $\alpha_{\vec{x}, \vec{a}}$ is a real coefficient and $p(\vec{a} | \vec{x})$ is the conditional probability distribution of the outcomes given the settings.

When the parties operate in a definite causal order, the probability distribution $p(\vec{a} | \vec{x})$ is subject to a set of linear constraints [2, 12, 16]. In the case of $N = 2$ parties, Alice and Bob, the constraints have a simple expression: if Alice's experiment precedes Bob's experiment, then the probability distribution of Alice's outcomes must be independent of Bob's settings, namely $p_A(a_1 | x_1, x_2) = p_A(a_1 | x_1, x'_2), \forall x_2, x'_2$, with $p_A(a_1 | x_1, x_2) := \sum_{a_2} p(a_1, a_2 | x_1, x_2)$. Vice-versa, if Bob's experiment precedes Alice's experiment, then Bob's outcomes must be independent of Alice's settings, namely $p_B(a_2 | x_1, x_2) = p_B(a_2 | x'_1, x_2), \forall x_1, x'_1$, with $p_B(a_2 | x_1, x_2) := \sum_{a_1} p(a_1, a_2 | x_1, x_2)$. A probability distribution $p(a_1, a_2 | x_1, x_2)$ is called *causal* if it is a random mixture of probability distributions corresponding to scenarios in which either Alice's experiment precedes Bob's or Bob's experiment precedes Alice's [2]. In the multipartite case, causal probability distributions can arise in a more general way, by dynamically controlling the order of some of the parties based on outcomes obtained by some of the other parties [12, 16].

A causal inequality is an upper bound on the correlations achievable by causal probability distributions; explicitly, it is a bound of the form $\mathcal{I}^{\text{causal}} \leq \beta$, where $\mathcal{I}^{\text{causal}}$ is the maximum correlation achieved by causal probability distributions and $\beta \in \mathbb{R}$ is some constant.

The first example of a causal inequality was introduced by OCB [2], who showed that quantum theory is in principle compatible with its violation.

In general, the violation of causal inequalities takes place when the experiments performed by the different parties are connected in an indefinite order. The connections are implemented by suitable processes, which are in principle compatible with the validity of quantum theory in the parties' local laboratories [2, 3, 33–35]. In the N -partite setting, a process of this kind is represented by a linear map \mathcal{S} that transforms the parties' local operations into the conditional probability distribution

$$p(\vec{a} | \vec{x}) = \mathcal{S} \left(\mathcal{M}_{a_1 | x_1}^{(1)}, \dots, \mathcal{M}_{a_N | x_N}^{(N)} \right), \quad (2)$$

where, for every $i \in \{1, \dots, N\}$, $\mathcal{M}_{a_i | x_i}^{(i)}$ is the quantum operation occurring in the i -th laboratory when an experiment with setting x_i produces the outcome a_i . The positivity and normalization of the probability distributions $p(\vec{a} | \vec{x})$ for every possible set of experiments performed in the parties' laboratories place constraints on the admissible maps \mathcal{S} (see Appendix A for details). The set of maps \mathcal{S} satisfying these constraints contains all possible processes with definite causal order, as well as other processes that are incompatible with any pre-defined causal order, hereafter referred to as quantum processes with indefinite causal order (ICO).

Given a correlation function \mathcal{I} , a fundamental problem is to determine the maximum of \mathcal{I} over all possible probability distributions (2) generated by local quantum experiments and by quantum processes with ICO. In the following, we will denote the maximum by \mathcal{I}^{ICO} , and call it the ICO bound. The ICO bound is an analogue of the Tsirelson bound [18] for causal inequalities: the value of \mathcal{I}^{ICO} determines whether quantum mechanics allows for a violation of the causal inequality for the correlation \mathcal{I} , and, in the affirmative case, it provides the maximum quantum violation.

Finding the maximum of a correlation over all local experiments and over all ICO processes is a non-convex optimization problem that in principle requires optimization over quantum systems of arbitrary dimension, as in the case of Bell correlations. Compared to the maximization of Bell correlations, however, the maximization of ICO correlations appears to be a harder problem, and the exact value of the ICO bound has remained unknown until now for all causal inequalities, except those that have been shown to be violated to their algebraic maximum.

We now introduce a class of N -partite correlations for which the maximum quantum violation can be computed explicitly. These correlations, called single-trigger correlations, can be thought as the score achieved by the parties in a game where each party is asked a question and the payoff depends on the party's answer only if the party receives one specific question, called the "trigger." The precise definition is as follows: an N -partite correlation (1) is *single-trigger* if for every party i there exists a setting ξ_i (the "trigger") such that $\alpha_{\vec{x}, \vec{a}}$ depends on a_i

only if $x_i = \xi_i$.

An example of single-trigger correlation in the $N = 2$ case arises from the game known as Lazy Guess Your Neighbor's Input [13]. This game involves two parties, Alice and Bob, each of which has to guess the other party's setting when her/his own setting is equal to 1. The probability of success is $P_{\text{succ}}^{11} := p(1,1|1,1)$ if both parties have input 1, $P_{\text{succ}}^{01} := p_B(0|0,1)$ or $P_{\text{succ}}^{10} := p_A(0|1,0)$ if only one of the parties has input 1, and is equal to one when both parties have input 0. Assuming uniform probabilities for the possible inputs, the average probability of success is $\mathcal{I}_{\text{LGYNI}} = (P_{\text{succ}}^{11} + P_{\text{succ}}^{01} + P_{\text{succ}}^{10} + 1)/4$.

Crucially, every correlation \mathcal{I} can be decomposed into a sum of single-trigger correlations, and the number of non-zero terms in the sum is at most equal to the total number of settings, namely $n_1 \cdot n_2 \cdots n_N$, where n_i is the number of settings for the i -th party. Indeed, the existence of a decomposition with this number of terms is immediate, as one can always write the correlation $\mathcal{I} = \sum_{\vec{x}, \vec{a}} \alpha_{\vec{x}, \vec{a}} p(\vec{a}|\vec{x})$ as $\mathcal{I} = \sum_{\vec{\xi}} \mathcal{I}_{\vec{\xi}}$, where $\mathcal{I}_{\vec{\xi}}$ is the single-trigger correlation with coefficients $\alpha_{\vec{x}, \vec{a}}^{\vec{\xi}} := \delta_{\vec{x}, \vec{\xi}} \alpha_{\vec{x}, \vec{a}}$. In general, there can be other decompositions with a smaller number of non-zero terms.

In the following, the causal inequalities associated to single-trigger correlations will be called ‘‘single-trigger causal inequalities.’’ For these inequalities, we will provide an explicit expression for the ICO bound. Since every possible correlation can be decomposed into a sum of single-trigger correlations, this bound will imply general upper bounds on the violation of arbitrary causal inequalities.

Maximum quantum violation of single-trigger causal inequalities. We now provide the central result of the paper: the ICO bound for single-trigger causal inequalities. A crucial feature of our result is that it reduces the original non-convex problem in the definition of the ICO bound to a semidefinite program (SDP) on a space of fixed dimension, depending only on the number of parties, settings, and outcomes.

In addition to the value of the ICO bound, our result provides a canonical choice of local experiments that achieve the bound. The canonical choice for the i -th party is to use as input a quantum system of dimension m_i , equal to the number of possible outcomes, and to append to it an auxiliary system of dimension n_i , equal to the number of possible settings. In other words, the canonical strategy uses an input system with Hilbert space $\mathcal{H}_{\text{in}}^{(i)} = \mathbb{C}^{m_i}$ and an output system with Hilbert space $\mathcal{H}_{\text{out}}^{(i)} = \mathcal{H}_{\text{in}}^{(i)} \otimes \mathcal{H}_i^{\text{aux}}$ with $\mathcal{H}_i^{\text{aux}} = \mathbb{C}^{n_i}$. The auxiliary system is used by the i -th party to communicate their setting to the outside world: explicitly, the value of the setting will be encoded in the pure state $|x_i\rangle\langle x_i|$. If the setting is equal the trigger, then the party will measure the input system on the canonical basis for $\mathcal{H}_{\text{in}}^{(i)}$. If the setting is not the trigger, the party will not perform any measurement and will

just output a random outcome. Mathematically, this strategy is described by quantum operations of the form $\mathcal{M}_{a_i|x_i}^{(i)*} = \mathcal{N}_{a_i|x_i}^{(i)*} \otimes |x_i\rangle\langle x_i|$, where $\mathcal{N}_{a_i|x_i}^{(i)*}$ is a quantum operation acting only on the input system. Explicitly, the action of this quantum operation on a generic density matrix ρ is $\mathcal{N}_{a_i|x_i}^{(i)*}(\rho) = \langle a_i|\rho|a_i\rangle |a_i\rangle\langle a_i|$ if $x_i = \xi_i$ and $\mathcal{N}_{a_i|x_i}^{(i)*}(\rho) = \rho/m_i$ if $x_i \neq \xi_i$.

The value of the ICO bound is expressed in terms of the canonical strategy. To this purpose, we define the linear map $\mathcal{M}_{\mathcal{I}}^* = \sum_{\vec{x}, \vec{a}} \alpha_{\vec{x}, \vec{a}} \mathcal{M}_{\vec{a}|\vec{x}}^*$, where $\{\alpha_{\vec{x}, \vec{a}}\}$ are the coefficient of the correlation \mathcal{I} [cf. Eq. (1)] and $\mathcal{M}_{\vec{a}|\vec{x}}^* := \mathcal{M}_{a_1|x_1}^{(1)*} \otimes \mathcal{M}_{a_2|x_2}^{(2)*} \otimes \cdots \otimes \mathcal{M}_{a_N|x_N}^{(N)*}$ are the parties' local operations. We then prove that the ICO bound is a measure of the deviation between the map $\mathcal{M}_{\mathcal{I}}^*$ and the set of N -partite no-signalling channels, that is, the set of quantum processes that do not allow to any subset of the parties to transmit information to any other subset (see Methods). Without loss of generality, here we focus our attention on the case where all the coefficients $\alpha_{\vec{x}, \vec{a}}$ are nonnegative. In this case, we show that the single-trigger ICO bound is

$$\mathcal{I}_{\text{single-trigger}}^{\text{ICO}} = 2^{D_{\text{max}}(\mathcal{M}_{\mathcal{I}}^* \parallel \text{NoSig})}, \quad (3)$$

where NoSig is the set of N -partite no-signalling channels, $D_{\text{max}}(\mathcal{A} \parallel \mathcal{B}) := \min\{\lambda \mid \mathcal{A} \leq 2^\lambda \mathcal{B}\}$ is the max relative entropy between two completely positive maps \mathcal{A} and \mathcal{B} (with the notation $\mathcal{A} \leq \mathcal{B}$ meaning that $\mathcal{B} - \mathcal{A}$ is completely positive), and, for a set of completely positive maps \mathcal{S} , $D_{\text{max}}(\mathcal{A} \parallel \mathcal{S}) := \max_{\mathcal{B} \in \mathcal{S}} D_{\text{max}}(\mathcal{A} \parallel \mathcal{B})$.

The quantity $D_{\text{max}}(\mathcal{M}_{\mathcal{I}}^* \parallel \text{NoSig})$ appearing in the ICO bound is also known as the max relative entropy of signalling [34]. Eq. (3) provides an SDP expression for the ICO bound of all single-trigger causal inequalities. In particular, it yields the ICO bound $\mathcal{I}_{\text{LGYNI}}^{\text{ICO}} \approx 0.8194$ for the single-trigger causal inequality corresponding to the LGYNI game. This result provides the answer to an open question raised in Ref. [13], where the value $\mathcal{I}_{\text{LGYNI}} \approx 0.8194$ was obtained by optimizing over local experiments and ICO processes under the assumption that all quantum systems in the parties' local laboratories are two-dimensional.

Bound for arbitrary causal inequalities. The ICO bound for single-trigger causal inequalities directly implies a general bound for arbitrary causal inequalities. The bound follows from the fact that every correlation \mathcal{I} can be decomposed as a sum of single-trigger correlations, and therefore its ICO bound cannot exceed the sum of the ICO bounds for the single-trigger correlations in its decomposition. Optimizing over all possible decompositions, one obtains the bound

$$\mathcal{I}^{\text{ICO}} \leq \min_{\{\mathcal{I}_j\}_{j=1}^n} \left\{ \sum_{j=1}^n \mathcal{I}_j^{\text{ICO}} \right\}, \quad (4)$$

where $\{\mathcal{I}_j\}_{j=1}^n$ are single-trigger correlations satisfying the condition $\sum_{j=1}^n \mathcal{I}_j = \mathcal{I}$.

Crucially, the evaluation of the right-hand-side of Eq. (4) is an SDP on a space of fixed dimension (See Appendix A). The solution of this SDP provides a computable upper bound on the violation of causal inequalities for every desired correlation \mathcal{I} . Mathematically, the upper bound can be seen as an SDP relaxation of the original problem of computing the ICO bound.

In the next sections, we show three important implications of our general results.

Maximum quantum violation of the OCB inequality. The first example of a causal inequality is the OCB inequality [2], which corresponds to a bipartite scenario in which a_1, a_2 , and x_1 are bits, while x_2 is a pair of bits, here denoted by b and c , respectively. The inequality arises from a game, in which either Alice is asked to guess Bob's bit b or Bob is asked to guess Alice's bit x_1 , depending on the value of bit c . When the order between Alice's and Bob's experiments is well-defined, only one player can communicate the value of his/her input to the other player, and therefore one of the two players must make a random guess of the other players' input. Hence, every causal probability distribution satisfies the constraint that the probability that Alice's guess is correct, plus the probability that Bob's guess is correct is upper bounded as $\mathcal{I}_{\text{OCB}}^{\text{causal}} := P(a_1 = b | c = 0) + P(a_2 = x_1 | c = 1) \leq \frac{3}{2}$, with $P(a_1 = b | c = 0) := \sum_{x_1, b} p_A(b | x_1, b, 0)/4$ and $P(a_2 = x_1 | c = 1) := \sum_{x_1, b} p_B(x_1 | x_1, b, 1)/4$.

OCB showed that the above causal inequality can be violated by a quantum ICO process and by suitable experiments in Alice's and Bob's laboratories, the combination of which reaches the value

$$\mathcal{I}_{\text{OCB}} = 1 + \frac{1}{\sqrt{2}}. \quad (5)$$

A fundamental question raised in the original OCB paper is whether higher violations are possible, and, in case they are, what is the maximum violation. The value (5) was shown to be maximum among the violations achievable with a restricted set of local measurements, involving measurement and reparations with traceless binary observables [29]. Whether more general types of measurements could lead to higher violations, however, remained as an open question until now. In fact, even the most basic question of whether the violation of the OCB inequality over arbitrary measurements and arbitrary ICO processes can reach its maximum algebraic value $\mathcal{I}_{\text{OCB}} = 2$ had remained unanswered so far.

We now solve the problem in full generality, proving that (5) is indeed the largest violation allowed by arbitrary quantum processes with indefinite causal order and by arbitrary operations in Alice's and Bob's laboratory. In fact, we derive the exact ICO bound for a version of the OCB correlation, called the tilted OCB correlation [36] and given by

$$\mathcal{I}_{\text{OCB}, \alpha} = P(a_1 = b | c = 0) + \alpha P(a_2 = x_1 | c = 1), \quad (6)$$

where α is an arbitrary real number. To derive the ICO bound, we observe that the tilted OCB correlation is a random mixture of two single-trigger correlations and we evaluate the ICO bound (3) for these correlations. Using these two ICO bounds, we provide an upper bound on the quantum violation, and show that it coincides with the value found in [36] with a specific example of ICO process (see Appendix B). All together, these results establish the ICO bound

$$\mathcal{I}_{\text{OCB}, \alpha}^{\text{ICO}} = \frac{1 + \alpha + \sqrt{1 + \alpha^2}}{2}, \quad (7)$$

thereby providing an analogue of the Tsirelson bound for indefinite causal order.




FIG. 1. Geometry of causal, ICO, and general probability distributions. The figure shows the two probabilities appearing in the OCB correlation. The inner square (in green) corresponds to the values achievable by causal probability distributions, while the circle (in blue) corresponds to the values achievable through quantum ICO processes. The outer square (in violet) corresponds to the values achievable by arbitrary, unconstrained probability distributions.

The quantum ICO set of correlations. Our results provide insights into the geometry of the set of probability distributions generated by quantum ICO processes. Let us consider the paradigmatic case of the OCB game, and visualize the possible values of Alice's and Bob's success probabilities $P_A := P(a_1 = b | c = 0)$ and $P_B := P(a_2 = x_1 | c = 1)$ by representing them in a two-dimensional plane. With this notation, the ICO bound Eq. (7) is equivalent to the condition $\cos \theta (P_A - 1/2) + \sin \theta (P_B - 1/2) \leq 1/2$ for $\theta = \arctan \alpha$. This condition identifies a circle of radius $1/2$ centred around the point $(1/2, 1/2)$, as illustrated in Figure 1. In the figure, we also show the area corresponding to causal probability distributions, which corresponds to a square inscribed inside the circle. Finally, arbitrary unconstrained probability distributions occupy the square circumscribing the circle of quantum ICO probabilities.

Intriguingly, Figure 1 coincides with the analogous picture in the case of the CHSH inequality, where the smaller

square is the set of probability distribution allowed by local realism, the circle is the set of probability distributions allowed by quantum mechanics, and the larger square is the set of general no-signalling probability distributions [37]. Later in this paper, we will show that, in the case of causal inequalities, the largest set can be achieved by probability distributions generated by a new class of operations that do not assume a definite direction of time outside Alice's and Bob's laboratories.

In Appendix C, we provide further insights into the geometry of quantum correlations with indefinite causal order by analyzing the set of correlations arising from the LGYNI game and a tilted version thereof.

Bound for Guess Your Neighbor's Input. Guess Your Neighbor's input (GYNI) [30] is a multipartite quantum game where each party has to guess the input of one of its neighbors. This game is well known in the study of quantum nonlocality, being the first example of a tight Bell inequality with no quantum violation. In the two-party setting, the probability of success is $\mathcal{I}_{\text{GYNI}} = P(a_1 = x_2, a_2 = x_1)$. GYNI has also been studied in the context of causal inequalities [13, 38], where it was observed that causal probability distributions satisfy the bound $\mathcal{I}_{\text{GYNI}}^{\text{causal}} \leq 1/2$ when the settings are uniformly random, while quantum ICO processes can violate this bound. The question about the maximum violation of the GYNI causal inequality has been tackled in Refs. [31, 38]. In particular, Ref. [31] proved that the quantum ICO violation cannot reach the algebraic maximum $\mathcal{I}_{\text{GYNI}} = 1$. However, no explicit bound on the maximum violation other than the algebraic maximum has been known up to now.

Using our general expression (4), we can now show that the ICO bound for the GYNI game satisfies the inequality $\mathcal{I}_{\text{GYNI}}^{\text{ICO}} \leq 0.7592$. At present, this inequality represents the state-of-the-art in upper bounding the quantum ICO violation of the GYNI inequality. Nevertheless, finding the exact value of $\mathcal{I}_{\text{GYNI}}^{\text{ICO}}$ remains an open problem. Previous numerical results based on a see-saw algorithm using quantum systems of dimension 5 showed that quantum ICO processes can reach the value $\mathcal{I}_{\text{GYNI}} \approx 0.6218$ [13]. An interesting question is whether increasing the dimension of the local quantum systems could yield correlations that go all the way to the upper bound 0.7592.

Classical processes with indefinite time direction. We now introduce a new type of processes that allow for the violation of all causal inequalities up to the maximum algebraic value. These processes are the classical version of the quantum processes with indefinite time direction introduced in Ref. [32]. They are in principle compatible with the validity of classical physics in the laboratories of the different parties, but do not assume a privileged direction of time outside each laboratory.

Recall that in the framework of causal inequalities the laboratory of the i -th party is shielded from the outside world at all times, except for two moments t_i and $t'_i \geq t_i$, in which a physical system is allowed to enter and exit

the laboratory, respectively. Typically, it is assumed that information flows through the laboratory from the past to the future, namely that the state of the system at the earlier time t_i can be chosen freely outside the party's laboratory, while the state of the system at the later time t'_i is obtained by applying the party's operation to the state at time t_i . In principle, however, also the opposite situation is logically conceivable: the state of the system at later time t'_i could be freely chosen outside the party's laboratory, and the state of the system at the earlier time t_i could be obtained by applying the party's operation to the state at time t'_i .

The lack of a privileged direction of time outside the parties' laboratories implies that the operations performed inside the laboratories must be compatible with two alternative directions of the information flow. In the quantum setting, it has been shown that compatibility requires the party's operations to be described by bistochastic quantum channels, that is, trace-preserving completely positive maps that preserve the maximally mixed state [32]. In the following, we will focus on the classical setting, which is equivalent to a quantum setting where all systems are completely decohered in a fixed basis. In this case, a classical channel can be described by a conditional probability distribution specifying the probability $q^{(i)}(s'_i | s_i) \geq 0$ that an input state s_i at time t_i is transformed into an output state s'_i at time t'_i . Compatibility with two alternative directions of the information flow requires the probability distribution to be bistochastic, that is, to satisfy both conditions $\sum_{s'_i} q^{(i)}(s'_i | s_i) = 1$ and $\sum_{s_i} q^{(i)}(s'_i | s_i) = 1$. The most general experiments performed by the i -th party then correspond to classical bistochastic instruments, mathematically described by subnormalized probability distributions $q_{a_i | x_i}^{(i)}(s'_i | s_i)$ satisfying the condition that $\sum_{a_i} q_{a_i | x_i}^{(i)}(s'_i | s_i)$ is a bistochastic probability distribution for every setting x_i .

The most general processes that could in principle connect the local operations performed by the parties are described by linear maps \mathcal{S} that transform the parties' local operations into conditional probability distributions of the form

$$p(\vec{a} | \vec{x}) = \mathcal{S} \left(q_{a_1 | x_1}^{(1)}, \dots, q_{a_N | x_N}^{(N)} \right). \quad (8)$$

As in the case of ICO, the positivity and normalization of the probability distributions $p(\vec{a} | \vec{x})$ for all possible operations place constraints on the admissible maps \mathcal{S} . An explicit characterization of these constraints in the bipartite case is provided in Appendix D. The set of maps \mathcal{S} satisfying these constraints contains all possible processes with definite causal order, all possible ICO processes, and a new type of processes that are incompatible with both a pre-defined causal order and a pre-defined direction of time. We call these processes as classical processes with indefinite causal order and time direction (ICOTD).

Reaching the algebraic maximum. We now show that classical ICOTD processes can reach the algebraic

maximum of every causal inequality. To prove this result, we start from the GYNI game, which can be won with certainty only if Alice and Bob can perfectly signal to one another.

Consider a scenario where all the classical systems entering and existing the parties' laboratories are bits. To win the GYNI game, Alice and Bob adopt the following strategy: First, they measure the values of the bits s_1 and s_2 entering their laboratories, respectively, and, depending on their settings, they either leave the bit unchanged or they flip them, obtaining the new values $s'_1 = s_1 \oplus x_1$ and $s'_2 = s_2 \oplus x_2$, respectively (here \oplus denotes addition modulo 2). The final value of the bit then becomes the outcome of their measurements, namely $a_1 = s'_1$ and $a_2 = s'_2$. Mathematically, this strategy corresponds to the classical bistochastic instruments $q_{a_1|x_1}^{(1)}(s'_1|s_1) = \delta_{s'_1, s_1 \oplus x_1} \delta_{a_1, s'_1}$ for Alice and $q_{a_2|x_2}^{(2)}(s'_2|s_2) = \delta_{s'_2, s_2 \oplus x_2} \delta_{a_2, s'_2}$ for Bob. To achieve unit probability of winning the GYNI game, Alice's and Bob's local experiments are connected by a deterministic process \mathcal{S} , which can be conveniently represented as a function that maps the final values of the bits (s'_1, s'_2) into their initial values (s_1, s_2) according to the rule

$$s_1 = s'_1 \oplus s'_2 \quad \text{and} \quad s_2 = s'_1 \oplus s'_2. \quad (9)$$

The action of the process \mathcal{S} on the parties' operations produces the probability distribution $\mathcal{S}(q_{a_1|x_1}^{(1)}, q_{a_2|x_2}^{(2)}) = \sum_{s'_1, s'_2} q_{a_1|x_1}^{(1)}(s'_1|s'_1 \oplus s'_2) q_{a_2|x_2}^{(2)}(s'_2|s'_1 \oplus s'_2)$. In Appendix E, we show that the process \mathcal{S} is logically consistent, in the sense that it gives rise to a normalized probability distribution for every choice of bistochastic instruments $q_{a_1|x_1}^{(1)}$ and $q_{a_2|x_2}^{(2)}$. For the specific choice described earlier in this paragraph, the process \mathcal{S} generates the perfect signalling distribution $p(a_1, a_2 | x_1, x_2) = \delta_{a_1, x_2} \delta_{a_2, x_1}$, which allows Alice and Bob to win the GYNI game with certainty.

Notice that the perfect signalling achieved by classical ICOTD processes can be used to win not only the GYNI game, but also to violate every bipartite causal inequality up to its algebraic maximum. This finding shows a striking difference between indefinite causal order alone and indefinite causal order combined with indefinite time direction: in the ICO case it is known that no classical bipartite process can violate any causal inequality [2], whereas in the ICOTD case there exists a bipartite process that can reach the algebraic maximum violation for all causal inequalities.

In Appendix E, we extend the above result to arbitrary causal inequalities with arbitrary numbers of parties, settings, and outcomes. Specifically, we prove that every multipartite conditional probability distribution can be generated by choices of classical bistochastic instruments and classical ICOTD processes. Again, this result shows a difference between ICO and ICOTD processes: while in the ICO case it is known that classical multipartite processes can sometimes violate causal inequalities and even

reach the algebraic maximum [14], in the ICOTD case there exists a classical process that attains the algebraic maximum for all possible causal inequalities.

III. DISCUSSION

Our general bound on the violation of causal inequalities is based on an SDP relaxation of the original problem of computing the quantum ICO bound. For certain correlations, such as the OCB correlation and all single-trigger correlations, we have shown that the SDP gives the exact value of the quantum ICO bound. In principle, one could ask our bound is tight for all possible causal inequalities. The analogy with Bell inequalities, however, suggests a negative answer. In Bell scenarios, a converging sequence of upper bounds on the value of maximal quantum violations is provided by the Navascués-Pironio-Acín SDP hierarchy [22, 39]. The analogy with this situation suggests that our SDP relaxation may be just the first level of a similar hierarchy of SDPs. Determining whether this analogy is correct, and, in the affirmative case, identifying the other levels of the hierarchy are among the most important research directions opened by our work. Another interesting direction is to extend our method for the calculation of the ICO bound to other type of inequalities with non-trivial causal structure, such as the inequalities recently studied in Refs. [40, 41].

A natural development of our research is to establish self-testing results for causal inequalities, in analogy to self-testing for Bell inequalities [42, 43]. For example, it is interesting to determine whether the OCB process is the only quantum process (up to local transformations) that achieves the maximum violation of the OCB inequality. More broadly, it is interesting to search for applications of the violation of causal inequalities to quantum technologies, in analogy to the applications of the violation of Bell inequalities in quantum cryptography [44–46], communication complexity [47], and control of quantum hardware [48]. While the physical realization of general quantum ICO processes is still an open problem, exploring the potential applications of causal inequalities contributes to a deeper information-theoretic understanding of indefinite causal order in quantum theory.

Finally, our results open up a search for physical principles capable of explaining why the violation of causal inequalities by quantum ICO processes is not equal, in general, to the algebraic maximum, and, of determining the exact value of the quantum violation. In the context of Bell inequalities, the analogue question was originally raised by Popescu and Rohrlich [19], and led to the discovery of new information theoretic principles, such as non-trivial communication complexity [20, 21, 23], non-trivial nonlocal computation [24], information causality [25], macroscopic locality [26], and local orthogonality [27]. The formulation of new principles that capture the quantum violation of causal inequalities is likely to be valuable for the understanding of quantum physics in

spacetime, and may offer guidance in the long-standing problem of unifying quantum theory with Einstein's theory of relativity.

IV. METHODS

Labelled projective instruments. We now develop a way to reduce the search over the set of all possible local experiments to a search over a smaller set, generated by ideal measurements and ideal state preparations. This reduction plays a similar role to the reduction to projective measurements in Bell scenarios.

In the standard framework of quantum theory, an experiment with m possible outcomes is described by a quantum instrument [49–51], that is, a collection of completely positive maps $(\mathcal{M}_a)_{a=0}^{m-1}$ satisfying the condition that their sum is trace-preserving. Each map \mathcal{M}_a describes a physical process transforming an input system with Hilbert space \mathcal{H}_{in} into a (possibly different) output system, with Hilbert space \mathcal{H}_{out} . In the special case $\mathcal{H}_{\text{in}} = \mathcal{H}_{\text{out}}$, an instrument $(\mathcal{M}_a)_{a=0}^{m-1}$ is called projective if each map \mathcal{M}_a is of the form $\mathcal{M}_a(\rho) = P_a \rho P_a$, where P_a is a projector and the projectors $(P_a)_{a=0}^{m-1}$ form a resolution of the identity.

We now introduce the notion of labelled projective instruments, that is, instruments that output a label describing their settings. For every setting x , let $(\mathcal{N}_{a|x})_{a=0}^{m-1}$ be a projective instrument on a quantum system with Hilbert space \mathcal{H}_{in} , and let $(\rho_x)_{x=0}^{n-1}$ be a set of perfectly distinguishable density matrices for an auxiliary quantum system with Hilbert space \mathcal{H}_{aux} . A labelled projective instrument is an instrument $(\mathcal{M}_{a|x})_{a=0}^{m-1}$ of the form $\mathcal{M}_{a|x} = \mathcal{N}_{a|x} \otimes \rho_x$.

Our first result is that the optimization of a conditional probability distribution $p(\vec{a}|\vec{x})$ over arbitrary instruments and arbitrary ICO processes can be restricted without loss of generality to an optimization over labelled projective instruments. The key result is the following theorem:

Theorem 1 *For every set of quantum instruments $(\mathcal{M}_{a_i|x_i}^{(i)})_{a_i=0}^{m_i-1}$, $x_i \in \{0, \dots, n_i - 1\}$, $i \in \{1, \dots, N\}$ and every process \mathcal{S} acting on them, there exists a set of labelled projective instruments $(\mathcal{M}_{a_i|x_i}^{(i)'})_{a_i=0}^{m_i-1}$, $x_i \in \{0, \dots, n_i - 1\}$, $i \in \{1, \dots, N\}$ (with possibly different input and output systems) and a process \mathcal{S}' acting on them such that*

$$\mathcal{S}(\mathcal{M}_{a_1|x_1}^{(1)}, \dots, \mathcal{M}_{a_N|x_N}^{(N)}) = \mathcal{S}'(\mathcal{M}_{a_1|x_1}^{(1)'}, \dots, \mathcal{M}_{a_N|x_N}^{(N)'}) \quad (10)$$

for all settings \vec{x} and all outcomes \vec{a} . In addition, the labelled projective measurements associated to each party i can be chosen without loss of generality to have projectors of the same rank r_i for every outcome a_i and every setting x_i .

The proof uses Ozawa's dilation theorem [50] for quantum instruments, combined with a use of auxiliary systems to include unitary state changes depending on the settings. The details of the proof are provided in Appendix F.

Theorem 1 can be used in every problem involving the joint optimization of local instruments and global ICO processes. In particular, it can be used to simplify the search for the ICO bound of arbitrary causal inequalities. In the next section, we will show that a strengthening of Theorem 1 can be provided in the case of single-trigger causal inequalities.

The canonical instrument. We now show that the ICO bound for single-trigger correlations can be achieved by the canonical choice of labelled projective instruments $\mathcal{M}_{a_i|x_i}^{(i)*}$ provided in the main text.

In general, the calculation of the ICO bound involves a double optimization, over the all possible local instruments and over all possible ICO processes. As an intermediate step, it is useful to consider the simpler problem where the local instruments are fixed and the optimization runs over the set of ICO processes. Using Eqs. (1) and (2), the correlation achieved by an ICO process \mathcal{S} can be written as $\mathcal{I} = \mathcal{S}(\mathcal{M}_{\mathcal{I}})$, with $\mathcal{M}_{\mathcal{I}} = \sum_{\vec{a}, \vec{x}} \alpha_{\vec{a}, \vec{x}} \mathcal{M}_{a_1|x_1}^{(1)} \otimes \dots \otimes \mathcal{M}_{a_N|x_N}^{(N)}$. The maximum correlation achieved by arbitrary processes is given by $v(\mathcal{M}_{\mathcal{I}}) := \max_{\mathcal{S}} \mathcal{S}(\mathcal{M}_{\mathcal{I}})$. The maximization is a semidefinite program, whose solution can be equivalently computed as [34]

$$v(\mathcal{M}_{\mathcal{I}}) = \min \left\{ \eta \in \mathbb{R} \mid \exists \mathcal{C} \in \text{Aff}(\text{NoSig}) : \eta \mathcal{C} \geq \mathcal{M}_{\mathcal{I}} \right\}, \quad (11)$$

where $\text{Aff}(\text{NoSig})$ is the set of affine combinations of no-signalling channels, and, for two linear maps \mathcal{A} and \mathcal{B} with the same input and output spaces, $\mathcal{A} \geq \mathcal{B}$ means that $\mathcal{A} - \mathcal{B}$ is completely positive. When the coefficients $\alpha_{\vec{x}, \vec{a}}$ are nonnegative, the minimization can be restricted without loss of generality to the set of no-signalling channels, and the minimum is given $2^{D_{\max}(\mathcal{M}_{\mathcal{I}}|\text{NoSig})}$, where $D_{\max}(\mathcal{M}_{\mathcal{I}}|\text{NoSig})$ is the max relative entropy distance between the operator $\mathcal{M}_{\mathcal{I}}$ and the set of no-signalling channels. In short, the maximum correlation achievable with a fixed set of local instruments is given by the deviation of the map $\mathcal{M}_{\mathcal{I}}$ from the set of no-signalling channels.

We are now ready to tackle the full problem of computing the ICO value. With the above notation, the ICO value can be written as $\mathcal{I}^{\text{ICO}} = \max_{\mathcal{M}_{\mathcal{I}}} v(\mathcal{M}_{\mathcal{I}})$, where the maximization is over all maps $\mathcal{M}_{\mathcal{I}}$ generated by all possible local quantum instruments with inputs and outputs of arbitrary dimensions. By Theorem 1, the maximization can be restricted without loss of generality to maps $\mathcal{M}_{\mathcal{I}}$ generated by labelled projective instruments. In Appendix G, we show that, for single-trigger correlations and labelled projective instruments, the map $\mathcal{M}_{\mathcal{I}}$ can be decomposed into a convex combination of maps

associated to instruments of an even simpler form, as stated by the following theorem:

Theorem 2 *Every map $\mathcal{M}_{\mathcal{I}}$ associated to a given single trigger correlation and a given set of labelled projective instruments can be decomposed into a convex combination $\mathcal{M}_{\mathcal{I}} = \sum_j p_j \mathcal{M}_{\mathcal{I},j}$, where (p_j) is a probability distribution and, for every j , $\mathcal{M}_{\mathcal{I},j}$ is the map associated to local instruments of the form*

$$\mathcal{M}_{a_i|x_i}^{(i,j)}(\rho) = \begin{cases} P_{a_i,\xi_i}^{(i)} \rho P_{a_i,\xi_i}^{(i)} \otimes |\xi_i\rangle\langle\xi_i| & x_i = \xi_i, \\ \frac{1}{m_i} U_{x_i}^{(i,j)} \rho U_{x_i}^{(i,j)\dagger} \otimes |x_i\rangle\langle x_i| & x_i \neq \xi_i. \end{cases} \quad (12)$$

where $(P_{a_i|x_i}^{(i)})_{a_i=0}^{m_i-1}$ are the projectors appearing in the original labelled projective instrument, and $U_{x_i}^{(i,j)}$ is a unitary operator for every i, j , and x_i .

Operationally, the instruments appearing in Theorem 2 describe experiments where the i -th player performs a labelled projective instrument when their setting is the trigger, and, for all the other settings, the player generates a uniformly random outcome, while performing a unitary gate on the system entering their laboratory. Since v is a convex function of the map $\mathcal{M}_{\mathcal{I}}$, Theorem 2 guarantees that its maximization can be restricted without loss of generality to maps generated by instruments of this form. Note that the canonical instruments $(\mathcal{M}^{(i*)})_{a_i|x_i}$ are a special case of instrument of the form (12), corresponding to the case where all projectors are rank-one, and all unitary operators are equal to the identity operator.

In Appendix H we show that every value of the function $v(\mathcal{M}_{\mathcal{I}})$ achievable by instruments of the form (12) can be

achieved by the canonical instrument. Intuitively, the argument is as follows: first, Theorem 1 guarantees that all the instruments associated to the same party have projectors of the same rank. For single-trigger correlations, Theorem 2 ensures that one can effectively consider only one projective instrument per party. But then, a single projective instrument with projectors of equal rank can be reduced to a rank-one projective instrument by appending an additional quantum system in the party's local laboratory, without performing any operation on it. Without loss of generality, the additional system can be incorporated in the definition of the process \mathcal{S} connecting the local laboratories. Similarly, arbitrary unitary operations depending on the parties' settings can be obtained from the identity operation, by appending to the process \mathcal{S} a controlled unitary operation, controlled by the state of the auxiliary systems that carry the values of the parties' settings (see Appendix H for the details). The above argument implies the bound $v(\mathcal{M}_{\mathcal{I}}^*) \geq v(\mathcal{M}_{\mathcal{I}})$ for every map $\mathcal{M}_{\mathcal{I}}$ generated by local instruments, and therefore $\mathcal{I}_{\text{single-trigger}}^{\text{ICO}} = v(\mathcal{M}_{\mathcal{I}}^*)$.

Using Eq. (11) we then obtain an explicit SDP expression for the ICO value. When the coefficients $\alpha_{\vec{a},\vec{x}}$ are nonnegative, this expression reduces to the max relative entropy distance, in agreement with Eq. (3) in the main text.

DATA AVAILABILITY

The authors declare that the data supporting the findings of this study are available within the paper and in the supplementary information files.

* giulio@cs.hku.hk

-
- [1] G. Chiribella, G. D'Ariano, P. Perinotti, and B. Valiron, arXiv preprint arXiv:0912.0195 (2009).
 - [2] O. Oreshkov, F. Costa, and Č. Brukner, *Nature Communications* **3**, 1 (2012).
 - [3] G. Chiribella, G. M. D'Ariano, P. Perinotti, and B. Valiron, *Physical Review A* **88**, 022318 (2013).
 - [4] Č. Brukner, *Nature Physics* **10**, 259 (2014).
 - [5] G. Chiribella, *Physical Review A* **86**, 040301 (2012).
 - [6] M. Araújo, F. Costa, and Č. Brukner, *Physical Review Letters* **113**, 250402 (2014).
 - [7] P. A. Guérin, A. Feix, M. Araújo, and Č. Brukner, *Physical Review Letters* **117**, 100502 (2016).
 - [8] D. Ebler, S. Salek, and G. Chiribella, *Physical Review Letters* **120**, 120502 (2018).
 - [9] X. Zhao, Y. Yang, and G. Chiribella, *Physical Review Letters* **124**, 190503 (2020).
 - [10] D. Felce and V. Vedral, *Physical Review Letters* **125**, 070603 (2020).
 - [11] L. Hardy, *Journal of Physics A: Mathematical and Theoretical* **40**, 3081 (2007).
 - [12] O. Oreshkov and C. Giarmatzi, *New Journal of Physics* **18**, 093020 (2016).
 - [13] C. Branciard, M. Araújo, A. Feix, F. Costa, and Č. Brukner, *New Journal of Physics* **18**, 013008 (2015).
 - [14] Ä. Baumeler, A. Feix, and S. Wolf, *Physical Review A* **90**, 042106 (2014).
 - [15] A. Feix, M. Araújo, and Č. Brukner, *New Journal of Physics* **18**, 083040 (2016).
 - [16] A. A. Abbott, C. Giarmatzi, F. Costa, and C. Branciard, *Physical Review A* **94**, 032131 (2016).
 - [17] J. Wechs, C. Branciard, and O. Oreshkov, *Nature Communications* **14**, 1471 (2023).
 - [18] B. S. Cirel'son, *Letters in Mathematical Physics* **4**, 93 (1980).
 - [19] S. Popescu and D. Rohrlich, *Foundations of Physics* **24**, 379 (1994).
 - [20] W. Van Dam, *Nonlocality and communication complexity*, Ph.D. thesis, University of Oxford (1999).
 - [21] G. Brassard, H. Buhrman, N. Linden, A. A. Méthot, A. Tapp, and F. Unger, *Physical Review Letters* **96**, 250401 (2006).

- [22] M. Navascués, S. Pironio, and A. Acín, *Physical Review Letters* **98**, 010401 (2007).
- [23] N. Brunner and P. Skrzypczyk, *Physical Review Letters* **102**, 160403 (2009).
- [24] N. Linden, S. Popescu, A. J. Short, and A. Winter, *Physical Review Letters* **99**, 180502 (2007).
- [25] M. Pawłowski, T. Paterek, D. Kaszlikowski, V. Scarani, A. Winter, and M. Żukowski, *Nature* **461**, 1101 (2009).
- [26] M. Navascués and H. Wunderlich, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* **466**, 881 (2010).
- [27] T. Fritz, A. B. Sainz, R. Augusiak, J. B. Brask, R. Chaves, A. Leverrier, and A. Acín, *Nature Communications* **4**, 2263 (2013).
- [28] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, *Physical Review Letters* **23**, 880 (1969).
- [29] Č. Brukner, *New Journal of Physics* **17**, 083034 (2015).
- [30] M. L. Almeida, J.-D. Bancal, N. Brunner, A. Acín, N. Gisin, and S. Pironio, *Physical Review Letters* **104**, 230404 (2010).
- [31] R. Kunjwal and O. Oreshkov, *arXiv preprint arXiv:2307.02565* (2023).
- [32] G. Chiribella and Z. Liu, *Communications Physics* **5**, 1 (2022).
- [33] M. Araújo, C. Branciard, F. Costa, A. Feix, C. Giarmatzi, and Č. Brukner, *New Journal of Physics* **17**, 102001 (2015).
- [34] G. Chiribella and D. Ebler, *New Journal of Physics* **18**, 093053 (2016).
- [35] A. Bisio and P. Perinotti, *Proceedings of the Royal Society A* **475**, 20180706 (2019).
- [36] S. S. Bhattacharya and M. Banik, *arXiv preprint arXiv:1509.02721* (2015).
- [37] N. Brunner, D. Cavalcanti, S. Pironio, V. Scarani, and S. Wehner, *Reviews of Modern Physics* **86**, 419 (2014).
- [38] J. Bavaresco, M. Araújo, Č. Brukner, and M. T. Quintino, *Quantum* **3**, 176 (2019).
- [39] M. Navascués, S. Pironio, and A. Acín, *New Journal of Physics* **10**, 073013 (2008).
- [40] S. Gogioso and N. Pinzani, *arXiv preprint arXiv:2303.09017* (2023).
- [41] T. van der Lugt, J. Barrett, and G. Chiribella, *Nature Communications* **14**, 5811 (2023).
- [42] D. Mayers and A. Yao, in *Proceedings 39th Annual Symposium on Foundations of Computer Science (Cat. No. 98CB36280)* (IEEE, 1998) pp. 503–509.
- [43] D. Mayers and A. Yao, *Quantum Info. Comput.* **4**, 273–286 (2004).
- [44] A. Acin, N. Gisin, and L. Masanes, *Physical Review Letters* **97**, 120405 (2006).
- [45] R. Colbeck, *Quantum and relativistic protocols for secure multi-party computation*, Ph.D. thesis, University of Cambridge (2006).
- [46] S. Pironio, A. Acín, S. Massar, A. B. de La Giroday, D. N. Matsukevich, P. Maunz, S. Olmschenk, D. Hayes, L. Luo, T. A. Manning, *et al.*, *Nature* **464**, 1021 (2010).
- [47] H. Buhrman, R. Cleve, S. Massar, and R. De Wolf, *Reviews of Modern Physics* **82**, 665 (2010).
- [48] B. W. Reichardt, F. Unger, and U. Vazirani, *Nature* **496**, 456 (2013).
- [49] E. B. Davies and J. T. Lewis, *Communications in Mathematical Physics* **17**, 239 (1970).
- [50] M. Ozawa, *Journal of Mathematical Physics* **25**, 79 (1984).
- [51] T. Heinosaari and M. Ziman, *The mathematical language of quantum theory: from uncertainty to entanglement* (Cambridge University Press, 2011).
- [52] A. Jamiołkowski, *Reports on Mathematical Physics* **3**, 275 (1972).
- [53] M.-D. Choi, *Linear algebra and its applications* **10**, 285 (1975).
- [54] G. Chiribella, G. M. D’Ariano, and P. Perinotti, *EPL (Europhysics Letters)* **83**, 30004 (2008).
- [55] G. Chiribella, G. M. D’Ariano, and P. Perinotti, *Physical Review A* **80**, 022339 (2009).
- [56] J. Watrous, *The theory of quantum information* (Cambridge university press, 2018).

ACKNOWLEDGMENTS

The authors thank Č. Brukner and M. Araújo for helpful comments. This work has been supported by the Hong Kong Research Grant Council through the Senior Research Fellowship Scheme SRFS2021-7S02 and through the Research Impact Grant R7035-21F. The work was also supported by the John Templeton Foundation through grant 62312, The Quantum Information Structure of Spacetime (qiss.fr). The opinions expressed in this publication are those of the authors and do not necessarily reflect the views of the John Templeton Foundation. Research at the Perimeter Institute is supported by the Government of Canada through the Department of Innovation, Science and Economic Development Canada and by the Province of Ontario through the Ministry of Research, Innovation and Science.

AUTHOR CONTRIBUTIONS

ZL introduced the notion of single-trigger correlations, derived the bounds on quantum ICO correlations, and showed the achievability of the algebraic maximum by classical ICOTD processes. GC supervised the research and proved the reduction to labelled projective measurements (Theorem 1). GC and ZL wrote the manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

Appendix A: SDP relaxation of the problem of computing the ICO bound

In this section, we introduce Choi-Jamiołkowski isomorphism which allows us to represent quantum processes and local quantum operations as operators. Then we explicitly present the SDPs for computing the ICO

bound of single-trigger correlations and for computing upper bounds for general correlations.

Choi-Jamiołkowski isomorphism

Through Choi-Jamiołkowski isomorphism [52, 53], a linear map \mathcal{K} on operators can be conveniently represented as a Choi operator, defined as follows:

$$\text{Choi}(\mathcal{K}) := \sum_{i,j} |i\rangle\langle j| \otimes \mathcal{K}(|i\rangle\langle j|). \quad (\text{A1})$$

If \mathcal{K} is a completely positive map with Kraus operators $\{K_i\}$, it holds that

$$\text{Choi}(\mathcal{K}) = \sum_i |K_i\rangle\rangle\langle\langle K_i|, \quad (\text{A2})$$

where we use the double-ket notation defined as $|M\rangle\rangle = \sum_j |j\rangle \otimes M|j\rangle$.

Quantum supermaps and process matrices

In an experiment without assumption of global causal structure, the local operations are described by quantum instruments, that is, collections of linear maps $(\mathcal{M}_j)_j$ labelled by possible measurement outcomes j . Each linear map \mathcal{M}_j , called a quantum operation, transforms density matrices on an input system into (generally sub-normalized) density matrices on an output system, and is required to be completely positive. In the following, we will denote the i -th party's instruments (conditional on the setting x_i) by $(\mathcal{M}_{a_i|x_i}^{(i)})_{a_i}$, each map $\mathcal{M}_{a_i|x_i}^{(i)}$ transforming quantum states of an input system $A_i^{(i)}$ into quantum states of an output system $A_o^{(i)}$. The most general rule for assigning probabilities to outcomes is provided by a process \mathcal{S} that transforms all local quantum operations $(\mathcal{M}^{(1)}, \dots, \mathcal{M}^{(N)})$ into probabilities $\mathcal{S}(\mathcal{M}^{(1)}, \dots, \mathcal{M}^{(N)})$. In this way, the joint conditional probability distribution of outcomes is given by

$$p(\vec{a} | \vec{x}) = \mathcal{S}(\mathcal{M}_{a_1|x_1}^{(1)}, \dots, \mathcal{M}_{a_N|x_N}^{(N)}). \quad (\text{A3})$$

Mathematically, a process is a higher-order map transforming a list of channels to the unit probability, which is a special case of quantum supermaps, the physically admissible operations transforming quantum channels [54, 55]. Given a supermap \mathcal{S} , we can induce a map $\widehat{\mathcal{S}}$ which transforms the Choi operators of channels,

$$\widehat{\mathcal{S}}: \text{Choi}(\mathcal{C}) \mapsto \text{Choi}(\mathcal{S}(\mathcal{C})), \quad (\text{A4})$$

and we define the Choi operator of \mathcal{S} to be the Choi operator of $\widehat{\mathcal{S}}$, i.e. $\text{Choi}(\mathcal{S}) = \text{Choi}(\widehat{\mathcal{S}})$. The condition

that \mathcal{S} is a CP map is equivalent to the condition that $\text{Choi}(\mathcal{S})$ is a positive operator, i.e. $\text{Choi}(\mathcal{S}) \geq 0$.

The constraints on the processes \mathcal{S} that are in principle compatible with the validity of quantum mechanics in all local laboratories can be completely characterized in terms of the Choi operator S of the map \mathcal{S} , called the *process matrix* [2], satisfying the conditions (i) $S \geq 0$, (ii) $S \in \text{DualAff}(\text{Choi}(\text{NoSig}))$ where $\text{Choi}(\text{NoSig})$ is the set of Choi operators of no-signalling channels and DualAff denotes the dual affine space. See Refs. [2, 33] for an explicit characterization of process matrices. The joint conditional probability in Eq. (A3) is then computed by

$$p(\vec{a} | \vec{x}) = \text{Tr} \left[S^T \cdot \bigotimes_{i=1}^N M_{a_i|x_i}^{(i)} \right]. \quad (\text{A5})$$

where $M_{a_i|x_i}^{(i)}$ are Choi operators of the corresponding instruments $\mathcal{M}_{a_i|x_i}^{(i)}$ and S^T represents transposition of the process matrix in the computational basis.

SDP duality

In Methods, with fixed instruments $(\mathcal{M}_{a_i|x_i}^{(i)})_{a_i}$, we associate every correlation $\mathcal{I} = \sum_{\vec{a}, \vec{x}} \alpha_{\vec{a}, \vec{x}} p(\vec{a} | \vec{x})$ with a map $\mathcal{M}_{\mathcal{I}}$. The Choi operator of $\mathcal{M}_{\mathcal{I}}$ will be referred to as the performance operator, which is given by

$$\Omega_{\mathcal{I}} := \text{Choi}(\mathcal{M}_{\mathcal{I}}) = \sum_{\vec{a}, \vec{x}} \alpha_{\vec{a}, \vec{x}} \bigotimes_{i=1}^N \text{Choi}(\mathcal{M}_{a_i|x_i}^{(i)}), \quad (\text{A6})$$

In particular, when \mathcal{I} is a single-trigger correlation with triggers $\vec{\xi}$, we associated it with a single-trigger operator, which is the performance operator $\Omega_{\mathcal{I}}^*$ with the canonical choice of instruments, and is in turn the Choi operator of the map $\mathcal{M}_{\mathcal{I}}^*$ in the main text. The single-trigger operator $\Omega_{\mathcal{I}}^*$ is given by

$$\Omega_{\mathcal{I}}^* = \sum_{\vec{a}, \vec{x}} \alpha_{\vec{a}, \vec{x}} P_{\vec{\xi}, \vec{a}, \vec{x}} \quad (\text{A7})$$

where $P_{\vec{\xi}, \vec{a}, \vec{x}}$ are projectors defined as

$$P_{\vec{\xi}, \vec{a}, \vec{x}} := \bigotimes_{i=1}^N P_{\xi_i, a_i, x_i}^{(i)} \quad (\text{A8})$$

with

$$P_{\xi_i, a_i, x_i}^{(i)} = \begin{cases} |a_i\rangle\langle a_i| \otimes |a_i\rangle\langle a_i| \otimes |\xi_i\rangle\langle \xi_i| & x_i = \xi_i, \\ |\Phi\rangle\langle \Phi| \otimes |x_i\rangle\langle x_i| & x_i \neq \xi_i, \end{cases} \quad (\text{A9})$$

where $|\Phi\rangle := \sum_j |j\rangle \otimes |j\rangle / \sqrt{m_i}$ is the canonical maximally entangled state.

With respect to a performance operator $\Omega_{\mathcal{I}}$, the optimization problem of the corresponding correlation can

be expressed through the following SDP [34]:

$$\begin{aligned} & \text{maximize} && \text{Tr}(\Omega_{\mathcal{I}} S) && \text{(A10)} \\ & \text{subject to} && S \in \text{DualAff}(\text{Choi}(\text{NoSig})) \\ & && S \geq 0. \end{aligned}$$

The corresponding dual problem is

$$\begin{aligned} & \text{minimize} && \eta && \text{(A11)} \\ & \text{subject to} && \eta C \geq \Omega_{\mathcal{I}} \\ & && C \in \text{Aff}(\text{Choi}(\text{NoSig})), \end{aligned}$$

where $\text{Aff}(\text{Choi}(\text{NoSig}))$ denotes the set of affine combinations of the Choi operators of no-signalling channels.

By the property of duality, every dual feasible point (η, C) provides an upper bound η of the SDP pair (A10, A11). Furthermore, strong duality holds because both primal optimal value and dual optimal value are finite and the primal feasible set contains a strictly positive operator (the process matrix proportional to the identity) [34, 56]. In particular, the dual problem (A11), with the performance operator being a single-trigger operator $\Omega_{\mathcal{I}}^*$, is the SDP corresponding to the ICO bound of the corresponding single-trigger correlation \mathcal{I} . When the coefficients of a single-trigger correlation \mathcal{I} are non-negative, the corresponding single-trigger operator $\Omega_{\mathcal{I}}^*$ is positive and thus the dual problem (A11) is equivalent to $\min_{C \in \text{Choi}(\text{NoSig})} \{\eta \in \mathbb{R} \mid \Omega_{\mathcal{I}}^* \leq \eta C\} = 2^{D_{\max}(\Omega_{\mathcal{I}}^* \parallel \text{Choi}(\text{NoSig}))}$, which is the explicit form of the formula $\mathcal{I}^{\text{ICO}} = 2^{D_{\max}(\mathcal{M}_{\mathcal{I}}^* \parallel \text{NoSig})}$ in the main text.

We will denote the value of the SDP pair (A10, A11) as a convex function $\eta(\Omega_{\mathcal{I}})$ of the performance operator $\Omega_{\mathcal{I}}$. The function η is equivalent to (via Choi-Jamiołkowski isomorphism) the function v in the main text defined on the set of maps $\mathcal{M}_{\mathcal{I}}$.

SDP relaxation in an explicit form

Now we present the SDP relaxation of the problem of computing the ICO bound in an explicit form. For given triggers $\vec{\xi}$, define the following operator for every setting vector \vec{x} and outcome vector \vec{a} :

$$Q_{\vec{\xi}, \vec{a}, \vec{x}} := \bigotimes_{i=1}^N Q_{\xi_i, a_i, x_i} \quad \text{(A12)}$$

with

$$Q_{\xi_i, a_i, x_i}^{(i)} = \begin{cases} |a_i\rangle\langle a_i| \otimes |a_i\rangle\langle a_i| \otimes |\xi_i\rangle\langle \xi_i| & x_i = \xi_i, \\ \frac{1}{m_i} |\Phi\rangle\langle \Phi| \otimes |x_i\rangle\langle x_i| & x_i \neq \xi_i. \end{cases} \quad \text{(A13)}$$

By the definition of single-trigger operator, a single-trigger operator $\Omega_{\vec{\xi}}$ with triggers $\vec{\xi}$ is a linear combination of the projectors $\left\{ P_{\vec{\xi}, \vec{a}, \vec{x}} \right\}_{\vec{a}, \vec{x}}$, which is in turn equivalent

to the following constraint:

$$\Omega_{\vec{\xi}} = \sum_{\vec{a}, \vec{x}} \text{Tr}[\Omega_{\vec{\xi}} P_{\vec{\xi}, \vec{a}, \vec{x}}] P_{\vec{\xi}, \vec{a}, \vec{x}}. \quad \text{(A14)}$$

and the coefficients of the corresponding single-trigger correlation is given by

$$\alpha_{\vec{a}, \vec{x}}^{\vec{\xi}} = \text{Tr}[\Omega_{\vec{\xi}} Q_{\vec{\xi}, \vec{a}, \vec{x}}]. \quad \text{(A15)}$$

Combining Eq. (A14), Eq. (A15) and the dual SDP (A11) for single-trigger correlations, the optimization over the decomposition of a general correlation $\mathcal{I} = \sum_{\vec{x}, \vec{a}} \alpha_{\vec{x}, \vec{a}} p(\vec{a} \mid \vec{x})$ into single-trigger correlations can be formulated into the following SDP:

$$\begin{aligned} & \text{minimize} && \frac{\sum_{\vec{\xi}} \text{Tr}(C_{\vec{\xi}})}{\prod_{i=1}^N m_i} && \text{(A16)} \\ & \text{subject to} && \forall \vec{\xi} \quad \Omega_{\vec{\xi}} = \sum_{\vec{a}, \vec{x}} \text{Tr}[\Omega_{\vec{\xi}} P_{\vec{\xi}, \vec{a}, \vec{x}}] P_{\vec{\xi}, \vec{a}, \vec{x}}, \\ & && \forall \vec{a} \forall \vec{x} \quad \alpha_{\vec{a}, \vec{x}} = \sum_{\vec{\xi}} \text{Tr}[\Omega_{\vec{\xi}} Q_{\vec{\xi}, \vec{a}, \vec{x}}], \\ & && \forall \vec{\xi} \quad C_{\vec{\xi}} \in \text{Span}(\text{Choi}(\text{NoSig})), \\ & && \forall \vec{\xi} \quad C_{\vec{\xi}} \geq \Omega_{\vec{\xi}}. \end{aligned}$$

Appendix B: Tilted OCB correlations

To show that the value $(1 + \alpha + \sqrt{1 + \alpha^2})/2$ is maximal for a tilted OCB correlation, we decompose the tilted OCB correlation into a random mixture of two single-trigger correlations, i.e.

$$\begin{aligned} & P(a_1 = b \mid c = 0) + \alpha P(a_2 = x_1 \mid c = 1) \\ & = \sum_{z \in \{0,1\}} \frac{1}{2} \left(P(a_1 = b \mid x_1 = z, c = 0) \right. \\ & \quad \left. + \alpha P(a_2 = x_1 \mid b = z, c = 1) \right). \quad \text{(B1)} \end{aligned}$$

The proof is then complete with the following lemma, which indicates that both correlations in the decomposition is upper bounded by $(1 + \alpha + \sqrt{1 + \alpha^2})/2$.

Lemma 1 For any $x_*, b_* \in \{0,1\}$ and $\alpha \in \mathbb{R}$, the ICO bound of the following single-trigger correlation

$$\begin{aligned} & P(a_1 = b \mid x_1 = x_*, c = 0) \\ & + \alpha P(a_2 = x_1 \mid b = b_*, c = 1) \quad \text{(B2)} \end{aligned}$$

is $(1 + \alpha + \sqrt{1 + \alpha^2})/2$.

Proof. To derive an upper bound of the single-trigger correlation, we claim that the following positive operator C on a 7-qubit Hilbert space $(\mathcal{H}_1 \otimes \mathcal{H}_{2,3}) \otimes (\mathcal{H}_4 \otimes \mathcal{H}_{5,6,7})$ is

equal to $((1 + \alpha) + \sqrt{1 + \alpha^2})/2$ times the Choi operator of a no-signalling channel

$$C := \Omega + \sum_{k,l} |\Psi_{k,l}\rangle\langle\Psi_{k,l}|, \quad (\text{B3})$$

where Ω is the single-trigger operator of the correlation $P(a_1 = b \mid x_1 = x_*, c = 0) + \alpha P(a_2 = x_1 \mid b = b_*, c = 1)$

$$\begin{aligned} \Omega := & \frac{1}{2} |00\rangle\langle 00| \otimes |x_*\rangle\langle x_*| \otimes |I\rangle\langle I| \otimes |0\rangle\langle 0| \otimes |0\rangle\langle 0| \\ & + \frac{1}{2} |11\rangle\langle 11| \otimes |x_*\rangle\langle x_*| \otimes |I\rangle\langle I| \otimes |1\rangle\langle 1| \otimes |0\rangle\langle 0| \\ & + \frac{\alpha}{2} |I\rangle\langle I| \otimes |0\rangle\langle 0| \otimes |00\rangle\langle 00| \otimes |b_*\rangle\langle b_*| \otimes |1\rangle\langle 1| \\ & + \frac{\alpha}{2} |I\rangle\langle I| \otimes |1\rangle\langle 1| \otimes |11\rangle\langle 11| \otimes |b_*\rangle\langle b_*| \otimes |1\rangle\langle 1| \end{aligned} \quad (\text{B4})$$

and

$$|\Psi_{k,l}\rangle = \sum_{i,j} c_{i,j,k,l} |i, i\rangle \otimes |k\rangle \otimes |j, j\rangle \otimes |l, 0\rangle. \quad (\text{B5})$$

with real coefficients $c_{i,j,k,l}$. The conditions that C is proportional to the Choi operator of a no-signalling channel are

$$\text{Tr}_{2,3} C = \frac{I_1}{2} \otimes \text{Tr}_{1,2,3} C, \quad \text{Tr}_{5,6,7} C = \frac{I_4}{2} \otimes \text{Tr}_{4,5,6,7} C, \quad (\text{B6})$$

which are equivalent to the following equations:

$$\forall j, l \quad \sum_k c_{1,j,k,l}^2 - c_{0,j,k,l}^2 = \frac{(-1)^l}{2}, \quad (\text{B7})$$

$$\forall i, k \quad \sum_l c_{i,1,k,l}^2 - c_{i,0,k,l}^2 = \frac{(-1)^k \alpha}{2}, \quad (\text{B8})$$

$$\forall l \quad \sum_k c_{1,0,k,l} c_{1,1,k,l} - c_{0,0,k,l} c_{0,1,k,l} = \frac{(-1)^l}{2}, \quad (\text{B9})$$

$$\forall k \quad \sum_l c_{0,1,k,l} c_{1,1,k,l} - c_{0,0,k,l} c_{1,0,k,l} = \frac{(-1)^k \alpha}{2}. \quad (\text{B10})$$

We find the following solution of Eq. (B7)-(B10)

$$c_{i,j,k,l} = \frac{(1 + \alpha^2)^{1/4}}{4} \left(1 - \frac{(-1)^{i+l}}{\sqrt{1 + \alpha^2}} - \frac{(-1)^{j+k} \alpha}{\sqrt{1 + \alpha^2}} \right), \quad (\text{B11})$$

With which $\text{Tr}(C) = \text{Tr}(\Omega) + \sum_{i,j,k,l} |c_{i,j,k,l}|^2 = 2(1 + \alpha) + 2\sqrt{1 + \alpha^2}$, which is $((1 + \alpha) + \sqrt{1 + \alpha^2})/2$ times the trace of the Choi operator of a no-signalling channel. Therefore, $((1 + \alpha) + \sqrt{1 + \alpha^2})/2$ is an upper bound of $\eta(\Omega)$ and thus an upper bound of the single-trigger correlation.

On the other hand, the bound can be achieved with the following process matrix which was introduced in Ref.

[36].

$$\begin{aligned} S_{\text{OCB}} = & \frac{1}{4} I^{\otimes 4} + \frac{\alpha}{4\sqrt{1 + \alpha^2}} I \otimes Z \otimes Z \otimes I \\ & + \frac{1}{4\sqrt{1 + \alpha^2}} Z \otimes I \otimes X \otimes Z. \end{aligned} \quad (\text{B12})$$

where X and Z are Pauli- X and Pauli- Z matrices, respectively. To attain the bound, Alice uses the instruments (in Choi representation)

$$M_{a_1|x_1} = |a_1\rangle\langle a_1| \otimes |x_1\rangle\langle x_1|, \quad (\text{B13})$$

and Bob uses the instruments (in Choi representation)

$$N_{a_2|b,0} = |X_{a_2}\rangle\langle X_{a_2}| \otimes |a_2 \oplus b\rangle\langle a_2 \oplus b|, \quad (\text{B14})$$

$$N_{a_2|b,1} = |a_2\rangle\langle a_2| \otimes \rho, \quad (\text{B15})$$

where ρ is an arbitrary density operator on a qubit and $|X_{a_2}\rangle = (|0\rangle + (-1)^{a_2}|1\rangle)/\sqrt{2}$ are the eigenstates of Pauli- X matrix. The joint conditional probability distribution between Alice's and Bob's outcomes is computed to be

$$p(a_1, a_2 \mid x_1, b, 0) = \frac{1}{4} + \frac{(-1)^{a_1+b}}{4\sqrt{1 + \alpha^2}}, \quad (\text{B16})$$

$$p(a_1, a_2 \mid x_1, b, 1) = \frac{1}{4} + \frac{(-1)^{a_2+x_1} \alpha}{4\sqrt{1 + \alpha^2}}. \quad (\text{B17})$$

It follows that the tilted OCB correlation achieves the bound:

$$\begin{aligned} & P(a_1 = b \mid x_1 = x_*, c = 0) + \alpha P(a_2 = x_1 \mid b = b_*, c = 1) \\ & = \frac{1}{2} \sum_{a_1, a_2, x_1, b} \delta_{a_1, b} \delta_{x_1, x_*} p(a_1, a_2 \mid x_1, b, 0) \\ & \quad + \frac{\alpha}{2} \sum_{a_1, a_2, x_1, b} \delta_{a_2, x_1} \delta_{b, b_*} p(a_1, a_2 \mid x_1, b, 1) \\ & = \frac{1}{2} \left(1 + \frac{1}{\sqrt{1 + \alpha^2}} \right) + \frac{\alpha}{2} \left(1 + \frac{\alpha}{\sqrt{1 + \alpha^2}} \right) \\ & = \frac{1}{2} \left(1 + \alpha + \sqrt{1 + \alpha^2} \right). \end{aligned} \quad (\text{B18})$$

■

Appendix C: Tilted LGYNI correlations

We now introduce a tilted version of the LGYNI game, corresponding to the correlation

$$\mathcal{I}_{\text{LGYNI}, \alpha} = \alpha P_{\text{succ}}^{11} + (1 - \alpha) \frac{P_{\text{succ}}^{01} + P_{\text{succ}}^{10}}{2}, \quad (\text{C1})$$

where α ranges between 0 and 1. Clearly, the correlation $\mathcal{I}_{\text{LGYNI}, \alpha}$ is single-trigger, and for $\alpha = 1/3$ its maximization is equivalent to the maximization of the probability of success in the original LGYNI game.




FIG. 2. **Visualizing correlations and probability distributions.** (a) Causal bound, ICO bound and algebraic maximal of tilted LGYNI. (b) Geometry of probability distributions on a 2-dimensional plane. The green area represents the projection of causal probability distributions; the blue area represents the projection of probability distributions exhibiting ICO. The remaining area (violet area) represents the non-causal probability distributions beyond the ICO ones.

For the tilted LGYNI game, one has the causal inequality

$$\mathcal{I}_{\text{LGYNI},\alpha} \leq \max \left\{ 1 - \alpha, \frac{1 + \alpha}{2} \right\}. \quad (\text{C2})$$

The proof is as follows: let p_A and p_B be the marginal probability distributions of Alice's and Bob's outcomes, defined as

$$p_A(a_1 | x_1, x_2) := \sum_{a_2} p(a_1, a_2 | x_1, x_2), \quad (\text{C3})$$

$$p_B(a_2 | x_1, x_2) := \sum_{a_1} p(a_1, a_2 | x_1, x_2) \quad (\text{C4})$$

respectively. If Alice's operation causally precedes Bob's, then

$$p(1, 1 | 1, 1) \leq p_A(1 | 1, 1) = p_A(1 | 1, 0) \quad (\text{C5})$$

In this case, tilted LGYNI is upper bounded as follows

$$\begin{aligned} \mathcal{I}_{\text{LGYNI},\alpha} &= \alpha P_{\text{succ}}^{11} + (1 - \alpha) \frac{P_{\text{succ}}^{01} + P_{\text{succ}}^{10}}{2} \\ &= \alpha p(1, 1 | 1, 1) + \frac{1 - \alpha}{2} p_A(0 | 1, 0) + \frac{1 - \alpha}{2} p_B(0 | 0, 1) \\ &\leq \alpha p_A(1 | 1, 0) + \frac{1 - \alpha}{2} p_A(0 | 1, 0) + \frac{1 - \alpha}{2} p_B(0 | 0, 1) \\ &= \alpha(1 - p_A(0 | 1, 0)) + \frac{1 - \alpha}{2} p_A(0 | 1, 0) \\ &\quad + \frac{1 - \alpha}{2} p_B(0 | 0, 1) \\ &\leq \max \left\{ \alpha, \frac{1 - \alpha}{2} \right\} + \frac{1 - \alpha}{2} \\ &= \max \left\{ \frac{1 + \alpha}{2}, 1 - \alpha \right\}, \end{aligned} \quad (\text{C6})$$

where the first inequality is due to (C5), and the second inequality is due to the monotonicity as a linear function of $p_A(0 | 1, 0)$ and the probability constraint $p_B \leq 1$. Vice versa, tilted LGYNI is upper bounded by the same value when Bob's operation precedes Alice's. Since a random mixture can not increase the bound, the causal bound of tilted LGYNI is no greater than (C6). The value (C6) can be saturated by the following causal process: Alice sends her classical bit x to Bob; she produces outcome 0 if $\alpha < \frac{1 - \alpha}{2}$ and produces outcome 1 otherwise; Bob produces his outcome to be x .

Quantum processes with ICO can violate the inequality (C2) for suitable values of the parameter α . Explicitly,

the ICO bound for the tilted LGYNI game can be computed as an SDP. The result is shown in Figure 2 for all possible values of α in the interval $[0, 1]$. Interestingly, it appears that the tilted LGYNI causal inequality cannot be violated by any quantum ICO process for $\alpha \leq 0.188$. (In the last part of this section, we provide a rigorous proof of this fact for $\alpha \leq \frac{4-\sqrt{5}}{11} \approx 0.16$.) To the best of our knowledge, tilted LGYNI with $\alpha \leq 0.188$ is the first example of a causal inequality with no quantum violation in the ICO framework.

At first sight, the inequality (C2) for $\alpha \leq 0.188$ may appear as a causal analogue of Bell inequalities with no quantum violations [27, 30]. A more in-depth analysis of the geometry of ICO correlations, however, reveals that this inequality is not tight, meaning that it does not identify a facet of the quantum set.

Let us visualize the success probabilities P_{succ}^{11} and $P_{\text{succ}}^{\text{ave}}$ in a two-dimensional plane. The values of P_{succ}^{11} and $P_{\text{succ}}^{\text{ave}}$ compatible with the causal inequality (C2) and with the ICO bound are shown in Figure 2, where they correspond to the green area and the blue area, respectively. The constraints arising from the causal inequality can be equivalently characterized by the condition

$$P_{\text{succ}}^{11} + 2P_{\text{succ}}^{\text{ave}} \leq 2, \quad (\text{C7})$$

The condition (C7) is the causal inequality associated with the canonical LGYNI game, which corresponds to a facet of the causal polytope [13].

The shape of the boundary of the quantum ICO set, instead, shows that the set of probability distributions generated by quantum ICO processes is not a polytope, *i.e.* it has infinite extreme points. Figure 2 also shows that the boundary of the quantum set is distinct from the boundary of the causal polytope for every value except for the extreme points of the causal polytope, corresponding to the values $p_{\text{succ}}^{11} = 0$ and $p_{\text{succ}}^{11} = 1$. The fact that the quantum and causal boundaries do not coincide implies that the causal inequality (C2) is not tight. An interesting open question is whether there exist examples of tight causal inequalities with no quantum ICO violation, namely whether there exists flat faces of the quantum set that coincide with faces of the causal polytope.

Now we rigorously prove that when $\alpha \in \left[0, \frac{4-\sqrt{5}}{11}\right]$, the ICO bound of tilted LGYNI is equal to its causal bound $1 - \alpha$. We claim that the follow positive operator C on a 6-qubit Hilbert space $(\mathcal{H}_1 \otimes \mathcal{H}_{2,3}) \otimes (\mathcal{H}_4 \otimes \mathcal{H}_{5,6})$ is equal to $(1 - \alpha)$ times the Choi operator of a no-signalling channel

$$C := \Omega + |\Psi_{01}\rangle\langle\Psi_{01}| + |\Psi_{10}\rangle\langle\Psi_{10}| + |\Psi_{11}\rangle\langle\Psi_{11}|, \quad (\text{C8})$$

where Ω is the single-trigger operator of tilted LGYNI

$$\begin{aligned} \Omega := & \alpha|11\rangle\langle 11| \otimes |1\rangle\langle 1| \otimes |11\rangle\langle 11| \otimes |1\rangle\langle 1| \\ & + \frac{1-\alpha}{2} |00\rangle\langle 00| \otimes |1\rangle\langle 1| \otimes |I\rangle\langle I| \otimes |0\rangle\langle 0| \\ & + \frac{1-\alpha}{2} |I\rangle\langle I| \otimes |0\rangle\langle 0| \otimes |00\rangle\langle 00| \otimes |1\rangle\langle 1|, \quad (\text{C9}) \end{aligned}$$

and

$$|\Psi_{01}\rangle := c_0|00\rangle \otimes |0\rangle \otimes |00\rangle \otimes |1\rangle + c_1|00\rangle \otimes |0\rangle \otimes |11\rangle \otimes |1\rangle \quad (\text{C10})$$

$$|\Psi_{10}\rangle := c_0|00\rangle \otimes |1\rangle \otimes |00\rangle \otimes |0\rangle + c_1|11\rangle \otimes |1\rangle \otimes |00\rangle \otimes |0\rangle \quad (\text{C11})$$

$$|\Psi_{11}\rangle := c_2|00\rangle \otimes |1\rangle \otimes |00\rangle \otimes |1\rangle + c_3|00\rangle \otimes |1\rangle \otimes |11\rangle \otimes |1\rangle + c_3|11\rangle \otimes |1\rangle \otimes |00\rangle \otimes |1\rangle, \quad (\text{C12})$$

with non-negative coefficients c_0, \dots, c_3 . The conditions that C is proportional to the Choi operator of a no-signalling channel are

$$\text{Tr}_{2,3} C = \frac{I_1}{2} \otimes \text{Tr}_{1,2,3} C, \quad \text{Tr}_{5,6} C = \frac{I_4}{2} \otimes \text{Tr}_{4,5,6} C, \quad (\text{C13})$$

which is equivalent to the following equations:

$$c_1^2 - c_0^2 = \alpha, \quad (\text{C14})$$

$$c_0^2 + c_2^2 - c_3^2 = \frac{1-\alpha}{2}, \quad (\text{C15})$$

$$c_1^2 + c_3^2 = \frac{1-\alpha}{2}. \quad (\text{C16})$$

$$c_0c_1 + c_2c_3 = \frac{1-\alpha}{2}, \quad (\text{C17})$$

We can solve c_1, c_2 and c_3 in terms of c_0 according to Eq. (C14)-(C16)

$$c_1 = \sqrt{c_0^2 + \alpha}, \quad (\text{C18})$$

$$c_2 = \sqrt{1 - 2\alpha - 2c_0^2}, \quad (\text{C19})$$

$$c_3 = \sqrt{\frac{1-3\alpha}{2} - c_0^2}. \quad (\text{C20})$$

It follows that

$$\frac{\text{Tr}(C)}{4} = \frac{\alpha + 2(1-\alpha) + 2(c_0^2 + c_1^2) + (c_2^2 + 2c_3^2)}{4} = 1 - \alpha \quad (\text{C21})$$

according to Eq. (C18)-(C20). Now it suffices to show the existence of c_0, \dots, c_3 satisfying Eq. (C14)-(C17). With the solutions in Eq. (C18)-(C20), we focus on the quantity $f(c_0) = c_0c_1 + c_2c_3 - \frac{1-\alpha}{2}$. when $\alpha \in \left[0, \frac{4-\sqrt{5}}{11}\right]$, it holds that

$$f(0) = \sqrt{\frac{(1-2\alpha)(1-3\alpha)}{2}} - \frac{1-\alpha}{2} \geq 0; \quad (\text{C22})$$

and

$$f\left(\sqrt{\frac{1-3\alpha}{2}}\right) = \sqrt{\frac{1-3\alpha}{2}} \sqrt{\frac{1-\alpha}{2}} - \frac{1-\alpha}{2} \leq 0; \quad (\text{C23})$$

By continuity of the function f , there exists $c_0 \in \left[0, \sqrt{\frac{1-3\alpha}{2}}\right]$ such that $f(c_0) = 0$ and thus Eq. (C17) is satisfied. Therefore, $(1 - \alpha)$ is an upper bound of $\eta(\Omega)$ and thus an upper bound of the tilted LGYNI. This completes the proof.

Appendix D: Classification of simplest bipartite classical processes

In this section, we introduce the characterization of quantum supermaps with indefinite time direction and present all deterministic bipartite classical processes without predefined time direction in which each party receives a classical bit and outputs a classical bit.

Let A_I (B_I) and A_O (B_O) be the Alice's (Bob's) input and output systems, respectively, both associated with Hilbert spaces of dimension d_A (d_B). A bipartite process with indefinite time direction is described by a quantum supermap on no-signalling bistochastic channels, which can be represented by an positive operator S satisfying the following constraints [32]:

$$\begin{aligned} S &\geq 0, \\ \text{Tr}(S) &= d_A d_B, \\ [(1-A_I)(1-A_O)B_I B_O]S &= 0, \\ [(1-B_I)(1-B_O)A_I A_O]S &= 0, \\ [(1-A_I)(1-A_O)(1-B_I)(1-B_O)]S &= 0, \end{aligned} \quad (\text{D1})$$

having use the notation $[_X]S := \text{Tr}_X[S] \otimes \frac{I_X}{d_X}$ for a system of dimension d_X and the shorthand notation $[\sum_X \alpha_X X]S := \sum_X \alpha_X \cdot [_X]S$.

In the simplest case of classical limit, the systems A_I , A_O , B_I and B_O are all bits. Classical processes satisfying the constraints in (D1) can be represented by positive diagonal matrices on $\mathcal{H}_{A_I} \otimes \mathcal{H}_{A_O} \otimes \mathcal{H}_{B_I} \otimes \mathcal{H}_{B_O}$

$$S = \sum_{i,j,k,l \in \{0,1\}} S_{i,j,k,l} |i\rangle\langle i| \otimes |j\rangle\langle j| \otimes |k\rangle\langle k| \otimes |l\rangle\langle l|, \quad (\text{D2})$$

satisfying

$$\begin{aligned} \sum_{i=j, k=l} S_{i,j,k,l} &= 1, \\ \sum_{i=j, k=l \oplus 1} S_{i,j,k,l} &= 1, \\ \sum_{i=j \oplus 1, k=l} S_{i,j,k,l} &= 1, \\ \sum_{i=j \oplus 1, k=l \oplus 1} S_{i,j,k,l} &= 1. \end{aligned} \quad (\text{D3})$$

It follows that the set of classical processes is a convex set with 2^8 extreme points:

$$\sum_{m=0}^3 |i_m\rangle\langle i_m| \otimes |j_m\rangle\langle j_m| \otimes |k_m\rangle\langle k_m| \otimes |l_m\rangle\langle l_m| \quad (\text{D4})$$

where

$$i_0 = j_0, k_0 = l_0, \quad (\text{D5})$$

$$i_1 = j_1, k_1 = l_1 \oplus 1, \quad (\text{D6})$$

$$i_2 = j_2 \oplus 1, k_2 = l_2, \quad (\text{D7})$$

$$i_3 = j_3 \oplus 1, k_3 = l_3 \oplus 1. \quad (\text{D8})$$

We provide a classification of the 2^8 extreme points of classical processes, corresponding to deterministic processes without pre-defined direction of time:

- No-signalling processes ($A \not\preceq B$ and $B \not\preceq A$). There are 2^4 points of this type, all equivalent to the circuit in Figure 3(a), up to interchange of input/output systems;
- Unidirectional signalling processes ($A \preceq B$ or $B \preceq A$) with fixed time direction. There are 2^5 points of this type, all equivalent to the circuit in Figure 3(b), up to interchange of input/output systems and causal orders;
- Unidirectional signalling processes ($A \preceq B$ or $B \preceq A$) with dynamically controlled time direction. There are 2^6 points of this type, all equivalent to the circuit in Figure 3(c) where the time direction of B is dynamically decided by the output of A , up to interchange of input/output systems and causal orders;
- Processes with ICOTD. There are $2^4 + 2^7$ points of this type. Up to interchange of input/output systems, 2^4 points of them can be expressed through process functions: $i_1 = o_1 \oplus o_2 \oplus \alpha$ and $i_2 = o_1 \oplus o_2 \oplus \beta$ where $\alpha, \beta \in \{0, 1\}$. The other 2^7 points, however, cannot be understood as stochastic processes.

Appendix E: Every probability distribution can be generated by a suitable classical process with ICOTD

Logical consistence of the classical process that enables perfect bidirectional signalling

According to Birkhoff-von Neumann theorem, every bistochastic classical channel can be written as a random mixture of permutations. In the case of the channel acting on a single bit, the permutation is either the identity function or the bit-flip function. That is to say, for every bistochastic instrument $q_{a_1|x_1}^{(1)}$ performed by Alice (similarly for Bob), $\sum_{a_1} q_{a_1|x_1}^{(1)}(s'_1 | s_1)$ is a random mixture of $\delta_{s'_1, s_1}$ (identity) and $\delta_{s'_1, s_1 \oplus 1}$ (bit-flip). Hence it suffices to check the cases in which the local interventions are either identity or bit-flip. Let $f^{(1)}(s'_1 | s_1) = \delta_{s'_1, s_1 \oplus h_1}$ be the deterministic probability distribution corresponding to the permutation performed by Alice and $f^{(2)}(s'_2 | s_2) = \delta_{s'_2, s_2 \oplus h_2}$ be the deterministic probability distribution corresponding to the permutation performed by Bob. For any local permutations performed by Alice and Bob (i.e., any values of h_1 and




FIG. 3. **Circuit realization of bipartite classical processes with definite causal order.** (a) no-signalling processes ($A \not\leq B$ and $B \not\leq A$); (b) unidirectional signalling processes ($A \leq B$ or $B \leq A$) with fixed time direction; (c) unidirectional signalling processes ($A \leq B$ or $B \leq A$) with dynamical time direction.

h_2), it holds that

$$\begin{aligned}
 \mathcal{S}(f^{(1)}, f^{(2)}) &= \sum_{s'_1, s'_2} f^{(1)}(s'_1 | s'_1 \oplus s'_2) f^{(2)}(s'_2 | s'_1 \oplus s'_2) \\
 &= \sum_{s'_1, s'_2} \delta_{s'_1, s'_1 \oplus s'_2 \oplus h_1} \delta_{s'_2, s'_1 \oplus s'_2 \oplus h_2} \\
 &= \sum_{s'_1, s'_2} \delta_{s'_2, h_1} \delta_{s'_1, h_2} = 1.
 \end{aligned} \tag{E1}$$

This completes the proof of logical consistence.

Generating an arbitrary multipartite probability distribution

In the N -party case, the set of joint conditional probability distributions form a convex polytope in a real vector space of dimension $\prod_{i=1}^N m_i n_i$, each of them represented as a vector with entries $p(\vec{a} | \vec{x})$ for specific values of \vec{a} and \vec{x} . The extreme points of the convex polytope are deterministic probability distributions defined by $p(\vec{a} | \vec{x}) := \prod_{i=1}^N \delta_{a_i, f_i(\vec{x})}$ where f_i are functions from settings \vec{x} to outcomes a_i , respectively, and δ is the Kronecker delta.

Let every two parties share a process described in the main text that enables perfect bidirectional signalling between them. This allows every party to broadcast his/her setting to the other parties. An arbitrary extreme point $p(\vec{a} | \vec{x}) := \prod_{i=1}^N \delta_{a_i, f_i(\vec{x})}$ of the set of probability distribution can be generated if every party i produces a deterministic outcome $a_i = f_i(\vec{x})$ according to a function f_i . In addition, if the N parties share a random variable, they can generate a random mixture of the ex-

treme points thus generating arbitrary probability distribution. Explicitly, the local classical channel performed every party i is a conditional deterministic probability distribution

$$q_{x_i}^{(i)}(s^{(i)'} | s^{(i)}) = \prod_j \delta_{s_j^{(i)'}, s_j^{(i)} \oplus x_i}, \tag{E2}$$

where $s^{(i)} = (s_1^{(i)}, \dots, s_N^{(i)})$ ($s^{(i)'} = (s_1^{(i)'}, \dots, s_N^{(i)'})$) is the input (output) of the i -th party, and each party produce their outcome with the decoder:

$$a_i = \begin{cases} f_i(s_1^{(i)'}, \dots, s_{i-1}^{(i)'}, x_i, s_{i+1}^{(i)'}, \dots, s_N^{(i)'}) & s_i^{(i)} = 0, \\ g_i(s_1^{(i)'}, \dots, s_{i-1}^{(i)'}, x_i, s_{i+1}^{(i)'}, \dots, s_N^{(i)'}) & s_i^{(i)} = 1, \end{cases} \tag{E3}$$

where the value of $s_i^{(i)}$ determines which function to compute. Define the N -partite process outside the laboratories to be

$$\begin{aligned}
 \mathcal{S}_N &\left(q_{a_1|x_1}^{(1)}, \dots, q_{a_N|x_N}^{(N)} \right) \\
 &= \sum_{\lambda} p(\lambda) \sum_{s^{(1)'}, \dots, s^{(N)'}} \prod_{i=1}^N q_{a_i|x_i}^{(i)} \left(s^{(i)'} | F_i \left(s^{(i)'} \right) \right),
 \end{aligned} \tag{E4}$$

where $F_i \left(s^{(i)'} \right)$ is the vector defined as

$$\left[F_i \left(s^{(i)'} \right) \right]_j = \begin{cases} \lambda & j = i, \\ s_j^{(i)'} \oplus s_j^{(j)'} & j \neq i. \end{cases} \tag{E5}$$

\mathcal{S}_N is logically consistent since it can be viewed as $\binom{N}{2}$ copies of the bipartite process \mathcal{S} that enables bidirectional signalling, plus a broadcasting of a classical variable λ . The overall correlation among the outcomes \vec{a} is

$p(\vec{a} | \vec{x}) = \lambda \prod_{i=1}^N \delta_{a_i, f_i(\vec{x})} + (1 - \lambda) \prod_{i=1}^N \delta_{a_i, g_i(\vec{x})}$ which is a random mixture of two extreme points. The settings, outcomes, inputs and outputs can be directly extended to any number of bits, thus any random mixture of deterministic processes can be generated in the above way.

Appendix F: Proof of Theorem 1 in Methods

Consider the realization of the instruments $\mathcal{M}_{a_i|x_i}^{(i)}$ by an isometry and a projective measurement. Let $\{K_{a_i|x_i,j}^{(i)}\}$ be a Kraus decomposition of the instrument $\mathcal{M}_{a_i|x_i}^{(i)}$ of the i -th party. For every classical label x_i , define $V_{x_i}^{(i)}$ to be the following isometry from $\mathcal{H}_{A_1^{(i)}}$ to $\mathcal{H}_L \otimes \mathcal{H}_{A_0^{(i)}} \otimes \mathcal{H}_J$

$$V_{x_i}^{(i)} := \sum_{a_i} |a_i\rangle_L \otimes \sum_j K_{a_i|x_i,j}^{(i)} \otimes |j\rangle_J, \quad (\text{F1})$$

where L is the classical register for the outcome a_i and J the classical register for the index j . Choosing $\mathcal{H}_L \simeq \mathbb{C}^{m_i}$ and $\mathcal{H}_J \simeq \mathcal{H}_{A_1^{(i)}} \otimes \mathcal{H}_{A_0^{(i)}}$, the two systems have no dependence on the setting x_i . Extending $V_{x_i}^{(i)}$ to an unitary $U_{x_i}^{(i)}$ from $\mathcal{H}_{A_1^{(i)}} \otimes \mathcal{H}_{E_1^{(i)}}$ to $\mathcal{H}_L \otimes \mathcal{H}_{A_0^{(i)}} \otimes \mathcal{H}_J \otimes \mathcal{H}_{E_0^{(i)}}$ with some auxiliary systems $E_1^{(i)}$ and $E_0^{(i)}$, the map $\mathcal{M}_{a_i|x_i}^{(i)}$ can be realized as

$$\rho \mapsto \text{Tr}_{L,J,E_0^{(i)}} \left[\Pi_{a_i} U_{x_i}^{(i)} \left(\rho \otimes |0\rangle\langle 0|_{E_1^{(i)}} \right) U_{x_i}^{(i)\dagger} \Pi_{a_i} \right], \quad (\text{F2})$$

where

$$\Pi_{a_i} = |a_i\rangle\langle a_i|_L \otimes I_{A_0^{(i)}} \otimes I_J \otimes I_{E_0^{(i)}}. \quad (\text{F3})$$

Define local processes $\mathcal{E}^{(i)}$ and $\mathcal{D}^{(i)}$ to be the channels

$$\forall \rho \quad \mathcal{E}^{(i)}(\rho) = \rho \otimes |0\rangle\langle 0|_{E_1^{(i)}}, \quad (\text{F4})$$

$$\forall \sigma \quad \mathcal{D}^{(i)}(\sigma) = \text{Tr}_{L,J,E_0^{(i)},F} \mathcal{U}^{(i)}(\sigma), \quad (\text{F5})$$

where $\mathcal{U}^{(i)}$ is the controlled unitary channel

$$\sum_{x_i} U_{x_i}^{(i)} \otimes |x_i\rangle\langle x_i|_F. \quad (\text{F6})$$

with the control system denoted by F .

We construct the projectors of the labelled projective instruments $\mathcal{M}_{a_i|x_i}^{(i)}$ to be $\left\{ U_{x_i}^{(i)\dagger} \Pi_{a_i} U_{x_i}^{(i)} \right\}_{a_i}$ (whose rank is independent of a_i and x_i), the label of $\mathcal{M}_{a_i|x_i}^{(i)}$ to be x_i , and the process \mathcal{S}' to be the composite of the local processes $\mathcal{E}^{(i)}$, $\mathcal{D}^{(i)}$ and the process \mathcal{S} , i.e.

$$\begin{aligned} \mathcal{S}' \left(\mathcal{M}_{a_1|x_1}^{(1)'} , \dots , \mathcal{M}_{a_N|x_N}^{(N)'} \right) \\ = \mathcal{S} \left(\mathcal{D}^{(1)} \circ \mathcal{M}_{a_1|x_1}^{(1)'} \circ \mathcal{E}^{(1)} , \dots , \mathcal{D}^{(N)} \circ \mathcal{M}_{a_N|x_N}^{(N)'} \circ \mathcal{E}^{(N)} \right) \end{aligned} \quad (\text{F7})$$

According to the realization in Eq. (F2), the local processes $\mathcal{E}^{(i)}$ and $\mathcal{D}^{(i)}$ transform the labelled projective instruments $\mathcal{M}_{a_i|x_i}^{(i)'} to $\mathcal{M}_{a_i|x_i}^{(i)}$. Hence $\mathcal{S}' \left(\mathcal{M}_{a_1|x_1}^{(1)'} , \dots , \mathcal{M}_{a_N|x_N}^{(N)'} \right)$ reproduces the joint conditional probability $\mathcal{S} \left(\mathcal{M}_{a_1|x_1}^{(1)} , \dots , \mathcal{M}_{a_N|x_N}^{(N)} \right)$.$

Appendix G: Proof of Theorem 2 in Methods

The key of the proof is the fact that a projective measurement followed by the measurement outcome being discarded is a uniform mixture of a collection of unitary channels. Let $\{P_j\}_{j=0}^{n-1}$ be the projectors of a projective measurement. Proposition 4.6 of Watrous' textbook [56] implies that there exist 2^n unitary gates U_i such that

$$\sum_{j=1}^n |P_j\rangle\langle P_j| = \sum_{i=1}^{2^n} \frac{1}{2^n} |U_i\rangle\langle U_i|. \quad (\text{G1})$$

Let Ω be a performance operator the single trigger correlation $\sum_{\vec{a}, \vec{x}} \alpha_{\vec{a}, \vec{x}} p(\vec{x} | \vec{a})$ corresponding to the map $\mathcal{M}_{\mathcal{I}}$ in the theorem. With triggers $\vec{\xi}$ and the labelled projective instruments $\mathcal{M}_{a_i|x_i}^{(i)}(\cdot) = P_{a_i, x_i}^{(i)} \cdot P_{a_i, x_i}^{(i)} \otimes |x_i\rangle\langle x_i|$, Ω is given by

$$\Omega := \sum_{\vec{a}, \vec{x}} \alpha_{\vec{a}, \vec{x}} \bigotimes_{i=1}^N M_{a_i|x_i}^{(i)}. \quad (\text{G2})$$

where $M_{a_i|x_i}^{(i)}$ are the Choi operators of the corresponding $\mathcal{M}_{a_i|x_i}^{(i)}$. We can replace $M_{a_i|x_i}^{(i)}$ with $\frac{1}{m_i} \sum_{k=0}^{m_i-1} M_{k|x_i}^{(i)}$ for any party i and any setting $x_i \neq \xi_i$. For example, let x_1 be a non-trigger setting of the first party, i.e. $x_1 \neq \xi_1$. By the property of single-trigger correlations, $\alpha_{\vec{x}, \vec{a}}$ does not depends on a_1 , we have

$$\begin{aligned} \sum_{a_1} \alpha_{\vec{a}, \vec{x}} M_{a_1|x_1}^{(1)} \otimes M_{a_2|x_2}^{(2)} \otimes \dots \otimes M_{a_N|x_N}^{(N)} \\ = \sum_{k=0}^{m_1-1} \sum_{a_1} \frac{\alpha_{\vec{a}, \vec{x}}}{m_1} M_{k|x_1}^{(1)} \otimes M_{a_2|x_2}^{(2)} \otimes \dots \otimes M_{a_N|x_N}^{(N)} \\ = \sum_{a_1} \alpha_{\vec{a}, \vec{x}} \left(\frac{\sum_{k=0}^{m_1-1} M_{k|x_1}^{(1)}}{m_1} \right) \otimes M_{a_2|x_2}^{(2)} \otimes \dots \otimes M_{a_N|x_N}^{(N)}. \end{aligned} \quad (\text{G3})$$

Hence replacing $M_{a_1|x_1}^{(1)}$ with $\frac{1}{m_1} \sum_{k=0}^{m_1-1} M_{k|x_1}^{(1)}$ does not change the value of the performance operator in Eq. (G2). Since local instruments are labelled projective instruments with the label being the corresponding setting, according to the property described in Eq. (G1), for every party i and every corresponding setting $x_i \neq \xi_i$, there exists unitary operators $U_{x_i}^{(i,j)}$ such that

$$\sum_{k=0}^{m_i-1} M_{k|x_i}^{(i)} = \frac{1}{2^{m_i}} \sum_{j=1}^{2^{m_i}} |U_{x_i}^{(i,j)}\rangle\langle U_{x_i}^{(i,j)}| \otimes |x_i\rangle\langle x_i|. \quad (\text{G4})$$

Iterating the above procedure for every party, Ω can be written as a uniform mixture of $(\Omega_j)_j$ where each Ω_j corresponds to the Choi operator of the map $\mathcal{M}_{\mathcal{I},j}$ defined in Theorem 2. This completes the proof.

Appendix H: Proof of ICO bound of single-trigger correlations

In Methods, the ICO bound of a correlation \mathcal{I} is written as $\mathcal{I}^{\text{ICO}} = \max_{\mathcal{M}_{\mathcal{I}}} v(\mathcal{M}_{\mathcal{I}})$. An alternative form of the formula is the maximum of the function $\eta(\Omega_{\mathcal{I}})$ (see Appendix A for the definition of η) for all choices of performance operator $\Omega_{\mathcal{I}}$ of the correlation, with no restriction on the dimension of the underlying Hilbert spaces.

Let \mathcal{I} be a single-trigger correlation with triggers $\vec{\xi}$, and let $\Omega_{\mathcal{I}}^*$ be the corresponding single-trigger operator. To show that $\eta(\Omega_{\mathcal{I}}^*)$ is an upper bound of the ICO bound, without loss of generality (according to Theorem 1), we restrict the local instruments to labelled projective ones such that the label of each instrument is the corresponding setting and the projectors of every projective measurement associated with the instruments are of equal rank. Theorem 2 implies that every performance operator of the single-trigger correlation is a random mixture of operators in the following form

$$\Omega_j := \sum_{\vec{a}, \vec{x}} \alpha_{\vec{a}, \vec{x}} \bigotimes_{i=1}^N Q_{a_i, x_i}^{(i,j)}. \quad (\text{H1})$$

with

$$Q_{a_i, x_i}^{(i,j)} = \begin{cases} |P_{a_i, x_i}^{(i)}\rangle\langle P_{a_i, x_i}^{(i)}| \otimes |\xi_i\rangle\langle \xi_i| & x_i = \xi_i, \\ \frac{1}{m_i} |U_{x_i}^{(i,j)}\rangle\langle U_{x_i}^{(i,j)}| \otimes |x_i\rangle\langle x_i| & x_i \neq \xi_i, \end{cases}$$

where $U_{x_i}^{(i,j)}$ are unitary operators.

We show that $\eta(\Omega_{\mathcal{I}}^*) \geq \eta(\Omega)$ for an arbitrary performance operator Ω of the single-trigger correlation. Since η is a convex function, it suffices to show that $\eta(\Omega_{\mathcal{I}}^*) \geq \eta(\Omega_j)$ for every j .

Given a choice of Ω_j , we show that there exists an isomorphism W_j from $\bigotimes_{i=1}^N (\mathbb{C}^{m_i} \otimes \mathbb{C}^{m_i n_i})$ to the Hilbert space on which Ω_j is defined, such that

- $W_j C W_j^\dagger$ is the Choi operator of a no-signalling channel whenever C is the Choi operator of a no-signalling channel, and
- $W_j \Omega_{\mathcal{I}}^* W_j^\dagger = \Omega_j$.

In fact, we construct a product isomorphism $W_j = \bigotimes_{i=1}^N W_j^{(i)}$. Since the projectors $\{P_{a_i, x_i}^{(i)}\}_{a_i}$ of every instrument are of equal rank, there exists a collection of unitary operators $\{V_{k, x_i}^{(i)}\}_{k=0}^{m_i-1}$ such that

$$V_{k, x_i}^{(i)} P_{a_i, x_i}^{(i)} V_{k, x_i}^{(i)\dagger} = P_{(a_i+k), x_i}^{(i)}. \quad (\text{H2})$$

Note that in this section, the indices are to be taken modulo the corresponding size. The construction of $W_j^{(i)}$ is

$$W_j^{(i)} : |a_i\rangle \otimes |(a_i+k)\rangle \otimes |x_i\rangle \mapsto |U_{x_i}^{(i,j)} V_{k, x_i}^{(i)} P_{a_i, x_i}^{(i)}\rangle \otimes |x_i\rangle. \quad (\text{H3})$$

where $U_{\xi_i}^{(i,j)}$ is defined to be identity. Immediately we have $W_j \Omega_{\mathcal{I}}^* W_j^\dagger = \Omega_j$. To check that the product isomorphism W_j preserves Choi operators of no-signalling channels, it suffices to check that partial tracing the output system on both sides of (H3) yields a unital map. The partial trace on the output systems gives the operators

$$|a_i\rangle\langle a'_i| \otimes |(a_i+k)\rangle\langle (a'_i+k')| \otimes |x_i\rangle\langle x'_i| \xrightarrow{\text{partial trace}} \delta_{x_i, x'_i} \delta_{(a_i+k), (a'_i+k')} |a_i\rangle\langle a'_i|, \quad (\text{H4})$$

and

$$|U_{x_i}^{(i,j)} V_{k, x_i}^{(i)} P_{a_i, x_i}^{(i)}\rangle\langle U_{j, x'_i}^{(i,j)} V_{k', x'_i}^{(i)} P_{a'_i, x'_i}^{(i)}| \otimes |x_i\rangle\langle x'_i| \xrightarrow{\text{partial trace}} \delta_{x_i, x'_i} \delta_{(a_i+k), (a'_i+k')} P_{a_i, x_i}^{(i)T} \overline{V_{a_i-a'_i, x_i} P_{a'_i, x_i}^{(i)}}. \quad (\text{H5})$$

We see that the map that transforms the operators $|a_i\rangle\langle a'_i|$ (corresponding to the output of (H4)) to $P_{a_i, x_i}^{(i)T} \overline{V_{a_i-a'_i, x_i} P_{a'_i, x_i}^{(i)}}$ (corresponding to the output of (H5)) is a unital map. Explicitly, $I = \sum_{a_i} |a_i\rangle\langle a_i|$ is mapped to $I = \sum_{a_i} P_{a_i, x_i}^{(i)T}$.

Now let $(\eta(\Omega_{\mathcal{I}}^*), C_0)$ be a dual optimal point of the SDP pair (A10, A11) with the performance operator being the single-trigger operator $\Omega_{\mathcal{I}}^*$. C_0 is the Choi operator of a no-signalling channel implies that $W_j C_0 W_j^\dagger$ is the Choi operator of a no-signalling channel on the corresponding Hilbert space. In addition, $\eta(\Omega_{\mathcal{I}}^*) C_0 \geq \Omega_{\mathcal{I}}^*$ implies $\eta(\Omega_{\mathcal{I}}^*) W_j C_0 W_j^\dagger \geq W_j \Omega_{\mathcal{I}}^* W_j^\dagger = \Omega_j$. Therefore, $(\eta(\Omega_{\mathcal{I}}^*), W_j C_0 W_j^\dagger)$ is a dual feasible point of the SDP pair (A10, A11) with the performance operator being Ω_j . Thus $\eta(\Omega_{\mathcal{I}}^*) \geq \eta(\Omega_j)$. This completes the proof.