

Solvability of a fourth order elliptic problem in a bounded sector, part II

Rabah Labbas, Stéphane Maingot & Alexandre Thorel

R. L., S. M. & A. T.: Université Le Havre Normandie, Normandie Univ, LMAH UR 3821, 76600 Le Havre, France.

rabah.labbas@univ-lehavre.fr, stephane.maingot@univ-lehavre.fr, alexandre.thorel@univ-lehavre.fr

Abstract

After different variables and functions changes, the generalized dispersal problem, recalled in (1) below and considered in part I, see Labbas, Maingot and Thorel [14], leads us to consider, to study and to invert the sum of linear operators (4) below in a suitable Banach space by using two strategies: namely the theory of sums of operators in Banach spaces as developed by Da Prato-Grisvard [4] and successfully improved by Dore-Venni [5].

Key Words and Phrases: Sum of linear operators, second and fourth order boundary value problem, functional calculus, bounded imaginary powers, maximal regularity

2020 Mathematics Subject Classification: 34G10, 35B65, 35C15, 35R20, 47A60.

1 Introduction and main result

This work is a natural continuation of Labbas, Maingot and Thorel [14], where we have considered, for $k > 0$, the following problem

$$\begin{cases} \Delta^2 u - k\Delta u = f & \text{in } S_{\omega,\rho} \\ u = \frac{\partial u}{\partial n} = 0 & \text{on } \Gamma_0 \cup \Gamma_{\omega,\rho} \\ u = \frac{\partial^2 u}{\partial n^2} = 0 & \text{on } \Gamma_\rho, \end{cases} \quad (1)$$

with

$$\begin{cases} S_{\omega,\rho} &= \{(x, y) = (r \cos \theta, r \sin \theta) : 0 < r < \rho \text{ and } 0 < \theta < \omega\} \\ \Gamma_0 &= (0, \rho) \times \{0\} \\ \Gamma_\omega &= \{(r \cos \omega, r \sin \omega) : 0 < r < \rho\} \\ \Gamma_\rho &= \{(\rho \cos \theta, \rho \sin \theta) : 0 < \theta < \omega\}, \end{cases}$$

for given $\rho > 0$ and $\omega \in (0, 2\pi]$.

In part I, see Labbas, Maingot and Thorel [14], to study problem (1), we have done many variables and functions changes to write it as a sum of linear operators. To this end, we have introduced the following functions for $t > 0$ and $(r, \theta) \in S_{\rho,\omega}$:

$$\begin{cases} v(r, \theta) = u(r \cos \theta, r \sin \theta) \\ G(t)(\theta) := G(t, \theta) = g(\rho e^{-t}, \theta) = f(\rho e^{-t} \cos \theta, \rho e^{-t} \sin \theta) \\ \phi(t)(\theta) := \phi(t, \theta) = \frac{v(\rho e^{-t}, \theta)}{\rho e^{-t}} \\ H(t)(\theta) := H(t, \theta) = e^{-3t} G(t)(\theta), \end{cases} \quad (2)$$

and the two abstract vector-valued functions

$$V(t) = \begin{pmatrix} e^{\nu t} \phi(t) \\ e^{\nu t} \phi''(t) \end{pmatrix}, \quad \mathcal{F}_\nu(t) = \begin{pmatrix} 0 \\ \rho^3 e^{\nu t} H(t) \end{pmatrix}, \quad \nu = 3 - \frac{2}{p} \in (1, 3) \text{ with } p > 1. \quad (3)$$

By considering the Banach space

$$X = W_0^{2,p}(0, \omega) \times L^p(0, \omega),$$

and after the changes indicated above in (2) and (3), we have rewritten problem (1) in the space $L^p(0, +\infty; X)$, see Labbas, Maingot and Thorel [14], in the following form

$$(\mathcal{L}_{1,\nu} + \mathcal{L}_2)V + k\rho^2(\mathcal{P}_1 + \mathcal{P}_{2,\nu})V = \mathcal{F}_\nu,$$

where

$$\begin{cases} D(\mathcal{L}_{1,\nu}) &= \{V \in W^{2,p}(0, +\infty; X) : V(0) = V(+\infty) = 0\} \\ [\mathcal{L}_{1,\nu}(V)](t) &= (\partial_t - \nu I)^2 V(t) = V''(t) - 2\nu V'(t) + \nu^2 V(t), \\ D(\mathcal{L}_2) &= \{V \in L^p(0, +\infty; X) : \text{for a.e. } t \in (0, +\infty), V(t) \in D(\mathcal{A})\} \\ [\mathcal{L}_2(V)](t) &= -\mathcal{A}V(t), \end{cases}$$

with

$$\begin{cases} D(\mathcal{A}) &= [W^{4,p}(0, \omega) \cap W_0^{2,p}(0, \omega)] \times W_0^{2,p}(0, \omega) \subset X \\ \mathcal{A} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} &= \begin{pmatrix} \psi_2 \\ -\left(\frac{\partial^2}{\partial \theta^2} + 1\right)^2 \psi_1 - 2\left(\frac{\partial^2}{\partial \theta^2} - 1\right) \psi_2 \end{pmatrix}, \quad \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \in D(\mathcal{A}), \end{cases}$$

and

$$\begin{cases} D(\mathcal{P}_1) &= \{V \in L^p(0, +\infty; X) : \text{for a.e. } t \in (0, +\infty), V(t) \in D(\mathcal{A}_0)\} \\ [\mathcal{P}_1(V)](t) &= -e^{-2t} \mathcal{A}_0 V(t), \end{cases}$$

with

$$\begin{cases} D(\mathcal{A}_0) &= W_0^{2,p}(0, \omega) \times L^p(0, \omega) = X \\ \mathcal{A}_0 \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} &= \begin{pmatrix} 0 \\ \left(\frac{\partial^2}{\partial \theta^2} + 1\right) \psi_1 + \psi_2 \end{pmatrix}, \quad \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \in D(\mathcal{A}_0), \end{cases}$$

and

$$\begin{cases} D(\mathcal{P}_{2,\nu}) &= W^{1,p}(0, +\infty; X) \\ [\mathcal{P}_{2,\nu}(V)](t) &= -e^{-2t} (\mathcal{B}_{2,\nu} V)(t), \end{cases}$$

with

$$\mathcal{B}_{2,\nu} = \begin{pmatrix} 0 & 0 \\ -2(\partial_t - \nu I) & 0 \end{pmatrix}.$$

In the present paper, we will focus ourselves on the resolution of the following abstract equation

$$(\mathcal{L}_{1,\mu} + \mathcal{L}_2)V + k\rho^2(\mathcal{P}_1 + \mathcal{P}_{2,\mu})V = \mathcal{F}, \quad (4)$$

where $\mu \in \mathbb{R}$ is a general parameter and $\mathcal{F} \in L^p(0, +\infty; X)$.

Note that, in part I, see Labbas, Maingot and Thorel [14], subsection 3.4, we have worked with $\mu = \nu = 3 - \frac{2}{p}$ which comes from the variable change concerning the weighted Sobolev space. Here, in this second part, we consider a more general $\mu \in \mathbb{R}$.

The aim of this work is to show that there exists a unique classical solution of (4) that is a function V such that

$$V \in W^{2,p}(0, +\infty; X) \cap L^p(0, +\infty; D(\mathcal{A})).$$

This regularity is necessary to deduce all those of the function v stated in Theorem 2.2 in Labbas, Maingot and Thorel [14].

To this end, we will use the Da Prato-Grisvard sum theory in order to invert $\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2}$. Then, we solve the following equation

$$\left(\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2}\right) V + k\rho^2 (\mathcal{P}_1 + \mathcal{P}_{2,\mu}) V = \mathcal{F},$$

by using a perturbation argument. Next, we use some convexity inequalities to prove that V belongs to a more suitable space, more precisely

$$V \in W^{1,p}(0, +\infty; X) \cap L^p\left(0, +\infty; \left[W^{3,p}(0, \omega) \cap W_0^{2,p}(0, \omega)\right] \times L^p(0, \omega)\right).$$

At this step, V is the unique strong solution of (4), see (46). To obtain that V is a classical solution, we use the Dore-Venni sum theory, see Section 6.

Among the results that we will use is the fact that the roots of the equation

$$(\sinh(z) + z)(\sinh(z) - z) = 0,$$

in $\mathbb{C}_+ := \{w \in \mathbb{C} : \operatorname{Re}(w) > 0\}$, constitute a family of complex numbers $(z_j)_{j \geq 1}$ such that

$$\tau := \min_{j \geq 1} |\operatorname{Im}(z_j)| > 0 \quad \text{and} \quad |z_j| \longrightarrow +\infty.$$

These roots are computed in Fädle [7] with $\tau \simeq 4.21239$.

Our main result is the following.

Theorem 1.1. Let $\mathcal{F} \in L^p(0, +\infty; X)$ and assume that

$$\omega\mu < \tau. \tag{5}$$

Then, there exists $\rho_0 > 0$ such that for all $\rho \in (0, \rho_0]$, the abstract equation

$$(\mathcal{L}_{1,\mu} + \mathcal{L}_2) V + k\rho^2 (\mathcal{P}_1 + \mathcal{P}_{2,\mu}) V = \mathcal{F},$$

has a unique classical solution $V \in L^p(0, +\infty; X)$, that is

$$V \in W^{2,p}(0, +\infty; X) \cap L^p(0, +\infty; D(\mathcal{A})).$$

In particular, $\mathcal{L}_{1,\mu} + \mathcal{L}_2$ is closed and $V \in D(\mathcal{L}_{1,\mu} + \mathcal{L}_2)$.

This second part is organized as follows. Section 2 is devoted to recalling some needed results. In Section 3, we analyze the spectral properties of operators $\mathcal{L}_{1,\mu}$ and \mathcal{L}_2 in view to study the invertibility of $\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2}$ in Section 4. In Section 5, by considering that operator $k\rho^2 (\mathcal{P}_1 + \mathcal{P}_{2,\mu})$ is a perturbation, we deduce the existence and the uniqueness of a strong solution of equation (4). Finally, Section 6 is devoted to the proof of our main result given in Theorem 1.1.

2 Definitions and prerequisites

2.1 The class of Bounded Imaginary Powers of operators

Definition 2.1. A Banach space E is a UMD space if and only if for all $p \in (1, +\infty)$, the Hilbert transform is bounded from $L^p(\mathbb{R}, E)$ into itself (see Bourgain [2] and Burkholder [3]).

Definition 2.2. Let $\alpha \in (0, \pi)$. $\operatorname{Sect}(\alpha)$ denotes the space of closed linear operators T_1 which satisfying

- i) $\sigma(T_1) \subset \overline{S_\alpha}$,
- ii) $\forall \alpha' \in (\alpha, \pi), \quad \sup \left\{ \|\lambda(\lambda I - T_1)^{-1}\|_{\mathcal{L}(X)} : \lambda \in \mathbb{C} \setminus \overline{S_{\alpha'}} \right\} < +\infty,$

where

$$S_\alpha := \begin{cases} \{z \in \mathbb{C} : z \neq 0 \text{ and } |\arg(z)| < \alpha\} & \text{if } \alpha \in (0, \pi) \\ (0, +\infty) & \text{if } \alpha = 0, \end{cases} \tag{6}$$

see Haase [9], p. 19. Such an operator T_1 is called sectorial operator of angle α .

Remark 2.3. From Komatsu [11], p. 342, we know that any injective sectorial operator T_1 admits imaginary powers T_1^{is} for all $s \in \mathbb{R}$; but in general, T_1^{is} is not bounded.

Definition 2.4. Let $\kappa \in [0, \pi)$. We denote by $\text{BIP}(E, \kappa)$, the class of sectorial injective operators T_2 such that

- i) $\overline{D(T_2)} = \overline{R(T_2)} = E$,
- ii) $\forall s \in \mathbb{R}, \quad T_2^{is} \in \mathcal{L}(E)$,
- iii) $\exists C \geq 1, \forall s \in \mathbb{R}, \quad \|T_2^{is}\|_{\mathcal{L}(E)} \leq Ce^{|s|\kappa}$,

see Prüss and Sohr [19], p. 430.

2.2 Recall on the sum of linear operators

Let us fix a pair of two closed linear densely defined operators \mathcal{M}_1 and \mathcal{M}_2 in a general Banach space \mathcal{E} . We note their domains by $D(\mathcal{M}_1)$ and $D(\mathcal{M}_2)$ respectively. Then we can define their sum by

$$\begin{cases} \mathcal{M}_1 w + \mathcal{M}_2 w \\ w \in D(\mathcal{M}_1) \cap D(\mathcal{M}_2). \end{cases}$$

We assume the following hypotheses

(H₁) There exist $\theta_{\mathcal{M}_1} \in [0, \pi)$, $\theta_{\mathcal{M}_2} \in [0, \pi)$, $C > 0$ and $R > 0$ such that

$$\begin{cases} \rho(\mathcal{M}_1) \supset \Sigma_{1,R} = \{z \in \mathbb{C} \setminus \{0\} : |z| \geq R \text{ and } |\arg(z)| < \pi - \theta_{\mathcal{M}_1}\} \\ \forall z \in \Sigma_{1,R}, \quad \|(\mathcal{M}_1 - zI)^{-1}\| \leq \frac{C}{|z|}, \end{cases}$$

and

$$\begin{cases} \rho(\mathcal{M}_2) \supset \Sigma_{2,R} = \{z \in \mathbb{C} \setminus \{0\} : |z| \geq R \text{ and } |\arg(z)| < \pi - \theta_{\mathcal{M}_2}\} \\ \forall z \in \Sigma_{2,R}, \quad \|(\mathcal{M}_2 - zI)^{-1}\| \leq \frac{C}{|z|}, \end{cases}$$

with

$$\theta_{\mathcal{M}_1} + \theta_{\mathcal{M}_2} < \pi.$$

(H₂) $\sigma(\mathcal{M}_1) \cap \sigma(-\mathcal{M}_2) = \emptyset$.

(H₃) The resolvents of \mathcal{M}_1 and \mathcal{M}_2 commute, that is

$$(\mathcal{M}_1 - \lambda_1 I)^{-1} (\mathcal{M}_2 - \lambda_2 I)^{-1} = (\mathcal{M}_2 - \lambda_2 I)^{-1} (\mathcal{M}_1 - \lambda_1 I)^{-1},$$

for all $\lambda_1 \in \rho(\mathcal{M}_1)$ and all $\lambda_2 \in \rho(\mathcal{M}_2)$.

Remark 2.5. Note that from (H₂), we have $\rho(\mathcal{M}_1) \cup \rho(-\mathcal{M}_2) = \mathbb{C}$ and in particular \mathcal{M}_1 or \mathcal{M}_2 is boundedly invertible.

Theorem 2.6 (Da Prato and Grisvard [4], Grisvard [8]). Assume that (H₁), (H₂) and (H₃) hold. Then, operator $\mathcal{M}_1 + \mathcal{M}_2$ is closable. Its closure $\overline{\mathcal{M}_1 + \mathcal{M}_2}$ is boundedly invertible and

$$\left(\overline{\mathcal{M}_1 + \mathcal{M}_2}\right)^{-1} = \frac{-1}{2i\pi} \int_{\Gamma} (\mathcal{M}_1 - zI)^{-1} (\mathcal{M}_2 + zI)^{-1} dz; \quad (7)$$

where Γ is a path which separates $\sigma(\mathcal{M}_1)$ and $\sigma(-\mathcal{M}_2)$ and joins $\infty e^{-i\theta_0}$ to $\infty e^{i\theta_0}$ with θ_0 such that

$$\theta_{\mathcal{M}_1} < \theta_0 < \pi - \theta_{\mathcal{M}_2}.$$

This Theorem is proved in Da Prato and Grisvard [4] (Theorem 3.7, p. 324), when $R = 0$ and has been extended to the case $R \geq 0$ in Grisvard [8] (Theorem 2.1, p. 7). In this last case, the curve Γ does not need to be connected.

Corollary 2.7. Assume that (H_1) , (H_2) and (H_3) hold. Let $i = 1, 2$ and \mathcal{E}_i a Banach space with $D(\mathcal{M}_i) \hookrightarrow \mathcal{E}_i \hookrightarrow \mathcal{E}$. We suppose that there exist $C > 0$ and $\delta \in (0, 1)$ such that

$$\begin{cases} \|w\|_{\mathcal{E}_i} \leq C \left(\|w\|_{\mathcal{E}} + \|w\|_{\mathcal{E}}^{1-\delta} \|\mathcal{M}_i w\|_{\mathcal{E}}^{\delta} \right) \\ \text{for every } w \in D(\mathcal{M}_i). \end{cases} \quad (8)$$

Then $D(\overline{\mathcal{M}_1 + \mathcal{M}_2}) \subset \mathcal{E}_i$.

Proof. It is enough to prove that the integral in (7) converges in \mathcal{E}_1 . For all $\xi \in \mathcal{E}$, we have

$$\left\| \int_{\Gamma} (\mathcal{M}_1 - zI)^{-1} (\mathcal{M}_2 + zI)^{-1} \xi dz \right\|_{\mathcal{E}_1} \leq \int_{\Gamma} \left\| (\mathcal{M}_1 - zI)^{-1} (\mathcal{M}_2 + zI)^{-1} \xi \right\|_{\mathcal{E}_1} |dz|,$$

then, applying (8), we obtain

$$\begin{aligned} & \left\| (\mathcal{M}_1 - zI)^{-1} (\mathcal{M}_2 + zI)^{-1} \xi \right\|_{\mathcal{E}_1} \\ & \leq C \left\| (\mathcal{M}_1 - zI)^{-1} (\mathcal{M}_2 + zI)^{-1} \xi \right\|_{\mathcal{E}} \\ & + C \left\| (\mathcal{M}_1 - zI)^{-1} (\mathcal{M}_2 + zI)^{-1} \xi \right\|_{\mathcal{E}}^{1-\delta} \left\| \mathcal{M}_1 (\mathcal{M}_1 - zI)^{-1} (\mathcal{M}_2 + zI)^{-1} \xi \right\|_{\mathcal{E}}^{\delta}. \end{aligned}$$

Now, for all $z \in \Gamma$, we have

$$\begin{aligned} & \left\| (\mathcal{M}_1 - zI)^{-1} (\mathcal{M}_2 + zI)^{-1} \xi \right\|_{\mathcal{E}}^{1-\delta} \left\| \mathcal{M}_1 (\mathcal{M}_1 - zI)^{-1} (\mathcal{M}_2 + zI)^{-1} \xi \right\|_{\mathcal{E}}^{\delta} \\ & \leq \frac{[C_1(\theta_{\mathcal{M}_1})C_2(\theta_{\mathcal{M}_2})]^{1-\delta} [C_1(\theta_{\mathcal{M}_1})]^{\delta} [C_2(\theta_{\mathcal{M}_2})]^{\delta}}{|z|^{2(1-\delta)} |z|^{\delta}} \|\xi\|_{\mathcal{E}} = \frac{C_1(\theta_{\mathcal{M}_1})C_2(\theta_{\mathcal{M}_2})}{|z|^{1+(1-\delta)}} \|\xi\|_{\mathcal{E}}, \end{aligned}$$

from which we deduce the convergence of the integral in (7). \square

3 Spectral study of operators

In all the sequel, in view to apply the above results, we will consider the following particular Banach space

$$\mathcal{E} = L^p(0, +\infty; X) \quad \text{with} \quad X = W_0^{2,p}(0, \omega) \times L^p(0, \omega),$$

equipped with its natural norm.

3.1 Study of operator $\mathcal{L}_{1,\mu}$

We study the spectral equation

$$\mathcal{L}_{1,\mu} V - \lambda V = R,$$

where $V \in D(\mathcal{L}_{1,\mu})$, $R \in \mathcal{E}$ and $\lambda \in \mathbb{C}$ (which will be precised below), that is

$$\begin{cases} V''(t) - 2\mu V'(t) + (\mu^2 - \lambda)V(t) = R(t), & t > 0 \\ V(0) = 0, \quad V(+\infty) = 0. \end{cases} \quad (9)$$

We set

$$\Pi_{\mu} = \{z \in \mathbb{C} \setminus \mathbb{R}_- : \operatorname{Re}(\sqrt{z}) > \mu\}.$$

Now, let us specify Π_{μ} . For all $z = x + iy \in \mathbb{C} \setminus \mathbb{R}_-$, we have

$$\operatorname{Re}(\sqrt{z}) > \mu \iff \sqrt{\frac{|z| + \operatorname{Re}(z)}{2}} > \mu \iff \sqrt{x^2 + y^2} > 2\mu^2 - x.$$

- First case : if $x > \mu^2$, we have $\sqrt{x^2 + y^2} + x \geq 2x > 2\mu^2$, then $\operatorname{Re}(\sqrt{z}) > \mu$.
- Second case : if $x \leq \mu^2$, then $y^2 + 4\mu^2x - 4\mu^4 > 0$. Thus, we deduce that Π_μ is strictly outside the parabola of equation

$$y^2 + 4\mu^2x - 4\mu^4 = 0,$$

turned towards the negative real axis and passing through the points $(\mu^2, 0)$, $(0, 2\mu^2)$ and $(0, -2\mu^2)$.




Figure 1: In this figure, Π_μ is the entire uncolored area.

Now, let $\varepsilon_{\mathcal{L}_{1,\mu}}$ be a small fixed positive number and consider the following set

$$\Sigma_{\mathcal{L}_{1,\mu}} := \left\{ \lambda \in \Pi_\mu, \quad |\arg(\lambda)| \leq \pi - 2\varepsilon_{\mathcal{L}_{1,\mu}} \quad \text{and} \quad |\lambda| \geq \frac{4\mu^2}{\sin^2(\varepsilon_{\mathcal{L}_{1,\mu}})} \right\}. \quad (10)$$

We then obtain the following proposition.

Proposition 3.1. The linear operator $\mathcal{L}_{1,\mu}$ is closed and densely defined in \mathcal{E} . Moreover, there exists a constant $M_{\mathcal{L}_{1,\mu}} > 0$ such that for all $\lambda \in \Sigma_{\mathcal{L}_{1,\mu}}$, operator $\mathcal{L}_{1,\mu} - \lambda I$ is invertible and

$$\|(\mathcal{L}_{1,\mu} - \lambda I)^{-1}\|_{\mathcal{L}(\mathcal{E})} \leq \frac{M_{\mathcal{L}_{1,\mu}}}{|\lambda|}.$$

Therefore, assumption (H_1) in Section 2.2 is verified for $\mathcal{L}_{1,\mu}$ with

$$\theta_{\mathcal{L}_{1,\mu}} = 2\varepsilon_{\mathcal{L}_{1,\mu}}. \quad (11)$$

Proof. Let $\lambda \in \Pi_\mu$. From Eltaief and Maingot [6], Theorem 2, p. 712, there exists a unique solution $V \in W^{2,p}(0, +\infty; X)$ of problem (9), given by

$$\begin{aligned} V(t) &= \frac{e^{t(\mu - \sqrt{\lambda})}}{2\sqrt{\lambda}} \int_0^{+\infty} e^{-s(\mu + \sqrt{\lambda})} R(s) ds \\ &\quad - \frac{1}{2\sqrt{\lambda}} \left(\int_0^t e^{(t-s)(\mu - \sqrt{\lambda})} R(s) ds + \int_t^{+\infty} e^{-(s-t)(\mu + \sqrt{\lambda})} R(s) ds \right), \end{aligned} \quad (12)$$

see formula (15) in Eltaief and Maingot [6] where $L_1 := -\mu I - \sqrt{\lambda} I$ and $L_2 := \mu I - \sqrt{\lambda} I$. It follows that $\Pi_\mu \subset \rho(\mathcal{L}_{1,\mu})$. This proves that $\mathcal{L}_{1,\mu}$ is closed. The boundary conditions are verified by using Lemma 8, p. 718 in Eltaief and Maingot [6].

Moreover, from (12), we obtain

$$\begin{aligned} \|V\|_{\mathcal{E}} &\leq \frac{1}{2\sqrt{|\lambda|}} \left(\int_0^{+\infty} e^{-tp(\operatorname{Re}(\sqrt{\lambda})-\mu)} dt \right)^{1/p} \int_0^{+\infty} e^{-s(\mu+\operatorname{Re}(\sqrt{\lambda}))} \|R(s)\|_X ds \\ &\quad + \sup_{t \in \mathbb{R}_+} \left(\int_0^t |e^{(t-s)(\mu-\sqrt{\lambda})}| ds + \int_t^{+\infty} |e^{-(s-t)(\mu+\sqrt{\lambda})}| ds \right) \frac{\|R\|_{\mathcal{E}}}{2\sqrt{|\lambda|}}, \end{aligned}$$

hence, noting q the conjugate exponent of p , we have

$$\begin{aligned} \|V\|_{\mathcal{E}} &\leq \left(\frac{1}{p(\operatorname{Re}(\sqrt{\lambda})-\mu)} \right)^{1/p} \left(\int_0^{+\infty} e^{-sq(\mu+\operatorname{Re}(\sqrt{\lambda}))} ds \right)^{1/q} \frac{\|R\|_{\mathcal{E}}}{2\sqrt{|\lambda|}} \\ &\quad + \sup_{t \in \mathbb{R}_+} \left(\frac{1 - e^{-t(\operatorname{Re}(\sqrt{\lambda})-\mu)}}{\operatorname{Re}(\sqrt{\lambda})-\mu} + \frac{1}{\operatorname{Re}(\sqrt{\lambda})+\mu} \right) \frac{\|R\|_{\mathcal{E}}}{2\sqrt{|\lambda|}} \\ &\leq \frac{1}{p^{1/p}(\operatorname{Re}(\sqrt{\lambda})-\mu)^{1/p}} \frac{1}{q^{1/q}(\operatorname{Re}(\sqrt{\lambda})+\mu)^{1/q}} \frac{\|R\|_{\mathcal{E}}}{2\sqrt{|\lambda|}} \\ &\quad + \frac{2}{\operatorname{Re}(\sqrt{\lambda})-\mu} \frac{\|R\|_{\mathcal{E}}}{2\sqrt{|\lambda|}} \\ &\leq \frac{2}{\operatorname{Re}(\sqrt{\lambda})-\mu} \frac{\|R\|_{\mathcal{E}}}{\sqrt{|\lambda|}}. \end{aligned}$$

Let $\lambda = |\lambda|e^{i\arg(\lambda)} \in \Sigma_{\mathcal{L}_{1,\mu}}$, then $|\arg(\lambda)| \leq \pi - 2\varepsilon_{\mathcal{L}_{1,\mu}}$. Thus

$$\begin{aligned} \operatorname{Re}(\sqrt{\lambda}) - \mu &\geq \sqrt{|\lambda|} \cos\left(\frac{\arg(\lambda)}{2}\right) - \mu \geq \sqrt{|\lambda|} \cos\left(\frac{\pi}{2} - \varepsilon_{\mathcal{L}_{1,\mu}}\right) - \mu \\ &\geq \sqrt{|\lambda|} \sin(\varepsilon_{\mathcal{L}_{1,\mu}}) - \frac{\sqrt{|\lambda|}}{2} \sin(\varepsilon_{\mathcal{L}_{1,\mu}}) \geq \frac{\sqrt{|\lambda|}}{2} \sin(\varepsilon_{\mathcal{L}_{1,\mu}}). \end{aligned}$$

We then obtain

$$\|V\|_{\mathcal{E}} \leq \frac{M_{\mathcal{L}_{1,\mu}}}{|\lambda|} \|R\|_{\mathcal{E}}, \quad (13)$$

where $M_{\mathcal{L}_{1,\mu}} = \frac{4}{\sin(\varepsilon_{\mathcal{L}_{1,\mu}})}$. □

3.2 Study of operator \mathcal{L}_2

The spectral properties of \mathcal{L}_2 are the same as those of its realization $-\mathcal{A}$.

In all this subsection, we assume that

$$\lambda \leq 0.$$

We have to solve the following spectral equation in X

$$\mathcal{A}\Psi - \lambda\Psi = F, \quad (14)$$

that is

$$\left(\begin{array}{cc} 0 & 1 \\ -\left(\frac{\partial^2}{\partial\theta^2} + 1\right)^2 & -2\left(\frac{\partial^2}{\partial\theta^2} - 1\right) \end{array} \right) \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} - \lambda \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = \begin{pmatrix} F_1 \\ F_2 \end{pmatrix},$$

with

$$F_1 \in W_0^{2,p}(0, \omega) \quad \text{and} \quad F_2 \in L^p(0, \omega).$$

We have to find the unique couple $(\psi_1, \psi_2) \in \left(W^{4,p}(0, \omega) \cap W_0^{2,p}(0, \omega)\right) \times W_0^{2,p}(0, \omega)$, which satisfies the following system

$$\begin{cases} \psi_2 - \lambda\psi_1 & = F_1 \\ -\psi_1^{(4)} - 2\psi_1'' - \psi_1 - 2\psi_2'' + 2\psi_2 - \lambda\psi_2 & = F_2. \end{cases}$$

Thus, we first have to solve

$$\begin{cases} -\psi_1^{(4)} - 2\psi_1'' - \psi_1 - 2(\lambda\psi_1 + F_1)'' + (2 - \lambda)(\lambda\psi_1 + F_1) = F_2 \\ \psi_1 \in W^{4,p}(0, \omega) \cap W_0^{2,p}(0, \omega), \end{cases}$$

that is

$$\begin{cases} -\psi_1^{(4)} - 2(1 + \lambda)\psi_1'' - (\lambda - 1)^2\psi_1 = F_2 + 2F_1'' + (\lambda - 2)F_1 \\ \psi_1(0) = \psi_1(\omega) = \psi_1'(0) = \psi_1'(\omega) = 0. \end{cases}$$

Set $G_\lambda = -F_2 - 2(F_1'' - F_1) - \lambda F_1$, it follows that the previous system writes

$$\begin{cases} \psi_1^{(4)} + 2(\lambda + 1)\psi_1'' + (\lambda - 1)^2\psi_1 = G_\lambda \\ \psi_1(0) = \psi_1(\omega) = \psi_1'(0) = \psi_1'(\omega) = 0. \end{cases} \quad (15)$$

Then, the characteristic equation

$$\chi^4 + 2(1 + \lambda)\chi^2 + (\lambda - 1)^2 = 0,$$

admits, for $\lambda < 0$, the following four distinct solutions

$$\begin{cases} \alpha_1 = \sqrt{-\lambda} + i, & \alpha_3 = -\alpha_1 \\ \alpha_2 = \sqrt{-\lambda} - i, & \alpha_4 = -\alpha_2, \end{cases} \quad (16)$$

and for $\lambda = 0$, two double solutions i and $-i$.

We have to distinguish below two cases : $\lambda = 0$ and $\lambda < 0$.

3.2.1 Case $\lambda = 0$: Invertibility of \mathcal{A}

Proposition 3.2. \mathcal{A} is boundedly invertible. Then : $\exists \varepsilon_0 > 0 : \overline{B(0, \varepsilon_0)} \subset \rho(\mathcal{A})$.

Proof. Here $\lambda = 0$. We have to solve (14). This is equivalent to solve (15) that is

$$\begin{cases} \psi_1^{(4)} + 2\psi_1'' + \psi_1 = -F_2 - 2(F_1'' - F_1) \\ \psi_1(0) = \psi_1(\omega) = \psi_1'(0) = \psi_1'(\omega) = 0. \end{cases} \quad (17)$$

From Thorel [22], Theorem 2.8, statement 2., there exists a unique classical solution of problem (17) that is $(\psi_1, \psi_2) = (\psi_1, F_1) \in \left(W^{4,p}(0, \omega) \cap W_0^{2,p}(0, \omega)\right) \times W_0^{2,p}(0, \omega)$. We then deduce that there exists $C_1 > 0$ such that

$$\|\psi_1''\|_{L^p(0, \omega)} \leq C_1 \left(\|F_2\|_{L^p(0, \omega)} + 2\|F_1\|_{W^{2,p}(0, \omega)} \right) \leq 2C_1 \|F\|_X,$$

and from the Poincaré inequality, there exists $C_\omega > 0$ such that

$$\|\psi_1\|_{W_0^{2,p}(0, \omega)} \leq C_\omega \|\psi_1''\|_{L^p(0, \omega)} \leq 2C_1 C_\omega \|F\|_X.$$

Finally, since $\psi_2 = F_1$, we have

$$\|\psi_2\|_{L^p(0, \omega)} = \|F_1\|_{L^p(0, \omega)} \leq \|F_2\|_{L^p(0, \omega)} + \|F_1\|_{W^{2,p}(0, \omega)} = \|F\|_X.$$

□

3.2.2 Case $\lambda < 0$: Spectral study of \mathcal{A}

In order to prove Proposition 3.9, we first have to state the following technical results.

Lemma 3.3. Let $\alpha \in \mathbb{C} \setminus \{0\}$, $a, b \in \mathbb{R}$ with $a < b$ and $f \in W_0^{2,p}(a, b)$. For all $x \in [a, b]$, we set

$$K(x) = \int_a^x e^{-(x-s)\alpha} f(s) ds + \int_x^b e^{-(s-x)\alpha} f(s) ds.$$

Then, we have

$$K(x) = \frac{2}{\alpha} f(x) + \frac{1}{\alpha^2} \int_a^x e^{-(x-s)\alpha} f''(s) ds + \frac{1}{\alpha^2} \int_x^b e^{-(s-x)\alpha} f''(s) ds.$$

Proof. The result is easily obtained by two integrations by parts. \square

Now, let us solve explicitly problem (15) for $\lambda < 0$.

Proposition 3.4. Problem (15) has a unique solution which can be written in the following form

$$\begin{aligned} \psi_1(\theta) &:= e^{-\theta\alpha_2}(\beta_1 + \beta_2 + \beta_3 + \beta_4) + e^{-(\omega-\theta)\alpha_2}(\beta_3 + \beta_4 - \beta_1 - \beta_2) + S(\theta) \\ &+ \left(e^{-\theta\alpha_1} - e^{-\theta\alpha_2} \right) (\beta_2 + \beta_4) + \left(e^{-(\omega-\theta)\alpha_1} - e^{-(\omega-\theta)\alpha_2} \right) (\beta_4 - \beta_2), \end{aligned} \quad (18)$$

where the constants β_i , $i = 1, 2, 3, 4$, and the particular solution S will be explicitly given below.

Proof. In order to apply results obtained in Labbas, Lemrabet, Maingot and Thorel [12] and Labbas, Maingot, Manceau and Thorel [13], we set

$$L_- = -\alpha_1 I, \quad M = -\alpha_2 I, \quad r_- = \alpha_1^2 - \alpha_2^2, \quad a = 0 \quad \text{and} \quad b = \omega.$$

Then, problem (15) can be written as

$$\begin{cases} \psi_1^{(4)} - (L_-^2 + M^2)\psi_1'' + L_-^2 M^2 \psi_1 = G_\lambda \\ \psi_1(0) = \psi_1(\omega) = \psi_1'(0) = \psi_1'(\omega) = 0. \end{cases} \quad (19)$$

From Labbas, Maingot, Manceau and Thorel [13], there exists a unique classical solution ψ_1 of problem (19). Its representation formula, in the form (18), is explicitly given in Labbas, Lemrabet, Maingot and Thorel [12] by (14)-(15)-(16) or also in a more general framework in Labbas, Maingot and Thorel [15] by (22)-(30)-(31).

For the reader convenience, we will prove, by a long calculus, that

$$\begin{cases} \beta_1 = \frac{1}{4i} U_1^{-1} \frac{1 - e^{-\omega\alpha_1}}{1 - e^{-\omega\alpha_2}} (J(0) - J(\omega)) \\ \beta_2 = -\frac{1}{4i} U_1^{-1} (J(0) - J(\omega)) \\ \beta_3 = -\frac{1}{4i} U_2^{-1} \frac{1 + e^{-\omega\alpha_1}}{1 + e^{-\omega\alpha_2}} (J(0) + J(\omega)) \\ \beta_4 = \frac{1}{4i} U_2^{-1} (J(0) + J(\omega)), \end{cases} \quad (20)$$

with

$$\begin{cases} U_1 := 1 - e^{-2\omega\sqrt{-\lambda}} - 2\omega\sqrt{-\lambda} e^{-\omega\sqrt{-\lambda}} \\ U_2 := 1 - e^{-2\omega\sqrt{-\lambda}} + 2\omega\sqrt{-\lambda} e^{-\omega\sqrt{-\lambda}}, \end{cases} \quad (21)$$

and for all $\theta \in [0, \omega]$

$$\begin{aligned} S(\theta) &:= \frac{e^{-\theta\alpha_2}}{2\alpha_2(1-e^{-2\omega\alpha_2})} (J(0) - e^{-\omega\alpha_2}J(\omega)) - \frac{\lambda}{\alpha_1^2\alpha_2^2} F_1(\theta) \\ &\quad + \frac{e^{-(\omega-\theta)\alpha_2}}{2\alpha_2(1-e^{-2\omega\alpha_2})} (J(\omega) - e^{-\omega\alpha_2}J(0)) - \frac{1}{2\alpha_2} J(\theta), \end{aligned} \quad (22)$$

with

$$J(\theta) := \int_0^\theta e^{-(\theta-s)\alpha_2} v(s) ds + \int_\theta^\omega e^{-(s-\theta)\alpha_2} v(s) ds, \quad (23)$$

where

$$\begin{aligned} v(\theta) &:= \frac{e^{-\theta\alpha_1}}{2\alpha_1(1-e^{-2\omega\alpha_1})} (I(0) - e^{-\omega\alpha_1}I(\omega)) + \frac{\lambda}{\alpha_1^2\alpha_2^2} F_1''(\theta) \\ &\quad + \frac{e^{-(\omega-\theta)\alpha_1}}{2\alpha_1(1-e^{-2\omega\alpha_1})} (I(\omega) - e^{-\omega\alpha_1}I(0)) - \frac{1}{2\alpha_1} I(\theta), \end{aligned} \quad (24)$$

and

$$\begin{aligned} I(\theta) &= \int_0^\theta e^{-(\theta-s)\alpha_1} \left(-F_2 - 2(F_1'' - F_1) + \frac{\lambda}{\alpha_1^2} F_1'' \right) (s) ds \\ &\quad + \int_\theta^\omega e^{-(s-\theta)\alpha_1} \left(-F_2 - 2(F_1'' - F_1) + \frac{\lambda}{\alpha_1^2} F_1'' \right) (s) ds. \end{aligned} \quad (25)$$

Let us begin our proof. Here, we are inspired by the formulas given by (15) and (16), p. 2943, in Labbas, Lemrabet, Maingot and Thorel [12], where the authors have used the notations F_- , f_- , U_- and V_- which are replace respectively here by S , G_λ , U_1 and U_2 . Consequently, we have

$$\begin{aligned} S(\theta) &= \frac{Ze^{-\theta\alpha_2}}{2\alpha_2} \int_0^\omega e^{-s\alpha_2} v_0(s) ds + \frac{Ze^{-(\omega-\theta)\alpha_2}}{2\alpha_2} \int_0^\omega e^{-(\omega-s)\alpha_2} v_0(s) ds \\ &\quad - \frac{1}{2\alpha_2} \int_0^\theta e^{-(\theta-s)\alpha_2} v_0(s) ds - \frac{1}{2\alpha_2} \int_\theta^\omega e^{-(s-\theta)\alpha_2} v_0(s) ds \\ &\quad - \frac{Ze^{-\omega\alpha_2}e^{-\theta\alpha_2}}{2\alpha_2} \int_0^\omega e^{-(\omega-s)\alpha_2} v_0(s) ds \\ &\quad - \frac{Ze^{-\omega\alpha_2}e^{-(\omega-\theta)\alpha_2}}{2\alpha_2} \int_0^\omega e^{-s\alpha_2} v_0(s) ds, \end{aligned} \quad (26)$$

where

$$\begin{aligned} v_0(\theta) &:= \frac{We^{-\theta\alpha_1}}{2\alpha_1} \int_0^\omega e^{-s\alpha_1} G_\lambda(s) ds + \frac{We^{-(\omega-\theta)\alpha_1}}{2\alpha_1} \int_0^\omega e^{-(\omega-s)\alpha_1} G_\lambda(s) ds \\ &\quad - \frac{We^{-\omega\alpha_1}e^{-\theta\alpha_1}}{2\alpha_1} \int_0^\omega e^{-(\omega-s)\alpha_1} G_\lambda(s) ds \\ &\quad - \frac{We^{-\omega\alpha_1}e^{-(\omega-\theta)\alpha_1}}{2\alpha_1} \int_0^\omega e^{-s\alpha_1} G_\lambda(s) ds - \frac{1}{2\alpha_1} I_1(\theta), \end{aligned} \quad (27)$$

with $Z := (1 - e^{-2\omega\alpha_2})^{-1}$, $W := (1 - e^{-2\omega\alpha_1})^{-1}$ and

$$I_1(\theta) = \int_0^\theta e^{-(\theta-s)\alpha_1} G_\lambda(s) ds + \int_\theta^\omega e^{-(s-\theta)\alpha_1} G_\lambda(s) ds. \quad (28)$$

Then, since $G_\lambda = -F_2 - 2(F_1'' - F_1) - \lambda F_1$, we have

$$\begin{aligned} I_1(\theta) &= \int_0^\theta e^{-(\theta-s)\alpha_1} (-F_2 - 2(F_1'' - F_1)) (s) ds \\ &\quad + \int_\theta^\omega e^{-(s-\theta)\alpha_1} (-F_2 - 2(F_1'' - F_1)) (s) ds \\ &\quad - \lambda \int_0^\theta e^{-(\theta-s)\alpha_1} F_1(s) ds - \lambda \int_\theta^\omega e^{-(s-\theta)\alpha_1} F_1(s) ds. \end{aligned}$$

From Lemma 3.3 and the fact that $F_1 \in W_0^{2,p}(0, \omega)$, it follows that

$$\begin{aligned}
I_1(\theta) &= \int_0^\theta e^{-(\theta-s)\alpha_1} (-F_2 - 2(F_1'' - F_1))(s) ds \\
&\quad + \int_\theta^\omega e^{-(s-\theta)\alpha_1} (-F_2 - 2(F_1'' - F_1))(s) ds \\
&\quad - \frac{2\lambda}{\alpha_1} F_1(\theta) + \frac{\lambda}{\alpha_1^2} \left(\int_0^\theta e^{-(\theta-s)\alpha_1} F_1''(s) ds + \int_\theta^\omega e^{-(s-\theta)\alpha_1} F_1''(s) ds \right) \\
&= -\frac{2\lambda}{\alpha_1} F_1(\theta) + \int_0^\theta e^{-(\theta-s)\alpha_1} \left(-F_2 - 2(F_1'' - F_1) + \frac{\lambda}{\alpha_1^2} F_1'' \right) (s) ds \\
&\quad + \int_\theta^\omega e^{-(s-\theta)\alpha_1} \left(-F_2 - 2(F_1'' - F_1) + \frac{\lambda}{\alpha_1^2} F_1'' \right) (s) ds.
\end{aligned}$$

Thus I , given by (25), satisfies

$$I(\theta) = I_1(\theta) + \frac{2\lambda}{\alpha_1} F_1(\theta). \quad (29)$$

Note that, from (28) and (29), we have

$$\int_0^\omega e^{-s\alpha_1} G_\lambda(s) ds = I_1(0) = I(0) \quad \text{and} \quad \int_0^\omega e^{-(\omega-s)\alpha_1} G_\lambda(s) ds = I_1(\omega) = I(\omega).$$

Therefore, from (27), we deduce that for all $\theta \in [0, \omega]$

$$\begin{aligned}
v_0(\theta) &= \frac{W e^{-\theta\alpha_1}}{2\alpha_1} (I(0) - e^{-\omega\alpha_1} I(\omega)) \\
&\quad + \frac{W e^{-(\omega-\theta)\alpha_1}}{2\alpha_1} (I(\omega) - e^{-\omega\alpha_1} I(0)) - \frac{1}{2\alpha_1} I_1(\theta) \\
&= \frac{W e^{-\theta\alpha_1}}{2\alpha_1} (I(0) - e^{-\omega\alpha_1} I(\omega)) + \frac{\lambda}{\alpha_1^2} F_1(\theta) \\
&\quad + \frac{W e^{-(\omega-\theta)\alpha_1}}{2\alpha_1} (I(\omega) - e^{-\omega\alpha_1} I(0)) - \frac{1}{2\alpha_1} I(\theta).
\end{aligned}$$

Set

$$v_1 = v_0 - \frac{\lambda}{\alpha_1^2} F_1, \quad (30)$$

and

$$J_1(\theta) = \int_0^\theta e^{-(\theta-s)\alpha_2} v_0(s) ds + \int_\theta^\omega e^{-(s-\theta)\alpha_2} v_0(s) ds; \quad (31)$$

then, due to Lemma 3.3, we obtain

$$\begin{aligned}
J_1(\theta) &= \int_0^\theta e^{-(\theta-s)\alpha_2} v_1(s) ds + \int_\theta^\omega e^{-(s-\theta)\alpha_2} v_1(s) ds \\
&\quad + \frac{\lambda}{\alpha_1^2} \left(\int_0^\theta e^{-(\theta-s)\alpha_2} F_1(s) ds + \int_\theta^\omega e^{-(s-\theta)\alpha_2} F_1(s) ds \right) \\
&= \int_0^\theta e^{-(\theta-s)\alpha_2} \left(v_1(s) + \frac{\lambda}{\alpha_1^2 \alpha_2} F_1''(s) \right) ds \\
&\quad + \int_\theta^\omega e^{-(s-\theta)\alpha_2} \left(v_1(s) + \frac{\lambda}{\alpha_1^2 \alpha_2} F_1''(s) \right) ds + \frac{2\lambda}{\alpha_1^2 \alpha_2} F_1(\theta).
\end{aligned}$$

From (30), for all $\theta \in [0, \omega]$, we deduce that v given by (24) and J given by (23), satisfy

$$v(\theta) = v_1(\theta) + \frac{\lambda}{\alpha_1^2 \alpha_2^2} F_1''(\theta),$$

and

$$J(\theta) = J_1(\theta) - \frac{2\lambda}{\alpha_1^2 \alpha_2} F_1(\theta). \quad (32)$$

Note that, from (31) and (32), we have

$$\int_0^\omega e^{-s\alpha_2} v_0(s) ds = J_1(0) = J(0) \quad \text{and} \quad \int_0^\omega e^{-(\omega-s)\alpha_2} v_0(s) ds = J_1(\omega) = J(\omega).$$

Finally, from (23), (26), (32) and for all $\theta \in [0, \omega]$, we deduce that

$$\begin{aligned} S(\theta) &= \frac{Ze^{-\theta\alpha_2}}{2\alpha_2} (J(0) - e^{-\omega\alpha_2} J(\omega)) \\ &\quad + \frac{Ze^{-(\omega-\theta)\alpha_2}}{2\alpha_2} (J(\omega) - e^{-\omega\alpha_2} J(0)) - \frac{1}{2\alpha_2} J_1(\theta), \end{aligned}$$

which is (22).

Now, in order to compute β_i , $i = 1, \dots, 4$, we must explicit U_1 , U_2 and $S'(0) \pm S'(\omega)$.

$$\begin{aligned} U_1 &= 1 - e^{-\omega(\alpha_1 + \alpha_2)} - (\alpha_1^2 - \alpha_2^2)^{-1} (\alpha_1 + \alpha_2)^2 (e^{-\omega\alpha_2} - e^{-\omega\alpha_1}) \\ &= 1 - e^{-2\omega\sqrt{-\lambda}} + i\sqrt{-\lambda} (e^{-\omega(\sqrt{-\lambda}-i)} - e^{-\omega(\sqrt{-\lambda}+i)}) \\ &= 1 - e^{-2\omega\sqrt{-\lambda}} - 2\sqrt{-\lambda} e^{-\omega\sqrt{-\lambda}} \sin(\omega), \end{aligned}$$

and

$$\begin{aligned} U_2 &= 1 - e^{-\omega(\alpha_1 + \alpha_2)} + (\alpha_1^2 - \alpha_2^2)^{-1} (\alpha_1 + \alpha_2)^2 (e^{-\omega\alpha_2} - e^{-\omega\alpha_1}) \\ &= 1 - e^{-2\omega\sqrt{-\lambda}} - i\sqrt{-\lambda} (e^{-\omega(\sqrt{-\lambda}-i)} - e^{-\omega(\sqrt{-\lambda}+i)}) \\ &= 1 - e^{-2\omega\sqrt{-\lambda}} + 2\sqrt{-\lambda} e^{-\omega\sqrt{-\lambda}} \sin(\omega). \end{aligned}$$

From (22), it follows that

$$\begin{aligned} S'(\theta) &= -\frac{Ze^{-\theta\alpha_2}}{2} (J(0) - e^{-\omega\alpha_2} J(\omega)) - \frac{\lambda}{\alpha_1^2 \alpha_2^2} F_1'(\theta) \\ &\quad + \frac{Ze^{-(\omega-\theta)\alpha_2}}{2} (J(\omega) - e^{-\omega\alpha_2} J(0)) - \frac{1}{2\alpha_2} J'(\theta) \\ &= -\frac{Ze^{-\theta\alpha_2}}{2} (J(0) - e^{-\omega\alpha_2} J(\omega)) + \frac{Ze^{-(\omega-\theta)\alpha_2}}{2} (J(\omega) - e^{-\omega\alpha_2} J(0)) \\ &\quad + \frac{1}{2} \left(\int_0^\theta e^{-(\theta-s)\alpha_2} v(s) ds - \int_\theta^\omega e^{-(s-\theta)\alpha_2} v(s) ds \right) - \frac{\lambda}{\alpha_1^2 \alpha_2^2} F_1'(\theta), \end{aligned}$$

then

$$S'(0) + S'(\omega) = -\frac{J(0) - J(\omega)}{(1 - e^{-\omega\alpha_2})} \quad \text{and} \quad S'(0) - S'(\omega) = -\frac{J(0) + J(\omega)}{(1 + e^{-\omega\alpha_2})}.$$

from which we deduce the constants β_i , $i = 1, 2, 3, 4$, written in (20). \square

Remark 3.5. Since $0 < \sin(\omega) < \omega$, for all $\omega > 0$, then we have

$$U_1 = 1 - e^{-2\omega\sqrt{-\lambda}} - 2\sqrt{-\lambda} e^{-\omega\sqrt{-\lambda}} \sin(\omega) \geq 1 - e^{-2\omega\sqrt{-\lambda}} - 2\omega\sqrt{-\lambda} e^{-\omega\sqrt{-\lambda}},$$

and

$$U_2 = 1 - e^{-2\omega\sqrt{-\lambda}} + 2\sqrt{-\lambda}e^{-\omega\sqrt{-\lambda}}\sin(\omega) \geq 1 - e^{-2\omega\sqrt{-\lambda}} - 2\omega\sqrt{-\lambda}e^{-\omega\sqrt{-\lambda}}.$$

For $x > 0$, we consider the following function

$$h(x) = 1 - e^{-2x} - 2xe^{-x};$$

we then have

$$h'(x) = 2e^{-2x} - 2e^{-x} + 2xe^{-x} = 2e^{-x}(e^{-x} + x - 1) > 0.$$

It follows that $h(x) > h(0) = 0$, for all $x > 0$. Finally, we deduce that

$$U_1 \geq 1 - e^{-2\omega\sqrt{-\lambda}} - 2\omega\sqrt{-\lambda}e^{-\omega\sqrt{-\lambda}} = h(\omega\sqrt{-\lambda}) > 0, \quad (33)$$

and

$$U_2 \geq 1 - e^{-2\omega\sqrt{-\lambda}} - 2\omega\sqrt{-\lambda}e^{-\omega\sqrt{-\lambda}} = h(\omega\sqrt{-\lambda}) > 0. \quad (34)$$

Lemma 3.6. For all $\lambda \in (-\infty, -\varepsilon_0]$, where ε_0 is defined in Proposition 3.2, functions J , v and I , given by (23), (24) and (25), satisfy the following estimates

1. $\|I\|_{L^p(0,\omega)} \leq \frac{2}{\sqrt{-\lambda}} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right).$
2. $|I(0)| + |I(\omega)| \leq \frac{2}{\sqrt{-\lambda}^{1-1/p}} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right).$
3. $\|v\|_{L^p(0,\omega)} \leq \frac{M_1}{-\lambda} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right),$
where $M_1 = 2 + \frac{2}{1 - e^{-2\omega\sqrt{-\lambda}}}$.
4. $\|J\|_{L^p(0,\omega)} \leq \frac{2}{\sqrt{-\lambda}} \|v\|_{L^p(0,\omega)}.$
5. $|J(0)| + |J(\omega)| \leq \frac{2}{\sqrt{-\lambda}^{1-1/p}} \|v\|_{L^p(0,\omega)}.$

Proof.

1. From (16) and (25), we obtain

$$\begin{aligned} \|I\|_{L^p(0,\omega)} &\leq \sup_{\theta \in [0,\omega]} \left(\int_0^\theta e^{-(\theta-s)\sqrt{-\lambda}} ds + \int_\theta^\omega e^{-(s-\theta)\sqrt{-\lambda}} ds \right) \|2F_1 - F_2\|_{L^p(0,\omega)} \\ &\quad + \sup_{\theta \in [0,\omega]} \left(\int_0^\theta e^{-(\theta-s)\sqrt{-\lambda}} ds + \int_\theta^\omega e^{-(s-\theta)\sqrt{-\lambda}} ds \right) \left\| \left(\frac{\lambda}{\alpha_1^2} - 2 \right) F_1'' \right\|_{L^p(0,\omega)} \\ &\leq \sup_{\theta \in [0,\omega]} \left(\frac{1 - e^{-\theta\sqrt{-\lambda}}}{\sqrt{-\lambda}} + \frac{1 - e^{-(\omega-\theta)\sqrt{-\lambda}}}{\sqrt{-\lambda}} \right) \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} \right) \\ &\quad + 3 \sup_{\theta \in [0,\omega]} \left(\frac{1 - e^{-\theta\sqrt{-\lambda}}}{\sqrt{-\lambda}} + \frac{1 - e^{-(\omega-\theta)\sqrt{-\lambda}}}{\sqrt{-\lambda}} \right) \|F_1''\|_{L^p(0,\omega)} \\ &\leq \frac{2}{\sqrt{-\lambda}} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right). \end{aligned}$$

2. Due to (16), (25) and the Hölder inequality, it follows

$$\begin{aligned}
|I(0)| + |I(\omega)| &\leq \int_0^\omega e^{-s\sqrt{-\lambda}} \left| -F_2(s) - 2(F_1''(s) - F_1(s)) + \frac{\lambda}{\alpha_1^2} F_1''(s) \right| ds \\
&\quad + \int_0^\omega e^{-(\omega-s)\sqrt{-\lambda}} \left| -F_2(s) - 2(F_1''(s) - F_1(s)) + \frac{\lambda}{\alpha_1^2} F_1''(s) \right| ds \\
&\leq \left(\int_0^\omega e^{-q(\omega-s)\sqrt{-\lambda}} ds \right)^{1/q} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right) \\
&\quad + \left(\int_0^\omega e^{-sq\sqrt{-\lambda}} ds \right)^{1/q} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right) \\
&\leq \frac{2(1 - e^{-\omega q\sqrt{-\lambda}})^{1/q}}{q^{1/q}\sqrt{-\lambda}^{1/q}} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right) \\
&\leq \frac{2}{\sqrt{-\lambda}^{1-1/p}} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right).
\end{aligned}$$

3. From (16), (24) and the fact that $|\alpha_1| = |\alpha_2| = \sqrt{1-\lambda} > \sqrt{-\lambda}$, we have

$$\begin{aligned}
\|v\|_{L^p(0,\omega)} &\leq \frac{|I(0)| + |I(\omega)|}{2\sqrt{1-\lambda}(1 - e^{-2\omega\sqrt{\varepsilon_0}})} \left(\int_0^\omega e^{-p\theta\sqrt{-\lambda}} d\theta \right)^{1/p} + \frac{-\lambda}{(1-\lambda)^2} \|F_1''\|_{L^p(0,\omega)} \\
&\quad + \frac{|I(0)| + |I(\omega)|}{2\sqrt{1-\lambda}(1 - e^{-2\omega\sqrt{\varepsilon_0}})} \left(\int_0^\omega e^{-p(\omega-\theta)\sqrt{-\lambda}} d\theta \right)^{1/p} + \frac{1}{2\sqrt{1-\lambda}} \|I\|_{L^p(0,\omega)} \\
&\leq \frac{(|I(0)| + |I(\omega)|)}{\sqrt{1-\lambda}\sqrt{-\lambda}^{1/p} p^{1/p} (1 - e^{-2\omega\sqrt{\varepsilon_0}})} + \frac{1}{2\sqrt{1-\lambda}} \|I\|_{L^p(0,\omega)} \\
&\quad + \frac{1}{1-\lambda} \|F_1''\|_{L^p(0,\omega)},
\end{aligned}$$

hence

$$\begin{aligned}
\|v\|_{L^p(0,\omega)} &\leq \frac{2 \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right)}{\sqrt{1-\lambda}\sqrt{-\lambda}^{1/p+1/q} q^{1/q} p^{1/p} (1 - e^{-2\omega\sqrt{\varepsilon_0}})} \\
&\quad + \frac{\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)}}{\sqrt{1-\lambda}\sqrt{-\lambda}} + \frac{1}{1-\lambda} \|F_1''\|_{L^p(0,\omega)} \\
&\leq \frac{M_1}{\sqrt{1-\lambda}\sqrt{-\lambda}} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right).
\end{aligned}$$

4. From (16) and (23), we have

$$\begin{aligned}
\|J\|_{L^p(0,\omega)} &\leq \sup_{\theta \in [0,\omega]} \left(\int_0^\theta e^{-(\theta-s)\sqrt{-\lambda}} ds + \int_\theta^\omega e^{-(s-\theta)\sqrt{-\lambda}} ds \right) \|v\|_{L^p(0,\omega)} \\
&\leq \sup_{\theta \in [0,\omega]} \left(\frac{1 - e^{-\theta\sqrt{-\lambda}}}{\sqrt{-\lambda}} + \frac{1 - e^{-(\omega-\theta)\sqrt{-\lambda}}}{\sqrt{-\lambda}} \right) \|v\|_{L^p(0,\omega)} \\
&\leq \frac{2}{\sqrt{-\lambda}} \|v\|_{L^p(0,\omega)}.
\end{aligned}$$

5. Due to (16), (23) and the Hölder inequality, we deduce that

$$\begin{aligned}
|J(0)| + |J(\omega)| &\leq \int_0^\omega e^{-s\sqrt{-\lambda}} |v(s)| ds + \int_0^\omega e^{-(\omega-s)\sqrt{-\lambda}} |v(s)| ds \\
&\leq \left(\left(\int_0^\omega e^{-sq\sqrt{-\lambda}} ds \right)^{1/q} + \left(\int_0^\omega e^{-(\omega-s)q\sqrt{-\lambda}} ds \right)^{1/q} \right) \|v\|_{L^p(0,\omega)} \\
&\leq \frac{2 \left(1 - e^{-\omega q \sqrt{-\lambda}} \right)^{1/q}}{q^{1/q} \sqrt{-\lambda}^{1/q}} \|v\|_{L^p(0,\omega)} \\
&\leq \frac{2}{\sqrt{-\lambda}^{1-1/p}} \|v\|_{L^p(0,\omega)}.
\end{aligned}$$

□

Lemma 3.7. Let $\lambda < 0$. Then, we have

$$\left(\int_0^\omega \left| e^{-\theta\alpha_1} - e^{-\theta\alpha_2} \right|^p d\theta \right)^{1/p} \leq \frac{4}{\sqrt{-\lambda}^{1+1/p}},$$

and

$$\left(\int_0^\omega \left| e^{-(\omega-\theta)\alpha_1} - e^{-(\omega-\theta)\alpha_2} \right|^p d\theta \right)^{1/p} \leq \frac{4}{\sqrt{-\lambda}^{1+1/p}}.$$

Proof. For $x \geq 0$, we have $e^{-\frac{px}{2}} x^p < 1$, so

$$\int_0^{+\infty} e^{-px} x^p dx = \int_0^{+\infty} e^{-\frac{px}{2}} e^{-\frac{px}{2}} x^p dx \leq \int_0^{+\infty} e^{-\frac{px}{2}} dx = \frac{2}{p}.$$

Then, from (16), we have

$$\int_0^\omega \left| e^{-\theta\alpha_1} - e^{-\theta\alpha_2} \right|^p d\theta = \int_0^\omega \left| e^{-\theta\sqrt{-\lambda}} \left(e^{-\theta i} - e^{\theta i} \right) \right|^p d\theta = 2^p \int_0^\omega e^{-p\theta\sqrt{-\lambda}} |\sin(\theta)|^p d\theta,$$

hence, setting $x = \theta\sqrt{-\lambda}$, it follows that

$$\begin{aligned}
\int_0^\omega \left| e^{-\theta\alpha_1} - e^{-\theta\alpha_2} \right|^p d\theta &= 2^p \int_0^{\omega\sqrt{-\lambda}} e^{-px} \left| \sin\left(\frac{x}{\sqrt{-\lambda}}\right) \right|^p \frac{dx}{\sqrt{-\lambda}} \\
&\leq \frac{2^p}{\sqrt{-\lambda}} \int_0^{\omega\sqrt{-\lambda}} e^{-px} \left(\frac{x}{\sqrt{-\lambda}} \right)^p dx \\
&\leq \frac{2^p}{\sqrt{-\lambda}^{p+1}} \int_0^{+\infty} e^{-px} x^p dx \\
&\leq \frac{2^{p+1}}{p\sqrt{-\lambda}^{p+1}} \leq \frac{2^{2p}}{\sqrt{-\lambda}^{p+1}}.
\end{aligned}$$

The second estimate is obtained by change of variable, taking $\omega - \theta$ instead of θ . □

Lemma 3.8. For all $\lambda \in (-\infty, -\varepsilon_0]$, where ε_0 is defined in Proposition 3.2, the constants $\beta_1, \beta_2, \beta_3$ and β_4 , defined by (20), satisfy

$$\max(|\beta_1 + \beta_2|, |\beta_3 + \beta_4|) \leq \frac{M_1 \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right)}{\omega(-\lambda)\sqrt{-\lambda}^{2-1/p} \left(1 - e^{-2\omega\sqrt{\varepsilon_0}} - 2\omega\sqrt{\varepsilon_0} e^{-\omega\sqrt{\varepsilon_0}} \right) \left(1 - e^{-\omega\sqrt{\varepsilon_0}} \right)},$$

and

$$\max(|\beta_2|, |\beta_4|) \leq \frac{M_1 \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right)}{2(-\lambda)\sqrt{-\lambda}^{1-1/p} \left(1 - e^{-2\omega\sqrt{\varepsilon_0}} - 2\omega\sqrt{\varepsilon_0} e^{-\omega\sqrt{\varepsilon_0}} \right)},$$

where $M_1 = 2 + \frac{2}{1 - e^{-2\omega\sqrt{\varepsilon_0}}}$.

Proof. Recall that β_i , $i = 1, 2, 3, 4$, depends on U_1^{-1} and U_2^{-1} . From (33) and (34), it follows

$$U_1 \geq h(\omega\sqrt{-\lambda}) \geq h(\omega\sqrt{\varepsilon_0}) = 1 - e^{-2\omega\sqrt{\varepsilon_0}} - 2\omega\sqrt{\varepsilon_0} e^{-\omega\sqrt{\varepsilon_0}} > 0,$$

and

$$U_2 \geq h(\omega\sqrt{-\lambda}) \geq h(\omega\sqrt{\varepsilon_0}) = 1 - e^{-2\omega\sqrt{\varepsilon_0}} - 2\omega\sqrt{\varepsilon_0} e^{-\omega\sqrt{\varepsilon_0}} > 0.$$

Thus, we deduce

$$U_1^{-1} \leq \frac{1}{1 - e^{-2\omega\sqrt{\varepsilon_0}} - 2\omega\sqrt{\varepsilon_0} e^{-\omega\sqrt{\varepsilon_0}}} \quad \text{and} \quad U_2^{-1} \leq \frac{1}{1 - e^{-2\omega\sqrt{\varepsilon_0}} - 2\omega\sqrt{\varepsilon_0} e^{-\omega\sqrt{\varepsilon_0}}}.$$

Therefore, from (20) and Lemma 3.6, we have

$$\begin{aligned} |\beta_1 + \beta_2| &\leq \frac{|J(0)| + |J(\omega)|}{4 \left(1 - e^{-2\omega\sqrt{\varepsilon_0}} - 2\omega\sqrt{\varepsilon_0} e^{-\omega\sqrt{\varepsilon_0}} \right)} \left| \frac{1 - e^{-\omega\alpha_1}}{1 - e^{-\omega\alpha_2}} - 1 \right| \\ &\leq \frac{\|v\|_{L^p(0,\omega)}}{2\sqrt{-\lambda}^{1-1/p} \left(1 - e^{-2\omega\sqrt{\varepsilon_0}} - 2\omega\sqrt{\varepsilon_0} e^{-\omega\sqrt{\varepsilon_0}} \right)} \left| \frac{e^{-\omega\alpha_2} - e^{-\omega\alpha_1}}{1 - e^{-\omega\alpha_2}} \right| \\ &\leq \frac{M_1 \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right)}{2(-\lambda)\sqrt{-\lambda}^{1-1/p} \left(1 - e^{-2\omega\sqrt{\varepsilon_0}} - 2\omega\sqrt{\varepsilon_0} e^{-\omega\sqrt{\varepsilon_0}} \right)} \frac{2e^{-\omega\sqrt{-\lambda}}}{1 - e^{-\omega\sqrt{\varepsilon_0}}} \\ &\leq \frac{M_1 \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right)}{\omega(-\lambda)\sqrt{-\lambda}^{2-1/p} \left(1 - e^{-2\omega\sqrt{\varepsilon_0}} - 2\omega\sqrt{\varepsilon_0} e^{-\omega\sqrt{\varepsilon_0}} \right) \left(1 - e^{-\omega\sqrt{\varepsilon_0}} \right)} \end{aligned}$$

and similarly

$$\begin{aligned} |\beta_3 + \beta_4| &\leq \frac{|J(0)| + |J(\omega)|}{4 \left(1 - e^{-2\omega\sqrt{\varepsilon_0}} - 2\omega\sqrt{\varepsilon_0} e^{-\omega\sqrt{\varepsilon_0}} \right)} \left| 1 - \frac{1 + e^{-\omega\alpha_1}}{1 + e^{-\omega\alpha_2}} \right| \\ &\leq \frac{M_1 \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right)}{\omega(-\lambda)\sqrt{-\lambda}^{2-1/p} \left(1 - e^{-2\omega\sqrt{\varepsilon_0}} - 2\omega\sqrt{\varepsilon_0} e^{-\omega\sqrt{\varepsilon_0}} \right) \left(1 - e^{-\omega\sqrt{\varepsilon_0}} \right)}. \end{aligned}$$

In the same way, we obtain

$$|\beta_2| \leq \frac{M_1 \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right)}{2(-\lambda)\sqrt{-\lambda}^{1-1/p} \left(1 - e^{-2\omega\sqrt{\varepsilon_0}} - 2\omega\sqrt{\varepsilon_0} e^{-\omega\sqrt{\varepsilon_0}} \right)},$$

and

$$|\beta_4| \leq \frac{M_1 \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right)}{2(-\lambda)\sqrt{-\lambda}^{1-1/p} \left(1 - e^{-2\omega\sqrt{\varepsilon_0}} - 2\omega\sqrt{\varepsilon_0} e^{-\omega\sqrt{\varepsilon_0}} \right)}.$$

□

Proposition 3.9. \mathcal{A} is closed and densely defined in X . Moreover, there exists a constant $M > 0$ such that for all $\lambda \leq 0$, operator $\mathcal{A} - \lambda I$ is invertible with bounded inverse and

$$\|(\mathcal{A} - \lambda I)^{-1}\|_{\mathcal{L}(X)} \leq \frac{M}{1 + |\lambda|}.$$

Remark 3.10. This result implies, in particular, that $-\sqrt{\mathcal{A}}$ is well-defined and generates a uniformly bounded analytic semigroup $(e^{-s\sqrt{\mathcal{A}}})_{s \geq 0}$.

Proof. Recall that

$$D(\mathcal{A}) = [W^{4,p}(0, \omega) \cap W_0^{2,p}(0, \omega)] \times W_0^{2,p}(0, \omega).$$

It is clear that

$$\mathcal{D}(0, \omega) \times \mathcal{D}(0, \omega) \subset D(\mathcal{A}) \subset X = W_0^{2,p}(0, \omega) \times L^p(0, \omega),$$

where $\mathcal{D}(0, \omega)$ is the set of C^∞ -functions with compact support in $(0, \omega)$. Since $\mathcal{D}(0, \omega)$ is dense in each spaces $W_0^{2,p}(0, \omega)$ and $L^p(0, \omega)$ for their respective norms, then $D(\mathcal{A})$ is dense in X .

From Proposition 3.2, $0 \in \rho(\mathcal{A})$, thus \mathcal{A} is closed. From Proposition 3.4, for all $\lambda < 0$, there exists a unique couple

$$(\psi_1, \psi_2) \in (W^{4,p}(0, \omega) \cap W_0^{2,p}(0, \omega)) \times W_0^{2,p}(0, \omega)$$

which satisfies

$$\begin{cases} \psi_2 & = \lambda \psi_1 + F_1 \\ \psi_1^{(4)} + 2(\lambda + 1)\psi_1'' + (\lambda - 1)^2\psi_1 & = G_\lambda, \end{cases} \quad (35)$$

where $G_\lambda = -F_2 - 2(F_1'' - F_1) - \lambda F_1$. Then $\mathbb{R}_- \subset \rho(\mathcal{A})$ and

$$\Psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = (\mathcal{A} - \lambda I)^{-1} \begin{pmatrix} F_1 \\ F_2 \end{pmatrix} = (\mathcal{A} - \lambda I)^{-1} F,$$

where ψ_1 is given by (18)-(20)-(22) and

$$\begin{aligned} \psi_2(\theta) &:= \lambda e^{-\theta\alpha_2}(\beta_1 + \beta_2 + \beta_3 + \beta_4) + \lambda e^{-(\omega-\theta)\alpha_2}(\beta_3 + \beta_4 - \beta_1 - \beta_2) \\ &+ \lambda(e^{-\theta\alpha_1} - e^{-\theta\alpha_2})(\beta_2 + \beta_4) + \lambda(e^{-(\omega-\theta)\alpha_1} - e^{-(\omega-\theta)\alpha_2})(\beta_4 - \beta_2) \\ &+ \lambda S(\theta) + F_1(\theta), \end{aligned} \quad (36)$$

with β_i , $i = 1, 2, 3, 4$ are defined in (20)-(21). From (22), we have

$$\begin{aligned} \lambda S(\theta) + F_1(\theta) &= \frac{\lambda}{2\alpha_2(1 - e^{-2\omega\alpha_2})} e^{-\theta\alpha_2} (J(0) - e^{-\omega\alpha_2} J(\omega)) \\ &+ \frac{\lambda}{2\alpha_2(1 - e^{-2\omega\alpha_2})} e^{-(\omega-\theta)\alpha_2} (J(\omega) - e^{-\omega\alpha_2} J(0)) \\ &- \frac{\lambda^2}{\alpha_1^2 \alpha_2^2} F_1(\theta) + F_1(\theta) - \frac{\lambda}{2\alpha_2} J(\theta), \end{aligned} \quad (37)$$

where $J(\theta)$ is given by (23). Our aim is to prove that, for all $\lambda \leq 0$, there exists $M > 0$, such that

$$\|(\mathcal{A} - \lambda I)^{-1} F\|_{\mathcal{L}(X)} \leq \frac{M}{1 + |\lambda|} \|F\|_X,$$

with

$$\|F\|_X = \left\| \begin{pmatrix} F_1 \\ F_2 \end{pmatrix} \right\|_X = \|F_1\|_{W_0^{2,p}(0, \omega)} + \|F_2\|_{L^p(0, \omega)}. \quad (38)$$

To this end, we consider that $\lambda \in (-\infty, -\varepsilon_0]$, where ε_0 is defined in Proposition 3.2. We first study S'' and ψ_1'' . From (22), for *a.e.* $\theta \in [0, \omega]$, we have

$$\begin{aligned} S''(\theta) &= \frac{\alpha_2 e^{-\theta\alpha_2}}{2(1 - e^{-2\omega\alpha_2})} (J(0) - e^{-\omega\alpha_2}J(\omega)) - \frac{\lambda}{\alpha_1^2\alpha_2^2} F_1''(\theta) \\ &\quad + \frac{\alpha_2 e^{-(\omega-\theta)\alpha_2}}{2(1 - e^{-2\omega\alpha_2})} (J(\omega) - e^{-\omega\alpha_2}J(0)) - \frac{1}{2\alpha_2} J''(\theta), \end{aligned}$$

and from (23), we obtain $J''(\theta) = \alpha_2^2 J(\theta) - 2\alpha_2 v(\theta)$, hence

$$\begin{aligned} S''(\theta) &= \frac{\alpha_2 e^{-\theta\alpha_2}}{2(1 - e^{-2\omega\alpha_2})} (J(0) - e^{-\omega\alpha_2}J(\omega)) - \frac{\lambda}{\alpha_1^2\alpha_2^2} F_1''(\theta) \\ &\quad + \frac{\alpha_2 e^{-(\omega-\theta)\alpha_2}}{2(1 - e^{-2\omega\alpha_2})} (J(\omega) - e^{-\omega\alpha_2}J(0)) - \frac{\alpha_2}{2} J(\theta) + v(\theta). \end{aligned}$$

Then, since $\alpha_1 = \sqrt{-\lambda} + i$ and $\alpha_2 = \sqrt{-\lambda} - i$, we have $|e^{-\omega\alpha_1}| = |e^{-\omega\alpha_2}| = e^{-\omega\sqrt{-\lambda}} \leq 1$ with $-\lambda \geq \varepsilon_0$, thus

$$\begin{aligned} \|S''\|_{L^p(0,\omega)} &\leq \frac{\sqrt{1-\lambda} (|J(0)| + |J(\omega)|)}{2(1 - e^{-2\omega\sqrt{\varepsilon_0}})} \left(\int_0^\omega e^{-p\theta\sqrt{-\lambda}} d\theta \right)^{1/p} \\ &\quad + \frac{\sqrt{1-\lambda} (|J(0)| + |J(\omega)|)}{2(1 - e^{-2\omega\sqrt{\varepsilon_0}})} \left(\int_0^\omega e^{-p(\omega-\theta)\sqrt{-\lambda}} d\theta \right)^{1/p} \\ &\quad + \frac{-\lambda}{(1-\lambda)^2} \|F_1''\|_{L^p(0,\omega)} + \frac{\sqrt{1-\lambda}}{2} \|J\|_{L^p(0,\omega)} + \|v\|_{L^p(0,\omega)} \\ &\leq \frac{\sqrt{1-\lambda} (|J(0)| + |J(\omega)|)}{\sqrt{-\lambda}^{1/p} (1 - e^{-2\omega\sqrt{\varepsilon_0}})} + \frac{1}{1-\lambda} \|F_1''\|_{L^p(0,\omega)} \\ &\quad + \frac{\sqrt{1-\lambda}}{2} \|J\|_{L^p(0,\omega)} + \|v\|_{L^p(0,\omega)}. \end{aligned}$$

From Lemma 3.6, we have

$$\begin{aligned} \|S''\|_{L^p(0,\omega)} &\leq \frac{2\sqrt{1-\lambda}}{\sqrt{-\lambda}^{1-1/p+1/p} (1 - e^{-2\omega\sqrt{\varepsilon_0}})} \|v\|_{L^p(0,\omega)} \\ &\quad + \frac{\sqrt{1-\lambda}}{\sqrt{-\lambda}} \|v\|_{L^p(0,\omega)} + \|v\|_{L^p(0,\omega)} + \frac{1}{1-\lambda} \|F_1''\|_{L^p(0,\omega)} \\ &\leq \frac{2M_1}{-\lambda (1 - e^{-2\omega\sqrt{\varepsilon_0}})} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right) \\ &\quad + \frac{2M_1}{-\lambda} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right) + \frac{1}{1-\lambda} \|F_1''\|_{L^p(0,\omega)}. \end{aligned}$$

Finally, we obtain

$$\|S''\|_{L^p(0,\omega)} \leq \frac{M_2}{-\lambda} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right), \quad (39)$$

where $M_2 = \frac{2M_1}{1 - e^{-2\omega\sqrt{\varepsilon_0}}} + 2M_1 + 1$.

Now, we have

$$\begin{aligned}\psi_1''(\theta) - S''(\theta) &= \alpha_2^2 e^{-\theta\alpha_2}(\beta_1 + \beta_2 + \beta_3 + \beta_4) + \alpha_2^2 e^{-(\omega-\theta)\alpha_2}(\beta_3 + \beta_4 - \beta_1 - \beta_2) \\ &\quad + \alpha_2^2 \left(e^{-\theta\alpha_1} - e^{-\theta\alpha_2} \right) (\beta_2 + \beta_4) + \alpha_2^2 \left(e^{-(\omega-\theta)\alpha_1} - e^{-(\omega-\theta)\alpha_2} \right) (\beta_4 - \beta_2).\end{aligned}$$

Then

$$\begin{aligned}\|\psi_1'' - S''\|_{L^p(0,\omega)} &\leq (1-\lambda)(|\beta_1 + \beta_2| + |\beta_3 + \beta_4|) \left(\int_0^\omega e^{-p\theta\sqrt{-\lambda}} d\theta \right)^{1/p} \\ &\quad + (1-\lambda)(|\beta_1 + \beta_2| + |\beta_3 + \beta_4|) \left(\int_0^\omega e^{-p(\omega-\theta)\sqrt{-\lambda}} d\theta \right)^{1/p} \\ &\quad + (1-\lambda)(|\beta_2| + |\beta_4|) \left(\int_0^\omega |e^{-\theta\alpha_1} - e^{-\theta\alpha_2}|^p d\theta \right)^{1/p} \\ &\quad + (1-\lambda)(|\beta_2| + |\beta_4|) \left(\int_0^\omega |e^{-(\omega-\theta)\alpha_1} - e^{-(\omega-\theta)\alpha_2}|^p d\theta \right)^{1/p}.\end{aligned}$$

Using the fact that

$$\frac{1-\lambda}{-\lambda} = 1 + \frac{1}{-\lambda} \leq 1 + \frac{1}{\varepsilon_0},$$

with Lemma 3.7 and Lemma 3.8, we obtain

$$\begin{aligned}\|\psi_1'' - S''\|_{L^p(0,\omega)} &\leq \frac{2(1-\lambda)(|\beta_1 + \beta_2| + |\beta_3 + \beta_4|)}{\sqrt{-\lambda}^{1/p}} + \frac{8(1-\lambda)(|\beta_2| + |\beta_4|)}{\sqrt{-\lambda}^{1+1/p}} \\ &\leq \frac{M_3 \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right)}{-\lambda},\end{aligned}$$

where $M_3 = \frac{4M_1(1+\frac{1}{\omega})\left(1+\frac{1}{\varepsilon_0}\right)}{(1-e^{-2\omega\sqrt{\varepsilon_0}}-2\omega\sqrt{\varepsilon_0}e^{-\omega\sqrt{\varepsilon_0}})(1-e^{-\omega\sqrt{\varepsilon_0}})}$.

Due to (39), it follows that

$$\begin{aligned}\|\psi_1''\|_{L^p(0,\omega)} &\leq \|\psi_1'' - S''\|_{L^p(0,\omega)} + \|S''\|_{L^p(0,\omega)} \\ &\leq \frac{M_3 + M_2}{-\lambda} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right) \\ &\leq \frac{3(M_2 + M_3)}{-\lambda} \|F\|_X.\end{aligned}$$

From the Poincaré inequality, there exists $C_\omega > 0$ such that

$$\|\psi_1\|_{W_0^{2,p}(0,\omega)} \leq C_\omega \|\psi_1''\|_{L^p(0,\omega)} \leq \frac{3C_\omega(M_2 + M_3)}{-\lambda} \|F\|_X. \quad (40)$$

Now, we focus ourselves on $\|\psi_2\|_{L^p(0,\omega)}$. As previously, we obtain

$$\begin{aligned}\|\psi_2\|_{L^p(0,\omega)} &\leq |\lambda|(|\beta_1 + \beta_2| + |\beta_3 + \beta_4|) \left(\int_0^\omega e^{-p\theta\sqrt{-\lambda}} d\theta \right)^{1/p} \\ &\quad + |\lambda|(|\beta_1 + \beta_2| + |\beta_3 + \beta_4|) \left(\int_0^\omega e^{-p(\omega-\theta)\sqrt{-\lambda}} d\theta \right)^{1/p} \\ &\quad + |\lambda|(|\beta_2| + |\beta_4|) \left(\int_0^\omega |e^{-\theta\alpha_1} - e^{-\theta\alpha_2}|^p d\theta \right)^{1/p} \\ &\quad + |\lambda|(|\beta_4| + |\beta_2|) \left(\int_0^\omega |e^{-(\omega-\theta)\alpha_1} - e^{-(\omega-\theta)\alpha_2}|^p d\theta \right)^{1/p} + \|\lambda S + F_1\|_{L^p(0,\omega)} \\ &\leq \frac{M_2 + M_3}{-\lambda} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right) \\ &\quad + \|\lambda S + F_1\|_{L^p(0,\omega)}.\end{aligned}$$

Moreover, from (37) and Lemma 3.6, we deduce that

$$\begin{aligned}
\|\lambda S + F_1\|_{L^p(0,\omega)} &\leq \frac{-\lambda(|J(0)| + |J(\omega)|)}{2\sqrt{1-\lambda}(1 - e^{-2\omega\sqrt{\varepsilon_0}})} \left(\int_0^\omega e^{-p\theta\sqrt{-\lambda}} d\theta \right)^{1/p} \\
&\quad + \frac{-\lambda(|J(0)| + |J(\omega)|)}{2\sqrt{1-\lambda}(1 - e^{-2\omega\sqrt{\varepsilon_0}})} \left(\int_0^\omega e^{-p(\omega-\theta)\sqrt{-\lambda}} d\theta \right)^{1/p} \\
&\quad + \left| 1 - \frac{\lambda^2}{\alpha_1^2 \alpha_2^2} \right| \|F_1\|_{L^p(0,\omega)} + \frac{-\lambda}{2\sqrt{1-\lambda}} \|J\|_{L^p(0,\omega)} \\
&\leq \frac{-\lambda(|J(0)| + |J(\omega)|)}{\sqrt{-\lambda}^{1+1/p}(1 - e^{-2\omega\sqrt{\varepsilon_0}})} + \left| 1 - \frac{\lambda^2}{(1-\lambda)^2} \right| \|F_1\|_{L^p(0,\omega)} + \|v\|_{L^p(0,\omega)} \\
&\leq \frac{-2\lambda}{\sqrt{-\lambda}^2(1 - e^{-2\omega\sqrt{\varepsilon_0}})} \|v\|_{L^p(0,\omega)} + \frac{1-2\lambda}{(1-\lambda)^2} \|F_1\|_{L^p(0,\omega)} + \|v\|_{L^p(0,\omega)} \\
&\leq \left(\frac{2}{1 - e^{-2\omega\sqrt{\varepsilon_0}}} + 1 \right) \|v\|_{L^p(0,\omega)} + \left(\frac{1}{\lambda^2} + \frac{1}{-\lambda} \right) \|F_1\|_{L^p(0,\omega)} \\
&\leq \left(\frac{2}{1 - e^{-2\omega\sqrt{\varepsilon_0}}} + 1 \right) \|v\|_{L^p(0,\omega)} + \frac{\frac{1}{\varepsilon_0} + 1}{-\lambda} \|F_1\|_{L^p(0,\omega)}.
\end{aligned}$$

Then, from Lemma 3.6, we obtain

$$\|\lambda S + F_1\|_{L^p(0,\omega)} \leq \frac{M_4}{-\lambda} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right),$$

where $M_4 = \left(\frac{2}{1 - e^{-2\omega\sqrt{\varepsilon_0}}} + 1 \right) M_1 + \frac{1}{\varepsilon_0} + 1$.

Thus, it follows that

$$\begin{aligned}
\|\psi_2\|_{L^p(0,\omega)} &\leq \frac{M_2 + M_3 + M_4}{-\lambda} \left(\|F_2\|_{L^p(0,\omega)} + 2\|F_1\|_{L^p(0,\omega)} + 3\|F_1''\|_{L^p(0,\omega)} \right) \\
&\leq \frac{3(M_2 + M_3 + M_4)}{-\lambda} \|F\|_X.
\end{aligned}$$

Finally, from (40), we have

$$\|(\mathcal{A} - \lambda I)^{-1} F\|_X = \|\psi_1\|_{W_0^{2,p}(0,\omega)} + \|\psi_2\|_{L^p(0,\omega)} \leq \frac{M}{|\lambda|} \|F\|_X,$$

where $M = 3((C_\omega + 1)(M_2 + M_3) + M_4)$. □

Since $-\mathcal{A}$ is the realization of \mathcal{L}_2 , we deduce the following corollary.

Corollary 3.11. There exist $\varepsilon_{\mathcal{L}_2} \in (0, \pi)$ small enough and $M_{\mathcal{L}_2} > 0$ such that

$$\forall z \in \Sigma_{\mathcal{L}_2} := \overline{B(0, \varepsilon_0)} \cup \{z \in \mathbb{C} \setminus \{0\} : |\arg(z)| \leq \varepsilon_{\mathcal{L}_2}\},$$

we have

$$\left\| (\mathcal{L}_2 - zI)^{-1} \right\|_{\mathcal{L}(X)} \leq \frac{M_{\mathcal{L}_2}}{1 + |z|}.$$

Therefore, assumption (H_1) in Section 2.2 is verified for \mathcal{L}_2 with

$$\theta_{\mathcal{L}_2} = \pi - \varepsilon_{\mathcal{L}_2}. \tag{41}$$

Remark 3.12. \mathcal{A} is anti-compact; since $\sigma(-\mathcal{L}_2) = \sigma(\mathcal{A})$ then $\sigma(-\mathcal{L}_2)$ is uniquely composed by isolated eigenvalues $(\lambda_j)_{j \geq 1}$ such that $|\lambda_j| \rightarrow +\infty$, see Kato [10], Theorem 6.29, p. 187. More precisely, the calculus of the resolvent operator $(\mathcal{A} - \lambda I)^{-1}$ requires that, for all $\lambda \in \mathbb{C} \setminus \mathbb{R}_+$, U_1 and U_2 defined by (21) do not vanish. Since $U_1 U_2 = 0$ is equivalent to

$$\left(\sinh(\omega\sqrt{-\lambda}) - \omega\sqrt{-\lambda} \right) \left(\sinh(\omega\sqrt{-\lambda}) + \omega\sqrt{-\lambda} \right) = 0,$$

then, using $(z_j)_{j \geq 1}$ defined in Section 1, we deduce that

$$\forall j \geq 1, \quad \lambda_j = -\frac{z_j^2}{\omega^2} \in \mathbb{C} \setminus \mathbb{R}_+.$$

Now, we prove that operator \mathcal{A} has Bounded Imaginary Powers, see Definition 2.4.

Proposition 3.13. $\mathcal{A} \in \text{BIP}(X, \theta_{\mathcal{A}})$, for any $\theta_{\mathcal{A}} \in (0, \pi)$.

Proof. We will be inspired by the method used in Labbas and Moussaoui [16] or in Labbas and Sadallah [17].

Let $\varepsilon > 0$ and $r \in \mathbb{R}$. For all $\lambda > 0$, $F_1 \in W_0^{2,p}(0, \omega)$ and $F_2 \in L^p(0, \omega)$, we have

$$\begin{aligned} \left[(\mathcal{A} + I)^{-\varepsilon+ir} \begin{pmatrix} F_1 \\ F_2 \end{pmatrix} \right] (\theta) &= \frac{1}{\Gamma_{\varepsilon,r}} \int_0^{+\infty} \lambda^{-\varepsilon+ir} \left[(\mathcal{A} + I + \lambda I)^{-1} \begin{pmatrix} F_1 \\ F_2 \end{pmatrix} \right] (\theta) d\lambda \\ &= \frac{1}{\Gamma_{\varepsilon,r}} \int_0^{+\infty} \lambda^{-\varepsilon+ir} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} (\theta) d\lambda \end{aligned}$$

where $\Gamma_{\varepsilon,r} = \Gamma(1 - \varepsilon + ir)\Gamma(\varepsilon - ir)$, see for instance Triebel [23], (6), p. 100.

Now, let us focus ourselves on the first component ψ_1 for instance. Due to Seeley [21], we only consider the convolution term in ψ_1 , which is the most singular term. In our case, this term is given by

$$\begin{aligned} I_S(\theta) &= \frac{1}{4\alpha_1\alpha_2} \int_0^\omega e^{-|\theta-s|\alpha_2} \int_0^\omega e^{-|s-t|\alpha_1} G_{\lambda+1}(t) dt ds \\ &= \frac{1}{4\alpha_1\alpha_2} \int_0^\omega e^{-|\theta-s|\alpha_2} \int_0^\omega e^{-|s-t|\alpha_1} (-F_2 - 2(F_1'' - F_1) + (\lambda + 1)F_1)(t) dt ds, \end{aligned}$$

see (26) and (27). We will use the two following extensions

$$\widetilde{G}_0(\theta) = \begin{cases} -F_2(\theta) - 2F_1''(\theta) + 2F_1(\theta), & \text{if } \theta \in [0, \omega] \\ 0, & \text{else,} \end{cases} \quad \text{with} \quad \widetilde{F}_1(\theta) = \begin{cases} F_1(\theta), & \text{if } \theta \in [0, \omega] \\ 0, & \text{else,} \end{cases}$$

and

$$E_\alpha(\theta) = e^{-|\theta|\alpha}.$$

Now, we will use the Fourier transform denoted by \mathcal{F}_t . We then have

$$\begin{aligned} I_{\varepsilon,r}(\theta) &:= \frac{1}{\Gamma_{\varepsilon,r}} \int_0^{+\infty} \lambda^{-\varepsilon+ir} I_S(\theta) d\lambda \\ &= \frac{1}{\Gamma_{\varepsilon,r}} \int_0^{+\infty} \frac{\lambda^{-\varepsilon+ir}}{4\alpha_1\alpha_2} \left(E_{\alpha_2} \star \left(E_{\alpha_1} \star \left(\widetilde{G}_0 + (\lambda + 1)\widetilde{F}_1 \right) \right) \right) (\theta) d\lambda \\ &= \frac{1}{\Gamma_{\varepsilon,r}} \int_0^{+\infty} \frac{\lambda^{-\varepsilon+ir}}{4\alpha_1\alpha_2} \mathcal{F}_t^{-1} \left(\mathcal{F}_t \left(E_{\alpha_2} \star \left(E_{\alpha_1} \star \left(\widetilde{G}_0 + (\lambda + 1)\widetilde{F}_1 \right) \right) \right) (\xi) \right) (\theta) d\lambda, \end{aligned}$$

hence

$$\begin{aligned}
I_{\varepsilon,r}(\theta) &= \mathcal{F}_t^{-1} \left(\frac{1}{\Gamma_{\varepsilon,r}} \int_0^{+\infty} \frac{\lambda^{-\varepsilon+ir}}{4\alpha_1\alpha_2} \mathcal{F}_t(E_{\alpha_2})(\xi) \mathcal{F}_t(E_{\alpha_1})(\xi) \mathcal{F}_t(\widetilde{G}_0 + (\lambda+1)\widetilde{F}_1)(\xi) d\lambda \right) (\theta) \\
&= \mathcal{F}_t^{-1} \left(\frac{1}{\Gamma_{\varepsilon,r}} \int_0^{+\infty} \frac{\lambda^{-\varepsilon+ir}}{4\alpha_1\alpha_2} \mathcal{F}_t(E_{\alpha_2})(\xi) \mathcal{F}_t(E_{\alpha_1})(\xi) d\lambda \mathcal{F}_t(\widetilde{G}_0)(\xi) \right) (\theta) \\
&\quad + \mathcal{F}_t^{-1} \left(\frac{1}{\Gamma_{\varepsilon,r}} \int_0^{+\infty} \frac{\lambda^{-\varepsilon+ir}(\lambda+1)}{4\alpha_1\alpha_2} \mathcal{F}_t(E_{\alpha_2})(\xi) \mathcal{F}_t(E_{\alpha_1})(\xi) d\lambda \mathcal{F}_t(\widetilde{F}_1)(\xi) \right) (\theta).
\end{aligned}$$

We recall that

$$\mathcal{F}_t(E_\alpha)(\xi) = \frac{2\alpha}{\alpha^2 + 4\pi^2\xi^2},$$

here $\alpha_1 = \sqrt{\lambda+1} + i$ and $\alpha_2 = \sqrt{\lambda+1} - i$. Hence

$$\begin{aligned}
\frac{\lambda^{-\varepsilon+ir}}{4\alpha_1\alpha_2} \mathcal{F}_t(E_{\alpha_2})(\xi) \mathcal{F}_t(E_{\alpha_1})(\xi) &= \frac{\lambda^{-\varepsilon+ir}}{4\alpha_1\alpha_2} \frac{4\alpha_1\alpha_2}{(\alpha_1^2 + 4\pi^2\xi^2)(\alpha_2^2 + 4\pi^2\xi^2)} \\
&= \frac{\lambda^{-\varepsilon+ir}}{\alpha_1^2\alpha_2^2 + 4\pi^2\xi^2(\alpha_1^2 + \alpha_2^2) + 16\pi^4\xi^4} \\
&= \frac{\lambda^{-\varepsilon+ir}}{\lambda^2 + 4(1 + 2\pi^2\xi^2)\lambda + 4(1 + 4\pi^4\xi^4)} \\
&= \frac{\lambda^{-\varepsilon+ir}}{(\lambda + \lambda_1)(\lambda + \lambda_2)},
\end{aligned}$$

where

$$\begin{cases} \lambda_1 = 2 + 4\pi\xi + 4\pi^2\xi^2 = 4\pi^2 \left(\xi - \frac{(1+i)}{2\pi} \right) \left(\xi - \frac{(1-i)}{2\pi} \right) \\ \lambda_2 = 2 - 4\pi\xi + 4\pi^2\xi^2 = 4\pi^2 \left(\xi + \frac{(1+i)}{2\pi} \right) \left(\xi + \frac{(1-i)}{2\pi} \right). \end{cases}$$

Thus, since

$$\frac{1}{(\lambda + \lambda_1)(\lambda + \lambda_2)} = \frac{1}{\lambda_1 - \lambda_2} \left(-\frac{1}{\lambda + \lambda_1} + \frac{1}{\lambda + \lambda_2} \right),$$

it follows that

$$\frac{\lambda^{-\varepsilon+ir}}{4\alpha_1\alpha_2} \mathcal{F}_t(E_{\alpha_2})(\xi) \mathcal{F}_t(E_{\alpha_1})(\xi) = \frac{1}{8\pi\xi} \left(-\frac{\lambda^{-\varepsilon+ir}}{\lambda + \lambda_1} + \frac{\lambda^{-\varepsilon+ir}}{\lambda + \lambda_2} \right).$$

Then, setting

$$\sigma_1 = \frac{\lambda}{\lambda_1} \quad \text{and} \quad \sigma_2 = \frac{\lambda}{\lambda_2},$$

we obtain

$$\begin{aligned}
\int_0^{+\infty} \frac{\lambda^{-\varepsilon+ir}}{4\alpha_1\alpha_2} \mathcal{F}_t(E_{\alpha_2})(\xi) \mathcal{F}_t(E_{\alpha_1})(\xi) d\lambda &= \frac{1}{8\pi\xi} \left(-\int_0^{+\infty} \frac{\lambda^{-\varepsilon+ir}}{\lambda + \lambda_1} d\lambda + \int_0^{+\infty} \frac{\lambda^{-\varepsilon+ir}}{\lambda + \lambda_2} d\lambda \right) \\
&= -\frac{\lambda_1^{-\varepsilon+ir}}{8\pi\xi} \int_0^{+\infty} \frac{\sigma_1^{-\varepsilon+ir}}{\sigma_1 + 1} d\sigma_1 \\
&\quad + \frac{\lambda_2^{-\varepsilon+ir}}{8\pi\xi} \int_0^{+\infty} \frac{\sigma_2^{-\varepsilon+ir}}{\sigma_2 + 1} d\sigma_2.
\end{aligned}$$

Moreover, for all $z \in \mathbb{C} \setminus \mathbb{N}^-$, where \mathbb{N}^- is the set of negative integer, we have

$$\int_0^{+\infty} \frac{\sigma^{-z}}{\sigma + 1} d\sigma = \Gamma(z)\Gamma(1-z). \quad (42)$$

It follows that

$$\int_0^{+\infty} \frac{\sigma^{-\varepsilon+ir}}{\sigma+1} d\sigma = \Gamma(\varepsilon - ir)\Gamma(1 - \varepsilon + ir) = \Gamma_{\varepsilon,r}, \quad (43)$$

hence

$$\begin{aligned} \frac{1}{\Gamma_{\varepsilon,r}} \int_0^{+\infty} \frac{\lambda^{-\varepsilon+ir}}{4\alpha_1\alpha_2} \mathcal{F}_t(E_{\alpha_2})(\xi) \mathcal{F}_t(E_{\alpha_1})(\xi) d\lambda &= \frac{1}{8\pi\xi \Gamma_{\varepsilon,r}} \left(\lambda_2^{-\varepsilon+ir} - \lambda_1^{-\varepsilon+ir} \right) \Gamma_{\varepsilon,r} \\ &= \frac{1}{8\pi\xi} \left(\lambda_2^{-\varepsilon+ir} - \lambda_1^{-\varepsilon+ir} \right). \end{aligned}$$

In the same way, we have

$$\begin{aligned} \frac{\lambda^{-\varepsilon+ir}(\lambda+1)}{4\alpha_1\alpha_2} \mathcal{F}_t(E_{\alpha_2})(\xi) \mathcal{F}_t(E_{\alpha_1})(\xi) &= \frac{1}{\lambda_1 - \lambda_2} \left(-\frac{(\lambda+1)\lambda^{-\varepsilon+ir}}{\lambda + \lambda_1} + \frac{(\lambda+1)\lambda^{-\varepsilon+ir}}{\lambda + \lambda_2} \right) \\ &= \frac{1}{8\pi\xi} \left(-\frac{\lambda^{1-\varepsilon+ir}}{\lambda + \lambda_1} + \frac{\lambda^{1-\varepsilon+ir}}{\lambda + \lambda_2} \right) \\ &\quad + \frac{1}{8\pi\xi} \left(-\frac{\lambda^{-\varepsilon+ir}}{\lambda + \lambda_1} + \frac{\lambda^{-\varepsilon+ir}}{\lambda + \lambda_2} \right). \end{aligned}$$

Then, setting

$$\sigma_1 = \frac{\lambda}{\lambda_1} \quad \text{and} \quad \sigma_2 = \frac{\lambda}{\lambda_2},$$

we obtain

$$\begin{aligned} \Upsilon &= \int_0^{+\infty} \frac{\lambda^{-\varepsilon+ir}(\lambda+1)}{4\alpha_1\alpha_2} \mathcal{F}_t(E_{\alpha_2})(\xi) \mathcal{F}_t(E_{\alpha_1})(\xi) d\lambda \\ &= -\frac{1}{8\pi\xi} \int_0^{+\infty} \frac{\lambda^{-\varepsilon+ir}}{\lambda + \lambda_1} d\lambda + \frac{1}{8\pi\xi} \int_0^{+\infty} \frac{\lambda^{-\varepsilon+ir}}{\lambda + \lambda_2} d\lambda \\ &\quad - \frac{1}{8\pi\xi} \int_0^{+\infty} \frac{\lambda^{1-\varepsilon+ir}}{\lambda + \lambda_1} d\lambda + \frac{1}{8\pi\xi} \int_0^{+\infty} \frac{\lambda^{1-\varepsilon+ir}}{\lambda + \lambda_2} d\lambda \\ &= -\frac{\lambda_1^{-\varepsilon+ir}}{8\pi\xi} \int_0^{+\infty} \frac{\sigma_1^{-\varepsilon+ir}}{\sigma_1+1} d\sigma_1 + \frac{\lambda_2^{-\varepsilon+ir}}{8\pi\xi} \int_0^{+\infty} \frac{\sigma_2^{-\varepsilon+ir}}{\sigma_2+1} d\sigma_2 \\ &\quad - \frac{\lambda_1^{1-\varepsilon+ir}}{8\pi\xi} \int_0^{+\infty} \frac{\sigma_1^{1-\varepsilon+ir}}{\sigma_1+1} d\sigma_1 + \frac{\lambda_2^{1-\varepsilon+ir}}{8\pi\xi} \int_0^{+\infty} \frac{\sigma_2^{1-\varepsilon+ir}}{\sigma_2+1} d\sigma_2. \end{aligned}$$

Moreover, from (42) and (43), we deduce that

$$\begin{aligned} \Upsilon &= \left(\frac{\lambda_2^{-\varepsilon+ir} - \lambda_1^{-\varepsilon+ir}}{8\pi\xi} \right) \Gamma_{\varepsilon,r} + \left(\frac{\lambda_2^{-\varepsilon+ir} - \lambda_1^{-\varepsilon+ir}}{8\pi\xi} \right) \Gamma(\varepsilon - ir - 1)\Gamma(1 - (\varepsilon - ir - 1)) \\ &= \left(\frac{\lambda_2^{-\varepsilon+ir} - \lambda_1^{-\varepsilon+ir}}{2\pi\xi} \right) (\Gamma_{\varepsilon,r} + \Gamma(\varepsilon - ir - 1)\Gamma(1 - (\varepsilon - ir - 1))). \end{aligned}$$

For all $z \in \mathbb{C} \setminus \mathbb{Z}$, we have

$$\Gamma(z-1)\Gamma(1-(z-1)) = \frac{\pi}{\sin(\pi(z-1))} = -\frac{\pi}{\sin(\pi z)} = -\Gamma(z)\Gamma(1-z),$$

Setting $z = \varepsilon - ir$, with $\varepsilon \in (0, 1)$, it follows that

$$\Gamma(\varepsilon - ir - 1)\Gamma(1 - (\varepsilon - ir - 1)) = -\Gamma(\varepsilon - ir)\Gamma(1 - \varepsilon + ir) = -\Gamma_{\varepsilon,r},$$

hence $\Upsilon = 0$. Finally, we obtain that

$$I_{\varepsilon,r}(\theta) = \mathcal{F}_t^{-1} \left(m_\varepsilon(\xi) \mathcal{F}_t \left(\widetilde{G}_0 \right) (\xi) \right) (\theta),$$

where

$$m_\varepsilon(\xi) = \frac{\lambda_2^{-\varepsilon+ir} - \lambda_1^{-\varepsilon+ir}}{8\pi\xi}.$$

Setting

$$m(\xi) := \lim_{\varepsilon \rightarrow 0} m_\varepsilon(\xi) = \frac{\lambda_2^{ir} - \lambda_1^{ir}}{8\pi\xi},$$

due to the Lebesgue's dominated convergence Theorem, it follows that

$$I_{0,r}(\theta) := \lim_{\varepsilon \rightarrow 0} I_{\varepsilon,r}(\theta) = \mathcal{F}_t^{-1} \left(m(\xi) \mathcal{F}_t \left(\widetilde{G}_0 \right) (\xi) \right) (\theta).$$

Moreover, for all $x_1, x_2 \in \mathbb{R}$, we have

$$\left| e^{ix_1} - e^{ix_2} \right| \leq |x_1 - x_2|,$$

then, for all $\xi \in \mathbb{R} \setminus \{0\}$, we deduce that

$$|m(\xi)| = \frac{|\lambda_2^{ir} - \lambda_1^{ir}|}{8\pi|\xi|} = \frac{|e^{ir \ln(\lambda_2)} - e^{ir \ln(\lambda_1)}|}{8\pi|\xi|} \leq \frac{|r| |\ln(\lambda_2) - \ln(\lambda_1)|}{8\pi|\xi|} \leq \frac{|r| \left| \ln \left(\frac{2+\sqrt{2}}{2-\sqrt{2}} \right) \right|}{8\pi\xi}.$$

Thus

$$\sup_{\xi \in \mathbb{R}} |m(\xi)| = \lim_{\xi \rightarrow 0} |m(\xi)| = \left| \lim_{\xi \rightarrow 0} \frac{\lambda_2^{ir} - \lambda_1^{ir}}{8\pi\xi} \right|,$$

and

$$\begin{aligned} \lim_{\xi \rightarrow 0} \frac{\lambda_2^{ir} - \lambda_1^{ir}}{8\pi\xi} &= \lim_{\xi \rightarrow 0} 2^{ir} \left(\frac{1 + ir(-2\pi\xi + 2\pi^2\xi^2) + 2ir(ir-1)\pi^2\xi^2 + o(\xi^2)}{8\pi\xi} \right) \\ &\quad - \lim_{\xi \rightarrow 0} 2^{ir} \left(\frac{1 + ir(2\pi\xi + 2\pi^2\xi^2) + 2ir(ir-1)\pi^2\xi^2 + o(\xi^2)}{8\pi\xi} \right) \\ &= \lim_{\xi \rightarrow 0} 2^{ir} \left(\frac{-4ir\pi\xi + o(\xi^2)}{8\pi\xi} \right) \\ &= -2^{ir-1}ir. \end{aligned}$$

Then

$$\sup_{\xi \in \mathbb{R}} |m(\xi)| = \frac{|r|}{2}.$$

We have

$$\begin{aligned} \xi m'(\xi) &= \frac{2\pi\xi^2 \left(ir\lambda_2^{ir-1}(-4\pi + 8\pi^2\xi) - ir\lambda_1^{ir-1}(4\pi + 8\pi^2\xi) \right) - 2\pi\xi(\lambda_2^{ir} - \lambda_1^{ir})}{64\pi^2\xi^2} \\ &= \frac{ir}{32} \left(\lambda_2^{ir-1}(-1 + 2\pi\xi) - \lambda_1^{ir-1}(1 + 2\pi\xi) \right) - \frac{\lambda_2^{ir} - \lambda_1^{ir}}{32\pi\xi}, \end{aligned}$$

and in the same way we obtain

$$\sup_{\xi \in \mathbb{R}} |\xi m'(\xi)| = \lim_{\xi \rightarrow 0} |\xi m'(\xi)| = \left| \lim_{\xi \rightarrow 0} \xi m'(\xi) \right|,$$

with

$$\lim_{\xi \rightarrow 0} \xi m'(\xi) = \frac{ir}{32} \lim_{\xi \rightarrow 0} \left(\lambda_2^{ir-1}(-1 + 2\pi\xi) - \lambda_1^{ir-1}(1 + 2\pi\xi) \right) - \frac{\lambda_2^{ir} - \lambda_1^{ir}}{32\pi\xi},$$

where

$$\lim_{\xi \rightarrow 0} \lambda_2^{ir-1} = \lim_{\xi \rightarrow 0} 2^{ir-1} \left(1 + (ir-1) \left(-2\pi\xi + 2\pi^2\xi^2 \right) + 4ir(ir-1)\pi^2\xi^2 + o(\xi^2) \right) = 2^{ir-1},$$

and

$$\lim_{\xi \rightarrow 0} \lambda_1^{ir-1} = \lim_{\xi \rightarrow 0} 2^{ir-1} \left(1 + (ir-1) \left(2\pi\xi + 2\pi^2\xi^2 \right) + 4ir(ir-1)\pi^2\xi^2 + o(\xi^2) \right) = 2^{ir-1}.$$

Thus

$$\begin{aligned} \lim_{\xi \rightarrow 0} \xi m'(\xi) &= \frac{1}{4} \lim_{\xi \rightarrow 0} \frac{ir}{8} \left(- \left(\lambda_2^{ir-1} + \lambda_1^{ir-1} \right) + 2\pi\xi \left(\lambda_2^{ir-1} - \lambda_1^{ir-1} \right) \right) - \frac{\lambda_2^{ir} - \lambda_1^{ir}}{8\pi\xi} \\ &= \frac{1}{4} \left(-\frac{2^{ir}ir}{8} + 2^{ir-1}ir \right) \\ &= 3 \times 2^{ir-5}ir. \end{aligned}$$

Then

$$\sup_{\xi \in \mathbb{R}} |\xi m'(\xi)| = \left| 3 \times 2^{ir-5}ir \right| = \frac{3}{32}|r|.$$

Therefore, we deduce that

$$\sup_{\xi \in \mathbb{R}} |m(\xi)| + \sup_{\xi \in \mathbb{R}} |\xi m'(\xi)| = \frac{|r|}{2} + \frac{3}{32}|r| = \frac{19}{32}|r|,$$

From the Mihlin Theorem, see Mihlin [18], for all $\gamma > 0$, there exists $C_{\gamma,p} > 0$, such that, for all $r \in \mathbb{R}$, we have

$$\|I_{0,r}(\cdot)\|_{\mathcal{L}(X)} = \left\| \mathcal{F}_t^{-1} \left(m(\xi) \mathcal{F}_t(\widetilde{G}_0)(\xi) \right) (\cdot) \right\|_{\mathcal{L}(X)} \leq C_{\gamma,p} e^{\gamma|r|}.$$

Finally, for all $\gamma > 0$, there exists a constant $C_{\gamma,p} > 0$ such that for all $r \in \mathbb{R}$, we obtain

$$\left\| (\mathcal{A} + I)^{ir} \right\|_{\mathcal{L}(X)} \leq C_{\gamma,p} e^{\gamma|r|}.$$

Therefore, taking $\theta_{\mathcal{A}} = \gamma > 0$, we have $\mathcal{A} + I \in \text{BIP}(X, \theta_{\mathcal{A}})$ and from Theorem 2.3, p. 69 in Arendt, Bu and Haase [1], we deduce that $\mathcal{A} = \mathcal{A} + I - I \in \text{BIP}(X, \theta_{\mathcal{A}})$. \square

4 Study of the sum $\mathcal{L}_{1,\mu} + \mathcal{L}_2$

4.1 Invertibility of the closure of the sum

In this section, we will apply the results described in Section 2.2. We take $\mathcal{L}_{1,\mu} = \mathcal{M}_1$ and $\mathcal{L}_2 = \mathcal{M}_2$.

Theorem 4.1. Assume that (5) holds. Then $\mathcal{L}_{1,\mu} + \mathcal{L}_2$ is closable and its closure $\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2}$ is invertible.

Proof. Assumption (H_1) is satisfied from Proposition 3.1 and Corollary 3.11, with

$$\theta_{\mathcal{M}_1} + \theta_{\mathcal{M}_2} = \varepsilon_{\mathcal{L}_{1,\mu}} + \pi - \varepsilon_{\mathcal{L}_2},$$

where it suffices to take $\varepsilon_{\mathcal{L}_2} > \varepsilon_{\mathcal{L}_{1,\mu}}$ in order to obtain $\theta_{\mathcal{M}_1} + \theta_{\mathcal{M}_2} < \pi$.

For assumption (H_2) , due to Proposition 3.2, it follows that $0 \notin \sigma(\mathcal{L}_{1,\mu}) \cap \sigma(-\mathcal{L}_2)$. Moreover, from Proposition 3.1, we have

$$\sigma(\mathcal{L}_{1,\mu}) = \{\lambda \in \mathbb{C} : |\arg(\lambda)| < \pi \text{ and } \operatorname{Re}(\sqrt{\lambda}) \leq \mu\},$$

and from Remark 3.12, it follows that

$$\sigma(-\mathcal{L}_2) = \{\lambda \in \mathbb{C} \setminus \mathbb{R}_+ : \sinh(\omega\sqrt{-\lambda}) = \pm\omega\sqrt{-\lambda}\} = \left\{ -\frac{z_j^2}{\omega^2} \in \mathbb{C} \setminus \mathbb{R}_+ : j \in \mathbb{N} \setminus \{0\} \right\}.$$

Then, since

$$\operatorname{Re} \left(\sqrt{-\frac{z_j^2}{\omega^2}} \right) = \frac{1}{\omega} |\operatorname{Im}(z_j)|,$$

the condition $\sigma(\mathcal{L}_{1,\mu}) \cap \sigma(-\mathcal{L}_2) = \emptyset$ is fulfilled if (5) holds.

The commutativity assumption (H_3) is clearly verified since the actions of operators $\mathcal{L}_{1,\mu}$ and \mathcal{L}_2 are independent.

Now, applying Theorem 2.6, we obtain the result. \square

Remark 4.2. We can conjecture that, for the critical case $\omega\mu = \tau$, the sum $\mathcal{L}_{1,\mu} + \mathcal{L}_2$ is not closable.

4.2 Convexity inequalities

In view to apply Corollary 2.7, we are going to verify inequality (8) in two situations.

Proposition 4.3. Let

$$\mathcal{E}_1 = W^{1,p}(0, +\infty; X) \subset \mathcal{E} = L^p(0, +\infty; X),$$

and

$$\mathcal{E}_2 = L^p \left(0, +\infty; \left[W^{3,p}(0, \omega) \cap W_0^{2,p}(0, \omega) \right] \times L^p(0, \omega) \right) \subset \mathcal{E}.$$

Then, we have

$$D(\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2}) \subset \mathcal{E}_1 \cap \mathcal{E}_2. \quad (44)$$

Proof. Let $V \in D(\mathcal{L}_{1,\mu})$. We must prove that there exists $\delta \in (0, 1)$ such that

$$\|V\|_{\mathcal{E}_1} \leq C \left[\|V\|_{\mathcal{E}} + \|V\|_{\mathcal{E}}^{1-\delta} \|\mathcal{L}_{1,\mu}(V)\|_{\mathcal{E}}^{\delta} \right].$$

For all $V \in W^{2,p}(0, +\infty; X)$, from Kato [10], inequality (1.15), p. 192, we have the convexity inequality

$$\|V'\|_{\mathcal{E}} \leq 2\sqrt{2} \|V\|_{\mathcal{E}}^{1/2} \|V''\|_{\mathcal{E}}^{1/2}.$$

Thus, we deduce that

$$\|V\|_{\mathcal{E}_1} = \|V\|_{\mathcal{E}} + \|V'\|_{\mathcal{E}} \leq \|V\|_{\mathcal{E}} + 2\sqrt{2} \|V\|_{\mathcal{E}}^{1/2} \|V''\|_{\mathcal{E}}^{1/2}.$$

Since $\mathcal{L}_{1,\mu}$ is not invertible, we will estimate $\|V''\|_{\mathcal{E}}$ by $\|\mathcal{L}_{1,\mu}(V) - \lambda_0 V\|_{\mathcal{E}}$, where $\lambda_0 \in \rho(\mathcal{L}_{1,\mu})$. We have

$$V'' - 2\mu V' + (\mu^2 - \lambda_0)V = \mathcal{L}_{1,\mu}(V) - \lambda_0 V.$$

Then, there exists a constant $C > 0$ such that

$$\|V''\|_{\mathcal{E}} + \|V'\|_{\mathcal{E}} + \|V\|_{\mathcal{E}} \leq C \|\mathcal{L}_{1,\mu}(V) - \lambda_0 V\|_{\mathcal{E}},$$

hence

$$\|V''\|_{\mathcal{E}} \leq C \|\mathcal{L}_{1,\mu}(V) - \lambda_0 V\|_{\mathcal{E}} \leq C \|\mathcal{L}_{1,\mu}(V)\|_{\mathcal{E}} + |\lambda_0| C \|V\|_{\mathcal{E}}.$$

Thus

$$\begin{aligned}
\|V\|_{\mathcal{E}_1} = \|V\|_{\mathcal{E}} + \|V'\|_{\mathcal{E}} &\leq \|V\|_{\mathcal{E}} + 2\sqrt{2}\|V\|_{\mathcal{E}}^{1/2}\|V''\|_{\mathcal{E}}^{1/2} \\
&\leq \|V\|_{\mathcal{E}} + 2\sqrt{2C}\|V\|_{\mathcal{E}}^{1/2}(\|\mathcal{L}_{1,\mu}(V)\|_{\mathcal{E}} + |\lambda_0|\|V\|_{\mathcal{E}})^{1/2} \\
&\leq \|V\|_{\mathcal{E}} + 2\sqrt{2C}\|V\|_{\mathcal{E}}^{1/2}\left(\|\mathcal{L}_{1,\mu}(V)\|_{\mathcal{E}}^{1/2} + |\lambda_0|^{1/2}\|V\|_{\mathcal{E}}^{1/2}\right) \\
&\leq \left(1 + 2\sqrt{2C}|\lambda_0|^{1/2}\right)\|V\|_{\mathcal{E}} + 2\sqrt{2C}\|V\|_{\mathcal{E}}^{1/2}\|\mathcal{L}_{1,\mu}(V)\|_{\mathcal{E}}^{1/2}.
\end{aligned}$$

Therefore, inequality (8) is satisfied for $\delta = 1/2$ and $\mathcal{M}_1 = \mathcal{L}_{1,\mu}$. Using Corollary 2.7, we obtain

$$D\left(\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2}\right) \subset \mathcal{E}_1.$$

Now, we must show that, for all $V \in D(\mathcal{L}_2)$, we have

$$\|V\|_{\mathcal{E}_2} \leq C \left[\|V\|_{\mathcal{E}} + \|V\|_{\mathcal{E}}^{1/2} \|\mathcal{L}_2(V)\|_{\mathcal{E}}^{1/2} \right].$$

To this end, it suffices to do it for \mathcal{A} . Set

$$\mathcal{G}_1 = \left[W^{3,p}(0, \omega) \cap W_0^{2,p}(0, \omega) \right] \times L^p(0, \omega) \subset X.$$

We must prove that

$$\forall \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \in D(\mathcal{A}), \quad \left\| \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \right\|_{\mathcal{G}_1} \leq C \left[\left\| \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \right\|_X + \left\| \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \right\|_X^{1/2} \left\| \mathcal{A} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \right\|_X^{1/2} \right].$$

Here, we have

$$\begin{aligned}
\left\| \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \right\|_{\mathcal{G}_1} &= \|\psi_1\|_{W^{3,p}(0,\omega)} + \|\psi_2\|_{L^p(0,\omega)} \\
&= \|\psi_1\|_{L^p(0,\omega)} + \|\psi_1'\|_{L^p(0,\omega)} + \|\psi_1''\|_{L^p(0,\omega)} + \|\psi_1'''\|_{L^p(0,\omega)} + \|\psi_2\|_{L^p(0,\omega)}.
\end{aligned}$$

Set $\varphi = \psi_1''$. Then, for all $\eta > 0$, from Kato [10], inequality (1.12), p. 192, taking $n = \eta + 1$ and $b - a = \omega$, we obtain

$$\|\varphi'\|_{L^p(0,\omega)} \leq \frac{\omega}{\eta} \|\varphi''\|_{L^p(0,\omega)} + \frac{2}{\omega} \left(\eta + 3 + \frac{2}{\eta} \right) \|\varphi\|_{L^p(0,\omega)}.$$

It is not difficult to see that the second member is minimal when

$$\eta = \frac{\sqrt{2} \left(\|\varphi''\|_{L^p(0,\omega)} + 4\|\varphi\|_{L^p(0,\omega)} \right)^{1/2}}{2 \|\varphi\|_{L^p(0,\omega)}^{1/2}}.$$

Therefore, we deduce that

$$\begin{aligned}
\|\varphi'\|_{L^p(0,\omega)} &\leq \frac{\omega\sqrt{2}\|\varphi\|_{L^p(0,\omega)}^{1/2}\|\varphi''\|_{L^p(0,\omega)}}{\left(\|\varphi''\|_{L^p(0,\omega)} + 4\|\varphi\|_{L^p(0,\omega)}\right)^{1/2}} + \frac{4}{\omega} \frac{\sqrt{2}\|\varphi\|_{L^p(0,\omega)}^{1/2}\|\varphi\|_{L^p(0,\omega)}}{\left(\|\varphi''\|_{L^p(0,\omega)} + 4\|\varphi\|_{L^p(0,\omega)}\right)^{1/2}} \\
&\quad + \frac{\sqrt{2}\left(\|\varphi''\|_{L^p(0,\omega)} + 4\|\varphi\|_{L^p(0,\omega)}\right)^{1/2}\|\varphi\|_{L^p(0,\omega)}}{\omega\|\varphi\|_{L^p(0,\omega)}^{1/2}} + \frac{6}{\omega}\|\varphi\|_{L^p(0,\omega)} \\
&\leq \frac{\sqrt{2}}{\omega} \left(\|\varphi''\|_{L^p(0,\omega)} + 4\|\varphi\|_{L^p(0,\omega)} \right)^{1/2} \|\varphi\|_{L^p(0,\omega)}^{1/2} + \frac{6}{\omega} \|\varphi\|_{L^p(0,\omega)} \\
&\quad + \left(\frac{4}{\omega} \|\varphi\|_{L^p(0,\omega)} + \omega \|\varphi''\|_{L^p(0,\omega)} \right) \frac{\sqrt{2}\|\varphi\|_{L^p(0,\omega)}^{1/2}}{\left(\|\varphi''\|_{L^p(0,\omega)} + 4\|\varphi\|_{L^p(0,\omega)}\right)^{1/2}} \\
&\leq C \left(\|\varphi\|_{L^p(0,\omega)} + \|\varphi\|_{L^p(0,\omega)}^{1/2} \|\varphi''\|_{L^p(0,\omega)}^{1/2} \right).
\end{aligned}$$

Then, we have

$$\|\psi_1'''\|_{L^p(0,\omega)} \leq C_\omega \left(\|\psi_1''\|_{L^p(0,\omega)} + \|\psi_1''\|_{L^p(0,\omega)}^{1/2} \|\psi_1^{(4)}\|_{L^p(0,\omega)}^{1/2} \right).$$

Hence

$$\begin{aligned} \left\| \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \right\|_{\mathcal{G}_1} &\leq \|\psi_1\|_{L^p(0,\omega)} + \|\psi_1'\|_{L^p(0,\omega)} + \|\psi_1''\|_{L^p(0,\omega)} \\ &\quad + C_\omega \left(\|\psi_1''\|_{L^p(0,\omega)} + \|\psi_1''\|_{L^p(0,\omega)}^{1/2} \|\psi_1^{(4)}\|_{L^p(0,\omega)}^{1/2} \right) + \|\psi_2\|_{L^p(0,\omega)} \\ &\leq (1 + C_\omega) \|\psi_1\|_{W_0^{2,p}(0,\omega)} + C_\omega \|\psi_1\|_{W_0^{2,p}(0,\omega)}^{1/2} \|\psi_1^{(4)}\|_{L^p(0,\omega)}^{1/2} + \|\psi_2\|_{L^p(0,\omega)} \end{aligned}$$

Now, since \mathcal{A} is invertible, see Proposition 3.2, we have proved that there exists a constant C'_ω depending only on ω such that

$$\|\psi_1^{(4)}\|_{L^p(0,\omega)} \leq C'_\omega \left\| \mathcal{A} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \right\|_X.$$

Moreover, it follows

$$\left\| \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \right\|_{\mathcal{G}_1} \leq (1 + C_\omega) \left\| \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \right\|_X + C_\omega C'_\omega \left\| \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \right\|_X^{1/2} \left\| \mathcal{A} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \right\|_X^{1/2}.$$

Therefore, inequality (8) is satisfied for $\delta = 1/2$ and $\mathcal{M}_2 = \mathcal{L}_2$. Using Corollary 2.7, we obtain

$$D(\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2}) \subset \mathcal{E}_2,$$

which gives the expected result. \square

5 Back to the abstract problem

Now, we are in position to solve the following equation

$$\left(\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2} \right) V + k\rho^2 (\mathcal{P}_1 + \mathcal{P}_{2,\mu}) V = \mathcal{F}. \quad (45)$$

Theorem 5.1. Let $\mathcal{F} \in L^p(0, +\infty; X)$ and assume that (5) holds. Then, there exists $\rho_0 > 0$ such that for all $\rho \in (0, \rho_0]$, equation (45) has a unique strong solution $V \in L^p(0, +\infty; X)$, that is

$$\left\{ \begin{array}{l} \exists (V_n)_{n \geq 0} \in D(\mathcal{L}_{1,\mu}) \cap D(\mathcal{L}_2) : \\ V_n \xrightarrow{n \rightarrow +\infty} V \text{ in } L^p(0, +\infty; X) \text{ and} \\ (\mathcal{L}_{1,\mu} + \mathcal{L}_2) V_n + k\rho^2 (\mathcal{P}_1 + \mathcal{P}_{2,\mu}) V_n \xrightarrow{n \rightarrow +\infty} \mathcal{F} \text{ in } L^p(0, +\infty; X), \end{array} \right. \quad (46)$$

satisfying

$$V \in W^{1,p}(0, +\infty; X) \cap L^p(0, +\infty; [W^{3,p}(0, \omega) \cap W_0^{2,p}(0, \omega)] \times L^p(0, \omega)). \quad (47)$$

Proof. Due to Theorem 4.1, if (5) holds, then $\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2}$ is invertible. Thus, it follows that

$$\left[I + k\rho^2 (\mathcal{P}_1 + \mathcal{P}_{2,\mu}) \left(\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2} \right)^{-1} \right] \left(\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2} \right) V = \mathcal{F}.$$

From (44), we deduce that $V \in D(\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2}) \subset \mathcal{E}_1 \cap \mathcal{E}_2$, that is (47) which involves that

$$(\mathcal{P}_1 + \mathcal{P}_{2,\mu})(\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2})^{-1} \in \mathcal{L}(X).$$

Then, there exists $\rho_0 > 0$ small enough such that, for all $\rho \in (0, \rho_0]$, we have

$$V = \left(\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2}\right)^{-1} \left[I + k\rho^2 (\mathcal{P}_1 + \mathcal{P}_{2,\mu}) \left(\overline{\mathcal{L}_{1,\mu} + \mathcal{L}_2}\right)^{-1} \right]^{-1} \mathcal{F}, \quad (48)$$

which means that V is the unique strong solution of (45). \square

6 Proof of Theorem 1.1

From Theorem 5.1, there exists $\rho_0 > 0$ such that for all $\rho \in (0, \rho_0]$, equation (45) has a unique strong solution $V \in L^p(0, +\infty; X)$ satisfying (47). Then, due to (46), there exists a sequence $(V_n)_{n \in \mathbb{N}} \in D(\mathcal{L}_{1,\mu} + \mathcal{L}_2)$ such that $V_n \xrightarrow[n \rightarrow +\infty]{} V$ and

$$\lim_{n \rightarrow +\infty} (\mathcal{L}_{1,\mu} + \mathcal{L}_2) V_n + k\rho^2 (\mathcal{P}_1 + \mathcal{P}_{2,\mu}) V_n = \mathcal{F}.$$

Since $(V_n)_{n \in \mathbb{N}} \in D(\mathcal{L}_{1,\mu} + \mathcal{L}_2)$, then the previous equality can be written as

$$\begin{cases} \lim_{n \rightarrow +\infty} (V_n''(t) - \mathcal{A}V_n(t) - \mathcal{F}_n(t)) = 0 \\ \lim_{n \rightarrow +\infty} V_n(0) = 0, \quad \lim_{n \rightarrow +\infty} V_n(+\infty) = 0, \end{cases} \quad (49)$$

where

$$\mathcal{F}_n(t) = k\rho^2 e^{-2t} \mathcal{A}_0 V_n(t) + k\rho^2 e^{-2t} [(\mathcal{B}_{2,\mu} V_n)](t) + 2\mu V_n'(t) - \mu^2 V_n(t) + \mathcal{F}(t).$$

Since $V_n \xrightarrow[n \rightarrow +\infty]{} V$ in \mathcal{E} and V satisfies (47), we deduce that

$$\lim_{n \rightarrow +\infty} V_n(0) = V(0) = 0 \quad \text{with} \quad \lim_{n \rightarrow +\infty} V_n(+\infty) = V(+\infty) = 0,$$

and

$$\lim_{n \rightarrow +\infty} \mathcal{F}_n(t) = \mathcal{F}_\infty(t) \in L^p(0, +\infty; X),$$

where

$$\mathcal{F}_\infty(t) = k\rho^2 e^{-2t} \mathcal{A}_0 V(t) + k\rho^2 e^{-2t} [(\mathcal{B}_{2,\mu} V)](t) + 2\mu V'(t) - \mu^2 V(t) + \mathcal{F}(t).$$

Thus, problem (49) can be written as follows

$$\begin{cases} \lim_{n \rightarrow +\infty} (V_n''(t) - \mathcal{A}V_n(t)) = \mathcal{F}_\infty(t) \\ V(0) = 0, \quad V(+\infty) = 0. \end{cases}$$

Moreover, from Proposition 3.13, $\mathcal{A} \in \text{BIP}(X, \theta_{\mathcal{A}})$, with $\theta_{\mathcal{A}} \in (0, \pi)$ and due to Haase [9], Proposition 3.2.1, e), p. 71, it follows that $\sqrt{\mathcal{A}} \in \text{BIP}(X, \theta_{\mathcal{A}}/2)$ with $\theta_{\mathcal{A}}/2 \in (0, \pi/2)$. Therefore, due to Eltaief and Maingot [6], Theorem 2, p. 712, with $L_1 = L_2 = -\sqrt{\mathcal{A}}$, there exists a unique classical solution to the following problem

$$\begin{cases} \mathcal{V}''(t) - \mathcal{A}\mathcal{V}(t) = \mathcal{F}_\infty(t) \\ \mathcal{V}(0) = 0, \quad \mathcal{V}(+\infty) = 0, \end{cases}$$

that is

$$\mathcal{V} \in W^{2,p}(0, +\infty; X) \cap L^p(0, +\infty; D(\mathcal{A})).$$

Thus, it follows that

$$\lim_{n \rightarrow +\infty} (V_n''(t) - \mathcal{A}V_n(t)) = \mathcal{V}''(t) - \mathcal{A}\mathcal{V},$$

hence

$$\lim_{n \rightarrow +\infty} [(V_n(t) - \mathcal{V}(t))'' - \mathcal{A}(V_n(t) - \mathcal{V}(t))] = 0.$$

Now, set

$$\begin{cases} D(\delta_2) &= \{\varphi \in W^{2,p}(0, +\infty; X) : \varphi(0) = \varphi(+\infty) = 0\} \\ \delta_2\varphi &= \varphi'', \quad \varphi \in D(\delta_2). \end{cases}$$

Then

$$0 = \lim_{n \rightarrow +\infty} [(V_n(t) - \mathcal{V}(t))'' - \mathcal{A}(V_n(t) - \mathcal{V}(t))] = \lim_{n \rightarrow +\infty} -(-\delta_2 + \mathcal{A})(V_n(t) - \mathcal{V}(t)). \quad (50)$$

From Prüss and Sohr [20], Theorem C, p. 166-167, it follows that $-\delta_2 \in \text{BIP}(X, \theta_{\delta_2})$, for every $\theta_{\delta_2} \in (0, \pi)$ and due to Proposition 3.13, $\mathcal{A} \in \text{BIP}(X, \theta_{\mathcal{A}})$, for all $\theta_{\mathcal{A}} \in (0, \pi)$. Thus, since $-\delta_2$ and \mathcal{A} are resolvent commuting with $\theta_{\delta_2} + \theta_{\mathcal{A}} < \pi$, from Prüss and Sohr [19], Theorem 5, p. 443, we obtain that

$$-\delta_2 + \mathcal{A} \in \text{BIP}(X, \theta), \quad \theta = \max(\theta_{\delta_2}, \theta_{\mathcal{A}}).$$

Moreover, due to Proposition 3.2, we have $0 \in \rho(\mathcal{A})$, then we deduce from Prüss and Sohr [19], remark at the end of p. 445, that $0 \in \rho(\delta_2 + \mathcal{A})$. Therefore, due to (50), we obtain that

$$\lim_{n \rightarrow +\infty} V_n(t) - \mathcal{V}(t) = 0,$$

hence, since $V_n \xrightarrow[n \rightarrow +\infty]{} V$, by uniqueness of the limit, we deduce that

$$V = \mathcal{V} \in W^{2,p}(0, +\infty; X) \cap L^p(0, +\infty; D(\mathcal{A})).$$

This prove that $\mathcal{L}_{1,\mu} + \mathcal{L}_2$ is closed and that $V \in D(\mathcal{L}_{1,\mu} + \mathcal{L}_2)$.

Acknowledgments

We would like to thank the referee for its valuable comments and corrections which have helped us to improve this second part.

Conflict of interest

On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

- [1] W. ARENDT, S. BU, M. HAASE, “Functional calculus, variational methods and Liapunov’s theorem”, *Arch. Math.*, 77, 2001, pp. 65-75.
- [2] J. BOURGAIN, “Some remarks on Banach spaces in which martingale difference sequences are unconditional”, *Ark. Mat.*, vol. 21,1983, pp. 163-168.
- [3] D.L. BURKHOLDER, “A geometrical characterisation of Banach spaces in which martingale difference sequences are unconditional”, *Ann. Probab.*, vol. 9, 1981, pp. 997-1011.
- [4] G. DA PRATO & P. GRISVARD, “Somme d’opérateurs linéaires et équations différentielles opérationnelles”, *J. Math. Pures et Appl.*, 54, 1975, pp. 305-387.

- [5] G. DORE & A. VENNI, “On the closedness of the sum of two closed operators”, *Math. Z.*, 196, 1987, pp. 189-201.
- [6] A. ELTAIEF & S. MAINGOT, “Second Order Abstract Differential Equations of Elliptic Type Set in \mathbb{R}_+ ”, *Demonstratio Mathematica*, XLVI, 4, 2013, pp. 709-727.
- [7] J. FÄDLE, "Die Selbstspannungs-Eigenwertfunktionen der Quadratischen Scheibe", *Ing. Arch.*, 11, 1940, pp. 125-149.
- [8] P. GRISVARD, “Singular behavior of elliptic problems in non Hilbertian Sobolev spaces”, *Prépublication n°321, Université de Nice*, 1992.
- [9] M. HAASE, *The functional calculus for sectorial Operators*, Birkhauser, 2006.
- [10] T. KATO, *Perturbation theory for linear operators*, Springer, 1980.
- [11] H. KOMATSU, “Fractional powers of operators”, *Pacific Journal of Mathematics*, Vol. 19, No. 2, 1966, pp. 285-346.
- [12] R. LABBAS, K. LEMRABET, S. MAINGOT & A. THOREL, “Generalized linear models for population dynamics in two juxtaposed habitats”, *Discrete and Continuous Dynamical Systems - A*, Vol. 39, 5, 2019, pp. 2933-2960.
- [13] R. LABBAS, S. MAINGOT, D. MANCEAU & A. THOREL, “On the regularity of a generalized diffusion problem arising in population dynamics set in a cylindrical domain”, *Journal of Mathematical Analysis and Applications*, 450, 2017, pp. 351-376.
- [14] R. LABBAS, S. MAINGOT & A. THOREL, “Solvability of a fourth order elliptic problem in a bounded sector, part I”, *Boll. Unione Mat. Ital.*, (2023).
- [15] R. LABBAS, S. MAINGOT & A. THOREL, “Generation of analytic semigroup for some generalized diffusion operators in L^p -spaces”, *Math. Ann.*, 384, 2022, pp. 1-49.
- [16] R. LABBAS & M. MOUSSAOUI, “On the resolution of the heat equation with discontinuous coefficients”, *Semigroup Forum*, 60, 2000, pp. 187-201.
- [17] R. LABBAS & B.-K. SADALLAH, “Smoothness of the solution of a fourth order parabolic equation in a polygonal domain”, *Int. J. Appl. Math.*, Vol. 1, No. 1, 1999, pp. 75-90.
- [18] S.G. MIHLIN, “On the multipliers of Fourier integrals”, *Dokl. Akad. Nauk SSSR*, N.S., 109, 1956, pp. 701-703.
- [19] J. PRÜSS & H. SOHR, “On operators with bounded imaginary powers in Banach spaces”, *Mathematische Zeitschrift, Springer-Verlag, Math. Z.*, 203, 1990, pp. 429-452.
- [20] J. PRÜSS & H. SOHR, “Imaginary powers of elliptic second order differential operators in L^p -spaces”, *Hiroshima Math. J.*, 23, no. 1, 1993, pp. 161-192.
- [21] R. SEELEY, “Norms and domains of the complex powers A_B^z ”, *American Journal of Mathematics*, Vol. 93, No. 2, 1971, pp. 299-309.
- [22] A. THOREL, “Operational approach for biharmonic equations in L^p -spaces”, *Journal of Evolution Equations*, 20, 2020, pp. 631-657.
- [23] H. TRIEBEL, *Interpolation theory, function Spaces, differential Operators*, North-Holland publishing company Amsterdam New York Oxford, 1978.