The summatory function of the Möbius function involving the greatest common divisor

Isao Kiuchi and Sumaia Saad Eddin

Abstract

Let gcd(m, n) denote the greatest common divisor of the positive integers mand n, and let μ represent the Möbius function. For any real number x > 5, we define the summatory function of the Möbius function involving the greatest common divisor as $S_{\mu}(x) := \sum_{mn \leq x} \mu(gcd(m, n))$. In this paper, we present an asymptotic formula for $S_{\mu}(x)$. Assuming the Riemann Hypothesis, we delve further into the asymptotic behavior of $S_{\mu}(x)$ and derive a mean square estimate for its error term. Our proof employs the Perron formula, Parseval's theorem, complex integration techniques, and the properties of the Riemann zeta-function.

1 Introduction and main results

Let $\mu(n)$ denote the Möbius function, namely

 $\mu(n) = \begin{cases} 1 & \text{if } n = 1, \\ (-1)^k & \text{if } n \text{ is squarefree and } n = p_1 p_2 \cdots p_k, \\ 0 & \text{if } n \text{ is not squarefree.} \end{cases}$

For any integer $k \geq 2$, let $gcd(d_1, \ldots, d_k)$ denote the greatest common divisor of the positive integers d_1, \ldots, d_k . We define the Dirichlet convolution of two arithmetic functions f and g as $f * g(n) = \sum_{d|n} f(d)g(n/d)$ for all positive integers n. The arithmetic functions $\mathbf{1}(n)$ and id(n) are defined by $\mathbf{1}(n) = 1$ and id(n) = n, respectively. The function τ_k is the k-factors Piltz divisor function given by $\mathbf{1} * \mathbf{1} * \cdots \mathbf{1}$. Specifically, for k = 2, we denote τ_2 as τ . For an arbitrary arithmetic function g, we define the sum

$$f_{g,k}(n) := \sum_{d_1 \cdots d_k = n} g\left(\gcd(d_1, \dots, d_k)\right)$$

Mathematics Subject Classification 2020: 11N37, 11A25, 11M99,

Keywords: gcd-sum function, Möbius function, Dirichlet series, Riemann hypothesis, Asymptotic results on arithmetical functions

and its summatory function as

$$S_{g,k}(x) := \sum_{n \le x} f_{g,k}(n) = M_{g,k}(x) + E_{g,k}(x),$$
(1)

where $M_{g,k}(x)$ and $E_{g,k}(x)$ denote the main term and the error term of $S_{g,k}(x)$, respectively. In 2012, Krätzel, Nowak and Tóth [5] provided asymptotic formulas for a class of arithmetic functions that describe the value distribution of the greatest common divisor function, generated by a Dirichlet series whose analytic behavior is determined by the factor $\zeta^2(s)\zeta(2s-1)$, where $\zeta(s)$ is the Riemann zeta-function. They proved that

$$f_{g,k}(n) = \sum_{a^k b = n} \mu * g(a)\tau_k(b).$$
⁽²⁾

The identity (2) was instrumental in establishing asymptotic formulas for $f_{g,k}(n)$ for specific choices of g, such as the identity function id and the sum of divisors function $\sigma = id * 1$. Furthermore, they showed that the Dirichlet series generated by $f_{g,k}(n)$ is expressed by

$$\sum_{n=1}^{\infty} \frac{f_{g,k}(n)}{n^s} = \frac{\zeta^k(s)}{\zeta(ks)} \sum_{n=1}^{\infty} \frac{g(n)}{n^{ks}},$$
(3)

which converges absolutely in the half-plane $\Re(s) > \sigma_0$, where σ_0 depends on g and k.

Now, we set $g = \mu$, and use the identity $\sum_{n=1}^{\infty} \frac{\mu(n)}{n^s} = \frac{1}{\zeta(s)}$ to derive

$$S_{\mu,k}(x) = \sum_{m \le x^{\frac{1}{k}}} \mu * \mu(m) \sum_{n \le \frac{x}{m^k}} \tau_k(n)$$
(4)

and that

$$\sum_{n=1}^{\infty} \frac{f_{\mu,k}(n)}{n^s} = \frac{\zeta^k(s)}{\zeta^2(ks)},$$
(5)

where both series converge absolutely in the half-plane $\Re(s) > 1$. Throughout the article when k = 2, we adopt simplified notation: $f_{\mu,2}(n) = f_{\mu}(n), S_{\mu,2}(x) = S_{\mu}(x), M_{\mu,2}(x) = M_{\mu}(x)$, and $E_{\mu,2}(x) = E_{\mu}(x)$.

The first purpose of this paper is to establish the following result which provides an asymptotic formula of $S_{\mu,2}(x)$.

Theorem 1. For any real number x > 5, we have

$$S_{\mu}(x) = \frac{1}{\zeta^{2}(2)} \left(\log x + 2\gamma - 1 - 4\frac{\zeta'(2)}{\zeta(2)} \right) x + O\left(x^{1/2}\log^{2} x\right),$$

where γ is the Euler-Mascheroni constant and $\zeta'(s)$ is the first derivative of the zetafunction $\zeta(s)$. The proof is based on applying the Perron formula (which gives a direct link between the summatory function $\sum_{n \leq x} a_n$ and the corresponding Dirichlet series $\alpha(x) = \sum_{n \geq 1} a_n/n^s$, for more details see Section 2), contour integration to the generating Dirichlet series and using the properties of the Riemann zeta-function.

In the following result, we focus on estimating the error term and employ elementary methods, including the use of a specific Lemma 2 below (in Section 2), to achieve a better bound.

Theorem 2. For any real number x > 5, we have

$$E_{\mu}(x) = O\left(x^{1/2} \exp\left(-A\left(\log x \log\log x\right)^{1/3}\right)\right)$$
(6)

with a positive constant A.

Next, we assume that the Riemann Hypothesis is true. By using the complex integration approach and Montgomery–Vaughan's method [6], we obtain the following result.

Theorem 3. If the Riemann hypothesis is true, then, for any real number x > 5, we have

$$E_{\mu}(x) = \sum_{\ell \le Y} \mu * \mu(\ell) \Delta\left(\frac{x}{\ell^2}\right) + O\left(x^{\frac{1}{2} + \varepsilon} Y^{\frac{1-4\beta}{4-4\beta} + \varepsilon}\right)$$

with $x^{\varepsilon} \ll Y \ll x^{\frac{1}{2}-\varepsilon}$, and a fixed number β such that $\frac{1}{4} < \beta < \frac{1}{2}$. Here $\Delta(x)$ denote the error term in the Dirichlet divisor problem.

By setting $Y = x^{\frac{77}{258}}$ and $\beta = \frac{2}{5}$ in the above equation and applying the given estimation of $\Delta(x)$ from (11), we deduce the following result.

Corollary 1. We have

 $E_{\mu}(x) \ll x^{\frac{439}{1032} + \varepsilon}.$

Note that $\frac{439}{1032} = \frac{1}{2} - \frac{77}{1032} = 0.4253875 \cdots < \frac{1}{2} = 0.5$. This implies that the latest estimation of the error term yields a slight improvement in the order of magnitude of (6).

Now, we define the integral of the mean square of $E_{\mu}(x)$ as follows

$$I_{\mu}(T) := \int_1^T E_{\mu}^2(t)dt$$

for any real number T > 5. Lastly, we use Parseval's identity (which establishes a connection between the sum or integral of squares of a function and the sum or integral of squares of its Fourier transform, see (A.5) in [4]) along with certain properties of the Riemann zeta-function to derive the mean square estimates for $E_{\mu}(t)$. Further, we shall investigate the integral $I_{\mu}(T)$ assuming the Riemann Hypothesis.

Theorem 4. If the Riemann Hypothesis is true, then, for any real number T > 5, we have

$$I_{\mu}(T) \ll T^{3/2} \exp\left(A \frac{\log T}{\log \log T}\right),$$

with a positive constant A.

Remark 1. Under the Riemann Hypothesis, if $I_{\mu}(T)$ can be evaluated as

$$c_{\mu}T^{3/2}(\log T)^{\ell} + O\left(T^{3/2}(\log T)^{\ell-\delta}\right)$$

as $x \to \infty$ with a constant $c_{\mu} > 0$ depending on μ and numbers $\ell > 0$, $\delta > 0$, then we can derive the Omega-result for $E_{\mu}(x)$, namely

$$E_{\mu}(x) = \Omega\left(x^{1/4} (\log x)^{\ell/2}\right)$$

Sums of squarefree numbers

We set $g = |\mu|$ in (2), yielding

$$S_{|\mu|,k}(x) = \sum_{m \le x^{1/k}} \mu * |\mu|(m) \sum_{n \le x/m^k} \tau_k(n).$$
(7)

Using (3) and the Dirichlet series $\sum_{n=1}^{\infty} \frac{|\mu(n)|}{n^s} = \frac{\zeta(s)}{\zeta(2s)}$, we deduce

$$\sum_{n=1}^{\infty} \frac{f_{|\mu|,k}(n)}{n^s} = \frac{\zeta^k(s)}{\zeta(2ks)},\tag{8}$$

which converges absolutely in the half-plane $\Re(s) > 1$. From (8), we derive an alternative expression for (7), as follows:

$$S_{|\mu|,k}(x) = \sum_{m \le x^{1/(2k)}} \mu(m) \sum_{n \le x/m^{2k}} \tau_k(n).$$
(9)

Taking k = 2 in (9) and using (10) and (11) along with the Dirichlet series $\sum_{n=1}^{\infty} \frac{\mu(n)}{n^s} = \frac{1}{\zeta(s)}$ and $\sum_{n=1}^{\infty} \frac{\mu(n)}{n^s} \log n = \frac{\zeta'(s)}{\zeta^2(s)}$, we obtain

$$S_{|\mu|,2}(x) = \sum_{m \le x^{1/4}} \mu(m) \sum_{n \le x/m^4} \tau(n)$$

= $x (\log x + 2\gamma - 1) \sum_{m \le x^{1/4}} \frac{\mu(m)}{m^4} - 2x \sum_{m \le x^{1/4}} \frac{\mu(m)}{m^4} \log m + \sum_{m \le x^{1/4}} \mu(m) \Delta\left(\frac{x}{m^4}\right)$
= $\frac{x}{\zeta(4)} \left(\log x + 2\gamma - 1 - 2\frac{\zeta'(4)}{\zeta(4)}\right) + O\left(x^{\theta + \varepsilon}\right).$

Therefore, we have

$$S_{|\mu|,2}(x) = \frac{x}{\zeta(4)} \left(\log x + 2\gamma - 1 - 2\frac{\zeta'(4)}{\zeta(4)} \right) + O\left(x^{\theta+\varepsilon}\right)$$

with $\theta = \frac{131}{416}$ and any real number $\varepsilon > 0$.

2 Some known results

In this section, we recall the material we need to prove our results. Throughout this section and in subsequent ones, we adhere to the traditional notation $s = \sigma + it$.

Lemma 1. For $t \ge t_0 > 0$ uniformly in σ , we have

$$\zeta(\sigma + it) \ll \begin{cases} t^{(3-4\sigma)/6} \log t & (0 \le \sigma \le 1/2), \\ t^{(1-\sigma)/3} \log t & (1/2 \le \sigma \le 1), \\ \log t & (1 \le \sigma < 2). \end{cases}$$

Proof. The above formulas follow from [9, Theorem II.3.8], and [4, Theorem 8.5 and (8.112)].

Lemma 2. For any number x > 5, we have

$$\sum_{n \le x} \frac{\mu * \mu(n)}{n^2} = \frac{1}{\zeta^2(2)} + O\left(\frac{\delta(x)}{x}\right)$$

and

$$\sum_{n \le x} \frac{\mu * \mu(n)}{n^2} \log n = 2\frac{\zeta'(2)}{\zeta^3(2)} + O\left(\frac{\delta(x)}{x}\right).$$

where $\delta(x) = \exp\left(-c\left(\log x \log \log x\right)^{1/3}\right)$ with c being a positive constant. Proof. The proof of this result can be found in [3, Lemma 3.3].

The following two well-known results about the properties of the Riemann Zetafunction, which provides the functional equation of the Riemann Zeta-function and its bounds under the Riemann Hypothesis. The proof of these results can be found in [4], [7], and [10].

Lemma 3. The Riemann zeta-function $\zeta(s)$ can be analytically continued to a meromorphic function in the whole complex plane \mathbb{C} , its only singularity being a simple pole at s = 1 with residue 1. It satisfies a functional equation

$$\zeta(s) = \chi(s)\zeta(1-s),$$

where $\chi(s) = 2^s \pi^{s-1} \sin \frac{\pi s}{2} \Gamma(1-s)$. Also, in any bounded vertical strip, using the Stirling formula, we deduce

$$\chi(s) = \left(\frac{t}{2\pi}\right)^{\frac{1}{2}-\sigma-it} e^{i(t+\frac{\pi}{4})} \left(1+O\left(\frac{1}{t}\right)\right) \qquad (t \ge t_0 > 0).$$

Lemma 4. Assume that the Riemann Hypothesis is true. For $t \ge t_0 \ge 0$, uniformly in σ , we have

$$\zeta(\sigma + it) \ll \begin{cases} t^{\varepsilon} & (1/2 \le \sigma \le 2), \\ 1 & (\sigma > 2), \end{cases}$$

and

$$\frac{1}{\zeta(\sigma+it)} \ll \begin{cases} t^{\varepsilon} & \left(\frac{1}{2} + \frac{1}{\log\log t} \le \sigma \le 2\right), \\ 1 & (\sigma > 2). \end{cases}$$

Perron's formula

The Perron formula is used to express a function defined by a Dirichlet series $\alpha(s) = \sum_{n=1}^{\infty} a_n/n^s$, with abscissa of convergence σ_c , in terms of its complex integral. That means

$$\sum_{n \le x} a_n = \frac{1}{2\pi i} \int_{\sigma_0 - i\infty}^{\sigma_0 + i\infty} \alpha(s) \frac{x^s}{s} \, ds,$$

for $\sigma_0 > \max(\sigma_c, 0)$.

Theorem 5 (Perron's formula, Theorem 5.1 in [7]). If $\sigma_0 > \max(\sigma_c, 0)$ and x > 0, then

$$\sum_{n \le x}' a_n = \lim_{T \to \infty} \frac{1}{2\pi i} \int_{\sigma_0 - iT}^{\sigma_0 + iT} \alpha(s) \frac{x^s}{s} \, ds,$$

where \sum' indicates that if x is an integer, then the last term into the sum becomes $a_x/2$ instead of a_x .

Theorem 6 (Theorem 5.2 in [7]). If $\sigma_0 > \max(\sigma_c, 0)$ and x > 0, then

$$\sum_{n \le x}' a_n = \frac{1}{2\pi i} \int_{\sigma_0 - iT}^{\sigma_0 + iT} \alpha(s) \frac{x^s}{s} \, ds + R,$$

where

$$R = \frac{1}{\pi} \sum_{x/2 < n < x} a_n \operatorname{si}\left(T \log \frac{x}{n}\right) - \frac{1}{\pi} \sum_{x < n < 2x} a_n \operatorname{si}\left(T \log \frac{x}{n}\right) + O\left(\frac{4^{\sigma_0} + x^{\sigma_0}}{T} \sum_n \frac{|a_n|}{n^{\sigma_0}}\right).$$

Here si(x) is the sine integral and equal to $-\int_x^\infty \frac{\sin u}{u}$.

Corollary 2 (Corollary 5.3 in [7]). Under the conditions stated above, we have

$$R \ll \sum_{\substack{x < n < 2x, \\ n \neq x}} |a_n| \min\left(1, \frac{x}{T|x-n|}\right) + \frac{4^{\sigma_0} + x^{\sigma_0}}{T} \sum_{n=1}^{\infty} \frac{|a_n|}{n^{\sigma_0}}.$$

3 Proofs

3.1 Proof of Theorem 1

Without loss of generality we may suppose that $x \in \mathbb{Z} + \frac{1}{2}$. Applying Theorem 6 and Corollary 2 and using (5) with k = 2, we obtain

$$S_{\mu,2}(x) = \frac{1}{2\pi i} \int_{\alpha-iT}^{\alpha+iT} \frac{\zeta^2(s)}{\zeta^2(2s)} \frac{x^s}{s} ds + O\left(\frac{x^{1+\varepsilon}}{T}\right),$$

where $\alpha = 1 + \frac{1}{\log x}$ and T is a real parameter at our disposal. Let $\Gamma(\alpha, \beta, T)$ denote the contour consisting of the line segments $[\alpha - iT, \beta - iT]$, $[\beta - iT, \beta + iT]$ and $[\beta + iT, \alpha + iT]$. We denote the integrals over the horizontal line segments as K_1 and K_3 , and the integral over the vertical line segment as K_2 . Setting $\beta = \frac{1}{2}$, and using Lemma 1 along with the estimate $\frac{1}{|\zeta(\sigma+it)|} \ll \log(1+|t|)$ for $|t| \ge t_0 > 0$ and $\sigma \ge 1$, we find

$$K_{1}, K_{3} \ll \frac{\log^{2} T}{T} \int_{\frac{1}{2}}^{\alpha} |\zeta(\sigma + iT)|^{2} x^{\sigma} d\sigma$$
$$\ll \frac{\log^{4} T}{T^{1/3}} \int_{\frac{1}{2}}^{\alpha} \left(\frac{x}{T^{2/3}}\right)^{\sigma} d\sigma$$
$$\ll \log^{4} T \left(\frac{x}{T} + \frac{x^{1/2}}{T^{2/3}}\right).$$

For the integral K_2 , using the estimate given in [11] that

$$\int_{1}^{T} \frac{|\zeta(\frac{1}{2} + it)|^2}{|\zeta(1+2it)|^2} dt \ll T \log T$$

and applying integration by parts, we obtain

$$\int_{1}^{T} \frac{|\zeta(\frac{1}{2}+it)|^{2}}{|\zeta(1+2it)|^{2}} \frac{dt}{t} \ll T\log T + \int_{T/2}^{T} \frac{\log t}{t} dt \ll \log^{2} T.$$

Thus, employing Lemma 1, we have

$$K_2 \ll x^{1/2} + x^{1/2} \int_{t_0 \le |t| \le T} \frac{|\zeta(\frac{1}{2} + it)|^2}{|\zeta(1+2it)|^2} \frac{dt}{t}$$
$$\ll x^{1/2} \log^2 T.$$

To complete the proof, we calculate the residue at the double pole s = 1 with the residue

$$\operatorname{Res}_{s=1} \frac{\zeta^2(s)}{\zeta^2(2s)} \frac{x^s}{s},$$

which equals

$$\frac{1}{\zeta^2(2)} \left(\log x + 2\gamma - 1 - 4 \frac{\zeta'(2)}{\zeta(2)} \right) x.$$

Therefore, we conclude that

$$S_{\mu,2}(x) = \frac{1}{\zeta^2(2)} \left(\log x + 2\gamma - 1 - 4\frac{\zeta'(2)}{\zeta(2)} \right) x + O\left(x^{1/2} \log^2 x\right),$$

and by setting T = x, the proof is complete.

3.2 Proof of Theorem 2

Let $\tau(n)$ denote the divisor function, defined by 1 * 1, and let $\Delta(x)$ represent the error term in the asymptotic formula for the sum

$$D(x) := \sum_{n \le x} \tau(n) = (\log x + 2\gamma - 1) x + \Delta(x)$$
(10)

valid for any real number x > 2. It is known the Dirichlet divisor problem involves studying the properties of $\Delta(x)$. The best known estimate

$$\Delta(x) \ll x^{\theta + \varepsilon},\tag{11}$$

with $\theta = \frac{131}{416}$, was proved by Huxley [2]. Using (10) and (4) with k = 2, we obtain

$$S_{\mu}(x) = \sum_{m \le x^{1/2}} \mu * \mu(m) \sum_{n \le x/m^2} \tau(n)$$

= $x (\log x + 2\gamma - 1) \sum_{m \le x^{1/2}} \frac{\mu * \mu(m)}{m^2} - 2x \sum_{m \le x^{1/2}} \frac{\mu * \mu(m)}{m^2} \log m + U(x),$

where

$$U(x) := \sum_{m \le x^{1/2}} \mu * \mu(m) \Delta\left(\frac{x}{m^2}\right).$$

Applying Lemma 2, we have

$$S_{\mu}(x) = \frac{x}{\zeta^{2}(2)} \left(\log x + 2\gamma - 1 - 4\frac{\zeta'(2)}{\zeta(2)} \right) + U(x) + O\left(x^{1/2}\delta(x)\right),$$

where

$$\delta(x) = \exp\left(-c\left(\log x \log\log x\right)^{1/3}\right)$$

with c being a positive constant. We divide U(x) into two sums as follows:

$$U(x) = \left(\sum_{n \le x^{1/2} \delta(x)} + \sum_{x^{1/2} \delta(x) < n \le x^{1/2}}\right) \mu * \mu(n) \Delta\left(\frac{x}{n^2}\right)$$

:= $U_1(x) + U_2(x)$, say. (12)

Using the well-known result $\Delta(x) = O(x^{1/3})$ and partial summation, we obtain

$$U_{1}(x) \ll \sum_{n \leq x^{1/2} \delta(x)} \tau(n) \Delta\left(\frac{x}{n^{2}}\right)$$
$$\ll x^{1/3} \sum_{n \leq x^{1/2} \delta(x)} \frac{\tau(n)}{n^{2/3}}$$
$$\ll x^{1/3} \left(x^{1/2} \delta(x)\right)^{1/3} \log x^{1/2} \delta(x)$$
$$\ll x^{1/2} \exp\left(-c'(\log x \log \log x)^{1/3}\right)$$
(13)

with c' being a positive constant. Next, employing the formula $\int_1^y \Delta(\xi) d\xi = \frac{y}{4} + O(y^{3/4})$ for any y > 1 (see [1]), we deduce

$$U_{2}(x) \ll \max_{x^{1/2}\delta(x) < n \le x^{1/2}} |\mu * \mu(n)| \sum_{x^{1/2}\delta(x) < n \le x^{1/2}} \Delta\left(\frac{x}{n^{2}}\right)$$
$$\ll x^{\varepsilon} \int_{1}^{1/\delta^{2}(x)} \Delta(\xi) d\xi$$
$$\ll \frac{x^{\varepsilon}}{\delta^{2}(x)} \ll x^{1/3+\varepsilon}.$$
(14)

Substituting (13) and (14) into (12), we establish the assertion of Theorem 2.

3.3 Proof of Theorem 3

From (5) with k = 2, we observe that

$$\sum_{n=1}^{\infty} \frac{f_{\mu,2}(n)}{n^s} = \sum_{n=1}^{\infty} \left(\sum_{d^2m=n} \mu * \mu(d)\tau(m) \right) \frac{1}{n^s} = \frac{\zeta^2(s)}{\zeta^2(2s)},\tag{15}$$

which converges absolutely for $\Re(s) > 1$. Without loss of generality, we can assume that $x \in \mathbb{Z} + \frac{1}{2}$. From (15), we derive

$$S_{\mu}(x) := \sum_{n \le x} f_{\mu,2}(n) = \sum_{n \le x} \sum_{n = m\ell^2} \mu * \mu(\ell)\tau(m).$$

Let $x^{\varepsilon} < Y < x^{\frac{1}{2}-\varepsilon}$. Then the above formula can be rewritten as follows

$$S_{\mu}(x) = \left(\sum_{\substack{m\ell^{2} \leq x \\ \ell \leq Y}} + \sum_{\substack{m\ell^{2} \leq x \\ \ell > Y}}\right) \mu * \mu(\ell)\tau(m)$$

$$:= S_{1}(x;Y) + S_{2}(x;Y) \quad \text{say.}$$
(16)

We define

$$g_Y(s) := \sum_{n > Y} \frac{\mu * \mu(n)}{n^s} = \frac{1}{\zeta^2(s)} - \sum_{n \le Y} \frac{\mu * \mu(n)}{n^s}$$

for $\Re(s) > 1$. Following a similar approach as in Lemma 3 in [8], we find that

$$g_Y(s) = O\left(Y^{\frac{1}{2}-\sigma+\varepsilon}(1+|t|)^{\varepsilon}\right)$$
(17)

is valid for $\sigma \geq \frac{1}{2} + \varepsilon$ under the Riemann Hypothesis. Applying (10), we have

$$S_1(x;Y) = \sum_{\ell \le Y} \mu * \mu(\ell) D\left(\frac{x}{\ell^2}\right)$$
$$= x \sum_{\ell \le Y} \frac{\mu * \mu(\ell)}{\ell^2} \log \frac{x}{\ell^2} + (2\gamma - 1)x \sum_{\ell \le Y} \frac{\mu * \mu(\ell)}{\ell^2} + \sum_{\ell \le Y} \mu * \mu(\ell) \Delta\left(\frac{x}{\ell^2}\right).$$
(18)

To estimate $S_2(x; Y)$, we employ the methodology outlined in the proof of Theorem 1. Let T be a real parameter at our disposal. Employing Perron's formula (see Section 2) and (17), we deduce

$$S_2(x;Y) = \frac{1}{2\pi i} \int_{\alpha-iT}^{\alpha+iT} \zeta^2(s) g_Y(2s) \frac{x^s}{s} ds + O\left(\frac{x^{1+\varepsilon}}{T}\right),$$

with $\alpha = 1 + \frac{1}{\log x}$. Now, let $\Gamma(\alpha, \beta, T)$ denote the contour consisting of the line segments $[\alpha - iT, \beta - iT]$, $[\beta - iT, \beta + iT]$ and $[\beta + iT, \alpha + iT]$. We shift the integration with respect to s to $\Gamma(\alpha, \beta, T)$ with $\frac{1}{4} < \beta < \frac{1}{2}$. The integrals over the horizontal line segments are denoted by $S_{2,1}$ and $S_{2,3}$, and the integral over the vertical line segment by $S_{2,2}$. The main term arises from the sum of the residue at the double pole s = 1. Hence, by the Cauchy residue theorem, we obtain

$$S_2(x;Y) = \frac{1}{2\pi i} \left\{ \int_{\beta+iT}^{\alpha+iT} + \int_{\beta-iT}^{\beta+iT} + \int_{\alpha-iT}^{\beta-iT} \right\} \zeta^2(s) g_Y(2s) \frac{x^s}{s} ds + \operatorname{Res}_{s=1} \zeta^2(s) g_Y(2s) \frac{x^s}{s} + O\left(\frac{x^{1+\varepsilon}}{T}\right).$$
(19)

By applying Lemmas 3 and 4, we deduce that

$$S_{2,1}, S_{2,3} \ll \frac{1}{T} \left(\int_{\beta}^{\frac{1}{2}} + \int_{\frac{1}{2}}^{\alpha} \right) |\zeta(\sigma + iT)|^2 |g_Y(2(\sigma + iT))| x^{\sigma} d\sigma$$
$$\ll Y^{1/2+\varepsilon} T^{2\varepsilon} \int_{\beta}^{\frac{1}{2}} \left(\frac{x}{T^2 Y^2} \right)^{\sigma} d\sigma + Y^{1/2+\varepsilon} T^{-1+2\varepsilon} \int_{\frac{1}{2}}^{\alpha} \left(\frac{x}{Y^2} \right)^{\sigma} d\sigma$$
$$\ll x T^{-1+2\varepsilon} Y^{-3/2+\varepsilon} + x^{1/2} T^{-1+2\varepsilon} Y^{-1/2+\varepsilon} + x^{\beta} T^{-2\beta+2\varepsilon} Y^{1/2-2\beta+\varepsilon}.$$
(20)

Consider the integral $S_{2,2}$. Let $T \ge T_0$, where T_0 is a sufficiently large positive number. Using (17), and Lemmas 3 and 4, we deduce

$$S_{2,2} \ll x^{\beta} Y^{1/2 - 2\beta + \varepsilon} + x^{\beta} \int_{T_0 \le |t| \le T} \frac{|\chi(\beta + it)|^2 |\zeta(1 - \beta - it)|^2 |g_Y(2\beta + 2it)|}{1 + |t|} dt$$

$$\ll x^{\beta} T^{1 - 2\beta + 2\varepsilon} Y^{\frac{1}{2} - 2\beta + \varepsilon}.$$
(21)

Alternative expressions for the first and second sums on the right-hand side of (18) are

$$\operatorname{Res}_{s=1} \frac{\zeta^2(s)}{\zeta^2(2s)} \frac{x^s}{s} - \operatorname{Res}_{s=1} \zeta^2(s) g_Y(2s) \frac{x^s}{s}.$$

Therefore, by substituting (18)–(20) and (21) into (16) and setting $T = x^{\frac{1}{2}}Y^{\frac{4\beta-1}{4-4\beta}-\varepsilon}$, we obtain

$$S_{\mu}(x) = \frac{x}{\zeta^{2}(2)} \left(\log x + 2\gamma - 1 - 4\frac{\zeta'(2)}{\zeta(2)} \right) + \sum_{\ell \le Y} \mu * \mu(\ell) \Delta\left(\frac{x}{\ell^{2}}\right) + O\left(x^{\frac{1}{2} + \varepsilon} Y^{\frac{1 - 4\beta}{4 - 4\beta} + \varepsilon}\right),$$

where $x^{\varepsilon} < Y < x^{\frac{1}{2}-\varepsilon}$. This completes the proof.

3.4 Proof of Corollary 1

Given that

$$\frac{1-4\beta}{4-4\beta} = -\frac{1}{4}, \quad \left(\text{taking } \beta = \frac{2}{5}\right),$$

the error term in Theorem 3 can be estimated as

$$\ll x^{1/2+\varepsilon}Y^{-1/4+\varepsilon}.$$
(22)

This allows us to directly estimate the sum in Theorem 3, which is

$$\sum_{\ell \le Y} \mu * \mu(\ell) \Delta\left(\frac{x}{\ell^2}\right) \ll x^{\theta + \varepsilon} Y^{1 - 2\theta + \varepsilon}, \tag{23}$$

by applying (11). From (22) and (23), setting $Y = x^{\frac{2-4\theta}{5-8\theta}}$, with $\theta = \frac{131}{416}$ we obtain

$$E_{\mu}(x) \ll x^{\frac{439}{1032} + \varepsilon}.$$

This completes the proof.

3.5 Proof of Theorem 4

We aim to explore the mean square estimate for $E_{\mu}(x)$ under the Riemann Hypothesis. Our analysis begins with (5), where setting k = 2 yields the following relation

$$\sum_{n\geq 1} \frac{f_{\mu}(n)}{n^s} = \frac{\zeta^2(s)}{\zeta^2(2s)}$$

Employing the Perron formula, as detailed in Section 2, we derive that

$$S_{\mu}(x) = \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \frac{\zeta^2(s)}{\zeta^2(2s)} \frac{x^s}{s} ds.$$
 (24)

Upon shifting the line of integration in (24) to some $1/2 < \alpha < 1$, we encounter a pole at s = 1 within the integrand. Using the Cauchy residue theorem, we get

$$S_{\mu}(x) = \operatorname{Res}_{s=1} \frac{\zeta^{2}(s)}{\zeta^{2}(2s)} \frac{x^{s}}{s} + \frac{1}{2\pi i} \int_{\alpha-i\infty}^{\alpha+i\infty} \frac{\zeta^{2}(s)}{\zeta^{2}(2s)s} \left(\frac{1}{x}\right)^{-s} ds,$$

As previously discussed, the residue at s = 1 is given by

$$\operatorname{Res}_{s=1} \frac{\zeta^2(s)}{\zeta^2(2s)} \frac{x^s}{s} = \frac{x}{\zeta^2(2)} \left(\log x + 2\gamma - 1 - 4\frac{\zeta'(2)}{\zeta(2)} \right).$$

We have

$$E_{\mu}(x) = \frac{1}{2\pi i} \lim_{T \to \infty} \int_{\alpha - iT}^{\alpha + iT} \frac{\zeta^2(s)}{\zeta^2(2s)s} \left(\frac{1}{x}\right)^{-s} ds.$$
 (25)

We define α_0 as the infimum of $\alpha > 0$ for which

$$\int_{-\infty}^{\infty} \frac{|\zeta(\alpha + it)|^4}{|\zeta(2\alpha + 2it)|^4 |\alpha + it|^2} dt \ll 1.$$
 (26)

Our goal is to show that for $\alpha > \alpha_0$ and approaching to 1, the following holds:

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{|\zeta(\alpha+it)|^4}{|\zeta(2\alpha+2it)|^4 |\alpha+it|^2} dt = \int_0^{\infty} E_{\mu}^2(x) x^{-2\alpha-1} dx.$$
(27)

From (25), since $\zeta^2(s)\zeta^{-2}(2s)s^{-1} \to 0$ uniformly in the strip as $t \to \pm \infty$, and by integrating over the rectangular path passing through the vertices, $\alpha' - iT$, $\alpha - iT$, $\alpha + iT$ and $\alpha' + iT$ where $\alpha_0 < \alpha' < \alpha < 1$, we confirm that (25) is valid for any $\alpha > \alpha_0$. By substituting x with 1/x, and applying Parseval's identity, we achieve the desired equality in (27). Consequently, if (26) is satisfied for some α ($\alpha_0 < \alpha < 1$), then for any $T \ge 5$ we have

$$\int_{T/2}^{T} E_{\mu}^{2}(x) dx \ll T^{2\alpha+1}.$$

To finalize the proof of our theorem, we must demonstrate that (26) is attainable. Assuming $\alpha_0 < \alpha < 1/2$ as a parameter to be precisely chosen, and for any $T \ge 5$, we employ Lemma 3 to obtain

$$\begin{split} L(T) &:= \int_{T/2}^{T} \frac{|\zeta(\alpha+it)|^4}{|\zeta(2\alpha+2it)|^4 |\alpha+it|^2} dt \\ &= \int_{T/2}^{T} \frac{|\chi(\alpha+it)|^4 |\zeta(1-\alpha+it)|^4}{|\zeta(2\alpha+2it)|^4 |\alpha+it|^2} dt \\ &\ll \int_{T/2}^{T} \frac{t^{2-4\alpha} |\zeta(1-\alpha+it)|^4}{|\zeta(2\alpha+2it)|^4 |\alpha+it|^2} dt. \end{split}$$

By setting $\alpha = 1/4 + 3\varepsilon$, for any sufficiently small $\varepsilon > 0$ (choosing $\varepsilon := \frac{c}{\log \log T}$ with a positive constant c), and applying Lemma 4, we deduce

$$L(T) \ll \int_{T/2}^{T} \frac{|\zeta(\frac{3}{4} - 3\varepsilon - it)|^4}{|\zeta(\frac{1}{2} + 6\varepsilon + 2it)|^4 t^{1+12\varepsilon}} dt \ll \int_{T/2}^{T} \frac{1}{t^{1+4\varepsilon}} dt \ll 1.$$

Therefore, we conclude that

$$\begin{split} \int_{1}^{T} E_{\mu}^{2}(u) du &= \sum_{k \leq \frac{\log T}{\log 2}} \int_{2^{k-1}}^{2^{k}} E_{\mu}^{2}(u) du \ll T^{3/2 + 6\varepsilon} \\ &\ll T^{3/2} \exp\left(A \frac{\log T}{\log \log T}\right), \end{split}$$

with a positive real number A.

Acknowledgement

The first author is supported by JSPS Grant-in-Aid for Scientific Research (C)(21K03205). The second author is supported by the Austrian Science Fund (FWF): Project N. 35863.

References

- [1] J. L. Hafner and A. Ivić, On the mean-square of the Riemann zeta-function on the critical line, *J. Number Theory*, 32 (1989), 151–191.
- [2] M. N. Huxley, Exponential sums and lattice points. III, Proc. London Math. Soc., 87 (2003) 591–609.
- [3] S. Inoue and I. Kiuchi, On sums of gcd-sum functions, Int. J. Number Theory, 19 (2023) 593-619.
- [4] A. Ivić, *The Riemann Zeta-Function*, Theory and Applications Dover publications, Inc. Mineola, New York (1985).
- [5] E. Krätzel, W. G. Nowak and L. Tóth, On certain arithmetic functions involving the greatest common divisor, *Cent. Eur. J. Math.*, 10 (2012), 761–774.
- H. L. Montgomery and R. C. Vaughan, *The distribution of squarefree numbers*, in;
 H. Halberstam, C. Hooley (Eds.), Recent Progress in Analytic Number Theory, vol.1, Academic Press, London, 1981, pp.247–256.
- [7] H. L. Montgomery and R. C. Vaughan, *Multiplicative Number Theory I. Classi*cal Theory, Cambridge Studies in Advanced Mathematics, Cambridge University press, 2007.
- [8] W. G. Nowak, On the average order of arithmetic functions connected with the Riemann zeta-function, in Analysis, Geometry and Groups: A Riemann Legacy Volume (editted by H. M. Srivastava and Th. M. Rassias.), page 449– 470, Hadronic Press Collect. Orig. Artic. Hadronic Press, Palm Harbor, Florida, 1993.

- [9] G. Tenenbaum, Introduction to Analytic and Probabilistic Number Theory, Graduate Studies in Mathematics, 163 AMS, 2008.
- [10] E. C. Titchmarsh, The Theory of the Riemann Zeta-Function (revised by D. R. Heath-Brown), Oxford University Press, 1986.
- [11] D. Yang, Mean values of ratios of the Riemann zeta function, arXiv:2307.08091v1 [math.NT] 16 Jul 2023.

Isao Kiuchi: Department of Mathematical Sciences, Faculty of Science, Yamaguchi University, Yoshida 1677-1, Yamaguchi 753-8512, Japan. e-mail: kiuchi@yamaguchi-u.ac.jp

Sumaia Saad Eddin: Johann Radon Institute for Computational and Applied Mathematics, Austrian Academy of Sciences, Altenbergerstrasse 69, 4040 Linz, Austria. e-mail: sumaia.saad-eddin@ricam.oeaw.ac.at