Quanyuan Chen School of Information and Engineering, Jingdezhen Ceramic University, Jingdezhen 333000. China. cqy0798@163.com Yaqi Li School of Information and Engineering, Jingdezhen Ceramic University, Jingdezhen 333000. China. 1yq8532@163.com

March 19, 2024

ABSTRACT

Let X be a Banach algebra and B(X) be the set of all bounded linear operators on X. Suppose that $\alpha : B(X) \to B(X)$ is an automorphism. We say that a mapping δ from B(X) into itself is derivable at $G \in B(X)$ if $\delta(G) = \alpha(A) \delta(B) + \delta(A) \alpha(B)$ for all $A, B \in B(X)$ with AB = G. We say that an element $G \in B(X)$ is an (α, α) -all derivable point of B(X) if every (α, α) -derivable mapping δ at G is an (α, α) - derivation. In this paper, we show that every (α, α) -derivable mapping at a nonzero element in B(X) is an (α, α) -derivation.

1 Introduction

Let \mathcal{A} be an algebra over \mathbb{F} , where \mathbb{F} is either the real field \mathbb{R} or the complex field \mathbb{C} , and $\delta : \mathcal{A} \to \mathcal{A}$ be an additive (linear) mapping. Recall that δ is called a derivation if $\delta(AB) = \delta(A)B + A\delta(B)$ for all $A, B \in \mathcal{A}$, and δ is called a Jordan derivation if $\delta(A^2) = \delta(A)A + A\delta(A)$ for all $A \in \mathcal{A}$. Suppose that α and β are ring automorphisms on \mathcal{A} . An additive mapping $\delta : \mathcal{A} \to \mathcal{A}$ is called an additive (α, β) -derivation if $\delta(AB) = \alpha(A)\delta(B) + \delta(A)\beta(B)$ for all $A, B \in \mathcal{A}$. An additive mapping $\delta : \mathcal{A} \to \mathcal{A}$ is called an additive (α, β) -derivation if $\delta(AB) = \alpha(A)\delta(B) + \delta(A)\beta(B)$ for all $A, B \in \mathcal{A}$.

Derivations are very important mappings both in theory and applications and have been studied extensively. In [1], Jacobson and Rickart proved that every Jordan derivation on an unital ring R is a derivation. In [2], Herstein proved that every Jordan derivation on a 2-torsion free prime ring is a derivation. In [3], Xue et al. proved that if $\mathcal{A} = Alg\mathcal{N}$ is a nest algebra and $\mathcal{M} = B(X)$, then a linear mapping $\delta : \mathcal{A} \to \mathcal{M}$ is Jordan derivable at a nontrivial projection $P \in Alg\mathcal{N}$ if and only if it is derivation. In [4], Jing and Lu studied (α, β) -derivations in operator algebras and characterized (α, β) -derivations on B(X).

In recent years, some scholars have been interested in characterizing the mapping as derivable at some point. Suppose that G is an element in \mathcal{A} . We say that δ is a derivable mapping at G if $\delta(G) = \delta(A)B + A\delta(B)$ for all $A, B \in \mathcal{A}$ with AB = G. An element $G \in \mathcal{A}$ is called an all-derivable point in \mathcal{A} if every derivable mapping at G is a derivation. Jing et al. in [5] proved that every linear mapping derivable at zero point with $\delta(I) = 0$ is an inner derivation of the nest algebra on Hilbert spaces. Zhu proved in [6] that every invertible operator on the nest algebra $Alg\mathcal{N}$ is an all-derivable point of the nest algebra under the strong operator topology. In [7], Jing proved that the zero point is a generalized Jordan all-derivable point of B(H) if H is infinite-dimensional and I is a Jordan all-derivable point of B(H) of a Hilbert space. He et al. proved in [8] that every derivable mapping at a nonzero element is a derivation in B(H).

Inspired by [8], we consider (α, α) -all derivable points for an automorphism α on \mathcal{A} . We say that δ is (α, α) -derivable at point $G \in \mathcal{A}$ if $\delta(G) = \alpha(A)\delta(B) + \delta(A)\alpha(B)$ for all $A, B \in \mathcal{A}$ with AB = G. It is obvious that (I, I)-derivations are usual derivations, and (α, α) -all derivable points are all derivable points if α is the identity mapping. In [9], Jiao and Hou showed that if δ is generalized (α, β) -derivable at the zero, then δ is an additive generalized (α, β) -derivation, where $Alg\mathcal{N}$ is a nest algebra on a (real or complex) Banach space.

The purpose of this paper is to characterize the mapping (α, α) -derivable at a nonzero element in B(X). In section 2, we prove that every (α, α) -derivable mapping at a nonzero element in B(X) is an (α, α) -derivation. Moreover, we also prove that every (α, α) -generalized derivable mapping at a nonzero element G in B(X) (that is, $\delta(G) = \delta(A)\alpha(B) + \alpha(A)\delta(B) - \alpha(A)\delta(I)\alpha(B)$ for all $A, B \in \mathcal{A}$ with AB = G.) is a (α, α) -general derivation.

The following lemma will get repeated use.

Lemma 1 (Lemma 1.1 in [8]). Let \mathcal{R} be a semiprime ring. *b*, *p*, *q* are fixed elements in \mathcal{R} .

(1) If babab = 0 for all $a \in \mathcal{R}$, then b = 0.

(2) If paqbpaq = 0 for all $a \in \mathcal{R}$, then paq = 0.

Lemma 2 (Corollary 1 in [10]). Let \mathcal{R} be a 2-torsion free semiprime ring and let α , β be an automorphism of \mathcal{R} . Then every Jordan (α, β) -derivation of \mathcal{R} is an (α, β) -derivation.

2 Main theorem

In this section, X is a Banach space on \mathbb{F} , where \mathbb{F} is either the real field \mathbb{R} or the complex field \mathbb{C} . B(X) is an algebra of all bounded linear operators on X. For convenience, we use the symbol \mathcal{A} to represent B(X) in the following proof. I is the identity element of \mathcal{A} , and X^* is the set of all bounded linear functionals from X into \mathbb{F} .

For any $x \in X$, $f \in X^*$, the rank one operator on X is defined as $x \otimes f(y) = f(y)x$ for all $y \in X$. Clearly, $x \otimes f$ is a rank one idempotent operator when f(x) = 1. In this case, we have $(x \otimes f) A (x \otimes f) = \lambda x \otimes f$ for some $\lambda = f(AX)$.

According to the Peirce decomposition, for a nontrivial idempotent P_1 in \mathcal{A} and $P_2 = I - P_1$, we denote $\mathcal{A}_{11} = P_1 \mathcal{A} P_1$, $\mathcal{A}_{12} = P_1 \mathcal{A} P_2$, $\mathcal{A}_{21} = P_2 \mathcal{A} P_1$, $\mathcal{A}_{22} = P_2 \mathcal{A} P_2$, respectively, then \mathcal{A} can be decomposed as $\mathcal{A} = \mathcal{A}_{11} + \mathcal{A}_{12} + \mathcal{A}_{21} + \mathcal{A}_{22}$. For any operator $A \in \mathcal{A}$, it can be decomposed as $A = A_{11} + A_{12} + A_{21} + A_{22}$, where $A_{ij} \in \mathcal{A}_{ij}$, i, j = 1, 2. Inspired by [10], we consider (α, α) -all derivable points for an automorphism α on \mathcal{A} . Let $Q_1 = \alpha (P_1)$ and $Q_2 = \alpha (P_2)$, then Q_1 and Q_2 are nontrivial idempotents in \mathcal{A} with $Q_2 = I - Q_1$.

The following are our main results of the paper.

Theorem 2.1. Suppose that G is a nonzero element in $\mathcal{A} = B(X)$, α is an automorphism on \mathcal{A} , and δ is a linear mapping on \mathcal{A} . If δ is (α, α) -derivable at G, which means that $\delta(G) = \delta(A)\alpha(B) + \alpha(A)\delta(B)$ for all $A, B \in \mathcal{A}$ with AB = G, then δ is an (α, α) -derivation.

Proof. If $\overline{ran\alpha(G)} = X$, where $\overline{ran\alpha(G)}$ is the closure of the range of $\alpha(G)$, then $\alpha(G)$ is a right separating point of B(X). By [11], δ is a Jordan (α, α) -derivation, then δ is an (α, α) -derivation by Lemma 2.

Next, it is only necessary to prove the case in which $\overline{ran\alpha(G)} \neq X$. Take a nonzero element $\alpha(x_0) \in X$ which is not in $\alpha(G)$. According to the Hahn-Banach Theorem, consider $f_0 \in \alpha(X^*)$ such that $f_0(\alpha(x_0)) = 1$ and $f_0(\operatorname{ran} \alpha(G)) = 0$. Let $Q_2 = \alpha(x_0) \otimes f_0$, $Q_1 = I - Q_2$, then we have $Q_2\alpha(G) = 0$, $Q_1\alpha(G) = \alpha(G)$.

Assume that A, B are arbitrary elements in \mathcal{A} , A_{ij} , B_{ij} are any element in \mathcal{A}_{ij} , i, j = 1, 2, and t, s are any nonzero elements in \mathbb{F} . According to the condition that Q_1 is surjective on $\alpha(\mathcal{A}_{11})$, it is known that A_{11} is invertible in \mathcal{A}_{11} , which provides the next decomposition of G.

First, we define a new linear mapping $\phi : \mathcal{A} \to \mathcal{A}$ by $\phi(\mathcal{A}) = \delta(\mathcal{A}) + [\alpha(\mathcal{A}), D]$, where $D = \alpha(P_1) \delta(P_2) \alpha(P_2) - \alpha(P_2) \delta(P_2) \alpha(P_1)$. According to the assumption that $\delta : \mathcal{A} \to \mathcal{A}$ is (α, α) -derivable at the point G, ϕ is also an (α, α) -derivable at the point G by linear operations and satisfies $\phi(P_2) \in \alpha(\mathcal{A}_{11}) + \alpha(\mathcal{A}_{22})$. Now, it is only necessary to prove that ϕ is an (α, α) -derivation to obtain that δ is an (α, α) -derivation. The point G is written as:

$$(A_{11} + tsA_{11}A_{12})(A_{11}^{-1}G - sA_{12}A_{2i} + t^{-1}A_{2i}) = G, i = 1, 2.$$

According to ϕ is (α, α) -derivable at the point G gives:

$$\phi(G) = \phi \left(A_{11} + tsA_{11}A_{12}\right) \alpha \left(A_{11}^{-1}G - sA_{12}A_{2i} + t^{-1}A_{2i}\right) + \alpha \left(A_{11} + tsA_{11}A_{12}\right) \phi \left(A_{11}^{-1}G - sA_{12}A_{2i} + t^{-1}A_{2i}\right).$$
(1)

According to the linear property, we expand Equation 1 and simplify the coefficients of t^{-1} , s, and ts^2 , respectively. We obtain the following three equations:

$$\phi(A_{11}) \alpha(A_{2i}) + \alpha(A_{11}) \phi(A_{2i}) = 0$$
(2)

$$(A_{11}A_{12})\alpha(A_{2i}) + \alpha(A_{11}A_{12})\phi(A_{2i}) = \phi(A_{11})\alpha(A_{12}A_{2i}) + \alpha(A_{11})\phi(A_{12}A_{2i})$$
(3)

$$\phi(A_{11}A_{12})\,\alpha(A_{12}A_{2i}) + \alpha(A_{11}A_{12})\,\phi(A_{12}A_{2i}) = 0. \tag{4}$$

By the linear property of ϕ and the decomposition of arbitrary element A and B in A, to prove that ϕ is an (α, α) derivation it is sufficient to show that the following equation holds:

$$\phi(A_{ij}B_{sk}) = \phi(A_{ij}) \alpha(B_{sk}) + \alpha(A_{ij}) \phi(B_{sk})$$
(5)

where $1 \leq i, j, s, k \leq 2$.

 ϕ

Claim 1: $\phi(A_{22}) = 0$.

We consider $A_{11} = P_1$ and $A_{2i} = P_2$ in Equation 2, and obtain $\phi(P_1) \alpha(P_2) + \alpha(P_1) \phi(P_2) = 0$. We multiply the right-hand side of the above equation by $Q_1 = \alpha(P_1)$ and obtain $\alpha(P_1) \phi(P_2)$ $\alpha(P_1) = 0$. By the known condition $\phi(P_2) \in \alpha(A_{11}) + \alpha(A_{22})$, we obtain $\phi(P_2) \in \alpha(A_{22})$, so $\phi(P_2) = \lambda \alpha(P_2)$ for some $\lambda \in \mathbb{F}$ and $\phi(P_1) \alpha(P_2) = 0$. We consider $A_{2i} = P_2$ in Equation 2 again and obtain

$$\phi(A_{11})\,\alpha(P_2) = 0. \tag{6}$$

Letting $A_{11} = P_1$ and $A_{2i} = A_{21}$ in Equation 2, we obtain that

$$\alpha(P_1)\phi(A_{21}) = 0. (7)$$

Cunsidering that $A_{11} = P_1$ and $A_{2i} = P_2$ in Equation 3 and $\phi(P_2) = \lambda \alpha(P_2), \lambda \in \mathbb{F}$, we get that

$$\phi(A_{12}) \alpha(P_2) + \lambda \alpha(A_{12}) = \phi(P_1) \alpha(A_{12}) + \alpha(P_1) \phi(A_{12}).$$
(8)

Multiply the right-hand side of Equation 8 by $Q_1 = \alpha(P_1)$, we obtain that

$$\alpha\left(P_{1}\right)\phi\left(A_{21}\right)\alpha\left(P_{1}\right) = 0.$$
(9)

We multiply the left-hand side of Equation 8 by $Q_1 = \alpha(P_1)$, and associated with Equation 9, we obtain $\alpha(P_1)\phi(P_1)\alpha(A_{12}) = \lambda\alpha(A_{12})$. It is known that \mathcal{A} is simple, so the $\alpha(\mathcal{A})$ is also simple, and we get:

$$\alpha(P_1)\phi(P_1)\alpha(P_1) = \lambda\alpha(P_1).$$
(10)

As IG = G and ϕ is an (α, α) -derivable at point G, we have $\phi(I)\alpha(G) + \alpha(I)\phi(G) = \phi(G)$, then $\phi(I)\alpha(G) = 0$. Since $Q_2\alpha(G) = \alpha(P_2)\alpha(G) = 0$ and $\phi(P_2) = \lambda\alpha(P_2)$, then we have $\phi(P_1)\alpha(G) = \phi(I)\alpha(G) - \phi(P_2)\alpha(G) = 0$ $-\lambda \alpha(P_2) \alpha(G) = 0$. We multiply the righthand side of Equation 10 by $\alpha(G)$, due to $Q_1 \alpha(G) = \alpha(P_1) \alpha(G) = \alpha(Q_1) \alpha(G)$ $\alpha(G)$, we get $\lambda\alpha(G) = \lambda\alpha(p_1)\overline{\alpha(G)} = \lambda\alpha(P_1)\phi(P_1)\alpha(G) = 0$. Since $G \neq 0$ and $\alpha(G) \neq 0$, which in turn gives $\lambda = 0$. Now we prove:

$$\alpha(P_1)\phi(P_1)\alpha(P_1) = \lambda\alpha(P_1) = 0 \tag{11}$$

and $\phi(P_2) = \lambda \alpha(P_2) = 0$, so we obtain $\phi(A_{22}) = 0$.

Claim 2:
$$\phi(A_{11}) \subseteq \alpha(A_{11})$$

We consider $A_{11} = P_1$ and $A_{2i} = P_2$ in Equation 4, and we get

$$\phi(A_{12}) \alpha(A_{12}) + \alpha(A_{12}) \phi(A_{12}) = 0.$$
(12)

The left multiplication of Equation 12 by $Q_1 = \alpha (P_1)$ and associated with Equation 9, then obtain:

$$\alpha(A_{12})\phi(A_{12}) = 0 \tag{13}$$

and

$$\phi(A_{12})\,\alpha(A_{12}) = 0. \tag{14}$$

According to Equation 13, we can obtain $\alpha(B_{12}) \phi(B_{12}) = 0$ and $\alpha(A_{12} + B_{12}) \phi(A_{12} + B_{12}) = 0$, then we have:

$$\alpha(A_{12})\phi(B_{12}) + \alpha(B_{12})\phi(A_{12}) = 0.$$
(15)

We multiply the right-hand side of Equation 15 by $\alpha(B_{12})$ and associated with Equation 14, we can obtain $\alpha(B_{12})\phi(A_{12})\alpha(B_{12}) = 0$. By Lemma 1 we have $\alpha(P_2)\phi(A_{12})\alpha(P_1) = 0$. Associating with Equation 9, we get:

$$\phi(A_{12})\,\alpha(P_1) = 0. \tag{16}$$

We consider $A_{2i} = P_2$ in Equation 3 then associated with $\phi(P_2) = 0$ and Equation 16 yields:

$$\phi(A_{11}A_{12}) = \phi(A_{11}) \alpha(A_{12}) + \alpha(A_{11}) \phi(A_{12}).$$
(17)

So, we have:

$$\phi (A_{11}B_{11}A_{12}) = \phi (A_{11}B_{11}) \alpha (A_{12}) + \alpha (A_{11}B_{11}) \phi (A_{12}) = \phi (A_{11}) \alpha (B_{11}A_{12}) + \alpha (A_{11}) \phi (B_{11}A_{12}) = \phi (A_{11}) \alpha (B_{11}A_{12}) + \alpha (A_{11}) \phi (B_{11}) \alpha (A_{12}) + \alpha (A_{11}B_{11}) \phi (A_{12}).$$

Since $\phi(A_{11}) \alpha(P_2) = 0$, the above equation implies that

$$\phi(A_{11}B_{11}) = \phi(A_{11})\alpha(B_{11}) + \alpha(A_{11})\phi(B_{11}).$$
(18)

Letting $A_{11} = P_1$ in Equation 18 we obtain that

$$\phi(B_{11}) = \phi(P_1) \alpha(B_{11}) + \alpha(P_1) \phi(B_{11}).$$
(19)

We consider $A_{11} = P_1$ and $A_{2i} = A_{21}$ in Equation 4, associated with Equation 16 we obtain $\alpha(A_{12}) \phi(A_{12}A_{21}) = 0$ and associated with Equation 19 we obtain $\alpha(A_{12}) \phi(P_1) \alpha(A_{12}A_{21}) = 0$. Since $\alpha(\mathcal{A})$ is simple, we have $\alpha(A_{12}) \phi(P_1) \alpha(A_{12}) = 0$. By Lemma 1 and α is an automorphism on \mathcal{A} we have that $\alpha(P_2) \phi(P_1) \alpha(P_1) = 0$. We multiply the left-hand side of Equation 19 by $Q_2 = \alpha(P_2)$ obtains $\alpha(P_2) \phi(B_{11}) = 0$. Together with $\phi(B_{11}) \alpha(P_2) = 0$, this implies that $\phi(\mathcal{A}_{11}) \subseteq \alpha(\mathcal{A}_{11})$.

Claim 3:
$$\phi(\mathcal{A}_{12}) \subseteq \alpha(\mathcal{A}_{12})$$

We consider $A_{11} = P_1$ in Equation 17 and multiply the left-hand side by $Q_2 = \alpha(P_2)$, associated with $\phi(A_{11}) \subseteq \alpha(A_{11})$, then we wet that $\alpha(P_2)\phi(A_{12}) = 0$. Togeter with Equation 6: $\phi(A_{12})\alpha(P_1) = 0$, we obtain that $\phi(A_{12}) \subseteq \alpha(A_{12})$.

Claim 4: $\phi(\mathcal{A}_{21}) \subseteq \alpha(\mathcal{A}_{21})$.

Since $\phi(P_1) \subseteq \alpha(A_{11})$ and $\alpha(P_1)\phi(P_1)\alpha(P_1) = 0$ according to Equation 11 yields $\phi(P_1) = 0$. We consider $A_{11} = P_1$ and $A_{2i} = A_{21}$ in Equation 3, associated with $\phi(A_{11}) \subseteq \alpha(A_{11})$ and $\phi(P_1) = 0$, we obtain that

$$\phi(A_{12}A_{21}) = \phi(A_{12}) \alpha(A_{21}) + \alpha(A_{12}) \phi(A_{21}).$$
(20)

We multiply the right-hand side of Equation 20 by $Q_2 = \alpha (P_2)$ and yield $\alpha (A_{12}) \phi (A_{21}) \alpha (P_2) = 0$, which in turn gives $\alpha (P_2) \phi (A_{21}) \alpha (P_2) = 0$. $\alpha (P_1) \phi (A_{21}) = 0$ and Equation 7, implies that $\phi (A_{21}) \subseteq \alpha (A_{21})$.

Equation 5 obviously holds when $j \neq s$ according to Claims 1-4. Next, it is only necessary to prove the case when j = s, which is the following eight cases:

$$(1)\phi (A_{11}B_{11}) = \phi (A_{11}) \alpha (B_{11}) + \alpha (A_{11}) \phi (B_{11});$$

$$(2)\phi (A_{11}B_{12}) = \phi (A_{11}) \alpha (B_{12}) + \alpha (A_{11}) \phi (B_{12});$$

$$(3)\phi (A_{12}B_{21}) = \phi (A_{12}) \alpha (B_{21}) + \alpha (A_{12}) \phi (B_{21});$$

$$(4)\phi (A_{12}B_{22}) = \phi (A_{12}) \alpha (B_{22}) + \alpha (A_{12}) \phi (B_{22});$$

$$(5)\phi (A_{22}B_{21}) = \phi (A_{22}) \alpha (B_{21}) + \alpha (A_{22}) \phi (B_{21});$$

$$(6)\phi (A_{22}B_{22}) = \phi (A_{22}) \alpha (B_{22}) + \alpha (A_{22}) \phi (B_{22});$$

$$(7)\phi (A_{21}B_{11}) = \phi (A_{21}) \alpha (B_{11}) + \alpha (A_{21}) \phi (B_{11});$$

$$(8)\phi (A_{21}B_{12}) = \phi (A_{21}) \alpha (B_{12}) + \alpha (A_{21}) \phi (B_{12}).$$

Case (1), case (2), and case (3) can be obtained from Equation 18, Equation 17, and Equation 20, respectively.

Since $B_{22} = \mu Q_2$ for some $\mu \in \mathbb{F}$ and $\phi(B_{22}) = 0$, then case (4) holds. Similarly, it follows that case (5) holds. Additionally, case (6) is trivial because $\phi(A_{22}) = \{0\}$.

According to case (1) and case (2) we can get:

$$\phi (A_{12}A_{21}B_{11}) = \phi (A_{12}) \alpha (A_{21}B_{11}) + \alpha (A_{12}) \phi (A_{21}B_{11}) = \phi (A_{12}A_{21}) \alpha (B_{11}) + \alpha (A_{12}A_{21}) \phi (B_{11}) = \phi (A_{12}) \alpha (A_{21}B_{11}) + \alpha (A_{12}) \phi (A_{21}) \alpha (B_{11}) + \alpha (A_{12}A_{21}) \phi (B_{11})$$

Claim 4 implies case (7).

Case (8) can be derived similarly from the following equation:

$$\begin{split} \phi \left(A_{12}A_{21}B_{12} \right) &= \phi \left(A_{12} \right) \alpha \left(A_{21}B_{12} \right) + \alpha \left(A_{12} \right) \phi \left(A_{21}B_{12} \right) \\ &= \phi \left(A_{12}A_{21} \right) \alpha \left(B_{12} \right) + \alpha \left(A_{12}A_{21} \right) \phi \left(B_{12} \right) \\ &= \phi \left(A_{12} \right) \alpha \left(A_{21}B_{12} \right) + \alpha \left(A_{12} \right) \phi \left(A_{21} \right) \alpha \left(B_{12} \right) + \alpha \left(A_{12}A_{21} \right) \phi \left(B_{12} \right) . \end{split}$$

So ϕ is an (α, α) -derivation on \mathcal{A} , and δ is an (α, α) -derivation on \mathcal{A} .

Corollary 2.1. Suppose that G is a nonzero element in $\mathcal{A} = B(X)$, α is an automorphism on \mathcal{A} , and δ is a linear mapping on \mathcal{A} . If δ is (α, α) -generalized derivable at G, which means that $\delta(G) = \delta(A)\alpha(B) + \alpha(A)\delta(B) - \alpha(A)\delta(I)\alpha(B)$ for all $A, B \in \mathcal{A}$ with AB = G, then δ is an (α, α) -generalized derivation.

Proof. We define a new linear mapping $\phi : \mathcal{A} \to \mathcal{A}$ by $\phi(A) = \delta(A) - \delta(I)\alpha(A)$. From the equations

$$\phi(AB) = \delta(AB) - \delta(I)\alpha(AB)$$

and

 $\phi(A)\alpha(B) + \alpha(A)\phi(B) = (\delta(A) - \delta(I)\alpha(A))\alpha(B) + \alpha(A)(\delta(B) - \delta(I)\alpha(B)),$

 δ is (α, α) -generalized derivable at G if and only if ϕ is (α, α) -derivable at G. Then, δ is (α, α) -generalized derivable if and only if ϕ is (α, α) -derivable.

Because ϕ is (α, α) -derivable at G, by Theorem 2.1, ϕ is (α, α) -derivable. Then δ is an (α, α) -generalized derivation on \mathcal{A} .

References

- N. Jacobson and C. E. Rickart, "Jordan homomorphisms of rings," *Transactions of the American Mathematical Society*, vol. 69, pp. 479–502, 1950.
- [2] I. N. Herstein, "Jordan derivations of prime rings," *Proceedings of the American Mathematical Society*, vol. 8, no. 6, pp. 1104–1110, 1957.
- [3] X. Tianjiao, A. Runling, and H. Jinchuan, "Characterization of derivations on b (x) by local actions," *Acta Mathematica Scientia*, vol. 37, no. 3, pp. 668–678, 2017.
- [4] W. Jing and S. Lu, "Topological reflexivity of the spaces of (α, β)-derivations on operator algebras," Studia Mathematica, vol. 2, no. 156, pp. 121–131, 2003.
- [5] W. Jing, S. Lu, and P. Li, "Characterisations of derivations on some operator algebras," *Bulletin of the Australian Mathematical Society*, vol. 66, no. 2, pp. 227–232, 2002.
- [6] J. Zhu, "All-derivable points of operator algebras," *Linear algebra and its applications*, vol. 427, no. 1, pp. 1–5, 2007.
- [7] W. Jing, "On jordan all-derivable points of b (h)," Linear Algebra Appl, vol. 430, no. 4, pp. 941–946, 2009.
- [8] J. He, H. Zhao, and G. An, "Generalized derivable mappings and centralizable mappings on b (x)," *Linear and Multilinear Algebra*, vol. 71, no. 17, pp. 2719–2727, 2023.
- [9] M. Jiao and J. Hou, "Additive maps derivable or jordan derivable at zero point on nest algebras," *Linear Algebra and its Applications*, vol. 432, no. 11, pp. 2984–2994, 2010.
- [10] C.-K. Liu and W.-K. Shiue, "Generalized jordan triple (θ , ϕ) -derivations on semiprime rings," *Taiwanese Journal of Mathematics*, vol. 11, no. 5, pp. 1397–1406, 2007.
- [11] J. Li and J. Zhou, "Characterizations of jordan derivations and jordan homomorphisms," *Linear and Multilinear Algebra*, vol. 59, no. 2, pp. 193–204, 2011.