# QUATERNIONIC MAHLER MEASURE

WEIJIA WANG AND HAO ZHANG

ABSTRACT. We introduce the quaternionic Mahler measure for non-commutative polynomials, extending the classical complex Mahler measure. We establish the existence of quaternionic Mahler measure for slice regular polynomials in one and two variables. We study the quaternionic Mahler measure for real and slice regular polynomials, and consider the associated Lehmer problem. Various formulas of quaternionic Mahler measures are proved.

# 1. INTRODUCTION

Given a non-zero complex Laurent polynomial  $P \in \mathbb{C}[x_1^{\pm}, \ldots, x_n^{\pm}]$ , the complex (logarithmic) Mahler measure of P is defined as

$$\mathbf{m}_{\mathbb{C}}(P) \coloneqq \int_{\mathbb{T}^n(\mathbb{C})} \log |P(x_1, \dots, x_n)| \, \mathrm{d}\mu(\mathbb{T}^n(\mathbb{C})),$$

where  $\mathbb{T}^n(\mathbb{C}) = \{(x_1, \dots, x_n) \in \mathbb{C}^n \mid |x_1| = \dots = |x_n| = 1\}$  is the unit complex torus and  $d\mu(\mathbb{T}^n(\mathbb{C})) = (2\pi i)^{-n} \prod_{\ell=1}^n \frac{dx_\ell}{x_\ell}$  is its probability Haar measure.

The origins of Mahler measure can date back to the early 20th century when Lehmer [24] introduced it (of one variable) in his study of finding large prime numbers. Lehmer proposed what came to be known as the *Lehmer's conjecture*, which concerns the smallest non-zero Mahler measure that an irreducible polynomial in  $\mathbb{Z}[x]$  could attain. Later, Mahler [25] extended the concept to several complex variables in utilization to explore the heights of complex commutative polynomials.

In the early 1980s, Smyth [32] discovered the notable identity,

$$m_{\mathbb{C}}(1+x+y) = \frac{3\sqrt{3}}{4\pi}L(\chi_{-3},2) = L'(\chi_{-3},1),$$

where  $L(\chi_{-3}, s)$  is the Dirichlet *L*-function of the odd Dirichlet character of conductor 3. The machinery of relating the Mahler measure with *L*-functions and regulator, continues to be explored through the work of Boyd [4], Deninger [5] and

<sup>(</sup>Weijia Wang) MORNINGSIDE CENTER OF MATHEMATICS, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, P. R. CHINA

<sup>(</sup>Hao Zhang) School of Mathematics, Hunan University, Changsha 410082, P. R. China

E-mail addresses: weijiawang@amss.ac.cn, zhanghaomath@hnu.edu.cn.

Rodriguez-Villegas [31]. For a thorough survey about complex Mahler measure, the readers are also recommended to consult the book [3] of Brunault and Zudilin.

Variants of Mahler measure have also been proposed in many literatures. For example, the definition of Elliptic Mahler measure by Everest and ní Fhlathúin [8], Akatsuka's Zeta Mahler measure [1], and Pritsker's examination of the areal Mahler measure [28] by replacing the integration with area measure on the open unit complex disk (see also Pritsker [29], Lalín–Roy [22, 23]).

In this manuscript, let  $\mathbb{H}$  be the algebra of quaternion. For a non-commutative polynomial  $P(x_1, \ldots, x_n)$  with coefficients in  $\mathbb{H}$ , we consider the following quaternionic analogue of Mahler measure

$$\mathbf{m}_{\mathbb{H}}(P) \coloneqq \int_{\mathbb{T}^n(\mathbb{H})} \log |P(x_1, \dots, x_n)| \, \mathrm{d}\mu(\mathbb{T}^n(\mathbb{H})),$$

where  $\mathbb{T}^n(\mathbb{H}) = \{(x_1, \ldots, x_n) \in \mathbb{H}^n | |x_1| = \cdots = |x_n| = 1\}$  becomes the unit quaternion torus and  $d\mu(\mathbb{T}^n(\mathbb{H}))$  is again the probability Haar measure. Following [1], we may also define the corresponding quaternionic zeta Mahler measure of  $P(x_1, \ldots, x_n)$  by

$$Z_{\mathbb{H}}(P;s) \coloneqq \int_{\mathbb{T}^n(\mathbb{H})} |P(x_1,\ldots,x_n)|^s \, \mathrm{d}\mu(\mathbb{T}^n(\mathbb{H})).$$

The quaternionic Mahler measure shares many properties analogous to its complex counterpart. For example, given two quaternionic polynomials P and Q, one has  $m_{\mathbb{H}}(PQ) = m_{\mathbb{H}}(P) + m_{\mathbb{H}}(Q)$ , where PQ must be the *non-commutative product* of P and Q. It's worth noting that even for real polynomials, the quaternionic Mahler measure is not merely the sum of Mahler measures of monomials.

Our first result on monomials and real polynomials is summarized as follows.

**Theorem 1.1.** In the following formulas, let  $\alpha \in \mathbb{H}$  and  $\phi = \arccos(\operatorname{Re} \alpha/|\alpha|)$ . 1. For  $m_{\mathbb{H}}(x - \alpha)$ , we have

$$m_{\mathbb{H}}(x-\alpha) = \begin{cases} \frac{1}{4} |\alpha|^2 & \text{if } |\alpha| \le 1, \\ \frac{1}{4} |\alpha|^{-2} + \log |\alpha| & \text{if } |\alpha| > 1. \end{cases}$$

2. For  $m_{\mathbb{H}}(x^2 - 2\operatorname{Re}\alpha \cdot x + |\alpha|^2)$ , we have

$$m_{\mathbb{H}}(x^2 - 2\operatorname{Re}\alpha \cdot x + |\alpha|^2) = \begin{cases} \frac{1}{2} |\alpha|^2 \cos 2\phi & \text{if } |\alpha| \le 1\\ \frac{1}{2} |\alpha|^{-2} \cos 2\phi + 2\log |\alpha| & \text{if } |\alpha| > 1. \end{cases}$$

3. For  $m_{\mathbb{H}}(P(x))$ , where P(x) is a real polynomial completely factorized as

(1) 
$$P(x) = c \prod (x - \alpha_h)^{n_h} \prod (x^2 - 2\operatorname{Re}\beta_\ell \cdot x + |\beta_\ell|^2)^{m_\ell},$$

we have

$$m_{\mathbb{H}}(P(x)) = \log |c| + \sum n_h m_{\mathbb{H}}(x - \alpha_h) + \sum m_\ell m_{\mathbb{H}}(x^2 - 2 \operatorname{Re} \beta_\ell \cdot x + |\beta_\ell|^2)$$

In addition, we calculate the zeta Mahler measure of quaternionic monomials

$$Z_{\mathbb{H}}(x-\alpha;s) \coloneqq \int_{\mathbb{T}^1(\mathbb{H})} |x-\alpha|^s \, \mathrm{d}\mu(\mathbb{T}^1(\mathbb{H})),$$

as listed in Theorem 4.3.

Additionally, we also consider the Mahler measure of a class of more general quaternionic univariate polynomials, called *slice regular polynomials*. These polynomials are a special kind of entire *slice regular functions*. The theory of slice regular functions, which extends the theory of holomorphic functions in complex variables, has been systematically developed over the past two decades. Initially introduced within the context of quaternion algebra by Gentili and Struppa [13, 14] in 2006, slice regular theory has been successfully broadened to the real alternative \*-algebras [16] and several variables [17]. (See also the monograph [12] for an overview on the theory of slice regular functions in one quaternionic variable).

For slice regular polynomial of one variable, we prove the existence of its Mahler measure and establish also one lower bound.

**Theorem 1.2.** Let P(x) be a slice regular quaternionic polynomial. Then the quaternionic Mahler measure  $m_{\mathbb{H}}(P)$  exists as an improper Riemann integral. Moreover, we have always

$$m_{\mathbb{H}}(P) \ge \frac{1}{2} m_{\mathbb{H}}(P^s),$$

where the equality holds if and only if P is a real polynomial.

This theorem has also been extended to slice regular polynomials in two-variable cases, see Section 8.

Our next results concern the \*-*products* of quaternionic monomials (see Section 3 for definition).

**Theorem 1.3.** Let  $n \in \mathbb{N}$  and let  $(x - i)^{*n}$  denote the iterated \*-product. Then we have

$$m_{\mathbb{H}}((x-i)^{*n}) = \frac{2n}{\pi} \int_0^\infty t \coth nt \tanh^2 t \operatorname{sech} t \, \mathrm{d}t - \frac{2+n}{4}.$$

In particular, this leads to the following

$$m_{\mathbb{H}}(x^2 - 2xi - 1) = \frac{5}{6},$$
$$m_{\mathbb{H}}(x^3 - 3x^2i - 3x + i) = \frac{31}{4} - 2\pi.$$

In the following theorem, we examine examples of Mahler measures for noncommutative linear polynomials in the Sylvester form.

**Theorem 1.4.** Let  $a, b \in \mathbb{H}$  be non-zero and L(x) = ax + xb. Then

$$m_{\mathbb{H}}(L) = 2\pi^2 \left( \frac{\lambda^+ \log \lambda^+ - \lambda^- \log \lambda^-}{\lambda^+ - \lambda^-} - 1 \right),$$

where we put  $\lambda^{\pm} = (\operatorname{Re} a + \operatorname{Re} b)^2 + (|\operatorname{Im} a| \pm |\operatorname{Im} b|)^2$ .

Additionally, we provide several instances of Mahler measures in multiple variables. As for Smyth's example 1 + x + y, its quaternionic Mahler measure is as follows.

Theorem 1.5. We have

$$m_{\mathbb{H}}(1+x+y) = \frac{1}{2} - \frac{11\sqrt{3}}{16\pi} + \frac{3\sqrt{3}}{4\pi}L(\chi_{-3},2)$$

*Remark.* The formula here can be compared with Smyth's formula on  $m_{\mathbb{C}}(1+x+y)$ and Lalín–Roy's [22] evaluation of the areal Mahler measure  $m_{\mathbb{D}}(1+x+y)$ . We have indeed the following relation

$$m_{\mathbb{H}}(1+x+y) - m_{\mathbb{D}}(1+x+y) = \frac{1}{3}.$$

The resemblance of the Jensen formulas of monomials in complex case, the areal case and quaternionic case might serve as a foundational explanation.

Generalizing the argument of Maillot [26] to quaternion, we also derive the following formula.

**Theorem 1.6.** If |a|, |b|, |c| do not form a triangle, then we have

$$m_{\mathbb{H}}(ax + by + c) = \log \max\{|a|, |b|, |c|\} + \frac{|a|^2 + |b|^2 + |c|^2}{4\max\{|a|, |b|, |c|\}^2} - \frac{1}{4}.$$

This article is structured as follows. Section 2 presents preliminary properties on quaternions. In Section 3, a concise introduction to quaternion slice regular functions is provided. The quaternion Mahler measure of real polynomials is computed in Section 4, with a generalization to slice regular polynomials explored in Section 5. Section 6 offers examples of non-commutative univariate polynomials, while Section 7 provides some examples of multivariable quaternion Mahler measures. Finally, we introduce the quaternion analogue of the Lehmer problem in Section 8, along with numerical evidence.

#### Acknowledgements

The research of the second author was supported by Fundamental Research Funds for the Central Universities (Grant No. 531118010622). The authors thank François Brunault and Matilde Lalín for meaningful discussion concerning the paper.

#### QUATERNIONIC MAHLER MEASURE

### 2. Preliminaries on quaternions

The quaternion  $\mathbb{H}$  is a real non-commutative algebra with the canonical basis  $\{1, i, j, k\}$ . A quaternion number q in  $\mathbb{H}$  can be written uniquely as the form  $q = x_0 + ix_1 + jx_2 + kx_3$ , where  $x_\ell$  are real numbers and i, j, k share the multiplication rule

$$i^{2} = j^{2} = k^{2} = -1$$
,  $ij = k = -ji$ ,  $jk = i = -kj$ ,  $ki = j = -ik$ .

Every quaternion number x comprises two parts, the real part  $\operatorname{Re} q = x_0$  and the *imaginary part*  $\operatorname{Im} q = ix_1 + jx_2 + kx_3$ . The conjugate of q is  $\overline{q} = \operatorname{Re} x - \operatorname{Im} x$  and the norm of q is defined as  $|q|^2 = q\overline{q} = x_0^2 + x_1^2 + x_2^2 + x_3^2$ . Denote S by the unit 2-sphere of purely imaginary quaternion, that is,

$$\mathbb{S} = \{ q = ix_1 + jx_2 + kx_3 \in \mathbb{H} \, | \, x_1^2 + x_2^2 + x_3^2 = 1 \},\$$

these are exactly the quaternion numbers I such that  $I^2 = -1$ . Then every quaternion number can be rewritten as  $q = \alpha + I\beta$  for some  $\alpha, \beta \in \mathbb{R}$  and  $I \in \mathbb{S}$ . For a fixed  $I \in \mathbb{S}$ , the set of all quaternions in this form is defined as a *slice*, denoted by  $\mathbb{C}_I = \mathbb{R} + \mathbb{R}I$ . Note that the quaternion algebra  $\mathbb{H}$  can be decomposed as the union of all the slices  $\mathbb{C}_I$  where  $I \in \mathbb{S}$ 

$$\mathbb{H} = \bigcup_{I \in \mathbb{S}} \mathbb{C}_I, \quad \mathbb{R} = \bigcap_{I \in \mathbb{S}} \mathbb{C}_I.$$

Two quaternions  $p, q \in \mathbb{H}$  are called *similar* if there exists non-zero  $r \in \mathbb{H}$  such that  $q = r^{-1}pr$ . It is well known that two quaternions p, q are similar precisely if  $\operatorname{Re} p = \operatorname{Re} q$  and  $|\operatorname{Im} p| = |\operatorname{Im} q|$ . All the quaternions which are similar to p is thusly the 2-sphere given by

$$\mathbb{S}_p \coloneqq \{q \in \mathbb{H} \mid q \text{ is similar to } p\} = \operatorname{Re} p + |\operatorname{Im} p| \mathbb{S}.$$

Let  $I, J \in \mathbb{S}$  such that  $I \perp J$ . Then every quaternion can be written uniquely as  $q = z + Jw \in \mathbb{H}$  with  $z, w \in \mathbb{C}_I$ . This gives the canonical isomorphism

$$\mathbb{T}^{1}(\mathbb{H}) \simeq \mathrm{SU}(2) = \left\{ \begin{pmatrix} z & w \\ -\overline{w} & \overline{z} \end{pmatrix} \middle| z, w \in \mathbb{C} \text{ such that } |z|^{2} + |w|^{2} = 1 \right\}.$$

The probability Haar measure on SU(2) is given by  $\frac{1}{4\pi^2} \frac{dz}{z} \wedge dw \wedge d\overline{w}$ . Thus, the quaternionic Mahler measure may be expressed as

(2) 
$$m_{\mathbb{H}}(P) = \frac{1}{(4\pi)^n} \int_{\mathrm{SU}(2)^n} \log |P(x_1, \dots, x_n)| \prod_{\ell=1}^n \frac{\mathrm{d}z_\ell}{z_\ell} \,\mathrm{d}w_\ell \,\mathrm{d}\overline{w_\ell},$$

where each  $x_{\ell} = \begin{pmatrix} \frac{z_{\ell}}{-w_{\ell}} & \frac{w_{\ell}}{z_{\ell}} \end{pmatrix}$  is in SU(2).

The unit quaternion  $\mathbb{T}^1(\mathbb{H})$  is also canonically isomorphic to the 3-sphere  $\mathbb{S}^3$ . This yields the formula

(3) 
$$\mathbf{m}_{\mathbb{H}}(P) = \int_{(\mathbb{S}^3)^n} \log |P(x_1, \dots, x_n)| \,\mathrm{d}^n \mu(\mathbb{S}^3).$$

In particular, given an orthonormal basis  $\{I, J, K = IJ\}$  in  $\mathbb{S}$ , then every unit q can be represented in a hyperspherical form  $q = \cos \theta + \sin \theta \cos \theta_1 I + \sin \theta \sin \theta_1 \cos \theta_2 J + \sin \theta \sin \theta_1 \sin \theta_2 K$ . Then the Haar measure of  $\mathbb{S}^3$  is given by

$$d\mu(\mathbb{S}^3) = \frac{1}{2\pi^2} \sin^2 \theta \sin \theta_1 \, d\theta \, d\theta_1 \, d\theta_2.$$

# 3. QUATERNION SLICE REGULAR FUNCTIONS

In the sequel, we offer here a minimalistic introduction for slice regular functions. We will concentrate on functions within the open ball B(0, R) of radius R > 0 in  $\mathbb{H}$ , while the general theory can be found in aforementioned references. For each  $I \in \mathbb{S}$ , let  $B_I = B \cap \mathbb{C}_I$  denote the slices of the set B.

**Definition 3.1.** Let f be a quaternion-valued function defined on B = B(0, R). For each  $I \in \mathbb{S}$ , let  $f_I = f|_{B_I}$  be the restriction of f on  $B_I$ . Then f is called slice (left) regular if  $f_I$  is holomorphic for all  $I \in \mathbb{S}$ , i.e. it has continuous partial derivatives and

$$\overline{\partial}_I f(x+yI) = \frac{1}{2} \left( \frac{\partial}{\partial x} + I \frac{\partial}{\partial y} \right) f_I(x+yI) = 0.$$

As is proved in [13], a slice regular function f on B(0, R) has a quaternion series representation

$$f(q) = \sum_{n \in \mathbb{N}} q^n a_n$$
, where  $a_n \in \mathbb{H}$ .

The converse is also true, a quaternion series of the form  $\sum_{n \in \mathbb{N}} q^n a_n$  also defines a slice regular function within its domain of convergence. By splitting the quaternion coefficients by complex numbers, the following lemma associates the slice regular functions with complex holomorphic functions.

**Lemma 3.2** (Splitting Lemma). Let f be a slice regular function on B(0, R). Let  $I, J \in \mathbb{S}$  such that  $I \perp J$ . Then there exist two complex holomorphic functions  $F, G: B_I \to \mathbb{C}_I$  such that for any z = x + Iy in  $B_I$ , we have

$$f_I(z) = F(z) + G(z)J.$$

Just as holomorphic functions, there is also an identity principle holds for slice regular functions.

**Lemma 3.3** (Identity principle). Let f and g be two slice regular functions on B(0, R). If there are some  $I \in S$  such that the functions f and g coincide on a subset of  $B_I$  with one accumulation point in  $B_I$ , then  $f \equiv g$  holds on B(0, R).

The values of a slice regular function can be determined exactly by the values of its restrictions on a complex line. In particular, a holomorphic function  $f_I : B_I \to \mathbb{H}$  admits a unique regular extension  $ext(f_I)$  which has restriction  $f_I$  on  $B_I$ .

**Lemma 3.4** (Representation Formula). Let f be a slice regular function on the ball B = B(0, R) and let  $I, J \in \mathbb{S}$ . Then for all  $x + yI \in B$  we have

$$f(x+yJ) = \frac{1-JI}{2}f(x+yI) + \frac{1+JI}{2}f(x-yI).$$

Moreover, let  $K \in \mathbb{S}$  and  $I \neq J$ , we have

$$f(x+yK) = (I-J)^{-1}[If(x+yI) - Jf(x+yJ)] + K(I-J)^{-1}[f(x+yI) - f(x+yJ)].$$

Unfortunately, the product of two slice regular functions is usually no longer regular. A surrogate multiplication called the *\*-product*, which preserves slice regularity, is defined as below.

**Definition 3.5.** Let  $f = \sum_{n \in \mathbb{N}} q^n a_n$  and  $g = \sum_{n \in \mathbb{N}} q^n b_n$  be two slice regular functions on B(0, R). The \*-product of f and g is defined as the slice regular function f \* g on B(0, R) with the power series

$$(f * g)(q) = \sum_{n \in \mathbb{N}} q^n \sum_{l=0}^n a_l b_{n-l}.$$

Note that the \*-products are not commutative in general. Also, the \*-product (f \* g)(q) is not equal to the product f(q)g(q). However, we can still connect \*-product and pointwise product by

**Lemma 3.6.** Let f and g be two slice regular functions on B(0, R). Then we have

$$(f * g)(p) = \begin{cases} f(p)g(f(p)^{-1}pf(p)) & \text{if } f(p) \neq 0\\ 0 & \text{if } f(p) = 0 \end{cases}$$

We can associate a slice regular function f with a particular power series  $f^s$  with real coefficients, which vanishes at all zeros of f. In fact, if f has one zero  $a \in \mathbb{H}$ , then  $f^s$  vanishes on the 2-sphere  $\mathbb{S}_a$  (see [10, Theorem 4.3]).

**Definition 3.7.** Let  $f = \sum_{n \in \mathbb{N}} q^n a_n$  be a slice regular function on B(0, R). Then we define *regular conjugate* of f as the slice regular function

$$f^c(q) \coloneqq \sum_{n \in \mathbb{N}} q^n \overline{a}_n,$$

and the symmetrization of f as

$$f^{s}(q) \coloneqq (f * f^{c})(q) = (f^{c} * f)(q) = \sum_{n \in \mathbb{N}} q^{n} \sum_{l=0}^{n} a_{l} \overline{a}_{n-l}.$$

With the notations of regular conjugate and symmetrization, we give the following definition. **Definition 3.8.** Let f be a slice regular function on B = B(0, R). Let the zero set of its symmetrization  $\mathcal{Z}_{f^s}$  be a proper subset of B. The regular reciprocal of f is defined as the function  $f^{-*}(q) : B \setminus \mathcal{Z}_{f^s} \to \mathbb{H}$  by

$$f^{-*}(q) \coloneqq (f^s(q))^{-1} f^c(q).$$

If a slice regular function  $f = \sum_{\ell=0}^{n} q^{\ell} a_{\ell}$  is given a finite series, then f is called a *slice regular polynomial*. The following well-known theorem, also recognized as *Eilenberg–Niven theorem*, given precise factorizations of slice regular polynomials.

**Theorem 3.9** (Fundamental Theorem of Algebra for Quaternion [6, 10, 15, 16]). Let f(q) be a non-zero slice regular polynomial on  $\mathbb{H}$ . Then its zero set  $\mathcal{Z}_f$  is non-empty. More precisely, the zero set  $\mathcal{Z}_f$  is a finite union of spherical roots (of the form  $\mathbb{S}_{\alpha}$  for some  $\alpha \in \mathbb{H}$ ) and isolated roots. Also, the polynomial f(q) can be factorized completely as

$$f(q) = (q^2 - 2\operatorname{Re}\alpha_1 \cdot q + |\alpha_1|^2)^{m_1} * \dots (q^2 - 2\operatorname{Re}\alpha_\ell \cdot q + |\alpha_\ell|^2)^{m_\ell} * \prod^* (q - \beta_h)c,$$
  
for some  $\alpha_\ell, \beta_h, c \in \mathbb{H}$ .

*Remark.* Note that the factorization of the slice regular polynomial f(q) is usually not unique. Also, the quaternions  $\beta_h$  (h > 1) are in general not the root of f(q). For example, the quadratic polynomial f(q) = (q - i) \* (q - j) has only one root q = i.

The following two kinds of functions are of particular importance in our computations.

**Definition 3.10.** Let f be a slice regular function on B = B(0, R). Then f is called *slice preserving* if  $f(B_I) \subset \mathbb{C}_I$  for any  $I \in \mathbb{S}$ . It is called *one-slice preserving* on  $\mathbb{C}_I$  if  $f(B_I) \subset \mathbb{C}_I$  for one certain  $I \in \mathbb{S}$ .

It is possible to characterize these two kinds of functions by their power series.

**Theorem 3.11.** Let  $f(q) = \sum_{n \in \mathbb{N}} q^n a_n$  be a slice regular function on B(0, R). Then we have

- 1. The function f is slice preserving precisely if  $a_n \in \mathbb{R}$  for all n.
- 2. The function f is one-slice preserving on  $\mathbb{C}_I$  precisely if  $a_n \in \mathbb{C}_I$  for all n.

*Proof.* The first claim is given in [12, Section 1.4] and the second claim is a corollary of [2, Proposition 1.4].  $\Box$ 

Here we give an example of slice regular function which is slice preserving.

**Example 3.12.** The quaternion exponential function  $\exp(q)$  is defined by the slice regular power series

$$e^q \coloneqq \sum_{n=0}^{\infty} \frac{q^n}{n!}, \quad for \ q \in \mathbb{H}.$$

It has all real coefficients, and is thus slice preserving. In fact, when  $I^2 = -1$  is an imaginary unit, we have Euler's formula  $e^{I\theta} = \cos \theta + I \sin \theta$  for all  $\theta \in \mathbb{R}$ . So it holds that

$$e^{q} = e^{\operatorname{Re} q} \left( \cos \left| \operatorname{Im} q \right| + \frac{\operatorname{Im} q}{\left| \operatorname{Im} q \right|} \sin \left| \operatorname{Im} q \right| \right).$$

The logarithm (principal branch) is its inverse

$$\log q = \log |q| + \frac{\operatorname{Im} q}{|\operatorname{Im} q|} \operatorname{arccos} \frac{\operatorname{Re} q}{|q|}.$$

In the SU(2)-model of the unit quaternions, the quaternion exponential function is given by

$$\exp:\mathfrak{su}(2) \to \operatorname{SU}(2) \\ \begin{pmatrix} i\theta_1 & \theta_2 + i\theta_3 \\ -\theta_2 + i\theta_3 & -i\theta_1 \end{pmatrix} \mapsto \begin{pmatrix} \cos\theta + i\frac{\sin\theta}{\theta}\theta_1 & \frac{\sin\theta}{\theta}(\theta_2 + i\theta_3) \\ \frac{\sin\theta}{\theta}(-\theta_2 + i\theta_3) & \cos\theta - i\frac{\sin\theta}{\theta}\theta_1 \end{pmatrix},$$
  
where  $\theta = \sqrt{\theta_1^2 + \theta_2^2 + \theta_3^2}.$ 

**Lemma 3.13.** Let  $\mathbb{D} = \{i\theta_1 + j\theta_2 + k\theta_3 \in \mathbb{H} \mid \theta = \sqrt{\theta_1^2 + \theta_2^2 + \theta_3^2} \in [0, \pi)\}$ . Then the quaternion functions exp and log define a bijection between

$$\mathbb{D} \xrightarrow[\log]{\operatorname{log}} \mathbb{T}^1(\mathbb{H}) \backslash \{-1\}$$

*Proof.* Let  $q \in \mathbb{D}$ . Notice first that  $|e^q| = e^{\operatorname{Re} q}$ . Since we have  $\theta \neq \pi$ , it follows that  $e^q \neq -1$  is in  $\mathbb{T}^1(\mathbb{H})$ . For the other side, let  $q \in \mathbb{T}^1(\mathbb{H})$  and  $q \neq -1$ . We have  $\operatorname{Re} \log q = 0$  and  $\operatorname{arccos} \frac{\operatorname{Re} q}{|q|} \in [0, \pi)$ , which indicates that  $\log q \in \mathbb{D}$ . This means that both two maps are well-defined. A direct calculation shows that  $\exp \circ \log$  and  $\log \circ \exp$  are identity maps.

We end this section by giving the formulas of Mahler measure in terms of quaternionic exponential.

**Theorem 3.14.** Let P be any quaternionic polynomial and let  $\mathbb{D}$  be defined as before, then we have

$$m_{\mathbb{H}} \left( P(x_1, \dots, x_n) \right)$$

$$= \frac{1}{\left(2\pi^2\right)^n} \int_{\mathbb{D}^n} \log \left| P(e^{i\theta_{11} + j\theta_{12} + k\theta_{13}}, \dots, e^{i\theta_{n1} + j\theta_{n2} + k\theta_{n3}}) \right| \prod_{i=1}^n \frac{\sin^2 \theta_i}{\theta_i^2} \, \mathrm{d}\theta_{i1} \, \mathrm{d}\theta_{i2} \, \mathrm{d}\theta_{i3},$$

and

$$Z_{\mathbb{H}}\left(P(x_1,\ldots,x_n),s\right)$$

$$=\frac{1}{(2\pi^2)^n}\int_{\mathbb{D}^n}\left|P(e^{i\theta_{11}+j\theta_{12}+k\theta_{13}},\ldots,e^{i\theta_{n1}+j\theta_{n2}+k\theta_{n3}})\right|^s\prod_{i=1}^n\frac{\sin^2\theta_i}{\theta_i^2}\,\mathrm{d}\theta_{i1}\,\mathrm{d}\theta_{i2}\,\mathrm{d}\theta_{i3},$$

where we denote  $\theta_n = \sqrt{\theta_{n1}^2 + \theta_{n2}^2 + \theta_{n3}^2}$ .

*Proof.* In (2), putting  $z = \cos \theta + i \frac{\sin \theta}{\theta} \theta_1$  and  $w = \frac{\sin \theta}{\theta} (\theta_2 + i \theta_3)$ , one finds that

$$\frac{\mathrm{d}z}{z} \wedge \mathrm{d}w \wedge \mathrm{d}\overline{w} = 2 \,\frac{\sin^2\theta}{\theta^2} \,\mathrm{d}\theta_1 \wedge \mathrm{d}\theta_2 \wedge \mathrm{d}\theta_3$$

Then using Lemma 3.13 we derive the desired formulas.

## 4. JENSEN FORMULA AND MAHLER MEASURES OF REAL POLYNOMIALS

As a preparatory step, we give some formulas that we will be using to compute the quaternionic Mahler measures.

**Lemma 4.1** (Jensen's formula). Let  $I \in \mathbb{S}$  and let  $\alpha = ae^{I\phi}$  with  $a, \phi \in \mathbb{R}$ . Define the function

$$J(\alpha) \coloneqq \frac{1}{\pi} \int_0^\pi \log\left(1 - 2a\cos(\theta - \phi) + a^2\right) \sin^2\theta \,\mathrm{d}\theta.$$

Then for  $|\alpha| \leq 1$  we have

$$J(\alpha) = \frac{|\alpha|^2}{4} \cos 2\phi - \frac{1}{2\pi I} \operatorname{Im}\left[\left(\alpha - \frac{1}{\alpha}\right) - \left(\alpha^2 - \frac{1}{\alpha^2}\right) \operatorname{arctanh} \alpha + 4\operatorname{Li}_2(\alpha) - \operatorname{Li}_2(\alpha^2)\right].$$

And for  $|\alpha| > 1$  we have

$$J(\alpha) = J(\alpha/|\alpha|^2) + \log |\alpha|,$$

where the functions  $Li_2$  and arctanh are regarded as the regular extensions of their complex counterparts.

*Proof.* Denote by  $J(a, \phi) = J(\alpha)$ . Note that we have

$$J(a,\phi) - \log|a| = \frac{1}{\pi} \int_0^{\pi} \log\left(1 - \frac{2}{a}\cos(\theta - \phi) + \frac{1}{a^2}\right)\sin^2\theta \,\mathrm{d}\theta.$$

The latter is nothing but  $J(1/a, \phi)$ , allowing us to assume that  $|a| \leq 1$ . Consider the following expansion

(4) 
$$\log(1 - 2a\cos(\theta - \phi) + a^2) = -2\sum_{n \ge 1} \frac{a^n}{n} \cos n(\theta - \phi).$$

This, combined with the given integral

$$\int_{0}^{\pi} \cos n(\theta - \phi) \sin^{2} \theta \, \mathrm{d}\theta = \begin{cases} -\frac{4}{(n^{2} - 4)n} \sin n\phi & n \text{ odd,} \\ 0 & n \text{ even, } n \neq 2, \\ -\frac{\pi}{4} \cos 2\phi & n = 2, \end{cases}$$

yields

$$J(a,\phi) = \frac{a^2}{4}\cos 2\phi + \sum_{n \text{ odd}, n \ge 1} \frac{8a^n}{n^2(n^2 - 4)\pi} \sin n\phi.$$

Through partial fraction decomposition, we obtain

$$\frac{1}{n^2(n^2-4)} = \frac{1}{16} \left( \frac{1}{n-2} - \frac{1}{n+2} - \frac{4}{n^2} \right).$$

It is also worth noting that for  $|\alpha| < 1$ , one has

(5) 
$$\operatorname{arctanh} \alpha = \sum_{n \text{ odd, } n \ge 1} \frac{\alpha^n}{n}.$$

 $\operatorname{So}$ 

$$J(\alpha) = \frac{|\alpha|^2}{4} \cos 2\phi - \frac{1}{2\pi I} \operatorname{Im} \left[ \left( \alpha - \frac{1}{\alpha} \right) - \left( \alpha^2 - \frac{1}{\alpha^2} \right) \operatorname{arctanh} \alpha + 4 \operatorname{Li}_2(\alpha) - \operatorname{Li}_2(\alpha^2) \right].$$

In fact, the Mahler measures become much easier to compute if the polynomial is slice preserving or one-slice preserving.

**Proposition 4.2.** Let  $P(x) = \sum_{\ell=0}^{n} x^{\ell} a_{\ell}$  be a slice regular polynomial.

1. If P is slice preserving, then we have

$$m_{\mathbb{H}}(P) = \frac{2}{\pi} \int_0^{\pi} \log |P(e^{i\theta})| \sin^2 \theta \, \mathrm{d}\theta,$$
$$Z_{\mathbb{H}}(P,s) = \frac{2}{\pi} \int_0^{\pi} |P(e^{i\theta})|^s \sin^2 \theta \, \mathrm{d}\theta.$$

2. If P is one-slice preserving on  $\mathbb{C}_I$ , and let  $I \perp J$ . Then

$$m_{\mathbb{H}}(P) = \frac{1}{\pi} \int_0^{\pi} \int_0^{\pi} \log |P(\cos\theta + Ie^{J\theta_1}\sin\theta)| \sin^2\theta \sin\theta_1 \,\mathrm{d}\theta \,\mathrm{d}\theta_1,$$
$$Z_{\mathbb{H}}(P,s) = \frac{1}{\pi} \int_0^{\pi} \int_0^{\pi} |P(\cos\theta + Ie^{J\theta_1}\sin\theta)|^s \sin^2\theta \sin\theta_1 \,\mathrm{d}\theta \,\mathrm{d}\theta_1.$$

*Proof.* If P is slice preserving, then the coefficients of P are real. Since  $e^{i\theta_1+j\theta_2+k\theta_3}$  is similar to  $e^{i\theta}$ , we have  $|P(e^{i\theta_1+j\theta_2+k\theta_3})| = |P(e^{i\theta})|$ . Let again  $\theta = \sqrt{\theta_1^2 + \theta_2^2 + \theta_3^2}$ . Changing the variables with spherical coordinates  $\theta_1 = \theta \cos \psi_1$ ,  $\theta_2 = \theta \sin \psi_1 \cos \psi_2$ 

and  $\theta_3 = \theta \sin \psi_1 \sin \psi_2$ , one obtains from Theorem 3.14 that

$$m_{\mathbb{H}}(P) = \frac{1}{2\pi^2} \int_{\theta_1^2 + \theta_2^2 + \theta_3^2 \le \pi^2} \log |P(e^{i\theta})| \frac{\sin^2 \theta}{\theta^2} \, \mathrm{d}\theta_1 \, \mathrm{d}\theta_2 \, \mathrm{d}\theta_3$$
$$= \frac{1}{\pi^2} \int_0^{2\pi} \int_0^{\pi} \int_0^{\pi} \log |P(e^{i\theta})| \sin^2 \theta \sin \phi_1 \, \mathrm{d}\theta \, \mathrm{d}\psi_1 \, \mathrm{d}\psi_2$$
$$= \frac{2}{\pi} \int_0^{\pi} \log |P(e^{i\theta})| \sin^2 \theta \, \mathrm{d}\theta.$$

The proof of  $Z_{\mathbb{H}}(P, s)$  is the same. This proves the first claim.

For the second claim, let P be one-slice preserving on  $\mathbb{C}_I$ , then the coefficients of P is in  $\mathbb{C}_I$ . We may represent x in the hyperspherical form

$$x = \cos\theta + \sin\theta\cos\theta_1 I + \sin\theta\sin\theta_1\cos\theta_2 J + \sin\theta\sin\theta_1\sin\theta_2 K,$$

where  $I \perp J$  and K = IJ. Let  $\phi = \theta_2/2 - \pi/4$  and  $q = e^{I\phi} \in \mathbb{C}_I$ , then a direct calculation shows that x is similar to  $q^{-1}xq = \cos \theta + Ie^{J\theta_1}\sin \theta$ . So for any  $x \in \mathbb{T}^1(\mathbb{H})$ , we have  $|P(x)| = |P(\cos \theta + Ie^{J\theta_1}\sin \theta)|$ . Again changing the variable, we get

$$m_{\mathbb{H}}(P) = \frac{1}{2\pi^2} \int_0^{2\pi} \int_0^{\pi} \int_0^{\pi} \log |P(x)| \sin^2 \theta \sin \theta_1 \, \mathrm{d}\theta \, \mathrm{d}\theta_1 \, \mathrm{d}\theta_2$$
$$= \frac{1}{\pi} \int_0^{\pi} \int_0^{\pi} \log |P(\cos \theta + Ie^{J\theta_1} \sin \theta)| \sin^2 \theta \sin \theta_1 \, \mathrm{d}\theta \, \mathrm{d}\theta_1. \qquad \Box$$

We now come to the proof of the part (1) in Theorem 1.1.

Proof of (1) in Theorem 1.1. Let us consider  $\alpha \in \mathbb{H}$  with the most generality. Then there exists  $q_0 \in \mathbb{H}$  such that  $\alpha$  is similar to  $\alpha_0 = q_0^{-1} \alpha q_0 \in \mathbb{C}$ . So

$$m_{\mathbb{H}}(x-\alpha) = \int_{\mathbb{T}^{1}(\mathbb{H})} \log |q_{0}^{-1}xq_{0} - \alpha_{0}| \,\mathrm{d}\mu$$
$$= \int_{\mathbb{T}^{1}(\mathbb{H})} \log |x-\alpha_{0}| \,\mathrm{d}\mu = m_{\mathbb{H}}(x-\alpha_{0}).$$

This means we may always assume that  $\alpha \in \mathbb{C}$ . For the next step, we set  $\alpha = |\alpha|e^{i\theta}$ . Writing the integral by the SU(2)-model of unit quaternions, we get

$$m_{\mathbb{H}}(x-\alpha) = \frac{1}{8\pi^2} \int_{|z|^2 + |w|^2 = 1} \log(|z-|\alpha|e^{i\theta}|^2 + |w|^2) \frac{\mathrm{d}z}{z} \,\mathrm{d}w \,\mathrm{d}\overline{w}$$
$$= \frac{1}{8\pi^2} \int_{|z|^2 + |w|^2 = 1} \log(|ze^{-i\theta} - |\alpha||^2 + |w|^2) \frac{\mathrm{d}z}{z} \,\mathrm{d}w \,\mathrm{d}\overline{w} = m_{\mathbb{H}}(x-|\alpha|).$$

Now we can restrict our attention only to the case where  $\alpha \in \mathbb{R}$ . In this scenario, the polynomial  $x - \alpha$  is slice-preserving. From Proposition 4.2, we find

$$m_{\mathbb{H}}(x-\alpha) = \frac{2}{\pi} \int_0^{\pi} \log |e^{i\theta} - \alpha| \sin^2 \theta \, \mathrm{d}\theta$$
$$= \frac{1}{\pi} \int_0^{\pi} \log |1 - 2\alpha \cos \theta + \alpha^2| \sin^2 \theta \, \mathrm{d}\theta.$$

Thus, the formula follows directly from Jensen's formula 4.1.

We can also evaluate the quaternionic zeta Mahler measure of  $x - \alpha$  for  $\alpha \in \mathbb{H}$ . This offers a quaternion analogue of Akatsuka's results [1, Theorem 1, 2].

**Theorem 4.3.** Let  $\alpha \in \mathbb{H}$ . The quaternionic zeta Mahler measure of  $x - \alpha$  is given below.

1. If  $|\alpha| = 1$  and  $\operatorname{Re} s > -3$ , we have

$$Z_{\mathbb{H}}(x-\alpha;s) = \frac{2^{2+s}}{\sqrt{\pi}} \frac{\Gamma(s+\frac{3}{2})}{\Gamma(s+3)}.$$

2. If  $|\alpha| < 1$  and  $s \in \mathbb{C}$ , then  $Z_{\mathbb{H}}(x - \alpha; s)$  is

$$A(\alpha, s)_2 F_1 \begin{pmatrix} -\frac{s}{2}, & -\frac{s}{2} \\ 1 & ; |\alpha|^2 \end{pmatrix} + B(\alpha, s)_2 F_1 \begin{pmatrix} 1 - \frac{s}{2}, & -\frac{s}{2} \\ 1 & ; |\alpha|^2 \end{pmatrix},$$

3. If  $|\alpha| > 1$ , and  $s \in \mathbb{C}$ , then  $Z_{\mathbb{H}}(x - \alpha; s)$  is

$$|\alpha|^{s} A(\alpha, s)_{2} F_{1} \begin{pmatrix} -\frac{s}{2}, & -\frac{s}{2} \\ 1 \end{pmatrix} + |\alpha|^{s} B(\alpha, s)_{2} F_{1} \begin{pmatrix} 1 - \frac{s}{2}, & -\frac{s}{2} \\ 1 \end{pmatrix},$$

where we denote

$$A(\alpha, s) = \frac{(8+6s)|\alpha|^2 + 2s}{(s+2)(s+4)|\alpha|^2}, \quad B(\alpha, s) = \frac{-2s(1-|\alpha|^2)}{(s+2)(s+4)|\alpha|^2}.$$

*Proof.* Using the same reasoning as in the previous proof, we can safely assume that  $\alpha \in \mathbb{R}^+$ . Referring to Proposition 4.2, we have

$$Z_{\mathbb{H}}(x-\alpha;s) = \frac{2}{\pi} \int_0^\pi |1-2\alpha\cos\theta + \alpha^2|^{\frac{s}{2}}\sin^2\theta \,\mathrm{d}\theta.$$

Let  $\alpha = 1$ , then for  $\operatorname{Re} s > -3$ , one has

$$Z_{\mathbb{H}}(x-\alpha;s) = \frac{2^{1+\frac{s}{2}}}{\pi} \int_{0}^{\pi} (1-\cos\theta)^{\frac{s}{2}} \sin^{2}\theta \,\mathrm{d}\theta$$
  
$$= \frac{2^{3+\frac{s}{2}}}{\pi} \int_{0}^{\pi} \left(2\sin^{2}\frac{\theta}{2}\right)^{\frac{s}{2}} \sin^{2}\frac{\theta}{2} \left(1-\sin^{2}\frac{\theta}{2}\right) \mathrm{d}\theta$$
  
$$= \frac{2^{4+s}}{\pi} \int_{0}^{\frac{\pi}{2}} \sin^{s+2}\theta (1-\sin^{2}\theta) \,\mathrm{d}\theta$$
  
$$= \frac{2^{2+s}}{\sqrt{\pi}} \frac{\Gamma(s+\frac{3}{2})}{\Gamma(s+3)}.$$

If  $0 < \alpha < 1$ , we recall the result of Akatsuka's results [1, Theorem 2] on the complex Mahler measures  $Z_{\mathbb{C}}(x - \alpha; s)$ , which can be rewritten as

(6)  

$$Z_{\mathbb{C}}(x-\alpha;s) = \frac{1}{2\pi} \int_{-\pi}^{\pi} (1-2\alpha\cos\theta+\alpha^2)^{\frac{s}{2}} d\theta$$

$$= \frac{1}{\pi} \int_{0}^{\pi} (1-2\alpha\cos\theta+\alpha^2)^{\frac{s}{2}} d\theta$$

$$= {}_{2}F_{1} \left( -\frac{s}{2}, -\frac{s}{2}; |\alpha|^{2} \right).$$

Differentiating with respect to  $\alpha$ , one infers that

(7) 
$$\frac{\partial}{\partial \alpha} Z_{\mathbb{C}}(x-\alpha;s) = \frac{s}{2\pi} \int_0^\pi (1-2\alpha\cos\theta + \alpha^2)^{\frac{s}{2}} (2\alpha - 2\cos\theta) \,\mathrm{d}\theta.$$

Additionally, it can be seen that

$$Z_{\mathbb{C}}(x-\alpha;s+4) = \frac{1}{\pi} \int_0^{\pi} (1-2\alpha\cos\theta + \alpha^2)^{\frac{s}{2}} \left( (1+\alpha^2)^2 + 4\alpha^2\cos^2\theta - 4\alpha(1+\alpha^2)\cos\theta \right) d\theta.$$

The linear combination of (6), (7) and (8) will give us the desired formula for

$$Z_{\mathbb{H}}(x-\alpha;s) = \frac{1}{\pi} \int_0^{\pi} (1-2\alpha\cos\theta + \alpha^2)^{\frac{s}{2}} \sin^2\theta \,\mathrm{d}\theta.$$

If  $\alpha > 1$ , we have the functional equation  $Z_{\mathbb{H}}(x - \alpha^{-1}; s) = \alpha^{-s} Z_{\mathbb{H}}(x - \alpha; s)$ . This reduces to the case before.

We arrive at proof of the part (2) in Theorem 1.1.

Proof of (2) in Theorem 1.1. Again, since every quaternion is similar to a complex number, without loss of generality we may assume that  $\alpha = ae^{i\phi} \in \mathbb{C}$ . By Proposition 4.2, we have

$$m_{\mathbb{H}}(x^2 - 2\operatorname{Re}\alpha \cdot x + |\alpha|^2) = \frac{2}{\pi} \int_0^\pi \log|x^2 - 2\operatorname{Re}\alpha \cdot x + |\alpha|^2|\sin^2\theta \,\mathrm{d}\theta$$
$$= \frac{2}{\pi} \int_0^\pi (\log|e^{i\theta} - ae^{i\phi}| + \log|e^{i\theta} - ae^{-i\phi}|)\sin^2\theta \,\mathrm{d}\theta.$$

It is evident from Lemma 4.1 that the last integral is  $J(\alpha) + J(\overline{\alpha})$ .

To give the final proof of the part (3) in Theorem 1.1, we start with the following theorem.

**Theorem 4.4.** Let  $P(x) \in \mathbb{R}[x]$  be a real polynomial and Q(x) be any quaternion slice regular polynomial. Then we have

$$m_{\mathbb{H}}(P * Q) = m_{\mathbb{H}}(Q * P)$$
  
=  $m_{\mathbb{H}}(PQ) = m_{\mathbb{H}}(QP) = m_{\mathbb{H}}(P) + m_{\mathbb{H}}(Q).$ 

*Proof.* Since P are slice preserving and having real coefficients, we know everything commutes. It turns out that P\*Q = Q\*P (the \*-product) is exactly the PQ = QP (the non-commutative product).

*Remark.* We point out that  $m_{\mathbb{H}}(P * Q) = m_{\mathbb{H}}(P) + m_{\mathbb{H}}(Q)$  is usually false if P, Q are not real polynomials. For instance, for P(x) = x - i and Q(x) = x + i it follows

$$m_{\mathbb{H}}((x-i)*(x+i)) = m_{\mathbb{H}}(x^2+1) = -\frac{1}{2},$$

while

$$m_{\mathbb{H}}(x-i) + m_{\mathbb{H}}(x+i) = \frac{1}{2}$$

We will see in Section 5 it still fails even when P = Q. A counterexample is

$$m_{\mathbb{H}}((x-i)*(x-i)) = \frac{5}{6} \neq 2m_{\mathbb{H}}(x-i).$$

Proof of (3) in Theorem 1.1. This is an immediate corollary of Theorem 4.4.  $\Box$ 

### 5. Mahler measures of slice regular polynomials

In this section, we will focus on Mahler measures of slice regular polynomials. We commence by establishing the existence of quaternionic Mahler measures. To validate the proof, we introduce an important lemma which is essential for demonstrating the existence of the complex Mahler measure.

**Lemma 5.1** (Lemma 3.8 in Everest–Ward [9]). Let  $P \in \mathbb{C}[x_1^{\pm}, \ldots, x_n^{\pm}]$  be a complex Laurent polynomial. Then for any  $\varepsilon > 0$ , there exists  $C, \delta > 0$  depending only on P such that

$$\mu_{\mathbb{C}}\left(\left\{(x_1,\ldots,x_n)\in\mathbb{T}^n(\mathbb{C})\,\middle|\,|P(x_1,\ldots,x_n)|\leq\varepsilon\right\}\right)\leq C\varepsilon^{\delta},$$

where  $\mu_{\mathbb{C}}$  represents the probability Haar measure on  $\mathbb{T}^n(\mathbb{C})$ .

The following lemma introduces a quaternionic analogue for analysing the singularities in one single variable, similar to its complex counterpart shown previously.

**Lemma 5.2.** Let P(x) be a slice regular quaternionic polynomial. Let  $\mu_{\mathbb{H}}$  be the probability Haar measure of the unit quaternion. Then for any  $\varepsilon > 0$ , there exists  $C, \delta > 0$  depending only on P such that

$$\mu_{\mathbb{H}}\left(\left\{q\in\mathbb{T}^{1}(\mathbb{H})\,\big|\,|P(q)|\leq\varepsilon\right\}\right)\leq C\varepsilon^{\delta}.$$

*Proof.* Let us firstly prove the assertion when P is a real polynomial. Let  $q \in \mathbb{T}^1(\mathbb{H})$  be with  $|P(q)| \leq \varepsilon$ . Then for any  $p \in \mathbb{S}_q$ , we have still  $|P(p)| \leq \varepsilon$ . Thus

$$\mu_{\mathbb{H}}\left(\left\{q \in \mathbb{T}^{1}(\mathbb{H}) \mid |P(q)| \leq \varepsilon\right\}\right) = \mu_{\mathbb{C}}\left(\left\{q \in \mathbb{T}^{1}(\mathbb{C}) \mid |P(q)| \leq \varepsilon\right\}\right),$$

where  $\mu_{\mathbb{C}}$  is probability Haar measure of the unit complex torus  $\mathbb{T}^1(\mathbb{C})$ . By Lemma 5.1, we have the bound

$$\mu_{\mathbb{C}}\left(\left\{q\in\mathbb{T}^{1}(\mathbb{C})\,\big|\,|P(q)|\leq\varepsilon\right\}\right)\leq C\varepsilon^{\delta}.$$

Secondly, we prove the assertion when P is an arbitrary slice regular polynomial. Let  $I, J \in \mathbb{S}$  be two orthogonal imaginary units. By the Splitting Lemma 3.2, there are two polynomials  $F, G : \mathbb{C}_I \to \mathbb{C}_I$  such that

$$P(q) = F(q) + G(q)J$$
 and  $|P(q)|^2 = |F(q)|^2 + |G(q)|^2$ ,

Meanwhile, its symmetrization is a real polynomial with

$$P^{s}(q) = F(q)\overline{F(\overline{q})} + G(q)\overline{G(\overline{q})}$$
 and  $|P^{s}(q)|^{2} = |F(q)\overline{F(\overline{q})} + G(q)\overline{G(\overline{q})}|^{2}$ .

Via the Cauchy–Schwarz inequality, one finds that  $|P^s(q)| \leq |P(q)||P(\overline{q})|$ . So that if  $|P(q)| \leq \varepsilon$ , one has

$$|P^{s}(q)| \leq M\varepsilon$$
, where  $M = \max_{q \in \mathbb{T}^{1}(\mathbb{H})} |P(q)|$ .

However the measure of the latter is controlled by  $\leq C\varepsilon^{\delta}$ . We know thusly

$$\mu_{\mathbb{H}}\left(\left\{q \in \mathbb{T}^1(\mathbb{H}) \,\middle|\, |P(q)| \le \varepsilon\right\}\right) \le C'\varepsilon^{\delta}.$$

At this stage, we are now prepared to demonstrate Theorem 1.2.

Proof of Theorem 1.2. From Lemma 5.2 we see

$$\int_{|P| \le \varepsilon} \log |P| \, \mathrm{d}\mu(\mathbb{T}^1(\mathbb{H})) = O(\varepsilon^{\delta} \log \varepsilon),$$

as  $\varepsilon \to 0$ . Therefore, the integral converges absolutely as an improper Riemann integral.

Next we prove the asserted inequality. Let  $z = \cos \theta + I \sin \theta$ , then again by the Splitting Lemma 3.2, there are two polynomials  $F, G : \mathbb{C}_I \to \mathbb{C}_I$  such that

$$P(z) = F(z) + G(z)J$$
 and  $|P(z)|^2 = |F(z)|^2 + |G(z)|^2$ ,

and then

$$m_{\mathbb{H}}(P) = \frac{1}{2\pi} \int_0^{2\pi} \int_{\mathbb{S}} \log |f(\cos\theta + I\sin\theta)| \,\mathrm{d}\mu(I) \,\mathrm{d}\theta$$
$$= \frac{1}{4\pi} \int_0^{2\pi} \int_{\mathbb{S}} \log(|F(z)|^2 + |G(z)|^2) \,\mathrm{d}\mu(I) \,\mathrm{d}\theta.$$

Using Cauchy–Schwarz once more we find

$$2 \operatorname{m}_{\mathbb{H}}(P) \ge \frac{1}{4\pi} \int_{0}^{2\pi} \int_{\mathbb{S}} \log |F(z)\overline{F(\overline{z})} + G(z)\overline{G(\overline{z})}|^2 \,\mathrm{d}\mu(I) \,\mathrm{d}\theta = \operatorname{m}_{\mathbb{H}}(P^s),$$

where the inequality holds if and only if  $\overline{F(\overline{z})} = \lambda F(z)$  and  $G(\overline{z}) = \overline{\lambda}G(z)$  for any  $z \in \mathbb{H}$  (using the Identity Principle 3.3) with a fixed  $\lambda \in \mathbb{C}_I$ . As I, J varies on  $\mathbb{S}$ , we know that  $\lambda = 1$  and  $\overline{G(\overline{z})} = 0$ . This is exactly  $P(z) = \overline{P(\overline{z})}$ , that is, P is slice preserving.

In light of Theorem 1.2, we now turn to the task of determining the Mahler measure of slice regular polynomials. Using the Fundamental Theorem of Algebra for Quaternion 3.9, we reach at the following problem.

**Problem 5.3.** Given a slice regular quaternionic polynomial P(x) with its complete factorization

$$(q^2 - 2\operatorname{Re}\alpha_1 \cdot q + |\alpha_1|^2)^{m_1} * \dots (q^2 - 2\operatorname{Re}\alpha_\ell \cdot q + |\alpha_\ell|^2)^{m_\ell} * \prod^* (q - \beta_h)c$$

Determine its Mahler measure  $m_{\mathbb{H}}(P)$ .

In view of Theorem 4.4, our objective is narrowed down to find the Mahler measure of the \*-product of monomials. In contrast to real polynomials, there are no straightforward solutions, since  $m_{\mathbb{H}}(P * Q) = m_{\mathbb{H}}(P) + m_{\mathbb{H}}(Q)$  does not hold in general.

While the general task can indeed be quite challenging, there are specific cases where the complexity diminishes. For examples, when P is one-slice preserving, the computation is more approachable. The following proof provides us with an example illustrating how to obtain the Mahler measures of  $(x - i)^{*n}$ .

Proof of Theorem 1.3. Let q = a + rI with  $I = i \cos \theta + j \sin \theta \cos \psi + k \sin \theta \sin \psi$ . By the Representation Formula 3.4, we get

$$P^{*n}(q) = \frac{1 - Ii}{2}P(a + ri)^n + \frac{1 + Ii}{2}P(a - ri)^n.$$

Then by Proposition 4.2 we have

$$m(P^{*n}) = \frac{1}{\pi} \int_0^1 \int_0^\pi \log |P^{*,n}(a+rI)|^2 \frac{r^2 \sin \theta}{\sqrt{1-r^2}} \,\mathrm{d}\theta \,\mathrm{d}r$$
$$= \frac{n}{2} \log 2 + \frac{1}{\pi} \int_0^1 \int_0^\pi \log \left(\sum_{\ell \text{ even}} \binom{n}{\ell} r^\ell - \sum_{\ell \text{ odd}} \binom{n}{\ell} r^\ell \cos \theta\right) \frac{r^2 \sin \theta}{\sqrt{1-r^2}} \,\mathrm{d}\theta \,\mathrm{d}r.$$

Note that for any real numbers a > b > 0, we have

(9) 
$$\int_0^\pi \log(a - b\cos\theta)\sin\theta \,\mathrm{d}\theta = \log(a^2 - b^2) + \frac{a}{b}\log\left(\frac{a + b}{a - b}\right) - 2.$$

Let  $A^+(r) = \sum_{\ell \text{ even }} \binom{n}{\ell} r^{\ell}$  and  $A^-(r) = \sum_{\ell \text{ odd }} \binom{n}{\ell} r^{\ell}$ . Then we have

$$m(P^{*n}) = \frac{n}{2}\log 2 + \frac{1}{\pi} \int_0^1 \left(\log(A^+(r)^2 - A^-(r)^2) - 2\right) \frac{r^2}{\sqrt{1 - r^2}} dr + \frac{1}{\pi} \int_0^1 \left(\frac{A^+(r)}{A^-(r)}\log\frac{A^+(r) + A^-(r)}{A^+(r) - A^-(r)}\right) \frac{r^2}{\sqrt{1 - r^2}} dr.$$

Changing variable with  $r = \sin \theta$  in the first integral, we derive that

$$A^{+}(r)^{2} - A^{-}(r)^{2} = \cos^{2n}\theta.$$

So we know the first line turns out to be  $-\frac{1}{4}(n+2)$ . In the second integral, substituting  $r = \tanh t$  we find

$$\frac{A^+(r)}{A^-(r)} = \coth nt, \quad \log \frac{A^+(r) + A^-(r)}{A^+(r) - A^-(r)} = 2nt.$$

These yield the expected formula.

Note that if we set  $u = e^t$ , then

$$\operatorname{coth} nt \tanh^2 t \operatorname{sech} t = 2u \frac{(u^n + 1)(u^2 - 1)^2}{(u^n - 1)(u^2 + 1)^3}.$$

By partial decomposition we can always interpret  $m_{\mathbb{H}}((x-i)^{*n})$  with polylogarithms. In particular,

$$m_{\mathbb{H}}(x-i) = \frac{1}{4}, \quad m_{\mathbb{H}}((x-i)^{*2}) = \frac{5}{6},$$
$$m_{\mathbb{H}}((x-i)^{*3}) = \frac{31}{4} - \frac{12}{\pi} \left( \text{Li}_2(\zeta_6) - \text{Li}_2(-\zeta_6) - \text{Li}_2(\zeta_6^2) + \text{Li}_2(-\zeta_6^2) \right)$$
$$= \frac{31}{4} - 2\pi,$$

where  $\zeta_6 = e^{\frac{2\pi i}{6}}$  is the sextic root of unity. And

$$m_{\mathbb{H}}((x-i)^{*4}) = \frac{13}{3} - \frac{2\sqrt{2}}{\pi} \left( \text{Li}_2(\zeta_8) - \text{Li}_2(-\zeta_8) + \text{Li}_2(\zeta_8^3) - \text{Li}_2(-\zeta_8^3) \right)$$
$$= \frac{13}{3} - \frac{\sqrt{2}}{2}\pi,$$

where  $\zeta_8 = e^{\frac{2\pi i}{8}}$  is the octic root of unity.

Theorem 1.3 can be extended to the slice regular rational functions as well.

**Proposition 5.4.** Let  $n \in \mathbb{N}$  and let  $(x-i)^{*-n}$  denote the regular reciprocal of the iterated \*-product. Then

$$m_{\mathbb{H}}((x-i)^{*-n}) = \frac{2n}{\pi} \int_0^\infty t \coth nt \tanh^2 t \operatorname{sech} t \, \mathrm{d}t - \frac{2-n}{4}.$$

*Proof.* Let P(x) = x - i. Its conjugate and symmetrization are

$$P^{c}(x) = x + i, \quad P^{s}(x) = x^{2} + 1,$$

so we have

$$P^{*-n}(x) = (x^2 + 1)^{-n}(x+i)^{*n}.$$

Therefore

$$m_{\mathbb{H}}(P^{*-n}) = m_{\mathbb{H}}((x+i)^{*n}) - n m_{\mathbb{H}}(x^2+1)$$
  
=  $m_{\mathbb{H}}((x-i)^{*n}) - n m_{\mathbb{H}}(x^2+1).$ 

**Corollary 5.5.** It can be seen that as  $n \to \pm \infty$ , we have

$$\lim_{n \to \infty} \frac{m_{\mathbb{H}}((x-i)^{*n})}{n} = \frac{1+2G}{\pi} - \frac{1}{4},$$

where  $G = L(\chi_{-4}, 2)$  is the Catalan's constant.

*Proof.* Notice that  $\operatorname{coth} nt \sim 1 + \frac{1}{t}O(\frac{1}{n})$  as  $n \to \pm \infty$  and

$$\int_0^\infty t \tanh^2 t \operatorname{sech} t \, \mathrm{d}t = \frac{1}{2} + G$$

So the result follows directly from Theorem 1.3.

In order to generalize the above formula, we first introduce the  $\sigma$ -distance which appears in Gentili–Stoppato [11].

**Definition 5.6.** Let  $p, q \in \mathbb{H}$ , the  $\sigma$ -distance is given by

$$\sigma(q,p) = \begin{cases} |q-p| & \text{if } p, q \in \mathbb{C}_I \text{ for some } I \in \mathbb{S}, \\ \sqrt{(\operatorname{Re} q - \operatorname{Re} p)^2 + (|\operatorname{Im} q| + |\operatorname{Im} p|)^2} & \text{otherwise.} \end{cases}$$

In particular, if  $p \in \mathbb{C}_I$ , then

$$\sigma(q, p) = \max_{z \in \mathbb{C}_I \cap \mathbb{S}_q} \left\{ |z - p|, \, |\overline{z} - p| \right\}.$$

The following lemma is a strengthen version of the inequality as presented in Gentili–Stoppato [11, Theorem 6].

**Lemma 5.7.** Let  $I, J \in \mathbb{S}$  and  $p \in \mathbb{C}_I$ . Then for any  $q \in \mathbb{C}_J$ , we have

$$\frac{1}{2n} \log \left( \frac{1}{2} (1 - |\langle I, J \rangle|) \right) \le \frac{1}{n} \log |(q - p)^{*n}| - \sigma(q, p) \le 0$$

Proof. For convenience, we can put  $p \in \mathbb{C}$  and I = i. Let  $J = i \cos \theta + j \sin \theta \cos \theta_1 + k \sin \theta \sin \theta_1$ . If  $q \in \mathbb{C}$ , the inequalities become obvious because  $|(q - p)^{*n}|^{\frac{1}{n}} = |q - p| = \sigma(q, p)$ . If  $q \in \mathbb{C}_J$  with  $J \neq \pm i$ , then by Representation Formula 3.4 for  $z \in \mathbb{C}_I \cap \mathbb{S}_q$  we have

$$(q-p)^{*n} = \frac{1-JI}{2}(z-p)^n + \frac{1+JI}{2}(\overline{z}-p)^n.$$

If  $|z-p| \leq |\overline{z}-p|$  then the distance  $\sigma(p,q) = |\overline{z}-p|$ . Hence, we obtain  $\frac{1}{n} \log |(q-p)^{*n}| - \sigma(q,p) = \frac{1}{n} \log \left| \frac{1-JI}{2} \left( \frac{z-p}{\overline{z}-p} \right)^n + \frac{1+JI}{2} \right|$   $= \frac{1}{2n} \log \left( \frac{1}{2} (1-\cos\theta) + (1+\cos\theta) \left| \frac{z-p}{\overline{z}-p} \right|^n \right).$ 

One finds thusly

$$\frac{1}{2n}\log\left(\frac{1}{2}(1-\cos\theta)\right) \le \frac{1}{n}\log|(q-p)^{*n}| - \sigma(q,p) \le 0.$$

The case  $|z - p| > |\overline{z} - p|$  can be treated in the same manner.

**Theorem 5.8.** Let  $\alpha \in \mathbb{H}$  and let I be such that  $\alpha = \operatorname{Re} \alpha - I |\operatorname{Im} \alpha|$ . Then

$$\lim_{n \to +\infty} \frac{m_{\mathbb{H}}((x-\alpha)^{*n})}{n} = \frac{|\alpha|^2}{4} \cos 2\phi - \frac{1}{2\pi I} \operatorname{Im}\left[\left(\alpha - \frac{1}{\alpha}\right) - \left(\alpha^2 - \frac{1}{\alpha^2}\right) \operatorname{arctanh} \alpha + 4\operatorname{Li}_2(\alpha) - \operatorname{Li}_2(\alpha^2)\right].$$

where  $\phi = -\arccos \frac{\operatorname{Re} \alpha}{|\alpha|} \in (-\pi, 0]$  and the functions Li<sub>2</sub> and arctanh are regarded as the regular extensions of their complex holomorphic counterparts.

Proof of Theorem 5.8. By similarity, we may assume that  $\alpha \in \mathbb{C}$  and I = i. By Lemma 5.7, we have

$$\left|\frac{m_{\mathbb{H}}(P^{*n})}{n} - \int_{\mathbb{T}^1(\mathbb{H})} \log \sigma(q,\alpha) \, \mathrm{d}\mu(q)\right| \le \frac{1}{2n} \left|\int_0^{\pi} \int_{\mathbb{S}} \log\left(\frac{1 - |\langle i, J \rangle|}{2}\right) \, \mathrm{d}J \, \mathrm{d}\theta\right|.$$

This implies that

$$\lim_{n \to \infty} \frac{m_{\mathbb{H}}(P^{*n})}{n} = \int_{\mathbb{T}^1(\mathbb{H})} \log \sigma(q, \alpha) \, \mathrm{d}\mu(q).$$

For any  $q \in \mathbb{H} \setminus \mathbb{C}$  and  $p \in \mathbb{H}$ , we have  $\sigma(q, \alpha) = \sigma(p^{-1}qp, \alpha)$  is also "slice preserving". Therefore, using the same method as in Proposition 4.2 it can be deduced that

$$\int_{\mathbb{T}^1(\mathbb{H})} \log \sigma(q,\alpha) \, \mathrm{d}\mu(q) = \frac{1}{\pi} \int_0^\pi \log \left| (\cos \theta - \operatorname{Re} \alpha)^2 + (\sin \theta + |\operatorname{Im} \alpha|)^2 \right| \sin^2 \theta \, \mathrm{d}\theta$$
$$= \frac{1}{\pi} \int_0^\pi \log((1+|\alpha|^2 - 2|\alpha|\cos(\theta - \phi))\sin^2 \theta \, \mathrm{d}\theta.$$

The formula follows immediately from Lemma 4.1.

The method herein can be applied to arbitrary one-slice preserving polynomial.

**Theorem 5.9.** Let P(x) be a slice regular polynomial which preserves  $\mathbb{C}_I$ . Then as  $n \to +\infty$  one has

$$\frac{1}{n} \operatorname{m}_{\mathbb{H}}(P^{*n}) \sim \frac{2}{\pi} \int_0^{\pi} \log \max\{|P(e^{I\theta})|, |P(e^{-I\theta})|\} \sin^2 \theta \,\mathrm{d}\theta,$$

and as  $n \to -\infty$  one has,

$$\frac{1}{n} \operatorname{m}_{\mathbb{H}}(P^{*n}) \sim \frac{2}{\pi} \int_0^{\pi} \log \min\{|P(e^{I\theta})|, |P(e^{-I\theta})|\} \sin^2 \theta \,\mathrm{d}\theta.$$

*Proof.* If  $n \ge 0$ , the proof is exactly the same as the proof of Theorem 5.8. If n < 0, like Proposition 5.4, we get

$$\mathbf{m}_{\mathbb{H}}(P^{*-n}) = \mathbf{m}_{\mathbb{H}}((P^c)^{*n}) - n \,\mathbf{m}_{\mathbb{H}}(P^s).$$

The conjugate  $P^c$  and symmetrization  $P^s$  both preserve  $\mathbb{C}_I$ , and

$$P^{c}(e^{I\theta}) = \overline{P(e^{-I\theta})}, \quad P^{s}(e^{I\theta}) = P(e^{I\theta})\overline{P(e^{-I\theta})}.$$

These combines to give us the formula when n < 0.

Once more, we offer the following Mahler measures for one-slice preserving polynomial.

**Proposition 5.10.** We have

$$m_{\mathbb{H}}((x-i)*(x-2i)) = \frac{9}{16} + \log 2.$$

*Proof.* By Proposition 4.2, we find that

$$m_{\mathbb{H}}((x-i)*(x-2i)) = \frac{1}{2\pi} \int_0^{\pi} \int_0^{\pi} \log \left[ 2(5+4\sin^2\theta - 9\sin\theta\cos\theta_1) \right] \sin^2\theta\sin\theta_1 \,\mathrm{d}\theta \,\mathrm{d}\theta_1.$$

Using (9) we get

$$H(r) \coloneqq \int_0^\pi \log\left(5 + 4r^2 - 9r\cos\theta_1\right)\sin\theta_1\,\mathrm{d}\theta_1$$
  
=  $\frac{1}{9r}(4r^2 + 9r + 5)\log(r+1)(4r+5) - \frac{1}{9r}(4r^2 - 9r + 5)\log(r-1)(4r-5) - 2$ 

where  $r = \sin \theta$ . For the next step, we mention that H(r) = H(-r). This implies that

$$m_{\mathbb{H}}((x-i)*(x-2i)) = \frac{1}{4\pi} \int_0^{2\pi} H(\cos\theta) \cos^2\theta \,\mathrm{d}\theta + \frac{1}{2}\log 2.$$

The integral of H is in fact  $\frac{9}{16} + \frac{1}{2} \log 2$ , which can be computed by

$$\frac{1}{4\pi} \int_0^{2\pi} H(\cos\theta) \cos^2\theta \,\mathrm{d}\theta$$
  
=  $\frac{1}{18} (4A^+(3) + 9A^+(2) + 5A^+(1) + 4B^+(3) + 9B^+(2) + 5B^+(1)) - \frac{1}{2}$   
 $- \frac{1}{18} (4A^-(3) - 9A^-(2) + 5A^-(1) + 4B^-(3) - 9B^-(2) + 5B^-(1)),$ 

where  $A^{\pm}(n)$  and  $B^{\pm}(n)$  are given by the following formulas

$$A^{\pm}(n) \coloneqq \frac{1}{2\pi} \int_0^{2\pi} \log(1 \pm 1 \cos \theta) \cos^n \theta \, \mathrm{d}\theta = \begin{cases} \pm \frac{5}{6} & n = 3, \\ -\frac{1}{4} - \frac{1}{2} \log 2 & n = 2, \\ \pm 1 & n = 1, \end{cases}$$

and

$$B^{\pm}(n) \coloneqq \frac{1}{2\pi} \int_{0}^{2\pi} \log(5 \pm 4\cos\theta) \cos^{n}\theta \,\mathrm{d}\theta = \begin{cases} \pm \frac{37}{96} & n = 3, \\ -\frac{1}{16} + \log 2 & n = 2, \\ \pm \frac{1}{2} & n = 1. \end{cases}$$

One can use Cauchy's theorem to deduce these formulas, for example,

$$\frac{1}{2\pi} \int_0^{2\pi} \log(5+4\cos\theta) \cos^3\theta \,\mathrm{d}\theta = \frac{1}{4} \operatorname{Im} \oint_{|z|=1} \log(z+2) \left(z+\frac{1}{z}\right)^3 \frac{\mathrm{d}z}{z} \\ = \frac{\pi i}{2} \operatorname{Res} \frac{\log(z+2)}{z} \left(z+\frac{1}{z}\right)^3 = \frac{37}{96}.$$

Finally, we provide an example of Mahler measure of a slice regular polynomial which does not preserve any slice.

Proposition 5.11. We have

$$m_{\mathbb{H}}((x-i)*(x-j)) = \frac{\pi}{2} - 1.$$

*Proof.* Let us write x in the hyperspherical form

 $x = \sin\theta\sin\theta_1\cos\theta_2 + k\sin\theta\sin\theta_1\sin\theta_2 + i\cos\theta + j\sin\theta\cos\theta_1.$ 

In this representation, the Mahler measure takes the form as an integral

$$\mathbf{m}_{\mathbb{H}}((x-i)*(x-j)) = \frac{1}{4\pi^2} \int_0^{2\pi} \int_0^{\pi} \int_0^{\pi} \log A(\theta, \theta_1, \theta_2) \sin^2 \theta \sin \theta_1 \, \mathrm{d}\theta \, \mathrm{d}\theta_1 \, \mathrm{d}\theta_2$$

where

$$A(\theta, \theta_1, \theta_2) = A_1 \cdot A_2 = 4(1 - \cos \theta) \cdot \left(1 + \cos^2 \frac{\theta}{2} \sin^2 \theta_1 \sin 2\theta_2\right).$$

To compute it, we need to evaluate two separate integrals. The first integral, involving  $A_1$ , can be computed using Lemma 4.1

$$\int_{\mathbb{S}^3} \log A_1 \, \mathrm{d}\mu(\mathbb{S}^3) = \frac{1}{\pi} \int_0^\pi \log(4 - 4\cos\theta) \sin^2\theta \, \mathrm{d}\theta = \frac{1}{4} + \frac{1}{2}\log 2.$$

The second integral, involving  $A_2$ , requires expanding the log as

$$\log A_2 = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{1}{n} \cos^{2n} \frac{\theta}{2} \sin^{2n} \theta_1 \sin^n 2\theta_2.$$

It is not hard to find

$$\int_0^\pi \cos^{2n} \frac{\theta}{2} \sin^2 \theta \, \mathrm{d}\theta = 2\sqrt{\pi} \frac{\Gamma(n+\frac{3}{2})}{\Gamma(n+3)},$$
$$\int_0^\pi \sin^{2n+1} \theta_1 \, \mathrm{d}\theta_1 = \sqrt{\pi} \frac{\Gamma(n+1)}{\Gamma(n+\frac{3}{2})},$$

and

 $\int_0^{2\pi} \sin^n 2\theta_2 \,\mathrm{d}\theta_2 = 2\sqrt{\pi} \frac{\Gamma(\frac{n+1}{2})}{\Gamma(\frac{n}{2}+1)} \quad \text{if } n \text{ is even, otherwise } 0.$ 

Integrating term by term, we obtain

$$\int_{\mathbb{S}^3} \log A_2 \, \mathrm{d}\mu(\mathbb{S}^3) = -\frac{1}{8} \sum_{m=1}^{\infty} \frac{\left(\frac{1}{2}\right)_m}{m\left(m + \frac{1}{2}\right)(m+1)!}$$

Finally, we simplify our task to proving a specific identity

$$S \coloneqq \frac{1}{4} \sum_{m=1}^{\infty} \frac{\left(\frac{1}{2}\right)_m}{m\left(m + \frac{1}{2}\right)(m+1)!} = \frac{5}{2} + \log 2 - \pi.$$

Observe first that the expression

$$\frac{1}{4} \sum_{m=0}^{\infty} \frac{\left(\frac{1}{2}\right)_m}{\left(m + \frac{1}{2}\right)(m+1)!} x^m = \sqrt{x} \arcsin\sqrt{x} + (1-x)^{\frac{1}{2}} - 1,$$

implies that

$$S_1 \coloneqq \frac{1}{4} \sum_{m=1}^{\infty} \frac{\left(\frac{1}{2}\right)_m}{\left(m + \frac{1}{2}\right)(m+1)!} = \frac{\pi}{2} - \frac{3}{2}.$$

Meanwhile, we find that

$$S_2 \coloneqq \frac{1}{4} \sum_{m=0}^{\infty} \frac{\left(\frac{1}{2}\right)_m}{m(m+1)!} = \int_0^1 \left(\frac{1-\sqrt{1-x}}{2x^2} - \frac{1}{4x}\right) \mathrm{d}x = \frac{1}{2}\log 2 - \frac{1}{2}$$

Consequently, the sum S is

$$S = 2(S_2 - S_1) = \frac{5}{2} + \log 2 - \pi,$$

thus establishing the desired identity.

### 6. Examples on non-commutative linear polynomials

In this section, we give some examples of Mahler measure of non-commutative linear polynomials.

**Definition 6.1.** Let  $L : \mathbb{H} \to \mathbb{H}$  be an operator on quaternion. The operator is said to be *real-linear* if for any  $x, y \in \mathbb{H}$ ,

$$L(x+y) = L(x) + L(y),$$
$$L(qx) = qL(x), \text{ for } q \in \mathbb{R}.$$

A well-known fact is that every real-linear operator can be depicted as a noncommutative linear polynomial

$$L(x) = \sum_{i=1}^{n} p_i x q_i$$
, where  $p_i, q_i \in \mathbb{H}$  are non-zero.

For example, the quaternion conjugation can be written as

$$\overline{q} = -\frac{1}{2}(q + iqi + jqj + kqk).$$

Let  $x = x_1 + x_2 i + x_3 j + x_4 k$  and  $v = (x_1, x_2, x_3, x_4)^T$ , then there exist a real symmetric  $(4 \times 4)$ -matrix P such that

$$(10) |L(x)|^2 = v^T P v.$$

Let  $p_1, p_2, q_1, q_2 \in \mathbb{H}$  be non-zero quaternions. Within this section, we examine a particular type of linear polynomials by sum of two monomials

$$L(x) = p_1 x q_1 + p_2 x q_2.$$

Upon direct simplification, we find

$$m_{\mathbb{H}}(L) = \log |p_2| + \log |q_1| + m_{\mathbb{H}}(p_2^{-1}p_1x + xq_2q_1^{-1}).$$

Thus, our attention is directed towards those kinds of polynomials under the Sylvester form L(x) = ax + xb.

Before the proof of the Theorem 1.4, we prepare a simple lemma.

**Lemma 6.2.** Let  $a, b \in \mathbb{H}$  be non-zero and  $x = x_1 + x_2i + x_3j + x_4k$ . Then the four eigenvalues of the matrix P in Eq. (10) are

$$\lambda_1^+ = \lambda_2^+ = (\operatorname{Re} a + \operatorname{Re} b)^2 + (|\operatorname{Im} a| + |\operatorname{Im} b|)^2,$$
  
$$\lambda_1^- = \lambda_2^- = (\operatorname{Re} a + \operatorname{Re} b)^2 + (|\operatorname{Im} a| - |\operatorname{Im} b|)^2.$$

*Proof.* The eigenvalues can be calculated through direct computation. Refer, for example, to [30, Theorem 2.3.3].  $\Box$ 

Proof of Theorem 1.4. Let  $v = (x_1, x_2, x_3, x_4)^T$ , then Lemma 6.2 shows that there exists an orthogonal matrix Q such that

$$|ax + xb|^2 = v^T Q^T \Lambda Q v,$$

where  $\Lambda = \text{diag}(\lambda_1^+, \lambda_2^+, \lambda_1^-, \lambda_2^-)$ . Upon changing the variables, we have

$$m_{\mathbb{H}}(L) = \int_{\mathbb{T}^{1}(\mathbb{H})} \log |ax + xb| \, \mathrm{d}\mu(x)$$
  
$$= \frac{1}{2} \int_{\mathrm{SU}(2)} \log(\lambda^{+}|z|^{2} + \lambda^{-}|w|^{2}) \frac{\mathrm{d}z}{z} \, \mathrm{d}w \, \mathrm{d}\overline{w}$$
  
$$= \pi i \int_{0 \le |w| \le 1} \log(\lambda^{+} + (\lambda^{+} - \lambda^{-})|w|^{2}) \, \mathrm{d}w \, \mathrm{d}\overline{w}$$

The last integral gives

$$m_{\mathbb{H}}(L) = 2\pi^2 \left( \frac{\lambda^+ \log \lambda^+ - \lambda^- \log \lambda^-}{\lambda^+ - \lambda^-} - 1 \right).$$

Here we also exhibit a special case when a = -b. This in particular gives the quaternionic Mahler measure of the commutator [a, x] = ax - xa.

**Theorem 6.3.** Let  $a \in \mathbb{H}$  and  $c \in \mathbb{R}$ . Let L(x) = ax - xa + c. Then we have

$$m_{\mathbb{H}}(L) = \log |\mathrm{Im}\,a| + \frac{1}{8} \left( (4 + c_0^2) \log(4 + c_0^2) - 2c_0^2 \log c_0 - 4 \right),$$

where  $c_0 = |c/\operatorname{Im} a|$ .

*Proof.* Let us put  $a = \operatorname{Re} a + I |\operatorname{Im} a|$ . Then we get

$$m_{\mathbb{H}}(ax - xa + b) = m_{\mathbb{H}}(I|\operatorname{Im} a|x - xI|\operatorname{Im} a| + c)$$
$$= \log |\operatorname{Im} a| + m_{\mathbb{H}}(Ix - xI + c_0).$$

Observe that I commutes with  $\mathbb{C}_I$ , then letting I, J, K = IJ be an orthogonal basis of S we have

$$\log |I(z + Jw) - (z + Jw)I + c_0| = \log |2Kw + c_0|.$$

Let w = u - Iv, then the integral can be simplified to an integral on the unit disk

$$m_{\mathbb{H}}(Ix - xI + c_0) = \frac{1}{4\pi^2} \int_{|z|^2 + |w|^2 = 1} \log |2Kw + c_0| \frac{\mathrm{d}z}{z} \,\mathrm{d}w \,\mathrm{d}\overline{w}$$
$$= \frac{i}{2\pi} \int_{0 \le |w| \le 1} \log |2Kw + c_0| \,\mathrm{d}w \,\mathrm{d}\overline{w}.$$

Expanding this expression explicitly, we get

$$m_{\mathbb{H}}(Ix - xI + c_0) = \frac{1}{2\pi} \int_{u^2 + v^2 \le 1} \log\left(c_0^2 + 4u^2 + 4v^2\right) du dv$$
$$= \frac{1}{8} \left( (4 + c_0^2) \log(4 + c_0^2) - 2c_0^2 \log c_0 - 4 \right).$$

#### 7. Multivariable quaternionic Mahler measures

This section is dedicated to provide some examples of multivariable quaternionic Mahler measures. Recall for a non-commutative multivariable quaternionic polynomial  $P(x_1, \ldots, x_n)$ , its Mahler measure is the integral

$$\mathbf{m}_{\mathbb{H}}(P) = \int_{\mathbb{T}^n(\mathbb{H})} \log |P(x_1, \dots, x_n)| \, \mathrm{d}\mu(\mathbb{T}^n(\mathbb{H})).$$

The theoretical framework concerning slice regular functions of multiple quaternionic variables is elucidated in Ghiloni–Perotti [17]. Multivariate slice regular polynomials, as a special kind of slice regular functions, are polynomial functions  $P(x_1, \ldots, x_n)$  on  $\mathbb{H}^n$  under the form

$$P(x_1, \dots, x_n) = \sum_{\ell_1 + \dots + \ell_n = 0}^{\deg(P)} x_1^{\ell_1} \dots x_n^{\ell_n} a_{\ell_1, \dots, \ell_n},$$

where  $a_{\ell_1,\ldots,\ell_n} \in \mathbb{H}$ . In the recent papers [18, 19], Gori, Sarfatti and Vlacci study the zero sets and Hilbert Nullstellensatz of slice regular polynomials in several variables.

In the theorem presented below, we confirm the existence of Mahler measures for slice regular polynomials of two quaternion variables.

**Theorem 7.1.** Let P(x, y) be a slice regular quaternionic polynomial of two variables, then its Mahler measure  $m_{\mathbb{H}}(P)$  exists as an improper Riemann integral.

*Proof.* As in the proof of Theorem 1.2, we need to ascertain the estimation

$$\mu_{\mathbb{H}}\left(\left\{(x,y)\in\mathbb{T}^{2}(\mathbb{H})\mid|P(x,y)|\leq\varepsilon\right\}\right)\leq C\varepsilon^{\delta}$$

Let Q(x, y) be the following one-variable symmetrization of P(x, y) with respect to the variable x,

$$Q(x,y) = \sum_{\ell} x^{\ell} \sum_{h} \sum_{h_1,h_2} y^{h_1} \overline{y}^{h_2} a_{h,h_1} \overline{a}_{\ell-h,h_2}.$$

Since P(x, y) is left slice regular in x, using the Cauchy–Schwarz estimation as in Lemma 5.2, it is enough to settle

$$\mu_{\mathbb{H}}\left(\left\{(x,y)\in\mathbb{T}^2(\mathbb{H})\,\big|\,|Q(x,y)|\leq\varepsilon\right\}\right)\leq C\varepsilon^{\delta}.$$

The function Q(x, y) is a real polynomial with respect to the variable x. We start by induction on the degree n of Q(x, y) in x. We may factorize Q(x, y) by

$$Q(x,y) = A(y,\overline{y}) \prod_{\ell=1}^{m_1} (x^2 - 2\alpha_\ell(y,\overline{y})x + \beta_\ell(y,\overline{y})) \prod_{h=1}^{m_2} (x - \gamma_h(y,\overline{y})),$$

where  $A, \alpha_{\ell}, \beta_h$  and  $\gamma_h$  are real-valued algebraic functions with respect to y and  $\overline{y}$ . Note that these functions rely only on Re y and |Im y|, not taking into account

the imaginary unit within  $y = \operatorname{Re} y + I |\operatorname{Im} y|$ . Thus, if  $|Q(x,y)| < \varepsilon$ , then either

 $|A(y,\overline{y})| \le \varepsilon^{\frac{1}{n+1}},$ 

or one of the real factors

$$|x^2 - 2\alpha_\ell(y,\overline{y})x + \beta_\ell(y,\overline{y})|$$
 or  $|x - \gamma_h(y,\overline{y})| \le \varepsilon^{\frac{1}{n+1}}$ .

In the first case,  $A(y, \overline{y}) = A(y, y^{-1})$  is a Laurent polynomial which relies only on Re y and |Im y|. So we can deduce that

$$\mu_{\mathbb{H}}\left(\left\{y \in \mathbb{T}^{1}(\mathbb{H}) \mid |A(y, y^{-1})| \leq \varepsilon\right) = \mu_{\mathbb{C}}\left(\left\{y \in \mathbb{T}^{1}(\mathbb{C}) \mid |A(y, y^{-1})| \leq \varepsilon\right).\right.$$

The latter is thusly controlled by  $C_1 \varepsilon^{\delta_1}$  with the help of Lemma 5.1.

In the latter cases, by the same fashion, we have

$$\mu_{\mathbb{H}}\big(\big\{(x,y)\in\mathbb{T}^2(\mathbb{H})\,\big|\,|x-\gamma_h(y,\overline{y})|\leq\varepsilon\big)\\ =\mu_{\mathbb{C}}\,\big(\big\{(x,y)\in\mathbb{T}^1(\mathbb{C})\,\big|\,|x-\gamma_h(y,y^{-1})|\leq\varepsilon\big)\,.$$

For fixed  $y \in \mathbb{C}$ , using again Lemma 5.1 for  $P(x) = x - \gamma_h(y, y^{-1})$ , it is apparent that

$$\mu_{\mathbb{C}}\left(\left\{x \in \mathbb{T}^{1}(\mathbb{C}) \mid |x - \gamma_{h}(y, y^{-1})| \leq \varepsilon\right\} \leq C_{2}\varepsilon^{\delta_{2}}.$$

This makes us realize that

$$\mu_{\mathbb{H}}\left(\left\{(x,y)\in\mathbb{T}^{1}(\mathbb{H})\mid |x-\gamma_{h}(y,y^{-1})|\leq\varepsilon\right)\leq C_{2}\varepsilon^{\delta_{2}}\right)$$

For the quadratic factor  $x^2 - 2\alpha_\ell(y, \overline{y})x + \beta_\ell(y, \overline{y})$ , using the same approach yields one same bound.

Put everything together, we find at last there exist  $C, \delta > 0$  so that

$$\mu_{\mathbb{H}}\left(\left\{(x,y)\in\mathbb{T}^2(\mathbb{H})\mid |P(x,y)|\leq\varepsilon\right\}\right)\leq C\varepsilon^{\delta}.$$

*Remark.* In [18], Gori–Sarfatti–Vlacci offer the following example, the slice regular polynomial P(x, y) = xy - k has its *multivariable symmetrization* 

$$P^s(x,y) = x^2 y^2 + 1.$$

However, it holds P(i, j) = 0 while  $P^s(i, j) = 2$ . It can be seen that the zero of a polynomial P might no longer be the zero of its symmetrization, rendering the proof in one variable ineffective in the multivariable case.

Subsequently, we compute the quaternionic Mahler measure  $m_{\mathbb{H}}(1 + x + y)$ .

Proof of Theorem 1.5. By changing the variable  $y \to yx$ , we get  $m_{\mathbb{H}}(x+y+1) = m_{\mathbb{H}}(x+yx+1)$ . By definition, we have

$$m_{\mathbb{H}}((1+y)x+1) = \int_{\mathbb{T}^{2}(\mathbb{H})} \log |(1+y)x+1| \,\mathrm{d}\mu(x) \,\mathrm{d}\mu(y)$$
$$= \frac{2}{\pi} \int_{0}^{\pi} \int_{\mathbb{T}^{1}(\mathbb{H})} \log |(1+e^{i\theta})x+1| \sin^{2}\theta \,\mathrm{d}\mu(x) \,\mathrm{d}\theta$$

Since  $|1 + e^{i\theta}| \ge 1$  if and only if  $\theta \in [0, \frac{2}{3}\pi]$ , by Lemma 4.1, we have

$$m_{\mathbb{H}}(x+y+1) = \frac{1}{2\pi} \int_{\frac{2}{3}\pi}^{\pi} |1+e^{i\theta}|^2 \sin^2\theta \,\mathrm{d}\theta + \frac{1}{2\pi} \int_0^{\frac{2}{3}\pi} \frac{\sin^2\theta}{|1+e^{i\theta}|^2} \,\mathrm{d}\theta + \frac{2}{\pi} \int_0^{\frac{2}{3}\pi} \log|1+e^{i\theta}| \sin^2\theta \,\mathrm{d}\theta.$$

Denote the above integrals by  $I_1, I_2$  and  $I_3$  respectively. The integrals  $I_1, I_2$  are easy to calculate via  $|1 + e^{i\theta}|^2 = 2 + 2\cos\theta$ . A direct calculation gives

$$I_1 = \frac{1}{6} - \frac{\sqrt{3}}{4\pi}$$
 and  $I_2 = \frac{1}{6} - \frac{\sqrt{3}}{8\pi}$ .

For the third integral, we have

$$I_3 = \frac{2}{\pi} \operatorname{Re} \int_0^{\frac{2}{3}\pi} \log(1+e^{i\theta}) \sin^2\theta \, \mathrm{d}\theta = -\frac{1}{2\pi i} \operatorname{Im} \int_1^{\rho} \log(1+z) \left(z - \frac{1}{z}\right)^2 \frac{\mathrm{d}z}{z},$$

where  $\rho = e^{\frac{2}{3}\pi i}$ . Since

$$\int \left(z + \frac{1}{z^3}\right) \log(1+z) \, \mathrm{d}z$$
  
=  $\frac{1}{4z^2} \left(2(z^4 - 1)\log(1+z) - 2z^2\log(z) - 2z + 2z^3 - z^4\right),$ 

it can be deduced that

$$\operatorname{Im} \int_{1}^{\rho} \left( z + \frac{1}{z^{3}} \right) \log(1+z) \, \mathrm{d}z = \left( \frac{5\sqrt{3}}{8} - \frac{\pi}{3} \right) i.$$

Finally, for the integral of  $\log(1+z)/z$ , by expanding  $\log(1+z)$  by Taylor series, we have

$$\operatorname{Im} \int_{1}^{\rho} \frac{\log(1+z)}{z} \, \mathrm{d}z = \operatorname{Im} \int_{1}^{\rho} \sum_{n \ge 1} \frac{(-1)^{n-1} z^{n-1}}{n} \, \mathrm{d}z$$
$$= \sum_{n \ge 1} \frac{(-1)^{n-1} i \sin \frac{2n\pi}{3}}{n^2} = \frac{3\sqrt{3} i}{4} L(\chi_{-3}, 2).$$

Combining the results together, we get the desired result.

More generally, we can consider the Mahler measure of ax + by + c. If |a|, |b|, |c| forms a triangle, the above calculation still works, yet a general formula would be of much difficulty. If |a|, |b|, |c| do not form a triangle, following the same argument as Maillot [26], we are able to give the proof of 1.6.

Proof of Theorem 1.6. With the same argument as in Theorem 1.1, we may assume that  $a, b, c \in \mathbb{R}^+$ . Without loss of generality, suppose that  $a \ge b \ge c$ . By our

assumption, |a|, |b|, |c| do not form a triangle, so for any  $y \in \mathbb{T}^1(\mathbb{H})$ , we have  $|a+by| \geq a-b \geq c$ . For a fixed  $y \in \mathbb{T}^1(\mathbb{H})$ , by Theorem 1.1, we have

$$\int_{\mathbb{T}^1(\mathbb{H})} \log |(a+by)x+c| \, \mathrm{d}\mu(x) = \log |a+by| + \frac{c^2}{4|a+by|^2}.$$

So we get

$$m_{\mathbb{H}}(ax+by+c) = \int_{\mathbb{T}^1(\mathbb{H})} \int_{\mathbb{T}^1(\mathbb{H})} \log|ax+byx+c|\,\mathrm{d}\mu(x)\,\mathrm{d}\mu(y)$$
$$= \int_{\mathbb{T}^1(\mathbb{H})} \log|a+by| + \frac{c^2}{4|a+by|^2}\,\mathrm{d}\mu(y)$$
$$= \frac{2}{\pi} \int_0^{\pi} \left(\log|a+be^{i\theta}| + \frac{c^2}{4|a+be^{i\theta}|^2}\right) \sin^2\theta \,\mathrm{d}\theta.$$

By applying Lemma 4.1, the first term equals to  $\log a + \frac{b^2}{4a^2}$ . Following from Theorem 4.3, the second term equals to  $\frac{c^2}{4a^2}$ . These combines to complete the formula.

In the following theorem, we illustrate an example of the Mahler measure for a slice regular polynomial in two variables.

**Theorem 7.2.** We have the identity

$$\mathbf{m}_{\mathbb{H}}(x^2 - y^2) = 0,$$

*Proof.* The Mahler measure can be reshaped to

$$m_{\mathbb{H}}(x^2 - y^2) = m_{\mathbb{H}}((xy)^2 - y^2) = m_{\mathbb{H}}(xy - yx^{-1}).$$

Applying Theorem 1.4 in the variable y yields

$$m_{\mathbb{H}}(xy - yx^{-1}) = 2\pi^2 \int_{\mathbb{T}^1(\mathbb{H})} \log(4|\mathrm{Im}\,x|^2) - 1\,\mathrm{d}\mu(x).$$

Observe that the integrand depends only on |Im x|. Following the same pattern as Proposition 4.2, we get

$$\int_{\mathbb{T}^1(\mathbb{H})} \log |\operatorname{Im} x| \, \mathrm{d}\mu(x) = \frac{2}{\pi} \int_0^\pi \log(\sin\theta) \sin^2\theta \, \mathrm{d}\theta.$$

The latter integral can be computed in the same manner as Lemma 4.1. Via the expansion

$$\log(2\sin\theta) = -\sum_{n=1}^{\infty} \frac{\cos 2n\theta}{2n},$$

one finds that

$$\frac{2}{\pi} \int_0^\pi \log(\sin\theta) \sin^2\theta \,\mathrm{d}\theta = \frac{2}{\pi} \int_0^\pi \log(2\sin\theta) \sin^2\theta \,\mathrm{d}\theta - \log 2$$
$$= \frac{1}{2} - \log 2.$$

Thus

$$m_{\mathbb{H}}(x^2 - y^2) = 4\pi^2 \int_{\mathbb{T}^1(\mathbb{H})} \log |\operatorname{Im} x| \, \mathrm{d}\mu(x) - 2\pi^2(\log 4 - 1) = 0.$$

Below here are two examples of non-slice-regular polynomials in two variables we present for consideration.

**Theorem 7.3.** We have the following identities

$$m_{\mathbb{H}}(xy - yx) = 0,$$
$$m_{\mathbb{H}}(xy - yx + 1) = \frac{1}{4} \left( 1 - \sqrt{5} + 3\log\frac{3 + \sqrt{5}}{2} \right).$$

*Proof.* Using Theorem 6.3 in the variable y, we get

$$m_{\mathbb{H}}(xy - yx) = \int_{\mathbb{T}^1(\mathbb{H})} \log |\operatorname{Im} x| \, \mathrm{d}\mu(x) + \log 2 - \frac{1}{2}.$$

Recall the proof of Theorem 7.2 where we get

$$\int_{\mathbb{T}^1(\mathbb{H})} \log |\operatorname{Im} x| \, \mathrm{d}\mu(x) = \frac{1}{2} - \log 2.$$

This proves the given identity. It is analogous that Mahler measure of xy - yx + 1 can be reduced to

$$m_{\mathbb{H}}(xy - yx + 1) = \frac{1}{2} - \log 2 + \frac{1}{\pi} \int_0^{\pi} M(\theta) \, \mathrm{d}\theta,$$

where

$$M(\theta) = \left(\frac{1}{4} + \sin^2\theta\right)\log(5 - 4\cos^2\theta) - 2\log(\sin\theta)\sin^2\theta - \sin^2\theta.$$

The calculation uses exactly the same trick as Theorem 7.2 and Lemma 4.1 before, so we omit the detail.  $\hfill \Box$ 

# 8. QUATERNION LEHMER PROBLEM

In [24], Lehmer mentioned the question, what is the smallest Mahler measure for monic integer polynomials. In addition to his question, he gave the following example of degree 10, irreducible reciprocal polynomials

$$L_{10}(x) = x^{10} + x^9 - x^7 - x^6 - x^5 - x^4 - x^3 + x + 1,$$

where we have  $m_{\mathbb{C}}(L_{10}) = 0.16235...$  are conjectured as the possible smallest Mahler measures.

A parallel question is, what is the minimal quaternionic Mahler measure that a monic irreducible integer polynomial f could achieve? The requirement of irreducibility here is crucial. As the quaternion Mahler measures may not always be positive, consider the following example

$$m_{\mathbb{H}}((x^2+1)^n) = -\frac{1}{2}n \to -\infty \text{ as } n \to \infty.$$

**Conjecture 8.1** (Quaternionic Lehmer conjecture). There is a constant c such that for all monic, irreducible polynomial  $P(x) \in \mathbb{Z}[x]$ , we have

$$m_{\mathbb{H}}(P) \ge c.$$

One starts by observing the quaternionic Mahler measure of cyclotomic polynomials.

**Proposition 8.2.** Let  $\Phi_n(x)$  be the n-th cyclotomic polynomial. Then we have

$$m_{\mathbb{H}}(\Phi_n) = \begin{cases} \frac{1}{4}\mu(n) & \text{if } 2 \nmid n\\ \frac{1}{4}\mu(n/2) & \text{if } 2 \mid n, \ 4 \nmid n\\ \frac{1}{2}\mu(n/2) & \text{if } 4 \mid n, \end{cases}$$

where  $\mu(n)$  is the Möbius function.

*Proof.* Since  $\Phi_n(x) = \prod_{(d,n)=1} (x - \zeta_n^d)$ , by Theorem 1.1, we have

$$m_{\mathbb{H}}(\Phi_n) = \frac{1}{2} \sum_{(d,n)=1} m_{\mathbb{H}}((x-\zeta_n^d)(x-\zeta_n^{-d})) = \frac{1}{4} \sum_{(d,n)=1} \cos\frac{4d\pi}{n}.$$

The right-hand side is the well-known Ramanujun's sum, see for instance [20, Theorem 272].  $\hfill \Box$ 

The following proposition offers a trivial bound for quaternionic Mahler measure.

**Proposition 8.3.** If P(x) is a monic real polynomial of degree n, then we have

$$\mathbf{m}_{\mathbb{H}}(P) \ge -\frac{1}{2} \left\lfloor \frac{n}{2} \right\rfloor,$$

where the equality holds if and only if  $P(x) = (x^2 + 1)^m$  when n = 2m is even, and  $P(x) = x(x^2 + 1)^m$  when n = 2m + 1 is odd.

*Proof.* With (1) and (2) in Theorem 1.1, one finds the smallest possible Mahler measure of one monomial happens when

$$m_{\mathbb{H}}(x-\alpha) \ge m_{\mathbb{H}}(x) = 0,$$

the smallest possible Mahler measure of a quadratic polynomial happens when

$$m_{\mathbb{H}}(x^2 - 2\operatorname{Re}\alpha \cdot x + |\alpha|^2) \ge m_{\mathbb{H}}(x^2 + 1) = -\frac{1}{2}$$

Then using (3) of Theorem 1.1 we get the lower bound of Mahler measures.  $\Box$ 

Employing the Cauchy–Schwarz inequality as in Theorem 1.2, we find the following lower bound for slice regular polynomials. It is natural to wonder if the lower bound in Proposition 8.3 still holds when n is odd.

**Corollary 8.4.** Let P(x) be a monic slice regular quaternionic polynomial of degree n, then we have

$$\mathrm{m}_{\mathbb{H}}(P) \ge -\frac{n}{4},$$

where the equality holds only if n = 2m is even and  $P(x) = (x^2 + 1)^m$ .

The formula of Mahler measures simplifies considerably when all roots of the polynomial are located either outside or inside the unit circle.

**Proposition 8.5.** Let  $P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_0$  be a real polynomial with all roots inside the unit circle, then we have

$$m_{\mathbb{H}}(P) = \frac{a_{n-1}^2 - 2a_{n-2}a_n}{4a_n^2} + \log|a_n|.$$

If all the roots of P lie outside the unit circle, then we have

$$m_{\mathbb{H}}(P) = \frac{a_1^2 - 2a_2a_0}{4a_0^2} + \log|a_0|.$$

*Proof.* Note that the roots of  $\widetilde{P}(x) := x^n P(1/x)$  are inside the unit circle if all roots of P(x) are outside the unit circle, so we only need to prove the first assertion. Suppose  $\beta_\ell$  is a complex root of P(x), then by Theorem 1.1, we have

$$m_{\mathbb{H}}(x^2 - 2\operatorname{Re}\beta_{\ell} \cdot x + |\beta_{\ell}|^2) = \frac{1}{2}|\beta_{\ell}|^2 \cos 2\phi_{\ell} = \frac{1}{4}(\beta_{\ell}^2 + \overline{\beta}_{\ell}^2).$$

So by factoring P as in (1) and applying the Theorem 1.1 again, we get

$$m_{\mathbb{H}}(P) = \log|a_n| + \frac{1}{4} \sum_{h,\ell} (\alpha_h^2 + \beta_\ell^2 + \overline{\beta}_\ell^2) = \log|a_n| + \frac{a_{n-1}^2 - 2a_{n-2}a_n}{4a_n^2}.$$

As a direct consequence, this allows us to explore the quaternionic Mahler measure of a real polynomial under the power change of the variable. While the classical Mahler measure is known to remain invariant under the transformation  $x \mapsto x^r$ , this property does not hold in our context.

**Proposition 8.6.** Let P(x) be a real polynomial and r > 2 be a positive integer, then

$$m_{\mathbb{H}}(P(x^r)) = m_{\mathbb{C}}(P).$$

Proof. Suppose P(x) factors completely as in (1). Since the roots of  $x^r - \alpha_h$  and  $x^{2r} - 2 \operatorname{Re} \beta_{\ell} \cdot x + |\beta_{\ell}|^2$  are inside or outside the unit circle depending respectively on either  $|\alpha_h|, |\beta_{\ell}| \leq 1$  or  $\geq 1$ , so by Proposition 8.5, we have  $m_{\mathbb{H}}(x^r - \alpha_h) = \log^+ |\alpha_h|$  and  $m_{\mathbb{H}}(x^{2r} - 2 \operatorname{Re} \beta_{\ell} \cdot x + |\beta_{\ell}|^2) = \log^+ |\beta_{\ell}|^2$ . Then the result is clear.

Let P(z) be a monic irreducible polynomial of degree N with a complex factorization

$$P(z) = \prod_{\ell=1}^{N} (z - z_{\ell}) = z^{n} + a_{N-1} z^{N-1} + \dots + a_{0}.$$

Numerical tests show that the polynomials with negative quaternionic Mahler measure are highly rare. The quaternionic Mahler measure of P becomes much smaller when the roots  $z_j$  are closer to the imaginary units  $\pm i$  on the unit circle. For an arc I on the unit circle, let N(I; P) denote the number of complex zeros  $z_j$  with arg  $z_j \in I$ . The discrepancy of P is defined by

$$\mathcal{D}(P) = \max_{I} \left| N(I; P) - \frac{I}{2\pi} N \right|.$$

Recall the famous result of Erdös–Turán [7] (see also Mignotte [27])

$$\mathcal{D}(P) \le \frac{8}{\pi} \sqrt{Nh(P)},$$

where

$$h(P) = \frac{1}{2\pi} \int_0^{2\pi} \log^+ \frac{|P(e^{i\theta})|}{\sqrt{|a_0|}} \,\mathrm{d}\theta \quad \text{with } \log^+(x) = \max\{0, \log(x)\}.$$

Note the constant  $8/\pi$  is given by Soundararajan [33]. It indicates that the equidistribution of the angles of roots if the length of P is not too large.

**Proposition 8.7.** Let  $P(z) \in \mathbb{Z}[x]$  be a monic irreducible polynomial. Then we have

$$\mathbf{m}_{\mathbb{H}}(P) \ge \mathbf{m}_{\mathbb{C}}(P) - \frac{8}{\pi}\sqrt{Nh(P)}.$$

*Proof.* Let  $\nu(\theta) = N([0, \theta]; P)$  denote the number of roots within the interval  $[0, \theta]$ . Then by Theorem 1.1, one get

$$m_{\mathbb{H}}(P) \ge m_{\mathbb{C}}(P) - \frac{1}{4} \sum_{\ell=1}^{N} \cos(2\phi_{\ell})$$
$$= m_{\mathbb{C}}(P) - \frac{1}{4} \int_{0}^{2\pi} \cos(2\theta) \, \mathrm{d}\nu(\theta)$$
$$\ge m_{\mathbb{C}}(P) - \frac{8}{\pi} \sqrt{Nh(P)}.$$

Here the last line follows by the Erdös–Turán inequality and the total variation of  $\cos 2x$  is 4.

Following Hughes–Nikeghbali [21], we derive the following result the same way as Proposition 8.7.

**Proposition 8.8.** Let  $(P_N)$  be a sequence of polynomials, with

$$P_N(z) = \sum_{k=0}^N a_{N,k} z^k,$$

such that  $a_{N,0}a_{N,N} \neq 0$  for all N, let

$$L_N(P_N) = \log\left(\sum_{k=0}^N |a_{N,k}|\right) - \frac{1}{2}\log|a_{N,0}| - \frac{1}{2}\log|a_{N,N}|.$$

Then if  $L_N(P_N) = O(1)$ , we have

$$\frac{1}{\sqrt{N}} |\mathbf{m}_{\mathbb{H}}(P_N) - \mathbf{m}_{\mathbb{C}}(P_N)| = O(1) \qquad as \ N \to \infty.$$

**Example 8.9.** By examining the polynomial of degree  $\leq 10$  with coefficients in [-3,3], we found the minimal quaternionic Mahler measure is reached by the following polynomial of degree 8

$$Q_8(x) = x^8 - x^7 + x^6 - 2x^5 + x^4 - 2x^3 + x^2 - x + 1.$$

Its Mahler measure is  $-0.26975\cdots$ . This example is related to cyclotomic polynomial by

$$Q_8(x) = \Phi_5(x)\Phi_{10}(x) + x\Phi_3(x)\Phi_4(x)\Phi_6(x).$$

Meanwhile, the quaternionic Mahler measure of Lehmer's polynomial  $L_{10}$  is

$$m_{\mathbb{H}}(L_{10}) = 0.247131\cdots$$

**Example 8.10.** Inspired by the observation of  $Q_8$ , by testing millions of combinations involving cyclotomic polynomials, we find the following two examples with Mahler measure less than -0.5. Let

$$Q_{66}(x) = \prod_{h=3}^{10} \Phi_h(x) \Phi_{14}(x) \Phi_{18}(x) \Phi_{90}(x) + x \Phi_{204}(x),$$

then  $Q_{66}$  is of degree 66 and

$$m_{\mathbb{H}}(Q_{66}) = -0.5829\cdots$$

Let

$$Q_{50}(x) = \Phi_4(x)\Phi_5(x)\Phi_6(x)\Phi_7(x)\Phi_{10}(x)\Phi_{80}(x) + x\Phi_{140}(x),$$

then  $Q_{50}$  is of degree 50 and

$$m_{\mathbb{H}}(Q_{50}) = -0.5145\cdots$$

It can thus be inferred that, if the Conjecture 8.1 is true, we have at least

 $c \leq -0.5829 \cdots$ .

#### QUATERNIONIC MAHLER MEASURE

#### References

- [1] H. Akatsuka, Zeta Mahler measures, Journal of Number Theory 129 (2009), pp. 2713-2734.
- [2] A. Altavilla, C. de Fabritiis, s-Regular functions which preserve a complex slice, Annali di Matematica 197 (2018), pp. 1269–1294.
- [3] F. Brunault, W. Zudilin, Many variations of Mahler measures—a lasting symphony, Australian Mathematical Society Lecture Series, Volume 28, Cambridge University Press, Cambridge, 2020.
- [4] D. W. Boyd, Mahler's measure and special values of L-function, Experimental Mathematics 7 (1998), pp. 341–349.
- [5] C. Deninger, Deligne periods of mixed motives, K-theory and the entropy of certain Z<sup>n</sup>actions, Journal of the American Mathematical Society 10 (1997), pp. 259–281.
- [6] S. Eilenberg, I. Niven, The "fundamental theorem of algebra" for quaternions, Bulletin of the American Mathematical Society (4) 50 (1944), pp. 246–248.
- [7] P. Erdös, P. Turán, On the distribution of roots of polynomials, Annals of Mathematics 51 (1950), pp. 105–119.
- [8] G. Everest, B. ní Fhlathúin, The elliptic Mahler measure, Mathematical Proceedings of the Cambridge Philosophical Society 120 (1996), pp. 13–25.
- [9] G. Everest, T. Ward, *Heights of Polynomials and Entropy in Algebraic Dynamics*, Universitext, Springer-Verlag London (1999).
- [10] G. Gentili, C. Stoppato, Zeros of regular functions and polynomials of a quaternionic variable, Michigan Mathematical Journal 56 (2008), pp. 655-667.
- G. Gentili, C. Stoppato, Power series and analyticity over the quaternions, Mathematische Annalen 352 (2012), pp. 113–131.
- [12] G. Gentili, C. Stoppato, D.C. Struppa, Regular functions of a quaternionic variable, Berlin: Springer (2013).
- [13] G. Gentili, D.C. Struppa, A new approach to Cullen-regular functions of a quaternionic variable, Comptes Rendus de l'Académie des Sciences Paris 342 (2006), no. 10, pp. 741–744.
- [14] G. Gentili, D.C. Struppa, A new theory of regular functions of a quaternionic variable, Advances in Mathematics 216 (2007), pp. 279–301.
- [15] G. Gentili, D.C. Struppa, On the Multiplicity of Zeroes of Polynomials with Quaternionic Coefficients, Milan Journal of Mathematics 76 (2008), pp. 15–25.
- [16] R. Ghiloni, A. Perotti, Slice regular functions on real alternative algebras, Advances in Mathematics 226 (2010), pp. 1662-1691.
- [17] R. Ghiloni, A. Perotti, Slice regular functions in several variables, Mathematische Zeitschrift 302 (2022), pp. 295–351.
- [18] A. Gori, G. Sarfatti, F. Vlacci, Zero sets and Nullstellensatz type theorems for slice regular quaternionic polynomials, Linear Algebra and its Applications Volume 685 (2024), pp. 162-181.
- [19] A. Gori, G. Sarfatti, F. Vlacci, A Strong Version of the Hilbert Nullstellensatz for slice regular polynomials in several quaternionic variables, arXiv preprint (2024), arXiv:2402.16784.
- [20] G. Hardy, and E. Wright, An Introduction to the Theory of Numbers, Oxford, Fourth edition, (1975).
- [21] C. P. Hughes, A. Nikeghbali, The zeros of random polynomials cluster uniformly near the unit circle, Compositio Mathematica 144 (3) (2008), pp. 734-746.
- [22] M. N. Lalín, S. Roy, Evaluations of the areal Mahler measure of multivariable polynomials, Journal of Number Theory 254 (2024), pp. 103-145.
- [23] M. N. Lalín, S. Roy, *The areal Mahler measure under a power change of variables*, To appear, Illinois Journal of Mathematics.

- [24] D. H. Lehmer, Factorization of certain cyclotomic functions, Annals of Mathematics (2) 34 (1933), no. 3, pp. 461–479.
- [25] K. Mahler, On some inequalities for polynomials in several variables, Journal of the London Mathematical Society 37 (1962), pp. 341–344.
- [26] V. Maillot, Géométrie d'Arakelov des variétés toriques et fibrés en droites intégrables, Mémoires de la Société Mathématique de France, Serie 2, no. 80 (2000).
- [27] M. Mignotte, Remarque sur une question relative à des fonctions conjuguées, Comptes Rendus de l'Académie des Sciences Series I Mathematics 315 (8) (1992), pp. 907–911.
- [28] I. E. Pritsker, An areal analog of Mahler's measure, llinois Journal of Mathematics, Volume 52, Number 2 (2008), pp. 347–363.
- [29] I. E. Pritsker, *Polynomial inequalities*, Mahler's measure and multipliers, Number theory and polynomials, LMS Lecture Notes, Volume 352, Cambridge University Press, Cambridge (2008), pp. 255–276.
- [30] L. Rodman, *Topics in quaternion linear algebra*, Princeton Series in Applied Mathematics; Princeton University Press: Princeton, NJ, USA, 2014.
- [31] F. Rodriguez-Villegas, Modular Mahler measures, I, Topics in number theory (University Park, PA, 1997), Mathematical Applications, Volume 467, Kluwer Academic Publishers, pp. 17–48.
- [32] C. J. Smyth, On measures of polynomials in several variables, Bulletin of the Australian Mathematical Society 23 (1981), no. 1, pp. 49–63.
- [33] K. Soundararajan, Equidistribution of Zeros of Polynomials, The American Mathematical Monthly 126 (3) (2019), pp. 226-236.