# BOUNDS FOR THE INDEPENDENCE AND CHROMATIC NUMBERS OF LOCALLY SPARSE GRAPHS

ABHISHEK DHAWAN

ABSTRACT. In this note we consider a more general version of local sparsity introduced recently by Anderson, Kuchukova, and the author. In particular, we say a graph G = (V, E) is (k, r)-locallysparse if for each vertex  $v \in V(G)$ , the subgraph induced by its neighborhood contains at most kcliques of size r. For  $r \ge 3$  and  $\varepsilon \in [0, 1]$ , we show that a  $(\Delta^{\varepsilon r}, r)$ -locally-sparse graph G of maximum degree  $\Delta$  satisfies  $\alpha(G) = \Omega\left(\frac{n}{\gamma\Delta}\right)$  and  $\chi(G) = O(\gamma\Delta)$ , where  $\gamma \coloneqq \max\left\{\varepsilon, \frac{r\log\log\Delta}{\log\Delta}\right\}$ . As  $K_{r+1}$ -free graphs are (k, r)-locally-sparse for any k, we asymptotically recover classical results of Shearer and Johansson by setting  $\varepsilon = 0$ . We prove a stronger bound on the independence number in terms of the average degree, and establish a local version of the coloring result in the more general setting of correspondence coloring.

**Basic Notation.** All graphs considered here are finite, undirected, and simple. For  $n \in \mathbb{N}$ , we let  $[n] := \{1, \ldots, n\}$ . For a graph G, its vertex and edge sets are denoted V(G) and E(G) respectively. We say G is **complete** if every pair of vertices in G form an edge, and **edgeless** if G contains no edges.

For a vertex  $v \in V(G)$ ,  $N_G(v)$  denotes the neighbors of v and  $\deg_G(v) := |N_G(v)|$  denotes the degree of v. We let  $\Delta(G)$ ,  $\delta(G)$ , and d(G) denote the maximum, minimum, and average degrees of G, respectively. Let  $N_G[v] := N_G(v) \cup \{v\}$  denote the closed neighborhood of v. For a subset  $U \subseteq V(G)$ , the subgraph induced by U is denoted by G[U], and  $N_G(U)$  is the set of all vertices adjacent to a vertex in U, i.e.,  $N_G(U) = \bigcup_{u \in U} N_G(u)$ . We drop the subscript G when the context is clear.

A set  $I \subseteq V(G)$  is *independent* if G[I] is edgeless, and a set  $K \subseteq V(G)$  forms a *clique* if G[K] is complete. A proper coloring of G is a function  $\varphi : V(G) \to \mathbb{N}$  such that no edge  $uv \in E(G)$  satisfies  $\varphi(u) = \varphi(v)$ . The *independence number* of G (denoted  $\alpha(G)$ ) is the size of the largest independent set in G, the *clique number* (denoted  $\omega(G)$ ) is the size of the largest clique in G, and the *chromatic number* of G (denoted  $\chi(G)$ ) is the minimum value q such that G admits a proper coloring  $\varphi$  where the image of  $\varphi$  satisfies  $|im(\varphi)| = q$ .

We let  $\log^{(1)} x = \log x$  and  $\log^{(s)} x = \log(\log^{(s-1)} x)$  for  $s \ge 2$ .

### 1. INTRODUCTION

#### 1.1. Background and Results

It is well-known that a graph G satisfies  $\alpha(G) = \Omega(n/\Delta(G))$  and  $\chi(G) = O(\Delta(G))$ . These bounds are often referred to as the greedy bounds. For  $\alpha(G)$ , one can obtain a stronger bound of  $\Omega(n/d(G))$ which we refer to as the random greedy bound (see Algorithm 3.1). A natural question is the following: under what structural constraints can we obtain improved bounds on  $\alpha(\cdot)$  and  $\chi(\cdot)$ ? In this note, we consider the constraint of local sparsity. For  $k \in \mathbb{R}$ , we say a graph G is k-locally-sparse if for each  $v \in V(G)$ , the subgraph G[N(v)] contains at most  $\lfloor k \rfloor$  edges. Such a notion has been

School of Mathematics, Georgia Institute of Technology, Atlanta, GA, USA

*E-mail address*: abhishek.dhawan@math.gatech.edu.

This research was partially supported by the Georgia Tech ARC-ACO Fellowship, NSF grant DMS-2053333 (PI: Cheng Mao), and the NSF CAREER grant DMS-2239187 (PI: Anton Bernshteyn).

considered extensively (see for example [AKS99; FV06; Dav+20b; PH21]). We consider the following generalization of this notion introduced by Anderson, Kuchukova, and the author in [ADK24]:

**Definition 1.1.** For graphs F and G, a **copy** of F in G is a subgraph  $H \subseteq G$  which is isomorphic to F. Let F be a graph and let  $k \in \mathbb{R}$ . A graph G is (k, F)-**sparse** if G contains at most  $\lfloor k \rfloor$  copies of F (not necessarily vertex-disjoint). A graph G is (k, F)-**locally-sparse** if, for every  $v \in V(G)$ , the induced subgraph G[N(v)] is (k, F)-sparse.

When  $F = K_r$ , we simply call such graphs (k, r)-sparse or (k, r)-locally-sparse. Note that  $K_{r+1}$ -free graphs are (0, r)-locally-sparse. There has been a lot of work in this regime. Ajtai, Komlós, and Szemerédi first considered the independence number of  $K_3$ -free graphs. They showed that an *n*-vertex  $K_3$ -free graph G of average degree d has independence number  $\alpha(G) = \Omega(n \log d/d)$  [AKS80]. For  $r \ge 3$ , the same group along with Erdős showed that  $\alpha(G) = \Omega(n \log (\log d/r)/(rd))$  for  $K_{r+1}$ -free graphs G [Ajt+81]. Shearer improved the constant factor in the bound for  $K_3$ -free graphs [She83] and went on to prove the following celebrated result for  $r \ge 3$ :

**Theorem 1.2** ([She95]). Let  $r \in \mathbb{N}$  such that  $r \ge 3$ , and let n, d be sufficiently large. For any  $K_{r+1}$ -free graph G on n vertices such that d(G) = d, we have the following:

$$\alpha(G) = \Omega\left(\frac{n\,\log d}{r\,d\,\log^{(2)}d}\right)$$

Our first result is a generalization of the above. In particular, rather than considering  $K_{r+1}$ -free graphs, we consider the case where G contains "few" copies of  $K_{r+1}$ . Equivalently, on average each vertex is contained in "few" copies of  $K_{r+1}$ .

**Theorem 1.3.** There exists  $\rho > 0$  such that the following holds for n, d sufficiently large. Let  $\varepsilon, \gamma \in [0, 1], r \in \mathbb{N}$ , and  $k \in \mathbb{R}$  be such that

$$3 \leqslant r \leqslant \frac{\rho \log^{(2)} d}{\log^{(3)} d}, \quad k = \frac{n d^{\varepsilon r}}{r+1}, \quad and \quad \gamma = \max\left\{\varepsilon, \frac{r \log^{(2)} d}{\log d}\right\}.$$

For any *n*-vertex (k, r + 1)-sparse graph G of average degree d, we have

$$\alpha(G) = \Omega\left(\frac{n}{\gamma d}\right).$$

A few remarks are in order. First, we note that the value k above implies that on average a vertex  $v \in V(G)$  is contained in at most  $d(G)^{\varepsilon r}$  copies of  $K_{r+1}$  for such graphs. Additionally, for  $\varepsilon < r \log^{(2)} d / \log d$ , we obtain the same asymptotic bound as Theorem 1.2 with the weaker constraint of (k, r + 1)-sparsity as opposed to  $K_{r+1}$ -freeness. Finally, for  $\varepsilon \ge r \log^{(2)} d / \log d$  we improve upon the random greedy bound of  $\Omega(n/d)$  by a factor of  $1/\varepsilon$ .

Theorem 1.3 follows as a corollary to a stronger result on regular graphs. For a graph G, let I(G) denote the set of independent sets of G and let i(G) := |I(G)|. We define the **average** *independence number* (denoted  $\hat{\alpha}(G)$ ) and the *median independence number* (denoted  $\overline{\alpha}(G)$ ) as follows:

$$\hat{\alpha}(G) \coloneqq \frac{1}{i(G)} \sum_{I \in I(G)} |I|,$$
  
$$\overline{\alpha}(G) \coloneqq \max\left\{ \ell \in \mathbb{N} \ : \ |\{I \in I(G) \ : \ |I| \ge \ell\}| \ge \frac{i(G)}{2} \right\}.$$

The problem of computing lower bounds on the above parameters has been considered by Shearer [She95] and Molloy [Mol19] for  $K_{r+1}$ -free graphs, and by Davies, Jenssen, Perkins, and Roberts for  $K_3$ -free graphs [Dav+18]. The following result extends the one of Shearer's [She95, Theorem 1]:

**Theorem 1.4.** There exists  $\rho > 0$  such that the following holds for  $n, \Delta$  sufficiently large. Let  $\varepsilon, \gamma \in [0, 1], r \in \mathbb{N}$ , and  $k \in \mathbb{R}$  be such that

$$3 \leqslant r \leqslant \frac{\rho \log^{(2)} \Delta}{\log^{(3)} \Delta}, \quad k = \Delta^{\varepsilon r}, \quad and \quad \gamma = \max\left\{\varepsilon, \frac{r \log^{(2)} \Delta}{\log \Delta}\right\}.$$

For any *n*-vertex  $\Delta$ -regular (k, r)-locally-sparse graph G, we have

$$\hat{\alpha}(G) = \Omega\left(\frac{n}{\gamma\Delta}\right).$$

We note that  $k \in [1, \Delta^r]$ . In particular, as a vertex can contain at most  $\binom{\Delta}{r}$  cliques of size r in its neighborhood, this range covers all possible values of k. As a corollary, we obtain the following result, which implies Theorem 1.3 (see §2 for the proofs).

**Corollary 1.5.** There exists  $\rho > 0$  such that the following holds for  $n, \Delta$  sufficiently large. Let  $\varepsilon, \gamma \in [0, 1], r \in \mathbb{N}$ , and  $k \in \mathbb{R}$  be such that

$$3 \leqslant r \leqslant \frac{\rho \log^{(2)} \Delta}{\log^{(3)} \Delta}, \quad k = \Delta^{\varepsilon r}, \quad and \quad \gamma = \max\left\{\varepsilon, \frac{r \log^{(2)} \Delta}{\log \Delta}\right\}.$$

For any n-vertex (k, r)-locally-sparse graph G of maximum degree  $\Delta$ , we have

$$\alpha(G) = \Omega\left(\frac{n}{\gamma\Delta}\right).$$

When considering colorings, the best known asymptotic bound on the chromatic number for  $K_{r+1}$ free graphs is due to Johansson. He proved the following result employing a complex randomized
coloring procedure:

**Theorem 1.6** ([Joh96a; Joh96b]). For  $r \ge 2$ , let G be a  $K_{r+1}$ -free graph of maximum degree  $\Delta$  sufficiently large. Then,

$$\chi(G) = \begin{cases} O\left(\frac{r\Delta\log^{(2)}\Delta}{\log\Delta}\right) & r \ge 3, \\ O\left(\frac{\Delta}{\log\Delta}\right) & r = 2. \end{cases}$$

We note that the result holds in the more general setting of *list coloring* described in §1.2. Molloy provided a simpler proof of Theorem 1.6 by employing the so-called *entropy compression method* [Mol19]. Bernshteyn simplified the proof further by employing the *lopsided local lemma* and extended the result to *correspondence coloring* (also described in §1.2) [Ber19]. Both Molloy's and Bernshteyn's proofs for  $r \ge 3$  rely on the result of Theorem 1.2. We are similarly able to achieve a bound on the chromatic number of locally sparse graphs as follows:

**Theorem 1.7.** There exists  $\rho > 0$  such that the following holds for  $\Delta$  sufficiently large. Let  $\varepsilon, \gamma \in [0, 1], r \in \mathbb{N}$ , and  $k \in \mathbb{R}$  be such that

$$3 \leqslant r \leqslant \frac{\rho \log^{(2)} \Delta}{\log^{(3)} \Delta}, \quad k = \Delta^{\varepsilon r}, \quad and \quad \gamma = \max\left\{\varepsilon, \frac{r \log^{(2)} \Delta}{\log \Delta}\right\}.$$

For any (k, r)-locally-sparse graph G of maximum degree  $\Delta$ , we have  $\chi(G) = O(\gamma \Delta)$ .

For  $\varepsilon < r \log^{(2)} \Delta / \log \Delta$ , we obtain the same asymptotic bound as Theorem 1.6 (for  $r \ge 3$ ) with the weaker assumption of local sparsity. For larger  $\varepsilon$ , our results improve upon the greedy bound of  $O(\Delta)$  by a factor of  $\varepsilon$ . Additionally, our results hold for list and correspondence colorings as well (see Corollary 1.12). Note that rather than having a strict bound on the clique number of G (as is the case in Theorem 1.6), we consider the case that vertices are not contained in "too many" small cliques. To see the versatility of our results, consider a (k, r)-locally-sparse graph G of maximum degree  $\Delta$  sufficiently large. It is not too difficult to see that  $\omega(G) = O(rk^{1/r})$  by considering the neighborhood of a vertex contained in a maximum clique. In particular, for  $k = \Delta^{\varepsilon r}$ , Theorem 1.6 provides the bound  $\chi(G) = O(r\Delta^{1+\varepsilon}\log^{(2)}\Delta/\log\Delta)$ , which is significantly larger than that provided by Theorem 1.7. A result of Bonamy, Kelly, Nelson, and Postle [Bon+22, Theorem 1.6] provides a stronger bound of  $O\left(\Delta\sqrt{\max\{\varepsilon,\log r/\log\Delta\}}\right)$ , which is still weaker than our result.

#### 1.2. List and Correspondence Coloring

Introduced independently by Vizing [Viz76] and Erdős, Rubin, and Taylor [ERT79], *list coloring* is a generalization of graph coloring in which each vertex is assigned a color from its own predetermined list of colors. Formally,  $L: V(G) \to 2^{\mathbb{N}}$  is a *list assignment* for G, and an *L*-coloring of G is a proper coloring of G such that each vertex  $v \in V(G)$  receives a color from its list L(v). When  $|L(v)| \ge q$  for each  $v \in V(G)$ , where  $q \in \mathbb{N}$ , we say L is *q*-fold. The *list-chromatic number* of G, denoted  $\chi_{\ell}(G)$ , is the smallest q such that G has an *L*-coloring for every q-fold list assignment L for G.

Correspondence coloring (also known as DP-coloring) is a generalization of list coloring introduced by Dvořák and Postle [DP18]. Just as in list coloring, each vertex is assigned a list of colors, L(v); in contrast to list coloring, though, the identifications between the colors in the lists are allowed to vary from edge to edge. That is, each edge  $uv \in E(G)$  is assigned a matching  $M_{uv}$  (not necessarily perfect and possibly empty) from L(u) to L(v). A proper correspondence coloring is a mapping  $\varphi : V(G) \to \mathbb{N}$  satisfying  $\varphi(v) \in L(v)$  for each  $v \in V(G)$  and  $\varphi(u)\varphi(v) \notin M_{uv}$  for each  $uv \in E(G)$ . Formally, we describe correspondence colorings in terms of an auxiliary graph known as a correspondence cover of G. The definition below appears in earlier works of the author along with Anderson and Bernshteyn [ABD22; ABD23].

**Definition 1.8** (Correspondence Cover). A correspondence cover (also known as a **DP**-cover) of a graph G is a pair  $\mathcal{H} = (L, H)$ , where H is a graph and  $L : V(G) \to 2^{V(H)}$  such that:

- (DP1) The set  $\{L(v) : v \in V(G)\}$  forms a partition of V(H).
- (DP2) For each  $v \in V(G)$ , L(v) is an independent set in H.
- (DP3) For each  $u, v \in V(G)$ , the edges of the induced subgraph  $H[L(u) \cup L(v)]$  form a matching; this matching is empty whenever  $uv \notin E(G)$ .

We call the vertices of H colors. If two colors  $c, c' \in V(H)$  are adjacent in H, we say that they **correspond** to each other. An  $\mathcal{H}$ -coloring is a mapping  $\varphi \colon V(G) \to V(H)$  such that  $\varphi(v) \in L(v)$  for all  $v \in V(G)$ . An  $\mathcal{H}$ -coloring  $\varphi$  is **proper** if the image of  $\varphi$  is an independent set in H.

A correspondence cover  $\mathcal{H} = (L, H)$  is q-fold if  $|L(v)| \ge q$  for all  $v \in V(G)$ . The correspondence chromatic number of G, denoted by  $\chi_c(G)$ , is the smallest q such that G admits a proper  $\mathcal{H}$ coloring for every q-fold correspondence cover  $\mathcal{H}$ . As correspondence coloring generalizes list coloring, which in turn generalizes ordinary coloring, we have

$$\chi(G) \leqslant \chi_{\ell}(G) \leqslant \chi_{c}(G). \tag{1.9}$$

Our main coloring result is on *local correspondence colorings* of locally sparse graphs. A correspondence cover is *local* if for each vertex  $v \in V(G)$ , the size of its list L(v) is a function of the local structure of G with respect to v. This notion has gathered much interest in recent years in both vertex coloring [Dav+20a; Dav+20c; Bon+22] and edge coloring [Bon+20; Dha23]. Before we state our result, we introduce the following definition, which is a local version of Definition 1.1 for  $F = K_r$ .

**Definition 1.10.** Let G be a graph and let  $\mathbf{k} : V(G) \to \mathbb{R}$  and  $\mathbf{r} : V(G) \to \mathbb{N}$ . A graph G is  $(\mathbf{k}, \mathbf{r})$ -locally-sparse if, for every  $v \in V(G)$ , the induced subgraph G[N(v)] is  $(\mathbf{k}(v), \mathbf{r}(v))$ -sparse.

We are now ready to state our result.

**Theorem 1.11.** There exist  $C, \rho > 0$  such that the following holds for  $\Delta$  sufficiently large. Let G be a graph of maximum degree  $\Delta$  and let  $\mathbf{k} : V(G) \to \mathbb{R}$ ,  $\mathbf{r} : V(G) \to \mathbb{N}$ , and  $\boldsymbol{\varepsilon} : V(G) \to [0,1]$  be such that G is  $(\mathbf{k}, \mathbf{r})$ -locally-sparse and the following hold for each  $v \in V(G)$ :

- (S1)  $\deg(v) \ge \log^2 \Delta$ , and
- (S2)  $3 \leq \mathbf{r}(v) \leq \rho \log^{(2)} \deg(v) / \log^{(3)} \deg(v)$  and  $\mathbf{k}(v) = \deg(v)^{\varepsilon(v)\mathbf{r}(v)}$ .

For any correspondence cover  $\mathcal{H} = (L, H)$  of G satisfying

$$|L(v)| \ge C \operatorname{deg}(v) \max\left\{ \boldsymbol{\varepsilon}(v), \frac{\mathbf{r}(v) \log^{(2)} \operatorname{deg}(v)}{\log \operatorname{deg}(v)} \right\},\$$

G admits a proper  $\mathcal{H}$ -coloring.

We note that a local correspondence coloring version of Theorem 1.6 for  $r \ge 3$  appeared in work of Bonamy, Kelly, Nelson, and Postle [Bon+22, Corollary 1.9]. As a corollary, we obtain a correspondence coloring version of Theorem 1.7 (see §2 for the proof), which implies Theorem 1.7 as a result of (1.9).

**Corollary 1.12.** There exists  $\rho > 0$  such that the following holds for  $\Delta$  sufficiently large. Let  $\varepsilon, \gamma \in [0, 1], r \in \mathbb{N}$ , and  $k \in \mathbb{R}$  be such that

$$3 \leqslant r \leqslant \frac{\rho \log^{(2)} \Delta}{\log^{(3)} \Delta}, \quad k = \Delta^{\varepsilon r}, \quad and \quad \gamma = \max\left\{\varepsilon, \frac{r \log^{(2)} \Delta}{\log \Delta}\right\}.$$

For any (k, r)-locally-sparse graph G of maximum degree  $\Delta$ , we have  $\chi_c(G) = O(\gamma \Delta)$ .

## 1.3. Proof Overview

Our proof closely follows the strategy employed by Shearer in [She95] and Molloy in [Mol19]. While Molloy's result focused on improving the constant factor in Theorem 1.6, his approach provided a simpler proof of Theorem 1.2 as well. The crux of the proof lies in the following lemma:

**Lemma 1.13** ([Mol19, Lemma 13]). For any  $r \ge 3$ , let G be a  $K_r$ -free graph on n vertices. Then,

$$n \ge \log_2(i(G)) \ge n^{1/(r-1)}.$$

The upper bound is trivial as  $i(G) \leq 2^n$  where equality holds if and only if G is edgeless. For the lower bound, it is enough to show that G contains an independent set of size at least  $n^{1/(r-1)}$ (which is precisely Molloy's strategy). As every (k, r)-sparse graph G contains a  $K_r$ -free subgraph H such that  $|E(H)| \geq |E(G)| - k$ , we may conclude the following:

**Lemma 1.14.** For any  $r \ge 3$ , let G be a (k, r)-sparse graph on n vertices. Then,

$$n \ge \log_2(i(G)) \ge n^{1/(r-1)} - k.$$

However, this result is not strong enough to prove Theorems 1.3 and 1.7. In fact, with this lemma, we may prove the following weaker result:

**Theorem 1.15.** There exists  $\rho > 0$  such that the following holds for  $n, \Delta$  sufficiently large. Let  $\varepsilon, \gamma \in [0, 1], r \in \mathbb{N}$ , and  $k \in \mathbb{R}$  be such that

$$3 \,\leqslant\, r \leqslant \frac{\rho\,\log^{(2)}\Delta}{\log^{(3)}\Delta}, \quad k \,=\, \Delta^{\varepsilon\,r}, \quad and \quad \gamma = \max\left\{\varepsilon\,r^2,\, \frac{r\log^{(2)}\Delta}{\log\Delta}\right\}.$$

For any n-vertex (k, r)-locally-sparse graph G of maximum degree  $\Delta$ , we have

$$\alpha(G) = \Omega\left(\frac{n}{\gamma\Delta}\right) \quad and \quad \chi_c(G) = O\left(\gamma\Delta\right)$$

We do not include a proof of Theorem 1.15, however, it can be inferred by our arguments. Note that for r = O(1), the above bounds match those of Corollary 1.5 and Theorem 1.7. However, for larger r, the results of Corollary 1.5 and Theorem 1.7 are much stronger.

The key part of our proof, therefore, is a stronger version of Lemma 1.14 (see Lemma 3.2). With this in hand, we prove lower bounds on  $\hat{\alpha}(\cdot)$  and  $\overline{\alpha}(\cdot)$  for (k, r)-sparse graphs by employing a similar strategy to Shearer and Molloy. To prove Theorem 1.4, we show that the expected size of a uniformly random independent set in I(G) is bounded from below by the desired quantity. For Theorem 1.11, we apply a result of Bonamy, Kelly, Nelson, and Postle in conjunction with our bound on the median independence number of (k, r)-sparse graphs.

#### 1.4. Concluding Remarks

In this paper, we generalize classical results of Shearer and Johansson on  $K_{r+1}$ -free graphs to (k, r)-locally-sparse-graphs. Furthermore, we prove local versions of our coloring result in the setting of correspondence coloring, which extends work of Bonamy, Kelly, Nelson, and Postle. In the entire paper, we do not attempt to optimize the constant factors involved and leave it as an open problem to do so. While not explicitly stated in their papers, Theorems 1.2 and 1.6 hold for  $r = O(\log^{(2)} d)$  and  $r = O(\log^{(2)} \Delta)$ , respectively. For Theorem 1.6, this range was extended to  $r = O(\log \Delta)$  by Bonamy, Kelly, Nelson, and Postle [Bon+22]. Our proof fails for larger r and we leave it as an open problem to extend this range.

We conclude this section with a discussion for arbitrary graphs F. A simple counting argument shows the following fact:

**Fact 1.16.** Let G be a graph,  $k \in \mathbb{R}$  and let F be an arbitrary graph on r vertices. If G is (k, F)-locally-sparse, then G is  $(\tilde{k}, r)$ -locally-sparse, where

$$\tilde{k} = \frac{|k| |\operatorname{Aut}(F)|}{r!}.$$

Here, Aut(F) is the set of automorphisms of F.

In particular, our results in this section imply the following:

**Theorem 1.17.** There exists  $\rho > 0$  such that the following holds for  $n, \Delta$  sufficiently large. Let  $\varepsilon, \gamma \in [0, 1], r \in \mathbb{N}$ , and  $k \in \mathbb{R}$  be such that

$$3 \leqslant r \leqslant \frac{\rho \log^{(2)} \Delta}{\log^{(3)} \Delta}, \quad k = \Delta^{\varepsilon r}, \quad and \quad \gamma = \max\left\{\varepsilon, \frac{r \log^{(2)} \Delta}{\log \Delta}\right\}.$$

Let F be an arbitrary graph on r vertices. Then for any n-vertex (k, F)-locally-sparse graph G of maximum degree  $\Delta$ , we have

$$\alpha(G) = \Omega\left(\frac{n}{\gamma\Delta}\right) \quad and \quad \chi_c(G) = O\left(\gamma\Delta\right),$$

where the constant factors in the  $\Omega(\cdot)$  and  $O(\cdot)$  may depend on F.

Note that for any graph F on r vertices, we have  $\binom{\Delta}{r}r!/|\operatorname{Aut}(F)| \leq \Delta^r$ . In particular, the above result covers all possible values of k. Anderson, Kuchukova, and the author proved a better asymptotic bound on  $\chi_c(G)$  when F is bipartite and k is not too large.

**Theorem 1.18** ([ADK24]). Let F be a bipartite graph and let  $\Delta$  be sufficiently large. For  $k \in \mathbb{R}$  satisfying  $1/2 \leq k \leq \Delta^{|V(F)|/10}$ , the following holds. For any (k, F)-locally-sparse graph G of maximum degree  $\Delta$ , we have

$$\chi_c(G) \leqslant \frac{8\,\Delta}{\log\left(\Delta\,k^{-1/|V(F)|}\right)}$$

This led the authors to make the following conjecture which Theorem 1.17 constitutes some progress toward:

**Conjecture 1.19** ([ADK24, Conjecture 1.20]). For every graph F, the following holds for  $\Delta \in \mathbb{N}$  sufficiently large. Let

$$1/2 \leqslant k < \Delta^{|V(F)|}$$

and let G be a (k, F)-locally-sparse graph of maximum degree  $\Delta$ . Then,

$$\chi_c(G) = O\left(\frac{\Delta}{\log\left(\Delta k^{-1/|V(F)|}\right)}\right),$$

where the constant factor in the  $O(\cdot)$  may depend on F.

The rest of the paper is structured as follows: in \$2, we prove some of the corollaries mentioned in this section; in \$3, we prove Theorem 1.4; and in \$4, we prove Theorem 1.11.

### 2. PROOF OF COROLLARIES

The following proposition will be key in proving the corollaries in this section.

**Proposition 2.1.** Let G be a graph of maximum degree  $\Delta$  and let  $\mathbf{k} : V(G) \to \mathbb{R}$  and  $\mathbf{r} : V(G) \to \mathbb{N}$  be such that G is  $(\mathbf{k}, \mathbf{r})$ -locally-sparse. For any  $\delta \leq \Delta$ , there is a graph G' and functions  $\tilde{\mathbf{k}} : V(G') \to \mathbb{R}$  and  $\tilde{\mathbf{r}} : V(G') \to \mathbb{N}$  such that G' is  $(\tilde{\mathbf{k}}, \tilde{\mathbf{r}})$ -locally-sparse and the following hold for  $j := \max\{\delta - \delta(G), 0\}$ :

- (I1)  $\delta(G') \ge \delta$ ,
- (I2)  $\Delta(G') = \Delta$ ,
- (I3)  $|V(G')| = |V(G)| 2^j$ , and
- (I4) for  $\ell := 2^j$ , there exist homomorphisms  $\varphi_1, \ldots, \varphi_\ell : G \to G'$  such that
  - for each  $v \in V(G)$  and  $\ell' \in [\ell]$ , we have  $\mathbf{k}(v) = \tilde{\mathbf{k}}(\varphi_{\ell'}(v))$  and  $\mathbf{r}(v) = \tilde{\mathbf{r}}(\varphi_{\ell'}(v))$ , and
  - for  $\ell_1, \ell_2 \in [\ell]$  such that  $\ell_1 \neq \ell_2$ , the images  $\operatorname{im}(\varphi_{\ell_1})$  and  $\operatorname{im}(\varphi_{\ell_2})$  are vertex-disjoint.

*Proof.* If  $\delta(G) \ge \delta$ , the claim holds for G' = G. If not, we will define G' iteratively as follows:

- (1) Let  $G_0 := G$ .
- (2) For  $i \ge 0$ , define  $G_{i+1}$  by taking two vertex-disjoint copies of  $G_i$ , say  $G_i^{(1)}$  and  $G_i^{(2)}$ . For each  $v \in V(G_i)$  such that  $\deg_{G_i}(v) < \delta$ , draw an edge between the corresponding vertices in  $G_i^{(1)}$  and  $G_i^{(2)}$ .
- (3) Let  $G' = G_j$ , where j is as defined in the statement of the proposition.

Note that (I3) holds by construction. We will first show that the graphs  $G_i$  satisfy  $\Delta(G_i) = \Delta$  for all  $0 \leq i \leq j$ . For i = 0, this holds trivially. Suppose it holds for some  $0 \leq i < j$ . Let  $G_i^{(1)}$  and  $G_i^{(2)}$ be the copies of  $G_i$  in  $G_{i+1}$ . For  $s \in \{1, 2\}$  and any vertex  $v \in V(G_i^{(s)})$ ,  $\deg_{G_{i+1}}(v) \neq \deg_{G_i^{(s)}}(v)$  if and only if  $\deg_{G_i^{(s)}}(v) < \delta \leq \Delta$ . Furthermore, by construction, we have the following:

$$\deg_{G_i^{(s)}}(v) < \delta \implies \deg_{G_{i+1}}(v) = \deg_{G_i^{(s)}}(v) + 1 \le \delta.$$

$$(2.2)$$

As any vertex of degree  $\Delta \ge \delta$  in  $G_i$  has the same degree in  $G_{i+1}$ , it follows that  $\Delta(G_{i+1}) = \Delta$ . Setting i = j completes the proof of (I2).

We will define functions  $\mathbf{k}_i : V(G_i) \to \mathbb{R}$  and  $\mathbf{r}_i : V(G_i) \to \mathbb{N}$  such that  $G_i$  is  $(\mathbf{k}_i, \mathbf{r}_i)$ -locallysparse to assist with the proof of (I4). Let  $\mathbf{k}_0 := \mathbf{k}$  and  $\mathbf{r}_0 := \mathbf{r}$ . Suppose we have defined  $\mathbf{k}_i$ ,  $\mathbf{r}_i$  for some  $i \ge 0$ . Let  $G_i^{(1)}$  and  $G_i^{(2)}$  be the copies of  $G_i$  in  $G_{i+1}$ . For  $s \in \{1, 2\}$  and any vertex  $v \in V(G_i^{(s)})$ , the graph  $G_{i+1}[N_{G_{i+1}}(v)]$  is either isomorphic to  $G_i^{(s)}[N_{G_i^{(s)}}(v)]$  or contains one additional isolated vertex. In particular, for  $\mathbf{k}_{i+1}(v) = \mathbf{k}_i(v)$  and  $\mathbf{r}_{i+1}(v) = \mathbf{r}_i(v)$ , the graph  $G_{i+1}$  is  $(\mathbf{k}_{i+1}, \mathbf{r}_{i+1})$ -locally-sparse.

Let us now consider (I4). We will prove the following more general statement which implies (I4) for  $\tilde{\mathbf{k}} := \mathbf{k}_j$  and  $\tilde{\mathbf{r}} := \mathbf{r}_j$ .

**Claim 2.2.1.** For each  $0 \leq i \leq j$ , there exist  $\ell_i := 2^i$  homomorphisms  $\varphi_1, \ldots, \varphi_{\ell_i} : G \to G_i$  such that

- (1) for each  $v \in V(G)$  and  $\ell' \in [\ell_i]$ , we have  $\mathbf{k}(v) = \tilde{\mathbf{k}}_i(\varphi_{\ell'}(v))$  and  $\mathbf{r}(v) = \tilde{\mathbf{r}}_i(\varphi_{\ell'}(v))$ , and
- (2) for  $\ell_1, \ell_2 \in [\ell_i]$  such that  $\ell_1 \neq \ell_2$ , the images  $\operatorname{im}(\varphi_{\ell_1})$  and  $\operatorname{im}(\varphi_{\ell_2})$  are vertex-disjoint.

*Proof.* We will prove this by induction on *i*. For i = 0, the claim is trivial. Suppose the claim holds for some  $0 \leq i < j$ . Let  $G_i^{(1)}$  and  $G_i^{(2)}$  be the copies of  $G_i$  in  $G_{i+1}$ , and for  $s \in \{1, 2\}$  let  $\varphi_1^{(s)}, \ldots, \varphi_{\ell_i}^{(s)} : G \to G_i^{(s)}$  be the homomorphisms guaranteed by the induction hypothesis. By definition of  $\mathbf{k}_{i+1}$  and  $\mathbf{r}_{i+1}$  and since the graphs  $G_i^{(1)}$  and  $G_i^{(2)}$  are vertex-disjoint, it follows that the homomorphisms

$$\{\varphi_{\ell}^{(s)} : s \in \{1, 2\}, 1 \leq \ell \leq \ell_i\},\$$

 $\neg$ 

satisfy the conditions of the claim. As  $\ell_{i+1} = 2\ell_i$ , this completes the proof.

Finally, note that by (2.2), we may conclude the following for  $i \ge 0$ :

$$\delta(G_i) < \delta \implies \delta(G_{i+1}) = \delta(G_i) + 1.$$

In particular, (I1) holds for G' by definition of j, completing the proof.

With this proposition in hand, we are ready to prove the corollaries stated in §1.

Proof of Corollary 1.5. For each  $v \in V(G)$ , let  $\mathbf{k}(v) \coloneqq k$  and  $\mathbf{r}(v) \coloneqq r$ . If G is not regular, we form G' by applying Proposition 2.1 with  $\delta = \Delta$ . We may now apply Theorem 1.4 to G'. As a result of (I3) and (I4), if an independent set in G' contains an

$$s \coloneqq \Omega\left(\frac{1}{\gamma\Delta}\right)$$

fraction of the vertices of G, it must contain an s fraction of vertices from some copy of G in G', completing the proof.

Let us now show how Theorem 1.3 follows from Corollary 1.5.

*Proof of Theorem 1.3.* Consider the following sets of vertices:

$$V_1 := \{ v \in V(G) : \deg(v) \leq 3d \},$$
  

$$V_2 := \{ v \in V(G) : G[N(v)] \text{ is } (3d^{\varepsilon r}, r) \text{-sparse} \},$$
  

$$V_3 := V_1 \cap V_2.$$

Let us show that  $|V_3| \ge n/3$ . First, we note the following:

$$n d = \sum_{v \in V(G)} \deg_G(v) \ge 3d(n - |V_1|) \implies |V_1| \ge 2n/3.$$

For each  $v \in V(G)$ , let  $n_v$  denote the number of copies of  $K_r$  in  $G[N_G(v)]$ . By definition of (k, r+1)-sparsity, we have the following

$$(r+1) k = \sum_{v \in V(G)} n_v \ge 3d^{\varepsilon r} (n - |V_2|) \implies |V_2| \ge 2n/3.$$

With the above in hand, we conclude

$$|V_3| = |V_1| + |V_2| - |V_1 \cup V_2| \ge 2n/3 + 2n/3 - n = n/3,$$

as claimed.

Let  $G' \subseteq G$  be the subgraph induced by the vertices in  $V_3$ . Since an independent set in G' is also independent in G, it is enough to show that G' contains an independent set of the desired size. Suppose  $\Delta(G') \leq \gamma d$ . Then we have the following as a result of the greedy bound:

$$\alpha(G') = \Omega\left(\frac{|V_3|}{\Delta(G')}\right) = \Omega\left(\frac{n}{\gamma d}\right),$$

as desired. If not, by definition of  $V_1$ , we have  $\gamma d < \Delta(G') \leq 3d$ . Furthermore, by definition of  $V_2$ , G' is  $(\Delta(G')^{\tilde{\varepsilon}r}, r)$ -locally-sparse, where

$$\tilde{\varepsilon} := \min\left\{1, \frac{\varepsilon \log d + \log 3}{\log\left(\Delta(G')\right)}\right\}.$$

As a result of the range of  $\Delta(G')$ , we may conclude that  $\tilde{\varepsilon} = \Theta(\varepsilon)$  for d large enough. We may now apply Corollary 1.5 to G' to complete the proof.

Finally, let us prove Corollary 1.12 under the assumption that Theorem 1.11 is true.

Proof of Corollary 1.12. For each  $v \in V(G)$ , define

$$\mathbf{r}(v) \coloneqq r$$
, and  $\mathbf{k}(v) \coloneqq k$ .

Note that G is  $(\mathbf{k}, \mathbf{r})$ -locally-sparse. Form G' by applying Proposition 2.1 with  $\delta = \Delta$ . Let  $\varepsilon(v) = \varepsilon$  for each  $v \in V(G')$ . Note that G' satisfies the conditions of Theorem 1.11. In particular, for any correspondence cover  $\mathcal{H} = (L, H)$  of G' satisfying

$$|L(v)| \ge C \deg_{G'}(v) \max\left\{ \boldsymbol{\varepsilon}(v), \ \frac{\mathbf{r}(v)\log^{(2)}\deg_{G'}(v)}{\log\deg_{G'}(v)} \right\},\$$

G' admits a proper  $\mathcal{H}$ -coloring. As G' is  $\Delta$ -regular,  $\varepsilon(v) = \varepsilon$  for all  $v \in V(G')$ , and G is isomorphic to a subgraph of G' by (I4), the claim follows.

## 3. PROOF OF THEOREM 1.4

As mentioned in §1, it is a well-known fact that an *n*-vertex graph G contains an independent set of size at least  $n/(1 + d(G)) = \Omega(n/d(G))$ . This lower bound is achieved by what some refer to as the Random Greedy Algorithm. For completeness, we describe the algorithm here.

Algorithm 3.1 The Random Greedy Algorithm

Input: A graph G. Output: An independent set I in G. 1: Let  $x_v$  be i.i.d.  $\mathcal{U}(0, 1)$  for each  $v \in V(G)$ . 2:  $I \leftarrow \emptyset$ . 3: for  $v \in V(G)$  do 4: if  $x_v > \max_{u \in N_G(v)} x_u$  then 5:  $I \leftarrow I \cup \{v\}$ 6: end if 7: end for 8: return I

The following lemma bounds the expected size of the independent set I output by Algorithm 3.1. The proof follows by a standard application of Jensen's inequality (see for example [BD23, Lemma 8.2]).

**Lemma 3.1.** Consider the output I of Algorithm 3.1 on input G. We have

$$\mathbb{E}[|I|] \ge \frac{|V(G)|}{1+d(G)}.$$

The next lemma provides a general lower bound on the independence number of (k, r)-sparse graphs and will be key in proving the main results of this paper as mentioned in §1.3. We note that our main results only consider the case that  $r \ge 3$ , however, including the case r = 2 below provides for a simpler proof.

**Lemma 3.2.** Let  $r, n \in \mathbb{N}$  and  $k \in \mathbb{R}$  such that  $r \ge 2$ ,  $n \ge r^{2r}$ , and  $1 \le k \le {n \choose r}$ . Let G be an *n*-vertex (k, r)-sparse graph. Then,

$$n \, \geqslant \, \log_2\left(i(G)\right) \, \geqslant \, \frac{1}{r} \left(\frac{n}{k^{1/r}}\right)^{1/(r-1)}$$

*Proof.* The upper bound follows as  $i(G) \leq 2^n$  trivially. For the lower bound, it is enough to show that G contains an independent set of size at least  $\frac{1}{r} \left(\frac{n}{k^{1/r}}\right)^{1/(r-1)}$ . We will prove this by induction on r.

As a base case, let r = 2. We remark that it is enough to have  $n \ge 3$  here, however, by the assumptions of the lemma, we have  $n \ge 16$ . By Lemma 3.1, we have an independent set of size at least

$$\frac{n}{1+d(G)}$$

As G is (k, 2)-sparse, we have  $|E(G)| \leq k$ . It follows that

$$\frac{n}{1+d(G)} \ge \frac{n}{1+2k/n}$$

It is now enough to show that the above is at least  $n/(2\sqrt{k})$ . To this end, we consider the following function:

$$f(k) = 2\sqrt{k} - 2k/n - 1.$$

Note that  $f(k) \ge 0$  for  $1 \le k \le {n \choose 2}$  implies the desired result. Consider the following for  $n \ge 16$ :

$$f(1) = 2 - 2/n - 1 > 0,$$
  

$$f(n^2/4) = n - n/2 - 1 > 0,$$
  

$$f(n^2/2) = (\sqrt{2} - 1)n - 1 > 0,$$

$$f'(k) = k^{-1/2} - 2/n \ge 0 \iff k \le \frac{n^2}{4}.$$

It follows that f is increasing for  $1 \le k \le n^2/4$  and decreasing for  $n^2/4 < k \le n^2/2$ . As  $f(1), f(n^2/4), f(n^2/2) > 0$ , we conclude f(k) > 0 for  $1 \le k \le \binom{n}{2}$ , as desired.

Now let us consider  $r \ge 3$ . We define the following parameters:

$$\begin{aligned} \alpha_r &\coloneqq \left(\frac{r-1}{r}\right)^{r-2-\frac{1}{r-1}}, & \beta_r &\coloneqq r-1, \\ X &\coloneqq \alpha_r \, n^{\frac{r-2}{r-1}} \, k^{\frac{1}{r(r-1)}}, & B &\coloneqq \{v \in V(G) \, : \, \deg_G(v) \ge X\}. \end{aligned}$$

The following bound on X follows as  $r \ge 3$ ,  $n \ge r^{2r}$ , and  $k \ge 1$ :

$$\begin{aligned} X &= \left(\frac{r-1}{r}\right)^{r-2-\frac{1}{r-1}} n^{\frac{r-2}{r-1}} k^{\frac{1}{r(r-1)}} \\ &\geqslant \left(\frac{r-1}{r}\right)^{r-2-\frac{1}{r-1}} r^{\frac{2r(r-2)}{r-1}} \\ &= (r-1)^{2(r-1)-r-\frac{1}{r-1}} r^{\frac{2r(r-2)+1}{r-1}-(r-2)} \\ &= (r-1)^{2(r-1)} (r-1)^{-r-\frac{1}{r-1}} r^{\frac{(r+1)(r-2)+1}{r-1}} \\ &= (r-1)^{2(r-1)} (r-1)^{-r-\frac{1}{r-1}} r^{\frac{r^2-r-1}{r-1}} \\ &= (r-1)^{2(r-1)} \left(\frac{r}{r-1}\right)^r \frac{1}{(r(r-1))^{1/(r-1)}}. \end{aligned}$$

For r = 3, it can be verified that

$$\left(\frac{r}{r-1}\right)^r \frac{1}{(r(r-1))^{1/(r-1)}} \ge 1.$$

For  $r \ge 4$ , we have

$$\log(r(r-1)) \leq 2\log r \leq r-1.$$

It follows that

$$(r(r-1))^{1/(r-1)} \le e.$$

Furthermore, we have

$$(1-1/r)^r \leq \frac{1}{e} \implies \frac{1}{(1-1/r)^r} \geq e,$$

In particular, we may conclude that

$$X \ge (r-1)^{2(r-1)}.$$
(3.3)

We will consider two cases based on the size of B.

(Case1)  $|B| \ge \beta_r k^{1/r}$ . For each  $v \in B$ , let  $n_v$  denote the number of copies of  $K_r$  in G containing v. By definition of (k, r)-sparsity, we have:

$$\sum_{v \in B} n_v \leqslant rk,$$

implying there is some  $v \in B$  such that  $n_v \leq rk/|B| \leq rk^{1-1/r}/\beta_r$ . In particular, the graph H := G[N(v)] is  $(\tilde{k}, r-1)$ -sparse, where

$$\tilde{k} \coloneqq \min\left\{\frac{r\,k^{1-1/r}}{\beta_r}, \, \binom{|V(H)|}{r-1}\right\}.$$

Note the following:

$$\frac{r\,k^{1-1/r}}{\beta_r} \ge \frac{r}{(r-1)} > 1.$$

Furthermore,  $X \ge (r-1)^{2(r-1)}$  by (3.3) and so, we have

$$\binom{|V(H)|}{r-1} \ge \binom{X}{r-1} > 1.$$

Therefore, we conclude  $1 \leq \tilde{k} \leq {\binom{|V(H)|}{r-1}}$  and  $|V(H)| \geq (r-1)^{2(r-1)}$ . By the induction hypothesis, there is an independent set in H of size at least

$$\begin{aligned} \frac{1}{r-1} \left(\frac{|V(H)|}{\tilde{k}^{1/(r-1)}}\right)^{1/(r-2)} &\ge \frac{1}{r-1} \left(\frac{X}{\left(r \, k^{1-1/r}/\beta_r\right)^{1/(r-1)}}\right)^{1/(r-2)} \\ &= \frac{1}{r-1} \left(\frac{n}{k^{1/r}}\right)^{1/(r-1)} \left(\frac{\alpha_r^{r-1}\beta_r}{r}\right)^{1/((r-1)(r-2))} \\ &= \frac{1}{r} \left(\frac{n}{k^{1/r}}\right)^{1/(r-1)}, \end{aligned}$$

as desired. (We note that as  $r \ge 3$ , 1/(r-2) is well defined.) As this set is also independent in G, this concludes (**Case1**).

(Case2)  $|B| < \beta_r k^{1/r}$ . First, we note the following:

$$\frac{|B|}{X} < \frac{\beta_r \, k^{1/r}}{X} = \frac{\beta_r}{\alpha_r} \left(\frac{k^{1/r}}{n}\right)^{\frac{r-2}{r-1}} \leqslant \frac{\beta_r}{\alpha_r},$$

where we use the fact that  $k \leq \binom{n}{r} \leq n^r$ . As a result, we have

$$d(G) \leq \frac{1}{n} \left( |B|n + (n - |B|)X \right) \leq |B| + X$$
$$\leq \frac{X}{\alpha_r} \left(\beta_r + \alpha_r\right)$$
$$= \frac{rX}{\alpha_r} - \frac{X}{\alpha_r} \left(1 - \alpha_r\right).$$

Let us consider the term on the right. We have

$$\frac{X}{\alpha_r} (1 - \alpha_r) = n^{\frac{r-2}{r-1}} k^{\frac{1}{r(r-1)}} \left( 1 - \left(1 - \frac{1}{r}\right)^{r-2-\frac{1}{r-1}} \right)$$
$$\ge n^{1/2} \left( 1 - \exp\left(-\frac{1}{r}\left(r-2 - \frac{1}{r-1}\right)\right) \right)$$
$$= n^{1/2} \left( 1 - \exp\left(-1 + \frac{2}{r} + \frac{1}{r(r-1)}\right) \right)$$
$$\ge n^{1/2} \left( 1 - \exp(-1/6) \right) > 1,$$

where we use the fact that  $n \ge r^{2r}$  and  $r \ge 3$ . In particular, we have

$$d(G) \leqslant \frac{rX}{\alpha_r} - 1$$

By Lemma 3.1, it follows that G has an independent set of size at least

$$\frac{n}{1+d(G)} \geqslant \frac{\alpha_r n}{rX} = \frac{1}{r} \left(\frac{n}{k^{1/r}}\right)^{1/(r-1)},$$

as desired.

This covers all cases, completing the proof.

Next, we prove bounds on the average size and median size of an independent set in a (k, r)-sparse graph. This will play a key role in the proofs of Theorems 1.4 and 1.11.

**Lemma 3.4.** Let  $r, n \in \mathbb{N}$  and  $k \in \mathbb{R}$  such that  $r \ge 2, n \ge r^{2r}$ , and  $1 \le k \le {n \choose r}$ . Let G be an *n*-vertex (k, r)-sparse graph. Then there is a constant c > 0 such that,

$$\overline{\alpha}(G), \, \widehat{\alpha}(G) \ge \frac{c \, \log\left(i(G)\right)}{r \log\left(r \, k^{1/(r(r-1))} \log\left(i(G)\right)\right)}.$$

*Proof.* Let us first show that at most half of the independent sets in i(G) have size at most

$$\ell := \frac{\log_2(i(G))}{2(r-1)\log_2(r\,k^{1/(r(r-1))}\log_2(i(G)))}.$$

This will complete the proof for the bound on  $\overline{\alpha}(G)$ . In fact, we will show something stronger, i.e.,

$$\sum_{i=1}^{\ell} \binom{n}{i} \leqslant \frac{i(G)}{2}.$$

If  $\ell < 1$ , the claim is trivial, so we may assume  $\ell \ge 1$ . Note the following as a result of Lemma 3.2:

$$\log_2(i(G)) \leq n \leq k^{1/r} (r \log_2(i(G)))^{r-1}.$$

In particular, the first inequality implies that  $\ell \leq n/2$  (assuming  $i(G) \geq 2$ , which holds for  $n \geq 2$ ). With this in hand, we have

$$\sum_{i=1}^{\ell} \binom{n}{i} \leq \sum_{i=1}^{\ell} n^{i} \leq \ell n^{\ell} \leq \frac{1}{2} n^{2\ell}$$
$$\leq \frac{1}{2} \left( k^{1/r} \left( r \log_{2} \left( i(G) \right) \right)^{r-1} \right)^{2\ell}$$
$$\leq \frac{1}{2} x^{2(r-1)\ell},$$

for  $x := r k^{1/(r(r-1))} \log_2{(i(G))}$ . From here, it follows that

$$x^{2(r-1)\ell} = 2^{2(r-1)\ell \log_2 x} = i(G),$$

completing the proof for  $\overline{\alpha}(G)$ . For  $\hat{\alpha}(G)$ , we note the following:

$$\hat{\alpha}(G) = \frac{1}{i(G)} \sum_{I \in I(G)} |I| \ge \frac{1}{i(G)} \sum_{\substack{I \in I(G), \\ |I| \ge \ell}} |I| \ge \frac{\ell}{2},$$

as desired.

We are now ready to prove Theorem 1.4.

Proof of Theorem 1.4. Let I be a uniformly random independent set in I(G). We will show that  $\hat{\alpha}(G) = \mathbb{E}[|I|]$  is bounded from below by the desired quantity. To this end, we define the following for a vertex v:

$$p_v \coloneqq \Pr[v \in I], \qquad \overline{p}_v \coloneqq \frac{\mathbb{E}[|I \cap N_G(v)|]}{\Delta}$$

Let  $H := G - N_G[v]$  be the graph obtained by removing the vertices in  $N_G[v]$ . Note that every independent set in H can be extended to one in G containing v. For each  $S \subseteq N_G(v)$ , we define the following:

$$J_S := \{ J \in I(H) : N_G(v) \setminus N_G(J) = S \},\$$

i.e.,  $J_S$  contains the independent sets J of H for which every vertex in S is not adjacent to a vertex in J and every vertex in  $N_G(v) \setminus S$  is adjacent to some vertex in J. We may conclude the following:

$$i(G) = i(H) + \sum_{S \subseteq N_G(v)} |J_S| i(G[S]),$$

where the first term corresponds to independent sets containing v and the second corresponds to those not containing v. In particular, for  $j_S := |J_S|/i(H)$ , we have

$$p_v = \frac{1}{1 + \sum\limits_{S \subseteq N_G(v)} j_S i(G[S])}, \quad \overline{p}_v = \frac{\sum\limits_{S \subseteq N_G(v)} j_S i(G[S]) \hat{\alpha}(G[S])}{\Delta \left(1 + \sum\limits_{S \subseteq N_G(v)} j_S i(G[S])\right)}$$

The expression for  $\overline{p}_v$  above follows due to the following chain of equalities:

$$\sum_{I \in I(G)} |I \cap N_G(v)| = \sum_{S \subseteq N_G(v)} \sum_{\substack{I \in I(G), \\ I \cap N_G(v) = S}} |S|$$
$$= \sum_{S \subseteq N_G(v)} \sum_{J \in J_S} \sum_{I' \in I(G[S])} |I'|$$
$$= \sum_{S \subseteq N_G(v)} \sum_{J \in J_S} i(G[S]) \hat{\alpha}(G[S])$$
$$= \sum_{S \subseteq N_G(v)} |J_S| i(G[S]) \hat{\alpha}(G[S]).$$

Let  $\lambda \coloneqq \Delta / \log \Delta$ , and let  $\eta$  be defined as follows:

$$\eta \coloneqq \sum_{\substack{S \subseteq N_G(v), \\ i(G[S]) \ge \lambda}} j_S i(G[S]).$$

Recall the bounds on r in the statement of Theorem 1.4. In particular, the following holds for  $\rho$  small enough and S such that  $i(G[S]) \ge \lambda$ :

$$r^{2r} \leqslant \frac{\log \Delta}{2} \leqslant \log_2 \lambda \leqslant |S|.$$

As G is (k, r)-locally-sparse, G[S] is  $(\tilde{k}, r)$ -sparse, where  $\tilde{k} := \min\left\{k, \binom{|S|}{r}\right\}$ . Therefore, by Lemma 3.4, we have the following for some constant c > 0:

$$\hat{\alpha}(G[S]) \geq \frac{c \log \lambda}{r \log \left(r \, \tilde{k}^{1/(r(r-1))} \log \lambda\right)} \geq \frac{c \log \lambda}{r \log \left(r \, k^{1/(r(r-1))} \log \lambda\right)},$$

whenever  $i(G[S]) \ge \lambda$ . Let  $\ell$  denote the expression above. It follows that

$$p_v \ge \frac{1}{1+\lambda+\eta},\tag{3.5}$$

$$\overline{p}_v \ge \frac{\eta \ell}{\Delta \left(1 + \lambda + \eta\right)}.\tag{3.6}$$

Note that (3.5) is decreasing in  $\eta$  while (3.6) is increasing in  $\eta$ . Therefore, we have

$$p_v + \overline{p}_v \ge \max\{p_v, \overline{p}_v\} \ge \max\left\{\frac{1}{1+\lambda+\eta}, \frac{\eta\ell}{\Delta} \frac{1}{(1+\lambda+\eta)}\right\} \ge \frac{1}{1+\lambda+\Delta/\ell}.$$

Note the following:

$$\hat{\alpha}(G) = \sum_{v \in V(G)} p_v = \sum_{v \in V(G)} \overline{p}_v \implies \hat{\alpha}(G) = \frac{1}{2} \sum_{v \in V(G)} (p_v + \overline{p}_v) \ge \frac{n}{2(1 + \lambda + \Delta/\ell)}.$$
 (3.7)

Plugging in  $\lambda = \Delta / \log \Delta$ , we have the following for some constants  $c_1, c_2 > 0$ :

$$\begin{split} \lambda + \frac{\Delta}{\ell} &\leq \frac{\Delta}{\log \Delta} + \Delta \, \frac{c_1 r \log \left( r \, k^{1/(r(r-1))} \log \Delta \right)}{\log \Delta} \\ &\leq \Delta \, \frac{c_2 r \log \left( r \, k^{1/(r(r-1))} \log \Delta \right)}{\log \Delta}. \end{split}$$

Plugging this into (3.7), we may conclude the following for some constant c > 0:

$$\begin{split} \hat{\alpha}(G) & \geq \frac{c \, n \, \log \Delta}{r \, \Delta \log \left( r \, k^{1/(r(r-1))} \log \Delta \right)} \\ & \geq \frac{c \, n \log \Delta}{3r \, \Delta \max \left\{ \log r, \, \frac{1}{r(r-1)} \log k, \, \log^{(2)} \Delta \right\}} \\ & \geq \frac{c \, n}{3\Delta \max \left\{ \frac{\log k}{(r-1) \log \Delta}, \, \frac{r \log^{(2)} \Delta}{\log \Delta} \right\}}, \end{split}$$

where the last inequality follows due to the bound on r. As  $k = \Delta^{\varepsilon r}$  and  $r - 1 \ge r/2$ , this completes the proof.

### 4. Proof of Theorem 1.11

We will apply the following result of Bonamy, Kelly, Nelson, and Postle.

**Theorem 4.1** ([Bon+22, Theorem 1.13]). There exists  $\Delta_0 \in \mathbb{N}$  such that the following holds. Let G be a graph of maximum degree at most  $\Delta \ge \Delta_0$  with a correspondence cover  $\mathcal{H} = (L, H)$  and let  $\varepsilon \in (0, 1/2)$ . Let  $\ell, t : V(G) \to \mathbb{N}$ , and for each  $v \in V(G)$ , define  $\alpha_{\min}(v)$  as follows:

$$\alpha_{\min}(v) \coloneqq \min_{\substack{S \subseteq N_G(v), \\ i(G[S]) \ge t(v)}} \overline{\alpha}(G[S]).$$
(4.2)

If for each  $v \in V(G)$ ,

$$|L(v)| \ge \max\left\{\frac{2\deg(v)}{(1-\varepsilon)^2\alpha_{\min}(v)}, \frac{2t(v)\,\ell(v)}{\varepsilon}\right\},\,$$

and

(C1)  $\varepsilon(1-\varepsilon)\ell(v)t(v) \ge 18\log\Delta + 6\log 16$ ,

- (C2)  $\ell(v) \ge 36 \log \Delta + 12 \log 16$ , and
- (C3)  $\binom{\deg(v)}{\ell(v)}/\ell(v)! < \Delta^{-3}/8.$

Then G admits a proper  $\mathcal{H}$ -coloring.

Let  $\varepsilon = 1/4$ ,  $\ell(v) := \deg(v)^{1/2+\eta}$ , and  $t(v) := \deg(v)^{1/2-2\eta}$  for some constant  $\eta \in (0, 1/10)$  to be determined. For each  $v \in V(G)$ , let  $S(v) \subseteq N_G(v)$  be the minimizer of (4.2). As G is  $(\mathbf{k}, \mathbf{r})$ -locally-sparse, it follows that G[S(v)] is  $(\tilde{\mathbf{k}}(v), \mathbf{r}(v))$ -sparse, where

$$\tilde{\mathbf{k}}(v) \coloneqq \min\left\{\mathbf{k}(v), \left( \begin{matrix} |S(v)| \\ \mathbf{r}(v) \end{matrix} \right) \right\}.$$

Furthermore, we note the following as a result of (S1) and (S2) for  $\rho$  small enough:

$$|S(v)| \ge \log_2 t(v) \ge \frac{\log \deg(v)}{4} \ge \mathbf{r}(v)^{2\mathbf{r}(v)}$$

Therefore, by Lemma 3.4, we have the following for appropriate constants C, C' > 0:

$$\begin{aligned} \alpha_{\min}(v) &\geq \frac{C \log t(v)}{\mathbf{r}(v) \log \left(\mathbf{r}(v) \tilde{\mathbf{k}}(v)^{\frac{1}{\mathbf{r}(v)(\mathbf{r}(v)-1)}} \log t(v)\right)} \\ &\geq \frac{C' \log \deg(v)}{\mathbf{r}(v) \log \left(\mathbf{r}(v) \mathbf{k}(v)^{\frac{1}{\mathbf{r}(v)(\mathbf{r}(v)-1)}} \log \deg(v)\right)} \\ &\geq \frac{C' \log \deg(v)}{3\mathbf{r}(v) \max \left\{ \log \mathbf{r}(v), \frac{1}{\mathbf{r}(v)(\mathbf{r}(v)-1)} \log \mathbf{k}(v), \log^{(2)} \deg(v) \right\}} \\ &\geq \frac{C'}{6 \max \left\{ \varepsilon(v), \frac{\mathbf{r}(v) \log^{(2)} \deg(v)}{\log \deg(v)} \right\}}, \end{aligned}$$

where the last step follows by the bounds on  $\mathbf{r}(v)$  and the definition of  $\boldsymbol{\varepsilon}(v)$ . From here, we may conclude for some constant C > 0 and  $\deg(v)$  large enough (which follows for  $\Delta$  large enough by (S1)) that

$$C \deg(v) \max\left\{ \boldsymbol{\varepsilon}(v), \, \frac{\mathbf{r}(v) \log^{(2)} \deg(v)}{\log \deg(v)} \right\} \geq \max\left\{ \frac{2 \deg(v)}{(1-\varepsilon)^2 \alpha_{\min}(v)}, \frac{2 t(v) \ell(v)}{\varepsilon} \right\}.$$

In particular, it is now enough to show that conditions (C1)-(C3) are satisfied. Let us first consider (C1). We have

$$\varepsilon(1-\varepsilon)\ell(v)t(v) = \frac{3\deg(v)^{1-\eta}}{16} \ge \frac{3}{16}(\log\Delta)^{2-2\eta} \ge 18\log\Delta + 6\log 16,$$

where the first inequality follows by (S1) and the second for  $\Delta$  large enough.

Similarly, we have

$$\ell(v) = \deg(v)^{1/2+\eta} \ge (\log \Delta)^{1+2\eta} \ge 36 \log \Delta + 12 \log 16,$$

completing the proof of (C2).

To prove (C3), we will use the following inequality due to Stirling for n large enough:

$$n! \ge n^{n+1/2} e^{-n}$$

With this in hand, we have

$$\binom{\deg(v)}{\ell(v)} / \ell(v)! \leq \frac{\deg(v)^{\ell(v)}}{(\ell(v)!)^2} \leq \left(\frac{e^2 \deg(v)}{\ell(v)^{2+1/\ell(v)}}\right)^{\ell(v)} = \left(\frac{e^2}{\deg(v)^{2\eta + (1+2\eta)/\ell(v)}}\right)^{\ell(v)}$$

Taking logs on both sides, it is enough to show that

$$\ell(v)\left((2\eta + (1+2\eta)/\ell(v))\log\deg(v) - 2\right) \ge 3\log(2\Delta).$$

As  $\ell(v) \ge (\log \Delta)^{1+2\eta}$ , it is enough to have

$$(2\eta + (1+2\eta)/\ell(v))\log\deg(v) > 2,$$

which follows by (S1) for  $\Delta$  large enough and  $\eta = 1/100$ .

## Acknowledgements

We thank Anton Bernshteyn for his helpful comments.

#### References

- [Ajt+81] M. AJTAI, P. ERDŐS, J. KOMLÓS, and E. SZEMERÉDI. On Turán's theorem for sparse graphs, Combinatorica, 1 (1981), 313–317 (cit. on p. 2)
- [AKS80] M. AJTAI, J. KOMLÓS, and E. SZEMERÉDI. A note on Ramsey numbers, Journal of Combinatorial Theory, Series A, **29** (3) (1980), 354–360 (cit. on p. 2)
- [AKS99] N. ALON, M. KRIVELEVICH, and B. SUDAKOV. Coloring graphs with sparse neighborhoods, Journal of Combinatorial Theory, Series B, 77 (1) (1999), 73–82 (cit. on p. 2)
- [ABD22] J. ANDERSON, A. BERNSHTEYN, and A. DHAWAN. Coloring graphs with forbidden almost bipartite subgraphs, https://arxiv.org/abs/2203.07222 (preprint), 2022 (cit. on p. 4)
- [ABD23] J. ANDERSON, A. BERNSHTEYN, and A. DHAWAN. Colouring graphs with forbidden bipartite subgraphs, Combinatorics, Probability and Computing, **32** (1) (2023), 45–67 (cit. on p. 4)
- [ADK24] J. ANDERSON, A. DHAWAN, and A. KUCHUKOVA. Coloring locally sparse graphs, https://arxiv.org/abs/2402.19271 (preprint), 2024 (cit. on pp. 2, 7)
- [Ber19] A. BERNSHTEYN. The Johansson-Molloy theorem for DP-coloring, Random Structures & Algorithms, 54 (4) (2019), 653–664 (cit. on p. 3)
- [BD23] A. BERNSHTEYN and A. DHAWAN. Fast algorithms for Vizing's theorem on bounded degree graphs, https://arxiv.org/abs/2303.05408 (preprint), 2023 (cit. on p. 10)
- [Bon+20] M. BONAMY, M. DELCOURT, R. LANG, and L. POSTLE. *Edge-colouring graphs with* local list sizes, https://arxiv.org/abs/2007.14944 (preprint), 2020 (cit. on p. 4)
- [Bon+22] M. BONAMY, T. KELLY, P. NELSON, and L. POSTLE. Bounding  $\chi$  by a fraction of  $\Delta$  for graphs without large cliques, Journal of Combinatorial Theory, Series B, **157** (2022), 263–282 (cit. on pp. 4–6, 15)
- [Dav+18] E. DAVIES, M. JENSSEN, W. PERKINS, and B. ROBERTS. On the average size of independent sets in triangle-free graphs, Proceedings of the American Mathematical Society, 146 (1) (2018), 111–124 (cit. on p. 2)
- [Dav+20a] E. DAVIES, R. DE JOANNIS DE VERCLOS, R.J. KANG, and F. PIROT. Coloring triangle-free graphs with local list sizes, Random Structures & Algorithms, 57 (3) (2020), 730–744 (cit. on p. 4)
- [Dav+20b] E. DAVIES, R.J. KANG, F. PIROT, and J-S. SERENI. An algorithmic framework for colouring locally sparse graphs, https://arxiv.org/abs/2004.07151 (preprint), 2020 (cit. on p. 2)
- [Dav+20c] E. DAVIES, R.J. KANG, F. PIROT, and J-S. SERENI. Graph structure via local occupancy, https://arxiv.org/abs/2003.14361 (preprint), 2020 (cit. on p. 4)
- [Dha23] A. DHAWAN. Multigraph edge-coloring with local list sizes, https://arxiv.org/abs/2307.
   12094 (preprint), 2023 (cit. on p. 4)
- [DP18] Z. DVOŘÁK and L. POSTLE. Correspondence coloring and its application to list-coloring planar graphs without cycles of lengths 4 to 8, Journal of Combinatorial Theory, Series B, 129 (2018), 38–54 (cit. on p. 4)
- [ERT79] P. ERDŐS, A.L. RUBIN, and H. TAYLOR. *Choosability in graphs*, Congr. Numer, **26** (4) (1979), 125–157 (cit. on p. 4)
- [FV06] A. FRIEZE and J. VERA. On randomly colouring locally sparse graphs, Discrete Mathematics & Theoretical Computer Science, 8 (2006) (cit. on p. 2)

- [Joh96a] A. JOHANSSON. Asymptotic choice number for triangle free graphs. Technical Report 91–95. DIMACS, 1996 (cit. on p. 3)
- [Joh96b] A. JOHANSSON. The choice number of sparse graphs, https://www.cs.cmu.edu/~anupamg/ down/johansson-choice-number-of-sparse-graphs-coloring-kr-free.pdf (preprint), 1996 (cit. on p. 3)
- [Mol19] M. MOLLOY. The list chromatic number of graphs with small clique number, Journal of Combinatorial Theory, Series B, **134** (2019), 264–284 (cit. on pp. 2, 3, 5)
- [PH21] F. PIROT and E. HURLEY. Colouring locally sparse graphs with the first moment method, https://arxiv.org/abs/2109.15215 (preprint), 2021 (cit. on p. 2)
- [She83] J.B. SHEARER. A note on the independence number of triangle-free graphs, Discrete Mathematics, 46 (1) (1983), 83–87 (cit. on p. 2)
- [She95] J.B. SHEARER. On the independence number of sparse graphs, Random Structures & Algorithms, 7 (3) (1995), 269–271 (cit. on pp. 2, 5)
- [Viz76] V.G. VIZING. Coloring the vertices of a graph in prescribed colors, Diskret. Analiz, **29** (3) (1976), 10 (cit. on p. 4)