BOGOMOLOV PROPERTY AND GALOIS REPRESENTATIONS

FRANCESCO AMOROSO⁽¹⁾ AND LEA TERRACINI⁽²⁾

⁽¹⁾Dipartimento di Matematica Università di Torino Via Carlo Alberto, 10 - 10123 Torino, Italy

⁽²⁾ Dipartimento di Informatica Università di Torino Corso Svizzera, 185 - 10149 Torino, Italy

ABSTRACT. In 2013 P. Habegger proved the Bogomolov property for the field generated over \mathbb{Q} by the torsion points of a rational elliptic curve. We explore the possibility of applying the same strategy of proof to the case of field extensions cut out by Galois representations arising from more general modular forms.

2020 Mathematical Subject Classification: 11F03, 11F80, 11G50

Keywords. Weil height, Bogomolov property, modular forms, Galois representations.

1. INTRODUCTION

Let α be an algebraic number of degree d. We denote by $h(\alpha)$ the absolute and logarithmic Weil height. Assuming that α is not a root of unity, D. H. Lehmer suggested that $dh(\alpha)$ can be bounded below uniformly in α . Known as "Lehmer's problem", this question is still open. Nevertheless, in some special cases, not only Lehmer's conjecture is true, but it can also be sharpened. This different point of view was considered for the first time in [5], where the authors introduced the notion of fields with Bogomolov property, (property (B) for short), where the height is uniformly bounded below outside torsion points. By Northcott's Theorem, number fields are trivial examples of fields with (B). The compositum \mathbb{Q}^{tr} of all totally real fields is a less trivial example (apply [24, Corollary 1'] to the linear polynomial $P(z) = z - \alpha$ to get the optimal lower bound). In [5] the authors also show that fields with bounded local degrees at some finite place has Property (B). Note that by a result of [9] a Galois extension has uniformly bounded local degrees at every finite place if and only if its Galois group has finite exponent. Thus (infinite) Galois extensions with Galois group of bounded exponent have Property (B).

There are still other several interesting examples of subfields of the algebraic numbers with Property (B). Zannier conjectured that the maximal abelian extension \mathbb{Q}^{ab} of the rational field also satisfies property (B), and this was proved in [2]. By a special case of [3, Theorem 1.1], the same is true for \mathbb{K}^{ab} , for any number field \mathbb{K} . Moreover we can combine this result with the quoted result on Galois extensions of bounded exponent. Let \mathbb{K} be a number field and let \mathbb{L}/\mathbb{K} be an infinite Galois extension with Galois group G with G/Z(G) of finite exponent. Then \mathbb{L} has Property (B) ([1, Corollary 1.7]).

Another class of fields with Property (B) has been exhibited in [15]. Let \mathcal{E} be an elliptic curve defined over \mathbb{Q} . Then the field $\mathbb{Q}(\mathcal{E}_{tors})$ obtained by adjoining all torsion points of \mathcal{E} has the Bogomolov property. If \mathcal{E} has complex multiplication then $\mathbb{Q}(\mathcal{E}_{tors})$ is an abelian extension of a quadratic number field and thus it has the property (B) by the quoted result of [3]. However for a non-CM elliptic curve there are no number fields \mathbb{K} such that $\mathbb{Q}(\mathcal{E}_{tors})$ is contained in the maximal abelian extension of \mathbb{K} .

Note that by Kronecker-Weber theorem \mathbb{Q}^{ab} is the field obtained by adjoining all torsion points of the multiplicative group. A conjecture due to S. David (see [13, Conjecture, p.114]) suggests that $\mathbb{Q}(A_{tors})$ has Property (B), when A is an abelian variety defined on a number field, according as it is already known for the algebraic groups \mathbb{G}_m and $\mathcal{E} =$ elliptic curve. Note that for a CM abelian variety A the field $\mathbb{Q}(A_{tors})$ is still an abelian extension of a (CM) number field, and thus property (B) holds, again by the quoted result of [3].

Habegger's method provides a lower bound $h(\alpha) \geq c(p)$ with c(p) > 0 whenever we have a prime $p \geq 5$ such that \mathcal{E} has supersingular reduction at p (assumption (P1)) and the Galois representation $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \to \operatorname{Aut}(\mathcal{E}[p])$ is surjective (assumption (P2)). See [13, Theorem 1.3, p.114] for an explicit result. By a result of Elkies there exist infinitely many primes satisfying (P1); and by a result of Serre (P2)holds outside a finite set of primes. Thus we can find a prime p satisfying both the above conditions. Let now A be an abelian variety. Even under similar assumptions for a prime p, an analogue of the result of Habegger is not known.

One of the main purposes of this article is to re-interpret some known results on property (B) in terms of Galois representations, with the aim of providing a unifying framework for a part of the theory.

Definition 1.1. Let Ω be a separated topological group, \mathbb{F} be a subfield of $\overline{\mathbb{Q}}$ and $G_{\mathbb{F}} = \operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{F})$. We say that a continuous homomorphism

$$o: G_{\mathbb{F}} \longrightarrow \Omega,$$

has property (B) if $\mathbb{F}(\rho) := \overline{\mathbb{Q}}^{\ker(\rho)}$ does.

From this point of view, the main result of [2] implies (by Galois theory) that every homomorphism of $G_{\mathbb{Q}}$ taking values in an abelian group has Property (B).

In this vein, [1, Corollary 1.7] has the following consequence. Let \mathbb{K} be a number field, $\rho: G_{\mathbb{K}} \to \operatorname{GL}_n(\mathbb{F})$ be an absolutely irreducible representation, and $\mathbb{L} = \mathbb{K}(\rho)$. Schur's Lemma asserts that every intertwining automorphism of ρ is a scalar operator. In particular the image of the center of $G = \operatorname{Gal}(\mathbb{L}/\mathbb{K})$ is contained in the subgroup of scalar matrices of $\operatorname{GL}_n(\mathbb{F})$, so that the reduction map $\tilde{\rho}: G \to \operatorname{PGL}_n(\mathbb{F})$ factors through G/Z(G). Thus, if the image of $\tilde{\rho}$ has finite exponent, the same is true for G/Z(G) and [1, Corollary 1.7] implies that ρ has property (B).

Habegger theorem [15, Theorem 1] establishes Property (B) for the map

$$\rho_{\mathcal{E}}: G_{\mathbb{Q}} \longrightarrow \operatorname{GL}_2(\widehat{\mathbb{Z}})$$

which is the product of all the ℓ -adic representations

$$\rho_{\mathcal{E},\ell} : \operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \longrightarrow Aut(T_{\ell}(\mathcal{E})) = \operatorname{GL}_2(\mathbb{Z}_{\ell})$$

where \mathcal{E} is an elliptic curve defined over \mathbb{Q} and $T_{\ell}(\mathcal{E})$ is the corresponding Tate module. Indeed, by Galois correspondence $\mathbb{Q}(\mathcal{E}[\ell^n])$ is the subfield of $\overline{\mathbb{Q}}$ fixed by the kernel of the action of $G_{\mathbb{Q}}$ on the ℓ^n -torsion points of \mathcal{E} ; by passing to the projective limit we see that $\mathbb{Q}(\mathcal{E}[\ell^{\infty}]) = \mathbb{Q}(\rho_{\mathcal{E},\ell})$. Since $\mathbb{Q}(\mathcal{E}_{tors})$ is the compositum of $\mathbb{Q}(\mathcal{E}[\ell^{\infty}])$, for every prime ℓ , then it coincides with the field fixed by $\bigcap_{\ell} \ker(\rho_{\mathcal{E},\ell}) = \ker(\rho_{\mathcal{E}})$.

By the Shimura-Taniyama-Weil conjecture, proved by Wiles, all elliptic curves defined over \mathbb{Q} are associated to a cuspidal eigenform of weight 2 on $\Gamma_0(N)$, whose

eigenvalues for the Hecke operators are rational numbers. A question that naturally arises starting from Habegger's theorem is whether and to what extent it can be generalized to other Galois representations associated with modular forms. Let $k \geq 2$ be a positive integer and $f \in S_k(\Gamma_0(N))$ be a normalized eigenform for the Hecke algebra. Let \mathbb{K}_f be the number field generated by the Hecke eigenvalues of f, and let \mathcal{O}_f be its ring of integers. A construction due to Shimura and Deligne attaches to f a family of compatible Galois representations

$$\rho_{f,v}: G_{\mathbb{Q}} \longrightarrow \mathrm{GL}_2(\mathcal{O}_{f,v})$$

indexed by the finite places v of \mathcal{O}_f . For a rational prime p, we let

$$\rho_{f,p}: G_{\mathbb{Q}} \longrightarrow \prod_{v|p} \operatorname{GL}_2(\mathcal{O}_{f,v})$$

Moreover, let

$$\rho_f = \prod_v \rho_{f,v} : G_{\mathbb{Q}} \longrightarrow \mathrm{GL}_2(\widehat{\mathcal{O}}_f),$$

where $\widehat{\mathcal{O}}_f$ is the profinite completion of \mathcal{O}_f . Inspired by Habegger's theorem, we venture the following:

Conjecture 1.2. Let $f \in S_k(\Gamma_0(N))$ be a normalized eigenform. Then,

- a) for every v, $\rho_{f,v}$ has property (B);
- b) ρ_f has property (B).

Notice that Conjecture 1.2a) does not imply Conjecture 1.2b), as property (B) is not preserved under field composition (for instance, \mathbb{Q}^{tr} and obviously also $\mathbb{Q}(i)$ have property (B) but not $\mathbb{Q}^{tr}(i)$, see [1, Theorem 5.3, p. 1902]).

If f is a CM modular form, then $\mathbb{Q}(\rho_f)$ is an abelian extension of a quadratic field, (see for instance [20, Proposition 4.4]). Thus ρ_f has property (B) by the special case D = 1 of [3, Theorem 1.1].

When k = 2, Shimura [28] associated to f an abelian variety A defined over \mathbb{Q} , of dimension $[\mathbb{K}_f : \mathbb{Q}]$ equipped with an action of \mathcal{O}_f ; such abelian varieties have been called of type GL₂ in [23] and a generalization of Wiles' theorem [17, Corollary 10.2] establishes that all of them arise from modular forms.

Let A be the abelian variety associated to a modular form f of weight k = 2. Then $\mathbb{Q}(\rho) = \mathbb{Q}(A_{tors})$. Indeed by costruction, for every prime p the representation $\rho_{f,p} = \prod_{v|p} \rho_{f,v}$ is given by the action of $G_{\mathbb{Q}}$ on the p-adic Tate module $T_p(A) = \lim_{n \to \infty} A[p^n]$. Then it is immediate to see that $\mathbb{Q}(A[p^{\infty}])$ is the fixed field by ker $\rho_{f,p}$, so that $\mathbb{Q}(\rho_{f,p}) = \mathbb{Q}(A[p^{\infty}])$. Passing to ρ_f , we have

$$\mathbb{Q}\left(\rho_{f}\right) = \mathbb{Q}\left(\prod_{p} \rho_{f,p}\right) = \prod_{p} \mathbb{Q}\left(A[p^{\infty}]\right) = \mathbb{Q}\left(\bigcup_{p} A[p^{\infty}]\right) = \mathbb{Q}\left(\sum_{p} A[p^{\infty}]\right),$$

where the last equality holds because the sum in A is given by rational functions over \mathbb{Q} .

Thus Conjecture 1.2 intersects the already mentioned conjecture of David, which predicts that $\mathbb{Q}(A_{tors})$ has Property (B), when A is an abelian variety defined on a number field.

Assume that $\sum_{n\geq 1} a_n q^n$ is the q-expansion of f. Let p be a rational prime. Having fixed f, we consider the following assumption on f and p:

Assumption 1.3.

$$(P0) \ p \nmid N;$$

$$(P1) \ a_p = 0;$$

$$(P2) \ the \ reduction \ G_{\mathbb{Q}} \to \operatorname{GL}_2(\mathcal{O}_f/p\mathcal{O}_f) \ of \ \rho_{f,p} \ has \ image^1$$

$$\widehat{G}(\mathcal{O}_f/p\mathcal{O}_f) = \{\gamma \in \operatorname{GL}_2(\mathcal{O}_f/p\mathcal{O}_f) \mid \det \gamma \in (\mathbb{F}_p^{\times})^{k-1}\}$$

$$(P3)^2 \ p \ge 5, \ p \nmid k-1 \ and \ \frac{p+1}{2} \nmid k-1.$$

Our main result is the following:

Theorem 1.4. Let $f = \sum_{n\geq 1} a_n q^n \in S_k(\Gamma_0(N))$ be a normalized eigenform. Assume that there exists a rational prime p which satisfies Assumption 1.3. Let \mathbb{F}/\mathbb{Q} be a Galois extension unramified at p with property (B). Then $\rho_{f,p}|_{G_{\mathbb{F}}}$ has property (B).

As a consequence (see section 6 for the proof), we obtain the following direct generalization of Habegger's result:

Theorem 1.5. Let $f = \sum_{n\geq 1} a_n q^n \in S_k(\Gamma_0(N))$ be a normalized eigenform. Assume that there exists a rational prime p satisfying Assumption 1.3. Then ρ_f has property (B).

Recall that a modular form f is supersingular at a prime \mathfrak{p} of \mathcal{O}_f above p if $a_p \in \mathfrak{p}$. When $a_p \in \mathbb{Z}$ (as in the case $\mathbb{K}_f = \mathbb{Q}$) and $p \geq 5$, this implies $a_p = 0$, by the Deligne bound: $|a_p| \leq 2\sqrt{p}$. Therefore in the case of elliptic curves, Habegger's assumption $p \geq 5$ supersingular is equivalent to $a_p = 0$. Note that when $\mathbb{K}_f \neq \mathbb{Q}$ the condition $a_p = 0$, required in our proof, is in general stronger than supersingularity.

Our condition (P2) generalizes Habegger's assumption on the surjectivity of the Galois representation $G_{\mathbb{Q}} \to \operatorname{Aut} \mathcal{E}[p]$. It has been extensively investigated in the literature (see [22, 23, 4, 14, 18]); we summarize here the most relevant results for the purposes of this work.

In the case of modular forms associated to a non CM elliptic curve, a theorem of Serre [27] establishes property (P2) for every p except finitely many. This result, together with the aforementioned Elkies'theorem, ensures the non conditional character of Habegger's theorem.

For a modular form such that $\mathbb{K}_f \neq \mathbb{Q}$ the situation concerning the image of $\rho_{f,p}$ is complicated by the possible of presence of *inner twists*; i.e. the elements of the subgroup $\Gamma \subseteq Aut(\mathbb{K}_f)$ consisting of those automorphisms σ for which f^{σ} is equal to the twist of f by some Dirichlet character. Let \mathbb{K}_1 be the subfield of \mathbb{K} fixed by Γ . It is proved in [22, Theorem 3.1] that there is an abelian number field E such that for p large enough relatively to f, the restriction $\rho_{f,p}|_{G_E}$ has image $\widehat{G}(\mathcal{O}_{\mathbb{K}_1} \otimes \mathbb{Z}_p)$.

Therefore, all we can say is that condition (P2) holds for p large enough when f has no inner twists. In the case of trivial nebentypus (the one we are dealing with) it is easy to see that inner twists must arise from a quadratic Dirichlet character; and Ribet proved that they do not appear when N is square free [21, 3.9].

In particular, the results in [22] imply (P2) when k = 2, N is a prime, $p \nmid 6(N-1)$, \mathbb{K}_f is unramified at p and \mathcal{O}_f is generated over \mathbb{Z} by the coefficients a_n .

Some explicit examples with k = 2 of such modular forms satisfying also (P0) and (P1) are the following:

-
$$N = 73, p = 59, \mathbb{K}_f = \mathbb{Q}(\sqrt{13})$$
 (form 73.2.a.c in [29]).

¹The image is always contained in $\widehat{G}(\mathcal{O}_f/p\mathcal{O}_f)$, see the remark above Assumption 1.3. ²Note that all these conditions are satisfied if $p \geq 2k - 1$.

- $N = 167, p = 11, \mathbb{K}_f = \mathbb{Q}(\sqrt{5})$ (form 167.2.a.a in [29]). - $N = 383, p = 13, \mathbb{K}_f = \mathbb{Q}(\sqrt{5})$ (form 383.2.a.a in [29].
- $N = 151, p = 41, [\mathbb{K}_f : \mathbb{Q}] = 3$ (form 151.2.a.a in [29]).

Some other examples in weight k > 2 with $a_p = 0$, can be produced when $\mathbb{K} = \mathbb{Q}$. In this case in fact conditions (i),(ii), (iii) in [22, page 192] are automatically fulfilled, and if $a_p = 0$, then condition (iv) is also satisfied, because the residual representation $\bar{\rho}_v$ is irreducible, for every v|p. There are several examples of this type with k = 4, for example

- N = 186, p = 11, (form 186.4.a.a in [29]).
- N = 210, p = 11 or p = 23, (form 210.4.a.e in [29]).
- N = 1265, p = 53, (form 1265.4.a.c in [29]).

Some other examples with $k \ge 6$ are:

- N = 390, p = 7, k = 6 ((form 390.6.a.c in [29]).
- N = 66, p = 5, k = 8 (form 66.8.a.a in [29]).

We were not able to find examples with weight > 8.

Indeed, the simultaneous occurrence of conditions (P0) and (P1) for a non CM modular form is expected to be a fairly rare phenomenon. In this direction, Calegari and Sardari [7] proved that, fixed a pair of coprime integers p, N, there is only a finite number of eigenforms for $\Gamma_1(N)$ having $a_p = 0$.

Plan of the paper.

In Section 2 we introduce most of the notations we need and we make some preliminary remarks.

Section 3 is devoted to two general statements on lower bounds for the height. The proofs rely on the techniques developed in [2], [3], [15] and [1]. We fix two (possibly infinite) Galois extensions \mathbb{F}/\mathbb{Q} , \mathbb{L}/\mathbb{Q} . We fix a rational prime p. Usually, lower bounds for the height come from a dicotomy depending on whether p is ramified or not in \mathbb{L} . In the second case a Frobenius argument is generally enough to conclude. In the first case we need a descent argument. Here we change the point of view, assuming \mathbb{F} not ramified at p, with the residual degrees at p not growing from \mathbb{F} to \mathbb{L} . Moreover, we avoid the equidistribution argument of [15], by using instead the acceleration lemma of [1]. This simplifies the proofs, although it leads to worse lower bounds, as remarked in [12, Section 4.2.2]. As a toy example we deduce a generalization (Theorem 3.5) of the main result of [2].

One of the crucial steps in applying the height-machinery above is to have a precise as possible description of the sequence of the local higher ramification groups. This is done in [15, Lemma 3.3] using the fact that the formal group associated to the elliptic curve is a Lubin-Tate module over \mathbb{Z}_p . In Section 4 we show that, by local Langlands correspondence and the classification of the two dimensional crystalline representations of $G_{\mathbb{Q}_p}$ given by Breuil, our local representation at p is a twist of a power of a fundamental character, so that Lubin Tate theory is also applicable in our more general case.

Section 5 is devoted to prove Proposition 5.1, which is the analogous of Lemma 6.2 of [15] on the normal closure of the last ramification group in the field cut out by the reduction modulo p^n of our modular representation. This was possible thanks to condition (P2) in Assumption 1.3. For the proof we needed a group theoretical study of the linear group of a finite product of local rings. This is done in Appendix A.

The concluding Section 6 contains the proof of the results announced in the introduction.

2. NOTATIONS AND PRELIMINARY RESULTS

In the following

- local fields will be denoted by F, K, E, L, \ldots
- global fields will be denoted by $\mathbb{F}, \mathbb{K}, \mathbb{E}, \mathbb{L}, \dots$

For a field k, \overline{k} denotes an algebraic closure of k, and $G_k = \text{Gal}(\overline{k}/k)$ the absolute Galois group of k, endowed with the profinite topology.

Let Ω be a separated topological group and $\rho : G_{\mathbb{F}} \to \Omega$ be a continuous homomorphism; we shall denote by $k(\rho) = \overline{k}^{\ker \rho}$ the field cut out by ρ . Then $k(\rho)/k$ is a Galois extension whose Galois group is isomorphic to $\operatorname{Im}(\rho)$. Moreover, since ker ρ is closed, $G_{k(\rho)} = \ker \rho$.

When k is a local or global field and v is a place of k, we shall say that ρ is unramified (resp. totally ramified) at v if $k(\rho)$ is. Note that ρ induces an isomorphism $\operatorname{Gal}(k(\rho)/k) \to \operatorname{Im}(\rho)$ which as usual we denote by the same letter ρ .

From now on we fix a normalized new eigenform $f = \sum_{n} a_n q^n \in S_k(\Gamma_0(N))$ for the Hecke algebra of weight $k \geq 2$; then k is an even integer. We recall the relevant definitions. We denote by K the number field generated by the a_n 's and by \mathcal{O} its integer ring. A construction due to Shimura and Deligne associates to f a family of compatible Galois representations:

$$\rho_v: G_{\mathbb{O}} \longrightarrow \operatorname{GL}_2(\mathcal{O}_v), \quad v \text{ finite place of } \mathbb{K}.$$

Each ρ_v is characterized by the following two properties (where we denote by p_v the rational prime divided by v):

- (2.1) ρ_v is continuous and unramified at primes ℓ with $\ell \nmid Np_v$;
- (2.2) for $\ell \nmid Np_v$, the characteristic polynomial of the image by ρ_v of a Frobenius element at ℓ is $X^2 - a_\ell X + \ell^{k-1}$.

Remark 2.1. Let p be a prime and v be any place of \mathbb{K} dividing p. Then, by (2.2) and by Chebotarev lemma, the determinant of ρ_v is ε_p^{k-1} , where $\varepsilon_p \colon G_{\mathbb{Q}} \to \mathbb{Z}_p^{\times}$ is the p-adic cyclotomic character. (Indeed, for a Frobenius element $\operatorname{Frob}_{\ell}$ with $\ell \nmid Np$, we have: det $(\rho_v(\operatorname{Frob}_{\ell})) = \ell^{k-1} = \varepsilon_p^{k-1}(\operatorname{Frob}_{\ell})$.)

We let

$$\rho = \prod_{v} \rho_{v} : G_{\mathbb{Q}} \longrightarrow \mathrm{GL}_{2}(\widehat{\mathcal{O}}),$$

where $\widehat{\mathcal{O}}$ is the profinite completion of \mathcal{O} . For a prime p and n > 1, we also denote by $\rho(p^n)$ the reduction $G_{\mathbb{Q}} \to \operatorname{GL}_2(\widehat{\mathcal{O}}/p^n\widehat{\mathcal{O}}) = \operatorname{GL}_2(\mathcal{O}/p^n\mathcal{O})$ of ρ .

Remark 2.2. Let p be a prime which does not divide N and assume that $p = \prod_{v|p} \mathfrak{P}_v^{e_v}$

in \mathcal{O} . Let $n \geq 1$ be an integer. Then



where e_v is the ramification index at v. Thus, by Remark 2.1, the determinant of

 $\rho(p^n)$ is $\varepsilon_p^{k-1} \mod p^n$, where ε_p is the p-adic cyclotomic character. In particular the determinant takes values in $(\mathbb{Z}/p^n\mathbb{Z})^{\times}$.



We let

$$\widehat{G}(\mathcal{O}/p\mathcal{O}) = \{ \alpha \in \mathrm{GL}_2(\mathcal{O}/p\mathcal{O}) \mid \det(\alpha) \in (\mathbb{F}_p^{\times})^{k-1} \}.$$

Notice that the image of $\rho(p)$ is clearly contained in $G(\mathcal{O}/p\mathcal{O})$ by Remark 2.2.

For the reader convenience, we also recall a statement from [15, Lemma 2.1].

Lemma 2.3. Let F/\mathbb{Q}_p be a finite extension. Let K, L Galois extensions of F with K/F totally ramified and L/F finite and unramified. Then the restriction map $G_i(KL/L) \to G_i(K/F)$ is a well defined isomorphism.

Remark 2.4. In [15, Lemma 2.1](*iii*) K/F is also assumed finite, but this is unnecessary. Moreover in op.cit. it is stated that $\operatorname{Gal}(KL/L) \cap G_i(KL/F) \simeq G_i(K/F)$, but $\operatorname{Gal}(KL/L) \cap G_i(KL/F) = G_i(KL/L)$ since L/F is unramified.

3. DIOPHANTINE RESULTS.

We start with a simple result, which is a special case of [1, Theorem 1.5]. Here we allow finite ramification at p in \mathbb{F} .

Proposition 3.1. Let $G = \operatorname{Gal}(\mathbb{F}/\mathbb{Q})$ and let Z be the center. We assume that the ramification index of p in \mathbb{F} and the order of a lift of a Frobenius σ over p in G/Z are both finite. Then \mathbb{F} has property (B).

Proof. Let e and r be respectively the the ramification index and the order of σ in G/Z. Let \mathbb{F}_0/\mathbb{Q} be a finite Galois extension and $\alpha \in \mathbb{F}_0$. Since σ^r lies in the center of $\operatorname{Gal}(\mathbb{F}_0/\mathbb{Q})$, we have

$$|\sigma^r(\beta) - \beta^{p^r}|_v \le p^{-1/e}$$

for any $\beta \in \mathcal{O}_{\mathbb{F}_0}$ and for any place v dividing p. By [1, Lemma 2.1], there exists a positive integer λ which is explicitly bounded in terms of p and of e such that

$$|\sigma^r(\beta^{p^{\lambda}}) - \beta^{p^{r+\lambda}}|_v \le p^{-1}$$

We now apply [1, Lemma 2.2], taking into account that $\sigma^r(\alpha^{p^{\lambda}}) \neq \alpha^{p^{r+\lambda}}$, since otherwise α would be a root of unity.

Let \mathbb{L} be a (possible infinite) Galois extension. We now make the first assumption on \mathbb{L} which allows to globalise the metric inequality at ramified primes. We follow some ideas from [3, Lemma 4.2] and [15, Proof of Proposition 6.1].

Assumption 3.2. There exists $\tau \in G_{\mathbb{Q}} \cap I$, where I is an inertia group³ over p, such that

(3.1)
$$\epsilon_p(\tau) = g \quad with \ g \in \mathbb{Z}, g > 1,$$

and $\tau|_{\mathbb{L}} \in Z(\operatorname{Gal}(\mathbb{L}/\mathbb{Q})).$

³That is, the inverse limit of inertia groups.

The second assumption on \mathbb{L} concerns ramification groups. Although of technical nature, it is quite general to be satisfied in several cases, namely for the extensions cut out by our Galois representations. Given a finite Galois extension \mathbb{E}/\mathbb{Q} and $i \geq 0$ we recall that the *i*-th ramification group G_i at a place v over p is the subgroup of $\sigma \in \operatorname{Gal}(\mathbb{E}/\mathbb{Q})$ such that $|\sigma(\gamma) - \gamma|_v \leq p^{-(i+1)/e}$ for $\gamma \in \mathcal{O}_{\mathbb{E}}$. Here e is the ramification index of \mathbb{E} over p. The group G_i is the isomorphic image, by restriction, of the corresponding local *i*-th ramification group of the completed extension $\mathbb{E}_v/\mathbb{Q}_p$. As it is well known, the *i*-th ramification groups at different places of \mathbb{E} over p are conjugate.

Assumption 3.3. There exist C_1 , $C_2 \ge 1$ with the following properties. There exists a sequence of finite Galois extensions $(\mathbb{L}_n/\mathbb{Q})_{n>0}$ with

$$\mathbb{Q} = \mathbb{L}_0 \subsetneq \mathbb{L}_1 \subsetneq \cdots \subsetneq \mathbb{L}_n \subsetneq \cdots \subsetneq \mathbb{L}, \text{ and } \mathbb{L} = \bigcup_n \mathbb{L}_n.$$

such that for $n \ge 1$ the extension \mathbb{L}_n/\mathbb{Q} is ramified over p. Moreover, let $e_n \ge 1$ be the ramification index, $(G_i^n)_i$ the sequence of ramification groups of \mathbb{L}_n/\mathbb{Q} at a place over p. We further denote by $H_n = G_{i_n}^n$ the last non trivial ramification group and $\overline{H_n}$ its normal closure in $\operatorname{Gal}(\mathbb{L}_n/\mathbb{Q})$. Then

$$\frac{e_n}{i_n+1} \le C_1, \qquad |\overline{H_n}| \le C_2$$
$$\mathbb{L}_n^{\overline{H_n}} = \mathbb{L}_{n-1}.$$

and

Even we are not interested in explicit computations, let us remark that the above assumptions can be weakened in some cases (as for abelian extensions) to get sharper lower bounds.

We can now state the main result of this section. We denote by μ the set of roots of unity and by $\mu_{p^{\infty}}$ the set of roots of unity of order a power of p in $\overline{\mathbb{Q}}$.

Proposition 3.4. Let \mathbb{F}/\mathbb{Q} and \mathbb{L}/\mathbb{Q} two (possibly infinite) Galois extensions. We suppose that:

- \mathbb{F} is unramified at p and it satisfies property (B);
- L satisfies Assumptions 3.2 and 3.3.
- Then \mathbb{FL} has property (B) as well.

Proof. Let $I \subseteq \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ be an inertia group over $p, \tau \in I$ and $g \in \mathbb{Z}$ as in Assumption 3.2.

By definition of inertia group and by Assumption 3.2, there exists a prime ideal \mathfrak{P} of \mathbb{F} over p such that $\tau_{|\mathbb{F}} \in I(\mathfrak{P}|p)$. Since \mathbb{F} is not ramified at p, this last group is trivial, thus τ is trivial on \mathbb{F} . Again by Assumption 3.2, $\tau|_{\mathbb{L}} \in Z(\operatorname{Gal}(\mathbb{L}/\mathbb{Q}))$. Thus

(3.2)
$$\tau|_{\mathbb{FL}} \in Z(\operatorname{Gal}(\mathbb{FL}/\mathbb{Q})).$$

Since \mathbb{F} has property (B), there exists $c_0 > 0$ such that

(3.3)
$$\forall \beta \in \mathbb{F}^* \backslash \boldsymbol{\mu}, \quad h(\beta) \ge c_0$$

Let $\alpha \in (\mathbb{FL})^* \setminus \mu$. We define

(3.4)
$$\beta = \frac{\tau(\alpha)}{\alpha^g}$$

We remark that $\beta \notin \mu$. Otherwise $gh(\alpha) = h(\alpha^g) = h(\tau(\alpha)) = h(\alpha)$ which fails since α is not a root of unity and since g > 1. Moreover, $h(\beta) \leq (g+1)h(\alpha)$. Thus it is enough to show that there exists a c > 0 depending only on p, c_0, C_1, C_2 such that $h(\beta) \ge c$.

We fix a finite Galois extension \mathbb{F}_0/\mathbb{Q} such that $\mathbb{F}_0 \subseteq \mathbb{F}$ and $\beta \in \mathbb{F}_0\mathbb{L}$. Let $\{\mathbb{L}_n/\mathbb{Q}\}_{n\geq 0}$ as in Assumption 3.3. We define $n \geq 0$ as the smallest integer such that $\beta \in \mathbb{F}_0\mathbb{L}_n$. If n = 0 then $\beta \in \mathbb{F}$ and $h(\beta) \geq c_0$ by (3.3), since $\beta \notin \mu$. Thus we can assume $n \geq 1$.

We now fix a place v of $\mathbb{F}_0\mathbb{L}_n$ over p. Since \mathbb{F}_0 is not ramified at p, the ramification index e_n of $v_{|\mathbb{L}_n}$ coincide with the ramification index of v. Let $G_i^n \subseteq \operatorname{Gal}(\mathbb{L}_n/\mathbb{Q})$ and $G_i'^n \subseteq \operatorname{Gal}(\mathbb{F}_0\mathbb{L}_n/\mathbb{Q})$ be respectively the *i*-th ramification subgroups at $v_{|\mathbb{L}_n}$ and at v. The restriction map $\operatorname{Gal}(\mathbb{F}_0\mathbb{L}_n/\mathbb{F}_0) \to \operatorname{Gal}(\mathbb{L}_n/\mathbb{Q})$ sends $H'_n := G'^n_{i_n}$ bijectively in $H_n = G^n_{i_n}$, by Lemma 2.3 applied to the completions at v of the involved fields. Therefore the normal closure $\overline{H'_n}$ of H'_n in $\operatorname{Gal}(\mathbb{F}_0\mathbb{L}_n/\mathbb{Q})$ is sent bijectively in the normal closure $\overline{H_n}$ of H_n in $\operatorname{Gal}(\mathbb{L}_n/\mathbb{Q})$. By Assumption 3.3 we have

(3.5)
$$\frac{e_n}{i_n+1} \le C_1, \qquad |\overline{H'_n}| \le C_2.$$

Since $\mathbb{L}_n^{\overline{H_n}} = \mathbb{L}_{n-1}$ by the last claim of Assumption 3.3, Galois theory shows that

(3.6)
$$(\mathbb{F}_0 \mathbb{L}_n)^{\overline{H'_n}} = \mathbb{F}_0 \mathbb{L}_{n-1}.$$

Thus $\beta \notin (\mathbb{F}_0 \mathbb{L}_n)^{\overline{H'_n}}$ by minimality of n. Since $\overline{H'_n}$ is generated by the conjugates of $G'^n_{i_n}$, by Galois theory $\beta \notin (\mathbb{F}_0 \mathbb{L}_n)^{\phi G'^n_{i_n} \phi^{-1}}$ for some $\phi \in \text{Gal}(\mathbb{F}_0 \mathbb{L}_n/\mathbb{Q})$. Thus there exists $\sigma \in G'^n_{i_n}$ such that $\phi \sigma \phi^{-1}(\beta) \neq \beta$. We fix ϕ, σ with the above property and we let $\beta' = \phi^{-1}(\beta)$. Thus $\sigma(\beta') \neq \beta'$. Moreover, if $\tilde{\phi}$ is a lifting of ϕ to \mathbb{FL} , we have $\beta' = \frac{\tau(\alpha')}{\alpha'^g}$ with $\alpha' = \tilde{\phi}^{-1}(\alpha)$, by definition (3.4) and since the restriction of τ is in the center of $\text{Gal}(\mathbb{FL}/\mathbb{Q})$ (*cf.* (3.2)). Note also that $\beta' \in \mathbb{F}_0\mathbb{L}_n$ and $h(\beta') = h(\beta)$ since β' is a conjugate of β . Summing up, we have found $\sigma \in G'^n_{i_n}$ and $\alpha' \in \mathbb{FL}$ such that

(3.7)
$$\beta' = \frac{\tau(\alpha')}{\alpha'^g} \text{ satisfies } \frac{\sigma(\beta')}{\beta'} \neq 1$$

and we need to prove that there exists a c > 0 depending only on p, c_0, C_1, C_2 such that $h(\beta') \ge c$.

Fact.
$$rac{\sigma(eta')}{eta'}
ot\in oldsymbol{\mu}_{p^{\infty}}$$

Proof. Let us assume by contradiction that $\frac{\sigma(\beta')}{\beta'} \in \mu_{p^{\infty}}$. Let $\tilde{\sigma}$ be a lift of σ to \mathbb{FL} . By (3.7) and since τ is central,

$$\frac{\sigma(\beta')}{\beta'} = \frac{\tilde{\sigma}\tau(\alpha')}{\tau(\alpha')} \left(\frac{\tilde{\sigma}(\alpha'^g)}{\alpha'^g}\right)^{-1} = \frac{\tau(\eta)}{\eta^g}.$$

with $\eta = \tilde{\sigma}(\alpha')/\alpha' \in \mathbb{FL}$. Hence

$$gh(\eta) = h(\eta^g) = h(\tau(\eta)) = h(\eta),$$

which implies $h(\eta) = 0$ since g > 1. Thus η is a root of unity. Write η as $\eta = \eta_1 \eta_2$ with $\eta_1 \in \boldsymbol{\mu}_{p^{\infty}}$ and with η_2 of order not divisible by p. Hence $\frac{\tau(\eta_1)}{\eta_1^g} = 1$ by (3.1). Thus $\frac{\sigma(\beta')}{\beta'} = \frac{\tau(\eta_2)}{\eta_2^g}$ has order not divisible by p. But $\frac{\sigma(\beta')}{\beta'} \in \boldsymbol{\mu}_{p^{\infty}}$ and $\frac{\sigma(\beta')}{\beta'} \neq 1$ by (3.7). Contradiction.

By the fact above,

(3.8) For any *p*-power
$$p^{\lambda}$$
 we have $\sigma(\beta')^{p^{\lambda}} \neq \beta'^{p^{\lambda}}$.

Since $\sigma \in G_{i_n}^{\prime n}$, for any $\gamma \in \mathcal{O}_{\mathbb{F}_0 \mathbb{L}_n}$ we have

$$|\sigma(\gamma) - \gamma|_v \le p^{-(i_n+1)/e_n}.$$

By [1, Lemma 2.1], there exists a positive integer λ which is explicitly bounded in terms of p, C_1 and C_2 , such that

(3.9)
$$|\sigma(\gamma^{p^{\lambda}}) - \gamma^{p^{\lambda}}|_{v} \le p^{-|\overline{H'_{n}}|}$$

We consider the action of $\operatorname{Gal}(\mathbb{F}_0\mathbb{L}_n/\mathbb{Q})$ on itself by conjugation and its stabiliser $C(\sigma) = \{\tau \in \operatorname{Gal}(\mathbb{F}_0\mathbb{L}_n/\mathbb{Q}) \mid \tau\sigma\tau^{-1} = \sigma\}$. Let $\tau \in C(\sigma)$. We use (3.9) with $\tau^{-1}\gamma$ instead of γ . Since $\sigma\tau^{-1} = \tau^{-1}\sigma$ and $|\tau^{-1}(\star)|_w = |\star|_{\tau(v)}$, we get

$$|\sigma(\gamma^{p^{\lambda}}) - \gamma^{p^{\lambda}}|_{w} \le p^{-|\overline{H'_{n}}|}$$

with $w = \tau(v)$ and thus for any w in the orbit S of v under the the action of $C(\sigma)$ on the places above p. Using [2, Lemma 1] as in the proof of [1, Lemma 2.1] we find

$$|\sigma(\beta'^{p^{\lambda}}) - \beta'^{p^{\lambda}}|_{w} \le c(w) \max(1, |\sigma(\beta')|_{w})^{p^{\lambda}} \max(1, |\beta'|_{w})^{p^{\lambda}}, \ \forall w \in S$$

with $c(w) = p^{-|\overline{H'_n}|}$. This last inequality also holds for an arbitrary place w of \mathbb{L} not dividing p, with

$$c(w) = \begin{cases} 1, & \text{if } w \nmid \infty; \\ 2, & \text{if } w \mid \infty. \end{cases}$$

Let $d_n = [\mathbb{F}_0 \mathbb{L}_n : \mathbb{Q}]$ and $d_{n,w} = [(\mathbb{F}_0 \mathbb{L}_n)_w : \mathbb{Q}_w]$ with w a place of $\mathbb{F}_0 \mathbb{L}_n$. Using the Product Formula to $\sigma(\beta')^{p^{\lambda}} - \beta'^{p^{\lambda}}$, which is $\neq 0$ by (3.8), as in the proof of [1, Lemma 2.1] we get:

$$0 = \sum_{w} \frac{d_{n,w}}{d_n} \log |\sigma(\beta'^{p^{\lambda}}) - \beta'^{p^{\lambda}}|_w$$

$$\leq \sum_{w} \frac{d_{n,w}}{d_n} \left(\log c(w) + p^{\lambda} \log \max\{1, |\sigma(\beta')|_w\} + p^{\lambda} \log \max\{1, |\beta'|_w\}\right)$$

$$= \left(\sum_{w\mid\infty} \frac{d_{n,w}}{d_n}\right) \log 2 - \left(\sum_{w\in S} \frac{d_{n,w}}{d_n}\right) |\overline{H'_n}| \log p + 2p^{\lambda}h(\beta').$$

Since $\sum_{w\mid\infty} \frac{d_{n,w}}{d_n} = 1$ and since $d_{n,w} = d_{n,v}$ for any place $w \mid p$ of \mathbb{L}_n , and thus in particular for any $w \in S$, we get

(3.10)
$$2p^{\lambda}h(\beta') \ge \frac{d_{n,v}}{d_n}|S| \cdot |\overline{H'_n}|\log p - \log 2.$$

To conclude, we need a lower bound for S. Note that $\sigma \in \overline{H'_n} \triangleleft \operatorname{Gal}(\mathbb{F}_0 \mathbb{L}_n / \mathbb{Q})$, thus the orbit O of σ is contained in $\overline{H'_n}$. Thus:

$$|C(\sigma)| = \frac{d_n}{|O|} \ge \frac{d_n}{|\overline{H'_n}|}.$$

The stabiliser of v under the action of $\operatorname{Gal}(\mathbb{F}_0\mathbb{L}_n/\mathbb{Q})$ is by definition the decomposition group D(v|p) of v over p which has (again by definition) cardinality d_v . Thus,

$$|S| = \frac{|C(\sigma)|}{|D(v|p) \cap C(\sigma)|} \ge \frac{d_n}{d_{n,v}} |\overline{H'_n}|^{-1}.$$

10

From (3.10) we then get $h(\beta') \ge c$ with

$$c = \frac{\log(p/2)}{2p^{\lambda}} > 0. \quad \Box$$

As a toy example we deduce from Proposition 3.4 the following statement which generalizes the main result of [2].

Theorem 3.5. Let \mathbb{F}/\mathbb{Q} be a Galois extension unramified at p with property (B). Then $\mathbb{F}(\mu_{p^{\infty}})$ has property (B) as well.

Proof. Let $\mathbb{L} = \mathbb{Q}(\boldsymbol{\mu}_{p^{\infty}})$. Since \mathbb{L}/\mathbb{Q} is totally ramified at p and abelian, Assumption 3.2 is trivially satisfied. We consider the filtration given by $\mathbb{L}_n = \mathbb{Q}(\zeta_{p^n})$. Then it is easy to see that $G_i^n = \operatorname{Gal}(\mathbb{Q}(\zeta_{p^n})/\mathbb{Q}(\zeta_{p^j}))$, for $p^{j-1} \leq i \leq p^j - 1$ (see [19, Ch. II, §10, Ex. 1]). Thus, in the notation of Assumption (3.3), $e_n = (p-1)p^{n-1}$, $i_n = p^{n-1} - 1$ and $\overline{H_n} = G_{i_n}^n$. We then have $e_n/(i_n + 1) \leq p$ and $|\overline{H_n}| \leq p$. Clearly, $\mathbb{L}_n^{\overline{H_n}} = \mathbb{L}_{n-1}$. The Assumption 3.3 therefore holds with $C_1 = C_2 = p$.

4. The local representation at $v \mid p$.

In what follows, we recall some results on crystalline representations. We fix a prime p and an embedding $\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_p$, which allows to identify $G_{\mathbb{Q}_p} = \operatorname{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$ with a subgroup of $G_{\mathbb{Q}}$. Note that this is helpful but harmful in our context, as some results may depend on the chosen embedding. To simplify notations and to avoid future conflicts, we simply denote by ε the p-adic cyclotomic character ε_p . Let $q = p^2$ and denote by \mathbb{Q}_q the unique unramified quadratic extension of \mathbb{Q}_p . Let v be a place of \mathbb{K} over p. The two dimensional irreducible crystalline representations of $G_{\mathbb{Q}_p}$ have been classified in [6, Proposition 3.1]. They are of the form $V_{k',a,\chi} = V_{k',a} \otimes \chi$, where k' is an integer ≥ 2 , a is an element in the maximal ideal of the valuation ring of $\overline{\mathbb{Q}}_p$, and χ is a crystalline character, that is the product of an unramified character by an integral power of the cyclotomic character. The only isomorphisms between them are $V_{k',a,\chi} \simeq V_{k',-a,\chi\mu_{-1}}$, where μ_{-1} is the unramified quadratic character of $G_{\mathbb{Q}_p}$. Up to a twist by a power of the cyclotomic character, we can suppose that χ is unramified. In this case $V_{k',a,\chi}$ has Hodge-Tate weights (0, k' - 1), and the φ -filtered module $D = D_{k',a,\chi}$ associated to $V_{k',a,\chi}$ is $\overline{\mathbb{Q}}_p e_1 + \overline{\mathbb{Q}}_p e_2$ with φ -action given by

$$\varphi(e_1) = p^{k'-1} \chi(p^{-1})^2 e_2$$

$$\varphi(e_2) = -e_1 + a \chi(p^{-1}) e_2$$

and the filtration is given by e

$$Fil^{i}(D) = \begin{cases} D & \text{if } i \leq 0\\ \overline{\mathbb{Q}}_{p}e_{1} & \text{if } 1 \leq i \leq k'-1\\ 0 & \text{if } i \geq k' \end{cases}$$

In particular, the characteristic polynomial of φ on the associated φ -filtered module is equal to $X^2 - aX + \chi(p^{-1})^2 p^{k'-1}$.

Let's come back to our representations ρ_v , with $v \mid p$. Let as assume $p \nmid N$ (Assumption 1.3 (P0)). By [25, Theorem 1.2.4 (ii)], $\rho_{v\mid G_{\mathbb{Q}_p}}$ is crystalline and the characteristic polynomial of φ on the associated φ -filtered module is equal to $X^2 - a_p X + p^{k-1}$. Moreover, representations associate to modular forms of weight k have Hodge-Tate weight (0, k-1), (see [11]). It follows that the filtered module associated to $\rho_{v\mid G_{\mathbb{Q}_p}}$ is $D_{k',a,\chi}$ with k' = k, $a = a_p$, and $\chi(p) = 1$ or -1. Now assume also $a_p = 0$ (Assumption 1.3 (P1)). Since $V_{k',a,\chi} \simeq V_{k',-a,\chi\mu_{-1}}$, where μ_{-1} is the unramified quadratic character of $G_{\mathbb{Q}_p}$, in both cases $\rho_v|_{G_{\mathbb{Q}_p}}$ is isomorphic to $V_{k,0}$. Taking into account the explicit description of $V_{k,0}$ in [6, Prop. 3.2] we obtain:

Proposition 4.1. Let us assume 1.3 (P0) and (P1). Then for every prime v of \mathcal{O} above p, $\rho_{v|G_{\Omega_n}}$ is isomorphic to the crystalline representation

$$V_{k,0} \simeq \left(\operatorname{ind}_{G_{\mathbb{Q}_q}}^{G_{\mathbb{Q}_p}} \epsilon_2^{k-1} \right) \otimes \mu_{\sqrt{-1}}$$

where:

- $\epsilon_2: G_{\mathbb{Q}_q} \to \mathbb{Z}_q^{\times}$ is one of the two continuous characters characterized by the property that their composition with the injection $\mathbb{Q}_q^{\times} \hookrightarrow G_{\mathbb{Q}_q}^{ab}$ (given by local class field theory) is trivial on p and is one of the two natural embeddings when restricted to \mathbb{Z}_q^{\times} ; - $\mu_{\sqrt{-1}}$ is one of the two unramified characters of $G_{\mathbb{Q}_p}$ sending Frob_p in $\sqrt{-1}$.

We also need the following proposition.

Proposition 4.2. Let us assume 1.3 (P0)-(P1)-(P3). Let v be a place of K over p. Then, up to conjugation by an element of $\operatorname{GL}_2(\mathcal{O}_v)$, the restriction $\rho_{v|G_{\mathbb{Q}_p}}$ takes values in $\operatorname{GL}_2(\mathbb{Z}_p)$.

Proof. Let \mathfrak{P} be the prime ideal corresponding to v. Denote by $\rho_v(\mathfrak{P})$ the reduction $G_{\mathbb{Q}} \to \operatorname{GL}_2(\mathcal{O}_v/\mathfrak{P}\mathcal{O}_v) = \operatorname{GL}_2(\mathcal{O}/\mathfrak{P})$ of ρ_v . Since $p+1 \nmid k-1$ (which holds a fortiori since $\frac{p+1}{2} \nmid k-1$ by Assumption 1.3 (P3)), [7, Lemma 2.1 (1)]) ensures that the restriction of $\rho_v(\mathfrak{P})$ to a decomposition group⁴ at p is absolutely irreducible, so that ρ_v a fortiori is absolutely irreducible. By Proposition 4.1, the representations $\rho_{v|G_{\mathbb{Q}_p}} \simeq V(k,0)$ take values in $\operatorname{GL}_2(\mathbb{Q}_p)$, so that the traces lie in \mathbb{Z}_p . By [8, Theorem 1], up to conjugation by an element of $\operatorname{GL}_2(\mathcal{O}_v)$, the restriction $\rho_{v|G_{\mathbb{Q}_p}}$ takes values in $\operatorname{GL}_2(\mathbb{Z}_p)$.

Remark 4.3. From the proof of Proposition 4.1 we see that ker $\rho_{v|G_{\mathbb{Q}_p}}$ does not depend on v|p.

Let p be a rational prime satisfying Assumptions 1.3 (P0), (P1). By Propositions 4.1, for every prime v of \mathcal{O} above p, $\rho_{v|G_{\mathbb{Q}_p}}$ is a twist by an unramified character of the induced representation of a power of a character arising from the Tate module of a Lubin-Tate formal group over \mathbb{Z}_q (see [10, Théorème I.2.1]). This is exactly the same situation occuring for the representation of $G_{\mathbb{Q}_p}$ on the p-adic Tate module of an elliptic curve \mathcal{E} over \mathbb{Q} having p as a supersingular prime, and this is the key point of the fundamental Galois theoretic analysis of the p-power torsion points on E in [15, Lemma 3.3]. The proof of this lemma rests on the theory of Lubin-Tate modules and its relation to local class field theory and thus only depends on the local p-adic representation. We provide here a reformulation suitable for our situation, without any reference to the theory of elliptic curves.

We first recall some simple facts concerning twists of representations. Let L be a local field, A be an abelian group, $\nu, \omega : G_L \to A$ be homomorphisms. Since ker $\nu \cap \ker \omega = \ker \nu \omega \cap \ker \omega$, we know that

$$L \subseteq L(\nu), L(\omega), L(\nu\omega) \subseteq L(\nu)L(\omega) = L(\nu\omega)L(\omega)$$

⁴By Proposition 4.1 this restriction does not depend on the decomposition group but only on p.

Lemma 4.4. Assume that $im(\omega) \subseteq im(\nu)$ and the extensions $L(\nu)/L$, $L(\omega)/L$ are linearly disjoint; then $im(\nu\omega) = im(\nu)$ and the restriction

(4.1)
$$\operatorname{Gal}(L(\nu)L(\omega)/L(\nu\omega)) \to \operatorname{Gal}(L(\omega)/L)$$

is an isomorphism.

Proof. Since the two extensions $L(\nu)/L$ and $L(\omega)/L$ are linearly disjoint, the product of restrictions induces an isomorphism

(4.2)
$$\operatorname{Gal}(L(\nu)L(\omega)/L) \simeq \operatorname{Gal}(L(\nu)/L) \times \operatorname{Gal}(L(\omega)/L).$$

This formula implies that the image of ν restricted to $\operatorname{Gal}(\overline{L}/L(\omega)) = \ker \omega$ coincides with $\operatorname{im}(\nu)$. Since $\nu = \nu \omega$ over $\ker \omega$, this prove that $\operatorname{im}(\nu \omega) = \operatorname{im}(\nu)$. We now prove the second assertion. The restriction (4.1) is clearly injective. Let $\tau \in \operatorname{Gal}(L(\omega)/L)$. Since $\operatorname{im}(\omega) \subseteq \operatorname{im}(\nu)$, there exists an element $\sigma \in \operatorname{Gal}(L(\nu)L(\omega)/L)$ such that $\nu(\sigma) = \omega(\tau)^{-1}$ and σ restricts to τ in $\operatorname{Gal}(L(\omega)/L)$. Then $\nu(\sigma)\omega(\sigma) = 1$ so that $\sigma \in \operatorname{Gal}(L(\nu)L(\omega)/L(\nu\omega))$.

Lemma 4.5. Assume that $im(\omega) \subseteq im(\nu)$. If ν is totally ramified and ω unramified with finite image, then

(i) $\operatorname{im}(\nu\omega) = \operatorname{im}(\nu);$

(ii) $\nu\omega$ is totally ramified.

(iii) For $i \geq 0$ let G_i denote the *i*-th higher ramification group. Then

$$G_i(L(\nu\omega)/L) \simeq G_i(L(\nu)/L).$$

In particular

$$\operatorname{Gal}(L(\nu\omega)/L) \simeq \operatorname{Gal}(L(\nu)/L).$$

Proof. A diagram may be of help



Here the two extensions on the left top and right top are unramified by [19, Ch. II, Prop. 7.2].

Let us prove (i) and (ii). Since ν is totally ramified and ω is unramified, the two extensions $L(\nu)/L$ and $L(\omega)/L$ are linearly disjoint. Lemma 4.4 then shows that $\operatorname{im}(\nu\omega) = \operatorname{im}(\nu)$ and

(4.3)
$$[E:L(\nu\omega)] = [L(\omega):L].$$

Denoting by f the residual degree, we then have:

$$f(E/L) = f(E/L(\nu)) = [E:L(\nu)] \le [L(\omega):L]$$

by looking to the right side of the diagram. Moreover, by looking to the left part of the diagram

$$f(E/L) \ge f(E/L(\nu\omega)) = [E:L(\nu\omega)] = [L(\omega):L]$$

where the last equality comes from (4.3). Thus $f(E/L) = [E : L(\nu\omega)]$ and then $L(\nu\omega)/L$ is totally ramified.

We now prove (*iii*). The second assertion follows from the first one for i = 0. For the first, using (i) and Lemma 2.3 twice (with $L = L(\omega)$ and with $K = L(\nu)$ the first time, $K = L(\nu\omega)$ the second time) we get

$$G_i(L(\nu)/L) = G_i(E/L(\omega)) = G_i(L(\nu\omega)/L).$$

We are now ready to completely describe the ramification of the local representation $\rho_p|_{G_{\mathbb{Q}_p}}$, as in [15, Lemma 3.3]. Recall that $q = p^2$. We put:

$$\delta = \frac{q-1}{(q-1,k-1)}.$$

Notice that $\delta \neq 1$ because k is even.

Lemma 4.6. Let $n \ge 1$. We denote by $\epsilon_{2,n}$ the composition of ϵ_2 with the reduction modulo p^n .

(i) The extension $\mathbb{Q}_q(\epsilon_{2,n}^{k-1})/\mathbb{Q}_q$ is totally ramified and abelian of degree δq^{n-1} , and

$$\operatorname{Gal}(\mathbb{Q}_q(\epsilon_{2,n}^{k-1})/\mathbb{Q}_q) \simeq \{g^{k-1} \mid g \in (\mathbb{Z}_q/p^n \mathbb{Z}_q)^{\times}\} \simeq \mathbb{Z}/\delta\mathbb{Z} \times (\mathbb{Z}/p^{n-1}\mathbb{Z})^2.$$

(ii) Let j and i be integers with $1 \le j \le n$ and $q^{j-1} \le (q-1, k-1)i \le q^j - 1$. The higher ramification groups are given by

$$G_i(\mathbb{Q}_q(\epsilon_{2,n}^{k-1})/\mathbb{Q}_q)) = \operatorname{Gal}(\mathbb{Q}_q(\epsilon_{2,n}^{k-1})/\mathbb{Q}_q(\epsilon_{2,j}^{k-1})).$$

Proof. By [10, Théorème I.2.1], ϵ_2 is up to isogeny the Tate module of a Lubin–Tate formal group \mathcal{F} over \mathbb{Z}_q . Then $\mathbb{Q}_q(\epsilon_{2,n}) = \mathbb{Q}_q(\mathcal{F}(n))$, where $\mathcal{F}(n)$ is the group of p^n -division points of \mathcal{F} . Then in the case k = 2 (i) and (ii) follow from [19, Theorem V.5.4] and [19, Proposition V.6.1].

Assume now k > 2. The extension $\mathbb{Q}_q(\epsilon_{2,n}^{k-1})/\mathbb{Q}_q$ is the subextension of $\mathbb{Q}_q(\epsilon_{2,n})/\mathbb{Q}_q$ cut out by the (k-1)-th power of $\epsilon_{2,n}$. By assertion (i) in the special case k = 2, and since $p \nmid k - 1$ by Assumption 1.3 (P3), we see that $\operatorname{Gal}(\mathbb{Q}_q(\epsilon_{2,n})/\mathbb{Q}_q(\epsilon_{2,n}^{k-1}))$ is the subgroup of $\mathbb{Z}/(q-1)\mathbb{Z}$ of index d := (q-1, k-1). Assertion (i) follows.

We now prove (ii). The group $\operatorname{Gal}(\mathbb{Q}_q(\epsilon_{2,n})/\mathbb{Q}_q(\epsilon_{2,n}^{k-1}))$ has order d. By Herbrand Theorem [19, Ch. II, Theorem 10.7], $G_r(\mathbb{Q}_q(\epsilon_{2,n})/\mathbb{Q}_q)$ is mapped onto $G_{\eta(r)}(\mathbb{Q}_q(\epsilon_{2,n}^{k-1})/\mathbb{Q}_q)$ by the restriction map

res:
$$\operatorname{Gal}(\mathbb{Q}_q(\epsilon_{2,n})/\mathbb{Q}_q) \longrightarrow \operatorname{Gal}(\mathbb{Q}_q(\epsilon_{2,n}^{k-1})/\mathbb{Q}_q).$$

Here (cf. [19, Ch. II, Proposition 10.6])

$$\eta(r) = \frac{1}{d} \sum_{\sigma \in \operatorname{Gal}(\mathbb{Q}_q(\epsilon_{2,n})/\mathbb{Q}_q(\epsilon_{2,n}^{k-1}))} \min\{i(\sigma), r+1\} - 1$$

with $i(\sigma) = v(\sigma(x) - x)$ for a generator x of the ring of integers of $\mathbb{Q}_q(\epsilon_{2,n})$ over $\mathbb{Q}_q(\epsilon_{2,n})$. For $\sigma \in \operatorname{Gal}(\mathbb{Q}_q(\epsilon_{2,n})/\mathbb{Q}_q(\epsilon_{2,n}^{k-1}))$ we have

$$i(\sigma) = \begin{cases} \infty & \text{if } \sigma = id \\ 1 & \text{if } \sigma \neq id \end{cases}$$

because $G_1(\mathbb{Q}_q(\epsilon_{2,n})/\mathbb{Q}_q)$ is a *p*-group and $\operatorname{Gal}(\mathbb{Q}_q(\epsilon_{2,n})/\mathbb{Q}_q(\epsilon_{2,n}^{k-1}))$ has order dividing q-1 and thus $G_1(\mathbb{Q}_q(\epsilon_{2,n})/\mathbb{Q}_q) \cap \operatorname{Gal}(\mathbb{Q}_q(\epsilon_{2,n})/\mathbb{Q}_q(\epsilon_{2,n}^{k-1}))$ is trivial. Therefore

$$\eta(r) = \frac{1}{d}(r+1+(d-1)) - 1 = \frac{r}{d}.$$

It follows that, for j and i integers such that $1 \le j \le n$ and $q^{j-1} \le d i \le q^j - 1$,

$$G_i(\mathbb{Q}_q(\epsilon_{2,n}^{k-1})/\mathbb{Q}_q)) = \operatorname{res} \left(G_{d\,i}(\mathbb{Q}_q(\epsilon_{2,n})/\mathbb{Q}_q)\right)$$

= $\operatorname{res} \left(\mathbb{Q}_q(\epsilon_{2,n})/\mathbb{Q}_q(\epsilon_{2,j})\right)$ (by the case $k = 2$)
= $\operatorname{Gal}(\mathbb{Q}_q(\epsilon_{2,n}^{k-1})/\mathbb{Q}_q(\epsilon_{2,j}^{k-1})).$

Now (ii) follows from (i).

Proposition 4.7. Let $n \ge 1$.

(i) The extension $\mathbb{Q}_q(p^n)/\mathbb{Q}_q$ is totally ramified and abelian of degree δq^{n-1} . Moreover

$$\operatorname{Gal}\left(\mathbb{Q}_q(p^n)/\mathbb{Q}_q\right) \simeq \mathbb{Z}/\delta\mathbb{Z} \times \left(\mathbb{Z}/p^{n-1}\mathbb{Z}\right)^2$$

(ii) Let j and i be integers with $1 \le j \le n$ and $q^{j-1} \le (q-1, k-1)i \le q^j - 1$. The higher ramification groups are given by

$$G_i(\mathbb{Q}_q(p^n)/\mathbb{Q}_q) = \operatorname{Gal}(\mathbb{Q}_q(p^n)/\mathbb{Q}_q(p^j))$$

In particular, the last non trivial higher ramification group has index

$$i_n := \frac{q^{n-1} - 1}{(q - 1, k - 1)}$$

and

$$G_{i_n} = \begin{cases} (\mathbb{Z}/p\mathbb{Z})^2 & \text{if } n \ge 2\\ \mathbb{Z}/\delta\mathbb{Z} & \text{if } n = 1. \end{cases}$$

(iii) If M is an integer prime to p and $v \mid p$, then the image of $\rho_{v|G_{\mathbb{Q}_q}}$ contains the scalar matrix $\begin{pmatrix} M^{k-1} & 0 \\ 0 & M^{k-1} \end{pmatrix}$.

Proof. By Proposition 4.1

$$\rho_{v|G_{\mathbb{Q}_q}} \simeq \left(\operatorname{ind}_{G_{\mathbb{Q}_q}}^{G_{\mathbb{Q}_p}} \epsilon_2^{k-1} \right) \otimes \mu_{\sqrt{-1}}.$$

By Mackey's restriction formula, we see that

(4.4)
$$\left(\operatorname{ind}_{G_{\mathbb{Q}_{q}}}^{G_{\mathbb{Q}_{p}}}\epsilon_{2}^{k-1}\right)_{|G_{\mathbb{Q}_{q}}} = \epsilon_{2}^{k-1} \oplus \epsilon_{2}^{\prime k-1},$$

where ϵ'_2 is the character obtained from ϵ_2 by composing with the algebraic conjugation of $\mathbb{Q}_q/\mathbb{Q}_p$. Let for short $\nu = \epsilon_2^{k-1}$, $\omega = \mu_{\sqrt{-1}|G_{\mathbb{Q}_q}}$ and remark that ω takes values in $\{\pm 1\}$ and thus it is stable by the algebraic conjugation of $\mathbb{Q}_q/\mathbb{Q}_p$. Notice also that $-1 \in \operatorname{im}(\nu) = (\mathbb{Z}_q^{\times})^{k-1}$, because the weight k is even. Then $\rho_{v|G_{\mathbb{Q}_q}}$ is conjugate to

(4.5)
$$\begin{aligned} G_{\mathbb{Q}_q} &\longrightarrow \mathrm{GL}_2(\mathbb{Z}_q) \\ \sigma &\to \begin{pmatrix} \nu\omega & 0 \\ 0 & \nu\omega & \nu \end{pmatrix}. \end{aligned}$$

Since the kernel is normal,

$$\ker \rho(p^n)_{|G_{\mathbb{Q}_q}} = \{ \sigma \in G_{\mathbb{Q}_q} \mid (\nu\omega)(\sigma) \equiv (\nu\omega)'(\sigma) \equiv 1 \pmod{p^n} \} \\ = \{ \sigma \in G_{\mathbb{Q}_q} \mid (\nu\omega)(\sigma) \equiv 1 \pmod{p^n} \} = \ker(\nu_n\omega_n),$$

where we have denoted by ν_n, ω_n , the composition of ν, ω respectively with the reduction modulo p^n . Thus

(4.6)
$$\mathbb{Q}_q(p^n) = \mathbb{Q}_q(\nu_n \omega_n)$$

By its definition, ν_n is totally ramified; on the other hand ω is unramified with finite image contained in im(ν). By (4.6), by Lemma 4.5 (iii) with $L = \mathbb{Q}_q$ and with ν , ω replaced by ν_n , ω_n respectively, and by Lemma 4.6

$$\operatorname{Gal}(\mathbb{Q}_q(p^n)/\mathbb{Q}_q) = \operatorname{Gal}(\mathbb{Q}_q(\nu_n\omega_n)/\mathbb{Q}_q) \simeq \operatorname{Gal}(\mathbb{Q}_q(\nu_n)/\mathbb{Q}_q) \simeq \mathbb{Z}/\delta\mathbb{Z} \times (\mathbb{Z}/p^{n-1}\mathbb{Z})^2$$
which proves (i), and

$$G_i(\mathbb{Q}_q(p^n)/\mathbb{Q}_q) = G_i(\mathbb{Q}_q(\nu_n\omega_n)/\mathbb{Q}_q) \simeq G_i(\mathbb{Q}_q(\nu_n)/\mathbb{Q}_q) = \operatorname{Gal}(\mathbb{Q}_q(\nu_n)/\mathbb{Q}_q(\nu_j)).$$

Moreover, again by Lemma 4.5 (iii), with $L = \mathbb{Q}_q$ and with ν , ω replaced by ν_i , ω_i (i = j, n), and by (4.6)

$$Gal(\mathbb{Q}_q(\nu_n)/\mathbb{Q}_q(\nu_j)) = \ker \left(Gal(\mathbb{Q}_q(\nu_n)/\mathbb{Q}_q) \to Gal(\mathbb{Q}_q(\nu_j)/\mathbb{Q}_q) \right)$$
$$= \ker \left(Gal(\mathbb{Q}_q(\nu_n\omega_n)/\mathbb{Q}_q) \to Gal(\mathbb{Q}_q(\nu_j\omega_j)/\mathbb{Q}_q) \right)$$
$$= Gal(\mathbb{Q}_q(\nu_n\omega_n)/\mathbb{Q}_q(\nu_j\omega_j)) = Gal(\mathbb{Q}_q(p^n)/\mathbb{Q}_q(p^j).$$

The last two displayed lines prove (ii).

To prove (iii), we recall that $\nu \omega \colon G_{\mathbb{Q}_q} \to (\mathbb{Z}_q^{\times})^{k-1}$ is surjective by Lemma 4.5 (i). Scalar matrices with coefficients in $(\mathbb{Z}_p^{\times})^{k-1}$ are central elements, invariant by the algebraic conjugation of $\mathbb{Q}_q/\mathbb{Q}_p$ and thus, by (4.5), are in the image of $\rho|_{G_{\mathbb{Q}_q}}$. \Box

5. The Normal Closure Lemma.

Proposition 5.1. Let p be a rational prime satisfying Assumptions 1.3. The normal closure of $\operatorname{Gal}(\mathbb{Q}_q(p^n)/\mathbb{Q}_q(p^{n-1}))$ in $\operatorname{Gal}(\mathbb{Q}(p^n)/\mathbb{Q})$ is $\operatorname{Gal}(\mathbb{Q}(p^n)/\mathbb{Q}(p^{n-1}))$.

Proof. We closely follow the proof of [15, Lemma 6.2], using the result in Appendix A instead of [15, Lemma 6.1].

Since $\operatorname{Gal}(\mathbb{Q}_q(p^n)/\mathbb{Q}_q(p^{n-1})) \subseteq \operatorname{Gal}(\mathbb{Q}(p^n)/\mathbb{Q}(p^{n-1}))$, and since the latter is normal, the normal closure of $\operatorname{Gal}(\mathbb{Q}_q(p^n)/\mathbb{Q}_q(p^{n-1}))$ in $\operatorname{Gal}(\mathbb{Q}(p^n)/\mathbb{Q})$ is contained in $\operatorname{Gal}(\mathbb{Q}(p^n)/\mathbb{Q}(p^{n-1}))$. We want to prove equality.

Let for short

$$\begin{array}{ccc} H_n & & & & & \\ H_n & & & & \\ \parallel & & & \parallel \\ & & & \parallel \\ & & & \parallel \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & & \\ & & & & \\ & & & \\ & & & & \\ & & & \\ & & & & \\ & & & & \\ & & & & \\$$

Since $\mathcal{O}/p\mathcal{O}$ is a finite product of local \mathbb{F}_p -algebras and $p \geq 5$, we can freely use the results of the Appendix A with $\mathcal{A} = \mathcal{O}/p\mathcal{O}$. We also recall that

$$\widehat{G}(\mathcal{O}/p\mathcal{O}) = \{ \alpha \in \operatorname{GL}_2(\mathcal{O}/p\mathcal{O}) \mid \det(\alpha) \in (\mathbb{F}_p^{\times})^{k-1} \}$$

Assume first n = 1.

By Proposition 4.2, $\bar{\rho}_1(H_1)$ is a subgroup of $\operatorname{GL}_2(\mathbb{F}_p)$; by Proposition 4.7 (ii), its cardinality is $\delta = \frac{q-1}{(q-1,k-1)}$. Let $s = \frac{p-1}{(p-1,k-1)}$. By Assumption 1.3 (P3),

 $\frac{p+1}{2} \nmid k-1$. Thus $\frac{p+1}{(p+1,k-1)} > 2$ and $\delta > 2s$. The determinant maps surjectively $\bar{\rho}_1(H_1)$ onto $(\mathbb{F}_p^{\times})^{k-1} = (\mathbb{F}_p^{\times})^{(k-1,p-1)}$, because its image is that of the (k-1)-th power $G_{\mathbb{Q}_p} \to (\mathbb{F}_p^{\times})^{k-1}$ of the cyclotomic character modulo p. Thus $\bar{\rho}_1(H_1)$ satisfies the hypotheses of Proposition A.1.1, so that the normal closure of $\bar{\rho}_1(H_1)$ in $\widehat{G}(\mathcal{O}/p\mathcal{O})$ is $\widehat{G}(\mathcal{O}/p\mathcal{O})$ itself. Since by the hypothesis 1.3 (P2), $\bar{\rho}_1$ is an isomorphism between G_1 and $\widehat{G}(\mathcal{O}/p\mathcal{O})$, the normal closure of H_1 in G_1 is G_1 itself.

Now we turn to the case $n \geq 2$. Let $\sigma \in \Omega_n$. As in the proof of [15, Lemma 6.2], $\bar{\rho}_n(\sigma)$ is the reduction mod p^n of a matrix of the shape $I + p^{n-1}A_{\sigma}$, where I denotes the 2-by-2 identity matrix and where the "log" matrix $A_{\sigma} \in M_2(\mathcal{O})$ is well-defined mod $p\mathcal{O}$. Moreover, $\det(\bar{\rho}_n(\sigma)) \in (\mathbb{Z}/p^n\mathbb{Z})^{\times}$ by Remark 2.2. Since $\det(I + p^{n-1}A_{\sigma}) \equiv 1 + \operatorname{trace}(A_{\sigma})p^{n-1} \mod p^n$, the matrix A_{σ} has trace in \mathbb{F}_p . Let $\widehat{M}_2(\mathcal{O}/p\mathcal{O})$ be the subgroup of $M_2(\mathcal{O}/p\mathcal{O})$ consisting of matrices having trace in \mathbb{F}_p . We denote by \mathcal{L}_n the map $\Omega_n \to \widehat{M}_2(\mathcal{O}/p\mathcal{O})$ which sends σ to A_{σ} . Thus we have the following commutative diagram :



Since $\bar{\rho}_n$ and "log" are both injective, \mathcal{L}_n is injective as well.

The group $M_2(\mathcal{O}/p\mathcal{O})$ is a direct sum

$$\widehat{M}_2(\mathcal{O}/p\mathcal{O}) = \mathrm{M}_2(\mathcal{O}/p\mathcal{O})^0 \oplus \mathbb{F}_p \cdot \mathrm{I}$$

where $M_2(\mathcal{O}/p\mathcal{O})^0$ is the set of trace zero matrices. We also remark that $\mathcal{L}_n(H_n) \subseteq M_2(\mathbb{F}_p)$, by Proposition 4.2. As in the proof of [15, Lemma 6.2], for $\tau \in \Omega_n$ and $\sigma \in \text{Gal}(\mathbb{Q}(p^n)/\mathbb{Q})$ one has

(5.1)
$$\tilde{\rho}_n(\sigma)\mathcal{L}_n(\tau)\tilde{\rho}_n(\sigma)^{-1} = \mathcal{L}_n(\sigma\tau\sigma^{-1}),$$

where

and "res" denotes the restriction. Let $\overline{H_n}$ be the normal closure of H_n in $\operatorname{Gal}(\mathbb{Q}(p^n)/\mathbb{Q})$. Recall that $\overline{H_n} \subseteq \Omega_n$. We want to show that $\mathcal{L}_n(\overline{H_n}) = \widehat{M_2}(\mathcal{O}/p\mathcal{O})$. This will imply that $\overline{H_n} = \Omega_n$ coincides with Ω_n as we want (and that \mathcal{L}_n is surjective). Note that, by property 1.3 (P2) $\overline{\rho_1}$ is surjective. Since "res" is obviously surjective, $\widetilde{\rho_n}$ is surjective as well. Thus (5.1) says in particular that $\mathcal{L}_n(\overline{H_n})$ is invariant by the adjoint action⁵ of $\widehat{G}(\mathcal{O}/p\mathcal{O})$. Note that $M_2(\mathcal{O}/p\mathcal{O})^0$ is also invariant by this action. Thus

$$V = \mathcal{L}_n(\overline{H_n}) \cap \mathcal{M}_2(\mathcal{O}/p\mathcal{O})^0$$

⁵That is, the action by conjugation.

is an \mathbb{F}_p -subspace of $M_2(\mathcal{O}/p\mathcal{O})^0$ invariant by the adjoint action of $\widehat{G}(\mathcal{O}/p\mathcal{O})$, and in particular by that of $\mathrm{SL}_2(\mathcal{O}/p\mathcal{O}) \subseteq \widehat{G}(\mathcal{O}/p\mathcal{O})$. Moreover

$$V \cap \mathcal{M}_2(\mathbb{F}_p)^0 = \mathcal{L}_n(\overline{H_n}) \cap \mathcal{M}_2(\mathbb{F}_p)^0 \neq \{0\}.$$

Indeed, $\mathcal{L}_n(H_n)$ is contained in $M_2(\mathbb{F}_p)$ by Proposition 4.2; since $|H_n| = p^2$ by Lemma 4.7(i), its intersection with $M_2(\mathbb{F}_p)^0$ is already not trivial.

Then, by Proposition A.2.5, $V = M_2(\mathcal{O}/p\mathcal{O})^0$ and thus $\mathcal{L}_n(\overline{H_n})$ contains $M_2(\mathcal{O}/p\mathcal{O})^0$. The cyclotomic character $H_n \to (1 + p^{n-1}\mathbb{Z}_p)/(1 + p^n\mathbb{Z}_p)$ is surjective. Indeed by Proposition 4.1 and Mackey formula

$$\rho_v|_{G_{\mathbb{Q}_q}} \simeq (\epsilon_2 \oplus \epsilon'_2) \otimes \mu_{\sqrt{-1}}|_{G_{\mathbb{Q}_q}}$$

(see the proof of Lemma 4.7), its image coincides with $N_{\mathbb{Q}_q/\mathbb{Q}_p}(1+p^{n-1}\mathbb{Z}_q) \mod 1+p^n\mathbb{Z}_p$, and the norm map from $1+p^{n-1}\mathbb{Z}_q \to 1+p^{n-1}\mathbb{Z}_p$ is surjective (see for example [26, Ch. V, §2, Prop. 3]). By condition 1.3 (P3) $p \nmid k-1$ and thus the (k-1)-th power of the cyclotomic character is surjective. Therefore also the trace from $\mathcal{L}_n(\overline{H_n})$ to \mathbb{F}_p is surjective.

Thus $\mathcal{L}_n(\overline{H_n}) = \widehat{M}_2(\mathcal{O}/p\mathcal{O})$ as claimed.

Remark 5.2. We have seen in the previous proof that $\overline{H_n} = \Omega_n$ can be identified to a subgroup of $\operatorname{GL}_2(\mathcal{O}/p\mathcal{O})$ if n = 1 and to a subgroup of $\operatorname{M}_2(\mathcal{O}/p\mathcal{O})$ if $n \ge 2$. Thus $|\overline{H_n}| \le p^{4[\mathbb{K}:\mathbb{Q}]}$, where we recall that \mathbb{K} is the number field generated by the coefficients of our modular form.

6. PROOF OF THE MAIN RESULTS.

Proposition 3.1 allows us to shortly settle the unramified case. We define

$$\rho_p = \prod_{v|p} \rho_v : G_{\mathbb{Q}} \longrightarrow \mathrm{GL}_2(\widehat{\mathcal{O}})$$

and

$$\rho^{(p)} = \prod_{v \nmid p} \rho_v : G_{\mathbb{Q}} \longrightarrow \mathrm{GL}_2(\widehat{\mathcal{O}}).$$

Proposition 6.1. If Assumption 1.3 (P0) and (P1) hold, then $\rho^{(p)}$ has property (B).

Proof. Let $\mathbb{F} = \mathbb{Q}(\rho^{(p)})$. By our assumptions and by (2.1) and (2.2), \mathbb{F} is unramified at p and the characteristic polynomial of $\rho^{(p)}(\operatorname{Frob}_p)$ is $X^2 + p^{k-1}$. It follows that Frob_p^2 lies in the center of $\operatorname{Gal}(\mathbb{F}/\mathbb{Q})$. Then Property (B) for \mathbb{F} is ensured by Proposition 3.1.

Proof of Theorem 1.4. We want to apply Proposition 3.4 with $\mathbb{L} = \mathbb{Q}(p^{\infty})$. We first check Assumption 3.2. By Lemma 4.7 (iii) (with M = 2) there exists an element $\tau \in \operatorname{Gal}(\mathbb{Q}_q(p^{\infty})/\mathbb{Q}_q)$ whose image by ρ_v is the scalar matrix $\binom{2^{k-1}}{0} \binom{0}{2^{k-1}}$. Since $\rho_v|_{G_{\mathbb{Q}_q}}$ is totally ramified and $\mathbb{Q}_q/\mathbb{Q}_p$ is unramified, τ can be lifted to an element in the inertia subgroup of $G_{\mathbb{Q}_p}$. As its image is a scalar, τ lies in the center of $\operatorname{Gal}(\mathbb{L}/\mathbb{Q})$. Its action on the p^{∞} -th roots of unity is given by the *p*-adic cyclotomic character, i.e. the determinant of $\rho_v(\tau)$, which is 4^{k-1} . Then $\tau(\zeta) = \zeta^{4^{k-1}}$ for every $\zeta \in \mu_{p^{\infty}}$.

We now check Assumption 3.3. We consider the filtration given by $\mathbb{L}_n = \mathbb{Q}(p^n)$. Let $n \geq 1$. Fix a place of $\mathbb{Q}(p^\infty)$ over p corresponding to an embedding $\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_p$. Then, by Lemma 4.7 (i) and (ii), and in the notation of Assumption (3.3), $e_n = \delta q^{n-1} = \frac{(q-1)q^{n-1}}{(q-1,k-1)}$, $i_n = \frac{q^{n-1}-1}{(q-1,k-1)}$ and $H_n \simeq \operatorname{Gal}(\mathbb{Q}_q(p^n)/\mathbb{Q}_q(p^{n-1}))$. Thus

$$\frac{e_n}{i_n+1} = \frac{(q-1)q^{n-1}}{q^{n-1}-1+(q-1,k-1)} \le q-1.$$

By Proposition 5.1, $\overline{H_n} = \text{Gal}(\mathbb{L}_n/\mathbb{L}_{n-1})$, so that $\mathbb{L}_n^{\overline{H_n}} = \mathbb{L}_{n-1}$. Moreover, by Remark 5.2, $|\overline{H_n}| \leq p^{4[\mathbb{K}:\mathbb{Q}]}$.

The Assumption 3.3 therefore holds with $C_1 = p^2$ and $C_2 = p^{4[\mathbb{K}:\mathbb{Q}]}$.

Proof of Theorem 1.5. Let again $\mathbb{F} = \mathbb{Q}(\rho^{(p)})$. Then \mathbb{F} is unramified at p, and it has Property (B) by Proposition 6.1. By Theorem 1.4, $\mathbb{F}(\rho_p)$ has Property (B). But

$$\mathbb{F}(\rho_p) = \mathbb{Q}(\rho^{(p)} \times \rho_p) = \mathbb{Q}(\rho).$$

APPENDIX A. ON THE LINEAR GROUPS OF A FINITE PRODUCT OF LOCAL RINGS

Let $\mathcal{A} = \prod_{i=1}^{r} \mathcal{A}_i$ be a finite product of local commutative \mathbb{F}_p -algebras. In this appendix we bring together some results on the linear groups overs \mathcal{A} .

A.1. A result on the normal closure. Let p be a prime number and k be a positive integer. We define

$$\widehat{G}(\mathcal{A}) = \{ \gamma \in \operatorname{GL}_2(\mathcal{A}) \mid \det(\gamma) \in (\mathbb{F}_p^{\times})^{k-1} \}$$

It is the semidirect product $\operatorname{SL}_2(\mathcal{A}) \rtimes (\mathbb{F}_p^{\times})^{k-1}$, where $x \in (\mathbb{F}^{\times})^{k-1}$ acts on $\operatorname{SL}_2(\mathcal{A})$ by conjugation by $\begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$. The aim of this subsection is to prove the following proposition.

Proposition A.1.1. Let us assume $p \ge 5$. Let $s = \frac{p-1}{(k-1,p-1)}$ and let H be a subgroup of $\operatorname{GL}_2(\mathbb{F}_p)$ such that |H| > 2s and such that the determinant maps surjectively onto $(\mathbb{F}_p^{\times})^{k-1}$. Then the normal closure of H in $\widehat{G}(\mathcal{A})$ is $\widehat{G}(\mathcal{A})$ itself.

To start with, we remark that since \mathcal{A} is a semilocal ring, $SL_2(\mathcal{A})$ is generated by elementary matrices, i.e. matrices of the form

$$S(\alpha) = \begin{pmatrix} 1 & \alpha \\ 0 & 1 \end{pmatrix}, \qquad T(\alpha) = \begin{pmatrix} 1 & 0 \\ \alpha & 1 \end{pmatrix}, \qquad \alpha \in \mathcal{A},$$

(see [16, Theorem 4.3.9]).

Lemma A.1.2. Let us assume $p \geq 5$. Then \mathcal{A} is generated as a \mathbb{F}_p -vector space by the squares of all its invertible elements.

Proof. Let \mathcal{B} be the \mathbb{F}_p -vector subspace of \mathcal{A} generated by the squares of all the invertible elements of \mathcal{A} . Note that \mathcal{B} is a subring with unity of \mathcal{A} which contains \mathbb{F}_p . For every $i = 1, \ldots, k$ denote by $e_i \in \mathcal{A}$ the idempotent projecting to the summand \mathcal{A}_i . Firstly we show that $e_i \in \mathcal{B}$ for every i. Indeed, choose two elements a, b in \mathbb{F}_p^{\times} such that $a^2 \neq b^2$ (which is possible because $|\mathbb{F}_p^{\times}| > 2$) and let $\alpha = ae_i + \sum_{j \neq i} e_j$, $\beta = be_i + \sum_{j \neq i} e_j$. Then $\alpha^2 - \beta^2 = (a^2 - b^2)e_i$ is a non-zero \mathbb{F}_p -multiple of e_i belonging to \mathcal{B} .

Now we prove that every \mathcal{A}_i coincides with the subring \mathcal{B}_i generated over \mathbb{F}_p by the squares of the invertible elements. Assume that $\alpha \in \mathcal{A}_i^{\times}$. Since $p \neq 2$ and since \mathcal{A}_i is local, one of $\alpha \pm 1$ is in \mathcal{A}_i^{\times} . Then $(\alpha \pm 1)^2 - \alpha^2 = \pm 2\alpha + 1 \in \mathcal{B}_i$, so that $\alpha \in \mathcal{B}_i$. It follows $\mathcal{A}_i^{\times} \subseteq \mathcal{B}_i$. Now consider an element π in the maximal ideal \mathfrak{m}_i of \mathcal{A}_i . then $(1 + \pi)^2 - 1 = 2\pi + \pi^2 = \pi(2 + \pi) \in \mathcal{B}_i$. Since $(2 + \pi)^{-1} \in \mathcal{A}_i^{\times} \subseteq \mathcal{B}_i$, we deduce $\pi \in \mathcal{B}$. Then \mathcal{B}_i is a subring containing all units of \mathcal{A}_i and \mathfrak{m}_i , so that $\mathcal{B}_i = \mathcal{A}_i$.

Finally, assume $\alpha \in \mathcal{A}$ and $\alpha = \sum_{i=1}^{s} \alpha_i e_i$ with $\alpha_i \in \mathcal{A}_i$. For $i = 1, \ldots, s$ we can now write $\alpha_i = \sum_j a_{i,j} \beta_{ij}^2$, with $a_{i,j} \in \mathbb{F}_p$ and $\beta_{ij} \in \mathcal{A}_i^{\times}$. Put $\tilde{\beta}_{ij} = \beta_{ij} e_i + \sum_{r \neq i} e_r$; then $\tilde{\beta}_{ij} \in \mathcal{A}^{\times}$ and

$$\alpha = \sum_{i,j} a_{ij} \tilde{\beta}_{ij}^2 e_i \in \mathcal{B}.$$

Lemma A.1.3. Let us assume $p \geq 5$. Then the normal closure of $SL_2(\mathbb{F}_p)$ in $SL_2(\mathcal{A})$ is $SL_2(\mathcal{A})$ itself.

Proof. We have

$$S(\alpha)S(\beta) = S(\alpha + \beta), \quad S(\alpha)^{-1} = S(-\alpha),$$

$$T(\alpha)T(\beta) = T(\alpha + \beta), \quad T(\alpha)^{-1} = T(-\alpha).$$

Let N be the normal closure of $\mathrm{SL}_2(\mathbb{F}_p)$ in $\mathrm{SL}_2(\mathcal{A})$. It suffices to prove that N contains all matrices $S(\alpha)$ and $T(\alpha)$ with α varying in a set of generators of \mathcal{A} as a \mathbb{F}_p -vector space. If $\alpha \in \mathcal{A}^{\times}$, then

$$\begin{pmatrix} \alpha & 0 \\ 0 & \alpha^{-1} \end{pmatrix} S(1) \begin{pmatrix} \alpha^{-1} & 0 \\ 0 & \alpha \end{pmatrix} = S(\alpha^2), \qquad \begin{pmatrix} \alpha & 0 \\ 0 & \alpha^{-1} \end{pmatrix} T(1) \begin{pmatrix} \alpha^{-1} & 0 \\ 0 & \alpha \end{pmatrix} = T(\alpha^2).$$

The claim now follows from Lemma A.1.2, since S(1), T(1) generate $SL_2(\mathbb{F}_p)$. \Box

Lemma A.1.4. Let us assume $p \geq 5$. Let $s = \frac{p-1}{(k-1,p-1)}$ and let H be a subgroup of $\widehat{G}(\mathbb{F}_p)$ such that |H| > 2s and such that the determinant maps surjectively onto $(\mathbb{F}_p^{\times})^{k-1}$. Then the normal closure of H in $\widehat{G}(\mathbb{F}_p)$ is $\widehat{G}(\mathbb{F}_p)$ itself.

Proof. Let N be the normal closure of H in $\widehat{G}(\mathbb{F}_p)$. Since |N| > 2s, the intersection $N \cap \operatorname{SL}_2(\mathbb{F}_p)$ is a non trivial normal subgroup of $\operatorname{SL}_2(\mathbb{F}_p)$ of order strictly greater than 2. Since $|\mathbb{F}| \ge 4$, the only proper normal subgroup of $SL_2(\mathbb{F})$ is $\{\pm Id\}$ (see for example [29, Theorem 8.3 and Lemma 8.6, p.539]). Thus $\operatorname{SL}_2(\mathbb{F}_p) \subseteq N$. Since the projection over $(\mathbb{F}_p^{\times})^{k-1}$ is surjective, the result follows. \Box

We can now prove Proposition A.1.1. Let N be the normal closure of H in $\widehat{G}(\mathcal{A})$. By Lemma A.1.4, N contains $\widehat{G}(\mathbb{F}_p)$, so that it contains $\mathrm{SL}_2(\mathcal{A})$ by Lemma A.1.3. The surjectivity of the determinant then shows that $N = \widehat{G}(\mathcal{A})$.

A.2. Irreducibility of the adjoint representation. We consider the adjoint representation of $SL_2(\mathcal{A})$, which is realized as its action by conjugation on the \mathbb{F}_p -vector space $M_2(\mathcal{A})^0$ of 2×2 matrices with coefficients in \mathcal{A} that have trace zero. We want to prove:

Proposition A.2.5. Let us assume $p \geq 3$. Let V be a \mathbb{F}_p -subspace of $M_2(\mathcal{A})^0$ such that $V \cap M_2(\mathbb{F}_p)^0 \neq \{0\}$ and invariant by the adjoint action of $SL_2(\mathcal{A})$. Then $V = M_2(\mathcal{A})^0$.

The following results are well known, we report it for convenience of the reader. Observe that $M_2(\mathbb{F}_p)^0$ is generated as an \mathbb{F}_p -vector space by the three matrices

$$A = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \qquad B = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \qquad C = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

Moreover if we put

$$X = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad Y = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \qquad Z = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix},$$

we see that

(A.1)
$$YAY^{-1} - A = -2B, \quad XBX^{-1} = -C, \quad YCY^{-1} - C + B = A.$$

Lemma A.2.6. Let us assume $p \geq 3$. Then the adjoint representation of $SL_2(\mathbb{F}_p)$ is irreducible.

Proof. The argument is standard. By (A.1), any invariant subspace containing one of A, B, C will also contain the others. Given some arbitrary (non-zero) matrix M = aA + bB + cC of trace 0, put

$$U = YMY^{-1} - M, \qquad V = ZMZ^{-1} - M.$$

Then

$$YUY^{-1} - U = -2cB, \qquad ZVZ^{-1} - V = -2bC.$$

It follows that any invariant \mathbb{F}_p -subspace of $M_2(\mathbb{F}_p)^0$ containing M must also contain one of A, B, C. The claim follows.

We can now prove Proposition A.2.5. Firstly we notice that $M_2(\mathbb{F}_p)^0 \subseteq V$ by Lemma A.2.6. Since $M_2(\mathcal{A})^0 = M_2(\mathbb{F}_p)^0 \otimes_{\mathbb{F}_p} \mathcal{A}$, it suffices to show that V is closed under the multiplication by elements in \mathcal{A} . Let α be an invertible element of \mathcal{A} and let $B = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. We have

$$\begin{pmatrix} \alpha & 0 \\ 0 & \alpha^{-1} \end{pmatrix} B \begin{pmatrix} \alpha^{-1} & 0 \\ 0 & \alpha \end{pmatrix} = \alpha^2 B.$$

Therefore $\alpha^2 B \in V$, and by Lemma A.1.2 we get that $\mathcal{A}B \subseteq V$. Then (A.1) proves that also $\mathcal{A}A$ and $\mathcal{A}C$ lie in V, so that $\mathcal{A}V \subseteq V$.

Acknowledgements

We are grateful to Andrea Conti, Samuel Le Fourn and Davide Lombardi for helpful discussions.

We thank the INdAM GNSAGA for support.

References

- F. Amoroso, S. David, and U. Zannier. On fields with Property (B). Proc. Amer. Math. Soc., 142(6):1893–1910, 2014.
- [2] F. Amoroso and R. Dvornicich. A lower bound for the height in abelian extensions. J. Number Theory, 80(2):260-272, 2000.
- [3] F. Amoroso and U. Zannier. A relative Dobrowolski lower bound over abelian extensions. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 29(3):711–727, 2000.
- [4] K. Arai. Galois images and modular curves. In Algebraic number theory and related topics 2010, volume B32 of RIMS Kôkyûroku Bessatsu, pages 145–161. Res. Inst. Math. Sci. (RIMS), Kyoto, 2012.
- [5] E. Bombieri and U. Zannier. A note on heights in certain infinite extensions of Q. Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl., 12:5–14 (2002), 2001.
- [6] C. Breuil. Sur quelques représentations modulaires et *p*-adiques de $GL_2(\mathbf{Q}_p)$. II. J. Inst. Math. Jussieu, 2(1):23–58, 2003.
- [7] F. Calegari and N. Talebizadeh Sardari. Vanishing Fourier coefficients of Hecke eigenforms. Math. Ann., 381(3-4):1197–1215, 2021.

- [8] H. Carayol. Formes modulaires et représentations galoisiennes à valeurs dans un anneau local complet. In p-adic monodromy and the Birch and Swinnerton-Dyer conjecture (Boston, MA), volume 165 of Contemp. Math., pages 213–237. 1991.
- [9] S. Checcoli. Fields of algebraic numbers with bounded local degrees and their properties. Trans. Amer. Math. Soc., 365(4):2223–2240, 2013.
- [10] P. Colmez. Périodes des variétés abéliennes à multiplication complexe. Ann. of Math. (2), 138(3):625–683, 1993.
- [11] G. Faltings. Hodge-Tate structures and modular forms. Math. Ann., 278(1-4):133–149, 1987.
- [12] L. Frey. Lower Height Bounds in Non-Abelian Extensions. Phd thesis, University of Basel, 2018.
- [13] L. Frey. Explicit small heights in infinite non-abelian extensions. Acta Arith., 199(2):111–133, 2021.
- [14] E. Ghate and P. Parent. On uniform large Galois images for modular abelian varieties. Bull. Lond. Math. Soc., 44(6):1169–1181, 2012.
- [15] P. Habegger. Small height and infinite nonabelian extensions. Duke Math. J., 162(11):2027– 2076, 2013.
- [16] A. J. Hahn and O. T. O'Meara. The classical groups and K-theory, volume 291 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1989. With a foreword by J. Dieudonné.
- [17] C. Khare and J.-P. Wintenberger. Serre's modularity conjecture. I. Invent. Math., 178(3):485– 504, 2009.
- [18] D. Lombardo. Explicit surjectivity of Galois representations for abelian surfaces and GL₂varieties. J. Algebra, 460:26–59, 2016.
- [19] J. Neukirch. Algebraic number theory, volume 322 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1999.
- [20] K. A. Ribet. Galois representations attached to eigenforms with Nebentypus. In Modular functions of one variable, V (Proc. Second Internat. Conf., Univ. Bonn, Bonn, 1976), volume Vol. 601 of Lecture Notes in Math., pages 17–51. Springer, Berlin-New York, 1977.
- [21] K. A. Ribet. Twists of modular forms and endomorphisms of abelian varieties. Math. Ann., 253(1):43–62, 1980.
- [22] K. A. Ribet. On *l*-adic representations attached to modular forms. II. Glasgow Math. J., 27:185–194, 1985.
- [23] K. A. Ribet. Images of semistable Galois representations. Number Special Issue, pages 277– 297. 1997. Olga Taussky-Todd: in memoriam.
- [24] A. Schinzel. Addendum to the paper: "On the product of the conjugates outside the unit circle of an algebraic number" (Acta Arith. 24 (1973), 385–399). Acta Arith., 26(3):329–331, 1974/75.
- [25] A. J. Scholl. Motives for modular forms. Invent. Math., 100(2):419–430, 1990.
- [26] J.-P. Serre. Corps locaux, volume No. VIII of Publications de l'Université de Nancago. Hermann, Paris, 1968. Deuxième édition.
- [27] J.-P. Serre. Propriétés galoisiennes des points d'ordre fini des courbes elliptiques. Invent. Math., 15(4):259–331, 1972.
- [28] G. Shimura. On the factors of the Jacobian variety of a modular function field. J. Math. Soc. Japan, 25:523–544, 1973.
- [29] The LMFDB Collaboration. The L-functions and modular forms database. https://www.lmfdb.org, 2024.