

# SPECTRUM OF RANDOM CENTROSYMMETRIC MATRICES; CLT AND CIRCULAR LAW

INDRAJIT JANA AND SUNITA RANI

**ABSTRACT.** We analyze the asymptotic fluctuations of linear eigenvalue statistics of random centrosymmetric matrices with i.i.d. entries. We prove that for a complex analytic test function, the centered and normalized linear eigenvalue statistics of random centrosymmetric matrices converge to a normal distribution. We find the exact expression of the variance of the limiting normal distribution via combinatorial arguments. Moreover, we also argue that the limiting spectral distribution of properly scaled centrosymmetric matrices follows the circular law.

keywords: Centrosymmetric matrix, Linear eigenvalue statistics, Central limit theorem, Circular law.

## 1. INTRODUCTION

Let  $M$  be an  $n \times n$  matrix with real or complex entries. The empirical spectral distribution (ESD) is defined by the measure

$$\mu_n(\cdot) = \frac{1}{n} \sum_{i=1}^n \delta_{\lambda_i}(\cdot),$$

where  $\lambda_1, \lambda_2, \dots, \lambda_n$  are the eigenvalues of the matrix  $M$ , and  $\delta_x(\cdot)$  is the point measure at  $x \in \mathbb{C}$ . The linear eigenvalue statistics (LES) of  $M$  corresponding to a test function  $f$  is defined by

$$L(f) = \sum_{k=1}^n f(\lambda_k).$$

The function  $f$  is referred to as the test function. The asymptotic behaviour of  $\mu_n(\cdot)$  and  $a_n(L(f) - b_n)$  for some sequences  $\langle a_n \rangle$  and  $\langle b_n \rangle$  have been studied in depth for different types of random matrices. We give a brief overview below.

Jonsson [20] established the central limit theorem (CLT) for LES in Wishart matrices. Subsequently, researchers extensively studied the fluctuation of eigenvalues of various random matrices. Some of the key contributions include Johansson [19], Sinai and Soshnikov [32], Bai and Silverstein [5], Lytova and Pastur [24], Shcherbina [29]. Furthermore, the same has been studied for several structured matrices, such as band and sparse matrices [4, 22, 30, 18], Toeplitz and band Toeplitz matrices [10, 23], circulant matrices [1, 2, 25].

The CLT of LES for non-Hermitian matrices have been studied in different set ups as well; such as when the test function is analytic [27, 26, 17], the test function is non-analytic but the moments of the matrix entries match with that of a Ginibre ensemble up to fourth

---

Indrajit Jana's research is partially supported by INSPIRE Fellowship DST/INSPIRE/04/2019/000015, Dept. of Science and Technology, Govt. of India.

Sunita Rani's research is fully supported by the University Grant Commission (UGC), New Delhi.

moment [21]. Later, it was generalized for continuous bounded test functions [13, 11, 12]. A more comprehensive list of recent results can be found on the review article by Forrester [16].

In this article, we study the spectral properties of random centrosymmetric matrices. A centrosymmetric matrix is symmetric around its center which is formally defined in the Definition 3.1. Centrosymmetric matrices appeared in many different contexts in mathematics, such as solutions to chessboard separation problems [9], stochastic matrices [8], and in various other contexts [15, 33, 28, 6]. However, to the best of our knowledge the spectral properties of random centrosymmetric matrices have not been studied yet. In this paper, we investigate the limiting ESD of a random centrosymmetric matrix and the fluctuations of the LES.

## 2. NOTATIONS AND LAYOUT

This article is organized as follows. We state the main theorem in Section 3. The proof of the CLT is shown in Section 6, and the variance calculation of the limiting normal distribution is delegated to Section 7. In addition, it was also argued that the limiting spectral distribution follows the circular law, which was demonstrated in Section 5.

Throughout this article, the identity matrix is denoted by  $I$ , and the counter-identity matrix is denoted by  $J$ , as defined in Definition 4.1. The dimensions of such matrices should be understood from the context. The eigenvalues of an  $n \times n$  matrix  $M$  are denoted by  $\lambda_1(M), \lambda_2(M), \dots, \lambda_n(M)$ . The complex conjugate transpose of a matrix  $Y$  is denoted by  $Y^*$ , whereas  $\bar{Y}$  stands for only the complex conjugate of  $Y$  without taking transpose. The notation  $\mathcal{R}_z(M) = (zI - M)^{-1}$  represents the resolvent of a square matrix  $M$ . The notations  $\zeta_n \xrightarrow{d} \zeta$ ,  $\zeta_n \xrightarrow{p} \zeta$ , and  $\zeta_n \xrightarrow{a.s.} \zeta$  are used to indicate that the sequence of random variables  $\{\zeta_n\}_n$  converges to  $\zeta$  in distribution, in probability, and almost surely, respectively.

For any random variable  $\xi$ ,  $\xi^\circ$  denotes the centered random variable  $\xi - \mathbb{E}[\xi]$ . Moreover, if  $\xi$  depends on a random matrix, then  $\xi_k^\circ := \xi - \Xi_k[\xi]$ , where  $\Xi_k[\cdot]$  signifies averaging with respect to the  $k$ -th column of the underlying random matrix.

The vectors  $e_1, e_2, \dots, e_n$  denote the canonical basis vectors of  $\mathbb{R}^n$ . The disc of radius  $r$ , denoted by  $\mathbb{D}_r$ , is the set  $\{z \in \mathbb{C} : |z| \leq r\}$ , and its boundary is denoted by  $\partial\mathbb{D}_r$ . The notations  $K$  or  $C$  are used to denote a universal constant whose exact value may change from line to line.

## 3. MAIN RESULT

Let us first define the centrosymmetric matrix.

**Definition 3.1** (Centrosymmetric Matrix). A random Centrosymmetric matrix is a random matrix which is symmetric about its center. Formally, let  $M = [m_{i,j}]_{n \times n}$  be a random matrix, where  $m_{ij}$  are i.i.d. random variables subject to the following constraint

$$m_{i,j} = m_{n-i+1, n-j+1} \text{ for } i, j \in \{1, \dots, n\}.$$

Then  $M$  is called a random centrosymmetric matrix.

**Definition 3.2** (Poincaré inequality). A complex random variable  $\zeta$  is said to satisfy the Poincaré inequality with a positive constant  $\alpha$  if for any differentiable function  $g : \mathbb{C} \rightarrow \mathbb{C}$ , we have  $\text{Var}(g(\zeta)) \leq \frac{1}{\alpha} \mathbb{E}[|g'(\zeta)|^2]$ . Here,  $\mathbb{C}$  is identified with  $\mathbb{R}^2$  and  $|g'(\zeta)|$  represents the




FIGURE 1. Symmetry pattern of a centrosymmetric  $5 \times 5$  matrix.

$\ell^2$  norm of the two dimensional vector  $g'$ . Here are some characteristics of the Poincaré inequality.

- (1) If  $\zeta$  satisfies the Poincaré inequality with constant  $\alpha$ , then  $\kappa\zeta$  also satisfies the Poincaré inequality with constant  $\alpha/\kappa^2$  for any nonzero real constant  $\kappa$ .
- (2) If two independent random variables  $\zeta_1, \zeta_2$  satisfy the Poincaré inequality with the same constant  $\alpha$ , then  $\zeta = (\zeta_1, \zeta_2)$  also satisfies the Poincaré inequality with the same constant  $\alpha$ . Here  $\zeta$  is treated as a random vector and, accordingly,  $g'$  in the definition of Poincaré inequality will be replaced by  $\nabla g$ .
- (3) [3, Lemma 4.4.3] If  $\zeta \in \mathbb{C}^n$  satisfies the Poincaré inequality with a constant  $\alpha$ , then for any differentiable function  $g : \mathbb{C}^n \rightarrow \mathbb{C}$ ,

$$\mathbb{P}(|g(\zeta) - \mathbb{E}[g(\zeta)]| > t) \leq C \exp \left\{ -\frac{\sqrt{\alpha}t}{\sqrt{2} \|\|\nabla g\|_2\|_\infty} \right\},$$

where  $C = -2 \sum_{k=1}^{\infty} 2^k \log(1 - 2^{-2k-1})$ . Here  $\mathbb{C}^n$  is identified with  $\mathbb{R}^{2n}$ . Moreover,  $\|\nabla g(x)\|_2$  is the  $\ell^2$  norm of the  $2n$  dimensional vector  $\nabla g(x)$  at  $x \in \mathbb{C}^n$ , and  $\|\|\nabla g\|_2\|_\infty = \sup_{x \in \mathbb{C}^n} \|\nabla g(x)\|_2$ .

For instance, a Gaussian random variable satisfies the Poincaré inequality.

**Condition 3.3.** Let  $M = [m_{ij}]_{n \times n}$  be a centrosymmetric matrix, where

$$m_{ij} = \frac{1}{\sqrt{n}} x_{ij}.$$

Assume that  $\{x_{ij} : 1 \leq i, j \leq n\}$  are i.i.d. complex random variables subject to the constraint  $x_{ij} = x_{n+1-i, n+1-j}$ . In addition,  $x_{ij}$ s satisfy the following conditions.

- (i)  $\mathbb{E}[x_{ij}] = 0$ ,  $\mathbb{E}[x_{ij}^2] = 0$  and  $\mathbb{E}[|x_{ij}|^2] = 1$  for all  $1 \leq i, j \leq n$ ,
- (ii)  $x_{ij}$ s are continuous random variables with bounded density and satisfy the Poincaré inequality with some universal constant  $\alpha$ .

Before stating our main theorem, let us define the centered LES as

$$L_f^\circ(M) = \sum_{k=1}^n f(\lambda_k(M)) - \sum_{k=1}^n \mathbb{E}[f(\lambda_k(M))].$$

Our main theorem is stated as follows.

**Theorem 3.4.** *Suppose  $M$  is a centrosymmetric random matrix satisfying Condition 3.3. Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be an analytic function. Then  $L_f^\circ(M) \xrightarrow{d} N(0, \sigma_f^2)$ , where*

$$\sigma_f^2 = \frac{2}{\pi} \int_{\mathbb{D}_1} f(w) \overline{f'(w)} d\Re(w) d\Im(w).$$

*In particular, if  $f$  is a polynomial given by  $P_d(z) = \sum_{k=0}^d a_k z^k$ , then  $\sigma_f^2 = \sum_{k=1}^d 2k |a_k|^2$ .*

#### 4. TRUNCATION AND REDUCTION

Before proceeding to the technical proof of the main theorem, we need to recall one important theorem regarding centrosymmetric matrices.

**Definition 4.1** (Counter-identity Matrix). It is defined as the square matrix whose counter-diagonal elements are 1 and all the other elements are zero.

Throughout this article, we shall denote the counter-identity matrix by  $J$ . The size of  $J$  should be understood from the context. Notice that

$$J = J^T, \quad J^2 = I,$$

where  $I$  the identity matrix of the same dimension. Now, we invoke the following theorem, which reduces the centrosymmetric matrix to a block diagonal matrix.

**Theorem 4.2.** [35, Theorem 9]

(a) *If  $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$  is an  $n \times n$  centrosymmetric matrix with  $n = 2s$  and  $A, B, C$ , and  $D$  are  $s \times s$  matrices, then  $M$  is orthogonally similar to*

$$Q^T M Q = \begin{pmatrix} A + JC & 0 \\ 0 & A - JC \end{pmatrix},$$

where

$$Q = \sqrt{\frac{1}{2}} \begin{pmatrix} I & -J \\ J & I \end{pmatrix}.$$

(b) *If  $M = \begin{pmatrix} A & x & B \\ y & q & yJ \\ C & Jx & D \end{pmatrix}$  is an  $n \times n$  centrosymmetric matrix with  $n = 2s + 1$ , and  $A, B, C, D$  are  $s \times s$  matrices,  $x$  is  $s \times 1$  matrix,  $y$  is  $1 \times s$  matrix, and  $q$  is a scalar, then  $M$  is orthogonally similar to*

$$Q^T M Q = \begin{pmatrix} A + JC & \sqrt{2}x & 0 \\ \sqrt{2}y & q & 0 \\ 0 & 0 & A - JC \end{pmatrix},$$

where

$$Q = \sqrt{\frac{1}{2}} \begin{pmatrix} I & 0 & -J \\ 0 & \sqrt{2} & 0 \\ J & 0 & I \end{pmatrix}.$$

*In the definitions of  $Q$  above,  $I$  is an  $s \times s$  identity matrix and  $J$  is an  $s \times s$  counter-identity matrix. Moreover, notice that in both the cases  $Q^T Q = I = Q Q^T$ .*

Since the  $Q^T M Q$  is orthogonally similar to the matrix  $M$ , the eigenvalues of  $Q^T M Q$  are the same as those of  $M$ . The remaining part of this article analyses the eigenvalues of  $Q^T M Q$ . For convenience, we adopt the following simplified notation for  $Q^T M Q$ .

$$Q^T M Q = \begin{bmatrix} T_1 & 0 \\ 0 & T_2 \end{bmatrix},$$

where  $T_2 = A - JC$  and

$$T_1 = \begin{cases} A + JC & \text{if } n = 2s \\ \begin{pmatrix} A + JC & \sqrt{2}x \\ \sqrt{2}y & q \end{pmatrix} & \text{if } n = 2s + 1. \end{cases}$$

Though the above theorem reduces the centrosymmetric matrix to a block diagonal matrix, the block matrices are not independent. Therefore, we need to take care of the dependence structure in the course of the proof.

## 5. CIRCULAR LAW FOR RANDOM CENTROSYMMETRIC MATRIX

Let us first state the general circular law.

**Theorem 5.1** (Circular Law, [34]). *Consider a random matrix  $M = \frac{1}{\sqrt{n}}[x_{ij}]_{i,j=1}^n$ , where  $x_{ij}$ s are i.i.d. random variables with mean zero and variance 1. Then, as  $n \rightarrow \infty$ , the ESD of  $M$  converges almost surely to the uniform distribution on the unit disc of  $\mathbb{C}$  with density  $\frac{1}{\pi}1_{\{z \in \mathbb{C} \mid |z| \leq 1\}}$ .*

Now, let us state the circular law for random centrosymmetric matrix.

**Theorem 5.2** (Circular Law for Random Centrosymmetric Matrix). *Consider a random centrosymmetric matrix  $M = \frac{1}{\sqrt{n}}[x_{ij}]_{i,j=1}^n$ , where  $x_{ij}$ s are i.i.d. random variables for  $i \leq j$  with mean zero and variance 1. Then, as  $n \rightarrow \infty$ , the ESD of  $M$  converges almost surely to the uniform distribution on the unit disc of  $\mathbb{C}$  with density  $\frac{1}{\pi}1_{\{z \in \mathbb{C} \mid |z| \leq 1\}}$ .*




FIGURE 2. Scatter plot of real and imaginary parts of the eigenvalues of a  $5000 \times 5000$  random centrosymmetric matrix having standard Gaussian entries.

*Proof.* We have a centrosymmetric matrix  $M = \frac{1}{\sqrt{n}}[x_{ij}]_{i,j=1}^n$ , where  $x_{ij}$ s are i.i.d. centered random variables with variance 1.

From Theorem 4.2, it is evident that the eigenvalues of  $M$  are the eigenvalues of  $T_1$  and  $T_2$ . The entries of  $T_1$  and  $T_2$  are i.i.d. centered random variables having variance 2. However, sizes of  $T_1$  and  $T_2$  is  $\lceil n/2 \rceil \times \lceil n/2 \rceil$ , and they are scaled by  $1/\sqrt{n}$ . Hence, by Theorem 5.1 the ESD of  $T_1$  converges to the circular law. The same applies to  $T_2$  as well. Hence, ESD of the matrix  $M$  converges to the circular law.  $\square$

## 6. PROOF OF THE CENTRAL LIMIT THEOREM

In Section 5, we have seen that the limiting ESD follows the circular law. However, for any finite  $n$ , there might be some eigenvalues outside the unit disc. To avoid such situations, we shall work on the event where the eigenvalues are confined within a disc of finite radius. Keeping this in mind, we define the event

$$(6.1) \quad \Omega_n = \{\|T_1\| \leq \rho_1\} \cap \{\|T_2\| \leq \rho_2\},$$

where  $\rho_1$  and  $\rho_2$  are same as  $\rho$ , given in Lemma A.2. Additionally, Lemma A.2 also implies that

$$(6.2) \quad P(\Omega_n^c) \leq K \exp\left(-\sqrt{\frac{\alpha n}{2}} \frac{3\rho_m}{4}\right), \quad \text{where } \rho_m = \min\{\rho_1, \rho_2\}.$$

Let us denote  $\rho = \max\{\rho_1, \rho_2\}$ . If  $f$  is a complex analytic function on  $\partial\mathbb{D}_{\rho+\tau}$ , then we can express  $L_n(f)$  on the event  $\Omega_n$  as follows

$$\begin{aligned} L_f(M) &= \sum_{i=1}^n f(\lambda_i(M)) \\ &= \frac{1}{2\pi i} \oint_{\partial\mathbb{D}_{\rho+\tau}} \sum_{i=1}^n \frac{f(z)}{z - \lambda_i(M)} dz \\ &= \frac{1}{2\pi i} \oint_{\partial\mathbb{D}_{\rho+\tau}} f(z) \operatorname{Tr} \mathcal{R}_z(M) dz. \end{aligned}$$

The above represents  $L_f(M)$  in terms of the resolvent of the matrix. This brings down our task to the analysis of the resolvent. However, to keep the norm of the resolvent bounded, we need to restrict our attention to the event  $\Omega_n$ , where norms of  $T_1, T_2$  are bounded. From the probability estimate (6.2) and the Lemma A.2, we may express the centered linear eigenvalue statistic as follows.

$$L_f^\circ(M) = \frac{1}{2\pi i} \oint_{\partial\mathbb{D}_{\rho+\tau}} f(z) \operatorname{Tr} \hat{\mathcal{R}}_z^\circ(M) dz + o(1).$$

where  $\hat{\xi} = \xi 1_{\Omega_n}$ , and  $\xi^\circ = \xi - \mathbb{E}[\xi]$ . The  $o(1)$  at the end of the above expression can further be explained as follows.

We can rewrite

$$\operatorname{Tr} \mathcal{R}_z^\circ(M) = \operatorname{Tr} \hat{\mathcal{R}}_z^\circ(M) + \operatorname{Tr} \mathcal{R}_z(M) 1_{\Omega_n^c} - \mathbb{E}[\operatorname{Tr} \mathcal{R}_z(M) 1_{\Omega_n^c}].$$

We will demonstrate that  $\operatorname{Tr} \mathcal{R}_z(M) 1_{\Omega_n^c} - \mathbb{E}[\operatorname{Tr} \mathcal{R}_z(M) 1_{\Omega_n^c}] = o(1)$ . First, let us prove that  $\operatorname{Tr} \mathcal{R}_z(M) 1_{\Omega_n^c} = o(1)$ . Since we are dealing with continuous random variables, almost surely

there is no eigenvalue on the circle  $\partial\mathbb{D}_{\rho+\tau}$ . Therefore,  $\sup_{z \in \partial\mathbb{D}_{\rho+\tau}} |\text{Tr } \mathcal{R}_z(M)|$  is bounded almost surely. Consequently, we conclude

$$\mathbb{P} \left( \sup_{z \in \partial\mathbb{D}_{\rho+\tau}} |\text{Tr } \mathcal{R}_z(M) \mathbf{1}_{\Omega_n^c}| > \epsilon \right) \leq \mathbb{P}(\Omega_n^c) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Next, we prove that  $\mathbb{E}[\text{Tr } \mathcal{R}_z(M) \mathbf{1}_{\Omega_n^c}] \rightarrow 0$  asymptotically. We note that the assumptions listed in Condition 3.3 are consistent with the assumptions of the Lemma A.3. Hence, using the Lemma A.3, and the bound (6.2) we may conclude that

$$\begin{aligned} \mathbb{E}[\text{Tr } \mathcal{R}_z(M) \mathbf{1}_{\Omega_n^c}] &\leq \mathbb{E}[|\text{Tr } \mathcal{R}_z(M)|] \mathbb{P}(\Omega_n^c) \\ &\leq Cn^{7/2} \exp \left( -\sqrt{\frac{\alpha n}{2}} \frac{3\rho_m}{4} \right) \\ &\rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

Furthermore, standard use of complex analysis shows that

$$\begin{aligned} &\frac{1}{\pi} \int_{\mathbb{D}_1} f'(w) \overline{f'(w)} d\Re(w) d\Im(w) \\ &= \frac{1}{\pi} \int_{\mathbb{D}_1} \left[ \frac{1}{2\pi i} \oint_{\partial\mathbb{D}_1} \frac{f(z) dz}{(w-z)^2} \frac{1}{2\pi i} \oint_{\partial\mathbb{D}_1} \frac{f(\bar{\eta}) d\bar{\eta}}{(\bar{w}-\bar{\eta})^2} \right] d\Re(w) d\Im(w) \\ &= -\frac{1}{4\pi^2} \oint_{\partial\mathbb{D}_1} \oint_{\partial\mathbb{D}_1} f(z) \overline{f(\eta)} \left[ \frac{1}{\pi} \int_{\mathbb{D}_1} \frac{d\Re(w) d\Im(w)}{(w-z)^2 (\bar{w}-\bar{\eta})^2} \right] dz d\bar{\eta} \\ &= -\frac{1}{4\pi^2} \oint_{\partial\mathbb{D}_1} \oint_{\partial\mathbb{D}_1} \frac{f(z) \overline{f(\eta)}}{(1-z\bar{\eta})^2} dz d\bar{\eta}. \end{aligned}$$

Hence, the main Theorem 3.4 is essentially the same as proving Proposition 6.1.

**Proposition 6.1.** *The sequence  $\{\text{Tr } \hat{\mathcal{R}}_z^\circ(M)\}_n$  is tight in the space  $\mathcal{C}(\partial\mathbb{D}_{\rho+\tau})$ , which is the space of continuous function on  $\partial\mathbb{D}_{\rho+\tau}$ , and converges in distribution to a Gaussian process with covariance kernel given by  $2(1-z\bar{\eta})^{-2}$ .*

Our first goal is to reduce the above proposition to (6.5), (6.6), (6.7). First of all, by Cramer-Wold device, to show finite dimensional convergence it is sufficient to show that for any  $\{z_l; 1 \leq l \leq q\} \subset \partial\mathbb{D}_{\rho+\tau}$ , and  $\{\alpha_l, \beta_l; 1 \leq l \leq q\} \subset \mathbb{C}$  for which

$$(6.3) \quad \sum_{l=1}^q \left[ \alpha_l \text{Tr } \hat{\mathcal{R}}_{z_l}^\circ(M) + \beta_l \overline{\text{Tr } \hat{\mathcal{R}}_{z_l}^\circ(M)} \right] \mathbf{1}_{\Omega_n}$$

is real and converges to a Gaussian random variable with mean zero. In order to show this, we use the martingale CLT, which is stated in Lemma A.4. For notational convenience, we define the relevant conditional expectations with respect to the columns of the matrices  $T_1$  and  $T_2$ . Let  $\Xi_k[\xi]$  be the averaging of the random variable  $\xi$  over the  $k$ -th columns of matrices  $T_1$  and  $T_2$ , and  $\mathbb{E}_k[\cdot] = \Xi_{k+1} \Xi_{k+2} \dots \Xi_{\lfloor \frac{n}{2} \rfloor}[\cdot] = \Xi_{k+1} \mathbb{E}_{k+1}[\cdot]$ . For instance,  $\mathbb{E}_0[\cdot] = \mathbb{E}[\cdot]$  and  $\mathbb{E}_{\lfloor \frac{n}{2} \rfloor}[\cdot] = [\cdot]$ . Therefore, we can express

$$\begin{aligned} \text{Tr } \hat{\mathcal{R}}_z^\circ(M) &= \text{Tr } \hat{\mathcal{R}}_z^\circ(T_1) + \text{Tr } \hat{\mathcal{R}}_z^\circ(T_2) \\ &= \text{Tr } \hat{\mathcal{R}}_z(T_1) - \mathbb{E}[\text{Tr } \hat{\mathcal{R}}_z(T_1)] + \text{Tr } \hat{\mathcal{R}}_z(T_2) - \mathbb{E}[\text{Tr } \hat{\mathcal{R}}_z(T_2)] \end{aligned}$$

$$\begin{aligned}
&= \sum_{k=1}^{\lceil n/2 \rceil} (\mathbb{E}_k - \mathbb{E}_{k-1}) \operatorname{Tr} \hat{\mathcal{R}}_z(T_1) + \sum_{k=1}^{\lceil n/2 \rceil} (\mathbb{E}_k - \mathbb{E}_{k-1}) \operatorname{Tr} \hat{\mathcal{R}}_z(T_2) \\
&= \sum_{k=1}^{\lceil n/2 \rceil} \mathbb{E}_k \left\{ \left( \operatorname{Tr} \hat{\mathcal{R}}_z(T_1) \right)_k^\circ \right\} + \sum_{k=1}^{\lceil n/2 \rceil} \mathbb{E}_k \left\{ \left( \operatorname{Tr} \hat{\mathcal{R}}_z(T_2) \right)_k^\circ \right\} \\
&=: \sum_{k=1}^{\lceil n/2 \rceil} W_{z,k}(T_1) + \sum_{k=1}^{\lceil n/2 \rceil} W_{z,k}(T_2) \\
(6.4) \quad &= \sum_{k=1}^{\lceil n/2 \rceil} (W_{z,k}(T_1) + W_{z,k}(T_2)),
\end{aligned}$$

where  $\xi_k^\circ = \xi - \Xi_k[\xi]$ . Since  $\mathbb{E}_k[\cdot]$  is defined as the conditional expectation given the first  $k$  columns of  $T_1$  and  $T_2$ ,  $(\mathbb{E}_k - \mathbb{E}_{k-1})[\xi]$  forms a martingale difference sequence. Consequently, the sequences  $\{W_{z,k}(T_1)\}_{1 \leq k \leq \lceil n/2 \rceil}$  and  $\{W_{z,k}(T_2)\}_{1 \leq k \leq \lceil n/2 \rceil}$  are martingale difference sequences with respect to the filtration  $\mathcal{F}_{n,k} = \sigma\{t_j^1, t_j^2 : 1 \leq j \leq k\}$ , where  $t_j^1$  and  $t_j^2$  are the  $j$ th columns of matrices  $T_1$  and  $T_2$  respectively. Therefore, the sum  $W_{z,k}(T_1) + W_{z,k}(T_2)$  also forms a martingale difference sequence with respect to the same filtration  $\mathcal{F}_{n,k}$ .

Let us rewrite

$$\begin{aligned}
&\sum_{i=1}^q \left[ \alpha_i \operatorname{Tr} \hat{\mathcal{R}}_{z_i}^\circ(M) + \beta_i \overline{\operatorname{Tr} \hat{\mathcal{R}}_{z_i}^\circ(M)} \right] \\
&= \sum_{k=1}^{\lceil n/2 \rceil} \left\{ \sum_{i=1}^q (\alpha_i (W_{z_i,k}(T_1) + W_{z_i,k}(T_2)) + \beta_i (W_{\bar{z}_i,k}(T_1^*) + W_{\bar{z}_i,k}(T_2^*))) \right\} \\
&= \sum_{k=1}^{\lceil n/2 \rceil} \left\{ \sum_{i=1}^q \alpha_i W_{z_i,k}(T_1) + \beta_i W_{\bar{z}_i,k}(T_1^*) + \sum_{i=1}^q \alpha_i W_{z_i,k}(T_2) + \beta_i W_{\bar{z}_i,k}(T_2^*) \right\} \\
&=: \sum_{k=1}^{\lceil n/2 \rceil} (\psi_{n,k}^{(1)} + \psi_{n,k}^{(2)}) \\
&=: \sum_{k=1}^{\lceil n/2 \rceil} \psi_{n,k},
\end{aligned}$$

where  $X^*$  denotes the complex conjugate of the matrix  $X$ . As a result,  $\{\psi_{n,k}\}_{1 \leq k \leq \lceil n/2 \rceil}$  constitutes a martingale difference sequence with respect to the filtration  $\mathcal{F}_{n,k}$ . We notice that proving the following equation (6.5) is sufficient for the validity of the condition (i) of Lemma A.4;

$$(6.5) \quad \sum_{k=1}^{\lceil n/2 \rceil} \mathbb{E} \left[ \psi_{n,k}^2 \mathbf{1}_{|\psi_{n,k}| > \theta} \right] \leq \frac{1}{\theta^2} \sum_{k=1}^{\lceil n/2 \rceil} \mathbb{E} [\psi_{n,k}^4] \rightarrow 0.$$

Since  $\mathbb{E}[\cdot | \mathcal{F}_{n,k-1}] = \mathbb{E}_{k-1}[\cdot]$ , the condition (ii) of Lemma A.4 can be expressed as



$$(6.6) \quad \sum_{k=1}^{\lceil n/2 \rceil} \mathbb{E}_{k-1}[(W_{z_i,k}(T_1) + W_{z_i,k}(T_2))(W_{z_j,k}(T_1) + W_{z_j,k}(T_2))] \xrightarrow{P} 0,$$

and

$$(6.7) \quad \mathbb{E}_{k-1}[(W_{z_i,k}(T_1) + W_{z_i,k}(T_2))(W_{\bar{z}_j,k}(T_1^*) + W_{\bar{z}_j,k}(T_2^*))] \xrightarrow{P} \sigma(z_i, z_j),$$

where  $\sigma$  is a covariance kernel. We defer the calculation of the covariance kernel to Section 7.4. Hence, establishing Proposition 6.1 is equivalent to demonstrating the tightness of the sequence  $\{\text{Tr } \hat{\mathcal{R}}_z^\circ(T_1) + \text{Tr } \hat{\mathcal{R}}_z^\circ(T_2)\}_n$  and verifying the conditions outlined in (6.5), (6.6), and (6.7). Before proceeding any further, let us do some reductions.

For  $i = 1, 2$ , let  $T_i^{(k)}$  be the matrix obtained by zeroing out the  $k$ th column of  $T_i$ , and let  $\hat{\mathcal{R}}_z(T_i^{(k)}) := \mathcal{R}_z(T_i^{(k)})\mathbf{1}_{\Omega_{n,k}}$ , where  $\Omega_{n,k} = \left\{ \left\| T_1^{(k)} \right\| \leq \rho_1 \right\} \cap \left\{ \left\| T_2^{(k)} \right\| \leq \rho_2 \right\}$ . The space  $\Omega_{n,k}$  is different from  $\Omega_n$ , but asymptotically both the spaces are the same. In fact, we can check that

$$\begin{aligned} & \mathbb{E} \left| \sum_{i=1}^{\lceil n/2 \rceil} \Xi_i \left[ \text{Tr } \mathcal{R}_z(T_1^{(k)})\mathbf{1}_{\Omega_n} - \text{Tr } \mathcal{R}_z(T_1^{(k)})\mathbf{1}_{\Omega_{n,i}} \right]_i \right| \\ & \leq 2\mathbb{E} \left| \sum_{i=1}^{\lceil n/2 \rceil} \Xi_i \left[ \text{Tr } \mathcal{R}_z(T_1^{(k)})\mathbf{1}_{\Omega_n^c \cap \Omega_{n,i}} \right] \right| \\ & \leq \frac{n^2}{\tau} \mathbb{P}(\Omega_n^c) = o(1). \end{aligned}$$

The same is true for  $T_2^{(k)}$  as well. The important difference between  $\Omega_n$  and  $\Omega_{n,k}$  is that the latter is independent of the  $k$ -th columns of  $T_1$  and  $T_2$ , which will allow us to decouple some expectation calculations in future. Consequently, we shall proceed with the space  $\Omega_{n,k}$ .

Using Lemma A.5 and the resolvent identity, we have

$$\begin{aligned} \text{Tr } \hat{\mathcal{R}}_z(T_1) &= \text{Tr } \hat{\mathcal{R}}_z(T_1^{(k)}) + e_k^t \hat{\mathcal{R}}_z(T_1^{(k)}) \mathcal{R}_z(T_1) t_k^1 \\ &= \text{Tr } \hat{\mathcal{R}}_z(T_1^{(k)}) + \frac{e_k^t (\hat{\mathcal{R}}_z(T_1^{(k)}))^2 t_k^1}{1 + e_k^t \hat{\mathcal{R}}_z(T_1^{(k)}) t_k^1}. \end{aligned}$$

Similarly,

$$\text{Tr } \hat{\mathcal{R}}_z(T_2) = \text{Tr } \hat{\mathcal{R}}_z(T_2^{(k)}) + \frac{e_k^t (\hat{\mathcal{R}}_z(T_2^{(k)}))^2 t_k^2}{1 + e_k^t \hat{\mathcal{R}}_z(T_2^{(k)}) t_k^2},$$

where  $t_k^1$  and  $t_k^2$  are the  $k$ th columns of  $T_1$  and  $T_2$  respectively. Thus,

$$\begin{aligned} & \text{Tr } \hat{\mathcal{R}}_z(M) \\ &= \text{Tr } \hat{\mathcal{R}}_z(T_1^{(k)}) + \text{Tr } \hat{\mathcal{R}}_z(T_2^{(k)}) + \frac{e_k^t (\hat{\mathcal{R}}_z(T_1^{(k)}))^2 t_k^1}{1 + e_k^t \hat{\mathcal{R}}_z(T_1^{(k)}) t_k^1} + \frac{e_k^t (\hat{\mathcal{R}}_z(T_2^{(k)}))^2 t_k^2}{1 + e_k^t \hat{\mathcal{R}}_z(T_2^{(k)}) t_k^2} \\ &= \text{Tr } \hat{\mathcal{R}}_z(T_1^{(k)}) + \text{Tr } \hat{\mathcal{R}}_z(T_2^{(k)}) - \frac{\partial}{\partial z} \log(1 + \gamma_{1k}(z)) - \frac{\partial}{\partial z} \log(1 + \gamma_{2k}(z)), \end{aligned}$$

where  $\gamma_{1k}(z) := e_k^t \hat{\mathcal{R}}_z(T_1^{(k)}) t_k^1$  and  $\gamma_{2k}(z) := e_k^t \hat{\mathcal{R}}_z(T_2^{(k)}) t_k^2$ . Now, let us find the tail bounds of  $\gamma_{1k}(z)$  and  $\gamma_{2k}(z)$ . Note that  $\gamma_{1k}(z)$  is product of two independent random variables. Intuitively,  $\gamma_{1k}(z) \xrightarrow{a.s.} 0$  by conditioning on  $\{t_j^1, t_j^2 : j \in \{1, 2, \dots, \lceil n/2 \rceil\} \setminus \{k\}\}$ .

Recall that  $\sum_{j=1}^{\lceil n/2 \rceil} |\hat{\mathcal{R}}_z(T_1^{(k)})_{ij}| \leq \|\hat{\mathcal{R}}_z(T_1^{(k)})\|^2 \leq \tau^{-2}$  for  $z \in \partial\mathbb{D}_{\rho+\tau}$ . Now, treating  $\gamma_{1k}(z)$  as the function  $g$  in the property 3 of the Definition 3.2, we obtain

$$(6.8) \quad \mathbb{P}\left(|\gamma_{1k}(z)| > t \mid \hat{\mathcal{R}}_z(T_1^{(k)})\right) \leq K \exp\left(-\tau \sqrt{\frac{\alpha n}{2}} t\right).$$

In addition,

$$(6.9) \quad \mathbb{E}[|\gamma_{1k}(z)|^p] \leq Kp! \left(\frac{2}{\tau^2 \alpha n}\right)^{p/2} \text{ for all } z \in \partial\mathbb{D}_{\rho+\tau}.$$

Similarly, we have bound for  $\gamma_{2k}$  as follows.

$$(6.10) \quad \mathbb{E}[|\gamma_{2k}(z)|^p] \leq Kp! \left(\frac{2}{\tau^2 \alpha n}\right)^{p/2} \text{ for all } z \in \partial\mathbb{D}_{\rho+\tau}.$$

Since  $T_1^{(k)}, T_2^{(k)}$  are independent of the  $k$ th columns of  $T_1$  and  $T_2$  respectively, we can easily see that

$$(6.11) \quad \begin{aligned} & \left[ \text{Tr} \hat{\mathcal{R}}_z(T_1^{(k)}) + \text{Tr} \hat{\mathcal{R}}_z(T_2^{(k)}) \right]_k^\circ \\ &= \text{Tr} \hat{\mathcal{R}}_z(T_1^{(k)}) + \text{Tr} \hat{\mathcal{R}}_z(T_2^{(k)}) - \Xi_k[\text{Tr} \hat{\mathcal{R}}_z(T_1^{(k)}) + \text{Tr} \hat{\mathcal{R}}_z(T_2^{(k)})] = 0. \end{aligned}$$

Hence,

$$(6.12) \quad \begin{aligned} & W_{z,k}(T_1) + W_{z,k}(T_2) \\ &= \mathbb{E}_k \left[ \left( \text{Tr} \hat{\mathcal{R}}_z(T_1) + \text{Tr} \hat{\mathcal{R}}_z(T_2) \right)_k^\circ \right] \\ &= -\mathbb{E}_k \left[ \left( \frac{\partial}{\partial z} \log(1 + \gamma_{1k}(z)) + \frac{\partial}{\partial z} \log(1 + \gamma_{2k}(z)) \right)_k^\circ \right] \\ &= -\frac{\partial}{\partial z} \mathbb{E}_k [\log(1 + \gamma_{1k}(z)) + \log(1 + \gamma_{2k}(z))]_k^\circ \\ (6.13) \quad &= -\frac{\partial}{\partial z} \mathbb{E}_k [\log\{(1 + \gamma_{1k}(z))(1 + \gamma_{2k}(z))\}]_k^\circ. \end{aligned}$$

In the above,  $\mathbb{E}_k$  and  $\partial/\partial z$  are interchanged using the dominated convergence theorem. We can estimate  $W_{z,k}(T_1) + W_{z,k}(T_2)$  by the following fact from complex analysis. Let  $v$  be an analytic function on  $z + \mathbb{D}_{\tau/2}$ . Then

$$(6.14) \quad |v'(z)| = \left| \frac{1}{2\pi i} \oint_{z+\partial\mathbb{D}_{\tau/3}} \frac{v(t)dt}{(t-z)^2} \right| \leq \frac{5}{\pi\tau^2} \oint_{z+\partial\mathbb{D}_{\tau/3}} |v(t)|dt.$$

**6.1. Proof of tightness.** In this subsection, we show the tightness of  $\{\text{Tr } \hat{\mathcal{R}}_z^\circ(M)\}_n$  in the space  $\mathcal{C}(\partial\mathbb{D}_{\rho+\tau})$ . Since  $\text{Tr } \hat{\mathcal{R}}_z^\circ(M) = \text{Tr } \hat{\mathcal{R}}_z^\circ(T_1) + \text{Tr } \hat{\mathcal{R}}_z^\circ(T_2)$ , it is sufficient to show that both  $\{\text{Tr } \hat{\mathcal{R}}_z^\circ(T_1)\}_n$  and  $\{\text{Tr } \hat{\mathcal{R}}_z^\circ(T_2)\}_n$  are tight. To prove the tightness of  $\text{Tr } \hat{\mathcal{R}}_z^\circ(T_1)$  we need to show that for any  $\epsilon > 0$  there exist a compact set  $\mathcal{K}(\epsilon) \subset \mathcal{C}(\partial\mathbb{D}_{\rho+\tau})$  such that

$$(6.15) \quad \mathbb{P} \left( \text{Tr } \hat{\mathcal{R}}_z^\circ(T_1) \in \mathcal{K}(\epsilon)^c \right) < \epsilon.$$

However by Arzelà-Ascoli theorem, the compact sets in space of continuous functions are space of equicontinuous functions. Let us define,

$$\Phi_n := \{|\gamma_{1k}(z)| < n^{-1/8}, \text{ for all } 1 \leq k \leq \lceil n/2 \rceil\}.$$

Hence from (6.8),

$$\mathbb{P}(\Phi_n^c) \leq K \exp \left( \log \left( \frac{n}{2} \right) - \tau \sqrt{\frac{\alpha}{2}} n^{\frac{3}{8}} \right) = o(1).$$

Consequently,

$$(6.16) \quad \begin{aligned} \mathbb{P} \left( \text{Tr } \hat{\mathcal{R}}_z^\circ(T_1) \in \mathcal{K}(\epsilon)^c \right) &\leq \mathbb{P} \left( \frac{|\text{Tr } \hat{\mathcal{R}}_z^\circ(T_1) - \text{Tr } \hat{\mathcal{R}}_\eta^\circ(T_1)|}{|z - \eta|} > K \right) \\ &\leq \mathbb{P} \left( \frac{|\text{Tr } \hat{\mathcal{R}}_z^\circ(T_1) - \text{Tr } \hat{\mathcal{R}}_\eta^\circ(T_1)|}{|z - \eta|} \mathbf{1}_{\Phi_n} > K \right) + o(1), \end{aligned}$$

for some  $K > 0$ . Now, the following equation (6.17) is sufficient to prove (6.15);

$$(6.17) \quad \mathbb{E} \left\{ \left| \frac{\text{Tr } \hat{\mathcal{R}}_z^\circ(T_1) - \text{Tr } \hat{\mathcal{R}}_\eta^\circ(T_1)}{z - \eta} \right|^2 \mathbf{1}_{\Phi_n} \right\} \leq C,$$

uniformly for all  $z, \eta \in \partial\mathbb{D}_{\rho+\tau}$  and  $n \in \mathbb{N}$ . Using the resolvent identity, we can write

$$\text{Tr } \hat{\mathcal{R}}_z(T_1) - \text{Tr } \hat{\mathcal{R}}_\eta(T_1) = (z - \eta) \text{Tr} \left[ \hat{\mathcal{R}}_z(T_1) \hat{\mathcal{R}}_\eta(T_1) \right].$$

Therefore, using (6.4) we have

$$(6.18) \quad (z - \eta)^{-1} \left[ \text{Tr } \hat{\mathcal{R}}_z^\circ(T_1) - \text{Tr } \hat{\mathcal{R}}_\eta^\circ(T_1) \right] = \sum_{k=1}^{\lceil n/2 \rceil} \mathbb{E}_k \left\{ \text{Tr} \left[ \hat{\mathcal{R}}_k(T_1) \hat{\mathcal{R}}_\eta(T_1) \right]_k^\circ \right\}.$$

Using Lemma A.5 and the resolvent identity, we have

$$\begin{aligned} &\text{Tr} \left[ \hat{\mathcal{R}}_z(T_1) \hat{\mathcal{R}}_\eta(T_1) - \hat{\mathcal{R}}_z \left( T_1^{(k)} \right) \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right) \right] \\ &= \text{Tr} \left[ \left\{ \hat{\mathcal{R}}_z(T_1) - \hat{\mathcal{R}}_z \left( T_1^{(k)} \right) \right\} \left\{ \hat{\mathcal{R}}_\eta(T_1) - \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right) \right\} \right] \\ &\quad + \text{Tr} \left[ \hat{\mathcal{R}}_z \left( T_1^{(k)} \right) \left\{ \hat{\mathcal{R}}_\eta(T_1) - \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right) \right\} \right] + \text{Tr} \left[ \left\{ \hat{\mathcal{R}}_z(T_1) - \hat{\mathcal{R}}_z \left( T_1^{(k)} \right) \right\} \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right) \right] \\ &= \text{Tr} \frac{\hat{\mathcal{R}}_z \left( T_1^{(k)} \right) t_k^1 e_k^t \hat{\mathcal{R}}_z \left( T_1^{(k)} \right) \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right) t_k^1 e_k^t \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right)}{1 + \gamma_{k1}(z)} \frac{1}{1 + \gamma_{k1}(\eta)} \end{aligned}$$

$$\begin{aligned}
& + \operatorname{Tr} \frac{\hat{\mathcal{R}}_z \left( T_1^{(k)} \right) \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right) t_k^1 e_k^t \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right)}{1 + \gamma_{k1}(\eta)} + \operatorname{Tr} \frac{\hat{\mathcal{R}}_z \left( T_1^{(k)} \right) t_k^1 e_k^t \hat{\mathcal{R}}_z \left( T_1^{(k)} \right) \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right)}{1 + \gamma_{k1}(z)} \\
& = \frac{\left( e_k^t \hat{\mathcal{R}}_z \left( T_1^{(k)} \right) \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right) t_k^1 \right)^2}{(1 + \gamma_{k1}(z))(1 + \gamma_{k1}(\eta))} + \frac{e_k^t \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right) \hat{\mathcal{R}}_z \left( T_1^{(k)} \right) \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right) t_k^1}{1 + \gamma_{k1}(\eta)} \\
& \quad + \frac{e_k^t \hat{\mathcal{R}}_z \left( T_1^{(k)} \right) \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right) \hat{\mathcal{R}}_z \left( T_1^{(k)} \right) t_k^1}{1 + \gamma_{k1}(z)} \\
& =: \vartheta_1(k) + \vartheta_2(k) + \vartheta_3(k).
\end{aligned}$$

By (6.11), we have  $\left[ \operatorname{Tr} \left( \hat{\mathcal{R}}_z \left( T_1^{(k)} \right) \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right) \right) \right]_k^\circ = 0$ . Therefore, we can express (6.18) as

$$(z - \eta)^{-1} \left[ \operatorname{Tr} \hat{\mathcal{R}}_z(T_1) - \operatorname{Tr} \hat{\mathcal{R}}_\eta^\circ(T_1) \right] = \sum_{k=1}^{\lceil n/2 \rceil} \mathbb{E}_k \{ [\vartheta_1(k) + \vartheta_2(k) + \vartheta_3(k)]_k^\circ \}.$$

Since  $\{[\vartheta_i(k)]_k^\circ\}$  is a martingale difference sequence,

$$\mathbb{E} \left| \sum_{k=1}^{\lceil n/2 \rceil} \mathbb{E}_k \{ [\vartheta_1(k) + \vartheta_2(k) + \vartheta_3(k)]_k^\circ \} \right|^2 = \sum_{k=1}^{\lceil n/2 \rceil} \mathbb{E} |\mathbb{E}_k \{ [\vartheta_1(k) + \vartheta_2(k) + \vartheta_3(k)]_k^\circ \}|^2.$$

Hence, proving (6.17) is equivalent to verifying the validity of

$$(6.19) \quad n \mathbb{E} [|\vartheta_i(k)|^2 \mathbf{1}_{\Phi_n}] \leq C, \text{ for all } 1 \leq k \leq \lceil n/2 \rceil, \quad i = 1, 2, \dots$$

uniformly for all  $n \in \mathbb{N}$  and  $z, \eta \in \partial \mathbb{D}_{\rho+\tau}$ . By adopting the same approach as demonstrated in (6.9), similar tail estimates for  $e_k^t \hat{\mathcal{R}}_z \left( T_1^{(k)} \right) \hat{\mathcal{R}}_\eta \left( T_1^{(k)} \right) t_k^1$  can be obtained possibly with  $\tau^2$  or  $\tau^3$  in the rhs of (6.9). Noting that  $|\gamma_{1k}(z)| < n^{-1/8}$  on  $\Phi_n$ , leveraging the estimate  $|(1 + \gamma_{1k}(z))|^{-1} \leq 2$  in  $\mathbb{E}[|\vartheta_i(k)|^2]$ , we have (6.19). Similarly, we can prove the tightness of  $\operatorname{Tr} \hat{\mathcal{R}}_z^\circ(T_2)$ .

**6.2. Proof of (6.5).** To estimate the right hand side of (6.5), we first estimate the term  $|W_{z,k}(T_1) + W_{z,k}(T_2)|^4$ . We proceed by expanding  $\log(1 + \gamma_{k1}(z))$  and  $\log(1 + \gamma_{k2}(z))$  up to two terms first, and then using the estimates from (6.9),(6.10),(6.13), and (6.14) we get the bound

$$\mathbb{E}[|W_{z,k}(T_1) + W_{z,k}(T_2)|^4] = O(n^{-2}).$$

By substituting the above into (6.5), we get

$$\begin{aligned}
\sum_{k=1}^{\lceil n/2 \rceil} \mathbb{E}[\psi_{n,k}^2 \mathbf{1}_{|\psi_{n,k}| > \theta}] & \leq \frac{(2q)^3}{\theta^2} \sum_{k=1}^{\lceil n/2 \rceil} \sum_{i=1}^q (|\alpha_i|^4 \mathbb{E}[|W_{z_i,k}(T_1) + W_{z_i,k}(T_2)|^4]) \\
& \quad + |\beta_i|^4 \mathbb{E}[|W_{\bar{z}_i,k}(T_1^*) + W_{\bar{z}_i,k}(T_2^*)|^4]) \\
& = \theta^{-2} O(n^{-1}),
\end{aligned}$$

which implies (6.5).

**6.3. Proof of (6.6).** Likewise, we can expand  $\log(1 + \gamma_{k1}(z))$  and  $\log(1 + \gamma_{k2}(z))$  up to two terms and use (6.9), (6.10) to get the following estimate

$$\Xi_k \{ \mathbb{E}_k [(\log(1 + \gamma_{1k}(z)) + \log(1 + \gamma_{2k}(z)))_k^\circ] \mathbb{E}_k [(\log(1 + \gamma_{1k}(\eta)) + \log(1 + \gamma_{2k}(\eta)))_k^\circ] \} = O(n^{-2}).$$

Thus, using (6.13), (6.14) and the above we have

$$\sum_{k=1}^{\lceil n/2 \rceil} \mathbb{E}_{k-1} [(W_{z_i,k}(T_1) + W_{z_i,k}(T_2))(W_{z_j,k}(T_1) + W_{z_j,k}(T_2))] = O(n^{-1}),$$

which implies (6.6).

**6.4. Proof of (6.7).** Expanding  $\log(1 + \gamma_{k1}(z))$  and  $\log(1 + \gamma_{k2}(z))$  up to two terms and using (6.9), (6.10), and (6.13) we have

$$\mathbb{E}_{k-1} [(W_{z,k}(T_1) + W_{z,k}(T_2))(W_{\bar{\eta},k}(T_1^*) + W_{\bar{\eta},k}(T_2^*))] = \frac{\partial^2}{\partial z \partial \bar{\eta}} \Theta_k(z, \bar{\eta}),$$

where

$$\begin{aligned} \Theta_k(z, \bar{\eta}) &= \Xi_k \{ \mathbb{E}_k [(\log(1 + \gamma_{1k}(z)) + \log(1 + \gamma_{2k}(z)))_k^\circ] \mathbb{E}_k [(\log(1 + \overline{\gamma_{1k}(\eta)}) + \log(1 + \overline{\gamma_{2k}(\eta)}))_k^\circ] \} \\ &= \frac{1}{n} \left( e_k^t \mathbb{E}_k [\hat{\mathcal{R}}_z(T_1^{(k)})] \text{Var}((T_1^{(k)})_{ki}) I \mathbb{E}_k [\hat{\mathcal{R}}_{\bar{\eta}}(T_1^{(k)*}) e_k] \right) \\ &\quad + \frac{1}{n} \left( e_k^t \mathbb{E}_k [\hat{\mathcal{R}}_z(T_2^{(k)})] \text{Var}((T_2^{(k)})_{ki}) I \mathbb{E}_k [\hat{\mathcal{R}}_{\bar{\eta}}(T_2^{(k)*}) e_k] \right) + O(n^{-2}) \\ &=: \frac{1}{n} \Gamma_k(z, \bar{\eta}) + O(n^{-2}), \end{aligned}$$

where  $(T_1^{(k)})_{ij}$  and  $(T_2^{(k)})_{ij}$  are the  $ij$ th element of  $T_1^{(k)}$  and  $T_2^{(k)}$  respectively.

Consequently, (6.7) becomes

$$\begin{aligned} &\sum_{k=1}^{\lceil n/2 \rceil} \mathbb{E}_{k-1} [(W_{z,k}(T_1) + W_{z,k}(T_2))(W_{\bar{\eta},k}(T_1) + W_{\bar{\eta},k}(T_2))] \\ &= \frac{\partial^2}{\partial z \partial \bar{\eta}} \left[ \frac{1}{n} \sum_{k=1}^{\lceil n/2 \rceil} \Gamma_k(z, \bar{\eta}) \right] + O(n^{-1}) \\ (6.20) \quad &=: \frac{\partial^2}{\partial z \partial \bar{\eta}} \chi_k(z, \bar{\eta}) + O(n^{-1}). \end{aligned}$$

Since  $\|\hat{\mathcal{R}}_z(T_1^{(k)}) + \hat{\mathcal{R}}_z(T_2^{(k)})\| \leq 4\tau^{-1}$  on  $\mathbb{D}_{\rho+\tau/2}^c$ , the sequence  $\chi_k(z, \bar{\eta})$  consists of uniformly bounded analytic functions on  $\mathbb{D}_{\rho+\tau/2}^c$ . Thus, by Vitali's theorem (see [14, Section 2.9]), proving (6.7) is equivalent to show that  $\chi_k(z, \bar{\eta})$  converges in probability.

**Lemma 6.2.** *Let  $\chi_k(z, \bar{\eta})$  be the same as defined in (6.20), that is,  $\chi_k(z, \bar{\eta}) = \frac{1}{n} \sum_{k=1}^{\lceil n/2 \rceil} \Gamma_k(z, \bar{\eta})$ . Then  $\text{Var}(\chi_k(z, \bar{\eta})) \rightarrow 0$  as  $n \rightarrow \infty$ .*

*Proof.* Since  $\{W_{z,k}(T_1) + W_{z,k}(T_2)\}_n$  is a martingale difference sequence, we have

$$\mathbb{E}[(W_{z,k}(T_1) + W_{z,k}(T_2))(W_{\bar{\eta},l}(T_1^*) + W_{\bar{\eta},l}(T_2^*))] = 0 \text{ if } k \neq l.$$

This implies,

$$\begin{aligned}
& \mathbb{E} \left\{ \sum_{k=1}^{\lceil n/2 \rceil} \mathbb{E}_{k-1} [(W_{z,k}(T_1) + W_{z,k}(T_2))(W_{\bar{\eta},k}(T_1^*) + W_{\bar{\eta},k}(T_2^*)))] \right\} \\
&= \mathbb{E} \left\{ \left( \sum_{k=1}^{\lceil n/2 \rceil} (W_{z,k}(T_1) + W_{z,k}(T_2)) \right) \left( \sum_{k=1}^{\lceil n/2 \rceil} (W_{\bar{\eta},k}(T_1^*) + W_{\bar{\eta},k}(T_2^*)) \right) \right\} \\
&= \mathbb{E} \left\{ \left( \text{Tr } \hat{\mathcal{R}}_z^\circ(T_1) + \text{Tr } \hat{\mathcal{R}}_z^\circ(T_2) \right) \left( \text{Tr } \hat{\mathcal{R}}_{\bar{\eta}}^\circ(T_1^*) + \text{Tr } \hat{\mathcal{R}}_{\bar{\eta}}^\circ(T_2^*) \right) \right\} \\
&= \mathbb{E} \left\{ \text{Tr } \hat{\mathcal{R}}_z^\circ(M) \text{Tr } \hat{\mathcal{R}}_{\bar{\eta}}^\circ(M^*) \right\}.
\end{aligned}$$

Let us rewrite  $\Gamma_k(z, \bar{\eta})$  as

$$\begin{aligned}
& \Gamma_k(z, \bar{\eta}) \\
&= 4e_k^t \mathbb{E}_k[\hat{\mathcal{R}}_z(T_1^{(k)})] \mathbb{E}_k[\hat{\mathcal{R}}_{\bar{\eta}}(T_1^{(k)*})] e_k + 4e_k^t \mathbb{E}_k[\hat{\mathcal{R}}_z(T_2^{(k)})] \mathbb{E}_k[\hat{\mathcal{R}}_{\bar{\eta}}(T_2^{(k)*})] e_k \\
&:= \Gamma_k^1(z, \bar{\eta}) + \Gamma_k^2(z, \bar{\eta}).
\end{aligned}$$

Consequently, we may decompose

$$\begin{aligned}
& \text{Var} \left( \frac{1}{n} \Gamma_k(z, \bar{\eta}) \right) \\
&= \text{Var} \left[ \frac{1}{n} \Gamma_k^1(z, \bar{\eta}) + \frac{1}{n} \Gamma_k^2(z, \bar{\eta}) \right] \\
&\leq 2 \text{Var} \left( \frac{1}{n} \Gamma_k^1(z, \bar{\eta}) \right) + 2 \text{Var} \left( \frac{1}{n} \Gamma_k^2(z, \bar{\eta}) \right).
\end{aligned}$$

Therefore, proving the lemma is equivalent to showing

$$\text{Var} \left( \frac{1}{n} \Gamma_k^1(z, \bar{\eta}) \right) + \text{Var} \left( \frac{1}{n} \Gamma_k^2(z, \bar{\eta}) \right) \rightarrow 0.$$

We show that  $\text{Var} \left( \frac{1}{n} \Gamma_k^1(z, \bar{\eta}) \right) \rightarrow 0$ . We shall use the Poincaré inequality to get estimates on  $\text{Var} \left( \frac{1}{n} \Gamma_k^1(z, \bar{\eta}) \right)$ . This is shown in the equation (6.22). However, since we are working on the event  $\Omega_{n,k}$ , where norms of the matrices are bounded, some terms may not be differentiable with respect to the matrix entries. To circumvent such a problem, we define a smoothing function as follows.

Let  $\epsilon_n = 4/n$ , and  $\omega_{\epsilon_n, k} : \mathbb{R}^{\lceil n^2/4 \rceil} \rightarrow [0, 1]$  be a smooth function such that

$$\begin{aligned}
& \omega_{\epsilon_n, k} \Big|_{\{\|T_1^{(k)}\| \leq \rho_1\}} \equiv 1, \\
& \omega_{\epsilon_n, k} \Big|_{\{\|T_1^{(k)}\| \geq \rho_1 + \epsilon_n\}} \equiv 0, \\
& \left| \frac{\partial}{\partial x_i} \omega_{\epsilon_n, k} \right| \leq \frac{2}{\epsilon_n}, \quad \forall 1 \leq i \leq \lceil n^2/4 \rceil.
\end{aligned}$$

Let us define the smoothed version of  $\mathcal{R}$  as

$$\tilde{\mathcal{R}}_z(T_1^{(k)}) = \mathcal{R}_z(T_1^{(k)}) \omega_{\epsilon_n, k},$$

and

$$\tilde{\Gamma}_k^1(z, \bar{\eta}) = 4e_k^t \mathbb{E}_k[\tilde{\mathcal{R}}_z(T_1^{(k)})] \mathbb{E}_k[\tilde{\mathcal{R}}_{\bar{\eta}}(T_1^{(k)*})] e_k.$$

Since the smoothed version and the original one match everywhere except the narrow annulus  $\{\rho_1 < \|T_1^{(k)}\| < \rho_1 + \epsilon_n\}$ , we can easily obtain

$$|\Gamma_k^1(z, \bar{\eta}) - \tilde{\Gamma}_k^1(z, \bar{\eta})| \leq \frac{K}{(\tau - \epsilon_n)^2} \mathbf{1}_{\{\rho_1 < \|T_1^{(k)}\| < \rho_1 + \epsilon_n\}},$$

where  $K$  is some universal constant. Now, by Lemma A.2, we have,

$$(6.21) \quad \mathbb{E}[|\Gamma_k^1(z, \bar{\eta}) - \tilde{\Gamma}_k^1(z, \bar{\eta})|^2] \leq \frac{K}{(\tau - \epsilon_n)^4} \exp\left(-\tau \sqrt{\frac{\alpha n}{2}} \frac{3\rho_1}{4}\right).$$

Thus, we see that the smoothing function  $\omega$  did not alter the original quantities by significant amount. For the notational convenience, let us denote

$$\begin{aligned} \Lambda &:= \mathbb{E}_k[\mathcal{R}_z(T_1^{(k)})], & \Upsilon &:= \mathbb{E}_k[\mathcal{R}_\eta(T_2^{(k)*})], \\ \tilde{\Lambda} &:= \mathbb{E}_k[\tilde{\mathcal{R}}_z(T_1^{(k)})], & \tilde{\Upsilon} &:= \mathbb{E}_k[\tilde{\mathcal{R}}_\eta(T_2^{(k)*})]. \end{aligned}$$

So,  $\tilde{\Gamma}_k^1(z, \bar{\eta}) = \sum_{s=0}^{\lceil n/2 \rceil} \tilde{\Lambda}_{ks} \tilde{\Upsilon}_{sk}$ . Since  $x_{ij}$ s satisfy Poincaré inequality, we have

$$(6.22) \quad \text{Var}(\tilde{\Gamma}_k^1(z, \bar{\eta})) \leq \frac{1}{n\alpha} \sum_{i,j=1}^{k-1} \left[ \mathbb{E} \left| \frac{\partial \tilde{\Gamma}_k^1(z, \bar{\eta})}{\partial t_{ij}^1} \right|^2 + \mathbb{E} \left| \frac{\partial \tilde{\Gamma}_k^1(z, \bar{\eta})}{\partial t_{ij}^{\bar{1}}} \right|^2 \right].$$

The sum is divided by  $n$  because the matrix  $T_1$  is scaled by  $1/\sqrt{n}$ . Moreover, the sum stops at  $k-1$  because  $\tilde{\Gamma}_k^1(z, \bar{\eta})$  is a constant for  $i, j = k, k+1, \dots, \lceil n/2 \rceil$ . On the other hand, we have

$$\frac{\partial \mathcal{R}_z(T_1^{(k)})_{ks}}{\partial t_{ij}^1} = \mathcal{R}_z(T_1^{(k)})_{ki} \mathcal{R}_z(T_1^{(k)})_{js} \mathbf{1}_{\{j \neq k\}}, \quad \overline{\frac{\partial \mathcal{R}_z(T_1^{(k)})_{ks}}{\partial t_{ij}^1}} = 0,$$

As a result,

$$\begin{aligned} \frac{\partial \tilde{\Lambda}_{ks}}{\partial t_{ij}^1} &= \tilde{\Lambda}_{ki} \tilde{\Lambda}_{js} \mathbf{1}_{\{j \neq k\}} + \Lambda_{ks} \frac{\partial \omega_{\epsilon_n, k}}{\partial t_{ij}^1} \mathbf{1}_{\{\rho_1 < \|T_1^{(k)}\| < \rho_1 + \epsilon_n\}}, \\ \frac{\partial \tilde{\Upsilon}_{sk}}{\partial t_{ij}^1} &= \Upsilon_{sk} \frac{\partial \omega_{\epsilon_n, k}}{\partial t_{ij}^1} \mathbf{1}_{\{\rho_1 < \|T_1^{(k)}\| < \rho_1 + \epsilon_n\}}, \\ \frac{\partial (\tilde{\Gamma}_k^1(z, \bar{\eta}))}{\partial t_{ij}^1} &= \sum_{s=0}^{\lceil n/2 \rceil} \tilde{\Lambda}_{ki} \tilde{\Lambda}_{js} \tilde{\Upsilon}_{sk} + \sum_{s=0}^{n/2} (\Lambda_{ks} \tilde{\Upsilon}_{sk} + \tilde{\Lambda}_{ks} \Upsilon_{sk}) \frac{\partial \omega_{\epsilon_n, k}}{\partial t_{ij}^1} \mathbf{1}_{\{\rho_1 < \|T_1^{(k)}\| < \rho_1 + \epsilon_n\}}. \end{aligned}$$

Let us denote  $\tilde{y}_{jk} = \sum_{s=0}^{\lceil n/2 \rceil} \tilde{\Lambda}_{js} \tilde{\Upsilon}_{sk}$ . Then using the facts that  $\|\tilde{\Lambda}\|, \|\tilde{\Upsilon}\| \leq (\tau - \epsilon_n)^{-1}$ , we have

$$\sum_{i,j=1}^{k-1} \left| \sum_{s=0}^{\lceil n/2 \rceil} \tilde{\Lambda}_{ki} \tilde{\Lambda}_{js} \tilde{\Upsilon}_{sk} \right|^2 = \sum_{i,j=1}^{k-1} |\tilde{\Lambda}_{ki} \tilde{y}_{jk}|^2 \leq \|\tilde{\Lambda}_k\|_2^2 \|\tilde{y}_k\|_2^2 \leq (\tau - \epsilon_n)^{-6},$$

$$\begin{aligned}
& \sum_{i,j=1}^{k-1} \left| \sum_{s=0}^{\lceil n/2 \rceil} \left( \Lambda_{ks} \tilde{\Upsilon}_{sk} + \tilde{\Lambda}_{ks} \Upsilon_{sk} \right) \frac{\partial \omega_{\epsilon_n, k}}{\partial a_{ij}} \mathbf{1}_{\{\rho_1 < \|T_1^{(k)}\| < \rho_1 + \epsilon_n\}} \right|^2 \\
& \leq \frac{10}{\epsilon_n^2} \sum_{i,j=1}^{k-1} \left\| \tilde{\Lambda}_k \right\|_2^2 \left\| \tilde{\Upsilon}_k \right\|_2^2 \mathbf{1}_{\{\rho < \|T_1^{(k)}\| < \rho + \epsilon_n\}} \leq 10n^4 (\tau - \epsilon_n)^{-4} \mathbf{1}_{\{\rho_1 < \|T_1^{(k)}\| < \rho_1 + \epsilon_n\}}.
\end{aligned}$$

Using the above estimates, (6.1), we have

$$\begin{aligned}
& \text{Var} \left( \frac{1}{n} \sum_{k=1}^{\lceil n/2 \rceil} \tilde{\Gamma}_k^1(z, \bar{\eta}) \right) \\
& \leq \frac{1}{n} \sum_{k=1}^{\lceil n/2 \rceil} \text{Var} \left( \tilde{\Gamma}_k^1(z, \bar{\eta}) \right) \\
& \leq \frac{2}{\alpha n (\tau - \epsilon_n)^6} + \frac{Kn^3}{(\tau - \epsilon_n)^4} \exp \left( -\sqrt{\frac{\alpha n}{2}} \frac{3\rho_1}{4} \right).
\end{aligned}$$

Therefore from (6.21), we obtain

$$\begin{aligned}
& \text{Var} \left( \frac{1}{n} \sum_{k=1}^{\lceil n/2 \rceil} \Gamma_k^1(z, \bar{\eta}) \right) \\
& \leq \frac{1}{n} \sum_{k=1}^{\lceil n/2 \rceil} \mathbb{E} \left[ \left| \Gamma_k^1(z, \bar{\eta}) - \tilde{\Gamma}_k^1(z, \bar{\eta}) \right|^2 \right] + \text{Var} \left( \frac{1}{n} \sum_{k=1}^{\lceil n/2 \rceil} \tilde{\Gamma}_k^1(z, \bar{\eta}) \right) \\
& \leq \frac{2}{\alpha n (\tau - \epsilon_n)^6} + \frac{Kn^3}{(\tau - \epsilon_n)^4} \exp \left( -\sqrt{\frac{\alpha n}{2}} \frac{3\rho_1}{4} \right) \rightarrow 0 \text{ as } n \rightarrow \infty.
\end{aligned}$$

Similarly,  $\text{Var} \left( \frac{1}{n} \sum_{k=1}^{\lceil n/2 \rceil} \Gamma_k^2(z, \bar{\eta}) \right) \rightarrow 0$  as  $n \rightarrow \infty$ . Thus,

$$\begin{aligned}
& \text{Var}(\chi_k(z, \bar{\eta})) \\
& \leq 2 \text{Var} \left( \frac{1}{n} \sum_{k=1}^{\lceil n/2 \rceil} \Gamma_k^1(z, \bar{\eta}) \right) + 2 \text{Var} \left( \frac{1}{n} \sum_{k=1}^{\lceil n/2 \rceil} \Gamma_k^2(z, \bar{\eta}) \right) \rightarrow 0, \text{ as } n \rightarrow \infty.
\end{aligned}$$

□

This completes the proof of the central limit theorem. Now, we demonstrate the calculation of the covariance kernel in Section 7. Before that, let us verify this result through a numerical simulation; see Figure 3.






FIGURE 3. Histogram of  $L^\circ(P_d)$  of a  $4000 \times 4000$  random centrosymmetric matrix sampled 1000 times. The entries of the matrix are drawn from the Gaussian distribution, where  $P_d(x) = 2x^3 + x^4$ .

## 7. CALCULATION OF VARIANCE

This section is dedicated to calculating the covariance kernel. We proceed with the expression of the variance in the case of polynomial test functions.

**Proposition 7.1.** *Let  $M$  be the random matrix as defined in Definition 3.1 satisfying the Condition 3.3. Then we have*

$$\text{Var}(\text{Tr}(P_d(M))) = \sum_{k=1}^d 2k|a_k|^2 + o(1),$$

where  $P_d(x) = \sum_{k=1}^d a_k x^k$  is a polynomial of degree  $d \geq 1$ .

Notice that, the variance expression does not contain the constant term of the polynomial  $P_d(x)$ . The constant term of a matrix polynomial will be a constant multiplied by the identity matrix, and this term will not affect the fluctuation result of the LES of a random matrix, as we center the LES by its mean.

We recognize that the essential ingredients are  $\mathbb{E}[\text{Tr}(M^k) \text{Tr}(\bar{M}^l)]$  and  $\mathbb{E}[\text{Tr}(M^k)]$ , where  $\bar{M}$  is obtained by taking complex conjugates of all the entries of  $M$ . We compute these quantities using combinatorial arguments. Before proving the Proposition 7.1, we introduce and analyze some required notions such as pair partitions and chain merging in the following subsections.

**7.1. Analyzing expected product.** We evaluate  $\mathbb{E}[x_1 x_2 \cdots x_n]$ , where  $x_i$ s are random variables from Condition 3.3. Note that  $\mathbb{E}[x_i^2] = 0$  if  $x_i$  satisfies Condition 3.3.

In the next calculations, we need to evaluate the terms of the forms  $\mathbb{E}[\text{Tr}(M^k)]$  and  $\mathbb{E}[\text{Tr}(M^k) \text{Tr}(\bar{M}^l)]$ . Upon expansion, the term  $\text{Tr}(M^k)$  is represented as  $\sum_{i_1, i_2, \dots, i_k=1}^n x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_k i_1}$ . Notice that here the random variables  $x_{i_1 i_2}, x_{i_2 i_3}, \dots, x_{i_k i_1}$  are linked via their indices  $i_1, i_2, \dots, i_k$ . The arrangement of these indices plays an important role in evaluating expectations. We formalize this using the notation of *index chain* as defined below.

**Definition 7.2** (Index chains). The index chain of the product of random variables  $x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_k i_1}$  is the unique sequence  $i_1 \rightarrow i_2 \rightarrow \cdots \rightarrow i_k \rightarrow i_1$  of indices.

- (a) Single chain: In the calculation of  $\mathbb{E}[\text{Tr}(M^k)]$ , the indices of the product are chained as  $i_1 \rightarrow i_2 \rightarrow \cdots \rightarrow i_k \rightarrow i_1$  which will be referred as single chain.
- (b) Double chain: In the calculation of  $\mathbb{E}[\text{Tr}(M^k) \text{Tr}(\bar{M}^l)]$ , there are two chains of indices  $i_1 \rightarrow i_2 \rightarrow \cdots \rightarrow i_k \rightarrow i_1$  and  $j_1 \rightarrow j_2 \rightarrow \cdots \rightarrow j_l \rightarrow j_1$  which will be referred as double chain.

In principle, the indices of an index chain can take values from the set  $\{1, 2, 3, \dots, n\}$ . However, while calculating  $\mathbb{E}[\text{Tr}(M^k)]$  and  $\mathbb{E}[\text{Tr}(M^k) \text{Tr}(\bar{M}^l)]$ , we shall see that not all the indices are free to be chosen from  $\{1, 2, 3, \dots, n\}$ . Some indices will be determined by the choices of the other indices. In this context, we shall use the terms ‘degree of freedom’ and ‘free indices’, which are defined in Definition 7.3.

**Definition 7.3** (Degrees of freedom and free indices). In a given index chain, free indices are the indices which are not constrained by any other index. The degree of freedom represents the number of free indices in a chain.

This will be relevant while calculating the  $\mathbb{E}[\text{Tr} M^k]$  or  $\mathbb{E}[\text{Tr} M^k \bar{M}^l]$ . We shall use the notion of free indices in the remaining part of this section; in particular in the proofs of Lemma 7.4, Lemma 7.5, and Lemma 7.6.

Now, we proceed to the calculation of  $\mathbb{E}[\text{Tr}(M^k)]$  and  $\mathbb{E}[\text{Tr}(M^k) \text{Tr}(\bar{M}^l)]$ , which are discussed in Sections 7.2 and 7.3.

**7.2. Single Chain Expectation.** In this subsection, we focus on computing the expected value of the trace of a matrix raised to the power of  $k$ , that is, the expectation of a single chain. Specifically, we are interested in evaluating the expressions of the form.

$$\mathbb{E}[\text{Tr}(M^k)] = \frac{1}{n^{k/2}} \sum_{i_1, i_2, \dots, i_k=1}^n \mathbb{E}[x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_{k-1} i_k} x_{i_k i_1}].$$

We state the first lemma in this regard.

**Lemma 7.4.** *Let  $M$  be the random matrix as defined in Definition 3.1 satisfying the Condition 3.3. Then we have the following*

$$\begin{aligned} \mathbb{E}[\text{Tr}(M^k)] &\leq n^{-k/6} k^{2k} \left( \frac{4}{3\alpha} \right)^k \\ &\leq k^{3k} n^{-k/6}, \quad \text{if } k > 4/3\alpha. \end{aligned}$$

Here  $\alpha$  is the constant of Poincaré inequality. In particular, if  $k \leq n^\beta$  for some  $\beta < 1/18$ , then  $\mathbb{E}[\text{Tr}(M^k)] = o(1)$ .

*Proof.* This lemma evaluates the limit expectation of a single chain through intra chain merging, which is defined in Section 7.3. The expectation of the trace of  $M^k$  can be expressed as

$$\mathbb{E}[\text{Tr}(M^k)] = \frac{1}{n^{k/2}} \sum_{i_1, i_2, \dots, i_k=1}^n \mathbb{E}[x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_{k-1} i_k} x_{i_k i_1}].$$

To evaluate  $\mathbb{E}[x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_{k-1} i_k} x_{i_k i_1}]$ , we need to determine the number of ways in which  $x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_{k-1} i_k} x_{i_k i_1}$  simplifies to a product where each  $x_{ij}$  may appear with some multiplicity. However, given that  $\mathbb{E}[x_{ij}] = 0 = \mathbb{E}[x_{ij}^2]$  according to Condition 3.3, we should focus on the cases where more than two random variables are equal. Since the product has  $k$  random variables, essentially the task boils down to considering the integer partitions of  $k$ , where group sizes in each partition is at least 3. For example,  $22 = 10 + 5 + 4 + 3$  is an integer partition of 22 having four groups, where group sizes are 10, 5, 4, 3.

Suppose in an integer partition of  $k$ , we have  $g$  groups, where each group contains  $k_1, k_2, \dots, k_g$  random variables respectively, and the random variables within any particular group are equal. For example, a group of size  $k_i$  consists of  $k_i$  equal random variables, and no two groups share the same random variables. Additionally, the cardinality of each group is at least 3, and  $\sum_{l=1}^g k_l = k$ .

Let us analyze a generic group of size  $k_l$ . To reduce notional complexity, here we denote the group as  $x_{\alpha_1} x_{\alpha_2} \cdots x_{\alpha_{k_l}}$ , where each  $\alpha_p$  is of the form  $(i_{\alpha_p}, i_{\alpha_p+1})$ . As per the construction,  $x_{\alpha_1} = x_{\alpha_2} = \cdots = x_{\alpha_{k_l}}$ . Therefore, using the fact that  $x_{ij}$ s satisfy Poincaré inequality, we have

$$\mathbb{E}[x_{\alpha_1} x_{\alpha_2} \cdots x_{\alpha_{k_l}}] = \mathbb{E}[|x_{11}|^{k_l}] \leq k_l! \left(\frac{2}{\alpha}\right)^{k_l/2}.$$

Hence, we conclude that if we fix a partition  $\sum_{l=1}^g k_l = k$ , and fix random variables in each group of the partition, then

$$\begin{aligned} \mathbb{E}[x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_{k-1} i_k} x_{i_k i_1}] &\leq \prod_{l=1}^g k_l! \left(\frac{2}{\alpha}\right)^{k_l/2} \\ &= \left(\frac{2}{\alpha}\right)^{k/2} \prod_{l=1}^g k_l! \\ &\leq \left(\frac{2}{\alpha}\right)^{k/2} k!, \end{aligned}$$

where the last inequality follows from the fact that  $m!n! \leq (m+n)!$ . However, if we fix a partition of  $k$ , but do not fix the random variables in the partition groups, then we have some choices for the indices  $\alpha_1, \alpha_2, \dots, \alpha_{k_l}$  in the product  $x_{\alpha_1} x_{\alpha_2} \cdots x_{\alpha_{k_l}}$ , which is a group of size  $k_l$ . Now, since all the random variables in this group are equal, we only have the freedom to choose  $\alpha_1$ . Once  $\alpha_1$  is fixed, the remaining  $\alpha_p$ s in the same group can either be  $\alpha_1$  or  $n+1-\alpha_1$ . Hence, for a given  $\alpha_1$ , we have at most  $2^{k_l-1}$  many possibilities of the indices  $\alpha_2, \alpha_3, \dots, \alpha_{k_l}$  for that group.

The index  $\alpha_1 = (i_{\alpha_1}, i_{\alpha_1+1})$  consists of two indices. However, since this group  $x_{\alpha_1} x_{\alpha_2} \cdots x_{\alpha_{k_l}}$  is a part of the full chain  $x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_{k-1} i_k} x_{i_k i_1}$ , one of the indices among  $i_{\alpha_1}, i_{\alpha_1+1}$  must also appear in some other group. As a result, either  $i_{\alpha_1}$  or  $i_{\alpha_1+1}$  must be determined by some other group. Therefore, we have the freedom to choose only one of the indices  $i_{\alpha_1}, i_{\alpha_1+1}$ , which can be done in at most  $n$  possible ways. Hence, we have an additional factor  $n2^{k_l-1}$  in the  $l$ th group of the partition. And overall, we have a factor of  $n^g 2^{\sum_{l=1}^g (k_l-1)} = n^g 2^{k-g}$ .

Finally, combining all the above arguments we have

$$\begin{aligned}
\mathbb{E}[\text{Tr}(M^k)] &= \frac{1}{n^{k/2}} \sum_{i_1, i_2, \dots, i_k=1}^n \mathbb{E}[x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_{k-1} i_k} x_{i_k i_1}] \\
&= \frac{1}{n^{k/2}} \sum_{g=1}^{k/3} \sum_{\substack{k_1+k_2+\dots+k_g=k \\ k_l \geq 3}} n^g 2^{k-g} \left(\frac{2}{\alpha}\right)^{k/2} k! \\
&\leq \frac{1}{n^{k/2}} \left(\frac{k}{3}\right)^k n^{k/3} 2^{2k/3} \left(\frac{2}{\alpha}\right)^{k/2} k! \\
&\leq n^{-k/6} k^{2k} \left(\frac{4}{3\alpha}\right)^k \\
&\leq n^{-k/6} k^{3k},
\end{aligned}$$

for  $k > 4/3\alpha$ . The third last inequality follows from the fact that  $n^g 2^{k-g} \leq n^{k/3} 2^{2k/3}$  for  $1 \leq g \leq k/3$  and the number of partitions of  $k$  into at most  $k/3$  many groups is  $(k/3)^k$ . This completes the proof.  $\square$

Now, we shall proceed with the calculation of the covariance between  $\text{Tr } M^k$ ,  $\text{Tr } M^l$ .

**7.3. Double chain Expectation.** In this section, our focus is on computing the expected value of the product of traces of matrices raised to different powers, which requires the computation of the expectation of a double chain. Specifically, we are interested in expressions of the form

$$\sum_{k,l=1}^d \frac{1}{n^{(k+l)/2}} \sum_{i_1, i_2, \dots, i_k, j_1, j_2, \dots, j_l=1}^n \mathbb{E}[(x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_k i_1})(\bar{x}_{j_1 j_2} \bar{x}_{j_2 j_3} \cdots \bar{x}_{j_l j_1})].$$

Here,  $k$  and  $l$  are positive integers denoting the lengths of the respective chains. From the Condition 3.3, we know that  $\mathbb{E}[x_{ij}] = 0 = \mathbb{E}[x_{ij}^2]$ . Therefore as explained in Lemma 7.4, we need to choose the indices of the chains in such a way that each random variable occurs with a multiplicity. However, it is quite evident that if we make more random variables equal to each other within a chain or across different chains, then we lose more degrees of freedom in the index chains. So, we aim to make fewer random variables equal, preferably 2, across two different chains. Since in this case, we have the term  $\mathbb{E}[|x_{ij}|^2]$ , which is 1. Thus, when we have a product of two chains such as  $(x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_k i_1})(\bar{x}_{j_1 j_2} \bar{x}_{j_2 j_3} \cdots \bar{x}_{j_l j_1})$ , we can pair random variables from the first chain  $x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_k i_1}$  with the second chain  $\bar{x}_{j_1 j_2} \bar{x}_{j_2 j_3} \cdots \bar{x}_{j_l j_1}$  in the following three different ways:

- (1) All the random variables of one chain find their pair within the chain itself - *intra chain merging*.
- (2) All random variables in one chain find their pair in the other chain - *cross chain merging*.
- (3) Some of the random variables of one chain find their pairs in the chain itself, and others find in the other chain - *partial chain merging*.

Let us explain each case as follows.

**Case 1 (Intra chain merging):** As we have seen in the Lemma 7.4, the single chain expectation is asymptotically zero. Thus, intra chain merging gives us zero asymptotic contribution.

**Case 2 (Cross chain merging):** To evaluate  $\mathbb{E}[\text{Tr}(M^k) \text{Tr}(\bar{M}^l)]$ , let us consider a simple example with  $k = l = 5$ .

$$\mathbb{E}[\text{Tr}(M^5) \text{Tr}(\bar{M}^5)] = \frac{1}{n^5} \sum_{\substack{i_1, i_2, \dots, i_5, \\ j_1, j_2, \dots, j_5=1}}^n \mathbb{E}[(x_{i_1 i_2} x_{i_2 i_3} x_{i_3 i_4} x_{i_4 i_5} x_{i_5 i_1})(\bar{x}_{j_1 j_2} \bar{x}_{j_2 j_3} \bar{x}_{j_3 j_4} \bar{x}_{j_4 j_5} \bar{x}_{j_5 j_1})].$$

We shall see in the proof of Lemma 7.5, we get maximum contribution if every random variable from the first chain gets paired with a random variable from the second chain.

One way of cross-chain merging is that if  $x_{i_t i_l}$  gets paired with  $\bar{x}_{j_p j_q}$ , then the pairing for  $x_{i_{t+1} i_{l+1}}$  will be  $\bar{x}_{j_{p+1} j_{q+1}}$  for all  $t$  and  $l$  such that  $1 \leq t, l \leq 5$ . In other words, the random variables from the first chain get matched with the random variables in the second chain in the same order as they appear in the first chain, see Figure 4, and we demonstrate that it gives us a nonzero asymptotic contribution.




FIGURE 4. Sequential cross chain merging

For any pair of random variables  $(x_{ij}, \bar{x}_{kl})$ , where  $x_{ij}$  and  $\bar{x}_{kl}$  are from the chains  $x_{i_1 i_2} x_{i_2 i_3} x_{i_3 i_4} x_{i_4 i_5} x_{i_5 i_1}$ , and  $\bar{x}_{j_1 j_2} \bar{x}_{j_2 j_3} \bar{x}_{j_3 j_4} \bar{x}_{j_4 j_5} \bar{x}_{j_5 j_1}$  respectively,  $\mathbb{E}[x_{ij} \bar{x}_{kl}] \neq 0$  if and only if  $(i, j) = (k, l)$  or  $(i, j) = (n + 1 - k, n + 1 - l)$ . This leads to the following possible pairings in the above example.

$$\left\{ \begin{array}{l} j_1 = n + 1 - i_1 \\ j_2 = n + 1 - i_2 \end{array} \right. \quad \text{or} \quad \left\{ \begin{array}{l} j_1 = i_1 \\ j_2 = i_2 \end{array} \right.$$

$$\left\{ \begin{array}{l} j_2 = n + 1 - i_2 \\ j_3 = n + 1 - i_3 \end{array} \right. \quad \text{or} \quad \left\{ \begin{array}{l} j_2 = i_2 \\ j_3 = i_3 \end{array} \right.$$

$$\left\{ \begin{array}{l} j_3 = n + 1 - i_3 \\ j_4 = n + 1 - i_4 \end{array} \right. \quad \text{or} \quad \left\{ \begin{array}{l} j_3 = i_3 \\ j_4 = i_4 \end{array} \right.$$

$$\left\{ \begin{array}{l} j_4 = n + 1 - i_4 \\ j_5 = n + 1 - i_5 \end{array} \right. \quad \text{or} \quad \left\{ \begin{array}{l} j_4 = i_4 \\ j_5 = i_5 \end{array} \right.$$

$$\left\{ \begin{array}{l} j_5 = n + 1 - i_5 \\ j_1 = n + 1 - i_1 \end{array} \right. \quad \text{or} \quad \left\{ \begin{array}{l} j_5 = i_5 \\ j_1 = i_1 \end{array} \right.$$

If we start with  $j_1 = n + 1 - i_1$ , then the chain has to follow the constraints listed in the left column; not the ones which are listed on the right column. Similarly, if we start with the condition  $j_1 = i_1$ , then we have to follow the constraints that are listed on the right column.

For example, if we take  $(i_1, i_2) = (n + 1 - j_1, n + 1 - j_2)$  and decide to take  $(i_2, i_3) = (j_2, j_3)$ , then to keep the continuity of the chain, we must have  $j_2 = i_2 = n + 1 - j_2$  which enforces one extra constraint  $j_2 = n + 1 - j_2$ . Thus, we have lost a degree of freedom there. Now, if the indices follow the constraints listed on the first column, then there are  $n^5$  many choices for choosing all the indices. For each such choice, the chain will reduce to a product of the terms of the form  $\mathbb{E}[|x_{i_j}|^2](= 1)$ .

Thus, we conclude that when we do the cross chain merging for the first and the second chain, namely  $i_1 \rightarrow i_2 \rightarrow i_3 \rightarrow i_4 \rightarrow i_5 \rightarrow i_1$  and  $j_1 \rightarrow j_2 \rightarrow j_3 \rightarrow j_4 \rightarrow j_5 \rightarrow j_1$ , one way for cross chain merging is to copy the first chain in place of the second chain, resulting in:  $i_1 \rightarrow i_2 \rightarrow i_3 \rightarrow i_4 \rightarrow i_5 \rightarrow i_1 \rightarrow i_2 \rightarrow i_3 \rightarrow i_4 \rightarrow i_5 \rightarrow i_1$ . This represents one possibility with 5 free indices. Additionally, we can start the second chain with  $i_2$  or  $i_3$  or  $\dots$  or  $i_5$  like  $i_2 \rightarrow i_3 \rightarrow \dots \rightarrow i_5 \rightarrow i_1 \rightarrow i_2$ , resulting in 4 more choices. Therefore, we have  $5n^5$  combinations for cross chain merging of the first and the second chain so far. Moreover, since  $x_{(i,j)} = x_{(n+1-i, n+1-j)}$ , we can perform a cross chain merging by changing the second chain to  $n + 1 - i_1 \rightarrow n + 1 - i_2 \rightarrow \dots \rightarrow n + 1 - i_5 \rightarrow n + 1 - i_1$ , which will contribute another  $5n^5$ . Hence, the total number of combinations for cross chain merging of the first and second chain is  $2 \times 5n^5$  from type of merging in Figure 4.

Now, let us consider the another type of case as demonstrated in Figure 5, where the first chain is not merging with the second chain in the same sequence. We shall argue that such mergings will give negligible contributions.




FIGURE 5. Non sequential cross chain merging

For this case, we need to impose more constraints that will reduce the degree of freedom and leads to zero asymptotic contribution. Here,  $x_{i_1 i_2} = \bar{x}_{j_3 j_4}$  if and only if  $(i_1, i_2) = (j_3, j_4)$  or the dual case  $(i_2, i_3) = (n + 1 - j_3, n + 1 - j_4)$ .

If  $j_3 = i_1$ , this implies  $j_4 = i_2$ . Furthermore, since we are pairing  $x_{i_2 i_3}$  and  $\bar{x}_{j_5 j_1}$ , we shall have  $j_5 = i_2$  and  $i_3 = j_1$ , or the dual case. Now, let us look at the consecutive random variables of those already paired up. We observe that to maintain the continuity of the chain, we must impose restrictions that will significantly reduce the number of free indices and lead to zero asymptotic contribution.

In conclusion, the total number of combinations that provide a nonzero asymptotic contribution is  $2 \times 5n^5$ . We obtain that contribution from the cases as demonstrated in the Figure 4. Consequently,

$$\lim_{n \rightarrow \infty} \mathbb{E}[\text{Tr}(M^5) \text{Tr}(\bar{M}^5)] = \lim_{n \rightarrow \infty} \frac{2 \times 5n^5 + o(n^5)}{n^5} = 2 \times 5.$$

Now, we generalize the above result in the following lemma.

**Lemma 7.5.** *Let  $M$  the random matrix as defined in Definition 3.1 satisfying the Condition 3.3. Then we have*

$$\mathbb{E}[\text{Tr}(M^k) \text{Tr}(\bar{M}^k)] = 2k + o(1),$$

if  $k < n^\gamma$  for  $\gamma < 1/20$ .

*Proof.* Let us expand  $\mathbb{E}[\text{Tr}(M^k) \text{Tr}(\bar{M}^k)]$  as follows;

$$\begin{aligned} \mathbb{E}[\text{Tr}(M^k) \text{Tr}(\bar{M}^k)] &= \frac{1}{n^k} \mathbb{E} \left[ \sum_{\substack{i_1, i_2, \dots, i_k \\ j_1, j_2, \dots, j_k=1}}^n (x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_k i_1}) (\bar{x}_{j_1 j_2} \bar{x}_{j_2 j_3} \cdots \bar{x}_{j_k j_1}) \right] \\ &= \frac{1}{n^k} \sum_{\substack{i_1, i_2, \dots, i_k \\ j_1, j_2, \dots, j_k=1}}^n \mathbb{E} [(x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_k i_1}) (\bar{x}_{j_1 j_2} \bar{x}_{j_2 j_3} \cdots \bar{x}_{j_k j_1})]. \end{aligned}$$

The main contribution will come from two possible cases, such as (a) the chain  $i_1 \rightarrow i_2 \rightarrow \cdots \rightarrow i_k \rightarrow i_1$  overlaps completely with the chain  $j_1 \rightarrow j_2 \rightarrow \cdots \rightarrow j_k \rightarrow j_1$ , or (b) it overlaps with the dual version of the second chain, which is  $n+1-j_1 \rightarrow n+1-j_2 \rightarrow \cdots \rightarrow n+1-j_k \rightarrow n+1-j_1$ . In either cases, we have  $k$  many possible ways to start the second chain. Once the chains overlap, it reduces to the product of terms of the form  $\mathbb{E}[|x_{ij}|^2] (= 1)$ . Such complete overlapping will contribute  $2kn^k$ .

Now, we argue that if the chains do not overlap completely, then the contribution is negligible. Let us assume that only  $p$  random variables from the the first chain overlap with  $p$  random variables in the second chain. In that case, the overlapping part contributes  $2pn^p$ . Additionally, the extra  $k-p$  random variables from the first and the second chain may contribute at most  $n^{(k-p)/3}(k-p)^{3(k-p)}$  each. This is a consequence of the result stated in the Lemma 7.4 with a chain of length  $k-p$ . Note that in the Lemma 7.4, we had the factor  $n^{-k/6}$ , where  $k$  was the length of the chain, because the trace sum was already divided by  $n^{k/2}$ . Here, length of the chain is  $k-p$ , but the factor  $n^{-(k-p)/2}$  is not taken into account yet. Consequently, we would get a contribution of  $n^{(k-p)/3}$ . However, to keep the continuity of the full chain (e.g.  $x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_k i_1}$ ), one of the index among  $k-p$  variables is already determined by the remaining  $p$  variables which are matched with  $p$  variables of the another chain (e.g.  $x_{j_1 j_2} x_{j_2 j_3} \cdots x_{j_k j_1}$ ). As a result, we have the factor  $n^{(k-p-1)/3}$ . Hence in such a case, the combined contribution turns out to be at most

$$(7.1) \quad 2pn^p n^{2(k-p-1)/3} (k-p)^{6(k-p)} = n^k \times 2pn^{-(k-p+2)/3} (k-p)^{6(k-p)}.$$

Additionally, the starting points of the  $p$ -chains can be set to one of indices among  $i_1, i_2, \dots, i_k$  (for the first chain) or  $j_1, j_2, \dots, j_k$  (for the second chain). Therefore, by multiplying a factor of  $k^2$  and summing up over all possible values of  $p$ , we see that if the full chains do not overlap, then we have the following contribution

$$(7.2) \quad n^k \sum_{p=1}^{k-3} 2pk^2 n^{-(k-p+2)/3} (k-p)^{6(k-p)}.$$

Note that  $p$  can not be  $k-1$  or  $k-2$ . Because in that case, the remaining chain would be of length one or two, and from the Condition 3.3, we have  $\mathbb{E}[x_{ij}] = 0 = \mathbb{E}[x_{ij}^2]$ . The above sum is  $o(n^k)$  if  $k < n^\gamma$  for  $\gamma < 1/20$ . Thus, we have

$$\begin{aligned}
& \mathbb{E}[\text{Tr}(M^k) \text{Tr}(\bar{M}^k)] \\
&= \frac{1}{n^k} \sum_{\substack{i_1, i_2, \dots, i_k, \\ j_1, j_2, \dots, j_k=1}}^n \mathbb{E}[(x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_k i_1})(\bar{x}_{j_1 j_2} \bar{x}_{j_2 j_3} \cdots \bar{x}_{j_k j_1})] \\
&= \frac{2kn^k}{n^k} + o(1) \\
&= 2k + o(1).
\end{aligned}$$

□

**Case 3 (Partial chain merging):** If the chain lengths are not equal, then we have a partial chain merging, which is explained in the following lemma.

**Lemma 7.6.** *Let  $M$  be the random matrix as defined in Definition 3.1, and satisfying the Condition 3.3. Then for  $k \neq l$ , we have*

$$\mathbb{E}[\text{Tr}(M^k) \text{Tr}(\bar{M}^l)] \leq 2kln^{-|l-k|/6}|l-k|^{3|l-k|} + o(1),$$

if  $k, l < n^\gamma$  for  $\gamma < 1/20$ .

*Proof.* Without loss of generality, let us assume that  $k < l$ . We adopt the methodology from Lemma 7.5. Let a sub chain of length  $p$  from the first chain  $x_{i_1 i_2} x_{i_2 i_3} \cdots x_{i_k i_1}$  gets matched to a sub chain of length  $p$  from the second chain  $x_{j_1 j_2} x_{j_2 j_3} \cdots x_{j_l j_1}$ . In that case, following the equations (7.1) and (7.2), we have

$$\begin{aligned}
\mathbb{E}[\text{Tr} M^k \bar{M}^l] &\leq \frac{1}{n^{(k+l)/2}} \sum_{p=1}^{k-3} 2pn^p \times kn^{(k-p-1)/3}(k-p)^{3(k-p)} \times ln^{(l-p-1)/3}(l-p)^{3(l-p)} \\
&+ \frac{1}{n^{(k+l)/2}} 2kn^k \times ln^{(l-k-1)/3}(l-k)^{3(l-k)} \\
&\leq n^{-(l+k)/2} \sum_{p=1}^k 2pkl n^{(k+l+p-2)/3}(k-p)^{3(k-p)}(l-p)^{3(l-p)} \\
&+ 2kln^{-(l-k)/6}(l-k)^{3(l-k)} \\
&\leq 2kln^{-(l-k)/6}(l-k)^{3(l-k)} + o(1).
\end{aligned}$$

The  $o(1)$  in the last inequality is a consequence of  $k, l \leq n^\gamma$ ,  $\gamma < 1/20$ . □

We summarise all the above cases as follows.

**Remark 7.7.** *While calculating  $\mathbb{E}[\text{Tr}(M^k) \text{Tr}(\bar{M}^l)]$ , the significant contribution comes only when  $k = l$ , see Lemma 7.5. From Lemma 7.4 and Lemma 7.6, we conclude that if  $k \neq l$ , although intra-chain merging and partial-chain merging are possible strategies, asymptotically they provide zero contributions. In particular,  $\mathbb{E}[\text{Tr}(M^k) \text{Tr}(\bar{M}^l)] = O(|l-k|^{3|l-k|} n^{-|l-k|/6})$  for  $k, l \leq n^{1/20}$ .*

Now, we are ready to summarize the above results into the proof of Proposition 7.1.



*Proof of Proposition 7.1.* We may write the variance as

$$\begin{aligned}
& \text{Var}(\text{Tr}(P_d(M))) \\
&= \text{Var}\left(\sum_{k=1}^d a_k \text{Tr}(M^k)\right) \\
&= \sum_{k,l=1}^d a_k \bar{a}_l \text{Cov}(\text{Tr}(M^k), \text{Tr}(M^l)) \\
&= \sum_{k,l=1}^d a_k \bar{a}_l (\mathbb{E}[\text{Tr}(M^k) \text{Tr}(\bar{M}^l)] - \mathbb{E}[\text{Tr}(M^k)] \mathbb{E}[\text{Tr}(\bar{M}^l)]) \\
(7.3) \quad &= \sum_{\substack{k \neq l \\ k,l=1}}^d 0 + \sum_{k=1}^d 2k |a_k|^2 + o(1) \\
&= \sum_{k=1}^d 2k |a_k|^2 + o(1).
\end{aligned}$$

The equality in (7.3) follows from the Lemmas 7.4, 7.5, and Remark 7.7.  $\square$

**7.4. Covariance kernel of  $\text{Tr} \hat{\mathcal{R}}_z(M)$ .** In this subsection, we find the covariance kernel of the Gaussian process as mentioned in Proposition 6.1. As defined at the beginning of Section 6, let  $\rho = \max\{\rho_1, \rho_2\}$ . Now, on the event  $\Omega_n$ , we can expand  $\text{Tr} \hat{\mathcal{R}}_z(M)$  on the boundary  $\partial \mathbb{D}_{\rho+\tau}$  as follows

$$\text{Tr} \hat{\mathcal{R}}_z(M) = \frac{n}{z} + \sum_{k=1}^{\lfloor \log^2 n \rfloor - 1} z^{-k-1} \text{Tr} M^k + \frac{1}{z^{\lfloor \log^2 n \rfloor + 1}} \text{Tr}[\hat{\mathcal{R}}_z(M) M^{\lfloor \log^2 n \rfloor}].$$

We see that on the event  $\Omega_n$ , the last term is bounded by  $\frac{n}{\tau}(\rho/|z|)^{\lfloor \log^2 n \rfloor} = o(1)$ . Hence, using Remark 7.7, we have

$$\mathbb{E}[\text{Tr} M^k \text{Tr} \bar{M}^l] = O(|k-l|^{3|k-l|} n^{-|k-l|/6}).$$

Furthermore, since  $(x^3 n^{-1/6})^x$  is a decreasing function of  $x$  for  $1 \leq x \leq n^{1/18}/e$ , we can estimate

$$\begin{aligned}
& \sum_{k \neq l}^{\lfloor \log^2 n \rfloor} |k-l|^{3|k-l|} n^{-|k-l|/6} \\
& \leq 2 \log^2 n \sum_{m=1}^{\lfloor \log^2 n \rfloor} m^{3m} n^{-m/6} \\
& \leq \frac{2 \log^4 n}{n^{1/6}} = o(1).
\end{aligned}$$

As a result,  $\sum_{k \neq l=1}^{\lfloor \log^2 n \rfloor} z^{(-k-1)} \bar{z}^{(-l-1)} \mathbb{E}[\text{Tr } M^k \text{Tr } \bar{M}^l] = o(1)$ . Hence, by using Lemma 7.5, we obtain

$$\begin{aligned}
& \mathbb{E}[\text{Tr } \hat{\mathcal{R}}_z^\circ(M) \text{Tr } \hat{\mathcal{R}}_{\bar{\eta}}^\circ(\bar{M})] \\
&= \left\{ \mathbb{E}[\text{Tr } \hat{\mathcal{R}}_z(M) \text{Tr } \hat{\mathcal{R}}_{\bar{\eta}}(\bar{M})] - \frac{n^2}{z\bar{\eta}} \right\} + o(1) \\
&= \sum_{k=1}^{\lfloor \log^2 n \rfloor} (z\bar{\eta})^{-k-1} \mathbb{E}[M^k \bar{M}^k] + o(1) \\
&= 2 \sum_{k=1}^{\lfloor \log^2 n \rfloor} k (z\bar{\eta})^{-k-1} + o(1) \\
&= 2 \left( \frac{1}{z\bar{\eta}} \right)^2 \left( 1 - \frac{1}{z\bar{\eta}} \right)^{-2} \\
&= 2 (1 - z\bar{\eta})^{-2}.
\end{aligned}$$

This completes the proof of the Proposition 6.1.

#### APPENDIX A.

**Lemma A.1.** [17, Lemma A.3] *Let  $M$  be a periodic random band matrix with bandwidth  $b_n$  and variance profile  $\omega$ . The matrix  $M$  is of the form  $M = \frac{1}{\sqrt{2b_n+1}}(X \circ W)$ , where  $X \circ W$  denotes the entrywise hadamarad product between the matrices  $X$  and  $W$ . Assume that the entries of  $X$  are i.i.d. random variables with mean zero variance one, and satisfy the Poincaré inequality with constant  $\alpha$ . Moreover,  $W_{ij} = \sqrt{\omega\left(\frac{i-j}{2b_n+1}\right)}$ , where  $\omega : \mathbb{R} \rightarrow [0, 1]$  is a piece-wise continuous periodic function with periodicity 1. In addition,  $\omega$  is continuous at 0, and  $\int_{-1/2}^{1/2} \omega(x) dx = 1$ . Then there exists  $\rho \geq 1$  such that*

$$\mathbb{P}(\|M\| > \rho/4 + t) \leq K \exp\left(-\sqrt{\frac{\alpha(2b_n+1)}{2\omega}}t\right), \quad \forall t > 0,$$

where  $K$  is a constant, which does not depend on  $n$ .

In our case, we have a matrix with i.i.d. entries whose entries satisfy the Poincaré inequality. Therefore, by taking the variance profile  $\omega \equiv 1$ , and using the above lemma we can have the following.

**Lemma A.2.** *Let  $T$  be a random matrix with  $m \times m$  dimension having i.i.d. entries with mean zero and variance one. Moreover, assume that the entries satisfy the Poincaré inequality with constant  $\alpha$ . Then there exists  $\rho \geq 1$  such that*

$$\mathbb{P}(\|T/\sqrt{m}\| > \rho/4 + t) \leq K \exp\left(-\sqrt{\frac{\alpha m}{2}}t\right), \quad \forall t > 0,$$

where  $K > 0$  is a universal constant, which does not depend on the size of the matrix and the bandwidth.

**Lemma A.3.** [27, Lemma 2.2] *Let  $M = [m_{ij}]$  be an  $n \times n$  random matrix with i.i.d. complex valued random variables satisfying (i)  $\mathbb{E}[m_{ij}] = 0 = \mathbb{E}[m_{ij}^2]$ , (ii)  $\mathbb{E}[|m_{ij}|^k] \leq k^{\alpha k}$  for some  $\alpha > 0$  and for all  $k > 2$ , (iii)  $\Re(m_{ij})$  and  $\Im(m_{ij})$  have a bounded joint density. Then for any  $p \in [1, 2)$ , and any  $z \in \mathbb{C}$ ,*

$$\mathbb{E}[|\operatorname{Tr} R_z(M)|^p] \leq C(p)n^{3p/2+2},$$

where the constant  $C(p)$  depends on  $p$ , but  $C(p) \rightarrow \infty$  as  $p \uparrow 2$ .

We note down the essential ingredient of this article, which martingale CLT as follows.

**Lemma A.4.** [7, Theorem 35.12] *Let  $\{\psi_{n,k}\}_{1 \leq k \leq n}$  be a martingale difference sequence with respect to a filtration  $\{\mathcal{F}_{k,n}\}_{1 \leq k \leq n}$ . Suppose for any  $\gamma > 0$ ,*

- (i)  $\lim_{n \rightarrow \infty} \sum_{k=1}^n \mathbb{E}[\psi_{n,k}^2 \mathbf{1}_{|\psi_{n,k}| > \gamma}] = 0$ .
  - (ii)  $\sum_{k=1}^n \mathbb{E}[\psi_{n,k}^2 | \mathcal{F}_{n,k-1}] \xrightarrow{p} \sigma^2$  as  $n \rightarrow \infty$ .
- Then  $\sum_{k=1}^n \psi_{n,k} \xrightarrow{d} \mathcal{N}_1(0, \sigma^2)$ .

**Lemma A.5** (Sherman-Morrison formula, [31]). *Let  $M$  and  $M + ve_k^t$  be two invertible matrices, where  $v \in \mathbb{C}^n$ . Then*

$$(M + ve_k^t)^{-1}v = \frac{M^{-1}v}{1 + e_k^t M^{-1}v}.$$

## REFERENCES

- [1] K. Adhikari and K. Saha. Fluctuations of eigenvalues of patterned random matrices. *Journal of Mathematical Physics*, 58(6):063301, 2017.
- [2] K. Adhikari and K. Saha. Universality in the fluctuation of eigenvalues of random circulant matrices. *Statistics & Probability Letters*, 138:1–8, 2018.
- [3] G. W. Anderson, A. Guionnet, and O. Zeitouni. *An introduction to random matrices*. Number 118. Cambridge university press, 2010.
- [4] G. W. Anderson and O. Zeitouni. A clt for a band matrix model. *Probability Theory and Related Fields*, 134(2):283–338, 2006.
- [5] Z. D. Bai and J. W. Silverstein. Clt for linear spectral statistics of large-dimensional sample covariance matrices. In *Advances In Statistics*, pages 281–333. World Scientific, 2008.
- [6] Z.-J. Bai. The inverse eigenproblem of centrosymmetric matrices with a submatrix constraint and its approximation. *SIAM Journal on Matrix Analysis and Applications*, 26(4):1100–1114, 2005.
- [7] P. Billingsley. *Probability and measure*. John Wiley & Sons, 2017.
- [8] L. Cao, D. McLaren, and S. Plosker. Centrosymmetric stochastic matrices. *Linear and Multilinear Algebra*, 70(3):449–464, 2022.
- [9] R. D. Chatham, M. Doyle, R. J. Jeffers, W. A. Kosters, R. D. Skaggs, and J. Ward. Centrosymmetric solutions to chessboard separation problems. *Bulletin of the Institute of Combinatorics and its Applications*, 65:6–26, 2012.
- [10] S. Chatterjee. Fluctuations of eigenvalues and second order poincaré inequalities. *Probability Theory and Related Fields*, 143(1):1–40, 2009.
- [11] G. Cipolloni. Fluctuations in the spectrum of non-hermitian iid matrices. *Journal of Mathematical Physics*, 63(5), 2022.
- [12] G. Cipolloni, L. Erdős, and D. Schröder. Central limit theorem for linear eigenvalue statistics of non-hermitian random matrices. *Communications on Pure and Applied Mathematics*, 76(5):946–1034, 2023.
- [13] G. Cipolloni, L. Erdős, and D. Schröder. Fluctuation around the circular law for random matrices with real entries. *Electronic Journal of Probability*, 26(none):1 – 61, 2021.
- [14] J. B. Conway. *Functions of one complex variable II*, volume 159. Springer Science & Business Media, 2012.

- [15] A. B. Cruse. Some combinatorial properties of centrosymmetric matrices. *Linear Algebra and its Applications*, 16(1):65–77, 1977.
- [16] P. J. Forrester. A review of exact results for fluctuation formulas in random matrix theory. *Probability Surveys*, 20:170–225, 2023.
- [17] I. Jana. Clt for non-hermitian random band matrices with variance profiles. *Journal of Statistical Physics*, 187(2):13, 2022.
- [18] I. Jana, K. Saha, and A. Soshnikov. Fluctuations of linear eigenvalue statistics of random band matrices. *Theory of Probability & Its Applications*, 60(3):407–443, 2016.
- [19] K. Johansson. On fluctuations of eigenvalues of random Hermitian matrices. *Duke Mathematical Journal*, 91(1):151 – 204, 1998.
- [20] D. Jonsson. Some limit theorems for the eigenvalues of a sample covariance matrix. *Journal of Multivariate Analysis*, 12:1–38, 1982.
- [21] P. Kopel. Linear statistics of non-hermitian matrices matching the real or complex ginibre ensemble to four moments. *arXiv preprint arXiv:1510.02987*, 2015.
- [22] L. Li and A. Soshnikov. Central limit theorem for linear statistics of eigenvalues of band random matrices. *Random Matrices: Theory and Applications*, 02(04):1350009, 2013.
- [23] D. Liu, X. Sun, and Z. Wang. Fluctuations of eigenvalues for random toeplitz and related matrices. *Electronic Journal of Probability*, 17:1–22, 2012.
- [24] A. Lytova and L. Pastur. Central limit theorem for linear eigenvalue statistics of random matrices with independent entries. *The Annals of Probability*, 37(5):1778 – 1840, 2009.
- [25] S. N. Maurya and K. Saha. Fluctuations of linear eigenvalue statistics of reverse circulant and symmetric circulant matrices with independent entries. *Journal of Mathematical Physics*, 62(4):043506, 2021.
- [26] S. O’Rourke and D. Renfrew. Central limit theorem for linear eigenvalue statistics of elliptic random matrices. *Journal of Theoretical Probability*, 29:1121–1191, 2016.
- [27] B. Rider and J. W. Silverstein. Gaussian fluctuations for non-Hermitian random matrix ensembles. *The Annals of Probability*, 34(6):2118 – 2143, 2006.
- [28] O. Rojo and H. Rojo. Some results on symmetric circulant matrices and on symmetric centrosymmetric matrices. *Linear algebra and its applications*, 392:211–233, 2004.
- [29] M. Shcherbina. Central limit theorem for linear eigenvalue statistics of the wigner and sample covariance random matrices. *Journal of Mathematical Physics, Analysis, Geometry*, 7(2):176–192, 2011.
- [30] M. Shcherbina. On fluctuations of eigenvalues of random band matrices. *Journal of Statistical Physics*, 161(1):73–90, 2015.
- [31] J. Sherman and W. J. Morrison. Adjustment of an inverse matrix corresponding to a change in one element of a given matrix. *The Annals of Mathematical Statistics*, 21(1):124–127, 1950.
- [32] Y. Sinai and A. Soshnikov. Central limit theorem for traces of large random symmetric matrices with independent matrix elements. *Boletim da Sociedade Brasileira de Matemática-Bulletin/Brazilian Mathematical Society*, 29(1):1–24, 1998.
- [33] D. Tao and M. Yasuda. A spectral characterization of generalized real symmetric centrosymmetric and generalized real symmetric skew-centrosymmetric matrices. *SIAM journal on matrix analysis and applications*, 23(3):885–895, 2002.
- [34] T. Tao and V. Vu. Random matrices: the circular law. *Communications in Contemporary Mathematics*, 10(02):261–307, 2008.
- [35] J. R. Weaver. Centrosymmetric (cross-symmetric) matrices, their basic properties, eigenvalues, and eigenvectors. *The American Mathematical Monthly*, 92(10):711–717, 1985.

INDIAN INSTITUTE OF TECHNOLOGY, BHUBANESWAR  
 Email address: [ijana\[at\]iitbbs.ac.in](mailto:ijana[at]iitbbs.ac.in)

INDIAN INSTITUTE OF TECHNOLOGY, BHUBANESWAR  
 Email address: [s21ma09007\[at\]iitbbs.ac.in](mailto:s21ma09007[at]iitbbs.ac.in)