

Mass and decay width of T_{ccs} from symmetries

Mitsuru Tanaka,^{1,*} Yasuhiro Yamaguchi,^{1,2,3,†} and Masayasu Harada^{2,1,4,‡}

¹*Department of Physics, Nagoya University, Nagoya 464-8602, Japan*

²*Kobayashi-Maskawa Institute for the Origin of Particles and the Universe, Nagoya University, Nagoya, 464-8602, Japan*

³*Meson Science Laboratory, Cluster for Pioneering Research,
RIKEN, Hirosawa, Wako, Saitama 351-0198, Japan*

⁴*Advanced Science Research Center, Japan Atomic Energy Agency, Tokai 319-1195, Japan*

(Dated: September 4, 2024)

We analyze the mass and width of the doubly heavy tetraquark T_{ccs} composed of a heavy diquark and a light-quark cloud with strangeness with assuming that a color antitriplet heavy diquark is a dominant component of the doubly charmed tetraquarks T_{cc} and T_{ccs} . We construct an effective Lagrangian for masses of heavy hadrons based on the superflavor symmetry between the doubly heavy tetraquarks and the singly heavy baryons by including the terms that simultaneously break the heavy-quark and light-flavor symmetries, and predict the mass of T_{ccs} as $M(T_{ccs}) = 4047 \pm 11$ MeV. The comparison of this prediction with future experimental observation will give a clue to understand the color structure of the heavy diquark. We also predict the mass of Ω_{cc} as $M(\Omega_{cc}) = 3706_{-15}^{+14}$ MeV. We next calculate the decay width of T_{ccs} , based on solely the light-flavor symmetry, as $\Gamma(T_{ccs}) = 42 \pm 24$ MeV.

Keywords: superflavor symmetry, heavy quark symmetry, flavor symmetry, diquark, doubly heavy tetraquark

I. INTRODUCTION

The discovery of $X(3872)$ [1] marked the beginning of numerous exotic hadron discoveries in the heavy-quark sectors, yet their structure remains poorly understood. Hadrons with exotic structures beyond ordinary baryons (qqq) and mesons ($q\bar{q}$) were already indicated by Gell-Mann and Zweig in the 1960s [2–6]. Possible structures of the multi-quark state being a color singlet have been discussed in the literatures (See, for reviews, e.g. Refs. [7, 8].) The compact multi-quark has been investigated as a color singlet state of few-body multi-quark systems by the constituent quark model, etc. (e.g., Refs. [9–16].) The emergence of the hadronic molecules as a deuteronlike state, discussed as a deuson in Ref. [17], is expected near the thresholds. In fact many candidates of a hadronic molecule have been reported in the experimental studies as the XYZ tetraquarks being a meson-meson state and the P_c pentaquarks being a meson-baryon one. Investigating the exotic structures would lead to an understanding of the QCD phenomena such as color confinement.

The doubly charmed tetraquark T_{cc}^+ was reported in the LHCb experiment in 2021 [18, 19]. The reported state is consistent with a genuine exotic hadron having a flavor structure $cc\bar{u}\bar{d}$. The spin and parity of T_{cc}^+ are determined to be $J^P = 1^+$, and the LHCb considers T_{cc}^+ as an isoscalar. The mass of T_{cc}^+ is 3874.817 MeV close to the D^0D^{*+} threshold. The decay to $D^0D^0\pi^+$ has been confirmed, with a decay width of 410 keV [18] or 48 keV [19]. Furthermore, the LHCb analysis supports

that T_{cc}^+ decays to $D^0D^0\pi^+$ via the intermediate state D^{*+} .

Since the discovery of T_{cc}^+ , research on the doubly heavy tetraquarks (DHTs) has been actively conducted [7]. However at present, no clear answer has been obtained regarding the structure of DHTs. For example, analyses based on the hadronic molecular model, which assumes that T_{cc}^+ is a loosely bound state with D and D^* , have been conducted [20–32]. This is due to the fact that the mass of T_{cc}^+ resides in the vicinity of the DD^* threshold. On the other hand, a compact tetraquark structure of DHT is considered [33], based on the diquark picture proposed by Jaffe [34]. In addition, DHTs have been discussed in various approaches such as the string model [35], the QCD sum rules [36–38], and the lattice QCD [39–42].

Symmetries such as flavor symmetry and chiral symmetry play a crucial role in the classification of hadronic states. For exotic hadrons including heavy-quarks, symmetries that emerge in the heavy quark limit, such as heavy-quark symmetry (HQS) (see, e.g., Ref. [43]) and superflavor symmetry [44–46], are considered potentially useful for understanding the structures of exotics. The superflavor symmetry emerges in an exchange of an anti-heavy quark and a heavy diquark with the same color configuration $\mathbf{\bar{3}}$ in the heavy-quark limit. Because of the sufficiently large mass of the heavy diquark, the spin-dependent color magnetic force is negligible, and does not contribute in the heavy-quark limit. In this context, the heavy diquark and an antiheavy quark behave as the static color $\mathbf{\bar{3}}$ source and are equivalent in terms of color interaction. The property of hadrons remains invariant under the interchange of the heavy diquark and antiheavy quark. We will refer to hadrons related under this superflavor symmetry as superflavor partners. For instance, an anti-heavy meson(HM) $\bar{Q}q$ and a doubly heavy baryon(DHB) QQq are superflavor partners.

* tanaka@hken.phys.nagoya-u.ac.jp

† yamaguchi@hken.phys.nagoya-u.ac.jp

‡ harada@hken.phys.nagoya-u.ac.jp

Another color representation for diquarks, 6, is also allowed in DHTs but not in ordinary hadrons. If diquarks take the color 6 representation, the superflavor symmetry does not arise. Hence, superflavor symmetry might shed light on the color configuration of exotic hadrons. We think that understanding the color configuration of the diquark in DHTs is important because the interaction changes according to the color representation.

To analyze the mass spectrum of DHTs in terms of superflavor symmetry, this study assumes that DHTs consist of a color $\bar{3}$ heavy diquark, treated as a spatially compact object, and a light-quark cloud surrounding it. Although it may exist in a mixed state within a DHT, we also assume that the color antitriplet state is the dominant state of the diquark because the color $\bar{3}$ diquark is likely realized in the ground state [7, 47, 48]. Here, “being spatially compact” means that the heavy diquark can be approximated as a pointlike particle, with no radial excitation occurring between two heavy quarks. The analyses of T_{bb} , being the bottom counterpart of T_{cc} , based on the quark model [48–50] suggest that the distance between the two bottom quarks is shorter compared to other quark distances. This observation appears to support the notion that T_{bb} is composed of a heavy diquark and a light-quark cloud. In the present analysis, we assume that T_{cc} also holds such heavy diquark structure. For simplicity, this analysis focuses solely on diquarks where both heavy quarks are of the same flavor. Under these conditions, a color $\bar{3}$ diquark possesses spin one ($S_{QQ} = 1$), whereas a color 6 diquark has spin zero ($S_{QQ} = 0$).

If T_{cc}^+ is the superflavor partner of Λ_c^+ , we can naturally expect the existence of T_{ccs} as the superflavor partner of Ξ_c , which belongs to the same flavor multiplet as Λ_c^+ . We consider that investigation of T_{ccs} is useful to understand the nature not only of T_{ccs} itself but also of T_{cc}^+ . We analyze T_{ccs} by using the experimental result of T_{cc}^+ as an input. Thus if the results obtained in this paper are eventually consistent with results in future experiments, it can be interpreted that the color antitriplet state is dominant in DHTs T_{cc}^+ and T_{ccs} . In the following, we first derive simple mass relations assuming that the heavy diquark is spatially compact and color antitriplet state together with superflavor symmetry. We note that the relations agree with the ones derived in Ref. [51]. However, we find a discrepancy between the prediction by the simple mass relation and the recent experimental data of T_{cc}^+ . Thus, we invent the improved mass relations with correction terms violating the symmetries and a mixing term of color $\bar{3}$ and 6 states of the cc diquark. Then, we obtain new relations among HMs, DHBs, singly heavy baryons (SHBs) and DHTs, and predict the mass of T_{ccs} . Furthermore, we also predict the decay width of T_{ccs} from the SU(3) flavor symmetry.

This paper is organized as follows. In Sec. II, we derive the simple mass relations among singly heavy and doubly heavy hadrons based on the heavy quark and superflavor symmetries. We point out the existence of a discrepancy

between the theoretical formulas and the experimental results. In Sec. III, we construct the effective Lagrangians including corrections and obtain the improved mass relations. The mass of T_{ccs} and in addition, the one of Ω_{cc} are predicted. In Sec. IV, the decay width of T_{ccs} is predicted by using the effective Lagrangian approach respecting the flavor symmetry. Finally, Sec. V is devoted to the summary.

II. SIMPLE MASS RELATIONS FROM SUPERFLAVOR AND LIGHT-FLAVOR SYMMETRIES

In this section, we first introduce simple mass relations among superflavor partners, primarily derived from superflavor, heavy quark, and light-flavor symmetries. Then, we demonstrate that these mass relations are somewhat broken among real hadrons, indicating the need for improvements to the mass relations. Under the superflavor symmetry, a DHT is related to an anti-SHB, and a DHB to an anti-HM. Considering that T_{cc}^+ has isospin $I = 0$, the superflavor partners of T_{cc}^+ , T_{ccs}^+ , and T_{ccs}^{++} are, respectively, the antibaryons of Λ_c^+ , Ξ_c^+ , and Ξ_c^0 . Similarly, the superflavor partners of DHBs such as Ξ_{cc} and Ω_{cc} are the HMs, \bar{D} and \bar{D}_s , respectively.

We first study the relation between the masses of the DHBs and HMs in terms of the superflavor symmetry, and then extend the analysis to the DHTs and SHBs. Based on the HQS, we divide a heavy hadron into a heavy object and light-quark cloud which includes the interaction between them. Therefore, the dynamics of these hadrons are determined by the properties of the light-quark cloud. As a result, the masses of heavy hadrons treated in the present analysis are expressed as a sum of the mass of heavy objects and the energy of the light-quark cloud in the heavy-quark limit. Let us first estimate the masses of HMs. Here, we consider the spin average of the doublet under the HQS. We note that due to the spin average, the first-order term in $1/m_Q$ expansion that breaks only the heavy-quark spin symmetry does not appear in the mass formulas. As a result, the spin-averaged mass can be expressed as

$$M_{\text{ave}}(\bar{Q}q) = M(\bar{Q}) + E(q), \quad (1)$$

where $M(\bar{Q})$ ($\bar{Q} = \bar{c}, \bar{b}$) is the mass of antiheavy quark and $E(q)$ ($q = u, d, s$) denotes the contribution from the light-quark cloud. From Eq. (1), we obtain the meson mass difference between flavor partners as

$$M_{\text{ave}}(\bar{Q}s) - M_{\text{ave}}(\bar{Q}n) = E(s) - E(n), \quad (2)$$

where $n = u, d$.

In the DHBs, the heavy diquark QQ takes the color $\bar{3}$ representation, so that the anti-HMs and the DHBs include the common light-quark cloud in the heavy-quark limit. Thus, the spin-averaged mass of the doublet members of DHBs is expressed in a similar formula as for the

HMs:

$$M_{\text{ave}}(QQq) = M(QQ) + E(q) . \quad (3)$$

We stress that the term $E(q)$ is common in Eqs. (1) and (3). The mass difference between flavor partners is $E(\bar{q}_1) - E(\bar{q}_2)$, where $q_1, q_2 = u, d, s$. This leads to the following mass relation [52]:

$$\begin{aligned} M_{\text{ave}}(QQq_1) - M_{\text{ave}}(QQq_2) \\ = M_{\text{ave}}(\bar{Q}q_1) - M_{\text{ave}}(\bar{Q}q_2) . \end{aligned} \quad (4)$$

This implies that the mass differences between flavor partners are the same in the superflavor partners.

Next, we consider the masses of anti-SHBs and DHTs. By a similar argument as above, the mass of an anti-SHB ($Q\bar{q}\bar{q}$) is expressed as

$$M(\bar{Q}\bar{q}_1\bar{q}_2) = M(\bar{Q}) + E(\bar{q}_1\bar{q}_2) . \quad (5)$$

As we stated above, we assume that two charm quarks inside T_{cc} form a compact diquark and that the diquark belonging to the color $\bar{3}$ representation is dominated. Therefore, in the heavy-quark limit, the DHT shares a common light-quark cloud with the anti-SHB according to the superflavor symmetry. Consequently, the mass of DHT is expressed as

$$M(QQ\bar{q}_1\bar{q}_2) = M(QQ) + E(\bar{q}_1\bar{q}_2) . \quad (6)$$

From the mass formulas in Eqs. (5) and (6), we obtain the following mass relation corresponding to the mass relation (4):

$$\begin{aligned} M(QQ\bar{q}_1\bar{q}_2) - M(QQ\bar{q}_3\bar{q}_4) \\ = M(\bar{Q}\bar{q}_1\bar{q}_2) - M(\bar{Q}\bar{q}_3\bar{q}_4) . \end{aligned} \quad (7)$$

where $\bar{q}_i = \bar{u}, \bar{d}, \bar{s}$ ($i = 1, 2, 3, 4$).

We further combine Eqs. (1), (3), (5) and (6) to derive the following simple mass relation:

$$\begin{aligned} M(QQ\bar{q}_1\bar{q}_2) - M(\bar{Q}\bar{q}_1\bar{q}_2) \\ = M_{\text{ave}}(QQq) - M_{\text{ave}}(\bar{Q}q) . \end{aligned} \quad (8)$$

Now, we compare the obtained simple mass relations with existing experimental data. We first note that the relation (2) implies that the mass difference between D and D_s is equal to that between B and B_s , namely

$$M_{\text{ave}}(D_s) - M_{\text{ave}}(D) = M_{\text{ave}}(B_s) - M_{\text{ave}}(B) , \quad (9)$$

because the energy difference between light clouds, $E(s) - E(n)$, is independent of the heavy flavor. Similarly, we obtain the following mass relation for the masses of SHBs:

$$M(\Xi_c) - M(\Lambda_c) = M(\Xi_b) - M(\Lambda_b) . \quad (10)$$

However, from the experimental values of masses shown

Hadrons	Mass [MeV]	Input value [MeV]
T_{cc}^+	3874.74	3875
Ξ_{cc}^+	3623.0	3622
Ξ_{cc}^{++}	3621.55	3622
D^0	1864.84	1867
D^\pm	1869.66	1867
D^{*0}	2006.85	2009
$D^{*\pm}$	2010.26	2009
D_s^\pm	1968.35	1968
$D_s^{*\pm}$	2112.2	2112
Λ_c^+	2286.46	2286
Ξ_c^+	2467.71	2469
Ξ_c^0	2470.44	2469
$D^{*0}D_s^\pm$	3975.20	3979
$D^{*\pm}D_s^\pm$	3978.61	3979
$D^0D_s^{*\pm}$	3977.0	3980
$D^\pm D_s^{*\pm}$	3981.9	3980
B^\pm	5279.34	5280
B^0	5279.66	5280
B^*	5324.71	5325
B_s^0	5366.92	5367
B_s^*	5415.4	5415
Λ_b^+	5619.60	5620
Ξ_b^-	5797.0	5794
Ξ_b^0	5791.9	5794

TABLE I. Experimental values [18, 19, 53] and input values in this study.

in Table I, the mass differences are obtained as

$$M_{\text{ave}}(D_s) - M_{\text{ave}}(D) = 103 \text{ MeV}, \quad (11)$$

$$M_{\text{ave}}(B_s) - M_{\text{ave}}(B) = 90 \text{ MeV}. \quad (12)$$

and

$$M(\Xi_c) - M(\Lambda_c) = 183 \text{ MeV}, \quad (13)$$

$$M(\Xi_b) - M(\Lambda_b) = 175 \text{ MeV}. \quad (14)$$

This discrepancy implies the necessity of considering correction terms that break both the heavy-quark flavor symmetry and the light-flavor symmetry.

Similarly, applying the mass relation (8) to T_{cc} , we obtain the following mass relation:

$$M(T_{cc}^+) - M(\Lambda_c^+) = M_{\text{ave}}(\Xi_{cc}) - M_{\text{ave}}(D) . \quad (15)$$

Since Ξ_{cc}^* has not been experimentally confirmed yet, we estimate its mass using the following mass relation obtained from the superflavor symmetry [46, 54]:

$$M(\Xi_{cc}^*) - M(\Xi_{cc}) = \frac{3}{4} (M(D^*) - M(D)) , \quad (16)$$

leading to $M(\Xi_{cc}^*) = 3728 \text{ MeV}$, and thus $M_{\text{ave}}(\Xi_{cc}) = 3693 \text{ MeV}$. Substituting this value into Eq. (15), we get $M(T_{cc}^+) = 3970 \text{ MeV}$. This value is clearly different from the experimental value, 3875 MeV, which motivates us to consider the mixing between a state with a heavy diquark in the color $\bar{3}$ representation and one in the color 6 representation.

III. IMPROVED MASS RELATIONS

In this section, we construct effective Lagrangian terms for solving the problems raised in the previous section for heavy mesons and the T_{cc} . As we stated in the previous section, we need to include the terms that simultaneously break the heavy-quark flavor symmetry and SU(3)-flavor symmetry for light quarks to cure the problem of masses of heavy mesons and SHBs. For the problem of the mass of T_{cc} , we include a term leading to the mixing between the states constructed from the heavy diquark in the color $\bar{3}$ representation and the one in the color 6 representation. In addition, we include terms which break the heavy-quark spin symmetry for the HQS doublet.

At first, we define the effective DHB(QQq) fields with quantum numbers $J^P = \frac{1}{2}^+$ and $J^P = \frac{3}{2}^+$ in the heavy-quark limit by combining the heavy diquark with $J_{\text{heavy}}^P = 1^+$ to the light-quark cloud with $J_{\text{light}}^P = \frac{1}{2}^+$. This doubly heavy baryon field \mathcal{B} is expressed as

$$[\mathcal{B}]_{hh'l} = [P_+ \gamma_\mu C P_+^T]_{hh'} \psi_l^\mu, \quad (17)$$

where h and h' are spinor indices for heavy-quarks and l is the spinor index for the light quark cloud, $C = i\gamma_2\gamma_0$ is the charge conjugation matrix, and ψ^μ is the field for the heavy quark spin doublet with $J^P = (1/2^+, 3/2^+)$. We should note that the field \mathcal{B} carries the index to specify the heavy quark flavor Q . But here and henceforth, we omit the index to avoid too many indices for one field. The projection operator for heavy quark P_+ is defined as $P_+ = (1 + v^\mu \gamma_\mu)/2$, where v^μ is the velocity of the heavy-diquark. The ψ^μ field satisfies the following constraint:

$$v^\nu \gamma_\nu \psi^\mu = \psi^\mu. \quad (18)$$

It is convenient to further decompose the field ψ^μ into the $J^P = 1/2^+$ field $\psi_{1/2}$ and the $J^P = 3/2^+$ field $\psi_{3/2}^\mu$ as

$$\psi^\mu = \frac{1}{\sqrt{3}} \sigma^{\mu\nu} v_\nu \psi_{1/2} + \psi_{3/2}^\mu, \quad (19)$$

where

$$\sigma^{\mu\nu} = \frac{i}{2} [\gamma^\mu, \gamma^\nu], \quad (20)$$

and the spin-3/2 Rarita-Schwinger field $\psi_{3/2}^\mu$ satisfies

$$v_\mu \psi_{3/2}^\mu = \gamma_\mu \psi_{3/2}^\mu = 0. \quad (21)$$

For later use, we define the conjugate field $\bar{\mathcal{B}}$ for DHB as

$$[\bar{\mathcal{B}}]_{hh'l} = [\gamma_0]_{hh_1} [\mathcal{B}^\dagger]_{h_1 h_2 l_1} [\gamma_0]_{h_2 h'} [\gamma_0]_{l_1 l}. \quad (22)$$

For realizing the superflavor symmetry, we take the effective DHB field \mathcal{B} and the effective anti-HM field $\bar{H} (\sim \bar{Q}q)$ into a unified field Ψ as

$$\Psi = \begin{pmatrix} \bar{H} \\ \mathcal{B} \end{pmatrix}. \quad (23)$$

The parity transformations are given by

$$\bar{H} \rightarrow \gamma_0 \bar{H} \gamma_0, \quad (24)$$

$$[\mathcal{B}]_{hh'l} \rightarrow [\gamma_0]_{ll_1} [\gamma_0]_{hh_1} [\mathcal{B}]_{h_1 h_2 l_1} [\gamma_0]_{h_2 h'}, \quad (25)$$

where h, h', h_1, h_2 are heavy-quark spinor indices and l, l_1 are spinor indices for a light cloud. Since the field Ψ belongs to 3 representation of the SU(3)_f light flavor, the SU(3)_f transformation is given by

$$\Psi^i \rightarrow U_j^i \Psi^j, \quad (26)$$

where i, j are the light-flavor indices and $U \in \text{SU}(3)_f$.

Next, we define the effective DHT($\sim \bar{Q}Qq$) fields. As we stated in the previous section, we include two DHT fields: One is constructed from the heavy diquark with the color $\bar{3}$ representation carrying $J_{\text{heavy}}^P = 1^+$, which is combined with the light-quark cloud with $J_{\text{light}}^P = 0^+$ to make the DHT with $J^P = 1^+$. Another one is constructed from the heavy diquark carrying the color 6 representation and $J_{\text{heavy}}^P = 0^+$ combined with $J_{\text{light}}^P = 1^+$ to make the one with $J^P = 1^+$. The former one is denoted as $T_\mu^{(3)}$ and the latter as $T_\mu^{(6)}$. These fields are defined as

$$[T^{(3)}]_{hh'} = [P_+ \gamma^\mu C P_+^T]_{hh'} \phi_\mu, \quad (27)$$

$$[T_\mu^{(6)}]_{hh'} = [P_+ \gamma_5 C P_+^T]_{hh'} \varphi_\mu, \quad (28)$$

where h and h' are spinor indices for heavy quarks, and the upper indices of T , ($\bar{3}$) and (6), represent the color representation of heavy diquarks in DHTs. We note that ϕ and φ_μ stand for the annihilation operators and the same applies to T . We also note that these fields have the index of a light quark-flavor. As we said in the Introduction, we consider heavy diquarks made from two heavy quarks with the same flavor. The light-quark cloud is made from two anti light quarks in the flavor antisymmetric representation. Then, the fields T belong to 3 representation of the SU(3)_f light-flavor symmetry.

Based on the superflavor symmetry, the field $T_\mu^{(3)}$ and the effective anti-SHB field $\bar{S} (\sim \bar{Q}\bar{q})$ are arranged into a unified field Φ as

$$\Phi = \begin{pmatrix} \bar{S} \\ T^{(3)} \end{pmatrix}. \quad (29)$$

The parity transformations are given by

$$\bar{S} \rightarrow \bar{S} \gamma_0, \quad (30)$$

$$[T]_{hh'} \rightarrow [\gamma_0]_{hh_1} [T]_{h_1 h_2} [\gamma_0]_{h_2 h'}. \quad (31)$$

The conjugate fields are defined as

$$S = \gamma_0 \bar{S}^\dagger, \quad (32)$$

$$[\bar{T}]_{hh'} = [\gamma_0]_{hh_1} [\bar{T}^\dagger]_{h_1 h_2} [\gamma_0]_{h_2 h'}. \quad (33)$$

The SU(3)_f transformation is given by

$$\bar{\Phi}_{ij} \rightarrow (U^*)^k_i \bar{\Phi}_{kl} (U^\dagger)^l_j \quad (34)$$

Let us construct the effective Lagrangian for the DHT, SHB, DHB and HM. As we said in the previous section, we need to include the terms that break simultaneously the heavy-quark flavor symmetry and the light-flavor symmetry. We first note that the terms that break the heavy-quark flavor symmetry are inversely proportional to the heavy-quark mass, i.e., $\propto 1/m_Q$. We also include the term for generating the mixing between $\bar{T}_\mu^{(3)}$ and $\bar{T}_\mu^{(6)}$, which must include $1/m_Q$ since the mixing vanishes at the heavy-quark limit. For the light-flavor symmetry breaking, we use the spurion field corresponding to the light-quark mass matrix \mathcal{M} as

$$\mathcal{M} = \begin{pmatrix} m_u & & \\ & m_d & \\ & & m_s \end{pmatrix}, \quad (35)$$

where m_u , m_d and m_s are the current quark masses of u , d and s quarks, respectively. This \mathcal{M} transforms as

$$(\mathcal{M})^i_j \rightarrow (U)^i_k (\mathcal{M})^k_\ell (U^\dagger)^\ell_j. \quad (36)$$

Now, the kinetic and mass terms invariant under the heavy-quark symmetry and SU(3) light-flavor symmetry are given by

$$\begin{aligned} \mathcal{L}_0 = & -\text{tr} [\bar{\Psi} i v \cdot \partial \Psi - \Lambda_\Psi \bar{\Psi} \Psi] \\ & - \text{Tr} [\bar{\Phi} i v \cdot \partial \Phi - \Lambda_\Phi \bar{\Phi} \Phi], \end{aligned} \quad (37)$$

where Λ_Ψ and Λ_Φ are constants with mass dimension one and tr indicates that the traces in spinor space and heavy-spin space are taken, while Tr implies that the traces in the light-flavor space, in addition to the spinor space and heavy-spin space, are taken. In the following, we explicitly write the indices for light flavor while we omit the indices for heavy-quark flavor and spins of heavy quarks and light quarks.

A possible term that breaks the SU(3) light-flavor symmetry for the Ψ field is written as

$$\mathcal{L}_{\Psi\text{-br}} = + c_\Psi \text{tr} [\bar{\Psi}_i (\mathcal{M})^i_j \Psi^j], \quad (38)$$

where c_Ψ is a constant with mass dimension zero.

By noting that two light quarks in the Φ field are antisymmetric in flavor, the light-flavor breaking term is expressed as

$$\mathcal{L}_{\Phi\text{-br}} = + c_\Phi \text{tr} [\bar{\Phi}_{ij} (\mathcal{M})^j_k \Phi^{ki}], \quad (39)$$

where c_Φ is a constant with mass dimension zero.

We next construct possible terms that break both the SU(3) light-flavor symmetry and the heavy-quark flavor symmetry, as well as the heavy quark spin symmetry. We note that the terms also break the superflavor symmetry, so that the terms should be written separately for superflavor partners. The terms for \bar{H} (anti-HM) and B

(DHB) included in Ψ are expressed as

$$\begin{aligned} \mathcal{L}_{\text{HB-br}} = & - \frac{\Lambda_f}{m_Q} \text{tr} [H_i (\mathcal{M})^i_j \bar{H}^j] \\ & - \frac{\Lambda'_f}{2m_Q} \text{tr} [\bar{\mathcal{B}}_i (\mathcal{M})^i_j \mathcal{B}^j] \\ & - \frac{\Lambda_\sigma^2}{8m_Q} \text{tr} \left[H \left(\sigma_{heavy}^{\mu\nu} \right)^T \bar{H} \sigma_{\mu\nu}^{light} \right] \\ & - \frac{\Lambda_{\sigma f}}{8m_Q} \text{tr} \left[H_i \left(\sigma_{heavy}^{\mu\nu} \right)^T (\mathcal{M})^i_j \bar{H}^j \sigma_{\mu\nu}^{light} \right] \\ & - \frac{(\Lambda'_\sigma)^2}{8m_Q} \left[[\bar{\mathcal{B}}]_{h_1 h_2 l_1} \left(\sigma_{heavy}^{\mu\nu} \right)_{h_2 h_3} [\mathcal{B}]_{h_3 h_1 l_2} \left(\sigma_{\mu\nu}^{light} \right)_{l_2 l_1} \right] \\ & - \frac{\Lambda'_{\sigma f}}{8m_Q} \left[[\bar{\mathcal{B}}_i]_{h_1 h_2 l_1} \left(\sigma_{heavy}^{\mu\nu} \right)_{h_2 h_3} (\mathcal{M})^i_j \right. \\ & \quad \left. [\mathcal{B}^j]_{h_3 h_2 l_2} \left(\sigma_{\mu\nu}^{light} \right)_{l_2 l_1} \right], \end{aligned} \quad (40)$$

where Λ_f , Λ'_f , Λ_σ , $\Lambda_{\sigma f}$, Λ'_σ , $\Lambda'_{\sigma f}$ are constants with mass dimension one. We put an extra factor 1/2 in the second term since the DHB includes two heavy quarks. We note that, if we set $\Lambda_\sigma = \Lambda'_\sigma$ and $\Lambda_{\sigma f} = \Lambda'_{\sigma f}$, these two terms reproduce the mass relation (16).

Similarly, possible breaking terms for \bar{S} (anti-SHB) and T (DHT) in Φ field are expressed as

$$\begin{aligned} \mathcal{L}_{\text{ST-br}} = & - \frac{\Lambda_{\text{ff}}}{m_Q} \text{tr} [\bar{S}_{ij} (\mathcal{M})^j_k S^{ki}] \\ & - \frac{\Lambda'_{\text{ff}}}{2m_Q} \text{tr} \left[\left(T^{(\bar{3})} \right)_{ij} (\mathcal{M})^j_k \left(\bar{T}^{(\bar{3})} \right)^{ki} \right], \end{aligned} \quad (41)$$

where Λ_{ff} and Λ'_{ff} are constants with mass dimension one.

Finally, we consider a DHT field $T_\mu^{(6)}$ constructed from the heavy diquark with color $\mathbf{6}$ representation. The kinetic and mass terms are written as

$$\mathcal{L}_{T^{(6)}} = -\text{Tr} \left[\bar{T}_\mu^{(6)} (i v \cdot \partial - \Lambda_6) T^{(6)\mu} \right], \quad (42)$$

where Λ_6 is a constant with mass dimension one. The term for the mixing between two DHT fields $T_\mu^{(3)}$ and $T_\mu^{(6)}$ is expressed as

$$\mathcal{L}_{\text{mix}} = - \frac{\Lambda_{\text{mix}}^2}{2m_Q} \text{Tr} \left[\left(\bar{T}_\mu^{(6)} \right) \gamma_5 \gamma^\mu \left(T^{(3)} \right) + h.c. \right]. \quad (43)$$

From the above Lagrangian terms, the masses of HM,

DHB and SHB are modified from Eqs. (1), (3) and (5) as

$$M(\bar{Q}q) = M(\bar{Q}) + E(q) + \frac{m_q}{m_Q} \Lambda_f - \frac{3}{4} \frac{\Lambda_\sigma^2}{m_Q} - \frac{3}{4} \frac{m_q}{m_Q} \Lambda_{\sigma f}, \quad (44)$$

$$M^*(\bar{Q}q) = M(\bar{Q}) + E(q) + \frac{m_q}{m_Q} \Lambda_f + \frac{1}{4} \frac{\Lambda_{\sigma f}^2}{m_Q} + \frac{1}{4} \frac{m_q}{m_Q} \Lambda_{\sigma f}, \quad (45)$$

$$M(QQq) = M(QQ) + E(q) + \frac{m_q}{2m_Q} \Lambda'_f - \frac{1}{2} \frac{(\Lambda'_\sigma)^2}{m_Q} - \frac{1}{2} \frac{m_q}{m_Q} \Lambda'_{\sigma f}, \quad (46)$$

$$M^*(QQq) = M(QQ) + E(q) + \frac{m_q}{2m_Q} \Lambda'_f + \frac{1}{4} \frac{(\Lambda'_\sigma)^2}{m_Q} + \frac{1}{4} \frac{m_q}{m_Q} \Lambda'_{\sigma f}, \quad (47)$$

$$M(\bar{Q}\bar{q}_1\bar{q}_2) = M(\bar{Q}) + E(\bar{q}_1\bar{q}_2) + \frac{m_{q_1} + m_{q_2}}{m_Q} \Lambda_f, \quad (48)$$

where

$$E(q) = \Lambda_\Psi + c_\Psi m_q, \quad E(\bar{q}_1\bar{q}_2) = \Lambda_\Phi + c_\Phi (m_{q_1} + m_{q_2}). \quad (49)$$

The square mass matrix for two DHT fields is expressed as

$$\begin{pmatrix} M_3^2 & \frac{\Lambda_{\text{mix}}^2}{2m_Q} \sqrt{M_6 M_3} \\ \frac{\Lambda_{\text{mix}}^2}{2m_Q} \sqrt{M_6 M_3} & M_6^2 \end{pmatrix}, \quad (50)$$

where

$$M_3 = M(QQ) + E(\bar{q}_1\bar{q}_2) + \frac{m_{q_1} + m_{q_2}}{2m_Q} \Lambda'_f, \quad (51)$$

and M_6 is the mass of the $T_\mu^{(6)}$ field before mixing. By diagonalizing this matrix up to $1/m_Q$ order, the mass of the lightest DHT is obtained as

$$M(QQ\bar{q}_1\bar{q}_2) = M(QQ) + E(\bar{q}_1\bar{q}_2) + \frac{m_{q_1} + m_{q_2}}{2m_Q} \Lambda'_f - \frac{M_6}{2(M_6^2 - M_3^2)} \left(\frac{\Lambda_{\text{mix}}^2}{2m_Q} \right)^2. \quad (52)$$

By combining the above mass formulas, the simple mass

Quarks Mass [MeV]	
u, d	3.415
s	93.40
c	1270
b	4180

TABLE II. Mass of quarks [53] used as inputs.

relations (4), (7) and (8) are modified as:

$$M_{\text{ave}}(QQs) - M_{\text{ave}}(QQn) - \frac{m_s - m_n}{2m_Q} \Lambda'_f = M_{\text{ave}}(\bar{Q}s) - M_{\text{ave}}(\bar{Q}n) - \frac{m_s - m_n}{m_Q} \Lambda_f, \quad (53)$$

$$M(QQ\bar{s}\bar{n}) - M(QQ\bar{u}\bar{d}) - \frac{m_s - m_n}{2m_Q} \Lambda'_{ff} = M(\bar{Q}\bar{s}\bar{n}) - M(\bar{Q}\bar{u}\bar{d}) - \frac{m_s - m_n}{m_Q} \Lambda_{ff}, \quad (54)$$

$$M(QQ\bar{u}\bar{d}) - M(\bar{Q}\bar{u}\bar{d}) + \frac{M_6}{2(M_6^2 - M_3^2)} \left(\frac{\Lambda_{\text{mix}}^2}{2m_Q} \right)^2 - \frac{m_n}{2m_Q} (\Lambda'_{ff} - 2\Lambda_{ff}) = M_{\text{ave}}(QQn) - M_{\text{ave}}(\bar{Q}n) - \frac{m_n}{2m_Q} (\Lambda'_f - 2\Lambda_f), \quad (55)$$

where $n = u, d$ and $m_n = (m_u + m_d)/2$ is the isospin-averaged mass of up and down quarks. We note that other mass relations such as the ones between non-spin-averaged masses are obtained by recombining Eqs. (44)-(48), and (52). Furthermore, from Eqs. (44), (45) and (48), we obtain the following mass relation:

$$[M_{\text{ave}}(D_s) - M_{\text{ave}}(D)] - [M_{\text{ave}}(B_s) - M_{\text{ave}}(B)] = (m_s - m_n) \left(\frac{1}{m_c} - \frac{1}{m_b} \right) \Lambda_f, \quad (56)$$

$$[M(\Xi_c) - M(\Lambda_c)] - [M(\Xi_b) - M(\Lambda_b)] = (m_s - m_n) \left(\frac{1}{m_c} - \frac{1}{m_b} \right) \Lambda_{ff} \quad (57)$$

where $n = u, d$. Using the masses shown in Table I together with the current quark masses shown in Table II, we determine the value of parameters as

$$\Lambda_f = 272.2 \pm 70.9 \text{ MeV}, \quad (58)$$

$$\Lambda_{ff} = 157.4 \pm 70.9 \text{ MeV}, \quad (59)$$

$$\Lambda_\sigma = 423.5 \pm 25.1 \text{ MeV}, \quad (60)$$

$$\Lambda_{\sigma f} = 36.70 \pm 70.87 \text{ MeV}. \quad (61)$$

The error bars are estimated by

$$\delta\Lambda_f = \delta\Lambda_{ff} = \delta\Lambda_{\sigma f} = \frac{\Lambda_{\text{QCD}}^2}{m_Q}, \quad \delta\Lambda_\sigma = \frac{\Lambda_{\text{QCD}}^3}{2\Lambda_\sigma m_Q}, \quad (62)$$

Hadrons Mass [MeV]		
T_{ccs}	4047 ± 11	Our result
	4106	NQM [55]
T_{ccs}^+	3969 ± 8	Lattice QCD [40]

TABLE III. Theoretical predictions for the masses of T_{ccs} . NQM imply a nonrelativistic quark model.

where Λ_{QCD} is the typical scale of QCD and we set $\Lambda_{\text{QCD}} = 300 \text{ MeV}$ in this paper. The parameters Λ_σ and $\Lambda_{\sigma f}$ are determined to fit the hyperfine splittings between charmed mesons, and we confirmed that the hyperfine splittings between bottom mesons are within error bars. The other parameters Λ'_f and Λ'_{ff} cannot be determined due to lack of input hadron masses. We therefore assume that they are of the same order as Λ_f and Λ_{ff} . Specifically, we set

$$\Lambda'_f = 0 \pm 378.4 \text{ MeV}, \quad (63)$$

$$\Lambda'_{ff} = 0 \pm 263.6 \text{ MeV}, \quad (64)$$

$$\Lambda'_\sigma = 0 + 448.6 \text{ MeV}, \quad (65)$$

$$\Lambda'_{\sigma f} = 0 + 107.9 \text{ MeV}. \quad (66)$$

The conditions Λ'_σ and $\Lambda'_{\sigma f}$ are required to be non-negative, based on the assumption that the $J = \frac{3}{2}$ state does not have a smaller mass than the $J = \frac{1}{2}$ state. By applying the improved mass relation corresponding to Eq. (7) to T_{cc}^+ , we derive

$$\begin{aligned} M(T_{ccs}) - M(T_{cc}^+) - \frac{m_s - m_n}{2m_c} \Lambda'_{ff} \\ = M(\Xi_c) - M(\Lambda_c^+) - \frac{m_s - m_n}{m_c} \Lambda_{ff}, \end{aligned} \quad (67)$$

and we obtain the mass of T_{ccs} as

$$M(T_{ccs}) = 4047 \pm 11 \text{ MeV}. \quad (68)$$

We note that the above value corresponds to the isospin-averaged masses of T_{ccs}^+ and T_{ccs}^{++} . If this result agrees with future experimental data, it means that the color antitriplet state of the heavy diquark is dominant in T_{cc}^+ and T_{ccs} . We compare our result with other results in Table III.

From the mass relation in Eq. (53), we further obtain

$$\begin{aligned} M(\Omega_{cc}) - M(\Xi_{cc}) - \frac{m_s - m_n}{2m_c} (\Lambda'_f - \Lambda'_{\sigma f}) \\ = M_{\text{ave}}(D_s) - M_{\text{ave}}(D) - \frac{m_s - m_n}{m_c} \Lambda_f, \end{aligned} \quad (69)$$

where we used spin-averaged masses for D_s and D to reduce the ambiguity from the correction of the $\Lambda_{\sigma f}$ term. From this relation, the mass of Ω_{cc} , which has not been experimentally reported so far, is also predicted as

$$M(\Omega_{cc}) = 3706_{-15}^{+14} \text{ MeV}. \quad (70)$$

Hadrons Mass [MeV]		
	3706_{-15}^{+14}	Our result
	3766 ± 2	NQM [56]
	3715	RQM [57]
Ω_{cc}	3778	RQM [58]
	3590	NQM [59]
	3815	NQM [60]
	3733 ± 13	Lattice QCD [61]
	3704 ± 17	Lattice QCD [62]

TABLE IV. Several theoretical predictions of the mass of Ω_{cc} . NQM and RQM imply nonrelativistic and relativistic quark models, respectively.

Hadrons	Mass [MeV]	
	172 ± 3	Our result
$M(T_{bbs}) - M(T_{bb})$	$178 \pm 12 \pm 4$	Lattice QCD [65]
	143 ± 47	Lattice QCD [66]
	$104 \pm 23 \pm 10$	Lattice QCD [63, 64]
	84_{-5}^{+4}	Our result
$M_{\text{ave}}(\Omega_{bb}) - M_{\text{ave}}(\Xi_{bb})$	$96 \pm 13_{-21}^{+17}$	Lattice QCD [67]
	$130 \pm 25 \pm 23$	Lattice QCD [68]
	99 ± 23	Lattice QCD [69]

TABLE V. Several theoretical predictions of the mass differences of bottom hadrons. For tetraquarks, we estimated the masses in the lattice QCD by subtracting binding energies from physical thresholds, i.e. $M(T_{bb}) = M(B^*) + M(B) - \Delta E(T_{bb})$, $M(T_{bbs}) = M(B^*) + M(B_s) - \Delta E(T_{bbs})$, where $\Delta E(T_{bb})$ and $\Delta E(T_{bbs})$ are the binding energies of T_{bb} and T_{bbs} shown in Refs.[63–66], respectively.

This result serves as another indicator for assessing whether our mass relations are correct. If this result does not match future experimental results, one possible reason might be that Λ'_f is larger than Λ_f . Additionally, the second-order effects of $1/m_Q$ might also contribute. We show comparison with other results in Table IV. Since hadrons containing two b quarks have not yet been experimentally discovered, we cannot predict in the bottom sector. However, we can predict mass differences as

$$M(T_{bbs}) - M(T_{bb}) = 172 \pm 3 \text{ MeV}, \quad (71)$$

and

$$M(\Omega_{bb}) - M(\Xi_{bb}) = 84_{-5}^{+4} \text{ MeV}. \quad (72)$$

We compare the above results with the results obtained in some lattice analyses in Table V. This shows that our results are consistent with lattice QCD results except for those in Refs. [63, 64].

IV. WIDTH OF T_{ccs}

In this section, we construct the Lagrangian solely from the SU(3) light-flavor symmetry and predict the decay width of T_{ccs} using the decay width of T_{cc}^+ as an input.

Our approach hinges on the fact that both T_{ccs} and T_{cc}^+ belong to the same light-flavor representation. Notably, this method is independent of both superflavor symmetry and color representation.

Since T_{cc}^+ locates just below the DD^* threshold, we consider the decay process of $T_{cc}^+ \rightarrow D^0 D^0 \pi^+$ and $T_{cc}^+ \rightarrow D^0 D^+ \pi^0$ with D^* as an intermediate state, as depicted in Fig. 1. While the decay $T_{cc}^+ \rightarrow D^0 D^+ \pi^0$ has not been observed in experiments [18, 19], it is not prohibited kinematically. Hence, in this study, we also incorporate Figs. 1(b) and 1(c). For T_{ccs} , we consider the decay processes $T_{ccs} \rightarrow DD_s^*$ and $T_{ccs} \rightarrow D^* D_s$ as shown in Fig. 2, because the mass of T_{ccs} predicted in the previous section is above the DD_s^* and $D^* D_s$ thresholds.

Let us construct an effective Lagrangian for the interaction among the tetraquarks and charmed mesons based on just light-flavor symmetry. In the following we use relativistic field T^μ for the mass eigenstates of flavor triplet tetraquarks:

$$T^\mu = \begin{pmatrix} 0 & T_{cc}^+ & T_{ccs}^+ \\ -T_{cc}^+ & 0 & T_{ccs}^{++} \\ -T_{ccs}^+ & -T_{ccs}^{++} & 0 \end{pmatrix}, \quad (73)$$

While the fields for charmed mesons with $J^P = (0^-, 1^-)$ are represented by relativistic fields $(D, D^{*\mu})$:

$$D = \begin{pmatrix} D^0 \\ D^+ \\ D_s \end{pmatrix}, \quad D^* = \begin{pmatrix} D^{*0} \\ D^{*+} \\ D_s^* \end{pmatrix}. \quad (74)$$

From Eqs. (26) and (34), the SU(3) light-flavor transformations of these fields are given as

$$\begin{aligned} T_{ij}^\mu &\rightarrow (U^*)_i^k T_{kl}^\mu (U^\dagger)^l_j, \\ D_i &\rightarrow D_j (U^\dagger)^j_i, \quad D_i^{*\mu} \rightarrow D_j^{*\mu} (U^\dagger)^j_i. \end{aligned} \quad (75)$$

The Lagrangian for $T_{cc(s)}$ and heavy mesons invariant under this transformation is given by

$$\mathcal{L} = g_{TDD} \bar{D}_\mu^{*i} T_{ij}^\mu \bar{D}^j. \quad (76)$$

As we said above, we determine the value of g_{TDD} from the decay width of T_{cc} , 48 keV, assuming that the decay is dominated by the process shown in Fig. 1. By using the value of the $D^* D \pi$ coupling constant determined from the decay of the D^* meson (see, e.g., Ref [43]), the value of g_{TDD} is calculated as

$$g_{TDD} = (4.2 \pm 1.2) \times 10^3 \text{ MeV}. \quad (77)$$

We estimate the error to be 30% based on $\frac{ms}{\Lambda_{\text{QCD}}}$. We note that this value is natural when nondimensionalized by the mass of the charm quark. Now, based on the processes shown in Fig. 2, the formula of the decay width of T_{ccs} is calculated as

$$\Gamma(T_{ccs}) = g_{TDD}^2 \frac{|P_1| + |P_2|}{6\pi M(T_{ccs})^2}, \quad (78)$$

where P_1 and P_2 are phase space momenta corresponding to Figs. 2(a) and 2(b), respectively. By using the mass of T_{ccs} predicted in Eq. (68) together with the value of g_{TDD} in Eq. (77), the decay width is predicted as

$$\Gamma(T_{ccs}) = 42 \pm 24 \text{ MeV}. \quad (79)$$

We consider this decay width to be sufficiently small as to be experimentally observable.

V. SUMMARY

In this study, we investigated the mass and decay width of the doubly heavy tetraquark T_{ccs} from the superflavor and SU(3) light-flavor symmetries. We assumed that doubly charmed tetraquarks are constructed from a color antitriplet cc diquark and thus they are the superflavor partners of the singly heavy baryons. First, we derived the simple mass relations under heavy quark and superflavor symmetries. However, we found a discrepancy between predictions of the obtained mass relations and the experimental data. Then, we constructed an effective Lagrangian based on these symmetries by including correction terms violating simultaneously the heavy-quark symmetry and the light-flavor symmetry. From the Lagrangian, we obtained the improved mass relations among heavy mesons, doubly heavy baryons, singly heavy baryons, and doubly heavy tetraquarks. Based on the relations, we predicted the mass of unobserved tetraquark T_{ccs} as $M(T_{ccs}) = 4047 \pm 11 \text{ MeV}$. We also predicted the mass of unobserved Ω_{cc} as $M(\Omega_{cc}) = 3706_{-15}^{+14} \text{ MeV}$.

We then constructed an effective Lagrangian term for the decay of T_{ccs} based on SU(3) light-flavor symmetry. The unknown coupling constant was determined by using the T_{cc} decay data. Incorporating the predicted mass, we derived the decay width of T_{ccs} as $\Gamma(T_{ccs}) = 42 \pm 24 \text{ MeV}$.

The obtained masses and widths will be useful to understand the color configuration of DHTs. If these results agree with future experimental data, it means that the color antitriplet state in T_{cc}^+ and T_{ccs} is dominant.

The isovector counterpart of the isoscalar T_{cc}^+ can also exist independently from T_{cc}^+ , and we expect that the analyses can be done separately. It is interesting to study the isovector state by extending the present analysis, which we leave to the future work.

VI. ACKNOWLEDGMENT

This work was supported in part by JSPS KAKENHI Grants Nos. JP20K03927, No. JP23H05439, No. JP24K007045 (M.H.), and No. JP20K14478 (Y.Y.).




FIG. 1. Feynmann diagrams of the $T_{cc}^+ \rightarrow DD\pi$ decays.




FIG. 2. Feynmann diagrams of the $T_{ccs} \rightarrow DD_s^*$ and D^*D_s decays.

-
- [1] S. K. Choi et al. (Belle), Observation of a narrow charmonium-like state in exclusive $B^\pm \rightarrow K^\pm \pi^+ \pi^- J/\psi$ decays, Phys. Rev. Lett. **91**, 262001 (2003), arXiv:hep-ex/0309032.
- [2] M. Gell-Mann, A Schematic Model of Baryons and Mesons, Phys. Lett. **8**, 214 (1964).
- [3] G. Zweig, An SU(3) model for strong interaction symmetry and its breaking. Version 1, DEVELOPMENTS IN THE QUARK THEORY OF HADRONS. VOL. 1 (1964).
- [4] G. Zweig, An SU(3) model for strong interaction symmetry and its breaking. Version 2 (Hadronic Press, 1964) pp. 22–101.
- [5] E. S. Swanson, The New heavy mesons: A Status report, Phys. Rept. **429**, 243 (2006), arXiv:hep-ph/0601110.
- [6] Y. Yamaguchi, A. Hosaka, S. Takeuchi, and M. Takizawa, Heavy hadronic molecules with pion exchange and quark core couplings: a guide for practitioners, J. Phys. G **47**, 053001 (2020), arXiv:1908.08790 [hep-ph].
- [7] H.-X. Chen, W. Chen, X. Liu, Y.-R. Liu, and S.-L. Zhu, An updated review of the new hadron states, Rept. Prog. Phys. **86**, 026201 (2023), arXiv:2204.02649 [hep-ph].
- [8] L. Meng, B. Wang, G.-J. Wang, and S.-L. Zhu, Chiral perturbation theory for heavy hadrons and chiral effective field theory for heavy hadronic molecules, Phys. Rept. **1019**, 1 (2023), arXiv:2204.08716 [hep-ph].
- [9] J. Vijande, F. Fernandez, A. Valcarce, and B. Silvestre-Brac, Tetraquarks in a chiral constituent quark model, Eur. Phys. J. A **19**, 383 (2004), arXiv:hep-ph/0310007.
- [10] G. Yang, J. Ping, and F. Wang, Structure of pentaquarks P_c^+ in the chiral quark model, Phys. Rev. D **95**, 014010 (2017).
- [11] E. Hiyama, A. Hosaka, M. Oka, and J.-M. Richard, Quark model estimate of hidden-charm pentaquark resonances, Phys. Rev. C **98**, 045208 (2018), arXiv:1803.11369 [nucl-th].
- [12] Q. Meng, E. Hiyama, K. U. Can, P. Gubler, M. Oka, A. Hosaka, and H. Zong, Compact $sssc\bar{c}$ pentaquark states predicted by a quark model, Phys. Lett. B **798**, 135028 (2019), arXiv:1907.00144 [nucl-th].
- [13] Q. Zhang, B.-R. He, and J.-L. Ping, Pentaquarks with the $qqs\bar{Q}Q$ configuration in the Chiral Quark Model (2020), arXiv:2006.01042 [hep-ph].
- [14] G.-J. Wang, Q. Meng, and M. Oka, S-wave fully charmed tetraquark resonant states, Phys. Rev. D **106**, 096005 (2022), arXiv:2208.07292 [hep-ph].
- [15] G. Yang, J. Ping, and J. Segovia, Hidden-Charmed Pentaquarks with Strangeness in a Chiral Quark Model, Symmetry **16**, 354 (2024), arXiv:2311.01044 [hep-ph].
- [16] H. Huang, C. Deng, X. Liu, Y. Tan, and J. Ping, Tetraquarks and Pentaquarks from Quark Model Perspective, Symmetry **15**, 1298 (2023).
- [17] N. A. Tornqvist, From the deuteron to deusons, an analysis of deuteron-like meson meson bound states, Z. Phys. C **61**, 525 (1994), arXiv:hep-ph/9310247.
- [18] R. Aaij et al. (LHCb), Observation of an exotic narrow doubly charmed tetraquark, Nature Phys. **18**, 751 (2022), arXiv:2109.01038 [hep-ex].

- [19] R. Aaij et al. (LHCb), Study of the doubly charmed tetraquark T_{cc}^+ , Nature Commun. **13**, 3351 (2022), arXiv:2109.01056 [hep-ex].
- [20] A. V. Manohar and M. B. Wise, Exotic Q Q anti-q anti-q states in QCD, Nucl. Phys. B **399**, 17 (1993), arXiv:hep-ph/9212236.
- [21] D. Janc and M. Rosina, The $T_{cc} = DD^*$ molecular state, Few Body Syst. **35**, 175 (2004), arXiv:hep-ph/0405208.
- [22] S. Ohkoda, Y. Yamaguchi, S. Yasui, K. Sudoh, and A. Hosaka, Exotic mesons with double charm and bottom flavor, Phys. Rev. D **86**, 034019 (2012), arXiv:1202.0760 [hep-ph].
- [23] N. Li, Z.-F. Sun, X. Liu, and S.-L. Zhu, Coupled-channel analysis of the possible $D^{(*)}D^{(*)}$, $\bar{B}^{(*)}\bar{B}^{(*)}$ and $D^{(*)}\bar{B}^{(*)}$ molecular states, Phys. Rev. D **88**, 114008 (2013), arXiv:1211.5007 [hep-ph].
- [24] J.-B. Cheng, Z.-Y. Lin, and S.-L. Zhu, Double-charm tetraquark under the complex scaling method, Phys. Rev. D **106**, 016012 (2022), arXiv:2205.13354 [hep-ph].
- [25] F.-L. Wang and X. Liu, Investigating new type of doubly charmed molecular tetraquarks composed of charmed mesons in the H and T doublets, Phys. Rev. D **104**, 094030 (2021), arXiv:2108.09925 [hep-ph].
- [26] F.-L. Wang, R. Chen, and X. Liu, A new group of doubly charmed molecule with T-doublet charmed meson pair, Phys. Lett. B **835**, 137502 (2022), arXiv:2111.00208 [hep-ph].
- [27] H. Ren, F. Wu, and R. Zhu, Hadronic Molecule Interpretation of Tcc+ and Its Beauty Partners, Adv. High Energy Phys. **2022**, 9103031 (2022), arXiv:2109.02531 [hep-ph].
- [28] Q. Xin and Z.-G. Wang, Analysis of the doubly-charmed tetraquark molecular states with the QCD sum rules, Eur. Phys. J. A **58**, 110 (2022), arXiv:2108.12597 [hep-ph].
- [29] B. Wang and L. Meng, Revisiting the DD^* chiral interactions with the local momentum-space regularization up to the third order and the nature of Tcc+, Phys. Rev. D **107**, 094002 (2023), arXiv:2212.08447 [hep-ph].
- [30] Q. Xin, Z.-G. Wang, and X.-S. Yang, Analysis of the X(3960) and related tetraquark molecular states via the QCD sum rules, AAPPS Bull. **32**, 37 (2022), arXiv:2207.09910 [hep-ph].
- [31] T. Asanuma, Y. Yamaguchi, and M. Harada, Analysis of DD^* and $\bar{D}^{(*)}\Xi_{cc}^{(*)}$ molecule by one boson exchange model based on Heavy quark symmetry (2023), arXiv:2311.04695 [hep-ph].
- [32] M. Sakai and Y. Yamaguchi, Analysis of Tcc and Tbb based on the hadronic molecular model and their spin multiplets, Phys. Rev. D **109**, 054016 (2024), arXiv:2312.08663 [hep-ph].
- [33] Y. Kim, M. Oka, and K. Suzuki, Doubly heavy tetraquarks in a chiral-diquark picture, Phys. Rev. D **105**, 074021 (2022), arXiv:2202.06520 [hep-ph].
- [34] R. L. Jaffe, Multi-Quark Hadrons. 1. The Phenomenology of (2 Quark 2 anti-Quark) Mesons, Phys. Rev. D **15**, 267 (1977).
- [35] O. Andreev, $QQq\bar{q}$ potential in string models, Phys. Rev. D **105**, 086025 (2022), arXiv:2111.14418 [hep-ph].
- [36] F. S. Navarra, M. Nielsen, and S. H. Lee, QCD sum rules study of QQ - anti-u anti-d mesons, Phys. Lett. B **649**, 166 (2007), arXiv:hep-ph/0703071.
- [37] M.-L. Du, W. Chen, X.-L. Chen, and S.-L. Zhu, Exotic $QQ\bar{q}\bar{q}$, $QQ\bar{q}\bar{s}$ and $QQ\bar{s}\bar{s}$ states, Phys. Rev. D **87**, 014003 (2013), arXiv:1209.5134 [hep-ph].
- [38] S. S. Agaev, K. Azizi, and H. Sundu, Newly observed exotic doubly charmed meson Tcc+, Nucl. Phys. B **975**, 115650 (2022), arXiv:2108.00188 [hep-ph].
- [39] Y. Ikeda, B. Charron, S. Aoki, T. Doi, T. Hatsuda, T. Inoue, N. Ishii, K. Murano, H. Nemura, and K. Sasaki, Charmed tetraquarks T_{cc} and T_{cs} from dynamical lattice QCD simulations, Phys. Lett. B **729**, 85 (2014), arXiv:1311.6214 [hep-lat].
- [40] P. Junnarkar, N. Mathur, and M. Padmanath, Study of doubly heavy tetraquarks in Lattice QCD, Phys. Rev. D **99**, 034507 (2019), arXiv:1810.12285 [hep-lat].
- [41] M. Padmanath and S. Prelovsek, Signature of a Doubly Charm Tetraquark Pole in DD^* Scattering on the Lattice, Phys. Rev. Lett. **129**, 032002 (2022), arXiv:2202.10110 [hep-lat].
- [42] Y. Lyu, S. Aoki, T. Doi, T. Hatsuda, Y. Ikeda, and J. Meng, Doubly Charmed Tetraquark Tcc+ from Lattice QCD near Physical Point, Phys. Rev. Lett. **131**, 161901 (2023), arXiv:2302.04505 [hep-lat].
- [43] A. V. Manohar and M. B. Wise, Heavy quark physics, Vol. 10 (Cambridge University Press, 2000).
- [44] H. Georgi and M. B. Wise, Superflavor Symmetry for Heavy Particles, Phys. Lett. B **243**, 279 (1990).
- [45] M. J. Savage and M. B. Wise, Spectrum of baryons with two heavy quarks, Phys. Lett. B **248**, 177 (1990).
- [46] S. Fleming and T. Mehen, Doubly heavy baryons, heavy quark-diquark symmetry and NRQCD, Phys. Rev. D **73**, 034502 (2006), arXiv:hep-ph/0509313.
- [47] J. Carlson, L. Heller, and J. A. Tjon, Stability of Dimesons, Phys. Rev. D **37**, 744 (1988).
- [48] Q. Meng, E. Hiyama, A. Hosaka, M. Oka, P. Gubler, K. U. Can, T. T. Takahashi, and H. S. Zong, Stable double-heavy tetraquarks: spectrum and structure, Phys. Lett. B **814**, 136095 (2021), arXiv:2009.14493 [nucl-th].
- [49] Q. Meng, M. Harada, E. Hiyama, A. Hosaka, and M. Oka, Doubly heavy tetraquark resonant states, Phys. Lett. B **824**, 136800 (2022), arXiv:2106.11868 [hep-ph].
- [50] B.-R. He, M. Harada, and B.-S. Zou, Quark model with hidden local symmetry and its application to Tcc, Phys. Rev. D **108**, 054025 (2023), arXiv:2306.03526 [hep-ph].
- [51] E. J. Eichten and C. Quigg, Heavy-quark symmetry implies stable heavy tetraquark mesons $Q_i Q_j \bar{q}_k \bar{q}_l$, Phys. Rev. Lett. **119**, 202002 (2017).
- [52] Y.-L. Ma and M. Harada, Chiral partner structure of doubly heavy baryons with heavy quark spin-flavor symmetry, J. Phys. G **45**, 075006 (2018), arXiv:1709.09746 [hep-ph].
- [53] Particle Data Group, Review of Particle Physics, Prog. Theor. Exp. Phys. **2023**, 083C01 (2023).
- [54] N. Brambilla, A. Vairo, and T. Rosch, Effective field theory Lagrangians for baryons with two and three heavy quarks, Phys. Rev. D **72**, 034021 (2005), arXiv:hep-ph/0506065.
- [55] M. Karliner and J. L. Rosner, Doubly charmed strange tetraquark, Phys. Rev. D **105**, 034020 (2022), arXiv:2110.12054 [hep-ph].
- [56] E. Ortiz-Pacheco and R. Bijker, Masses and radiative decay widths of S- and P-wave singly, doubly, and triply heavy charm and bottom baryons, Phys. Rev. D **108**, 054014 (2023), arXiv:2307.04939 [hep-ph].

- [57] Q.-F. Lü, K.-L. Wang, L.-Y. Xiao, and X.-H. Zhong, Mass spectra and radiative transitions of doubly heavy baryons in a relativized quark model, *Phys. Rev. D* **96**, 114006 (2017), arXiv:1708.04468 [hep-ph].
- [58] D. Ebert, R. N. Faustov, V. O. Galkin, and A. P. Martynenko, Mass spectra of doubly heavy baryons in the relativistic quark model, *Phys. Rev. D* **66**, 014008 (2002), arXiv:hep-ph/0201217.
- [59] S. S. Gershtein, V. V. Kiselev, A. K. Likhoded, and A. I. Onishchenko, Spectroscopy of doubly heavy baryons, *Phys. Rev. D* **62**, 054021 (2000).
- [60] W. Roberts and M. Pervin, Heavy baryons in a quark model, *Int. J. Mod. Phys. A* **23**, 2817 (2008), arXiv:0711.2492 [nucl-th].
- [61] H. Bahtiyar, K. U. Can, G. Erkol, P. Gubler, M. Oka, and T. T. Takahashi (TRJQCD Collaboration), Charmed baryon spectrum from lattice qcd near the physical point, *Phys. Rev. D* **102**, 054513 (2020).
- [62] Y. Namekawa et al. (PACS-CS), Charmed baryons at the physical point in 2+1 flavor lattice QCD, *Phys. Rev. D* **87**, 094512 (2013), arXiv:1301.4743 [hep-lat].
- [63] S. Meinel, M. Pflaumer, and M. Wagner, Search for b^-b^-us and b^-c^-ud tetraquark bound states using lattice QCD, *Phys. Rev. D* **106**, 034507 (2022), arXiv:2205.13982 [hep-lat].
- [64] M. Wagner, C. Alexandrou, J. Finkenrath, T. Leontiou, S. Meinel, and M. Pflaumer, Lattice QCD study of antiheavy-antiheavy-light-light tetraquarks based on correlation functions with scattering interpolating operators both at the source and at the sink, *PoS LATTICE2022*, 270 (2023), arXiv:2210.09281 [hep-lat].
- [65] A. Francis, R. J. Hudspith, R. Lewis, and K. Maltman, Lattice prediction for deeply bound doubly heavy tetraquarks, *Phys. Rev. Lett.* **118**, 142001 (2017).
- [66] P. Junnarkar, N. Mathur, and M. Padmanath, Study of doubly heavy tetraquarks in lattice qcd, *Phys. Rev. D* **99**, 034507 (2019).
- [67] R. Lewis and R. M. Woloshyn, Bottom baryons from a dynamical lattice qcd simulation, *Phys. Rev. D* **79**, 014502 (2009).
- [68] Z. S. Brown, W. Detmold, S. Meinel, and K. Orginos, Charmed bottom baryon spectroscopy from lattice QCD, *Phys. Rev. D* **90**, 094507 (2014), arXiv:1409.0497 [hep-lat].
- [69] P. Mohanta and S. Basak, Heavy baryon spectrum on the lattice with NRQCD bottom and HISQ lighter quarks, *Phys. Rev. D* **101**, 094503 (2020), arXiv:1911.03741 [hep-lat].