RANDOM COVARIANT QUANTUM CHANNELS

ION NECHITA AND SANG-JUN PARK

ABSTRACT. The group symmetries inherent in quantum channels often make them tractable and applicable to various problems in quantum information theory. In this paper, we introduce natural probability distributions for covariant quantum channels. Specifically, this is achieved through the application of "twirling operations" on random quantum channels derived from the Stinespring representation that use Haar-distributed random isometries. We explore various types of group symmetries, including unitary and orthogonal covariance, hyperoctahedral covariance, diagonal orthogonal covariance (DOC), and analyze their properties related to quantum entanglement based on the model parameters. In particular, we discuss the threshold phenomenon for positive partial transpose and entanglement breaking properties, comparing thresholds among different classes of random covariant channels. Finally, we contribute to the PPT² conjecture by showing that the composition between two random DOC channels is generically entanglement breaking.

CONTENTS

1. Introduction	1
2. Random covariant quantum channels	5
2.1. Covariant quantum channels	5
2.2. Random quantum channels	7
3. Random orthogonal covariant channels	8
4. Random hyperoctahedral covariant channels	13
5. Random diagonal orthogonal covariant channels	17
6. The PPT ² conjecture holds for random DOC channels	24
Appendix A. Technical lemmas and proofs	26
A.1. General results about the combinatorics of permutations and Weingarten functions	26
A.2. Scalar parameters	32
A.3. Matrix parameters	35
References	43

1. INTRODUCTION

Quantum channels are essential mathematical constructs in quantum theory, modeling the most general evolution open quantum systems can undergo. Generalizing the unitary quantum dynamics described by Schrödinger's equation, quantum channels have been studied extensively, from different perspectives [NC10, Wil17, Wat18]. In physics, quantum channels describe the evolution of quantum states as they propagate through space and/or time, and model usually the noise coming from the environment. In quantum information theory, quantum channels model the transmission of quantum information between distant parties, akin to a classical communication channel but with quantum properties. Mathematically, quantum channels are represented by completely positive, trace preserving maps.

The study of *covariant quantum evolutions* was initiated by Holevo [Hol93, Hol96]. This type of covariance encodes usually a symmetry present in the system Hamiltonian and can be mathematically formulated as follows:

$$\forall g \in G, \qquad \Phi\Big(\pi_A(g) \cdot \pi_A(g)^*\Big) = \pi_B(g)\Phi(\cdot)\pi_B(g)^*.$$

Above, π_A and π_B are unitary representations of the physical symmetry group G. A channel $\Phi: \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$ having the property above is called a (π_A, π_B) -covariant channel. This notion is not only important in physically-relevant settings where symmetry is present, but also as a tool in several areas of quantum information theory, such as quantum cloning [KW99], programmability [GBW21], quantum Shannon theory [WH02, Kin03, Hol05, DHS06, KW09, DTW16, WTB17, BCLY20, etc. The importance of this notion stems also from the fact that group symmetries inherent in quantum channels often make them tractable and applicable to various problems in quantum information theory, especially related to quantum entanglement VW01, COS18, CMHW18, ?, BCS20, SN21, SN22b, PJPY23, PY23]. Their general structure and mathematical properties have been extensively studied in recent years [AN14, MSD17, MS22, LY22].

Another way to probe the structure of quantum channels is to consider their generic behavior, that is to study random quantum channels. Broadly speaking, random quantum channels model typical behavior of quantum operations [BCSZ09], hence it is of crucial importance to understand their physical and mathematical properties, such as: entanglement, information transmission capacity, spectral properties, etc. A systematic study of the different models of random quantum channels and of their properties can be found in [KNP⁺21a]. Moreover, random constructions for quantum channels have played a crucial role in many areas of quantum Shannon theory, mainly as a source of (counter-)examples [HW08, Has09, BCN16, CN15, CP22]; this situation is parallel to classical information theory, where random constructions of codes and channels are ubiquitous.

In this work, we consider random covariant quantum channels. We introduce and study probability distributions on the set of covariant quantum channels, for different choices of group symmetry. We find in which asymptotic regimes such channels are *PPT* or *entanglement breaking*. In particular, we explain the *threshold phenomenon* for these two properties. We show that in the case of diagonal symmetry, random covariant channels satisfy the PPT^2 conjecture. Our work sets up the framework for the study of random covariant channels, by defining the relevant probabilistic models in the most common settings of group symmetry. We expect the models and the results in this paper to find further applications in quantum information theory, in the study of problems having a specific type of symmetry and also as a rich source of examples of channels with a tractable form.

We shall consider several cases of (random) covariant channels, corresponding to the standard representations of:

- the unitary group $\mathcal{U}_d = \{ U \in \mathcal{M}_d(\mathbb{C}) : U^*U = I_d \}$
- the orthogonal group $\mathcal{O}_d = \{ U \in \mathcal{M}_d(\mathbb{R}) : O^\top O = I_d \}$
- the hyperoctahedral group $\mathcal{H}_d = \{\pm 1\} \wr \mathcal{S}_d$ of signed permutation matrices the diagonal unitary group $\mathcal{DU}_d = \{ \operatorname{diag}(e^{\mathrm{i}\theta_1}, \dots, e^{\mathrm{i}\theta_d}) : \theta_1, \dots, \theta_d \in \mathbb{R} \}$
- the diagonal orthogonal group $\mathcal{DO}_d = \{ \operatorname{diag}(\varepsilon_1 \dots, \varepsilon_d) : \varepsilon_1, \dots, \varepsilon_d = \pm 1 \}.$

We depict in Fig. 1 the groups (left) and the corresponding classes of covariant channels (right). Edges correspond to set inclusion relations, meaning that the source is contained in the target. Top elements correspond to the largest groups (reps. channel sets), while bottom elements correspond to the smallest groups (resp. channel sets). The notation is self-evident, with the exception of (see Section 5):

DUC channels	\leftrightarrow	$(\mathcal{DU}_d \times \mathcal{DU}_d)$ covariance
CDUC channels	\leftrightarrow	$(\mathcal{DU}_d \times \overline{\mathcal{DU}}_d)$ covariance
DOC channels	\leftrightarrow	$(\mathcal{DO}_d \times \mathcal{DO}_d)$ covariance.



FIGURE 1. The groups we consider (left) and the corresponding class of covariant channels (right). Edges correspond to set inclusion relations, meaning that the source is contained in the target. Top elements correspond to the largest groups (reps. channel sets), while bottom elements correspond to the smallest groups (resp. channel sets).

The contributions of our work are two-fold:

- Entanglement theory: we find the range of parameters s for which random covariant channels are asymptotically PPT, respectively EB. In particular, we specify the threshold s_0 such that, in the limit $d \to \infty$, the probability that random covariant channels are PPT/EB tends to 1 (resp. < 1) whenever whenever $s \gg s_0$ (resp. $s \ll s_0$).
- PPT^2 conjecture: we generalize the main result of [SN22b]. Furthermore, we show that random DOC channels satisfy the PPT² conjecture, under very general conditions.

We recall that a quantum channel is PPT (positive partial transpose) if the composition of the channel with the (non-physical) transpose map is still a quantum channel. On the other hand, a quantum channel is called *entanglement breaking* (EB) if, when applied to one system of a bipartite quantum state, the output bipartite state is separable [HSR03]. Our results on the PPT and EB properties of random covariant tables are informally presented in the table below. The rows of the table are ordered, starting with the largest groups on the top, and finishing with the smallest group at the bottom. The last row of the table corresponds to no symmetry whatsoever, hence to random channels before the twirling operation. Here s_0 denotes the threshold for corresponding properties. For example, $s_0 \sim \text{const.}$ for PPT means that we have the property with probability 1 (resp. < 1) whenever $d \to \infty$ and $s = s(d) \to \infty$ (resp. s is fixed). Note that the two last columns are devoted to the entanglement breaking (EB) property of channels: since deciding whether a channels is EB is an NP-hard problem, it is difficult in general to obtain sharp thresholds for this property. All the results present in the table are new, with the exception of the last row.

Group symmetry	PPT property	EB property	Not EB property
(U, U)-cov.	Always	Always	Never
(U,\overline{U}) -cov.	$s_0 \sim \text{const.}$	$s_0 \sim \text{const.}$	$s_0 \sim \text{const.}$
(O, O)-cov.	$s_0 \sim \text{const.}$	$s_0 \sim \text{const.}$	$s_0 \sim \text{const.}$
(H,H)-cov.	$s_0 \sim \text{const.}$	$s_0 \sim \text{const.}$	$s_0 \sim \text{const.}$
DUC	$s_0 \sim \text{const.}$	$s\gtrsim d^{2+\varepsilon}$	s = O(1)
CDUC	$s_0 \sim 4d$	$s\gtrsim d^{2+\varepsilon}$	$s \le (4 - \varepsilon)d$
DOC	$s_0 \sim 4d$	$s\gtrsim d^{2+\varepsilon}$	$s \le (4 - \varepsilon)d$
No symmetry	$s_0 \sim 4d^2$ [Aub12]	$s \gtrsim d^3 \log^2 d $ [ASY14]	$s \lesssim d^3 $ [ASY14]

The formal statements for the unitary and orthogonal cases are given in Theorem 3.5; the ones for the hyperoctahedral case are given in Theorem 4.4. We present below the statement of a formal result, corresponding to the CDUC, DUC, and DOC rows of the table above; see Theorems 5.10 and 5.13 for the details.

Theorem A. Let $\Phi_{\text{DUC}} \sim \mu_{d,s}^{\text{DUC}}$, $\Phi_{\text{CDUC}} \sim \mu_{d,s}^{\text{CDUC}}$, and $\Phi_{\text{DOC}} \sim \mu_{d,s}^{\text{DOC}}$ be, respectively, random DUC, CDUC, and DOC channels. Then:

- (1) In the regime s is fixed and $d \to \infty$, we have $\lim_{d\to\infty} \mathbb{P}(\Phi_{\mathsf{DUC}} \text{ is } \mathsf{PPT}) = 0$.
- (2) In the regime $s \gtrsim d^{\varepsilon}$ for some $\varepsilon > 0$, Φ_{DUC} is almost surely PPT as $d \to \infty$.
- (3) In the regime $s/d \to 0$, Φ_{CDUC} and Φ_{DOC} are almost surely not PPT as $d \to \infty$.
- (4) Consider the regime $s \sim cd$.
 - If c < 4, then Φ_{CDUC} and Φ_{DOC} are almost surely not PPT as $d \to \infty$.
 - If c > 4, then both Φ_{CDUC} and Φ_{DOC} have Choi matrices whose partial transpositions converge, almost surely as $d \to \infty$, in moments, to a probability measure supported on the positive real line.
- (5) In the regime $s \gtrsim d^{1+\varepsilon}$ for some $\varepsilon > 0$, Φ_{CDUC} and Φ_{DOC} are almost surely PPT as $d \to \infty$. (6) In the regime $s \gtrsim d^{2+\varepsilon}$ for some $\varepsilon > 0$, Φ_{DUC} , Φ_{CDUC} , and Φ_{DOC} are almost surely EB as $d \to \infty$.

Note that in item (5) above we show that the smallest eigenvalue of the partial transpose of the Choi matrix of the random channels we consider converges to a positive value. We also obtain results about the *realignment criterion* [CW03, Rud05] applied to the Choi matrix of DUC, CDUC, and DOC channels in Proposition 5.12.

Our second contribution concerns the PPT^2 conjecture for DOC channels. Motivated by the theory of quantum repeaters, this conjecture [Chr12, CMHW18] has received a lot of attention in the recent years. The conjecture states that the composition of two arbitrary PPT quantum channels is entanglement breaking. Recent works establish the conjecture in several restricted scenarios, or with weaker conclusions [LG15, KMP17, RJP18, CMHW18, CYT19, HRF20, GKS20, JY20, SN22b]. Moreover, in the case of random quantum channels (without any covariance property), it has been proven [CYZ18] that two *independent* random quantum channels that are PPT satisfy the conjecture.

Regarding the quantum channels with diagonal orthogonal symmetry, it has been shown in [SN22b] that the PPT² conjecture holds for any two DUC / CDUC channels, while the same question for DOC channels has been left open. We tackle this question from both *deterministic* and random perspectives. The results can be summarized as follows; see Corollary 6.4 and Theorem 6.5for the details.

(1) The PPT^2 conjecture holds for the composition between any DOC and Theorem B. (C)DUC channels. That is, if $\Phi_1, \Phi_2 : \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$ are two PPT channels such that Φ_1 is DOC and Φ_2 is DUC or CDUC, then both $\Phi_1 \circ \Phi_2$ and $\Phi_2 \circ \Phi_1$ are entanglement breaking.

(2) Consider two random DOC channels $\Phi_i \sim \mu_{d,s_i}^{\text{DOC}}$, i = 1, 2, in the asymptotic regime where $d \to \infty$ and $s_i \gtrsim d^{t_i}$ for some constants $t_i > 0$. Then, almost surely as $d \to \infty$, the composition $\Phi_1 \circ \Phi_2$ is entanglement breaking. In particular, in the asymptotic regime above, any two random DOC channels satisfy the PPT² conjecture.

Note that Theorem B (1) generalizes the main result of [SN22b] by allowing one of the channels to belong to the larger class of DOC channels. Moreover, it is worth mentioning that the hypotheses of Theorem B (2) are very weak when compared to those of [CYZ18]: the channels need not be PPT, nor independent.

Our paper is organized as follows. In Section 2 we recall the theory of covariant quantum channels and that of random quantum channels; we also give the general recipe for sampling random covariant quantum channels. Sections 3 to 5 deal, respectively, with the three main classes of symmetry groups: unitary/orthogonal, hyperoctahedral, and diagonal unitary/orthogonal. These sections present the general structure theorems for the corresponding covariant channels, and then gather our main results regarding the PPT and EB properties of the random covariant quantum channels. Finally, Section 6 contains our result about DOC channels satisfying the PPT² conjecture.

2. RANDOM COVARIANT QUANTUM CHANNELS

In this section we recall two of the fundamental notions discussed in this paper: that of symmetric (covariant) quantum channels, and that of random quantum channels. In the rest of the paper, we shall combine these two points of view by analyzing the properties of *random covariant quantum channels*.

2.1. Covariant quantum channels. For a compact group G, consider (finite-dimensional) unitary representations $\pi : G \to \mathcal{U}_d$, $\pi_A : G \to \mathcal{U}_{d_A}$, and $\pi_B : G \to \mathcal{U}_{d_B}$ of G. Then we call

- (1) a matrix $X \in \mathcal{M}_d(\mathbb{C})$ π -invariant if $\pi(x)X\pi(x)^* = X$ for all $x \in G$,
- (2) a linear map $\Phi : \mathcal{M}_{d_A}(\mathbb{C}) \to \mathcal{M}_{d_B}(\mathbb{C}) \ (\pi_A, \pi_B)$ -covariant if $\Phi(\pi_A(x)Z\pi_A(x)^*) = \pi_B(x)\Phi(Z)\pi_B(x)^*$ for all $x \in G$ and $Z \in \mathcal{M}_{d_A}(\mathbb{C})$.

Let us denote by $\operatorname{Inv}(\pi)$ (resp. $\operatorname{Cov}(\pi_A, \pi_B)$) the set of all π -invariant matrices (resp. (π_A, π_B) covariant linear maps). π is called *irreducible* if $\operatorname{Inv}(\pi) = \mathbb{C}I_d$ (which coincides with usual definition of irreducibility in representation theory, thanks to Schur's lemma). If π is irreducible, so is the *contragredient representation* $\overline{\pi} : G \to \mathcal{U}_d$ of π which is defined by $\overline{\pi}(x) = \overline{\pi}(x)$ for $x \in G$. For unitary representations $\pi_A : G \to \mathcal{U}_{d_A}$ and $\pi_B : G \to \mathcal{U}_{d_B}$, the *direct sum representation* $\pi_A \oplus \pi_B : G \to \mathcal{U}_{d_A + d_B}$ (resp. *tensor representation* $\pi_A \otimes \pi_B : G \to \mathcal{U}_{d_A d_B}$) is given by $(\pi_A \oplus \pi_B)(x) =$ $\pi_A(x) \oplus \pi_B(x)$ (resp. $(\pi_A \otimes \pi_B)(x) = \pi_A(x) \otimes \pi_B(x)$) for $x \in G$. Two unitary representations $\pi_1 : G \to \mathcal{U}_{d_1}$ and $\pi_2 : G \to \mathcal{U}_{d_2}$ are said to be *unitarily equivalent* if there exists a unitary matrix $U : \mathbb{C}^{d_1} \to \mathbb{C}^{d_2}$ such that $\pi_2(x) = U\pi_1(x)U^*$ for all $x \in G$, where we denote by $\pi_1 \cong \pi_2$.

The following is a well-known fact in representation theory.

Proposition 2.1 ([Sim95]). Suppose that a unitary representation $\pi : G \to U_d$ has an irreducible decomposition of the form

$$\pi \cong \bigoplus_{i=1}^{l} \sigma_i \otimes I_{m_i},\tag{1}$$

where $\sigma_i: G \to \mathcal{U}_{n_i}$ (i = 1, 2, ..., l) are mutually inequivalent irreducible sub-representations of π with multiplicity m_i , respectively. Then we have

$$\operatorname{Inv}(\pi) \cong \bigoplus_{i=1}^{\circ} I_{n_i} \otimes \mathcal{M}_{m_i}(\mathbb{C}) \subseteq \mathcal{M}_{n_i}(\mathbb{C}) \otimes \mathcal{M}_{m_i}(\mathbb{C}).$$
(2)

In this section, we are interested in two types of twirling operations: for unitary representations π, π_A, π_B of G,

- (1) π -twirling $\mathcal{T}_{\pi}X := \mathbb{E}_G[\mathrm{Ad}_{\pi(\cdot)}(X)] = \mathbb{E}_G[\pi(\cdot)X\pi(\cdot)^*]$ for a matrix $X \in \mathcal{M}_d(\mathbb{C})$,
- (2) (π_A, π_B) -twirling $\mathcal{T}_{\pi_A, \pi_B} \Phi := \mathbb{E}_G[\mathrm{Ad}_{\pi_B(\cdot)^*} \circ \Phi \circ \mathrm{Ad}_{\pi_A(\cdot)}]$ for a linear map $\Phi : \mathcal{M}_{d_A}(\mathbb{C}) \to \mathcal{M}_{d_B}(\mathbb{C}),$

where $\operatorname{Ad}_U : Z \mapsto UZU^*$ and the expectation is defined with respect to the (normalized) Haar measure of the compact group G.

Let us collect some useful properties of the twirling operations for the next section. First of all, \mathcal{T}_{π} is a *conditional expectation* onto the *-subalgebra $\operatorname{Inv}(\pi)$ of $\mathcal{M}_d(\mathbb{C})$. Note that for any (finitedimensional) von Neumann subalgebra \mathcal{M} of $\mathcal{M}_d(\mathbb{C})$, there is a unique TP conditional expectation of $\mathcal{M}_d(\mathbb{C})$ onto \mathcal{M} [BO08, Lemma 1.5.11]. For example, the map $X \in \mathcal{M}_n(\mathbb{C}) \otimes \mathcal{M}_m(\mathbb{C}) \mapsto$ $\frac{1}{n}I_n \otimes \operatorname{Tr}_n(X)$ is the unique TP conditional expectation onto $\mathcal{M} = I_n \otimes \mathcal{M}_m(\mathbb{C})$. This observation allow us to get the following explicit formula of the twirling map \mathcal{T}_{π} for the case $\mathcal{M} = \operatorname{Inv}(\pi)$.

Proposition 2.2. In Proposition 2.1, let Π_i be the orthogonal projection from \mathbb{C}^d onto $\mathbb{C}^{n_i} \otimes \mathbb{C}^{m_i}$. Then the twirling $\mathcal{T}_{\pi}(X)$ of $X \in \mathcal{M}_d(\mathbb{C})$ is given by

$$\mathcal{T}_{\pi}(X) = \bigoplus_{i=1}^{l} \frac{1}{n_i} I_{n_i} \otimes \operatorname{Tr}_{n_i}(\Pi_i X \Pi_i).$$
(3)

In particular, if the irreducible decomposition of π is multiplicity-free, i.e., if $m_i \equiv 1$ for all $i = 1, 2, \dots, l$, then

$$\mathcal{T}_{\pi}(X) = \sum_{i=1}^{l} \frac{\operatorname{Tr}(\Pi_i X)}{n_i} \Pi_i.$$
(4)

Proposition 2.3. Let $\pi_0, \pi_1 : G \to U_d$ be two unitary representations of G such that $\operatorname{Ran} \pi_1 \subset \operatorname{Ran} \pi_0$. Then

$$\mathcal{T}_{\pi_0} \circ \mathcal{T}_{\pi_1} = \mathcal{T}_{\pi_1} \circ \mathcal{T}_{\pi_0} = \mathcal{T}_{\pi_0}.$$

Proof. Recall that each \mathcal{T}_{π_i} is a TP conditional expectation onto $\operatorname{Inv}(\pi_i)$. Since $\operatorname{Inv}(\pi_0) \subset \operatorname{Inv}(\pi_1)$, we obviously have $\mathcal{T}_{\pi_1} \circ \mathcal{T}_{\pi_0} = \mathcal{T}_{\pi_0}$. Next, observe that

$$(\mathcal{T}_{\pi_0} \circ \mathcal{T}_{\pi_1}) \circ (\mathcal{T}_{\pi_0} \circ \mathcal{T}_{\pi_1}) = \mathcal{T}_{\pi_0} \circ (\mathcal{T}_{\pi_1} \circ \mathcal{T}_{\pi_0}) \circ \mathcal{T}_{\pi_1} = \mathcal{T}_{\pi_0} \circ \mathcal{T}_{\pi_1} \circ \mathcal{T}_{\pi_1} = \mathcal{T}_{\pi_0} \circ \mathcal{T}_{\pi_1}.$$

Since $\mathcal{T}_{\pi_0} \circ \mathcal{T}_{\pi_1}$ is a unital CPTP map and $\operatorname{Ran}(\mathcal{T}_{\pi_0} \circ \mathcal{T}_{\pi_1}) = \operatorname{Inv}(\pi_0), \ \mathcal{T}_{\pi_0} \circ \mathcal{T}_{\pi_1}$ is a TP conditional expectation onto $\operatorname{Inv}(\pi_0)$. By the uniqueness, we conclude that $\mathcal{T}_{\pi_0} \circ \mathcal{T}_{\pi_1} = \mathcal{T}_{\pi_0}$.

Recall that the (unnormalized) Choi-Jamiołkowski matrix $J(\Phi)$ of a linear map $\Phi : \mathcal{M}_{d_A}(\mathbb{C}) \to \mathcal{M}_{d_B}(\mathbb{C})$ is defined by

$$J(\Phi) := d_A[\Phi \otimes \mathrm{id}](|\Omega_{d_A}\rangle \langle \Omega_{d_A}|) = \sum_{i,j=1}^{d_A} \Phi(|i\rangle \langle j|) \otimes |i\rangle \langle j| \in \mathcal{M}_{d_B}(\mathbb{C}) \otimes \mathcal{M}_{d_A}(\mathbb{C}),$$
(5)

where $|\Omega_d\rangle$ is the maximally entangled state

$$|\Omega_d\rangle := \frac{1}{\sqrt{d}} \sum_{i=1}^d |ii\rangle \in \mathbb{C}^d \otimes \mathbb{C}^d, \tag{6}$$

and, for future reference, F_d is the flip (or swap) unitary operator

$$F_d := \sum_{i,j=1}^d |ij\rangle\!\langle ji| \in \mathcal{U}_{d^2}.$$
(7)

Importantly, the PPT and entanglement breaking (EB) properties of a channel can be read on its Choi matrix:

$$\Phi \text{ is PPT } \iff J(\Phi) \text{ is PPT}$$

$$\Phi \text{ is EB } \iff J(\Phi) \text{ is separable.}$$

We recall that a positive semidefinite matrix $X \in \mathcal{M}_{d_A}(\mathbb{C}) \otimes \mathcal{M}_{d_B}(\mathbb{C})$ is called

PPT if
$$X^{\Gamma} = [\mathrm{id} \otimes \top](X) \ge 0$$

separable if $X = \sum_{i} A_i \otimes B_i$ for positive semidefinite matrices $A_i \in \mathcal{M}_{d_A}(\mathbb{C}), B_i \in \mathcal{M}_{d_B}(\mathbb{C}).$

The following Proposition 2.4 establishes the connection between group symmetries of quantum channels and those of their Choi matrices.

Proposition 2.4 ([PJPY23]). Let $\pi_A : G \to \mathcal{U}_{d_A}$ and $\pi_B : G \to \mathcal{U}_{d_B}$ be unitary representations of G. Then, for any linear map $\Phi : \mathcal{M}_{d_A}(\mathbb{C}) \to \mathcal{M}_{d_B}(\mathbb{C})$,

$$J(\mathcal{T}_{\pi_A,\pi_B}\Phi) = \mathcal{T}_{\pi_B \otimes \overline{\pi_A}}(J(\Phi)).$$

In particular, $\Phi \in \text{Cov}(\pi_A, \pi_B)$ if and only if $J(\Phi) \in \text{Inv}(\pi_B \otimes \overline{\pi_A})$. Moreover, if Φ is PPT / EB, then so is $\mathcal{T}_{\pi_A, \pi_B} \Phi$.

2.2. Random quantum channels. We shall now endow the set of all quantum channels with a probability measure, laying down the foundation for the study of symmetric (covariant) channels in the subsequent sections. The set of quantum channels (i.e. completely positive, trace preserving maps) $\Phi : \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$ is a compact convex subset of the set of linear maps $\mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$. Actually, since quantum channels preserve Hermitian matrices, one can see it as a subset of hermiticity preserving maps $\mathcal{M}_d^{sa}(\mathbb{C}) \to \mathcal{M}_d^{sa}(\mathbb{C})$. The trace preservation condition adds a number of linear restrictions, easily seen in the Choi-Jamiołkowski picture from Eq. (5): a linear map Φ is trace preserving iff

$$[\operatorname{Tr} \otimes \operatorname{id}](J(\Phi)) = I_d$$

Let V be a Haar-distributed random isometry of size $ds \times d$ and define the random Stinespring channel $\Phi_V : \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$ by

$$\Phi_V(\rho) := (\mathrm{id}_d \otimes \mathrm{Tr}_s)(V\rho V^*). \tag{8}$$

The probability distribution of Φ_V depends on the choice of dimension parameters d and s, and we shall denote it by $\mu_{d,s}$. Let us recall that a Haar-distributed random isometry $V : \mathbb{C}^d \to \mathbb{C}^{ds}$ can be obtained by truncating a random Haar-distributed unitary matrix $U \in \mathcal{U}_{ds}$.

According to [KNP⁺21b], we can give an equivalent definition for a random Stinespring channel Φ_V . We first consider a random Wishart matrix $W = GG^*$ of parameter (d^2, s) and set $H := (\operatorname{Tr}_d \otimes \operatorname{id}_d)W$. Then H is almost surely invertible, and the probability distribution of $J(\Phi_V)$ is the same as that of

$$\tilde{J} := (I_d \otimes H^{-1/2}) W(I_d \otimes H^{-1/2}).$$
 (9)

Note that the matrix \tilde{J} is PPT (resp. separable) if and only if W is PPT (resp. separable). We gather below some of the foundational results from [Aub12, ASY14, CYZ18]. In our work, we shall prove similar results in the case of covariant channels.

Proposition 2.5. Consider the random Stinespring channel $\Phi_{d,s}$ with parameter (d, s).

- (1) (Threshold phenomenon of PPT property for Φ) For any $\varepsilon > 0$
 - if $s < (4 \varepsilon)d^2$, $\lim_{d \to \infty} \mathbb{P}(\Phi_{d,s} \text{ is } PPT) = 0$,
 - if $s > (4 + \varepsilon)d^2$, $\lim_{d \to \infty} \mathbb{P}(\Phi_{d,s} \text{ is } PPT) = 1$.

(2) (Threshold phenomenon of EB property for Φ) There exist constants $C_1, C_2 > 0$ and a sequence s_d satisfying

$$C_1 d^3 \le s_d \le C_2 d^3 (\log d)^2$$

such that for any $\varepsilon > 0$,

- if $s < (1 \varepsilon)s_d$, $\lim_{d \to \infty} \mathbb{P}(\Phi_{d,s} \text{ is } EB) = 0$,
- if $s > (1 + \varepsilon)s_d$, $\lim_{d \to \infty} \mathbb{P}(\Phi_{d,s} \text{ is } EB) = 1$.
- (3) (PPT² conjecture for Φ) Let $\tilde{\Phi}_{d,s}$ be a random Stinespring channel independent to $\Phi_{d,s}$. For any $\varepsilon > 0$, if

$$(4+\varepsilon)d^2 < s < (1-\varepsilon)s_d$$

so that $\Phi_{d,s}$ and $\widetilde{\Phi}_{d,s}$ are generically PPT and not EB, then $\lim_{d\to\infty} \mathbb{P}(\Phi_{d,s} \circ \widetilde{\Phi}_{d,s} \text{ is } EB) = 1.$

Let us end this introductory section by providing the general recipe that shall be used to produce *random covariant channels* in the following sections:

- (1) first, we sample a random quantum channel Φ_V , without any symmetry, from the ensemble $\mu_{d,s}$, see (8)
- (2) we then symmetrize Φ using the twirling operation below, to obtain a random (π_A, π_B) covariant channel Φ_{π_A, π_B} :

$$\Phi_{\pi_A,\pi_B}(X) := \mathcal{T}_{\pi_A,\pi_B}(\Phi_V)(X) = \int_G \pi_B(g)^* \Phi_V \Big(\pi_A(g) X \pi_A(g)^* \Big) \pi_B(g) \,\mathrm{d}\mathfrak{h}(g), \tag{10}$$

where \mathfrak{h} is the Haar measure on the group G.

3. RANDOM ORTHOGONAL COVARIANT CHANNELS

In this section we study three basic and important classes of linear maps in quantum information theory, those having a unitary or orthogonal covariance group. Unitary covariance is a very strong form of symmetry, since input states can be diagonalized, the effect of the unitary rotation being transferred to the output of the channel. Such strong symmetry makes very suitable for analysis in various settings related to quantum Shannon theory. We recall below their definition.

(1) The depolarizing maps characterized by (U, U)-covariance:

$$\Phi(UZU^*) = U\Phi(Z)U^*, \ Z \in \mathcal{M}_d(\mathbb{C}), \ U \in \mathcal{U}_d,$$

(2) The transpose depolarizing maps characterized by (U, \overline{U}) -covariance:

$$\Phi(UZU^*) = \overline{U}\Phi(Z)U^{\top}, \ Z \in \mathcal{M}_d(\mathbb{C}), \ U \in \mathcal{U}_d,$$

(3) The orthogonal covariant maps characterized by (O, O)-covariance:

$$\Phi(OZO^{\top}) = O\Phi(Z)O^{\top}, \ Z \in \mathcal{M}_d(\mathbb{C}), \ O \in \mathcal{O}_d,$$

Proposition 3.1. Let $\Phi : \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$ be a quantum channel. Then, the (U,U), (U,\overline{U}) , (O,O) twirlings of Φ are given, respectively, by:

$$\mathcal{T}_{U,U}(\Phi) = p \cdot \mathrm{id} + (1-p) \cdot \Delta$$

$$\mathcal{T}_{U,\overline{U}}(\Phi) = q \cdot \top + (1-q) \cdot \Delta$$

$$\mathcal{T}_{O,O}(\Phi) = p' \cdot \mathrm{id} + q' \cdot \top + (1-p'-q') \cdot \Delta,$$

(11)

where id, \top, Δ are, respectively, the identity, the transposition, and the completely depolarizing linear maps. Here the coefficients p, q, p', q' are explicitly related to Φ by

$$p = \frac{\lambda_1 - 1}{d^2 - 1},$$

$$q = \frac{\lambda_2 - 1}{d^2 - 1},$$

$$p' = \frac{(d+1)\lambda_1 - \lambda_2 - d}{d(d+2)(d-1)},$$

$$q' = \frac{(d+1)\lambda_2 - \lambda_1 - d}{d(d+2)(d-1)},$$

where the parameters $\lambda_{1,2}$ associated to the channel are defined (see also Figure 2):

$$\lambda_1(\Phi) := d \langle \Omega_d | J(\Phi) | \Omega_d \rangle$$

$$\lambda_2(\Phi) := \operatorname{Tr}(J(\Phi) F_d).$$

Above we have used maximally entangled state $|\Omega_d\rangle$ and the flip operator F_d defined respectively in Eq. (6) and Eq. (7).

Proof. All the formulas follow from Proposition 2.2. Regarding the (U, U)-twirling $\mathcal{T}_{U,U}$, note that the representation $U \in \mathcal{U}_d \mapsto U \otimes \overline{U}$ can be decomposed into two irreducible sub-representations whose corresponding projections are

$$|\Omega_d\rangle\langle\Omega_d|, \quad I_{d^2} - |\Omega_d\rangle\langle\Omega_d|.$$

Therefore, Proposition 2.2 and 2.4 implies that

$$\begin{split} J(\mathcal{T}_{U,U}\Phi) &= \mathcal{T}_{U\otimes\overline{U}}(J(\Phi)) \\ &= \mathrm{Tr}(|\Omega_d\rangle\langle\Omega_d|J(\Phi))|\Omega_d\rangle\langle\Omega_d| + \mathrm{Tr}((I_{d^2} - |\Omega_d\rangle\langle\Omega_d|)J(\Phi))\frac{I_{d^2} - |\Omega_d\rangle\langle\Omega_d|}{d^2 - 1} \\ &= \frac{\lambda_1}{d}|\Omega_d\rangle\langle\Omega_d| + \left(d - \frac{\lambda_1}{d}\right)\frac{I_{d^2} - |\Omega_d\rangle\langle\Omega_d|}{d^2 - 1} \\ &= \frac{\lambda_1 - 1}{d^2 - 1}d|\Omega_d\rangle\langle\Omega_d| + \frac{d^2 - \lambda_1}{d^2 - 1}\frac{I_{d^2}}{d}, \end{split}$$

where we used $\operatorname{Tr}(J(\Phi)) = d$ since Φ is TP. Therefore, we have the formula $\mathcal{T}_{U,U}(\Phi)$ as in Eq. (11).

The remaining formulas can be derived analogously by noting that the representations $U \in \mathcal{U}_d \mapsto U \otimes U$ and $O \in \mathcal{O}_d \mapsto O \otimes O$ have the irrep decompositions corresponding to the projections (see [TiB16, Appendix A] and [Bra37])

$$\Pi_{\text{sym}} := \frac{I_{d^2} + F_d}{2}, \quad \Pi_{\text{anti}} := \frac{I_{d^2} - F_d}{2},$$

and

$$|\Omega_d\rangle\langle\Omega_d|, \quad \Pi_{\rm sym} - |\Omega_d\rangle\langle\Omega_d|, \quad \Pi_{\rm anti},$$

respectively.

The channel parameters
$$\lambda_{1,2}(\Phi)$$
 can take values in specific intervals, as follows.

Proposition 3.2. For a channel $\Phi : \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$, we have

$$0 \le \lambda_1(\Phi) \le d^2$$
$$-d \le \lambda_2(\Phi) \le d,$$

with the inequalities being tight.

$$\lambda_{1}(\Phi) = \left(\begin{array}{c} J(\Phi) \end{array} \right) = \sum_{i,j=1}^{d} \langle i | \Phi(|i\rangle \langle j|) | j \rangle$$
$$\lambda_{2}(\Phi) = \left(\begin{array}{c} J(\Phi) \end{array} \right) = \sum_{i,j=1}^{d} \langle i | \Phi(j\rangle \langle i|) | j \rangle$$

FIGURE 2. The quantities $\lambda_{1,2}(\Phi)$ associated to a quantum channel $\Phi : \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$.

Proof. The map $\Phi \mapsto \mathcal{T}_{U,U}(\Phi)$ is onto the set of channels of the form $p \cdot \mathrm{id} + (1-p) \cdot \Delta$. The range of p for which the latter maps are quantum channels is known to be

$$p \in \Big[\frac{-1}{d^2 - 1}, 1\Big],$$

hence the claim for $\lambda_1(\Phi)$ follows. The second claim can be proven in a similar manner, since the allowed range of q is

$$q \in \left[\frac{-1}{d-1}, \frac{1}{d+1}\right].$$

The PPT and entanglement breaking properties of (conjugate) unitary and orthogonal covariant channels are fully characterized by the parameters $\lambda_{1,2}(\Phi)$, as follows.

Proposition 3.3 ([VW01]). For the parameters $\lambda_1(\Phi)$ and $\lambda_2(\Phi)$ defined above,

- (1) $\mathcal{T}_{U,U}(\Phi)$ is PPT iff it is EB iff $0 \le \lambda_1(\Phi) \le d$; (2) $\mathcal{T}_{U,\overline{U}}(\Phi)$ is PPT iff it is EB iff $0 \le \lambda_2(\Phi) \le d$;
- (3) $\mathcal{T}_{O,O}(\Phi)$ is PPT iff it is EB iff $0 \leq \lambda_1(\Phi), \lambda_2(\Phi) \leq d$.

We define now random covariant quantum channels by twirling the random quantum channels introduced in Section 2.2. More precisely, we introduce the following image distributions:

$$\mu_{d,s}^{U,U} := (\mathcal{T}_{U,U})_{\#} \mu_{d,s}$$
$$\mu_{d,s}^{U,\overline{U}} := (\mathcal{T}_{U,\overline{U}})_{\#} \mu_{d,s}$$
$$\mu_{d,s}^{O,O} := (\mathcal{T}_{O,O})_{\#} \mu_{d,s}.$$

Recall that $\mu_{d,s}$ is the probability distribution of the random channel Φ_V associated to the Haar random isometry $V : \mathbb{C}^d \to \mathbb{C}^d \otimes \mathbb{C}^s$. The distributions above are characterized by the parameters p, q, p', q' from Proposition 3.1. In turn, these are related to the isometry V defining the channel Φ via the following quantities:

$$\lambda_1^{(d,s)} := \lambda_1(\Phi_V) = d\langle \Omega_d | J(\Phi_V) | \Omega_d \rangle$$

$$\lambda_2^{(d,s)} := \lambda_2(\Phi_V) = \operatorname{Tr}(J(\Phi_V)F_d),$$

see Figure 3 for a graphical representations of these scalars. Above, the random channel Φ_V is defined via the Stinespring dilation from the Haar-distributed random isometry $V : \mathbb{C}^d \to \mathbb{C}^d \otimes \mathbb{C}^s$:

$$\Phi_V(\rho) = \operatorname{Tr}_s \left[V \rho V^* \right].$$

In order to understand the PPT and the entanglement breaking behavior of random (conjugate) unitary and orthogonal covariant channels, we need first to understand the large d limit of the



FIGURE 3. The quantities $\lambda_{1,2}^{(d,s)}$ associated to a Haar-distributed random isometry $V : \mathbb{C}^d \to \mathbb{C}^d \otimes \mathbb{C}^s$.

random variables $\lambda_{1,2}^{(d,s)}$. To start with, it follows from Lemma A.11 that the random variables $\lambda_{1,2}^{(d,s)}$ have average and variance given by

$$\mathbb{E}\left[\lambda_{1,2}^{(d,s)}\right] = 1 \quad \text{and} \quad \operatorname{Var}\left[\lambda_{1,2}^{(d,s)}\right] = \frac{s(d^2 - 1)}{(ds)^2 - 1}.$$

Note that although $\lambda_{1,2}^{(d,s)}$ have the same first two moments, they do not have the same distribution, see Lemma A.11 for the exact moments of all orders. In fact, they have very different behavior as $d \to \infty$, as we shall see next. We first recall the definitions of the *normal* (or *Gaussian*) distribution of mean μ and variance σ^2 :

$$X \sim \mathcal{N}(m, \sigma^2) \qquad \iff \qquad \mathrm{d}X = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(\frac{-(x-m)^2}{2\sigma^2}\right) \mathrm{d}x,$$

and that of the *Gamma* distribution of parameters (α, β)

$$Y \sim \Gamma(\alpha, \beta) \qquad \iff \qquad \mathrm{d}Y = \frac{\beta^{\alpha}}{\Gamma(\alpha)} y^{\alpha-1} \exp(-\beta y) \mathbf{1}_{y>0} \mathrm{d}y$$

Note that the Gamma distribution of parameters $(1, \lambda)$ is also called the *exponential* distribution of parameter λ , for which we denote by $\mathsf{Exp}(\lambda) := \Gamma(1, \lambda)$.

Proposition 3.4. As $d \to \infty$, the random variables $\lambda_{1,2}^{(d,s)}$ behave as follows:

- (1) If s is fixed, then
 - the law of the random variable $s\lambda_1^{(d,s)}$ converges in moments to the Gamma distribution $\Gamma(s,1)$
 - the law of the random variable $s\lambda_2^{(d,s)}$ converges in moments to the normal distribution $\mathcal{N}(s,s)$.
- (2) If $s = s(d) \rightarrow \infty$, then

$$\lambda_{1,2}^{(d,s)} \to 1 \text{ in } L^p \text{ for all } p < \infty.$$

Furthermore, the same convergence holds almost surely if $s \gtrsim d^t$ for some t > 0. (3) Moreover, for any t > 0, we have

$$d^{-t}(\lambda_1^{(d,s)}-1) \to 0 \qquad and \qquad d^{-t}(\lambda_2^{(d,s)}-1) \to 0$$

almost surely as $d \to \infty$ or $s \to \infty$ (more precisely, this means that " $d \to \infty$ and s = s(d) arbitrary" or " $s \to \infty$ and d = d(s) arbitrary").

Proof. See Appendix A.2.

We plot numerical realizations of the random variables $\lambda_{1,2}^{(d,s)}$ against the theoretical curves in Figs. 4 and 5.

Theorem 3.5. Let $\Phi_{U,U} \sim \mu_{d,s}^{U,U}$, $\Phi_{U,\overline{U}} \sim \mu_{d,s}^{U,\overline{U}}$, and $\Phi_{O,O} \sim \mu_{d,s}^{O,O}$ be, respectively, random (U,U)-covariant, (U,\overline{U}) -covariant, and (O,O)-covariant quantum channels.



FIGURE 4. Histograms of 10^4 realizations of the random variable $s\lambda_1^{(d,s)}$ for d = 100 and, respectively, s = 1, 2, 5, 10. In blue, the probability density of the $\Gamma(s, 1)$ distribution.



FIGURE 5. Histograms of 10⁴ realizations of the random variable $s\lambda_2^{(d,s)}$ for d = 100 and, respectively, s = 1, 5. In blue, the probability density of the $\mathcal{N}(s, s)$ distribution.

- (1) Almost surely, as $d \to \infty$, $\Phi_{U,U}$ is PPT and EB.
- (2) If s is fixed, then

 $\lim_{d\to\infty} \mathbb{P}(\Phi_{U,\overline{U}} \text{ is } PPT \Leftrightarrow EB) = \lim_{d\to\infty} \mathbb{P}(\Phi_{O,O} \text{ is } PPT \Leftrightarrow EB) = \mathbb{P}(Z \ge -\sqrt{s}) < 1,$ where $Z \sim \mathcal{N}(0,1)$ is a standard normal random variable.

(3) If $s = s(d) \to \infty$ as $d \to \infty$, then

$$\lim_{d \to \infty} \mathbb{P}(\Phi_{U,\overline{U}} \text{ is } PPT \Leftrightarrow EB) = \lim_{d \to \infty} \mathbb{P}(\Phi_{O,O} \text{ is } PPT \Leftrightarrow EB) = 1,$$

Furthermore, if $s \gtrsim d^t$ for some t > 0, then almost surely as $d \to \infty$, $\Phi_{U,\overline{U}}$ and $\Phi_{O,O}$ are PPT and EB.

We remark that by the statement ' $\Phi = \Phi_d$ is almost surely PPT/EB as $d \to \infty$ ', we mean the eventual property of Φ_d : there exists an event set Ω of probability 1 such that for every $\omega \in \Omega$, there exists a number $N = N(\omega) \ge 1$ such that $\Phi_d(\omega)$ is PPT/EB for all $d \ge N$. In particular, this implies that $\mathbb{P}(\Phi \text{ is PPT/EB}) \to 1$ as $d \to \infty$.

Proof. We start with the first item, related to the unitary covariant channel $\Phi_{U,U}$ and the random variable $\lambda_1^{(d,s)} \ge 0$. It follows from Proposition 3.4, (3) with t = 1, that, almost surely,

$$\lim_{d\to\infty}\frac{\lambda_1^{(d,s)}}{d}=0$$

which yields (1) by Proposition 3.3.

Let us now move to the second point, corresponding to the regime s fixed and the channels $\Phi_{U,\overline{U}}$ and $\Phi_{O,O}$. From the previous point, we have that, almost surely as $d \to \infty$, $\lambda_1^{(d,s)} \leq d$; similarly for $\lambda_2^{(d,s)}$. The only constraint in Proposition 3.3 that we need to ensure is $\lambda_2^{(d,s)} \geq 0$, and the conclusion follows from Proposition 3.4 (1): if $\tilde{Z} \sim \mathcal{N}(s, s)$, then

$$\mathbb{P}(\tilde{Z} \ge 0) = \mathbb{P}(\sqrt{sZ} + s \ge 0) = \mathbb{P}(Z \ge -\sqrt{s}),$$

for a standard normal random variable $Z \sim \mathcal{N}(0, 1)$.

Finally, regarding the proof of (3), the only constraint again is whether $\lambda_2^{(d,s)} \ge 0$. Then the assertions follow directly from Proposition 3.4 (2): for the general regime $s(d) \to \infty$, $\lambda_2^{(d,s)} \to 1$ in probability, so we have

$$\lim_{d \to \infty} \mathbb{P}(\Phi_{O,O} \text{ is } \text{PPT}) = \lim_{d \to \infty} \mathbb{P}(\lambda_2^{(d,s)} \ge 0) = 1,$$

and analogously $\lim_{d\to\infty} \mathbb{P}(\Phi_{U,\overline{U}} \text{ is PPT}) = 1$. Furthermore, when $s \gtrsim d^t$ for some t > 0, we have $\lambda_{1,2}^{(d,s)} \to 1$ hence $0 \leq \lambda_{1,2}^{(d,s)} \leq d$ almost surely.

Remark 3.6. One can also show, by slightly modifying the previous arguments, that if $s \to \infty$, then, independently of the behavior of d = d(s), all the three random channels $\Phi_{U,U}$, $\Phi_{U,\overline{U}}$, and $\Phi_{O,O}$ are almost surely PPT and entanglement breaking.

In conclusion, we have shown that the equivalent PPT and EB properties for large $(d \to \infty)$ random (conjugate) unitary and orthogonal covariant quantum channels are as follows:

- random unitary covariant channels are asymptotically PPT and EB (for any s)
- random conjugate unitary and orthogonal covariant channels admit a phase transition between the regimes s fixed and $s \to \infty$.

4. RANDOM HYPEROCTAHEDRAL COVARIANT CHANNELS

Here we consider a random behavior of the channels which are covariant with respect to hyperoctahedral group. Recall that one can realize the hyperoctahedral group of degree d as the subgroup \mathcal{H}_d of the orthogonal group \mathcal{O}_d which is generated by diagonal orthogonal matrices and permutation matrices. The group \mathcal{H}_d is also known as the group of signed permutations and it has order $2^d \cdot d!$. It is recently shown [PJPY23, Proposition 4.4] that a linear map $\Phi : \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$ is covariant with respect to the fundamental representation $\pi_{\mathcal{H}_d} : H \in \mathcal{H}_d \mapsto H \in \mathcal{O}_d$ if and only if

$$\Phi(Z) = p \frac{\operatorname{Tr}(Z)}{d} I_d + qZ + rZ^\top + s \operatorname{diag}(Z), \ Z \in \mathcal{M}_d(\mathbb{C}),$$
(12)

for some $p, q, r, s \in \mathbb{C}$. In other words, the vector space of hyperoctahedral covariant linear maps is spanned by three quantum channels (the completely depolarizing, the identity, and the diagonal conditional expectation) and the transposition map. Note that the formula above covers the (conjugate) unitary and orthogonal covariant quantum channels discussed in Section 3. Let us simply denote $\operatorname{Cov}(H, H) := \operatorname{Cov}(\pi_{\mathcal{H}_d}, \pi_{\mathcal{H}_d})$ and $\operatorname{Inv}(H, H) := \operatorname{Inv}(\pi_{\mathcal{H}_d} \otimes \pi_{\mathcal{H}_d}) = J(\operatorname{Cov}(H, H))$ (by Proposition 2.4). Note that the Choi matrix of the map Φ given in (12) is

$$J(\Phi) = p \frac{I_{d^2}}{d} + dq |\Omega_d\rangle \langle \Omega_d| + rF_d + s\Pi_{\text{diag}} \in \text{Inv}(H \otimes H),$$
(13)

where $|\Omega_d\rangle = \frac{1}{\sqrt{d}} \sum_i |ii\rangle$ and F_d are defined in Eqs. (6) and (7), and

$$\Pi_{\rm diag} := \sum_{i=1}^d |ii\rangle \langle ii$$

is the Choi matrix of the "diagonalization" (or completely dephasing) map $Z \mapsto \text{diag}(Z)$. From this we can check that the space Inv(H, H) = J(Cov(H, H)) is spanned by four mutually orthogonal projections

$$\begin{split} \Pi_0 &= |\Omega_d \rangle \langle \Omega_d |, \\ \Pi_1 &= \Pi_{\text{sym}} - \Pi_{\text{diag}} = \frac{I_{d^2} + F}{2} - \Pi_{\text{diag}}, \\ \Pi_2 &= \Pi_{\text{anti}} = \frac{I_{d^2} - F}{2}, \\ \Pi_3 &= \Pi_{\text{diag}} - |\Omega_d \rangle \langle \Omega_d |. \end{split}$$

Therefore, Proposition 2.2 gives the exact formula of $\mathcal{T}_{H\otimes H} := \mathcal{T}_{\pi_{\mathcal{H}_d}\otimes\pi_{\mathcal{H}_d}}$,

$$\mathcal{T}_{H\otimes H}X = \sum_{i=0}^{3} \operatorname{Tr}(\Pi_{i}X) \frac{\Pi_{i}}{d_{i}}, \quad X \in \mathcal{M}_{d}(\mathbb{C}) \otimes \mathcal{M}_{d}(\mathbb{C}),$$
(14)

where the dimensions read $d_0 = 1$, $d_1 = d_2 = \frac{d^2 - d}{2}$, and $d_3 = d - 1$. We can now give the formula for the (H, H)-twirling of a quantum channel.

Proposition 4.1. Let $\Phi : \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$ be a quantum channel. Then, the (H, H) twirling of Φ is given by:

$$\mathcal{T}_{H,H}(\Phi) = p \cdot \Delta + q \cdot \mathrm{id} + r \cdot \top + (1 - p - q - r) \cdot \mathrm{diag}$$

where $id, \top, diag, \Delta$ are, respectively, the identity, the transposition, the diagonalization, and the completely depolarizing linear maps. Here the coefficients p, q, r are explicitly related to Φ by

$$p = \frac{d - \lambda_3(\Phi)}{d - 1}$$
$$q = \frac{\lambda_1(\Phi) - \lambda_3(\Phi)}{d^2 - d}$$
$$r = \frac{\lambda_2(\Phi) - \lambda_3(\Phi)}{d^2 - d},$$

where the parameters $\lambda_{1,2,3}(\Phi)$ associated to the channel are defined by (see also Figs. 2 and 6):

$$\lambda_1(\Phi) = d\langle \Omega_d | J(\Phi) | \Omega_d \rangle$$

$$\lambda_2(\Phi) = \operatorname{Tr}(J(\Phi)F_d)$$

$$\lambda_3(\Phi) := \operatorname{Tr}(J(\Phi)\Pi_{\text{diag}}).$$

$$\lambda_{3}(\Phi) = \bigcup_{i=1}^{d} \langle i | \Phi(|i\rangle\langle i|) | i \rangle \qquad \qquad \lambda_{3}^{(d,s)} = \bigcup_{s}^{d} \bigcup_{s} V^{*}$$

FIGURE 6. The quantity $\lambda_3(\Phi)$ associated to a quantum channel $\Phi : \mathcal{M}_d(\mathbb{C}) \to$ $\mathcal{M}_d(\mathbb{C})$ and the corresponding random variable $\lambda_3^{(d,s)} = \lambda_3(\Phi_V)$ for a random channel obtained from a Haar-distributed random isometry $V : \mathbb{C}^d \to \mathbb{C}^d \otimes \mathbb{C}^s$.

Proof. By Proposition 2.4 and Equation (14), we have

$$J(\mathcal{T}_{H,H}\Phi) = \mathcal{T}_{H\otimes H}(J(\Phi)) = \sum_{i=0}^{3} \operatorname{Tr}(\Pi_{i}J(\Phi))\frac{\Pi_{i}}{d_{i}}$$

$$= \frac{\lambda_{1}}{d}|\Omega_{d}\rangle\langle\Omega_{d}| + \left(\frac{d+\lambda_{2}}{2} - \lambda_{3}\right)\frac{I_{d^{2}} + F_{d} - 2\Pi_{\text{diag}}}{d^{2} - d}$$

$$+ \frac{d-\lambda_{2}}{2}\frac{I_{d^{2}} - F_{d}}{d^{2} - d} + \left(\lambda_{3} - \frac{\lambda_{1}}{d}\right)\frac{\Pi_{\text{diag}} - |\Omega_{d}\rangle\langle\Omega_{d}|}{d - 1}$$

$$= \frac{d-\lambda_{3}}{d - 1}\frac{I_{d^{2}}}{d} + \frac{\lambda_{1} - \lambda_{3}}{d^{2} - d}d|\Omega_{d}\rangle\langle\Omega_{d}| + \frac{\lambda_{2} - \lambda_{3}}{d^{2} - d}F_{d} + \frac{(d+2)\lambda_{3} - \lambda_{1} - \lambda_{2} - d}{d^{2} - d}\Pi_{\text{diag}}.$$
we used the fact $\operatorname{Tr}(J(\Phi)) = d$ since Φ is TP.

Here we used the fact $Tr(J(\Phi)) = d$ since Φ is TP.

One can now state the precise conditions for a (H, H)-twirling of a quantum channel to be PPT; the PPT property has been shown in PJPY23, Theorem 4.1 to be equivalent to the entanglement breaking property for hyperoctahedral covariant channels.

Proposition 4.2. Given a quantum channel $\Phi : \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$, its (H, H)-twirling $\mathcal{T}_{H,H}(\Phi)$ is PPT iff it is EB iff

$$0 \le \lambda_3(\Phi) \le d,$$

$$0 \le \lambda_1(\Phi), \lambda_2(\Phi) \le d\lambda_3(\Phi),$$

$$2\lambda_3(\Phi) - d \le \lambda_1(\Phi), \lambda_2(\Phi) \le d.$$
(15)

Proof. The conditions in the statement are simply the following inequalities from [PJPY23, Eq. (4.17)] reformulated in terms of the parameters $\lambda_{1,2,3}(\Phi)$:

$$0 \leq p \leq \frac{d}{d-1},$$

$$\frac{p}{d} - \frac{1}{d-1} \leq q, r \leq 1 - \frac{d-1}{d}p,$$

$$-\frac{p}{d} \leq q, r \leq \frac{p}{d}.$$

Now we consider a random Stinespring channel $\Phi: \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$ with environment dimension s and define the random hyperoctahedral covariant channel

$$\Phi_{H,H} := \mathcal{T}_{H,H}(\Phi) \sim \mu_{d,s}^{H,H}.$$

Since the PPT (and therefore the EB) property of the twirled channel $\mathcal{T}_{H,H}(\Phi)$ depends only on the three parameters $\lambda_{1,2,3}(\Phi)$, we shall consider the corresponding random variables associated to the random channel:

$$\lambda_1^{(d,s)} = d\langle \Omega_d | J(\Phi) | \Omega_d \rangle, \quad \lambda_2^{(d,s)} = \operatorname{Tr}(J(\Phi)F_d), \quad \lambda_3^{(d,s)} := \operatorname{Tr}(J(\Phi)\Pi_{\operatorname{diag}}).$$

The asymptotic behavior of the first two variables has been studied in the previous section, Proposition 3.4; we now turn to the study of $\lambda_3^{(d,s)}$. The average and the variance of this random variable are given by (Lemma A.11)

$$\mathbb{E}\left[\lambda_3^{(d,s)}\right] = 1$$
 and $\operatorname{Var}\left[\lambda_3^{(d,s)}\right] = \frac{1}{ds+1}$

Proposition 4.3. The following almost sure convergence holds, independently of the scaling of the parameter s:

$$\lim_{d \to \infty} \lambda_3^{(d,s)} = 1.$$

Proof. See Appendix A.2.

We can now prove the main result of this section, giving the PPT and EB properties of a hyperoctahedral covariant random channel, in the $d \to \infty$ regime.

Theorem 4.4. Let $\Phi_{H,H} \sim \mu_{d,s}^{H,H}$ be a random (H,H)-covariant quantum channel. Then:

(1) If s is fixed,

$$\lim_{h \to \infty} \mathbb{P}(\Phi_{H,H} \text{ is } PPT \Leftrightarrow EB) = \mathbb{P}(Z \ge -\sqrt{s}) < 1,$$

where $Z \sim \mathcal{N}(0,1)$ is a standard normal random variable. (2) If $s = s(d) \to \infty$ as $d \to \infty$, then

$$\lim_{d\to\infty} \mathbb{P}(\Phi_{H,H} \text{ is } PPT \Leftrightarrow EB) = 1.$$

Furthermore, if $s \gtrsim d^t$ for some t > 0, then almost surely as $d \to \infty$, $\Phi_{H,H}$ is PPT and EB.

Proof. Since $\lambda_3^{(d,s)} \to 1$ as $d \to \infty$, the equations (15) reduce to the conditions from Proposition 3.3 (3). The proof is now the same as Theorem 3.5 (2) and (3).

Note that

$$\mathcal{T}_{U,\overline{U}} \circ \mathcal{T}_{H,H} = \mathcal{T}_{U,\overline{U}}, \quad \mathcal{T}_{U,U} \circ \mathcal{T}_{H,H} = \mathcal{T}_{U,U}, \quad \text{and} \quad \mathcal{T}_{O,O} \circ \mathcal{T}_{H,H} = \mathcal{T}_{O,O},$$

by Proposition 2.3 and Proposition 2.4. Since the twirling operations preserve both PPT and entanglement breaking properties, Theorem 4.4 directly implies the asymptotic PPT and EB behavior of random depolarizing / transpose depolarizing / orthogonal covariant channels, in the regime where $s \gtrsim d^t$ with $d \to \infty$, recovering part of Theorem 3.5.

Corollary 4.5. Suppose $s \gtrsim d^t$ for some constants t > 0. Then almost surely, the random depolarizing channel $\mathcal{T}_{U,\overline{U}}\Phi$, the random transpose depolarizing channel $\mathcal{T}_{U,U}\Phi$, and the random orthogonal covariant channel $\mathcal{T}_{O,O}\Phi$ are PPT and EB, as $d \to \infty$.

In conclusion, we have shown that the equivalent PPT and EB properties for large $(d \to \infty)$ random hyperoctahedral covariant quantum channels admit a phase transition between the regimes s fixed and $s \to \infty$, which is completely parallel with random conjugate unitary and orthogonal covariant channels.

5. Random diagonal orthogonal covariant channels

In this section, we consider random quantum channels which are covariant with respect to the actions of the diagonal unitary group

$$\mathcal{DU}_d := \left\{ \operatorname{diag}(z) : z \in \mathbb{C}^d, \, |z_i| = 1 \quad \forall i \in [d] \right\}$$

and the diagonal orthogonal group

$$\mathcal{DO}_d := \left\{ \operatorname{diag}(\varepsilon) : \varepsilon \in \{\pm 1\}^d \right\}$$

Bipartite quantum states that are invariant with respect to the actions of the groups above, as well as channels that are covariant for these actions have appeared in the literature in the last 20 years [DSS⁺⁰⁰, CK06, Liu15, LS15, SBSGS22], but have been studied in detail in [SN21]. Since, they have found many applications, see e.g. [SN22a, SN22b, SDN22]. We recall from [SN21, Definition 6.3] that a quantum channel $\Phi : \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$ is called:

• conjugate diagonal unitary covariant (CDUC) if

$$\forall Z \in \mathcal{M}_d(\mathbb{C}) \text{ and } U \in \mathcal{DU}_d: \qquad \Phi(UZU^*) = U^* \Phi(Z)U,$$

• diagonal unitary covariant (DUC) if

$$\forall Z \in \mathcal{M}_d(\mathbb{C}) \text{ and } U \in \mathcal{DU}_d: \quad \Phi(UZU^*) = U\Phi(Z)U^*,$$

• diagonal orthogonal covariant (DOC) if

$$\forall Z \in \mathcal{M}_d(\mathbb{C}) \text{ and } O \in \mathcal{DO}_d : \Phi(OZO) = O\Phi(Z)O.$$

Note that we employ a different naming convention than the one in [SN21], by interchanging the DUC and CDUC terms. In the present work, the emphasis is on the covariance property, hence we prefer to use the "conjugate" adjective for the case where $\pi_B = \overline{\pi_A}$.

The structure of linear maps satisfying the covariance conditions above has been obtained in [SN21, Eqs. (36)-(38)]: a linear map $\Phi: \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$ is

• CDUC if there exists a pair $A, C \in \mathcal{M}_d(\mathbb{C})$ such that diag A = diagC and

$$\forall Z \in \mathcal{M}_d(\mathbb{C}) \qquad \Phi(Z) = \Phi_{A,C}^{(1)}(Z) = \operatorname{diag}(A | \operatorname{diag}(Z) \rangle) + \mathring{C} \odot Z^\top$$

• DUC if there exists a pair $A, B \in \mathcal{M}_d(\mathbb{C})$ such that diag A = diagB and

$$\forall Z \in \mathcal{M}_d(\mathbb{C}) \qquad \Phi(Z) = \Phi_{A,B}^{(2)}(Z) = \operatorname{diag}(A |\operatorname{diag}(Z)\rangle) + \mathring{B} \odot Z.$$

• DOC if there exists a triple $A, B, C \in \mathcal{M}_d(\mathbb{C})$ such that diag A = diagB = diagC and

$$\forall Z \in \mathcal{M}_d(\mathbb{C}) \qquad \Phi(Z) = \Phi_{A,B,C}^{(3)}(Z) = \operatorname{diag}(A | \operatorname{diag}(Z) \rangle) + \mathring{B} \odot Z + \mathring{C} \odot Z^\top.$$

Above, the notation $\overset{\circ}{X}$, for a matrix $X \in \mathcal{M}_d(\mathbb{C})$, denotes the matrix X without its diagonal, i.e.

$$\mathring{X} := X - \operatorname{diag} X \quad \text{or} \quad (\mathring{X})_{ij} = \begin{cases} X_{ij} & \text{if } i \neq j \\ 0 & \text{if } i = j. \end{cases}$$

Note that the maps $\Phi^{(1,2)}$ can be seen as special cases of the DOC map $\Phi^{(3)}$ by setting, respectively, B = diagB and C = diagC. A linear DOC map $\Phi_{A,B,C}^{(3)}$ is completely positive (CP) iff

- A is entrywise nonnegative: $A_{ij} \ge 0$ for all $i, j \in [d]$
- B is positive semidefinite: B ≥ 0
 C is selfadjoint and |C_{ij}|² ≤ A_{ij}A_{ji} for all i, j ∈ [d].

A linear DOC map $\Phi_{A,B,C}^{(3)}$ is trace preserving (TP) iff A is column stochastic, i.e.

$$\forall j \in [d] \qquad \sum_{i=1}^d A_{ij} = 1.$$

It is important to note at this point that the sets of CDUC, DUC, and DOC linear maps or quantum channels are parametrized by a pair or a triple of $d \times d$ matrices satisfying certain conditions; the number of parameters in these models is growing with the dimension d. This fact is in stark contrast with the symmetries investigated in Section 3 (full-unitary and full-orthogonal) and in Section 4 (hyperoctahedral). It is the richness of these models that allows for the existence of PPT symmetric channels that are not entanglement breaking, as we show next. Let us first discuss the PPT property for diagonal symmetric channels.

Proposition 5.1 ([SN21]). For any triple of matrices $A, B, C \in \mathcal{M}_d(\mathbb{C})$ having the same diagonal, we have

$$\Phi_{A,C}^{(1)} \circ \top = \Phi_{A,C}^{(2)}, \qquad \Phi_{A,B}^{(2)} \circ \top = \Phi_{A,B}^{(1)}, \qquad \Phi_{A,B,C}^{(3)} \circ \top = \Phi_{A,C,B}^{(3)}$$

Hence, we have the following characterizations of the PPT property:

- A CDUC quantum channel Φ⁽¹⁾_{A,C} is PPT iff C is positive semidefinite (C ≥ 0).
 A DUC quantum channel Φ⁽²⁾_{A,B} is PPT iff |B_{ij}|² ≤ A_{ij}A_{ji} for all i, j ∈ [d].
 A DOC quantum channel Φ⁽³⁾_{A,B,C} is PPT iff C is positive semidefinite (C ≥ 0) and |B_{ij}|² ≤ $A_{ii}A_{ji}$ for all $i, j \in [d]$.

To characterize the entanglement breaking property. we recall the following two important definitions from [JM19] and [NS21]:

• A pair of matrices $A, B \in \mathcal{M}_d(\mathbb{C})$ having the same diagonal is said to be *pairwise completely* positive (PCP) if there exist matrices $V, W \in \mathcal{M}_{d \times k}(\mathbb{C})$, for some $k \in \mathbb{N}$, such that the following decomposition holds:

$$A = (V \odot \overline{V})(W \odot \overline{W})^*, \qquad B = (V \odot W)(V \odot W)^*.$$

• A triple of matrices $A, B, C \in \mathcal{M}_d(\mathbb{C})$ having the same diagonal is said to be triplewise completely positive (TCP) if there exist matrices $V, W \in \mathcal{M}_{d \times k}(\mathbb{C})$, for some $k \in \mathbb{N}$, such that the following decomposition holds:

$$A = (V \odot \overline{V})(W \odot \overline{W})^*, \qquad B = (V \odot W)(V \odot W)^*, \qquad C = (V \odot \overline{W})(V \odot \overline{W})^*.$$

Proposition 5.2 ([SN21]). For any triple of matrices $A, B, C \in \mathcal{M}_d(\mathbb{C})$ having the same diagonal, we have

- A CDUC quantum channel $\Phi_{A,C}^{(1)}$ is EB iff the pair (A,C) is pairwise completely positive (PCP).
- A DUC quantum channel $\Phi_{A,B}^{(2)}$ is EB iff the pair (A,B) is pairwise completely positive (PCP).
- A DOC quantum channel $\Phi_{A,B,C}^{(3)}$ is EB iff the triple (A,B,C) is triplewise completely positive (TCP).

Importantly, for the diagonal covariant channels discussed in this section, the EB property is not equivalent to the PPT property: there exist PPT channels that are not entanglement breaking. Such examples can be found in a simple way starting from the PPT entangled states of size 3×3 in [KKL11, KO12] or from nonnegative and positive semidefinite matrices of size $d \ge 5$ which are not completely positive [BSM03, TAQ⁺18, Yu16, SN21].



FIGURE 7. The three matrices A, B, C associated to a linear map Φ through its Choi matrix $J(\Phi)$.

Let us now provide the formulas for the diagonal twirlings of a quantum channels. We would like to point out again that our naming convention here is different from the one in [SN21], in the sense that the terms DUC and CDUC are swapped.

Proposition 5.3 ([NS21, Section 6]). Let $\Phi : \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$ be a linear map. Then, its twirlings with respect to the diagonal groups \mathcal{DU}_d and \mathcal{DO}_d are given by

$$\begin{split} \mathcal{T}_{\mathsf{CDUC}} \Phi &:= \mathcal{T}_{\mathcal{D}\mathcal{U}_d, \overline{\mathcal{D}\mathcal{U}}_d} \Phi = \Phi_{A(\Phi), C(\Phi)}^{(1)} \\ \mathcal{T}_{\mathsf{DUC}} \Phi &:= \mathcal{T}_{\mathcal{D}\mathcal{U}_d, \mathcal{D}\mathcal{U}_d} \Phi = \Phi_{A(\Phi), B(\Phi)}^{(2)} \\ \mathcal{T}_{\mathsf{DOC}} \Phi &:= \mathcal{T}_{\mathcal{D}\mathcal{O}_d, \mathcal{D}\mathcal{O}_d} \Phi = \Phi_{A(\Phi), B(\Phi), C(\Phi)}^{(3)}, \end{split}$$

where the matrices A, B, C are defined by (See Fig. 7 for a graphical representation):

$$A(\Phi)_{ij} = \langle ij|J(\Phi)|ij\rangle, \quad B(\Phi)_{ij} = \langle ii|J(\Phi)|jj\rangle, \quad and \quad C(\Phi)_{ij} = \langle ij|J(\Phi)|ji\rangle \qquad \forall i, j \in [d].$$

Let us now consider random diagonal covariant channels. To this end, let $\Phi_V : \mathcal{M}_d(\mathbb{C}) \to \mathcal{M}_d(\mathbb{C})$ be a random Stinespring channel with environment dimension s and define the following classes of random symmetric channels:

• random CDUC channels

$$\Phi_{\mathsf{CDUC}} := \mathcal{T}_{\mathsf{CDUC}}(\Phi_V) \sim \mu_{d,s}^{\mathsf{CDUC}}.$$

• random DUC channels

$$\Phi_{\mathsf{DUC}} := \mathcal{T}_{\mathsf{DUC}}(\Phi_V) \sim \mu_{d,s}^{\mathsf{DUC}}.$$

• random DOC channels

$$\Phi_{\mathsf{DOC}} := \mathcal{T}_{\mathsf{DOC}}(\Phi_V) \sim \mu_{d,s}^{\mathsf{DOC}}.$$

Since the PPT and EB properties of the random channels we consider here are determined by the associated A, B, C matrices, we shall study the corresponding random matrices, derived from the Haar-distributed random isometry $V : \mathbb{C}^d \to \mathbb{C}^d \otimes \mathbb{C}^s$:

$$A = \sum_{k=1}^{s} V^{(k)} \odot \overline{V^{(k)}}, \quad B = \sum_{k=1}^{s} |\operatorname{diag} V^{(k)}\rangle \langle \operatorname{diag} V^{(k)}|, \quad C = \sum_{k=1}^{s} V^{(k)} \odot (V^{(k)})^*, \tag{16}$$

where V is represented in the block matrix form $V = \begin{pmatrix} V^{(1)} \\ \vdots \\ V^{(s)} \end{pmatrix}$ for $V^{(k)} \in \mathcal{M}_d(\mathbb{C})$. Moreover, we can

draw the diagrams of A, B, C as in Fig. 8. Note that these $A, B, C \in \mathcal{M}_d(\mathbb{C})$ and their distribution depends on the integer parameter $s \geq 1$.



FIGURE 8. The three random matrix parameters A, B, and C for random CDUC, DUC, and DOC channels, defined via a Haar-distributed random isometry $V : \mathbb{C}^d \to \mathbb{C}^d \otimes \mathbb{C}^s$.

Remark 5.4. It is useful to connect the random parameters A, B, C discussed in this section to the scalars $\lambda_{1,2,3}^{(d,s)}$ from Sections 3 and 4; graphically, compare Figs. 3 and 6 with Fig. 8. We have the following relations:

$$\lambda_1(\Phi) = \sum_{i,j=1}^d \langle i|B(\Phi)|j\rangle = \operatorname{Tr}(J_d B(\Phi)),$$

$$\lambda_2(\Phi) = \sum_{i,j=1}^d \langle i|C(\Phi)|j\rangle = \operatorname{Tr}(J_d C(\Phi)),$$

$$\lambda_3(\Phi) = \operatorname{Tr} A(\Phi) = \operatorname{Tr} B(\Phi) = \operatorname{Tr} C(\Phi),$$
(17)

where $J_d \in \mathcal{M}_d(\mathbb{C})$ is the matrix in which all entries are equal to 1.

As in the previous two sections, we study now the asymptotic behavior of three random matrix parameters A, B, C in the regime $d \to \infty$. The following results will be used to establish, later in this section, the PPT and the EB properties of random CDUC, DUC, and DOC channels. Since both the entrywise and the spectral distributions of the matrices A, B, C are needed (see e.g. Proposition 5.1), the analysis in this section is more involved. We defer the technical proofs of the following three results, Propositions 5.5, 5.8 and 5.9, to Appendix A.3.

We start by giving the large-*d* limit spectrum of the self-adjoint matrix $C \in \mathcal{M}_d^{sa}(\mathbb{C})$, whose positive semi-definiteness is needed for PPT property of DOC channels. We recall that the *semicircular* distribution with average *m* and variance σ^2 is given by

$$\mathrm{dSC}_{m,\sigma} = \frac{\sqrt{4\sigma^2 - (x-m)^2}}{2\pi\sigma} \mathbf{1}_{[-2\sigma+m,2\sigma+m]}(x) \,\mathrm{d}x.$$

Proposition 5.5. The asymptotic spectral behavior of the random matrix C, as $d \to \infty$, is given as follows:

(1) In the regime $s/d \rightarrow 0$,

- in probability, the empirical spectral distribution of $(ds)^{1/2}C$ converges, in moment, to a semicircular distribution with mean 0 and variance 1,
- almost surely, we have the following convergence of the first three moments of $(ds)^{1/2}C$:

$$\frac{1}{d}\operatorname{Tr}\left((ds)^{1/2}C\right) \to 0, \quad \frac{1}{d}\operatorname{Tr}\left(dsC^{2}\right) \to 1, \quad \frac{1}{d}\operatorname{Tr}\left((ds)^{3/2}C^{3}\right) \to 0$$

- (2) In the regime $s \sim cd$ for a constant c > 0, the empirical spectral distribution of sC converges, in moments, to a semicircular distribution with mean c and variance c almost surely.
- (3) In the regime $s/d \to \infty$, the empirical spectral distribution of dC converges, in moments, to the probability mass δ_1 almost surely.

Proof. See Appendix A.3.

21

Under some slightly stronger hypothesis, the convergence in moments in item (3) above can be upgraded to a convergence in operator norm.

Proposition 5.6. When $s \gtrsim d^{1+t}$ for some constant t > 0, $||dC - I_d||_{\infty} \to 0$ almost surely as $d \to \infty$. In particular, the minimum eigenvalue $\lambda_{\min}(dC)$ of dC converges to 1 almost surely as $d \to \infty$.

Proof. See Appendix A.3.

Remark 5.7. Although it is not needed in our work, we can show the convergence of matrix B in various regimes, using the same method. For example, in the regime $s \sim cd$ and $d \rightarrow \infty$, the the empirical spectral distribution of sB converges, almost surely, to a Marčenko-Pastur distribution of parameter c: the distribution MP_c given by

$$dMP_{c} = \max(1-c,0)\delta_{0} + \frac{\sqrt{(b-x)(x-a)}}{2\pi x} \mathbf{1}_{[a,b]}(x) dx,$$

with $a = (1 - \sqrt{c})^2$ and $b = (1 + \sqrt{c})^2$.

We now turn to the distribution of the elements of the matrices A, B, C.

Proposition 5.8. When s is fixed, we have as $d \to \infty$,

- (1) the law of dsA_{ij} converges, in moments, to the Gamma distribution $\Gamma(s, 1)$,
- (2) both the laws of $|dsB_{ij}|^2$ and $|dsC_{ij}|^2$ converge, in moments, to that of the product of two independent random variables $X \sim \Gamma(s, 1)$ and $Y \sim \mathsf{Exp}(1)$,
- (3) for any fixed $N \leq d/2$ and pairwise distinct indices $i_1, \ldots, i_N, j_1, \ldots, j_N \in [d]$, the 3N random variables

$$dsA_{i_1j_1}, dsA_{j_1i_1}, |dsB_{i_1j_1}|^2, dsA_{i_2j_2}, dsA_{j_2i_2}, |dsB_{i_2j_2}|^2, \dots, dsA_{i_Nj_N}, dsA_{j_Ni_N}, |dsB_{i_Nj_N}|^2$$

are asymptotically independent.

Proof. See Appendix A.3.

Proposition 5.9. When $s \gtrsim d^t$ for some constant t > 0, we have the following almost sure convergences as $d \to \infty$:

(1) $\max_{\substack{i,j \\ i \neq j}} |dA_{ij} - 1| \to 0 \text{ (in particular, } \max_{\substack{i,j \\ i,j \\ i \neq j}} dA_{ij} \to 1 \text{ and } \min_{\substack{i,j \\ i,j \\ i \neq j}} dA_{ij} \to 1).$ (2) $\max_{\substack{i \neq j \\ i \neq j}} d^{1+t'/2} |B_{ij}| \to 0 \text{ whenever } t' < t.$ (3) $\max_{\substack{i \neq j \\ i \neq j}} d^{1+t'/2} |C_{ij}| \to 0 \text{ whenever } t' < t.$

Proof. See Appendix A.3.

The previous three results allow us to establish the PPT property for diagonal unitary and orthogonal covariant random channels. Note the qualitatively different behaviors for the DUC case versus the CDUC and DOC symmetries.

Theorem 5.10. Let $\Phi_{\mathsf{DUC}} \sim \mu_{d,s}^{\mathsf{DUC}}$ be a random DUC channel. Then:

(1) In the regime s is fixed and $d \to \infty$, we have

$$\lim_{d\to\infty} \mathbb{P}(\Phi_{\mathsf{DUC}} \text{ is } PPT) = 0.$$

(2) In the regime $s \gtrsim d^t$ for some constant t > 0, Φ_{DUC} is almost surely PPT as $d \to \infty$. Let $\Phi_{\mathsf{CDUC}} \sim \mu_{d,s}^{\mathsf{CDUC}}$, and $\Phi_{\mathsf{DOC}} \sim \mu_{d,s}^{\mathsf{DOC}}$ be, respectively, random CDUC and DOC channels. Then:

(3) In the regime $d \to \infty$ and $s/d \to 0$, almost surely, neither Φ_{CDUC} nor Φ_{DOC} are PPT.

- (4) Consider the regime $s \sim cd$. If c < 4, then almost surely as $d \to \infty$, neither Φ_{CDUC} nor Φ_{DOC} are PPT. If c > 4, then almost surely as $d \to \infty$, both Φ_{CDUC} and Φ_{DOC} have Choi matrices whose partial transpositions converge, in moments, to a probability measure supported on the positive real line.
- (5) In the regime $s \gtrsim d^{1+t}$ for some constant t > 0, both Φ_{CDUC} and Φ_{DOC} are almost surely PPT as $d \to \infty$.

Proof. Let us start with the first and second points. By Proposition 5.1, we know that

$$\Phi_{\mathsf{DUC}}$$
 is PPT $\iff A_{ij}A_{ji} \ge |B_{ij}|^2 \quad \forall i \neq j.$

If s is fixed, then by Proposition 5.8, the limit probability can be estimated, for any fixed $N \ge 1$, as follows:

$$\begin{split} \limsup_{d \to \infty} \mathbb{P}(\Phi_{\mathsf{DUC}} \text{ is PPT }) &\leq \limsup_{d \to \infty} \mathbb{P}(A_{2n-1,2n} A_{2n,2n-1} \geq |B_{2n-1,2n}|^2 \; \forall \, n = 1, \dots, N) \\ &= \prod_{n=1}^{N} \lim_{d \to \infty} \mathbb{P}((ds A_{2n-1,2n}) \cdot (ds A_{2n,2n-1}) \geq |ds B_{2n-1,2n}|^2) \\ &= (\mathbb{P}(X_1 X_2 \geq X_3 Y))^N \,, \end{split}$$

where X_1, X_2, X_3, Y are four independent random variables such that $X_1, X_2, X_3 \sim \Gamma(s, 1)$ and $Y \sim \mathsf{Exp}(1)$. Moreover, the last equation can be more explicitly written as an integral

$$\mathbb{P}(X_1X_2 \ge X_3Y) = \int_{\mathbb{R}^4} \mathbf{1}_{\left\{\substack{x_1x_2 \ge x_3y, \\ x_1, x_2, x_3, y \ge 0}\right\}} \frac{(x_1x_2x_3)^{s-1}}{((s-1)!)^3} e^{-(x_1+x_2+x_3+y)} dx_1 \, dx_2 \, dx_3 \, dy < 1.$$

Therefore, we have (1) by taking $N \to \infty$. When $s \gtrsim d^t$ for some t > 0, we can show that $d^2 \min_{i \neq j} (A_{ij}A_{ji} - |B_{ij}|^2) \to 1$, almost surely, from the inequality

$$\left(\min_{i,j} dA_{ij}\right)^2 - \left(\max_{i \neq j} d|B_{ij}|\right)^2 \le d^2 \min_{i \neq j} (A_{ij}A_{ji} - |B_{ij}|^2) \le \left(\max_{i,j} dA_{ij}\right)^2$$

and Proposition 5.9, and hence (2) is shown.

Now let us focus on (3) - (5), namely the PPT behaviors of CDUC and DOC channels. Note that for both two channels, their PPT property requires the condition $C \ge 0$, and in particular, this implies that

 $\operatorname{Tr} C \operatorname{Tr} C^3 \ge (\operatorname{Tr} C^2)^2$

by Cauchy-Schwarz inequality. However, when $s/d \to 0$ as $d \to \infty$, we have by Proposition 5.5 (1)

$$s^{2} \left(\operatorname{Tr} C \operatorname{Tr} C^{3} - (\operatorname{Tr} C^{2})^{2} \right) = \frac{1}{d} \operatorname{Tr} \left((ds)^{1/2} C \right) \cdot \frac{1}{d} \operatorname{Tr} \left((ds)^{3/2} C^{3} \right) - \left(\frac{1}{d} \operatorname{Tr} \left(ds C^{2} \right) \right)^{2} \to -1 < 0$$

almost surely, which proves (3). On the other hand, when $s \gtrsim d$ we again have

 $d^2 \min_{i \neq j} (A_{ij} A_{ji} - |B_{ij}|^2) \to 1$ almost surely.

Therefore, it is enough to consider the condition $C \ge 0$ for PPT properties by Proposition 5.1. Now Proposition 5.5 (2) and (3), combined with Proposition 5.9, and 5.6, imply the assertions (4) and (5), respectively.

Remark 5.11. Note that in the case of CDUC and DOC channels, in the regime $d \to \infty$, $s \sim cd$ where c > 4, we cannot prove exactly that Φ_{CDUC} and Φ_{DOC} are asymptotically PPT. This is due to the fact that the absence of negative support of the limiting measure of the partial transpositions of the respective Choi matrices does not exclude the absence of negative outliers. To reach the desired conclusion, one may show the strong convergence [HT05] of the random matrices to their corresponding limit. For the partial transpositions of Wishart random matrices, Aubrun [Aub12] computed very large moments of the random matrices in order to show the norm convergence, as we do in the proof of Proposition 5.6 for the last assertion in Theorem 5.10.

The next result describes the asymptotic behavior of *computable cross norm* or *realignment criterion* [CW03, Rud05] for (the Choi matrix of) random DOC channels. The realignment criterion is known to be, in general, *incomparable* to the PPT criterion. Its form for diagonal invariant bipartite matrices was derived in [SN21, Lemma 2.12], namely

$$\Phi_{A,B,C}^{(3)} \text{ is entanglement breaking } \implies \sum_{i,j=1}^d A_{ij} - \|A\|_1 \ge \sum_{i \neq j=1}^d \max(|B_{ij}|, |C_{ij}|)$$

The result below can be interpreted as the fact the realignment criterion is *weaker* than the PPT criterion for random DOC channels, from a random perspective. Note that the same phenomenon was proven for random bipartite quantum states in [AN12].

Proposition 5.12. In every regime $s \gtrsim d^t$ for some constants t > 0 and $d \to \infty$, random DOC channels satisfy the realignment criterion almost surely,

$$\sum_{i,j=1}^{d} A_{ij} - \|A\|_1 \ge \sum_{i \neq j=1}^{d} \max(|B_{ij}|, |C_{ij}|).$$

The same conclusion holds for random CDUC and DUC channels.

Proof. This follows from the three almost sure limits:

$$\frac{1}{d}\sum_{i,j}A_{ij} \to 1, \quad \frac{1}{d}\|A\|_1 \to 0, \quad \text{and} \quad \frac{1}{d}\sum_{i\neq j}\max(|B_{ij}|, |C_{ij}|) \to 0.$$

The limits above follow from Proposition 5.9 and the following three inequalities:

$$\begin{split} \min_{i,j} dA_{ij} &\leq \frac{1}{d} \sum_{i,j} A_{ij} \leq \max_{i,j} dA_{ij}, \\ \frac{1}{d} \|A\|_{1} \leq \frac{1}{d} \|I_{d}\|_{2} \|A\|_{2} = \frac{1}{\sqrt{d}} \left(\sum_{i,j} A_{ij}^{2} \right)^{1/2} \leq \frac{1}{\sqrt{d}} (\max_{i,j} dA_{ij}), \\ \frac{1}{d} \sum_{i \neq j} \max(|B_{ij}|, |C_{ij}|) \leq \frac{1}{d} \sum_{i \neq j} (|B_{ij}| + |C_{ij}|) \leq (d-1) (\max_{i \neq j} |B_{ij}| + \max_{i \neq j} |C_{ij}|). \end{split}$$

We now turn to the entanglement breaking (EB) property for random DOC channels.

Theorem 5.13. Under the regime $s \gtrsim d^{2+t}$ for some constants t > 0 and $d \to \infty$, the random DOC channel Φ_{DOC} is almost surely entanglement breaking.

Proof. Let us denote by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}_{i,j\in[d]} := a|i\rangle\langle i| + b|i\rangle\langle j| + c|j\rangle\langle i| + |j\rangle\langle j| \in \mathcal{M}_d(\mathbb{C})$$
(18)

for simplicity, and let us decompose $(A, B, C) = \sum_{1 \le i < j \le d} (A^{(ij)}, B^{(ij)}, C^{(ij)})$ where

$$A^{(ij)} = \begin{pmatrix} \frac{1}{d-1}A_{ii} & A_{ij} \\ A_{ji} & \frac{1}{d-1}A_{jj} \end{pmatrix}_{i,j\in[d]}, \quad B^{(ij)} = \begin{pmatrix} \frac{1}{d-1}A_{ii} & B_{ij} \\ B_{ji} & \frac{1}{d-1}A_{jj} \end{pmatrix}_{i,j\in[d]}, \quad C^{(ij)} = \begin{pmatrix} \frac{1}{d-1}A_{ii} & C_{ij} \\ C_{ji} & \frac{1}{d-1}A_{jj} \end{pmatrix}_{i,j\in[d]}.$$

Let $\Phi_{ij} := \Phi_{A_{ij},B_{ij},C_{ij}}^{(3)}$ be the DOC map associated to the matrix triple $(A^{(ij)}, B^{(ij)}, C^{(ij)})$. Note that in the regime $s \gtrsim d^{2+t}$ and $d \to \infty$, Proposition 5.9 implies

$$\min_{i,j} dA_{ij} \to 1, \quad \max_{i < j} d^2 |B_{ij}| \to 0, \quad \max_{i < j} d^2 |C_{ij}| \to 0 \text{ almost surely.}$$

Therefore, we obtain almost surely,

$$\frac{1}{(d-1)^2} A_{ii} A_{jj} \ge |B_{ij}|^2, \quad \frac{1}{(d-1)^2} A_{ii} A_{jj} \ge |C_{ij}|^2, \quad A_{ij} A_{ji} \ge |B_{ij}|^2, \quad \forall i < j.$$

In particular, this implies that the map Φ_{ij} is PPT for all i < j according to Proposition 5.1. However, since each triple $(A^{(ij)}, B^{(ij)}, C^{(ij)})$ is supported on a 2-dimensional subspace, this implies that Φ_{ij} is EB for all i < j, and hence $\Phi_{\mathsf{DOC}} = \sum_{i < j} \Phi_{ij}$ is EB.

Note that the proof does not work anymore for the regime $s \sim cd^2$. For example, when c = 1, we can show that d^2B_{ij} converges in moment to the exponential distribution $\mathsf{Exp}(1)$ using Lemma A.13, and thus

$$\mathbb{P}\left(\frac{1}{(d-1)^2}A_{ii}A_{jj} \ge |B_{ij}|^2\right) \to \mathbb{P}(Z_{\mathsf{Exp}(1)} \le 1) < 1$$

as $d \to \infty$.

Since channels having diagonal (conjugate) unitary covariance can be obtained from DOC channels by twirling, we have the following simple consequence of Theorem 5.13.

Corollary 5.14. Under the same hypothesis as above, the same conclusion holds for random DUC and CDUC channels.

6. The PPT^2 conjecture holds for random DOC channels

Originally motivated by the theory of quantum repeaters, the PPT^2 conjecture is now a central open question in quantum information theory, with recent ramifications into operator theory and operator algebras. We recall below its statement.

Conjecture 6.1 ([Chr12, CMHW18]). The composition of two arbitrary PPT quantum channels is entanglement breaking.

The conjecture holds trivially in dimension d = 2, and it has been shown to hold also in dimension d = 3 [CMHW18, CYT19]. A plethora of papers establish the conjecture in several restricted scenarios, or with weaker conclusions (such as the need to compose a PPT map with itself several times) [LG15, KMP17, RJP18, HRF20, GKS20, JY20]. Importantly for us, Collins, Yin, and Zhong showed [CYZ18] that two *independent* random quantum channels that are PPT satisfy the conjecture.

In the case of quantum channels with diagonal unitary symmetry, is has been shown in [SN22b] that any two (C)DUC channels satisfy the PPT² conjecture. The question whether the same holds for DOC channels has been left open [SN22b, Conjecture 5.1]. In this section, we contribute to this conjecture in two perspectives: *deterministic* and *random*. First of all, we show that one of the conditions for two PPT (C)DUC channels in [SN22b] can be relaxed to DOC. Furthermore, we answer in the affirmative this question for *random* DOC channels, under much weaker conditions. We refer to Corollary 6.4 and Theorem 6.5 for the details.

The results rely on the following condition for the composition of two matrix triples to be TCP. Recall from [SN21, Lemmas 9.3 and 9.7] that the composition of two DOC channels is DOC:

$$\Phi_{A,B,C}^{(3)} \circ \Phi_{D,E,F}^{(3)} = \Phi_{\mathcal{A},\mathcal{B},\mathcal{C}}^{(3)}$$

Proposition 6.2. If (A, B, C) and (D, E, F) are triples of matrices such that the corresponding DOC linear maps $\Phi_{A,B,C}^{(3)}$, $\Phi_{D,E,F}^{(3)}$ are completely positive, and, moreover, for all $i \neq j \in [d]$,

$$\left(\frac{A_{ii}D_{ii}}{d-1} + A_{ij}D_{ji}\right)\left(\frac{A_{jj}D_{jj}}{d-1} + A_{ji}D_{ij}\right) \ge \max(|B_{ij}E_{ij} + C_{ij}\overline{F_{ij}}|^2, |B_{ij}F_{ij} + C_{ij}\overline{E_{ij}}|^2), \quad (19)$$

$$(AD)_{ij}(AD)_{ji} \ge \max(|B_{ij}E_{ij} + C_{ij}\overline{F_{ij}}|^2, |B_{ij}F_{ij} + C_{ij}\overline{E_{ij}}|^2).$$
 (20)

Then, their composition $(A, B, C) \circ (D, E, F)$ is TCP.

Proof. The proof is a slight modification of the proof of [SN22b, Theorem 4.5] in straightforward way. Using the notation in Eq. (18), we can decompose $(\mathcal{A}, \mathcal{B}, \mathcal{C}) = \sum_{1 \leq i < j \leq d} (\mathcal{A}^{(ij)}, \mathcal{B}^{(ij)}, \mathcal{C}^{(ij)})$ where

$$\mathcal{A}^{(ij)} = \begin{pmatrix} \frac{1}{d-1}A_{ii}D_{ii} + A_{ij}D_{ji} & (AD)_{ij} \\ (AD)_{ji} & \frac{1}{d-1}A_{jj}D_{jj} + A_{ji}D_{ij} \end{pmatrix}_{i,j \in [d]}, \\ \mathcal{B}^{(ij)} = \begin{pmatrix} \frac{1}{d-1}A_{ii}D_{ii} + A_{ij}D_{ji} & B_{ij}E_{ij} + C_{ij}\overline{F_{ij}} \\ B_{ji}E_{ji} + C_{ji}\overline{F_{ji}} & \frac{1}{d-1}A_{jj}D_{jj} + A_{ji}D_{ij} \end{pmatrix}_{i,j \in [d]}, \\ \mathcal{C}^{(ij)} = \begin{pmatrix} \frac{1}{d-1}A_{ii}D_{ii} + A_{ij}D_{ji} & B_{ij}F_{ij} + C_{ij}\overline{E_{ij}} \\ B_{ji}F_{ji} + C_{ji}\overline{E_{ji}} & \frac{1}{d-1}A_{jj}D_{jj} + A_{ji}D_{ij} \end{pmatrix}_{i,j \in [d]},$$

since $E^T = \overline{E}$ and $F^T = \overline{F}$. The two conditions (19) and (20), combined with Proposition 5.1, guarantee that each triple $(\mathcal{A}^{(ij)}, \mathcal{B}^{(ij)}, \mathcal{C}^{(ij)})$ is a "PPT" triple in 2-dimensional space, and hence TCP. This shows that $(\mathcal{A}, \mathcal{B}, \mathcal{C})$ is TCP.

Remark 6.3. We actually show something stronger: the composition has factor width 2, see [SN22b, Definition 3.2]:

$$(A, B, C) \circ (D, E, F) \in \mathsf{TCP}_d^{(2)}$$

Note that the conditions Eqs. (19) and (20) are already weak enough to improve the main result of [SN22b, Theorem 4.5]. Indeed, suppose that $\Phi_{A,B,C}^{(3)}$ and $\Phi_{D,E,F}^{(3)}$ are PPT, and that one of B, C, E, or F is diagonal (corresponding to either $\Phi_{A,B,C}^{(3)}$ or $\Phi_{D,E,F}^{(3)}$ being DUC / CDUC). Then by Proposition 5.1 we have the entrywise inequalities

$$A_{ij}A_{ji} \ge \max(|B_{ij}|^2, |C_{ij}|^2), \quad D_{ij}D_{ji} \ge \max(|E_{ij}|^2, |F_{ij}|^2), \quad i \ne j \in [d],$$

which yields Eq. (19) since any of the diagonal conditions imply that

 $\max(|B_{ij}E_{ij} + C_{ij}\overline{F_{ij}}|^2, |B_{ij}F_{ij} + C_{ij}\overline{E_{ij}}|^2) \le \max(|B_{ij}E_{ij}|^2, |C_{ij}\overline{F_{ij}}|^2, |B_{ij}F_{ij}|^2, |C_{ij}\overline{E_{ij}}|^2).$

Furthermore, Eq. (20) holds since $\Phi_{\mathcal{A},\mathcal{B},\mathcal{C}}^{(3)} = \Phi_{A,B,C}^{(3)} \circ \Phi_{D,E,F}^{(3)}$ is PPT. Consequently, $(\mathcal{A},\mathcal{B},\mathcal{C}) \in \mathsf{TCP}_d^{(2)}$, and hence $\Phi_{\mathcal{A},\mathcal{B},\mathcal{C}}^{(3)}$ is entanglement breaking. We summarize the argument in the following corollary.

Corollary 6.4. If $\Phi_{A,B,C}^{(3)}$ and $\Phi_{D,E}^{(i)}$ $(i \in \{1,2\})$ are both PPT maps, then both compositions $\Phi_{A,B,C}^{(3)} \circ \Phi_{D,E}^{(i)}$ and $\Phi_{D,E}^{(i)} \circ \Phi_{A,B,C}^{(3)}$ are entanglement breaking.

The conditions in Proposition 5.1 also allow us to obtain another main result for random DOC channels.

Theorem 6.5. Consider two random DOC channels $\Phi_i \sim \mu_{d,s_i}^{\text{DOC}}$, i = 1, 2, in the asymptotic regime where $d \to \infty$ and $s_i \gtrsim d^{t_i}$ for some constants $t_i > 0$. Then almost surely as $d \to \infty$, the composition $\Phi_1 \circ \Phi_2$ is entanglement breaking.

In particular, in the asymptotic regime above, any two random DOC channels satisfy the PPT^2 conjecture.

Proof. Let $\Phi_1 = \Phi_{A,B,C}^{(3)}$ and $\Phi_2 = \Phi_{D,E,F}^{(3)}$. The result follows from Proposition 6.2 and Proposition 5.9: almost surely, we have

$$\lim_{d \to \infty} d^4 \min_{i \neq j} \left(\frac{A_{ii}D_{ii}}{d-1} + A_{ij}D_{ji} \right) \left(\frac{A_{jj}D_{jj}}{d-1} + A_{ji}D_{ij} \right) = 1,$$
$$\liminf_{d \to \infty} d^4 \min_{i \neq j} (AD)_{ij} (AD)_{ji} \ge \liminf_{d \to \infty} \min_{i \neq j} d^4 (A_{ii}D_{ij}A_{jj}D_{ji} + A_{ij}D_{jj}A_{ji}D_{ii}) = 2,$$

while $d^4 \max_{i \neq j} |B_{ij}E_{ij} + C_{ij}\overline{F_{ij}}|^2 \to 0$ and $d^4 \max_{i \neq j} |B_{ij}F_{ij} + C_{ij}\overline{E_{ij}}|^2 \to 0$. Therefore, the conditions (19) and (20) hold asymptotically.

Remark 6.6. We would like to emphasize that the hypothesis of Theorem 6.5 are very weak:

- The two channels need not be PPT; actually, from Theorem 5.10, if $s_i \sim c_i d^{t_i}$ with $0 < t_i < 1$ and $c_i > 0$, or $t_i = 1$ and $c_i < 4$, Φ_i is asymptotically not PPT.
- There is no assumption whatsoever about the correlations of the two random variables Φ_1 and Φ_2 : the two DOC channels can be independent, equal, complex conjugate to one another, or have any possible correlation.

The fact that we can show that the PPT² conjecture holds for random DOC quantum channels under such weak conditions (compared to [CYZ18] where the channels are assumed to be PPT and independent) is due to the presence of the \mathcal{DO}_d covariance.

Conflict of interest. The authors declare no competing interests.

Data availability statement. Data sharing is not applicable to this article as no new data were created or analyzed in this study.

Acknowledgments. Both authors were supported by the ANR project ESQuisses, grant number ANR-20-CE47-0014-01, and by the PHC program *Star* (Applications of random matrix theory and abstract harmonic analysis to quantum information theory). I.N. also received support from the ANR project STARS, grant number ANR-20-CE40-0008.

APPENDIX A. TECHNICAL LEMMAS AND PROOFS

We gather in this appendix some technical background material, as well as the proofs of some of the results in the main text.

A.1. General results about the combinatorics of permutations and Weingarten functions. Our main approach to understanding the entanglement properties of the different models of random covariant channels is the *moment method* in random matrix theory. Specifically, we shall apply graphical Weingarten calculus [Col03, CS06, CN10] to compute many kinds of integrals involving Haar random isometries for the precise analysis. In this section, we introduce basic notions on permutations, partitions, and the Weingarten formula. Moreover, we present several combinatorial lemmas which will be useful throughout the appendix for the actual computation later on. The main combinatorial objects that appear in the resolution of integrals with respect to the Haar measure on the unitary group are *permutations* and, in the limit of large dimensions, *non-crossing partitions*. We refer the reader to the excellent collection of lecture notes [NS06] for an introduction to the relation between these concepts, free probability theory, and random matrix theory.

We shall denote by S_p the symmetric group of order p, that is the permutations of the set

$$[p] := \{1, 2, \dots, p\}.$$

For a permutation $\alpha \in S_p$, we denote by $\#\alpha$ the number of cycles in its cycle decomposition, including singletons. We have #id = p, whereas for the full cycle permutation $\#\gamma = 1$, were

$$\gamma := (p \ p - 1 \cdots 3 \ 2 \ 1) \in S_p.$$

The length $|\alpha|$ of a permutation $\alpha \in S_p$ is the minimal number of transpositions that multiply to α . We have the important relation [NS06, Proposition 23.11]

$$\forall \alpha \in S_p, \qquad \#\alpha + |\alpha| = p.$$

Let us discuss now the Weingarten formula [Col03, CS06], the main technical tool that will allow us to compute various average quantities related to the Haar measure on the set of isometries. Below, the integral is taken either to the Haar measure on the set of isometries $V : \mathbb{C}^k \to \mathbb{C}^d$ for $k \leq d$. The isometry V can be thought of as the truncation (keeping the first k columns) of a Haar-distributed random unitary matrix $U \in \mathcal{U}_d$.

Theorem A.1. Let $k \leq d$ be positive integers and $\mathbf{i} = (i_1, \ldots, i_p)$, $\mathbf{i}' = (i'_1, \ldots, i'_p)$, $\mathbf{j} = (j_1, \ldots, j_p)$, $\mathbf{j}' = (j'_1, \ldots, j'_p)$ be p-tuples of positive integers with $\mathbf{i}, \mathbf{i}' \in [d]^p$ and $\mathbf{j}, \mathbf{j}' \in [k]^p$. Then

$$\int V_{i_1 j_1} \cdots V_{i_p j_p} \overline{V}_{i'_1 j'_1} \cdots \overline{V}_{i'_p j'_p} \, \mathrm{d}V = \sum_{\sigma, \tau \in S_p} \delta_{i_1 i'_{\sigma(1)}} \cdots \delta_{i_p i'_{\sigma(p)}} \delta_{j_1 j'_{\tau(1)}} \cdots \delta_{j_p j'_{\tau(p)}} \, \mathrm{Wg}_d(\tau \sigma^{-1}).$$
(21)

Above, $Wg_d(\sigma)$ is a rational function of d that depends only on the cycle type of the permutation $\sigma \in S_p$. The large d behavior of this function will be important for us:

$$\operatorname{Wg}_{d}(\sigma) = (1 + O(d^{-2})) \operatorname{M\ddot{o}b}(\sigma) d^{-p-|\sigma|},$$

where Möb is the *Möbius function*

$$\operatorname{M\"ob}(\sigma) := \prod_{c \in \sigma} (-1)^{|c|-1} \operatorname{Cat}_{|c|-1}$$

with the Catalan numbers

$$\operatorname{Cat}_n := \frac{1}{n+1} \binom{2n}{n}.$$

We shall mainly use the graphical formulation of this formula, introduced in [CN10]; we refer the reader to the review article [CN15] for a pedagogical presentation.

Lemma A.2 ([Stal1, Proposition 1.3.7],[CN10]). We have the following two identities for all $p \ge 1$: (1) $\sum_{\alpha \in S_p} n^{\#\alpha} = n(n+1)\cdots(n+p-1),$ (2) $\sum_{\alpha \in S_p} \operatorname{Wg}_n(\alpha) = (n(n+1)\cdots(n+p-1))^{-1}.$

The following combinatorial formula will be crucially used throughout the appendix.

Lemma A.3 ([Col03, Eq.(2.61)]). For any integer $n \ge 1$ and any polynomial F of degree < n, we have

$$\sum_{p=0}^{n} (-1)^{n-p} F(p) \binom{n}{p} = 0.$$

Lemma A.4. (1) Let us write the power series expansion

$$s^{1-p}(s+1)\cdots(s+p-1) = \sum_{k=0}^{\infty} a_{k,p}s^{-k}$$

in s^{-1} . Then for each k, the coefficient $a_{k,p}$ is a polynomial in p of degree 2k. Moreover, $a_{k,p} = 0$ for all $k \ge p$.

(2) If we write the power series expansion

$$\sum_{z \in \mathcal{P}_{1,2}(p)} s^{-|\alpha|} = \sum_{k=0}^{\infty} b_{k,p} s^{-k},$$

then $b_{0,p} \equiv 1$ and $b_{k,p} = \frac{p(p-1)\cdots(p-2k+1)}{2^k k!}$ for $k \geq 1$. (3) If we write the power series expansion

$$\sum_{\alpha \in NC_{1,2}(p)} s^{-|\alpha|} = \sum_{k=0}^{\infty} c_{k,p} s^{-k},$$

then
$$c_{0,p} \equiv 1$$
 and $c_{k,p} = {p \choose 2k} \operatorname{Cat}_k = \frac{p(p-1)\cdots(p-2k+1)}{(k+1)!k!}$ for $k \ge 1$.

Proof. (1) The second assertion is obvious since $s(s+1)\cdots(s+p-1)$ is a polynomial in s of degree p with zero constant. On the other hand, we can easily see that $a_{0,p} \equiv 1$, and $a_{k,p}$ satisfy the recurrence

$$a_{k,p} = (p-1)a_{k-1,p-1} + a_{k,p-1}, \ p \ge 1, k \ge 1,$$

from the relation

$$s(s+1)\cdots(s+p-1) = s^p \sum_{k=0}^{\infty} a_{k,p} s^{-k} = (s+p-1)s^{p-1} \sum_{k=0}^{\infty} a_{k,p-1} s^{-k}$$

Therefore, we have $a_{k,p} = \sum_{j=0}^{p-1} j a_{k-1,j} + a_{k,0} = \sum_{j=0}^{p-1} j a_{k-1,j}$. By induction hypothesis, $pa_{k-1,p}$ is a polynomial in p of degree 2k - 1, and hence the first assertion follows.

(2) We leave its proof to the reader, just by noting that $b_{k,p}$ is equal to the number of permutations consisting of exactly k transpositions (and p - 2k singletons) in their cycles.

(3) Similarly with (2), $c_{k,p}$ is the number of permutations consisting of k transpositions which are non-crossing. We can find this by first choosing 2k elements in [p] and then count the number of non-crossing pair partitions (= Cat_k) of these elements.

Lemma A.5. Given two permutations $\sigma_1, \sigma_2 \in S_p$, the map

$$S_p \ni \alpha \mapsto (|\sigma_1^{-1}\alpha| + |\alpha^{-1}\sigma_2|) \mod 2 \in \{0,1\}$$

is constant. In particular, for all $\alpha \in S_p$, there exists a nonnegative integer g such that

$$|\sigma_1^{-1}\alpha| + |\alpha^{-1}\sigma_2| = |\sigma_1^{-1}\sigma_2| + 2g.$$

Proof. The first assertion follows from the observations that every permutation α can be written as a product of transpositions and $|\sigma\tau| = |\sigma| \pm 1$ for every $\sigma \in S_p$ and a transposition τ [NS06, Lemma 23.10]. The second point then follows from the first point and the triangle inequality $|\sigma_1^{-1}\alpha| + |\alpha^{-1}\sigma_2| \ge |\sigma_1^{-1}\sigma_2|$.

We shall denote by $\mathcal{P}(p)$ the set of partitions of $[p] = \{1, 2, \ldots, p\}$. Importantly, we denote by NC(p) the set of non-crossing partitions of [p], that is partitions π for which there do not exist $i < j < k < l \in [p]$ such that $i \stackrel{\pi}{\sim} k$ and $j \stackrel{\pi}{\sim} l$ (in which case we say that the blocks of π containing i and k and, respectively, j and l are crossing). Let us denote by $S_{NC}(\sigma) := \{\alpha \in S_p : |\alpha| + |\alpha^{-1}\sigma| = |\sigma|\}$ the set of permutations which lie on a geodesic from id to σ . In what follows, we will provide

a complete description of the elements of $S_{NC}(\sigma)$ for arbitrary permutation $\sigma \in S_p$. First, we introduce the projection map $\pi : S_p \to \mathcal{P}(p)$ which naturally identifies each of the cycles of a permutation α as blocks of $\pi(\alpha)$. Then the following proposition states that we can interpret the elements of $S_{NC}(\gamma)$ as non-crossing partitions when γ is a full cycle.

Proposition A.6 ([NS06, Proposition 23.23]). Let $\gamma = (p \ p - 1 \cdots 1) \in S_p$. Then the restriction map $\pi \mid_{S_{NC}(\gamma)}$ of the projection map $\pi : S_p \to \mathcal{P}(p)$ defines a natural bijection between $S_{NC}(\gamma)$ and NC(p), which preserves the partial order structure. More explicitly,

(1) $\alpha \in S_{NC}(\gamma)$ if and only if $\pi(\alpha)$ forms a non-crossing partition and every cycle of α can be written in the form

 $(j_1 j_2 \cdots j_r), \quad j_1 > j_2 > \cdots > j_r,$

(2) if $\alpha \in S_{NC}(\gamma)$ and $\beta \in S_{NC}(\alpha)$, then $\beta \in S_{NC}(\gamma)$ and $\pi(\beta) \leq \pi(\alpha)$ as partitions [NS06, Definition 9.14].

Proposition A.7. Let $\sigma = c_1 \cdots c_k$ be the cycle decomposition of $\sigma \in S_p$. If $\alpha \in S_{NC}(\sigma)$, then α can be decomposed into disjoint permutations $\alpha_1 \cdots \alpha_k$ such that, for each $i = 1, \ldots, k$, $\alpha_i \leq c_i$ as partitions and $\alpha_i \in S_{NC}(c_i)$. Furthermore, if $\alpha' \in S_{NC}(\sigma)$ and $\alpha \equiv \alpha'$ as partitions, then $\alpha = \alpha'$ as permutations.

Proof. By re-indexing (i.e. conjugation $\delta\sigma\delta^{-1}$ for some $\delta\in S_p$), we may assume that

$$c_i = (p_i \ (p_i - 1) \ \cdots \ (p_{i-1} + 1)), \quad i = 1, \dots, k,$$

for some $1 = p_0 < p_1 < \cdots < p_k = p$. Then Proposition A.6 implies that $\sigma, \alpha \in S_{NC}(\gamma)$ and $\alpha \leq \sigma$ as partitions. In particular, each block $V_i = \{p_{i-1} + 1, \dots, p_i - 1, p_i\}$ is invariant under α , so the restriction $\alpha_i := \alpha \mid_{V_i} \in S(V_i)$ is well-defined. Now we can write $\alpha = \alpha_1 \cdots \alpha_k$ and $|\alpha| = \sum_{i=1}^k |\alpha_i|$. Moreover, the condition $\alpha \in S_{NC}(\sigma)$ implies that

$$|\sigma| = \sum_{i=1}^{\kappa} |\sigma_i| \le \sum_{i=1}^{\kappa} (|\alpha_i| + |\alpha_i^{-1}\sigma_i|) = \sum_i |\alpha_i| + \sum_i |\sigma_i^{-1}\alpha_i| = |\alpha| + |\alpha^{-1}\sigma| = |\sigma|,$$

and therefore, the equality condition forces that $\alpha_i \in S_{NC}(\sigma_i)$ for all *i*. Finally, the last statement follows from Proposition A.6(1): $\alpha, \alpha' \in S_{NC}(\gamma)$ and $\pi(\alpha) = \pi(\alpha')$ imply that $\alpha = \alpha'$.

Proposition A.7 implies that we have the natural bijection

$$S_{NC}(\sigma) \cong S_{NC}(c_1) \times \dots \times S_{NC}(c_k).$$
⁽²²⁾

Now let us denote by $S_{NC_{1,2}}(\sigma)$ the set of permutations in $S_{NC}(\sigma)$ having only cycles of size 1 or 2. The set can be defined equivalently by

$$\alpha \in S_{NC_{1,2}}(\sigma) \iff \alpha \in S_{NC}(\sigma) \text{ and } \alpha = \alpha^{-1} \iff \alpha, \alpha^{-1} \in S_{NC}(\sigma).$$
 (23)

since any cycle having size larger than 2 is distinct from its inverse, and since $\alpha, \alpha^{-1} \in S_{NC}(\sigma)$ implies $\alpha = \alpha^{-1}$ by Proposition A.7. Furthermore, note that the set $S_{NC_{1,2}}(\gamma)$ can be identified with $NC_{1,2}(p)$, the set of non-crossing partitions having only blocks of size 1 or 2, in a trivial way. Finally, if $\sigma = c_1 \cdots c_k$ is the cycle decomposition of σ , then the bijection Eq. (22) reduces to

$$S_{NC_{1,2}}(\sigma) \cong S_{NC_{1,2}}(c_1) \times \dots \times S_{NC_{1,2}}(c_k).$$

$$(24)$$

For two partitions $\pi_1, \pi_2 \in \mathcal{P}(p)$, the join $\pi_1 \vee \pi_2$ is defined as the supremum of π_1 and π_2 in $\mathcal{P}(p)$ [NS06, Definition 9.15]. Since the length function $|\alpha| = p - \# \alpha$ only depends on the corresponding partition $\pi(\alpha)$, we can define $|\alpha \vee \beta| := p - \#(\pi(\alpha) \vee \pi(\beta))$ for arbitrary permutations $\alpha, \beta \in S_p$. Note that $|\alpha \vee \beta| \ge \max(|\alpha|, |\beta|)$ in general since the join operation \vee does not increase the number of blocks as partitions. Moreover, if $|\alpha \vee \beta| = |\alpha| = |\beta|$, or equivalently $\#(\pi(\alpha) \vee \pi(\beta)) = \#(\pi(\alpha)) = \#(\pi(\beta))$, then we have $\pi(\alpha) = \pi(\beta)$. We now present several permutation inequalities that will be applied later to prove Proposition 5.5.

Lemma A.8. Let $\sigma \in S_p$ be a permutation.

(1) For all $\alpha, \beta \in S_p$, we have

$$|\alpha| + 2|\alpha\beta^{-1}| + |(\sigma^{-1}\alpha) \vee (\sigma^{-1}\beta^{-1})| \ge |\sigma|,$$

and the equality holds if and only if $\alpha = \beta \in S_{NC_{1,2}}(\sigma)$. Furthermore, whenever the equality does not hold, we have

$$|\alpha| + 2|\alpha\beta^{-1}| + |(\sigma^{-1}\alpha) \vee (\sigma^{-1}\beta^{-1})| \ge |\sigma| + 2.$$

(2) For all $\alpha, \beta \in S_p$, we have

$$|\alpha\beta^{-1}| + |(\sigma^{-1}\alpha) \vee (\sigma^{-1}\beta^{-1})| \ge \frac{1}{2}|\sigma^2|,$$

and the equality holds if and only if $\alpha = \beta \in S_{NC_{1,2}}(\sigma)$ and $\sigma^{-1}\alpha \in S_{NC}(\sigma^{-2})$. Furthermore, when $\sigma = \gamma = (p \ p - 1 \cdots 1)$, $\alpha \in NC_{1,2}(p)$ is 'maximally paired', i.e.,

$$p = even \implies \alpha \text{ consists of } \frac{p}{2} \text{ transpositions i.e., } \alpha \in NC_2(p),$$
 (25)

$$p = odd \implies \alpha \text{ consists of one singleton and } \frac{p-1}{2} \text{ transpositions.}$$
 (26)

Proof. (1) Since $|(\sigma^{-1}\alpha) \vee (\sigma^{-1}\beta^{-1})| \ge \max(|\sigma^{-1}\alpha|, |\sigma^{-1}\beta^{-1}|)$, we have $IHS \ge |\alpha| + |\sigma^{-1}\alpha| \ge |\sigma|$

LHS
$$\geq |\alpha| + |\sigma^{-1}\alpha| \geq |\sigma|,$$

LHS $\geq |\alpha| + |\alpha\beta^{-1}| + |\sigma^{-1}\beta^{-1}| \geq |\alpha^{-1}| + |\sigma^{-1}\alpha^{-1}| \geq |\sigma|.$

In particular, the equality conditions for both two inequalities imply that $\alpha = \beta$ and $\alpha, \alpha^{-1} \in S_{NC}(\sigma)$, and therefore $\alpha \in S_{NC_{1,2}}$ by Eq. (23). On the other hand, if the equality does not hold, then either $\alpha \neq \beta$ or $\alpha = \beta \notin S_{NC_{1,2}}$. For the former case, we have

LHS
$$\geq (|\alpha| + |\sigma^{-1}\alpha|) + 2|\alpha\beta^{-1}| \geq |\sigma| + 2,$$

and for the latter case, we again have

LHS
$$\ge \max(|\alpha| + |\sigma^{-1}\alpha|, |\alpha^{-1}| + |\sigma^{-1}\alpha^{-1}|) \ge |\sigma| + 2$$

by Lemma A.5.

(2) Similarly as above, we have

LHS
$$\geq |\alpha\beta^{-1}| + \max(|\sigma^{-1}\alpha|, |\sigma^{-1}\beta^{-1}|) \geq \frac{1}{2} \Big(|\alpha\beta^{-1}| + (|\sigma^{-1}\alpha| + |\alpha^{-1}\beta| + |\beta^{-1}\sigma^{-1}|) \Big) \geq \frac{1}{2} |\sigma^{2}|.$$

All the equalities above hold if and only if $\alpha = \beta$, $\sigma^{-1}\alpha \equiv \sigma^{-1}\alpha^{-1}$ as partitions, and $|\sigma^{-1}\alpha| + |\sigma^{-1}\alpha^{-1}| = |\sigma^{-2}|$. Note that the last condition implies that $\sigma^{-1}\alpha, \sigma^{-1}\alpha^{-1} \in S_{NC}(\sigma^{-2})$, and therefore, $\alpha = \alpha^{-1}$ by Proposition A.7. In particular, every cycle of α has size 1 or 2.

We claim that $\alpha \in S_{NC_{1,2}}(\sigma)$ in this case. Indeed, by considering the cycle decomposition of σ and applying Proposition A.7 for $\sigma^{-1}\alpha \in S_{NC}(\sigma^{-2})$, we may assume that $\sigma = \gamma = (p \ p - 1 \cdots 1)$. Then we have

$$|\gamma^{-1}\alpha| = |\gamma^{-1}\alpha^{-1}| = \frac{1}{2}|\gamma^{-2}| = \begin{cases} p/2 - 1 & \text{if } p \text{ is even,} \\ (p-1)/2 & \text{if } p \text{ is odd.} \end{cases}$$

Therefore, it suffices to show $|\alpha| = \begin{cases} p/2 & \text{if } p \text{ is even,} \\ (p-1)/2 & \text{if } p \text{ is odd} \end{cases}$ to conclude $\alpha \in S_{NC_{1,2}}(\gamma) \cong NC_{1,2}(p)$, which is equivalent to Eqs. (25) and (26). First, when p is even, we have

$$\gamma^{-2} = (1 \ 3 \cdots p - 1)(2 \ 4 \cdots p),$$

and in particular, the condition $\gamma^{-1}\alpha \in S_{NC}(\gamma^{-2})$ implies that every cycle $\gamma^{-1}\alpha$ consists of the elements having the same parity. This is impossible when there is any $i \in [p]$ such that $\alpha(i) = i$. Therefore, we have Eq. (25). On the other hand, when p is odd, we can write

$$\gamma^{-2} = (1 \ 3 \ \cdots \ p \ 2 \ 4 \cdots \ p - 1).$$

If Eq. (26) is not true, then we have at least three singletons inside α , and in particular, we can find two distinct integers i < i' with the same parity such that $\alpha(i) = i$ and $\alpha(i') = i'$. Then $\gamma^{-1}\alpha(i) = i + 1$, $\gamma^{-1}\alpha(i') = i' + 1$, and we can easily see that the pairs (i, i + 1) and (i', i' + 1) are crossing inside the full cycle γ^{-2} . Moreover, i, i + 1, i', i' + 1 cannot appear, in this order, in the same cycle of $\gamma^{-1}\alpha$ since any cycle inside $\gamma^{-1}\alpha \in S_{NC}(\gamma^{-2})$ must be of the form

$$(i_1 \cdots i_r j_1 \cdots j_{r'})$$

where $i_1 < \cdots < i_r$ are odd integers and $j_1 < \cdots < j_{r'}$ are even integers, according to Proposition A.6 (1). Consequently, we have a contradiction, and hence Eq. (26) holds.

Lemma A.9. Let $\gamma := (p \ p - 1 \ \cdots \ 1) \in S_p$ and $\tilde{\gamma} := (p \ p - 1 \ \cdots \ 1)(2p \ 2p - 1 \ \cdots \ p + 1) \in S_{2p}$. For p = 2, 3 we have

$$\alpha, \beta \in S_p, \ |\alpha| + |\alpha\beta^{-1}| > \frac{p}{2} \implies |\gamma^{-1}\alpha \lor \gamma^{-1}\beta^{-1}| + |\alpha\beta^{-1}| \ge \frac{p}{2} + \frac{1}{2}, \tag{27}$$

$$\alpha, \beta \in S_{2p}, \ |\alpha| + |\alpha\beta^{-1}| > p \implies |\tilde{\gamma}^{-1}\alpha \vee \tilde{\gamma}^{-1}\beta^{-1}| + |\alpha\beta^{-1}| \ge p.$$

$$(28)$$

Proof. Let us first consider the case p = 2. The proof of Eq. (27) is simple: $|\alpha| + |\alpha\beta^{-1}| > 1$ holds only if $\alpha = (12)$ and $\beta = id$, and $|\gamma^{-1}\alpha \vee \gamma^{-1}\beta^{-1}| + |\alpha\beta^{-1}| = 2$ in this case. Regarding Eq. (28), note that we always have $|\tilde{\gamma}^{-1}\alpha \vee \tilde{\gamma}^{-1}\beta^{-1}| \ge 1$: otherwise, $\alpha = \beta = \tilde{\gamma} = (12)(34)$, but this contradicts the condition $|\alpha| + |\alpha\beta^{-1}| > 2$. Now if we assume $\alpha = \beta$ and $|\tilde{\gamma}^{-1}\alpha \vee \tilde{\gamma}^{-1}\alpha^{-1}| = 1$, then $\max(|\tilde{\gamma}^{-1}\alpha|, |\tilde{\gamma}^{-1}\alpha^{-1}|) = 1$. Both the cases $|\tilde{\gamma}^{-1}\alpha| = 0$ and $|\tilde{\gamma}^{-1}\alpha| = 0$ imply that $\alpha = \beta = \tilde{\gamma}$ for which we already had a contradiction. If $|\tilde{\gamma}^{-1}\alpha| = |\tilde{\gamma}^{-1}\alpha^{-1}| = 1 = |\tilde{\gamma}^{-1}\alpha \vee \tilde{\gamma}^{-1}\alpha^{-1}|$, then we actually have $\tilde{\gamma}^{-1}\alpha = \tilde{\gamma}^{-1}\alpha^{-1}$ and $\alpha = \alpha^{-1}$. This again induces a contradiction since $|\alpha| + |\alpha\beta^{-1}| = |\alpha| \le 2$.

Now consider the case p = 3. By Lemma A.8 (2), we already know

$$\gamma^{-1} \alpha \vee \gamma^{-1} \beta^{-1} |+ |\alpha \beta^{-1}| \ge \frac{1}{2} |\gamma^2| = 1$$

for all $\alpha, \beta \in S_p$ and the equality holds only if $\alpha = \beta$ and $|\alpha| = 1$ (Eq. (26)) which does not satisfy $|\alpha| + |\alpha\beta^{-1}| > 3/2$. This implies Eq. (27). Similarly, we always have

$$|\tilde{\gamma}^{-1}\alpha \vee \tilde{\gamma}^{-1}\beta^{-1}| + |\alpha\beta^{-1}| \ge \frac{1}{2}|\tilde{\gamma}^2| = 2,$$

and the equality condition implies that $\alpha = \beta \in S_{NC_{1,2}}(\tilde{\gamma})$ and

$$|\alpha| = |\tilde{\gamma}| - |\tilde{\gamma}^{-1}\alpha| = |\tilde{\gamma}| - \frac{1}{2}|\tilde{\gamma}^2| = 2,$$

which contradicts the condition $|\alpha| + |\alpha\beta^{-1}| > 3$. This implies Eq. (28).

Remark A.10. By repeating the same arguments above, we can actually show that Eqs. (27) and (28) hold for all odd integers p. However, they are not true for even $p \ge 4$ in general: when $\alpha = \beta = (1234) \in S_4$,

$$|\gamma^{-1}\alpha \vee \gamma^{-1}\beta^{-1}| + |\alpha\beta^{-1}| = |(13)(24)| = 2 < \frac{5}{2}$$

and when $\alpha = \beta = (1\,2\,3\,4)(5\,6)(7\,8) \in S_8$,

$$|\tilde{\gamma}^{-1}\alpha \vee \tilde{\gamma}^{-1}\beta^{-1}| + |\alpha\beta^{-1}| = |(1\,3)(2\,4)(5\,7)| = 3 < 4.$$

A.2. Scalar parameters. This section contains results about the real parameters $\lambda_{1,2,3}^{(d,s)}$ which appear in the study of random (conjugate) unitary, orthogonal, and hyperoctahedral quantum channels from Sections 3 and 4.

The following moment formulae characterize the distributions of these random variables.

Lemma A.11. The moments of the random variables $\lambda_{1,2,3}^{(d,s)}$ are given by

$$\mathbb{E}\left[\left(\lambda_{1}^{(d,s)}\right)^{p}\right] = s(s+1)\cdots(s+p-1)\sum_{\beta\in S_{p}}d^{\#\beta}\mathrm{Wg}_{ds}(\beta)$$
$$\mathbb{E}\left[\left(\lambda_{2}^{(d,s)}\right)^{p}\right] = \sum_{\alpha,\beta\in S_{p}}s^{\#\alpha}d^{\#(\alpha\beta)}\mathrm{Wg}_{ds}(\alpha\beta^{-1})$$
$$\mathbb{E}\left[\left(\lambda_{3}^{(d,s)}\right)^{p}\right] = \sum_{\alpha,\beta\in S_{p}}s^{\#\alpha}d^{\#(\alpha\vee\beta)}\mathrm{Wg}_{ds}(\alpha\beta^{-1}).$$

Proof. To obtain the three formulas in the statement, we apply the graphical Weingarten formula for the diagram consisting of p (disconnected) copies of the diagram for $\lambda_{1,2,3}^{(d,s)}$ (see Figs. 3 and 6). In the case of $\lambda_1^{(d,s)}$, we obtain

$$\mathbb{E}\Big[\big(\lambda_1^{(d,s)}\big)^p\Big] = \sum_{\alpha,\beta\in S_p} d^{\#(\alpha^{-1}\beta)} s^{\#\alpha} \operatorname{Wg}_{ds}(\alpha^{-1}\beta) = \sum_{\sigma,\pi\in S_p} d^{\#\pi} s^{\#\sigma} \operatorname{Wg}_{ds}(\pi),$$

since there are precisely $\#(\alpha^{-1}\beta)$ loops of dimension d and $\#\alpha$ loops of dimension s. We use Lemma A.2 to compute explicitly the sum over σ .

Counting the loops in a similar fashion gives the formulas for $\lambda_2^{(d,s)}$. Regarding the integration of $(\lambda_3^{(d,s)})^p$, we can understand the diagram of $\lambda_3^{(d,s)}$ by joining the two separated lines representing the tensor component of dimension d as in Fig. 9. Note that after the graphical calculus in the diagram for $\alpha, \beta \in S_p$, the upper (external) and lower (internal) lines contributes $\# \alpha$ and $\# \beta$ loops of dimension d, respectively. However, we have to "join" them because we identify the the two kinds of loops (thick lines in the figure). Consequently, all the blocks in $\alpha \vee \beta$ contribute the same loop in the final diagram, and hence there are exactly $\#(\alpha \vee \beta)$ loops of dimension d. The term $s^{\#\alpha}$ appears similarly with the case $\lambda_{1,2}^{(d,s)}$, and this concludes the proof.



FIGURE 9. The way the permutations α and β connect the V and the V^{*} boxes in the diagram for $\mathbb{E}\left[\left(\lambda_3^{(d,s)}\right)^p\right]$. Note the thick vertical lines appearing in each diagram of $\lambda_3^{(d,s)}$ that are responsible for joining the otherwise disjoint α and β loops.

In the case s = 1, the random variable $\lambda_1^{(d,1)}$ has an interesting interpretation. Indeed, in that case the random isometry V is a Haar-distributed random unitary matrix $U \in \mathcal{U}(d)$, and thus

$$\lambda_1^{(d,s=1)} = |\operatorname{Tr} U|^2.$$

Then, for any $p \leq d/2$, we have [DS94, Theorem 2]

$$\mathbb{E}\Big[\big|\operatorname{Tr} U\big|^{2p}\Big] = p!.$$

In the general case. if we apply graphical Weingarten formula and use the fact that $\sigma \mapsto Wg_d(\sigma)$ is the pseudo-inverse of $\sigma \mapsto d^{\#\sigma}$ in $\mathbb{C}[S_p]$, we can extend the same formula to all d, and in general we have

$$\mathbb{E}\left[|\operatorname{Tr} U|^{2p}\right] = p! \sum_{\sigma \in S_p} d^{\#\sigma} \operatorname{Wg}_d(\sigma) = \sum_{\lambda \vdash p, \ l(\lambda) \leq d} d_\lambda^2,$$

where d_{λ} is the dimension of the irreducible representation of S_p associated to the partition λ , with the condition that the length of the diagram λ is smaller than the dimension d.

We now prove the claims about the convergence of the random variables $\lambda_{1,2}^{(d,s)}$ from Section 3.

Proof of Proposition 3.4. We start with the case where s is fixed and we treat the $\lambda_1^{(d,s)}$ random variable. We have

$$\mathbb{E}\Big[\big(\lambda_1^{(d,s)}\big)^p\Big] = s(s+1)\cdots(s+p-1)\sum_{\beta\in S_p} d^{\#\beta} \mathrm{Wg}_{ds}(\beta)$$

= $s(s+1)\cdots(s+p-1)\sum_{\beta\in S_p} d^{p-|\beta|}(ds)^{-p-|\beta|} \mathrm{M\ddot{o}b}(\beta)(1+O((ds)^{-2}))$
= $s^{-p}\cdot s(s+1)\cdots(s+p-1)+O(d^{-1}).$

The assertion follows now from the observation that

$$s(s+1)\cdots(s+p-1) = \int_0^\infty \frac{x^{s+p-1}}{(s-1)!} e^{-x} dx$$

is the *p*-th moment of the Gamma distribution $\Gamma(s, 1)$.

For the assertion regarding $\lambda_2^{(d,s)}$, compute

$$\mathbb{E}\left[(\lambda_{2}^{(d,s)})^{p} \right] = \sum_{\alpha,\beta \in S_{p}} s^{\#\alpha} d^{\#(\alpha\beta)} \operatorname{Wg}_{ds}(\alpha\beta^{-1})$$

$$= \sum_{\alpha,\beta \in S_{p}} s^{p-|\alpha|} d^{p-|\alpha\beta|} (ds)^{-p-|\alpha\beta^{-1}|} (\operatorname{M\"ob}(\alpha\beta^{-1}) + O((ds)^{-2}))$$

$$= \sum_{\alpha,\beta \in S_{p}} s^{-|\alpha|} + \sum_{\substack{\alpha,\beta \in S_{p} \\ |\alpha\beta| = |\alpha\beta^{-1}| = 0}} s^{-|\alpha|} d^{-1} \operatorname{M\"ob}(\alpha\beta^{-1}) + O(d^{-2})$$

$$= \sum_{\substack{\alpha \in S_{p} \\ \alpha^{2} = \operatorname{id}}} s^{-|\alpha|} + \sum_{\substack{\alpha,\beta \in S_{p} \\ |\alpha\beta| = 1, |\alpha\beta^{-1}| = 0}} s^{-|\alpha|} d^{-1} - \sum_{\substack{\alpha,\beta \in S_{p} \\ |\alpha\beta| = 0, |\alpha\beta^{-1}| = 1}} s^{-|\alpha|-1} d^{-1} + O(d^{-2})$$

$$= \sum_{\substack{\alpha \in S_{p} \\ \alpha^{2} = \operatorname{id}}} s^{-|\alpha|} + O(d^{-2}).$$

Here we used the fact that there is no pair (α, β) such that $|\alpha\beta| = 1, |\alpha\beta^{-1}| = 0$ or $|\alpha\beta| = 0, |\alpha\beta^{-1}| = 1$. Indeed, for both two cases we have $|\alpha^2| = 1$ or equivalently, α^2 is a transposition. However, this is impossible since $\operatorname{sgn}(\alpha^2) = \operatorname{sgn}(\alpha)^2 = 1$. We have thus the moment limit

$$\lim_{d \to \infty} \mathbb{E}[(s\lambda_2^{(d,s)})^p] = \sum_{\pi \in \mathcal{P}_{1,2}(p)} s^{\#\pi},$$

where we have identified idempotent permutations $\alpha \in S_p$ with partitions $\pi = \pi(\alpha) \in \mathcal{P}(p)$ having only blocks of size 1 or 2. Recall now that the normal distribution $\mathcal{N}(s, s)$ has classical cumulants

$$c_n = \begin{cases} s & \text{if } n = 1, 2, \\ 0 & \text{otherwise.} \end{cases}$$

Hence the moment limit above corresponds to the *p*-th moment of the normal distribution $\mathcal{N}(s, s)$ by the classical moment-cumulant formula, finishing the proof of the case *s* fixed.

Let us now prove the statement (2) of the proposition, regarding the convergence $\lambda_{1,2}^{(d,s)} \to 1$ as $d \to \infty$ and $s(d) \to \infty$. We start with the case of $\lambda_1^{(d,s)}$. By Lemma A.11 and A.4, we have

$$\mathbb{E}\left[\left(\lambda_{1}^{(d,s)}\right)^{p}\right] = s^{p} \sum_{k=0}^{\infty} a_{k,p} s^{-k} \sum_{\beta} d^{p-|\beta|} (ds)^{-p-|\beta|} (\text{M\"ob}(\beta) + O((ds)^{-2}))$$
$$= \sum_{k=0}^{\infty} a_{k,p} s^{-k} \left(1 - \frac{p(p-1)}{2} d^{-2} s^{-1} + O((ds)^{-2})\right)$$
$$= \sum_{k=0}^{\infty} a_{k,p} s^{-k} + O(d^{-2})$$
$$= \sum_{k=0}^{N-1} a_{k,p} s^{-k} + O(s^{-N}) + O(d^{-2})$$

for any integers $p, N \ge 0$. Then by Lemma A.3, we can further estimate

$$\mathbb{E}\left[(\lambda_1^{(d,s)} - 1)^{2N}\right] = \sum_{p=0}^{2N} (-1)^{2N-p} \mathbb{E}[(\lambda_1^{(d,s)})^p] \binom{2N}{p}$$
$$= \sum_{k=0}^{N-1} \sum_{p=0}^{2N} (-1)^{2N-p} a_{k,p} \binom{2N}{p} s^{-k} + O(s^{-N}) + O(d^{-2})$$
$$= O(s^{-N}) + O(d^{-2}).$$

In particular, this implies $\|\lambda_1^{(d,s)} - 1\|_{L^{2N}} \to 0$ for all integers $N \ge 1$, and hence $\|\lambda_1^{(d,s)} - 1\|_{L^p} \to 0$ for all $p \in (0,\infty)$, whenever $s, d \to \infty$. Now assuming $s \gtrsim d^t$ for some t, we can choose N such that $Nt \ge 2$ so that above estimation again implies

$$\mathbb{E}\left[(\lambda_1^{(d,s)} - 1)^{2N}\right] = O(d^{-2}).$$

Therefore, we have for any $\varepsilon > 0$,

$$\sum_{d=1}^{\infty} \mathbb{P}\left(|\lambda_1^{(d,s)} - 1| \ge \varepsilon \right) \le \sum_{d=1}^{\infty} \frac{\mathbb{E}[(\lambda_1^{(d,s)} - 1)^{2N}]}{\varepsilon^{2N}} < \infty$$

by the Markov inequality. Then the Borel-Cantelli lemma implies that $\lambda_1^{(d,s)} \to 1$ almost surely.

The proof for $\lambda_2^{(d,s)}$ is identical to the one above, up to replacing the constants $a_{k,p}$ by the explicit constants $b_{k,p}$ from Lemma A.4.

Finally, let us prove the statement (3) of Proposition 3.4. We show only the assertions for $\lambda_1^{(d,s)}$ since the one for $\lambda_2^{(d,s)}$ can be analogously shown. Lemma A.11 implies that for every $\varepsilon > 0$,

$$\mathbb{P}(|d^{-t}\lambda_1^{(d,s)}| > \varepsilon) \le \frac{1}{d^{Nt}\varepsilon^N} \mathbb{E}[(\lambda_1^{(d,s)})^N] = s^{1-p}(s+1)\cdots(s+N-1)O(d^{-Nt}) = O(d^{-2}),$$

for any choice of large N such that $Nt \ge 2$; this proves the claim in the regime $d \to \infty$. On the other hand, for the regime $s \to \infty$, Lemma A.11 again implies

$$\mathbb{E}[(\lambda_1^{(d,s)})^p] = 1 + \frac{p(p-1)}{2}(1-d^{-2})s^{-1} + O(s^{-2}).$$

Thus, we have

$$\begin{split} \mathbb{P}(|d^{-t}(\lambda_1^{(d,s)} - 1)| > \varepsilon) &\leq \frac{1}{d^{4t}\varepsilon^4} \mathbb{E}[(\lambda_1^{(d,s)} - 1)^4] \\ &= \sum_{p=0}^4 (-1)^{4-p} \left(1 + \frac{p(p-1)}{2}(1 - d^{-2}s^{-1})\right) + O(s^{-2}) = O(s^{-2}) \end{split}$$

by Lemma A.3. Now the usual Borel-Cantelli arguments again show the almost sure convergence $d^{-t}(\lambda_1^{(d,s)}-1) \to 0$ both for the cases $d \to \infty$ and $s \to \infty$, and the proof is complete.

We now show the convergence result for the random variable $\lambda_3^{(d,s)}$.

Proof of Proposition 4.3. The proof proceeds by a moment computation using the formula for $\lambda_3^{(d,s)}$ from Lemma A.11:

$$\mathbb{E}\left[(\lambda_{3}^{(d,s)})^{p}\right] = \sum_{\alpha,\beta\in S_{p}} s^{\#\alpha} d^{\#(\alpha\vee\beta)} \operatorname{Wg}_{ds}(\alpha\beta^{-1}) = 1 + \sum_{\substack{\alpha,\beta\in S_{p} \\ (\alpha,\beta)\neq(\mathrm{id},\mathrm{id})}} s^{\#\alpha} d^{\#(\alpha\vee\beta)} \operatorname{Wg}_{ds}(\alpha\beta^{-1}) = 1 + \sum_{\substack{\alpha,\beta\in S_{p} \\ (\alpha,\beta)\neq(\mathrm{id},\mathrm{id})}} s^{p-|\alpha|} d^{p-|\alpha\vee\beta|} (ds)^{-p-|\alpha\beta^{-1}|} (\operatorname{M\"ob}(\alpha\beta^{-1}) + O((ds)^{-2})) = 1 + \sum_{\substack{\alpha,\beta\in S_{p} \\ (\alpha,\beta)\neq(\mathrm{id},\mathrm{id}) \\ |\alpha\vee\beta|+|\alpha\beta^{-1}|=1}} s^{-|\alpha|-|\alpha\beta^{-1}|} d^{-1} \operatorname{M\"ob}(\alpha\beta^{-1}) + O(d^{-2}) = 1 + \sum_{\substack{\alpha,\beta\in S_{p} \\ (\alpha,\beta)\neq(\mathrm{id},\mathrm{id}) \\ |\alpha\vee\beta|=1, |\alpha\beta^{-1}|=0}} s^{-|\alpha|} d^{-1} + O(d^{-2}) = 1 + \frac{p(p-1)}{2} (ds)^{-1} + O(d^{-2}).$$

Using Lemma A.3, we can show

$$\mathbb{E}\left[(\lambda_3^{(d,s)} - 1)^4\right] = O(d^{-2}).$$

The almost sure convergence $\lambda_3^{(d,s)} \to 1$ follows by the Markov inequality and the Borel-Cantelli lemma.

A.3. Matrix parameters. This section contains results about the matrix parameters A, B, C which appear in the study of random CDUC, DUC, and DOC quantum channels from Section 5.

We begin with the description of average moments. Note that for our analysis, only the asymptotic eigenvalue distribution and moments of the matrix C is needed. We give, for the sake of completeness, the formulas for the three matrices.

Lemma A.12. The average pth moment of the random matrices A, B, C are given by

$$\mathbb{E}[\operatorname{Tr}(A^{p})] = \sum_{\alpha,\beta\in S_{p}} s^{\#\alpha} d^{\#((\gamma^{-1}\alpha\gamma)\vee\beta)} \operatorname{Wg}_{ds}(\alpha\beta^{-1}),$$
$$\mathbb{E}[\operatorname{Tr}(B^{p})] = \sum_{\alpha,\beta\in S_{p}} s^{\#\alpha} d^{\#((\gamma^{-1}\alpha)\vee(\gamma^{-1}\beta))} \operatorname{Wg}_{ds}(\alpha\beta^{-1}),$$
$$\mathbb{E}[\operatorname{Tr}(C^{p})] = \sum_{\alpha,\beta\in S_{p}} s^{\#\alpha} d^{\#((\gamma^{-1}\alpha)\vee(\gamma^{-1}\beta^{-1}))} \operatorname{Wg}_{ds}(\alpha\beta^{-1}).$$

where $\gamma = (p \ p - 1 \ \cdots \ 2 \ 1) \in S_p$ is the full cycle.

Proof. We shall prove the result only for the matrix C, since it is the only result we shall use later. The proofs for A, B are similar and we leave them to the reader.

To show the third formula above, we need to compute the expectation of the diagram in Fig. 10, which is obtained by tracing the product of p copies of the diagram for C from Fig. 8.



FIGURE 10. The diagram that corresponds to the quantity $Tr(C^p)$.

To compute the expectation value of the diagram, we use the graphical Weingarten formula from [CN10, CN15]. We obtain a sum indexed by a pair of permutations $\alpha, \beta \in S_p$ of diagrams $\mathcal{D}_{\alpha,\beta}$, weighted by the Weingarten function $Wg_{ds}(\alpha\beta^{-1})$. Each diagram $\mathcal{D}_{\alpha,\beta}$ in the sum will be a collection of loops corresponding to the \mathbb{C}^d and \mathbb{C}^s vector spaces. Our goal is to count how many loops of each type appear in the diagram $\mathcal{D}_{\alpha,\beta}$. We refer to Fig. 11 for the construction of the diagram $\mathcal{D}_{\alpha,\beta}$, obtained by erasing the V and the V* boxes, and connecting the loose wires using α for the left side of the V boxes and β for the right side of the V boxes. Note that in the Weingarten formula (21), the indices of V and \overline{V} are matched; in our setting, we connect to the opposite sides of $V^* = \overline{V}^\top$.



FIGURE 11. The way the *i*-th V and V^{*} boxes are connected, using the permutations $\alpha, \beta \in S_p$, to the rest of the diagram.

The number of loops of type s in $\mathcal{D}_{\alpha,\beta}$ is easily computed: only the permutation α is involved in connecting the *s*-type wires, and it is easy to see that each cycle of α yields a connected component (loop) in the end. We have thus a contribution of $s^{\#\alpha}$. The computation of the loops of size d is more complicated, because of the presence of the "vertical" connection between the *i*-th V and V^* boxes. In the absence of this vertical connection, each V box would have been connected to

the previous V^* box, resulting in a total number of $\#(\gamma^{-1}\alpha) + \#(\gamma^{-1}\beta^{-1})$ d-loops. The vertical wire connects some of the resulting loops, yielding the announced number of connected components (loops).

We now prove the formulas for the limiting eigenvalue distributions of the random matrix C.

Proof of Proposition 5.5. First of all, suppose $s/d \to 0$ as $d \to \infty$. Our method is to show that every mean p-th moment of $(ds)^{1/2}C$ converges to that of the semicircular distribution $SC_{0,1}$, and its variance converges to 0. First, by applying Lemma A.12,

$$\begin{aligned} \frac{1}{d} \mathbb{E} \left[\operatorname{Tr} \left((ds)^{p/2} C^p \right) \right] &= \sum_{\alpha, \beta \in S_p} s^{p/2 - |\alpha| - |\alpha\beta^{-1}|} d^{p/2 - 1 - |\gamma^{-1}\alpha \vee \gamma^{-1}\beta^{-1}| - |\alpha\beta^{-1}|} \left(\operatorname{M\"ob}(\alpha\beta^{-1}) + O((ds)^{-2}) \right) \\ &= \left[\sum_{|\alpha| + |\alpha\beta^{-1}| \le p/2} \left(\frac{s}{d} \right)^{p/2 - |\alpha| - |\alpha\beta^{-1}|} d^{p-1 - |\alpha| - |\gamma^{-1}\alpha \vee \gamma^{-1}\beta^{-1}| - 2|\alpha\beta^{-1}|} \right. \\ &+ \left. \sum_{|\alpha| + |\alpha\beta^{-1}| > p/2} s^{p/2 - |\alpha| - |\alpha\beta^{-1}|} d^{p/2 - 1 - |\gamma^{-1}\alpha \vee \gamma^{-1}\beta^{-1}| - |\alpha\beta^{-1}|} \right] \left(\operatorname{M\"ob}(\alpha\beta^{-1}) + O((ds)^{-2}) \right) \end{aligned}$$

Note that by Lemma A.8,

$$\frac{p}{2} - 1 - |\gamma^{-1}\alpha \vee \gamma^{-1}\beta^{-1}| - |\alpha\beta^{-1}| \le \frac{p}{2} - 1 - \frac{1}{2}|\gamma^{-2}| = \begin{cases} 0 & \text{if } p \text{ is even} \\ -1/2 & \text{if } p \text{ is odd,} \end{cases}$$

and when p is even, the equality condition further implies $\alpha = \beta \in NC_2(p)$ (Eq. (25)) which contradicts the condition $|\alpha| > p/2$. Therefore, the second sum becomes $O(s^{-1/2}d^{-1/2})$ regardless of p. Moreover, regarding the first sum,

$$p - 1 - |\alpha| - |\gamma^{-1}\alpha \vee \gamma^{-1}\beta^{-1}| - 2|\alpha\beta^{-1}| \le p - 1 - |\gamma| = 0,$$

and the equality holds iff $\alpha = \beta \in S_{NC}(\gamma) \cong NC_{1,2}(p)$. Therefore, we can further estimate

$$\frac{1}{d} \mathbb{E} \left[\operatorname{Tr} \left((ds)^{p/2} C^p \right) \right] = \sum_{\alpha \in NC_{1,2}(p)} \left(\frac{s}{d} \right)^{p/2 - |\alpha|} + O(d^{-2}) + O((ds)^{-1/2})$$
$$\rightarrow \begin{cases} \sum_{\alpha \in NC_2(p)} 1 = \operatorname{Cat}_{p/2} & \text{if } p \text{ is even,} \\ 0 & \text{if } p \text{ is odd,} \end{cases}$$

as $d \to \infty$. On the other hand, if we define $\tilde{\gamma} := (2p \, 2p - 1 \, \dots \, p + 1)(p \, p - 1 \, \dots \, 2 \, 1) \in S_{2p}$ and apply the Weingarten calculus, we have

$$\mathbb{E}\left[\frac{1}{d^2}\left(\operatorname{Tr}\left((ds)^{p/2}C^p\right)\right)^2\right] = \sum_{\alpha,\beta\in S_{2p}} s^{p-|\alpha|-|\alpha\beta^{-1}|} d^{p-2-|\tilde{\gamma}^{-1}\alpha\vee\tilde{\gamma}^{-1}\beta^{-1}|-|\alpha\beta^{-1}|} \left(\operatorname{M\"ob}(\alpha\beta^{-1}) + O((ds)^{-2})\right).$$

By repeating the prior argument, we can show that

$$\mathbb{E}\left[\frac{1}{d^2} \left(\operatorname{Tr}\left((ds)^{p/2} C^p\right) \right)^2 \right] = \sum_{\alpha \in S_{NC_{1,2}}(\tilde{\gamma})} \left(\frac{s}{d}\right)^{p-|\alpha|} + O(d^{-2}) + O((ds)^{-1})$$
$$= \left(\sum_{\pi \in NC_{1,2}(p)} \left(\frac{s}{d}\right)^{p/2-|\pi|} \right)^2 + O(d^{-1}),$$

where the last equality follows from $S_{NC_{1,2}}(\tilde{\gamma}) \cong N_{1,2}(p) \times N_{1,2}(p)$ (Eq. (24)). Therefore, we obtain

$$\mathbb{E}\left[\left(\frac{1}{d}\operatorname{Tr}\left((ds)^{p/2}C^{p}\right)\right)^{2}\right] - \left(\frac{1}{d}\mathbb{E}\left[\operatorname{Tr}\left((ds)^{p/2}C^{p}\right)\right]\right)^{2} = O(d^{-1/2}) \to 0$$

for all p, hence proving that for every $p \ge 1$,

$$\frac{1}{d}\operatorname{Tr}\left((ds)^{p/2}C^p\right) \to p\text{-th moment of } SC_{0,1}$$

in probability.

Now let us show that the convergence above holds almost surely for p = 1, 2, 3. First, note that Eq. (17) and Proposition 4.3 implies that

$$\frac{1}{d}\operatorname{Tr}\left((ds)^{1/2}C\right) = \left(\frac{s}{d}\right)^{1/2}\lambda_3^{(d,s)} \to 0$$

almost surely, regardless of the scale of s. Furthermore, by combining the previous arguments with Lemma A.9, we have that

$$\frac{1}{d} \mathbb{E}\left[\operatorname{Tr}\left((ds)^{p/2} C^p\right)\right] = \sum_{\alpha \in NC_{1,2}(p)} \left(\frac{s}{d}\right)^{p/2-|\alpha|} + O(d^{-3/2}),$$
$$\mathbb{E}\left[\frac{1}{d^2}\left(\operatorname{Tr}\left((ds)^{p/2} C^p\right)\right)^2\right] = \left(\sum_{\alpha \in NC_{1,2}(p)} \left(\frac{s}{d}\right)^{p/2-|\alpha|}\right)^2 + O(d^{-2})$$

hold for p = 2, 3. Therefore, the usual Borel-Cantelli technique shows the wanted almost sure convergences.

The remaining convergences can be obtained in analogous way. For the regime $s \sim cd$ for c > 0, Lemma A.12 and A.8 again imply

$$\frac{1}{d} \mathbb{E}\left[\operatorname{Tr}(s^{p}C^{p})\right] = \sum_{\alpha,\beta\in S_{p}} \left(\frac{s}{d}\right)^{\#\alpha-|\alpha\beta^{-1}|} d^{p-1-|\alpha|-|\gamma^{-1}\alpha\vee\gamma^{-1}\beta^{-1}|-2|\alpha\beta^{-1}|} \left(\operatorname{M\"ob}(\alpha\beta^{-1}) + O(d^{-4})\right) \\
= \sum_{\pi\in NC_{1,2}(p)} \left(\frac{s}{d}\right)^{\#\pi} + O(d^{-2}) \\
\to \sum_{\pi\in NC_{1,2}(p)} c^{\#\pi} \text{ as } d \to \infty,$$

$$\mathbb{E}\left[\left(\frac{1}{d}\operatorname{Tr}(s^{p}C^{p})\right)^{2}\right] = \sum_{\alpha,\beta\in S_{2p}} \left(\frac{s}{d}\right)^{\#\alpha-|\alpha\beta^{-1}|} d^{2p-2-|\alpha|-|\tilde{\gamma}^{-1}\alpha\vee\tilde{\gamma}^{-1}\beta^{-1}|-2|\alpha\beta^{-1}|} \left(\operatorname{M\"ob}(\alpha\beta^{-1}) + O(d^{-4})\right)$$
$$= \left(\sum_{\pi\in NC_{1,2}(p)} \left(\frac{s}{d}\right)^{\#\pi}\right)^{2} + O(d^{-2}),$$

and therefore,

$$\mathbb{E}\left[\left(\frac{1}{d}\operatorname{Tr}\left((ds)^{p/2}C^p\right)\right)^2\right] - \left(\frac{1}{d}\mathbb{E}\left[\operatorname{Tr}\left((ds)^{p/2}C^p\right)\right]\right)^2 = O(d^{-2}).$$

This shows that $\frac{1}{d} \operatorname{Tr}(s^p C^p) \to \sum_{\pi \in NC_{1,2}(p)} c^{\#\pi}$ almost surely for every $p \ge 0$. Note that this limit value is the *p*-th moment of the semicircular distribution $\operatorname{SC}_{c,c}$.

Fianlly, in the regime $d \to \infty$ and $s/d \to \infty$, we similarly have

$$\frac{1}{d}\mathbb{E}\left[\operatorname{Tr}(d^{p}C^{p})\right] = \sum_{\alpha,\beta\in S_{p}} \left(\frac{s}{d}\right)^{-|\alpha|-|\alpha\beta^{-1}|} d^{p-1-|\alpha|-|\gamma^{-1}\alpha\vee\gamma^{-1}\beta^{-1}|-2|\alpha\beta^{-1}|} \left(\operatorname{M\ddot{o}b}(\alpha\beta^{-1}) + O((ds)^{-2})\right) \\
= \sum_{\alpha\in NC_{1,2}(p)} \left(\frac{s}{d}\right)^{-|\alpha|} + O((ds)^{-1}) \\
\to 1 \text{ as } d \to \infty,$$
(29)

$$\mathbb{E}\left[\left(\frac{1}{d}\operatorname{Tr}(d^{p}C^{p})\right)^{2}\right] = \sum_{\alpha,\beta\in S_{2p}} \left(\frac{s}{d}\right)^{-|\alpha|-|\alpha\beta^{-1}|} d^{2p-2-|\alpha|-|\tilde{\gamma}^{-1}\alpha\vee\tilde{\gamma}^{-1}\beta^{-1}|-2|\alpha\beta^{-1}|} \left(\operatorname{M\"ob}(\alpha\beta^{-1}) + O((ds)^{-2})\right)$$
$$= \left(\sum_{\pi\in NC_{1,2}(p)} \left(\frac{s}{d}\right)^{-|\pi|}\right)^{2} + O(d^{-2}),$$
and therefore

and therefore,

$$\mathbb{E}\left[\left(\frac{1}{d}\operatorname{Tr}(d^p C^p)\right)^2\right] - \left(\frac{1}{d}\mathbb{E}\left[\operatorname{Tr}(d^p C^p)\right]\right)^2 = O(d^{-2}).$$

Proof of Proposition 5.6. By the Borel-Cantelli argument, it suffices to show that for any $\varepsilon > 0$,

$$\mathbb{P}\left(\|dC - I_d\|_{\infty} > \varepsilon\right) = O(d^{-(1+t)}).$$

Let us begin with the mean moment formula Eq. (29) of the random matrix dC. Applying Lemma A.4 (3), we further obtain

$$\frac{1}{d} \mathbb{E} \left[\text{Tr}(d^p C^p) \right] = \sum_{k=0}^{\infty} c_{k,p} \left(\frac{s}{d} \right)^{-k} + O((ds)^{-1}) \\ = \sum_{k=0}^{N-1} c_{k,p} \left(\frac{s}{d} \right)^{-k} + O\left((s/d)^{-N} \right) + O((ds)^{-1})$$

For arbitrary $N \ge 1$, where $c_{k,p} = \operatorname{Cat}_k {p \choose 2k}$. If we choose N such that $Nt \ge 2 + t$, then

$$\mathbb{E}\left[\operatorname{Tr}(d^{p}C^{p})\right] = \sum_{k=0}^{N-1} c_{k,p} d\left(\frac{s}{d}\right)^{-k} + O(d^{-(1+t)}),$$

Therefore, Lemma A.3 implies that

$$\mathbb{E}\left[\operatorname{Tr}\left((dC - I_d)^{2N}\right)\right] = \sum_{p=0}^{2N} (-1)^{2N-p} \mathbb{E}\left[\operatorname{Tr}(dC)^p\right] \binom{2N}{p}$$
$$= \sum_{k=0}^{N-1} d\left(\frac{s}{d}\right)^{-k} \sum_{p=0}^{2N} (-1)^{2N-p} c_{k,p} \binom{2N}{p} + O(d^{-(1+t)})$$
$$= O(d^{-(1+t)}).$$

Consequently, we obtain by Markov's inequality and the matrix norms $\|\cdot\|_{\infty} \leq \|\cdot\|_{2N}$,

$$\mathbb{P}(\|dC - I_d\|_{\infty} > \varepsilon) \le \mathbb{P}\left(\operatorname{Tr}\left((dC - I_d)^{2N}\right) > \varepsilon^{2N}\right) \le \frac{\mathbb{E}\left[\operatorname{Tr}\left((dC - I_d)^{2N}\right)\right]}{\varepsilon^{2N}} = O(d^{-(1+t)}).$$

Let us now consider the matrix elements of the three parameters A, B, C. We first compute the moments.

Lemma A.13. For i, j, the pth moment of the entry A_{ij} is

$$\mathbb{E}\left[(A_{ij})^p\right] = \frac{s(s+1)\cdots(s+p-1)}{ds(ds+1)\cdots(ds+p-1)}.$$

Moreover, for $i \neq j$, the pth moment of the values $|B_{ij}|^2$ and $|C_{ij}|^2$ are

$$\mathbb{E}\left[|B_{ij}|^{2p}\right] = \mathbb{E}\left[|C_{ij}|^{2p}\right] = p!s(s+1)\cdots(s+p-1)\sum_{\beta\in\mathcal{S}}\mathrm{Wg}_{ds}(\beta),$$

where $S = \{ \alpha \in S_{2p} : \alpha(i) \equiv i \mod 2 \text{ for all } i \}$. In particular, $|B_{ij}| \text{ and } |C_{ij}| \text{ have the same distribution.}$



FIGURE 12. Diagrams for the for the moments of the entries of the random matrices A, B, C. The first diagram contains p connected sub-diagrams, while the ones for B and C contain 2p connected sub-diagrams.

Proof. (Figure 12) By Weingarten calculus and Lemma A.2, we have

$$\mathbb{E}\left[(A_{ij})^p\right] = \sum_{\alpha,\beta\in S_p} s^{\#\alpha} \operatorname{Wg}_{ds}(\alpha\beta^{-1}) = \sum_{\alpha\in S_p} s^{\#\alpha} \sum_{\beta\in S_p} \operatorname{Wg}_{ds}(\beta) = \frac{s(s+1)\cdots(s+p-1)}{ds(ds+1)\cdots(ds+p-1)}$$

Next, since $B_{ij} = \overline{B_{ji}}$, we have

$$\mathbb{E}\left[|B_{ij}|^{2p}\right] = \mathbb{E}\left[(B_{ij}B_{ji})^p\right] = \sum_{\alpha,\beta\in\mathcal{T}} s^{\#\alpha} \mathrm{Wg}_{ds}(\alpha\beta^{-1}) = \sum_{\alpha\in\mathcal{T}} s^{\#\alpha} \sum_{\beta\in\mathcal{S}} \mathrm{Wg}_{ds}(\beta),$$

where $\mathcal{T} = \{ \alpha \in S_{2p} : \alpha(i) \equiv i + 1 \mod 2 \text{ for all } i \}$. Note the last equality follows from the observation that for any $\alpha \in \mathcal{T}$, the map $\beta \in \mathcal{T} \mapsto \alpha \beta^{-1} \in \mathcal{S}$ is a bijection. We claim that the map $\varphi : \alpha \in \mathcal{T} \mapsto \alpha^2|_{\{1,3,\ldots,2p-1\}} \in S(\{1,3,\ldots,2p-1\})$

is well-defined p!-to-1 surjection and $\#(\varphi(\alpha)) = \#\alpha$. Indeed, the fact that the map φ is well-defined is clear from the condition $\alpha(i) \equiv i+1 \mod 2$. Moreover, for arbitrary $\sigma \in S(\{1, 3, \ldots, 2p-1\})$ and a pair partition $\pi \in \mathcal{P}_2(2p)$ with $\pi(i) \equiv i+1 \mod 2$ for all i, we can construct a unique $\alpha_{\sigma,\pi} \in \mathcal{T}$ satisfying

 $\alpha_{\sigma,\pi}(2i-1) = \pi(2i-1), \quad \alpha_{\sigma,\pi}(\pi(2i-1)) = \sigma(2i-1)$

so that $\varphi(\alpha_{\sigma,\pi}) = \sigma$. One can easily check

$$\varphi^{-1}(\sigma) = \{\alpha_{\sigma,\pi} : \pi \in \mathcal{P}_2(2p) \text{ and } \pi(i) \equiv i+1 \mod 2 \text{ for all } i\}$$

and $\#(\alpha_{\sigma,\pi}) = \#\sigma$ regardless of the choice of π , and therefore the claim is shown. Now from the claim and Lemma A.2, we obtain that

$$\sum_{\alpha \in \mathcal{T}} s^{\#\alpha} = \sum_{\alpha \in \mathcal{T}} s^{\#\varphi(\alpha)} = p! \sum_{\sigma \in S_p} s^{\#\sigma} = p! s(s+1) \cdots (s+p-1),$$

hence proving the moment formula $\mathbb{E}[|B_{ij}|^{2p}]$.

The moment $\mathbb{E}[|C_{ij}|^{2p}]$ is calculated by the same method. These moments satisfy Carleman's condition [Akh20, Chapter 2, Addenda and Problems, 11], so the distributions of $|B_{ij}|$ and $|C_{ij}|$ are the same by uniqueness of the moments.

This allows us to prove the limit result for the entries of A, B, C.

Proof of Proposition 5.8. (1) and (2) is clear from

$$\lim_{d \to \infty} \mathbb{E}[(dsA_{ij})^p] = \lim_{d \to \infty} \frac{(ds)^p s(s+1) \cdots (s+p-1)}{ds(ds+1) \cdots (ds+p-1)} = s(s+1) \cdots (s+p-1),$$
$$\lim_{d \to \infty} \mathbb{E}[|dsB_{ij}|^{2p}] = \lim_{d \to \infty} \mathbb{E}[|dsC_{ij}|^{2p}] = \lim_{d \to \infty} p!(ds)^p s(s+1) \cdots (s+p-1)(ds)^{-p}(1+O((ds)^{-1})))$$
$$= p!s(s+1) \cdots (s+p-1),$$

by Lemma A.13, and the fact that $s(s+1)\cdots(s+p-1)$ and p! are p-th moment of the Gamma distribution $\Gamma(s, 1)$ and the exponential distribution $\mathsf{Exp}(1)$, as noted in the proof of Proposition 3.4.

For the proof of (3), let us recall from Eq. (16) that the random variables

$$A_{i_n j_n} = \sum_{k=1}^s |V_{i_n j_n}^{(k)}|^2, \quad A_{j_n i_n} = \sum_{k=1}^s |V_{i_n j_n}^{(k)}|^2, \quad B_{i_n j_n} = \sum_{k=1}^s V_{i_n i_n}^{(k)} \overline{V_{j_n j_n}^{(k)}},$$

for n = 1, ..., N, are functions of the $2s \times 2N$ -submatrix $V^{(2s \times 2N)}$ of V, where $V^{(2s \times 2N)}$ can be written as a block matrix $\left(V_{i_1j_1}^{(2s \times 2)} V_{i_2j_2}^{(2s \times 2)} \cdots V_{i_Nj_N}^{(2s \times 2)}\right)$ for

$$V_{ij}^{(2s\times2)} := \begin{pmatrix} V_{ii}^{(1)} & V_{ij}^{(1)} \\ \vdots & \vdots \\ V_{ii}^{(s)} & V_{ij}^{(s)} \\ V_{ji}^{(1)} & V_{jj}^{(1)} \\ \vdots & \vdots \\ V_{ji}^{(s)} & V_{jj}^{(s)} \end{pmatrix}$$

However, it is known [PR04, MT07] that if s and N are fixed, the random matrix $\sqrt{ds}V^{(2s\times 2N)}$ converges in distribution to the $(2s\times 2N)$ Ginibre ensemble whose entries are standard normal random variable as $d \to \infty$. In particular, the entries of $\sqrt{ds}V^{(2s\times 2N)}$ are asymptotically independent. Now the assertion follows from the observation that all the summands defining $\{A_{i_nj_n}, A_{j_ni_n}, B_{i_nj_n}\}_{n=1}^N$ are distinct.

Proof of Proposition 5.9. (1) The proof is similar with that of Proposition 3.4 (2). Applying Lemma A.13 and Lemma A.4,

$$\mathbb{E}\left[(dA_{ij})^{p}\right] = \frac{d^{p}s(s+1)\cdots(s+p-1)}{ds(ds+1)\cdots(ds+p-1)}$$
$$= \left(\sum_{k=0}^{\infty} a_{k,p}s^{-k}\right) \left(1 + \sum_{k=1}^{\infty} a_{k,p}(ds)^{-k}\right)^{-1}$$
$$= \left(\sum_{k=0}^{\infty} a_{k,p}s^{-k}\right) \left(1 - a_{1,p}(ds)^{-1} + (a_{1,p}^{2} - a_{2,p})(ds)^{-2} + O((ds)^{-3})\right)$$

for $p \ge 0$. If we choose large N such that $Nt \ge 3 + t$, then we can further proceed

$$\mathbb{E}\left[(dA_{ij})^p\right] = \sum_{k=0}^{N-1} s^{-k} a_{k,p} \left(1 - a_{1,p}(ds)^{-1} + (a_{1,p}^2 - a_{2,p})(ds)^{-2}\right) + O(d^{-(3+t)}).$$

Therefore, Lemma A.3 implies that

$$\mathbb{E}\left[(dA_{ij}-1)^{2N+4}\right] = \sum_{p=0}^{2N+4} (-1)^{2N+4-p} \mathbb{E}\left[(dA_{ij})^p\right] \binom{2N+4}{p}$$
$$= \sum_{k=0}^{N-1} s^{-k} \sum_{p=0}^{2N+4} (-1)^{2N+4-p} a_{k,p} F(p) \binom{2N+4}{p} + O(d^{-(3+t)})$$
$$= O(d^{-(3+t)}),$$

for all i, j, where $F(p) = 1 - a_{1,p}(ds)^{-1} + (a_{1,p}^2 - a_{2,p})(ds)^{-2}$ is a polynomial in p of degree 4. Now by Markov's inequality,

$$\mathbb{P}\left(\max_{i,j}|dA_{ij}-1| \ge \varepsilon\right) \le \sum_{i,j} \mathbb{P}\left(|dA_{ij}-1| \ge \varepsilon\right) \le \sum_{i,j} \frac{\mathbb{E}\left[(dA_{ij}-1)^{2N+4}\right]}{\varepsilon^{2N+4}} = O\left(d^{-(1+t)}\right)$$

for all $\varepsilon > 0$. Therefore, $\max_{i,j} |dA_{ij} - 1| \to 0$ (and hence $\max_{i,j} dA_{ij} \to 1$ and $\min_{i,j} dA_{ij} \to 1$) almost surely by the Borel-Cantelli lemma.

For (2) and (3), we again apply Lemma A.13 to have

$$\mathbb{E}\left[|d^{1+t'/2}B_{ij}|^{2p}\right] = \mathbb{E}\left[|d^{1+t'/2}C_{ij}|^{2p}\right] = d^{p(2+t')}s(s+1)\cdots(s+p-1)\sum_{\beta\in\mathcal{S}}\mathrm{Wg}_{ds}(\beta)$$
$$= d^{p(2+t')}p!s^{p}(1+O(s^{-1}))(ds)^{-2p}(1+O((ds)^{-2}))$$
$$= O(d^{pt'}s^{-p}) = O(d^{-p(t-t')}).$$

for $p \ge 0$. If we choose N such that $N(t - t') \ge 4$, we further have

$$\mathbb{P}\left(\max_{i,j}|d^{1+t'/2}B_{ij}| \ge \varepsilon\right) \le \sum_{i,j} \mathbb{P}\left(|d^{1+t'/2}B_{ij}| \ge \varepsilon\right)$$
$$\le \sum_{i,j} \frac{\mathbb{E}\left[|d^{1+t'/2}B_{ij}|^{2N}\right]}{\varepsilon^{2N}} = d^2 O(d^{-N(t-t')}) = O(d^{-2}),$$

and similarly $\mathbb{P}\left(\max_{i,j} |d^{1+t'/2}C_{ij}| \ge \varepsilon\right) = O(d^{-2})$ for all $\varepsilon > 0$. This concludes the proof. \Box

References

- [Akh20] Naum Ilich Akhiezer. The classical moment problem and some related questions in analysis. SIAM, 2020. 41
- [AN12] Guillaume Aubrun and Ion Nechita. Realigning random states. Journal of Mathematical Physics, 53(10):102210, 2012. 23
- [AN14] Muneerah Al Nuwairan. The extreme points of SU(2)-irreducibly covariant channels. Internat. J. Math., 25(6):1450048, 30, 2014. 2
- [ASY14] Guillaume Aubrun, Stanisław J. Szarek, and Deping Ye. Entanglement thresholds for random induced states. Comm. Pure Appl. Math., 67(1):129–171, 2014. 4, 7
- [Aub12] Guillaume Aubrun. Partial transposition of random states and non-centered semicircular distributions. Random Matrices Theory Appl., 1(2):1250001, 29, 2012. 4, 7, 22
- [BCLY20] Michael Brannan, Benoît Collins, Hun Hee Lee, and Sang-Gyun Youn. Temperley–lieb quantum channels. Communications in Mathematical Physics, 376(2):795–839, Jun 2020. 2
- [BCN16] Serban T. Belinschi, Benoît Collins, and Ion Nechita. Almost one bit violation for the additivity of the minimum output entropy. Comm. Math. Phys., 341(3):885–909, 2016. 2
- [BCS20] Ivan Bardet, Benoît Collins, and Gunjan Sapra. Characterization of equivariant maps and application to entanglement detection. Ann. Henri Poincaré, 21(10):3385–3406, 2020. 2
- [BCSZ09] Wojciech Bruzda, Valerio Cappellini, Hans-Jürgen Sommers, and Karol Życzkowski. Random quantum operations. Physics Letters A, 373(3):320–324, 2009. 2
- [BO08] Nathanial P. Brown and Narutaka Ozawa. C^{*}-algebras and finite-dimensional approximations, volume 88 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2008. 6
- [Bra37] Richard Brauer. On algebras which are connected with the semisimple continuous groups. Annals of Mathematics, 38(4):857–872, 1937. 9
- [BSM03] Abraham Berman and Naomi Shaked-Monderer. Completely positive matrices. World Scientific, 2003. 18

[Chr12] M. Christandl. PPT square conjecture. Banff International Research Station Workshop: Operator Structures in Quantum Information Theory, 2012. 4, 24

- [CK06] Dariusz Chruściński and Andrzej Kossakowski. Class of positive partial transposition states. Physical Review A, 74(2):022308, 2006. 17
- [CMHW18] Matthias Christandl, Alexander Müller-Hermes, and Michael M. Wolf. When do composed maps become entanglement breaking? *Annales Henri Poincaré*, 20:2295–2322, 2018. 2, 4, 24
- [CN10] Benoît Collins and Ion Nechita. Random quantum channels I: Graphical calculus and the Bell state phenomenon. *Communications in Mathematical Physics*, 297(2):345–370, 2010. 26, 27, 36
- [CN15] Benoît Collins and Ion Nechita. Random matrix techniques in quantum information theory. Journal of Mathematical Physics, 57(1):015215, 12 2015. 2, 27, 36
- [Col03] Benoît Collins. Moments and cumulants of polynomial random variables on unitarygroups, the Itzykson-Zuber integral, and free probability. International Mathematics Research Notices, 2003(17):953–982, 2003. 26, 27
- [COS18] Benoît Collins, Hiroyuki Osaka, and Gunjan Sapra. On a family of linear maps from $M_n(\mathbb{C})$ to $M_{n^2}(\mathbb{C})$. Linear Algebra Appl., 555:398–411, 2018. 2
- [CP22] Benoît Collins and Félix Parraud. Concentration estimates for random subspaces of a tensor product and application to quantum information theory. *Journal of Mathematical Physics*, 63(10):102202, 10 2022. 2
- [CS06] Benoît Collins and Piotr Śniady. Integration with respect to the Haar measure on unitary, orthogonal and symplectic group. *Communications in Mathematical Physics*, 264(3):773–795, 2006. 26, 27
- [CW03] Kai Chen and Ling-An Wu. A matrix realignment method for recognizing entanglement. Quantum Information & Computation, 3(3):193–202, 2003. 4, 23
- [CYT19] Lin Chen, Yu Yang, and Wai-Shing Tang. Positive-partial-transpose square conjecture for n = 3. Phys. Rev. A, 99:012337, Jan 2019. 4, 24
- [CYZ18] Benoît Collins, Zhi Yin, and Ping Zhong. The PPT square conjecture holds generically for some classes of independent states. Journal of Physics A: Mathematical and Theoretical, 51(42):425301, 2018. 4, 5, 7, 24, 26
- [DHS06] NILANJANA DATTA, ALEXANDER S. HOLEVO, and YURI SUHOV. Additivity for transpose depolarizing channels. International Journal of Quantum Information, 04(01):85–98, 2006. 2
- [DS94] Persi Diaconis and Mehrdad Shahshahani. On the eigenvalues of random matrices. Journal of Applied Probability, 31(A):49–62, 1994. 33
- [DSS⁺00] David P. DiVincenzo, Peter W. Shor, John A. Smolin, Barbara M. Terhal, and Ashish V. Thapliyal. Evidence for bound entangled states with negative partial transpose. *Phys. Rev. A*, 61:062312, May 2000. 17

ION NECHITA AND SANG-JUN PARK

- [DTW16] Nilanjana Datta, Marco Tomamichel, and Mark M Wilde. On the second-order asymptotics for entanglement-assisted communication. *Quantum Information Processing*, 15(6):2569–2591, 2016. 2
- [GBW21] Martina Gschwendtner, Andreas Bluhm, and Andreas Winter. Programmability of covariant quantum channels. *Quantum*, 5:488, June 2021. 2
- [GKS20] Mark Girard, Seung-Hyeok Kye, and Erling Størmer. Convex cones in mapping spaces between matrix algebras. *Linear Algebra and its Applications*, 608:248–269, 2020. 4, 24
- [Has09] M. B. Hastings. Superadditivity of communication capacity using entangled inputs. Nature Physics, 5(4):255–257, Apr 2009. 2
- [Hol93] Alexander S Holevo. A note on covariant dynamical semigroups. *Reports on mathematical physics*, 32(2):211–216, 1993. 2
- [Hol96] Alexander S Holevo. Covariant quantum markovian evolutions. *Journal of Mathematical Physics*, 37(4):1812–1832, 1996. 2
- [Hol05] A. S. Holevo. Additivity conjecture and covariant channels. International Journal of Quantum Information, 03(01):41–47, 2005. 2
- [HRF20] Eric P Hanson, Cambyse Rouzé, and Daniel Stilck França. Eventually entanglement breaking markovian dynamics: Structure and characteristic times. Ann. Henri Poincaré, 21:1517–1571, 2020. 4, 24
- [HSR03] Michael Horodecki, Peter W Shor, and Mary Beth Ruskai. Entanglement breaking channels. Reviews in Mathematical Physics, 15(06):629–641, 2003. 3
- [HT05] Uffe Haagerup and Steen Thorbjørnsen. A new application of random matrices: $\text{Ext}(C^*_{red}(\mathbb{F}_2))$ is not a group. Annals of Mathematics, pages 711–775, 2005. 22
- [HW08] Patrick Hayden and Andreas Winter. Counterexamples to the maximal p-norm multiplicativity conjecture for all <math>p > 1. Communications in Mathematical Physics, 284(1):263–280, Nov 2008. 2
- [JM19] Nathaniel Johnston and Olivia MacLean. Pairwise completely positive matrices and conjugate local diagonal unitary invariant quantum states. *Electronic Journal of Linear Algebra*, 35:156–180, 2019. 18
- [JY20] Ryan Jin and Yu Yang. Investigation of the PPT squared conjecture for high dimensions. *preprint* arXiv:2010.15554, 2020. 4, 24
- [Kin03] C. King. The capacity of the quantum depolarizing channel. IEEE Transactions on Information Theory, 49(1):221–229, 2003. 2
- [KKL11] Young-Hoon Kiem, Seung-Hyeok Kye, and Jungseob Lee. Existence of product vectors and their partial conjugates in a pair of spaces. *Journal of mathematical physics*, 52(12), 2011. 18
- [KMP17] Matthew Kennedy, Nicholas Manor, and Vern Paulsen. Composition of PPT maps. Quantum Information and Computation, 18, 10 2017. 4, 24
- [KNP⁺21a] Ryszard Kukulski, Ion Nechita, Łukasz Pawela, Zbigniew Puchała, and Karol Życzkowski. Generating random quantum channels. *Journal of Mathematical Physics*, 62(6):062201, 2021. 2
- [KNP⁺21b] Ryszard Kukulski, Ion Nechita, Łukasz Pawela, Zbigniew Puchała, and Karol Życzkowski. Generating random quantum channels. *Journal of Mathematical Physics*, 62(6):062201, 06 2021. 7
- [KO12] Seung-Hyeok Kye and Hiroyuki Osaka. Classification of bi-qutrit positive partial transpose entangled edge states by their ranks. *Journal of mathematical physics*, 53(5), 2012. 18
- [KW99] Michael Keyl and Reinhard F Werner. Optimal cloning of pure states, testing single clones. Journal of Mathematical Physics, 40(7):3283–3299, 1999. 2
- [KW09] Robert Koenig and Stephanie Wehner. A strong converse for classical channel coding using entangled inputs. *Physical Review Letters*, 103(7):070504, 2009. 2
- [LG15] L. Lami and V. Giovannetti. Entanglement-breaking indices. Journal of Mathematical Physics, 56(9):092201, Sep 2015. 4, 24
- [Liu15] Chaobin Liu. Unitary conjugation channels with continuous random phases. Quantum Studies: Mathematics and Foundations, 2(2):177–181, 2015. 17
- [LS15] Artur O Lopes and Marcos Sebastiani. Generic properties for random repeated quantum iterations. Quantum Studies: Mathematics and Foundations, 2(4):389–402, 2015. 17
- [LY22] Hun Hee Lee and Sang-Gyun Youn. Quantum channels with quantum group symmetry. Comm. Math. Phys., 389(3):1303–1329, 2022. 2
- [MS22] Laleh Memarzadeh and Barry C. Sanders. Group-covariant extreme and quasiextreme channels. *Phys. Rev. Res.*, 4:033206, Sep 2022. 2
- [MSD17] Marek Mozrzymas, Michał Studziński, and Nilanjana Datta. Structure of irreducibly covariant quantum channels for finite groups. *Journal of Mathematical Physics*, 58(5):052204, 05 2017. 2
- [MT07] Christian Mastrodonato and Roderich Tumulka. Elementary proof for asymptotics of large haardistributed unitary matrices. *Letters in Mathematical Physics*, 82(1):51–59, Oct 2007. 41
- [NC10] Michael A Nielsen and Isaac L Chuang. *Quantum computation and quantum information*. Cambridge University Press, 2010. 1

- [NS06] Alexandru Nica and Roland Speicher. Lectures on the Combinatorics of Free Probability. London Mathematical Society Lecture Note Series. Cambridge University Press, 2006. 27, 28, 29
- [NS21] Ion Nechita and Satvik Singh. A graphical calculus for integration over random diagonal unitary matrices. Linear Algebra and its Applications, 613:46 – 86, 2021. 18, 19
- [PJPY23] Sang-Jun Park, Yeong-Gwang Jung, Jeongeun Park, and Sang-Gyun Youn. A universal framework for entanglement detection under group symmetry, 2023. 2, 7, 13, 15
- [PR04] Dénes Petz and Júlia Réffy. On asymptotics of large haar distributed unitary matrices. *Periodica Mathematica Hungarica*, 49(1):103–117, Sep 2004. 41
- [PY23] Sang-Jun Park and Sang-Gyun Youn. *k*-positivity and schmidt number under orthogonal group symmetries. Preprint, arXiv:2306.00654, 2023. 2
- [RJP18] Mizanur Rahaman, Samuel Jaques, and Vern Paulsen. Eventually entanglement breaking maps. Journal of Mathematical Physics, 59, 01 2018. 4, 24
- [Rud05] Oliver Rudolph. Further results on the cross norm criterion for separability. Quantum Information Processing, 4:219–239, 2005. 4, 23
- [SBSGS22] Marcel Seelbach Benkner, Jens Siewert, Otfried Gühne, and Gael Sentís. Characterizing generalized axisymmetric quantum states in $d \times d$ systems. *Phys. Rev. A*, 106:022415, Aug 2022. 17
- [SDN22] Satvik Singh, Nilanjana Datta, and Ion Nechita. Ergodic theory of diagonal orthogonal covariant quantum channels. arXiv preprint arXiv:2206.01145, 2022. 17
- [Sim95] Barry Simon. Representations of Finite and Compact Groups (Graduate Studies in Mathematics; V. 10). American Mathematical Society, 12 1995. 5
- [SN21] Satvik Singh and Ion Nechita. Diagonal unitary and orthogonal symmetries in quantum theory. *Quantum*, 5:519, 2021. 2, 17, 18, 19, 23, 24
- [SN22a] Satvik Singh and Ion Nechita. Diagonal unitary and orthogonal symmetries in quantum theory II: Evolution operators. Journal of Physics A: Mathematical and Theoretical, 2022. 17
- [SN22b] Satvik Singh and Ion Nechita. The PPT² conjecture holds for all Choi-type maps. Annales Henri Poincaré, 23(9):3311–3329, 2022. 2, 3, 4, 5, 17, 24, 25
- [Sta11] Richard P. Stanley. Enumerative Combinatorics: Volume 1 (Cambridge Studies in Advanced Mathematics, Series Number 49). Cambridge University Press, paperback edition, 12 2011. 27
- [TAQ⁺18] Jordi Tura, Albert Aloy, Ruben Quesada, Maciej Lewenstein, and Anna Sanpera. Separability of diagonal symmetric states: a quadratic conic optimization problem. *Quantum*, 2:45, 2018. 18
- [TiB16] Jordi Tura i Brugués. Characterizing entanglement and quantum correlations constrained by symmetry. Springer, 2016. 9
- [VW01] K. G. H. Vollbrecht and R. F. Werner. Entanglement measures under symmetry. Phys. Rev. A, 64:062307, Nov 2001. 2, 10
- [Wat18] John Watrous. The Theory of Quantum Information. Cambridge University Press, 2018. 1
- [WH02] R. F. Werner and A. S. Holevo. Counterexample to an additivity conjecture for output purity of quantum channels. J. Math. Phys., 43(9):4353–4357, 2002. Quantum information theory. 2
- [Wil17] Mark M Wilde. *Quantum information theory*. Cambridge University Press, 2017. 1
- [WTB17] Mark M Wilde, Marco Tomamichel, and Mario Berta. Converse bounds for private communication over quantum channels. *IEEE Transactions on Information Theory*, 63(3):1792–1817, 2017. 2
- [Yu16] Nengkun Yu. Separability of a mixture of dicke states. *Physical Review A*, 94(6):060101, 2016. 18

Email address: nechita@irsamc.ups-tlse.fr

LABORATOIRE DE PHYSIQUE THÉORIQUE, UNIVERSITÉ DE TOULOUSE, CNRS, UPS, FRANCE

Email address: spark@irsamc.ups-tlse.fr

LABORATOIRE DE PHYSIQUE THÉORIQUE, UNIVERSITÉ DE TOULOUSE, CNRS, UPS, FRANCE