# Spectral Algorithms on Manifolds through Diffusion<sup>†</sup>

Weichun Xia and Lei Shi

School of Mathematical Sciences and Shanghai Key Laboratory for Contemporary Applied Mathematics, Fudan University, Shanghai 200433, China

#### Abstract

The existing research on spectral algorithms, applied within a Reproducing Kernel Hilbert Space (RKHS), has primarily focused on general kernel functions, often neglecting the inherent structure of the input feature space. Our paper introduces a new perspective, asserting that input data are situated within a low-dimensional manifold embedded in a higher-dimensional Euclidean space. We study the convergence performance of spectral algorithms in the RKHSs, specifically those generated by the heat kernels, known as diffusion spaces. Incorporating the manifold structure of the input, we employ integral operator techniques to derive tight convergence upper bounds concerning generalized norms, which indicates that the estimators converge to the target function in strong sense, entailing the simultaneous convergence of the function itself and its derivatives. These bounds offer two significant advantages: firstly, they are exclusively contingent on the intrinsic dimension of the input manifolds, thereby providing a more focused analysis. Secondly, they enable the efficient derivation of convergence rates for derivatives of any k-th order, all of which can be accomplished within the ambit of the same spectral algorithms. Furthermore, we establish minimax lower bounds to demonstrate the asymptotic optimality of these conclusions in specific contexts. Our study confirms that the spectral algorithms are practically significant in the broader context of high-dimensional approximation.

**Keywords:** Spectral algorithms, Heat kernel, Diffusion space, High-dimensional approximation, Convergence analysis

### **1** Introduction

Suppose that we are given a dataset  $D := \{(x_i, y_i)\}_{i=1}^n$  independently and identically distributed from an unknown distribution P defined on the product space  $X \times Y$  encompassing both the input space X and the output space Y. Our goal is to utilize this dataset to approximate the regression function, denoted as  $f^* : X \to Y$ , which minimizes the mean-squared error

$$\int_{X \times Y} \left( y - f(x) \right)^2 dP(x, y)$$

over all  $\nu$ -measurable functions  $f: X \to Y$  with  $\nu$  representing the marginal distribution of P on X.

<sup>&</sup>lt;sup>†</sup> The work described in this paper is supported by the National Natural Science Foundation of China (Grants Nos.12171039 and 12061160462) and Shanghai Science and Technology Program [Project No. 21JC1400600]. Email addresses: 22210180107@m.fudan.edu.cn (W. Xia), leishi@fudan.edu.cn (L. Shi).

To address this non-parametric regression problem, a widely adopted strategy is kernelbased regularization methods. This approach has been extensively explored and validated in many studies, e.g., [31, 4, 39, 32, 23, 17]. In this paper, we focus specifically on the spectral algorithms, which can be effectively implemented by acting a filter function on the spectra of a finite-dimensional kernel matrix. These algorithms demonstrate efficient performance, primarily attributed to the regularization imposed through a select group of filter functions [1, 14]. The choice of the regularization family is pivotal, as it enables the spectral algorithm to cover an extensive range of widely employed regression algorithms, including kernel ridge regression (also referred to as Tikhonov regularization), kernel principal component regression (often known as spectral cut-off), and various gradient methods. Such versatility is exemplified in the literature, see, e.g., [16, 11, 3, 26, 22, 5], which offers further insights into the applications of these methodologies. This adaptability also underscores the importance of spectral algorithms in regression analysis.

In many critical areas of modern data analysis, such as image recognition and DNA analysis, the prevalence of high-dimensional data is a notable challenge. This scenario is characterized by the dimensionality of the data significantly surpassing the number of available samples. To address this, researchers frequently turn to various dimensionality reduction techniques as a preliminary step. Among these techniques, some are based on the premise that data inherently resides on low-dimensional feature manifolds, where the intrinsic dimension is markedly lower than the ambient dimension. This concept is supported by evidence in various studies, such as [29, 2, 8, 36]. Building on this premise, it is plausible to hypothesize that the input space X is a low-dimensional manifold embedded within a higher-dimensional Euclidean space  $\mathbb{R}^d$ . This suggests that the intrinsic dimension m of the input manifold, now denoted as  $\mathcal{M}$ , is substantially lower than the ambient dimension d. Such a hypothesis aligns with the commonly accepted low-dimensional manifold hypothesis prevalent in related research, as illustrated in studies like [15, 25, 18]. This understanding forms a foundational aspect of our approach, providing an essential prior for the methodologies and techniques employed in our analysis.

Prior research on spectral algorithms within the framework of Reproducing Kernel Hilbert Space (RKHS) has predominantly concentrated on the utilization of general kernel functions. This approach frequently overlooks the intrinsic structure of the input feature space, which is a critical aspect of comprehending and enhancing the performance of algorithms. In our study, we particularly choose to utilize the specific heat kernel  $H_t$  with a fixed t > 0 on the input manifold  $\mathcal{M}$  as the generating kernel function for the RKHS. Our objective is to leverage intrinsic properties of the manifold, such as curvature, to enhance the performance of spectral algorithms. This approach represents a novel angle, one that has been relatively unexplored in existing studies on manifold regression. A particularly compelling theoretical challenge in this regression problem is how to derive the convergence rates in the "hard learning" scenarios, also known as model misspecifications [28, 27]. In this situation, the regression function may not necessarily be contained within the RKHS traditionally relied upon. Our focus is on comprehending the generalization error, represented by metrics like the  $\alpha$ -power norm (see (2.7))

$$\|f^* - f_{D,\lambda}\|_{\alpha} \tag{1.1}$$

in these demanding contexts, where  $f_{D,\lambda}$  is given by (2.10), denoting the estimator of spectral algorithm with regularization parameter  $\lambda$ . This aspect of our research is aligned with and generalizes several previous studies [13, 24, 40], seeking to provide a deeper understanding

and more effective solutions for this hard learning scenario.

Our foremost contribution is the derivation of convergence rates for (1.1) in hard learning scenarios, where the regression function is defined as  $f^* \in \mathcal{H}_t^\beta$  with  $\beta < 1$  and  $\mathcal{H}_t^\beta$  is given by (2.6). We achieve this result by leveraging the rapid decay property of the eigenvalues of the heat kernel, which demonstrates exponential decay attributable to the structure of the input manifold. This attribute facilitates a convergence rate as described in

$$\|f_{D,\lambda_n} - f^*\|_{\alpha}^2 \lesssim \left(\frac{(\log n)^{\frac{m}{2}}}{n}\right)^{\frac{\beta-\alpha}{\beta}}.$$
(1.2)

In this context,  $A \leq B$  means that there exists a constant C > 0 such that  $A \leq CB$ . This finding is outlined in Theorem 1. Notably, these results are predicated exclusively on the intrinsic dimension m of the input manifold  $\mathcal{M}$ , echoing findings from previous research [25]. However, our research expands those results to broader scenarios, offering improved convergence rates. Moreover, our work improves the convergence rate results obtained in prior studies that presupposed polynomial decay [13, 40]. As a consequential benefit, owing to the embedding property of the RKHS generated by the heat kernel, also referred to as the diffusion space, we also ascertain  $C^k$ -convergence rates for any k-th order derivatives, given by

$$\|f_{D,\lambda_n} - f^*\|_{C^k}^2 \lesssim \left(\frac{(\log n)^{\frac{m}{2}}}{n}\right)^{1-\varepsilon}$$

This is achieved with  $\epsilon > 0$  selected arbitrarily small, a result that may be intriguing in its own right.

Our second contribution is the demonstration of the optimality of the spectral algorithm using the heat kernel in terms of minimax rates, as elucidated in Theorem 2. We specifically focus on the special case when  $\alpha = 0$ , corresponding to the  $L^2$ -norm error scenario. The minimax lower bound in this situation is given by

$$\|\hat{f} - f^*\|_{L^2}^2 \gtrsim \frac{(\log n)^{\frac{m}{2}}}{n}.$$

In this context, our derived upper bound successfully aligns with the minimax lower bound. This alignment showcases the effectiveness of the spectral algorithm in this specific setting, underlining its adaptability and efficiency in handling  $L^2$ -norm error scenarios within the framework of minimax rates. Moreover, it extends previous research on lower bound results for exponential-type decaying eigenvalues (see, e.g., [11]) to manifold settings with dimension  $m \geq 2$ . However, in the more general error case  $\alpha > 0$ , we encounter challenges due to the complex nature of the exponential function. One may refer to Section 3 for details. As a result, our method cannot derive a lower bound that aligns precisely with the upper bound indicated by (1.2). Instead, we establish a lower bound that is slightly inferior in the power term of n, a finding also evident in Theorem 2. Given the potential interest this phenomenon might attract from future researchers, we deliberately underscore it, anticipating that subsequent studies might tackle this challenge in a more refined manner. We will also consider this problem in our future study.

As our third contribution, to the best of our knowledge, our work represents the first exploration into the use of a specific kernel function (the heat kernel) in spectral algorithms for misspecified models. This exploration utilizes a methodology that combines integral operator techniques with the embedding property. Integral operator techniques were initially employed in the study of kernel ridge regression (see, e.g., [4, 31]) and have recently been further developed for scalable kernel-based methods on massive data [16, 26, 30, 22]. However, investigation on integral operator techniques when the target function does not belong to the corresponding reproducing kernel Hilbert space and has low regularity has been scarce until recent work [13], which utilized integral operator techniques in conjunction with the embedding property to investigate the optimal convergence of kernel ridge regression in hard learning scenarios. However, the existing work has focused on using integral operator methods to study a general kernel function. In contrast to several prior works on spectral algorithms for misspecified models, which predominantly employ general polynomial decaying kernels, our investigation focuses on the heat kernels on manifolds. This particular kernel possesses superior eigenvalue decay properties (exponential decay) and nice space embedding properties, reflecting the intrinsic properties of the manifolds and thus leading to improved convergence rates and minimax rates. Notably, our convergence rate is optimal even when the target function exhibits low regularity. This improvement may encourage future research to explore specific kernels with superior performance.

The rest of this paper is structured as follows. In Section 2, we elucidate the fundamental properties of the heat kernel, diffusion spaces, integral operators, and spectral algorithms. In Section 3, we present our primary results concerning convergence rates and the minimax lower bounds, alongside some direct corollaries. In Section 4, we provide comprehensive proofs of our upper bounds results, which take into account the approximation error in Subsection 4.1 and the estimation error in Subsection 4.2 with a final proof for Theorem 1 furnished in Subsection 4.3. We ultimately complete the proof of Theorem 2 in Section 5.

# 2 Preliminaries

We now turn to a detailed description of our settings. We postulate that the input space X is an *m*-dimensional compact, connected, and immersed submanifold of  $\mathbb{R}^d$ , which we refer to as  $\mathcal{M}$ . First, we introduce several fundamental concepts related to the Riemannian manifold  $\mathcal{M}$ , as discussed in various literatures, including [12, 20]. We specify that the submanifold  $\mathcal{M}$  is endowed with the Radon measure  $\mu_g$ , which is derived from the integral operator on  $C_c(\mathcal{M})$ , as elucidated by the expression

$$\int_{\mathcal{M}} f d_{\mu_g} = \mu_g(f) = \sum_j \int_{\varphi_j(U_j)} \left( \psi_j f \sqrt{\det(g_{ij})} \right) \circ \varphi_j^{-1} dx_j^1 \cdots dx_j^m.$$

In this context,  $g = (g_{ij})$  signifies the Riemannian metric,  $\{U_j, \varphi_j; x_j\}_j$  denotes a smooth atlas, and  $\{\psi_j\}_j$  represents the partition of unity subordinate to the covering  $\{U_j\}_j$ . It is important to note that if  $\mathcal{M}$  is also orientable, a smooth, non-zero *m*-form, commonly known as a volume form, can be naturally selected on  $\mathcal{M}$  at the local level, as defined by

$$\omega = \sqrt{\det(g_{ij})} dx^1 \wedge \dots \wedge dx^m.$$

This volume form is characterized by the property that its integral over  $\mathcal{M}$  is positive, establishing the orientation of  $\mathcal{M}$  as per  $[\omega]$ . Consequently, this leads to the result

$$\mu_g(f) = \int_{\mathcal{M}} f\omega.$$

We now focus on the Laplace-Beltrami operator on  $\mathcal{M}$ , which is defined as

$$\Delta f = -\operatorname{div}(\nabla f).$$

The formulation of this definition is grounded in  $C_c^{\infty}(\mathcal{M})$  and is further extended to  $L^2(\mathcal{M}; \mu_g)$  through the application of Friedrich's extension, a detailed discussion of which can be found in [10]. Particularly, in the scenario where  $\mathcal{M} = \mathbb{R}^m$ , the Laplace-Beltrami operator aligns with the conventional second-order derivatives as indicated in  $-\sum_{i=1}^m \partial_i^2$ . To facilitate ease of reference in forthcoming discussions, we will henceforth denote  $\Delta$  as the Laplacian.

Utilizing the classical Sturm-Liouville decomposition approach for the Laplacian  $\Delta$ , we are able to delineate an orthonormal basis  $\{\varphi_k\}_{k\in\mathbb{N}}$  for  $L^2(\mathcal{M};\mu_g)$ . This approach and its underpinnings are thoroughly elaborated in [6]. Within this framework, each  $\varphi_k$  is characterized by its smoothness and adheres to the condition

$$\Delta \varphi_k = \lambda_k \varphi_k$$

for every integer k. Correspondingly,

$$0 = \lambda_1 \le \lambda_2 \le \dots \le \lambda_k \le \dots$$

represents the eigenvalues associated with the Laplacian  $\Delta$ , where  $\lambda_k \to +\infty$  as  $k \to \infty$ .

Fixing a specific t > 0, our attention is directed towards the diffusion space on  $\mathcal{M}$ , represented by

$$\mathcal{H}_t = e^{-\frac{t}{2}\Delta} L^2(\mathcal{M}; \mu_g).$$

This space is endowed with an inner product, defined as

$$\langle f,g\rangle_{\mathcal{H}_t} = \left\langle e^{\frac{t}{2}\Delta}f, e^{\frac{t}{2}\Delta}g\right\rangle_{L^2(\mathcal{M};\mu_g)},$$

thereby constituting a Hilbert space. Furthermore,  $\mathcal{H}_t$  serves as an RKHS, in which the reproducing kernel, known as the heat kernel, is designated by  $H_t(u, u')$ . The definition of this kernel is given by

$$\int_{\mathcal{M}} H_t(u, u') f(u') d\mu_g(u') = \left(e^{-t\Delta}f\right)(u).$$

In addition, we assume that the heat kernel is bounded, i.e., there exists a constant  $\kappa \geq 1$ , such that

$$\sup_{u \in \mathcal{M}} H_t(u, u) \le \kappa^2.$$
(2.1)

This assumption plays a crucial role in the subsequent analysis and applications of the heat kernel within our framework. To illustrate, let us consider the specific case where  $\mathcal{M} = S^m$  represents the m-dimensional unit sphere. In this context, the heat kernel on  $S^m$  (for  $m \geq 3$ ) could be characterized as follows, detailed in [7] as

$$H_t^{S^m}(u, u') = e^{\frac{(m-1)^2 t}{8}} \left(\frac{r}{\sin r}\right)^{\frac{m-1}{2}} \frac{e^{-r^2/2t}}{(2\pi t)^{m/2}} \cdot \mathbb{E}_r \exp\left(-\frac{(m-1)(m-3)}{8} \int_0^t \left(\frac{1}{\sin^2 R_s} - \frac{1}{R_s^2}\right) ds\right).$$

Here,  $r = d_{S^m}(u, u')$ ,  $R_s$  is indicative of an m-dimensional Bessel process, and  $\mathbb{E}_r$  symbolizes the expectation conditioned on  $R_t = r$ . It is important to note that the original theorem presented in [7] pertains to the hyperbolic space  $H^m$ , yet it can be suitably adapted for application to the sphere  $S^m$ . We have chosen to omit these details, as they fall outside the primary scope of our discussion. Given that the exponent in the integrands within the expectation is consistently negative, we are justified in setting  $\kappa^2 = e^{\frac{(m-1)^2 t}{8}}/(2\pi t)^{m/2}$ . In fact, for a generic compact manifold  $\mathcal{M}$ , its Ricci curvature and sectional curvature are intrinsically bounded. This inherent boundedness assures that the condition on the boundedness of heat kernel invariably satisfied, which follows from the heat kernel comparison theorems. We refer to [19] for more details.

A pivotal attribute of the diffusion space  $\mathcal{H}_t$  is its ability to be embedded, as outlined in [10], which is given by

$$\mathcal{H}_{t'} \hookrightarrow \mathcal{H}_t \hookrightarrow W^s(\mathcal{M}). \tag{2.2}$$

This embedding holds true for all values of 0 < t < t' and s > 0, where  $W^s(\mathcal{M})$  signifies the s-order Sobolev space on  $\mathcal{M}$  under the  $L^2_{\mu_g}$ -norm. For every  $s > \frac{m}{2}$ , the Sobolev space  $W^s(\mathcal{M})$  is compactly embedded into  $C(\mathcal{M})$ , which has been thoroughly discussed in [34]. Given that  $\mathcal{M}$  is compact, it follows that any diffusion space can be effectively embedded into  $L^{\infty}(\mathcal{M}; \mu_g)$ . This implies that for all values of t > 0, there exists a constant A (which depends on t) such that

$$\|i: \mathcal{H}_t \hookrightarrow L^\infty\| \le A.$$

In this context,  $\|\cdot\|$  represents the operator norm. This result plays a crucial role in our analysis of upper bounds. The practical implementation and significance of this embedding property have been historically traced to the work presented in [13].

Let the output space Y be the real line  $\mathbb{R}$ . We postulate that the unknown probability distribution P on  $\mathcal{M} \times \mathbb{R}$  adheres to the condition

$$P|_2^2 = \int_{\mathcal{M} \times \mathbb{R}} y^2 dP(x, y) < \infty.$$

The marginal distribution of P on  $\mathcal{M}$  is denoted by  $\nu$ , and the regular conditional probability of P given  $x \in \mathcal{M}$  is represented as  $P(\cdot|x)$ . Furthermore, we presume that the marginal distribution  $\nu$  corresponds to the uniform distribution on  $\mathcal{M}$ . In other words,  $\nu$  varies from  $\mu_g$  merely by multiplying a constant factor  $p = \frac{1}{\text{vol}\mathcal{M}}$ . This congruence ensures that  $L^2(\mathcal{M}; \mu_g)$ aligns with  $L^2(\mathcal{M}; \nu)$ , and hence, we will omit the symbol  $\mathcal{M}$  in subsequent discussions for the sake of conciseness. Additionally, through direct computation, the regression function is formulated as

$$f^*(x) = \int_{\mathbb{R}} y \ dP(y|x).$$

We now turn our attention to the inclusion map  $I_{\nu}$  from  $\mathcal{H}_t$  to  $L^2(\nu)$ . The adjoint of this operator, denoted as  $I_{\nu}^*$ , functions as an integral operator mapping from  $L^2(\nu)$  to  $\mathcal{H}_t$ , as defined by

$$(I_{\nu}^*f)(x) = \int_{\mathcal{M}} H_t(x, z) f(z) d\nu(z).$$

Both  $I_{\nu}$  and  $I_{\nu}^*$  qualify as Hilbert-Schmidt operators, which inherently implies their compactness. The Hilbert-Schmidt norm of these operators conforms to the relationship

$$\|I_{\nu}^{*}\|_{\mathscr{L}^{2}(L^{2}(\nu),\mathcal{H}_{t})} = \|I_{\nu}\|_{\mathscr{L}^{2}(\mathcal{H}_{t},L^{2}(\nu))} = \|H_{t}\|_{L^{2}(\nu)} = \left(\int_{\mathcal{M}} H_{t}(x,x)d\nu(x)\right)^{\frac{1}{2}} \leq \kappa$$

In addition, we introduce two supplementary integral operators, represented by

$$L_{\nu} = I_{\nu}I_{\nu}^* : L^2(\nu) \to L^2(\nu)$$

and

$$T_{\nu} = I_{\nu}^* I_{\nu} : \mathcal{H}_t \to \mathcal{H}_t.$$

$$(2.3)$$

These operators,  $L_{\nu}$  and  $T_{\nu}$ , are characterized as self-adjoint, positive-definite, and fall within the trace class. This classification suggests that they are also Hilbert-Schmidt and compact, with their trace norm adhering to

$$\|T_{\nu}\|_{\mathscr{L}^{1}(\mathcal{H}_{t})} = \|L_{\nu}\|_{\mathscr{L}^{1}(L^{2}(\nu))} = \|I_{\nu}^{*}\|_{\mathscr{L}^{2}(L^{2}(\nu),\mathcal{H}_{t})}^{2} = \|I_{\nu}\|_{\mathscr{L}^{2}(\mathcal{H}_{t},L^{2}(\nu))}^{2} \leq \kappa^{2}.$$

It is critical to highlight that  $L_{\nu}$  is the integral operator corresponding to the heat kernel  $H_t$ under the uniform distribution  $\nu$ . Moreover, by definition, the integral operator associated with  $H_t$  under the Radon measure  $\mu_g$  is explicitly expressed as  $e^{-t\Delta}$ . Given that  $\nu = p\mu_g$ holds, it is straightforward to verify that  $\{pe^{-t\lambda_k}, p^{-\frac{1}{2}}\varphi_k\}_{k\in\mathbb{N}}$  constitutes an eigensystem of  $L_{\nu}$  in  $L^2(\nu)$ . Consequently,  $\{p^{-\frac{1}{2}}\varphi_k\}_{k\in\mathbb{N}}$  serves as an orthonormal basis of  $L^2(\nu)$ , and we denote them as  $f_k = p^{-\frac{1}{2}}\varphi_k$ .

Given that both  $L_{\nu}$  and  $T_{\nu}$  are self-adjoint and compact operators, the spectral theorem enables us to deduce their respective spectral decompositions, which are expressed as

$$L_{\nu} = \sum_{k=1}^{\infty} p e^{-t\lambda_k} \langle \cdot, f_k \rangle_{L^2(\nu)} \cdot f_k$$

and

$$T_{\nu} = \sum_{k=1}^{\infty} p e^{-t\lambda_k} \langle \cdot, p^{\frac{1}{2}} e^{-\frac{t\lambda_k}{2}} f_k \rangle_{\mathcal{H}_t} \cdot p^{\frac{1}{2}} e^{-\frac{t\lambda_k}{2}} f_k.$$
(2.4)

Moreover, we have the capability to articulate  $I_{\nu}^*$  in the form of

$$I_{\nu}^{*} = \sum_{k=1}^{\infty} p^{\frac{1}{2}} e^{-\frac{t\lambda_{k}}{2}} \langle \cdot, f_{k} \rangle_{L^{2}(\nu)} \cdot p^{\frac{1}{2}} e^{-\frac{t\lambda_{k}}{2}} f_{k}.$$

Crucially, by applying Mercer's theorem, the heat kernel can be expanded, both uniformly and absolutely (as elaborated in [9, 32]), as

$$H_t(x,z) = \sum_{k=1}^{\infty} p e^{-t\lambda_k} f_k(x) f_k(z).$$
 (2.5)

Furthermore,  $\{p^{\frac{1}{2}}e^{-\frac{t\lambda_k}{2}}f_k\}_{k\in\mathbb{N}}$  establishes an orthonormal basis for  $\mathcal{H}_t$ , thereby providing a foundational structure for further analysis and application within this framework.

We will now turn our attention to the  $\alpha$ -power space denoted by  $\mathcal{H}_t^{\alpha} = \operatorname{Ran}(L_{\nu}^{\frac{\alpha}{2}})$ , which is defined as

$$\mathcal{H}_t^{\alpha} = \left\{ \sum_{k=1}^{\infty} a_k p^{\frac{\alpha}{2}} e^{-\frac{\alpha \lambda_k}{2} t} f_k : \{a_k\}_{k=1}^{\infty} \in l^2(\mathbb{N}) \right\}.$$
(2.6)

This space,  $\mathcal{H}_t^{\alpha}$ , will be equipped with an  $\alpha$ -power norm, specified as

$$\left\|\sum_{k=1}^{\infty} a_k p^{\frac{\alpha}{2}} e^{-\frac{\alpha\lambda_k}{2}t} f_k\right\|_{\alpha} := \left\|\sum_{k=1}^{\infty} a_k p^{\frac{\alpha}{2}} e^{-\frac{\alpha\lambda_k}{2}t} f_k\right\|_{\mathcal{H}^{\alpha}_t} = \|(a_k)\|_{l^2(\mathbb{N})} = \left(\sum_{k=1}^{\infty} a_k^2\right)^{\frac{1}{2}}.$$
 (2.7)

Consequently, this leads us to the following relationship

$$\|L_{\nu}^{\overline{2}}(f)\|_{\alpha} = \|f\|_{L^{2}(\nu)}.$$

With the introduction of this  $\alpha$ -power norm,  $\mathcal{H}_t^{\alpha}$  is constituted as a Hilbert space, complete with an orthonormal basis  $\{p^{\frac{\alpha}{2}}e^{-\frac{\alpha\lambda_k}{2}t}\}_{k\in\mathbb{N}}$ . For ease of reference in our subsequent discussions, we will employ the abbreviation  $\|\cdot\|_{\alpha}$  to represent the  $\alpha$ -power norm. It is essential to acknowledge that both  $\mathcal{H}_t^{\alpha} = \mathcal{H}_{\alpha t}$  and  $\mathcal{H}_t^0 = L^2(\nu)$  hold true. Further, due to the embedding property of the diffusion space, we can establish that for all  $0 < \beta < \alpha$ , the following is applicable:

$$\mathcal{H}^{\alpha}_t \hookrightarrow \mathcal{H}^{\beta}_t \hookrightarrow L^2(\nu).$$

This embedding property can also be deduced directly from the definition of  $\mathcal{H}_t^{\alpha}$ . In fact, the  $\alpha$ -power space  $\mathcal{H}_t^{\alpha}$  can be effectively described using the real interpolation method, as detailed in [33], given by

$$\mathcal{H}_t^{\alpha} = [L^2(\nu), \mathcal{H}_t]_{\alpha, 2}.$$

To provide an in-depth understanding of the spectral algorithm, it is imperative to first introduce the concept of the regularization family, often referred to as filter functions. A set of functions denoted by  $\{g_{\lambda} : \mathbb{R}^+ \to \mathbb{R}^+\}_{\lambda>0}$  is recognized as a regularization family if it satisfies specific criteria outlined in

$$\sup_{\substack{0 < t \le \kappa^2}} |tg_{\lambda}(t)| < 1, \\
\sup_{\substack{0 < t \le \kappa^2}} |1 - tg_{\lambda}(t)| < 1, \\
\sup_{\substack{0 < t \le \kappa^2}} |g_{\lambda}(t)| < \lambda^{-1}.$$
(2.8)

For a given regularization family  $g_{\lambda}$ , we define its qualification  $\xi$  as the supremum of the set

$$\xi = \sup\left\{\alpha > 0: \sup_{0 < s \le \kappa^2} |1 - sg_\lambda(s)| s^\alpha < \lambda^\alpha\right\}.$$
(2.9)

To exemplify these concepts, let us consider a few commonly utilized regularization families. The first example is  $g_{\lambda}(t) = \frac{1}{\lambda+t}$ , which possesses a qualification of  $\xi = 1$  and aligns with kernel ridge regression (also known as Tikhonov regularization). Another notable instance is  $g_{\lambda}(t) = \frac{1}{t} \mathbb{1}_{\{t \geq \lambda\}}$ , characterized by a qualification of  $\xi = \infty$ , and is associated with kernel principal component regularization (or spectral cut-off). These examples offer a more comprehensive insight into the functionality of different regularization families within the framework of spectral algorithms. For additional examples and a deeper exploration of this topic, please refer to [14].

In our final formulation of the spectral algorithm, we start with a dataset  $D = \{(x_i, y_i)\}_{i=1}^n$ that is independently and identically distributed from the distribution P. Initially, we define the sampling operator  $H_{t,x} : y \mapsto yH_t(x, \cdot)$  that maps from  $\mathbb{R}$  to  $\mathcal{H}_t$ , as well as its adjoint operator  $H_{t,x}^* : f \mapsto f(x)$  that maps from  $\mathcal{H}_t$  to  $\mathbb{R}$ . Further, we establish the sample covariance operator  $T_{\delta} : \mathcal{H}_t \to \mathcal{H}_t$ , which is described by

$$T_{\delta} = \frac{1}{n} \sum_{i=1}^{n} H_{t,x_i} H_{t,x_i}^*.$$

It is important to note that  $T_{\delta}$  is in alignment with the integral operator defined in (2.3), but with respect to the empirical marginal distribution  $\delta = \frac{1}{n} \sum_{i=1}^{n} \delta_{x_i}$ . In addition, we introduce the sample basis function as

$$g_D = \frac{1}{n} \sum_{i=1}^n y_i H_t(x_i, \cdot) \in \mathcal{H}_t.$$

Utilizing the sample function and operator outlined above, we can now define the spectral algorithm estimator, which is given by

$$f_{D,\lambda} = g_{\lambda}(T_{\delta})g_D. \tag{2.10}$$

In this setup,  $g_{\lambda}(T_{\delta})$  suggests that  $g_{\lambda}$  interacts with  $T_{\delta}$  through functional calculus. Significantly, (2.10) offers a finite, computable representation, the calculation of which entails an eigen-decomposition of the kernel matrix. This provides a practical approach to implementing the spectral algorithm in computational settings.

It is noteworthy that deriving an explicit representation of the heat kernel  $H_t$  for a general Riemannian manifold  $\mathcal{M}$  presents a significant challenge. This complexity poses a potential obstacle in the development of our spectral algorithm, which depends on the kernel function  $H_t(x, \cdot)$ . To circumvent this limitation, we propose employing graph Laplacian techniques as a means to approximate the heat kernel. We plan to elaborate on this approach and its implementation in detail in our forthcoming research.

# 3 Main Results

**Assumption 1.** There exists positive constants  $\sigma$ , L such that

$$\int_{\mathbb{R}} |y - f^*(x)|^m dP(y|x) \le \frac{1}{2} m! \sigma^2 L^{m-2}$$
(3.1)

for  $\nu$ -almost every  $x \in \mathcal{M}$  and all  $m \geq 2$ .

The moment condition we adopt here is a widely accepted assumption, principally designed to manage the discrepancy between observed data and the true underlying value. This condition guarantees that the tail probability of the noise diminishes at a sufficiently rapid pace, a practice that has been embraced in several scholarly works, as documented in [37]. Crucially, this moment condition is typically met in scenarios involving Gaussian noise characterized by bounded variance, as well as in distributions that are concentrated within certain bounded intervals on the real line. This makes it a mild and reliable assumption in a variety of statistical contexts.

**Assumption 2.** For  $\beta > 0$ , there exists a constant R > 0 such that  $f^* \in \mathcal{H}_t^{\beta}$  and

$$\|f^*\|_{\beta} \le R. \tag{3.2}$$

The source condition serves as a conventional metric for assessing the smoothness of the regression function, a methodology that has gained widespread adoption, as evidenced by the literature [1, 4]. It is important to recognize that lower values of  $\beta$  correspond to a decrease

in the smoothness of the regression function. This reduction in smoothness presents greater challenges for algorithms in terms of achieving accurate estimations. In instances where  $\beta < 1$ , the situation is classified as a hard learning scenario. This particular scenario has recently garnered considerable interest among researchers, as reflected in studies and discussions found in Section 1.

We are now prepared to unveil our principal findings. In light of our emphasis on the hard learning scenario, which is indicated by  $0 < \beta < 1$ , our theorem will be specifically tailored to the range encapsulated by  $0 < \beta \leq 1$ . Nonetheless, it is important to note that similar results could be obtained for  $\beta > 1$  with minimal adjustments. This approach ensures that our conclusions are both relevant and adaptable to varying scenarios within the scope of our study.

**Theorem 1.** Suppose that the regularization family  $\{g_{\lambda}\}_{\lambda>0}$  has qualification  $\xi \geq \frac{1}{2}$ , the moment condition (3.1) in Assumption 1 holds for some positive constants  $\sigma, L$ , and the source condition (3.2) in Assumption 2 holds for some R > 0 and  $0 < \beta \leq 1$ . For all  $0 \leq \gamma < \beta$ , suitably choose a regularization parameter sequence  $\{\lambda_n\}$  such that

$$\lambda_n \sim \left(\frac{(\log n)^{\frac{m}{2}}}{n}\right)^{\frac{1}{\beta}}.$$

Then, for all t > 0,  $\tau \ge 1$ , the spectral algorithm estimators  $f_{D,\lambda_n}$  with respect to RKHS  $\mathcal{H}_t$  satisfies

$$\|f_{D,\lambda_n} - f^*\|_{\gamma}^2 \le C\tau^2 \left(\frac{(\log n)^{\frac{m}{2}}}{n}\right)^{\frac{\beta-\gamma}{\beta}}$$

for sufficiently large  $n \ge 1$  with probability at least  $1 - 4e^{-\tau}$ , where C > 0 is a constant independent of  $n, \tau$ .

If the regression function is such that  $f^* \in \mathcal{H}_s$  holds for some s > 0, we can appropriately select and fix an  $t_o > s$  and employ the spectral algorithm estimator (2.10) to estimate the regression function in the RKHS  $\mathcal{H}_{t_0}$ . Drawing from Theorem 1, for any t < s, we are able to establish the following convergence rates in  $\mathcal{H}_t$ :

$$\|f_{D,\lambda_n} - f^*\|_{\mathcal{H}_t}^2 \lesssim \left(\frac{(\log n)^{\frac{m}{2}}}{n}\right)^{\frac{s-t}{s}}.$$
(3.3)

It is significant to note that this  $\mathcal{H}_t$ -convergence rate (3.3) is not contingent upon the specific RKHS  $\mathcal{H}_{t_0}$  previously selected; the only requirement being placed on the parameter  $t_o$  is  $t_o > s$ . Additionally, given that

$$\mathcal{H}_t \hookrightarrow W^s(\mathcal{M}) \hookrightarrow C^k(\mathcal{M})$$

holds true for all t > 0 and  $s > k + \frac{m}{2}$ , we also obtain  $C^k$ -convergence rates for any k-th order derivatives:

$$\|f_{D,\lambda_n} - f^*\|_{C^k}^2 \lesssim \left(\frac{(\log n)^{\frac{m}{2}}}{n}\right)^{1-\varepsilon}.$$

In this context, the parameter  $0 < \varepsilon < 1$  is derived from the stipulations of  $\beta > \gamma$  in Theorem 1. This finding underscores the versatility and robustness of the proposed spectral algorithm in various analytical contexts.

We are now prepared to introduce the minimax lower bound.

**Theorem 2.** Suppose that  $\nu$  is the uniform distribution on  $\mathcal{M}$ . Then, for all positive  $\sigma$ , L, R, and  $0 \leq \gamma < \beta \leq 1$ , there exists a constant C > 0 such that, for all estimators  $\hat{f} :=$  $\{(x_i, y_i)\}_{i=1}^n \mapsto \hat{f}, all \tau \in (0, 1), and all sufficiently large <math>n \geq 1$ , there exists a probability distribution P on  $\mathcal{M} \times \mathbb{R}$  with marginal distribution  $\nu$  on  $\mathcal{M}$  satisfying the moment condition (3.1) in Assumption 1 with respect to  $\sigma$ , L and the source condition (3.2) in Assumption 2 with respect to  $R, \beta$  such that, with  $P^n$ -probability at least  $1 - \tau^2$ , it holds

$$\|\hat{f} - f^*\|_{\gamma}^2 \ge C\tau^2 \frac{(\log n)^{\frac{m}{2}}}{n^s}$$

with s = 1 for  $\gamma = 0$  and  $s > \frac{(\beta - \gamma)2^{\frac{2}{m}}}{\gamma + (\beta - \gamma)2^{\frac{2}{m}}}$  for  $\gamma > 0$ .

Integrating the insights from Theorem 1 and Theorem 2, we discern that the spectral algorithm estimator (2.10) aligns with the minimax lower bound, particularly in the scenario defined by  $\gamma = 0$ . This scenario, described as the L<sup>2</sup>-convergence, is articulated by

$$||f_{D,\lambda_n} - f^*||_{L^2(\nu)}^2 \sim \frac{(\log n)^{\frac{m}{2}}}{n}$$

Such an alignment not only corroborates but also extends previous findings related to exponentially decayed or Gaussian decayed eigenvalues. These earlier results, with decay rates of  $O\left(\frac{\log n}{n}\right)$  corresponding to m = 2 and rates of  $O\left(\frac{\sqrt{\log n}}{n}\right)$  corresponding to m = 1 (as explored in [11]), are now applicable to manifolds of higher dimensions  $m \ge 2$ .

However, in the more general setting of  $\gamma > 0$ , challenges emerge due to the exponential function's rapid growth property, particularly highlighted in (5.6) of our proof. Specifically, the issue arises because  $e^{2x}$  cannot be effectively controlled by a constant multiple of  $e^x$ . This contrasts with approaches in [3, 13], where eigenvalues demonstrate polynomial decay. Consequently, our methods fall short in providing a precise bound for the constructed Gaussian type probability distribution. This shortfall is evident in the proof of Theorem 2 when aimed at establishing optimality for the general case  $\gamma > 0$ . To effectively manage the sum of eigenvalues of the integral operator in (5.6) and thereby derive a more stringent lower bound, a more advanced analytical approach is required. It is our hope that future research will be able to bridge this gap and offer a comprehensive solution to this complex problem.

# 4 Upper Bounds Analysis

Let us initially examine the continuous version of the estimator represented by  $f_{D,\lambda}$ , which we denote as  $f_{P,\lambda}$ . To define this, we consider

$$f_{P,\lambda} = g_{\lambda}(T_{\nu})g_P. \tag{4.1}$$

Here,  $g_P$  signifies the continuous variant of the sample function  $g_D$ . The expression for this continuous version is given by

$$g_P = I_{\nu}^* f^* = \int_{\mathcal{M} \times \mathbb{R}} y H_t(x, \cdot) dP(x, y).$$

The  $\gamma$ -norm error can be effectively decomposed into two distinct components:

$$||f_{D,\lambda} - f^*||_{\gamma} \le ||f_{D,\lambda} - f_{P,\lambda}||_{\gamma} + ||f_{P,\lambda} - f^*||_{\gamma}.$$
(4.2)

In this decomposition, the first term encapsulates the estimation error, while the second term pertains to the approximation error. We intend to undertake a detailed estimation of these two terms separately, with the estimation error being addressed in section 4.1 and the approximation error in section 4.2. This structured approach allows for a comprehensive analysis of the error components inherent in the estimator.

#### 4.1 Bounding approximation error

Recalling the spectral decomposition (2.4) of  $T_{\nu}$ , we have

$$T_{\nu} = \sum_{k=1}^{\infty} p e^{-t\lambda_k} \langle \cdot, p^{\frac{1}{2}} e^{-\frac{t\lambda_k}{2}} f_k \rangle_{\mathcal{H}_t} \cdot p^{\frac{1}{2}} e^{-\frac{t\lambda_k}{2}} f_k.$$

Through spectral calculus, this leads us to

$$g_{\lambda}(T_{\nu}) = \sum_{k=1}^{\infty} g_{\lambda}(pe^{-t\lambda_k}) \langle \cdot, p^{\frac{1}{2}}e^{-\frac{t\lambda_k}{2}} f_k \rangle_{\mathcal{H}_t} \cdot p^{\frac{1}{2}}e^{-\frac{t\lambda_k}{2}} f_k.$$

Further, if we expand  $f^*$  utilizing the orthonormal basis  $\{f_k\}_{k\in\mathbb{N}}$  within  $L^2(\nu)$ , it results in the expression

$$f^* = \sum_{k=1}^{\infty} a_k f_k.$$
 (4.3)

This expansion allows us to deduce the subsequent elaboration of  $f_{P,\lambda}$ , given by

$$f_{P,\lambda} = \sum_{k=1}^{\infty} g_{\lambda} (p e^{-t\lambda_k}) p^{\frac{1}{2}} e^{-\frac{t\lambda_k}{2}} a_k \cdot p^{\frac{1}{2}} e^{-\frac{t\lambda_k}{2}} f_k.$$
(4.4)

Consequently, this brings us to the conclusion encapsulated in

$$f^* - f_{P,\lambda} = \sum_{k=1}^{\infty} \left( 1 - g_{\lambda}(pe^{-t\lambda_k})pe^{-t\lambda_k} \right) a_k \cdot f_k.$$
(4.5)

**Lemma 1.** Suppose that  $f^* \in \mathcal{H}_t^{\beta}$  for some  $\beta > 0$ ,  $\{g_{\lambda}\}_{\lambda>0}$  has qualification  $\xi > 0$ . Then, for all  $\gamma \ge 0$ ,  $0 < \frac{\beta - \gamma}{2} \le \xi$ , and  $\lambda > 0$ , it holds

$$||f_{P,\lambda} - f^*||_{\gamma}^2 \le ||f^*||_{\beta}^2 \cdot \lambda^{\beta - \gamma}.$$
 (4.6)

*Proof.* Starting with the expansion (4.5) of the approximation error, we can write

$$\begin{split} \|f_{P,\lambda} - f^*\|_{\gamma}^2 &= \left\|\sum_{k=1}^{\infty} \left[1 - g_{\lambda}(pe^{-t\lambda_k})pe^{-t\lambda_k}\right] p^{-\frac{\gamma}{2}} e^{\frac{\gamma t\lambda_k}{2}} a_k \cdot p^{\frac{\gamma}{2}} e^{-\frac{\gamma t\lambda_k}{2}} f_k\right\|_{\gamma}^2 \\ &= \sum_{k=1}^{\infty} \left( \left[1 - g_{\lambda}(pe^{-t\lambda_k})pe^{-t\lambda_k}\right] p^{-\frac{\gamma}{2}} e^{\frac{\gamma t\lambda_k}{2}} a_k \right)^2 \\ &= \sum_{k=1}^{\infty} \left( \left[1 - g_{\lambda}(pe^{-t\lambda_k})pe^{-t\lambda_k}\right] p^{\frac{\beta-\gamma}{2}} e^{-\frac{(\beta-\gamma)t\lambda_k}{2}} \right)^2 p^{-\beta} e^{\beta t\lambda_k} a_k^2 \end{split}$$

Given that  $\{g_{\lambda}\}$  possesses a qualification of  $\xi \geq \frac{\beta - \gamma}{2}$ , and referring to (2.9), we arrive at

$$\left| \left[ 1 - g_{\lambda}(pe^{-t\lambda_k})pe^{-t\lambda_k} \right] p^{\frac{\beta-\gamma}{2}} e^{-\frac{(\beta-\gamma)t\lambda_k}{2}} \right| < \lambda^{\frac{\beta-\gamma}{2}}.$$

Thus, we can deduce

$$\|f_{P,\lambda} - f^*\|_{\gamma}^2 \le \sum_{k=1}^{\infty} \lambda^{\beta-\gamma} p^{-\beta} e^{\beta t \lambda_k} a_k^2 = \lambda^{\beta-\gamma} \|f^*\|_{\beta}^2.$$

The last equation is derived from combining the expansion (4.3) with the definition of the  $\beta$ -norm, as outlined in (2.7). The proof is then finished.

Utilizing the spectral decomposition (2.5) of the heat kernel  $H_t$ :

$$H_t(x,z) = \sum_{k=1}^{\infty} p e^{-t\lambda_k} f_k(x) f_k(z).$$

our preceding boundedness assumption, (2.1), can be reformulated as

$$\sum_{k=1}^{\infty} p e^{-t\lambda_k} f_k^2(x) \le \kappa^2.$$

This inequality serves as the foundation for introducing the concept of  $\alpha$ -boundedness of the heat kernel  $H_t$ , a topic explored in greater depth in [33]. Specifically, we define  $H_t$  as being  $\alpha$ -bounded by a constant A if the condition

$$\sum_{k=1}^{\infty} p^{\alpha} e^{-\alpha t \lambda_k} f_k^2(x) \le A^2$$
(4.7)

is met for  $\nu$ -almost every  $x \in \mathcal{M}$ . Additionally, the smallest constant A that satisfies (4.7) is denoted as  $||H_t^{\alpha}||_{\infty}$ . A key aspect of  $\alpha$ -boundedness is its connection to the embedding from the corresponding RKHS  $\mathcal{H}_t^{\alpha}$  into  $L^{\infty}(\nu)$ . This relationship is elaborated upon in the following lemma, which is extracted from [13]. This lemma plays a pivotal role in understanding the underlying dynamics of the heat kernel within the framework of RKHS.

**Lemma 2.** For all  $\alpha > 0$ , consider the inclusion map  $i_{\alpha} : \mathcal{H}_t^{\alpha} \to L^{\infty}(\nu)$ , then we have

$$\|i_{\alpha}\| = \|H_t^{\alpha}\|_{\infty}$$

Given the established fact that  $\mathcal{H}_t^{\alpha}$  is continuously embedded into  $L^{\infty}(\nu)$  for all t > 0and  $\alpha > 0$ , as inferred from the embedding property of the diffusion space (2.2), we are at liberty to select and fix a value for  $\alpha$  such that  $0 < \alpha < \beta$  holds. Subsequently, we denote the corresponding smallest  $\alpha$ -bound  $||\mathcal{H}_t^{\alpha}||_{\infty}$  as  $A_{\alpha}$ . As a result of this setup, and by integrating the insights from Lemma 1 and Lemma 2, we can further constrain the approximation error in the  $L^{\infty}$ -norm, which is articulated as

$$\|f_{P,\lambda} - f^*\|_{L^{\infty}(\nu)}^2 \le A_{\alpha}^2 \|f^*\|_{\beta}^2 \cdot \lambda^{\beta - \alpha}.$$
(4.8)

This step is crucial in effectively managing the estimation error, a task that we will delve into in Subsection 4.2. The ability to bound the approximation error in this manner plays a pivotal role in enhancing the precision and reliability of our overall analysis in the subsequent section.

Finally, we turn our attention to the effective dimension, represented by

$$N_{\nu}(\lambda) = \operatorname{tr}\left((T_{\nu} + \lambda)^{-1}T_{\nu}\right) = \sum_{k=1}^{\infty} \frac{pe^{-t\lambda_k}}{\lambda + pe^{-t\lambda_k}}.$$
(4.9)

The decay rate of this dimension as  $\lambda \to 0$  is a critical factor in our analysis of the estimation error. To establish an upper bound for the effective dimension, it is necessary to apply Weyl's law. This application is encapsulated in the following Lemma 3, which provides a means to control the growth rate of the eigenvalues  $\lambda_k$  of the Laplacian on the *m*-dimensional manifold  $\mathcal{M}$ . For a comprehensive understanding of Weyl's law, especially in the context of classical Riemannian manifolds, one can refer to [21, 6]. These references offer in-depth discussions and insights into the application and implications of Weyl's law, enriching the understanding of its role in controlling the eigenvalue growth and, consequently, in bounding the effective dimension in our analysis.

**Lemma 3.** For the eigenvalue  $\lambda_k$  of the Laplacian  $\Delta$  on a compact, connected Riemann manifold  $\mathcal{M}$  with dimension m, we have the following asymptotic estimation

$$C_{low}k^{\frac{2}{m}} \le \lambda_k \le C_{up}k^{\frac{2}{m}} \tag{4.10}$$

for all  $k \in \mathbb{N}$  with  $C_{low}$  and  $C_{up}$  two absolute constants depending only on  $\mathcal{M}$ .

Utilizing Weyl's law, as previously discussed, allows us to derive the necessary upper bound for the effective dimension  $N_{\nu}(\lambda)$  that is integral to our analysis. This upper bound is articulated in the following Lemma 4.

**Lemma 4.** Suppose that  $m \ge 2$ . Then, there exists a constant D such that, for all  $0 < \lambda < \frac{1}{2}$ , it holds

$$N_{\nu}(\lambda) \le D(\log \lambda^{-1})^{\frac{m}{2}}$$

In this context, the constant D only rely on  $\mathcal{M}$  and t.

*Proof.* Given that the function represented by  $t \mapsto t/(t+\lambda)$  is increasing in t, and by combining (4.9) with (4.10), we arrive at

$$N_{\nu}(\lambda) = \sum_{k=1}^{\infty} \frac{p e^{-t\lambda_k}}{\lambda + p e^{-t\lambda_k}} \le \sum_{k=1}^{\infty} \frac{p e^{-tC_{low}k^{\frac{2}{m}}}}{\lambda + p e^{-tC_{low}k^{\frac{2}{m}}}} = \sum_{k=1}^{\infty} \frac{1}{1 + \lambda p^{-1} e^{tC_{low}k^{\frac{2}{m}}}}.$$

Considering that the function  $x \mapsto 1/(1 + \lambda p^{-1} e^{tC_{low} x^{\frac{2}{m}}})$  is decreasing in x and remains positive, we can set an upper bound for the summation using an integral, expressed as

$$N_{\nu}(\lambda) \leq \int_{0}^{\infty} \frac{1}{1 + \lambda p^{-1} e^{tC_{low}x^{\frac{2}{m}}}} dx = \frac{m}{2} (C_{low}t)^{-\frac{m}{2}} \int_{0}^{\infty} \frac{u^{\frac{m}{2}-1}}{1 + \lambda p^{-1} e^{u}} du$$

In above equation, the remaining integral on the right-hand side represents the complete Fermi-Dirac integral. To evaluate this integral, we utilize the polylogarithm function

$$\operatorname{Li}_s(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^s}.$$

Drawing from [38], we can transform the complete Fermi-Dirac integral into an expression involving the polylogarithm function and Gamma function, resulting in

$$\int_0^\infty \frac{u^{\frac{m}{2}-1}}{1+p^{-1}\lambda e^u} du = -\text{Li}_{\frac{m}{2}}(-p\lambda^{-1}) \cdot \Gamma(\frac{m}{2}).$$

Given that the polylogarithm exhibits a specific limit behavior, as also referenced in [38], we deduce

$$\operatorname{Li}_{\frac{m}{2}}(-p\lambda^{-1}) \sim -\frac{1}{\Gamma(\frac{m}{2}+1)} \left(\log(p\lambda^{-1})\right)^{\frac{m}{2}}$$

as  $\lambda \to 0$ . Consequently, since  $\Gamma(s+1) = s\Gamma(s)$ , there exists a constant  $\lambda_0 \ll 1$  such that, for all  $0 < \lambda \leq \lambda_0$  it holds

$$N_{\nu}(\lambda) \leq 1.1 (C_{low} t)^{-\frac{m}{2}} \left( \log(p\lambda^{-1}) \right)^{\frac{m}{2}}$$
  
$$\leq 1.1 (C_{low} t)^{-\frac{m}{2}} \cdot 2^{\frac{m}{2}-1} \left( (\log\lambda^{-1})^{\frac{m}{2}} + (\log p)^{\frac{m}{2}} \right)$$
  
$$\leq 2^{\frac{m}{2}+1} (C_{low} t)^{-\frac{m}{2}} \cdot (\log\lambda^{-1})^{\frac{m}{2}}.$$

As for  $\lambda_0 < \lambda < 1/2$ , due to the decreasing property of  $N_{\nu}(\lambda)$ , we can find a constant  $D_0$ (which only depends on  $\mathcal{M}, t$ ) such that

$$N_{\nu}(\lambda) \le N_{\nu}(\lambda_0) \le D_0(\log 2)^{\frac{m}{2}} \le D_0(\log \lambda^{-1})^{\frac{m}{2}}.$$

Finally, taking  $D = \max\{2^{\frac{m}{2}+1}(C_{low}t)^{-\frac{m}{2}}, D_0\}$  yields the desired result and completes the proof.

#### 4.2 Bounding estimation error

Firstly, we will demonstrate that the  $\gamma$ -norm can be appropriately transformed into the  $\mathcal{H}_t$ -norm. This transformation is crucial as it allows for a more versatile and applicable analysis by bridging different norms.

**Lemma 5.** For all  $0 \leq \gamma \leq 1$ ,  $f \in \mathcal{H}_t$  we have

$$\|f\|_{\gamma} = \left\|T_{\nu}^{\frac{1-\gamma}{2}}f\right\|_{\mathcal{H}_{t}}.$$
(4.11)

Proof. Let's denote

$$f = \sum_{k=1}^{\infty} b_k p^{\frac{1}{2}} e^{-\frac{t\lambda_k}{2}} f_k$$

with  $b_k = \langle f, p^{\frac{1}{2}} e^{-\frac{t\lambda_k}{2}} f_k \rangle_{\mathcal{H}_t}$ . Given that  $\{p^{\frac{\gamma}{2}} e^{-\frac{\gamma t\lambda_k}{2}} f_k\}_{k \in \mathbb{N}}$  constitutes an orthonormal basis of  $\mathcal{H}_t^{\gamma}$ , we can apply the Parseval identity, leading us to

$$||f||_{\gamma}^{2} = \left\| \sum_{k=1}^{\infty} b_{k} p^{-\frac{\gamma-1}{2}} e^{(\gamma-1)\frac{t\lambda_{k}}{2}} \cdot p^{\frac{\gamma}{2}} e^{-\gamma\frac{t\lambda_{k}}{2}} f_{k} \right\|_{\gamma}^{2} = \sum_{k=1}^{\infty} b_{k}^{2} p^{1-\gamma} e^{(\gamma-1)t\lambda_{k}}.$$

Similarly, when we incorporate (2.4) into our calculations, it results in

$$\left\| T_{\nu}^{\frac{1-\gamma}{2}} f \right\|_{\mathcal{H}_{t}}^{2} = \left\| \sum_{k=1}^{\infty} p^{\frac{1-\gamma}{2}} e^{-t\lambda_{k}\frac{1-\gamma}{2}} b_{k} \cdot p^{\frac{1}{2}} e^{-\frac{t\lambda_{k}}{2}} f_{k} \right\|_{\mathcal{H}_{t}}^{2} = \sum_{k=1}^{\infty} b_{k}^{2} p^{1-\gamma} e^{(\gamma-1)t\lambda_{k}}$$

as desired. Then we complete the proof.

We are now ready to tackle the estimation error term. Starting with (4.11), we decompose the estimation error into three distinct terms, as outlined below:

$$\|f_{D,\lambda} - f_{P,\lambda}\|_{\gamma}^{2} = \left\|T_{\nu}^{\frac{1-\gamma}{2}}(f_{D,\lambda} - f_{P,\lambda})\right\|_{\mathcal{H}_{t}}^{2}$$

$$= \left\|T_{\nu}^{\frac{1-\gamma}{2}}(T_{\nu} + \lambda)^{-\frac{1}{2}}(T_{\nu} + \lambda)^{\frac{1}{2}}(T_{\delta} + \lambda)^{-\frac{1}{2}}(T_{\delta} + \lambda)^{\frac{1}{2}}(f_{D,\lambda} - f_{P,\lambda})\right\|_{\mathcal{H}_{t}}^{2}$$

$$\leq \left\|T_{\nu}^{\frac{1-\gamma}{2}}(T_{\nu} + \lambda)^{-\frac{1}{2}}\right\|_{\mathscr{B}(\mathcal{H}_{t})}^{2} \cdot \left\|(T_{\nu} + \lambda)^{\frac{1}{2}}(T_{\delta} + \lambda)^{-\frac{1}{2}}\right\|_{\mathscr{B}(\mathcal{H}_{t})}^{2}$$

$$(4.12a)$$

$$= \left\|(T_{\nu} + \lambda)^{\frac{1}{2}}(f_{D,\lambda} - f_{D,\lambda})\right\|^{2}$$

$$\cdot \left\| (T_{\delta} + \lambda)^{\frac{1}{2}} (f_{D,\lambda} - f_{P,\lambda}) \right\|_{\mathcal{H}_t}^2.$$
(4.12b)

We provide an explanation for the symbols adopted here. Let  $(\mathcal{H}, \langle \cdot, \cdot \rangle_{\mathcal{H}})$  and  $(\mathcal{H}', \langle \cdot, \cdot \rangle_{\mathcal{H}'})$ be two Hilbert spaces. Hereinafter, the collection of bounded linear operators from  $\mathcal{H}$  to  $\mathcal{H}'$  forms a Banach space, symbolized as  $\mathscr{B}(\mathcal{H}, \mathcal{H}')$ , when considered under operator norm  $\|A\|_{\mathscr{B}(\mathcal{H}, \mathcal{H}')} = \sup_{\|f\|_{\mathcal{H}}=1} \|Af\|_{\mathcal{H}'}$ . When  $\mathcal{H} = \mathcal{H}'$ , the space is then denoted by  $\mathscr{B}(\mathcal{H})$  with the corresponding norm given by  $\|A\|_{\mathscr{B}(\mathcal{H})}$ . Our approach will involve establishing upper bounds for these three terms in (4.12a) and (4.12b), which will be addressed separately.

Let us begin with the estimation of the first term in (4.12a). For this term, we utilize the spectral decomposition outlined in (2.4), combined with direct calculations on the function  $t \mapsto t^{1-\gamma}/(\lambda+t)$ . This process leads us to the conclusion that

$$\left\| T_{\nu}^{\frac{1-\gamma}{2}} (T_{\nu} + \lambda)^{-\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_{t})}^{2} = \sup_{k \ge 1} \frac{p^{1-\gamma} e^{-t\lambda_{k}(1-\gamma)}}{p e^{-t\lambda_{k}} + \lambda} \le \lambda^{-\gamma}, \tag{4.13}$$

which is valid for all  $0 \le \gamma \le 1$ . Moving on to the second term in (4.12a), we can assert

$$\left\| (T_{\nu} + \lambda)^{\frac{1}{2}} (T_{\delta} + \lambda)^{-\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_{t})}^{2} = \left\| (T_{\nu} + \lambda)^{\frac{1}{2}} (T_{\delta} + \lambda)^{-1} (T_{\nu} + \lambda)^{\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_{t})}.$$
(4.14)

This conclusion is derived from the fact that both  $T_{\nu}$  and  $T_{\delta}$  are self-adjoint operators. Additionally, it is important to note the transformation represented by

$$(T_{\nu} + \lambda)^{\frac{1}{2}} (T_{\delta} + \lambda)^{-1} (T_{\nu} + \lambda)^{\frac{1}{2}} = \left( (T_{\nu} + \lambda)^{-\frac{1}{2}} (T_{\delta} + \lambda) (T_{\nu} + \lambda)^{-\frac{1}{2}} \right)^{-1}$$
$$= \left( (T_{\nu} + \lambda)^{-\frac{1}{2}} (T_{\delta} - T_{\nu} + T_{\nu} + \lambda) (T_{\nu} + \lambda)^{-\frac{1}{2}} \right)^{-1}$$
$$= \left( (T_{\nu} + \lambda)^{-\frac{1}{2}} (T_{\delta} - T_{\nu}) (T_{\nu} + \lambda)^{-\frac{1}{2}} + I \right)^{-1}.$$

Combining with the Taylor series of function  $x \mapsto \frac{1}{1-x}$  at origin, we derive that

$$\begin{split} \left\| (T_{\nu} + \lambda)^{\frac{1}{2}} (T_{\delta} + \lambda)^{-1} (T_{\nu} + \lambda)^{\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_{t})} \\ &= \left\| \left( I - (T_{\nu} + \lambda)^{-\frac{1}{2}} (T_{\nu} - T_{\delta}) (T_{\nu} + \lambda)^{-\frac{1}{2}} \right)^{-1} \right\|_{\mathscr{B}(\mathcal{H}_{t})} \\ &= \sum_{k=0}^{\infty} \left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} (T_{\nu} - T_{\delta}) (T_{\nu} + \lambda)^{-\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_{t})}^{k}. \end{split}$$
(4.15)

Therefore, our analysis for the second term reduces to the estimation of the operator norm of the difference between  $T_{\nu}$  and its empirical counterpart  $T_{\delta}$ . We employ Lemma 6 to accomplish it. This lemma is grounded in Bernstein's type concentration inequality, which is applicable to random variables valued in Hilbert spaces. For a detailed discussion on this Bernstein's type inequality, one may refer to [31].

#### Lemma 6. Denote

$$p_{\lambda} = \log \left( 2eN_{\nu}(\lambda) \frac{\lambda + \|T_{\nu}\|}{\|T_{\nu}\|} \right).$$

Then, for all  $n \ge 1$ ,  $\lambda > 0$ , and  $\tau \ge 1$ , the following operator norm bound holds with probability not less than  $1 - 2e^{-\tau}$ :

$$\left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} (T_{\nu} - T_{\delta}) (T_{\nu} + \lambda)^{-\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_t)} \leq \frac{4A_{\alpha}^2 \tau p_{\lambda}}{3n\lambda^{\alpha}} + \sqrt{\frac{2A_{\alpha}^2 \tau p_{\lambda}}{n\lambda^{\alpha}}}.$$

The proof of Lemma 6 is detailed in [13]. Building upon the previous lemma, we consolidate our findings into Lemma 7 that provides an upper bound for the second term in (4.12a).

**Lemma 7.** Suppose that  $n \geq 8A_{\alpha}^2 \tau p_{\lambda} \lambda^{-\alpha}$ . Then, for all  $\tau \geq 1$  and  $\lambda > 0$ , the following operator norm bound holds with probability not less than  $1 - 2e^{-\tau}$ :

$$\left\| (T_{\nu} + \lambda)^{\frac{1}{2}} (T_{\delta} + \lambda)^{-\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_{t})}^{2} \leq 3.$$
(4.16)

*Proof.* Given that  $n \geq 8A_{\alpha}^2 \tau p_{\lambda} \lambda^{-\alpha}$ , from Lemma 6 we have

$$\left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} (T_{\nu} - T_{\delta}) (T_{\nu} + \lambda)^{-\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_t)} \leq \frac{4A_{\alpha}^2 \tau p_{\lambda}}{3n\lambda^{\alpha}} + \sqrt{\frac{2A_{\alpha}^2 \tau p_{\lambda}}{n\lambda^{\alpha}}} \leq \frac{2}{3}$$

Incorporating with our preceding calculations (4.14) and (4.15), it is straightforward to observe that, under our specified assumptions, the following holds true

$$\left\| (T_{\nu} + \lambda)^{\frac{1}{2}} (T_{\delta} + \lambda)^{-\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_{t})}^{2} = \sum_{k=0}^{\infty} \left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} (T_{\nu} - T_{\delta}) (T_{\nu} + \lambda)^{-\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_{t})}^{k} \le \sum_{k=0}^{\infty} \left( \frac{2}{3} \right)^{k} = 3,$$

which is the desired result. The proof is then finished.

Now, let us revisit the third term in (4.12b). We recall the expressions (2.10) and (4.1), given by

$$f_{D,\lambda} = g_{\lambda}(T_{\delta})g_D,$$
  
$$f_{P,\lambda} = g_{\lambda}(T_{\nu})g_P$$

with  $g_P = I_{\nu}^* f^*$ . Let  $h_{\lambda}(t) = 1 - tg_{\lambda}(t)$ . Given that  $h_{\lambda}(t) + tg_{\lambda}(t) = 1$  holds, we arrive at the following result

$$f_{D,\lambda} - f_{P,\lambda} = g_{\lambda}(T_{\delta})g_D - (h_{\lambda}(T_{\delta}) + T_{\delta}g_{\lambda}(T_{\delta})) f_{P,\lambda}$$
  
=  $g_{\lambda}(T_{\delta})(g_D - T_{\delta}f_{P,\lambda}) - h_{\lambda}(T_{\delta})f_{P,\lambda}.$ 

Consequently, this allows us to further decompose (4.12b) into two separate components:

$$\left\| (T_{\delta} + \lambda)^{\frac{1}{2}} (f_{D,\lambda} - f_{P,\lambda}) \right\|_{\mathcal{H}_{t}}^{2}$$

$$\leq 2 \left\| (T_{\delta} + \lambda)^{\frac{1}{2}} (g_{\lambda}(T_{\delta})(g_{D} - T_{\delta}f_{P,\lambda})) \right\|_{\mathcal{H}_{t}}^{2} + 2 \left\| (T_{\delta} + \lambda)^{\frac{1}{2}} (h_{\lambda}(T_{\delta})f_{P,\lambda}) \right\|_{\mathcal{H}_{t}}^{2}.$$

$$(4.17)$$

Our immediate focus will be on addressing the second term in (4.17). The result concerning the upper bound for this term is encapsulated in the subsequent Lemma 8.

**Lemma 8.** Suppose that  $\{g_{\lambda}\}_{\lambda>0}$  has a qualification  $\xi \geq \frac{1}{2}$ . Then, for all  $\lambda > 0$  and  $0 < \beta \leq 1$  it holds

$$\left| (T_{\delta} + \lambda)^{\frac{1}{2}} (h_{\lambda}(T_{\delta}) f_{P,\lambda}) \right|_{\mathcal{H}_{t}}^{2} \leq 4\lambda^{\beta} \|f^{*}\|_{\beta}^{2}.$$

$$(4.18)$$

*Proof.* Given that  $\{g_{\lambda}\}_{\lambda>0}$  possesses a qualification of  $\xi \geq \frac{1}{2}$ , we can apply (2.9) to obtain

$$\left\| (T_{\delta} + \lambda)^{\frac{1}{2}} h_{\lambda}(T_{\delta}) \right\|_{\mathscr{B}(\mathcal{H}_{t})}^{2} \leq \left( \sup_{0 \leq t \leq \kappa^{2}} (t + \lambda)^{\frac{1}{2}} h_{\lambda}(t) \right)^{2}$$
$$\leq \left( \sup_{0 \leq t \leq \kappa^{2}} (t^{\frac{1}{2}} + \lambda^{\frac{1}{2}}) h_{\lambda}(t) \right)^{2}$$
$$\leq \left( \lambda^{\frac{1}{2}} + \lambda^{\frac{1}{2}} \right)^{2} = 4\lambda.$$
(4.19)

Here, the second inequality is derived from  $(a + b)^p \leq a^p + b^p$ , applicable whenever  $0 \leq p \leq 1$  holds. Regarding the remaining  $\mathcal{H}_t$ -norm of  $f_{P,\lambda}$ , we refer to the expansion (4.4), given by

$$f_{P,\lambda} = \sum_{k=1}^{\infty} g_{\lambda}(pe^{-t\lambda_k}) p^{\frac{1}{2}} e^{-\frac{t\lambda_k}{2}} a_k \cdot p^{\frac{1}{2}} e^{-\frac{t\lambda_k}{2}} f_k.$$

This leads us to

$$\begin{aligned} \|f_{P,\lambda}\|_{\mathcal{H}_{t}}^{2} &= \left\|\sum_{k=1}^{\infty} g_{\lambda}(pe^{-t\lambda_{k}})p^{\frac{1}{2}}e^{-\frac{t\lambda_{k}}{2}}a_{k} \cdot p^{\frac{1}{2}}e^{-\frac{t\lambda_{k}}{2}}f_{k}\right\|_{\mathcal{H}_{t}}^{2} \\ &= \sum_{k=1}^{\infty} \left(g_{\lambda}(pe^{-t\lambda_{k}})p^{\frac{1}{2}}e^{-\frac{t\lambda_{k}}{2}}\right)^{2}a_{k}^{2} \\ &= \sum_{k=1}^{\infty} g_{\lambda}(pe^{-t\lambda_{k}})^{1-\beta} \cdot g_{\lambda}(pe^{-t\lambda_{k}})^{\beta+1}p^{\beta+1}e^{-(\beta+1)t\lambda_{k}} \cdot p^{-\beta}e^{\beta t\lambda_{k}}a_{k}^{2} \\ &\leq \sum_{k=1}^{\infty} \lambda^{\beta-1} \cdot p^{-\beta}e^{\beta t\lambda_{k}}a_{k}^{2} \\ &= \lambda^{\beta-1}\|f^{*}\|_{\beta}^{2}. \end{aligned}$$
(4.20)

In this context, the inequality is a result of the properties (2.8) associated with the regularization family. By synthesizing (4.19) with (4.20), we can achieve the desired result and complete the proof.

Now, we shift our focus to the first term in (4.17). To analyze this term, we will use the following decomposition:

$$\begin{aligned} \left\| (T_{\delta} + \lambda)^{\frac{1}{2}} \left( g_{\lambda}(T_{\delta}) (g_D - T_{\delta} f_{P,\lambda}) \right) \right\|_{\mathcal{H}_{t}}^{2} \\ &= \left\| (T_{\delta} + \lambda)^{\frac{1}{2}} g_{\lambda}(T_{\delta}) (T_{\delta} + \lambda)^{\frac{1}{2}} \cdot (T_{\delta} + \lambda)^{-\frac{1}{2}} (T_{\nu} + \lambda)^{\frac{1}{2}} \cdot (T_{\nu} + \lambda)^{-\frac{1}{2}} (g_D - T_{\delta} f_{P,\lambda}) \right\|_{\mathcal{H}_{t}}^{2} \\ &\leq \left\| (T_{\delta} + \lambda)^{\frac{1}{2}} g_{\lambda}(T_{\delta}) (T_{\delta} + \lambda)^{\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_{t})}^{2} \cdot \left\| (T_{\delta} + \lambda)^{-\frac{1}{2}} (T_{\nu} + \lambda)^{\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_{t})}^{2} \tag{4.21a}$$

$$\cdot \left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} \left( g_D - T_{\delta} f_{P,\lambda} \right) \right\|_{\mathcal{H}_t}^2.$$
(4.21b)

In addressing the first two terms in (4.21a), we will leverage the properties (2.8) of the regularization family and the upper bound (4.16). These enable us to establish the following estimates, given by

$$\left\| (T_{\delta} + \lambda)^{\frac{1}{2}} g_{\lambda}(T_{\delta}) (T_{\delta} + \lambda)^{\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_{t})}^{2} = \left\| (T_{\delta} + \lambda) g_{\lambda}(T_{\delta}) \right\|_{\mathscr{B}(\mathcal{H}_{t})}^{2}$$
$$\leq \left( \sup_{0 \le t \le \kappa^{2}} (t + \lambda) g_{\lambda}(t) \right)^{2} \le 4$$
(4.22)

and

$$\left\| (T_{\delta} + \lambda)^{-\frac{1}{2}} (T_{\nu} + \lambda)^{\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_{t})}^{2} = \left\| (T_{\nu} + \lambda)^{\frac{1}{2}} (T_{\delta} + \lambda)^{-1} (T_{\nu} + \lambda)^{\frac{1}{2}} \right\|_{\mathscr{B}(\mathcal{H}_{t})} \le 3.$$
(4.23)

Moving on, we address the final term in (4.21b). To establish an upper bound for this term, we will rely on Lemma 9 that offers control over the error resulting from sampling. This kind of lemma is commonly found in statistical literature and is typically derived through the application of concentration inequalities. This lemma comes specifically from [13].

Lemma 9. Denote

$$L_{\lambda} = \max\{L, \|f_{P,\lambda} - f^*\|_{L^{\infty}(\nu)}\}$$

Then, for all  $0 < \alpha \leq 1$ ,  $\lambda > 0$ ,  $\tau \geq 1$ , and  $n \geq 1$ , the following operator norm bound holds with probability not less than  $1 - 2e^{-\tau}$ :

$$\left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} \left( (g_D - T_{\delta} f_{P,\lambda}) - (g_P - T_{\nu} f_{P,\lambda}) \right) \right\|_{\mathcal{H}_t}^2$$
  
$$\leq \frac{64\tau^2}{n} \left( \sigma^2 N_{\nu}(\lambda) + A_{\alpha}^2 \frac{\|f^* - f_{P,\lambda}\|_{L^2(\nu)}^2}{\lambda^{\alpha}} + 2A_{\alpha}^2 \frac{L_{\lambda}^2}{n\lambda^{\alpha}} \right)$$

Now, we will present our upper bound for the final term in (4.21b) as detailed in the following Lemma 10.

**Lemma 10.** For all  $0 < \alpha \le 1$ ,  $0 < \beta \le 2\xi$ ,  $\lambda > 0$ ,  $\tau \ge 1$ , and  $n \ge 1$ , the following operator norm bound holds with probability not less than  $1 - 2e^{-\tau}$ :

$$\left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} (g_D - T_{\delta} f_{P,\lambda}) \right\|_{\mathcal{H}_t}^2$$
  
$$\leq \frac{128\tau^2}{n} \left( \sigma^2 N_{\nu}(\lambda) + A_{\alpha}^2 \|f^*\|_{\beta}^2 \lambda^{\beta-\alpha} + 2A_{\alpha}^2 \frac{L_{\lambda}^2}{n\lambda^{\alpha}} \right) + 2\|f^*\|_{\beta}^2 \lambda^{\beta}.$$
(4.24)

*Proof.* Initially, we establish

$$\begin{split} \left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} (g_D - T_{\delta} f_{P,\lambda}) \right\|_{\mathcal{H}_t}^2 \\ &= \left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} ((g_D - T_{\delta} f_{P,\lambda}) - (g_P - T_{\nu} f_{P,\lambda})) + (T_{\nu} + \lambda)^{-\frac{1}{2}} (g_P - T_{\nu} f_{P,\lambda}) \right\|_{\mathcal{H}_t}^2 \\ &\leq 2 \left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} ((g_D - T_{\delta} f_{P,\lambda}) - (g_P - T_{\nu} f_{P,\lambda})) \right\|_{\mathcal{H}_t}^2 \tag{4.25a} \\ &+ 2 \left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} (g_P - T_{\nu} f_{P,\lambda}) \right\|_{\mathcal{H}_t}^2. \tag{4.25b}$$

For (4.25a), by applying Lemma 9, we can derive

$$\left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} \left( (g_D - T_{\delta} f_{P,\lambda}) - (g_P - T_{\nu} f_{P,\lambda}) \right) \right\|_{\mathcal{H}_t}^2$$
  
$$\leq \frac{64\tau^2}{n} \left( \sigma^2 N_{\nu}(\lambda) + A_{\alpha}^2 \frac{\|f^* - f_{P,\lambda}\|_{L^2(\nu)}^2}{\lambda^{\alpha}} + 2A_{\alpha}^2 \frac{L_{\lambda}^2}{n\lambda^{\alpha}} \right).$$
(4.26)

This result is obtained with a probability of not less than  $1 - 2e^{-\tau}$ . When considering (4.25b), noting that  $g_P = I_{\nu}^* f^*$  and  $T_{\nu} = I_{\nu}^* I_{\nu}$ , we arrive at

$$\left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} (g_P - T_{\nu} f_{P,\lambda}) \right\|_{\mathcal{H}_t}^2 = \left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} (I_{\nu}^* f^* - I_{\nu}^* I_{\nu} f_{P,\lambda}) \right\|_{\mathcal{H}_t}^2$$
$$= \left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} I_{\nu}^* (f^* - f_{P,\lambda}) \right\|_{\mathcal{H}_t}^2.$$

On one hand, we have

$$\left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} I_{\nu}^* \right\|_{\mathscr{B}(L^2(\nu),\mathcal{H}_t)}^2 = \sup_{k \ge 0} \frac{p e^{-t\lambda_k}}{\lambda + p e^{-t\lambda_k}} \le 1.$$

On the other hand, by integrating our previous approximation error result from Lemma 1 with  $\gamma = 0$ , we obtain

$$\|f^* - f_{P,\lambda}\|_{L^2(\nu)}^2 \le \|f^*\|_{\beta}^2 \lambda^{\beta}.$$
(4.27)

From these analyses, we can conclude an upper bound for (4.25b) as

$$\left\| (T_{\nu} + \lambda)^{-\frac{1}{2}} (g_P - T_{\nu} f_{P,\lambda}) \right\|_{\mathcal{H}_t}^2 \le \|f^*\|_{\beta}^2 \lambda^{\beta}.$$
(4.28)

By substituting (4.27) back into (4.26) and integrating it with (4.28), we are able to achieve the desired result and complete the proof.  $\Box$ 

Finally, by merging (4.13), (4.16), (4.18), (4.22), (4.23), and (4.24) we are able to summarize the estimation error bound as follows.

**Lemma 11.** Suppose that  $n \ge 8A_{\alpha}^2 \tau p_{\lambda} \lambda^{-\alpha}$ . Then, for all  $0 < \alpha \le 1$ ,  $0 \le \gamma \le 1$ ,  $\gamma < \beta \le 1$ ,  $\xi \ge \frac{1}{2}$ ,  $\tau \ge 1$ , and  $\lambda > 0$ , with probability not less than  $1 - 4e^{-\tau}$ , the following estimation error bound holds:

$$\|f_{P,\lambda} - f_{D,\lambda}\|_{\gamma}^{2} \leq \frac{9216\tau^{2}}{n\lambda^{\gamma}} \left( \sigma^{2} N_{\nu}(\lambda) + A_{\alpha}^{2} \|f^{*}\|_{\beta}^{2} \lambda^{\beta-\alpha} + 2A_{\alpha}^{2} \frac{L_{\lambda}^{2}}{n\lambda^{\alpha}} \right) + 168 \|f^{*}\|_{\beta}^{2} \lambda^{\beta-\gamma}.$$
(4.29)

### 4.3 Proof of Theorem 1

At this juncture, we are fully prepared to prove Theorem 1.

Proof of Theorem 1. Given the choice of  $\lambda_n$  as

$$\lambda_n \sim \left(\frac{(\log n)^{\frac{m}{2}}}{n}\right)^{\frac{1}{\beta}},$$

we first consider the condition on n proposed in Lemma 11, which relates specifically to the derivation of the sample lower bound. Remember that from our notations, we have

$$p_{\lambda} = \log\left(2eN_{\nu}(\lambda)\frac{\lambda + \|T_{\nu}\|}{\|T_{\nu}\|}\right)$$

Therefore, by applying Lemma 4, we can derive that there exists an  $n_0$  such that, for all  $n \ge n_0$ , the following inequalities hold true:

$$\begin{split} 8A_{\alpha}^{2}\tau p_{\lambda_{n}}\lambda_{n}^{-\alpha} &= 8A_{\alpha}^{2}\tau \log\left(2eN_{\nu}(\lambda_{n})\frac{\lambda_{n}+\|T_{\nu}\|}{\|T_{\nu}\|}\right)\lambda_{n}^{-\alpha} \\ &\leq 8A_{\alpha}^{2}\tau \log\left(2.2eD(\log\lambda_{n}^{-1})\frac{m}{2}\right)\lambda_{n}^{-\alpha} \\ &\leq 16A_{\alpha}^{2}\tau \log\left((\log\lambda_{n}^{-1})\frac{m}{2}\right)\lambda_{n}^{-\alpha} \\ &= 8mA_{\alpha}^{2}\tau \log(\log\lambda_{n}^{-1})\lambda_{n}^{-\alpha}. \end{split}$$

Notice that

$$\log(\log \lambda_n^{-1})\lambda_n^{-\alpha} \sim \log\left(\frac{1}{\beta}(\log n - \frac{m}{2}\log\log n)\right)(\log n)^{-\frac{\alpha m}{2\beta}} \cdot n^{\frac{\alpha}{\beta}} \lesssim n^{\frac{\alpha}{\beta}}.$$

By fixing an  $\alpha$  that satisfies condition  $0 < \alpha < \beta$ , there exists another  $n_1 \ge n_0$  such that, for all  $n \ge n_1$ , it holds

$$n > 8mA_{\alpha}^2 \tau \log(\log \lambda_n^{-1})\lambda_n^{-\alpha}.$$

Therefore, we can utilize (4.29) for  $n \ge n_1$  to derive

$$\|f_{P,\lambda_{n}} - f_{D,\lambda_{n}}\|_{\gamma}^{2} \leq \frac{9216\tau^{2}}{n\lambda_{n}^{\gamma}} \left(\sigma^{2}N_{\nu}(\lambda_{n}) + A_{\alpha}^{2}\|f^{*}\|_{\beta}^{2}\lambda_{n}^{\beta-\alpha} + 2A_{\alpha}^{2}\frac{L_{\lambda}^{2}}{n\lambda_{n}^{\alpha}}\right) + 168\|f^{*}\|_{\beta}^{2}\lambda_{n}^{\beta-\gamma}.$$
(4.30)

Moreover, the sample lower bound  $n_1$  (and  $n_0$ ) only depends on constants  $m, D, ||T_{\nu}||$ , and parameters  $\alpha, \beta, \tau$ .

Without any loss of generality, we assume that  $\lambda_n \leq 0.1$  for  $n \geq n_1$ . Regarding the third term in the bracket of (4.30), combining (4.8) with Assumption 2 we obtain

$$A_{\alpha}^{2} \frac{L_{\lambda_{n}}^{2}}{n\lambda_{n}^{\alpha}} = \frac{A_{\alpha}^{2}}{n\lambda_{n}^{\alpha}} \max\{L^{2}, \|f_{P,\lambda_{n}} - f^{*}\|_{L^{\infty}(\nu)}^{2}\}$$
$$\leq \frac{A_{\alpha}^{2}}{n\lambda_{n}^{\alpha}} \max\{L^{2}, A_{\alpha}^{2}R^{2}\lambda_{n}^{\beta-\alpha}\} \leq \frac{K_{1}}{n\lambda_{n}^{\alpha}}$$

with  $K_1 = \max\{A_{\alpha}^2 L^2, A_{\alpha}^4 R^2\}$ . Taking use of Lemma 4 and Assumption 2, we can derive that

$$\|f_{P,\lambda_n} - f_{D,\lambda_n}\|_{\gamma}^2 \le \frac{9216\tau^2}{n\lambda_n^{\gamma}} \left( D\sigma^2 (\log \lambda_n^{-1})^{\frac{m}{2}} + A_\alpha^2 R^2 \lambda_n^{\beta-\alpha} + \frac{2K_1}{n\lambda_n^{\alpha}} \right) + 168R^2 \lambda_n^{\beta-\gamma}$$
(4.31)

Now we turn our attention to those terms in the bracket of (4.31). Recalling the choice of  $\lambda_n$ , we have

$$\begin{aligned} (\log \lambda_n^{-1})^{\frac{m}{2}} + \lambda_n^{\beta-\alpha} + \frac{1}{n\lambda_n^{\alpha}} &\sim \left(\frac{1}{\beta}(\log n - \frac{m}{2}\log\log n)\right)^{\frac{m}{2}} + (\log n)^{\frac{m}{2}(1-\frac{\alpha}{\beta})}n^{\frac{\alpha}{\beta}-1} \\ &+ (\log n)^{-\frac{\alpha m}{2\beta}}n^{\frac{\alpha}{\beta}-1} \\ &\lesssim \left(\frac{1}{\beta}(\log n - \frac{m}{2}\log\log n)\right)^{\frac{m}{2}} &\sim (\log \lambda_n^{-1})^{\frac{m}{2}}. \end{aligned}$$

Therefore, we can choose  $K_2 = 27648 \max\{D\sigma^2, A_{\alpha}^2 R^2, 2K_1\}$  such that, for all  $n \ge n_1$ , we have

$$\|f_{P,\lambda_n} - f_{D,\lambda_n}\|_{\gamma}^2 \le K_2 \tau^2 \frac{(\log \lambda_n^{-1})^{\frac{m}{2}}}{n\lambda_n^{\gamma}} + 168R^2 \lambda_n^{\beta-\gamma}$$
(4.32)

Finally, by combining (4.32) with the approximation error bound (4.6) derived from Lemma 1, we arrive at our targeted  $\gamma$ -norm error bound as follows

$$\begin{aligned} \|f_{D,\lambda_n} - f^*\|_{\gamma}^2 &\leq K_2 \tau^2 \frac{\left(\log \lambda_n^{-1}\right)^{\frac{m}{2}}}{n\lambda_n^{\gamma}} + 169R^2 \lambda_n^{\beta-\gamma} \\ &\leq \tau^2 \lambda_n^{\beta-\gamma} \left(K_2 \frac{\left(\log \lambda_n^{-1}\right)^{\frac{m}{2}}}{n\lambda_n^{\beta}} + 169R^2\right). \end{aligned}$$

A direct computation indicates that

$$\frac{(\log \lambda_n^{-1})^{\frac{m}{2}}}{n\lambda_n^\beta} \sim \frac{\left(\frac{1}{\beta} (\log n - \frac{m}{2} \log \log n)\right)^{\frac{m}{2}}}{(\log n)^{\frac{m}{2}}} \leq \left(\frac{1}{\beta}\right)^{\frac{m}{2}}$$

Consequently, we finally obtain an upper bound as

$$\|f_{D,\lambda_n} - f^*\|_{\gamma}^2 \lesssim \tau^2 \lambda_n^{\beta-\gamma} \le C\tau^2 \left(\frac{(\log n)^{\frac{m}{2}}}{n}\right)^{\frac{\beta-\gamma}{\beta}},$$

which valids for all  $n \ge n_1$ . Here, the constant C only depends on constants  $m, D, \sigma, R, L$ , and parameters  $\alpha, \beta$ . The proof is then completed.

## 5 Lower Bound Analysis

In this section, we follow the approach presented in [35] to derive the minimax lower bound. This methodology involves leveraging Lemma 12 from [35], which is instrumental in our analysis. In our notations, the Kullback–Leibler divergence between two probability measures  $\rho_1$  and  $\rho_2$  on  $(\Omega, \mathcal{F})$ , denoted by  $D_{KL}(\rho_1 \parallel \rho_2)$ , is defined as

$$D_{KL}(\rho_1 \parallel \rho_2) = \int_{\Omega} \log\left(\frac{\mathrm{d}\rho_1}{\mathrm{d}\rho_2}\right) \mathrm{d}\rho_1,$$

provided that  $\rho_1$  is absolutely continuous with respect to  $\rho_2$ , and  $D_{KL}(\rho_1 \parallel \rho_2) = \infty$  otherwise.

**Lemma 12.** Suppose that there is a non-parametric class of functions  $\Theta$  and a semi-distance  $d(\cdot, \cdot)$  on  $\Theta$ ,  $\{P_{\theta} : \theta \in \Theta\}$  is a family of probability distributions indexed by  $\Theta$ . Assume that  $K \geq 2$  and  $\Theta$  contains elements  $\theta_0, \theta_1, \cdots, \theta_K$  such that

- (1)  $\forall 0 \le i < j \le K$ ,  $d(\theta_i, \theta_j) \ge 2s > 0.$  (5.1)
- (2)  $P_j \ll P_0, \ \forall j = 1, ..., K$  and

$$\frac{1}{K} \sum_{j=1}^{K} D_{KL}(P_j || P_0) \le a \log K$$
(5.2)

with  $0 < a < \frac{1}{8}$  and  $P_j = P_{\theta_j}, \ j = 0, 1, \dots, K$ .

Then it holds

$$\inf_{\hat{\theta}} \sup_{\theta \in \Theta} P_{\theta} \left( d(\hat{\theta}, \theta) \ge s \right) \ge \frac{\sqrt{K}}{1 + \sqrt{K}} \left( 1 - 2a - \sqrt{\frac{2a}{\log K}} \right).$$

Our current objective is to construct a family of probability distributions with elements  $P_0, P_1, \ldots, P_K$ , which adhere to the Kullback-Leibler divergence condition outlined in (5.2). Let  $\bar{\sigma} = \min\{\sigma, L\}$  be given. For any measurable function  $f: \mathcal{M} \to \mathbb{R}$  and for any  $x \in \mathcal{M}$ , we define the conditional distribution given x as  $P_f(\cdot|x) = \mathcal{N}(f(x), \bar{\sigma}^2)$ . This distribution is a normal distribution with a mean of f(x) and a variance of  $\bar{\sigma}^2$ . Further, we construct a probability distribution  $P_f$  related to f on  $\mathcal{M} \times \mathbb{R}$ . This distribution is designed such that its marginal distribution on  $\mathcal{M}$  is  $\nu$ , and its conditional distribution given  $x \in \mathcal{M}$  on  $\mathbb{R}$  is  $P_f(\cdot|x)$ . In practical terms, this means that we sample x from  $\nu$  and set  $y = f(x) + \varepsilon$ , where  $\varepsilon \sim \mathcal{N}(0, \bar{\sigma}^2)$  represents an independent Gaussian random error. It can be straightforwardly verified that for all  $f \in L^2(\nu)$ , the condition  $|P_f|_2^2 = ||f||_{L^2(\nu)}^2 + \bar{\sigma}^2 < \infty$  is satisfied. Furthermore, the conditional mean function of this distribution, which corresponds to the regression function  $f^*$ , is identical to f. It is important to note that probability distributions of this nature inherently satisfy the moment condition (3.1) outlined in Assumption 1, as specified in Lemma 13. This compliance is deduced from a straightforward computation involving the high-order moments of Gaussian variables. For a more detailed explanation of this calculation, we refer to [13]. This construction lays the foundation for our analysis, ensuring that the distributions we work with are both theoretically sound and aligned with the assumptions underpinning our approach.

**Lemma 13.** For a measurable function  $f : \mathcal{M} \to \mathbb{R}$  with its corresponding probability measure  $P_f$  defined above, the moment condition (3.1) is satisfied for  $\sigma = L = \bar{\sigma}$ .

Given that  $f^* = f$  holds for the probability distribution  $P_f$ , we can focus on constructing a specific family of functions. This family, comprising elements  $f_0, f_1, \ldots, f_K$ , must satisfy the source condition (3.2) as stipulated in Assumption 2. To develop these function elements, we define

$$f_{i} = \epsilon^{\frac{1}{2}} \sum_{j=1}^{k} \omega_{j}^{(i)} p^{\frac{\gamma}{2}} e^{-\frac{\gamma}{2}\lambda_{k+j}t} f_{k+j}$$
(5.3)

for  $0 \leq i \leq K$ , where  $\omega^{(i)} = (\omega_1^{(i)}, \dots, \omega_k^{(i)}) \in \{0, 1\}^k$  are binary strings and  $\epsilon, k$  are to be determined later. Our approach, which hinges on the use of binary strings, needs to fulfill the separation condition (5.1) as stated in Lemma 12. Ensuring that a sufficient number of binary strings  $\omega$  exists, with significant separation between them, is crucial for our analysis. To substantiate the presence of these adequately distanced binary strings, we will invoke Lemma 14, commonly referred to as the Gilbert-Varshamov Bound. This lemma is a wellknown result in coding theory and provides a means to guarantee the existence of binary strings with the required properties. A detailed discussion of this lemma and its implications also can be found in [35]. This construction strategy is pivotal as it forms the basis of our function family, ensuring that it adheres to the necessary conditions and assumptions for our analysis to be valid.

**Lemma 14.** For  $k \ge 8$ , there exists  $K \ge 2^{\frac{k}{8}}$  and binary strings  $w^{(0)}, \ldots, \omega^{(K)} \in \{0, 1\}^m$  with  $\omega^{(0)} = (0, \ldots, 0), \ \omega^{(j)} = (\omega_1^{(j)}, \ldots, \omega_k^{(j)})$  such that

$$\sum_{i=1}^{k} \left(\omega_i^{(l)} - \omega_i^{(j)}\right)^2 \ge \frac{k}{8}$$
(5.4)

for all  $0 \leq l \neq j \leq K$ .

In order to meet the Kullback-Leibler divergence condition specified in (5.2) as per Lemma 12, it's necessary to perform direct computations pertaining to the Gaussian-type probability distribution we have constructed. These calculations, as outlined in Lemma 15, are crucial for ensuring that the probability distribution aligns with the required divergence condition, which is a fundamental aspect of establishing the minimax lower bound. For a more in-depth understanding and additional context, one can refer to [3].

**Lemma 15.** For  $f, f' \in L^2(\nu)$  and  $n \ge 1$ , the corresponding probability distribution  $P_f, P_{f'}$  defined above satisfy

$$D_{KL}(P_f^n || P_{f'}^n) = \frac{n}{2\bar{\sigma}^2} || f - f' ||_{L^2(\nu)}^2.$$
(5.5)

Having completed all the preparatory works, we now proceed to furnish the proof for Theorem 2.

Proof of Theorem 2. Our strategic approach involves a careful selection of  $\epsilon, k$ , aimed at bounding the  $\beta$ -norm of our constructed  $f_i$  from (5.3). This is crucial to ensure that the source condition (3.2) is satisfied. Simultaneously, we will also focus on bounding the Kullback-Leibler divergence of the corresponding probability distributions to adhere to the condition (5.2). Once these bounds are established and both the source condition and KL-divergence requirements are met, we will then turn to Lemma 12. Utilizing this lemma, we will derive the desired minimax lower bound. This culmination of steps represents a comprehensive method, bringing together various facets of our analysis to ultimately reach the sought-after minimax lower bound result. This careful and methodical approach is essential for the robustness and validity of our proof of Theorem 2.

Setting:

$$f_i = \epsilon^{\frac{1}{2}} \sum_{j=1}^k \omega_j^{(i)} p^{\frac{\gamma}{2}} e^{-\frac{\gamma}{2}\lambda_{k+j}t} f_{k+j}, \quad i = 0, 1, \dots, K$$

with binary strings  $\omega^{(i)}$  and a certain  $K \ge 2^{\frac{k}{8}}$  derived from Lemma 14, we define  $\epsilon = C_{\epsilon} n^{-s}$ and  $k = C_k (\log n)^{\frac{m}{2}}$  with constants  $s, C_{\epsilon}, C_k$  to be determined later. Given  $\gamma < \beta$ , and applying (4.10), we establish

$$\begin{aligned} \|f_i\|_{\beta}^2 &= \epsilon \sum_{j=1}^k p^{\gamma-\beta} e^{(\beta-\gamma)\lambda_{k+j}t} \left(\omega_j^{(i)}\right)^2 \\ &\leq \epsilon \sum_{j=1}^k p^{\gamma-\beta} e^{(\beta-\gamma)tC_{up}(k+j)\frac{2}{m}} \\ &\leq \epsilon k p^{\gamma-\beta} e^{(\beta-\gamma)tC_{up}(2k)\frac{2}{m}} \\ &= p^{\gamma-\beta} C_{\epsilon} C_k n^{-s} (\log n)^{\frac{m}{2}} n^{(\beta-\gamma)tC_{up}(2C_k)\frac{2}{m}}. \end{aligned}$$
(5.6)

To satisfy the source condition (3.2), we initially need to set

$$(\beta - \gamma)tC_{up}(2C_k)^{\frac{2}{m}} - s < 0.$$
(5.7)

Then, utilizing (5.5) from Lemma 15 and combining it with (4.10), we can determine

$$D_{KL}(P_{f_i}^n || P_{f_0}^n) = \frac{n}{2\bar{\sigma}^2} || f_i ||_{L^2(\nu)}^2$$
$$= \frac{n\epsilon}{2\bar{\sigma}^2} \sum_{j=1}^k p^{\gamma} e^{-\gamma \lambda_{k+j} t} \left( \omega_j^{(i)} \right)^2$$
$$\leq \frac{n\epsilon}{2\bar{\sigma}^2} k p^{\gamma} e^{-\gamma t C_{low} k^{\frac{2}{m}}}.$$

Recalling  $K \ge 2^{\frac{k}{8}}$  and fixing  $a \in (0, \frac{1}{8})$ , the Kullback-Leibler divergence condition (5.2) results in

$$D_{KL}(P_{f_i}^n || P_{f_0}^n) \le \frac{n\epsilon}{2\bar{\sigma}^2} k p^{\gamma} e^{-\gamma t C_{low} k^{\frac{2}{m}}} \le a \frac{\log 2}{8} k \le a \log K.$$

In essence, this implies

$$C_{\epsilon} n^{1-s} n^{-\gamma t C_{low} C_k^{\frac{2}{m}}} \le a p^{-\gamma} \frac{\bar{\sigma}^2 \log 2}{4}.$$
(5.8)

Thus, we should set

$$(1-s) - \gamma t C_{low} C_k^{\frac{2}{m}} \le 0.$$
(5.9)

When we merge (5.7) with (5.9), specifically for the case where  $\gamma = 0$ , we can let s = 1 and choose  $C_k$  such that

$$C_k^{\frac{2}{m}} < \frac{1}{\beta t C_{up} 2^{\frac{2}{m}}}$$

In the general scenario where  $\gamma > 0$ , we are limited to setting

$$s > \frac{(\beta - \gamma)tC_{up}2^{\frac{2}{m}}}{(\beta - \gamma)tC_{up}2^{\frac{2}{m}} + \gamma tC_{low}} \ge \frac{(\beta - \gamma)2^{\frac{2}{m}}}{(\beta - \gamma)2^{\frac{2}{m}} + \gamma}$$

and selecting  $C_k$  so that

$$\frac{1-s}{\gamma t C_{low}} \le C_k^{\frac{2}{m}} < \frac{s}{(\beta-\gamma)t C_{up} 2^{\frac{2}{m}}}.$$

Moreover, as our constructed  $f_i$  needs to satisfy both (5.6) and (5.8), we also need to pick a  $C_{\epsilon}$  that fulfills

$$C_{\epsilon} \leq \min\left\{ap^{-\gamma}\frac{\bar{\sigma}^2\log 2}{4}, p^{\beta-\gamma}\frac{R^2}{C_k}\right\}.$$

Then, for large n, both the source condition (3.2) and the KL-divergence condition (5.2) are satisfied. Now we consider the probability distribution family  $\{P_f^n : \|f\|_{\beta}^2 \leq R^2\}$  indexed by f, endowed with a distance defined by  $d(f_i, f_j) = \|f_i - f_j\|_{\gamma}$ . Applying the Gilbert-Varshamov bound (5.4) to this family yields

$$d(f_i, f_j)^2 = \|f_i - f_j\|_{\gamma}^2 = \epsilon \sum_{l=1}^k \left(\omega_l^{(i)} - \omega_l^{(j)}\right)^2 \ge \epsilon \frac{k}{8} = Ca \frac{(\log n)^{\frac{m}{2}}}{n^s},$$

where the constant C is independent of n, a. Finally, by applying Lemma 12, we obtain

$$\inf_{\hat{f}} \sup_{f^*} P_{f^*}^n \left( \|\hat{f} - f^*\|_{\gamma}^2 \ge C' a \frac{(\log n)^{\frac{m}{2}}}{n^s} \right) \ge \frac{\sqrt{K}}{1 + \sqrt{K}} \left( 1 - 2a - \sqrt{\frac{2a}{\log K}} \right)$$
(5.10)

with  $C' = \frac{1}{4}C$ . For sufficiently large n (and correspondingly large K), the right-hand side of (5.10) exceeds 1 - 3a. Therefore, for all  $\tau \in (0, 1)$  and any estimator  $\hat{f}$ , we can identify a function  $f^*$  meeting the source condition (3.2). Moreover, its corresponding probability distribution  $P_{f^*}$ , as defined above, will satisfy

$$\|\hat{f} - f^*\|_{\gamma}^2 \ge C'' \tau^2 \frac{(\log n)^{\frac{m}{2}}}{n^s}$$

with a probability of at least  $1 - \tau^2$ . In this context,  $C'' = \frac{1}{12}C$ . This confirmation substantiates our proposed result. The proof is finished.

# References

- Frank Bauer, Sergei Pereverzev, and Lorenzo Rosasco. On regularization algorithms in learning theory. *Journal of complexity*, 23(1):52–72, 2007.
- [2] Mikhail Belkin and Partha Niyogi. Laplacian eigenmaps for dimensionality reduction and data representation. *Neural computation*, 15(6):1373–1396, 2003.
- [3] Gilles Blanchard and Nicole Mücke. Optimal rates for regularization of statistical inverse learning problems. *Foundations of Computational Mathematics*, 18:971–1013, 2018.
- [4] Andrea Caponnetto and Ernesto De Vito. Optimal rates for the regularized least-squares algorithm. *Foundations of Computational Mathematics*, 7:331–368, 2007.
- [5] Alain Celisse and Martin Wahl. Analyzing the discrepancy principle for kernelized spectral filter learning algorithms. The Journal of Machine Learning Research, 22(1):3498– 3556, 2021.
- [6] Isaac Chavel. Eigenvalues in Riemannian geometry. Academic press, 1984.
- [7] Xue Cheng and Tai-Ho Wang. Bessel bridge representation for the heat kernel in hyperbolic space. Proceedings of the American Mathematical Society, 146(4):1781–1792, 2018.
- [8] Ronald R Coifman and Stéphane Lafon. Diffusion maps. Applied and computational harmonic analysis, 21(1):5–30, 2006.
- [9] Felipe Cucker and Ding Xuan Zhou. *Learning theory: an approximation theory viewpoint*, volume 24. Cambridge University Press, 2007.
- [10] Ernesto De Vito, Nicole Mücke, and Lorenzo Rosasco. Reproducing kernel hilbert spaces on manifolds: Sobolev and diffusion spaces. *Analysis and Applications*, 19(03):363–396, 2021.
- [11] Lee H Dicker, Dean P Foster, and Daniel Hsu. Kernel ridge vs. principal component regression: Minimax bounds and the qualification of regularization operators. *Electronic Journal of Statistics*, 11:1022–1047, 2017.
- [12] Manfredo P Do Carmo. Differential geometry of curves and surfaces: revised and updated second edition. Courier Dover Publications, 2016.
- [13] Simon Fischer and Ingo Steinwart. Sobolev norm learning rates for regularized least-squares algorithms. *The Journal of Machine Learning Research*, 21(1):8464–8501, 2020.
- [14] L Lo Gerfo, Lorenzo Rosasco, Francesca Odone, E De Vito, and Alessandro Verri. Spectral algorithms for supervised learning. *Neural Computation*, 20(7):1873–1897, 2008.
- [15] Rajarshi Guhaniyogi and David B Dunson. Compressed gaussian process for manifold regression. The Journal of Machine Learning Research, 17(1):2472–2497, 2016.
- [16] Zheng-Chu Guo, Shao-Bo Lin, and Ding-Xuan Zhou. Learning theory of distributed spectral algorithms. *Inverse Problems*, 33(7):074009, 2017.

- [17] Zheng-Chu Guo and Lei Shi. Optimal rates for coefficient-based regularized regression. Applied and Computational Harmonic Analysis, 47(3):662–701, 2019.
- [18] Thomas Hamm and Ingo Steinwart. Adaptive learning rates for support vector machines working on data with low intrinsic dimension. *The Annals of Statistics*, 49(6):3153–3180, 2021.
- [19] Elton P Hsu. Stochastic analysis on manifolds. Number 38. American Mathematical Soc., 2002.
- [20] John M Lee. Introduction to Riemannian manifolds, volume 2. Springer, 2018.
- [21] Peter Li and Shing-Tung Yau. On the schrödinger equation and the eigenvalue problem. Communications in Mathematical Physics, 88(3):309–318, 1983.
- [22] Junhong Lin and Volkan Cevher. Optimal convergence for distributed learning with stochastic gradient methods and spectral algorithms. *The Journal of Machine Learning Research*, 21(1):5852–5914, 2020.
- [23] Shao-Bo Lin and Ding-Xuan Zhou. Distributed kernel-based gradient descent algorithms. Constructive Approximation, 47(2):249–276, 2018.
- [24] Jiading Liu and Lei Shi. Statistical optimality of divide and conquer kernel-based functional linear regression. arXiv preprint arXiv:2211.10968, 2022.
- [25] Andrew McRae, Justin Romberg, and Mark Davenport. Sample complexity and effective dimension for regression on manifolds. Advances in Neural Information Processing Systems, 33:12993–13004, 2020.
- [26] Nicole Mücke and Gilles Blanchard. Parallelizing spectrally regularized kernel algorithms. The Journal of Machine Learning Research, 19(30):1–29, 2018.
- [27] Loucas Pillaud-Vivien, Alessandro Rudi, and Francis Bach. Statistical optimality of stochastic gradient descent on hard learning problems through multiple passes. Advances in Neural Information Processing Systems, 31, 2018.
- [28] Potluri Rao. Some notes on misspecification in multiple regressions. *The American Statistician*, 25(5):37–39, 1971.
- [29] Sam T Roweis and Lawrence K Saul. Nonlinear dimensionality reduction by locally linear embedding. *Science*, 290(5500):2323–2326, 2000.
- [30] Lei Shi. Distributed learning with indefinite kernels. Analysis and Applications, 17(06):947–975, 2019.
- [31] Steve Smale and Ding-Xuan Zhou. Learning theory estimates via integral operators and their approximations. *Constructive approximation*, 26(2):153–172, 2007.
- [32] Ingo Steinwart and Andreas Christmann. Support vector machines. Springer Science & Business Media, 2008.
- [33] Ingo Steinwart and Clint Scovel. Mercer's theorem on general domains: On the interaction between measures, kernels, and rkhss. *Constructive Approximation*, 35:363–417, 2012.

- [34] Hans Triebel. *Theory of Function Spaces II*. Modern Birkhäuser Classics. Birkhäuser Basel, 1992.
- [35] Alexandre B. Tsybakov. Introduction to Nonparametric Estimation. Springer series in statistics. Springer, 2009.
- [36] Laurens Van der Maaten and Geoffrey Hinton. Visualizing data using t-sne. The Journal of machine learning research, 9(11), 2008.
- [37] Martin J Wainwright. High-dimensional statistics: A non-asymptotic viewpoint, volume 48. Cambridge university press, 2019.
- [38] David C Wood. The computation of polylogarithms. Technical report. University of Kent, Canterbury, UK, 1992.
- [39] Yuan Yao, Lorenzo Rosasco, and Andrea Caponnetto. On early stopping in gradient descent learning. *Constructive Approximation*, 26(2):289–315, 2007.
- [40] Haobo Zhang, Yicheng Li, and Qian Lin. On the optimality of misspecified spectral algorithms. arXiv preprint arXiv:2303.14942, 2023.