

HOMOCLINIC CLASSES OF GEODESIC FLOWS ON RANK 1 MANIFOLDS

YURI LIMA AND MAURICIO POLETTI

ABSTRACT. We prove that the homoclinic class of every hyperbolic periodic orbit of a geodesic flow over a C^∞ closed rank 1 Riemannian manifold equals the unit tangent bundle. As an application, we give a proof using symbolic dynamics of the theorem of Knieper on the uniqueness of the measure of maximal entropy [Kni98] and theorems of Burns et al on the uniqueness of equilibrium states [BCFT18].

1. INTRODUCTION

Homoclinic classes are an efficient way of decomposing the dynamics of hyperbolic systems into smaller pieces. Introduced by Newhouse in [New72] as generalizations of the basic sets introduced by Smale in his spectral decomposition theorem [Sma67], homoclinic classes have been used in more complicated settings beyond uniform hyperbolicity, see e.g. [BC04].

Rodriguez Hertz et al introduced ergodic homoclinic classes of hyperbolic periodic points, and studied the uniqueness of SRB measures for surface diffeomorphisms [RHRHTU11]. Buzzi, Crovisier and Sarig introduced homoclinic classes of hyperbolic measures and proved that each class carries at most one measure of maximal entropy [BCS22]. Later, Buzzi, Crovisier and Lima studied homoclinic classes of measures for three-dimensional flows with positive speed [BCL23].

In this article, we consider homoclinic classes for geodesic flows over rank 1 manifolds, in any dimension. Let M be a C^∞ closed Riemannian manifold with nonpositive sectional curvature, and let $\varphi = \{\varphi^t\}_{t \in \mathbb{R}} : T^1M \rightarrow T^1M$ be its geodesic flow. We say that M is *rank 1* if there is a vector $x \in T^1M$ without a parallel Jacobi field perpendicular to the flow direction. This assumption implies various geometrical and dynamical properties of φ , see Section 2.1. Let $\text{HC}(\mathcal{O})$ denote the homoclinic class of the periodic orbit \mathcal{O} , see Section 2.2 for the definition.

Main Theorem. *Let φ be the geodesic flow over a C^∞ closed rank 1 Riemannian manifold. If \mathcal{O} is a hyperbolic periodic orbit, then $\text{HC}(\mathcal{O}) = T^1M$.*

Returning to the case of diffeomorphisms, Buzzi, Crovisier and Sarig proved that the constructions of countable Markov partitions of Sarig [Sar13] and Ben Ovadia [BO18] code homoclinic classes by *irreducible* countable topological Markov shifts,

Date: March 7, 2024.

2020 Mathematics Subject Classification. 37B10, 37C05, 37C83, 37D25, 37D35.

Key words and phrases. geodesic flow, Markov partition, Pesin theory, symbolic dynamics.

We thank Katrin Gelfert, Davi Obata, and Sergio Rom ana for fruitful conversations. MP was supported by CNPq and Instituto Serrapilheira, grant number Serra-R-2211-41879. YL was supported by CNPq and Instituto Serrapilheira, grant “Jangada Din amica: Impulsionando Sistemas Din amicos na Regi o Nordeste”.

and that each homoclinic class supports at most one equilibrium state for each admissible potential, see Theorem 3.1 and Corollary 3.3 in [BCS22]. Employing the same methods, we provide applications of the Main Theorem to give proofs, for rank 1 manifolds, of the theorem of Knieper on the uniqueness of the measure of maximal entropy [Kni98] and theorems of Burns et al on the uniqueness of equilibrium states for some classes of potentials [BCFT18]. Let ψ^u denote the geometric potential of φ and let Sing be the *singular set* of φ , see Section 2.1 for the definitions.

Theorem 1.1 (Theorem A of [BCFT18]). *Let M be a C^∞ closed rank 1 Riemannian manifold, and let $\psi : T^1M \rightarrow \mathbb{R}$ be Hölder continuous or of the form $\psi = q\psi^u$ for $q \in \mathbb{R}$. If $P(\text{Sing}, \psi) < P(\psi)$, then ψ has a unique equilibrium state μ . This measure is hyperbolic and fully supported.*

Above, $P(\psi)$ and $P(\text{Sing}, \psi)$ denote the topological pressure and topological pressure restricted to Sing respectively. We note that Theorem 1.1 is not the full statement proved in [BCFT18], since we do not characterize μ as the weak- $*$ limit of hyperbolic periodic orbits. We also remark that μ is Bernoulli, due to [LLS16] in dimension two and to [CT22, ALP20] in any dimension.

For multiples of the geometric potential in surfaces, we also recover part of Theorem C of [BCFT18].

Theorem 1.2 (Theorem C of [BCFT18]). *If M is a C^∞ closed rank 1 surface, then its geodesic flow φ has a unique equilibrium state μ_q for the potential $q\psi^u$ for each $q \in (-\infty, 1)$. This measure is hyperbolic and fully supported.*

Again, we are not able to characterize μ as the weak- $*$ limit of hyperbolic periodic orbits. Finally, using the pressure gap $P(\text{Sing}, 0) < P(0)$ proved in [BCFT18, Theorem B], we are also able to recover Knieper's theorem on the uniqueness of the measure of maximal entropy.

Theorem 1.3 ([Kni98]). *If M is a C^∞ closed rank 1 Riemannian manifold, then its geodesic flow φ has a unique measure of maximal entropy. This measure is hyperbolic and fully supported.*

Let us stress that, in dimension two, the pressure gap is automatic, hence we provide a self-contained proof of Knieper's theorem in dimension two, using homoclinic classes and symbolic dynamics.

2. PRELIMINARIES

2.1. Geodesic flows in nonpositive curvature. Let M be a C^∞ closed Riemannian manifold and T^1M be its unit tangent bundle, which is also a closed Riemannian manifold. Let $\varphi = \{\varphi^t\}_{t \in \mathbb{R}} : T^1M \rightarrow T^1M$ be the geodesic flow on M , and X be the vector field generating this flow. We assume that M is a *rank 1* manifold: it has nonpositive sectional curvature and there is a vector without parallel Jacobi fields perpendicular to the flow direction.

REGULAR AND SINGULAR SETS: The *regular set* of φ is defined by

$$\text{Reg} = \{x \in T^1M : \exists t \in \mathbb{R} \text{ s.t. } K(\pi) < 0 \text{ for every plane } \pi \subset T_{\gamma_x(t)}M\},$$

where γ_x is the geodesic starting at x . The *singular set* of φ is defined by

$$\text{Sing} = T^1M \setminus \text{Reg} = \{x \in T^1M : \forall t \in \mathbb{R}, \exists \pi \subset T_{\gamma_x(t)}M \text{ plane s.t. } K(\pi) = 0\}.$$

Clearly, Reg and Sing form a partition of T^1M , with Reg open and Sing closed. Geodesic flows on rank 1 manifolds have (weak) invariant directions and manifolds at every point. More specifically, there are continuous $d\varphi$ -invariant bundles $x \in T^1M \rightarrow E_x^{s/u}$ and continuous foliations $\{\widehat{W}^{ss/uu}(x) : x \in T^1M\}$ tangent to $E^{s/u}$ and invariant under φ such that:

- (1) E^s, E^u are orthogonal to X and $\dim(E^s) = \dim(E^u) = \dim(M) - 1$.
- (2) $x \in \text{Reg}$ if and only if $E_x^s \oplus \langle X_x \rangle \oplus E_x^u = T_x(T^1M)$.
- (3) Reg is dense in T^1M .
- (4) $\widehat{W}^{s/u}(x) := \bigcup_{t \in \mathbb{R}} \widehat{W}^{ss/uu}(\varphi^t(x))$ is a φ -invariant connected manifold of dimension $\dim(M)$ and tangent to $E^{s/u} \oplus \langle X \rangle$.
- (5) There is a universal constant $C = C(M)$ such that if d_x is the induced distance on $\widehat{W}^s(x)$ then $d_x(\varphi^t(x), \varphi^t(y)) \leq C d_x(x, y)$ for all $y \in \widehat{W}^s(x)$ and all $t \geq 0$; a similar statement holds for $\widehat{W}^u(x)$ and $t \leq 0$.
- (6) $\widehat{W}^{s/u}(x)$ is dense in T^1M for every $x \in T^1M$.

See [Ebe01] for proofs of (1)–(5), and [Bal82] for a proof of (6). For an orbit \mathcal{O} , define $\widehat{W}^{s/u}(\mathcal{O}) := \widehat{W}^{s/u}(x)$ for any $x \in \mathcal{O}$. We also recall the definition of the geometric potential.

THE GEOMETRIC POTENTIAL OF φ [BR75]: The *geometric potential* of φ is the function $\psi^u : T^1M \rightarrow \mathbb{R}$ defined as

$$\psi^u(x) = - \left. \frac{d}{dt} \right|_{t=0} \log \det(d\varphi_x^t|_{E_x^u}) = - \lim_{t \rightarrow 0} \frac{1}{t} \log \det(d\varphi_x^t|_{E_x^u}).$$

2.2. Homoclinic classes and symbolic dynamics. In this section, we combine results of [LS19, ALP20, BCS22, BCL23] on codings of hyperbolic measures. Let $\mathcal{G} = (V, E)$ be an oriented graph, where V, E are the vertex and edge sets. We denote edges by $v \rightarrow w$, and assume that V is countable.

TOPOLOGICAL MARKOV SHIFT (TMS): It is a pair (Σ, σ) where

$$\Sigma := \{\mathbb{Z}\text{-indexed paths on } \mathcal{G}\} = \{v = \{v_n\}_{n \in \mathbb{Z}} \in V^{\mathbb{Z}} : v_n \rightarrow v_{n+1}, \forall n \in \mathbb{Z}\}$$

is the symbolic space and $\sigma : \Sigma \rightarrow \Sigma$, $[\sigma(v)]_n = v_{n+1}$, is the *left shift*. We endow Σ with the distance $d(v, w) := \exp[-\inf\{|n| \in \mathbb{Z} : v_n \neq w_n\}]$. The *regular set* of Σ is

$$\Sigma^\# := \left\{ v \in \Sigma : \exists v, w \in V \text{ s.t. } \begin{array}{l} v_n = v \text{ for infinitely many } n > 0 \\ v_n = w \text{ for infinitely many } n < 0 \end{array} \right\}.$$

We will sometimes omit σ from the definition, referring to Σ as a TMS. We only consider TMS that are *locally compact*, i.e. for all $v \in V$ the number of ingoing edges $u \rightarrow v$ and outgoing edges $v \rightarrow w$ is finite. Although we use the same notation, the regular set of φ and the regular set of Σ are not related; we maintain the notation because they are classical in their contexts.

Given (Σ, σ) a TMS, let $r : \Sigma \rightarrow (0, +\infty)$ be a continuous function. For $n \geq 0$, let $r_n = r + r \circ \sigma + \dots + r \circ \sigma^{n-1}$ be the n -th *Birkhoff sum* of r , and extend this definition for $n < 0$ in the unique way such that the *cocycle identity* holds: $r_{m+n} = r_m + r_n \circ \sigma^m$, $\forall m, n \in \mathbb{Z}$.

TOPOLOGICAL MARKOV FLOW (TMF): The TMF defined by (Σ, σ) and *roof function* r is the pair (Σ_r, σ_r) where $\Sigma_r := \{(v, t) : v \in \Sigma, 0 \leq t < r(v)\}$ and $\sigma_r : \Sigma_r \rightarrow \Sigma_r$ is the flow on Σ_r given by $\sigma_r^t(v, t') = (\sigma^n(v), t' + t - r_n(v))$, where n is the unique integer such that $r_n(v) \leq t' + t < r_{n+1}(v)$. We endow Σ_r with a natural

metric $d_r(\cdot, \cdot)$, called the *Bowen-Walters metric*, such that σ_r is a continuous flow [BW72]. The *regular set* of (Σ_r, σ_r) is $\Sigma_r^\# = \{(\underline{v}, t) \in \Sigma_r : \underline{v} \in \Sigma^\#\}$.

Similarly, we will sometimes omit σ_r and refer to Σ_r as a TMF. The roof functions we consider will always be Hölder continuous, in which case $\exists \kappa, C > 0$ such that $d_r(\sigma_r^t(z), \sigma_r^t(z')) \leq Cd_r(z, z')^\kappa$ for all $|t| \leq 1$ and $z, z' \in \Sigma_r$, see [LS19, Lemma 5.8].

IRREDUCIBLE COMPONENT: If Σ is a TMS defined by an oriented graph $\mathcal{G} = (V, E)$, its *irreducible components* are the subshifts $\Sigma' \subset \Sigma$ defined over maximal subsets $V' \subset V$ satisfying the following condition:

$$\forall v, w \in V', \exists \underline{v} \in \Sigma \text{ and } n \geq 1 \text{ such that } v_0 = v \text{ and } v_n = w.$$

An *irreducible component* Σ'_r of Σ_r is a set of the form $\Sigma'_r = \{(\underline{v}, t) \in \Sigma_r : \underline{v} \in \Sigma'\}$ where Σ' is an irreducible component of Σ .

χ -HYPERBOLIC MEASURE: A φ -invariant probability measure μ on T^1M is χ -*hyperbolic* if for μ -a.e. $x \in T^1M$ all the Lyapunov exponents are greater than χ in absolute value, except for the zero exponent in the flow direction.

Let μ be a hyperbolic φ -invariant probability measure. For μ -a.e. $x \in T^1M$,

$$W^{ss}(x) = \left\{ y \in T^1M : \limsup_{t \rightarrow +\infty} \frac{1}{t} \log d(\varphi^t(x), \varphi^t(y)) < 0 \right\}$$

denotes the *strong stable manifold* of x and

$$W^s(x) = \bigcup_{t \in \mathbb{R}} \varphi^t[W^{ss}(x)]$$

denotes the *stable manifold* of x . We define similarly $W^{uu}(x)$ and $W^u(x)$ the *strong unstable* and *unstable manifolds* of x .

Given a hyperbolic periodic orbit \mathcal{O} , we let $W^{s/u}(\mathcal{O}) = W^{s/u}(x)$ denote the stable/unstable manifold of \mathcal{O} , for any $x \in \mathcal{O}$.

HOMOCLINIC CLASS OF HYPERBOLIC PERIODIC ORBIT: The *homoclinic class* of a hyperbolic periodic orbit \mathcal{O} is the set

$$\text{HC}(\mathcal{O}) = \overline{W^u(\mathcal{O}) \pitchfork W^s(\mathcal{O})}.$$

To simplify the notation, we will sometimes write $N := T^1M$.

HOMOCLINIC RELATION OF MEASURES [BCS22, BCL23]: We say that two ergodic hyperbolic measures μ, ν are *homoclinically related* if for μ -a.e. x and ν -a.e. y there exist transverse intersections $W^s(x) \pitchfork W^u(y) \neq \emptyset$ and $W^u(x) \pitchfork W^s(y) \neq \emptyset$, i.e. points $z_1 \in W^s(x) \cap W^u(y)$ and $z_2 \in W^u(x) \cap W^s(y)$ such that $T_{z_1}N = T_{z_1}W^s(x) + T_{z_1}W^u(y)$ and $T_{z_2}N = T_{z_2}W^u(x) + T_{z_2}W^s(y)$.

The homoclinic relation is an equivalence relation among ergodic hyperbolic measures, see [BCL23, Prop. 10.1]. Next, we state the coding theorem for non-uniformly hyperbolic flows that we will use to prove Theorems 1.1, 1.2, and 1.3. This theorem mixes the versions presented in [ALP20, Theorem 10.1] and [BCL23, Theorem 1.1].

Theorem 2.1. *Let X be a $C^{1+\beta}$ vector field with $X \neq 0$ everywhere, let $\varphi : N \rightarrow N$ be the flow generated by X , and let $\chi > 0$. If μ_1, μ_2 are homoclinically related χ -hyperbolic probability measures, then there is an irreducible topological Markov flow (Σ_r, σ_r) and a Hölder continuous map $\pi_r : \Sigma_r \rightarrow N$ s.t.:*

- (1) $r : \Sigma \rightarrow \mathbb{R}^+$ is Hölder continuous and bounded away from zero and infinity.
- (2) $\pi_r \circ \sigma_r^t = \varphi^t \circ \pi_r$ for all $t \in \mathbb{R}$.
- (3) $\pi_r[\Sigma_r^\#]$ has full measure with respect to μ_1 and μ_2 .
- (4) Every $x \in N$ has finitely many pre-images in $\Sigma_r^\#$.

Proof. If M has dimension two, then we can apply [BCL23, Theorem 1.1] directly. Since this latter theorem is not available in higher dimension, we have to argue differently. Let $\mathcal{O}_1, \mathcal{O}_2, \dots$ be the χ -hyperbolic periodic orbits homoclinically related to μ_1 (and hence to μ_2).

CLAIM: There is a global Poincaré section Λ such that if $f : \Lambda \rightarrow \Lambda$ is the Poincaré return map and \mathcal{S} is the *singular set* of f , then:

- (1) Λ is *adapted* for μ_1 and μ_2 , i.e.

$$\lim_{|t| \rightarrow \infty} \frac{1}{t} \log d(\varphi^t(x), \mathcal{S}) = 0 \quad (2.1)$$

for μ_i -a.e. $x \in \Lambda$, $i = 1, 2$.¹

- (2) For every n , there exists a compact, φ -invariant, transitive, locally maximal, χ -hyperbolic set K_n that contains $\mathcal{O}_1, \dots, \mathcal{O}_n$ and such that $K_n \cap \mathcal{S} = \emptyset$.

The singular set is $\mathcal{S} = \{x \in \Lambda : \{x, f(x), f^{-1}(x)\} \cap \partial\Lambda \neq \emptyset\}$.

Proof of the claim. Fix $x_i \in \mathcal{O}_i$ and $z_{ij} \in W^u(x_i) \pitchfork W^s(x_j)$, and let $\tau_{ij} \in \mathbb{R}$ such that $z_{ij} \in W^{uu}(x_i) \cap W^{ss}(\varphi^{\tau_{ij}}(x_j))$. We construct a one-parameter family of global Poincaré sections Λ_r , where r varies in an interval $[a, b]$ and each Λ_r is the union of finitely many codimension 1 disjoint balls $D_1(r), \dots, D_k(r) \subset N$ centered at points $y_1, \dots, y_k \in N$, each of them with radius r and almost orthogonal to X . The details of the construction can be found in [LS19, Section 2] for three-dimensional flows and in [ALP20, Section 10] for any dimension. We can assume that the roof functions of the Poincaré return maps to Λ_r are all larger than some $\varepsilon_0 > 0$. As the radius r varies, the boundary of Λ_r varies as well. Applying a double counting argument and the Borel-Cantelli lemma, for Lebesgue almost every choice $r \in [a, b]$ the section Λ_r satisfies (1) above, see [LS19, Theorem 2.8] for details. Hence we focus on showing that (2) holds in a set of parameters $r \in [a, b]$ of positive measure. We will prove this using parameter selection. Observe that if K is φ -invariant, then $K \cap \mathcal{S} = \emptyset$ if and only if $K \cap \partial\Lambda_r = \emptyset$.

Write $z_{ii} = x_i$, and let \mathcal{O}_{ij} denote the orbit of z_{ij} . Consider the countable union of orbits $\bigcup_{i,j} \mathcal{O}_{ij}$. Clearly, the set of parameters $r \in [a, b]$ such that $(\bigcup_{i,j} \mathcal{O}_{ij}) \cap \partial\Lambda_r \neq \emptyset$ is countable. Below, we consider r in the complement of this set. We will use the following notation: given two functions $g(\delta), h(\delta)$, write $g = \mathcal{O}_n(h)$ when there is a constant $C > 0$ that depends on n and $\delta_0 > 0$ such that $|g(\delta)| \leq C|h(\delta)|$ for all $|\delta| < \delta_0$.

Fix n and $\delta > 0$. In the sequel, we construct a δ -neighborhood of $\bigcup_{1 \leq i, j \leq n} \mathcal{O}_{ij}$. Let $C = C(n) > 0$ such that

$$\begin{aligned} d(\varphi^{-t}(x_i), \varphi^{-t}(z_{ij})) &\leq Ce^{-xt} \\ d(\varphi^{t+\tau_{ij}}(x_j), \varphi^t(z_{ij})) &\leq Ce^{-xt} \quad , \quad \forall 1 \leq i, j \leq n, \forall t \geq 0. \end{aligned}$$

¹This is equivalent to the projection of μ_i to Λ being adapted in the notation of [LS19].

Let $t_{ij} = O_n(|\log \delta|)$ positive such that $\varphi^{-t_{ij}}(z_{ij}) \in B_\delta(\mathcal{O}_i)$ and $\varphi^{t_{ij}}(z_{ij}) \in B_\delta(\mathcal{O}_j)$. The set $Z_n = \bigcup_i \mathcal{O}_i \cup \bigcup\{\varphi^t(z_{ij}) : 1 \leq i, j \leq n, |t| \leq t_{ij}\}$ is the union of finitely many pieces of orbits and $\bigcup_{1 \leq i, j \leq n} \mathcal{O}_{ij} \subset B_\delta(Z_n)$. Arguing as in [BCS22, Lemma 3.11], we can construct inside $B_\delta(Z_n)$ a compact, φ -invariant, transitive, locally maximal, χ -hyperbolic set that contains $\mathcal{O}_1, \dots, \mathcal{O}_n$. Thus, we estimate the measure of the set of $r \in [a, b]$ such that $B_\delta(Z_n) \cap \partial\Lambda_r \neq \emptyset$.

Denoting by $|\mathcal{O}_i|$ the length of \mathcal{O}_i , the total length of Z_n is

$$\sum_{1 \leq i \leq n} |\mathcal{O}_i| + \sum_{1 \leq i, j \leq n} 2t_{ij} = O_n(|\log \delta|).$$

Recalling that ε_0 is a lower bound for the roof function of Λ_r , the set $Z_n \cap \Lambda_r$ has at most $\frac{1}{\varepsilon_0} O_n(|\log \delta|) = O_n(|\log \delta|)$ elements. Then the intersection $B_\delta(Z_n) \cap \Lambda_b$ is contained in $O_n(|\log \delta|)$ balls of radius 2δ . The set of parameters $\text{Bad}(n) = \{r \in [a, b] : B_\delta(Z_n) \cap \partial\Lambda_r \neq \emptyset\}$ is thus contained in the union of $O_n(|\log \delta|)$ intervals of length 4δ , and so has Lebesgue measure $O_n(\delta |\log \delta|)$. Since $\lim_{\delta \rightarrow 0} \delta \log \delta = 0$, we can take δ_n such that $\text{Leb}[\text{Bad}(n)] < \varepsilon/n^2$ for small $\varepsilon > 0$ and so the complement $[a, b] \setminus \bigcup_{n \geq 1} \text{Bad}(n)$ has positive Lebesgue measure.

In summary, we can choose $r \in [a, b]$ such that:

- Condition (2.1) is satisfied: the space of $r \in [a, b]$ satisfying it has full measure.
- $(\bigcup_{i,j} \mathcal{O}_{ij}) \cap \partial\Lambda_r = \emptyset$: the space of $r \in [a, b]$ satisfying it is the complement of a countable set, hence has full measure.
- $B_{\delta_n}(Z_n) \cap \partial\Lambda_r = \emptyset$ for all $n \geq 1$: the space of $r \in [a, b]$ satisfying it has positive measure.

This concludes the proof of the claim. \square

Once the section $\Lambda = \Lambda_r$ is chosen, apply [ALP20, Theorem 10.1] to construct a TMF (Σ_r, σ_r) and a Hölder continuous map $\pi_r : \Sigma_r \rightarrow N$ satisfying (1)–(4) in Theorem 2.1, with the exception that Σ_r might not be irreducible. Since $K_n \cap \mathcal{S} = \emptyset$ and K_n is compact and χ -hyperbolic, [ALP20] implies that $K_n \subset \pi_r[\Sigma_r]$. Since K_n is uniformly hyperbolic, we actually have $K_n \subset \pi_r[\Sigma_r^\#]$.

The final step is to find an irreducible component Σ'_r of Σ_r that lifts both μ_1 and μ_2 . For that, we proceed as in [BCL23, Lemma 10.4]:

- For each $n \geq 1$, there is an invariant, compact, transitive set $X_n \subset \Sigma_r^\#$ that lifts K_n .
- Since \mathcal{O}_1 has finitely many lifts in $\Sigma_r^\#$, the sequence X_1, X_2, \dots has a subsequence X_{n_1}, X_{n_2}, \dots containing a same lift (\underline{v}, t) of \mathcal{O}_1 . Take Σ'_r to be the irreducible component of (\underline{v}, t) . Then \mathcal{O}_1, \dots all lift to periodic orbits in Σ'_r .

Now proceed as in [BCL23, Theorem 1.1] to lift generic points for μ_1 and μ_2 to Σ'_r . This completes the proof of the theorem. \square

3. PROOFS

Proof of Main Theorem. Let \mathcal{O} be a hyperbolic periodic orbit.

CLAIM: $W^{s/u}(\mathcal{O}) = \widehat{W}^{s/u}(\mathcal{O})$.

Proof of the claim. We prove the claim for $W^s(\mathcal{O})$ (the other proof is analogous). We first show that $W^s(\mathcal{O}) \subset \widehat{W}^s(\mathcal{O})$. Fix $\varepsilon > 0$ small enough so that

$$W_\varepsilon^s(\mathcal{O}) = \{y \in T^1M : d(\varphi^t(y), \mathcal{O}) < \varepsilon, \forall t \geq 0\} \subset W^s(\mathcal{O}).$$

Let $\widehat{W}_\delta^s(\mathcal{O}) \subset \widehat{W}^s(\mathcal{O})$ be the δ -neighborhood of \mathcal{O} in the d_x metric, for (any) $x \in \mathcal{O}$. If $y \in \widehat{W}_{\varepsilon/C}^s(\mathcal{O})$ with $d_x(y, x) < \varepsilon/C$ where $x \in \mathcal{O}$, then by property (5) of Section 2.1 we have

$$d(\varphi^t(y), \mathcal{O}) \leq d_x(\varphi^t(y), \varphi^t(x)) \leq Cd_x(y, x) < \varepsilon$$

for all $t \geq 0$, hence $\widehat{W}_{\varepsilon/C}^s(\mathcal{O}) \subset W_\varepsilon^s(\mathcal{O})$. Since $\widehat{W}_{\varepsilon/C}^s(\mathcal{O})$ and $W_\varepsilon^s(\mathcal{O})$ both have dimension $\dim(M)$, we conclude that $W_\delta^s(\mathcal{O}) \subset \widehat{W}_{\varepsilon/C}^s(\mathcal{O})$ for small $\delta > 0$. Therefore $W_\delta^s(\mathcal{O}) \subset \widehat{W}^s(\mathcal{O})$ and by the invariance of \widehat{W}^s the same occurs for $W^s(\mathcal{O}) = \bigcup_{t \geq 0} \varphi^{-t}[W_\delta^s(\mathcal{O})]$.

Now we prove that $W^s(\mathcal{O}) = \widehat{W}^s(\mathcal{O})$. Since $\widehat{W}^s(\mathcal{O})$ is connected (see e.g. [BP07, Prop. 12.2.4]), it is enough to show that $W^s(\mathcal{O})$ is open and dense in the induced topology of $\widehat{W}^s(\mathcal{O})$. Recall that $\varepsilon > 0$ is small such that $W^s(\mathcal{O}) = \bigcup_{t \geq 0} \varphi^{-t}[W_\varepsilon^s(\mathcal{O})]$, where $W_\varepsilon^s(\mathcal{O}) = \{y \in T^1M : d(\varphi^t(y), \mathcal{O}) < \varepsilon, \forall t \geq 0\}$. Since $W_\varepsilon^s(\mathcal{O})$ is open in the induced topology of $\widehat{W}^s(\mathcal{O})$, the same holds for $W^s(\mathcal{O})$. To prove that $W^s(\mathcal{O})$ is closed in the induced topology of $\widehat{W}^s(\mathcal{O})$, let $z \in \widehat{W}^s(\mathcal{O})$ and $z_k \in W^s(\mathcal{O})$ such that $d_x(z_k, z) \rightarrow 0$, for (any) $x \in \mathcal{O}$. We wish to show that $z \in W^s(\mathcal{O})$. By the triangle inequality and property (5) in Section 2.1, we have

$$\begin{aligned} d(\varphi^t z, \mathcal{O}) &\leq d(\varphi^t z, \varphi^t z_k) + d(\varphi^t z_k, \mathcal{O}) \\ &\leq d_x(\varphi^t z, \varphi^t z_k) + d(\varphi^t z_k, \mathcal{O}) \leq Cd_x(z, z_k) + d(\varphi^t z_k, \mathcal{O}). \end{aligned}$$

Let k such that $d_x(z_k, z) < \varepsilon/2C$, and then let $T_0 > 0$ such that $d(\varphi^t z_k, \mathcal{O}) < \varepsilon/2$ for all $t \geq T_0$. Hence $d(\varphi^t z, \mathcal{O}) < \varepsilon$ for all $t \geq T_0$, and so $\varphi^{T_0} z \in W_\varepsilon^s(\mathcal{O})$. This proves that $z \in W^s(\mathcal{O})$, as wished. \square

By the claim and property (6) of Section 2.1, we conclude that $W^{s/u}(\mathcal{O}) = \widehat{W}^{s/u}(\mathcal{O})$ is dense in T^1M . Fix $y \in \text{Reg}$, and let $U \subset \text{Reg}$ be an open set containing y . Let $\varepsilon, \delta > 0$ small such that:

- For every $z \in B_\varepsilon(y)$, $\widehat{W}^{s/u}(z)$ has internal radius at z larger than δ .
- If N^s, N^u are submanifolds tangent to $E^s \oplus \langle X \rangle, E^u \oplus \langle X \rangle$ respectively and if there is $z_{s/u} \in N^{s/u} \cap B_\varepsilon(y)$ such that $N^{s/u}$ has internal radius at $z_{s/u}$ larger than δ , then N^s, N^u have a transverse intersection at some $z \in U$.

The parameter $\varepsilon > 0$ exists because $E^{s/u}$ varies continuously and $E_y^s \oplus \langle X_y \rangle \oplus E_y^u = T_y(T^1M)$. Since $W^{s/u}(\mathcal{O})$ is dense, then $W^s(\mathcal{O}) \cap B_\varepsilon(y) \neq \emptyset$ and $W^u(\mathcal{O}) \cap B_\varepsilon(y) \neq \emptyset$, hence $W^s(\mathcal{O}) \pitchfork W^u(\mathcal{O})$ at some point $z \in U$. This implies that $y \in \text{HC}(\mathcal{O})$. Since $y \in \text{Reg}$ is arbitrary, $\text{HC}(\mathcal{O}) \supset \text{Reg}$. Finally, since Reg is dense in T^1M , we conclude that $\text{HC}(\mathcal{O}) = T^1M$. \square

Proof of Theorem 1.1. Let ψ be Hölder continuous or of the form $\psi = q\psi^u$ with $q \in \mathbb{R}$. In particular, ψ is continuous, hence the existence of an equilibrium state is guaranteed by the C^∞ regularity of φ [New89]. For the uniqueness, let μ_1, μ_2 be two ergodic equilibrium states. Since $P(\text{Sing}, \psi) < P(\psi)$, we have $\mu_1(\text{Reg}) = \mu_2(\text{Reg}) = 1$ and so μ_1, μ_2 are hyperbolic. Since μ_1, μ_2 are ergodic, we can take $\chi > 0$ small so that μ_1, μ_2 are χ -hyperbolic. The measures μ_1 and μ_2 are homoclinically related.

Indeed, let \mathcal{O} be a hyperbolic periodic orbit and $x \in \mathcal{O}$. Let $y_i \in \text{Reg}$ be a generic point for μ_i , $i = 1, 2$. Since $W^{s/u}(\mathcal{O}) = \widehat{W}^{s/u}(\mathcal{O})$ is dense in T^1M , we have $W_{\text{loc}}^s(y_1) \pitchfork W^u(x)$ and $W_{\text{loc}}^u(y_2) \pitchfork W^s(x)$. By the inclination lemma, we conclude that $W^s(y_1) \pitchfork W^u(y_2)$ near x . Interchanging the roles of y_1, y_2 , we also have $W^s(y_2) \pitchfork W^u(y_1)$.

Applying Theorem 2.1 to μ_1 and μ_2 , we get an irreducible TMF (Σ_r, σ_r) and a Hölder continuous map $\pi_r : \Sigma_r \rightarrow \mathbb{R}$ such that $\mu_i[\pi_r(\Sigma_r^\#)] = 1$ for $i = 1, 2$. Therefore, μ_1, μ_2 lift to ergodic measures $\widehat{\mu}_1, \widehat{\mu}_2$ on Σ_r . These measures are equilibrium states of the potential $\widehat{\psi} = \psi \circ \pi_r$.

CLAIM: $\widehat{\psi}$ is Hölder continuous.

Proof of the claim. When ψ is Hölder continuous, $\widehat{\psi} = \psi \circ \pi_r$ is the composition of two Hölder maps, hence Hölder continuous.

When $\psi = q\psi^u$ for some $q \in \mathbb{R}$, we show that $(\underline{v}, t) \in \Sigma_r \mapsto E_{\pi_r(\underline{v}, t)}^u$ is Hölder continuous, which obviously implies that $\widehat{\psi}$ is Hölder continuous. It is enough to show that $\underline{v} \in \Sigma \mapsto E_{\pi_r(\underline{v}, 0)}^u$ is Hölder continuous, since by $d\varphi$ -invariance this implies that $(\underline{v}, t) \in \Sigma_r \mapsto E_{\pi_r(\underline{v}, t)}^u$ is Hölder continuous.

As in the proof of Theorem 2.1, let $f : \Lambda \rightarrow \Lambda$ be the Poincaré return map. The map $\mathcal{F} : \underline{v} \in \Sigma \rightarrow F_x^u \subset T_x\Lambda$, where F_x^u is the unstable direction for f at $x = \pi_r(\underline{v}, 0)$, is Hölder continuous. As a matter of fact, employing the notation of [ALP20], F_x^u is the tangent direction of the u -admissible graph $V^u[\underline{v}] = \lim_{n \rightarrow \infty} (\mathcal{F}_{-1} \circ \dots \circ \mathcal{F}_{-n})(V_{-n})$, where \mathcal{F}_{-n} is the unstable graph transform defined by the edge $v_n \rightarrow v_{n+1}$ and V_{-n} is any u -admissible graph at v_{-n} , see [ALP20, Section 4.5].² Therefore, if two sequences $\underline{v}, \underline{w}$ are close then $V^u[\underline{v}], V^u[\underline{w}]$ are C^1 -close. For the details, see [BO18, Proposition. 3.12(5)], whose proof applies verbatim in the context of [ALP20].

Let $\mathbf{p}_x : T_x\Lambda \rightarrow X_x^\perp$ be the orthogonal projection. Such map exists and is an isomorphism because both $X_x^\perp, T_x\Lambda$ have dimension $\dim(T^1M) - 1$ and $T_x\Lambda$ is almost orthogonal to X_x . We have $E_x^u = \mathbf{p}_x(F_x^u)$. Since $x \in T^1M \mapsto X_x^\perp$ and $x \in \Lambda \mapsto T_x\Lambda$ are C^∞ , the map $\mathcal{P} : x \in \Lambda \mapsto \mathbf{p}_x$ is C^∞ . Therefore $\underline{v} \in \Sigma \mapsto E_{\pi_r(\underline{v}, 0)}^u$, being the composition $\mathcal{P} \circ \mathcal{F}$, is Hölder continuous. \square

The measures $\widehat{\mu}_1, \widehat{\mu}_2$ project to ergodic σ -invariant probability measures $\widehat{\nu}_1, \widehat{\nu}_2$ on the irreducible component Σ which are equilibrium states of the Hölder continuous potential $\widehat{\psi}_r - P_{\text{top}}(\psi)r$ where $\widehat{\psi}_r(\underline{v}) = \int_0^{r(\underline{v})} \widehat{\psi}(\underline{v}, t) dt$, see e.g. [PP90, Proposition 6.1]. By [BS03, Theorem 1.1], we conclude that $\widehat{\nu}_1 = \widehat{\nu}_2$, and so $\mu_1 = \mu_2$.

Finally, we show that the unique equilibrium state μ is fully supported. The proof is the same of [BCS22, Corollary 3.3]. Using the same notation of the previous paragraphs, $\widehat{\nu}$ has full support in Σ by [BS03] and so $\widehat{\mu}$ has full support in Σ_r . This implies that $\text{supp}(\mu) = \overline{\pi_r(\Sigma_r)}$, so it is enough to show that $\pi_r(\Sigma_r)$ is dense in T^1M . Let \mathcal{O} be a hyperbolic periodic orbit homoclinically related to μ .³ We thus have $\text{supp}(\mu) \subset \text{HC}(\mathcal{O}) = T^1M$, hence it is enough to show that $\pi_r(\Sigma_r)$ is dense in $\{W^u(\mathcal{O}) \pitchfork W^s(\mathcal{O})\}$. The proof of this fact in [BCS22, Corollary 3.3]

²Actually, the coding Σ is obtained from a refinement of a preliminary coding. The sequences on Σ lift to sequences on this preliminary coding, whose vertices are double Pesin charts and whose edges define stable/unstable graph transforms.

³The existence of this orbit is consequence of Katok's horseshoe theorem for flows; another way to obtain this is by the symbolic coding of [ALP20].

uses [BCS22, Proposition 3.7], which works equally well in our context. Hence we conclude that $\text{supp}(\mu) = \overline{\pi_r(\Sigma_r)} = \text{HC}(\mathcal{O}) = T^1M$. \square

Remark 3.1. When M is a surface and $\psi = q\psi^u$, [LLS16] proves uniqueness without using the Hölder continuity of $\widehat{\psi}$. For that, the authors show that $\widehat{\psi}_r$ above is cohomologous to the Hölder continuous function $\underline{v} \in \Sigma \mapsto -\log \det(df|_{F_{\pi_r(\underline{v},0)}^u})$, see [LLS16, Lemmas 8.1 and 8.2]. We note that the same proof applies here.

Proof of Theorem 1.2. The proof is the same of [BCFT18, Theorem C]. We include it for completeness. Let M be a surface and $\psi = q\psi^u$ for $q < 1$. Inside Sing all Lyapunov exponents are zero. Let μ be a φ -invariant measure supported on Sing. By the Ruelle inequality, $h(\mu) = 0$. Also, since $\psi^u = 0$ on Sing, it follows that

$$h(\mu) + q \int \psi^u d\mu = 0.$$

Therefore $P(\text{Sing}, \psi) = 0$. On the other hand, if λ is the Lebesgue measure, then by Pesin equality

$$h(\lambda) + q \int \psi^u d\mu = (1 - q)h(\lambda) > 0.$$

Therefore $P(\psi) > 0 = P(\text{Sing}, \psi)$. Now apply Theorem 1.1. \square

Proof of Theorem 1.3. When M is a surface we have, as in the previous proof, that $P(\text{Sing}, 0) = 0$. Since $P(0) = h_{\text{top}}(\varphi) > 0$, we get the pressure gap $P(\text{Sing}, 0) < P(0)$. Applying Theorem 1.1, we conclude the uniqueness of the measure of maximal entropy. In higher dimension, we invoke [BCFT18, Theorem B] to get the pressure gap $P(\text{Sing}, 0) < P(0)$. Again, applying Theorem 1.1, we obtain the uniqueness of the measure of maximal entropy. \square

REFERENCES

- [ALP20] Ermerson Araujo, Yuri Lima, and Mauricio Poletti, *Symbolic dynamics for nonuniformly hyperbolic maps with singularities in high dimension*, 2020. preprint arXiv:2010.11808.
- [Bal82] Werner Ballmann, *Axial isometries of manifolds of nonpositive curvature*, Math. Ann. **259** (1982), no. 1, 131–144.
- [BC04] Christian Bonatti and Sylvain Crovisier, *Réurrence et généricité*, Invent. Math. **158** (2004), no. 1, 33–104.
- [BCFT18] K. Burns, V. Climenhaga, T. Fisher, and D. J. Thompson, *Unique equilibrium states for geodesic flows in nonpositive curvature*, Geom. Funct. Anal. **28** (2018), no. 5, 1209–1259.
- [BCL23] Jérôme Buzzi, Sylvain Crovisier, and Yuri Lima, *Symbolic dynamics for large non-uniformly hyperbolic sets of three dimensional flows*, 2023. preprint arXiv:2307.14319.
- [BCS22] Jérôme Buzzi, Sylvain Crovisier, and Omri Sarig, *Measures of maximal entropy for surface diffeomorphisms*, Ann. of Math. (2) **195** (2022), no. 2, 421–508.
- [BO18] Snir Ben Ovadia, *Symbolic dynamics for non-uniformly hyperbolic diffeomorphisms of compact smooth manifolds*, Journal of Modern Dynamics **13** (2018), 43–113.
- [BP07] Luis Barreira and Yakov Pesin, *Nonuniform hyperbolicity*, Encyclopedia of Mathematics and its Applications, vol. 115, Cambridge University Press, Cambridge, 2007. Dynamics of systems with nonzero Lyapunov exponents. MR2348606 (2010c:37067)
- [BR75] Rufus Bowen and David Ruelle, *The ergodic theory of Axiom A flows*, Invent. Math. **29** (1975), no. 3, 181–202. MR0380889 (52 #1786)

- [BS03] Jérôme Buzzi and Omri Sarig, *Uniqueness of equilibrium measures for countable Markov shifts and multidimensional piecewise expanding maps*, Ergodic Theory Dynam. Systems **23** (2003), no. 5, 1383–1400. MR2018604 (2004k:37053)
- [BW72] Rufus Bowen and Peter Walters, *Expansive one-parameter flows*, J. Diff. Equations **12** (1972), 180–193.
- [CT22] Benjamin Call and Daniel J. Thompson, *Equilibrium states for self-products of flows and the mixing properties of rank 1 geodesic flows*, J. Lond. Math. Soc. (2) **105** (2022), no. 2, 794–824.
- [Ebe01] Patrick Eberlein, *Geodesic flows in manifolds of nonpositive curvature*, Smooth ergodic theory and its applications (Seattle, WA, 1999), 2001, pp. 525–571.
- [Kni98] Gerhard Knieper, *The uniqueness of the measure of maximal entropy for geodesic flows on rank 1 manifolds*, Ann. of Math. (2) **148** (1998), no. 1, 291–314. MR1652924 (2000b:37016)
- [LLS16] François Ledrappier, Yuri Lima, and Omri Sarig, *Ergodic properties of equilibrium measures for smooth three dimensional flows*, Comment. Math. Helv. **91** (2016), no. 1, 65–106.
- [LS19] Yuri Lima and Omri M. Sarig, *Symbolic dynamics for three-dimensional flows with positive topological entropy*, J. Eur. Math. Soc. (JEMS) **21** (2019), no. 1, 199–256.
- [New72] Sheldon E. Newhouse, *Hyperbolic limit sets*, Trans. Amer. Math. Soc. **167** (1972), 125–150.
- [New89] ———, *Continuity properties of entropy*, Ann. of Math. (2) **129** (1989), no. 2, 215–235. MR986792 (90f:58108)
- [PP90] William Parry and Mark Pollicott, *Zeta functions and the periodic orbit structure of hyperbolic dynamics*, Astérisque **187-188** (1990), 268. MR1085356 (92f:58141)
- [RHRHTU11] F. Rodriguez Hertz, M. A. Rodriguez Hertz, A. Tahzibi, and R. Ures, *Uniqueness of SRB measures for transitive diffeomorphisms on surfaces*, Comm. Math. Phys. **306** (2011), no. 1, 35–49.
- [Sar13] Omri M. Sarig, *Symbolic dynamics for surface diffeomorphisms with positive entropy*, J. Amer. Math. Soc. **26** (2013), no. 2, 341–426.
- [Sma67] S. Smale, *Differentiable dynamical systems*, Bull. Amer. Math. Soc. **73** (1967), 747–817.

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO CEARÁ (UFC), CAMPUS DO PICI, BLOCO 914, CEP 60455-760. FORTALEZA – CE, BRASIL
Email address: yurilima@gmail.com

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO CEARÁ (UFC), CAMPUS DO PICI, BLOCO 914, CEP 60455-760. FORTALEZA – CE, BRASIL
Email address: mpoletti@mat.ufc.br