

TOWARDS A SCHAUDER THEORY FOR FRACTIONAL VISCOUS HAMILTON–JACOBI EQUATIONS

ESPEN R. JAKOBSEN AND ARTUR RUTKOWSKI

ABSTRACT. We survey some results on Lipschitz and Schauder regularity estimates for viscous Hamilton–Jacobi equations with subcritical Lévy diffusions. The Schauder estimates, along with existence of smooth solutions, are obtained with the help of a Duhamel formula and L^1 bounds on the spatial derivatives of the heat kernel. Our results cover very general nonlocal and mixed local–nonlocal diffusions, including strongly anisotropic, nonsymmetric, mixed order, and spectrally one-sided models.

1. INTRODUCTION

The goal of this note is an elementary exposition of some existing results on classical well-posedness and Schauder regularity theory for viscous Hamilton–Jacobi (HJ) equations. We are interested in the following problem:

$$(1.1) \quad \begin{cases} \partial_t u - \mathcal{L}u + H(t, x, u, Du) = f(t, x) & \text{in } (0, T) \times \mathbb{R}^d, \\ u(0) = u_0 & \text{in } \mathbb{R}^d. \end{cases}$$

Here, \mathcal{L} is a translation invariant diffusion operator of the form:

$$(1.2) \quad \mathcal{L}u = B \cdot Du + \operatorname{div}(ADu) + \int_{\mathbb{R}^d} (u(\cdot+z) - u(\cdot) - Du(\cdot) \cdot z \mathbf{1}_{B(0,1)}(z)) \nu(dz),$$

with Lévy triplet $(B, A, \nu) \in \mathbb{R}^d \times \mathbb{R}^{d \times d} \times \mathcal{M}_+(\mathbb{R}^d)$ where $A \geq 0$ and the Lévy measure ν is a nonnegative Borel measure satisfying $\int_{\mathbb{R}^d} (1 \wedge |z|^2) \nu(dz) < \infty$. The function $H \in C((0, T) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d)$ is called the Hamiltonian. Throughout the paper we assume that the operator \mathcal{L} is subcritical, i.e. a negative operator of order $\alpha \in (1, 2]$. The order is 2 when $A > 0$, otherwise it depends on the behavior of ν near $z = 0$. The order condition is implicitly formulated through the heat kernel of \mathcal{L} , see Section 2 below. Because the linear diffusion term is the principal term, we call (1.1) a *viscous* HJ equation.

Definition 1.1. We say that u is a classical solution to (1.1) if $u \in C([0, T] \times \mathbb{R}^d)$, $\partial_t u, Du, \mathcal{L}u \in C((0, T) \times \mathbb{R}^d)$ and the equation and the initial condition are satisfied pointwise.

2020 *Mathematics Subject Classification.* 49L12, 35S10, 35A01, 35K08, 45K05, 35K61, 35B65, 35A09.

Key words and phrases. Viscous Hamilton–Jacobi equation, Schauder estimates, fractional operators, nonlocal operators, Duhamel formula, classical solutions, well-posedness.

For linear (possibly nonlocal) PDEs in non-divergence form, Schauder regularity theory loosely speaking states that if data and coefficients are β -Hölder continuous, then solutions are $\alpha + \beta$ Hölder continuous where α is the order of the equation [24, 36, 37, 39]. In this paper we will give two results of this type for the nonlinear equation (1.1). As opposed to many other papers on regularity theory, our method of proof also produces well-posedness of solutions with this regularity. In Section 4 we give uniform in time Schauder estimates for smooth initial conditions and Lipschitz f , and in Section 5 we have Schauder estimates for initial data and f with low Hölder regularity where gradients will blow-up for small times. For the first type of results, to handle locally Lipschitz Hamiltonians in the long time existence proof, we need uniform Lipschitz estimates on the solutions. Such estimates along with a comparison principle are proved in Section 3.

The viscous HJ equations arise in the optimal stochastic control problems or differential games. The unknown function u then represents the (upper or lower) value function for the optimally controlled process. Equation (1.1) corresponds to the case of controlled drift but not diffusion. When H is convex in Du , it corresponds to the control problem

$$u(T - t, x) = \inf_{\alpha_s \in \mathcal{A}} E \left(\int_t^T [L(X_s^{t,x}, \alpha_s, s) + f(s, X_s^{t,x})] ds + u_0(X_T^{t,x}) \right),$$

where L and f are running costs, u_0 a terminal cost, \mathcal{A} is the set of admissible controls, and $X_s = X_s^{t,x}$ is the solution of an SDE driven by a Lévy process N_s with generator \mathcal{L} ,

$$dX_s = b(\alpha_s, X_s) ds + dN_s, \quad s > t; \quad X_t = x.$$

We refer e.g. to Fleming and Soner [23, Chapter IV.3] for more details in the case of Gaussian noise. For non-Gaussian (Lévy) noise see Øksendal and Sulem [40], and for differential games see Carmona [11].

Early contributions to viscous HJ equations were given by Kružkov [35] and Crandall and Lions [18]. Nowadays they have an extensive literature and are still an active area of research. With hundreds of papers, we sample a few to give an idea of some of activity which encompasses well-posedness, regularity, boundary value problems, long time/ergodic/asymptotic problems, singular solutions, approximation and numerics, homogenization, control and large deviations, and stochastic perturbations [4, 5, 6, 13, 17, 19, 28, 38, 41]. The list is very far from being complete or even representative. Our own main motivation comes from the theory of mean field games with individual noises. There the Nash equilibria are characterized by a forward-backward system consisting of a viscous HJ equation and a Fokker–Planck equation, see [1, 10, 12] for the background and [15, 22] for mean field games with nonlocal diffusions.

In equation (1.1) the diffusion term dominates and induces a regularizing effect, which allows to solve the equation with the help of the heat kernel of \mathcal{L} and a Duhamel formula, yielding the so-called mild solutions. In the nonlocal

case, this approach to the viscous HJ equation was used by Imbert [29] and many other authors [3, 21, 22, 30, 32, 33], whereas the local case goes back at least to Ben-Artzi [7]. We note that for regular data the Duhamel formula for the viscous HJ equation works somewhat similar to the linear case, at least for the short-time results. In this connection we mention the work of Mikulevičius and Pragarauskas [39] who study regularization properties for the heat equation with quite general Lévy-type diffusions. More recently, Schauder estimates (without existence of classical solutions) were developed by Dong, Jin and Zhang [20] for fully nonlinear equations involving nonlocal operators with the Lévy measure comparable to the one of the fractional Laplacian. The results given below cover a large class of translation invariant diffusions which were not present in [20], including strongly anisotropic and nonsymmetric stable operators.

The long-time existence under *local* regularity assumptions for H is more delicate, as it requires global Lipschitz bounds for the solutions, which do not follow from the Duhamel formula. In general, these Lipschitz bounds can be obtained using the comparison principle and viscosity solution techniques, see Imbert [29] and Jakobsen and Karlsen [31]. If the solution is known to be more regular pointwise in time, one can also apply the Bernstein method, which is quite straightforward when one assumes that $f(t) \in W^{1,\infty}(\mathbb{R}^d)$ uniformly in t . A more sophisticated Bernstein-type argument can be applied in order to get bounds for u in terms of weaker norms of f , see e.g. Goffi [26], but this is beyond the scope of this note.

In case that $u_0, f(t) \in C_b^\beta(\mathbb{R}^d)$ for $t \in (0, T)$ with $\beta \in (0, 1)$ we can still obtain classical solutions, but $Du(t)$ and $\mathcal{L}u(t)$ can, in general, blow up for small t . There are many works concerning the viscous HJ equations with irregular initial data, see e.g., Gilding, Guedda, and Kersner [25], Benachour and Laurençot [8] for the local diffusions and [22, 29, 32, 43] for the nonlocal case. However, we are not aware of papers which explicitly study blow up rates for $u_0 \in C_b^\beta(\mathbb{R}^d)$ data in the vein of Theorem 5.1 below, obtained by the authors together with Amund Bergset, cf. his master thesis [9]. Admittedly, we only allow Hamiltonians with global Lipschitz regularity there so far; an extension to locally Lipschitz Hamiltonians seems non-trivial and interesting. In a future work we aim to study mean field game systems with Hölder payoff functions. The blowup for the viscous HJ equation means that the drift term in the corresponding Fokker–Planck equation may become large close to the terminal time, which poses a difficulty for compactness arguments.

2. ASSUMPTIONS

We adopt rather standard assumptions on H and f , which we will use in Sections 3 and 4. We note that by assuming regularity for higher order derivatives in (H1) and (F0) below, one could get higher regularity of solutions to the viscous HJ equation. Here, for the sake of clarity, we work with

quite weak assumptions, but strong enough to allow for the classical solution setting.

(H0): $H: [0, \infty) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$ is continuous and for every R there exists C_R such that for every $t, |u| \leq R$, $x, y \in \mathbb{R}^d$ and $p, q \in B_R$,

$$|H(t, x, u, p) - H(t, y, u, q)| \leq C_R(|x - y| + |p - q|).$$

(H1): H is smooth and for all $l \in \mathbb{N}^{2d+1}$ with $|l| \leq 2$, $\sup_{x \in \mathbb{R}^d} |D_{x,u,p}^l H(\cdot, x, \cdot, \cdot)|$ is locally bounded.

(H2): For every $R > 0$ there exists $C_R > 0$ such that for $x, y \in \mathbb{R}^d$, $t, |u| \leq R$, and $p \in \mathbb{R}^d$,

$$|H(t, x, u, p) - H(t, y, u, p)| \leq C_R(1 + |p|)|x - y|.$$

(H3): For every $T > 0$ there exists $\gamma = \gamma(T) \in \mathbb{R}$ such that for all $t \in [0, T]$, $x \in \mathbb{R}^d$, $u \leq v$, and $p \in \mathbb{R}^d$,

$$H(t, x, v, p) - H(t, x, u, p) \geq \gamma(v - u).$$

(F0): $f \in C_b([0, T] \times \mathbb{R}^d)$ and for every $T > 0$ there exists C such that for $t \in [0, T]$ and $x, y \in \mathbb{R}^d$

$$|f(t, x) - f(t, y)| \leq C|x - y|.$$

Note that (H1) is strictly stronger than (H0).

In the Schauder estimates we use the fact that the order of the operator is greater than 1. We express that through appropriate estimates for the heat kernel. If we let ψ be the Fourier multiplier corresponding to \mathcal{L} (see e.g. [34]), then the heat kernel of \mathcal{L} is defined as

$$K_0 = \delta_0, \quad \mathcal{F}(K_t)(\xi) = e^{-t\psi(\xi)}, \quad t > 0.$$

(K): There is $\mathcal{K} > 0$ and $\alpha \in (1, 2]$, such that the heat kernels K and K^* of \mathcal{L} and \mathcal{L}^* respectively are smooth densities of probability measures, and for $\tilde{K} = K, K^*$ and $\beta \in \mathbb{N}^d$ we have

$$\|D^\beta \tilde{K}(t, \cdot)\|_{L^1(\mathbb{R}^d)} \leq \mathcal{K}t^{-\frac{\beta}{\alpha}}.$$

Under (K) we have $K_t \in C_0^\infty(\mathbb{R}^d)$ for every $t > 0$ and K_t satisfies $(\partial_t - \mathcal{L})K_t = 0$ pointwise on $(0, \infty) \times \mathbb{R}^d$. Here are some examples of operators satisfying (K) (taken mostly from [22, Section 4]):

- (i) There is $c > 0$ such $\langle Ax, x \rangle \geq c|x|^2$ for $x \in \mathbb{R}^d$, e.g. $\mathcal{L} = \Delta$. Then (K) is satisfied with $\alpha = 2$.
- (ii) The Lévy measure ν has an absolutely continuous part with density $\nu_{ac}(z) \approx |z|^{-d-\alpha}$ for all $|z| \leq 1$ and $\alpha \in (1, 2)$, see [22, Theorem 4.3], based on the pointwise estimates in [27].
- (iii) $\mathcal{L} = -(\partial_{x_1}^2)^{\alpha_1/2} - (\partial_{x_2}^2)^{\alpha_2/2} - \dots - (\partial_{x_d}^2)^{\alpha_d/2}$, where $\alpha_i \in (1, 2]$. Then $\alpha = \min_i \alpha_i$.

- (iv) The family of Riesz–Feller operators on \mathbb{R} , including the spectrally one-sided model with $A = 0$ and $\nu(z) = |z|^{-1-\alpha}\mathbf{1}_{(0,\infty)}(z)$, see [2, Lemma 2.1 (G7) and Proposition 2.3].
- (v) The generator of the CGMY model in finance, see [22, Example 4.4].
- (vi) If \mathcal{L} satisfies (K) and L is any other Lévy operator, then $\mathcal{L} + L$ satisfies (K).

3. COMPARISON PRINCIPLE AND LIPSCHITZ ESTIMATES

In the next sections we will construct short-time solutions via fixed point arguments and we will get rather strong regularity estimates, but only on short time intervals. For the sake of obtaining long-time solutions we will establish a global in time Lipschitz bound using different methods in Theorem 3.3 below.

We first give a comparison principle in order to obtain supremum bounds. A similar result was given by Imbert [29, Theorem 2] for purely nonlocal \mathcal{L} in the (more general) viscosity solution setting; it extends rather easily to general Lévy operators.

Theorem 3.1. *Assume that (H0), (H2), (H3), and (F0) hold. Let u_0 be uniformly continuous and bounded and suppose that u and v are respectively classical sub- and supersolution to (1.1), i.e., $u(0, x) \leq u_0(x) \leq v(0, x)$ for $x \in \mathbb{R}^d$ and*

$$\begin{aligned} \partial_t u - \mathcal{L}u + H(t, x, u, Du) &\leq f(t, x) \quad \text{in } (0, T) \times \mathbb{R}^d, \\ \partial_t v - \mathcal{L}v + H(t, x, v, Dv) &\geq f(t, x) \quad \text{in } (0, T) \times \mathbb{R}^d. \end{aligned}$$

Then $u \leq v$ on $[0, T) \times \mathbb{R}^d$.

Proof. Note that under present assumptions f can be absorbed into H , so we can assume that $f = 0$. Since the diffusion operator \mathcal{L} is linear and translation invariant, it suffices to follow the proof of Imbert. First of all, since the Lévy measure ν is finite outside neighborhoods of 0, there exists a function $\varphi \in C^2(\mathbb{R}^d)$ such that

$$\|D\varphi\|_\infty + \|D^2\varphi\|_\infty \leq C, \quad \lim_{|x| \rightarrow \infty} \varphi(x) = \infty, \quad \mathcal{L}\varphi \in C(\mathbb{R}^d).$$

Since the sub/supersolutions are classical, we do not need to introduce the sub- and superdifferentials after [29, (14)], because the operator can be evaluated pointwise, and we have (in the language of [29]) $DU(\bar{t}, \bar{x}) - DV(\bar{s}, \bar{y}) = \alpha D\varphi(\bar{x})$. Then it is easy to obtain the following counterpart of inequality [29, (16)]:

$$\mathcal{L}V(\bar{s}, \bar{y}) - \mathcal{L}U(\bar{t}, \bar{x}) \leq -\alpha \mathcal{L}\varphi(\bar{x}).$$

The rest of the proof does not depend on the form of the diffusion operator. \square

Corollary 3.2. *Assume that (H0), (H2), (H3), and (F0) hold. If u_0 is uniformly continuous and bounded and u is a classical solution to (1.1), then*

$$(3.1) \quad \|u(t)\|_\infty \leq e^{-\gamma t} \|u_0\|_\infty + (1 - \frac{1}{\gamma} e^{-\gamma t})(\|f\|_\infty + \|H(\cdot, \cdot, 0, 0)\|_\infty), \quad t \in [0, T].$$

Proof. Let $w(t)$ denote the right hand side of (3.1). Then the proof follows by comparison since by (H3) w and $-w$ are super- and subsolutions of (1.1). \square

Theorem 3.3. *Assume that (H0), (H2), (H3), and (F0) hold, $u_0 \in C_b^1(\mathbb{R}^d)$ and let u be a classical solution to (1.1). Then*

$$\|Du\|_\infty \leq C(T, H, f)(1 + \|Du_0\|_\infty), \quad t \in [0, T].$$

Remark 3.4. We only give the proof in the case of regular data and solutions to showcase the elegant and more accessible Bernstein method. For less regular data one can apply the viscosity solution proof of Imbert [29]. In order to adapt the proof of Imbert to mixed local-nonlocal operators an appropriate viscosity solution formulation is needed, see e.g., Jakobsen and Karlsen [31] in this regard.

Proof of Theorem 3.3 for regular data and solutions. If (H1) is satisfied, $f \in C([0, T], C_b^2(\mathbb{R}^d))$, and $u \in C([0, T], C_b^3(\mathbb{R}^d))$, then we can differentiate the equation in x_i and get

$$(3.2) \quad \begin{aligned} \partial_t u_{x_i} - \mathcal{L}u_{x_i} + H_{x_i}(t, x, u, Du) + u_{x_i} H_u(t, x, u, Du) \\ + Du_{x_i} \cdot H_p(t, x, u, Du) = f_{x_i}. \end{aligned}$$

Now, let $w = \frac{1}{2}|Du|^2$. If we multiply (3.2) by u_{x_i} and sum over i we get

$$\begin{aligned} \partial_t w - \mathcal{L}w + \sum_i \Gamma(u_{x_i}) + Du \cdot H_x(t, x, u, Du) + 2w H_u(t, x, u, Du) \\ + Dw \cdot H_p(t, x, u, Du) = Df \cdot Du, \end{aligned}$$

where (recall that $A \geq 0$)

$$\Gamma(u) = \frac{1}{2}(\mathcal{L}u^2 - 2u\mathcal{L}u) = \frac{1}{2} \int_{\mathbb{R}^d} (u(x+y) - u(x))^2 \nu(dy) + Du \cdot ADu \geq 0.$$

Furthermore, we have $w \geq 0$ and $H_u \geq \gamma$ by (H3). Therefore, by (H2) and Corollary 3.2,

$$\begin{aligned} \partial_t w - \mathcal{L}w + Dw \cdot H_p(t, x, u, Du) \\ \leq -Du \cdot H_x(t, x, u, Du) - 2\gamma w + \frac{1}{2}\|Df\|_\infty^2 + \frac{1}{2}w \\ \leq C(T, H, f)(1 + w). \end{aligned}$$

Let $C = C(T, H, f)$ and note that $v = e^{Ct}(1 + \|w(0)\|_\infty) - 1$ satisfies

$$\partial_t v - \mathcal{L}v + Dv \cdot H_p(t, x, u, Du) = C(1 + v), \quad v(0) = \|w(0)\|_\infty \geq w(0).$$

Therefore by the comparison principle for the linear parabolic problems we get that $w \leq v$, which ends the proof in this case. \square

4. UNIFORM SCHAUDER ESTIMATES WITH REGULAR INITIAL DATA

In this section we give uniform in time Schauder estimates under the assumption that the initial data is regular. The result and proof were essentially given in [32]. We repeat the arguments here in order to demonstrate the technique where the regularity of the heat kernel is used to gain $\alpha - \varepsilon$ derivatives over the forcing term f . Similar results were given earlier by, e.g., Imbert [29] and Ersland and Jakobsen [22], but with only one derivative gain. It seems that in order to gain full α regularity one needs a stronger assumption than (K), see e.g. [20] or [14].

In the proof we first show the existence of a short-time mild solution using the Duhamel formula and the Banach fixed point theorem. Then we show that the mild solution is classical, which enables us to use the results of the previous section to obtain the long time existence. Finally, we get the uniqueness from the comparison principle.

Theorem 4.1. *Let $\varepsilon \in (0, \alpha - 1)$ and assume that (H1), (H2), (H3), (F0), and (K) are satisfied and let $u_0 \in C_b^{1+\alpha-\varepsilon}(\mathbb{R}^d)$. Then problem (1.1) has a unique classical solution u satisfying*

$$(4.1) \quad \|\partial_t u\|_\infty + \sup_{t \in [0, T]} \|u(t, \cdot)\|_{C_b^{1+\alpha-\varepsilon}(\mathbb{R}^d)} \\ \leq c(T, \mathcal{K}) \left(c(H) + \sup_{t \in [0, T]} \|f(t, \cdot)\|_{W^{1, \infty}(\mathbb{R}^d)} + \|u_0\|_{C_b^{1+\alpha-\varepsilon}(\mathbb{R}^d)} \right).$$

Proof. We denote $\sigma = \alpha - 1 - \varepsilon$, so that $1 + \alpha - \varepsilon = 2 + \sigma$.

1) Short-time mild solutions by Banach's fixed point theorem. Let

$$R = R(u) = \sup_t \|u(t)\|_{C_b^1(\mathbb{R}^d)}, \quad R_1 = 2(\|u_0\|_{C_b^{2+\sigma}} + \sup_t \|f(t)\|_{W^{1, \infty}(\mathbb{R}^d)}) + 1,$$

and define the space

$$X = \{u \in C_b((0, T], C_b^{2+\sigma-\varepsilon}(\mathbb{R}^d)) : \sup_{t \in [0, T]} \|u(t, \cdot)\|_{C_b^{2+\sigma}(\mathbb{R}^d)} \leq R_1\}.$$

Note that for all $u \in X$ we have $R \leq R_1$. For $u \in X$, $t \in [0, T]$, $x \in \mathbb{R}^d$ we define the fixed-point map:

$$S(u)(t, x) := K_t * u_0(x) + \int_0^t K_{t-s} * [H(s, \cdot, u, Du) + f(s)](x) ds.$$

We will now estimate the $C_b^{2+\sigma}$ norms of $S(u)$ in order to see that it belongs to X . The time continuity of the $C_b^{2+\sigma-\varepsilon}$ norms follows from the $C_b^{2+\sigma}$ estimates, see e.g. [32, Lemma B.2]. First note that for $k = 1, 2$ we have

$$(4.2) \quad D_x^k S(u)(t, x) \\ = K_t * D_x^k u_0(x) + \int_0^t D_x K_{t-s} * D_x^{k-1} [H(s, \cdot, u, Du) + f(s)](x) ds.$$

By (K) and interpolation for Hölder functions (e.g. [32, Remark 2.22 d)]),

$$\int_0^T \|D_x K_{t-s}\|_{L^1(\mathbb{R}^d)} ds \leq \mathcal{K} \int_0^T (t-s)^{-\frac{1}{\alpha}} ds = \frac{\alpha-1}{\alpha} \mathcal{K} T^{\frac{\alpha-1}{\alpha}} := c_{T,\alpha},$$

$$\int_0^T \frac{\|D_x K_{t-s}(\cdot+h) - D_x K_{t-s}(\cdot)\|_{L^1(\mathbb{R}^d)}}{|h|^\sigma} ds \leq \mathcal{K} \int_0^T (t-s)^{-\frac{1+\sigma}{\alpha}} ds =: c_{T,\alpha,\sigma},$$

for all $h \in \mathbb{R}^d \setminus \{0\}$. Therefore by Young's inequality for convolutions and (H1) we have for $k = 0, 1, 2$ (for $k = 0$ we use $\|K_{t-s}\|_{L^1(\mathbb{R}^d)} = 1$):

$$\begin{aligned} \|D^k S u(t)\|_\infty &\leq \|D^k u_0\|_\infty + (T \vee c_{T,\alpha}) \left(\sup_t \|f(t)\|_{W^{1,\infty}(\mathbb{R}^d)} \right. \\ &\quad \left. + C(H, R)(1 + \sup_t \|u(t, \cdot)\|_{C_b^2(\mathbb{R}^d)}) \right), \\ [D_x^2 S u(t)]_{C^\sigma(\mathbb{R}^d)} &\leq [D_x^2 u_0]_{C^\sigma(\mathbb{R}^d)} + c_{T,\alpha,\sigma} \left(\sup_t \|f(t)\|_{W^{1,\infty}(\mathbb{R}^d)} \right. \\ &\quad \left. + C(H, R)(1 + \sup_t \|u(t, \cdot)\|_{C_b^2(\mathbb{R}^d)}) \right). \end{aligned}$$

Since $c_{T,\alpha}$ and $c_{T,\alpha,\sigma}$ converge to 0 as $T \rightarrow 0^+$, by taking T small enough we get $\sup_t \|S u(t)\|_{C_b^{2+\sigma}(\mathbb{R}^d)} \leq R_1$. Thus S maps X to X .

Then it remains to show that S is a contraction. The computations are similar to the above, we only inspect the highest order term to see that the bounds in (H1) are sufficient: by (4.2) we have

$$\begin{aligned} &[D_x^2(S(u) - S(v))]_{C^\sigma(\mathbb{R}^d)} \\ &\leq \mathcal{K} \int_0^t (t-s)^{-\frac{\sigma+1}{\alpha}} \|D_x(H(s, \cdot, u, Du) - H(s, \cdot, v, Dv))\|_\infty ds \\ &\leq c_{T,\alpha,\sigma} C(H, R) \|u - v\|_{C_b^2(\mathbb{R}^d)}. \end{aligned}$$

Again, by taking T small enough we get that S is a contraction on X . Therefore, by the Banach fixed point theorem S has a unique fixed point on X , which we call the mild solution.

2) Mild solutions are classical solutions. The quickest way to prove that is by using the Duhamel formula on $(t, t+h)$ with $h > 0$:

$$u(t+h, x) = K_h * u(t)(x) + \int_t^{t+h} K_{t+h-s} * [H(s, \cdot, u, Du) + f(s)](x) ds.$$

Hence,

$$\begin{aligned} \frac{u(t+h, x) - u(t, x)}{h} &= \frac{K_h * u(t)(x) - u(t, x)}{h} \\ &\quad + \frac{1}{h} \int_t^{t+h} K_{t+h-s} * [H(s, \cdot, u, Du) + f(s)](x) ds. \end{aligned}$$

As $h \rightarrow 0^+$, the first term on the right-hand side converges to the generator of the semigroup associated with K_t , but since $u \in C_b^2(\mathbb{R}^d)$ this is equal to

$\mathcal{L}u(t, x)$, see e.g. Sato [42, Theorem 31.5].¹ The second term converges to $H(t, x, u, Du) + f(t, x)$ by the regularity of f, H and u , and the fact that $K_t \rightarrow \delta_0$ weakly as $t \rightarrow 0^+$. Since the right-hand side converges, the left-hand side converges too, and is equal to $\partial_t u$, which proves that u is a classical solution.

3) Long-time existence by gluing. The idea for extending the solution in time consists in repeating the fixed-point procedure starting from a later point in time, e.g., $T/2$ and then iterating. However, we need to find appropriate lower bounds for lengths of the short-time existence intervals. To this end we use the fact that the mild solutions are classical and apply Theorem 3.3. This gives us a locally uniform in time bound for the constant R which determines the length of the existence interval, letting us extend the solution to an arbitrarily long time interval.

4) Uniqueness follows from the comparison principle in Theorem 3.1. \square

5. SCHAUDER ESTIMATES WITH IRREGULAR INITIAL DATA

In an ongoing project with Amund Bergset we study viscous HJ equations with initial data $u_0 \in C_b^\beta(\mathbb{R}^d)$ for $\beta \in (0, 1)$. In this case $\|Du(t)\|_\infty$ blows up as $t \rightarrow 0^+$, and therefore we need more explicit and global control of growth/regularity of the Hamiltonian in p . For simplicity we consider the following global Lipschitz condition:

(HLip) $H = H(p)$ and $H, D_p H$ are globally Lipschitz in p .

It should be possible to consider e.g. power-type Hamiltonians, but the analysis would be more delicate in this case. For local diffusions with subquadratic Hamiltonians and measure initial data see [8]. Superquadratic Hamiltonians were studied e.g. in [16].

Our result shows that the β -Hölder continuity of the data is preserved by the solution, while higher order Hölder norms blow up with a specific rate as $t \rightarrow 0$. There are two cases depending on whether $\alpha + \beta$ is greater or smaller than 2.

Theorem 5.1. *Assume that $T > 0$, (HLip) holds, $f \in L^\infty([0, T], C_b^\beta(\mathbb{R}^d))$, and $u_0 \in C_b^\beta(\mathbb{R}^d)$. Then (1.1) has a unique classical solution u . Moreover for every small $\varepsilon > 0$ there exists C such that for all $t \in (0, T]$:*

$$(i) \quad \|u(t)\|_\infty + [u(t)]_{C^\beta(\mathbb{R}^d)} + t^{\frac{1-\beta}{\alpha}} \|Du(t)\|_\infty + t^{\frac{\alpha-\varepsilon}{\alpha}} [Du(t)]_{\alpha+\beta-\varepsilon} \leq C, \quad \text{if } \alpha + \beta \leq 2,$$

$$(ii) \quad \|u(t)\|_\infty + [u(t)]_{C^\beta(\mathbb{R}^d)} + t^{\frac{1-\beta}{\alpha}} \|Du(t)\|_\infty + t^{\frac{2-\beta}{\alpha}} \|D^2u(t)\|_\infty + t^{\frac{\alpha-\varepsilon}{\alpha}} [D^2u(t)]_{\alpha+\beta-\varepsilon} \leq C, \quad \text{if } \alpha + \beta > 2.$$

¹The result is given for $C_0^2(\mathbb{R}^d)$, but an easy cut-off argument suffices.

The proof uses a similar strategy as the proof of Theorem 4.1 above, we refer to [9] for details. Compared to earlier results, this result cover a larger class of nonlocal operators, gives the behaviour of solution smoothness near $t = 0$, and provide existence of solutions with the stated smoothness. Our ultimate goal is to use this result to study new classes of Mean Field Games.

ACKNOWLEDGEMENTS

E. R. Jakobsen received funding from the Research Council of Norway under Grant Agreement No. 325114 “IMod. Partial differential equations, statistics and data: An interdisciplinary approach to data-based modelling”. While this project was carried out A. Rutkowski was an ERCIM Alain Bensoussan fellow at NTNU.

REFERENCES

- [1] Y. Achdou, P. Cardaliaguet, F. Delarue, A. Porretta, and F. Santambrogio. *Mean field games*, volume 2281 of *Lecture Notes in Mathematics*. Springer, Cham; Centro Internazionale Matematico Estivo (C.I.M.E.), Florence, [2020] ©2020. Edited by Pierre Cardaliaguet and Alessio Porretta, Fondazione CIME/CIME Foundation Subseries.
- [2] F. Achleitner and C. Kuehn. Traveling waves for a bistable equation with nonlocal diffusion. *Adv. Differential Equations*, 20(9-10):887–936, 2015.
- [3] N. Alibaud and C. Imbert. Fractional semi-linear parabolic equations with unbounded data. *Trans. Amer. Math. Soc.*, 361(5):2527–2566, 2009.
- [4] A. Arapostathis, A. Biswas, and L. Caffarelli. On uniqueness of solutions to viscous HJB equations with a subquadratic nonlinearity in the gradient. *Comm. Partial Differential Equations*, 44(12):1466–1480, 2019.
- [5] S. N. Armstrong and P. Cardaliaguet. Quantitative stochastic homogenization of viscous Hamilton-Jacobi equations. *Comm. Partial Differential Equations*, 40(3):540–600, 2015.
- [6] G. Barles, A. Porretta, and T. T. Tchamba. On the large time behavior of solutions of the Dirichlet problem for subquadratic viscous Hamilton-Jacobi equations. *J. Math. Pures Appl. (9)*, 94(5):497–519, 2010.
- [7] M. Ben-Artzi. Global existence and decay for a nonlinear parabolic equation. *Nonlinear Anal.*, 19(8):763–768, 1992.
- [8] S. Benachour and P. Laurençot. Global solutions to viscous Hamilton-Jacobi equations with irregular initial data. *Comm. Partial Differential Equations*, 24(11-12):1999–2021, 1999.
- [9] A. Bergset. *Mean field games in Hölder spaces*. NTNU (Trondheim), 2023. Master thesis.
- [10] P. Cardaliaguet, F. Delarue, J.-M. Lasry, and P.-L. Lions. *The master equation and the convergence problem in mean field games*, volume 201 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2019.
- [11] R. Carmona. *Lectures on BSDEs, stochastic control, and stochastic differential games with financial applications*, volume 1 of *Financial Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2016.
- [12] R. Carmona and F. Delarue. *Probabilistic theory of mean field games with applications. I*, volume 83 of *Probability Theory and Stochastic Modelling*. Springer, Cham, 2018. Mean field FBSDEs, control, and games.
- [13] E. Chasseigne and N. Ichihara. Qualitative properties of generalized principal eigenvalues for superquadratic viscous Hamilton-Jacobi equations. *NoDEA Nonlinear Differential Equations Appl.*, 23(6):Art. 66, 17, 2016.

- [14] P.-E. Chaudru de Raynal, S. Menozzi, and E. Priola. Schauder estimates for drifted fractional operators in the supercritical case. *J. Funct. Anal.*, 278(8):108425, 57, 2020.
- [15] M. Cirant and A. Goffi. On the existence and uniqueness of solutions to time-dependent fractional MFG. *SIAM J. Math. Anal.*, 51(2):913–954, 2019.
- [16] M. Cirant and A. Goffi. Lipschitz regularity for viscous Hamilton-Jacobi equations with L^p terms. *Ann. Inst. H. Poincaré C Anal. Non Linéaire*, 37(4):757–784, 2020.
- [17] M. Cirant and A. Goffi. On the problem of maximal L^q -regularity for viscous Hamilton-Jacobi equations. *Arch. Ration. Mech. Anal.*, 240(3):1521–1534, 2021.
- [18] M. G. Crandall and P.-L. Lions. Viscosity solutions of Hamilton-Jacobi equations. *Trans. Amer. Math. Soc.*, 277(1):1–42, 1983.
- [19] A. Davini, H. Ishii, R. Iturriaga, and H. Sanchez Morgado. Discrete approximation of the viscous HJ equation. *Stoch. Partial Differ. Equ. Anal. Comput.*, 9(4):1081–1104, 2021.
- [20] H. Dong, T. Jin, and H. Zhang. Dini and Schauder estimates for nonlocal fully nonlinear parabolic equations with drifts. *Anal. PDE*, 11(6):1487–1534, 2018.
- [21] J. Droniou and C. Imbert. Fractal first-order partial differential equations. *Arch. Ration. Mech. Anal.*, 182(2):299–331, 2006.
- [22] O. Ersland and E. R. Jakobsen. On fractional and nonlocal parabolic mean field games in the whole space. *J. Differential Equations*, 301:428–470, 2021.
- [23] W. H. Fleming and H. M. Soner. *Controlled Markov processes and viscosity solutions*, volume 25 of *Stochastic Modelling and Applied Probability*. Springer, New York, second edition, 2006.
- [24] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Grundlehren der Mathematischen Wissenschaften, Vol. 224. Springer-Verlag, Berlin-New York, 1977.
- [25] B. H. Gilding, M. Guedda, and R. Kersner. The Cauchy problem for $u_t = \Delta u + |\nabla u|^q$. *J. Math. Anal. Appl.*, 284(2):733–755, 2003.
- [26] A. Goffi. A priori Lipschitz estimates for nonlinear equations with mixed local and nonlocal diffusion via the adjoint-Bernstein method. *Boll. Unione Mat. Ital.*, 2022.
- [27] T. Grzywny and K. Szczypkowski. Estimates of heat kernels of non-symmetric Lévy processes. *Forum Math.*, 33(5):1207–1236, 2021.
- [28] N. Ichihara. Criticality of viscous Hamilton-Jacobi equations and stochastic ergodic control. *J. Math. Pures Appl. (9)*, 100(3):368–390, 2013.
- [29] C. Imbert. A non-local regularization of first order Hamilton-Jacobi equations. *J. Differential Equations*, 211(1):218–246, 2005.
- [30] T. Iwabuchi and T. Kawakami. Existence of mild solutions for a Hamilton-Jacobi equation with critical fractional viscosity in the Besov spaces. *J. Math. Pures Appl. (9)*, 107(4):464–489, 2017.
- [31] E. R. Jakobsen and K. H. Karlsen. Continuous dependence estimates for viscosity solutions of integro-PDEs. *J. Differential Equations*, 212(2):278–318, 2005.
- [32] E. R. Jakobsen and A. Rutkowski. The master equation for mean field game systems with fractional and nonlocal diffusions. *arXiv:2305.18867*, 2023.
- [33] G. Karch and W. A. Woyczyński. Fractal Hamilton-Jacobi-KPZ equations. *Trans. Amer. Math. Soc.*, 360(5):2423–2442, 2008.
- [34] D. Khoshnevisan and R. Schilling. *From Lévy-type processes to parabolic SPDEs*. Advanced Courses in Mathematics. CRM Barcelona. Birkhäuser/Springer, Cham, 2016. Edited by Lluís Quer-Sardanyons and Frederic Utzet.
- [35] S. N. Kružkov. Generalized solutions of Hamilton-Jacobi equations of eikonal type. I. Statement of the problems; existence, uniqueness and stability theorems; certain properties of the solutions. *Mat. Sb. (N.S.)*, 98(140)(3(11)):450–493, 496, 1975.

- [36] O. A. Ladyženskaja, V. A. Solonnikov, and N. N. Uralceva. *Linear and quasilinear equations of parabolic type*. Translations of Mathematical Monographs, Vol. 23. American Mathematical Society, Providence, R.I., 1968. Translated from the Russian by S. Smith.
- [37] G. M. Lieberman. *Second order parabolic differential equations*. World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
- [38] P.-L. Lions and P. E. Souganidis. Stochastic homogenization of Hamilton-Jacobi and “viscous”-Hamilton-Jacobi equations with convex nonlinearities—revisited. *Commun. Math. Sci.*, 8(2):627–637, 2010.
- [39] R. Mikulevičius and H. Pragarauskas. On the Cauchy problem for certain integro-differential operators in Sobolev and Hölder spaces. *Liet. Mat. Rink.*, 32(2):299–331, 1992.
- [40] B. Øksendal and A. Sulem. *Applied stochastic control of jump diffusions*. Universitext. Springer, Cham, 2019. Third edition of [MR2109687].
- [41] A. Porretta and P. Souplet. Blow-up and regularization rates, loss and recovery of boundary conditions for the superquadratic viscous Hamilton-Jacobi equation. *J. Math. Pures Appl. (9)*, 133:66–117, 2020.
- [42] K. Sato. *Lévy processes and infinitely divisible distributions*, volume 68 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2013. Translated from the 1990 Japanese original, Revised edition of the 1999 English translation.
- [43] L. Silvestre. On the differentiability of the solution to the Hamilton-Jacobi equation with critical fractional diffusion. *Adv. Math.*, 226(2):2020–2039, 2011.

DEPARTMENT OF MATHEMATICAL SCIENCES, NORWEGIAN UNIVERSITY OF SCIENCE
AND TECHNOLOGY, TRONDHEIM, NORWAY
Email address: `espen.jakobsen@ntnu.no`

DEPARTMENT OF MATHEMATICAL SCIENCES, NORWEGIAN UNIVERSITY OF SCIENCE
AND TECHNOLOGY, TRONDHEIM, NORWAY

FACULTY OF PURE AND APPLIED MATHEMATICS, WROCLAW UNIVERSITY OF SCIENCE
AND TECHNOLOGY, WROCLAW, POLAND
Email address: `artur.rutkowski@pwr.edu.pl`