

Two-Person Adversarial Games are Zero-Sum: An Elaboration of a Folk Theorem*

M. Ali Khan[†], Arthur Paul Pedersen^{‡ §} and David Schritteser[¶]

July 2024

Abstract. The observation that every two-person adversarial game is an affine transformation of a zero-sum game is traceable to Luce-Raiffa (1957) and made explicit in Aumann (1987). Recent work of (ADP) Adler-Daskalakis-Papadimitriou (2009), and of Raimondo (2023) in increasing generality, proves what has so far remained a conjecture. We present two proofs of an even more general formulation: the first draws on multilinear utility theory developed by Fishburn-Roberts (1978); the second is a consequence of the ADP proof itself for a special case of a two-player game with a set of three actions. (92 words)

Journal of Economic Literature Classification Numbers: C72, D01.

2020 Mathematics Subject Classification Numbers: 91A05, 91A10, 91A30.

Key Words: Game theory, two-person games, strictly competitive, adversarial, utility theory.

Running Title: Two-Person Adversarial Games are Zero-Sum

* The authors are grateful to Marcelo Ariel Fernandez, Ani Ghosh, Michael Grossberg, Conor Mayo-Wilson, Andrew McKenzie, Lawrence S. Moss, Aristotelis Panagiotopolous, Rohit Parikh, Marcus Pivato, Andrew Powell, Ram Ramanujam, Maxwell Stinchcombe, and Metin Uyanik for conversation and continuing collaboration on the subject of this letter. Khan's work was supported by the 2022 Provost's JHU Discovery Award "Deception and Bad-Faith Communication." Aspects of this work were presented for the Research Seminar in Set Theory at the University of Vienna (March 14, 2024), the CUNY Graduate Center Seminar in Philosophy, Logic and Games (PHILOG; March 29, 2024), and *The 35th International Conference on Game Theory* at Stony Brook (16 July, 2024). The authors extend a final thanks to an anonymous reviewer for a careful reading of the manuscript.

[†] Department of Economics, The Johns Hopkins University, Baltimore, MD 21218. **email** akhan@jhu.edu

[‡] Remote Sensing Earth Systems Institute, The City University of New York (CUNY), New York, New York 10031. **email** app@arthurpaulpedersen.org

[§] Corresponding author

[¶] Institute for Advanced Study in Mathematics, Harbin Institute of Technology, Harbin 150001. **email** davschr@gmail.com

Highlights

- A theorem that formulates the Luce-Raiffa-Aumann (LRA) conjecture in its fullest form to date and has as its corollaries recent theorems of [Adler et al. \[2009\]](#) (ADP) and [Raimondo \[2023\]](#)
- A proof of the theorem that appeals to a result of [Fishburn and Roberts \[1978\]](#) on axiomatizations of multilinear utility theory; a connection not forged before
- An alternative proof of the theorem as a consequence of the ADP result for a game with two-players, each limited to three actions; an observation missed in earlier work
- An open question (to be pursued elsewhere) arising of the re-scaling result concerning the existence of Nash equilibria in games defined on mixture spaces.
- A potential connection of the second proof to the work of [Moulin and Vial \[1978\]](#) on strategically zero-sum games, and thereby to recent work in computer science on fast algorithms for the computation of Nash equilibria

In all of man's written record there has been a preoccupation with conflict of interest... [W]e find today that conflict of interest, both among individuals and among institutions, is one of the more dominant concerns of ... economics, sociology, political science, and other areas to a lesser degree.¹
Luce-Raiffa (1957)

Game theory, a mathematical theory that shares a common foundation in the assumption that actors must be strategic, or individualistically competitive against others, offers a unified methodology and a comprehensive understanding of purposive agency that rejects joint maximization and shared intention, and reduces preference satisfaction to narrow self-interest.²
S. M. Amadae (2015)

1 Introduction

A folk theorem of central importance to the theory of games states that every two-person adversarial game is an affine transformation of a zero-sum game. [Luce and Raiffa \[1957\]](#) appeal to this observation in their exposition of two-person non-cooperative games to justify confining attention

¹ The authors note that “possibly only the topics of God, love, and inner struggle have received comparable attention;” see [[Luce and Raiffa, 1957](#), paragraph 1, page 1]. The reader may also want to look at the excellent, treatment of the subject in [[Luce and Raiffa, 1957](#), Chapters 4 and 5].

² The author notes “Theorists apply the same tools and models to widely divergent fields of investigation: economics, politics, conflict resolution and evolution.” The epigraph is a composite taken from sentences in [[Amadae, 2015](#), pp. vxiii-xx]. The footnotes of her prologue give the most up-to-date bibliography of the interdisciplinary reach of the subject known to the authors.

to zero-sum games in their treatment of adversarial games — called by them *strictly competitive* games.

Decades later, [Aumann \[1987\]](#) gave the contours of a formal description of the folk theorem in his masterful 1987 survey of game theory:

Since randomized strategies are admitted, this condition applies also to mixed outcomes (probability mixtures of pure outcomes). From this it may be seen that a two-person game is strictly competitive if and only if, for an appropriate choice of utility functions, the utility payoffs of the players sum to zero in each square of the matrix (p. 19).

More recently, [Adler et al. \[2009\]](#), henceforth (ADP), have complained that the literature treating the folk theorem has sown confusion and further that no proof exists in full or in outline. They proceed to give two elementary proofs of the folk theorem for finite action sets, one in an algebraic register, the other in a combinatorial one, each being, they contend, “quite non-trivial” (p. 472).

Subsequent work by [Raimondo \[2023\]](#) takes up the folk theorem, who calls it the Luce-Raiffa-Aumann (LRA) conjecture, as we shall henceforth call the folk theorem.³ [Raimondo \[2023\]](#) proves the LRA conjecture for the case of closed interval action sets and continuous utilities, extending the result of ADP for finite action sets. He offers a functional analytic proof by necessity different from the previous ones.

This letter offers a synthetic treatment, and thereby contributes to the theory of two-person games in the following ways:

- (i) It offers a formulation of the LRA conjecture in the form of a general theorem that subsumes the central results of [Adler et al. \[2009\]](#) and [Raimondo \[2023\]](#) and may have interest also for its own sake.
- (ii) It derives the result as a simple consequence of a well-known results of [Fishburn \[1976\]](#) and [Fishburn and Roberts \[1978\]](#) on axiomatizations of multilinear utility theory for n -person games. This connection has so far escaped attention in the antecedent literature.
- (iii) It offers a second proof of the generalized theorem presented here as a simple corollary of the theorem in [Adler et al. \[2009\]](#) for the specific instance of games with two-players, each with three actions. This observation has also escaped attention.
- (iv) Since the work of [Moulin and Vial \[1978\]](#) has received substantial attention by computer scientists in the presentation of efficient algorithm, the relationship we draw to the rescaling theorem presented here has obvious implications for future work.

The material presented below is laid out as follows: Section 2 presents lays out the basic notation and terminology that allows us to present what we refer to as the Fishburn-Roberts theorem. Section 3 presents our main result and uses it to derive the recent theorems of ADP and Raimondo; it also offers two proofs of the result. Section 4 concludes the letter by drawing connections to [Moulin and Vial \[1978\]](#) on strategically zero-sum games, and through it, to recent work in both computer science and economics on computability of equilibria of two-player games.

³ For precise renderings of the LRA conjecture, see Theorem 3.1 below, and its earlier formalizations in [Aumann \[1987\]](#), [Adler et al. \[2009\]](#), [Raimondo \[2023\]](#), and more generally, the discussion in the references in Footnote 1.

2 Notation and Terminology

Given a binary relation \succsim on a set X , we denote its asymmetric and symmetric parts on X by \prec and \sim , respectively. Given an n -tuple (p_1, \dots, p_n) belonging to the n -fold Cartesian product of n sets and $i = 1, \dots, n$, we adopt the usual convention to denote by p_{-i} the $n-1$ -tuple $(p_1, \dots, p_{i-1}, p_{i+1}, \dots, p_n)$ and by (q, p_{-i}) the n -tuple $(p_1, \dots, p_{i-1}, q, p_{i+1}, \dots)$.

An **ordered bilinear mixture space** is a quintuple $\mathcal{M} = (M_1, M_2, \succsim, \oplus_1, \oplus_2)$ consisting of nonempty sets M_1 and M_2 , a binary relation \succsim on $M_1 \times M_2$, and functions \oplus_i for $i = 1, 2$:

$$\oplus_i : [0, 1] \times M_i \times M_i \rightarrow M_i \quad (\alpha, p, q) \mapsto \alpha p \oplus_i (1 - \alpha)q$$

satisfying the following five requirements for all $p, q, r, s \in M_1 \times M_2$, $\alpha, \beta \in [0, 1]$, and $i, j = 1, 2$:

MS1 \succsim is a total preorder on $M_1 \times M_2$;

$$\text{MS2} \quad (\alpha p_i \oplus_i (1 - \alpha)q_i, r_{-i}) \sim ((1 - \alpha)q_i \oplus_i \alpha p_i, r_{-i})$$

$$\text{MS3} \quad \left(\beta(\alpha p_i \oplus_i (1 - \alpha)q_i) \oplus_i (1 - \beta)q_i, r_{-i} \right) \sim (\alpha\beta p_i \oplus_i (1 - \alpha\beta)q_i, r_{-i})$$

MS4 If $p \succ q$ and $q \succ (r_i, p_{-i})$, then there are $\alpha, \beta \in (0, 1)$ such that:

$$\left(\alpha p_i \oplus_i (1 - \alpha)r_i, p_{-i} \right) \succ q \quad \text{and} \quad q \succ \left(\beta p_i \oplus_i (1 - \beta)r_i, p_{-i} \right)$$

MS5 If $p \succ q$ and $(r_i, p_{-i}) \sim (s_j, q_{-j})$, then $(\alpha p_i \oplus_i (1 - \alpha)r_i, p_{-i}) \succ (\alpha q_j \oplus_j (1 - \alpha)s_j, q_{-j})$.

Conditions MS2, MS3, MS4 are the usual mixture space axioms along each dimension, while condition MS5 is an interdimensional reformulation of the usual independence axiom.

A function $u : M_1 \times M_2 \rightarrow \mathbb{R}$ is said to be a **bilinear representation** of \succsim if it satisfies the following two properties for all $p, q, r \in M_1 \times M_2$, $\alpha \in [0, 1]$, and $i = 1, 2$:

$$\text{REP} \quad p \succsim q \quad \text{if and only if} \quad u(p) \leq u(q); \quad \text{and}$$

$$\text{BILIN} \quad u(\alpha p_i \oplus_i (1 - \alpha)q_i, r_{-i}) = \alpha u(p_i, r_{-i}) + (1 - \alpha)u(q_i, r_{-i}).$$

Let X be a set. Recall that a function $f : X \rightarrow \mathbb{R}$ is said to be a **positive affine transformation** of a function $g : X \rightarrow \mathbb{R}$ if there are $\alpha, \beta \in \mathbb{R}$ with $\alpha > 0$ such that $f(x) = \alpha g(x) + \beta$ for all $x \in X$.

We can now present:

Theorem 2.1 (Fishburn and Roberts 1978). *Suppose $\mathcal{M} = (M_1, M_2, \succsim, \oplus_1, \oplus_2)$ is an ordered bilinear mixture space. Then there is a bilinear representation u of \succsim such that any other bilinear representation of \mathcal{M} is a positive affine transformation of u — that is to say, the function u is unique up to a positive affine transformation.*

Proof. See the proof of Theorem 2 from Fishburn and Roberts [1978]. ■

3 The Luce-Raiffa-Aumann Conjecture Reformulated

A **two-person non-cooperative game** is a quadruple $\mathcal{G} = \langle \mathcal{P}_1, \mathcal{P}_2, u_1, u_2 \rangle$ consisting of nonempty sets \mathcal{P}_1 and \mathcal{P}_2 and real-valued functions u_1 and u_2 on the Cartesian product $\mathcal{P}_1 \times \mathcal{P}_2$. It is said to be **adversarial**, or **strictly competitive**, if for all $\sigma, \tau \in \mathcal{P}_1 \times \mathcal{P}_2$:

$$u_1(\sigma) \geq u_1(\tau) \iff u_2(\sigma) \leq u_2(\tau). \quad (1)$$

The game \mathcal{G} is said to be **zero-sum** if equality $u_1(\sigma) + u_2(\sigma) = 0$ obtains for all $\sigma \in \mathcal{P}_1 \times \mathcal{P}_2$.

If in addition each \mathcal{P}_i is convex, the game is said to be **bilinear**, or **bi-affine**, if each function u_i is linear in each coordinate — that is, for all $\sigma, \tau \in \mathcal{P}_1 \times \mathcal{P}_2$, $\alpha, \beta \in [0, 1]$, and $u \in \{u_1, u_2\}$:

$$\begin{aligned} u\left(\alpha\sigma_1 + (1-\alpha)\tau_1, \beta\sigma_2 + (1-\beta)\tau_2\right) &= \alpha u\left(\sigma_1, \beta\sigma_2 + (1-\beta)\tau_2\right) + (1-\alpha)u\left(\tau_1, \beta\sigma_2 + (1-\beta)\tau_2\right) \\ &= \beta u\left(\alpha\sigma_1 + (1-\alpha)\tau_1, \sigma_2\right) + (1-\beta)u\left(\alpha\sigma_1 + (1-\alpha)\tau_1, \tau_2\right). \end{aligned}$$

3.1 Two-Person Adversarial Games are Zero-Sum

Established forthwith is that *every two-person adversarial game is, up to a positive affine transformation, zero-sum*, by appeal to [Theorem 2.1](#).

Theorem 3.1. *Let $\mathcal{G} = \langle \mathcal{P}_1, \mathcal{P}_2, u_1, u_2 \rangle$ be a two-person non-cooperative game. Suppose \mathcal{P}_1 and \mathcal{P}_2 are convex and \mathcal{G} is bilinear. Then the following are equivalent:*

- (a) *Game \mathcal{G} is adversarial;*
- (b) *Function u_2 is a positive affine transformation of function $-u_1$;*
- (c) *There is a positive affine transformation v_1 of u_1 such that $\mathcal{Z} = \langle \mathcal{P}_1, \mathcal{P}_2, v_1, u_2 \rangle$ is zero-sum. ■*

Proof. To show that (a) implies (b), suppose \mathcal{G} is adversarial. Define a binary relation \succsim on $\mathcal{P}_1 \times \mathcal{P}_2$ by setting for all $\sigma, \tau \in \mathcal{P}_1 \times \mathcal{P}_2$:

$$\sigma \succsim \tau \iff -u_1(\sigma) \leq -u_1(\tau).$$

By stipulation, the function $-u_1$ is a bilinear representation of \succsim . Since \mathcal{G} is adversarial, the function u_2 is also a bilinear representation of \succsim . Thus, by [Theorem 2.1](#), it follows that $u_2 = \alpha(-u_1) + \beta$ for some $\alpha, \beta \in \mathbb{R}$ with $\alpha > 0$, whence (b).

To see that (b) implies (c), suppose u_2 is a positive affine transformation of $-u_1$, whereby $u_2 = -\alpha u_1 + \beta$ for some $\alpha, \beta \in \mathbb{R}$ with $\alpha > 0$. Observe that $\mathcal{Z} = \langle \mathcal{P}_1, \mathcal{P}_2, \alpha u_1 - \beta, u_2 \rangle$ is zero-sum, as desired.

For the implication from (c) to (a), it is straightforwardly verified that if $\mathcal{Z} = \langle \mathcal{P}_1, \mathcal{P}_2, v_1, u_2 \rangle$ is zero-sum for some positive affine transformation v_1 of u_1 , then game \mathcal{G} is a adversarial. ■

3.2 Antecedent Results as Corollaries

Given a finite set of actions S , let $\Delta(S)$ denote the set of all simple probability mass function on S . Given a function $\nu : S_1 \times S_2 \rightarrow \mathbb{R}$, let \mathbb{E}_ν denote expected utility $\mathbb{E}_\nu : \Delta(S_1) \times \Delta(S_2) \rightarrow \mathbb{R}$ given by

requiring for all $p \in \Delta(S_1) \times \Delta(S_2)$:

$$\mathbb{E}_\nu(p) := \sum_{(s_1, s_2) \in S_1 \times S_2} p_1(s_1)p_2(s_2)\nu(s_1, s_2)$$

Now consider $\mathcal{G} = \langle \Delta(S_1), \Delta(S_2), \mathbb{E}_{\nu_1}, \mathbb{E}_{\nu_2} \rangle$ for real functions ν_1, ν_2 on $S_1 \times S_2$, and observe that \mathcal{G} is a bilinear two-person non-cooperative game. An immediate corollary of [Theorem 3.1](#) is the main result reported in [[Adler et al., 2009](#), Theorem 1], which we state without proof.

Corollary 3.2 ([Adler et al. 2009](#)). *Let $\mathcal{G} = \langle \Delta(S_1), \Delta(S_2), \mathbb{E}_{\nu_1}, \mathbb{E}_{\nu_2} \rangle$ be an adversarial game based on finite action sets S_1 and S_2 , as above. Then ν_2 is an positive affine transformation of $-\nu_1$.*

We may adopt similar notation to formulate the central result reported in [[Raimondo, 2023](#)]. Let $\Delta_\lambda([0, 1])$ denote the set of probability measures on $[0, 1]$ which are absolutely continuous with respect to Lebesgue measure. Given a continuous function $\xi : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$, again denote expected utility by \mathbb{E}_ξ , so that $\mathbb{E}_\xi : \Delta_\lambda([0, 1]) \times \Delta_\lambda([0, 1]) \rightarrow \mathbb{R}$ and for all $p \in \Delta_\lambda([0, 1]) \times \Delta_\lambda([0, 1])$:

$$\mathbb{E}_\xi(p) := \int \xi(x_1, x_2) (p_1 \otimes p_2) (dx_1 dx_2).$$

As before, let $\mathcal{G} := \langle \Delta_\lambda([0, 1]), \Delta_\lambda([0, 1]), \mathbb{E}_{\xi_1}, \mathbb{E}_{\xi_2} \rangle$ and observe that \mathcal{G} is a bilinear two-person non-cooperative game. A corollary of [Theorem 3.1](#), stated next, is the main result reported in [Raimondo \[2023\]](#).

Corollary 3.3 ([Raimondo 2023](#)). *Suppose $\mathcal{G} = \langle \Delta_\lambda([0, 1]), \Delta_\lambda([0, 1]), \mathbb{E}_{\xi_1}, \mathbb{E}_{\xi_2} \rangle$ is an adversarial game based on a common action set $[0, 1]$ for continuous real functions ξ_1 and ξ_2 , as above. Then ξ_2 is an positive affine transformation of $-\xi_1$.*

Proof. By [Theorem 3.1](#) we obtain immediately that there are $\alpha, \beta \in \mathbb{R}$ with $\alpha > 0$ such that for every $p \in \Delta_\lambda([0, 1]) \times \Delta_\lambda([0, 1])$, $u_{\xi_1}(p) = -\alpha u_{\xi_2}(p) + \beta$. By continuity of ξ_1 and ξ_2 , in fact $\xi_1(p) = -\alpha \xi_2(p) + \beta$. \blacksquare

3.3 An Alternative Proof

Finally, we note that the ADP result, [Corollary 3.2](#), is essentially a special case of our main result, [Theorem 3.1](#), specialized to the case of finitely many actions. One might be curious as to how one could derive the full result from this special case; especially since the proof of the latter given in [Adler et al. 2009](#) is (at least on the surface) quite different from the one given here, as well as from the proofs in [Fishburn and Roberts \[1978\]](#). With this motivation, we give an alternative proof of our main result, [Theorem 3.1](#), presupposing its consequence for finite actions sets, i.e., [Corollary 3.2](#).

For this proof, the following terminology will be useful. Given $\mathcal{P}' \subseteq \mathcal{P}_1 \times \mathcal{P}_2$, call $(\alpha, \beta) \in \mathbb{R}^2$ **compatible with \mathcal{P}'** if $u_2(p) = -\alpha u_1(p) + \beta$ for all $p \in \mathcal{P}'$.

Proof. We will concentrate on the non-trivial implication, (a) \Rightarrow (b). To this end, we make the following observation based upon [Adler et al. 2009](#).

Claim. For any three strategy profiles $\{p^{(1)}, p^{(2)}, p^{(3)}\} \subseteq \mathcal{P}_1 \times \mathcal{P}_2$, there is $(\alpha, \beta) \in \mathbb{R}^2$ with $\alpha > 0$ that is compatible with $\{p^{(1)}, p^{(2)}, p^{(3)}\}$.

To see this claim, consider the game with finite action sets where we take $S_i = \{p_i^{(1)}, p_i^{(2)}, p_i^{(3)}\}$ and define utilities u'_1, u'_2 on $\Delta(S_1) \times \Delta(S_2)$ to be the restrictions of u_1, u_2 to $\Delta(S_1) \times \Delta(S_2)$. We thereby obtain an adversarial game $\langle \Delta(S_1), \Delta(S_2), u'_1, u'_2 \rangle$. By [Corollary 3.2](#), there is $(\alpha, \beta) \in \mathbb{R}^2$ with $\alpha > 0$ such that $u'_2(p) = -\alpha u'_1(p) + \beta$ for all $p \in \Delta(S_1) \times \Delta(S_2)$. In particular, (α, β) is compatible with $\{p^{(1)}, p^{(2)}, p^{(3)}\}$.

[Theorem 3.1](#) now is immediate: Consider a two-person adversarial game $\mathcal{G} = \langle \mathcal{P}_1, \mathcal{P}_2, u_1, u_2 \rangle$. We may assume u_1 is not constant. Fix $p^{(1)}, p^{(2)}$ with $u_1(p^{(1)}) \neq u_1(p^{(2)})$. There exists exactly one $(\alpha, \beta) \in \mathbb{R}^2$ which is compatible with $\{p^{(1)}, p^{(2)}\}$; let us denote this unique pair by (α^*, β^*) . Now let $p^{(3)} \in \mathcal{P}_1 \times \mathcal{P}_2$ be arbitrary. By the claim, there is (α, β) which is compatible with $\{p^{(1)}, p^{(2)}, p^{(3)}\}$. We have $(\alpha, \beta) = (\alpha^*, \beta^*)$ by the uniqueness property of the latter. As $u_2(p^{(3)}) = -\alpha^* u_1(p^{(3)}) + \beta^*$ and $p^{(3)}$ was arbitrary, we are done. ■

4 Concluding Remarks

Two directions stem from the connections between the game theory and the decision theory communities that have been forged in this letter. We leave both for future work.

The first is a consequence of the observation that the central results presented here for convex spaces may be suitably recast in terms of mixture spaces as originally pioneered by [Herstein and Milnor \[1953\]](#). This being said, the question arises as to whether the theory of normal form games articulated by [Nash \[1950, 1951\]](#) and [Debreu \[1952\]](#) can be set in mixture spaces. This would require suitable embedding theorems that take action sets in a mixture-space setting to topological vector spaces, and then bring back the existence results available there.⁴

Second, in an influential paper, [Moulin and Vial \[1978\]](#) introduce the notion of *strategically equivalent games* in the context of two-player games, and use it to characterize games that are equivalent in this sense to a zero-sum game. The connection to these ideas also merits further investigation and resolution, especially so since the Moulin-Vial notion plays a crucial role in the burst of recent activity investigating computational aspects of two-player games by both economists and computer scientists. The second proof of the main result of this letter has special relevance in this context.⁵

A final summary statement. In its focus on two-player games, the basic thrust of this work goes against the grain of the development of non-cooperative game theory in which the generalization of two-player games was sought in n -player games, and even in games with an uncountable continuum player games⁶ The adversarial aspect that is emphasized here concentrates on the them-versus-us aspect without any defensiveness.

⁴ In this connection, a direct proof of the results in Fishburn-Roberts may be useful for workers in the field. We are grateful to an anonymous reviewer for suggesting this.

⁵ For the economic literature, see [Stengel et al. \[2002\]](#), [Savani and Stengel \[2006\]](#), [Tewolde \[2023\]](#), and their references to earlier work: [Lemke and Howson Jr. \[1964\]](#), [Vorob'ev \[1958\]](#) and [Isaacson and Millham \[1980\]](#). For the literature in computer science, see [Kontogiannis and Spirakis \[2012\]](#), [Possieri and Hespanha \[2017\]](#), [Heyman \[2019\]](#), [Heyman and Gupta \[2023\]](#), [Daskalakis et al. \[2009\]](#), [Chen et al. \[2009\]](#) and [Nisan et al. \[2007\]](#).

⁶ These are games in which each agent from a continuum is strategically-negligible but a statistical summary of the plays of all the players has an impact on an individual decision; see [Khan and Sun \[2002\]](#) and their references.

References

- Ilan Adler, Constantinos Daskalakis, and Christos H Papadimitriou. A note on strictly competitive games. In *Internet and Network Economics: 5th International Workshop, WINE 2009, Rome, Italy, December 14-18, 2009. Proceedings 5*, pages 471–474. Springer, 2009.
- S. M. Amadae. *Prisoners of Reason: Game Theory and Neoliberal Political Economy*. MIT Press, 2015.
- Robert J. Aumann. Game theory. In John Eatwell, Murray Milgate, and Peter Newman, editors, *The New Palgrave Dictionary of Economics*, volume 2. Palgrave Macmillan: London, 1987.
- X. Chen, X. Deng, and Teng S. H. Settling the complexity of computing two-player Nash equilibria. *Journal of the ACM (JACM)*, 56(3):1–14, 2009.
- C. Daskalakis, P. W. Goldberg, and C. H. Papadimitriou. The complexity of computing a Nash equilibrium. *SIAM Journal on Computing*, 39:195–259, 2009.
- Gerard Debreu. A social equilibrium existence theorem. *Proceedings of the National Academy of Sciences*, 38(10):886–893, 1952.
- Peter C. Fishburn. Axioms for expected utility in n -person games. *International Journal of Game Theory*, 5(2/3):137–149, 1976.
- Peter C. Fishburn and Fred Roberts. Mixture axioms in linear and multilinear utility theories. *Theory and Decision*, 9(2):161–171, 1978.
- Israel N Herstein and John Milnor. An axiomatic approach to measurable utility. *Econometrica*, 21(2):291–297, 1953.
- Joseph Lee Heyman. *On the computation of strategically equivalent games*. Unpublished Ph.D. dissertation, 2019.
- Joseph Lee Heyman and A. Gupta. Rank reduction in bimatrix games. *International Game Theory Review*, 25(1):1–29, 2023.
- K. Isaacson and C. Millham. On a class of nash-solvable bimatrix games and some related Nash subsets. *Naval Research Logistics Quarterly*, 27(3):407–412, 1980.
- M Ali Khan and Sun. Non-cooperative games with many players. In Robert J. Aumann and Sergiu Hart, editors, *Handbook of Game Theory*, volume 3, Chapter 46, pages 1761–1808. Amsterdam: Elsevier Science, 2002.
- S. Kontogiannis and P Spirakis. On mutual concavity and strategically-zero-sum bimatrix games. *Theoretical Computer Science*, 432(3):64–76, 2012.
- C. E. Lemke and J. T. Howson Jr. Equilibrium points of bimatrix games. *SIAM Journal*, III(2): 413–423, 1964.

- Duncan Luce and Howard Raiffa. *Games and Decisions: An Introduction and Critical Survey*. John Wiley, 1957.
- Hervé Moulin and J P Vial. Strategically zero-sum games: the class of games whose completely mixed equilibria cannot be improved upon. *International Journal of Game Theory*, 7:201–221, 1978.
- John F Nash. Equilibrium points in n -person games. *Proceedings of the National Academy of Sciences*, 36(1):48–49, 1950.
- John F Nash. Non-cooperative games. *Annals of Mathematics*, 54(2):286–295, 1951.
- N. Nisan, T. Roughgarden, E. Tardos, and V. V. Vazirani. *Algorithmic Game Theory*. Cambridge University Press, 2007.
- C. Possieri and J. P. Hespanha. An algebraic geometry approach to compute strategically equivalent bimatrix games. *IFAC-Papers (OnLine)*, 50(1):6116–6121, 2017.
- Roberto Raimondo. Strictly competitive games with infinitely many strategies. *Economics Letters*, 233(6):on line, 2023.
- Rahul Savani and Bernhard von Stengel. Hard to solve bimatrix games. *Econometrica*, 74(2):397–429, 2006.
- Bernhard von Stengel, Antoon van den Elszén, and Dolf Talman. Computing normal form perfect equilibria for extensive two-person games. *Econometrica*, 70(2):693–715, 2002.
- Emmanuel Tewolde. Game transformations that preserve Nash equilibria or best response sets. *ArXiv:2111.00076v2*, pages 101–15, 2023.
- N. N. Vorob’ev. Equilibrium points in bimatrix games. *Theory of Probability and its Applications*, III(3):297–309, 1958.