# To Spend or to Gain: Online Learning in Repeated Karma Auctions

DAMIEN BERRIAUD, Automatic Control Laboratory, ETH Zürich, Switzerland

EZZAT ELOKDA, Automatic Control Laboratory, ETH Zürich, Switzerland

DEVANSH JALOTA, Institute for Computational and Mathematical Engineering, Stanford, United States of America

EMILIO FRAZZOLI, Institute for Dynamic Systems and Control, ETH Zürich, Switzerland

MARCO PAVONE, Institute for Computational and Mathematical Engineering, Stanford, United States of America

FLORIAN DÖRFLER, Automatic Control Laboratory, ETH Zürich, Switzerland

Recent years have seen a surge of artificial currency-based mechanisms in contexts where monetary instruments are deemed unfair or inappropriate, e.g., in allocating food donations to food banks, course seats to students, and, more recently, even for traffic congestion management. Yet the applicability of these mechanisms remains limited in repeated auction settings, as it is challenging for users to learn how to bid an artificial currency that has no value outside the auctions. Indeed, users must jointly learn the value of the currency in addition to how to spend it optimally. In this work, we study the problem of learning to bid in two prominent classes of artificial currency auctions: those in which currency, which users spend to obtain public resources, is only issued at the beginning of a finite period; and those where, in addition to the initial currency endowment, currency payments are redistributed to users at each time step. In the latter class, the currency has been referred to as karma, since users do not only spend karma to obtain public resources but also gain karma for yielding them. In both classes, we propose a simple learning strategy, called *adaptive karma pacing*, and show that this strategy a) is asymptotically optimal for a single user bidding against competing bids drawn from a stationary distribution; b) leads to convergent learning dynamics when all users adopt it; and c) constitutes an approximate Nash equilibrium as the number of users grows. Our results require a novel analysis in comparison to adaptive pacing strategies in monetary auctions, since we depart from the classical assumption that the currency has known value outside the auctions, and moreover consider that the currency is both spent and gained in the class of auctions with redistribution.

### **1** INTRODUCTION

In shared or public resource allocation contexts, the use of monetary instruments to regulate resource consumption is often deemed inequitable (e.g., to manage traffic congestion [2, 10, 15, 36]), inappropriate (e.g., for organ and food donations [27, 30, 35] or course allocations [11]), or simply undesired (e.g., for peer-to-peer file sharing [17, 38] or babysitting services [26]). As a consequence, significant attention has been devoted to the study of non-monetary mechanism design [34], which is known to be challenging due to issues of interpersonal comparability [31] and the lack of a general instrument to manipulate incentives [19, 33].

However, a number of mechanisms have seen some recent success in jointly achieving the objectives of fairness, efficiency, and strategy-proofness when resources are allocated *repeatedly over time* [3, 4, 7, 9, 20–22]. The core principle of these mechanisms is to restrict the number of times the resource can be consumed and let the users trade off when it is most beneficial for them to do so. To achieve these goals, many of these mechanisms employ *artificial currencies* [3, 4, 9, 13, 20, 21, 26, 30], which involves issuing a budget of non-tradable credits or currency to users which they may use to repeatedly bid for resources. In artificial currency mechanisms, users, who may have time-varying and stochastic valuations for the resources, must be strategic in their bidding to not deplete the budget too quickly, and to spare currency for periods when they have the highest valuation for the resources. Thus, artificial currencies serve the dual purpose of monitoring resource consumption and providing a means for users to express their time-varying preferences, resulting in fair and efficient allocations over time.

The literature on artificial currency mechanisms for repeated resource allocation can be broadly categorized in two classes. In the first class, artificial currency is issued at the beginning of a finite episode only to be spent during the episode. This encompasses the vast majority of existing works [3, 4, 9, 20, 21]. In the second class, artificial currency is issued at the beginning of the episode but can also be gained throughout it, typically by means of peer-to-peer exchanges [13, 17, 26, 38] or by redistributing the payments collected in each time step [14, 30]. Some works have referred to this class of artificial currencies as *karma* [13, 38, 39], loosely inspired by the popular notion stemming from Indian tradition: when users yield resources to others they gain karma, and when instead they consume resources they lose karma. This class of karma mechanisms offers some advantages because from the system perspective, the resource allocation can be infinitely repeated with no central intervention other than the initial endowment of karma; and from the user perspective, karma is forgiving as it enables users to immediately replenish their budgets by yielding resources. For simplicity, we adopt the term *karma* to refer to artificial currencies in both of the aforementioned mechanism classes in this paper.

In both classes of karma mechanisms, the focus thus far has been on analyzing equilibrium properties, including existence [13], strategy-proofness [20], and robustness [9, 21]. However, the problem of *learning how to optimally bid karma and other artificial currency in repeated auction settings*, and whether such a learning procedure converges to a Nash equilibrium, remains unaddressed. This problem holds both significant importance and challenge. The importance is two-fold: on one hand, the equilibrium-based analysis of previous studies is only meaningful if an equilibrium is reached; on the other, devising simple learning rules that align with users' self-interest is crucial to implement these mechanisms in practice. The challenge stems from the fact that, unlike traditional monetary instruments, karma does not have any value outside the resource allocation context for which the karma has been issued, and never (directly) enters the utilities of users. Therefore, users must jointly learn the value of karma as well as how to spend it optimally.

The problem of learning how to optimally bid in *monetary* auctions is a classical one [28, 29]. This problem has gained recent traction in the context of repeated, budget-constrained auctions [5, 6, 8,

12, 18, 40], most famously to automate the bidding in multi-period online ad campaigns. We draw inspiration from these works, and in particular [6], in deriving a learning strategy in karma-based auctions, which we call *adaptive karma pacing*. However, the karma setting requires a completely novel analysis in comparison to the monetary setting. The aforementioned feature of karma that it has no value outside the auctions and does not enter in the immediate utilities of the users leads to an important de-stabilizing effect: while in a monetary auction a user will never bid more than the monetary valuation of the resource, in a karma auction a user might deplete the whole budget before learning the correct value of the resource in karma. Moreover, we study a class of auctions in which payments are redistributed to the users in each time step, leading to new strategic opportunities that are not typically considered in a monetary setting. For instance, even in a second-price auction a user has an incentive to bid non-truthfully to maximize the karma gained upon losing. Furthermore, the redistribution leads to challenges in the simultaneous adoption of classical adaptive pacing strategies, as the total karma held by the users is preserved, which makes it impossible for all users to simultaneously deplete their budgets.

#### 1.1 Contribution

In this paper, we devise a simple learning strategy, called *adaptive karma pacing*, that learns to bid optimally in both of the aforementioned classes of karma mechanisms: those in which karma is issued at the beginning of a finite episode only; and those in which additionally, karma is gained throughout the episode by redistributing the collected payments. Adaptive karma pacing uses private observations in a simple online dual ascent scheme to find an optimal multiplier by which to *pace* the rate of karma expenditures. Our contributions are summarized as follows. In both mechanism classes, we prove under mild conditions that adaptive karma pacing satisfies the following desirable features. First, adaptive karma pacing is a best response strategy: it is asymptotically optimal for a single user bidding against competing bids drawn from a stationary distribution. Second, adaptive karma pacing leads to convergent learning dynamics: when all users adopt the strategy, the expected dynamics converge asymptotically to a unique stationary point. Finally, adaptive karma pacing constitutes an *approximate Nash equilibrium*: it is asymptotically in the best selfish interest of each user to follow the strategy when there is a large number of auctions and users. Moreover, as a direct consequence of our results, adaptive karma pacing is truth-revealing: in symmetric settings, it asymptotically leads to bids that are perfectly correlated with the users' time-varying private valuations, implying that resources are allocated efficiently to those who value them most.

The remainder of the paper is organized as follows. In Section 2 we introduce the problem formulation including key definitions and notation. Our main results are then included in Section 3 for the first class of karma mechanisms with no karma redistribution; and Section 4 for the second class of karma mechanisms with karma redistribution. Finally, Section 5 discusses the results, shedding light on the key assumptions made and providing directions for future work. Numerical computation results and detailed proofs are included in the appendix.

#### 2 PROBLEM SETUP

This section introduces the setting studied in the paper, inlcuding notation and important definitions.

### 2.1 Notation

We denote by [*N*] the set  $\{1, ..., N\}$ , by  $\mathbb{1}\{\cdot\}$  the indicator function, by  $(\cdot)^+$  the function  $x \mapsto \max\{x, 0\}$ , and by  $P_{[a,b]}(\cdot)$  the projection  $x \mapsto \min\{\max\{x, a\}, b\}$ . Scalars x are distinguished from vectors  $\mathbf{x} = (x_i)_{i \in I}$ , for some index set I, through the use of boldface. If x is a scalar, then  $\underline{x}$  (respectively,  $\overline{x}$ ) is a lower bound (respectively, upper bound) of x. If  $\mathbf{x}$  is a vector, then  $\underline{x} = \min_{i \in I} x_i$ 

(respectively,  $\overline{x} = \max_{i \in I} x_i$ ). Finally, for the vector x and an index  $i \in I$ , the vector  $x_{-i} = (x_j)_{j \in I, i \neq j}$  is constructed by dropping component *i*.

## 2.2 Setting

We study a general class of repeated resource allocation problems in which a limited number of resources must be repeatedly allocated to a population  $\mathcal{N} = [N]$  of agents. For the sake of presentation, we instantiate this class of problems using a stylized morning commute setting [1, 37], which is schematically illustrated in Figure 1. At discrete time steps  $t \in \mathbb{N}$  (e.g., days), the agents seek to commute from the suburb to the city center using one of two roads. The *general purpose road* is subject to congestion, while access to the *priority road* is limited to its free-flow capacity of  $\gamma \in [N - 1]$  agents per time step, and therefore it remains free of congestion. Travelling on the general purpose road takes time  $\tau \in \mathbb{R}_+$ , while travelling on the priority road takes a shorter time  $0 \le \tau - \Delta < \tau$ . This model can be interpreted as an abstraction of a multi-lane highway with a governed express lane and un-govered, congested general purpose lanes.



Fig. 1. Schematic representation of repeated resource allocation using karma.

At each time step  $t \in \mathbb{N}$ , each agent  $i \in \mathcal{N}$  is associated with a private *valuation of time*  $v_{i,t} \in [0, 1]$  drawn independently across time from fixed, exogenous distributions  $\mathcal{V}_i$ . The valuations represent the agents' time-varying sensitivities to travel delays, e.g., because they have flexible schedules on some days but must be punctual on other days, and are normalized to the interval [0, 1] without loss of generality. We denote by  $v_t = (v_{i,t})_{i \in \mathcal{N}}$  the vector of agents' valuations at time t, which are distributed according to  $\mathcal{V} = \prod_{i \in \mathcal{N}} \mathcal{V}_i$  with support over  $[0, 1]^N$ .

Access to the priority road is governed by means of an artificial currency called *karma*. Each agent  $i \in N$  is endowed with an initial karma budget  $k_{i,1} \in \mathbb{R}_+$ . Then, at each time step  $t \in \mathbb{N}$ , each agent places a sealed bid  $b_{i,t} \in \mathbb{R}_+$  smaller than its current budget  $k_{i,t} \in \mathbb{R}_+$ . The  $\gamma$ -highest bidders, referred to as the 'auction winners', are granted access to the priority road, and must pay  $p_t^{\gamma+1} := \gamma + 1^{\text{th}} - \max\{b_{i,t}\}_{i=1}^N$  in karma. The  $N - \gamma$  remaining agents, referred to as the 'auction losers', must instead take the general purpose road and make no payments. The price  $p_t^{\gamma+1}$  is set by the highest bid among the auction losers, i.e., it corresponds to the second price auction if  $\gamma = 1$ .

Let  $\mathbf{b}_{-i,t} = (b_{j,t})_{j \neq i}$  be the bid profile of agents other than *i*, and  $d_{i,t}^{\gamma} = \gamma^{\text{th}} - \max_{j:j \neq i} \{b_{j,t}\}$  be the associated *competing bid*, since agent *i* must bid higher than  $d_{i,t}^{\gamma}$  to be among the auction winners. We assume that ties in bids do not occur (as is common in the literature [6]; in practice ties could be settled randomly). Let  $x_{i,t} = \mathbb{1} \{b_{i,t} > d_{i,t}^{\gamma}\} \in \{0,1\}$  indicate whether agent *i* is an auction winner at time *t*. Then the agent suffers a cost  $c_{i,t} = v_{i,t}$  ( $\tau - x_{i,t}\Delta$ ) and pays  $z_{i,t} = x_{i,t} d_{i,t}^{\gamma}$  at that time. Notice that the karma payment does not enter in the cost of the agent, unlike in the classical monetary setting [6]. Moreover, after payments are settled, agents may gain an amount  $g_{i,t} \in \mathbb{R}_+$  of karma as

a result of redistributing the payments, as discussed in Section 4. Their budget for the next time step is hence determined by  $k_{i,t+1} = k_{i,t} - z_{i,t} + g_{i,t}$ .

We consider rational agents that aim to minimize their expected total cost over a time horizon *T*. At time *t*, the information available to agent *i* to formulate its bid is the *history*  $\mathcal{H}_{i,t} = \{(v_{i,s}, k_{i,s}, b_{i,s}, z_{i,s}, c_{i,s})_{s=1}^{t-1}, v_{i,t}, k_{i,t}\}$ . A bidding strategy  $\beta_i \in \mathcal{B}$  for agent *i* is thus a sequence of functions  $\beta_i = (\beta_{i,1}, \ldots, \beta_{i,T})$ , where  $\beta_{i,t}$  maps the history  $\mathcal{H}_{i,t}$  to a probability distribution over the set of feasible bids  $[0, k_{i,t}]$ . A profile of strategies for all agents is accordingly denoted by  $\boldsymbol{\beta} = (\beta_i)_{i \in \mathcal{N}}$ . For a fixed agent *i* following strategy  $\beta_i$ , it will be convenient to define three notions of expected total cost, given by

$$C_{i}^{\beta_{i}}\left(\boldsymbol{v}_{i},\boldsymbol{d}_{i}^{\boldsymbol{\gamma}}\right) = \mathbb{E}_{\boldsymbol{b}_{i}\sim\beta_{i}}\left[\sum_{t=1}^{T}c_{i,t}\right] = \mathbb{E}_{\boldsymbol{b}_{i}\sim\beta_{i}}\left[\sum_{t=1}^{T}v_{i,t}\left(\tau - \mathbb{1}\left\{b_{i,t} > d_{i,t}^{\boldsymbol{\gamma}}\right\}\Delta\right)\right],\tag{1}$$

$$C_{i}^{\beta_{i}} = \mathbb{E}_{\boldsymbol{v}_{i} \sim \prod_{l=1}^{T} \mathcal{V}_{i}, \, \boldsymbol{d}_{i}^{\gamma} \sim \prod_{l=1}^{T} \mathcal{D}_{i}^{\gamma}, \, \boldsymbol{b}_{i} \sim \beta_{i}} \left[ C_{i}^{\beta_{i}}(\boldsymbol{v}_{i}, \boldsymbol{d}_{i}^{\gamma}) \right],$$
(2)

$$C_{i}^{\boldsymbol{\beta}} = \mathbb{E}_{\boldsymbol{\nu} \sim \prod_{t=1}^{T} \boldsymbol{\mathcal{V}}, \, \boldsymbol{b}_{-i} \sim \boldsymbol{\beta}_{-i}} \left[ C_{i}^{\beta_{i}} \left( \boldsymbol{\upsilon}_{i}, \boldsymbol{d}_{i}^{\boldsymbol{\gamma}} \left( \boldsymbol{b}_{-i} \right) \right) \right].$$
(3)

Equation (1) defines the sample path cost  $C_i^{\beta_i}(v_i, d_i^{\gamma})$  for a fixed realization of valuations  $v_i = (v_{i,t})_{t \in [T]}$  and competing bids  $d_i^{\gamma} = (d_{i,t}^{\gamma})_{t \in [T]}$ . Equation (2) defines the stationary competition cost  $C_i^{\beta_i}$ , in which the competing bids  $d_{i,t}^{\gamma}$  are assumed to be drawn independently across time from a stationary distribution  $\mathcal{D}_i^{\gamma}$ . Finally, Equation (3) defines the strategic competition cost  $C_i^{\beta}$ , which is agent *i*'s expected total cost when all agents follow strategy profile  $\beta$ . With this, we will say that strategy profile  $\beta$  constitutes an  $\varepsilon$ -Nash equilibrium if its strategic competition cost satisfies, for all agents  $i \in \mathcal{N}$ ,

$$\lim_{T \to \infty} \frac{1}{T} \left( C_i^{\beta} - \inf_{\tilde{\beta}_i \in \mathcal{B}} C_i^{\tilde{\beta}_i, \beta_{-i}} \right) = 0.$$
(4)

Equation (4) implies that under strategy profile  $\beta$ , no single agent *i* can asymptotically improve its expected average cost per time step by unilaterally deviating to a strategy  $\tilde{\beta}_i \neq \beta_i$ .

## 3 KARMA ISSUED AT THE START OF AN EPISODE

In this section, we study the first class of karma mechanisms where agents do not gain karma other than at the beginning of an episode of length T, i.e.,  $g_{i,t} = 0$  for all  $i \in N$ ,  $t \in [T]$ . This episodic setting represents when karma is issued periodically, e.g., at the beginning of every month or year, and does not carry over episodes [3, 4, 9, 20, 21].

Our main goal is to devise a candidate bidding strategy that constitutes an  $\varepsilon$ -Nash equilibrium when all agents follow it. To achieve this goal, this section proceeds as follows. In Section 3.1, we devise a candidate strategy that learns to bid optimally, called *adaptive karma pacing*, by studying the optimal cost an agent can achieve with the benefit of hindsight. Sections 3.2–3.4 then take required steps in the analysis of adaptive karma pacing. Section 3.2 establishes that adaptive karma pacing is asymptotically optimal for a single agent bidding against competing bids drawn from a stationary distribution (Theorem 3.1). Section 3.3 establishes that the learning dynamics converge to a unique stationary point when all agents adopt adaptive karma pacing (Theorem 3.2). Finally, Section 3.4 combines these results to achieve the main goal of proving that adaptive karma pacing constitutes an  $\varepsilon$ -Nash equilibrium under suitable conditions (Theorem 3.3).

## 3.1 Derivation of Adaptive Karma Pacing

This section derives a candidate optimal bidding strategy using an *online dual gradient ascent* scheme, a classical optimization technique that has gained recent traction in the context of budget-constrained auctions [6] and other related problems [8, 23–25]. To elucidate our bidding strategy, we first introduce the optimization problem of a single agent  $i \in N$  who has the *benefit of hindsight*, i.e., who can make optimal bidding decisions with prior knowledge of the future realizations of valuations  $v_i$  and competing bids  $d_i^{\gamma}$ . Thus, the optimal cost of this problem serves as a theoretical benchmark for the lowest cost that the agent can hope to achieve. Then, since in practice the agent only observes the stochastic valuations and competing bids online as the auctions progress, we introduce a candidate bidding strategy, based on online gradient ascent, to approximate the agent's optimal bidding strategy with the benefit of hindsight.

**Optimal Cost with the Benefit of Hindsight.** For a fixed realization of valuations  $v_i$  and competing bids  $d_i^{\gamma}$ , agent *i*'s optimal cost with the benefit of hindsight is given by the following optimization problem

$$C_{i}^{H}(\boldsymbol{v}_{i}, \boldsymbol{d}_{i}^{\gamma}) = \min_{\boldsymbol{x}_{i} \in \{0,1\}^{T}} \sum_{t=1}^{T} v_{i,t}(\tau - x_{i,t}\Delta), \quad \text{s.t.} \sum_{t=1}^{T} x_{i,t} \boldsymbol{d}_{i,t}^{\gamma} \le \rho_{i} T.$$
(5)

In Problem (5), we use auction outcomes  $x_{i,t}$  directly as decision variables rather than bids  $b_{i,t}$ . This is because given competing bids  $d_{i,t}^{\gamma}$ , bids  $b_{i,t}$  affect the cost and the karma budget only through  $x_{i,t} = \mathbb{1}\left\{b_{i,t} > d_{i,t}^{\gamma}\right\}$ . We also define  $\rho_i = k_{i,1}/T$  as the agent's *target expenditure rate*, i.e., the average expenditure per time step that will fully deplete the initial budget by the end of the time horizon.

The Lagrangian dual problem associated with Problem (5) is given by

$$C_i^H(\boldsymbol{v}_i, \boldsymbol{d}_i^{\boldsymbol{\gamma}}) \ge \delta_i^H(\boldsymbol{v}_i, \boldsymbol{d}_i^{\boldsymbol{\gamma}}) = \sup_{\mu_i \ge 0} \min_{\boldsymbol{x}_i \in \{0,1\}^T} \sum_{t=1}^T \left( x_{i,t} (\mu_i d_{i,t}^{\boldsymbol{\gamma}} - \Delta \boldsymbol{v}_{i,t}) + v_{i,t} \tau - \mu_i \rho_i \right)$$
(6a)

$$= \sup_{\mu_i \ge 0} \sum_{t=1}^{T} v_{i,t} \tau - \mu_i \rho_i - (\Delta v_{i,t} - \mu_i d_{i,t}^{\gamma})^+ := \sup_{\mu_i \ge 0} \sum_{t=1}^{T} \delta_{i,t} (v_{i,t}, d_{i,t}^{\gamma}, \mu_i).$$
(6b)

Notice that for a fixed multiplier  $\mu_i \ge 0$ , the inner minimum in (6a) is obtained by winning all auctions satisfying  $\Delta v_{i,t} > \mu_i d_{i,t}^{\gamma}$ . This can be achieved by bidding  $b_{i,t} = \Delta v_{i,t}/\mu_i$ , and yields (6b).

**Adaptive Karma Pacing.** We perform a stochastic gradient ascent scheme in order to approximately solve the dual Problem (6) using online observations. Namely, the agent considers a candidate optimal multiplier  $\mu_{i,t}$  and places its bid accordingly with  $b_{i,t} = \Delta v_{i,t}/\mu_{i,t}$ . It then updates  $\mu_{i,t+1}$  using the subgradient given by  $\frac{\partial \delta_{i,t}}{\partial \mu_{i,t}}(v_{i,t}, d_{i,t}^{\gamma}, \mu_{i,t}) = d_{i,t}^{\gamma} \mathbb{1}\{b_{i,t} > d_{i,t}^{\gamma}\} - \rho_i = z_{i,t} - \rho_i$ . This yields the *adaptive karma pacing strategy*, which is denoted by *A* and summarized in Algorithm 1.

The term 'adaptive karma pacing' is in line with the literature on budget-constrained monetary auctions [6, 16], as the strategy attempts to *pace* the budget depletion rate to match the target rate  $\rho_i$ . Indeed, expenses  $z_{i,t} > \rho_i$  increase  $\mu_{i,t+1}$ , effectively reducing future bids; and vice versa for  $z_{i,t} < \rho_i$ . However, an important novelty in strategy *A* is that the denominator in the bid (*A-b*) is  $\mu_{i,t}$  instead of  $\mu_{i,t} + 1$ , as common in the monetary setting. This is a consequence of the fact that the valuation in karma is not known a-priori; and could lead to a rapid depletion of the budget if  $\mu_{i,t}$  becomes small during the learning process even for a short transient period. For this reason, it is necessary to introduce the lower bound  $\mu_i$  in Algorithm 1, which we note is unlike the corresponding adaptive pacing algorithms in the monetary setting [6].

### ALGORITHM 1: Adaptive Karma Pacing A

#### Input:

- Time horizon *T*, target expenditure rate  $\rho_i > 0$ , multiplier bounds  $\overline{\mu}_i > \underline{\mu}_i > 0$ , gradient step size  $\epsilon_i > 0$ . Initialize:

- Initial multiplier  $\mu_{i,1} \in [\mu_i, \overline{\mu_i}]$ , initial budget  $k_{i,1} = \rho_i T$ .

**for** *t*=1,..., *T*: **do** 

(1) Observe the realized valuation  $v_{i,t}$  and place bid

$$b_{i,t} = \min\left\{\frac{\Delta v_{i,t}}{\mu_{i,t}}, k_{i,t}\right\}; \tag{A-b}$$

(2) Observe the expenditure  $z_{i,t}$  and update the multiplier

$$\mu_{i,t+1} = P_{\left[\mu_{i,\overline{\mu}_{i}}\right]} \left(\mu_{i,t} + \epsilon_{i}(z_{i,t} - \rho_{i})\right), \qquad (A-\mu)$$

as well as the karma budget  $k_{i,t+1} = k_{i,t} - z_{i,t}$ .

end

## 3.2 Asymptotic Optimality under Stationary Competition

This is the first of three sections whose main goal is to establish that our previously derived adaptive karma pacing strategy *A* constitutes an  $\varepsilon$ -Nash equilibrium when adopted by all agents. In this section, we take the first step towards this goal. Namely, we establish that strategy *A* is *asymptotically optimal in a stationary competition setting*, where a single agent *i* bids against competing bids  $d_i^{\gamma} = (d_{i,t}^{\gamma})_{t \in [T]}$  drawn independently across time from a fixed distribution  $\mathcal{D}_i^{\gamma}$ .

This section, as well as Sections 3.3 and 3.4, are organized similarly as follows. We first introduce required definitions and preliminaries; then state our assumptions; before presenting and discussing the main result of the section (Theorem 3.1 in this case). Detailed proofs are included the appendix.

**Preliminaries.** To state the main result of this section, we must first define the *expected dual objective* and *expected expenditure* when agent *i* follows strategy *A*. For a fixed multiplier  $\mu_i > 0$ , these two quantities are respectively given by

$$\Psi_{i}(\mu_{i}) = \mathbb{E}_{v_{i}, d_{i}^{\gamma}} \left[ v_{i}\tau - \mu_{i}\rho_{i} - (\Delta v_{i} - \mu_{i}d_{i}^{\gamma})^{+} \right], \quad (7) \qquad \qquad Z_{i}(\mu_{i}) = \mathbb{E}_{v_{i}, d_{i}^{\gamma}} \left[ d_{i}^{\gamma} \,\mathbb{1}\{\Delta v_{i} > \mu_{i}d_{i}^{\gamma}\} \right], \quad (8)$$

where the expectation is with respect to the stationary distributions  $\mathcal{V}_i$  and  $\mathcal{D}_i^{\gamma}$ , and we omitted the subscript t in  $\mu_{i,t}$ ,  $v_{i,t}$  and  $d_{i,t}^{\gamma}$  for ease of notation. Notice that in Equations (7)–(8), it is assumed that the budget constraint in the bid (*A-b*) does not become active. This is without loss of generality as we prove that asymptotically, the budget constraint indeed never becomes active until the end of the horizon *T*. The maximum expected dual objective is denoted by  $\Psi_i(\mu_i^*)$ , where  $\mu_i^* > 0$  is a multiplier achieving the maximum. Finally, the *average expected regret* of strategy *A* is given by

$$\mathcal{R}_i^A = \frac{1}{T} \mathop{\mathbb{E}}_{\boldsymbol{v}_i, \, \boldsymbol{d}_i^{\boldsymbol{\gamma}}} \left[ C_i^A(\boldsymbol{v}_i, \boldsymbol{d}_i^{\boldsymbol{\gamma}}) - C_i^H(\boldsymbol{v}_i, \boldsymbol{d}_i^{\boldsymbol{\gamma}}) \right].$$
(9)

Notice that achieving  $\mathcal{R}_i^A = 0$  is a sufficient condition for the optimality of strategy A under stationary competition, since the expected optimal cost with the benefit of hindsight  $\mathbb{E}_{\boldsymbol{v}_i, \boldsymbol{d}_i^{\boldsymbol{\gamma}}} \left[ C_i^H(\boldsymbol{v}_i, \boldsymbol{d}_i^{\boldsymbol{\gamma}}) \right]$  provides a lower bound on the stationary competition cost (2) of any strategy.

**Assumptions.** To prove strategy *A*'s asymptotic regret bound in the stationary competition setting, we require a few technical assumptions which are listed and briefly discussed below.

ASSUMPTION 1 (CONTINUITY OF VALUATIONS). For all agents  $j \in N$ , the valuation distribution  $\mathcal{V}_j$  is absolutely continuous with bounded density  $v_j : [0, 1] \mapsto [\underline{v}, \overline{v}] \subset \mathbb{R}_{>0}$ .

Assumption 2 (Differentiability and Concavity). The following conditions hold:

2.1 The expected dual objective  $\Psi_i(\mu_i)$  is differentiable on  $\mathbb{R}_+$  with derivative  $\Psi'_i : \mu_i \mapsto Z_i(\mu_i) - \rho_i$ ; 2.2 On  $\left[\underline{\mu}, \overline{\mu}\right]$ , the expected expenditure  $Z_i(\mu_i)$  is strongly decreasing with parameter  $\lambda > 0$ , and twice differentiable with first and second derivatives bounded by  $\overline{Z'} > 0$  and  $\overline{Z''} > 0$ , respectively.

Assumption 3 (Parameter Design). The input parameters of strategy A satisfy  $\mu_i < \mu_i^* < \overline{\mu}_i$ ,

$$\underline{\mu_i} < \frac{\Delta}{\rho_i} < \overline{\mu}_i, \text{ and } \epsilon_i \le \min\left\{\Delta \underline{\mu_i}^2, \frac{1}{2\lambda}, \frac{1}{\overline{Z'}}, \frac{\mu_i^\star - \underline{\mu_i}}{2\rho_i}\right\}$$

ASSUMPTION 4 ( $\epsilon_i(T)$  UNDER STATIONARY COMPETITION). The gradient step size  $\epsilon_i$  is a function of the time horizon T satisfying  $\epsilon_i(T) \xrightarrow[T \to \infty]{} 0$  and  $T\epsilon_i(T) \xrightarrow[T \to \infty]{} \infty$ .

Assumption 1 is a fundamental assumption on the valuation distributions that is common in the literature [6, 12, 20] and will be used throughout the paper. The technical Assumption 2 is also common in the optimization literature [6, 7] and implies strong concavity of  $\Psi_i$ , which ensures the existence of a unique maximizer  $\mu_i^*$  and the convergence of gradient-based techniques. It is guaranteed to hold under mild conditions on the distributions  $\mathcal{V}_i$  and  $\mathcal{D}_i^{\gamma}$  which we include in Appendix D.1. Assumption 3 regards the choice of the *input parameters of strategy A* and can be satisfied by design, i.e., by setting  $\mu_i$  and  $\epsilon_i$  sufficiently low and  $\overline{\mu_i}$  sufficiently high. Finally, Assumption 4 requires to vary the gradient step size as a function of the time horizon *T*, which is needed to control the average expected regret bound of strategy *A*.

**Main Result.** With these assumptions, we are ready to state the main result of this section regarding the asymptotic optimality of strategy *A* under stationary competition.

THEOREM 3.1 (ASYMPTOTIC OPTIMALITY UNDER STATIONARY COMPETITION). Fix an agent  $i \in N$ , and let Assumptions 1, 2 and 3 hold. Then, there exist a constant  $C \in \mathbb{R}_+$  such that the agent's average expected regret for following strategy A satisfies

$$\frac{1}{T} \mathbb{E}_{\boldsymbol{v}_i, \boldsymbol{d}_i^{\boldsymbol{\gamma}}} \left[ C_i^A(\boldsymbol{v}_i, \boldsymbol{d}_i^{\boldsymbol{\gamma}}) - C_i^H(\boldsymbol{v}_i, \boldsymbol{d}_i^{\boldsymbol{\gamma}}) \right] \le C \left( \epsilon_i + \frac{1 + \epsilon_i}{\epsilon_i T} \right).$$

Let in addition Assumption 4 hold. Then, strategy A achieves sub-linear regret and asymptotically converges to the expected optimal cost, i.e.,

$$\lim_{T \to \infty} \frac{1}{T} \mathbb{E}_{\boldsymbol{v}_i, \boldsymbol{d}_i^{\boldsymbol{\gamma}}} \left[ C_i^A(\boldsymbol{v}_i, \boldsymbol{d}_i^{\boldsymbol{\gamma}}) - C_i^H(\boldsymbol{v}_i, \boldsymbol{d}_i^{\boldsymbol{\gamma}}) \right] = 0$$

Moreover, the convergence rate is at least  $O(T^{-1/2})$  with the choice of  $\epsilon_i \propto T^{-1/2}$ .

The detailed proof of Theorem 3.1 is included in Appendix C.2, and we provide here a sketch of the proof. The average expected optimal cost with the benefit of hindsight is lower-bounded by the maximum expected dual objective  $\Psi_i(\mu_i^*)$ . Meanwhile, the average stationary competition cost of strategy *A* is upper-bounded in terms of  $\Psi_i(\mu_i^*)$  through a Taylor expansion in  $\mu_i^*$ . Controlling this upper bound then yields the desired result. This requires to control a) the budget depletion time such that  $k_{i,1}$  is depleted towards the end of the horizon *T*, as assumed in the definition of  $\Psi_i$ ; and b) the expected distance of the multiplier iterates to the optimal multiplier  $\mu_{i,t} - \mu_i^*$  such that  $\mu_{i,t}$  converges asymptotically to  $\mu_i^*$ . These two objectives are achieved with a suitable choice of the gradient step size  $\epsilon_i$  as per Assumptions 3 and 4.

## 3.3 Convergence under Simultaneous Learning

In this section, we take the next step towards our main goal of establishing that adaptive karma pacing strategy A constitutes an  $\varepsilon$ -Nash equilibrium when adopted by all agents. Namely, we establish that the learning dynamics *converge in the simultaneous learning setting* in which all agents follow strategy A, denoted by joint strategy profile A. The exact notion of convergence considered is presented in the main result of the section, Theorem 3.2.

**Preliminaries.** To state the main result of this section, we must first introduce some definitions. Let  $\mu_t \in \mathbb{R}^N_+$  be the *multiplier profile* stacking the multipliers  $\mu_{i,t}$  of all agents  $i \in \mathcal{N}$ . We adapt the previous definitions of the *expected dual objective* and the *expected expenditure* respectively as

$$\Psi_{i}(\boldsymbol{\mu}) = \mathbb{E}_{\boldsymbol{v}} \left[ v_{i}\tau - \mu_{i}\rho_{i} - (\Delta v_{i} - \mu_{i}d_{i}^{\gamma})^{+} \right], \quad (10) \qquad \qquad Z_{i}(\boldsymbol{\mu}) = \mathbb{E}_{\boldsymbol{v}} \left[ d_{i}^{\gamma} \,\mathbb{1}\{\Delta v_{i} > \mu_{i}d_{i}^{\gamma}\} \right]. \tag{11}$$

Compared to (7)–(8), the expectation is now with respect to the profile of valuations  $\boldsymbol{v} \sim \boldsymbol{V}$ , as given  $\boldsymbol{v}$  and  $\boldsymbol{\mu}$  the competing bid  $d_i^{\gamma} = \gamma^{\text{th}} - \max_{j:j \neq i} \{\Delta v_j / \mu_j\}$  can be uniquely determined. We will aim to show that  $\boldsymbol{\mu}_t$  converges to a *stationary multiplier profile*, which is a multiplier profile  $\boldsymbol{\mu}^{\star} \in \boldsymbol{U} = \prod_{i \in \mathcal{N}} \left( \underline{\mu}_i, \overline{\mu}_i \right)$  satisfying  $Z_i(\boldsymbol{\mu}^{\star}) = \rho_i$ , for all agents  $i \in \mathcal{N}$ . This multiplier profile is stationary in the sense that in expectation, update rule  $(A - \mu)$  will yield  $\boldsymbol{\mu}_{i,t+1}^{\star} = \boldsymbol{\mu}_{i,t}^{\star}$  for all agents i, since the expected expenditures  $Z_i(\boldsymbol{\mu}^{\star})$  equal the target rates  $\rho_i$ .

**Assumptions.** To prove convergence of the learning dynamics under strategy profile *A*, we require to adapt a few of the assumptions of Section 3.2 for the multi-agent setting.

ASSUMPTION 5 (MONOTONICITY). The expected expenditure  $Z(\mu)$  is  $\lambda$ -strongly monotone over U with parameter  $\lambda > 0$ , i.e., for all  $\mu, \mu' \in U$ , it holds that  $(\mu - \mu')^{\top}(Z(\mu) - Z(\mu')) \leq -\lambda \|\mu - \mu'\|_{2}^{2}$ .

Assumption 6 ( $\epsilon(T)$  under simultaneous learning). The gradient step size profile  $\epsilon = (\epsilon_i)_{i \in N}$ is a function of the time horizon T satisfying  $\overline{\epsilon}^2(T)/\underline{\epsilon}(T) \xrightarrow[T \to \infty]{} 0$  and  $T\underline{\epsilon}^2(T)/\overline{\epsilon}(T) \xrightarrow[T \to \infty]{} \infty$ .

Assumption 5 extends Assumption 2.2 for the multi-agent setting. It similarly ensures that the stationary multiplier profile  $\mu^{\star}$  is unique if it exists, as shown in Appendix D.2. It is also common in the literature, in fact, [6] shows that it is implied by a *diagonal strict concavity* condition [32], and proves that it holds in symmetric settings. Assumption 6 in turn extends Assumption 4.

**Main Result.** With these assumptions, we are ready to state the main result of this section regarding the asymptotic convergence of strategy profile *A*. We consider two notions of convergence: the first is with respect to the multiplier profile iterates  $\mu_t$ ; and the second is with respect to the strategic competition costs  $C_i^A$  of all agents  $i \in N$ , defined in Equation (3).

THEOREM 3.2 (CONVERGENCE UNDER SIMULTANEOUS LEARNING). Let Assumptions 1 and 5 hold, and in addition let Assumption 3 hold for all agents  $i \in N$ . Then there exists a constant  $C_1 \in \mathbb{R}_+$  such that the average expected distance to the stationary multiplier profile  $\mu^*$  satisfies

$$\frac{1}{T}\sum_{t=1}^{T} \mathbb{E}_{\boldsymbol{v}}\left[\|\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}\|_{2}^{2}\right] \leq C_{1}N\left(\frac{\overline{\epsilon}^{2}}{\underline{\epsilon}} + \frac{\overline{\epsilon} + \underline{\epsilon}^{2}}{\underline{\epsilon}^{2}T}\right).$$
(12)

Moreover, there exists a constant  $C_2 \in \mathbb{R}_+$  such that the strategic competition cost of strategy profile A satisfies, for all agents  $i \in N$ ,

$$\frac{1}{T}C_i^A - \Psi_i(\boldsymbol{\mu^{\star}}) \le C_2\left(\frac{1}{T} + N\left(\frac{\overline{\epsilon}^{1/2}}{\underline{\epsilon}^{3/2}T} + \frac{\overline{\epsilon}}{\underline{\epsilon}^{1/2}}\right)\right).$$
(13)

Let in addition Assumption 6 hold. Then, the multiplier profile  $\mu_t$  converges in expectation to the stationary multiplier profile  $\mu^*$ , and the average strategic competition cost  $C_i^A$  converges to the optimal

expected dual objective  $\Psi_i(\mu^{\star})$  for all agents  $i \in N$ , i.e.,

$$\lim_{T \to \infty} \frac{1}{T} \sum_{t=1}^{T} \mathbb{E}\left[ \|\boldsymbol{\mu}_t - \boldsymbol{\mu}^{\star}\|_2^2 \right] = 0, \qquad \lim_{T \to \infty} \frac{1}{T} C_i^A - \Psi_i(\boldsymbol{\mu}^{\star}) = 0.$$
(14)

The detailed proof of Theorem 3.2 is included in Appendix C.3. It follows similar arguments as Theorem 3.1 to show bound (12). The main step from bound (12) to bound (13) is to show that the profiles of expected dual objectives  $\Psi(\mu)$  and expected expenditures  $Z(\mu)$  are Lipschitz continuous in  $\mu$ , which can be guaranteed by the continuity Assumption 1. Finally, bounds (12)–(13) straightforwardly imply Equation (14) under Assumption 6.

#### 3.4 Approximate Nash equilibrium in Parallel Auctions

In this section, we finally combine the results of the previous two sections to achieve the main goal of establishing that the profile of adaptive karma pacing strategies A constitutes an  $\varepsilon$ -Nash equilibrium under suitable conditions.

Notice that one cannot immediately conclude that strategy profile A constitutes an  $\varepsilon$ -Nash equilibrium despite of the asymptotic guarantee on the strategic competition costs in Equation (14). This is because  $\Psi_i(\mu^*)$  in Equation (14) lower bounds the *stationary competition cost* rather than the strategic competition cost. Under strategy profile A, the multiplier profile converges asymptotically to  $\mu^*$  by Equation (14), and thereby agent *i*'s distribution of competing bids indeed becomes stationary. However, there could be a possibility for agent *i* to improve its cost by unilaterally deviating to a strategy  $\beta_i \neq A$  that causes non-convergence of  $\mu_t$  and violation of the stationary competition. For this reason, we consider a natural extension of our setting in which the effect of any single agent on the multipliers of others becomes negligible as the number of agents grows. This is referred to as the *parallel auction setting*, which was considered previously in monetary auctions [6].

**Preliminaries: Parallel Auctions.** Let there be  $M \ge 1$  auctions that are held in parallel at each time step  $t \in [T]$ . The number M could represent different priority roads, or the same road accessed at different times of the day, or a combination thereof. Each agent  $i \in N$  participates in one auction  $m_{i,t} \in [M]$  per time step, where  $m_{i,t}$  is drawn independently across agents and time from a fixed distribution  $\pi_i = (\pi_{i,m})_{m \in [M]}$ ; each  $\pi_{i,m}$  denotes the probability for agent i to participate in auction m. We adapt the definition of competing bids accordingly as  $d_{i,t}^{\gamma} = \gamma^{\text{th}} - \max_{j:j \ne i} \{\mathbbm{1}\{m_{j,t} = m_{i,t}\}b_{j,t}\}$ . The distributions  $(\pi_i)_{i \in N}$  yield matching probabilities  $a_i = (a_{i,j})_{j \ne i}$ , where  $a_{i,j} = \mathbb{P}\{m_j = m_i\}$  denotes the probability that agent j is matched with agent i in the same auction. It is straightforward to show that the previous Theorems 3.1 and 3.2 also hold in the extended parallel auction setting. To prove our  $\varepsilon$ -Nash equilibrium result, we require an additional assumption on the matching probabilities that effectively limits the interactions between any pair of agents.

Assumption 7 (MATCHING PROBABILITIES). The matching probabilities  $(a_i)_{i \in N}$  satisfy:

7.1 The fraction of potential auction winners  $\frac{M\gamma}{N}$  is constant for all  $N \in \mathbb{N}$ ;

7.2 There exists a constant  $\kappa > 0$  such that  $\sqrt{N} \max_{i \in \mathcal{N}} ||a_i||_2 \le \kappa$  for all  $N \in \mathbb{N}$ ;

7.3 It holds that 
$$\frac{\overline{\epsilon}}{\underline{\epsilon}} \max_{i \in \mathcal{N}} \|\boldsymbol{a}_i\|_2^2 \xrightarrow[T,N,M\to\infty]{} 0.$$

Notice that Assumptions 7.2 and 7.2 are implied by Assumption 7.1 in many cases, e.g. when agents are assigned to auctions uniformly at random. In this case,  $a_{i,j} = 1/M^2$  for all  $i \neq j$ , and  $\sqrt{N} ||a_i||_2 = N/M$  is constant.

**Main Result.** With this assumption, we are ready to state the main result of this section, namely, that strategy profile A constitutes an  $\varepsilon$ -Nash equilibrium in the parallel auction setting.

THEOREM 3.3 ( $\varepsilon$ -NASH EQUILIBRIUM). Fix an agent  $i \in N$ . Let Assumptions 1 and 5 hold, and in addition let Assumption 3 hold for all other agents  $j \neq i$ . Then there exists a constant  $C \in \mathbb{R}_+$  such that agent i can decrease its average strategic competition cost by deviating from strategy A to any strategy  $\beta_i \in \mathcal{B}$  by at most

$$\frac{1}{T}\left(C_{i}^{A}-C_{i}^{\beta_{i},A_{-i}}\right) \leq C\left(\frac{1+\epsilon_{i}}{\epsilon_{i}T}+\|\boldsymbol{a}_{i}\|_{2}^{2}\frac{\overline{\epsilon}}{\underline{\epsilon}}+\sqrt{N}\|\boldsymbol{a}_{i}\|_{2}\left(\frac{\overline{\epsilon}^{1/2}}{\underline{\epsilon}^{3/2}T}+\frac{\overline{\epsilon}}{\underline{\epsilon}^{1/2}}\right)\right).$$
(15)

Let in addition Assumptions 6 and 7 hold. Then strategy profile A constitutes an  $\varepsilon$ -Nash equilibrium, i.e., it holds for all agents  $i \in N$ ,

$$\lim_{T,N,M\to\infty}\frac{1}{T}\left(C_i^A - \inf_{\beta_i\in\mathcal{B}}C_i^{\beta,A_{-i}}\right) = 0.$$

The detailed proof of Theorem 3.3 is included in Appendix C.4. It consists in lower-bounding the average expected cost under strategy  $\beta_i$  in terms of the optimal expected dual objective  $\Psi_i(\mu^*)$ , and showing that asymptotically agent *i* cannot affect the multiplier profile of the other agents which converges to  $\mu^*$ . This means that the competition becomes stationary, for which strategy *A* is indeed optimal.

This achieves our main goal for the first class of karma mechanisms in which agents do not gain karma other than at the beginning of the episode. We have devised the simple adaptive karma pacing strategy A and provided conditions in which it constitutes an  $\varepsilon$ -Nash equilibrium.

## 4 KARMA GAINED THROUGHOUT AN EPISODE

In this section, we study the second class of karma mechanisms where in addition to initial endowments, agents also gain karma throughout the episode. We focus specifically on karma gains resulting from *payment redistribution*. We consider a simple redistribution rule in which the total karma payment at each time step  $t \in [T]$  is uniformly redistributed to all the agents  $i \in N$ , yielding karma gains  $g_{i,t} = \gamma p_t^{\gamma+1}/N$ . Notice that under this scheme, agents that access the priority road have a net decrease in karma, while those using the general purpose road have a net increase in karma. Meanwhile, the total amount of karma in the system set by the initial endowments  $k_1$  is preserved over time.

This section follows a similar structure as Section 3: Section 4.1 derives an adaptive karma pacing strategy with gains; Section 4.2 establishes the asymptomatic optimality of this strategy in the stationary competition setting (Theorem 4.1); Section 4.3 proves convergence of the learning dynamics when all agents follow the strategy (Theorem 4.2); and finally Section 4.4 achieves the main goal of proving that the strategy constitutes an  $\varepsilon$ -Nash equilibrium in the parallel auction setting (Theorem 4.3).

## 4.1 Derivation of Adaptive Karma Pacing with Gains

In this section, we derive a candidate optimal bidding strategy in the setting with karma gains using an online dual gradient ascent scheme. As in Section 3.1, we first study the optimal cost with the benefit of hindsight for an agent *i*, before deriving a bidding strategy that approximately optimizes that cost using online stochastic dual ascent.

**Optimal Cost with the Benefit of Hindsight.** For a fixed realizations of valuations  $v_i$  and competing bids  $d_i := \left(d_{i,t}^{\gamma}, d_{i,t}^{\gamma+1}\right)_{t \in [T]}$ , agent *i*'s optimal cost with the benefit of hindsight is given

by the following optimization problem

$$C_{i}^{H}(\boldsymbol{v}_{i}, \boldsymbol{d}_{i}) = \min_{\boldsymbol{b}_{i} \in \mathbb{R}_{+}^{T}} \sum_{t=1}^{I} \boldsymbol{v}_{i,t} \left( \tau - \mathbb{1} \left\{ b_{i,t} > d_{i,t}^{\gamma} \right\} \Delta \right),$$
s.t. 
$$\sum_{t=1}^{s} \mathbb{1} \left\{ b_{i,t} > d_{i,t}^{\gamma} \right\} d_{i,t}^{\gamma} \le \rho_{i}T + \sum_{t=1}^{s-1} g_{i,t} \left( b_{i,t}, d_{i,t} \right), \text{ for all } s \in [T].$$
(16)

Problem (16) bears significant complexity in comparison to its counterpart Problem (5) with no karma gains. First, the possibility of gaining karma requires a budget constraint for each time step  $s \in [T]$  instead of only one at the end of the horizon. Second, the outcomes  $x_{i,t}$  cannot be used directly as decision variables since the gains  $g_{i,t}$  depend non-trivially on the bids  $b_{i,t}$ , as given by

$$g_{i,t}(b_{i,t}, d_{i,t}) = \frac{\gamma}{N} p_t^{\gamma+1}(b_{i,t}, d_{i,t}) = \frac{\gamma}{N} \begin{cases} d_{i,t}^{\gamma}, & d_{i,t}^{\gamma} < b_{i,t}, \\ b_{i,t}, & d_{i,t}^{\gamma+1} < b_{i,t} \le d_{i,t}^{\gamma}, \\ d_{i,t}^{\gamma+1}, & b_{i,t} \le d_{i,t}^{\gamma+1}. \end{cases}$$
(17)

Finally, notice that we adapted the definition of competing bids to include both  $d_{i,t}^{\gamma}$  and  $d_{i,t}^{\gamma+1}$ . This is also because the gains  $g_{i,t}$  depend on both of these quantities in Equation (17). If agent *i* is among the auction winners, the price  $p_t^{\gamma+1}$  and thereby the gain  $g_{i,t}$  is determined by  $d_{i,t}^{\gamma}$ ; if the agent is among the auction losers and is not the *price setter*, the gain is determined by  $d_{i,t}^{\gamma+1}$ ; and if the agent is among the auction losers but sets the price the gain is determined by its own bid  $b_{i,t}$ .

To address the complexity of Problem (16), we perform a relaxation that forms a lower bound on the optimal cost with the benefit of hindsight, given by

$$C_i^H(\boldsymbol{v}_i, \boldsymbol{d}_i) \ge \underline{C}_i^H(\boldsymbol{v}_i, \boldsymbol{d}_i^{\boldsymbol{\gamma}}) = \min_{\boldsymbol{x}_i \in \{0,1\}^T} \sum_{t=1}^T v_{i,t}(\tau - x_{i,t}\Delta), \quad \text{s.t.} \sum_{t=1}^T x_{i,t} d_{i,t}^{\boldsymbol{\gamma}} - \frac{\boldsymbol{\gamma}}{N} d_{i,t}^{\boldsymbol{\gamma}} \le \rho_i T.$$
(18)

This lower bound is obtained by a) allowing temporary negative balances of karma as long as it is non-negative at the end of the horizon *T*; and b) eliminating the dependency of  $g_{i,t}$  on  $b_{i,t}$  and  $d_{i,t}^{\gamma+1}$  by assuming that when the agent is among the auction losers, it always manages to be the price setter and impose the maximum gain  $\frac{\gamma}{N} d_{i,t}^{\gamma}$ .

The relaxed Problem (18) has similar structure as Problem (5), and we similarly proceed with the associated Lagrangian dual problem given by

$$\underline{C}_{i}^{H}(\boldsymbol{v}_{i},\boldsymbol{d}_{i}^{\gamma}) \geq \delta_{i}^{H}(\boldsymbol{v}_{i},\boldsymbol{d}_{i}^{\gamma}) = \sup_{\mu_{i}\geq0} \min_{\boldsymbol{x}_{i}\in\{0,1\}^{T}} \sum_{t=1}^{T} \left( x_{i,t}(\mu_{i}d_{i,t}^{\gamma} - \Delta \boldsymbol{v}_{i,t}) + \boldsymbol{v}_{i,t}\tau - \mu_{i}\left(\rho_{i} + \frac{\gamma}{N}d_{i,t}^{\gamma}\right) \right)$$
(19a)

$$= \sup_{\mu_i \ge 0} \sum_{t=1}^{T} v_{i,t} \tau - \mu_i \left( \rho_i + \frac{\gamma}{N} d_{i,t}^{\gamma} \right) - (\Delta v_{i,t} - \mu_i d_{i,t}^{\gamma})^+ := \sup_{\mu_i \ge 0} \sum_{t=1}^{T} \delta_{i,t} (v_{i,t}, d_{i,t}^{\gamma}, \mu_i).$$
(19b)

This dual problem is essentially identical to its counterpart with no karma gains Problem (6) with the exception that the target expenditure rate  $\rho_i$  is replaced by the time-varying term  $\rho_i + \frac{Y}{N} d_{i,t}^{Y}$ . This is intuitive as the agent now aims to deplete both its initial budget as well as the gains it receives. Notice that for a fixed  $\mu_i \ge 0$ , the inner minimum in (19a) is also obtained by bidding  $b_{i,t} = \Delta v_{i,t}/\mu_i$  in this setting.

Adaptive Karma Pacing with Gains. Analogously to Section 3.1, we perform a stochastic gradient ascent scheme on the relaxed dual problem (19). The subgradient in this problem is given by  $\frac{\partial \delta_{i,t}}{\partial \mu_{i,t}}(v_{i,t}, d_{i,t}^{\gamma}, \mu_{i,t}) = z_{i,t} - \rho_i - \frac{\gamma}{N} d_{i,t}^{\gamma}$ . However, adopting the this subgradient in an online algorithm

raises two issues compared to Section 3.1. First, notice that the term  $\frac{\gamma}{N} d_{i,t}^{\gamma}$  is only observed by auction winners, hence we use the observed gain  $g_{i,t}$  as a proxy instead. Second, if all agents are to adopt this subgradient, it is impossible for them to simultaneously track  $\rho_i$  and fully deplete their karma by the end of the horizon, as the total amount of karma in the system is preserved. For this reason we will omit term  $\rho_i$ , such that each agent attempts to match its expenditures to its gains. This yields the *adaptive karma pacing strategy with gains*, which is denoted by *G* and summarized in Algorithm 2.

ALGORITHM 2: Adaptive Karma Pacing with Gains G

#### Input:

- Time horizon *T*, initial budget  $k_{i,1} > 0$ , multiplier bounds  $\overline{\mu}_i > \mu_i > 0$ , gradient step size  $\epsilon_i > 0$ .

#### Initialize:

- Initial multiplier  $\mu_{i,1} \in [\mu_i, \overline{\mu}_i]$ .

#### **for** t = 1, ..., T **do**

(1) Observe the realized valuation  $v_{i,t}$  and place bid

$$b_{i,t} = \min\left\{\frac{\Delta v_{i,t}}{P_{[\underline{\mu}_i,\overline{\mu}_i]}(\mu_{i,t})}, k_{i,t}\right\}; \qquad (G-b)$$

(2) Observe the expenditure  $z_{i,t}$  and the gain  $g_{i,t}$ . Update the multiplier

$$\mu_{i,t+1} = \mu_{i,t} + \epsilon_i (z_{i,t} - g_{i,t}), \qquad (G-\mu)$$

as well as the karma budget  $k_{i,t+1} = k_{i,t} - z_{i,t} + g_{i,t}$ .

end

In contrast to strategy *A* of the setting with no karma gains, strategy *G* uses the time-varying gains  $g_{i,t}$  instead of the target expenditure rate  $\rho_i$  in the multiplier update (*G*- $\mu$ ). Moreover, the projection of multiplier  $\mu_i$  on  $[\underline{\mu_i}, \overline{\mu_i}]$  now occurs in the bid (*G*-*b*) instead of the multiplier update (*G*- $\mu$ ). The importance of this technical difference will be discussed in Section 4.3.

#### 4.2 Asymptotic Optimality under Stationary Competition

This section mirrors Section 3.2 in establishing that strategy *G* is asymptotically optimal in a stationary competition setting, where a single agent *i* bids against competing bids  $d_i = \left(d_{i,t}^{\gamma}, d_{i,t}^{\gamma+1}\right)_{t \in [T]}$  drawn independently across time from a fixed distribution  $\mathcal{D}_i$ .

This section, as well as subsequent Sections 4.3 and 4.4, are organized similarly to Sections 3.2–3.4 in the setting with no karma gains. After stating the required definitions and assumptions, we present the main result of the section (Theorem 4.1 in this case) and focus our discussion on the differences to its counterpart result with no karma gains.

**Preliminaries.** When agent *i* bids according to a fixed  $\mu_i > 0$ , the *expected dual objective*, the *expected expenditure*, the *expected gain*, and the *expected karma loss* are respectively given by

$$\Psi_{i}^{\rho_{i}}(\mu_{i}) = \mathop{\mathbb{E}}_{v_{i},d_{i}} \left[ v_{i}\tau - \mu_{i}(\rho_{i} + g_{i}) - (\Delta v_{i} - \mu_{i}d_{i}^{Y})^{+} \right],$$
(20)

$$Z_i(\mu_i) = \mathop{\mathbb{E}}_{v_i, d_i} \left[ d_i^{\gamma} \mathbb{1} \{ \Delta v_i > \mu_i d_i^{\gamma} \} \right], \qquad G_i(\mu_i) = \mathop{\mathbb{E}}_{v_i, d_i} \left[ g_i \right], \qquad L_i(\mu_i) = Z_i(\mu_i) - G_i(\mu_i), \tag{21}$$

where the expectation is with respect to the stationary distributions  $\mathcal{V}_i$  and  $\mathcal{D}_i$ . We make two observations on the definition of the expected dual objective. First, Equation (20) is parameterized in the target expenditure rate  $\rho_i$ , which lets us write  $\Psi_i^0(\mu_i)$  as the objective that strategy *G* aims to maximize. Second, in line with the multiplier update  $(G - \mu)$ , Equation (20) is defined in terms of the actual gain  $g_i$  rather than the maximum possible gain  $\frac{\gamma}{N} d_i^{\gamma}$  used in the relaxed problems (18)–(19).

This discrepancy implies that there could be a gap between the maximum  $\Psi_i^{\rho_i}$  and the expected optimal dual objective of Problem (19), stemming from the fact that if agent *i* is not among the auction winners, it can increase its gain by bidding  $d_i^{\gamma+1} < b_i \leq d_i^{\gamma}$ . For this reason, it is convenient to define the *residual gain* as  $\hat{\varepsilon}_i = \frac{\gamma}{N} \sum_{d_i} \left[ d_i^{\gamma} - d_i^{\gamma+1} \right]$ , that is, the expected maximum additional gain that agent *i* can get by becoming the price setter.

Moreover, we adapt the definition of the *stationary multiplier* and denote it by  $\mu_i^{\star}$  if it satisfies  $L_i(\mu_i^{\star}) = 0$ , i.e., it causes the expected expenditure to equal the expected gain. Finally, we define the notion of *hitting time* as

$$\mathfrak{T}_{i} = \min\left\{\mathfrak{T}_{i}^{k}, \mathfrak{T}_{i}^{\underline{\mu}}, \mathfrak{T}_{i}^{\overline{\mu}}\right\}, \quad \text{where } \mathfrak{T}_{i}^{k} = \arg\max_{t\in[T]}\left\{\forall s\in[t], k_{i,s}\geq\Delta/\underline{\mu_{i}}\right\}, \\
\mathfrak{T}_{i}^{\underline{\mu}} = \arg\max_{t\in[T]}\left\{\forall s\in[t], \mu_{i,s}\geq\underline{\mu_{i}}\right\}, \quad \mathfrak{T}_{i}^{\overline{\mu}} = \arg\max_{t\in[T]}\left\{\forall s\in[t], \mu_{i,s}\leq\overline{\mu_{i}}\right\}.$$
(22)

This is the latest time step which guarantees that  $b_{i,t} = \frac{\Delta v_{i,t}}{\mu_{i,t}}$  in (*G*-*b*) for any valuation  $v_{i,t} \in [0, 1]$ . By definition, the hitting time is a stricter notion that the *budget depletion time*  $\mathcal{T}_i^k$  used in the proofs of Section 3. This modification is needed since the projection occurs in the bid (*G*-*b*) instead of the multiplier update (*G*- $\mu$ ) in strategy *G*.

Assumptions. To prove strategy G's asymptotic regret bound in the stationary competition setting, we adapt the technical assumptions of Section 3.2 to the setting with karma gains and gather them in Section B.1. Moreover, we formulate two new assumptions in this setting.

Assumption 8 ( $k_{i,1}(T)$  under stationary competition). The initial budget  $k_{i,1}$  is a function of T satisfying  $k_{i,1}(T)/T \xrightarrow{T \to \infty} 0$ .

Assumption 9 (Control of Hitting time). The following conditions hold:

- 9.1 The valuation distribution  $\mathcal{V}_i$  has support in  $[\underline{v}_i, 1]$ , where  $0 < \underline{v}_i < 1$ ;
- 9.2 The distribution of competing bids  $\mathcal{D}_i$  has support in  $\left[\underline{d_i}, \overline{d_i}\right]^2$ , where  $0 < \underline{d_i} < \overline{d_i}$ ;
- 9.3 The algorithm parameters satisfy  $\underline{\mu_i} < \frac{\Delta v_i}{\overline{d_i}^2}$  and  $\overline{\mu_i} > \frac{\Lambda}{\underline{d_i}}$  and  $\epsilon_i < \min\left\{\frac{\Delta v_i}{\overline{d_i}^2} \underline{\mu_i}, \frac{1}{\overline{d_i}}\left(\overline{\mu_i} \frac{\Lambda}{\underline{d_i}}\right)\right\}$ ; 9.4 The initial budget satisfies  $k_{i,1} > \frac{\overline{\mu_i} - \mu_{i,1}}{\epsilon_i} + \frac{\Lambda}{\underline{\mu_i}}$ .

**Main Result.** With these assumptions, we are ready to state the main result of this section regarding the asymptotic optimality of strategy *G* under stationary competition.

THEOREM 4.1 (ASYMPTOTIC OPTIMALITY UNDER STATIONARY COMPETITION). Fix an agent  $i \in N$ , and let Assumptions 1, 13 and 14 hold. Then, there exists a constant  $C \in \mathbb{R}_+$  such that the agent's average expected regret for following strategy G in the stationary competition setting satisfies

$$\frac{1}{T} \mathop{\mathbb{E}}_{\boldsymbol{v}_i, \boldsymbol{d}_i} \left[ C_i^G(\boldsymbol{v}_i, \boldsymbol{d}_i) - C_i^H(\boldsymbol{v}_i, \boldsymbol{d}_i) \right] \le C \left( \epsilon + \frac{1 + k_1 \epsilon}{\epsilon T} + \frac{\mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}} \left[ T - \mathcal{T} \right]}{T} + \hat{\varepsilon} \right)$$
(23)

Let in addition Assumptions 4, 8 and 9 hold. Then strategy G asymptotically converges to an  $O(\hat{\epsilon})$ -neighborhood of the expected optimal cost with the benfit of hindsight, i.e.,

$$\lim_{T \to \infty} \frac{1}{T} \mathop{\mathbb{E}}_{\boldsymbol{v}_i, \boldsymbol{d}_i} \left[ C_i^G(\boldsymbol{v}_i, \boldsymbol{d}_i) - C_i^H(\boldsymbol{v}_i, \boldsymbol{d}_i) \right] = O(\hat{\varepsilon}).$$
(24)

The detailed proof of Theorem 4.1 is included in Appendix E.2. It follows similar steps as its counterpart with no gains (Theorem 3.1); there are however fundamental differences that we discuss next.

First, Theorem 4.1 differs from Theorem 3.1 in that it does not establish an asymptotic average regret of zero but rather in the order of the residual gain  $\hat{\varepsilon} = \frac{\gamma}{N} \mathbb{E}_{d_i} \left[ d_i^{\gamma} - d_i^{\gamma+1} \right]$ . The intuition for the  $O(\hat{\varepsilon})$  term is that, without the benefit of hindsight, agent *i* will always regret not setting the price to the a priori unknown maximum  $d_i^{\gamma}$ , instead of  $d_i^{\gamma+1}$ , when losing the auction. However, in cases where  $d_i^{\gamma} - d_i^{\gamma+1}$  correspond to the distance between two adjacent independent samples from a common distribution  $\mathcal{D}_{-i}$ , the residual gain  $\hat{\varepsilon}$  diminishes as the number of samples, or equivalently N - 1, grows. For a large number of agents *N*, the residual gain  $\hat{\varepsilon}$  is hence expected to be modest. Notice that term  $O(\hat{\varepsilon})$  does not feature in Theorem 3.1 because agent *i* can only affect its karma budget by bidding more or less than  $d_i^{\gamma}$  in the setting with no gains.

The second difference to Theorem 3.1 is that the asymptotic guarantee in Equation (24) requires the initial budget  $k_{i,1}$  to vary as a function of T in addition to the gradient step size  $\epsilon_i$ . By requiring that the initial budget grows sublinearly with respect to the time horizon, Assumption 8 effectively ensures that the target expenditure rate  $\rho_i$  tends to zero and that strategy G maximizes the correct expected dual objective  $\Psi^0(\mu_i)$ . However, the initial budget  $k_{i,1}$  cannot be kept constant. We show in the proof of Theorem 4.1 that the multiplier under strategy G is bounded at all time steps  $t \in [T]$ by  $\mu_{i,t} \leq \mu_{i,1} + \epsilon_i k_{i,1}$ . Since  $\epsilon_i$  diminishes to zero asymptotically under Assumption 4,  $\mu_{i,t}$  would not be able to converge to any value greater than the initial value  $\mu_{i,1}$  with a constant  $k_{i,1}$ . We refer to this phenomenon as the *vanishing box problem*.

Finally, it is necessary for the proof of Theorem 4.1 to establish the sublinear growth rate of  $\mathbb{E}_{v_i,d_i} [T - \mathcal{T}_i]$  with respect to T, which is harder to obtain since the hitting time  $\mathcal{T}_i$  defined in Equation (22) is a stricter notion than the budget depletion time used in the proof of Theorem 3.1. This difficulty is addressed by Assumption 9. By introducing a strictly positive minimum valuation  $v_i$ , Assumption 9.1 ensures that agent *i* always wins when the multiplier is close to  $\mu_i$ , whereas the strictly positive minimum competing bid  $d_i$  introduced in Assumption 9.2 ensures that the agent always loses when the multiplier is close to  $\mu_i$ . In practice, these assumptions imply that agents are always willing to access the priority road and participate actively in the auction.

### 4.3 Convergence under Simultaneous Learning

This section mirrors Section 3.3 in establishing that the learning dynamics converge in the simultaneous learning setting, when all agents simultaneously adopt strategy G, which we denote by the joint strategy profile G. We adapt previously introduced definitions and assumptions as required to establish the main result of the section (Theorem 4.2).

**Preliminaries.** For a fixed multiplier profile  $\mu \in \mathbb{R}^N_+$ , we extend the definitions of the *expected dual objective, expected expenditure, expected gain*, and *expected karma loss* to the multi-agent setting, respectively, as

$$\Psi_i^{\rho_i}(\boldsymbol{\mu}) = \mathbb{E}\left[v_i \tau - \mu_i (\rho_i + g_i) - (\Delta v_i - \mu_i d_i^{\gamma})^+\right],\tag{25}$$

$$Z_{i}(\boldsymbol{\mu}) = \mathop{\mathbb{E}}_{\boldsymbol{v}} \left[ d_{i}^{\gamma} \, \mathbb{1} \{ \Delta v_{i} > \mu_{i} d_{i}^{\gamma} \} \right], \qquad G_{i}(\boldsymbol{\mu}) = \mathop{\mathbb{E}}_{\boldsymbol{v}} \left[ \frac{\gamma}{N} \, p^{\gamma+1} \right], \qquad L_{i}(\boldsymbol{\mu}) = Z_{i}(\boldsymbol{\mu}) - G_{i}(\boldsymbol{\mu}), \tag{26}$$

where as before  $d_i^{\gamma} = \gamma^{\text{th}} - \max_{j:j \neq i} \{\Delta v_j / \mu_j\}$  and  $p^{\gamma+1} = \gamma + 1^{\text{th}} - \max_j \{\Delta v_j / \mu_j\}$  are functions of  $\boldsymbol{v}$ and  $\boldsymbol{\mu}$ . We also extend the definition of the stationary multiplier  $\mu_i^{\star}$  to a *stationary multiplier profile*  $\boldsymbol{\mu}^{\star} \in \mathbb{R}^N_+$  satisfying  $L_i(\boldsymbol{\mu}^{\star}) = 0$  for all  $i \in \mathcal{N}$ .

Notice that stationary multipliers  $\mu^{\star}$  are numerous: if multiplier  $\mu^{\star}$  is stationary, so is  $\eta \mu^{\star}$  for all  $\eta > 0$ , as the expenditures  $Z_i$  and gains  $G_i$  are equally scaled by  $1/\eta$ . This property of  $\mu^{\star}$  is novel to the setting with karma gains, as the scaling of bids was uniquely determined by the target expenditure rates  $\rho_i$  in the absence of gains. To fix the unique  $\mu^{\star}$  that strategy *G* converges to,

we perform two modifications with respect to strategy *A*. First, we move the projection from the multiplier update  $(A-\mu)$  to the bid (G-b) in strategy *G*. Second, we consider that all agents follow the same gradient step sizes, i.e.,  $\epsilon_i = \epsilon > 0$  for all  $i \in N$ . These modifications imply the following properties

$$\sum_{i\in\mathcal{N}}\mu_{i,t+1} = \sum_{i\in\mathcal{N}}\mu_{i,t} + \epsilon \sum_{i\in\mathcal{N}} \left(z_{i,t} - g_{i,t}\right) \stackrel{(a)}{=} \sum_{i\in\mathcal{N}}\mu_{i,t},\tag{27}$$

$$\boldsymbol{\mu}_{t} \in \boldsymbol{U}_{\boldsymbol{\mu}_{1}} = \left\{ \boldsymbol{\mu}_{t} \in \mathbb{R}^{N} \middle| \sum_{i \in \mathcal{N}} \left( \boldsymbol{\mu}_{i,t} - \boldsymbol{\mu}_{i,1} \right) = 0 \right\},$$
(28)

where equality (a) holds under any scheme that redistributes all the expenses, and Property (28) is implied by Equation (27). Property (28) uniquely fixes the scale of  $\mu^*$  that is feasible under strategy *G* with respect to the initial multiplier profile  $\mu_1$ , and implies that the *average multiplier*  $\mu_m = \sum_{i \in N} \mu_{i,1}/N$  is preserved over time. Notice that Equation (27) would not hold if the projection was in the multiplier update (*G*- $\mu$ ): intuitively, a projection there would cause agent *i* to 'forget' part of the history of expenses and gains, and affect the convergence of the whole population as a consequence by shifting the hyperplane of feasible profiles in Property (28).

Assumptions. To prove convergence of the learning dynamics under strategy profile G, we adapt the technical assumptions of Section 4.2 to the multi-agent setting. Those adjustments are summarized in Section B.2. In addition, we make the following new assumption.

Assumption 10 (Control of hitting time). The following conditions hold for all agents  $i \in N$ :

10.1 The multiplier bounds satisfy  $0 < \underline{\mu_i} < \frac{v}{2}\mu_m$  and  $\overline{\mu_i} \ge \mu_m \left(1 + \frac{2}{\underline{v}}\left(1 - \frac{y}{N}\right)^{-1} - \frac{v}{\underline{v}}\right);$ 10.2 The shared gradient step size satisfies  $0 < \epsilon < \mu_m \frac{\mu}{\Delta} \min\left\{\frac{v}{\underline{v}}\left(1 + (\underline{v}+1)\frac{y}{N}\right)^{-1}, \frac{1}{\underline{v}}\left(1 - \frac{y^2}{N^2}\right)^{-1}\right\};$ 10.3 The underlying learning dynamics are such that  $\frac{1}{T}\mathbb{E}_{\boldsymbol{v}}\left[\Upsilon_i^{\underline{\mu}}\right] \xrightarrow[T \to \infty, \epsilon(T) \to 0]{}$ 

**Main Result.** With this assumption, we are ready to state the main result of this section regarding the asymptotic convergence of strategy profile *G*, both with respect to the multiplier profile iterates  $\mu_t$  and the strategic competition costs  $C_i^G$  of all agents  $i \in N$ .

THEOREM 4.2 (CONVERGENCE UNDER SIMULTANEOUS LEARNING). Let Assumptions 1, 16 and 17 hold. Then there exists a constant  $C_1 \in \mathbb{R}_+$  such that the average expected distance to the stationary multiplier profile  $\mu^* \in U_{\mu_1}$  satisfies

$$\frac{1}{T}\sum_{t=1}^{T} \mathbb{E}_{\boldsymbol{v}} \left[ \|\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}\|_{2}^{2} \right] \leq C_{1} N \left( \boldsymbol{\epsilon} + \frac{1}{\boldsymbol{\epsilon}T} + \frac{\mathbb{E}_{\boldsymbol{v}} \left[ T - \underline{\mathcal{T}} \right]}{T} \right)$$
(29)

Moreover, there exists a constant  $C_2 \in \mathbb{R}_+$  such that the strategic competition cost of strategy profile G satisfies, for all agents  $i \in N$ ,

$$\frac{1}{T}C_i^G - \Psi_i^0(\boldsymbol{\mu^{\star}}) \le C_2\left(N\left(\epsilon^{1/2} + \frac{1}{\epsilon T}\right) + \frac{\mathbb{E}_{\boldsymbol{\nu}}\left[T - \underline{\mathcal{T}}\right]}{T}\right).$$
(30)

Let in addition Assumptions 4, 9.1, 9.4 and 10 hold. Then the multiplier profile  $\mu_t$  converges in expectation to the stable multiplier profile  $\mu^* \in U_{\mu_1}$ , and the average strategic competition cost  $C_i^G$  converges to the expected dual objective  $\Psi_i^0(\mu^*)$  for all agents  $i \in N$ , i.e.,

$$\lim_{T \to \infty} \frac{1}{T} \sum_{t=1}^{T} \mathbb{E}_{\boldsymbol{v}} \left[ \|\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}\|_{2}^{2} \right] = 0, \qquad \lim_{T \to \infty} \frac{1}{T} C_{i}^{G} - \Psi_{i}^{0}(\boldsymbol{\mu}^{\star 0}) = 0.$$
(31)

The detailed proof of Theorem 4.2 is included in Appendix E.3, and follows similar steps as its counterpart with no karma gain Theorem 3.2. As before, the main difficulty in comparison to Theorem 3.2 lies in ensuring that the expectation  $\mathbb{E}_{\sigma} \left[T - \underline{T}\right]$  in bounds (29)–(30) grows sublinearly with respect to *T*, since the hitting time is a stricter notion than the budget depletion time used in the proof of Theorem 3.2. This challenge is addressed analogously as in Section 4.2. Namely, Assumptions 9.1, 10.1 and 10.2 deterministically guarantee that  $\mathcal{T}_i^{\overline{\mu}} = T$ : any agent close to  $\overline{\mu_i}$  will lose the auction and transition away from  $\overline{\mu_i}$ . A similar deterministic guarantee cannot be derived at the lower bound  $\overline{\mu_i}$ , however, due to the preservation of the mean multiplier  $\mu_m$ . For this reason, Assumption 10.3 imposes a probabilistic condition on the lower bound  $\underline{\mu_i}$ . This assumption is discussed further in Section 5.

#### 4.4 Approximate Nash equilibrium in Parallel Auctions

This section mirrors Section 3.4 in achieving the main goal of our analysis of strategy profile *G*. Namely, we establish that strategy profile *G* constitutes an  $\varepsilon$ -Nash equilibrium in the parallel auction setting, in which each agent is matched randomly to one of *M* parallel auctions in each time step. We first adapt previously introduced definitions and assumptions that are required to establish the main result of the section (Theorem 4.3).

**Preliminaries: Parallel Auctions.** We recall the parallel auction setting introduced in Section 3.4, for which it remains to specify how payments are redistributed in the new setting with karma gains. We consider that the aggregate payment of all *M* auctions gets redistributed uniformly, leading to karma gains  $g_{i,t} = \frac{\gamma}{N} \sum_{m \in [M]} p_{m,t}^{\gamma+1}$ , where  $p_{m,t}^{\gamma+1}$  is the price of auction *m* defined as  $p_{i,t}^{\gamma+1} = \gamma + 1^{\text{th}} - \max_{i \in N} \{\mathbbm{1}\{m_{i,t} = m\}b_{i,t}\}$ . This aggregate redistribution scheme is advantageous over redistributing the payment of each auction among its agents, since the aggregation restricts the influence of a single agent over the gains, and thereby the multipliers, of others.

**Assumptions.** To prove the  $\varepsilon$ -Nash equilibrium result, the following assumptions are needed.

ASSUMPTION 11 ( $\epsilon(T, N)$  AND  $k_1(T, N)$  UNDER PARALLEL AUCTIONS). The shared gradient step size  $\epsilon$  and initial budget  $k_1$  are functions of the time horizon T and population size N satisfying  $N\epsilon \xrightarrow[T,N\to\infty]{} 0, \epsilon^{3/2}T \xrightarrow[T,N\to\infty]{} \infty$  and  $\overline{k_1}/T \xrightarrow[T,N\to\infty]{} 0$ .

Assumption 12 (Control of Hitting time). The underlying learning dynamics are such that  $\frac{1}{T}\mathbb{E}_{\boldsymbol{v}}\left[\mathcal{T}_{i}^{\underline{\mu}}+\mathcal{T}_{i}^{\overline{\mu}}\right] \xrightarrow[T \to \infty, \epsilon(T) \to 0]{} 1$ 

Assumption 11 strengthens previous Assumption 4 by a) requiring that the gradient step size and initial budget vary with N in addition to T and b) requiring  $\epsilon = \omega(T^{-2/3})$  instead of  $\epsilon = \omega(T^{-1})$ . Minimum valuations are of little help in a parallel auction setting where agents with the highest and smallest multipliers may never interact, Assumption 12 therefore extends Assumption 10.3 to  $T_i^{\overline{\mu}}$ .

**Main Result.** With these assumptions, we are ready to state the main result of this section, namely, that strategy profile *G* constitutes an  $\varepsilon$ -Nash equilibrium in the parallel auction setting.

THEOREM 4.3 ( $\varepsilon$ -NASH EQUILIBRIUM). Fix an agent  $i \in N$ . Let Assumptions 1, 16 and 17 hold. Then there exists a constant  $C \in \mathbb{R}_+$  such that agent i can decrease its average strategic competition cost by deviating from strategy G to any strategy  $\beta_i \in \mathcal{B}$  by at most

$$\frac{1}{T} \left( C_i^G - C_i^{\beta_i, G_{-i}} \right) \leq C \left( \left( \| \boldsymbol{a}_i \|_2 + \frac{M\gamma}{N} \right) \left( \sqrt{N\epsilon} \left( 1 + \frac{1}{\epsilon^{3/2}T} \right) + \| \boldsymbol{a}_i \|_2 + \frac{\gamma}{\sqrt{N}} \right) + \left( \frac{\gamma}{N} + \frac{\overline{k_1}}{T} \right) + \left( \frac{\overline{k_1}}{T} + \frac{M\gamma}{N} \right) \frac{\mathbb{E}_{\boldsymbol{v}, \boldsymbol{m}} \left[ T - \underline{\mathcal{T}} \right]}{T} \right)$$
(32)

Let in addition Assumptions 7.1, 7.3, 9.4, 11 and 12 hold. Then strategy profile G constitutes an  $\varepsilon$ -Nash equilibrium, i.e., it holds for all  $i \in N$ ,

$$\lim_{T,N,M\to\infty} \frac{1}{T} \left( C_i^G - \inf_{\beta\in\mathcal{B}} C_i^{\beta,G_{-i}} \right) = 0.$$
(33)

The detailed proof of Theorem 4.3 is included in Appendix E.4. Similar to Theorem 3.3, it consists of lower-bounding the average expected cost under strategy  $\beta_i$  in terms of the expected dual objective  $\Psi_i^0(\mu^*)$ , and showing that asymptotically agent *i* cannot affect the multiplier profile of the other agents which converges to  $\mu^*$ . This means that the competition becomes stationary, for which strategy *G* is indeed optimal. However, the resulting bound (32) is substantially different from its counterpart with no karma gains (15), and requires adapting the technical assumptions in order to establish the asymptotic guarantee in Equation (33).

This achieves our main goal also for the second class of karma mechanisms in which agents gain karma throughout the episode. We have devised the simple adaptive karma pacing strategy with gains G and provided conditions in which it constitutes an  $\varepsilon$ -Nash equilibrium.

#### 5 DISCUSSION

In this paper, we devised a learning strategy, called *adaptive karma pacing*, that learns to bid optimally in two classes of karma mechanisms: those in which karma is issued at the beginning of a finite episode only; and those in which karma is additionally gained throughout the episode by redistributing the collected payments. This simple strategy constitutes an  $\varepsilon$ -Nash equilibrium in large populations, and can hence be effectively employed to provide decision support in artificial currency mechanisms, which is an important step towards their practical implementation.

*Welfare implications.* Adaptive karma pacing results in bids that are linear in the agents' private valuations. In a symmetric setting with a common valuation distribution, the simultaneous adoption of adaptive karma pacing converges asymptotically to the same multiplier for all agents, and bids become *truth-revealing*. Therefore, resources will be allocated efficiently to the agents with the highest private valuation, and without the need to use money. In a non-symmetric setting, adaptive karma pacing will be similarly efficient among identical agents, and further interpersonal comparability assumptions [31] are needed to assess its welfare properties across agents with different valuation distributions.

Comparison to adaptive pacing in monetary auctions. In the class of karma mechanisms without redistribution, cf. Section 3, the most prominent difference between adaptive karma pacing and adaptive pacing in budget-constrained monetary auctions, e.g., [6], is that the optimal bidding of an agent *i* takes form  $\frac{v_i}{\mu_i}$  instead of  $\frac{v_i}{\mu_i+1}$ . This difference is important both conceptually–it arises from the fact that karma has no value outside the auctions and therefore there is no natural upper bound on bids; and technically–it leads to unbounded bids and requires a non-trivial extension of the analysis considering a lower bound on the multipliers  $\mu_i$ . In the class of karma mechanisms with redistribution, cf. Section 4, the possibility to gain karma and manipulate that gain, as well as the preservation of karma per capita, leads to several conceptual and technical novelties. These novelties

include the inevitable regret term with respect to the optimal cost with the benefit of hindsight in Theorem 4.1, stemming from the possibility to manipulate the gain upon losing; the difficulty that agents cannot simultaneously deplete their budgets and track their target expenditure rates due to the preservation of karma, causing stationary multiplier profiles to become scale-invariant, and requiring to move the multiplier projection from the multiplier update to the bid in Algorithm 2 to converge to a unique multiplier profile; and, as a consequence, the need to control a stricter notion of hitting time in the proofs, which in addition to the budget depletion time considered in monetary auctions considers the first time a multiplier is projected.

Discussion of assumptions. Our main results require a number of technical assumptions which, we argue, are not highly restrictive. These assumptions can be categorized as follows. The assumptions on *valuation and competing bid distributions* (e.g. Assumption 1, 18, 15) are mild continuity and differentiability-kind assumptions that are common in the literature, including in the well-studied monetary setting [6]. Nonetheless, we performed numerical experiments with discrete and unbounded valuation distributions to test the robustness of our results, cf. Figures 3c and 3d in Appendix A, which show examples in which Theorems 4.1 and Theorem 4.2 hold under discrete uniform and geometric distributions.

The assumptions on *input parameters* of the adaptive karma pacing strategy (e.g., Assumptions 3, 14, and 17) can be satisfied by design and/or tuning. Generally, setting the initial multiplier  $\mu_{i,1}$  close to the center of a sufficiently low  $\mu_i$  and a sufficiently high  $\overline{\mu_i}$ , and using a sufficiently small gradient step size  $\epsilon_i$ , suffices to satisfy these assumptions. These assumptions are provided for technical completeness and to provide insight on the structure of the problem. On the other hand, assumptions requiring to *vary parameters asymptotically* (e.g., Assumptions 6, 8, and 11) are less natural to interpret in practice since typically the time horizon *T* and number of agents *N* are fixed by the setting. For this reason, we provided bounds in all our theorems that give finite time and population guarantees.

Finally, assumptions needed to *control the hitting time* in the class of karma mechanisms with redistribution (Assumption 9, 10, 12) arise from our proof technique seeking to deterministically guarantee that the multipliers  $\mu_{i,t}$  will never reach their bounds  $\mu_i$  and  $\overline{\mu_i}$ . In fact, we performed numerical experiments verifying that the hitting time quickly approaches the end of the time horizon under many parameter combinations that extend beyond those assumed, cf. Figure 3a in Appendix A. However, relaxing these assumptions requires a continuous-space Markov chain-based analysis, which provides a promising avenue for future work.

*Future work.* We conclude with a discussion of future research directions. We hope that our paper inspires future interest in the class of karma mechanisms with redistribution or other forms of karma gains. In this paper, we studied a simple uniform redistribution scheme; however, there are rich possibilities to design karma gains that have not yet been fully explored. Moreover, in this same class of karma mechanisms, it will be important to address the *vanishing box problem*, cf. Section 4.2, which prevents convergence from being achieved with a fixed initial budget. Towards this direction, we performed a numerical experiment using a *variable gradient step size*  $\epsilon_t$ , cf. Figure 3b in Appendix A, which provides preliminary evidence that asymptotic convergence can be achieved even with a fixed initial budget. Therefore, extending the present analysis for variable gradient step sizes provides an exciting avenue for further investigation. Finally, armed with the simple adaptive karma pacing strategy, we hope to see many practical implementations of karma mechanisms that address societal challenges involving scarce resource allocations.

### REFERENCES

- Richard Arnott, André De Palma, and Robin Lindsey. 1990. Economics of a bottleneck. *Journal of urban economics* 27, 1 (1990), 111–130.
- [2] Richard Arnott, André De Palma, and Robin Lindsey. 1994. The welfare effects of congestion tolls with heterogeneous commuters. Journal of Transport Economics and Policy (1994), 139–161.
- [3] Moshe Babaioff, Tomer Ezra, and Uriel Feige. 2022. On best-of-both-worlds fair-share allocations. In International Conference on Web and Internet Economics. Springer, 237–255.
- [4] Moshe Babaioff, Tomer Ezra, and Uriel Feige. 2023. Fair-share allocations for agents with arbitrary entitlements. Mathematics of Operations Research (2023).
- [5] Santiago R Balseiro, Omar Besbes, and Gabriel Y Weintraub. 2015. Repeated auctions with budgets in ad exchanges: Approximations and design. *Management Science* 61, 4 (2015), 864–884.
- [6] Santiago R Balseiro and Yonatan Gur. 2019. Learning in repeated auctions with budgets: Regret minimization and equilibrium. *Management Science* 65, 9 (2019), 3952–3968.
- [7] Santiago R Balseiro, Huseyin Gurkan, and Peng Sun. 2019. Multiagent mechanism design without money. Operations Research 67, 5 (2019), 1417–1436.
- [8] Santiago R Balseiro, Haihao Lu, and Vahab Mirrokni. 2023. The best of many worlds: Dual mirror descent for online allocation problems. Operations Research 71, 1 (2023), 101–119.
- [9] Siddhartha Banerjee, Giannis Fikioris, and Eva Tardos. 2023. Robust Pseudo-Markets for Reusable Public Resources. In Proceedings of the 24th ACM Conference on Economics and Computation. 241.
- [10] Devi K Brands, Erik T Verhoef, Jasper Knockaert, and Paul R Koster. 2020. Tradable permits to manage urban mobility: market design and experimental implementation. *Transportation Research Part A: Policy and Practice* 137 (2020), 34–46.
- [11] Eric Budish and Estelle Cantillon. 2012. The multi-unit assignment problem: Theory and evidence from course allocation at Harvard. *American Economic Review* 102, 5 (2012), 2237–71.
- [12] Matteo Castiglioni, Andrea Celli, and Christian Kroer. 2022. Online learning with knapsacks: the best of both worlds. In *International Conference on Machine Learning*. PMLR, 2767–2783.
- [13] Ezzat Elokda, Saverio Bolognani, Andrea Censi, Florian Dörfler, and Emilio Frazzoli. 2023. A self-contained karma economy for the dynamic allocation of common resources. *Dynamic Games and Applications* (2023), 1–33.
- [14] Ezzat Elokda, Carlo Cenedese, Kenan Zhang, John Lygeros, and Florian Dörfler. 2022. CARMA: Fair and efficient bottleneck congestion management with karma. arXiv preprint arXiv:2208.07113 (2022).
- [15] Andrew W Evans. 1992. Road congestion pricing: when is it a good policy? Journal of transport economics and policy (1992), 213–243.
- [16] Huang Fang, Nick Harvey, Victor Portella, and Michael Friedlander. 2020. Online mirror descent and dual averaging: keeping pace in the dynamic case. In *Proceedings of the 37th International Conference on Machine Learning (Proceedings of Machine Learning Research, Vol. 119)*, Hal Daumé III and Aarti Singh (Eds.). PMLR, 3008–3017. https://proceedings.mlr.press/v119/fang20a.html
- [17] Eric J Friedman, Joseph Y Halpern, and Ian Kash. 2006. Efficiency and Nash equilibria in a scrip system for P2P networks. In ACM conference on Electronic commerce. 140–149.
- [18] Jason Gaitonde, Yingkai Li, Bar Light, Brendan Lucier, and Aleksandrs Slivkins. 2022. Budget pacing in repeated auctions: Regret and efficiency without convergence. arXiv preprint arXiv:2205.08674 (2022).
- [19] Allan Gibbard. 1973. Manipulation of voting schemes: a general result. Econometrica 41, 4 (1973), 587-601.
- [20] Artur Gorokh, Siddhartha Banerjee, and Krishnamurthy Iyer. 2021. From monetary to nonmonetary mechanism design via artificial currencies. *Mathematics of Operations Research* 46, 3 (2021), 835–855.
- [21] Artur Gorokh, Siddhartha Banerjee, and Krishnamurthy Iyer. 2021. The Remarkable Robustness of the Repeated Fisher Market. In Proceedings of the 22nd ACM Conference on Economics and Computation. 562–562.
- [22] Yingni Guo and Johannes Hörner. 2020. Dynamic allocation without money. TSE Working Paper (2020).
- [23] Elad Hazan. 2023. Introduction to Online Convex Optimization. arXiv:1909.05207 [cs.LG]
- [24] Devansh Jalota, Karthik Gopalakrishnan, Navid Azizan, Ramesh Johari, and Marco Pavone. 2023. Online Learning for Traffic Routing under Unknown Preferences. In Proceedings of The 26th International Conference on Artificial Intelligence and Statistics (Proceedings of Machine Learning Research, Vol. 206), Francisco Ruiz, Jennifer Dy, and Jan-Willem van de Meent (Eds.). PMLR, 3210–3229. https://proceedings.mlr.press/v206/jalota23a.html
- [25] Devansh Jalota and Yinyu Ye. 2023. Stochastic Online Fisher Markets: Static Pricing Limits and Adaptive Enhancements. arXiv:2205.00825 [cs.GT]
- [26] Kris Johnson, David Simchi-Levi, and Peng Sun. 2014. Analyzing scrip systems. Operations Research 62, 3 (2014), 524–534.
- [27] Jaehong Kim, Mengling Li, and Menghan Xu. 2021. Organ donation with vouchers. Journal of Economic Theory 191 (2021), 105–159.
- [28] V. Krishna. 2009. Auction Theory. Academic Press, Burlington MA.

- [29] Noam Nisan, Tim Roughgarden, Éva Tardos, and Vijay V Vazirani. 2007. Algorithmic Game Theory. Cambridge University Press, Cambridge.
- [30] Canice Prendergast. 2022. The allocation of food to food banks. Journal of Political Economy 130, 8 (2022), 1993-2017.
- [31] Kevin WS Roberts. 1980. Interpersonal comparability and social choice theory. The Review of Economic Studies 47, 2 (1980), 421–439.
- [32] J Ben Rosen. 1965. Existence and uniqueness of equilibrium points for concave N-person games. Econometrica: Journal of the Econometric Society (1965), 520–534.
- [33] Mark Allen Satterthwaite. 1975. Strategy-proofness and Arrow's conditions: Existence and correspondence theorems for voting procedures and social welfare functions. *Journal of Economic Theory* 10, 2 (1975), 187–217.
- [34] James Schummer and Rakesh V Vohra. 2007. Mechanism design without money. In Algorithmic Game Theory. 243-299.
- [35] Tayfun Sönmez, M Utku Ünver, and M Bumin Yenmez. 2020. Incentivized kidney exchange. American Economic Review 110, 7 (2020), 2198–2224.
- [36] Brian D Taylor and Rebecca Kalauskas. 2010. Addressing equity in political debates over road pricing: Lessons from recent projects. *Transportation research record* 2187, 1 (2010), 44–52.
- [37] William S Vickrey. 1969. Congestion theory and transport investment. The American Economic Review 59, 2 (1969), 251–260.
- [38] Vivek Vishnumurthy, Sangeeth Chandrakumar, and Emin Gun Sirer. 2003. Karma: A secure economic framework for peer-to-peer resource sharing. In Workshop on Economics of Peer-to-peer Systems.
- [39] Midhul Vuppalapati, Giannis Fikioris, Rachit Agarwal, Asaf Cidon, Anurag Khandelwal, and Éva Tardos. 2023. Karma: Resource Allocation for Dynamic Demands. In 17th USENIX Symposium on Operating Systems Design and Implementation (OSDI 23).
- [40] Qian Wang, Zongjun Yang, Xiaotie Deng, and Yuqing Kong. 2023. Learning to bid in repeated first-price auctions with budgets. In *International Conference on Machine Learning*. PMLR, 36494–36513.

## A NUMERICAL EXPERIMENTS

This appendix presents numerical computation results that complement our main theoretical results presented in the body of the paper. The following figures were generated by running one hundred simulations for each of the considered parameter combinations, whose average and estimate of the 95% confidence interval are reported. The figures use a log-log scale with the time horizon T in the x-axis and the quantity of interest in the y-axis. Therefore, a decreasing trend implies that the quantity of interest is converging asymptotically to zero, and the slope of decrease gives the rate of convergence as a power of T.



(a) Theorem 3.1: Stationary competition, no karma gains.



(c) Theorem 3.2: Simultaneous learning, no karma gains.



(b) Theorem 4.1: Stationary competition, with karma gains.



(d) Theorem 4.2: Simultaneous learning, with karma gains.

Fig. 2. Numerical validation of the main theorems. Figures 2a and 2b show the convergence of costs to the minimum with the benefit of hindsight, without and with karma gains. Figures 2c and 2d show the convergence of multipliers under simultaneous learning, without and with karma gains.

Unless specified otherwise, the experiments share the following parameters: valuations are drawn uniformly at random from [0, 1]; multipliers are bounded to the interval [0.1, 1000]; and the travel time saving of the priority road is  $\Delta = 5$ . In the stationary competition setting, the competing bids  $d_{i,t}^{\gamma}$  are also drawn uniformly at random from [0, 1], and we consider that the agent cannot set the price, i.e.,  $d_{i,t}^{\gamma+1} \approx d_{i,t}^{\gamma}$ . In the simultaneous learning setting, we consider N = 50 agents and  $\gamma = 5$  winners per period. The experiment-specific parameters are detailed below. In the following,



(a) Validation of Assumption 10. The hitting time  $\underline{\mathcal{T}}$  approaches *T* asymptotically.



(c) Stationary competition with karma gains and discrete distributions of valuations and competing bids.



(b) For a fixed initial budget, fixed step sizes  $\epsilon$  diverge and variable step sizes  $\epsilon_t$  converge under simultaneous learning with karma gains.



(d) Simultaneous learning with karma gains and discrete valuation distributions.

Fig. 3. Numerical validation of Assumption 1 and 10, and potential solution for the vanishing box problem (Figure 3b). Figure 3a shows that Assumption 10.3 is satisfied in practice. Figures 3c and 3d respectively show that the results of Theorems 4.1 and 4.2 hold also when Assumption 1 is not satisfied.

 $C_{\epsilon}$  and  $C_k$  denote the multiplicative constants such that  $\epsilon(T) = C_{\epsilon}T^x$  and  $k_1(T) = C_kT^y$ , for the indicated powers *x* and *y* in the figures.

- Figure 2a: the initial multiplier is  $\mu_{i,1} = 10$ ; the target expenditure rate is  $\rho = 0.2$ ; and  $C_{\epsilon} = 40$ ;
- *Figure 2b*: the initial multiplier is  $\mu_{i,1} = 5$ ; the karma gain is  $g_{i,t} = d_{i,t}^{\gamma}/10$ ; and  $C_{\epsilon} = 20$ ,  $C_k = 3$ ;
- *Figure 2c*: the target expenditure rate is  $\rho_i = 0.1$ ; the initial multipliers are  $\mu_{i,1} = 5$  and  $\mu_{i,1} = 6$  for the first and second halves of the population, respectively; and  $C_{\epsilon} = 10$ . The stationary multiplier profile  $\mu^{\star}$  is approximated by the mean profile that simulations with largest *T* converged to;
- *Figure 2d:* the initial multipliers are  $\mu_{i,1} = 5$  and  $\mu_{i,1} = 6$  for the first and second halves of the population, respectively; the stationary multiplier is  $\mu_i^* = 5.5$  by symmetry;  $C_{\epsilon} = 1$  and  $C_k = 3$ ;
- *Figure 3a:* parameters are the same as Figure 2d with the exception of  $C_k = 6$ ;
- *Figure 3b:* the fixed initial budget is  $k_{i,1} = 100$ ; the initial multipliers are  $\mu_{i,1} = 5$  and  $\mu_{i,1} = 6$  for the first and second halves of the population, respectively; and  $C_{\epsilon} = 1$ ;

- *Figure 3c:* the fixed initial budget is  $k_{i,1} = 100$ ; the initial multiplier is  $\mu_{i,1} = 55 > \mu_i^*$ ; the karma gain is  $g_{i,t} = d_{i,t}^{\gamma}/10$ ; and  $C_{\epsilon} = 20$ ; Valuations and competing bids are drawn either from the discrete uniform distribution  $\mathcal{U}(\{1, ..., 10\})$  or the geometric distribution  $\mathcal{G}(p)$ with parameter p = 0.3;
- Figure 3d: the initial multipliers are  $\mu_{i,1} = 5$  and  $\mu_{i,1} = 6$  for the first and second halves of the population, respectively; the initial budget is  $k_{i,1} = 100 \times T^{0.61}$ ; and  $C_{\epsilon} = 1$ =. Valuations are drawn either from the discrete uniform distribution  $\mathcal{U}(\{1,\ldots,10\})$  or the geometric distribution  $\mathcal{G}(p)$  with parameter p = 0.3.

## **B** ADDITIONAL ASSUMPTIONS IN SECTION 4

We gather in this section the adaptations of hypotheses from Section 3 to the setting with karma gains in Section 4.

#### **B.1** Stationary Competition Setting

This section contains two technical assumptions for the stationary competition setting with karma gains, and adapted from Section 3.2. Namely, Assumption 13 mirrors Assumption 2 and Assumption 14 is the equivalent of Assumption 3 in the setting with no karma gains.

ASSUMPTION 13 (DIFFERENTIABILITY AND CONCAVITY). The following conditions hold for all  $\mu_i \in \left| \mu_i, \overline{\mu_i} \right|$ :

- 13.1 The expected dual objective  $\Psi_i^0(\mu_i)$  is differentiable with derivative  $\Psi_i^{0'}: \mu_i \mapsto L_i(\mu_i) \mu_i G_i'(\mu_i);$
- 13.2 The expected expenditure  $Z_i(\mu_i)$  and the expected gain  $G_i(\mu_i)$  are twice differentiable with derivatives bounded by  $\overline{Z'} > 0$  and  $\overline{G'} > 0$  in the first order, respectively, and  $\overline{Z''} > 0$  and G'' > 0 in the second order, respectively;

13.3 The expected karma loss  $L_i(\mu_i) = Z_i(\mu_i) - G_i(\mu_i)$  is strongly decreasing with parameter  $\lambda > 0$ .

Assumption 14 (Parameter design under stationary competition). The following conditions hold:

- 14.1 The multiplier bounds are chosen such that the stationary multiplier satisfies  $\mu_i^{\star} \in (\mu_i, \overline{\mu_i})$ ;
- 14.2 The multiplier lower-bound satisfies  $\underline{\mu_i} < \frac{\Delta}{\sup_{i=0}^{\infty} [d_i^{\gamma}]}$ ;

14.3 The initial budget 
$$k_{i,1}$$
 satisfies  $\frac{\Delta}{\mu_i} < k_{i,1} \le T\left(1 - \frac{\gamma}{N}\right) \mathop{\mathbb{E}}_{d_i} \left[d_i^{\gamma}\right]$   
14.4 The gradient step size satisfies  $\epsilon_i \le \min\left\{\frac{1}{1-\gamma}, \frac{1}{1-\gamma}\right\};$ 

Note that similarly to Assumption 2, Assumption 13 is guaranteed to hold under n  
on the distributions 
$$V_i$$
 and  $D_i$  and the residual gain  $\hat{\epsilon}$  (c.f. Section F.1 for the proof). The

nild conditions nese conditions auai ga are next described in Assumption 15.

ASSUMPTION 15. The following conditions hold:

- 15.1 The valuation density  $v_i$  is differentiable with bounded derivative  $|v'_i(v)| \leq \overline{v}'$  for all  $v \in [0, 1]$ ; 15.2 The distribution of competing bids  $D_i$  is absolutely continuous with bounded density  $h_i$ :  $[0, \Delta/\mu_i]^2 \mapsto [\underline{h}, h] \subset \mathbb{R}_{>0};$
- 15.3 The residual gain satisfies  $\hat{\varepsilon} \leq \frac{\nu \underline{h} \Delta^4}{3 \overline{\nu} \overline{u}^4}$ .

Note that Assumption 15 mirrors Assumption 18 in the setting with no gains, with the addition of the bound on the residual gain  $\hat{\epsilon}$ .

#### **B.2 Simultaneous Learning Setting**

This section gathers two technical assumptions for the simultaneous learning setting with karma gains. Assumptions 16 and 17 are respectively adapted from Assumptions 5 and 3 in the setting with no gains.

ASSUMPTION 16 (MONOTONICITY). For all  $\mu, \mu' \in \prod_{i \in \mathcal{N}} \left[ \underline{\mu_i}, \overline{\mu_i} \right]$  satisfying  $\sum_{i \in \mathcal{N}} \mu_i - \mu'_i = 0$ , the expected karma loss  $L(\mu)$  satisfies

$$(\boldsymbol{\mu} - \boldsymbol{\mu}')^{\top} (\boldsymbol{L}(\boldsymbol{\mu}) - \boldsymbol{L}(\boldsymbol{\mu}')) \leq -\lambda \|\boldsymbol{\mu} - \boldsymbol{\mu}'\|_2^2.$$

Assumption 17 (Parameter design under simultaneous learning). The following conditions hold:

17.1 The initial multiplier profile and multiplier bounds are chosen such that the associated stationary multiplier profile  $\mu^{\star} \in U_{\mu_1}$  satisfies  $\mu^{\star} \in \prod_{i \in \mathcal{N}} \left( \underline{\mu_i}, \overline{\mu_i} \right);$ 

17.2 The shared gradient step size of the agents satisfies  $\epsilon \leq \frac{1}{2\lambda}$ .

## C PROOFS OF THE THEOREMS IN SECTION 3

This section contains the proofs of the main Theorems in Section 3. It is organized as follows: first, we prove in Section C.1 a few handful lemmas. The proofs of Theorems 3.1, 3.2 and 3.3 are then included in Sections C.2, C.3 and C.4 respectively. The proofs of the secondary results discussed in Section 3 are deferred to Section D.

### C.1 Helpful Lemmas

Depletion Time. We define the depletion time  $\mathcal{T}_i = \min \left\{ \inf \{t \ge 1 : k_{i,t+1} < \Delta/\underline{\mu_i}\}, T \right\}$  as the last period in which the remaining budget of agent *i* is larger than its maximum possible bid (or *T* when its budget is still greater at the end of the time horizon). We further define  $\underline{\mathcal{T}} = \min_{i \in \mathcal{N}} \mathcal{T}_i$  as the earliest depletion time among the whole population. The following lemma provides a lower bound on the budget depletion time.

LEMMA C.1 (LOWER BOUND ON BUDGET DEPLETION TIME). The budget depletion time  $T_i$  of each agent *i* in *N* satisfies

$$T - \mathfrak{T}_i \leq \frac{\Delta}{\rho_i \underline{\mu_i}} + \frac{\overline{\mu_i}}{\rho_i \epsilon_i}.$$

The earliest depletion time  $\underline{T}$  then satisfies:

$$T - \underline{\mathfrak{T}} \le \frac{\Delta}{\underline{\rho\mu}} + \frac{\overline{\mu}}{\underline{\rho\epsilon}}$$

PROOF. Consider agent  $i \in N$ . From its multiplier update rule  $\mu_{i,t+1} = P_{[\mu_i,\overline{\mu_i}]}(\mu_{i,t} + \epsilon_i(z_{i,t} - \rho_i))$ , we define the projection error  $P_{i,t} = \mu_{i,t} + \epsilon_i(z_{i,t} - \rho_i) - \mu_{t+1}$ . Summing the update rule over all periods up to  $\mathcal{T}_i$  gives

$$\sum_{t=1}^{\mathcal{T}_i} (z_{i,t} - \rho_i) = \sum_{t=1}^{\mathcal{T}_i} \frac{1}{\epsilon_i} (\mu_{i,t+1} - \mu_{i,t}) + \sum_{t=1}^{\mathcal{T}_i} \frac{P_{i,t}}{\epsilon_i}.$$
(34)

The left term can be lower-bounded using the definition of  $T_i$ 

$$\sum_{t=1}^{\mathcal{T}_{i}} (z_{i,t} - \rho_{i}) = k_{i,1} - k_{i,\mathcal{T}_{i}+1} - \rho_{i}\mathcal{T}_{i} \ge \rho_{i}(T - \mathcal{T}_{i}) - \frac{\Delta}{\underline{\mu_{i}}}.$$
(35)

Turning to the right-hand side, the telescopic sum in the first term satisfies:

$$\sum_{t=1}^{\mathcal{T}_i} \frac{1}{\epsilon_i} (\mu_{i,t+1} - \mu_{i,t}) = \frac{\mu_{i,\mathcal{T}_i+1} - \mu_{i,1}}{\epsilon_i} \le \frac{\overline{\mu_i}}{\epsilon_i}.$$
(36)

We show in the following that the second term is non-positive.

$$P_{i,t} \leq P_{i,t}^{+} = \left(\mu_{i,t} + \epsilon_{i}(z_{i,t} - \rho_{i}) - P_{[\underline{\mu_{i}},\overline{\mu_{i}}]}(\mu_{i,t} + \epsilon_{i}(z_{i,t} - \rho_{i}))\right)^{+}$$
$$= \left(\mu_{i,t} + \epsilon_{i}(z_{i,t} - \rho_{i}) - \overline{\mu_{i}}\right) \mathbb{1}\{\mu_{i,t} + \epsilon_{i}(z_{i,t} - \rho_{i}) > \overline{\mu_{i}}\}$$
$$= 0$$
(37)

Indeed, Assumption 3 ensures that  $\underline{\mu_i} > \sqrt{\epsilon_i \Delta}$ , hence the function  $f : \mu \to \mu + \epsilon_i \Delta/\mu$  is strictly increasing on  $[\mu_i, \overline{\mu_i}]$ . This gives:

$$\begin{split} \mu_{i,t} + \epsilon_i \left( z_{i,t} - \rho_i \right) &\stackrel{(a)}{\leq} \mu_{i,t} + \epsilon_i \left( \frac{\Delta}{\mu_{i,t}} - \rho_i \right) = f(\mu_{i,t}) - \epsilon_i \rho_i \\ &\leq f(\overline{\mu_i}) - \epsilon_i \rho_i = \overline{\mu_i} + \epsilon_i \left( \frac{\Delta}{\overline{\mu_i}} - \rho_i \right) \stackrel{(b)}{\leq} \overline{\mu_i} \end{split}$$

Inequality (a) holds by bounding the valuation by one; inequality (b) holds under Assumption 3.

Combining Equations 34, 35, 36 and 37, we finally get that strategy *A* does not run out of budget very early.

$$T - \mathcal{T}_i \leq \frac{\Delta}{\rho_i \mu_i} + \frac{\overline{\mu_i}}{\rho_i \epsilon_i}.$$

Taking the appropriate upper and lower bounds on the whole population gives the desired result.

Expected Mean Squared Error. For any fixed  $t \leq \underline{\mathcal{T}}$ , we consider the expected mean squared error  $s_t = \mathbb{E}_{\boldsymbol{v}} \left[ \|\boldsymbol{\mu}_t - \boldsymbol{\mu}^{\star}\|_2^2 \right] = \sum_{i=1}^N s_{i,t}$ , where  $s_{i,t} = \mathbb{E}_{\boldsymbol{v}} \left[ (\mu_{i,t} - \mu_i^{\star})^2 \right]$  is the individual expected error of agent *i*. The following Lemma bounds the expected mean squared error in different scenarios.

LEMMA C.2. When all agents follow strategy A, the expected mean squared error  $s_t$  verifies:

$$s_t \le N\overline{\mu}^2 \frac{\overline{\epsilon}}{\underline{\epsilon}} (1 - 2\lambda \underline{\epsilon})^{t-1} + \frac{N\Delta^2 \overline{\epsilon}^2}{2\lambda \underline{\mu}^2 \underline{\epsilon}}$$

Alternatively for an agent in the stationary competition settings, its individual expected error satisfies:

$$s_{i,t} \leq \overline{\mu}^2 (1 - 2\lambda \underline{\epsilon})^{t-1} + \frac{\Delta^2 \epsilon}{2\lambda \mu^2}$$

**PROOF.** Let  $\hat{s}_{i,t} = s_{i,t}/\epsilon_i$ , and  $\hat{s}_t = \sum_{i=1}^N \hat{s}_{i,t}$ . For any agent  $i \in N$ , the multiplier update rule gives:

$$\begin{aligned} |\mu_{i,t+1} - \mu_i^{\star}|^2 &= \left| P_{[\underline{\mu_i, \mu_i}]}(\mu_{i,t} + \epsilon_i (z_{i,t}^A - \rho_i)) - \mu_i^{\star} \right|^2 \\ &\stackrel{(a)}{\leq} \left| \mu_{i,t} - \mu_i^{\star} + \epsilon_i (z_{i,t}^A - \rho_i) \right|^2 \\ &= |\mu_{i,t} - \mu_i^{\star}|^2 + 2\epsilon_i (\mu_{i,t} - \mu_i^{\star}) (z_{i,t}^A - \rho_i) + \epsilon_i^2 |z_{i,t}^A - \rho_i|^2 \end{aligned}$$

where (*a*) holds because  $\mu_i^{\star}$  lies in  $[\underline{\mu_i}, \overline{\mu_i}]$  by definition of  $\mu^{\star}$  (respectively Assumption 3 in the stationary competition setting).

Taking the expectation on previous valuations of all agents (resp. previous valuations and competing prices) and dividing by  $\epsilon_i$  gives

$$\hat{s}_{i,t+1} \leq \hat{s}_{i,t} + 2\mathbb{E}_{\boldsymbol{v}} \left[ (\mu_{i,t} - \mu_{i}^{\star}) \left( z_{i,t} - \rho_{i} \right) \right] + \epsilon_{i} \mathbb{E}_{\boldsymbol{v}} \left[ |z_{i,t} - \rho_{i}|^{2} \right],$$

$$\stackrel{(a)}{=} \hat{s}_{i,t} + 2\mathbb{E}_{\boldsymbol{v}} \left[ (\mu_{i,t} - \mu_{i}^{\star}) \left( \mathbb{E}_{\boldsymbol{v}} [z_{i,t} | \boldsymbol{\mu}_{t}] - \rho_{i} \right) \right] + \epsilon_{i} \mathbb{E}_{\boldsymbol{v}} \left[ |z_{i,t} - \rho_{i}|^{2} \right], \qquad (38)$$

$$\stackrel{(b)}{=} \hat{s}_{i,t} + 2\mathbb{E}_{\boldsymbol{v}} \left[ (\mu_{i,t} - \mu_{i}^{\star}) \left( Z_{i}(\boldsymbol{\mu}_{t}) - \rho_{i} \right) \right] + \epsilon_{i} \mathbb{E}_{\boldsymbol{v}} \left[ |z_{i,t} - \rho_{i}|^{2} \right].$$

Equality (*a*) holds by linearity of expectation; equality (*b*) by definition of  $Z_i(\mu_t)$  (resp.  $Z_i(\mu_{i,t})$ ). First, we bound the sum over the whole population of the middle term in Equation 38.

$$\mathbb{E}_{\boldsymbol{v}}\left[\sum_{i=1}^{N}(\mu_{i,t}-\mu_{i}^{\star})(Z_{i}(\boldsymbol{\mu}_{t})-\rho_{i})\right] = \mathbb{E}_{\boldsymbol{v}}\left[\sum_{i=1}^{N}(\mu_{i,t}-\mu_{i}^{\star})\left(Z_{i}(\boldsymbol{\mu}_{t})-Z_{i}(\boldsymbol{\mu}^{\star})+Z_{i}(\boldsymbol{\mu}^{\star})-\rho\right)\right]$$

$$\stackrel{(a)}{\leq} -\mathbb{E}_{\boldsymbol{v}}\left[\lambda\|\boldsymbol{\mu}_{t}-\boldsymbol{\mu}^{\star}\|_{2}^{2}\right]$$

$$= -\lambda s_{t}$$
(39)

Inequality (*a*) uses the definition of  $\mu^{\star}$  (resp. Equation 49), as well as the  $\lambda$ -strong monotonicity of the expected expenditure function *Z* from Assumption 5 (resp. Assumption 2).

On the other hand, as  $z_{i,t}$  and  $\rho_i$  are positive and bounded by  $\Delta/\mu$  under Assumption 3, we similarly bound the right-most term of Equation 38:

$$\sum_{i=1}^{N} \epsilon_{i} \mathbb{E}_{\boldsymbol{\nu}} \left[ |z_{i,t} - \rho_{i}|^{2} \right] \leq \frac{N \Delta^{2} \overline{\epsilon}}{\underline{\mu}^{2}}.$$
(40)

Combining Equations 38, 39 and 40 and noting that  $s_t \ge \epsilon \hat{s}_t$  gives the following recursion

$$\hat{s}_{t+1} \le (1 - 2\lambda \underline{\epsilon})\hat{s}_t + \frac{N\Delta^2 \overline{\epsilon}}{\mu^2}.$$
(41)

Since  $2\lambda \epsilon \leq 1$  because Assumption 3 holds for all agents, [6, Lemma C.4] yields

$$\hat{s}_t \leq \hat{s}_1 (1 - 2\lambda \underline{\epsilon})^{t-1} + \frac{N \Delta^2 \overline{\epsilon}}{2\lambda \mu^2 \underline{\epsilon}}$$

Noting that  $s_t \leq \overline{\epsilon} \hat{s}_t$  and that  $\hat{s}_1 \leq N \overline{\mu}^2 / \underline{\epsilon}$  finally gives the desired bound.

However, if a single agent *i* in  $\mathcal{N}$  places bids with a strategy  $\beta \in \mathcal{B}$  different from A in a parallel auction setting, we consider the individual expected mean squared errors of all other agents. For the purpose of the analysis, we allow this strategy to use *complete information*, i.e., it can access the budgets  $k_s$ , auctions  $m_s$ , valuations  $v_s$ , bids  $b_s$ , and expenditures  $z_s$  of all agents at all previous time steps s < t, and in addition, it has knowledge of the valuation distributions  $\mathcal{V}$ . We denote the set of strategies that use complete information as  $\mathcal{B}^{Cl} \supset \mathcal{B}$  and suppose that  $\beta_i \in \mathcal{B}^{Cl}$ .

To keep notations consistent, we take the convention  $\mu_{i,t} = \mu_i^*$  for all  $t \in \mathbb{N}$ , ensuring that  $s_{i,t} = 0$ .

LEMMA C.3. If all agents but agent i follow strategy A, the expected mean squared error of the population verifies

$$s_t \leq N\overline{\mu}^2 \frac{\overline{\epsilon}}{\underline{\epsilon}} (1 - \lambda \underline{\epsilon})^{t-1} + \frac{N\Delta^2}{\lambda \mu^2} \frac{\overline{\epsilon}^2}{\underline{\epsilon}} + \frac{\Delta^2 \|a_i\|_2^2}{\lambda^2 \mu^2} \frac{\overline{\epsilon}^2}{\underline{\epsilon}^2}.$$

**PROOF.** For an agent  $j \neq i$ , Equation 38 (*a*) still holds and gives

$$\hat{s}_{j,t+1} \stackrel{(a)}{\leq} \hat{s}_{j,t} + 2\mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ (\mu_{j,t} - \mu_{j}^{\star}) \left( \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} [\boldsymbol{z}_{j,t} | \boldsymbol{\mu}_{t}] - \rho_{j} \right) \right] + \epsilon_{j} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ |\boldsymbol{z}_{j,t} - \rho_{j}|^{2} \right]$$

$$\stackrel{(b)}{\leq} \hat{s}_{j,t} + 2\mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ (\mu_{j,t} - \mu_{j}^{\star}) \left( Z_{j}(\boldsymbol{\mu}_{t}) - \rho_{j} \right) \right] + \epsilon_{j} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ |\boldsymbol{z}_{j,t} - \rho_{j}|^{2} \right]$$

$$+ 2\mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ |\mu_{j,t} - \mu_{j}^{\star}| \cdot \left| \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} [\boldsymbol{z}_{j,t} | \boldsymbol{\mu}_{t} \right] - Z_{j}(\boldsymbol{\mu}_{t}) \right| \right]$$

$$(42)$$

Inequality (*a*) directly follows from inequality 38 (*a*); inequality (*b*) follows from adding and subtracting  $Z_j(\mu_t)$  in the second term of (*a*), and noting that  $xy \ge -|x| \cdot |y|$  for  $x, y \in \mathbb{R}$ .

As Equations 39 and 40 hold when summing over all agents  $j \neq i$ , it only remains to show what happens for the term in  $\mathbb{E}_{v,m}[z_{j,t}|\mu_t]$ . Defining  $\tilde{d}_{j,t}^{\gamma} = \gamma^{\text{th}} - \max_{k:k \neq j} \left\{ \mathbb{1}\{m_{k,t} = m_{j,t}\} \frac{\Delta v_{k,t}}{\mu_{k,t}} \right\}$  as the competing price in the imaginary setting where *i* follows strategy *A*, we have the following

$$\begin{aligned} \left| \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} [\boldsymbol{z}_{j,t} | \boldsymbol{\mu}_t] - Z_j(\boldsymbol{\mu}_t) \right| \stackrel{(a)}{=} \left| \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \mathbbm{1} \left\{ \boldsymbol{m}_{j,t} = \boldsymbol{m}_{i,t} \right\} \left( \boldsymbol{d}_j^{\gamma} \mathbbm{1} \left\{ \frac{\Delta \boldsymbol{v}_j}{\boldsymbol{\mu}_j} \ge \boldsymbol{d}_i^{\gamma} \right\} - \tilde{\boldsymbol{d}}_j^{\gamma} \mathbbm{1} \left\{ \frac{\Delta \boldsymbol{v}_j}{\boldsymbol{\mu}_j} \ge \tilde{\boldsymbol{d}}_j^{\gamma} \right\} \right) \right] \right| \\ \stackrel{(b)}{\leq} \frac{\Delta}{\underline{\mu}} a_{i,j} \end{aligned} \tag{43}$$

Equality (a) uses that  $\tilde{d}_j^{\gamma} = d_j^{\gamma}$  when *i* plays in a different auction from *j*; inequality (b) holds because  $\Delta v_j / \mu_j \leq \Delta / \mu$ .

Summing Equation 43 over all  $j \neq i$  gives

$$2\sum_{j\neq i} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ |\mu_{j,t} - \mu_{j}^{\star}| \cdot \left| \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} [\boldsymbol{z}_{j,t} | \boldsymbol{\mu}_{t}] - Z_{j}(\boldsymbol{\mu}_{t}) \right| \right] \leq 2\frac{\Delta}{\underline{\mu}} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{j\neq i} a_{i,j} |\mu_{j,t} - \mu_{j}^{\star}| \right]$$

$$\stackrel{(a)}{\leq} 2\frac{\Delta}{\underline{\mu}} ||a_{i}||_{2} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ ||\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}||_{2} \right] \qquad (44)$$

$$\stackrel{(b)}{\leq} 2\frac{\Delta}{\underline{\mu}} ||a_{i}||_{2} s_{t}^{1/2},$$

where inequality (a) follows from Cauchy-Schwartz inequality and (b) from Jensen's inequality.

Combining Equations 39, 40, 42 and 44 and noting that  $\overline{\epsilon}\hat{s}_t \ge s_t \ge \underline{\epsilon}\hat{s}_t$  gives the following recursion, i.e.,

$$\hat{s}_{t+1} \le (1 - 2\lambda \underline{\epsilon}) \hat{s}_t + 2\frac{\Delta}{\underline{\mu}} \|a_i\|_2 \overline{\epsilon}^{1/2} \hat{s}_t^{1/2} + \frac{N\Delta^2 \overline{\epsilon}}{\underline{\mu}^2}.$$
(45)

Since  $2\lambda \epsilon \leq 1$  because Assumption 3 holds for all agents, [6, Lemma C.5] yields:

$$\hat{s}_t \leq \hat{s}_1 (1 - \lambda \underline{\epsilon})^{t-1} + \frac{N \Delta^2 \overline{\epsilon}}{2\lambda \underline{\mu}^2 \underline{\epsilon}} + \frac{\Delta^2 ||a_i||_2^2 \overline{\epsilon}}{\lambda^2 \underline{\mu}^2 \underline{\epsilon}^2}$$

Noting that  $s_t \leq \overline{\epsilon} \hat{s}_t$  and that  $\hat{s}_1 \leq N \overline{\mu}^2 / \underline{\epsilon}$  finally gives the desired result, i.e.,

$$s_t \le N\overline{\mu}^2 \frac{\overline{\epsilon}}{\underline{\epsilon}} (1 - \lambda \underline{\epsilon})^{t-1} + \frac{N\Delta^2}{\lambda \underline{\mu}^2} \frac{\overline{\epsilon}^2}{\underline{\epsilon}} + \frac{\Delta^2 ||a_i||_2^2}{\lambda^2 \underline{\mu}^2} \frac{\overline{\epsilon}^2}{\underline{\epsilon}^2}.$$

#### C.2 Proof of Theorem 3.1

In this section, we prove Theorem 3.1 by considering the perspective of a single agent  $i \in N$ . We drop the subscript *i* for simplicity of notation.

*Step 1: Lower bound on Total Expected Cost in Hindsight.* We lower-bound the expected performance in hindsight using the Lagrangian dual function as follows:

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{Y}} \left[ \boldsymbol{C}^{H}(\boldsymbol{v},\boldsymbol{d}^{\boldsymbol{Y}}) \right] \stackrel{(a)}{\geq} \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ \sup_{\boldsymbol{\mu} \geq 0} \sum_{t=1}^{T} \boldsymbol{v}_{t} \tau - \boldsymbol{\mu} \boldsymbol{\rho} - (\Delta \boldsymbol{v}_{t} - \boldsymbol{\mu} \boldsymbol{d}_{t}^{\boldsymbol{Y}})^{+} \right],$$

$$\stackrel{(b)}{\geq} \sup_{\boldsymbol{\mu} \geq 0} \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ \sum_{t=1}^{T} \boldsymbol{v}_{t} \tau - \boldsymbol{\mu} \boldsymbol{\rho} - (\Delta \boldsymbol{v}_{t} - \boldsymbol{\mu} \boldsymbol{d}_{t}^{\boldsymbol{Y}})^{+} \right],$$

$$\stackrel{(c)}{=} \sup_{\boldsymbol{\mu} \geq 0} \sum_{t=1}^{T} \Psi(\boldsymbol{\mu}),$$

$$\stackrel{(d)}{=} T \Psi(\boldsymbol{\mu}^{\star}).$$
(46)

Inequality (a) uses Equation 6b; inequality (b) holds since  $\mathbb{E}_X[\sup_y f_y(X)] \ge \sup_y \mathbb{E}_X[f_y(X)]$ . Indeed,  $\mathbb{E}_X[\sup_y f_y(X)] \ge \mathbb{E}_X[f_y(X)]$  holds for all y, hence taking  $\sup_y$  on the right hand side gives the desired result. Inequality (c) uses the linearity of expectation and (d) the definition of  $\mu^*$  as the maximizer of  $\Psi$  in  $\mathbb{R}_+$ , unique under Assumption 2.

*Step 2: Upper bound on the Total Expected Cost of Strategy A.* We upper-bound the total cost of strategy *A* by considering that the worst always happens after budget depletion.

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{\gamma}} \left[ C^{A}(\boldsymbol{v},\boldsymbol{d}^{\boldsymbol{\gamma}}) \right] \stackrel{(a)}{\leq} \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{\gamma}} \left[ \sum_{t=1}^{\mathcal{T}} c_{t}^{A} \right] + \tau \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{\gamma}} \left[ T - \mathfrak{T} \right]$$

$$\stackrel{(b)}{\leq} \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{\gamma}} \left[ \sum_{t=1}^{\mathcal{T}} c_{t}^{A} \right] + \tau \left( \frac{\Delta}{\rho \underline{\mu}} + \frac{\overline{\mu}}{\rho \epsilon} \right)$$

$$(47)$$

Inequality (*a*) is obtained by assuming that the agent always gets maximum utility  $v_t = 1$  after time T, but never manages to access the fast road. Inequality (*b*) follows from Lemma C.1.

Step 3: Upper bound on Expected Cost-per-Period. For the rest of the proof, we only consider  $t \leq T$ , where the remaining budget is greater than the maximum admissible expense. We first rewrite the cost-per-period in the following manner:

$$c_t = v_t \left( \tau - \Delta \mathbb{1} \{ \Delta v_t > \mu_t d_t^{\gamma} \} \right),$$
  
$$= v_t \tau - \mu_t \rho - \left( \Delta v_t - \mu_t d_t^{\gamma} \right)^+ + \mu_t \rho - \mu_t d_t^{\gamma} \mathbb{1} \{ \Delta v_t > \mu_t d_t^{\gamma} \}.$$
(48)

Noting that the first three terms are reminiscent of the dual objective, while the last one corresponds to the period's expenses  $z_t$ , we further define  $C(\mu) = \mathbb{E}_{v,dY}[c|\mu] = \Psi(\mu) + \mu(\rho - Z(\mu))$  the expected cost-per-period. Note that Assumption 2 ensures it is twice differentiable with derivatives  $C'(\mu) = -\mu Z'(\mu)$  and  $C''(\mu) = -(Z'(\mu) + \mu Z''(\mu))$ .

Since the maximizer  $\mu^*$  lies in the open set  $]\mu, \overline{\mu}[$  under Assumption 3, it must be a local maximum of the dual function  $\Psi$ . Moreover since the dual function is differentiable under Assumption 2, its derivative must be null in  $\mu^*$ , hence:

$$Z(\mu^{\star}) = \rho, \tag{49}$$

This implies that  $C(\mu^{\star}) = \Psi(\mu^{\star})$ .

Taking expectation on Equation 48 and making a Taylor expansion in  $\mu^*$  gives the following for some  $\zeta$  between  $\mu_t$  and  $\mu^*$ :

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{\gamma}}\left[c_{t}\right] = \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{\gamma}}\left[C(\mu_{t})\right],$$

$$= C(\mu^{\star}) + \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{\gamma}}\left[\mu_{t} - \mu^{\star}\right]C'(\mu^{\star}) + \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{\gamma}}\left[(\mu_{t} - \mu^{\star})^{2}\right]\frac{C''(\zeta)}{2},$$

$$\leq \Psi(\mu^{\star}) + \left|\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{\gamma}}\left[\mu_{t} - \mu^{\star}\right]\right|\mu^{\star}\overline{Z'} + \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{\gamma}}\left[(\mu_{t} - \mu^{\star})^{2}\right]\frac{\overline{\mu}\overline{Z''} + \overline{Z'}}{2},$$

$$= \Psi(\mu^{\star}) + r_{t}\overline{Z'} + s_{t}\frac{\overline{\mu}\overline{Z''} + \overline{Z'}}{2}.$$
(50)

We next upper bound the mean squared error  $s_t = \mathbb{E}_{v,dv} \left[ (\mu_t - \mu^*)^2 \right]$  and the absolute mean error  $r_t = \mu^* \left| \mathbb{E}_{v,dv} \left[ \mu_t - \mu^* \right] \right|$ , defined for all  $t \leq \mathcal{T}$ .

Step 4: Upper bound on Absolute Mean Error  $r_t$ . The projection error  $P_t = \mu_t + \epsilon(z_t^A - \rho) - P_{[\underline{\mu}, \overline{\mu}]}(\mu_t + \epsilon(z_t - \rho))$  satisfies:

$$P_{t} \stackrel{(a)}{\geq} \left(\mu_{t} + \epsilon \left(z_{t} - \rho\right) - \underline{\mu}\right) \mathbb{1} \left\{\mu_{t} + \epsilon \left(z_{t} - \rho\right) < \underline{\mu}\right\},$$

$$\stackrel{(b)}{\geq} -\epsilon\rho \mathbb{1} \left\{\mu_{t} + \epsilon \left(z_{t} - \rho\right) < \underline{\mu}\right\},$$

$$\stackrel{(c)}{\geq} -\epsilon\rho \mathbb{1} \{\mu_{t} < \underline{\mu} + \epsilon\rho\}.$$
(51)

Inequality (*a*) holds as the projection error is negative only when  $\mu_t + \epsilon(z_t - \rho) - \mu < 0$ ; inequality (*b*) uses  $\mu_t \ge \mu$  and the non-negativity of  $z_t$ ; inequality (*c*) finally holds since  $(\mu_t + \epsilon z_t < \mu + \epsilon \rho)$  implies  $(\mu_t < \overline{\mu} + \epsilon \rho)$  by non-negativity of  $z_t$ .

Let  $\xi = (\mu^* - \underline{\mu})/2$ , note that  $\xi$  is strictly positive under Assumption 3. We take expectation on Equation 51.

$$-\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}[P_t] \leq \epsilon \rho \mathbb{P}\{\mu_t < \underline{\mu} + \epsilon \rho\}$$

$$\stackrel{(a)}{\leq} \epsilon \rho \mathbb{P}\{\mu_t - \mu^{\star} \leq -\xi\}$$

$$\stackrel{(b)}{\leq} \epsilon \rho \mathbb{P}\{(\mu_t - \mu^{\star})^2 \geq \xi^2\}$$

$$\stackrel{(c)}{\leq} \epsilon \frac{\rho}{\xi^2} s_t$$
(52)

Inequality (a) holds since events  $\{\mu_t < \mu + \epsilon \rho\}$  and  $\{\mu_t - \mu^* < \mu + \epsilon \rho - \mu^* \le -\xi\}$  are equal because  $\mu^* \ge \mu + \epsilon \rho + \xi$  under Assumption 3, and since the latter is in turn a subset of  $\{\mu_t - \mu^* \le -\xi\}$ . Inequality (b) uses that  $\{\mu_t - \mu^* \le -\xi\} = \{\mu^* - \mu_t \ge \xi\} \subset \{(\mu_t - \mu^*)^2 \ge \xi^2\}$  since the square function is monotonic on  $\mathbb{R}_+$ . Inequality (c) finally follows from Markov inequality. We now proceed to bound the absolute mean error  $r_t = \mu^* |\mathbb{E}_{v,d^{\gamma}}[\mu_t - \mu^*]|$ . From the update rule  $\mu_{t+1} = \mu_t + \epsilon(z_t - \rho) - P_t$  we get the following:

$$\mu^{\star} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}^{\boldsymbol{\gamma}}} \left[ \mu_{t+1} - \mu^{\star} \right] = \mu^{\star} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}^{\boldsymbol{\gamma}}} \left[ \mu_{t} - \mu^{\star} \right] + \epsilon \mu^{\star} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}^{\boldsymbol{\gamma}}} \left[ z_{t} - \rho \right] - \mu^{\star} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}^{\boldsymbol{\gamma}}} \left[ P_{t} \right],$$

$$\stackrel{(a)}{=} \mu^{\star} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}^{\boldsymbol{\gamma}}} \left[ \mu_{t} - \mu^{\star} \right] + \epsilon \mu^{\star} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}^{\boldsymbol{\gamma}}} \left[ Z(\mu_{t}) - Z(\mu^{\star}) + Z(\mu^{\star}) - \rho \right] - \mu^{\star} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}^{\boldsymbol{\gamma}}} \left[ P_{t} \right],$$

$$\stackrel{(b)}{=} \mu^{\star} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}^{\boldsymbol{\gamma}}} \left[ \mu_{t} - \mu^{\star} \right] - \mu^{\star} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}} \left[ P_{t} \right],$$

$$+ \epsilon \mu^{\star} \left( Z'(\mu^{\star}) \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}^{\boldsymbol{\gamma}}} \left[ \mu_{t} - \mu^{\star} \right] + \frac{1}{2} Z''(\zeta) \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}^{\boldsymbol{\gamma}}} \left[ (\mu_{t} - \mu^{\star})^{2} \right] \right).$$
(53)

Equality (a) is obtained by taking expectation conditioned on  $\mu_t$  inside the expectation in the middle term. Equality (b) uses a Taylor expansion of Z in  $\mu^*$  for some  $\zeta$  between  $\mu_t$  and  $\mu^*$ .

Taking absolute values in Equation 53 combined with Equation 52 gives the following recursion:

$$\begin{aligned} r_{t+1} &\leq \left| 1 + \epsilon Z'(\mu^{\star}) \right| r_t + \epsilon \mu^{\star} \left( \frac{\overline{Z''}}{2} + \frac{\rho}{\xi^2} \right) s_t, \\ \stackrel{(a)}{=} \left( 1 + \epsilon Z'(\mu^{\star}) \right) r_t + \epsilon \mu^{\star} \left( \frac{\overline{Z''}}{2} + \frac{\rho}{\xi^2} \right) s_t, \\ \stackrel{(b)}{\leq} (1 - \epsilon \lambda) r_t + \epsilon \overline{\mu} \left( \frac{\overline{Z''}}{2} + \frac{\rho}{\xi^2} \right) s_t. \end{aligned}$$

Note that we have  $-\overline{Z}' \leq Z'(\mu) \leq -\lambda$  for all  $\mu \geq 0$  under Assumption 2. Hence Equality (*a*) holds because  $1 + \epsilon Z'(\mu^*) \geq 1 - \epsilon \overline{Z}' \geq 0$  under Assumption 3. Inequality (*b*) on the other hand uses that  $1 + \epsilon Z'(\mu^*) \leq 1 - \epsilon \lambda$  and the bound on  $\mu^*$  from Assumption 3.

Since  $\lambda \epsilon \leq 1$  and  $r_1 \leq \overline{\mu}^2$  under Assumption 3, [6, Lemma C.4] then yields:

$$r_t \leq \overline{\mu}^2 \left(1 - \epsilon \lambda\right)^{t-1} + \epsilon \overline{\mu} \left(\frac{\overline{Z''}}{2} + \frac{\rho}{\xi^2}\right) \sum_{j=1}^{t-1} (1 - \epsilon \lambda)^{t-1-j} s_j.$$

Noting that the partial sums in  $(1 - \epsilon \lambda)^t$  are smaller than their limit  $1/(\epsilon \lambda)$ , we finally get the following upper bound on the total absolute error:

$$\sum_{t=1}^{\mathcal{T}} r_t \leq \overline{\mu}^2 \sum_{t=1}^{\mathcal{T}} (1 - \epsilon \lambda)^{t-1} + \epsilon \overline{\mu} \left( \frac{\overline{Z''}}{2} + \frac{\rho}{\xi^2} \right) \sum_{t=1}^{\mathcal{T}} \sum_{j=1}^{t-1} (1 - \epsilon \lambda)^{t-1-j} s_j,$$

$$\leq \frac{\overline{\mu}^2}{\epsilon \lambda} + \epsilon \overline{\mu} \left( \frac{\overline{Z''}}{2} + \frac{\rho}{\xi^2} \right) \sum_{j=1}^{\mathcal{T}-1} s_j \sum_{t=0}^{\mathcal{T}-1-j} (1 - \epsilon \lambda)^t,$$

$$\leq \frac{\overline{\mu}^2}{\epsilon \lambda} + \left( \frac{\overline{\mu} \overline{Z''}}{2\lambda} + \frac{\overline{\mu} \rho}{\lambda \xi^2} \right) \sum_{j=1}^{\mathcal{T}} s_j.$$
(54)

The same trick combined with Lemma C.2 also gives:

$$\sum_{j=1}^{\mathcal{T}} s_j \leq \mathfrak{T} \frac{\Delta^2 \epsilon}{2\lambda \mu^2} + \overline{\mu}^2 \sum_{j=1}^{\mathcal{T}} (1 - 2\lambda \underline{\epsilon})^{t-1},$$
  
$$\leq \mathfrak{T} \frac{\Delta^2 \epsilon}{2\lambda \mu^2} + \frac{\overline{\mu}^2}{2\lambda \epsilon}.$$
(55)

Conclusion: Together, Equations 46, 47, 50, 54 and 55 finally give:

$$\begin{split} \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{Y}}\left[C^{A}(\boldsymbol{v},\boldsymbol{d}^{\boldsymbol{Y}})-C^{H}(\boldsymbol{v},\boldsymbol{d}^{\boldsymbol{Y}})\right] &\leq \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{Y}}\left[\sum_{t=1}^{\mathcal{T}}c_{t}\right] + \tau\left(\frac{\Lambda}{\rho\underline{\mu}}+\frac{\overline{\mu}}{\rho\epsilon}\right) - T\Psi(\mu^{\star}),\\ &\leq \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{Y}}\left[\overline{Z'}\sum_{t=1}^{\mathcal{T}}r_{t}+\frac{\overline{\mu}\overline{Z''}+\overline{Z'}}{2}\sum_{t=1}^{\mathcal{T}}s_{t}\right] + \tau\left(\frac{\Lambda}{\rho\underline{\mu}}+\frac{\overline{\mu}}{\rho\epsilon}\right),\\ &\leq \left(\frac{\overline{\mu}\overline{Z''}+\overline{Z'}}{2}+\frac{\overline{\mu}\overline{Z'}}{\lambda}\left(\frac{\overline{Z''}}{2}+\frac{\rho}{\xi^{2}}\right)\right)\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}\boldsymbol{Y}}\left[\sum_{t=1}^{\mathcal{T}}s_{t}\right]\\ &+\frac{\overline{\mu}^{2}\overline{Z'}}{\epsilon\lambda} + \tau\left(\frac{\Lambda}{\rho\underline{\mu}}+\frac{\overline{\mu}}{\rho\epsilon}\right),\\ &\leq \left(\frac{\overline{\mu}\overline{Z''}+\overline{Z'}}{2}+\frac{\overline{\mu}\overline{Z'}}{\lambda}\left(\frac{\overline{Z''}}{2}+\frac{\rho}{\xi^{2}}\right)\right)\left(\frac{\Lambda^{2}\epsilon T}{2\lambda\underline{\mu}^{2}}+\frac{\overline{\mu}^{2}}{2\lambda\epsilon}\right)\\ &+\frac{\overline{\mu}^{2}\overline{Z'}}{\epsilon\lambda} + \tau\left(\frac{\Lambda}{\rho\underline{\mu}}+\frac{\overline{\mu}}{\rho\epsilon}\right). \end{split}$$

Hence there exist a constant C in  $\mathbb{R}_+$  such that

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}^{\boldsymbol{\gamma}}}\left[C^{A}(\boldsymbol{v},\boldsymbol{d}^{\boldsymbol{\gamma}})-C^{H}(\boldsymbol{v},\boldsymbol{d}^{\boldsymbol{\gamma}})\right] \leq C\left(1+\frac{1}{\epsilon}+\epsilon T\right).$$

## C.3 **Proof of Theorem 3.2**

We first prove the convergence on average of the multiplier profile  $\mu_t$  to the optimal  $\mu^{\star}$ .

Step 1: Convergence on Average of the Multiplier Profile  $\mu_t$ . Combining Lemma C.1 and C.2 gives the following:

$$\frac{1}{T} \sum_{t=1}^{T} \mathbb{E}_{\boldsymbol{v}} \left[ \|\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}\|_{2}^{2} \right] = \frac{1}{T} \mathbb{E}_{\boldsymbol{v}} \left[ \sum_{t=1}^{T} \|\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}\|_{2}^{2} \mathbb{1} \{t \leq \underline{\Upsilon}\} + \|\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}\|_{2}^{2} \mathbb{1} \{t > \underline{\Upsilon}\} \right],$$

$$\overset{(a)}{\leq} \frac{1}{T} \mathbb{E}_{\boldsymbol{v}} \left[ \sum_{t=1}^{\underline{\Upsilon}} s_{t} \right] + \frac{N\overline{\mu}^{2}}{T} \mathbb{E}_{\boldsymbol{v}} \left[ T - \underline{\Upsilon} \right],$$

$$\overset{(b)}{\leq} N \overline{\mu}^{2} \frac{\overline{\epsilon}}{\underline{\epsilon}T} \sum_{t=1}^{T} (1 - 2\lambda \underline{\epsilon})^{t-1} + \frac{N\Delta^{2}\overline{\epsilon}^{2}}{2\lambda \underline{\mu}^{2} \underline{\epsilon}} + \frac{N\overline{\mu}^{2}}{T} \left( \frac{\Delta}{\underline{\rho}\underline{\mu}} + \frac{\overline{\mu}}{\underline{\rho}\underline{\epsilon}} \right),$$

$$\overset{(c)}{\leq} N \left( \frac{\overline{\mu}^{2}}{2\lambda} \frac{\overline{\epsilon}}{\underline{\epsilon}^{2}T} + \frac{\Delta^{2}\overline{\epsilon}^{2}}{2\lambda \underline{\mu}^{2} \underline{\epsilon}} + \frac{\overline{\mu}^{2}}{T} \left( \frac{\Delta}{\underline{\rho}\underline{\mu}} + \frac{\overline{\mu}}{\underline{\rho}\underline{\epsilon}} \right) \right).$$

Inequality (*a*) holds by assuming that the error is maximal after time  $\underline{T}$ ; inequality (*b*) uses both Lemma C.1 and C.2, and inequality (*c*) holds by bounding the partial geometric series in  $1 - 2\lambda \underline{\epsilon}$  by its limit  $1/(2\lambda \underline{\epsilon})$ .

Hence there exist some  $C_1$  in  $\mathbb{R}_+$  such that

$$\frac{1}{T}\sum_{t=1}^{T} \mathbb{E}_{\boldsymbol{v}}\left[\|\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}\|_{2}^{2}\right] \leq C_{1}N\left(\frac{\overline{\epsilon}^{2}}{\underline{\epsilon}} + \frac{\overline{\epsilon} + \underline{\epsilon}^{2}}{\underline{\epsilon}^{2}T}\right)$$

We now turn to the second part of the theorem and show that the convergence in multiplier profile implies the convergence in cost.

Step 2: Upper Bound on the Total Expected Cost of Strategy A. Similarly as in the second step of Section C.2, we bound the expected total cost of agent  $i \in N$  by considering it always gets maximum valuation after time  $\underline{T}$  but never accesses the priority road.

$$C_{i}^{A} \leq \mathbb{E}_{\boldsymbol{v}} \left[ \sum_{t=1}^{\mathcal{T}} c_{i,t} \right] + \tau \mathbb{E}_{\boldsymbol{v}} \left[ T - \underline{\mathcal{T}} \right]$$

$$\stackrel{(a)}{\leq} \mathbb{E}_{\boldsymbol{v}} \left[ \sum_{t=1}^{\mathcal{T}} c_{i,t} \right] + \tau \left( \frac{\Delta}{\underline{\rho\mu}} + \frac{\overline{\mu}}{\underline{\rho\underline{\epsilon}}} \right)$$
(56)

Inequality (a) uses Lemma C.1.

We now prove the Lipschitz continuity of  $\Psi_i$  and  $Z_i$  in the more general setting with parallel auctions. Choosing the probability of agent *i* and *j* to play in the same auction  $a_{i,j} = 1$  equal to one for all pairs gives the setting considered in Section 3.3.

Step 3: Lipschitz Continuity of the Expected Dual Objective  $\Psi_i$ . For an an agent  $i \in N$ , recall the definition of its dual function  $\Psi_i(\boldsymbol{\mu}) = \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ v_i \tau - \mu_i \rho_i - (\Delta v_i - \mu_i d_i^{\gamma})^+ \right]$ , where  $d_i^{\gamma} = \gamma^{\text{th}} - \max_{j:j \neq i} \left\{ \mathbb{1} \{ m_j = m_i \} \Delta v_j / \mu_j \right\}$ .

For any realized vectors  $\boldsymbol{v} = (v_j)_{j=1}^N$  and  $\boldsymbol{m} = (m_j)_{j=1}^N$ , the function  $\boldsymbol{\mu} \mapsto (\Delta v_i - \mu_i d_i^{\gamma})^+$  is differentiable in  $\mu_i$  with derivative bounded by  $\Delta/\mu$ , except in the set  $\{(\boldsymbol{v}, \boldsymbol{m}) : \Delta v_i = \mu_i d_i^{\gamma}\}$  of measure zero since valuations are independent and absolutely continuous with support in [0, 1] under Assumption 1. Leibniz's integral rule hence implies:

$$\frac{\partial \Psi_i}{\partial \mu_i}(\boldsymbol{\mu}) = \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ d_i^{\gamma} \mathbb{1} \{ \Delta v_i \ge \mu_i d_i^{\gamma} \} \right] - \rho_i.$$

Hence Assumption 3 ensures the derivative is bounded, i.e.,

$$\left|\frac{\partial \Psi_i}{\partial \mu_i}(\boldsymbol{\mu})\right| \le \frac{\Delta}{\underline{\mu}}.$$
(57)

Furthermore, consider  $j \neq i$  and let  $d_{i,j}^{\gamma} = \gamma^{\text{th}} - \max_{\ell \neq i,j} \{\mathbb{1}\{m_{\ell} = m_i\}\Delta v_{\ell}/\mu_{\ell}\}$  be the competing price for *i* before *j* places its own bid. We then rewrite:

$$(\Delta v_i - \mu_i d_i^{\gamma})^+ = \begin{cases} (\Delta v_i - \mu_i \Delta v_j / \mu_j)^+ & \text{if } m_j = m_i \text{ and } d_{i,j}^{\gamma-1} \ge \Delta v_j / \mu_j \ge d_{i,j}^{\gamma} \\ (\Delta v_i - \mu_i d_{i,j}^{\gamma})^+ & \text{otherwise.} \end{cases}$$

The function  $\boldsymbol{\mu} \mapsto (\Delta v_i - \mu_i d_i^{\gamma})^+$  is hence differentiable in  $\mu_j$  with derivative bounded by  $\Delta \overline{\mu_i} / \underline{\mu_j}^2$  outside of the sets  $\left\{ (\boldsymbol{v}, \boldsymbol{m}) : d_{i,j}^{\gamma-1} > \frac{\Delta v_i}{\mu_i} = \frac{\Delta v_j}{\mu_j} > d_{i,j}^{\gamma}, \ m_i = m_j \right\}, \left\{ (\boldsymbol{v}, \boldsymbol{m}) : \frac{\Delta v_i}{\mu_i} \ge \frac{\Delta v_j}{\mu_j} = d_{i,j}^{\gamma-1}, \ m_i = m_j \right\}$ 

and  $\left\{ (v, m) : \frac{\Delta v_i}{\mu_i} \ge \frac{\Delta v_j}{\mu_j} = d_{i,j}^{\gamma}, m_i = m_j \right\}$ , which are all of measure zero using the same argument. Leibniz's integral rule then implies that  $\Psi_i$  is differentiable with respect to  $\mu_j$ , i.e.,

$$\frac{\partial \Psi_i}{\partial \mu_j}(\boldsymbol{\mu}) = \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}}\left[\frac{\Delta v_j \mu_i}{\mu_j^2} \mathbb{1}\left\{\frac{v_i}{\mu_i} \geq \frac{v_j}{\mu_j}, \ d_{i,j}^{\gamma-1} \geq \frac{\Delta v_j}{\mu_j} \geq d_{i,j}^{\gamma}, \ m_i = m_j\right\}\right].$$

Using that  $v_i/\mu_i \ge v_j/\mu_j$  in the indicator function, we can bound the derivative as follows, i.e.,

$$\left|\frac{\partial \Psi_i}{\partial \mu_j}(\boldsymbol{\mu})\right| \leq \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}}\left[\frac{\Delta v_i}{\mu_j}\mathbb{1}\left\{m_i = m_j\right\}\right] \leq \frac{\Delta}{\underline{\mu}}a_{i,j}.$$
(58)

Together, Equations 57 and 58 finally imply the Lipschitz continuity of  $\Psi_i$  on U.

$$\left|\Psi_{i}(\boldsymbol{\mu})-\Psi_{i}(\boldsymbol{\mu}')\right| \leq \frac{\Delta}{\underline{\mu}} \left(|\mu_{i}-\mu_{i}'|+\sum_{j\neq i}a_{i,j}|\mu_{j}-\mu_{j}'|\right).$$
(59)

We hence define  $\mathcal{L}_{\Psi} = \frac{\Delta}{\mu}$  the Lipschitz constant of  $\Psi_i$  in the case of a single auction.

Step 4: Bound on the Derivative of the Cumulative Distribution Function of Competing Prices. In order to prove the Lipschitz continuity of  $Z_i$ , we first bound the derivative of the cumulative distribution function of competing prices  $H_i(x, \mu_{-i}) = \mathbb{P}\left[d_i^{\gamma} \leq x\right]$ . Since [6, Lemma C.2] proposes such a bound when  $\gamma = 1$ , we only consider the case where  $\gamma > 1$ .

Let  $\mathcal{V}_j$  denote the cumulative distribution function of valuations associated with the density of valuations  $v_j$  of agent  $j \in N$ , and let  $\overline{\mathcal{V}}_j$  denote the function  $y \mapsto 1 - \mathcal{V}_j(y)$ . We further define  $\mathcal{M}_i \in 2^{N \setminus \{i\}}$  as the set of agents different from *i* that play in the same auction as *i*. For a fixed  $\mu_{-i} \in \mathbb{R}^{N-1}_+$ , we write the cumulative distribution of competing prices by conditioning on each agent  $j \neq i$  being the price setter for agent *i*, an event that we denote by  $\{j \to i\}$ .

$$H_{i}(x, \boldsymbol{\mu}_{-i}) = \mathbb{P}\left[d_{i}^{Y} \leq x\right]$$

$$\stackrel{(a)}{=} \sum_{\mathcal{M}_{i}} \mathbb{P}\left[\mathcal{M}_{i}\right] \mathbb{P}\left[d_{i}^{Y} \leq x|\mathcal{M}_{i}\right]$$

$$\stackrel{(b)}{=} \sum_{\mathcal{M}_{i}} \mathbb{P}\left[\mathcal{M}_{i}\right] + \sum_{\mathcal{M}_{i}} \sum_{p \in \mathcal{M}_{i}} \mathbb{P}\left[\mathcal{M}_{i}\right] \mathbb{P}\left[\left\{\frac{\Delta v_{p}}{\mu_{p}} \leq x\right\} \cap \{p \to i\} \middle| \mathcal{M}_{i}\right]\right]$$

$$\stackrel{(c)}{=} \sum_{\substack{\mathcal{M}_{i} \\ |\mathcal{M}_{i}| \leq \gamma}} \mathbb{P}\left[\mathcal{M}_{i}\right] + \sum_{\substack{\mathcal{M}_{i} \\ |\mathcal{M}_{i}| \geq \gamma}} \sum_{p \in \mathcal{M}_{i}} \sum_{\substack{W \subset \mathcal{M}_{i} \setminus \{p\} \\ |W| = \gamma - 1}} \mathbb{P}\left[\mathcal{M}_{i}\right]$$

$$\times \mathbb{P}\left[\left\{\frac{\Delta v_{p}}{\mu_{p}} \leq x\right\} \bigcap_{w \in W} \left\{\frac{v_{w}}{\mu_{w}} \geq \frac{v_{p}}{\mu_{p}}\right\} \bigcap_{\ell \in \mathcal{M}_{i} \setminus W} \left\{\frac{v_{\ell}}{\mu_{\ell}} \leq \frac{v_{p}}{\mu_{p}}\right\} \middle| \mathcal{M}_{i}\right]$$

$$\stackrel{(d)}{=} \sum_{\substack{\mathcal{M}_{i} \\ |\mathcal{M}_{i}| \leq \gamma}} \prod_{k \in \mathcal{M}_{i}} a_{i,k} + \sum_{\substack{\mathcal{M}_{i} \\ |\mathcal{M}_{i}| \geq \gamma}} \sum_{p \in \mathcal{M}_{i}} \sum_{\substack{W \subset \mathcal{M}_{i} \setminus \{p\} \\ |W| = \gamma - 1}} \prod_{k \in \mathcal{M}_{i}} a_{i,k}$$

$$\times \int_{0}^{x\mu_{p}/\Delta} v_{p}(y) \prod_{w \in W} \overline{\mathcal{V}}_{w}\left(\frac{y\mu_{w}}{\mu_{p}}\right) \prod_{\ell \in \mathcal{M}_{i} \setminus W} \mathcal{V}_{\ell}\left(\frac{y\mu_{\ell}}{\mu_{p}}\right) dy$$

$$(60)$$

Equality (*a*), (*b*) and (*c*) each follow from the Law of total probability, by partitioning on realizations of  $M_i$ , then price setters *p*, and finally sets of auction winners *W*. Equality (*d*) holds since valuations  $\boldsymbol{v} = (v_j)_{i=1}^N$  and auctions  $\boldsymbol{m} = (m_i)_{i=1}^N$  are drawn independently across agents.

Now consider an agent  $j \neq i$ . For each term of the sums, note that the integrand is differentiable with respect to  $\mu_j$  almost everywhere with derivative bounded by  $y\overline{\nu\mu}/\mu^2$ , which is integrable on  $[0, x\overline{\mu}/\Delta]$  for all  $x \ge 0$ . Leibniz's integration rule hence implies that  $H_i(\overline{x}, \mu_{-i})$  is differentiable in  $\mu_j$ , i.e.,

$$\frac{\partial H_i(x, \boldsymbol{\mu}_{-i})}{\partial \mu_j} = \sum_{\substack{\mathcal{M}_i \\ |\mathcal{M}_i| \ge \gamma \\ j \in \mathcal{M}_i}} A(\mathcal{M}_i) + B(\mathcal{M}_i) + C(\mathcal{M}_i), \tag{61}$$

where we regroup in A the terms where  $j \in W$ , in B the terms verifying  $j \notin W \cup \{p\}$ , and in C the terms where p = j, with A, B, and C, given by

$$\begin{split} A(\mathcal{M}_{i}) &= -\sum_{\substack{p \in \mathcal{M}_{i} \ W \subset \mathcal{M}_{i} \setminus \{p\} \ k \in \mathcal{M}_{i}}} \sum_{\substack{k \in \mathcal{M}_{i} \ W \in \mathcal{M}_{i} \setminus \{p\} \ k \in \mathcal{M}_{i}}} \prod_{\substack{a_{i,k} \ \int_{0}^{x\mu_{p}/\Delta} \frac{y}{\mu_{p}} v_{j}\left(\frac{y\mu_{j}}{\mu_{p}}\right) v_{p}(y)} \\ &\prod_{\substack{w \in W \ W \neq j}} \overline{\nabla}_{w}\left(\frac{y\mu_{w}}{\mu_{p}}\right) \prod_{\substack{\ell \in \mathcal{M}_{i} \setminus W \ \ell \neq p}} \nabla_{\ell}\left(\frac{y\mu_{\ell}}{\mu_{p}}\right) dy, \\ B(\mathcal{M}_{i}) &= \sum_{\substack{p \in \mathcal{M}_{i} \ W \subset \mathcal{M}_{i} \setminus \{p\} \ k \in \mathcal{M}_{i}}} \sum_{\substack{k \in \mathcal{M}_{i} \ M \in \mathcal{M}_{i}}} \prod_{\substack{a_{i,k} \ \int_{0}^{x\mu_{p}/\Delta} \frac{y}{\mu_{p}} v_{j}\left(\frac{y\mu_{j}}{\mu_{p}}\right) v_{p}(y)} \\ &\prod_{\substack{w \in W \ W \subset \mathcal{M}_{i} \setminus \{p\} \ k \in \mathcal{M}_{i}}} \sum_{\substack{k \in \mathcal{M}_{i} \ M \in \mathcal{M}_{i}}} \nabla_{\ell}\left(\frac{y\mu_{\ell}}{\mu_{p}}\right) dy, \\ C(\mathcal{M}_{i}) &= \left(C_{1}(\mathcal{M}_{i}) + C_{2}(\mathcal{M}_{i})\right) \prod_{\substack{k \in \mathcal{M}_{i} \ \mu_{j}^{2}}} y_{k}\left(\frac{y\mu_{k}}{\mu_{j}}\right) v_{j}(y) \prod_{\substack{w \in W \ W \neq W}} \overline{\nabla}_{w}\left(\frac{y\mu_{k}}{\mu_{j}}\right) dy, \\ C_{2}(\mathcal{M}_{i}) &= -\sum_{\substack{k \in \mathcal{M}_{i} \ W \subset \mathcal{M}_{i} \setminus \{j\} \ M \in \mathcal{M}_{i}}} \sum_{\substack{y \neq \mu_{j} \ \mu_{j}^{2}}} \int_{0}^{x\mu_{j}/\Delta} \frac{y\mu_{k}}{\mu_{j}^{2}} v_{k}\left(\frac{y\mu_{k}}{\mu_{j}}\right) v_{j}(y) \prod_{\substack{w \in W \ W \notin W}} \overline{\nabla}_{w}\left(\frac{y\mu_{k}}{\mu_{j}}\right) dy, \\ c_{2}(\mathcal{M}_{i}) &= -\sum_{\substack{k \in \mathcal{M}_{i} \ W \subset \mathcal{M}_{i} \setminus \{j\} \ M \in \mathcal{M}_{i}}} \sum_{\substack{y \in \mathcal{M}_{i} \setminus \{j\} \ \mu_{j}^{2}}} \int_{0}^{x\mu_{j}/\Delta} \frac{y\mu_{k}}{\mu_{j}^{2}} v_{k}\left(\frac{y\mu_{k}}{\mu_{j}}\right) v_{j}(y) \prod_{\substack{w \in W \ W \notin W}} \overline{\nabla}_{w}\left(\frac{y\mu_{k}}{\mu_{j}}\right) dy. \\ c_{2}(\mathcal{M}_{i}) &= -\sum_{\substack{k \in \mathcal{M}_{i} \ W \subset \mathcal{M}_{i} \setminus \{j\} \ H \in \mathcal{M}_{i}}} \sum_{\substack{w \in \mathcal{M}_{i} \ \mu_{j}^{2}}} \sum_{\substack{w \in \mathcal{$$

The expression of *C* follows from the product rule. We regrouped in  $C_1$  the terms of the product rule where the running index *k* belongs to *W*, and in  $C_2$  those where *k* belongs to  $\mathcal{M}_i \setminus W \cup \{j\}$ .

We now proceed to bound the sum of the terms in  $A(\mathcal{M}_i)$  in Equation 61. First, we write the cumulative distribution of  $d_{i,j}^{\gamma-1} = \gamma - 1^{\text{th}} - \max_{\ell \neq i,j} \{\mathbb{1}\{m_\ell = m_i\} \Delta v_\ell / \mu_\ell\}$  in a similar manner as in

Equation 60.

$$\mathbb{P}\left[d_{i,j}^{\gamma-1} \leq x\right] = \sum_{\substack{\mathcal{M}_i \\ |\mathcal{M}_i| < \gamma-1 \\ j \notin \mathcal{M}_i}} \prod_{\substack{k \in \mathcal{M}_i \\ |\mathcal{M}_i| \geq \gamma-1 \\ j \notin \mathcal{M}_i}} a_{i,k} + \sum_{\substack{\mathcal{M}_i \\ |\mathcal{M}_i| \geq \gamma-1 \\ j \notin \mathcal{M}_i}} \sum_{\substack{P \in \mathcal{M}_i \\ |\mathcal{W}| = \gamma-2 \\ |W| = \gamma-2}} \prod_{\substack{k \in \mathcal{M}_i \\ |W| = \gamma-2}} a_{i,k} \\ \times \int_0^{x \overline{\mu}_p / \Delta} v_p(y) \prod_{\substack{W \in W}} \overline{\mathcal{V}}_w \left(\frac{y\mu_w}{\mu_p}\right) \prod_{\substack{\ell \in \mathcal{M}_i \setminus W \\ \ell \neq p}} \mathcal{V}_\ell \left(\frac{y\mu_\ell}{\mu_p}\right) dy \tag{63}$$

Now consider the term *S* in the expression of *A* defined by the realization ( $\mathcal{M}_i, p, W$ ), and denote  $\mathcal{M}'_i = \mathcal{M}_i \setminus \{j\}$  as well as  $W' = W \setminus \{j\}$ . Note that ( $\mathcal{M}'_i, p, W'$ ) defines a term *S'* in Equation 63. Looking at the expression of *S* and *S'*, we verify that multiplying the integrand of *S* by a factor  $-a_{i,j}\frac{y}{\mu_p}v_j\left(\frac{y\mu_j}{\mu_p}\right)$  gives the integrand of *S'*. Since  $y \leq x\mu_p/\Delta$ , this factor is bounded by  $a_{i,j}x\overline{v}/\Delta$  under Assumption 1, in turn bounded by  $a_{i,j}\overline{v}/\mu$  since  $x \leq \Delta/\mu$ . Since this bound holds for all terms *S*, it holds for the sum. Noting that  $\mathbb{P}\left[d_{i,j}^{Y^{-1}} \leq x\right] \leq 1$  gives the following bound on *A*,

$$\sum_{\substack{\mathcal{M}_i \\ |\mathcal{M}_i| \ge \gamma \\ j \in \mathcal{M}_i}} A(\mathcal{M}_i) \middle| \le a_{i,j} \frac{\overline{\nu}}{\underline{\mu}} \mathbb{P}\left[ d_{i,j}^{\gamma-1} \le x \right] \le a_{i,j} \frac{\overline{\nu}}{\underline{\mu}}.$$
(64)

Furthermore, note that rewriting k = p and making the change of variable  $z = y\mu_p/\mu_j$  gives the following expression for  $C_1(\mathcal{M}_i)$ , i.e.,

$$C_{1}(\mathcal{M}_{i}) = \sum_{\substack{p \in \mathcal{M}_{i} \\ p \neq j}} \sum_{\substack{W \subset \mathcal{M}_{i} \setminus \{j\} \\ |W| = \gamma - 1 \\ p \in W}} \int_{0}^{x\mu_{p}/\Delta} \frac{z}{\mu_{p}} \nu_{p}(z) \nu_{j}\left(\frac{z\mu_{j}}{\mu_{p}}\right) \prod_{\substack{w \in W \\ w \neq p}} \overline{\mathcal{V}}_{w}\left(\frac{z\mu_{w}}{\mu_{p}}\right) \prod_{\substack{\ell \in \mathcal{M}_{i} \setminus W \\ \ell \neq j}} \mathcal{V}_{\ell}\left(\frac{z\mu_{\ell}}{\mu_{p}}\right) dz.$$

Now consider the term *S* in the expression of  $C_1$  defined by the realization ( $\mathcal{M}_i, p, W$ ), and denote  $\mathcal{M}'_i = \mathcal{M}'_i \setminus \{j\}$  as well as  $W' = W \setminus \{p\}$ . Note that ( $\mathcal{M}_i, p, W'$ ) also defines a term *S'* in Equation 63. The rest of the argument is similar as for *A* and we obtain:

$$\left| \sum_{\substack{\mathcal{M}_i \\ |\mathcal{M}_i| \ge \gamma \\ j \in \mathcal{M}_i}} C_1(\mathcal{M}_i) \prod_{k \in \mathcal{M}_i} a_{i,k} \right| \le a_{i,j} \frac{\overline{\nu}}{\underline{\mu}}.$$
 (65)

We now tackle the terms in  $B(\mathcal{M}_i)$  in Equation 61. First, note that  $B(\mathcal{M}_i) = 0$  whenever  $|\mathcal{M}_i| = \gamma$ , since there are no set W of size  $\gamma - 1$  that does not contain either j or p. If we now consider the term S in the expression of B defined by the realization  $(\mathcal{M}_i, p, W)$  verifying  $|\mathcal{M}_i| \ge \gamma + 1$ , and denote  $\mathcal{M}'_i = \mathcal{M}_i \setminus \{j\}$  as well as  $W' = W \setminus \{j\}$ , it appears that  $(\mathcal{M}'_i, p, W')$  defines a term S'in the expression of  $\mathbb{P}\left[d_{i,j}^{\gamma-1} \le x\right]$ , written similarly as in Equation 63. Once again looking at the expression of S and S', we verify that multiplying the integrand of S by a factor  $a_{i,j} \frac{y}{\mu_p} v_j \left(\frac{y\mu_j}{\mu_p}\right)$ gives the integrand of S'. Following the same argument as for A, we obtain the following bound for terms in B, i.e.,

$$\left|\sum_{\substack{\mathcal{M}_i\\|\mathcal{M}_i|\geq\gamma\\j\in\mathcal{M}_i}} B(\mathcal{M}_i)\right| \le a_{i,j} \frac{\overline{\nu}}{\underline{\mu}}.$$
(66)

We do the same munipulation for  $C_2$  than for  $C_1$ , i.e., we rewrite k = p and make the change of variable  $z = y\mu_p/\mu_j$ , which gives,

$$C_{2}(\mathcal{M}_{i}) = \sum_{\substack{p \in \mathcal{M}_{i} \ W \subset \mathcal{M}_{i} \setminus \{j\} \\ p \neq j}} \sum_{\substack{W \subset \mathcal{M}_{i} \setminus \{j\} \\ |W| = \gamma - 1 \\ p \notin W}} \int_{0}^{x\mu_{p}/\Delta} \frac{z}{\mu_{p}} \nu_{p}(z) \nu_{j}\left(\frac{z\mu_{j}}{\mu_{p}}\right) \prod_{w \in W} \overline{\mathcal{V}}_{w}\left(\frac{z\mu_{w}}{\mu_{p}}\right) \prod_{\ell \in \mathcal{M}_{i} \setminus W \\ \ell \neq j, p} \mathcal{V}_{\ell}\left(\frac{z\mu_{\ell}}{\mu_{p}}\right) dz.$$

The same argument as for *B* holds, and we may only consider the cases where  $|\mathcal{M}_i| \ge \gamma + 1$ . For terms of  $C_2$  defined by realizations  $(\mathcal{M}_i, p, W)$  verifying this condition, we denote  $\mathcal{M}'_i = \mathcal{M}_i \setminus \{j\}$ . We again verify that  $(\mathcal{M}'_i, p, W)$  defines a term *S'* of  $\mathbb{P}\left[d_{i,j}^{\gamma-1} \le x\right]$  (see Equation 63 and replace  $\gamma - 1$  by  $\gamma$ ). The rest of the argument is similar as before and we obtain a bound on terms in  $C_2$ , i.e.,

$$\left| \sum_{\substack{\mathcal{M}_i \\ |\mathcal{M}_i| \ge \gamma \\ j \in \mathcal{M}_i}} C_2(\mathcal{M}_i) \prod_{k \in \mathcal{M}_i} a_{i,k} \right| \le a_{i,j} \frac{\overline{\nu}}{\underline{\mu}}.$$
(67)

Combining Equations 64, 66, 65 and 67 with 61 finally gives a bound on the derivatives of  $H_i$ , i.e.,

$$\left|\frac{\partial H_i}{\partial \mu_j}(x, \boldsymbol{\mu}_{-i})\right| \le 4a_{i,j}\frac{\overline{\nu}}{\underline{\mu}} \tag{68}$$

Step 5: Lipschitz Continuity of  $Z_i$ . We finally use the result of the previous paragraph to prove the Lipschitz continuity of agent *i*'s expected expenditure function  $Z_i(\boldsymbol{\mu}) = \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ d_i^{\boldsymbol{\gamma}} \mathbb{1} \{ \Delta v_i > \mu_i d_i^{\boldsymbol{\gamma}} \} \right]$ , where  $d_i^{\boldsymbol{\gamma}} = \boldsymbol{\gamma}^{\text{th}} - \max_{j:j \neq i} \left\{ \mathbb{1} \{ m_j = m_i \} \Delta v_j / \mu_j \right\}$ .

As values are drawn independently across agents, and since  $v_i$  is null outside [0, 1], we can write:

$$Z_{i}(\boldsymbol{\mu}) = \int_{0}^{\Delta/\mu_{i}} x\left(1 - \mathcal{V}_{i}\left(\frac{x\mu_{i}}{\Delta}\right)\right) dH_{i}(x, \boldsymbol{\mu}_{-i}).$$
(69)

Since the function  $\mu_i \mapsto x (1 - \mathcal{V}_i (x\mu_i/\Delta))$  is differentiable almost everywhere with derivative bounded by  $\Delta^2/\mu^2 \overline{\nu}$  under Assumption 1, Leibniz's integral rule implies that the expenditure function is differentiable with respect to  $\mu_i$ , i.e.,

$$\frac{\partial Z_i}{\partial \mu_i}(\boldsymbol{\mu}) = -\int_0^{\Delta/\mu_i} \frac{x^2}{\Delta} v_i\left(\frac{x\mu_i}{\Delta}\right) dH_i(x, \boldsymbol{\mu}_{-i})$$

Bounding  $x^2/\Delta v_i (x\mu_i/\Delta)$  by  $\Delta \overline{v}/\mu^2$  using Assumption 1, and integrating over the competing prices gives the following bound, i.e.,

$$\left|\frac{\partial Z_i}{\partial \mu_i}(\boldsymbol{\mu})\right| \le \frac{\Delta \overline{\nu}}{\underline{\mu}^2}.$$
(70)

Furthermore, an integration by part on Equation 69 gives the following alternate expression for  $Z_i(\mu)$ 

$$Z_{i}(\boldsymbol{\mu}) = \int_{0}^{\Delta/\mu_{i}} \left(\frac{x\mu_{i}}{\Delta}v_{i}\left(\frac{x\mu_{i}}{\Delta}\right) - \overline{\mathcal{V}}_{i}\left(\frac{x\mu_{i}}{\Delta}\right)\right) H_{i}(x,\boldsymbol{\mu}_{-i})dx.$$
(71)

Let  $j \neq i$ . Since  $H_i(x, \mu_{-i})$  is differentiable almost everywhere with respect to  $\mu_j$  with derivative bounded according to Equation 68, Leibniz's integral rule implies that:

$$\frac{\partial Z_i}{\partial \mu_j}(\boldsymbol{\mu}) = \int_0^{\Delta/\mu_i} \left(\frac{x\mu_i}{\Delta} v_i\left(\frac{x\mu_i}{\Delta}\right) - \overline{\mathcal{V}}_i\left(\frac{x\mu_i}{\Delta}\right)\right) \frac{\partial H_i}{\partial \mu_j}(x, \boldsymbol{\mu}_{-i}) dx$$

Using Equation 68, bounding  $\overline{\mathcal{V}}_i\left(\frac{x\mu_i}{\Delta}\right)$  by one on one hand, and bounding x by  $\Delta/\mu_i$  before integrating over  $\nu_i$  on the other hand, we get the following bound on the derivative of  $Z_i$ , i.e.,

$$\left|\frac{\partial Z_i}{\partial \mu_j}(\boldsymbol{\mu})\right| \le 8a_{i,j}\frac{\Delta \overline{\nu}}{\underline{\mu}^2}.$$
(72)

Together, Equations 70 and 72 finally imply for all  $\mu$ ,  $\mu'$  in  $\mathbb{R}^{N}_{+}$ :

$$\left|Z_{i}(\boldsymbol{\mu})-Z_{i}\left(\boldsymbol{\mu}'\right)\right| \leq \frac{\Delta \overline{\nu}}{\underline{\mu}^{2}}\left(|\mu_{i}-\mu_{i}'|+\sum_{j\neq i}8a_{i,j}|\mu_{j}-\mu_{j}'|\right).$$
(73)

We hence define  $\mathcal{L}_Z = \frac{8\Delta\overline{\nu}}{\mu^2}$  the Lipschitz constant of  $Z_i$  in the case of a single auction.

*Step 6: Upper bound on Expected Cost-per-Period.* We now return to the main proof and use the Lipschitz continuity of  $\Psi_i$  and  $Z_i$  to bound the expected cost per period.

Generalizing Equation 48 to a multi-agent setting, we rewrite the cost suffered by agent *i* at time  $t \leq \underline{T}$ .

$$c_{i,t} = v_{i,t} \left( \tau - \Delta \mathbb{1} \{ \Delta v_{i,t} > \mu_{i,t} d_{i,t}^{\gamma} \} \right) \\ = v_{i,t} \tau - \mu_{i,t} \rho_i - (\Delta v_{i,t} - \mu_{i,t} d_{i,t}^{\gamma})^+ + \mu_{i,t} \rho_i - \mu_{i,t} d_{i,t}^{\gamma} \mathbb{1} \{ \Delta v_{i,t} > \mu_{i,t} d_{i,t}^{\gamma} \}.$$

Recognizing an expense and a dual objective term, we take expectation and condition on  $\mu_t$  to get

$$\mathbb{E}_{\boldsymbol{v}}\left[c_{i,t}\right] = \mathbb{E}_{\boldsymbol{v}}\left[\Psi_{i}(\boldsymbol{\mu}_{t})\right] + \mathbb{E}_{\boldsymbol{v}}\left[\mu_{i,t}\left(\rho_{i} - Z_{i}(\boldsymbol{\mu}_{t})\right)\right].$$
(74)

On the one hand, we bound the first term in Equation 74 using the Lipschitz continuity of the dual objective function  $\Psi_i$ .

$$\Psi_i(\boldsymbol{\mu}_t) \le \Psi_i(\boldsymbol{\mu}^{\star}) + \mathcal{L}_{\Psi} \| \boldsymbol{\mu}_t - \boldsymbol{\mu}^{\star} \|_1$$
(75)

(-)

On the other hand, the Lipschitz continuity of the expected expenditure function  $Z_i$  allows to bound the second term in Equation 74.

$$\mu_{i,t}\left(\rho_{i}-Z_{i}(\boldsymbol{\mu}_{t})\right)=\mu_{i,t}\left(Z_{i}(\rho_{i}-Z_{i}(\boldsymbol{\mu}^{\star})+Z_{i}(\boldsymbol{\mu}^{\star})-Z_{i}(\boldsymbol{\mu}_{t})\right)\overset{(a)}{\leq}\overline{\mu}\mathcal{L}_{Z}\|\boldsymbol{\mu}_{t}-\boldsymbol{\mu}^{\star}\|_{1}$$
(76)

Inequality (*a*) makes use of the definition of  $\mu^{\star}$ .

Combining Equations 74, 75 and 76, we get

$$\mathbb{E}_{\boldsymbol{v}}\left[c_{i,t}\right] \leq \Psi_{i}(\boldsymbol{\mu}^{\star}) + \left(\mathcal{L}_{\Psi} + \overline{\boldsymbol{\mu}}\mathcal{L}_{Z}\right) \mathbb{E}_{\boldsymbol{v}}\left[\|\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}\|_{1}\right],$$

$$\stackrel{(a)}{\leq} \Psi_{i}(\boldsymbol{\mu}^{\star}) + \left(\mathcal{L}_{\Psi} + \overline{\boldsymbol{\mu}}\mathcal{L}_{Z}\right) \mathbb{E}_{\boldsymbol{v}}\left[\sqrt{N}\|\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}\|_{2}\right],$$

$$\stackrel{(b)}{\leq} \Psi_{i}(\boldsymbol{\mu}^{\star}) + \left(\mathcal{L}_{\Psi} + \overline{\boldsymbol{\mu}}\mathcal{L}_{Z}\right) \sqrt{N} \left(\mathbb{E}_{\boldsymbol{v}}\left[\|\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}\|_{2}^{2}\right]\right)^{1/2}.$$
(77)

Inequality (a) uses that  $\sum_{i=1}^{n} |y_i| \le \left(n \sum_{i=1}^{n} y_i^2\right)^{1/2}$ ; inequality (b) follows from Jensen's inequality.

Conclusion. Combining Equations 56 and 76 with Lemma C.2 finally gives

$$\begin{split} \frac{1}{T} \mathcal{C}_{i}^{A} - \Psi_{i}(\boldsymbol{\mu}^{\star}) &\leq \frac{1}{T} \mathbb{E}_{\boldsymbol{\sigma}} \left[ \sum_{t=1}^{\underline{T}} c_{i,t} \right] + \frac{\tau}{T} \left( \frac{\Delta}{\underline{\rho}\underline{\mu}} + \frac{\overline{\mu}}{\underline{\rho}\underline{\epsilon}} \right) - \Psi_{i}(\boldsymbol{\mu}^{\star}), \\ &\leq \left( \mathcal{L}_{\Psi} + \overline{\mu}\mathcal{L}_{Z} \right) \frac{\sqrt{N}}{T} \mathbb{E}_{\boldsymbol{\sigma}} \left[ \sum_{t=1}^{\underline{T}} s_{t}^{1/2} \right] + \frac{\tau}{T} \left( \frac{\Delta}{\underline{\rho}\underline{\mu}} + \frac{\overline{\mu}}{\underline{\rho}\underline{\epsilon}} \right), \\ &\stackrel{(a)}{\leq} \left( \mathcal{L}_{\Psi} + \overline{\mu}\mathcal{L}_{Z} \right) N \left( \frac{\Delta\overline{\epsilon}}{\sqrt{2\lambda}\underline{\mu}\underline{\epsilon}^{1/2}} + \overline{\mu}\frac{\overline{\epsilon}^{1/2}}{\underline{\epsilon}^{1/2}T} \sum_{t=1}^{T} (1 - 2\lambda\underline{\epsilon})^{(t-1)/2} \right) + \frac{\tau}{T} \left( \frac{\Delta}{\underline{\rho}\underline{\mu}} + \frac{\overline{\mu}}{\underline{\rho}\underline{\epsilon}} \right), \\ &\stackrel{(b)}{\leq} \left( \mathcal{L}_{\Psi} + \overline{\mu}\mathcal{L}_{Z} \right) N \left( \frac{\Delta\overline{\epsilon}}{\sqrt{2\lambda}\underline{\mu}\underline{\epsilon}^{1/2}} + \frac{\overline{\mu}\overline{\epsilon}^{1/2}}{\lambda\underline{\epsilon}^{3/2}T} \right) + \frac{\tau}{T} \left( \frac{\Delta}{\underline{\rho}\underline{\mu}} + \frac{\overline{\mu}}{\underline{\rho}\underline{\epsilon}} \right). \end{split}$$

Inequality (*a*) follows from Lemma C.2 with  $\sqrt{x+y} \le \sqrt{x} + \sqrt{y}$ . Inequality (*b*) holds by bounding the geometric sum by the series' limit and noting that  $\frac{1}{1-(1-\lambda \underline{\epsilon})^{1/2}} \le \frac{2}{\lambda \underline{\epsilon}}$  since  $1 - (1-x)^{1/2} \ge x/2$  for  $x \in [0, 1]$ .

Hence there exist a constant  $C_2$  in  $\mathbb{R}_+$  such that

$$\frac{1}{T}C_i^A - \Psi_i(\boldsymbol{\mu^{\star}}) \le C_2\left(\frac{1}{T} + N\left(\frac{\overline{\epsilon}^{1/2}}{\underline{\epsilon}^{3/2}T} + \frac{\overline{\epsilon}}{\underline{\epsilon}^{1/2}}\right)\right).$$

## C.4 Proof of Theorem 3.3

Consider a unilateral deviation of agent  $i \in N$  to some strategy  $\beta \in \mathcal{B}^{CI}$ . For the purpose of the analysis, we allow this strategy to use *complete information*, i.e., it can access the budgets  $k_s$ , auctions  $m_s$ , valuations  $v_s$ , bids  $b_s$ , and expenditures  $z_s$  of all agents at all previous time steps s < t, and in addition, it has knowledge of the valuation distributions  $\mathcal{V}$ . We denote the set of strategies that use complete information as  $\mathcal{B}^{CI} \supset \mathcal{B}$  and suppose that  $\beta \in \mathcal{B}^{CI}$ .

Since agent *i* does not follow strategy *A*,  $\mu_{i,t}$  is a priory not defined. For convenience, we take the convention  $\mu_{i,t} = \mu_i^*$  for all  $t \in \mathbb{N}$ .

Step 1: Lower Bound on the Total Expected Cost under Strategy  $\beta$ . We lower-bound the total expected cost suffered by agent *i* for following strategy  $\beta$  by considering it always gets a null cost after time  $\Im$ .

$$C_{i}^{\beta,A_{-i}} \geq \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{t=1}^{\underline{\Upsilon}} c_{i,t}^{\beta,A_{-i}} \right]$$

$$\stackrel{(a)}{\geq} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{t=1}^{\underline{\Upsilon}} c_{i,t}^{\beta,A_{-i}} \right] + \mu_{i}^{\star} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{t=1}^{T} z_{i,t}^{\beta,A_{-i}} - \rho_{i} \right]$$

$$\stackrel{(b)}{\geq} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{t=1}^{\underline{\Upsilon}} c_{i,t}^{\beta,A_{-i}} + \mu_{i}^{\star} (z_{i,t}^{\beta,A_{-i}} - \rho_{i}) \right] + \frac{\Delta \overline{\mu}}{\underline{\mu}} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ T - \underline{\Upsilon} \right]$$

$$\stackrel{(c)}{\geq} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{t=1}^{\underline{\Upsilon}} c_{i,t}^{\beta,A_{-i}} + \mu_{i}^{\star} (z_{i,t}^{\beta,A_{-i}} - \rho_{i}) \right] + \frac{\Delta \overline{\mu}}{\underline{\mu}} \left( \frac{\Delta}{\underline{\rho\mu}} + \frac{\overline{\mu}}{\underline{\rho}\underline{\epsilon}} \right)$$

$$(78)$$

Inequality (a) uses that strategy  $\beta$  respects the budget constraint on average; inequality (b) holds since  $z_{i,t}^{\beta,A_{-i}} \leq d_{i,t}^{\gamma} \leq \Delta/\mu$  and inequality (c) uses Lemma C.1.

Step 2: Lower Bound on the Lagrangian Expected Cost. We bound the Lagrangian cost of agent i for the t<sup>th</sup> period.

$$c_{i,t}^{\beta,A_{-i}} + \mu_{i}^{\star}(z_{i,t}^{\beta,A_{-i}} - \rho_{i}) = v_{i,t}\tau - \mathbb{1}\left\{b_{i,t}^{\beta} \ge d_{i,t}^{\gamma}\right\}(\Delta v_{i,t} - \mu_{i}^{\star}d_{i,t}^{\gamma}) - \mu_{i}^{\star}\rho_{i} \\ \ge v_{i,t}\tau - (\Delta v_{i,t} - \mu_{i}^{\star}d_{i,t}^{\gamma})^{+} - \mu_{i}^{\star}\rho_{i}$$
(79)

Taking expectation on Equation 79 and using the Lipschitz continuity of the dual objective function gives

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{m}}\left[c_{i,t}^{\beta,A_{-i}} + \mu_{i}^{\star}(\boldsymbol{z}_{i,t}^{\beta,A_{-i}} - \rho_{i})\right] \stackrel{(a)}{\geq} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}}\left[\Psi_{i}(\boldsymbol{\mu}^{\star}) - \frac{\Delta}{\underline{\mu}}\sum_{j\neq i}a_{i,j}|\mu_{j,t} - \mu_{j}^{\star}|\right],$$

$$\stackrel{(c)}{\geq}\Psi_{i}(\boldsymbol{\mu}^{\star}) - \frac{\Delta}{\underline{\mu}}\|\boldsymbol{a}_{i}\|_{2}\mathbb{E}_{\boldsymbol{v},\boldsymbol{m}}\left[\|\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}\|_{2}\right],$$

$$\stackrel{(d)}{\geq}\Psi_{i}(\boldsymbol{\mu}^{\star}) - \frac{\Delta}{\underline{\mu}}\|\boldsymbol{a}_{i}\|_{2}\boldsymbol{s}_{t}^{1/2}.$$
(80)

Inequality (*a*) follows from Equation 79 by definition of  $\mu_{i,t} = \mu_i^*$  for all period *t*; inequality (*b*) uses the Lipschitz continuity of the dual function (see Equation 59); inequality (*c*) follows from the Cauchy-Schwartz inequality and inequality (*d*) holds since  $\mathbb{E}\left[\|\boldsymbol{\mu}_t - \boldsymbol{\mu}^*\|_2\right] \leq \left(\mathbb{E}\left[\|\boldsymbol{\mu}_t - \boldsymbol{\mu}^*\|_2^2\right]\right)^{1/2}$  by Jensen's inequality.

*Step 3: Extension of the Proof of Theorem 3.2 to a Parallel Auction Setting.* Note that Equation 74 still hold for the cost-per-period of strategy *A* in the parallel auctions setting.

Similarly as before, we use the Lipschitz continuity of the dual and expenditure functions to bound the different terms in Equation 74. Using the more careful result of Equation 59 gives however

$$\Psi_{i}(\boldsymbol{\mu}_{t}) \stackrel{(a)}{\leq} \Psi_{i}(\boldsymbol{\mu}^{\star}) + \frac{\Delta}{\underline{\mu}} \left( |\mu_{i,t} - \mu_{i}^{\star}| + \sum_{j \neq i} a_{i,j} |\mu_{j,t} - \mu_{j}^{\star}| \right),$$

$$\stackrel{(b)}{\leq} \Psi_{i}(\boldsymbol{\mu}^{\star}) + \frac{\Delta}{\underline{\mu}} ||\boldsymbol{a}_{i}||_{2} ||\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}||_{2}.$$
(81)

Inequality (*a*) follows from Equation 59 and inequality (*b*) from Cauchy-Schwartz inequality combined with  $\mu_{i,t} = \mu_i^{\star}$ .

Similarly using Equation 73 gives, i.e.,

$$\mu_{i,t} \left( Z_i(\boldsymbol{\mu}_t) - \rho_i \right) = \mu_{i,t} \left( Z_i(\boldsymbol{\mu}_t) - Z_i(\boldsymbol{\mu}^{\star}) + Z_i(\boldsymbol{\mu}^{\star}) - \rho_i \right)$$

$$\stackrel{(a)}{\leq} \frac{\Delta \overline{\mu \nu}}{\underline{\mu}^2} \left( |\mu_{i,t} - \mu_i^{\star}| + \sum_{j \neq i} 8a_{i,j} |\mu_{j,t} - \mu_j^{\star}| \right),$$

$$\stackrel{(b)}{\leq} \frac{8\Delta \overline{\mu \nu}}{\underline{\mu}^2} ||\boldsymbol{a}_i||_2 ||\boldsymbol{\mu}_t - \boldsymbol{\mu}^{\star}||_2.$$
(82)

Inequality (*a*) makes use of the definition of  $\mu^*$  as well as Equation 73, inequality (*b*) follows from Cauchy-Schwartz inequality and  $\mu_{i,t} = \mu_i^*$ .

Combining Equations 74, 81 and 82, we get

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{m}}\left[c_{i,t}\right] \leq \Psi_{i}(\boldsymbol{\mu}^{\star}) + \frac{\Delta}{\underline{\mu}}\left(1 + \frac{8\overline{\mu}\overline{\nu}}{\underline{\mu}}\right) \|\boldsymbol{a}_{i}\|_{2}\mathbb{E}_{\boldsymbol{v},\boldsymbol{m}}\left[\|\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}\|_{2}\right],$$

$$\overset{(a)}{\leq} \Psi_{i}(\boldsymbol{\mu}^{\star}) + \frac{\Delta}{\underline{\mu}}\left(1 + \frac{8\overline{\mu}\overline{\nu}}{\underline{\mu}}\right) \|\boldsymbol{a}_{i}\|_{2}\left(\mathbb{E}_{\boldsymbol{v},\boldsymbol{m}}\left[\|\boldsymbol{\mu}_{t} - \boldsymbol{\mu}^{\star}\|_{2}^{2}\right]\right)^{1/2}.$$
(83)

Inequality (a) follows from Jensen's inequality.

We conclude in the same manner that

$$\frac{1}{T}C_{i}^{A} - \Psi_{i}(\boldsymbol{\mu}^{\star}) \leq \frac{1}{T}\mathbb{E}_{\boldsymbol{\nu}}\left[\sum_{t=1}^{\mathcal{T}}c_{i,t}\right] + \frac{\tau}{T}\left(\frac{\Delta}{\underline{\rho\mu}} + \frac{\overline{\mu}}{\underline{\rho\epsilon}}\right) - \Psi_{i}(\boldsymbol{\mu}^{\star}),$$

$$\leq \frac{\Delta}{\underline{\mu}}\left(1 + \frac{8\overline{\mu\nu}}{\underline{\mu}}\right) \|\boldsymbol{a}_{i}\|_{2}\frac{1}{T}\mathbb{E}_{\boldsymbol{\nu},\boldsymbol{m}}\left[\sum_{t=1}^{\mathcal{T}}s_{t}^{1/2}\right] + \frac{\tau}{T}\left(\frac{\Delta}{\underline{\rho\mu}} + \frac{\overline{\mu}}{\underline{\rho\epsilon}}\right),$$

$$\overset{(a)}{\leq} \frac{\Delta}{\underline{\mu}}\left(1 + \frac{8\overline{\mu\nu}}{\underline{\mu}}\right)\sqrt{N}\|\boldsymbol{a}_{i}\|_{2}\left(\frac{\Delta\overline{\epsilon}}{\sqrt{2\lambda}\underline{\mu}\underline{\epsilon}^{1/2}} + \frac{\overline{\mu}\overline{\epsilon}^{1/2}}{\lambda\underline{\epsilon}^{3/2}T}\right)$$

$$+ \frac{\tau}{T}\left(\frac{\Delta}{\underline{\rho\mu}} + \frac{\overline{\mu}}{\underline{\rho\underline{\epsilon}}}\right).$$
(84)

Inequality (*a*) follows from the same steps as in the proof of Theorem 3.2. In particular, it uses Lemma C.2, which still holds true since we can bound all  $a_{i,j}$  individually by one.

Conclusion.

On one hand, combining Equations 78 and 80 gives

$$C_{i}^{\beta,A_{-i}} \geq \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{t=1}^{\underline{\mathcal{T}}} c_{i,t}^{\beta,A_{-i}} + \mu_{i}^{\star} (z_{i,t}^{\beta,A_{-i}} - \rho_{i}) \right] + \frac{\Delta \overline{\mu}}{\underline{\mu}} \left( \frac{\Delta}{\underline{\rho\mu}} + \frac{\overline{\mu}}{\underline{\rho\underline{e}}} \right),$$

$$\geq T \Psi_{i}(\boldsymbol{\mu}^{\star}) - \frac{\Delta}{\underline{\mu}} \|\boldsymbol{a}_{i}\|_{2} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{t=1}^{\underline{\mathcal{T}}} s_{t}^{1/2} \right] + \frac{\Delta \overline{\mu}}{\underline{\mu}} \left( \frac{\Delta}{\underline{\rho\mu}} + \frac{\overline{\mu}}{\underline{\rho\underline{e}}} \right).$$
(85)

We bound the second term using Lemma C.3.

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{m}}\left[\sum_{t=1}^{\underline{\mathcal{T}}} s_t^{1/2}\right] \stackrel{(a)}{\leq} \overline{\mu} \left(N\frac{\overline{\epsilon}}{\underline{\epsilon}}\right)^{1/2} \sum_{t=1}^{T} (1-\lambda\underline{\epsilon})^{(t-1)/2} + \frac{\Delta N^{1/2}}{\sqrt{\lambda}\underline{\mu}} \frac{\overline{\epsilon}}{\underline{\epsilon}^{1/2}} T + \frac{\Delta \|\boldsymbol{a}_i\|_2}{\lambda\underline{\mu}} \frac{\overline{\epsilon}}{\underline{\epsilon}} T \\ \stackrel{(b)}{\leq} \frac{2\overline{\mu}N^{1/2}}{\lambda} \frac{\overline{\epsilon}^{1/2}}{\underline{\epsilon}^{3/2}} + \frac{\Delta N^{1/2}}{\sqrt{\lambda}\underline{\mu}} \frac{\overline{\epsilon}}{\underline{\epsilon}^{1/2}} T + \frac{\Delta \|\boldsymbol{a}_i\|_2}{\lambda\underline{\mu}} \frac{\overline{\epsilon}}{\underline{\epsilon}} T$$

$$(86)$$

Inequality (*a*) holds since  $\sqrt{x + y + z} \le \sqrt{x} + \sqrt{y} + \sqrt{z}$  for  $x, y, z \ge 0$  and  $\underline{\Upsilon} \le T$ ; inequality (*b*) follows from bounding the geometric sum by the series' limit and using that  $\frac{1}{1 - (1 - \lambda \underline{\epsilon})^{1/2}} \le \frac{2}{\lambda \underline{\epsilon}}$  since  $1 - (1 - x)^{1/2} \ge x/2$  for  $x \in [0, 1]$ .

Equations 85 and 86 finally lead to

$$\Psi_{i}(\boldsymbol{\mu}^{\star}) - \frac{1}{T}C_{i}^{\beta,A_{-i}} \leq \frac{2\Delta\overline{\mu}\|a_{i}\|_{2}N^{1/2}}{\lambda\underline{\mu}} \frac{\overline{\epsilon}^{1/2}}{\underline{\epsilon}^{3/2}T} + \frac{\Delta^{2}\|a_{i}\|_{2}N^{1/2}}{\sqrt{\lambda}\underline{\mu}^{2}} \frac{\overline{\epsilon}}{\underline{\epsilon}^{1/2}} + \frac{\Delta^{2}\|a_{i}\|_{2}^{2}}{\lambda\underline{\mu}^{2}} \frac{\overline{\epsilon}}{\underline{\epsilon}} + \frac{\Delta\overline{\mu}}{\underline{\mu}T} \left(\frac{\Delta}{\underline{\rho}\underline{\mu}} + \frac{\overline{\mu}}{\underline{\rho}\underline{\epsilon}}\right).$$

$$(87)$$

Combining Equations 84 and 87 finally proves the theorem, and we have for some constant C in  $\mathbb{R}^+$ 

$$\frac{1}{T}\left(C_{i}^{A}-C_{i}^{\beta,A_{-i}}\right) \leq C\left(\frac{1+\epsilon_{i}}{\epsilon_{i}T}+\|\boldsymbol{a}_{i}\|_{2}^{2}\frac{\overline{\epsilon}}{\underline{\epsilon}}+\sqrt{N}\|\boldsymbol{a}_{i}\|_{2}\left(\frac{\overline{\epsilon}^{1/2}}{\underline{\epsilon}^{3/2}T}+\frac{\overline{\epsilon}}{\underline{\epsilon}^{1/2}}\right)\right).$$

## D PROOFS OF THE SECONDARY RESULTS IN SECTION 3

In this section, we gather the proofs of secondary results discussed in the main text of the article, mainly discussions on the relaxation of certain assumptions.

### D.1 Sufficient Conditions for Assumption 2

In this section, we provide mild differentiability and continuity conditions on the valuation and competing bid distributions that guarantee that the technical Assumption 2 is satisfied. This verifies that the conditions of Theorem 3.1 are not restrictive.

ASSUMPTION 18. The following conditions hold:

18.1 The valuation density  $v_i$  is differentiable with bounded derivative  $|v'_i(v)| \leq \overline{v}'$  for all  $v \in [0, 1]$ ; 18.2 The distribution of competing bids  $\mathcal{D}_i^{\gamma}$  is absolutely continuous with bounded density  $h_i : [0, \Delta/\mu_i] \mapsto [h, \overline{h}] \subset \mathbb{R}_+^*$ .

In the following, we show that Assumption 2 is implied by Assumptions 1 and 18 in three different steps. We first show that the dual function  $\Psi : \mu \mapsto \mathbb{E}_{v,d^{\gamma}} [v\tau - \mu\rho - (\Delta v - \mu d^{\gamma})^+]$  is differentiable, then give upper and lower bounds on the derivative of the expenditure function  $Z : \mu \mapsto \mathbb{E}_{v,d^{\gamma}} [d^{\gamma} \mathbb{I} \{\Delta v > \mu d^{\gamma}\}]$ , and finally bounds the second order derivative of *Z*.

Differentiability of  $\Psi$ . The function  $\mu \mapsto (\Delta v - \mu d^{\gamma})^+$  is differentiable with derivative  $\mu \mapsto d^{\gamma} \mathbb{1}\{\Delta v > \mu d^{\gamma}\}$ , except in the set  $\{(v, d^{\gamma}) : \Delta v = \mu d^{\gamma}\}$  of measure zero since valuations and competing prices are both absolutely continuous with respective support in  $[0, \overline{\nu}]$  and  $[0, \Delta/\mu]$ . As the derivative is bounded by  $\Delta/\mu$ , which is integrable, Leibniz's integral rule ensures that  $\overline{\Psi}$  is differentiable and we get for all  $\mu \geq 0$ :

$$\Psi'(\mu) = d^{\gamma} \mathbb{1}\{\Delta v > \mu d^{\gamma}\} - \rho = Z(\mu) - \rho.$$
(88)

Bounds on the Derivative of Z. Let  $\mathcal{V}$  denote the cumulative distribution function of valuations associated with the density *v*. Since *v* is null outside of [0, 1], we can write the expenditure function as follows:

$$Z(\mu) = \int_0^{\Delta/\mu} x \left( 1 - \mathcal{V}\left(\frac{\mu x}{\Delta}\right) \right) h(x) dx.$$
(89)

Since  $\mu \mapsto x(1 - \mathcal{V}(\mu x / \Delta))$  is differentiable almost everywhere with derivative bounded by  $\frac{\Delta^2}{\underline{\mu}^2} \overline{\nu}$ , Leibniz's integral rule applied to Equation 89 gives:

$$Z'(\mu) = -\int_0^{\Delta/\mu} \frac{x^2}{\Delta} v\left(\frac{\mu x}{\Delta}\right) h(x) dx.$$
(90)

Using the bounded densities of Assumption 18, we finally get some bounds on the derivative, i.e.,

$$-\frac{\Delta}{\underline{\mu}^{2}}\overline{\nu} \leq Z^{'}(\mu) \leq -\underline{\nu}h \int_{0}^{\Delta/\overline{\mu}} \frac{x^{2}}{\Delta} dx = -\frac{\underline{\nu}h\Delta^{3}}{3\overline{\mu}^{3}}.$$
(91)

Bound on the Second Order Derivative of Z. Since  $\mu \mapsto \frac{x^2}{\Delta} v\left(\frac{\mu x}{\Delta}\right) h(x)$  is differentiable almost everywhere with derivative bounded by  $\frac{\Delta}{\mu^3} \overline{v}'$ , using Leibniz integral rule on Equation 90 gives the following, i.e.,

$$Z^{''}(\mu) = \frac{\Delta^2}{\mu^4} v(1) h\left(\frac{\Delta}{\mu}\right) - \int_0^{\Delta/\mu} \frac{x^3}{\Delta^2} v^{'}\left(\frac{\mu x}{\Delta}\right) h(x) dx.$$

Using Assumptions 1 and 18, we finally get the desired bound.

$$\left| Z^{''}(\mu) \right| \leq \frac{\Delta^2}{\underline{\mu}^4} \overline{\nu} \overline{h} + \frac{\Delta}{\underline{\mu}^3} \overline{\nu}'$$

### D.2 Uniqueness of Stationary Multiplier Profile

We prove in the following that Assumption 5 ensures the uniqueness of a stationary multiplier profile.

First, note that all vectors  $\boldsymbol{\mu} \in \boldsymbol{U}$  verifying  $\mu_i \geq \overline{\mu_i}$  for some  $i \in \mathcal{N}$  are such that  $Z_i(\boldsymbol{\mu}) \leq \Delta/\overline{\mu_i} < \rho_i$ under Assumption 3. Hence, no such  $\boldsymbol{\mu}$  can be a stationary multiplier profile according to the definition of  $\boldsymbol{\mu}^*$ .

Now suppose there are two different vectors  $\mu$  and  $\mu'$  in U satisfy the definition of a stationary multiplier. We have:

$$0 \stackrel{(a)}{>} \sum_{i=1}^{N} (\mu_i - \mu'_i) \left( Z_i(\boldsymbol{\mu}) - \rho_i + \rho_i - Z_i(\boldsymbol{\mu'}) \right) \stackrel{(b)}{=} 0.$$

Inequality (*a*) follows from the strong monotonicity of *Z*; (*b*) uses that both  $\mu$  and  $\mu'$  are stationary multipliers. This is a contradiction.

## E PROOF OF THE THEOREMS IN SECTION 4

This section contains the proofs of the main Theorems in Section 4. It is organized as follows: first, we prove in Section E.1 a bound on the expected mean squared error  $s_t$ . We then include the proofs of Theorems 4.1, 4.2 and 4.3 in the respective Sections E.2, E.3 and E.4 respectively. The proofs of the secondary results discussed in Section 4 are deferred to Section F.

### E.1 Helpful Lemma

In the following, we denote by  $\mu_{i,t}^{\pi} = P_{[\underline{\mu}_i, \overline{\mu}_i]}(\mu_{i,t})$  the projected multiplier used to formulate bids in (*G-b*). Note that, by definition of the hitting time  $\mathcal{T}$  in Equation 22, it holds for all  $1 \le t \le \underline{\mathcal{T}}$  that  $\mu_{i,t}^{\pi} = \mu_{i,t}$ .

For any fixed  $t \leq \underline{\Upsilon}$ , we consider the expected mean squared error  $s_t = \mathbb{E}_{\upsilon} \left[ \| \boldsymbol{\mu}_t^{\pi} - \boldsymbol{\mu}^{\star 0} \|_2^2 \right] = \sum_{i=1}^N s_{i,t}$ , where  $s_{i,t} = \mathbb{E}_{\upsilon} \left[ (\boldsymbol{\mu}_{i,t}^{\pi} - \boldsymbol{\mu}_i^{\star 0})^2 \right]$  is the individual expected error of agent *i*. Lemma E.1 gives bounds on the expected mean squared error in different scenarios.

LEMMA E.1 (UPPER BOUND ON THE EXPECTED MEAN SQUARED ERROR).

When all agents follow G strategies, the expected mean squared error  $s_t$  verifies for all  $1 \le t \le \underline{T}$ :

$$s_t \le N\overline{\mu}^2 (1-2\lambda\epsilon)^{t-1} + \frac{N\Delta^2\epsilon}{2\lambda\mu^2}$$

Taking N = 1 gives a bound for the individual error of an agent in the stationary competition setting.

**PROOF.** For any agent  $i \in N$  and time step  $t < \underline{T}$ , the multiplier update rule gives:

$$\begin{pmatrix} \mu_{i,t+1}^{\pi} - \mu_{i}^{\star 0} \end{pmatrix}^{2} \stackrel{(a)}{=} \begin{pmatrix} \mu_{i,t} - \mu_{i}^{\star 0} + \epsilon_{i}(z_{i,t} - g_{i,t}) \end{pmatrix}^{2} \\ \stackrel{(b)}{=} \begin{pmatrix} \mu_{i,t}^{\pi} - \mu_{i}^{\star 0} \end{pmatrix}^{2} + 2\epsilon(\mu_{i,t}^{\pi} - \mu_{i}^{\star 0})(z_{i,t} - g_{i,t}) + \epsilon_{i}^{2}(z_{i,t} - g_{i,t})^{2},$$

where equality (*a*) uses that  $\mu_{i,t+1}^{\pi} = \mu_{i,t+1}$  since  $t < \underline{T}$ , and equality (*b*) uses Assumption 17.2 as well as  $\mu_{i,t}^{\pi} = \mu_{i,t}$ .

Taking the expectation on previous valuations of all agents (resp. previous valuations and competing prices), gives the following, i.e.,

$$s_{i,t+1} \leq s_{i,t} + 2\epsilon \mathbb{E}_{\boldsymbol{v}} \left[ (\mu_{i,t}^{\pi} - \mu_{i}^{\star 0}) (z_{i,t} - g_{i,t}) \right] + \epsilon^{2} \mathbb{E}_{\boldsymbol{v}} \left[ (z_{i,t} - g_{i,t})^{2} \right],$$

$$\stackrel{(a)}{=} s_{i,t} + 2\epsilon \mathbb{E}_{\boldsymbol{v}} \left[ (\mu_{i,t}^{\pi} - \mu_{i}^{\star 0}) \mathbb{E}_{\boldsymbol{v}} [z_{i,t} - g_{i,t} | \boldsymbol{\mu}_{t}^{\pi}] \right] + \epsilon^{2} \mathbb{E}_{\boldsymbol{v}} \left[ (z_{i,t} - g_{i,t})^{2} \right],$$

$$\stackrel{(b)}{=} s_{i,t} + 2\epsilon \mathbb{E}_{\boldsymbol{v}} \left[ (\mu_{i,t}^{\pi} - \mu_{i}^{\star 0}) (L_{i}(\boldsymbol{\mu}_{t}^{\pi}) - L_{i}(\boldsymbol{\mu}^{\star 0})) \right] + \epsilon^{2} \mathbb{E}_{\boldsymbol{v}} \left[ (z_{i,t} - g_{i,t})^{2} \right].$$
(92)

Equality (*a*) holds by linearity of expectation; equality (*b*) by definition of  $L_i(\mu_t)$  and  $\mu^{\star 0}$  (resp.  $L_i(\mu_{i,t})$  and the definition of  $\mu_i^{\star 0}$  in Assumption 14.1).

We bound the sum over the whole population of the middle term in Equation 92.

$$\mathbb{E}_{\boldsymbol{\upsilon}}\left[\sum_{i=1}^{N}(\mu_{i,t}^{\pi}-\mu_{i}^{\star 0})(L_{i}(\boldsymbol{\mu}_{t}^{\pi})-L_{i}(\boldsymbol{\mu}^{\star 0}))\right] \leq -\lambda s_{t}$$

$$\tag{93}$$

The bound follows from Assumption 16 (resp. the  $\lambda$ -strong monotonicity of the expected karma loss function  $L_i$  in Assumption 13).

On the other hand, as all  $z_{i,t}$  and  $g_{i,t}$  are positive and bounded by  $\Delta/\mu$  under Assumption 14.2, we similarly bound the right-most term of Equation 92:

$$\sum_{i=1}^{N} \mathbb{E}_{\boldsymbol{v}} \left[ \left| \boldsymbol{z}_{i,t} - \boldsymbol{g}_{i,t} \right|^2 \right] \le \frac{N \Delta^2}{\underline{\mu}^2}.$$
(94)

Combining Equations 92, 93 and 94 gives the following recursion, i.e.,

$$s_{t+1} \leq (1-2\lambda\epsilon)s_t + \frac{N\Delta^2\epsilon^2}{\underline{\mu}^2}.$$

Since  $s_1 \le N\overline{\mu}^2$  under Assumption 14.1 and  $2\lambda \epsilon \le 1$  under Assumption 14.4, [6, Lemma C.4] finally yields the desired result.

$$s_t \le N\overline{\mu}^2 (1 - 2\lambda\epsilon)^{t-1} + \frac{N\Delta^2\epsilon}{2\lambda\underline{\mu}^2}$$

## E.2 Proof of Theorem 4.1

In this section, we again take the perspective of a single agent  $i \in N$ , but drop the subscript *i* for simplicity of notations.

Step 1: Lower Bound on Performances in Hindsight. Let  $\Psi_d^{\rho}$  denote the expected Lagrangian dual function associated with the lower bound on the total cost in hindsight  $\underline{C}^H$ , defined for all  $\mu \ge 0$  as  $\Psi_d^{\rho}(\mu) = \mathbb{E}_{v,d} \left[ v\tau - \mu(\rho + \frac{\gamma}{N}d^{\gamma}) - (\Delta v - \mu d^{\gamma})^+ \right]$ . Let  $\mu_d^{\star \rho}$  denote its maximizer.

We lower-bound the lowest cost in hindsight using  $\Psi_d^{\rho}$  in the following manner:

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ \boldsymbol{C}^{H}(\boldsymbol{v},\boldsymbol{d}) \right] \stackrel{(a)}{\geq} \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ \underline{\boldsymbol{C}}^{H}(\boldsymbol{v},\boldsymbol{d}) \right],$$

$$\stackrel{(b)}{\geq} \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ \sup_{\mu \geq 0} \sum_{t=1}^{T} \boldsymbol{v}_{t} \tau - \mu \left( \rho + \frac{\gamma}{N} \boldsymbol{d}_{t}^{\gamma} \right) - \left( \Delta \boldsymbol{v}_{t} - \mu \boldsymbol{d}_{t}^{\gamma} \right)^{+} \right],$$

$$\stackrel{(c)}{\geq} \sup_{\mu \geq 0} \sum_{t=1}^{T} \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ \boldsymbol{v}_{t} \tau - \mu \left( \rho + \frac{\gamma}{N} \boldsymbol{d}_{t}^{\gamma} \right) - \left( \Delta \boldsymbol{v}_{t} - \mu \boldsymbol{d}_{t}^{\gamma} \right)^{+} \right],$$

$$= T \Psi_{\boldsymbol{d}}^{\rho}(\mu_{\boldsymbol{d}}^{\star \rho}).$$
(95)

Inequality (*a*) follows from Equation 18 and inequality (*b*) from Equation 19a. Inequality (*c*) holds since  $\mathbb{E}_X[\sup_u f_u(X)] \ge \sup_u \mathbb{E}_X[f_u(X)]$  and by linearity of expectation.

*Step 2: Upper Bound on the Total Cost and Cost-per-Auction of strategy G.* We upper-bound the total cost of strategy *G* by considering that the worst always happens after the hitting time.

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[C^{G}(\boldsymbol{v},\boldsymbol{d})\right] \leq \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[\sum_{t=1}^{\mathcal{T}} c_{t}\right] + \tau \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[T - \mathcal{T}\right]$$
(96)

For the rest of the proof, we only consider  $t \leq \mathcal{T}$ , where the remaining budget is greater than the maximum expense. We first rewrite the cost-per-period in the following manner:

$$c_{t} = v_{t} \left( \tau - \Delta \mathbb{1} \{ \Delta v_{t} > \mu_{t}^{\pi} d_{t}^{Y} \} \right),$$
  
$$= v_{t} \tau - \mu_{t}^{\pi} g_{t} - \left( \Delta v_{t} - \mu_{t}^{\pi} d_{t}^{Y} \right)^{+} + \mu_{t}^{\pi} g_{t} - \mu_{t}^{\pi} d_{t}^{Y} \mathbb{1} \{ \Delta v_{t} > \mu_{t}^{\pi} d_{t}^{Y} \}.$$
(97)

Similarly as in the proof of Theorem 3.1, we define  $C(\mu) = \mathbb{E}_{v,d}[c|\mu] = \Psi^0(\mu) - \mu L(\mu)$  the expected cost-per-period. Note that Assumption 13 ensures it is twice differentiable with derivatives  $C'(\mu) = -\mu (L'(\mu) + G'(\mu)) = -\mu Z'(\mu)$  and  $C''(\mu) = -(Z'(\mu) + \mu Z''(\mu))$ . Note moreover that  $C(\mu^{\star 0}) = \Psi^0(\mu^{\star 0})$  by definition of  $\mu^{\star 0}$ .

Taking expectation on Equation 97 and making a Taylor expansion in  $\mu^{\star 0}$  gives the following for some  $\zeta$  between  $\mu_t$  and  $\mu^{\star 0}$ :

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ c_t \right] = \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ C(\mu_t^{\pi}) \right], \\ = C(\mu^{\star 0}) + \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ \mu_t^{\pi} - \mu^{\star 0} \right] C'(\mu^{\star 0}) + \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ (\mu_t^{\pi} - \mu^{\star 0})^2 \right] \frac{C''(\zeta)}{2}, \\ \stackrel{(a)}{\leq} \Psi^0(\mu^{\star 0}) + \left| \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ \mu_t^{\pi} - \mu^{\star 0} \right] \right| \mu^{\star 0} \overline{Z'} + \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ (\mu_t^{\pi} - \mu^{\star 0})^2 \right] \frac{\overline{\mu} \overline{Z''} + \overline{Z'}}{2}, \\ \stackrel{(b)}{=} \Psi^0(\mu^{\star 0}) + r_t \overline{Z'} + s_t \frac{\overline{\mu} \overline{Z''} + \overline{Z'}}{2}. \end{aligned}$$
(98)

Inequality (*a*) follows from the bounds on the derivative of *Z* in Assumption 13; Equality (*b*) finally follows from defining the mean squared error  $s_t = \mathbb{E}_{v,d} \left[ (\mu_t^{\pi} - \mu^{\star 0})^2 \right]$  and the absolute mean error  $r_t = \mu^{\star 0} \left| \mathbb{E}_{v,d} [\mu_t^{\pi} - \mu^{\star 0}] \right|$  for all  $t \leq \mathcal{T}$ .

Lemma E.1 provides an upper bound for the mean squared error, we next construct an equivalent bound for the absolute mean error.

Step 3: Upper Bound on the Absolute Mean Error. Note that for  $t < \mathcal{T}$ ,  $\mu_{t+1}^{\pi} = \mu_{t+1}$ . Taking expectation on the update rule  $\mu_{t+1} = \mu_t + \epsilon(z_t - \rho)$  and multiplying by  $\mu^{\star 0}$  hence gives:

$$\mu^{\star 0} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}} \left[ \mu_{t+1}^{\pi} - \mu^{\star 0} \right] = \mu^{\star 0} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}} \left[ \mu_{t}^{\pi} - \mu^{\star 0} \right] + \epsilon \mu^{\star 0} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}} \left[ z_{t} - g_{t} \right],$$

$$\stackrel{(a)}{=} \mu^{\star 0} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}} \left[ \mu_{t}^{\pi} - \mu^{\star 0} \right] + \epsilon \mu^{\star 0} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}} \left[ L(\mu_{t}^{\pi}) - L(\mu^{\star 0}) \right],$$

$$\stackrel{(b)}{=} \mu^{\star 0} \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}} \left[ \mu_{t}^{\pi} - \mu^{\star 0} \right] + \epsilon \mu^{\star 0} \left( L'(\mu^{\star 0}) \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}} \left[ \mu_{t}^{\pi} - \mu^{\star 0} \right] + \frac{1}{2} L''(\zeta) \mathbb{E}_{\boldsymbol{v}, \boldsymbol{d}} \left[ (\mu_{t}^{\pi} - \mu^{\star 0})^{2} \right] \right).$$
(99)

Equality (*a*) is obtained by taking expectation conditioned on  $\mu_t^{\pi}$  inside the expectation in the middle term and using the definition of  $\mu^{\star 0}$  in Assumption 14.1. Equality (*b*) uses a Taylor expansion of *L* in  $\mu^{\star 0}$  for some  $\zeta$  between  $\mu_t^{\pi}$  and  $\mu^{\star 0}$ .

Taking absolute values gives the following recursion, i.e.,

$$\begin{split} r_{t+1} &\leq \left| 1 + \epsilon L^{'}(\mu^{\star 0}) \right| r_{t} + \epsilon \frac{\mu^{\star 0}}{2} \left( \overline{Z^{''}} + \overline{G^{''}} \right) s_{t}, \\ &\stackrel{(a)}{=} \left( 1 + \epsilon L^{'}(\mu^{\star 0}) \right) r_{t} + \epsilon \frac{\mu^{\star 0}}{2} \left( \overline{Z^{''}} + \overline{G^{''}} \right) s_{t}, \\ &\stackrel{(b)}{\leq} (1 - \epsilon \lambda) r_{t} + \epsilon \frac{\overline{\mu}}{2} \left( \overline{Z^{''}} + \overline{G^{''}} \right) s_{t}. \end{split}$$

Note that we have  $-\left(\overline{Z}' + \overline{G}'\right) \leq L'(\mu) \leq -\lambda$  for all  $\mu \geq 0$  under Assumption 13. Hence Equality (*a*) holds because  $1 + \epsilon L'(\mu^{\star 0}) \geq 1 - \epsilon \left(\overline{Z}' + \overline{G}'\right) \geq 0$  under Assumption 14.4. Inequality (*b*) on the other hand uses that  $1 + \epsilon L'(\mu^{\star 0}) \leq 1 - \epsilon \lambda$  and the bound on  $\mu^{\star 0}$  from Assumption 14.1.

Since  $\lambda \epsilon \leq 1$  under Assumption 14.4 and  $r_1 \leq \overline{\mu}^2$  under Assumption 14.1, [6, Lemma C.4] then yields

$$r_t \leq \overline{\mu}^2 \left(1 - \epsilon \lambda\right)^{t-1} + \epsilon \frac{\overline{\mu}}{2} \left(\overline{Z''} + \overline{G''}\right) \sum_{j=1}^{t-1} (1 - \epsilon \lambda)^{t-1-j} s_j.$$

Using that the partial sums in  $(1 - \epsilon \lambda)^t$  are smaller than their limit  $1/(\epsilon \lambda)$ , we finally get the following upper bound on the total absolute error, i.e.,

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[\sum_{t=1}^{\mathcal{T}} r_t\right] \leq \overline{\mu}^2 \sum_{t=1}^{T} (1-\epsilon\lambda)^{t-1} + \epsilon \frac{\overline{\mu}}{2} \left(\overline{Z''} + \overline{G''}\right) \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[\sum_{t=1}^{\mathcal{T}} \sum_{j=1}^{t-1} (1-\epsilon\lambda)^{t-1-j} s_j\right],$$

$$\leq \frac{\overline{\mu}^2}{\epsilon\lambda} + \epsilon \frac{\overline{\mu}}{2} \left(\overline{Z''} + \overline{G''}\right) \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[\sum_{j=1}^{\mathcal{T}-1} s_j \sum_{t=0}^{\mathcal{T}-1-j} (1-\epsilon\lambda)^t\right],$$

$$\leq \frac{\overline{\mu}^2}{\epsilon\lambda} + \frac{\overline{\mu}}{2\lambda} \left(\overline{Z''} + \overline{G''}\right) \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[\sum_{j=1}^{\mathcal{T}} s_j\right].$$
(100)

Note that the same trick used in combination with Lemma E.1 leads to

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[\sum_{j=1}^{\mathcal{T}} s_j\right] \leq T \frac{\Delta^2 \epsilon}{2\lambda \underline{\mu}^2} + \overline{\mu}^2 \sum_{j=1}^{T} (1 - 2\lambda \underline{\epsilon})^{t-1},$$

$$\leq T \frac{\Delta^2 \epsilon}{2\lambda \underline{\mu}^2} + \frac{\overline{\mu}^2}{2\lambda \epsilon}.$$
(101)

Step 4: Bound on the Difference of Expected Dual Objectives. We first relate the properties known for  $\Psi_0$  to  $\Psi_d^{\rho}$ . By linearity of the differentiability operator, we first get that  $\Psi_d^{\rho}$  is differentiable under Assumption 13 since  $d^{\gamma}$  is bounded under Assumption 14.2. Its derivative is defined for all  $\mu \ge 0$  by  $\Psi_d^{\rho'}(\mu) = Z(\mu) - \frac{\gamma}{N} \mathbb{E}_d [d^{\gamma}] - \rho = L_d(\mu) - \rho$ , where we recognize the analogous expected karma loss function  $L_d$ .

Since -G' gives a positive contribution to the derivative of L for all  $\mu \ge 0$  under Assumption 1 (c.f. Equation 132 in Section F.1), Assumption 16 also implies that Z is strictly decreasing with parameter  $\lambda$ , hence  $L_d$  is both continuous and  $\lambda$ -strictly decreasing. Finally, note that Assumptions 14.1 and 14.3 imply that  $L_d(0) = (1 - \frac{\gamma}{N}) \mathbb{E}_{v,d} [d^{\gamma}]$  is strictly greater than  $\rho$ , whereas  $L_{\gamma}(\mu^{\star 0}) = L(\mu^{\star 0}) - \frac{\gamma}{N} \mathbb{E}_{v,d} [d^{\gamma} - p_t^{\gamma+1}]$  is negative, hence strictly smaller than  $\rho$ .

Then, the intermediate value theorem implies the existence of a  $\mu$  in the open interval  $]0, \mu^{\star 0}[$  such that  $L_d(\mu) = \rho$ . Since  $L_d$  is moreover strictly monotonic, this  $\mu$  is the unique root of  $\Psi_d^{\rho}$ , i.e.,  $\mu_d^{\star \rho}$ . Besides, the  $\lambda$ -strong monotonicity of  $L_d$  further implies that:

$$0 \leq \mu^{\star 0} - \mu^{\star \rho} \leq \frac{1}{\lambda} \left( L_d \left( \mu_d^{\star \rho} \right) - L_d (\mu^{\star 0}) \right),$$
  
$$\leq \frac{1}{\lambda} \left( \rho + \frac{\gamma}{N} \mathbb{E}_d [d^{\gamma} - d^{\gamma+1}] \right),$$
  
$$= \frac{1}{\lambda} \left( \rho + \hat{\varepsilon} \right).$$
(102)

We now turn to bound the difference of dual functions. First, note that:

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[\left(\Delta\boldsymbol{v}-\boldsymbol{\mu}_{d}^{\star\rho}\boldsymbol{d}^{\boldsymbol{\gamma}}\right)^{+}-\left(\Delta\boldsymbol{v}-\boldsymbol{\mu}^{\star0}\boldsymbol{d}^{\boldsymbol{\gamma}}\right)^{+}\right]=\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[\left(\boldsymbol{\mu}^{\star0}-\boldsymbol{\mu}_{d}^{\star\rho}\right)\boldsymbol{d}^{\boldsymbol{\gamma}}\mathbb{1}\left\{\frac{\Delta\boldsymbol{v}}{\boldsymbol{d}^{\boldsymbol{\gamma}}}\geq\boldsymbol{\mu}^{\star0}\geq\boldsymbol{\mu}_{d}^{\star\rho}\right\}\right],$$
$$+\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}^{\boldsymbol{\gamma}}}\left[\left(\Delta\boldsymbol{v}-\boldsymbol{\mu}_{d}^{\star\rho}\boldsymbol{d}^{\boldsymbol{\gamma}}\right)^{+}\mathbb{1}\left\{\boldsymbol{\mu}^{\star0}>\frac{\Delta\boldsymbol{v}}{\boldsymbol{d}^{\boldsymbol{\gamma}}}\geq\boldsymbol{\mu}_{d}^{\star\rho}\right\}\right],\qquad(103)$$
$$\leq\frac{\Delta}{\underline{\mu}}\left(\boldsymbol{\mu}^{\star0}-\boldsymbol{\mu}_{d}^{\star\rho}\right)+\Delta\mathbb{P}\left(\boldsymbol{\mu}^{\star0}>\frac{\Delta\boldsymbol{v}}{\boldsymbol{d}^{\boldsymbol{\gamma}}}\geq\boldsymbol{\mu}_{d}^{\star\rho}\right).$$

We express this probability explicitly by conditioning on the value of  $d^{\gamma}$ . Let  $H : d^{\gamma} \mapsto H(d^{\gamma})$  denote the cumulative probability function of the competing bid  $d^{\gamma}$ .

$$\mathbb{P}\left(\mu^{\star 0} > \frac{\Delta v}{dY} \ge \mu_{d}^{\star \rho}\right) = \int_{0}^{\Delta/\underline{\mu}} \int_{y\mu_{d}^{\star \rho}/\Delta}^{y\mu^{\star 0}/\Delta} v(x) dH(y) dx$$

$$\stackrel{(a)}{\le} \frac{\overline{v}}{\Delta} (\mu^{\star 0} - \mu_{d}^{\star \rho}) \int_{0}^{\Delta/\underline{\mu}} y dH(y)$$

$$\leq \frac{\overline{v}}{\underline{\mu}} (\mu^{\star 0} - \mu_{d}^{\star \rho})$$
(104)

Inequality (*a*) uses the change of variable  $z = \frac{x}{y}$  as well as Assumption 1.

We can now write:

$$\begin{split} \left| \Psi^{0}(\mu^{\star 0}) - \Psi^{\rho}_{d}(\mu^{\star \rho}) \right| &= \left| \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ \frac{\gamma}{N} (\mu_{d}^{\star \rho} d^{\gamma} - \mu^{\star 0} p) + \left( \Delta \boldsymbol{v} - \mu_{d}^{\star \rho} d^{\gamma} \right)^{+} - \left( \Delta \boldsymbol{v} - \mu^{\star 0} d^{\gamma} \right)^{+} + \mu_{d}^{\star \rho} \rho \right] \right| \\ &\stackrel{(a)}{\leq} \frac{\gamma}{N} \left( \mu^{\star 0} - \mu_{d}^{\star \rho} \right) \mathbb{E}_{\boldsymbol{d}} \left[ d^{\gamma} \right] + \frac{\gamma}{N} \mu^{\star 0} \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ d^{\gamma} - p \right] + \frac{\Delta}{\underline{\mu}} \left( 1 + \overline{\boldsymbol{v}} \right) \left( \mu^{\star 0} - \mu_{d}^{\star \rho} \right) + \mu_{d}^{\star \rho} \rho \\ &\stackrel{(b)}{\leq} \frac{\Delta}{\underline{\mu}} \left( 1 + \overline{\boldsymbol{v}} + \frac{\gamma}{N} \right) \left( \mu^{\star 0} - \mu_{d}^{\star \rho} \right) + \overline{\mu} \hat{\boldsymbol{\varepsilon}} + \overline{\mu} \rho \\ &\stackrel{(c)}{\leq} \left( \frac{\Delta}{\lambda \underline{\mu}} \left( 2 + \overline{\boldsymbol{v}} \right) + \overline{\mu} \right) \left( \rho + \hat{\boldsymbol{\varepsilon}} \right) \end{split}$$
(105)

Inequality (*a*) uses Equations 103 and 104; inequality (*b*) follows from Assumptions 14.1 and 14.2; inequality (*c*) finally uses Equation 102 and the fact that the fraction of auction winners is bounded by one.

Intermediate Conclusion. Together, Equations 95, 96, 98, 100, 101 and 105 finally lead to the following, i.e.,

$$\begin{split} \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[C^{G}(\boldsymbol{v},\boldsymbol{d})-C^{H}(\boldsymbol{v},\boldsymbol{d})\right] &\leq \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[\sum_{t=1}^{\mathcal{T}}c_{t}\right]+\tau\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[T-\mathcal{T}\right]-T\Psi^{\rho}(\mu^{\star\rho}),\\ &\leq T\left(\Psi^{0}(\mu^{\star0})-\Psi^{\rho}(\mu^{\star\rho})\right)+\tau\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[T-\mathcal{T}\right]+\frac{\overline{\mu}^{2}\overline{Z'}}{\epsilon\lambda}\\ &+\left(\frac{\overline{\mu}\overline{Z'}}{2\lambda}\left(\overline{Z''}+\overline{G''}\right)+\frac{\overline{\mu}\overline{Z''}+\overline{Z'}}{2}\right)\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[\sum_{t=1}^{\mathcal{T}}s_{t}\right],\\ &\leq T\left(\frac{\Delta}{\lambda\underline{\mu}}\left(2+\overline{\nu}\right)+\overline{\mu}\right)\left(\rho+\hat{\varepsilon}\right)+\tau\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[T-\mathcal{T}\right]+\frac{\overline{\mu}^{2}\overline{Z'}}{\epsilon\lambda}\\ &+\left(\frac{\overline{\mu}\overline{Z'}}{2\lambda}\left(\overline{Z''}+\overline{G''}\right)+\frac{\overline{\mu}\overline{Z''}+\overline{Z'}}{2}\right)\left(T\frac{\Delta^{2}\epsilon}{2\lambda\underline{\mu}^{2}}+\frac{\overline{\mu}^{2}}{2\lambda\epsilon}\right). \end{split}$$

Rewriting  $\rho = k_1/T$ , we hence proved the existence of a constant *C* in  $\mathbb{R}$  such that

$$\frac{1}{T}\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[C^{G}(\boldsymbol{v},\boldsymbol{d}) - C^{H}(\boldsymbol{v},\boldsymbol{d})\right] \leq C\left(\epsilon + \frac{1+k_{1}\epsilon}{\epsilon T} + \frac{\mathbb{E}_{\boldsymbol{v},\boldsymbol{d}}\left[T-\mathcal{T}\right]}{T} + \hat{\varepsilon}\right).$$
(106)

*Step 5: Control of Hitting Time.* We finally show that  $T_i = T$  under Assumption 9.

We first prove that  $\mathcal{T}_i^{\overline{\mu}} = T$ , i.e. that  $\mu_t \leq \overline{\mu}$  for all  $1 \leq t \leq T$ . By contradiction, suppose there exists a first time  $2 \leq t \leq T$  where  $\mu_t > \overline{\mu}$ . By the pigeonhole principle,  $\mu_{t-1}$  must belong in  $\left[\overline{\mu} - \epsilon \overline{d}, \overline{\mu}\right]$  and the agent must have won the auction. The latter is however not possible because we have:

$$b_{t-1} = \frac{\Delta v_{t-1}}{\mu_{t-1}} \le \frac{\Delta}{\overline{\mu} - \epsilon \overline{d}} < \underline{d},$$

where the last inequality follows from  $\epsilon < \frac{1}{\overline{d}} \left( \overline{\mu} - \underline{\underline{A}} \right)$  in Assumption 9.

Similarly, we prove that  $\mathfrak{T}_{i}^{\underline{\mu}} = T$ , i.e.  $\mu_{t} \ge \underline{\mu}$  for all  $1 \le t \le T$ . Indeed, suppose  $\mu_{t}$  belongs in  $\left[\underline{\mu}, \underline{\mu} + \epsilon \overline{d}\right]$ . We then have:

$$b_t = \frac{\Delta v_t}{\mu_t} \ge \frac{\Delta \underline{v}}{\mu + \epsilon \overline{d}} > \overline{d},$$

where the last inequality follows from  $\epsilon < \frac{\Delta v}{\overline{d}^2} - \mu$  in Assumption 9. Hence, the auction is lost, and  $\mu_{t+1} > \mu_t$ . The pigeonhole principle then implies the desired result.

We finally prove that  $\mathcal{T}_i^k = T$  and that the budget is never depleted. Indeed, suppose t is such that  $k_t \leq \Delta/\mu$ . Then we have:

$$\mu_t = \mu_1 + \epsilon (k_1 - k_t) > \overline{\mu},$$

using  $k_{i,1} > \frac{\overline{\mu_i} - \mu_{i,1}}{\epsilon} + \frac{\Lambda}{\mu_i}$  in Assumption 9. This is impossible since we previously proved that  $\mu_t \leq \overline{\mu}$  for all  $1 \leq t \leq T$ .

Given the definition of  $T_i$  in Equation 22, these three properties together imply that  $T_i = T$ .

## E.3 Proof of Theorem 4.2

We start the proof of Theorem 4.2 by showing that the multiplier profile  $\mu_t$  converges on average to the stationary  $\mu^*$ .

Step 1: Convergence on Average of the Multiplier Profile . Summing Lemma E.1 gives:

$$\begin{split} \frac{1}{T} \sum_{t=1}^{T} \mathbb{E}_{\boldsymbol{v}} \left[ \|\boldsymbol{\mu}_{t}^{\boldsymbol{\pi}} - \boldsymbol{\mu}^{\boldsymbol{\star}}\|_{2}^{2} \right] &= \frac{1}{T} \mathbb{E}_{\boldsymbol{v}} \left[ \sum_{t=1}^{T} \|\boldsymbol{\mu}_{t}^{\boldsymbol{\pi}} - \boldsymbol{\mu}^{\boldsymbol{\star}}\|_{2}^{2} \mathbb{1} \{ t \leq \underline{\Upsilon} \} + \|\boldsymbol{\mu}_{t}^{\boldsymbol{\pi}} - \boldsymbol{\mu}^{\boldsymbol{\star}}\|_{2}^{2} \mathbb{1} \{ t > \underline{\Upsilon} \} \right], \\ &\stackrel{(a)}{\leq} \frac{1}{T} \sum_{t=1}^{\underline{\Upsilon}} \left[ N \overline{\mu}^{2} (1 - 2\lambda\epsilon)^{t-1} + \frac{N \Delta^{2} \epsilon}{2\lambda \underline{\mu}^{2}} \right] + \frac{N \overline{\mu}^{2}}{T} \mathbb{E}_{\boldsymbol{v}} \left[ T - \underline{\Upsilon} \right], \\ &\stackrel{(b)}{\leq} \frac{N \overline{\mu}^{2}}{2\lambda\epsilon T} + \frac{N \Delta^{2} \epsilon}{2\lambda \mu^{2}} + \frac{N \overline{\mu}^{2}}{T} \mathbb{E}_{\boldsymbol{v}} \left[ T - \underline{\Upsilon} \right]. \end{split}$$

Inequality (*a*) holds by assuming that the error is maximal after time  $\underline{T}$ , and by applying Lemma E.1; Inequality (*c*) holds by bounding the partial geometric series in  $1 - 2\lambda\epsilon$  by its limit  $1/(2\lambda\epsilon)$ , as well as using  $\underline{T} \leq T$ .

Hence there exist some constant *C* in  $\mathbb{R}_+$  such that

$$\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}_{\boldsymbol{v}}\left[\|\boldsymbol{\mu}_{t}^{\boldsymbol{\pi}}-\boldsymbol{\mu}^{\boldsymbol{\star}}\|_{2}^{2}\right] \leq CN\left(\boldsymbol{\epsilon}+\frac{1}{\boldsymbol{\epsilon}T}+\frac{\mathbb{E}_{\boldsymbol{v}}\left[T-\underline{\mathfrak{I}}\right]}{T}\right).$$

Step 2: Lipschitz Continuity of the Expected Dual Objective and the Expected Karma Loss. Again, we consider a more general setting with M parallel auctions. Let  $a_{i,j}$  denote the probability for two agents i and j to play in the same auction; we consider the competing price  $d_i^{\gamma} = \gamma^{\text{th}} - \max_{j:j \neq i} \{\mathbbm{1}\{m_j = m_i\}\Delta v_j/\mu_j\}$ , as well as the gain  $g_i = \frac{\gamma}{N} \sum_{m=1}^{M} p_m^{\gamma+1}$ , where  $p_m^{\gamma+1} = \gamma + 1^{\text{th}} - \max_i \{\mathbbm{1}\{m_i = m\}\Delta v_i/\mu_i\}$  denotes the price of the  $m^{\text{th}}$  auction.

For  $\rho \ge 0$ , recall the definitions of the dual function  $\Psi_i^{\rho}(\boldsymbol{\mu}) = \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} [v_i \tau - \mu_i (g_i + \rho) - (\Delta v_i - \mu_i d_i^{\gamma})^+]$ , the expenditure function  $Z_i(\boldsymbol{\mu}) = \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} [d_i^{\gamma} \mathbb{I} \{\Delta v_i > \mu_i d_i^{\gamma}\}]$ , the gain function  $G_i(\boldsymbol{\mu}) = \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} [g_i]$ , as well as the karma loss function  $L_i(\boldsymbol{\mu}) = Z_i(\boldsymbol{\mu}) - G_i(\boldsymbol{\mu})$ .

Similarly as in F.1, note that  $\Psi_i^{\rho}(\mu) = \Psi_i(\mu) + \mu_i(\rho_i - \rho) - \mu_i G_i(\mu)$ , where  $\Psi$  refers to the dual function of Section 3.3. Since the expenditure function  $Z_i$  is defined independently of the framework, it is still Lipschitz continuous (c.f. Section 3.3). It hence only remains to bound the derivatives of

 $G_i$ . Since the gains  $g_i$  are the same for everyone, we drop the subscript *i* and study the expected gain function G.

For any  $m \in [M]$  and realized vectors v and m, the function  $\mu \mapsto p_{m,t}^{\gamma+1}$  is differentiable in  $\mu_i$  outside of sets of measure zero under Assumption 1. Note that the derivative is non-null only when agent i is the price setter in auction m, that it is bounded  $\Delta/\underline{\mu}^2$  which is integrable over  $[\underline{\mu}, \overline{\mu}]$ , hence Leibniz integral rule implies:

$$\begin{split} \frac{\partial G}{\partial \mu_i}(\boldsymbol{\mu}) &= -\mathbb{E}_{\boldsymbol{v},\boldsymbol{m}}\left[\frac{\gamma}{N}\sum_{m=1}^M \frac{\Delta v_i}{\mu_i^2}\mathbbm{1}\left\{m_i = m\right\}\mathbbm{1}\left\{d_i^{\gamma} > \frac{\Delta v_i}{\mu_i} > d_i^{\gamma+1}\right\}\right],\\ &= -\mathbb{E}_{\boldsymbol{v},\boldsymbol{m}}\left[\frac{\gamma}{N}\frac{\Delta v_i}{\mu_i^2}\mathbbm{1}\left\{d_i^{\gamma} > \frac{\Delta v_i}{\mu_i} > d_i^{\gamma+1}\right\}\right]. \end{split}$$

Note that  $\mathbb{1}\left\{d_i^{\gamma} > \frac{\Delta v_i}{\mu_i} > d_i^{\gamma+1}\right\}$  is exactly equal to one when agent *i* sets the price for its auction. Since it can only happen for *M* agents simultaneously, we bound the sum of the derivatives as follows, i.e.,

$$\left| \sum_{i=1}^{N} \frac{\partial G}{\partial \mu_{i}}(\boldsymbol{\mu}) \right| = \sum_{i \in \mathcal{N}} \mathbb{E}_{\boldsymbol{\nu}, \boldsymbol{m}} \left[ \frac{\gamma \Delta v_{i}}{N \mu_{i}^{2}} \mathbb{1} \left\{ d_{i}^{\gamma} > \frac{\Delta v_{i}}{\mu_{i}} > d_{i}^{\gamma+1} \right\} \right],$$

$$\stackrel{(a)}{=} \mathbb{E}_{\boldsymbol{\nu}, \boldsymbol{m}} \left[ \sum_{i \in \mathcal{N}} \frac{\gamma \Delta v_{i}}{N \mu_{i}^{2}} \mathbb{1} \left\{ d_{i}^{\gamma} > \frac{\Delta v_{i}}{\mu_{i}} > d_{i}^{\gamma+1} \right\} \right],$$

$$\leq \frac{\gamma \Delta}{N \mu^{2}} \mathbb{E}_{\boldsymbol{\nu}, \boldsymbol{m}} \left[ \sum_{i \in \mathcal{N}} \mathbb{1} \left\{ d_{i}^{\gamma} > \frac{\Delta v_{i}}{\mu_{i}} > d_{i}^{\gamma+1} \right\} \right],$$

$$\stackrel{(b)}{\leq} \frac{M \gamma \Delta}{N \mu^{2}}.$$
(107)

Equality (*a*) holds by linearity of expectation, and inequality (*b*) uses that at most *M* agents can be price setters simultaneously. This allows to prove the Lipschitz continuity of the dual function of agent  $i \in N$ :

$$\begin{aligned} \left|\Psi_{i}^{\rho}(\boldsymbol{\mu})-\Psi_{i}^{\rho}(\boldsymbol{\mu}')\right| &\leq \left|\Psi_{i}(\boldsymbol{\mu})-\Psi_{i}(\boldsymbol{\mu}')\right|+\left|\mu_{i}G(\boldsymbol{\mu})-\mu_{i}'G(\boldsymbol{\mu}')\right|,\\ &\stackrel{(a)}{\leq}\left|\Psi_{i}(\boldsymbol{\mu})-\Psi_{i}(\boldsymbol{\mu}')\right|+\left|\mu_{i}\sum_{j\in\mathcal{N}}\frac{\partial G}{\partial\mu_{j}}(\boldsymbol{\mu})\left|\mu_{j}-\mu_{j}'\right|\right|+\left|G(\boldsymbol{\mu})-G(\boldsymbol{\mu}')\right|\left|\mu_{i}-\mu_{i}'\right|,\\ &\stackrel{(b)}{\leq}\left|\Psi_{i}(\boldsymbol{\mu})-\Psi_{i}(\boldsymbol{\mu}')\right|+\left|\sum_{j\in\mathcal{N}}\frac{\partial G}{\partial\mu_{j}}(\boldsymbol{\mu})\right|\overline{\boldsymbol{\mu}}\|\boldsymbol{\mu}-\boldsymbol{\mu}'\|_{\infty}+\frac{M\gamma\Delta}{N\underline{\mu}}\left|\mu_{i}-\mu_{i}'\right|,\\ &\stackrel{(c)}{\leq}\frac{\Delta}{\underline{\mu}}\sum_{j\neq i}a_{i,j}\left|\mu_{j}-\mu_{j}'\right|+\frac{M\gamma\Delta\overline{\mu}}{N\underline{\mu}^{2}}\|\boldsymbol{\mu}-\boldsymbol{\mu}'\|_{\infty}+\frac{\Delta}{\underline{\mu}}\left(1+\frac{M\gamma}{N}\right)\left|\mu_{i}-\mu_{i}'\right|,\\ &\stackrel{(d)}{\leq}\frac{\Delta}{\underline{\mu}}\left(\|\boldsymbol{a}_{i}\|_{2}+\frac{M\gamma\overline{\mu}}{N\underline{\mu}}\right)\|\boldsymbol{\mu}-\boldsymbol{\mu}'\|_{2}+\frac{\Delta}{\underline{\mu}}\left(1+\frac{M\gamma}{N}\right)\left|\mu_{i}-\mu_{i}'\right|.\end{aligned}$$

$$(108)$$

Inequality (*a*) uses the product rule and the triangle inequality; inequality (*b*) uses the definition of the norm  $\|\mathbf{x}\|_{\infty} = \max_{i} x_{i}$  as well as the fact that all *M* prices  $p_{m}^{\gamma+1}$  can change by at most  $\Delta/\mu$ ;

inequality (c) follows from Equations 59 and 107, and inequality (d) uses Cauchy-Schwartz as well as the norm inequality  $\|\mathbf{x}\|_{\infty} \leq \|\mathbf{x}\|_{2}$ .

We finally establish the Lipschitz continuity of the karma loss function:

$$\begin{aligned} \left| L_{i}(\boldsymbol{\mu}) - L_{i}\left(\boldsymbol{\mu}'\right) \right| &\leq \left| Z_{i}(\boldsymbol{\mu}) - Z_{i}\left(\boldsymbol{\mu}'\right) \right| + \left| \sum_{j \in \mathcal{N}} \frac{\partial G}{\partial \mu_{j}}(\boldsymbol{\mu}) \left| \mu_{j} - \mu_{j}' \right| \right|, \\ &\stackrel{(a)}{\leq} \frac{\Delta \overline{\nu}}{\underline{\mu}^{2}} \left( \left| \mu_{i} - \mu_{i}' \right| + \sum_{j \neq i} 8a_{i,j} \left| \mu_{j} - \mu_{j}' \right| \right) + \frac{M \gamma \Delta}{N \underline{\mu}^{2}} \left\| \boldsymbol{\mu} - \boldsymbol{\mu}' \right\|_{\infty}, \end{aligned}$$

$$\stackrel{(b)}{\leq} \frac{\Delta}{\underline{\mu}^{2}} \left( 8 \overline{\nu} \| \boldsymbol{a}_{i} \|_{2} + \frac{M \gamma}{N} \right) \| \boldsymbol{\mu} - \boldsymbol{\mu}' \|_{2} + \frac{\Delta \overline{\nu}}{\underline{\mu}^{2}} | \mu_{i} - \mu_{i}' |. \end{aligned}$$

$$(109)$$

Inequality (*a*) uses Equations 73 and 107 with the definition of the norm  $\|\cdot\|_{\infty}$ ; inequality (*b*) follows from Cauchy-Schwartz and the norm inequality  $\|\mathbf{x}\|_{\infty} \leq \|\mathbf{x}\|_{2}$ .

Step 3: Upper Bound on the Total Expected Cost of Strategy G. Similarly as its counterpart in C.3, it mainly consists in expressing the expected cost per period in terms of the dual and the loss functions, and using their bounded variations to express the cost in terms of the mean squared error.

Like in the second step of Section E.2, we bound the expected total cost of agent  $i \in N$  by considering it always gets maximum valuation after time  $\underline{T}$  but never accesses the priority road.

$$C_{i}^{G} \leq \mathbb{E}_{\upsilon} \left[ \sum_{t=1}^{\underline{\Upsilon}} c_{i,t} \right] + \tau \mathbb{E}_{\upsilon} \left[ T - \underline{\Upsilon} \right]$$
(110)

Rewriting the cost suffered by agent *i* at time  $t \leq \underline{T}$  in a similar manner as in Equation 97, taking expectation and conditioning on  $\mu_t^{\pi}$  gives:

$$\mathbb{E}_{\boldsymbol{v}}\left[c_{i,t}\right] = \mathbb{E}_{\boldsymbol{v}}\left[\Psi_{i}^{0}(\boldsymbol{\mu}_{t}^{\boldsymbol{\pi}})\right] - \mathbb{E}_{\boldsymbol{v}}\left[\mu_{i,t}^{\boldsymbol{\pi}}L_{i}(\boldsymbol{\mu}_{t}^{\boldsymbol{\pi}})\right].$$
(111)

We again use the Lipschitz continuity of  $\Psi_i^0$  to express the expected cost in terms of  $\mu^{\star 0}$ .

$$\Psi_i(\boldsymbol{\mu}_t^{\boldsymbol{\pi}}) \le \Psi_i^0(\boldsymbol{\mu}^{\star 0}) + \mathcal{L}_{\Psi} \| \boldsymbol{\mu}_t^{\boldsymbol{\pi}} - \boldsymbol{\mu}^{\star 0} \|_1$$
(112)

Similarly, the Lipschitz continuity of  $L_i$  gives

$$-\mu_{i,t}^{\pi} L_{i}(\boldsymbol{\mu}_{t}^{\pi}) \stackrel{(a)}{=} \mu_{i,t}^{\pi} \left( L_{i}(\boldsymbol{\mu}^{\star 0}) - L_{i}(\boldsymbol{\mu}_{t}^{\pi}) \right) \stackrel{(b)}{\leq} \overline{\mu} \mathcal{L}_{L} \| \boldsymbol{\mu}_{t}^{\pi} - \boldsymbol{\mu}^{\star 0} \|_{1},$$
(113)

where equality (a) holds by definition of a stationary multiplier profile, and inequality (b) uses the upper bound induced by the projection.

Combining Equations 111, 112 and 113, we get:

$$\mathbb{E}_{\boldsymbol{v}}\left[c_{i,t}\right] \leq \Psi_{i}^{0}\left(\boldsymbol{\mu^{\star 0}}\right) + \left(\mathcal{L}_{\Psi} + \overline{\boldsymbol{\mu}}\mathcal{L}_{L}\right) \mathbb{E}_{\boldsymbol{v}}\left[\|\boldsymbol{\mu}_{t}^{\pi} - \boldsymbol{\mu^{\star 0}}\|_{1}\right],$$

$$\overset{(a)}{\leq} \Psi_{i}^{0}\left(\boldsymbol{\mu^{\star 0}}\right) + \left(\mathcal{L}_{\Psi} + \overline{\boldsymbol{\mu}}\mathcal{L}_{L}\right) \mathbb{E}_{\boldsymbol{v}}\left[\sqrt{N}\|\boldsymbol{\mu}_{t}^{\pi} - \boldsymbol{\mu^{\star 0}}\|_{2}\right],$$

$$\overset{(b)}{\leq} \Psi_{i}^{0}\left(\boldsymbol{\mu^{\star 0}}\right) + \left(\mathcal{L}_{\Psi} + \overline{\boldsymbol{\mu}}\mathcal{L}_{L}\right)\sqrt{N}s_{t}^{1/2}.$$
(114)

Inequality (a) uses that  $\sum_{i=1}^{n} |y_i| \le (n \sum_{i=1}^{n} y_i^2)^{1/2}$ , and inequality (c) follows from Jensen's inequality.

Intermediary Conclusion: Combining Equations 110 and 113 with Lemma E.1 finally gives:

$$\begin{split} \frac{1}{T}C_{i}^{G} - \Psi_{i}^{0}(\boldsymbol{\mu}^{\star}) &\leq \frac{1}{T}\mathbb{E}_{\boldsymbol{\nu}}\left[\sum_{t=1}^{\underline{\Upsilon}}c_{i,t}\right] + \frac{\tau}{T}\mathbb{E}_{\boldsymbol{\nu}}\left[T - \underline{\Upsilon}\right] - \Psi_{i}^{0}(\boldsymbol{\mu}^{\star}), \\ &\leq \left(\mathcal{L}_{\Psi} + \overline{\mu}\mathcal{L}_{L}\right)\frac{\sqrt{N}}{T}\mathbb{E}_{\boldsymbol{\nu}}\left[\sum_{t=1}^{\underline{\Upsilon}}s_{t}^{1/2}\right] + \frac{\tau}{T}\mathbb{E}_{\boldsymbol{\nu}}\left[T - \underline{\Upsilon}\right], \\ &\stackrel{(a)}{\leq} \left(\mathcal{L}_{\Psi} + \overline{\mu}\mathcal{L}_{Z}\right)N\left(\frac{\Delta\epsilon^{1/2}}{\sqrt{2\lambda\mu}} + \frac{\overline{\mu}}{T}\sum_{t=1}^{T}\left(1 - 2\lambda\epsilon\right)^{(t-1)/2}\right) + \frac{\tau}{T}\mathbb{E}_{\boldsymbol{\nu}}\left[T - \underline{\Upsilon}\right] \\ &\stackrel{(b)}{\leq} \left(\mathcal{L}_{\Psi} + \overline{\mu}\mathcal{L}_{Z}\right)N\left(\frac{\Delta\epsilon^{1/2}}{\sqrt{2\lambda\mu}} + \frac{2\overline{\mu}}{\lambda\epsilon T}\right) + \frac{\tau}{T}\mathbb{E}_{\boldsymbol{\nu}}\left[T - \underline{\Upsilon}\right]. \end{split}$$

Inequality (*a*) follows from Lemma C.2 with  $\sqrt{x+y} \le \sqrt{x} + \sqrt{y}$ . Inequality (*b*) holds by bounding the geometric sum by the series' limit and noting that  $\frac{1}{1-(1-\lambda\epsilon)^{1/2}} \le \frac{2}{\lambda\epsilon}$  since  $1 - (1-x)^{1/2} \ge x/2$  for  $x \in [0, 1]$ .

Rewriting  $\rho = \underline{k_1}/T$ , this proves the existence of constants *C* in  $\mathbb{R}_+$  such that

$$\frac{1}{T}C_i^G - \Psi_i^0(\boldsymbol{\mu^{\star 0}}) \le C\left(N\left(\epsilon^{1/2} + \frac{1}{\epsilon T}\right) + \frac{\mathbb{E}_{\boldsymbol{\nu}}\left[T - \underline{\mathcal{I}}\right]}{T}\right)$$

*Step 4: Control of Hitting Time.* It only remains to show that the term  $\frac{\mathbb{E}_{v}[T-\underline{T}]}{T}$  converges to zero under Assumption 10.

Note that Assumptions 9.4 and 10.3 respectively imply  $\mathfrak{T}_i^k = T$  and  $\mathfrak{T}_i^{\underline{\mu}} = T$ .

We prove in the following that the multiplier profile  $\mu_t$  remains in the set  $\prod_{i=1}^{N} [-\overline{\mu_i}, \overline{\mu_i}]$  at all time  $1 \le t \le T$ , effectively implying  $\underline{\mathcal{T}}^{\overline{\mu}} = T$ , and in turn  $\underline{\mathcal{T}} = T$ .

For  $t \in \mathbb{N}$ , we define  $\overline{\mu}_t = \max_{i \in \mathbb{N}} \mu_{i,t}$  and  $\underline{\mu}_t = \min_{i \in \mathbb{N}} \mu_{i,t}$ . For a certain realization  $\mu_{t+1}$ , we retrospectively consider a "big" agent  $b \in \mathbb{N}$  such that  $\mu_{b,t+1} = \overline{\mu}_{t+1}$  (not necessarily unique). We similarly consider a "small" agent *s* satisfying  $\mu_{s,t+1} = \underline{\mu}_{t+1}$ . Finally, we denote by  $\mathcal{W}_t$  the set of auction winners, by  $\mathcal{L}_t$  the set of the losers, and  $\sum_{i=1}^{N} \mu_{i,1} = \mathbb{N} \mu_m$  the constant sum of the multipliers.

To prove the Lemma, we show by induction that the difference  $\overline{\mu}_t - \underline{\mu}_t$  remains bounded for all  $1 \le t \le T$ .

We proceed by exhaustion and consider at first the case where it does not happen that agent b wins the auction while agent s loses it:

PROPOSITION E.2. Suppose it is not the case that  $b \in W_t$  and  $s \in \mathcal{L}_t$ . Then  $\overline{\mu}_{t+1} - \mu_{t+1} \leq \overline{\mu}_t - \mu_t$ .

PROOF. By exhaustion, we need to consider the following two cases:

(1) If *b* and *s* are both in  $\mathcal{L}_t$  or both in  $\mathcal{W}_t$ , we have:

$$\overline{\mu}_{t+1} - \underline{\mu}_{t+1} \stackrel{(a)}{=} \mu_{b,t+1} - \mu_{s,t+1},$$
$$\stackrel{(b)}{=} \mu_{b,t} - \mu_{s,t},$$
$$\stackrel{(c)}{\leq} \overline{\mu}_t - \mu_t.$$

Equality (*a*) holds by definition of *b* and *s*. Equality (*b*) uses that variations in multiplier only depend on the affiliation of an agent to  $W_t$  or to  $\mathcal{L}_t$ , i.e., all  $w \in W_t$  increase their multiplier by the same value  $\epsilon p_t$ , while all  $\ell \in \mathcal{L}_t$  decrease their multiplier by a common  $\epsilon g_t$ . Inequality (*c*) finally follows from the definition of  $\overline{\mu}_t$  and  $\mu_t$ .

(2) If  $b \in \mathcal{L}_t$  and  $s \in \mathcal{W}_t$ , we have:

$$\overline{\mu}_{t+1} - \underline{\mu_{t+1}} \stackrel{(a)}{=} \mu_{b,t+1} - \mu_{s,t+1},$$

$$\stackrel{(b)}{\leq} \mu_{b,t} - \mu_{s,t} - \epsilon \left(1 + \frac{M\gamma}{N}\right) \frac{\Delta \underline{v}}{\overline{\mu}},$$

$$\stackrel{(c)}{\leq} \overline{\mu}_t - \mu_t.$$

Equality (*a*) holds by definition of *b* and *s*, equality (*b*) uses the minimum variation step of multipliers for the price  $p_t = \Delta \underline{v} / \overline{\mu}$ . Inequality (*c*) finally follows from the definition of  $\overline{\mu}_t$  and  $\mu_t$ .

We now turn to analyse the remaining case where agent *b* wins the auction while agent *s* loses it.

PROPOSITION E.3. Suppose that  $b \in W_t$  and  $s \in \mathcal{L}_t$ . Then  $\overline{\mu}_t - \underline{\mu}_t \leq \mu_m \left(\frac{2}{\underline{v}} \left(1 - \frac{Y}{N}\right)^{-1} - \frac{\overline{v}}{\underline{v}}\right)$ .

PROOF. We consider an hypothetical framework where  $\overline{\mu} = \infty$ , and iteratively bound  $\overline{\mu}_{t+1}, \underline{\mu_{t+1}}, \overline{\mu}_t$  and  $\mu_t$ .

We first note that  $\mu_{b,t}$  cannot be too far below  $\mu_m$ , hence it is greater than  $\mu$ :

$$\mu_{b,t} \stackrel{(a)}{\geq} \mu_{b,t+1} - \epsilon \frac{\gamma}{N} \frac{\Delta \overline{v}}{\underline{\mu}},$$

$$\stackrel{(b)}{=} \overline{\mu}_{t+1} - \epsilon \frac{\gamma}{N} \frac{\Delta \overline{v}}{\underline{\mu}},$$

$$\stackrel{(c)}{\geq} \mu_m - \epsilon \frac{\gamma}{N} \frac{\Delta \overline{v}}{\underline{\mu}},$$

$$\stackrel{(d)}{\geq} \frac{v}{\overline{v}} \frac{\mu_m}{2},$$

$$\stackrel{(e)}{\geq} \underline{\mu}.$$

$$(115)$$

Equality (a) uses that the maximum multiplier increase is  $\epsilon \frac{\gamma}{N} \frac{\Delta \bar{v}}{\mu}$ ; equality (b) follows from the definition of b; inequality (c) uses that the maximum must be greater than the mean; inequality (d) and (e) respectively follow from the bound on  $\epsilon$  and that on  $\mu$  in the statement of the Lemma.

We then note that for all  $\ell \in \mathcal{L}_t$ , we must have:

$$\frac{\Delta \underline{v}}{\max\{\mu_{\ell,t},\underline{\mu}\}} \le b_{\ell,t} \le b_{b,t} \le \frac{\Delta}{\max\{\mu_{b,t},\underline{\mu}\}} = \frac{\Delta}{\mu_{b,t}}.$$
(116)

Using the above, we next show that  $\max\{\mu_{\ell,t},\mu\} = \mu_{\ell,t}$  holds for all  $\ell \in \mathcal{L}_t$ .

$$\max\{\mu_{\ell,t}, \underline{\mu}\} \stackrel{(a)}{\geq} \underline{v}\mu_{b,t},$$

$$\stackrel{(b)}{\geq} \underline{v}\left(\mu_m - \epsilon \frac{\gamma}{N} \frac{\Delta}{\underline{\mu}}\right),$$

$$\stackrel{(c)}{\geq} \underline{v} \frac{\mu_m}{2},$$

$$\stackrel{(d)}{\geq} \underline{\mu}.$$
(117)

Inequalities (*a*) and (*b*) use the inequalities in Equations 116 and 115 respectively; inequalities (*c*) and (*d*) respectively follow from the bounds on  $\epsilon$  and  $\mu$  in the statement of the Lemma.

As Equation 117 holds in particular for *s*, we propagate this lower-bound to  $\mu_t$  and  $\mu_{t+1}$ :

$$\min\{\underline{\mu_{t}}, \underline{\mu_{t+1}}\} \stackrel{(a)}{\geq} \mu_{s,t} - \epsilon \left(1 + \frac{\gamma}{N}\right) \frac{\Delta}{\underline{\mu}},$$

$$\stackrel{(b)}{\geq} \underline{v}\mu_{m} - \epsilon \left(1 + \left(1 + \underline{v}\right) \frac{\gamma}{N}\right) \frac{\Delta}{\underline{\mu}},$$

$$\stackrel{(c)}{\geq} \underline{v} \frac{\mu_{m}}{2},$$

$$\stackrel{(d)}{\geq} \mu.$$
(118)

Equality (a) uses that both  $\underline{\mu_t}$  and  $\underline{\mu_{t+1}}$  are at most  $\epsilon \left(1 + \frac{\gamma}{N}\right) \frac{\Delta}{\mu}$  away from  $\mu_{s,t}$  by definition of s; inequality (b) makes use of the the lower-bound in Equation 117 (b), inequalities (c) and (d) respectively follow from the bounds on  $\epsilon$  and  $\mu$  in the statement of the Lemma.

We now proceed to upper-bound  $\overline{\mu}_t$  and  $\overline{\mu}_{t+1}$ . By conservation of the sum of multipliers, we have:

$$N\mu_{m} = \mu_{b,t} + \sum_{\ell \in \mathcal{L}_{t}} \mu_{\ell,t} + \sum_{w \in \mathcal{W}_{t} \setminus \{b\}} \mu_{w,t},$$
  

$$\geq \mu_{b,t} \left( 1 + (N - \gamma)\underline{v} \right) + (\gamma - 1)\underline{\mu}.$$
(119)

The inequality follows from inequation 117 (*a*), as well as the definition of  $\mu_t$  and inequation 118. It then follows that:

$$\max\{\overline{\mu}_{t}, \overline{\mu}_{t+1}\} \stackrel{(a)}{\leq} \mu_{b,t} + \epsilon \left(1 + \frac{\gamma}{N}\right) \frac{\Delta}{\underline{\mu}},$$

$$\stackrel{(b)}{\leq} \frac{N\mu_{m} - (\gamma - 1)\underline{\mu}}{1 + (N - \gamma)\underline{v}} + \epsilon \left(1 + \frac{\gamma}{N}\right) \frac{\Delta}{\underline{\mu}},$$

$$< \frac{1}{\underline{v}} \mu_{m} \left(1 - \frac{\gamma}{N}\right)^{-1} + \epsilon \left(1 + \frac{\gamma}{N}\right) \frac{\Delta}{\underline{\mu}},$$

$$\stackrel{(c)}{\leq} \frac{2}{\underline{v}} \mu_{m} \left(1 - \frac{\gamma}{N}\right)^{-1}.$$
(120)

Equality (a) uses that both  $\overline{\mu_t}$  and  $\overline{\mu_{t+1}}$  are at most  $\epsilon \left(1 + \frac{\gamma}{N}\right) \frac{\Delta}{\mu}$  away from  $\mu_{b,t}$  by definition of *b*; inequality (*b*) follows from Equation 119 and inequality (*c*) from the bounds on  $\epsilon$  in the statement of the Lemma.

This finally proves that 
$$\overline{\mu}_{t+1} - \underline{\mu}_{t+1} \le \mu_m \left(\frac{2}{\underline{v}} \left(1 - \frac{\gamma}{N}\right)^{-1} - \frac{\overline{v}}{\underline{v}}\right).$$

Together, the two propositions show the induction and  $\overline{\mu}_t - \underline{\mu}_t \leq \mu_m \left(\frac{2}{\underline{v}} \left(1 - \frac{Y}{N}\right)^{-1} - \frac{\overline{v}}{2}\right)$  for all  $1 \leq t \leq T$ .

Finally, noting that  $\underline{\mu_t} \leq \mu_m \leq \overline{\mu}_t$  concludes the proof, and we have

$$\mu_t \in \left[\mu_m \left(1 + \frac{\underline{v}}{2} - \frac{2}{\underline{v}} \left(1 - \frac{Y}{N}\right)^{-1}\right), \mu_m \left(1 + \frac{2}{\underline{v}} \left(1 - \frac{Y}{N}\right)^{-1} - \frac{\underline{v}}{\underline{v}}\right)\right].$$

#### E.4 Proof of Theorem 4.3

In this section, we consider that all agents follow strategy G except for agent i in  $\mathcal{N}$  following strategy  $\beta \in \mathcal{B}^{CI}$  (we refer the reader to Section C.1 for the definition of  $\mathcal{B}^{CI}$ ). To simplify notations, we drop the superscript  $G_{-i}$  and write  $c_i^{\beta}, b_i^{\beta}, z_i^{\beta}$ , and  $g_i^{\beta}$  for the respective costs, bids, expenditures, and gains of agent i. This is not to be confused with  $c_i^G, b_i^G, z_i^G$  and  $g_i^G$ , which are the hypothetical variables of agent i, had it followed a G strategy (but all other agents still placing bids according to G strategies). Importantly, we use the superscripts  $\beta$  or G for the prices  $(p_m)_{m=1}^M$  of the different auctions instead of the superscript  $\gamma + 1$ . Since agent i does not follow  $G, \mu_{i,t}$  is a priory not defined. For convenience, we take the convention  $\mu_{i,t} = \mu_i^{*0}$  for all  $t \in \mathbb{N}$ .

Step 1: Lower Bound on the Total Expected Cost under Strategy  $\beta$ . We lower-bound the total expected cost suffered by agent *i* for following strategy  $\beta$  by considering it always gets a null cost after time  $\underline{T}$ .

$$C_{i}^{\beta,G_{-i}} \geq \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{t=1}^{\underline{T}} c_{i,t}^{\beta} \right]$$

$$\stackrel{(a)}{\geq} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{t=1}^{\underline{T}} c_{i,t}^{\beta} \right] + \mu_{i}^{\star 0} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{t=1}^{T} z_{i,t}^{\beta} - \rho_{i} - g_{i,t}^{\beta} \right]$$

$$\stackrel{(b)}{\geq} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{t=1}^{\underline{T}} c_{i,t}^{\beta} + \mu_{i}^{\star 0} (z_{i,t}^{\beta} - \rho_{i} - g_{i,t}^{\beta}) \right] - \left( \rho_{i} + \frac{M\gamma\Delta}{N\underline{\mu}} \right) \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ T - \underline{T} \right]$$

$$(121)$$

Inequality (a) uses that strategy  $\beta$  respects the budget constraint on average, and inequality (b) holds since  $g_{i,t}^{\beta} \leq \frac{M_Y \Delta}{N \mu}$ .

*Step 2: Lower Bound on the Lagrangian Expected Cost.* We bound the Lagrangian cost of agent *i* for the period  $t \leq \underline{\mathcal{T}}$ .

$$c_{i,t}^{\beta} + \mu_{i}^{\star 0}(z_{i,t}^{\beta} - \rho_{i} - g_{i,t}^{\beta}) = v_{i,t}\tau - \mathbb{1}\left\{b_{i,t}^{\beta} \ge d_{i,t}^{\gamma}\right\}(\Delta v_{i,t} - \mu_{i}^{\star 0}d_{i,t}^{\gamma}) - \mu_{i}^{\star 0}\rho_{i}$$

$$\ge v_{i,t}\tau - (\Delta v_{i,t} - \mu_{i}^{\star 0}d_{i,t}^{\gamma})^{+} - \mu_{i}^{\star 0}\rho_{i}$$
(122)

54

Taking expectation on Equation 79 and using the Lipschitz continuity of the dual objective function gives:

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \boldsymbol{c}_{i,t}^{\beta} + \boldsymbol{\mu}_{i}^{\star 0} (\boldsymbol{z}_{i,t}^{\beta} - \boldsymbol{\rho}_{i} - \boldsymbol{g}_{i,t}^{\beta}) \right] \stackrel{(a)}{\geq} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \Psi_{i}^{0} (\boldsymbol{\mu}_{t}^{\pi}) \right] + \boldsymbol{\mu}_{i}^{\star 0} \left( \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \boldsymbol{g}_{i,t}^{G} - \boldsymbol{g}_{i,t}^{\beta} \right] - \boldsymbol{\rho}_{i} \right), \\ \stackrel{(b)}{\geq} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \Psi_{i}^{0} (\boldsymbol{\mu}^{\star 0}) - \frac{\Delta}{\underline{\mu}} \left( \|\boldsymbol{a}_{i}\|_{2} + \frac{M\gamma\overline{\mu}}{N\underline{\mu}} \right) \|\boldsymbol{\mu}_{t}^{\pi} - \boldsymbol{\mu}^{\star 0}\|_{2} \right] \\ + \boldsymbol{\mu}_{i}^{\star 0} \left( \frac{\gamma}{N} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \boldsymbol{p}_{m,t}^{G} - \boldsymbol{p}_{m,t}^{\beta} \right] - \boldsymbol{\rho}_{i} \right), \\ \stackrel{(c)}{\geq} \Psi_{i}^{0} (\boldsymbol{\mu}^{\star 0}) - \frac{\Delta}{\underline{\mu}} \left( \|\boldsymbol{a}_{i}\|_{2} + \frac{M\gamma\overline{\mu}}{N\underline{\mu}} \right) \boldsymbol{s}_{t}^{1/2} - \overline{\mu} \left( \frac{\gamma\Delta}{N\underline{\mu}} + \boldsymbol{\rho}_{i} \right). \end{aligned}$$

$$(123)$$

Inequality (*a*) follows from Equation 122 by adding and subtracting  $g_{i,t}^G$  and using that  $\mu_{i,t} = \mu_i^*$  by convention. Inequality (*b*) uses the Lipschitz continuity of the dual function in Equation 108 and the expression of  $g_t = \frac{\gamma}{N} \sum_{m=1}^{M} p_{m,t}$ ; inequality (*c*) finally uses the contraction property of the projection under Assumption 17.1 combined with Jensen's inequality, as well as the bounds on the price  $p_m$  of any auction  $0 \le p_m \le \Delta/\mu$ .

*Step 3: Upper Bound on the Mean Squared Error.* For an agent  $j \neq i$ , Equation 92 (*a*) still holds and gives:

$$s_{j,t+1} \stackrel{(a)}{\leq} s_{j,t} + 2\epsilon \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \left( \mu_{j,t}^{\pi} - \mu_{j}^{\star 0} \right) \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ z_{j,t}^{\beta} - g_{j,t}^{\beta} | \boldsymbol{\mu}_{t}^{\pi} \right] \right] + \epsilon^{2} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ |z_{j,t}^{\beta} - g_{i,t}^{\beta}|^{2} \right],$$

$$\stackrel{(b)}{\leq} s_{j,t} + 2\epsilon \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \left( \mu_{j,t}^{\pi} - \mu_{j}^{\star 0} \right) L_{j}(\boldsymbol{\mu}_{t}^{\pi}) \right] + \epsilon^{2} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ |z_{j,t}^{\beta} - g_{i,t}^{\beta}|^{2} \right],$$

$$+ 2\epsilon \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \left| \mu_{j,t}^{\pi} - \mu_{j}^{\star 0} \right| \cdot \left| \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ z_{j,t}^{\beta} - g_{j,t}^{\beta} | \boldsymbol{\mu}_{t}^{\pi} \right] - L_{j}(\boldsymbol{\mu}_{t}^{\pi}) \right| \right].$$
(124)

Inequality (a) directly follows from inequality 92 (a); inequality (b) follows from adding and subtracting  $L_i(\mu_t^{\pi})$  in the second term of (a) and taking absolute value.

As Equations 93 and 94 are still valid when we sum over all agents  $j \neq i$ , it only remains to show what happens for the term in  $\mathbb{E}_{v,m}[z_{j,t}^{\beta} - g_{i,t}^{\beta}|\boldsymbol{\mu}_{t}^{\pi}]$ . Let  $d_{j,t}^{\gamma,G} = \gamma^{\text{th}} - \max_{k \neq j} \left\{ \mathbbm{1}\{m_{k,t} = m_{j,t}\} \frac{\Delta v_{k,t}}{\boldsymbol{\mu}_{k,t}^{\pi}} \right\}$ denote the competing price in the imaginary setting where agent *i* follows a strategy *G*, as opposed to the real competing price  $d_{i,t}^{\gamma,\beta}$ .

$$\left| \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \boldsymbol{z}_{j,t}^{\beta} - \boldsymbol{g}_{i,t}^{\beta} \big| \boldsymbol{\mu}_{t}^{\pi} \right] - L_{j}(\boldsymbol{\mu}_{t}^{\pi}) \right| \stackrel{(a)}{=} \left| \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \mathbbm{1} \left\{ \boldsymbol{m}_{j,t} = \boldsymbol{m}_{i,t} \right\} \left( \boldsymbol{d}_{j,t}^{\gamma,\beta} \mathbbm{1} \left\{ \frac{\Delta \boldsymbol{v}_{j,t}}{\boldsymbol{\mu}_{j,t}^{\pi}} \ge \boldsymbol{d}_{j,t}^{\gamma,\beta} \right\} \right. \\ \left. - \boldsymbol{d}_{j,t}^{\gamma,G} \mathbbm{1} \left\{ \frac{\Delta \boldsymbol{v}_{j,t}}{\boldsymbol{\mu}_{j,t}^{\pi}} \ge \boldsymbol{d}_{j,t}^{\gamma,G} \right\} \right) + \boldsymbol{g}_{j,t}^{G} - \boldsymbol{g}_{j,t}^{\beta} \right] \right|,$$

$$\left| \stackrel{(b)}{\leq} \frac{\Delta}{\mu} a_{i,j} + \left| \frac{\gamma}{N} \left( \boldsymbol{p}_{m_{j,t}}^{G} - \boldsymbol{p}_{m_{j,t}}^{\beta} \right) \right|,$$

$$\left| \stackrel{(c)}{\leq} \frac{\Delta}{\mu} a_{i,j} + \frac{\gamma\Delta}{N\mu}. \right|$$

$$(125)$$

Equality (a) uses that  $d_j^{\gamma,\beta} = d_j^{\gamma,G}$  when *i* plays in a different auction from *j*; inequality (b) holds since  $0 \le d_j^{\gamma} \le \Delta/\underline{\mu}$  and by rewriting  $g_t = \frac{\gamma}{N} \sum_{m=1}^{M} p_{m,t}$ . Inequality (c) uses that the price  $p_m$  of

any auction is bounded  $0 \le p_m \le \frac{\Delta}{\mu}$ . Summing Equation 125 over all  $j \ne i$  gives

$$\sum_{j\neq i} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \left| \mu_{j,t}^{\pi} - \mu_{j}^{\star 0} \right| \cdot \left| \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ z_{j,t}^{\beta} - g_{i,t}^{\beta} \right| \boldsymbol{\mu}_{t} \right] - L_{j}(\boldsymbol{\mu}_{t}) \right| \right] \leq \frac{\Delta}{\underline{\mu}} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{j\neq i} a_{i,j} \left| \mu_{j,t}^{\pi} - \mu_{j}^{\star 0} \right| \right] + \frac{\gamma \Delta}{N\underline{\mu}} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \left\| \boldsymbol{\mu}_{t}^{\pi} - \boldsymbol{\mu}^{\star} \right\|_{1} \right], \\ \stackrel{(a)}{\leq} \left( \frac{\Delta}{\underline{\mu}} \| \boldsymbol{a}_{i} \|_{2} + \frac{\gamma \Delta}{\sqrt{N\underline{\mu}}} \right), \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \left\| \boldsymbol{\mu}_{t}^{\pi} - \boldsymbol{\mu}^{\star} \right\|_{2} \right] \\ \stackrel{(b)}{\leq} \left( \frac{\Delta}{\underline{\mu}} \| \boldsymbol{a}_{i} \|_{2} + \frac{\gamma \Delta}{\sqrt{N\underline{\mu}}} \right) s_{t}^{1/2},$$

$$(126)$$

where inequality (*a*) follows from Cauchy-Schwartz and the relation between  $\|\cdot\|_1$  and  $\|\cdot\|_2$ , and (*b*) from Jensen's inequality.

Together Equations 93, 94, 124 and 126 give the following recursion:

$$s_{t+1} \leq (1-2\lambda\epsilon)s_t + 2\epsilon \left(\frac{\Delta}{\underline{\mu}} \|\boldsymbol{a}_i\|_2 + \frac{\gamma\Delta}{\sqrt{N}\underline{\mu}}\right)s_t^{1/2} + \frac{N\Delta^2\epsilon^2}{\underline{\mu}^2}$$

Since  $2\lambda \epsilon \leq 1$  under Assumption 17.2, and  $s_1 \leq N\overline{\mu}^2$  under Assumption 17.1, [6, Lemma C.5] yields

$$s_t \le N\overline{\mu}^2 (1 - \lambda\epsilon)^{t-1} + \frac{\Delta^2}{\lambda^2 \underline{\mu}^2} \left( \|\boldsymbol{a}_i\|_2^2 + \frac{\gamma^2}{N} \right) + \frac{N\Delta^2\epsilon}{\lambda \underline{\mu}^2}.$$
(127)

Summing up to  $\underline{T}$ , we finally get:

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{m}}\left[\sum_{t=1}^{\underline{\mathcal{T}}} s_{t}^{1/2}\right] \stackrel{(a)}{\leq} \sqrt{N\mu} \sum_{t=1}^{T} (1-\lambda\epsilon)^{(t-1)/2} + \left(\frac{\sqrt{N}\Delta\epsilon^{1/2}}{\sqrt{\lambda\mu}} + \frac{\Delta}{\lambda\mu} \left(\|\boldsymbol{a}_{i}\|_{2} + \frac{\gamma}{\sqrt{N}}\right)\right) T,$$

$$\stackrel{(b)}{\leq} \frac{2\sqrt{N\mu}}{\lambda\epsilon} + \left(\frac{\sqrt{N}\Delta\epsilon^{1/2}}{\sqrt{\lambda\mu}} + \frac{\Delta}{\lambda\mu} \left(\|\boldsymbol{a}_{i}\|_{2} + \frac{\gamma}{\sqrt{N}}\right)\right) T.$$
(128)

Inequality (a) holds since  $\sqrt{x+y} \le \sqrt{x} + \sqrt{y}$  for  $x, y \ge 0$ , and inequality (b) uses  $\underline{\mathcal{T}} \le T$ , bounds the geometric sum by the series' limit and uses that  $\frac{1}{1-(1-\lambda\epsilon)^{1/2}} \le \frac{2}{\lambda\epsilon}$  since  $1-(1-x)^{1/2} \ge x/2$  for  $x \in [0, 1]$ .

Step 4: Extension of the Proof of Theorem 4.2 to a Parallel Auction Setting. Note that Equations 110 and 111 still hold in a parallel auction setting if we also take expectation on *m*. Equation 108 further implies:

$$\Psi_i^0(\boldsymbol{\mu}_t^{\boldsymbol{\pi}}) \le \Psi_i^0(\boldsymbol{\mu}^{\star 0}) + \frac{\Delta}{\underline{\mu}} \left( \|\boldsymbol{a}_i\|_2 + \frac{M\gamma\overline{\mu}}{N\underline{\mu}} \right) \|\boldsymbol{\mu}_t^{\boldsymbol{\pi}} - \boldsymbol{\mu}^{\star 0}\|_2,$$
(129)

where we used that  $\mu_{i,t} = \mu_i^{\star 0}$  by convention. Similarly, using the definition of a stationary multiplier profile as well as Equation 109 gives:

$$-\mu_{i,t}^{\pi} L_{i}(\boldsymbol{\mu}_{t}^{\pi}) = \mu_{i,t}^{\pi} \left( L_{i}(\boldsymbol{\mu}^{\star 0}) - L_{i}(\boldsymbol{\mu}_{t}^{\pi}) \right),$$

$$\leq \frac{\Delta \overline{\mu}}{\underline{\mu}^{2}} \left( 8 \overline{\nu} \|\boldsymbol{a}_{i}\|_{2} + \frac{M \gamma}{N} \right) \|\boldsymbol{\mu}_{t}^{\pi} - \boldsymbol{\mu}^{\star 0}\|_{2}.$$
(130)

Combining Equations 111, 129 and 130 then gives the following bound on the expected cost per auction when all agents follow a strategy *G* in a parallel auction setting:

$$\mathbb{E}_{\boldsymbol{v},\boldsymbol{m}}\left[c_{i,t}^{G}\right] \leq \Psi_{i}^{0}(\boldsymbol{\mu}^{\star 0}) + \left(\frac{\Delta}{\underline{\mu}}\left(1 + \frac{8\overline{\nu\mu}}{\underline{\mu}}\right) \|\boldsymbol{a}_{i}\|_{2} + \frac{2M\gamma\Delta\overline{\mu}}{N\underline{\mu}^{2}}\right) \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}}\left[\|\boldsymbol{\mu}_{t}^{\boldsymbol{\pi}} - \boldsymbol{\mu}^{\star 0}\|_{2}\right],$$

$$\overset{(a)}{\leq} \Psi_{i}^{0}(\boldsymbol{\mu}^{\star 0}) + \left(\frac{\Delta}{\underline{\mu}}\left(1 + \frac{8\overline{\nu\mu}}{\underline{\mu}}\right) \|\boldsymbol{a}_{i}\|_{2} + \frac{2M\gamma\Delta\overline{\mu}}{N\underline{\mu}^{2}}\right) s_{t}^{1/2}.$$
(131)

Inequality (*a*) uses that all the  $\mu_i^{\star 0}$  lie inside the projection set under Assumption 17.1, Jensen's inequality, as well as the fact that the bound in Equation 127 is greater than that in Lemma E.1.

Conclusion. Together, Equations 110, 121, 123, 128 and 131 finally imply the following bound, i.e.,

$$\begin{split} C_{i}^{G} - C_{i}^{\beta,G_{-i}} &\stackrel{(a)}{\leq} \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{t=1}^{\underline{\Upsilon}} c_{i,t}^{G} - \left( c_{i,t}^{\beta} + \mu_{i}^{\star 0} (\boldsymbol{z}_{i,t}^{\beta} - \rho_{i} - \boldsymbol{g}_{i,t}^{\beta}) \right) \right] \\ &+ \left( \tau + \rho_{i} + \frac{M\gamma\Delta}{N\underline{\mu}} \right) \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ T - \underline{\Upsilon} \right], \\ &\stackrel{(b)}{\leq} \left( \frac{\Delta}{\underline{\mu}} \left( 2 + \frac{8\overline{\nu}\mu}{\underline{\mu}} \right) \|\boldsymbol{a}_{i}\|_{2} + \frac{3M\gamma\Delta\overline{\mu}}{N\underline{\mu}^{2}} \right) \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ \sum_{t=1}^{\underline{\Upsilon}} s_{t}^{1/2} \right] + \overline{\mu} \left( \frac{\gamma\Delta}{N\underline{\mu}} + \rho_{i} \right) T \\ &+ \left( \tau + \rho_{i} + \frac{M\gamma\Delta}{N\underline{\mu}} \right) \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ T - \underline{\Upsilon} \right], \\ &\stackrel{(c)}{\leq} \left( \frac{\Delta}{\underline{\mu}} \left( 2 + \frac{8\overline{\nu}\mu}{\underline{\mu}} \right) \|\boldsymbol{a}_{i}\|_{2} + \frac{3M\gamma\Delta\overline{\mu}}{N\underline{\mu}^{2}} \right) \left( \frac{2\sqrt{N}\overline{\mu}}{\lambda\epsilon} + \left( \frac{\sqrt{N}\Delta\epsilon^{1/2}}{\sqrt{\lambda}\underline{\mu}} \right) \\ &+ \frac{\Delta}{\lambda\underline{\mu}} \left( \|\boldsymbol{a}_{i}\|_{2} + \frac{\gamma}{\sqrt{N}} \right) \right) T \right) + \overline{\mu} \left( \frac{\gamma\Delta}{N\underline{\mu}} + \rho_{i} \right) T + \left( \tau + \rho_{i} + \frac{M\gamma\Delta}{N\underline{\mu}} \right) \mathbb{E}_{\boldsymbol{v},\boldsymbol{m}} \left[ T - \underline{\Upsilon} \right] \end{split}$$

where inequality (a) uses Equations 110 and 121, inequality (b) Equations 123 and 131, and inequality (b) Equation 128.

Rewriting  $\rho = \underline{k_1}/T$ , this proves the existence of a constant *C* in  $\mathbb{R}_+$  such that

$$\begin{split} \frac{1}{T} \left( C_i^G - C_i^{\beta, G_{-i}} \right) &\leq C \Biggl( \left( \| \boldsymbol{a}_i \|_2 + \frac{M\gamma}{N} \right) \left( \sqrt{N\epsilon} \left( 1 + \frac{1}{\epsilon^{3/2} T} \right) + \| \boldsymbol{a}_i \|_2 + \frac{\gamma}{\sqrt{N}} \right) \\ &+ \left( \frac{\gamma}{N} + \frac{\overline{k_1}}{T} \right) + \left( \frac{\overline{k_1}}{T} + \frac{M\gamma}{N} \right) \frac{\mathbb{E}_{\boldsymbol{v}, \boldsymbol{m}} \left[ T - \underline{\mathcal{T}} \right]}{T} \Biggr). \end{split}$$

## F PROOFS OF THE SECONDARY RESULTS OF SECTION 4

We gather in this section the proofs of the secondary results from Section 4.

### F.1 Sufficient Conditions for Assumption 13.

Similarly to Section D.1, we prove in this section that the technical Assumption 2 is satisfied under the mild differentiability and continuity conditions on the valuation and competing bid distributions of Assumptions 1 and 15. This shows that the conditions of Theorem 4.1 are not restrictive.

Recall the definitions of the dual function  $\Psi^0(\mu) = \mathbb{E}_{v,d} \left[ v\tau - \mu g - (\Delta v - \mu d^{\gamma})^+ \right]$ , the expenditure function  $Z(\mu) = \mathbb{E}_{v,d} \left[ d^{\gamma} \mathbb{1} \{ \Delta v > \mu d^{\gamma} \} \right]$ , the gain function  $G(\mu) = \mathbb{E}_{v,d} \left[ g \right]$ , as well as the loss function  $L(\mu) = Z(\mu) - G(\mu)$ .

Note first that  $\Psi^0(\mu) = \Psi(\mu) + \mu \rho - \mu G(\mu)$ , where  $\Psi$  refers to the dual function of Section 3.2. Since the expenditure function is moreover equivalent in both frameworks, the results of D.1 still hold. Hence we only need to prove the differentiability of *G*, bound its derivatives, and show the strong monotonicity of *L*.

The function  $\mu \mapsto \frac{\gamma}{N}p$  is differentiable with derivative  $\mu \mapsto -\frac{\gamma\Delta v}{N\mu^2}\mathbb{1}\left\{d^{\gamma} > \frac{\Delta v}{\mu} > d^{\gamma+1}\right\}$ , except in the sets  $\left\{d^{\gamma} = \Delta v/\mu \ge d^{\gamma+1}\right\}$  and  $\left\{d^{\gamma} \ge \Delta v/\mu = d^{\gamma+1}\right\}$  of measure zero since valuations and competing prices are absolutely continuous under Assumptions 1 and 15. Since the derivative is bounded by  $\frac{\gamma\Delta}{N\mu^2}$  which is integrable, Leibniz's integral rule implies that the gain function *G* is differentiable, i.e.,

$$G'(\mu) = -\frac{\gamma}{N} \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ \frac{\Delta \boldsymbol{v}}{\mu^2} \mathbb{1} \left\{ d^{\gamma} > \Delta \boldsymbol{v}/\mu > d^{\gamma+1} \right\} \right].$$
(132)

In particular, G' is negative on  $[\mu, \overline{\mu}]$ .

We now carefully bound G'. Let  $H : d^{\gamma} \times d^{\gamma+1} \mapsto H(d^{\gamma}, d^{\gamma+1})$  denote the cumulative probability function of competing bids.

$$\begin{aligned} \left| G'(\mu) \right| &= \frac{\gamma}{N} \int_{x=0}^{\Delta/\underline{\mu}} \int_{y=0}^{\Delta/\underline{\mu}} \int_{z=\mu y/\Delta}^{\mu x/\Delta} v(z) dH(x,y) dz \\ &\stackrel{(a)}{\leq} \frac{\gamma}{N} \frac{\overline{\mu} v}{\Delta} \int_{x=0}^{\Delta/\underline{\mu}} \int_{y=0}^{\Delta/\underline{\mu}} (x-y) dH(x,y) \\ &= \frac{\gamma}{N} \frac{\overline{\mu} v}{\Delta} \mathbb{E}_{v,d} \left[ d^{\gamma} - d^{\gamma+1} \right] \end{aligned}$$
(133)

where inequality (a) uses Assumption 1.

Combining Equation 91 with Equation 133 and the fact that G' is negative on  $[\mu, \overline{\mu}]$  yields

$$L^{'}(\mu) = Z^{'}(\mu) - G^{'}(\mu) \leq \frac{\overline{\mu\nu}}{\Delta} \frac{\gamma}{N} \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ d^{\gamma} - d^{\gamma+1} \right] - \frac{\underline{\nu} \underline{h} \Delta^{3}}{3\overline{\mu}^{3}} \stackrel{(a)}{<} 0,$$

where inequality (*a*) follows from Assumption 15. This proves the strong monotonicity of the loss function L.

Furthermore, the function  $\mu \mapsto -\frac{\gamma \Delta v}{N\mu^2} \mathbb{1} \left\{ d^{\gamma} > \Delta v/\mu > d^{\gamma+1} \right\}$  is differentiable outside of the same measure zero sets, and its derivative, bounded by  $\frac{3\gamma\Delta}{N\mu^3}$ , is still integrable. Leibniz's integral rule hence implies that G' is also differentiable, i.e.,

$$\left| G^{''}(\mu) \right| = \left| \frac{3\gamma\Delta}{N\mu^3} \mathbb{E}_{\boldsymbol{v},\boldsymbol{d}} \left[ \boldsymbol{v}\mathbbm{1} \left\{ d^{\gamma} > \Delta \boldsymbol{v}/\mu > d^{\gamma+1} \right\} \right] \right| \leq \frac{3\Delta}{\underline{\mu}^3}$$

where we used that the fraction of winning agents  $\frac{\gamma}{N}$  is smaller than one.

Combining Equation 88 with the fact that  $g_t$  is itself bounded and Leibniz's rule, we finally get that the dual function  $\Psi^0$  is differentiable with derivative

$$\begin{split} \Psi^{0'}(\mu) &= d^{\gamma} \mathbb{1}\{\Delta v > \mu d^{\gamma}\} - g_t - \mu \frac{\partial \mathbb{E}_{v,d}[g]}{\partial \mu}(\mu), \\ &= Z(\mu) - \mu \frac{\gamma \Delta v}{N\mu^2} \mathbb{P}\left[d^{\gamma} > \Delta v/\mu > d^{\gamma+1}\right]. \end{split}$$