

WILF-ZEILBERGER SEEDS: q -ANALOGUES

KAM CHEONG AU

ABSTRACT. We point out a correspondence between many classical and q -hypergeometric Wilf-Zeilberger seeds, it enables us to find q -analogues of many classical hypergeometric series identities, and offers a conceptual explanation to why such q -analogue should exist.

Finding q -analogue of classical hypergeometric identities is an active and interesting area of research. While q -analogues of classical hypergeometric summation formulas were well-known, only isolated examples are known for exotic hypergeometric sums such as

$$\sum_{n \geq 1} \left(-\frac{1}{2^{10}}\right)^n \frac{(1)_n^5}{n^5 (1/2)_n^5} (205n^2 - 160n + 32) = -2\zeta(3)$$

$$\sum_{n \geq 0} \frac{(-1)^n}{2^{2n}} \frac{(\frac{1}{2})_n^5}{(1)_n^5} (20n^2 + 8n + 1) = \frac{8}{\pi^2}$$

coming from Wilf-Zeilberger (WZ) pairs. After introducing q -version of a concept called *Wilf-Zeilberger seed* (whose classical version discussed in author's previous article [2]), we are able to derive q -analogue of such summations, for example, above two instances have q -analogue

$$\sum_{k \geq 0} \frac{q^k (1 + q^{1+k})}{(1 - q^{1+k})^3} = \sum_{n \geq 1} \frac{(-1)^{n-1} q^{5n(n-1)/2} (10q^{2n} + 10q^{3n} + 5q^{4n} + q^{5n} - 5q^{3n-1} - 15q^{4n-1} + q^{5n-2} - 24q^{5n-1} + 5q^{6n-2} - 15q^{6n-1} + 10q^{7n-2} - 5q^{7n-1} + 10q^{8n-2} + 5q^{9n-2} + q^{10n-2} + 5q^n + 1)(q; q)_n^{10}}{(1 - q^n)^5 (q; q)_{2n}^5}$$

for first equality and

$$\sum_{n \geq 0} \frac{(-1)^n q^{n^2} (q^{2n+1} - 4q^{4n+1} + q^{6n+1} + q^{8n+2} + 1)(q; q^2)_n^6}{(q^{2n+1} + 1)(q^2; q^2)_{2n} (q^2; q^2)_n^4} = \frac{(1 - q^2)^2}{\Gamma_{q^2}(1/2)^4}$$

for second equality.

Conceptually, they possess q -analogues because a q -version already exists at the level of WZ-seeds. This allows us to derive classical and q -hypergeometric identity in striking parallel: through numerous examples we will see many q -identities can be proved in much the same way as their classical counterparts.

Up to know, q -analogues of such formulas have been found in isolated instance with *ad hoc* principle. ([12], [9], [11], [18], [5]) We will soon see how many such examples can be unified using the language of WZ-seeds.

One motivation in q -analogues of these identities is that, via a technique called "creative microscoping", it could help us prove corresponding super-congruence, see [13] for an illustrative example, as well as [20], [14] and [10].

We put an emphasis on deriving q -analogues of Ramanujan's $1/\pi^k$ -formulas, although the recipe

2020 *Mathematics Subject Classification*. Primary: 11B65, 33D15, 33C20. Secondary: 33F10.

Key words and phrases. WZ-pair, basic hypergeometric series, q -series, q -analogues, Ramanujan $1/\pi$ series.

works well for many other classical constants. For example, using two different WZ-seeds, we derive two q -analogues for $\sum_{n \geq 0} (\frac{1}{4})^n \frac{(\frac{1}{2})_n^3}{(1)_n^3} (6n+1) = \frac{4}{\pi}$ (Examples IV, VII):

$$\begin{aligned} \sum_{n \geq 0} \frac{q^{n^2} (1 - q^{1+6n})(q; q^2)_n^2 (q^2; q^4)_n}{(q^4; q^4)_n^3} &= \frac{1 - q^4}{\Gamma_{q^4}(1/2)^2} \\ \sum_{n \geq 0} \frac{q^{2n^2} (1 + q^{2n+1} - 2q^{4n+1})(q; q^2)_n^4}{(q^{2n+1} + 1)(q^2; q^2)_{2n} (q^2; q^2)_n^2} &= \frac{1 - q^2}{\Gamma_{q^2}(1/2)^2} \end{aligned}$$

the first one is already known in [11]. Actually, WZ-seeds carry auxiliary variables that we can put into formulas, for instance, first identity above is specialization under $b = c = d = 0$ of the following:

$$\begin{aligned} (0.1) \quad \sum_{n \geq 0} \frac{q^{-2bn+cn+\frac{a^2}{2}} (1 - q^{c+2d+3n+\frac{1}{2}})(q^{\frac{1}{2}-c}, q^{c+\frac{1}{2}}, q^{c+2d+\frac{1}{2}}; q)_n (q^{2b+2d+1}; q^2)_n}{(q^2, q^{2d+2}, q^{2c+2d+2}; q^2)_n (q^{-2b+c+\frac{1}{2}}; q)_n} \\ = \frac{(1-q)(1+q)^{2d+1} \Gamma_{q^2}(1-b) \Gamma_{q^2}(d+1) \Gamma_q(-2b+c+\frac{1}{2}) \Gamma_{q^2}(c+d+1)}{\Gamma_{q^2}(\frac{1}{2}) \Gamma_q(1-2b) \Gamma_{q^2}(-b+c+\frac{1}{2}) \Gamma_q(c+2d+\frac{1}{2})} \end{aligned}$$

Even classical version of this identity seems not known, nonetheless, both classical and q -version can be derived in one go.

We shall also write down q -analogues for four Ramanujan $1/\pi^2$ -formulas in Examples XIV and XV.

Acknowledgements The author would like to express his sincere gratitude to W. Zudilin, J. Guillera and Z.W. Sun for valuable discussions. A special thank to Wadim for encouraging me to continue mathematical research during an uneasy phase of life.

1. q -VERSION OF WILF-ZEILBERGER SEEDS

1.1. A source of WZ-pairs. We juxtapose the two cases: classical hypergeometric and q -hypergeometric case. A term $f(n)$ is called hypergeometric (resp. q -hypergeometric) if $f(n+1)/f(n)$ is a rational function in n (resp. q^n), with coefficient in a understood field. A WZ-pair (F, G) is any two-variable function satisfying

$$F(n+1, k) - F(n, k) = G(n, k+1) - G(n, k)$$

For proper classical hypergeometric or q -hypergeometric term $F(n, k)$, there exists a well-known algorithm (Gosper's algorithm) to find its WZ-mate $G(n, k)$ if it exists, or prove it does not exist. It is not easy to come up with non-trivial WZ-pairs, we shall outline a such a method below.

First we need the following definition:

Definition 1.1. Let $f(a_1, \dots, a_m, k)$ be a (classical or q) hypergeometric term in a_i and k , it is called a *WZ-seed* in variables a_i if for each $A_i \in \mathbb{Z}$ and $K \in \mathbb{Z}$,

$$F(n, k) = f(a_1 + A_1 n, a_2 + A_2 n, \dots, a_m + A_m n, k + K n)$$

has a (classical or q) hypergeometric WZ-mate $G(n, k)$.

Classical version of following empirical observation was already explicated in [2]:

Problem 1.2 (Hypergeometric summation formula induces WZ-candidate). Let $f(a_1, \dots, a_m; k)$ be a proper (classical or q) hypergeometric term in a_i, k . If

$$\sum_{k \geq 0} f(a_1, \dots, a_m, k)$$

exists and is independent of a_i , then is it true that, $f(a_1, \dots, a_m; k)$ is a WZ-seed?

The answer seems to be affirmative for all examples we have computed. For example, consider the Gauss ${}_2F_1$ formula:

$${}_2F_1 \left[\begin{matrix} a & b \\ c \end{matrix}; 1 \right] = \sum_{k \geq 0} \frac{(a)_k (b)_k}{(1)_k (c)_k} = \frac{\Gamma(c-a-b)\Gamma(c)}{\Gamma(c-a)\Gamma(c-b)}$$

divide the gamma factor of RHS into LHS summand, we arrive:

$$\sum_{k \geq 0} \frac{\Gamma(c-a)\Gamma(a+k)\Gamma(c-b)\Gamma(b+k)}{\Gamma(a)\Gamma(b)\Gamma(k+1)\Gamma(c+k)\Gamma(-a-b+c)} = 1$$

If we denote the summand by $\text{Gauss2F1}(a, b, c, k)$, then the above observation would predict

$$F(n, k) = \text{Gauss2F1}(a-n, b, c+n, k+n)$$

has a WZ-mate G that is also proper hypergeometric, indeed, Gosper's algorithm returns

$$\begin{aligned} G(n, k) = F(n, k) \times & (-a^2b+a^2c+a^2n+2abc+abk+5abn+ab-2ac^2-2ack-8acn-ac-ak^2-4akn-7an^2-an \\ & -bc^2-2bck-6bcn-bc-bk^2-5bkn-8bn^2-2bn+c^3+2c^2k+7c^2n+c^2+ck^2+8ckn+ck+15cn^2+4cn+2k^2n+k^2 \\ & +8kn^2+3kn+10n^3+4n^2) \div ((c+k+2n)(a+b-c-2n-1)(a+b-c-2n)) \end{aligned}$$

satisfies $F(n+1, k) - F(n, k) = G(n, k+1) - G(n, k)$.

Now consider its q -version, let $0 < q < 1$ (an assumption we always enforce in this article):

$$\sum_{k \geq 0} \frac{(q^a, q^b; q)_k}{(q^c, q; q)_k} q^{k(c-b-a)} = \frac{(q^{c-a}, q^{c-b}; q)_\infty}{(q^c, q^{c-a-b}; q)_\infty}$$

rewrite it into q -gamma function:

$$\sum_{k \geq 0} \frac{q^{k(-a-b+c)} \Gamma_q(c-a) \Gamma_q(a+k) \Gamma_q(c-b) \Gamma_q(b+k)}{\Gamma_q(a) \Gamma_q(b) \Gamma_q(k+1) \Gamma_q(c+k) \Gamma_q(-a-b+c)} = 1$$

denote the summand by $\text{Gauss2F1}_q(a, b, c, k)$, note that $\lim_{q \rightarrow 1^-} \text{Gauss2F1}_q = \text{Gauss2F1}$. q -version of above observation would predict

$$F_q(n, k) = \text{Gauss2F1}_q(a-n, b, c+n, k+n)$$

has a WZ-mate G that is also proper q -hypergeometric, indeed, q -Gosper's algorithm gives

$$(1.1) \quad G_q(n, k) = F_q(n, k) \times \frac{-q^{2a+b+c+k+2n} - q^{a+b+c+k+3n} + q^{2a+b+c+2k+3n} - q^{a+b+c+2k+4n+1} + q^{2a+b} - q^{2a+c+k+2n} + q^{a+c+k+3n+1} + q^{a+2c+k+4n} + q^{a+2c+k+4n+1} - q^{a+c+2n+1} + q^{b+2c+2k+6n+1} - q^{3c+2k+7n+1}}{(1 - q^{c+k+2n})(q^{a+b} - q^{c+2n})(q^{a+b} - q^{c+2n+1})}$$

and we have $F_q(n+1, k) - F_q(n, k) = G_q(n, k+1) - G_q(n, k)$. Note that

$$\lim_{q \rightarrow 1^-} F_q(n, k) = F(n, k) \quad \lim_{q \rightarrow 1^-} G_q(n, k) = G(n, k)$$

define above.

We remark Problem 1.2 has a refinement that seems also true: if $\sum_{k \geq 0} f(a_1, \dots, a_m, k)$ is independent for general a_2, \dots, a_m but only for $a_1 \in \mathbb{N}$ and *the sum is terminating*, then $f(a_1, \dots, a_m, k)$ should also be a WZ-seed. As an example, we consider the terminating 1-balanced ${}_3\phi_2$ summation formula

$$\sum_{k \geq 0} \frac{(q^{-d}, q^{a-d}, q^{1+a-b-c}; q)_k}{(q, q^{1+a-b}, q^{1+a-c}; q)_k} q^k = \frac{(q^b, q^c; q)_d}{(q^{1+a-b}, q^{1+a-c}; q)_d} q^{d(1+a-b-c)} \quad d \in \mathbb{N}$$

the refinement predicts

$$f(a, b, c, d, k) = \frac{(q^{-d}, q^{a-d}, q^{1+a-b-c}; q)_k}{(q, q^{1+a-b}, q^{1+a-c}; q)_k} q^k \left(\frac{(q^b, q^c; q)_d}{(q^{1+a-b}, q^{1+a-c}; q)_d} q^{d(1+a-b-c)} \right)^{-1}$$

should be a WZ-seed with accessory parameters a, b, c, d . One can quickly convince himself this is indeed so by trying several sets of accessory parameters. This f is the WZ-seed labelled **Balanced3F2** below.

1.2. List of WZ-seeds. We remark that the following WZ-seeds listed should be treated as empirical: for all choices of accessory parameters A_i, K we have ever tried to run q -Gosper's algorithm on, the existence of $G(n, k)$ is always positive. A rigorous demonstration that they are really WZ-seeds will be postpone to a later article.

The list is also by no mean exhaustive, many other seeds which do not generate useful examples are not included. If Seed_q is one of seed listed below, then $\text{Seed} := \lim_{q \rightarrow 1^+} \text{Seed}_q$ exists and is a classical hypergeometric WZ-seed, it is then identical to those listed in [2].

(1) Gauss2F1 $_q(a, b, c, k)$

$$= \frac{q^{k(-a-b+c)} \Gamma_q(c-a) \Gamma_q(a+k) \Gamma_q(c-b) \Gamma_q(b+k)}{\Gamma_q(a) \Gamma_q(b) \Gamma_q(k+1) \Gamma_q(c+k) \Gamma_q(-a-b+c)}$$

Origin: q -analogue of ${}_2F_1$ summation formula.

(2) Dixon3F2 $_q(a, b, c, k)$

$$= \frac{q^{k(a-b-c+1)} (q^{a+k} + 1) \Gamma_q(a-b+1) \Gamma_q(a-c+1) \Gamma_q(2a+k) \Gamma_q(b+k) \Gamma_q(c+k) \Gamma_q(2a-b-c+1)}{\Gamma_q(a) \Gamma_q(b) \Gamma_q(c) \Gamma_q(k+1) \Gamma_q(a-b-c+1) \Gamma_q(2a-b+k+1) \Gamma_q(2a-c+k+1)}$$

Origin: q -analogue of Dixon's ${}_3F_2$ summation formula.

(3) Dougall5F4 $_q(a, b, c, d, k)$

$$= \frac{q^{k(a-b-c-d+1)} (1-q^{a+2k}) \Gamma_q(a+k) \Gamma_q(b+k) \Gamma_q(c+k) \Gamma_q(d+k) \Gamma_q(a-b-c+1) \Gamma_q(a-b-d+1) \Gamma_q(a-c-d+1)}{(1-q) \Gamma_q(b) \Gamma_q(c) \Gamma_q(d) \Gamma_q(k+1) \Gamma_q(a-b+k+1) \Gamma_q(a-c+k+1) \Gamma_q(a-d+k+1) \Gamma_q(a-b-c-d+1)}$$

Origin: q -analogue of very-well-poised ${}_5F_4$ summation formula.

(4) Dougall7F6 $_q(a, b, c, d, e, k)$

$$= \frac{(-1)^{a-e} (1-q^{a+2k}) q^{\frac{1}{2}(a-e)(a-e+1)+k} \Gamma_q(a+k) \Gamma_q(b+k) \Gamma_q(c+k) \Gamma_q(d+k) \Gamma_q(e+k) \Gamma_q(a-b-c+1)}{(1-q) \Gamma_q(b) \Gamma_q(c) \Gamma_q(d) \Gamma_q(e) \Gamma_q(k+1) \Gamma_q(-a+b+e) \Gamma_q(a-b+k+1) \Gamma_q(-a+c+e) \Gamma_q(a-c+k+1)} \\ \Gamma_q(-a+d+e) \Gamma_q(a-d+k+1) \Gamma_q(a-e+k+1) \Gamma_q(a-b-c-d+1) \Gamma_q(2a-b-c-d-e+1) \Gamma_q(-a+b+c+d+e+k)$$

Origin: q -analogue of very-well-poised 2-balanced terminating ${}_7F_6$ summation formula.

(5) Balanced3F2 $_q(a, b, c, d, k)$

$$= \frac{(-1)^{a+b+c-d} q^{\frac{1}{2}(a+b+c-d)(a+b+c-d+1)+k} \Gamma_q(d-a) \Gamma_q(a+k) \Gamma_q(d-b) \Gamma_q(b+k) \Gamma_q(d-c) \Gamma_q(c+k)}{\Gamma_q(a) \Gamma_q(b) \Gamma_q(c) \Gamma_q(k+1) \Gamma_q(d+k) \Gamma_q(-a-b+d) \Gamma_q(-a-c+d) \Gamma_q(-b-c+d) \Gamma_q(a+b+c-d+k+1)}$$

Origin: 1-balanced terminating ${}_3F_2$ summation formula.

(6) Seed $1_q(a, c, k)$

$$= \frac{q^{-2ak+2ck-k}(q+1)^{a-2c-2k}\Gamma_q(-a+2c-1)\Gamma_q(a+2k)}{\Gamma_q(a)\Gamma_{q^2}(k+1)\Gamma_{q^2}(-a+c-\frac{1}{2})\Gamma_{q^2}(c+k)}$$

Origin: put $a \rightarrow a/2, b \rightarrow a/2 + 1/2$ in Gauss2F 1_q .(7) Seed $2_q(a, c, d, k)$

$$= \frac{(-1)^{a+c-d}(q+1)^{-2c-2k}\Gamma_q(-a+2d-1)\Gamma_q(a+2k)\Gamma_{q^2}(d-c)\Gamma_{q^2}(c+k)q^{a^2+2ac-2ad+2a+c^2-2cd+2c+d^2-2d+2k}}{\Gamma_q(a)\Gamma_{q^2}(c)\Gamma_{q^2}(k+1)\Gamma_{q^2}(-a+d-\frac{1}{2})\Gamma_{q^2}(d+k)\Gamma_q(-a-2c+2d-1)\Gamma_{q^2}(a+c-d+k+\frac{3}{2})}$$

Origin: put $a \rightarrow a/2, b \rightarrow a/2 + 1/2$ in Balanced3F 2_q .(8) Seed $3_q(a, b, d, k)$

$$= \frac{(1+q)^{2d}(1-q^{2a+4k})q^{2ak-2bk-2dk+k}\Gamma_{q^2}(a-b+\frac{1}{2})\Gamma_{q^2}(a+k)\Gamma_q(b+2k)\Gamma_{q^2}(d+k)\Gamma_q(2a-b-2d+1)}{(1-q)\Gamma_q(b)\Gamma_{q^2}(d)\Gamma_{q^2}(k+1)\Gamma_{q^2}(a-b-d+\frac{1}{2})\Gamma_q(2a-b+2k+1)\Gamma_{q^2}(a-d+k+1)}$$

Origin: put $b \rightarrow b/2, c \rightarrow b/2 + 1/2$ in Dougall5F 4_q .(9) Seed $4_q(a, b, k)$

$$= \frac{(q^2+q+1)^{b+1}(1-q^{3a+6k})q^{3ak-3bk}\Gamma_q(3a-2b)\Gamma_{q^3}(a+k)\Gamma_q(b+3k)}{(1-q^3)\Gamma_q(b)\Gamma_{q^3}(k+1)\Gamma_{q^3}(a-b)\Gamma_q(3a-b+3k+1)}$$

Origin: put $b \rightarrow b/3, c \rightarrow b/3 + 1/3, d \rightarrow b/3 + 2/3$ in Dougall5F 4_q .(10) Seed $7_q(a, b, d, e, k)$

$$= \frac{(-1)^{a-e}q^{a^2-2ae+a+e^2-e+2k}(1-q^{2a+4k})\Gamma_{q^2}(a-b+\frac{1}{2})\Gamma_{q^2}(a+k)\Gamma_q(b+2k)}{(1-q^2)\Gamma_q(b)\Gamma_{q^2}(d)\Gamma_{q^2}(e)\Gamma_{q^2}(k+1)\Gamma_{q^2}(a-b-d+\frac{1}{2})\Gamma_q(-2a+b+2e)\Gamma_q(2a-b+2k+1)\Gamma_{q^2}(-a+d+e)} \frac{\Gamma_{q^2}(d+k)\Gamma_{q^2}(e+k)\Gamma_q(2a-b-2d+1)\Gamma_{q^2}(-a+b+e+\frac{1}{2})\Gamma_q(-2a+b+2d+2e)\Gamma_{q^2}(2a-b-d-e+k+\frac{1}{2})}{\Gamma_{q^2}(a-d+k+1)\Gamma_{q^2}(a-e+k+1)\Gamma_{q^2}(2a-b-d-e+\frac{1}{2})\Gamma_{q^2}(-a+b+d+e+k+\frac{1}{2})}$$

Origin: put $b \rightarrow b/2, c \rightarrow b/2 + 1/2$ in Dougall7F 6_q .(11) Seed $9_q(a, b, d, k)$

$$= \frac{(-1)^b q^{2ab-b^2-bd+2k}(1-q^{2(a+2k)})(q+1)^{4a-2b-2d}\Gamma_{q^2}(a-b+\frac{1}{2})\Gamma_{q^2}(a+k)\Gamma_q(b+2k)}{(1-q)\Gamma_q(b)\Gamma_q(d)\Gamma_{q^2}(k+1)\Gamma_{q^2}(-a+d+\frac{1}{2})\Gamma_{q^2}(2a-b-d)} \frac{\Gamma_q(d+2k)\Gamma_q(2a-b-d+1)\Gamma_q(-2a+2b+d+1)\Gamma_q(-2a+b+2d+1)\Gamma_{q^2}(2a-b-d+k)}{\Gamma_q(-2a+b+d+1)\Gamma_q(2a-b+2k+1)\Gamma_q(2a-d+2k+1)\Gamma_{q^2}(-a+b+d+k+1)}$$

Origin: put $b \rightarrow b/2, c \rightarrow b/2 + 1/2, d \rightarrow d/2, e \rightarrow d/2 + 1/2$ in Dougall7F 6_q .(12) Seed $10_q(a, d, k)$

$$= \frac{(-1)^{a-d}(q^2+q+1)^{-a-3k}q^{\frac{3a^2}{2}-3ad+\frac{9a}{2}+\frac{3d^2}{2}-\frac{9d}{2}+3k}\Gamma_q(-a+3d-2)\Gamma_q(a+3k)}{\Gamma_q(a)\Gamma_{q^3}(k+1)\Gamma_q(-2a+3d-3)\Gamma_{q^3}(d+k)\Gamma_{q^3}(a-d+k+2)}$$

Origin: put $a \rightarrow a/3, b \rightarrow a/3 + 1/3, c \rightarrow a/3 + 2/3$ in Balanced3F 2_q .

All classical seeds found in [2] have their q -generalization in above list. But this does not mean we can assert all WZ-seeds have q -analogues. Consider the following WZ-seed (with auxiliary variable a):

$$f(a, k) = \frac{5^{5a/2}(\sqrt{5}+1)^{-3a}(1-\sqrt{5})^k 2^{a-k}\Gamma(a+\frac{2}{5})\Gamma(a+\frac{3}{5})\Gamma(-2a+k+1)\Gamma(2a+k)}{\Gamma(1-2a)\Gamma(a+\frac{1}{2})\Gamma(k+1)\Gamma(3a+k)}$$

it comes from the ${}_2F_1$ summation formula (cf. [6] or [4]):

$${}_2F_1\left[\begin{matrix} 1-2a, 2a \\ 3a \end{matrix}; 1-\phi\right] = \frac{(\sqrt{5}+3)\pi 5^{\frac{1}{4}-\frac{5a}{2}}\phi^{3a-\frac{5}{2}}\Gamma(3a)}{\Gamma(a)\Gamma(a+\frac{2}{5})\Gamma(a+\frac{3}{5})} \quad \phi = (1+\sqrt{5})/2$$

due to the irrational exponent, it is hard to conceive what a possible q -analogue of $f(a, k)$ might look like.

Before we astonishing ourselves with wonderful consequences of WZ-seeds, it is important to say something on the computational aspect, since most of manipulations in this article is not hand do-able.

In all our examples, full expression for $G(n, k)$ is complicated, we choose not to write it out. Readers are suggested to either trust the correctness of computations or perform Gosper's algorithm themselves.

For finding WZ-mate $G(n, k)$, we use three Mathematica implementations that are part of a bundle developed by RISC:

- For classical hypergeometric case: `fastZeil` written by P. Paule, M. Schorn, and A. Riese ([17]).
- For q -case: `qZeil` by A. Riese ([17]) and `HolonomicFunctions` by C. Koutschan ([15]).¹

Mathematica codes allowing readers to replicate examples below (and discovering new ones) will be made available.

1.3. Two introductory examples. The following proposition is an important bridge between WZ-pairs and infinite series.

Proposition 1.3 ([2]). *Suppose $F, G : \mathbb{N}^2 \rightarrow \mathbb{C}$ are two functions such that*

$$F(n+1, k) - F(n, k) = G(n, k+1) - G(n, k)$$

such that

- $\sum_{k \geq 0} F(0, k)$ converges
- $\sum_{n \geq 0} G(n, 0)$ converges
- $\lim_{k \rightarrow \infty} G(n, k) = g(n)$ exists for each $n \in \mathbb{N}$ and $\sum_{n \geq 0} g(n)$ converges

then $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$ exists and is finite, also

$$\sum_{k \geq 0} F(0, k) + \sum_{n \geq 0} g(n) = \sum_{n \geq 0} G(n, 0) + \lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$$

We now give two simple examples, in which $g(n)$ and $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$ can be shown to be zero, they would be justified in next section.

Example I. Let a, b, c, d be in a neighbourhood of 0. We start with a hypergeometric WZ-seed

$$F(n, k) = \text{Gauss2F1}(a - d - n, b - d - n, c - d + 1, d + k + 2n + 1)$$

call Gosper's algorithm to compute $G(n, k)$ (which we do not display here) then applying 1.3, we obtain,

$$\sum_{k \geq 0} F(0, k) = \sum_{n \geq 0} G(n, 0)$$

, converting gamma functions into pochhammer symbols becomes:

¹`qZeil` is faster, but sometime raises cryptic errors for complicated input, and seems not to be actively maintained any more. `HolonomicFunctions` is bug-free and better developed, but computation with it is many hundred times slower than `qZeil` and `fastZeil`

$$\begin{aligned}
(\text{ex1}) \quad & \sum_{k \geq 0} \frac{(1+a)_k (1+b)_k}{(1+c)_{k+1} (1+d)_{k+1}} \\
& = \sum_{n \geq 1} \frac{(1+a)_{n-1} (1+b)_{n-1} (1-a+c)_{n-1} (1-b+c)_{n-1} (1-a+d)_{n-1} (1-b+d)_{n-1} P(n)}{(1+c)_{2n} (1+d)_{2n} (1-a-b+c+d)_{2n}}
\end{aligned}$$

here $P(n) = a^2 b^2 - a^2 b c - a b^2 c + a b c^2 - a^2 b d - a b^2 d + a c d + a^2 c d + b c d + 3 a b c d + b^2 c d - c^2 d - 2 a c^2 d - 2 b c^2 d + c^3 d + a b d^2 - c d^2 - 2 a c d^2 - 2 b c d^2 + 2 c^2 d^2 + c d^3 - 2 a^2 b n - 2 a b^2 n + 2 a c n + a^2 c n + 2 b c n + 6 a b c n + b^2 c n - 2 c^2 n - 3 a c^2 n - 3 b c^2 n + 2 c^3 n + 2 a d n + a^2 d n + 2 b d n + 6 a b d n + b^2 d n - 6 c d n - 11 a c d n - 11 b c d n + 10 c^2 d n - 2 d^2 n - 3 a d^2 n - 3 b d^2 n + 10 c d^2 n + 2 d^3 n + 4 a n^2 + a^2 n^2 + 4 b n^2 + 8 a b n^2 + b^2 n^2 - 8 c n^2 - 13 a c n^2 - 13 b c n^2 + 13 c^2 n^2 - 8 d n^2 - 13 a d n^2 - 13 b d n^2 + 29 c d n^2 + 13 d^2 n^2 - 8 n^3 - 14 a n^3 - 14 b n^3 + 28 c n^3 + 28 d n^3 + 21 n^4$. The formula holds as long as $\mathfrak{A}(1-a-b+c+d) > 0$. When $a = b = c = d = 0$, this becomes a well-known conclusion due to Zeilberger:

$$(\text{ex1S}) \quad \zeta(2) = \sum_{n \geq 1} 4^{-3n} \frac{(1)_n^3}{(1/2)_n^3} \left(\frac{21}{n^2} - \frac{8}{n^3} \right)$$

To derive a q -version of ex1, one simply uses the corresponding q -version of the WZ-candidate:

$$F(n, k) = \text{Gauss2F1}_q(a-d-n, b-d-n, c-d+1, d+k+2n+1) \quad 0 < q < 1$$

now compute $G(n, k)$, which we again do not display here, readers can imagine it looks like equation 1.1, albeit much longer. Proposition 1.3 still gives $\sum_{k \geq 0} F(0, k) = \sum_{n \geq 0} G(n, 0)$. Converting q -gamma functions into q -pochhammer symbols:

$$\begin{aligned}
(\text{ex1Q}) \quad & \sum_{k \geq 0} \frac{q^{k(1-a-b+c+d)} (q^{a+1}; q)_k (q^{b+1}; q)_k}{(q^{c+1}; q)_{k+1} (q^{d+1}; q)_{k+1}} \\
& = \sum_{n \geq 1} \frac{q^{-1-a-b-2c-2d+(-3-a-b+2c+2d)n+3n^2} (q^{a+1}, q^{b+1}, q^{-a+c+1}, q^{-a+d+1}, q^{-b+c+1}, q^{-b+d+1}; q)_{n-1} \times P_q(n)}{(q^{c+1}, q^{d+1}, q^{-a-b+c+d+1}; q)_{2n}}
\end{aligned}$$

where $P_q(n) = -q^{a+b+c+d+2n+1} - q^{2a+b+c+d+3n} - q^{a+2b+c+d+3n} + q^{2a+2b+c+d+4n} + q^{2a+2b+c+d+4n+1} + q^{a+b+2c+d+4n} + q^{a+b+2c+d+4n+1} + q^{a+b+c+2d+4n} + q^{a+b+c+2d+4n+1} - q^{a+b+2c+2d+6n+1} - q^{2a+b+2c+2d+7n} - q^{a+2b+2c+2d+7n} + q^{a+b+3c+2d+8n} + q^{a+b+2c+3d+8n} - q^{2a+2b+c+2n+1} - q^{2a+2b+d+2n+1} + q^{2a+2b+1} + q^{2a+2c+2d+6n} - q^{a+3c+2d+7n} - q^{a+2c+3d+7n} + q^{2b+2c+2d+6n} - q^{b+3c+2d+7n} - q^{b+2c+3d+7n} + q^{3c+3d+8n}$. This is a q -analogue of ex1: letting $q \rightarrow 1^-$ recovers exactly that formula. It holds as long as $\mathfrak{A}(1-a-b+c+d) > 0$.

The special case of $a = b = c = d = 0$ in ex1Q gives

$$\sum_{k \geq 0} \frac{q^k}{(1-q^{1+k})^2} = \sum_{n \geq 1} \frac{q^{3n^2-3n} (1-2q^{3n-1} - 3q^{4n-1} - 3q^{5n-1} + 3q^n + 3q^{2n} + q^{3n})(q; q)_n^6}{(1-q^n)^3 (q; q)_{2n}^3}$$

this is a q -analogue of ex1S.

Example II. Let a, b, c, d, e be in a neighbourhood of 0. Let

$$F(n, k) = \text{Dougall5F4}(2a-e-n, a+b-e-n, a+c-e-n, a+d-e-n, 1-a+e+k+n)$$

Applying Proposition 1.3 we obtain

$$\begin{aligned}
(\text{ex2}) \quad & \sum_{k \geq 0} \frac{(2+e+2k)(1+a)_k (1+b)_k (1+c)_k (1+d)_k}{(1-a+e)_{k+1} (1-b+e)_{k+1} (1-c+e)_{k+1} (1-d+e)_{k+1}} = \\
& \frac{(-1)^n P(n) (1-a-b+e)_{n-1} (1-a-c+e)_{n-1} (1-b-c+e)_{n-1}}{(1-a-d+e)_{n-1} (1-b-d+e)_{n-1} (1-c-d+e)_{n-1}} \\
& \sum_{n \geq 1} \frac{1}{(1-a+e)_n (1-b+e)_n (1-c+e)_n (1-d+e)_n (1-a-b-c-d+2e)_{2n}}
\end{aligned}$$

here $P(n) = a^2 b^2 - a^2 b c - a b^2 c + a b c^2 - a^2 b d - a b^2 d + a c d + a^2 c d + b c d + 3 a b c d + b^2 c d - c^2 d - 2 a c^2 d - 2 b c^2 d + c^3 d + a b d^2 - c d^2 - 2 a c d^2 - 2 b c d^2 + 2 c^2 d^2 + c d^3 - 2 a^2 b n - 2 a b^2 n + 2 a c n + a^2 c n + 2 b c n + 6 a b c n + b^2 c n - 2 c^2 n - 3 a c^2 n - 3 b c^2 n +$

$2c^3n + 2adn + a^2dn + 2bdn + 6abdn + b^2dn - 6cdn - 11acd - 11bcd + 10c^2dn - 2d^2n - 3ad^2n - 3bd^2n + 10cd^2n + 2d^3n + 4an^2 + a^2n^2 + 4bn^2 + 8abn^2 + b^2n^2 - 8cn^2 - 13acn^2 - 13bcn^2 + 13c^2n^2 - 8dn^2 - 13adn^2 - 13bdn^2 + 29cdn^2 + 13d^2n^2 - 8n^3 - 14an^3 - 14bn^3 + 28cn^3 + 28dn^3 + 21n^4$. This formula holds when $\Re(1 - a - b - c - d + 2e) < 0$.

When $a = b = c = d = e = 0$, this reduces to

$$(ex2S) \quad \frac{2}{5}\zeta(3) = \sum_{n \geq 1} \left(-\frac{1}{4}\right)^{n-1} \frac{(1)_n}{(1/2)_n} \frac{1}{n^3}$$

For a q -analogue, let $F(n, k) = \text{Dougall5F4}_q(2a-e-n, a+b-e-n, a+c-e-n, a+d-e-n, 1-a+e+k+n)$, then

$$(ex2Q) \quad \sum_{k \geq 0} \frac{q^{k(1-a-b-c-d+2e)}(1-q^{e+2k+2})(q^{a+1}, q^{b+1}, q^{c+1}, q^{d+1}; q)_k}{(q^{-a+e+1}, q^{-b+e+1}, q^{-c+e+1}, q^{-d+e+1}; q)_{k+1}} = \sum_{n \geq 1} \frac{(-1)^n q^{-1-2a-2b-2c-2d+en-e+n(n+1)/2} (q^{-a-b+e+1}, q^{-a-c+e+1}, q^{-a-d+e+1}, q^{-b-c+e+1}, q^{-b-d+e+1}, q^{-c-d+e+1}; q)_{n-1} \times P_q(n)}{(q^{1-a+e}, q^{1-b+e}, q^{1-c+e}, q^{1-d+e}; q)_n (q^{1-a-b-c-d+2e}; q)_{2n}}$$

where $P_q(n) = q^{a+b+c+d+2e+2n} + q^{2a+b+c+d+2e+2n} + q^{a+2b+c+d+2e+2n} + q^{a+b+2c+d+2e+2n} + q^{a+b+c+2d+2e+2n} - q^{a+b+c+d+3e+3n} - q^{2a+2b+2c+2d} - q^{a+b+c+3e+3n} - q^{a+b+d+3e+3n} - q^{a+c+d+3e+3n} - q^{b+c+d+3e+3n} + q^{5e+5n}$. This formula holds when $\Re(1 - a - b - c - d + 2e) < 0$. This is a q -analogue of ex2. When $a = b = c = d = e = 0$, this reduces to

$$\sum_{k \geq 0} \frac{q^k(1+q^{1+k})}{(1-q^{1+k})^3} = \sum_{n \geq 1} \frac{(-1)^{n-1}(1+3q^n+q^{2n})(q; q)_n^2}{(1-q^n)^3(q; q)_{2n}}$$

this is a q -analogue of ex2S.

We can devise the following strategy on finding a q -analogue of a hypergeometric identity:

$$(1.2) \quad \sum_{n \geq 0} z^n \frac{(a_1)_n \cdots (a_m)_n}{(b_1)_n \cdots (b_m)_n} R(n) = \text{a simple constant} \quad z \in \mathbb{Q} \quad a_i, b_i \in \mathbb{Q} \cap (0, 1] \quad R(n) \in \mathbb{Q}(n)$$

Finding q -analogues of a classical hypergeometric identity

- (1) Using procedure described in [2], try to search for a classical WZ-seed $\text{Seed}(a + An, b + Bn, \dots, k + k_0 + Kn)$ that can prove 1.2 after using Proposition 1.3. The strategy fails if we cannot find such a seed.
- (2) Translate this seed, using previously tabulated WZ-seeds, to its q -version $\text{Seed}_q(a + An, b + Bn, \dots, k + k_0 + Kn)$
- (3) Find q -WZ-mate for $F(n, k) = \text{Seed}_q(a + An, b + Bn, \dots, k + k_0 + Kn)$, then apply again Proposition 1.3.

Our compatibility between classical and q WZ-seeds via

$$\lim_{q \rightarrow 1^+} \text{Seed}_q = \text{Seed}$$

ensures we are finding q -analogue for our original equality 1.2. We will illustrate through various not-so-trivial examples that, as long as step (1) passes through, a q -analogue can always be found.

However, among the four terms appearing in

$$\sum_{k \geq 0} F(0, k) + \sum_{n \geq 0} g(n) = \sum_{n \geq 0} G(n, 0) + \lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$$

the two terms involving limits $\sum_{n \geq 0} g(n)$ and $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$ make our life a bit more difficult, a difficulty already present in classical hypergeometric case. We analyse them now.

1.4. Asymptotic of q -gamma function. We reiterate our tacit convention that $0 < q < 1$ throughout.

Lemma 1.4. *As $x \rightarrow +\infty$, we have $\Gamma_q(x) = (1-q)^{1/2-x} C_q (1 + O(q^x))$, here C_q is a non-zero constant that depends on q only.*

Proof. See [16]. In fact, one can give an explicit expression of C_q , but we won't need it. \square

Proposition 1.5. (a) *Let $n \geq 1$ be an integer,*

$$(C_{q^n})^n = \sqrt{\frac{1-q^n}{1-q}} C_q \prod_{i=1}^{n-1} \Gamma_{q^n}\left(\frac{i}{n}\right)$$

(b) *For $a \in \mathbb{R}$, then as $n \rightarrow \infty$ with $n \in \mathbb{N}$, we have*

$$\Gamma_q(a+n) = (1-q)^{1/2-a-n} C_q (1 + O(q^n))$$

and if $a \notin \mathbb{Z}$,

$$\Gamma_q(a-n) = q^{(1-2a+n)n/2} (1-q)^{1/2-a+n} \frac{\Gamma_q(1-a)\Gamma_q(a)}{C_q} (1 + O(q^n))$$

Proof. (a) Follows immediately from the multiplication formula:

$$\Gamma_q(nx) \prod_{i=1}^{n-1} \Gamma_{q^n}\left(\frac{i}{n}\right) = \left(\frac{1-q^n}{1-q}\right)^{nx-1} \prod_{i=0}^{n-1} \Gamma_{q^n}\left(x + \frac{i}{n}\right)$$

(b) The first statement follows immediately from above lemma. For second statement, let $\rho(x)$ defined by $\Gamma_q(x)\Gamma_q(1-x) = q^{x(x-1)/2} e^{\pi i x} \rho(x)$, then using $\Gamma_q(x+1) = \frac{1-q^x}{1-q} \Gamma_q(x)$, it is easily to check $\rho(x)$ is 1-periodic: $\rho(x+1) = \rho(x)$. Therefore

$$\Gamma_q(a-n) = \frac{q^{(a-n)(a-n-1)/2} e^{\pi i(a-n)} \rho(a-n)}{\Gamma_q(1-a+n)} = \frac{q^{(a-n)(a-n-1)/2} e^{\pi i(a-n)} \rho(a)}{\Gamma_q(1-a+n)}$$

using first statement concludes the proof. \square

Above proposition completely settled the term $g(n) := \lim_{k \rightarrow \infty} G(n, k)$. We shall only give two illustrative examples:

Example 1.6. Let us elaborate why $g(n) := \lim_{k \rightarrow \infty} G(n, k)$ in Example I vanishes. We relabel our auxiliary variables, write $F(n, k) = \text{Gauss2F1}_q(a-n, b-n, c, d+k+2n)$, i.e.

$$F(n, k) = \frac{\Gamma_q(-a+c+n)\Gamma_q(-b+c+n)\Gamma_q(a+d+k+n)\Gamma_q(b+d+k+n)q^{-(d+k+2n)(a+b-c-2n)}}{\Gamma_q(a-n)\Gamma_q(b-n)\Gamma_q(d+k+2n+1)\Gamma_q(-a-b+c+2n)\Gamma_q(c+d+k+2n)}$$

Using part (b) of above proposition, we have, for fixed n as $k \rightarrow \infty$

$$F(n, k) \sim \frac{(1-q)^{-a-b+c+2n+1} \Gamma_q(-a+c+n)\Gamma_q(-b+c+n)q^{-((d+k+2n)(a+b-c-2n))}}{\Gamma_q(a-n)\Gamma_q(b-n)\Gamma_q(-a-b+c+2n)} = O(q^{(2n+c-a-b)k})$$

The q -rational function $G(n, k)/F(n, k)$ is quite long, but if we really computed it, then we will see its $O(1)$ when n is fixed and $k \rightarrow \infty$. Therefore we conclude, $c-a-b > 0$ implies $g(n) = 0$ for $n \geq 0$.

Example 1.7. Let

$$F(n, k) = \frac{(-1)^n (q+1)^{-2(k+n)} \Gamma_{q^2}(c-b)\Gamma_q(-a+2c+n-1)\Gamma_q(a+2k+n)\Gamma_{q^2}(b+k+n)q^{-2an-2bn+2cn+2k+n^2}}{\Gamma_{q^2}(b)\Gamma_q(a-n)\Gamma_{q^2}(k+n+1)\Gamma_{q^2}(-a+c+n-\frac{1}{2})\Gamma_{q^2}(c+k+n)\Gamma_{q^2}(a+b-c+k+\frac{3}{2})\Gamma_q(-a-2b+2c+n-1)}$$

this comes from $\text{Seed2}_q(a \rightarrow a-n, b \rightarrow b, c \rightarrow c, k \rightarrow k+n)$. It has WZ-mate

$$G(n, k) = -F(n, k)q^{-2k} \frac{q^{2c} (1-q^{2a+2b-2c+2k+1}) (q^{a+1} - q^{2c+2k+3n})}{(q^{2a+1} - q^{2c+2n}) (q^{a+2b+1} - q^{2c+n})}$$

We wish to compute $g(n) := \lim_{n \rightarrow \infty} G(n, k)$. First we use (b) of above proposition, giving

$$F(n, k) = q^{2k} \frac{(-1)^n (q-1)^2 C_q (q+1)^{a-n+\frac{3}{2}} \Gamma_{q^2}(c-b) \Gamma_q(-a+2c+n-1) q^{n(-2a-2b+2c+n)}}{(C_{q^2})^2 \Gamma_{q^2}(b) \Gamma_q(a-n) \Gamma_{q^2}(-a+c+n-\frac{1}{2}) \Gamma_q(-a-2b+2c+n-1)} (1 + O(q^k))$$

now the ratio $(C_{q^2})^2/C_q$ can be converted back to q -gamma function using (a), which is $\sqrt{1+q}\Gamma_{q^2}(1/2)$. For $G(n, k)$, we have a cancellation of q^{2k} in asymptotic, hence $g(n)$ is non-zero in this case, and we obtained its expression in terms of q -gamma function. We will employ this WZ-pair $F(n, k)$ in Example IV below.

Remark 1.8. We make two remarks. First, the cancellation of $(C_{q^2})^2/C_q$ into q -gamma function above is not accidental, it is a general feature of our tabulated WZ-seeds.

Second, $g(n)$ is always non-zero if the corresponding WZ-seed comes from a hypergeometric summation formula that is *balanced*. In above example, Seed2 comes from Balanced3F2. Other WZ-seeds sharing this property are Dougall7F6, Seed7, Seed9, Seed10. When we use WZ-pair from these seeds on Proposition 1.3, we must deal with an additional sum $\sum_{n \geq 0} g(n)$.

Next we investigate the behaviour of

$$\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$$

appearing in Proposition 1.3. In both classical and q cases, this is the most intractable term in the formula. Rest of materials in this section is rather technical, and can be skipped at first reading, provided readers are willing to accept $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$ is really 0 unless we specified otherwise.

We take a similar approach to classical case (Section 2.4 of [2]). Consider a very general q -hypergeometric term, not-necessarily coming from a WZ-seed,

$$F(n, k) = q^{An^2+Bn+Cnk+Dk^2+Ek} R_1(q)^n R_2(q)^k \frac{\prod_{i=1}^m \Gamma_{q_{r_i}}(a_i + C_i k + D_i n)}{\prod_{i=1}^l \Gamma_{q_{t_i}}(b_i + E_i k + F_i n)}$$

here

$$A, B, C, D, E \in \mathbb{R} \quad C_i, D_i, E_i, F_i \in \mathbb{Z} \quad a_i, b_i \in \mathbb{R} \quad r_i, t_i \in \mathbb{Z}^{\geq 1}$$

and $R_1(q), R_2(q)$ are two rational function in q with $R_1(q=0)$ and $R_2(q=0)$ non-zero.

We shall assume this $F(n, k)$ satisfies the following two assumptions:

- For fixed n , asymptotic of $F(n, k)$ as $k \rightarrow \infty$ has the form

$$F(n, k) \sim q^{U_1 k + U_2 k^2} \times O(1)$$

- For fixed k , asymptotic of $F(n, k)$ as $n \rightarrow \infty$ has the form

$$F(n, k) \sim q^{V_1 n + V_2 n^2} \times O(1)$$

i.e., we require there should be no $(1 - q^i)^n$ or $(1 - q^i)^k$ appearing in asymptotic expansion. After some contemplation, readers will see all $F(n, k)$ derived from our list of WZ-seed has this property.

For $F(n, k)$ satisfying this assumption, let $x \geq 0$ be a fixed real number, we have

$$|F(n, nx)| = q^{\nu_2(x)n^2 + \nu_1(x)n} \times O(1) \quad n \rightarrow \infty, nx \in \mathbb{Z}$$

for some function $\nu_1(x), \nu_2(x)$. Using Proposition 1.5, it is not difficult to derive the following expression of ν_1 and ν_2 :

$$\begin{aligned}\nu_2(x) &= A + Cx + Dx^2 + \sum_{i=1}^m r_i \chi_2(C_i x + D_i) - \sum_{i=1}^l t_i \chi_2(E_i x + F_i) \\ \nu_1(x) &= B + Ex + \sum_{i=1}^m r_i \left(a_i - \frac{1}{2}\right) \chi_1(C_i x + D_i) - \sum_{i=1}^l t_i \left(b_i - \frac{1}{2}\right) \chi_1(E_i x + F_i)\end{aligned}$$

here

$$\chi_1(x) = \begin{cases} x & x < 0 \\ 0 & x \geq 0 \end{cases} \quad \chi_2(x) = \begin{cases} x^2/2 & x < 0 \\ 0 & x \geq 0 \end{cases}$$

Both ν_i are continuous and note that ν_2 does not depend on a_i, b_i in expression of $F(n, k)$.

The following criterion is q -version of corresponding classical version in [2].

Proposition 1.9. *Given $F(n, k)$ as in above, if*

$$\inf_{x \geq 0} \min(\nu_1(x), \nu_2(x)) > 0$$

then $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k) = 0$.

Proof. It suffices to prove, for each $x \geq 0$, there exists $\delta > 0$ such that

$$\sum_{k \in (x-\delta, x+\delta) \times n} |F(n, k)|$$

tends to 0 as $n \rightarrow \infty$. For an $x \geq 0$, if $\nu_2(x) > \varepsilon > 0$, then for some small $\delta > 0$, we have

$$|F(n, k)| = O(q^{\varepsilon n^2}) \quad (x - \delta)n < k < (x + \delta)n$$

so the sum is dominated by

$$\sum_{k \in (x-\delta, x+\delta) \times n} |F(n, k)| \ll nq^{\varepsilon n^2}$$

which tends to 0. For an $x \geq 0$, if $\nu_2(x) = 0$ but $\nu_1(x) > \varepsilon > 0$, then again for some small $\delta > 0$,

$$|F(n, k)| = O(q^{\varepsilon n}) \quad (x - \delta)n < k < (x + \delta)n$$

so the sum is dominated by

$$\sum_{k \in (x-\delta, x+\delta) \times n} |F(n, k)| \ll nq^{\varepsilon n}$$

this also tends to 0. □

Example 1.10. For $F(n, k) = \text{Gauss2F1}_q(a - d - n, b - d - n, c - d + 1, d + k + 2n + 1)$ in Example I, one computes $\nu_2(x) = 3 + 2x$, which is positive on $x \geq 0$. Therefore above proposition implies $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k) = 0$.

For $F(n, k) = \text{Seed2}_q(1/2 + a - n, b + 1/2, 1 + c, d + k + n)$ we will use in Example IV below, one computes $\nu_2(x) \equiv 1/2$, so above Proposition implies $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k) = 0$.

For $F(n, k) = \text{Seed10}_q(a + 3b - 2n + 1/2, 2b + 1, c + k + n)$ we will use in Example XII below, one computes

$$\nu_2(x) = \begin{cases} 4 & x \geq 1 \\ \frac{5}{2} + 3x - \frac{3x^2}{2} & 0 \leq x < 1 \end{cases}$$

so above Proposition again implies $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k) = 0$.

In above examples, we only need knowledge of $\nu_2(x)$, free variables a, b, c, d play no role here. There are some other examples which we need to use the finer function $\nu_1(x)$, which depends on these free variables.

Example 1.11. For

$$F(n, k) = \text{Gauss2F1}_q(-a - b + c + 1/2 - n, -a + c - n + 1/2, -a - b + 2c - n + 1, 2a + 2b - 2c + d + k + 2n)$$

in Example VII below, one computes

$$\nu_2(x) \equiv 0 \quad \nu_1(x) = 2a + b + d + ax$$

thus if b, d in a neighbourhood of 0 and $a > 0$, we have $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k) = 0$. We shall apply this restriction on a, b, d when we perform that example.

Example 1.12. Consider

$$F(n, k) = \text{Seed9}_q(1 - 2(a + n), -a - n + \frac{1}{2}, -a - n + \frac{1}{2}, a + k + n)$$

that we will use in Example XVII. We have

$$\nu_2(x) = \begin{cases} 1 - 2x^2 & x \leq 1/2 \\ 2 - 4x + 2x^2 & 1/2 < x \leq 1 \\ 0 & x \geq 1 \end{cases} \quad \nu_1(x) = \begin{cases} 2a - 4x & x \leq 1/2 \\ -4 + 4a + 4x - 4ax & 1/2 < x \leq 1 \\ -2 + 2x & x \geq 1 \end{cases}$$

so $\min(\nu_1(x), \nu_2(x))$ is zero at $x = 1$, so above proposition fails to apply. Indeed, we will see $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$ is *nonzero*.

We will only give two examples in this article in which $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$ is non-zero: Examples XVI and XVII. For all other examples, above proposition applies to conclude it is zero.

When the limit is non-zero, it is sometimes a daunting task to compute them: the classical hypergeometric case is more understood since we have a greater variety of tools from asymptotic analysis available (cf. last section of [2]). For the q -hypergeometric case, our choice of strategy is more limited. We are only able to evaluate this limit for Example XVI, but not for Example XVII.

2. MORE EXAMPLES

Example III. Using

$$F(n, k) = \text{Dougall5F4}_q(-n, -n, -n, -n, k + 2n + 1)$$

and Proposition 1.3 says (with vanishing limiting terms):

$$\sum_{k \geq 0} F(0, k) = \sum_{n \geq 0} G(n, 0)$$

which becomes

$$\sum_{k \geq 0} \frac{q^k(1 + q^{1+k})}{(1 - q^{1+k})^3} = \sum_{n \geq 1} \frac{(-1)^{n-1} q^{5n(n-1)/2} P_q(n) (q; q)_n^{10}}{(1 - q^n)^5 (q; q)_{2n}^5}$$

with $P_q(n) = 10q^{2n} + 10q^{3n} + 5q^{4n} + q^{5n} - 5q^{3n-1} - 15q^{4n-1} + q^{5n-2} - 24q^{5n-1} + 5q^{6n-2} - 15q^{6n-1} + 10q^{7n-2} - 5q^{7n-1} + 10q^{8n-2} + 5q^{9n-2} + q^{10n-2} + 5q^n + 1$. This is a q -analogue of a hypergeometric expression of $\zeta(3)$ (due to [1]):

$$\sum_{n \geq 1} \left(-\frac{1}{2^{10}}\right)^n \frac{(1)_n^5}{n^5 (1/2)_n^5} (205n^2 - 160n + 32) = -2\zeta(3)$$

The hypergeometric version, can of course, be proved directly using classical version of WZ-pair.

Example IV. The Ramanujan-type $1/\pi$ formula

$$(2.1) \quad \sum_{n \geq 0} \left(\frac{1}{4}\right)^n \frac{\left(\frac{1}{2}\right)_n^3}{(1)_n^3} (6n+1) = \frac{4}{\pi}$$

has a four-parameter deformation, from which above is obtained by specializing at $a = b = c = d = 0$:

$$\begin{aligned} & a \sum_{k \geq 0} \frac{2^{-2k} \left(b+d+\frac{1}{2}\right)_k \left(a+c+2d+\frac{1}{2}\right)_{2k}}{(d+1)_k (c+d+1)_k \left(a+b+d+\frac{3}{2}\right)_k} \\ & \frac{2^{a+c+2d+1/2} \Gamma(d+1) \Gamma(c+d+1) \Gamma\left(a+b+d+\frac{3}{2}\right)}{\Gamma\left(b+d+\frac{1}{2}\right) \Gamma\left(a+c+2d+\frac{1}{2}\right) (2a+4b-2c+1) \sqrt{\pi}} \times \sum_{n \geq 0} \frac{2^{-n} \left(-a-c+\frac{1}{2}\right)_n \left(-a+c+\frac{1}{2}\right)_n}{(1-a)_n \left(-a-2b+c+\frac{1}{2}\right)_n} \\ & = \frac{2a+2b+2d+1}{2a+4b-2c+1} \sum_{n \geq 0} \frac{2^{-2n-1} (2a-2c-4d-6n-1) \left(-a-c+\frac{1}{2}\right)_n \left(-a+c+\frac{1}{2}\right)_n \left(b+d+\frac{1}{2}\right)_n \left(a+c+2d+\frac{1}{2}\right)_n}{(1-a)_n (d+1)_n (c+d+1)_n \left(-a-2b+c+\frac{1}{2}\right)_n} \end{aligned}$$

this can be obtained by apply Proposition 1.3 to

$$F(n, k) = \text{Seed2}(1/2 + a - n, b + 1/2, 1 + c, d + k + n)$$

here $\sum_{n \geq 0} g(n)$ gives the $\sum_{n \geq 0}$ on LHS of equality, here we assume reader knows how to compute asymptotic of ratio of gamma functions. And $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$ vanishes. To derive a q -analogue of above formula, we simply apply Proposition 1.3 on its q -version:

$$F(n, k) = \text{Seed2}_q(1/2 + a - n, b + 1/2, 1 + c, d + k + n)$$

giving

$$\begin{aligned} & \sum_{k \geq 0} \frac{(1-q^{2a}) q^{2k} (q^{2b+2d+1}; q^2)_k (q^{a+c+2d+\frac{1}{2}}; q)_{2k}}{(1-q^{2a+2b+2d+2k+1}) (q^{2d+2}, q^{1+2a+2b+2d}, q^{2+2c+2d}; q^2)_k} + \\ & \frac{(1+q)^{a+c+2d+\frac{1}{2}} \Gamma_{q^2}(d+1) \Gamma_{q^2}(c+d+1) \Gamma_{q^2}\left(a+b+d+\frac{1}{2}\right)}{(q^{a+2b+\frac{1}{2}} - q^c) \Gamma_{q^2}\left(\frac{1}{2}\right) \Gamma_{q^2}\left(b+d+\frac{1}{2}\right) \Gamma_q\left(a+c+2d+\frac{1}{2}\right)} \sum_{n \geq 0} \frac{(1-q) q^{-an+a-2bn+cn-2d+\frac{n^2}{2}-\frac{1}{2}} (q^{-a-c+\frac{1}{2}}, q^{-a+c+1/2}; q)_n}{(q^{2-2a}; q^2)_n (q^{-a-2b+c+\frac{1}{2}}; q)_n} \\ & = \sum_{n \geq 0} \frac{(1-q^{-a+c+2d+3n+\frac{1}{2}}) q^{-an+a-2bn+cn-2d+\frac{n^2}{2}-\frac{1}{2}} (q^{-a-c+\frac{1}{2}}, q^{-a+c+\frac{1}{2}}, q^{a+c+2d+\frac{1}{2}}; q)_n (q^{2b+2d+1}; q^2)_n}{(q^{a+2b+\frac{1}{2}} - q^c) (q^{2-2a}, q^{2+2c+2d}, q^{2+2d}; q^2)_n (q^{-a-2b+c+\frac{1}{2}}; q)_n} \end{aligned}$$

we remark the term $\Gamma_{q^2}(1/2)$ in $\lim_{n \rightarrow \infty} g(n)$ comes from Proposition 1.5 when calculating $\lim_{k \rightarrow \infty} G(n, k) = g(n)$. Letting $a = 0$ kills the summation in k , the remaining summation in n on LHS can actually expressible in terms of q -gamma (Lemma A.1 in appendix), resulting in

$$(2.2) \quad \sum_{n \geq 0} \frac{q^{-2bn+cn+\frac{n^2}{2}} (1-q^{c+2d+3n+\frac{1}{2}}) (q^{\frac{1}{2}-c}, q^{c+\frac{1}{2}}, q^{c+2d+\frac{1}{2}}; q)_n (q^{2b+2d+1}; q^2)_n}{(q^2, q^{2d+2}, q^{2c+2d+2}; q^2)_n (q^{-2b+c+\frac{1}{2}}; q)_n} = \frac{(1-q)(1+q)^{2d+1} \Gamma_{q^2}(1-b) \Gamma_{q^2}(d+1) \Gamma_q\left(-2b+c+\frac{1}{2}\right) \Gamma_{q^2}(c+d+1)}{\Gamma_{q^2}\left(\frac{1}{2}\right) \Gamma_q(1-2b) \Gamma_{q^2}\left(-b+c+\frac{1}{2}\right) \Gamma_q\left(c+2d+\frac{1}{2}\right)}$$

When $b = c = d = 0$, this is a q -analogue of equation 2.1:

$$\sum_{n \geq 0} \frac{q^{n^2} (1-q^{1+6n}) (q; q^2)_n^2 (q^2; q^4)_n}{(q^4; q^4)_n^3} = \frac{1-q^4}{\Gamma_{q^4}(1/2)^2}$$

this was also derived in [12]. Equation 2.2 can be seen as a 3-parameter deformation of it. We will obtain another q -analogue for this $1/\pi$ formula in Example VII.

Example V. The formula

$$(2.3) \quad \sum_{n \geq 0} \left(\frac{1}{26}\right)^n \frac{\left(\frac{1}{2}\right)_n^3}{(1)_n^3} (42n + 5) = \frac{16}{\pi}$$

has a 3-parameter q -analogue as follows:

$$\begin{aligned} \sum_{n \geq 0} \frac{q^{5n^2/2-2bn+cn-4b-4} \times P_q(n) (q^{1-2b}, q^{1-2b+2c}; q^2)_n (q^{1/2-c}, q^{1/2+c}, q^{1/2+c+2d}; q)_n (q^{c+2d+1/2}; q)_{3n}}{(q^2; q^2)_n (q^{-2b+c+\frac{1}{2}}; q)_{3n+2} (q^{2d+2}, q^{2c+2d+2}; q^2)_{2n+1}} \\ = \frac{(1-q)(1+q)^{2d+1} \Gamma_{q^2}(1-b) \Gamma_{q^2}(d+1) \Gamma_q(-2b+c+\frac{1}{2}) \Gamma_{q^2}(c+d+1)}{\sqrt{q} \Gamma_{q^2}(\frac{1}{2}) \Gamma_q(1-2b) \Gamma_{q^2}(-b+c+\frac{1}{2}) \Gamma_q(c+2d+\frac{1}{2})} \end{aligned}$$

with $P_q(n) = q^{4b+7/2} - q^{2b+c+3n+5} - q^{2b+4n+9/2} - q^{2b+2c+4n+9/2} - q^{4b+2d+4n+11/2} - q^{4b+2c+2d+4n+11/2} + q^{2b+c+5n+5} - q^{4b+c+2d+5n+4} - q^{4b+c+2d+5n+5} + q^{2c+6n+11/2} + q^{4b+2d+6n+11/2} - q^{2b+2c+2d+6n+9/2} + q^{4b+2c+2d+6n+11/2} + q^{2b+c+2d+7n+5} + q^{2b+c+2d+7n+6} + q^{2b+c+2d+7n+7} - q^{4b+c+2d+7n+6} + q^{2b+3c+2d+7n+5} + q^{2b+3c+2d+7n+6} + q^{2b+3c+2d+7n+7} + q^{2b+2c+2d+8n+9/2} + q^{4b+2c+4d+8n+11/2} + q^{4b+2c+4d+8n+15/2} - q^{3c+2d+9n+7} + q^{4b+2c+4d+10n+13/2} - q^{3c+2d+11n+6} - q^{3c+2d+11n+8} - q^{2b+3c+4d+11n+9} - q^{2b+2c+4d+12n+13/2} - q^{2b+2c+4d+12n+15/2} - q^{2b+2c+4d+12n+17/2} + q^{4c+4d+12n+15/2} - q^{2b+4c+4d+12n+13/2} - q^{2b+4c+4d+12n+15/2} - q^{2b+4c+4d+12n+17/2} - q^{3c+4d+13n+8} + q^{2b+3c+4d+13n+9} - q^{5c+4d+13n+8} - q^{4b+3c+6d+13n+8} + q^{4c+4d+14n+17/2} + q^{4c+4d+14n+19/2} - q^{2b+4c+6d+14n+17/2} + q^{3c+4d+15n+8} + q^{5c+4d+15n+8} + q^{2b+3c+6d+15n+9} + q^{2b+5c+6d+15n+9} + q^{2b+4c+6d+16n+17/2} - q^{5c+6d+19n+10}$.

This can be proved *mutatis mutandis* as previous example with $F(n, k) = \text{Seed}2_q(1/2 + a + c - n, b + 1/2 - n, c + 1, k + d + 2n)$, setting $a = 0$ will kill $\sum_{k \geq 0} F(0, k)$, and $\sum_{n \geq 0} g(n)$ can be found using the same trick as in previous example (i.e. similar to Lemma A.1). Letting $b = c = d = 0$ yields

$$\begin{aligned} \sum_{n \geq 0} \frac{q^{5n^2} (2q^{4n+2} + q^{6n+1} - 2q^{8n+2} + q^{8n+4} - q^{10n+1} + 2q^{10n+3} - q^{12n+4} - 2q^{14n+3} - q^{18n+5} + 1) (q; q^2)_n^2 (q^2; q^4)_n^3}{(q^{4n+2} + 1)^2 (q^4; q^4)_n (q^4; q^4)_{2n}^2} \\ = \frac{1 - q^4}{\Gamma_{q^4}(1/2)^2} \end{aligned}$$

letting $q \rightarrow 1^-$ recovers the formula at the start. We will give another q -analogue of this formula in Example VIII below.

Example VI.

$$(2.4) \quad \sum_{n \geq 0} \left(-\frac{3^3}{2^9}\right)^n \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{6}\right)_n \left(\frac{5}{6}\right)_n}{(1)_n^3} (154n + 15) = \frac{32\sqrt{2}}{\pi}$$

has the following generalization:

$$\begin{aligned} a \sum_{k \geq 0} \frac{2^{-a-b-c-2k} \left(-a+b+c+\frac{1}{2}\right)_{2k}}{(b+1)_k (c+1)_k} \\ = \sum_{n \geq 0} \frac{(-1)^n 2^{-a-b-c-5n-6} \times P(n) \left(a+b-c+\frac{1}{2}\right)_n \left(a-b+c+\frac{1}{2}\right)_n \left(-a+b+c+\frac{1}{2}\right)_{3n}}{(a+1)_n (b+1)_{2n+1} (c+1)_{2n+1}} \end{aligned}$$

where $P(n) = 8a^3 - 8a^2b - 8a^2c - 8a^2n + 4a^2 - 8ab^2 + 48abc + 48abn + 24ab - 8ac^2 + 48acn + 24ac + 88an^2 + 104an + 30a + 8b^3 + 24b^2c + 88b^2n + 36b^2 + 24bc^2 + 240bcn + 72bc + 440bn^2 + 312bn + 46b + 8c^3 + 88c^2n + 36c^2 + 440cn^2 + 312cn + 46c + 616n^3 + 676n^2 + 214n + 15$, this can be proved using $F(n, k) = \text{Gauss}2F1(1/2 - a + b - c - n, 1 + b - c, c + k + 2n)$, here the analytic prerequisites of Proposition 1.3 require $\Re(a) > 0$ and b, c near 0, we still have $g(n) = 0$, $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k) = 0$. Now we want to set $a = 0$, but this cannot be done right away since series $\sum_{k \geq 0}$ on LHS diverges when $a = 0$, one needs a limiting process. First,

$$\frac{2^{-a-b-c-2k} \left(-a+b+c+\frac{1}{2}\right)_{2k}}{(b+1)_k (c+1)_k} = \frac{2^{-2a-\frac{1}{2}} \Gamma(b+1) \Gamma(c+1)}{\sqrt{\pi} \Gamma(-a+b+c+\frac{1}{2})} k^{-a-1} + O(k^{-2-a})$$

whence as $a \rightarrow 0^-$,

$$\lim_{a \rightarrow 0^-} \left(a \sum_{k \geq 0} \frac{2^{-a-b-c-2k} \left(-a+b+c+\frac{1}{2}\right)_{2k}}{(b+1)_k (c+1)_k} \right) = \frac{2^{-\frac{1}{2}} \Gamma(b+1) \Gamma(c+1)}{\sqrt{\pi} \Gamma\left(b+c+\frac{1}{2}\right)} \underbrace{\lim_{a \rightarrow 0^-} a \zeta(1+a)}_{=1}$$

letting $a = 0$ on RHS poses no issue, we then have the following two-variable identity:

$$\sum_{n \geq 0} \frac{(-1)^n 2^{-b-c-5n-6} \tilde{P}(n) (b-c+\frac{1}{2})_n (-b+c+\frac{1}{2})_n (b+c+\frac{1}{2})_{3n}}{(b+2n+1)(c+2n+1)(1)_n (b+1)_{2n} (c+1)_{2n}} = \frac{2^{-1/2} \Gamma(b+1) \Gamma(c+1)}{\sqrt{\pi} \Gamma\left(b+c+\frac{1}{2}\right)}$$

with $\tilde{P}(n) = 8b^3 + 24b^2c + 88b^2n + 36b^2 + 24bc^2 + 240bcn + 72bc + 440bn^2 + 312bn + 46b + 8c^3 + 88c^2n + 36c^2 + 440cn^2 + 312cn + 46c + 616n^3 + 676n^2 + 214n + 15$. Specializing to $b = c = 0$ recovers 2.4.

A q -version of above argument can be carried out in exact parallel. Using q -counterpart of the same WZ-seed, we arrive at

$$(1-q^{2a}) \sum_{k \geq 0} \frac{q^{2ak} (q^{-a+b+c+\frac{1}{2}}; q)_{2k}}{(q^{2b+2}; q^2)_k (q^{2c+2}; q^2)_k} = - \sum_{n \geq 0} \frac{(-1)^n q^{3an+bn+cn+7n^2/2-3/2} \times P_q(n) (q^{a+b-c+\frac{1}{2}}, q^{a-b+c+\frac{1}{2}}; q)_n (q^{-a+b+c+\frac{1}{2}}; q)_{3n}}{(1-q^{b+2n+1}) (q^{b+2n+1} + 1) (1-q^{c+2n+1}) (q^{c+2n+1} + 1) (q^{2a+2}; q^2)_n (q^{2b+2}, q^{2c+2}; q^2)_{2n}}$$

where $P_q(n) = q^{a+b+c+5n+2} + q^{a+b+c+5n+3} + q^{3a+b+c+7n+4} - q^{2a+2b+2c+10n+9/2} + q^{a+3b+3c+13n+6} - q^{2a+2b+6n+7/2} - q^{2a+2c+6n+7/2} - q^{2b+2c+8n+7/2} - q^{2b+2c+8n+11/2} + q^{2b+4n+7/2} + q^{2c+4n+7/2} - q^{3/2}$. Here the same restriction on a, b, c applies: $\Re(a) > 0$, b, c in a neighbourhood of 0.

We next find the limit of LHS as $a \rightarrow 0^-$. Using Proposition 1.5, we calculate asymptotic of summand:

$$\frac{q^{2ak} (q^{-a+b+c+\frac{1}{2}}; q)_{2k}}{(q^{2b+2}; q^2)_k (q^{2c+2}; q^2)_k} = q^{2ak} \underbrace{\frac{(1-q^2)^{b+c+1} \Gamma_{q^2}(b+1) \Gamma_{q^2}(c+1) (1-q)^{-b-c}}{\sqrt{1+q} \Gamma_{q^2}\left(\frac{1}{2}\right) \Gamma_q\left(b+c+\frac{1}{2}\right)}}_{:=A} + O(q^k)$$

hence

$$\lim_{a \rightarrow 0^-} \left((1-q^{2a}) \sum_{k \geq 0} \frac{q^{2ak} (q^{-a+b+c+\frac{1}{2}}; q)_{2k}}{(q^{2b+2}; q^2)_k (q^{2c+2}; q^2)_k} \right) = A \lim_{a \rightarrow 0^-} (1-q^{2a}) \sum_{k \geq 0} q^{2ak} = A$$

whence

$$\sum_{n \geq 0} \frac{(-1)^n q^{bn+cn+7n^2/2-3/2} \tilde{P}_q(n) (q^{b-c+\frac{1}{2}}, q^{-b+c+\frac{1}{2}}; q)_n (q^{b+c+\frac{1}{2}}; q)_{3n}}{(1-q^{b+2n+1}) (q^{b+2n+1} + 1) (1-q^{c+2n+1}) (q^{c+2n+1} + 1) (q^2, q^{2+2b}, q^{2+2c}; q^2)_n} = \frac{(1-q)(1+q)^{b+c+\frac{1}{2}} \Gamma_{q^2}(b+1) \Gamma_{q^2}(c+1)}{\Gamma_{q^2}\left(\frac{1}{2}\right) \Gamma_q\left(b+c+\frac{1}{2}\right)}$$

with $\tilde{P}_q(n) = -q^{b+c+5n+2} - q^{b+c+5n+3} - q^{b+c+7n+4} + q^{2b+2c+8n+7/2} + q^{2b+2c+8n+11/2} + q^{2b+2c+10n+9/2} - q^{3b+3c+13n+6} - q^{2b+4n+7/2} + q^{2b+6n+7/2} - q^{2c+4n+7/2} + q^{2c+6n+7/2} + q^{3/2}$. This is a q -analogue of above 2-variable hypergeometric identity. Setting $b = c = 0$ produces a q -analogue of 2.4 that was described by J. Guillera ([9]):

$$\sum_{n \geq 0} \frac{(-1)^n q^{7n^2} (2q^{2n+1} + 3q^{4n+2} + 4q^{6n+3} + 3q^{8n+4} - q^{10n+1} - q^{10n+3} + 2q^{10n+5} - 2q^{12n+2} + q^{12n+6} - 3q^{14n+3} - 3q^{16n+4} - 3q^{18n+5} - 2q^{20n+6} - q^{22n+7} + 1) (q; q^2)_{3n} (q; q^2)_n^2}{(q^{2n+1} + 1)^2 (q^{4n+2} + 1)^2 (q^4; q^4)_n (q^4; q^4)_{2n}^2} = \frac{(1-q^2) \sqrt{1+q^2}}{\Gamma_{q^2}\left(\frac{1}{2}\right) \Gamma_{q^4}\left(\frac{1}{2}\right)}$$

Example VII. Using $F(n, k) = \text{Gauss2F1}_q(-a - b + c - n + \frac{1}{2}, -a + c - n + \frac{1}{2}, -a - b + 2c - n + 1, 2a + 2b - 2c + d + k + 2n)$, we find, for $\Re(a) > 0$ and b, c, d in a neighbourhood of 0:

$$\sum_{n \geq 0} \frac{q^{ak} (q^{a+b-c+d+\frac{1}{2}}, q^{1/2+a+2b-c+d}; q)_k}{(q^{a+b+d+1}, q^{2a+2b-2c+d+1}; q)_k} = \sum_{n \geq 0} \frac{q^{2an+bn-2c+dn+n^2} P_q(n) (q^{a-c+\frac{1}{2}}, q, q^{a+b-c+\frac{1}{2}}, q^{a+b-c+d+\frac{1}{2}}, q^{a+2b-c+d+\frac{1}{2}}; q)_n}{(1-q^a)(1-q^{2a+2b-2c+d+2n+1})(q^{a+1}, q^{a+b+d+1}; q)_n (q^{2a+2b-2c+d+1}; q)_{2n}}$$

with $P_q(n) = -q^{2a+b+c+d+2n+\frac{1}{2}} - q^{2a+2b+c+d+2n+\frac{1}{2}} - q^{2a+2b+d+2n+1} + q^{3a+2b+d+3n+1} + q^{3a+3b+2d+3n+1} + q^{2c}$.

Intimating above example, letting $a \rightarrow 0$ from the right gives

$$\sum_{n \geq 0} \frac{q^{bn-2c+dn+n^2} \tilde{P}_q(n) (q^{1/2-c}, q^{1/2+b-c}, q^{1/2-b-c+d}; q)_n (q^{2b-c+d+\frac{1}{2}}; q)_n}{(1-q^{2b-2c+d+2n+1})(q, q^{1+b+d}; q)_n (q^{2b-2c+d+1}; q)_{2n}} = \frac{(1-q)\Gamma_q(b+d+1)\Gamma_q(2b-2c+d+1)}{\Gamma_q(b-c+d+\frac{1}{2})\Gamma_q(2b-c+d+\frac{1}{2})}$$

here $\tilde{P}_q(n) = -q^{b+c+d+2n+\frac{1}{2}} - q^{2b+c+d+2n+\frac{1}{2}} - q^{2b+d+2n+1} + q^{2b+d+3n+1} + q^{3b+2d+3n+1} + q^{2c}$. Specializing at $b = c = d = 0$ gives

$$\sum_{n \geq 0} \frac{q^{2n^2} (1 + q^{2n+1} - 2q^{4n+1})(q; q^2)_n^4}{(q^{2n+1} + 1)(q^2; q^2)_{2n} (q^2; q^2)_{2n}^2} = \frac{1 - q^2}{\Gamma_{q^2}(1/2)^2}$$

this is another q -analogue of equation 2.1 different from that in Example IV.

Example VIII. Being with $F(n, k) = \text{Gauss2F1}_q(-a - b + c - n + \frac{1}{2}, b - n + \frac{1}{2}, c + 1, d + k + 2n)$, and proceeding like previous example, then let $a \rightarrow 0^-$, we obtain

$$\sum_{n \geq 0} \frac{q^{-2b+cn+2dn+3n^2} (q^{1/2-b}, q^{1/2+b}, q^{1/2+b-c}, q^{1/2-b+c}, q^{1/2+b+d}, q^{1/2-b+c+d}; q)_n \tilde{P}_q(n)}{(1-q^{2n+1})(1-q^{d+2n+1})(1-q^{c+d+2n+1})(q, q^{1+d}, q^{1+c+d}; q)_{2n}} = \frac{(1-q)\Gamma_q(d+1)\Gamma_q(c+d+1)}{\Gamma_q(b+d+\frac{1}{2})\Gamma_q(-b+c+d+\frac{1}{2})}$$

where $\tilde{P}_q(n) = -q^{2b+c+d+2n+1} - q^{b+c+d+3n+\frac{1}{2}} + q^{2b+c+d+4n+1} + q^{2b+c+d+4n+2} + q^{2b+c+2d+4n+1} + q^{2b+c+2d+4n+2} - q^{2b+c+2d+6n+3} - q^{b+c+2d+7n+\frac{5}{2}} - q^{3b+c+2d+7n+\frac{5}{2}} - q^{b+2c+2d+7n+\frac{5}{2}} - q^{3b+c+3d+7n+\frac{5}{2}} - q^{b+2c+3d+7n+\frac{5}{2}} + q^{2b+c+2d+8n+3} + q^{2b+c+3d+8n+3} + q^{2b+2c+3d+8n+3} - q^{2b+d+2n+1} - q^{3b+d+3n+\frac{1}{2}} + q^{2b+d+4n+1} + q^{2b+d+4n+2} + q^{4b+2d+6n+2} - q^{3b+2d+7n+\frac{5}{2}} - q^{2b+2n+1} + q^{2b} + q^{2c+2d+6n+2}$. Let $b = c = d = 0$ gives

$$\sum_{n \geq 0} \frac{q^{6n^2} (3q^{2n+1} + 3q^{4n+2} - 2q^{6n+1} + q^{6n+3} - 3q^{8n+2} - 3q^{10n+3} + 1)(q; q^2)_n^6}{(q^{2n+1} + 1)^3 (q^2; q^2)_{2n}^3} = \frac{1 - q^2}{\Gamma_{q^2}(\frac{1}{2})^2}$$

this is another q -analogue of 2.3 in Example V.

Example IX. Proceeding exactly the same as in the previous two examples using $F(n, k) = \text{Seed1}_q(1/2 - a + b - n, 1 + b, c + k + n)$, we can find a two-variable q -analogue of $\sum_{n \geq 0} (-\frac{1}{8})^n \frac{(\frac{1}{2})_n^3}{(1)_n^3} (6n+1) = \frac{2\sqrt{2}}{\pi}$:

$$\sum_{n \geq 0} \frac{(-1)^n q^{bn+2cn+3n^2/2} (1 - q^{b+2c+3n+1/2}) (q^{1/2-b}, q^{1/2+b}, q^{b+2c+\frac{1}{2}}; q)_n}{(q^2, q^{2+2c}, q^{2+2b+2c}; q^2)_n} = \frac{(1-q)(1+q)^{b+2c+\frac{1}{2}} \Gamma_{q^2}(c+1) \Gamma_{q^2}(b+c+1)}{\Gamma_{q^2}(\frac{1}{2}) \Gamma_q(b+2c+\frac{1}{2})}$$

setting auxiliary parameters $b = c = 0$ will recover a formula found in [11].

Example X. Using $F(n, k) = \text{Gauss2F1}_q(-a - b + c + \frac{1}{2}, b - n + \frac{1}{2}, c + n + 1, d + k + n)$ and proceed exactly as in Example VI, letting $a \rightarrow 0^-$ produces the formula

$$\sum_{n \geq 0} \frac{(-1)^n q^{bn-2b+2dn+\frac{3n^2}{2}} P_q(n) (q^{\frac{1}{2}-b}, q^{b+\frac{1}{2}}, q^{-b+c+d+\frac{1}{2}}; q)_n (q^{-b+c+\frac{1}{2}}; q)_{2n}}{(1-q^{2n+1})(q, q^{c+d+1}; q)_{2n} (1-q^{c+d+2n+1})} = \frac{(1-q)\Gamma_q(d+1)\Gamma_q(c+d+1)}{\Gamma_q(b+d+\frac{1}{2})\Gamma_q(-b+c+d+\frac{1}{2})}$$

with $P_1(n) = -q^{2b+c+d+2n+1} - q^{b+c+d+3n+\frac{1}{2}} + q^{2b+c+2d+3n+1} + q^{2b+c+d+4n+1} + q^{2b+c+d+4n+2} - q^{b+c+2d+4n+\frac{3}{2}} - q^{b+2c+2d+7n+\frac{5}{2}} - q^{3b+d+2n+\frac{1}{2}} + q^{2b+d+3n+1} - q^{2b+2n+1} + q^{2b} + q^{2c+2d+6n+2}$.

Using another WZ-pair $F(n, k) = \text{Seed3}_q(a + b + c + \frac{1}{2}, b + \frac{1}{2}, c - n + \frac{1}{2}, d + k + n)$ and proceed exactly as in Example VI, letting $a \rightarrow 0^-$ produces another formula:

$$\sum_{n \geq 0} \frac{(-1)^n q^{2cn+2dn+n^2-3} P_2(n) (q^{b+\frac{1}{2}}, q^{b+2d+\frac{1}{2}}; q)_{2n} (q^{2b+2c+2d+1}, q^{1-2c}; q^2)_n}{(1-q^{b+d+2n+1})(q^{b+d+2n+1} + 1)(q^{2d+2}; q^2)_n (1-q^{b+2c+2d+2n+\frac{1}{2}})(q^2; q^2)_n (q^{2b+2d+2}; q^2)_{2n} (q^{b+2c+2d+\frac{1}{2}}; q)_{2n}} = \frac{(1-q^2)(1+q)^{-2c}\Gamma_{q^2}(d+1)\Gamma_{q^2}(b+d+1)\Gamma_q(b+2c+2d+\frac{1}{2})}{\Gamma_q(b+2d+\frac{1}{2})\Gamma_{q^2}(c+d+\frac{1}{2})\Gamma_{q^2}(b+c+d+\frac{1}{2})}$$

with $P_2(n) = -q^{2b+2c+2d+4n+4} - q^{2b+2c+4d+4n+4} + q^{2b+2c+4d+6n+4} + q^{3b+2c+4d+6n+\frac{9}{2}} + q^{3b+2c+4d+6n+\frac{11}{2}} - q^{4b+2c+6d+10n+6} - q^{b+2d+4n+\frac{7}{2}} - q^{b+2d+4n+\frac{9}{2}} - q^{2b+2d+4n+5} + q^{2b+2d+6n+5} + q^{2b+4d+6n+5} + q^3$. Making all auxiliary variables to be 0 in two formulas above, we arrive at respectively:

$$\sum_{n \geq 0} \frac{(-1)^n q^{3n^2} (2q^{2n+1} - q^{4n+1} + q^{4n+2} - q^{6n+1} - q^{8n+2} - q^{10n+3} + 1)(q; q^2)_{2n} (q; q^2)_n^3}{(q^{2n+1} + 1)^2 (q^2; q^2)_n (q^2; q^2)_{2n}^2} = \frac{1 - q^2}{\Gamma_{q^2}(\frac{1}{2})^2}$$

$$\sum_{n \geq 0} \frac{(-1)^n q^{2n^2} (q^{4n+1} + q^{4n+2} - q^{8n+1} - q^{8n+2} - q^{12n+3} + 1)(q; q^2)_{2n} (q^2; q^4)_n^2}{(q^{4n+2} + 1)(q^4; q^4)_{2n} (q^4; q^4)_n^2} = \frac{1 - q^4}{\Gamma_{q^4}(\frac{1}{2})^2}$$

they are both q -analogues of $\sum_{n \geq 0} (-\frac{1}{2})^n \frac{(\frac{1}{2})(\frac{1}{4})(\frac{3}{4})}{(1)_3^n} (20n + 3) = \frac{8}{\pi}$.

Example XI. We can derive a q -analogue of

$$\sum_{n \geq 0} \left(\frac{1}{9}\right)^n \frac{(\frac{1}{2})_n (\frac{1}{4})_n (\frac{3}{4})_n}{(1)_3^n} (8n + 1) = \frac{16\sqrt{3}}{\pi}$$

using $F(n, k) = \text{Seed10}_q(a + 3b - n + \frac{1}{2}, 2b + 1, k + c + n)$. Similar to Example IV, setting $a = 0$ will kill $\sum_{k \geq 0} F(0, k)$, leaving us with

$$\sum_{n \geq 0} \frac{q^{n^2} (1 - q^{3b+3c+4n+\frac{1}{2}})(q^{1/2-3b}, q^{1/2+3b}; q)_n (q^{3b+3c+1/2}; q)_{2n}}{(q; q)_{2n} (q^{3c+3}, q^{3+6b+3c}; q^3)_n} = \frac{(1-q)(q^2+q+1)^{3b+3c+1/2}\Gamma_{q^3}(c+1)\Gamma_{q^3}(b+c+\frac{1}{2})\Gamma_{q^3}(2b+c+1)}{\Gamma_{q^3}(\frac{2}{3})\Gamma_{q^3}(\frac{1}{3})\Gamma_q(3b+3c+\frac{1}{2})} \sum_{n \geq 0} \frac{q^{n^2} (q^{1/2-3b}, q^{1/2+3b}; q)_n}{(q; q)_{2n}}$$

we remark the terms $\Gamma_{q^3}(\frac{2}{3})\Gamma_{q^3}(\frac{1}{3})$ in denominator again comes from Proposition 1.5 when calculating $\lim_{k \rightarrow \infty} G(n, k) = g(n)$. We evaluate in the appendix (Lemma A.2) the last sum on RHS, whence

$$\sum_{n \geq 0} \frac{q^{n^2} (1 - q^{3b+3c+4n+\frac{1}{2}})(q^{1/2-3b}, q^{1/2+3b}; q)_n (q^{3b+3c+1/2}; q)_{2n}}{(q; q)_{2n} (q^{3c+3}, q^{3+6b+3c}; q^3)_n} = \frac{(1-q)(q^2+q+1)^{3b+3c+\frac{1}{2}}\Gamma_{q^3}(c+1)\Gamma_{q^3}(b+c+\frac{1}{2})\Gamma_{q^3}(2b+c+1)}{\Gamma_{q^3}(\frac{1}{2}-b)\Gamma_{q^3}(b+\frac{1}{2})\Gamma_q(3b+3c+\frac{1}{2})}$$

Setting $b = c = 0$ gives

$$\sum_{n \geq 0} \frac{q^{2n^2} (1 - q^{8n+1})(q; q^2)_{2n} (q; q^2)_n^2}{(q^2; q^2)_{2n} (q^6; q^6)_n^2} = \frac{(1 - q^2)\sqrt{q^4 + q^2 + 1}}{\Gamma_{q^2}(\frac{1}{2})\Gamma_{q^6}(\frac{1}{2})}$$

this is a q -analogue of $1/\pi$ -formula cited at beginning. A one-variable special case of above two-variable formula is recorded in [13].

Example XII. Using the WZ-pair $F(n, k) = \text{Seed10}_q(a + 3b - 2n + \frac{1}{2}, 2b + 1, c + k + n)$, and proceed like above example, one shows the formula

$$\sum_{n \geq 0} \left(-\frac{1}{48}\right)^n \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{(1)_n^3} (28n + 3) = \frac{16\sqrt{3}}{\pi}$$

has a two-variable q -analogue:

$$\begin{aligned} & \sum_{n \geq 0} \frac{(-1)^n q^{3bn-b+3cn-c+\frac{5n^2}{2}-\frac{5}{2}} P_q(n) (q^{\frac{1}{2}-3b}, q^{3b+\frac{1}{2}}; q)_{2n} (q^{3/2-3b-3c}; q^3)_n (q^{3b+3c+1/2}; q)_n}{(q^{2b+2c+1} - q^{-b-c+3n-\frac{1}{2}}) (q; q)_{4n+2} (q^{3c+3}, q^{6b+3c+3}; q^3)_n} \\ &= \frac{(1-q)(q^2+q+1)^{3b+3c+\frac{1}{2}} \Gamma_{q^3}(c+1) \Gamma_{q^3}(b+c+\frac{1}{2}) \Gamma_{q^3}(2b+c+1)}{\Gamma_{q^3}\left(\frac{1}{3}\right) \Gamma_{q^3}\left(\frac{2}{3}\right) \Gamma_q\left(3b+3c+\frac{1}{2}\right)} \\ & \times \sum_{n \geq 0} \frac{q^{4n^2} (-q^{3b+4n+\frac{5}{2}} - q^{6b+6n+2} + q^{3b+8n+\frac{5}{2}} + q^{3b+8n+\frac{7}{2}} + q^{3b+\frac{1}{2}} - q^{6n+2}) (q^{\frac{1}{2}-3b}, q^{3b+\frac{1}{2}}; q)_{2n}}{(q; q)_{4n+2}} \end{aligned}$$

where $P_q(n) = -q^{6b+3c+4n+6} - q^{9b+6c+5n+9/2} - q^{9b+6c+5n+11/2} - q^{3b+3c+6n+11/2} - q^{9b+3c+6n+11/2} + q^{6b+6c+7n+6} + q^{12b+6c+7n+6} + q^{6b+3c+8n+3} + q^{6b+3c+8n+4} + q^{6b+3c+8n+6} + q^{6b+3c+8n+7} + q^{9b+6c+9n+11/2} - q^{3b+3c+10n+9/2} - q^{9b+3c+10n+9/2} - q^{6b+3c+12n+4} - q^{9b+6c+13n+15/2} + q^{6b+3c+16n+6} + q^{6b+3c+4} - q^{3b+3n+5/2} + q^{3b+7n+9/2} + q^{6b+9n+4} - q^{3b+11n+9/2} - q^{3b+11n+11/2} + q^{9n+4}$. The sum in n on RHS will be evaluated in appendix (Lemma A.4), giving

$$\begin{aligned} & \sum_{n \geq 0} \frac{(-1)^n q^{3bn-b+3cn-c+\frac{5n^2}{2}} P_q(n) (q^{\frac{1}{2}-3b}, q^{3b+\frac{1}{2}}; q)_{2n} (q^{3/2-3b-3c}; q^3)_n (q^{3b+3c+1/2}; q)_n}{(q^{2b+2c+1} - q^{-b-c+3n-\frac{1}{2}}) (q; q)_{4n+2} (q^{3c+3}, q^{6b+3c+3}; q^3)_n} \\ &= \frac{(1-q)q^{3b+3} (q^2+q+1)^{3b+3c+\frac{1}{2}} \Gamma_{q^3}(c+1) \Gamma_{q^3}(b+c+\frac{1}{2}) \Gamma_{q^3}(2b+c+1)}{\Gamma_{q^3}\left(\frac{1}{2}-b\right) \Gamma_{q^3}\left(b+\frac{1}{2}\right) \Gamma_q\left(3b+3c+\frac{1}{2}\right)} \end{aligned}$$

Substituting $b = c = 0$:

$$\begin{aligned} & \sum_{n \geq 0} \frac{(-1)^n q^{5n^2} (q^{2n+1} + q^{4n+1} + q^{4n+2} + q^{6n+2} + q^{6n+3} + q^{8n+2} + q^{8n+3} - q^{10n+1} + q^{10n+4} - q^{12n+2} - q^{12n+3} - q^{14n+2} - q^{14n+3} - q^{16n+3} - q^{16n+4} - q^{18n+4} - q^{20n+5} + 1) (q; q^2)_n (q^3; q^6)_n (q; q^2)_{2n}^2}{(q^{2n+1} + 1) (q^{4n+1} + 1) (q^{4n+2} + 1) (q^2; q^2)_{4n} (q^6; q^6)_{2n}^2} \\ &= \frac{(1-q^2) \sqrt{q^4 + q^2 + 1}}{\Gamma_{q^2}\left(\frac{1}{2}\right) \Gamma_{q^6}\left(\frac{1}{2}\right)} \end{aligned}$$

this is a q -analogue of our initial formula.

Among 36 rational $1/\pi$ -series (tabulated in [3]), only eight can be proved using our listed WZ-seeds. We already found (at least one) q -analogues for seven of them, while that for the remaining one $\sum_{n \geq 0} (-1)^n \frac{\left(\frac{1}{2}\right)_n^3}{(1)_n^3} (4n + 1) = \frac{2}{\pi}$ is rather trivial and well-known.

Example XIII. The following example has a somewhat different nature than others in this article. Inspired by a conjecture of Z.W. Sun, the author proved in [2], that

$$(2.5) \quad \sum_{n \geq 0} \frac{2^{-3n} (4a^2 + 8an - 4c^2 + 4c + 20n^2 + 8n - 1) \left(-a - c + \frac{1}{2}\right)_n \left(a - c + \frac{1}{2}\right)_{3n}}{(-c + 2n + 1) (2a - 2c + 2n + 1) (1)_n (1 - c)_{2n} \left(a - c + \frac{1}{2}\right)_n} = 0 \quad a, c \in \mathbb{C}$$

It turns out this formula has a "formal" q -analogue:

$$(2.6) \quad \sum_{n \geq 0} \frac{q^{-2an-a-c-1/2} (-q^{a-c+4n+3/2} - q^{2a-2c+5n+3} + q^{2a-2c+7n+3} - q^{a+c+1/2} + q^{2a+n+1} + q^{n+1}) (q^{-a-c+1/2}; q)_n (q^{a-c+1/2}; q)_{3n}}{(1 - q^{-c+2n+1}) (q^{-c+2n+1} + 1) (1 - q^{a-c+n+1/2}) (q^2; q^2)_n (q^{2-2c}; q^2)_{2n} (q^{a-c+1/2}; q)_n} \\ = - \frac{(1+q)^{-a-c-1/2} \Gamma_{q^2}(-a) \Gamma_{q^2}(1-c)}{\Gamma_{q^2}(\frac{1}{2}) \Gamma_q(-a-c+\frac{1}{2})} \quad \Re(a) < 0, c \in \mathbb{C}, 0 < q < 1$$

note the summand in 2.6 tends to that of 2.5 when $q \rightarrow 1$, whereas RHS does not. As a result, 2.6 provides no sensible q -analogue for $\sum_{n \geq 0} \left(\frac{27}{32}\right)^n \frac{(1/6)_n (5/6)_n}{(1)_n^2} \frac{10n-1}{2n+1} = 0$, which is obtained by setting $a = c = 0$ in 2.5.

To establish 2.6, one simply uses the q -version of WZ-seed that proves 2.5, i.e.,

$$F(n, k) = \text{Seed1}_q \left(a + c - n + \frac{1}{2}, c - n + 1, k + 2n \right) \\ = \frac{q^{-2a(k+2n)} (q+1)^{a-c-2k-3n-\frac{3}{2}} \Gamma_q \left(-a + c - n + \frac{1}{2} \right) \Gamma_q \left(a + c + 2k + 3n + \frac{1}{2} \right)}{\Gamma_{q^2}(-a) \Gamma_{q^2}(k+2n+1) \Gamma_q \left(a + c - n + \frac{1}{2} \right) \Gamma_{q^2}(c+k+n+1)}$$

When $\Re(a) < 0$ and $c \in \mathbb{C}$, analytic prerequisites of Proposition 1.3 are met, and $g(n) = 0$, therefore

$$\sum_{k \geq 0} F(0, k) = \sum_{n \geq 0} G(n, 0) + \lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$$

in our case $\sum_{k \geq 0} F(0, k)$ can be summed using q -Gauss theorem, giving RHS of 2.5, and $\sum_{n \geq 0} G(n, 0)$ is RHS. In the classical case (i.e. $q = 1$), $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k) \neq 0$, whereas for $0 < q < 1$, the situation is different

Lemma 2.1. *For $\Re(a) < 0$ and $0 < q < 1$, $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$ vanishes.*

Proof sketch. Since we know the limit exists, we concentrate on n even, that is, we can replace n by $2n$, then $\sum_{k \geq 0} F(n, k)$ is, after dropping factors irrelevant to n or k :

$$q^{-8an} \frac{\Gamma_q \left(-a + c - 2n + \frac{1}{2} \right) \Gamma_{q^2} \left(\frac{a}{2} + \frac{c}{2} + 3n + \frac{1}{4} \right) \Gamma_{q^2} \left(\frac{a}{2} + \frac{c}{2} + 3n + \frac{3}{4} \right)}{\Gamma_{q^2} \left(\frac{1}{2} \right) \Gamma_{q^2}(-a) \Gamma_{q^2}(4n+1) \Gamma_{q^2}(c+2n+1) \Gamma_q \left(a + c - 2n + \frac{1}{2} \right)} \sum_{k \geq 0} \frac{(q^{3/2+a+c+6n}, q^{1/2+a+c+6n}; q^2)_k}{(q^{2+2c+4n}, q^{2+8n}; q^2)_k} q^{-2ak}$$

In the classical case, finding asymptotic leading term of $\sum_{k \geq 0} \dots$ as $n \rightarrow \infty$ is a tricky problem, but for q -case, the leading term is simply $\sum_{k \geq 0} q^{-2ak} = O(1)$. Asymptotic of remaining q -gamma terms can be found using Proposition 1.5, after some calculation, we found above is $O(q^{-12an})$, so when $\Re(a) < 0$, it tends to 0. \square

Example XIV. The Ramanujan-type $1/\pi^2$ series are among first discovered and proved ([7]):

$$\sum_{n \geq 0} \frac{(-1)^n}{2^{2n}} \frac{(\frac{1}{2})_n^5}{(1)_n^5} (20n^2 + 8n + 1) = \frac{8}{\pi^2} \\ \sum_{n \geq 0} \frac{(-1)^n}{2^{10n}} \frac{(\frac{1}{2})_n^5}{(1)_n^5} (820n^2 + 180n + 13) = \frac{128}{\pi^2} \\ \sum_{n \geq 0} \left(\frac{1}{2^4}\right)^n \frac{(\frac{1}{2})_n^3 (\frac{1}{4})_n (\frac{3}{4})_n}{(1)_n^5} (120n^2 + 34n + 3) = \frac{32}{\pi^2}$$

have respective q -deformations

$$\begin{aligned} \sum_{n \geq 0} \frac{(-1)^n q^{n^2} (q^{2n+1} - 4q^{4n+1} + q^{6n+1} + q^{8n+2} + 1)(q; q^2)_n^6}{(q^{2n+1} + 1)(q^2; q^2)_{2n} (q^2; q^2)_n^4} &= \frac{(1 - q^2)^2}{\Gamma_{q^2}(1/2)^4} \\ \sum_{n \geq 0} \frac{(-1)^n q^{5n^2} P_1(n)(q; q^2)_n^{10}}{(q^{2n+1} + 1)^5 (q^2; q^2)_{2n}^5} &= \frac{(1 - q^2)^2}{\Gamma_{q^2}(1/2)^4} \\ \sum_{n \geq 0} \frac{q^{2n^2} P_2(n)(q; q^2)_{2n} (q; q^2)_n^6}{(q^{2n+1} + 1)^3 (q^2; q^2)_n^2 (q^2; q^2)_{2n}^3} &= \frac{(1 - q^2)^2}{\Gamma_{q^2}(1/2)^4} \end{aligned}$$

where $P_1(n) = 5q^{2n+1} + 10q^{4n+2} - 5q^{6n+1} + 10q^{6n+3} - 15q^{8n+2} + 5q^{8n+4} + q^{10n+1} - 24q^{10n+3} + q^{10n+5} + 5q^{12n+2} - 15q^{12n+4} + 10q^{14n+3} - 5q^{14n+5} + 10q^{16n+4} + 5q^{18n+5} + q^{20n+6} + 1$, $P_2(n) = 3q^{2n+1} - 3q^{4n+1} + 3q^{4n+2} - 2q^{6n+1} - 3q^{6n+2} + q^{6n+3} + q^{8n+1} - 3q^{8n+2} - 2q^{8n+3} + 3q^{10n+2} - 3q^{10n+3} + 3q^{12n+3} + q^{14n+4} + 1$. They can be proved, just like their classical counterparts, using respectively the following $F(n, k)$

$$\text{Dougall5F4}_q(1/2 - n, 1/2 - n, 1/2 - n, 1/2 - n, k + n)$$

$$\text{Dougall5F4}_q(1/2 - n, 1/2 - n, 1/2 - n, 1/2 - n, k + 2n)$$

$$\text{Dougall5F4}_q(1/2 - n, 1/2 - n, 1/2 - n, 1/2, k + 2n)$$

with $\sum_{k \geq 0} F(a, k) = \sum_{n \geq 0} G(n + a, 0)$ and then proceed exactly like Example VI. All three have a four-parameter deformation, by putting parameters into Dougall5F4.

Example XV. The following $1/\pi^4$ were conjectured by J. Cullen and Y. Zhao respectively, and are proved recently by the author in [2].

$$\begin{aligned} \sum_{n \geq 0} \left(\frac{1}{2^{12}}\right)^n \frac{\left(\frac{1}{2}\right)_n^7 \left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{(1)_n^9} (43680n^4 + 20632n^3 + 4340n^2 + 466n + 21) &= \frac{2048}{\pi^4} \\ \sum_{n \geq 0} \left(-\frac{3^3}{2^8}\right)^n \frac{\left(\frac{1}{2}\right)_n^5 \left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n \left(\frac{1}{3}\right)_n \left(\frac{2}{3}\right)_n}{(1)_n^9} (4528n^4 + 3180n^3 + 972n^2 + 147n + 9) &= \frac{768}{\pi^4} \end{aligned}$$

They have respective q -deformations:

$$\begin{aligned} \sum_{n \geq 0} \frac{q^{4n^2} P_1(n)(q^2; q^2)_{4n} (q; q^2)_n^{16}}{(q^{4n} + 1)(q^{2n+1} + 1)^9 (q^2; q^2)_{2n}^{10}} &= \frac{(1 - q^2)^4}{\Gamma_{q^2}(1/2)^8} \\ \sum_{n \geq 0} \frac{(-1)^n q^{n^2} P_2(n)(q; q^2)_{2n} (q^2; q^2)_{3n} (q; q^2)_n^{10}}{(q^{2n} - q^n + 1)(q^{2n} + q^n + 1)(q^{2n+1} + 1)^5 (q^2; q^2)_n^5 (q^2; q^2)_{2n}^5} &= \frac{(1 - q^2)^4}{\Gamma_{q^2}(1/2)^8} \end{aligned}$$

where $P_1(n) = q^{4n} + 9q^{2n+1} + 36q^{4n+2} - 7q^{6n+1} + 84q^{6n+3} - 63q^{8n+2} + 126q^{8n+4} - 231q^{10n+3} + 126q^{10n+5} + 35q^{12n+2} - 448q^{12n+4} + 84q^{12n+6} + 187q^{14n+3} - 560q^{14n+5} + 36q^{14n+7} - 9q^{16n+2} + 509q^{16n+4} - 441q^{16n+6} + 9q^{16n+8} - 65q^{18n+3} + 805q^{18n+5} - 225q^{18n+7} + q^{18n+9} + q^{20n+2} - 225q^{20n+4} + 805q^{20n+6} - 65q^{20n+8} + 9q^{22n+3} - 441q^{22n+5} + 509q^{22n+7} - 9q^{22n+9} + 36q^{24n+4} - 560q^{24n+6} + 187q^{24n+8} + 84q^{26n+5} - 448q^{26n+7} + 35q^{26n+9} + 126q^{28n+6} - 231q^{28n+8} + 126q^{30n+7} - 63q^{30n+9} + 84q^{32n+8} - 7q^{32n+10} + 36q^{34n+9} + q^{34n+11} + 9q^{36n+10} + q^{38n+11} + 1$ and $P_2(n) = q^{2n} + q^{4n} + 5q^{2n+1} - 5q^{4n+1} + 10q^{4n+2} - 5q^{6n+1} - 16q^{6n+2} + 10q^{6n+3} - q^{8n+1} - 11q^{8n+2} - 25q^{8n+3} + 5q^{8n+4} + 19q^{10n+2} - 14q^{10n+3} - 15q^{10n+4} + q^{10n+5} + q^{12n+2} + 46q^{12n+3} - 4q^{12n+4} - 4q^{12n+5} - 4q^{14n+2} - 4q^{14n+3} + 46q^{14n+4} + q^{14n+5} + q^{16n+2} - 15q^{16n+3} - 14q^{16n+4} + 19q^{16n+5} + 5q^{18n+3} - 25q^{18n+4} - 11q^{18n+5} - q^{18n+6} + 10q^{20n+4} - 16q^{20n+5} - 5q^{20n+6} + 10q^{22n+5} - 5q^{22n+6} + q^{22n+7} + 5q^{24n+6} + q^{24n+7} + q^{26n+7} + 1$. They can be proved exactly like their classical counterparts (Examples I and II of [2]) using q -version of corresponding $F(n, k)$:

$$\text{Dougall7F6}(1/2 - n, 1/2 - n, 1/2 - n, 1/2 - n, 1/2 - n, k + 2n) \quad \text{Dougall7F6}(1/2 + n, 1/2, 1/2, 1/2, 1/2, k + n)$$

respectively. They both have a 5-parameter generalization, which is unbearably long even in classical case.

For the one example below, we shall require some transformation formula of very-well-poised ${}_8\phi_7$ series (Gasper & Rahman (2.10.1), (2.10.10) [5]): Let

$$V_q(a; b, c, d, e, f) = \sum_{k \geq 0} \frac{1 - q^{2k+a}}{1 - q^a} \frac{(q^a, q^b, q^c, q^d, q^e, q^f; q)_k}{(q, q^{1+a-b}, q^{1+a-c}, q^{1+a-d}, q^{1+a-e}, q^{1+a-f}; q)_k} q^{(2a+2-b-c-d-e-f)k}$$

then assuming convergence, we have $V_q(a; b, c, d, e, f)$ actually equals to both:

$$\frac{\Gamma_q(a-e+1)\Gamma_q(a-f+1)\Gamma_q(2a-b-c-d+2)\Gamma_q(2a-b-c-d-e-f+2)}{\Gamma_q(a+1)\Gamma_q(a-e-f+1)\Gamma_q(2a-b-c-d-e+2)\Gamma_q(2a-b-c-d-f+2)} \\ \times V_q(2a-b-c-d+1; a-c-d+1, a-b-d+1, a-b-c+1, e, f)$$

and

$$\frac{\Gamma_q(a-b+1)\Gamma_q(a-c+1)\Gamma_q(a-d+1)\Gamma_q(a-e+1)\Gamma_q(3a-b-c-d-e-2f+3)\Gamma_q(2a-b-c-d-e-f+2)}{\Gamma_q(a+1)\Gamma_q(f)\Gamma_q(2a-b-c-d-f+2)\Gamma_q(2a-b-c-e-f+2)\Gamma_q(2a-b-d-e-f+2)\Gamma_q(2a-c-d-e-f+2)} \\ \times V_q(3a-b-c-d-e-2f+2; a-b-f+1, a-c-f+1, a-d-f+1, a-e-f+1, 2a-b-c-d-e-f+2)$$

Example XVI. I claim the follow identity holds²:

$$\frac{(1-q^2)^4}{(1-q)^2\Gamma_{q^2}(1/2)^6} \sum_{k \geq 0} (-1)^k q^{3k(k+1)} \frac{1+q^{1+2k}}{1+q} \frac{1-q}{1-q^{1+2k}} \\ = \sum_{n \geq 0} \frac{q^{2n}(3q^{2n+1} - 4q^{4n+1} + 2q^{4n+2} - 4q^{6n+1} - 4q^{6n+2} + 2q^{8n+1} - 4q^{8n+2} + 3q^{10n+2} + 3q^{12n+3} + 3)(q^2; q^2)_{3n}(q; q^2)_n^6}{(q^{2n} + q^{4n} + 1)(q^{2n+1} + 1)^3(q^2; q^2)_n^3(q^2; q^2)_{2n}^3}$$

It seems possible to express the k -summation in terms of q -gamma, we don't do it here. This is a q -analogue of

$$\sum_{n \geq 0} \left(\frac{3}{4}\right)^n \frac{\left(\frac{1}{2}\right)_n^3 \left(\frac{1}{3}\right)_n \left(\frac{2}{3}\right)_n}{(1)_n^5} (74n^2 + 27n + 3) = \frac{48}{\pi^2}$$

This classical formula is proved by J. Guillera [8]. Let $F(n, k) = \text{Dougall5F4}_q(1/2 - n, 1/2 - n, 1/2 - n, n, 2n + k)$, then one finds out $g(n) = 0$ and applying 1.3 to $F(n + a, k)$, we have

$$\sum_{k \geq 0} \frac{q^{k/2} (1 - q^{3a+2k+\frac{1}{2}}) (q^{3a+1}; q)_k (q^{a+\frac{1}{2}}; q)_k^3}{(1 - q^{k+\frac{1}{2}}) (1 - q^{3a+k}) (\sqrt{q}; q)_k (q^{2a+1}; q)_k^3} \\ = \sum_{n \geq 0} \frac{q^{a+n} P_q(n) (q^{3a+1}; q)_{3n} (q^{a+\frac{1}{2}}; q)_n^6}{(1 - q^a)^3 (q^{a+n} + q^{2a+2n} + 1) (1 - q^{2a+2n+1})^3 (q^{a+1}; q)_n^3 (q^{2a+1}; q)_{2n}^3} + \frac{1}{A} \lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n + a, k)$$

here $P_q(n) = 6q^{a+n+\frac{1}{2}} - 2q^{2a+2n+1} + 4q^{2a+2n+\frac{1}{2}} - 8q^{3a+3n+1} + 4q^{3a+3n+\frac{1}{2}} - 8q^{4a+4n+1} - 3q^{4a+4n+\frac{1}{2}} + 3q^{5a+5n+1} + 8q^{5a+5n+\frac{1}{2}} + 2q^{5a+5n+\frac{5}{2}} + 8q^{6a+6n+2} - 4q^{6a+6n+\frac{5}{2}} + 2q^{7a+7n+2} - 4q^{7a+7n+\frac{5}{2}} - 6q^{8a+8n+\frac{5}{2}} + 3q^{9a+9n+3} - 3$ and $A = \frac{q^a(1-q^a)\Gamma_q(1-a)^2\Gamma_q(a+\frac{1}{2})^4\Gamma_q(3a+1)}{\Gamma_q(\frac{1}{2})^2\Gamma_q(\frac{1}{2}-a)^2\Gamma_q(a+1)\Gamma_q(2a+1)^3}$.

From this expression, it is immediately clear $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n + a, k)$ must not be 0: LHS is analytic at $a = 0$, but the summation in n has order 3 pole at $a = 0$.

Lemma 2.2. In above notation, $\lim_{n \rightarrow \infty} F(n + a, k)$ equals

$$\frac{q^a \Gamma_q(1-a)^2 \Gamma_q(a)^2}{\Gamma_q(\frac{3}{2})^2 \Gamma_q(\frac{1}{2}-a)^2 \Gamma_q(a+\frac{1}{2})^2} \times \sum_{k \geq 0} (-1)^k q^{3k(k+1)/2} \frac{1+q^{1/2+k}}{1+q^{1/2}} \frac{1-q^{1/2}}{1-q^{1/2+k}}$$

Assuming this, finding leading coefficient as $a \rightarrow 0$ yields the identity claimed at start.

²note that RHS is the only WZ-method-derived series in this article without quadratic factor q^{Cn^2}

Proof of lemma. The idea is to first rewrite $\sum_{k \geq 0} F(n+a, k)$ into a very-well-poised ${}_8\phi_7$ series, then apply various transformation formulas, hoping to find a form in which the limiting behaviour as $n \rightarrow \infty$ is clear. Firstly, one sees $\sum_{k \geq 0} F(n+a, k) =$ equals

$$\frac{q^{a+n}(1-q^{3a+3n+\frac{1}{2}})\Gamma_q(-a-n+1)^2\Gamma_q(3a+3n)\Gamma_q(a+n+\frac{1}{2})^4}{(1-q)\Gamma_q(\frac{1}{2})\Gamma_q(\frac{3}{2})\Gamma_q(-a-n+\frac{1}{2})^2\Gamma_q(a+n)\Gamma_q(2(a+n)+1)^3} \times V_q(3a+3n+\frac{1}{2}, 1, a+n+\frac{1}{2}, a+n+\frac{1}{2}, a+n+\frac{1}{2}, 3a+3n)$$

then using above transformation, this V_q series equals

$$\frac{\Gamma_q(\frac{1}{2})\Gamma_q(2a+2n+1)^3\Gamma_q(3a+3n+\frac{1}{2})V_q(1, \frac{1}{2}, \frac{1}{2}, -a-n+1, -a-n+1, -a-n+1)}{\Gamma_q(\frac{3}{2})\Gamma_q(a+n+1)^3\Gamma_q(3a+3n)\Gamma_q(3a+3n+\frac{3}{2})}$$

now this V_q series, i.e.,

$$\sum_{k \geq 0} \frac{1+q^{1/2+k}}{1+q^{1/2}} \frac{1-q^{1/2}}{1-q^{1/2+k}} \frac{(q^{1-a-n}; q)_k^3}{(q^{1+a+n}; q)_k^3} (q^{3a+3n})^k$$

has desired property: it tends to the series in statement of lemma as $n \rightarrow \infty$. Calculating explicit asymptotic expansion of q -gamma function we picked up along the way (using Proposition 1.5) gives the statement. \square

If one uses a more general $F(n, k) = \text{Dougall}5F_4(q^{1/2-n}, 1/2+b-n, 1/2+c-n, n+d, e+2n+k)$ instead, then one finds a 4-variable generalization of the identity at start, this method of finding the limit is insensitive to these auxiliary variables.

Example XVII. Using our philosophy on correspondence between classical and q -hypergeometric identities, a q -analogue of³

$$\sum_{n \geq 0} \left(\frac{1}{2^6}\right)^n \frac{(\frac{1}{2})_n^7}{(1)_n^7} (168n^3 + 76n^2 + 14n + 1) = \frac{32}{\pi^3}$$

should have the form

$$(2.7) \sum_{n \geq 0} \frac{q^{2n^2}(1-q^{6n+1})(1+q^{4n}+q^{2n+1}-6q^{6n+1}+q^{8n+2}+q^{10n+1}+q^{12n+2})(q^4; q^4)_{2n}(q; q^2)_n^8}{(q^{4n}+1)(q^{2n+1}+1)(q^2; q^2)_{2n}^2(q^4; q^4)_n^6} = ???$$

We explain our reasoning. Let a be in a neighbourhood of 0, let

$$F(n, k) = \text{Seed}9_q(1-2n, n+\frac{1}{2}, n+\frac{1}{2}, k+n)$$

its classical version can be used to prove the $32/\pi^3$ -identity (see Appendix). Using Proposition 1.3,

$$\sum_{k \geq 0} F(a, k) + \sum_{n \geq 0} g(n+a) = \sum_{n \geq 0} G(n+a, 0) + \lim_{n \rightarrow \infty} \sum_{k \geq 0} F(a+n, k)$$

becomes

$$\begin{aligned} & \sum_{k \geq 0} \frac{q^{2k}(1-q^{2k+1})(q^{2k+1}+1)(q^{2-2a}; q^2)_k^2 (q^{a+\frac{1}{2}}; q)_{2k}^2}{(1-q^{-a+2k+\frac{1}{2}})^2 (1-q^{-a+2k+\frac{3}{2}})^2 (q^{\frac{1}{2}-a}; q)_{2k}^2 (q^{2a+2}; q^2)_k^2} \\ & + \frac{2(q+1)^{4a} (q^{2a}+1) \Gamma_q(\frac{1}{2}-a)^2 \Gamma_{q^2}(a+1)^2}{(q-q^{2a})(q^{2a}-1)^3 \Gamma_{q^2}(1-a)^2 \Gamma_q(a+\frac{1}{2})^2} \times \sum_{n \geq 0} \frac{q^{3a+n-\frac{1}{2}}(q^{3a+3n+\frac{1}{2}}-1)(q^{4a+2}; q^2)_{2n}(q^{a+\frac{1}{2}}; q)_n^4}{(q^{a+n+\frac{1}{2}}+1)(q^{2a+2n}+1)(q^{2a+1}; q)_{2n}^2 (q^{2a+2}; q^2)_n^2} = \\ & \frac{(q^{2a}+1)q^{2an+4a+n^2-1}(1-q^{3a+3n+\frac{1}{2}})(q^{a+n+\frac{1}{2}}+q^{2a+2n}-6q^{3a+3n+\frac{1}{2}}+q^{4a+4n+1}+q^{5a+5n+\frac{1}{2}}+q^{6a+6n+1}+1)}{\sum_{n \geq 0} \frac{(q^{4a+2}; q^2)_{2n}(q^{a+\frac{1}{2}}; q)_n^8}{(1-q^{2a})^5 (q^{2a}-q)(q^{a+n+\frac{1}{2}}+1)(q^{2a+2n}+1)(q^{2a+1}; q)_{2n}^2 (q^{2a+2}; q^2)_n^6}} \end{aligned}$$

³this hypergeometric equality was a well-known conjecture due to B. Gourevich, this is recently proved by author in [2]

$$+ \frac{1}{A} \lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$$

with A again being a q -gamma factor. First term on LHS has no pole at $a = 0$, second term on LHS has pole of order 3, first term on RHS has pole of order 5, so $L := \lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n + a, k)$ cannot vanish. Extracting the principal part of series on RHS gives 2.7.

In the classical case (i.e. $q = 1$), L is evaluated in appendix, but that method does not extend to q -case. Nonetheless the author still believes L can be evaluated using q -gamma function.

Problem 2.3. Let $F(n, k)$ be as above, a be in a neighbourhood of 0, what is

$$\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(a + n, k)?$$

We know the limit is finite and non-zero.

Inability to evaluate this limit prevents us from figuring out the RHS of 2.7.

APPENDIX A. SEVERAL q -HYPERGEOMETRIC SUMMATIONS

We supply proofs of some hypergeometric summations used in Examples IV, XI, XII.

Lemma A.1. *The follow identity holds*

$$\sum_{n \geq 0} \frac{q^{-2bn+cn+n^2/2} (q^{\frac{1}{2}-c}; q)_n (q^{c+\frac{1}{2}}; q)_n}{(q^2; q^2)_n (q^{-2b+c+\frac{1}{2}}; q)_n} = \frac{(q+1)^{\frac{1}{2}-c} \Gamma_{q^2}(1-b) \Gamma_q(-2b+c+\frac{1}{2})}{\Gamma_q(1-2b) \Gamma_{q^2}(-b+c+\frac{1}{2})}$$

Proof. Replacing c by $c+1/2$, we need to prove

$$(A.1) \quad f(q^c) = \sum_{n \geq 0} \frac{q^{-2bn+cn+\frac{n^2}{2}+\frac{n}{2}} (q^{-c}; q)_n (q^{c+1}; q)_n}{(q^2; q^2)_n (q^{-2b+c+1}; q)_n} = \frac{(q+1)^{-c} \Gamma_{q^2}(1-b) \Gamma_q(-2b+c+1)}{\Gamma_q(1-2b) \Gamma_{q^2}(-b+c+1)}$$

creative telescoping in c gives

$$f(q^c) = \frac{1-q^{c-2b}}{1-q^{2c-2b}} f(q^{c-1})$$

the initial condition $f(q^0) = 1$ implies equation A.1 holds for $c \in \mathbb{N}$. This already says $f(q^c)$ holds for any c : $f(x)$ is an analytic function in a neighbourhood of $x = 0$, in which the sequence $\{q^1, q^2, q^3, \dots\}$ accumulates, so we have equality for all c , completing the proof. \square

We next prove the summations used in Examples XI and XII. We quickly recall some basic facts on Jacobi theta function. Let $\Im(\tau) > 0$ and $x \in \mathbb{C}$,

$$\theta(x|\tau) = 2 \sum_{n \geq 0} (-1)^n e^{\pi i \tau (n+1/2)^2} \sin(2n+1)\pi x = 2e^{2\pi i x/4} (\sin \pi x) (e^{2\pi i \tau}, e^{2\pi i(\tau+x)}, e^{2\pi i(\tau-x)}; e^{2\pi i \tau})_{\infty}$$

The following properties are well-known:

- $\theta(-x) = -\theta(x)$ $\theta(x+1|\tau) = -\theta(x|\tau)$ $\theta(x+\tau|\tau) = -e^{-\pi i(\tau+2x)} \theta(x|\tau)$
- $\theta(x|\tau)$ has simple zero at $\mathbb{Z} + \tau\mathbb{Z}$ and non-zero elsewhere.
- $(-i\tau)^{1/2} \theta(x|\tau) = ie^{-ix^2\pi/\tau} \theta(\frac{x}{\tau} | \frac{-1}{\tau})$

Lemma A.2. *For $b \in \mathbb{C}$, we have*

$$\sum_{n \geq 0} \frac{q^{n^2} (q^{1/2-3b}, q^{1/2+3b}; q)_n}{(q; q)_{2n}} = \frac{\Gamma_{q^3}(\frac{1}{3}) \Gamma_{q^3}(\frac{2}{3})}{\Gamma_{q^3}(\frac{1}{2}-b) \Gamma_{q^3}(b+\frac{1}{2})}$$

Remark A.3. The terse proof in style of Lemma A.1 no longer works, because if $f(q^b)$ denote the summation, $f(x)$ has an *essential singularity* at $x = 0$.

Proof of lemma. It suffices to find $S(b) = \sum_{n \geq 0} \frac{q^{n^2} (q^{1+3b}, q^{-3b}; q)_n}{(q; q)_{2n}}$. A straightforward creative telescoping on b says $S(b) = -q^{1-3b} S(b-1)$. Now $(q^{2b} \Gamma_{q^3}(b) \Gamma_{q^3}(1-b))^{-1}$ satisfies the same recurrence. So

$$f(b) = S(b) q^{2b} \Gamma_{q^3}(b) \Gamma_{q^3}(1-b)$$

satisfies $f(b+1) = f(b)$. For $\tau = \frac{-2\pi i}{3 \log q}$ in upper half-plane, because $q^{3\tau} = 1$, we also have $f(b+\tau) = q^{2\tau} f(b) = e^{2\pi i/3} f(b)$, so

$$g(b) = f(b) \frac{\theta(b|\tau)}{\theta(b-1/3|\tau)}$$

is an elliptic function with periods 1 and τ . Note that f has poles only at $\mathbb{Z} + \tau\mathbb{Z}$ coming from the q -gamma function, so g has at most a simple pole at point modulo $b = 1/3$, this already forces it to

be a constant. This constant is $g(2/3) = f(2/3) \frac{\theta(2/3)}{\theta(1/3)} = f(2/3)$ because $\theta(x) = \theta(1-x)$, note that $f(2/3)$ can be easily found using q -gamma function since $S(2/3)$ is a terminating sum. Now

$$f(b) = f(2/3) \frac{\theta(b-1/3|\tau)}{\theta(b|\tau)} = f\left(\frac{2}{3}\right) \exp\left[-\frac{i\pi}{\tau} \left(\left(b - \frac{1}{3}\right)^2 - b^2\right)\right] \frac{\theta\left(\frac{3b-1}{3\tau} \middle| \frac{-1}{\tau}\right)}{\theta\left(\frac{b}{\tau} \middle| \frac{-1}{\tau}\right)}$$

Using infinite product representation of $\theta(x|\tau)$, the last expression can be converted to q -series⁴, some simplification gives the result. \square

Lemma A.4. For $b \in \mathbb{C}$, we have

$$\sum_{n \geq 0} \frac{q^{4n^2} (-q^{3b+4n+\frac{5}{2}} - q^{6b+6n+2} + q^{3b+8n+\frac{5}{2}} + q^{3b+8n+\frac{7}{2}} + q^{3b+\frac{1}{2}} - q^{6n+2}) (q^{\frac{1}{2}-3b}, q^{3b+\frac{1}{2}}; q)_{2n}}{(q; q)_{4n+2}} = \frac{q^{3b+1/2} \Gamma_{q^3}\left(\frac{1}{3}\right) \Gamma_{q^3}\left(\frac{2}{3}\right)}{\Gamma_{q^3}\left(\frac{1}{2}-b\right) \Gamma_{q^3}\left(b+\frac{1}{2}\right)}$$

Proof. The proof proceeds almost exactly like above lemma. It suffices to find

$$S(b) = \sum_{n=0}^9 \frac{q^{4n^2+1} (-q^{3b+4n+2} - q^{6b+6n+2} + q^{3b+8n+2} + q^{3b+8n+3} + q^{3b} - q^{6n+1}) (q^{-3b}; q)_{2n} (q^{3b+1}; q)_{2n}}{(q; q)_{4n+2}}$$

creative telescoping says $S(b) = -q^{4-3b}S(b-1)$, so $f(b) = S(b)q^{-b}\Gamma_{q^3}(b)\Gamma_{q^3}(1-b)$ satisfies $f(b) = f(b+1) = e^{2\pi i/3}f(b+\tau)$ with τ as in previous lemma. Arguing like before, we again have

$$f(b) = f(2/3) \frac{\theta(b-1/3|\tau)}{\theta(b|\tau)}$$

\square

APPENDIX B. CLASSICAL COUNTERPART OF EXAMPLE XVII

To compensate our setback in Example XVII for not being able to calculate the limit $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$, we can at least do it in the classical case. We find $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$ using a technique not available for q -series, thereby obtaining a classical 4-variable extension of

$$\sum_{n \geq 0} \left(\frac{1}{2^6}\right)^n \frac{\left(\frac{1}{2}\right)_n^7}{(1)_n^7} (168n^3 + 76n^2 + 14n + 1) = \frac{32}{\pi^3}$$

This provides an alternative proof of this identity other than Example XI of author's previous article [2]. Let a, b, c, d be in a neighbourhood of 0,

$$F(n, k) = \frac{(-1)^n (2d+2k+1) 2^{-2(2a+b+c+2n)} \Gamma(-a-b-n+1) \Gamma(a+2b+c+n+\frac{1}{2}) \Gamma(a+b+2c+n+\frac{1}{2}) \Gamma(-a+d+k-n+1)}{\Gamma(-a+b-n+\frac{1}{2}) \Gamma(-a+c-n+\frac{1}{2}) \Gamma(a+c+n) \Gamma(2a+b+c+2n-1) \Gamma(a+b+2d+2k+n+\frac{1}{2}) \Gamma(a+c+2d+2k+n+\frac{1}{2}) \Gamma(-a-b-c+d+k-n+1)} \\ \Gamma(a+d+k+n+1) \Gamma(-a-b+2d+2k-n+\frac{5}{2}) \Gamma(-a-c+2d+2k-n+\frac{5}{2}) \Gamma(a+b+c+d+k+n+1)$$

which comes from Seed9($1-2(a+n)$, $-a+b-n+\frac{1}{2}$, $-a+c-n+\frac{1}{2}$, $a+d+k+n$). When $b=c=d=0$, this is the classical analogue of Example XVII. Apply Proposition 1.3,

$$\sum_{k \geq 0} F(0, k) + \sum_{n \geq 0} g(n) = \sum_{n \geq 0} G(n, 0) + \lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$$

becomes⁵

$$(B.1) \quad \sum_{k \geq 0} A(k) + B_1 \sum_{n \geq 0} B_2(n) = \sum_{n \geq 0} C(n) + \frac{1}{D} \lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$$

⁴without using modular transform at last step, we only recover q -pochhammer with $q = e^{2\pi i \tau} = e^{4\pi^2 / \log q}$, this is undesirable

⁵a Mathematica notebook containing this long formula can be found in <https://www.researchgate.net/publication/376812577>

with

$$A(k) = \frac{(2d+2k+1)(-a+d+1)_k (a+b+2d+\frac{1}{2})_{2k} (a+c+2d+\frac{1}{2})_{2k} (-a-b-c+d+1)_k}{(a+d+1)_k (a+b+c+d+1)_k (-a-b+2d+\frac{1}{2})_{2k+2} (-a-c+2d+\frac{1}{2})_{2k+2}}$$

$$B_1 = \frac{2^{4a+2b+2c-4} \Gamma(a+d+1) \Gamma(-a-b+2d+\frac{1}{2}) \Gamma(-a-c+2d+\frac{1}{2}) \Gamma(a+b+c+d+1)}{(a+b)(a+c)(2a+b+c-1)(2a+b+c) \Gamma(-a+d+1) \Gamma(a+b+2d+\frac{1}{2}) \Gamma(a+c+2d+\frac{1}{2}) \Gamma(-a-b-c+d+1)}$$

$$B_2(n) = \frac{(12a^2+12ab+12ac+24an+8a-4bc+12bn+4b+12cn+4c+12n^2+8n+1) \times (a-b+\frac{1}{2})_n (a-c+\frac{1}{2})_n (a+2b+c+\frac{1}{2})_n (a+b+2c+\frac{1}{2})_n}{(2a+b+c+2n+1)(a+b+1)_n (a+c+1)_n (2a+b+c+1)_{2n}}$$

$$D = \frac{2^{-2(2a+b+c)} (2a+b+c-1) \Gamma(-a-b+1) \Gamma(-a+d+1) \Gamma(a+2b+c+\frac{1}{2}) \Gamma(a+b+2c+\frac{1}{2}) \Gamma(a+b+2d+\frac{1}{2}) \Gamma(a+c+2d+\frac{1}{2}) \Gamma(-a-b-c+d+1)}{\Gamma(-a+b+\frac{1}{2}) \Gamma(-a+c+\frac{1}{2}) \Gamma(a+c) \Gamma(a+d+1) \Gamma(2a+b+c) \Gamma(-a-b+2d+\frac{1}{2}) \Gamma(-a-c+2d+\frac{1}{2}) \Gamma(a+b+c+d+1)}$$

$$C(n) = \frac{-2^{-4n-8} (a-b+\frac{1}{2})_n (a-c+\frac{1}{2})_n (a+2b+c+\frac{1}{2})_n (a+b+2c+\frac{1}{2})_n (a+b-2d+\frac{1}{2})_n (a+b+2d+\frac{1}{2})_n (a+c-2d+\frac{1}{2})_n (a+c+2d+\frac{1}{2})_n P(n)}{(a+b)(a+c)(a-d)(2a+b+c-1)(2a+b+c)(a+b+c-d)(2a+b+c+2n+1)(a+b+1)_n (a+c+1)_n (a-d+1)_n (a+d+1)_n (2a+b+c+1)_{2n} (a+b+c-d+1)_n (a+b+c+d+1)_n}$$

and $P(n) \in \mathbb{Z}[a, b, c, d, n, k]$ a long polynomial.

Evaluating the limit is non-trivial but possible:

Lemma B.1. For $F(n, k)$ above, $\lim_{n \rightarrow \infty} \sum_{k \geq 0} F(n, k)$ equals

$$\frac{1}{8} \cos(\pi(a-b)) \csc(\pi(a+b)) \cos(\pi(a-c)) \csc(\pi(a-d)) \csc(\pi(a+b+c-d))$$

Remark B.2. Note that the result is 1-periodic in each auxiliary variable a, b, c, d . This is actually a general phenomenon when $F(n, k)$ is a WZ-seed. However, knowing this periodicity alone does not fix the expression completely, since we still lack "initial condition".

Proof of lemma. We heavily use ideas in [2]. For $x \geq 0$, let $\mathcal{E}_F(x) \geq 0$ be characterized by

$$|F(n, xn)| = (\mathcal{E}_F(x))^n \times O(n^C) \quad n \rightarrow +\infty$$

then it can be seen

$$\mathcal{E}_F(x) = \frac{1}{64} |x-1|^{2x-2} |x+1|^{-2x-2} |2x-1|^{2-4x} |2x+1|^{4x+2}$$

this has a global maximum at $x = \sqrt{3}/2$ for $x \geq 0$, attaining value 1. Therefore only sums over

$$\sum_{k \in (\sqrt{3}/2 - \delta, \sqrt{3}/2 + \delta) \times n} F(n, k)$$

contributes to the limit. (Section 2.4 of [2]).

Now we proceed exactly as in Example XVI in [2]: for θ with $1/2 < \theta < 1$, we expand $F(n, \theta n)$ using Stirling's formula:

$$F(n, \theta n) = 2^{-6a-3b-3c+1} \underbrace{\cos(\pi(a-b)) \cos(\pi(a-c)) \csc(\pi(a+b)) \csc(\pi(a-d)) \csc(\pi(a+b+c-d))}_{:=T} \times \theta(1-\theta)^{-2a-b-c+2d+1} (\theta+1)^{-2a-b-c-2d-1} (2\theta-1)^{2a+b+c-4d-4} (2\theta+1)^{2a+b+c+4d} \times \frac{1}{\sqrt{n\pi}} \exp[nf(\theta)] (1+O(1/n))$$

where $f(\theta) = 2\theta \log(1 - \theta) - 2\log(1 - \theta) - 2\theta \log(\theta + 1) - 2\log(\theta + 1) - 4\theta \log(2\theta - 1) + 2\log(2\theta - 1) + 4\theta \log(2\theta + 1) + 2\log(2\theta + 1) - 6\log(2)$, it has a maximum at $\theta = \sqrt{3}/2$, as expected. Now let $r \in (-n^\gamma, n^\gamma) \cap \mathbb{Z}$ with $0 < \gamma < 1$ to be chosen later, writing $\theta = \sqrt{3}/2 + \frac{r}{n}$, we have

$$f(\theta) = -\frac{12r^2}{n^2} + O\left(\frac{r^3}{n^3}\right) = -\frac{12r^2}{n^2} + O(n^{3\gamma-3})$$

and

$$2^{-6a-3b-3c+1}\theta(1-\theta)^{-2a-b-c+2d+1}(\theta+1)^{-2a-b-c-2d-1}(2\theta-1)^{2a+b+c-4d-4}(2\theta+1)^{2a+b+c+4d} = \frac{\sqrt{3}}{4} + O(n^{\gamma-1})$$

therefore we have

$$F(n, r + \frac{\sqrt{3}}{2}n) = \frac{\sqrt{3}}{4} T \frac{1}{\sqrt{n\pi}} e^{-12r^2/n} [1 + O(n^{3\gamma-2})]$$

If we require $2\gamma - 1 > 0$, then

$$\begin{aligned} \sum_{r \in (-n^\gamma, n^\gamma)} e^{-12r^2/n} &= \sum_{r \in \mathbb{Z}} e^{-12r^2/n} + O(n^{-1}) && \text{(Poisson summation)} \\ &= \frac{\sqrt{n\pi}}{2\sqrt{3}} \sum_{m \in \mathbb{Z}} e^{-n\pi^2 m^2/12} + O(n^{-1}) = \frac{\sqrt{n\pi}}{2\sqrt{3}} + O(n^{-1}) \end{aligned}$$

therefore we see the sub-sum

$$\sum_{k \in (\sqrt{3}n/2 - n^\gamma, \sqrt{3}n/2 + n^\gamma)} F(n, k)$$

already contributes to all the limit provided that $3\gamma - 2 < 0$ and $2\gamma - 1 > 0$, for example, we can choose $\gamma = 7/12$. The limit is then $T/8$, as claimed. \square

Setting $a = b = c = d = 0$ in B.1, we re-established the $32/\pi^3$ -identity. We also want to know the nature of coefficient in a, b, c, d when both sides are expanded as Laurent series at $(a, b, c, d) = (0, 0, 0, 0)$. For $\sum_{k \geq 0} A(k), B_1, D$, their natures are evident: level 2 colored multiple zeta values (CMZVs). For $\sum_{n \geq 0} B_2(n)$, using Gauss2F1 $(-a+b-n+\frac{1}{2}, -a-b-2c-n+\frac{1}{2}, -a-c-n+1, 2a+b+c+k+2n)$ on Proposition 1.3, one sees easily

$$4(a+c) \sum_{k \geq 0} \frac{(a-c+\frac{1}{2})_k (a+2b+c+\frac{1}{2})_k}{(a+b+1)_k (2a+b+c+1)_k} = \sum_{n \geq 0} B_2(n)$$

whose coefficients are level 4 CMZVs by some hypergeometric transformation formulas.

By comparing coefficient, we can also prove the following extensions⁶: ($a_n = \frac{1}{2^{6n}} \frac{(1/2)_n^7}{(1)_n^7}$ and $p_n = 168n^3 + 76n^2 + 14n + 1$),

$$\begin{aligned} \sum_{n \geq 0} a_n p_n &= \frac{32}{\pi^3} \\ \sum_{n \geq 0} a_n [7p_n (H_{2n} - H_n) + 252n^2 + 76n + 7] &= \frac{320 \log 2}{\pi^3} \\ \sum_{n \geq 0} a_n [p_n (16H_{2n}^{(2)} - 5H_n^{(2)}) + 48n + 8] &= \frac{80}{3\pi} \\ \sum_{n \geq 0} a_n [p_n (128H_{2n}^{(4)} - 7H_n^{(4)}) + \frac{64}{2n+1}] &= \frac{976\pi}{45} \end{aligned}$$

they were conjectured by Z.W. Sun ([19]).

⁶a Mathematica notebook containing computational details can be found in <https://www.researchgate.net/publication/376812577>

REFERENCES

- [1] Tewodros Amdeberhan and Doron Zeilberger. Hypergeometric series derivation via the WZ method. *arXiv preprint math/9804121*, 1998.
- [2] Kam Cheong Au. Wilf-zeilberger seeds and non-trivial hypergeometric identities. *arXiv preprint arXiv:2312.14051*, 2023.
- [3] Henri Cohen and Jesús Guillera. Rational hypergeometric Ramanujan identities for $1/\pi^c$: Survey and generalizations. *arXiv preprint arXiv:2101.12592*, 2021.
- [4] Akihito Ebisu. *Special values of the hypergeometric series*, volume 248. American Mathematical Society, 2017.
- [5] George Gasper and Mizan Rahman. *Basic hypergeometric series*, volume 96. Cambridge university press, 2011.
- [6] Édouard Goursat. Sur l'équation différentielle linéaire, qui admet pour intégrale la série hypergéométrique. In *Annales scientifiques de l'École Normale Supérieure*, volume 10, pages 3–142, 1881.
- [7] Jesús Guillera. A new Ramanujan-like series for $1/\pi^2$. *arXiv preprint arXiv:1003.1915*, 2010.
- [8] Jesús Guillera. Bilateral sums related to Ramanujan-like series. *arXiv preprint arXiv:1610.04839*, 2016.
- [9] Jesús Guillera. Wz pairs and q -analogues of ramanujan series for $1/\pi$. *Journal of Difference Equations and Applications*, 24(12):1871–1879, 2018.
- [10] Victor JW Guo. Proof of a supercongruence conjectured by sun through a q -microscope. *Proceedings-Mathematical Sciences*, 130:1–7, 2020.
- [11] Victor JW Guo and Ji-Cai Liu. q -analogues of two ramanujan-type formulas for $1/\pi$. *Journal of Difference Equations and Applications*, 24(8):1368–1373, 2018.
- [12] Victor JW Guo and Wadim Zudilin. Ramanujan-type formulae for $1/\pi$: q -analogues. *Integral Transforms and Special Functions*, 29(7):505–513, 2018.
- [13] Victor JW Guo and Wadim Zudilin. A q -microscope for supercongruences. *Advances in Mathematics*, 346:329–358, 2019.
- [14] Victor JW Guo and Wadim Zudilin. Dwork-type supercongruences through a creative q -microscope. *Journal of Combinatorial Theory, Series A*, 178:105362, 2021.
- [15] Christoph Koutschan. *Advanced Applications of the Holonomic Systems Approach*. PhD thesis, RISC, Johannes Kepler University, Linz, Austria, 2009.
- [16] Daniel S Moak. The q -analogue of Stirling's formula. *The Rocky Mountain Journal of Mathematics*, pages 403–413, 1984.
- [17] Peter Paule and Markus Schorn. A mathematica version of zeilberger's algorithm for proving binomial coefficient identities. *Journal of symbolic computation*, 20(5-6):673–698, 1995.
- [18] Mizan Rahman. Some cubic summation and transformation formulas. *The Ramanujan Journal*, 1:299–308, 1997.
- [19] Zhi-Wei Sun. New series involving binomial coefficients. *arXiv preprint arXiv:2307.03086*, 2023.
- [20] Wadim Zudilin. Ramanujan-type supercongruences. *Journal of Number Theory*, 129(8):1848–1857, 2009.

RHEINISCHE FRIEDRICH-WILHELMS-UNIVERSITÄT BONN, MATHEMATICAL INSTITUTE, 53115 BONN, GERMANY
 Email address: s6kmauu@uni-bonn.de