

On $[1, 2]$ -Domination in Interval and Circle Graphs

Mohsen Alambardar Meybodi^{1*} and Abolfazl Poureidi²

^{1*}Department of Applied Mathematics and Computer Science, Faculty of Mathematics and Statistics, University of Isfahan, Isfahan, 81746-73441, Iran.

²Faculty of Mathematical Sciences, Shahrood University of Technology, Shahrood, Iran.

*Corresponding author(s). E-mail(s): m.alambardar@sci.ui.ac.ir;
Contributing authors: a.poureidi@shahroodut.ac.ir;

Abstract

A subset S of vertices in a graph $G = (V, E)$ is Dominating Set if each vertex in $V(G) \setminus S$ is adjacent to at least one vertex in S . Chellali et al. in 2013, by restricting the number of neighbors in S of a vertex outside S , introduced the concept of $[1, j]$ -dominating set. A set $D \subseteq V$ of a graph $G = (V, E)$ is called $[1, j]$ -Dominating Set of G if every vertex not in D has at least one neighbor and at most j neighbors in D . The Minimum $[1, j]$ -Domination problem is the problem of finding the minimum set D . Given a positive integer k and a graph $G = (V, E)$, the $[1, j]$ -Domination Decision problem is to decide whether G has $[1, j]$ -dominating set of cardinality at most k . A polynomial-time algorithm was obtained in split graphs for a constant j in contrast to the classic Dominating Set problem which is NP-hard in split graphs. This result motivates us to investigate the effect of restriction j on the complexity of $[1, j]$ -domination problem on various classes of graphs. Although for $j \geq 3$, it has been proved that the minimum of classical domination is equal to minimum $[1, j]$ -domination in interval graphs, the complexity of finding the minimum $[1, 2]$ -domination in interval graphs is still outstanding. In this paper, we propose a polynomial-time algorithm for computing a minimum $[1, 2]$ on interval graphs by a dynamic programming technique. Next, on the negative side, we show that the minimum $[1, 2]$ -dominating set problem on circle graphs is *NP*-complete.

Keywords: Dominating set, $[1, j]$ -dominating set, Interval graph, Circle graph.

MSC Classification: 05C69 , 68Q25

1 Introduction

One of the active research areas in graph theory is the study of the domination problem and its variant. Applying some restrictions on the domination problem leads to various extensions of the domination problem such as independent domination (Extra restriction is: each vertex in dominating set, like D , is not adjacent to a vertex in D), total domination (Extra restriction is: each vertex in D is adjacent to at least one vertex in D) and, $[1, j]$ -domination (Extra restriction is: each vertex not in D is adjacent to at least one and at most j vertices in D). Combinatorial and algorithmic results have been widely obtained on the various extensions of domination problems. A discussion of these results can be found in [9].

1.1 Definition and notation

We begin with some terminology and notation.

Graph

Let $G = (V, E)$ is a simple graph with vertex set V and edge set E . The order of graph G is defined as $n = |V(G)|$. The open neighborhood of a vertex $v \in V$, denoted by $N(v)$, is defined as $\{u : \{u, v\} \in E\}$. Also the closed neighborhood of v is defined as $N[v] = N(v) \cup \{v\}$. We use $N(S)$ and $N[S]$ to denote the open and closed neighborhood S , for a set $S \subseteq V$, respectively. That is, $N[S] = \bigcup_{v \in S} N[v]$ and $N(S) = N[S] \setminus S$. The complete bipartite graph $K_{1;3}$ is called claw and, a graph without a claw as an induced subgraph is called claw-free.

Domination and $[1, j]$ -Domination

A set $S \subseteq V$ is a dominating set if every vertex not in S is adjacent to at least one vertex in S . The cardinality of the smallest dominating set, denoted by $\gamma(G)$, is called domination number. Finding a minimum dominating set was one of the first problems shown NP-hard in [6]. A set $D \subseteq V$ is called a $[1, j]$ -dominating set of G if for each $v \in V \setminus D$ we have $1 \leq |N(v) \cap D| \leq j$, i.e. v is adjacent to at least one but not more than j vertices in D . Every graph has at least one $[1, j]$ -dominating set since the set of vertices V is itself a $[1, j]$ -dominating set. The size of the smallest $[1, j]$ -dominating sets of G is denoted by $\gamma[1, j](G)$. For special case, where $j = 2$, this domination is already known as Quasiperfect Domination. In the decision version of the $[1, j]$ -dominating set problem, the input is a graph G and a positive integer k and, the objective is to test whether there is a $[1, j]$ -dominating set of size at most k .

Interval Graph

Let $I = \{I_1, I_2, \dots, I_n\}$ be a set of intervals on real line. A graph $G = (V, E)$ is called an interval graph if each of its vertices can be associated with an interval in I and two vertices are adjacent if and only if the corresponding intervals have intersection. The set I is called the interval representation of an interval graph G . The intervals in I are indexed by increasing the right endpoints. Figure 1 shows an interval graph and its internal representation.




Fig. 1 Illustrating (a) an interval graph G of order 10, (b) a set of intervals corresponding to the vertex set of G

A linear time algorithm proposed in [2] to recognize an interval graph and the corresponding intervals of the vertices of an interval graph. The induced subgraphs of interval graphs are also interval [8] which is called hereditary property of the interval graphs. A proper interval graph is an interval graph such that no interval is completely contained within another interval. If all the intervals have the same length, its corresponding interval graph called a unit interval graph.

Circle Graph

A circle graph is a graph $G = (V, E)$ such that there is a one-to-one mapping between vertices in V and a set C of chords in a circle. Two vertices in V are adjacent if and only if the corresponding chords in C intersect. C is called the chord intersection model for G . Equivalently, the vertices of a circle graph can be placed in the one-to-one correspondence the intervals such that two vertices are adjacent if and only if the corresponding intervals overlap, but neither contains the other. I is called the interval representation of the circle graph. There exists a polynomial transformation between a representation of a circle graph, the set of its chords or its intervals representation. Moreover those are equivalent. (See Figure 2)




Fig. 2 (a) A circle graph G on 8 vertices (b) The circle representation of G (c) The interval representation of G

1.2 Short review of $[1, j]$ -Domination

Chellali et al. in [3] introduced the concept of $[1, j]$ -dominating set with name $(1, j)$ -set, Although this concept for the constant $j = 2$ already known as quasiperfect domination and defined by Dejter [4]. Also, $[1, j]$ -dominating set fits into the general framework of (ρ, σ) -sets of graphs, which was first introduced by Telle in [17]. A (ρ, σ) -set is a set $D_{(\rho, \sigma)} \subseteq V$ such that for every vertex $v \in V$, we have $|N(v) \cap D_{(\rho, \sigma)}| \in \rho$ if $v \in D_{(\rho, \sigma)}$, and $|N(v) \cap D_{(\rho, \sigma)}| \in \sigma$ if $v \notin D_{(\rho, \sigma)}$. Discussion about the complexity of

$[1, j]$ -dominating set in various graph have been done extensively. The authors in [3] raised several open problems. One major open question is in what classes of graphs the $\gamma[1, j](G)$ is equal to $\gamma(G)$. In [3], it was shown that in P_4 -free graphs, claw-free graphs, and caterpillars $\gamma[1, j](G)$ is equal to $\gamma(G)$. In [16] it was shown that $\gamma[1, j](G) = \gamma(G)$ when G is a grid graph. It was proved in [19], that the value of $\gamma_{[1,2]}(G)$ in the classes of planar, bipartite, and triangle-free graphs can be equal to the entire set of vertices. So, $\gamma_{[1,2]}$ necessarily is not equal to γ in these classes. In [5], the authors showed that in the interval graph and permutation graph $\gamma_{[1,3]}(G) = \gamma(G)$. Moreover, it was shown that there exists an interval graph H for which $\gamma(H) < \gamma_{[1,2]}(H)$. Although they prove that for every unit interval graph G we have $\gamma_{[1,2]}(G) = \gamma(G)$, the complexity of finding $\gamma_{[1,2]}$ in non-proper interval graphs remains unsolved.

Another open question in [3] was in which graph classes $[1, j]$ -dominating set is efficiently solvable. On the negative side, it was also proven that the problem is NP -complete even for bipartite graphs[3]. In [1], the $[1, j]$ -dominating sets problem was shown to be NP -hard even for chordal and planar graphs. In [5], the authors showed the complexity of the decision problem of whether the $\gamma_{[1,j]}(G) = \gamma(G)$ is an NP problem. On the positive side, The linear-time algorithms proposed for computing $\gamma_{[1,2]}(G)$ when G is a tree [7], G is a block graph [14] and G is a serial-parallel graph [15]. For a constant j , a polynomial-time algorithm running in roughly $O(n^j p(\log n))$ where p is a polynomial function, was obtained for n -vertex split graphs [1]. This is in contrast to the classic dominating set problem which is NP -hard for this class of graphs. Moreover in [10], a lower bound for computing $[1, j]$ -dominating set in the split graph shown, i.e. there is no algorithm with running time $O(n^{j-\epsilon})$ where $\epsilon > 0$ unless $P = NP$. The complexity of $[1, j]$ -dominating set problem in various graph classes is summarized in Figure 3. It should be noted that this summary is far from being comprehensive and several other classes of graphs exist. In this paper, we focus only on the circle and interval graphs.

2 Algorithm for interval graphs

Roberts [13] has shown that an interval graph is proper if and only if it contains no induced copy of $K_{1,3}$. In [4], it was shown that for any $K_{1,3}$ -free graph $\gamma(G) = \gamma_{[1,2]}(G)$. So, easily the following lemma is concluded.

Lemma 1. *For every proper interval graph G , $\gamma(G) = \gamma_{[1,2]}(G)$.*

Another proof for the above lemma was shown in [5]. The authors also show that for every interval graph G , $\gamma(G) = \gamma_{[1,3]}(G)$ and, there exists graph G such that $\gamma(G) \leq \gamma_{[1,2]}(G)$, but the complexity of finding $[1, 2]$ -dominating set in interval graphs remained open.

In this section, we propose an $O(n^4)$ -time algorithm for computing $\gamma_{[1,2]}(G)$, where G is an interval graph of order n . To present and prove the correctness of our algorithm, we need the following lemmas and notation. Let $G = (V, E)$ be an interval graph. Ramalingam and Rangan [12] proposed a numbering of vertices of G ; they stated the following result.




Fig. 3 Inclusion relations among well-studied classes of graphs [18] -Complexity of $[1, j]$ -dominating set restricted to various classes of graphs in a hierarchy of graphs - *NPc* indicate that the problem is *NP*-complete, *P* indicates the problem has polynomial algorithm, and question mark indicate open problem.

Corollary 1 (Ramalingam and Rangan [12]). *For each interval graph $G = (V, E)$ of order n , there is a numbering $(1, 2, \dots, n)$ of vertices in V such that if $ik \in E$ and $i < j < k$, then $jk \in E$.*

This numbering of vertices in a given interval graph $G = (V, E)$ can be computed in $O(|V| + |E|)$ time [12]. Let a and b be integers such that $1 \leq a \leq b \leq n$. We have to introduce the following notation.

- $[a, b] = \{a, a + 1, \dots, b\}$,
- $(a, b) = [a, b] \setminus \{a\}$,
- $[a, b) = [a, b] \setminus \{b\}$,
- $(a, b) = [a, b] \setminus \{a, b\}$,
- $G[a, b] = G[S]$, where $S = [a, b]$,
- $\text{low}(a) = \min N_G[a]$,
- $\text{low}(a, b) = \min\{\text{low}(k) : k \in [a, b]\}$,
- $\text{maxlow}(a) = \max\{\text{low}(k) : \text{low}(a) \leq k \leq a\}$.

Let $i \in [1, n]$, $j \in [1, i]$, $k \in [1, j]$ and $l \in [1, k]$.

- $\gamma_{[1,2]}(i) = \min\{|D| : D \text{ is an } [1, 2]\text{-dominating set of } G[1, i]\}$,
- $\gamma_{[1,2]}^0(i) = \min\{|D| : D \text{ is an } [1, 2]\text{-dominating set of } G[1, i] \text{ (if exists) such that } i \notin D\}$,
- $\gamma_{[1,2]}^0(j, i) = \min\{|D| : D \text{ is an } [1, 2]\text{-dominating set of } G[1, i] \text{ (if exists) such that } x \notin D \text{ for all } x \in [j, i]\}$,
- $\gamma_{[1,2]}^0(j, i : k') = \min\{|D| : D \text{ is an } [1, 2]\text{-dominating set of } G[1, i] \text{ (if exists) such that } k' \in D \text{ and } x \notin D \text{ for all } x \in [j, i] \setminus \{k'\}\}$, where $k' \in [j, i]$,

- $\gamma_{[1,2]}^1(i) = \min\{|D| : D \text{ is an } [1, 2]\text{-dominating set of } G[1, i] \text{ (if exists) such that } i \in D\}$,
- $\gamma_{[1,2]}^1(j, i : i) = \min\{|D| : D \text{ is an } [1, 2]\text{-dominating set of } G[1, i] \text{ (if exists) such that } i \in D \text{ and } x \notin D \text{ for all } x \in [j, i)\}$,
- $\gamma_{[1,2]}^1(k, i : i, j) = \min\{|D| : D \text{ is an } [1, 2]\text{-dominating set of } G[1, i] \text{ (if exists) such that } i, j \in D \text{ and } x \notin D \text{ for all } x \in [k, i) \setminus \{j\}\}$,
- $\gamma_{[1,2]}^1(l, i : i, j, k) = \min\{|D| : D \text{ is an } [1, 2]\text{-dominating set of } G[1, i] \text{ (if exists) such that } i, j, k \in D \text{ and } x \notin D \text{ for all } x \in [l, i) \setminus \{j, k\}\}$,
- $\gamma_{[1,2]}^{11}(l, i : i, j, k) = \min\{|D| : D \text{ is an } [1, 2]\text{-dominating set of } G[1, i] \text{ (if exists) such that } i, j, x \in D \text{ for all } x \in [l, k] \text{ and } y \notin D \text{ for all } y \in (k, i) \setminus \{j\}\}$.

We get that $\text{low}(i) \leq \text{maxlow}(i) \leq i$. An example of an interval graph is shown in Figure 4. Also, the values $\text{low}(i)$ and $\text{maxlow}(i)$ are computed for each vertex i of the interval graph illustrated in Figure 4.




Fig. 4 (a) a set of intervals corresponding to the vertex set of the interval graph G (b) a numbering $(1, 2, \dots, 10)$ of vertices of G satisfying the condition of Corollary 1, and (c) values $\text{low}(i)$ and $\text{maxlow}(i)$ for all $i \in [1, 10]$

In the rest of this section, we assume that $G = (V, E)$ is an interval graph of order n , and without loss of generality with a numbering $(1, 2, \dots, n)$ of vertices of G satisfying the condition of Corollary 1.

Corollary 2 (Poureidi [11]). *Let $i \in [1, n]$.*

- (i) *The set $[\text{maxlow}(i), i]$ is a clique of G .*
- (ii) *There is a vertex $k \in [\text{maxlow}(i), i]$ such that $kl \notin E$ for each vertex $l \in [1, \text{maxlow}(i)]$.*

Lemma 2. *Let $1 \leq i' \leq i \leq n$.*

- (i) $\gamma_{[1,2]}^0(i) = \gamma_{[1,2]}^0(i, i)$.

(ii) If $\text{maxlow}(i) < i'$, then $\gamma_{[1,2]}^0(i', i) = \min\{\gamma_{[1,2]}^1(\text{low}(j+1, i), j : j), \gamma_{[1,2]}^1(\text{low}(j+1, i), j : j, k) : j \in [\text{maxlow}(i), i'], k \in [\text{low}(j+1, i), j]\}$, otherwise, $\gamma_{[1,2]}^0(i', i)$ is not defined.

Proof. It is clear that the case (i) holds. Now, we prove the case (ii). Let D be an $[1, 2]$ -dominating set of $G[1, i]$ with minimum cardinality such that $a \notin D$ for all $a \in [i', i]$. So, $|D| = \gamma_{[1,2]}^0(i', i)$. Since $i \notin D$, at least one and at most two of vertices adjacent to i are in D . If $i' \leq \text{low}(i)$, then there is no vertex in D to dominate i , that is, D is not an $[1, 2]$ -dominating set of $G[1, i]$. Hence, $\text{low}(i) < i'$. By the case (ii) of Corollary 2, there is a vertex $x \in [\text{maxlow}(i), i]$ such that $xy \notin E$ for each vertex $y \in [1, \text{maxlow}(i)]$. Thus, if $a \notin D$ for all $a \in [\text{maxlow}(i), i]$, then D is not an $[1, 2]$ -dominating set of $G[1, i]$ and so at least one of vertices of $[\text{maxlow}(i), i]$ is in D . Since $a \notin D$ for all $a \in [i', i]$, we obtain that $\text{maxlow}(i) < i'$ and at least one of vertices of $[\text{maxlow}(i), i']$ is in D . Assume that $j \in [\text{maxlow}(i), i']$ is a vertex in D such that $a \notin D$ for all $a \in (j, i]$. We obtain that D is an $[1, 2]$ -dominating set of $G[1, j]$. Let $b \in (j, i]$ be a vertex such that $\text{low}(b) \leq \text{low}(a)$ for all $a \in (j, i]$, that is, $\text{low}(b) = \text{low}(j+1, i)$. See Figure 5. Clearly, $\text{low}(j+1, i) \leq \text{low}(i)$. Because $b \notin D$ and $j \in D$, at most one of vertices of $[\text{low}(b), j]$ is in D . Hence, either no vertex of $[\text{low}(b), j]$ is in D , see Figure 5(a), or exactly one vertex of $[\text{low}(b), j]$ is in D , see Figure 5(b). In the following we consider these two cases.




Fig. 5 Illustrating an $[1, 2]$ -dominating set D of $G[1, i]$ such that $a \notin D$ for all $a \in [i', i]$; (a) no vertex of $[\text{low}(j+1, i), j]$ is in D and (b) exactly one vertex $k \in [\text{low}(j+1, i), j]$ is in D . Note that a vertex with label 1 is in D and a vertex with label 0 is not in D .

(a) Assume that no vertex of $[\text{low}(b), j]$ is in D . Hence, D is an $[1, 2]$ -dominating set of $G[1, j]$ such that $j \in D$ and $a \notin D$ for all $a \in [\text{low}(b), j)$ and so $|D| \leq \gamma_{[1,2]}^1(\text{low}(j+1, i), j : j)$.

- (b) Assume that exactly one vertex $k \in [\text{low}(b), j]$ is in D . Hence, D is an $[1, 2]$ -dominating set of $G[1, j]$ such that $j, k \in D$ and $a \notin D$ for all $a \in [\text{low}(b), j] \setminus \{k\}$ and so $|D| \leq \gamma_{[1,2]}^1(\text{low}(j+1, i), j : j, k)$.

It follows from (a)–(b) that $\gamma_{[1,2]}^0(i', i) = |D| \leq \min\{\gamma_{[1,2]}^1(\text{low}(j+1, i), j : j), \gamma_{[1,2]}^1(\text{low}(j+1, i), j : j, k) : j \in [\text{maxlow}(i), i'], k \in [\text{low}(j+1, i), j]\}$.

Conversely, let $j \in [\text{maxlow}(i), i']$. Assume that S is an $[1, 2]$ -dominating set of $G[1, j]$ with minimum cardinality such that $j \in S$ and $a \notin S$ for all $a \in [\text{low}(j+1, i), j]$. So, $|S| = \gamma_{[1,2]}^1(\text{low}(j+1, i), j : j)$. Since $j \in [\text{maxlow}(i), i']$ and $a \notin S$ for all $a \in [\text{low}(j+1, i), j]$, each vertex of $[i', i]$ is exactly dominated by j . We obtain that S is an $[1, 2]$ -dominating set of $G[1, i]$ such that $a \notin S$ for all $a \in [i', i]$, that is, $|S| \leq \gamma_{[1,2]}^0(i', i)$. Let $k \in [\text{low}(j+1, i), j]$. Assume that S' is an $[1, 2]$ -dominating set of $G[1, j]$ with minimum cardinality such that $j, k \in S'$ and $a \notin S'$ for all $a \in [\text{low}(j+1, i), j] \setminus \{k\}$. So, $|S'| = \gamma_{[1,2]}^1(\text{low}(j+1, i), j : j, k)$. Since $j \in [\text{maxlow}(i), i']$, $k \in [\text{low}(j+1, i), j] \cap S'$ and $a \notin S'$ for all $a \in [\text{low}(j+1, i), j] \setminus \{k\}$, each vertex of $[i', i]$ is dominated by at least one and at most two vertices of S' . We obtain that S' is an $[1, 2]$ -dominating set of $G[1, i]$ such that $a \notin S'$ for all $a \in [i', i]$, that is, $|S'| \leq \gamma_{[1,2]}^0(i', i)$. Hence, $\min\{\gamma_{[1,2]}^1(\text{low}(j+1, i), j : j) = |S|, \gamma_{[1,2]}^1(\text{low}(j+1, i), j : j, k) = |S'| : j \in [\text{maxlow}(i), i'], k \in [\text{low}(j+1, i), j]\} \leq \gamma_{[1,2]}^0(i', i)$. This completes the proofs of the case (ii) and the lemma. \square

Lemma 3. *Let $i \in [1, n]$, $j \in [1, i]$, $k \in [j, i]$ and $b = \min\{\text{low}(k+1, i), j\}$. If $\text{maxlow}(i) \leq k$, then $\gamma_{[1,2]}^0(j, i : k) = \min\{\gamma_{[1,2]}^1(b, k : k), \gamma_{[1,2]}^1(b, k : k, l) : l \in [b, j]\}$, otherwise, $\gamma_{[1,2]}^0(j, i : k)$ is not defined.*

Proof. Let D be an $[1, 2]$ -dominating set of $G[1, i]$ with minimum cardinality such that $k \in D$ and $a \notin D$ for all $a \in [j, i] \setminus \{k\}$. So, $|D| = \gamma_{[1,2]}^0(j, i : k)$. Since $a \notin D$ for all $a \in (k, i]$, it obtains that D is an $[1, 2]$ -dominating set of $G[1, k]$. By the case (ii) of Corollary 2, there is a vertex $x \in [\text{maxlow}(i), i]$ such that $xy \notin E$ for each vertex $y \in [1, \text{maxlow}(i)]$. Thus, if $k < \text{maxlow}(i)$, then D is not an $[1, 2]$ -dominating set of $G[1, i]$, that is, $\gamma_{[1,2]}^0(j, i : k)$ is not defined. Assume that $\text{maxlow}(i) \leq k$. Let $c \in (k, i]$ be a vertex such that $\text{low}(c) \leq \text{low}(a)$ for all $a \in (k, i]$, that is, $\text{low}(c) = \text{low}(k+1, i)$. See Figure 6. Recall that $b = \min\{\text{low}(k+1, i), j\}$ and $a \notin D$ for all $a \in [j, i] \setminus \{k\}$. If $\text{low}(k+1, i) < j$, that is $b = \text{low}(k+1, i)$, then because $c \notin D$ and $k \in D$, at most one of vertices of $[b, j]$ is in D , otherwise, $b = j$ and no vertex of $[b, k]$ is in D . Hence, either no vertex of $[b, k]$ is in D , see Figure 6(a), or exactly one vertex of $[b, j]$ is in D (if $b < j$), see Figure 6(b). In the following we consider these two cases.

- (a) Assume that no vertex of $[b, k]$ is in D . Hence, D is an $[1, 2]$ -dominating set of $G[1, k]$ such that $k \in D$ and $a \notin D$ for all $a \in [b, k]$ and so $|D| \leq \gamma_{[1,2]}^1(b, k : k)$.
- (b) Assume that exactly one vertex of $[b, j]$ is in D (if $b < j$). Hence, D is an $[1, 2]$ -dominating set of $G[1, k]$ such that $l, k \in D$, $l \in [b, j]$ and $a \notin D$ for all $a \in [b, k] \setminus \{l\}$ and so $|D| \leq \gamma_{[1,2]}^1(b, k : k, l)$.

It follows from (a)–(b) that $\gamma_{[1,2]}^0(j, i : k) = |D| \leq \min\{\gamma_{[1,2]}^1(b, k : k), \gamma_{[1,2]}^1(b, k : k, l) : l \in [b, j]\}$.




Fig. 6 Illustrating an $[1,2]$ -dominating set D of $G[1, i]$ such that $k \in D$ and $a \notin D$ for all $a \in [j, i] \setminus \{k\}$; (a) no vertex of $[b, k]$ is in D and (b) exactly one vertex $l \in [b, j]$ is in D . Note that a vertex with label 1 is in D and a vertex with label 0 is not in D .

Conversely, assume that S is an $[1,2]$ -dominating set of $G[1, k]$ with minimum cardinality such that $k \in S$ and $a \notin S$ for all $a \in [b, k]$. So, $|S| = \gamma_{[1,2]}^1(b, k : k)$. Recall that $b = \min\{\text{low}(k+1, i), j\}$. Since $k \in [\text{maxlow}(i), i]$ and $a \notin S$ for all $a \in [b, k]$, each vertex of $(k, i]$ is exactly dominated by k . We obtain that S is an $[1,2]$ -dominating set of $G[1, i]$ such that $k \in S$ and $a \notin S$ for all $a \in [j, i] \setminus \{k\}$, that is, $|S| \leq \gamma_{[1,2]}^0(j, i : k)$.

Let $l \in [b, j]$. Assume that S' is an $[1,2]$ -dominating set of $G[1, k]$ with minimum cardinality such that $l, k \in S'$ and $a \notin S'$ for all $a \in [b, k] \setminus \{l\}$. So, $|S'| = \gamma_{[1,2]}^1(b, k : k, l)$. Since $l \in [b, j]$, $k \in [\text{maxlow}(i), i]$ and $a \notin S'$ for all $a \in [b, i] \setminus \{k, l\}$, each vertex of $(k, i]$ is dominated by at least one and at most two vertices of S' . We obtain that S' is an $[1,2]$ -dominating set of $G[1, i]$ such that $k \in S'$ and $a \notin S'$ for all $a \in [j, i] \setminus \{k\}$, that is, $|S'| \leq \gamma_{[1,2]}^0(j, i : k)$. Hence, $\gamma_{[1,2]}^0(j, i : k) = \min\{\gamma_{[1,2]}^1(b, k : k) = |S|, \gamma_{[1,2]}^1(b, k : k, l) = |S'| : l \in [b, j]\}$. This completes the proof of the lemma. \square

Lemma 4. *Let D be an $[1,2]$ -dominating set of $G[1, i]$ with minimum cardinality such that $i \in [1, n]$, $i \in D$, $i' = \max D \cap [1, i]$, $i'' = \max D \cap [1, i']$ and $\text{low}(i) < i''$. If both i' and i'' exist, then $x \in D$ for all $x \in [\text{low}(i), i'']$.*

Proof. Suppose that $j = \max[\text{low}(i), i''] \setminus D$. See Figure 7. Since $j \notin D$ and j is adjacent to i , the vertex j is adjacent to at most one vertex of $S = D \cap (j, i)$. If j is adjacent to no vertex of S , then $D' = D \setminus S$ is an $[1,2]$ -dominating set of $G[1, i]$ such that $i \in D'$, a contradiction. Let j be adjacent to exactly one vertex $k \in S$. We obtain that $D'' = D \setminus S \cup \{k\}$ is an $[1,2]$ -dominating set of $G[1, i]$ such that $i \in D''$, a contradiction. This completes the proof of the lemma. \square

Lemma 5. *Let $b = \text{low}(i)$.*

- (i). *If $b = i$, then $\gamma_{[1,2]}^1(i) = \gamma_{[1,2]}(i-1) + 1$.*
- (ii). *If $b < i$, then $\gamma_{[1,2]}^1(i) = \min\{\gamma_{[1,2]}^1(b, i : i), \gamma_{[1,2]}^1(b, i : i, j), \gamma_{[1,2]}^1(b, i : i, j', k'), \gamma_{[1,2]}^{11}(b, i : i, j', k') : b \leq j < i, b \leq k' < j' < i\}$.*

Proof. The proof of the case (i) is clear. Assume that $\text{low}(i) < i$. Let D be an $[1,2]$ -dominating set of $G[1, i]$ with minimum cardinality such that $i \in D$. So, $|D| = \gamma_{[1,2]}^1(i)$.




Fig. 7 Illustrating an $[1,2]$ -dominating set D of $G[1, i]$ such that $i \in D$. Note that a vertex with label 1 is in D and a vertex with label 0 is not in D .

We obtain that all vertices of $[b, i)$ are not in D , exactly one vertex of $[b, i)$ is in D , exactly two vertices of $[b, i)$ are in D or at least three vertices of $[b, i)$ are in D . If all vertices of $[b, i)$ are not in D , then

$$\gamma_{[1,2]}^1(b, i : i) \leq |D| = \gamma_{[1,2]}^1(i). \quad (1)$$

If exactly one vertex $j \in [b, i)$ is in D , then

$$\gamma_{[1,2]}^1(b, i : i, j) \leq |D| = \gamma_{[1,2]}^1(i). \quad (2)$$

If exactly two vertices $j, k \in [b, i)$ are in D such that $k < j$, then

$$\gamma_{[1,2]}^1(b, i : i, j, k) \leq |D| = \gamma_{[1,2]}^1(i). \quad (3)$$

Assume that at least three vertices of $[b, i)$ are in D and let $i' = \max D \cap [1, i)$ and $i'' = \max D \cap [1, i')$. By Lemma 4, $x \in D$ for all $x \in [b, i'')$. Hence,

$$\gamma_{[1,2]}^{11}(b, i : i, i', i'') \leq |D| = \gamma_{[1,2]}^1(i). \quad (4)$$

Clearly, $\gamma_{[1,2]}^1(i) \leq \min\{\gamma_{[1,2]}^1(b, i : i), \gamma_{[1,2]}^1(b, i : i, j), \gamma_{[1,2]}^1(b, i : i, j', k'), \gamma_{[1,2]}^{11}(b, i : i, j', k') : b \leq j < i, b \leq k' < j' < i\}$. This, together with inequalities (1)–(4), completes the proof of the lemma. \square

Lemma 6. Let $i \in [1, n]$, $1 < b = \text{low}(i)$ and $a = \max\{j, b\}$.

- (i) If $j \leq \text{low}(b-1)$, then $\gamma_{[1,2]}^1(j, i : i)$ is not defined.
- (ii) If $\text{low}(b-1) < j < \text{low}(b, i-1)$, then $\gamma_{[1,2]}^1(j, i : i) = \gamma_{[1,2]}^0(j, b-1) + 1$.
- (iii) If $\text{low}(b, i-1) \leq j$, then $\gamma_{[1,2]}^1(j, i : i) = \min\{\gamma_{[1,2]}^0(\text{low}(c), j-1), \gamma_{[1,2]}^0(\text{low}(c), j-1 : k), \gamma_{[1,2]}^1(\text{low}(c), j-1 : j-1) : k \in [\text{low}(c), j-1]\} + 1$, where $c \in [a, i-1)$ such that $\text{low}(c) \leq \text{low}(x)$ for all $x \in [a, i-1)$.

Proof. Let D be an $[1, 2]$ -dominating set of $G[1, i]$ with minimum cardinality such that $i \in D$ and $x \notin D$ for all $x \in [j, i)$. So, $|D| = \gamma_{[1,2]}^1(j, i : i)$. If $j \leq \text{low}(b-1)$, then there is no vertex in D to dominate $b-1$ and so D is not an $[1, 2]$ -dominating set of $G[1, i]$, that is, $\gamma_{[1,2]}^1(j, i : i)$ is not defined. See Figure 8(a). In the rest of the proof, we assume that $\text{low}(b-1) < j$. Recall that $a = \max\{j, b\}$. Let $c \in [a, i-1)$ be a vertex such that $\text{low}(c) \leq \text{low}(x)$ for all $x \in [a, i-1)$, that is, $\text{low}(c) = \text{low}(a, i-1)$. We distinguish two cases depending on $j < \text{low}(c)$ or $\text{low}(c) \leq j$. In the following we consider these cases.




Fig. 8 Illustrating an $[1, 2]$ -dominating set D of $G[1, i]$ such that $i \in D$ and $x \notin D$ for all $x \in [j, i)$; (a) $j \leq \text{low}(b-1)$, (b) $\text{low}(b-1) < j < \text{low}(c)$, (c) $\text{low}(c) \leq j \leq b = a \leq c < i$, and (d) $\text{low}(c) \leq b < j = a \leq c < i$, where $b = \text{low}(i)$, $a = \max\{j, b\}$ and $\text{low}(c) = \text{low}(b, i-1)$. Note that a vertex with label 1 is in D and a vertex with label 0 is not in D .

- Assume that $j < \text{low}(c)$. If $b \leq j$, that is, $a = j$, then $\text{low}(c) = \text{low}(j-1, i-1) \leq \text{low}(j) \leq j$, a contradiction. Hence, $j < b$. See Figure 8(b). Recall that $x \notin D$ for all $x \in [j, i)$. Let $y \in [j, b-1]$. Since $y \notin D$ and y is adjacent to i , there are at least one and at most two vertices in D that are adjacent to y , that is, $1 \leq |[1, j) \cap D| \leq 2$. Let $D' = D \setminus \{i\}$. We get that D' is an $[1, 2]$ -dominating set of $G[1, b-1]$ such that $x \notin D'$ for all $x \in [j, b-1]$. Thus, $\gamma_{[1,2]}^0(j, b-1) \leq |D'| = \gamma_{[1,2]}^1(j, i : i) - 1$. Conversely, assume that S is an $[1, 2]$ -dominating set of $G[1, b-1]$ with minimum cardinality such that $x \notin S$ for all $x \in [j, b-1]$. So, $|S| = \gamma_{[1,2]}^0(j, b-1)$. We see that

$S' = S \cup \{i\}$ is an $[1, 2]$ -dominating set of $G[1, i]$ such that $i \in S'$ and $x \notin D$ for all $x \in [j, i - 1]$. So, $\gamma_{[1,2]}^1(j, i : i) \leq |S'| = \gamma_{[1,2]}^0(j, b - 1) + 1$. This completes the proof of the case (ii).

- Assume that $\mathbf{low}(c) \leq j$. If $j \leq b$, then $\mathbf{low}(c) \leq j \leq b = a \leq c < i$, see Figure 8(c), otherwise, $\mathbf{low}(c) \leq b < j = a \leq c < i$, see Figure 8(d). Since $a = \max\{j, b\}$, we obtain that c is adjacent to i , where $i \in D$. Since $x \notin D$ for all $x \in [j, i)$, $c \notin D$ and so c is adjacent to at most one vertex of $D \setminus \{i\}$, that is, either (a) $x \notin D$ for all $x \in [\mathbf{low}(c), j - 1]$ or (b) exactly one vertex of $[\mathbf{low}(c), j - 1]$ is in D . In the following we consider these cases.

- (a) Assume that $x \notin D$ for all $x \in [\mathbf{low}(c), j - 1]$. We obtain that $D' = D \setminus \{i\}$ is an $[1, 2]$ -dominating set of $G[1, j - 1]$ such that $x \notin D'$ for all $x \in [\mathbf{low}(c), j - 1]$. Hence,

$$\gamma_{1,2}^0(\mathbf{low}(c), j - 1) \leq |D'| = \gamma_{1,2}^1(j, i : i) - 1. \quad (5)$$

- (b) Assume that exactly one vertex $k \in [\mathbf{low}(c), j - 1]$ is in D . If $k < j - 1$, then $D' = D \setminus \{i\}$ is an $[1, 2]$ -dominating set of $G[1, j - 1]$ such that $k \in D'$ and $x \notin D'$ for all $x \in [\mathbf{low}(c), j - 1]$. Hence,

$$\gamma_{1,2}^0(\mathbf{low}(c), j - 1 : k) \leq |D'| = \gamma_{1,2}^1(j, i : i) - 1. \quad (6)$$

If $k = j - 1$, then $D' = D \setminus \{i\}$ is an $[1, 2]$ -dominating set of $G[1, j - 1]$ such that $j - 1 \in D'$ and $x \notin D'$ for all $x \in [\mathbf{low}(c), j - 1]$. Hence,

$$\gamma_{1,2}^1(\mathbf{low}(c), j - 1 : j - 1) \leq |D'| = \gamma_{1,2}^1(j, i : i) - 1. \quad (7)$$

Conversely, assume that S_1 is an $[1, 2]$ -dominating set of $G[1, j - 1]$ with minimum cardinality such that $x \notin S_1$ for all $x \in [\mathbf{low}(c), j - 1]$. So, $|S_1| = \gamma_{[1,2]}^0(\mathbf{low}(c), j - 1)$. Because $\mathbf{low}(c) = \mathbf{low}(a, i - 1)$, each vertex of $[b, i]$ is adjacent to no vertex of S_1 . Hence, $S'_1 = S_1 \cup \{i\}$ is an $[1, 2]$ -dominating set of $G[1, i]$ such that $i \in S'_1$ and $x \notin S'_1$ for all $x \in [j - 1, i)$. So,

$$\gamma_{1,2}^1(j, i : i) \leq |S'_1| = \gamma_{[1,2]}^0(\mathbf{low}(c), j - 1) + 1. \quad (8)$$

Assume that S_2 is an $[1, 2]$ -dominating set of $G[1, j - 1]$ with minimum cardinality such that $k \in S_2$ for some $k \in [\mathbf{low}(c), j - 1]$ and $x \notin S_2$ for all $x \in [\mathbf{low}(c), j - 1] \setminus \{k\}$. So, $|S_2| = \gamma_{[1,2]}^0(\mathbf{low}(c), j - 1 : k)$. Because $\mathbf{low}(c) = \mathbf{low}(a, i - 1)$, each vertex of $[b, i]$ is adjacent to at most one vertex k . Hence, $S'_2 = S_2 \cup \{i\}$ is an $[1, 2]$ -dominating set of $G[1, i]$ such that $i \in S'_2$ and $x \notin S'_2$ for all $x \in [j - 1, i)$. So,

$$\gamma_{1,2}^1(j, i : i) \leq |S'_2| = \gamma_{[1,2]}^0(\mathbf{low}(c), j - 1 : k) + 1. \quad (9)$$

Similarly, we obtain that

$$\gamma_{1,2}^1(j, i : i) \leq |S'_2| = \gamma_{[1,2]}^1(\mathbf{low}(c), j-1 : j-1) + 1. \quad (10)$$

The proof of the case (iii) follows from inequalities (1)–(6).

This completes the proof of the lemma. \square

Lemma 7. *Let $i \in [1, n]$, $j \in [1, i]$, $b = \mathbf{low}(i)$, $c = \mathbf{low}(j)$, $d = \mathbf{low}(j+1, i-1)$, $d' = \mathbf{low}(b, j-1)$, $d'' = \mathbf{low}(b, i-1)$, $z = \max\{x \in (j, i) : d = \mathbf{low}(x)\}$, and $k' = \min\{k, b, d, d'\}$.*

(i) *If either $j < b$ and $j < \mathbf{low}(x)$ for some $x \in (j, b)$ or $k < \min\{b, c\}$ and $k \leq \mathbf{low}(x)$ for some $x \in [k, \min\{b, c\})$, then $\gamma_{[1,2]}^1(k, i : i, j)$ is not defined,*

(ii) *assume that $j < b$ and $z < b$,*

(a) *if $k = j \leq d$, then $\gamma_{[1,2]}^1(k, i : i, j) = \gamma_{[1,2]}^1(j) + 1$,*

(b) *if $k < j$ and $\min\{k, d''\} \leq d$, then $\gamma_{[1,2]}^1(k, i : i, j) = \gamma_{[1,2]}^1(\min\{k, d''\}, j : j) + 1$,*

(c) *if $d < \min\{k, d''\}$, then $\gamma_{[1,2]}^1(k, i : i, j) = \min\{\gamma_{[1,2]}^1(d, j : j), \gamma_{[1,2]}^1(d, j : j, x) : x \in [d, \min\{k, d''\})\} + 1$,*

(iii) *assume that $j < b$ and $z \geq b$,*

(a) *if $k = j \leq d$, then $\gamma_{[1,2]}^1(k, i : i, j) = \gamma_{[1,2]}^1(j) + 1$,*

(b) *if $k < j$ and $k \leq d$, then $\gamma_{[1,2]}^1(k, i : i, j) = \gamma_{[1,2]}^1(k, j : j) + 1$,*

(c) *if $d < k$, then $\gamma_{[1,2]}^1(k, i : i, j) = \gamma_{[1,2]}^1(d, j : j) + 1$,*

(iv) *assume $j \geq b$,*

(a) *if $b = k = j \leq d$, then $\gamma_{[1,2]}^1(k, i : i, j) = \gamma_{[1,2]}^1(j) + 1$,*

(b) *if $b \leq c$, then $\gamma_{[1,2]}^1(k, i : i) < \gamma_{[1,2]}^1(k, i : i, j)$,*

(c) *if $c < b$, then $\gamma_{[1,2]}^1(k, i : i, j) = \gamma_{[1,2]}^1(k', j : j) + 1$.*

Proof. Let D be an $[1, 2]$ -dominating set of $G[1, i]$ with minimum cardinality such that $i, j \in D$ and $x \notin D$ for all $x \in [k, i] \setminus \{j\}$. So, $|D| = \gamma_{[1,2]}^1(k, i : i, j)$. If either $j < b$ and $j < \mathbf{low}(x)$ for some $x \in (j, b)$, see Figure 9(a), or $k < \min\{b, c\}$ and $k \leq \mathbf{low}(x)$ for some $x \in [k, \min\{b, c\})$, then there is no vertex of D to dominate x , a contradiction. This proves the case (i). In the rest of the proof, we assume that the case (i) does not hold.

Recall that $d = \mathbf{low}(j+1, i-1)$ and $z = \max\{x \in (j, i) : d = \mathbf{low}(x)\}$. If $j \geq b$, then $z \geq b$. Hence, we have (I) $j < b$ and $z < b$, (II) $j < b$ and $z \geq b$, or (III) $j \geq b$. In the following we consider these cases.

Case I. Assume that $j < b$ and $z < b$. If $d'' \leq k$, then y is adjacent to j . Hence, $y \notin D$ is adjacent to vertices i, j and so all vertices in $[d'', j)$ are not in D . Since $k \leq j$, we distinguish the following cases:




Fig. 9 Illustrating an $[1,2]$ -dominating set D of $G[1,i]$ such that $i, j \in D$ and $x \notin D$ for all $x \in [k,i] \setminus \{j\}$; (a) $j < b$ and $j < \text{low}(x)$ for some $x \in (j,b)$, (b) $k = j = d$, (c) $k < j$ and $k \leq d$, and (d) $d < k$. Note that a vertex with label 1 is in D and a vertex with label 0 is not in D .

- Assume $k = j \leq d$. See Figure 9(b). We get that $D' = D \setminus \{i\}$ is an $[1,2]$ -dominating set of $G[1,j]$ such that $j \in D'$. Hence, $\gamma_{[1,2]}^1(j) \leq |D'| = \gamma_{[1,2]}^1(k, i : i, j) - 1$. Conversely, assume that S is an $[1,2]$ -dominating set of $G[1,j]$ with minimum cardinality such that $j \in S$. So, $|S| = \gamma_{[1,2]}^1(j)$. Since $j \leq d$ and the case (i) does not hold, each vertex of (j,i) is adjacent to at most two vertices i and j . Hence, $S' = S \cup \{i\}$ is an $[1,2]$ -dominating set of $G[1,i]$ such that $i, j \in S'$ and $x \notin S'$ for all $x \in [k,i] \setminus \{j\}$ and so $\gamma_{[1,2]}^1(k, i : i, j) \leq |S'| = \gamma_{[1,2]}^1(j) + 1$. This completes the proof of the case (ii.a).
- Assume $k < j$. See Figure 9(c). Clearly, y is adjacent to i . We get that $D' = D \setminus \{i\}$ is an $[1,2]$ -dominating set of $G[1,j]$ such that $j \in D'$ and $x \notin D'$ for all $x \in [\min\{k, d''\}, j)$. Hence, $\gamma_{[1,2]}^1(\min\{k, d''\}, j : j) \leq |D'| = \gamma_{[1,2]}^1(k, i : i, j) - 1$. Conversely, assume that S is an $[1,2]$ -dominating set of $G[1,j]$ with minimum cardinality such that $j \in S$ and $x \notin S$ for all $x \in [\min\{k, d''\}, j)$. So, $|S| = \gamma_{[1,2]}^1(\min\{k, d''\}, j : j)$. Since $x \notin S$ for all $x \in [\min\{k, d''\}, j)$, $k \leq d$ and the case (i) does not hold, each vertex of (j,i) is adjacent to at most two vertices i and j . Hence, $S' = S \cup \{i\}$ is an $[1,2]$ -dominating set of $G[1,i]$ such that $i, j \in S'$ and $x \notin S'$ for all $x \in [\min\{k, d''\}, i) \setminus \{j\}$ and so $\gamma_{[1,2]}^1(k, i : i, j) \leq |S'| = \gamma_{[1,2]}^1(\min\{k, d''\}, j : j) + 1$. This completes the proof of the case (ii.b).

- Assume $d < \min\{k, d''\}$. See Figure 9(d). Since $z \notin D$ and z is adjacent to j , the vertex z is adjacent to at most one vertex of D . Thus, at least one vertex of $[d, \min\{k, d''\})$ is in D , that is, either no vertex of $[d, \min\{k, d''\})$ is in D or exactly one vertex of $[d, \min\{k, d''\})$ is in D . In the following, we consider these cases. If no vertex of $[d, \min\{k, d''\})$ is in D , then we get that $D'_1 = D \setminus \{i\}$ is an $[1, 2]$ -dominating set of $G[1, j]$ such that $j \in D'_1$ and $x \notin D'_1$ for all $x \in [d, j)$. Hence,

$$\gamma_{[1,2]}^1(d, j : j) \leq |D'_1| = \gamma_{[1,2]}^1(k, i : i, j) - 1. \quad (11)$$

If exactly one vertex $w \in [d, \min\{k, d''\})$ is in D , then we get that $D'_2 = D \setminus \{i\}$ is an $[1, 2]$ -dominating set of $G[1, j]$ such that $j, w \in D'_2$ and $x \notin D'_2$ for all $x \in [d, j) \setminus \{w\}$. Assume that $x' \in [d, \min\{k, d''\})$. Hence,

$$\gamma_{[1,2]}^1(d, j : j, x') \leq |D'_2| = \gamma_{[1,2]}^1(k, i : i, j) - 1. \quad (12)$$

Conversely, assume that S_1 is an $[1, 2]$ -dominating set of $G[1, j]$ with minimum cardinality such that $j \in S_1$ and $x \notin S_1$ for all $x \in [d, j)$. So, $|S_1| = \gamma_{[1,2]}^1(d, j : j)$. Since $x \notin S_1$ for all $x \in [d, j)$, $d < k$ and the case (i) does not hold, each vertex of (j, i) is adjacent to at most two vertices i and j . Hence, $S'_1 = S_1 \cup \{i\}$ is an $[1, 2]$ -dominating set of $G[1, i]$ such that $i, j \in S'_1$ and $x \notin S'_1$ for all $x \in [k, i) \setminus \{j\}$ and so

$$\gamma_{[1,2]}^1(k, i : i, j) \leq |S'_1| = \gamma_{[1,2]}^1(d, j : j) + 1. \quad (13)$$

Let $x \in [d, \min\{k, d''\})$. Assume that S_2 is an $[1, 2]$ -dominating set of $G[1, j]$ with minimum cardinality such that $j, x \in S_2$ and $w \notin S_2$ for all $w \in [d, j) \setminus \{x\}$. So, $|S_2| = \gamma_{[1,2]}^1(d, j : j, x)$. We get that $S'_2 = S_2 \cup \{i\}$ is an $[1, 2]$ -dominating set of $G[1, i]$ such that $i, j \in S'_2$ and $w \notin S'_2$ for all $w \in [k, i) \setminus \{j\}$ and so

$$\gamma_{[1,2]}^1(k, i : i, j) \leq |S'_2| = \gamma_{[1,2]}^1(d, j : j, x) + 1. \quad (14)$$

Inequalities (11)–(14) completes the proof of the case (ii.c).

Case II. Assume that $j < b$ and $z \geq b$. The proofs of the cases (iii.a) and (iii.c) is similar to the proof of the case (ii.a) and the proof of the case (iii.b) is similar to the proof of the case (ii.b), respectively.

Case III. Assume that $j \geq b$. If $b = k = j \leq d$, then similar to the case (ii.a) we prove that $\gamma_{[1,2]}^1(k, i : i, j) = \gamma_{[1,2]}^1(j) + 1$ that completes the proof of the case (iv.a). Assume that the case $b = k = j \leq d$ does not hold. We distinguish two cases depending on $b \leq c$ or $c < b$.

- Assume that $b \leq c$. Recall that $b = \text{low}(i)$ and $c = \text{low}(j)$. It obtains that each vertex adjacent to j is also adjacent to i . We see that $D' = D \setminus \{j\}$ is an $[1, 2]$ -dominating set of $G[1, i]$ such that $i \in D'$ and $x \notin D'$ for all $x \in [k, i)$ and so $\gamma_{[1,2]}^1(k, i : i) \leq |D'| = \gamma_{[1,2]}^1(k, i : i, j) - 1$. This completes the proof of the case (iv.b).

- Assume that $c < b$. Recall that $k \leq j$, $d = \mathbf{low}(j + 1, i - 1)$, $d' = \mathbf{low}(b, j - 1)$ and $k' = \min\{k, d, d'\}$. Clearly, k' exists. Since $b \leq j$, $k' \leq b$. We claim that all vertices of $[k', j)$ are not in D . If $k' = k$, the claim is clear. Assume that $k' = d$. So, $d \leq b$. Recall that $z = \max\{x \in (j, i) : d = \mathbf{low}(x)\}$. We obtain that $z \notin D$ is adjacent to $i, j \in D$. Since D is an $[1, 2]$ -dominating set, all vertices of $[d, j)$ are not in D . Assume that $k' = d'$. Let $z' \in [b, j - 1]$ such that $\mathbf{low}(z') = d'$. Since $c < b$, we get that $z' \notin D$ is adjacent to $i, j \in D$ and so all vertices of $[d', j)$ are not in D . Hence, all vertices of $[k', j)$ are not in D . The proof of the case (iv.c) is similar to the proof of the case (ii.b).

This completes the proof of the lemma. \square

Similarly, we can prove the following lemmas.

Lemma 8. *Let $i \in [1, n]$, $j \in [1, i]$, $k \in [1, j]$, $b = \mathbf{low}(i)$, $c = \mathbf{low}(j)$, $d = \mathbf{low}(k)$, $e = \mathbf{low}(j + 1, i - 1)$, $e' = \mathbf{low}(b, i - 1)$ and $z = \max\{x \in (j, i) : e = \mathbf{low}(x)\}$.*

- (i) *If $j < b$ and $j < \mathbf{low}(x)$ for some $x \in (j, b)$, $k < \min\{b, c\}$ and $k < \mathbf{low}(x)$ for some $x \in (k, \min\{b, c\})$ or $l < \min\{b, c, d\}$ and $l \leq \mathbf{low}(x)$ for some $x \in [l, \min\{b, c, d\})$, then $\gamma_{[1,2]}^1(l, i : i, j, k)$ is not defined,*
- (ii) *assume that $j < b$ and $z < b$,*
 - (a) *if $j \leq e$, then $\gamma_{[1,2]}^1(l, i : i, j, k) = \gamma_{[1,2]}^1(l, j : j, k) + 1$,*
 - (b) *if $e < j$ and $e' \leq k$, then $\gamma_{[1,2]}^1(l, i : i, j, k)$ is not defined,*
 - (c) *if $e < j$ and $e' > k$, then $\gamma_{[1,2]}^1(l, i : i, j, k) = \gamma_{[1,2]}^1(\min\{l, e\}, j : j, k) + 1$,*
- (iii) *assume that $z \geq b$,*
 - (a) *if $e \leq k$, then $\gamma_{[1,2]}^1(l, i : i, j, k)$ is not defined,*
 - (b) *if $e > k$, then $\gamma_{[1,2]}^1(l, i : i, j, k) = \gamma_{[1,2]}^1(l, j : j, k) + 1$.*

Lemma 9. *Let $b = \mathbf{low}(i)$, $c = \mathbf{low}(j)$, $d = \mathbf{low}(k)$, $e = \mathbf{low}(j + 1, i - 1)$, $e' = \mathbf{low}(b, i - 1)$ and $z = \max\{x \in (j, i) : e = \mathbf{low}(x)\}$.*

- (i) *If either $j < b$ and $j < \mathbf{low}(x)$ for some $x \in (j, b)$ or $k < \min\{b, c\}$ and $k < \mathbf{low}(x)$ for some $x \in (k, \min\{b, c\})$, then $\gamma_{[1,2]}^{11}(l, i : i, j, k)$ is not defined,*
- (ii) *assume that $j < b$ and $z < b$,*
 - (a) *if $e < k$ or $e' \leq k$, then $\gamma_{[1,2]}^{11}(l, i : i, j, k)$ is not defined,*
 - (b) *if $j \leq e$ and $k - l = 1$, then $\gamma_{[1,2]}^{11}(l, i : i, j, k) = \gamma_{[1,2]}^1(l, j : j, k, l) + 1$,*
 - (c) *if $j \leq e$ and $k - l > 1$, then $\gamma_{[1,2]}^{11}(l, i : i, j, k) = \gamma_{[1,2]}^{11}(l, j : j, k, k - 1) + 1$,*
 - (d) *if $k \leq e < j$, $e' > k$ and $k - l = 1$, then $\gamma_{[1,2]}^{11}(l, i : i, j, k) = \gamma_{[1,2]}^1(l, j : j, k, l) + 1$,*
 - (e) *if $k \leq e < j$, $e' > k$ and $k - l > 1$, then $\gamma_{[1,2]}^{11}(l, i : i, j, k) = \gamma_{[1,2]}^{11}(l, j : j, k, k - 1) + 1$,*
- (iii) *assume that $z \geq b$,*
 - (a) *if $e \leq k$, then $\gamma_{[1,2]}^{11}(l, i : i, j, k)$ is not defined,*
 - (b) *if $e > k$ and $k - l = 1$, then $\gamma_{[1,2]}^{11}(l, i : i, j, k) = \gamma_{[1,2]}^1(l, j : j, k, l) + 1$,*
 - (c) *if $e > k$ and $k - l > 1$, then $\gamma_{[1,2]}^{11}(l, i : i, j, k) = \gamma_{[1,2]}^{11}(l, j : j, k, k - 1) + 1$,*

Now, we are ready to present our algorithm (Algorithm 1) for computing $\gamma_{[1,2]}(G)$, where G is an interval graph. Note that if $12 \in E$, then the case (iv.b) of Lemma 7 holds and so $\gamma_{[1,2]}^1(1, 2 : 2) < \gamma_{[1,2]}^1(1, 2 : 2, 1)$. By considering Lemmas 2, 3, 5, 6, 7 and 8 we see that if we use $\gamma_{[1,2]}^1(1, 2 : 2, 1)$ for computing a value, then we also use $\gamma_{[1,2]}^1(1, 2 : 2)$ for computing that value. Hence, in Line 12 we set $\gamma_{[1,2]}^1(1, 2 : 2, 1)$ to ∞ . We also use this note in Line 20 for computing $\gamma_{[1,2]}^1(k, i : i, j)$.

Algorithm 1 [1,2]-DOMINATION-INTERVAL-GRAPH(G)

Input: An interval graph $G = (V, E)$ of order n .

Output: $\gamma_{[1,2]}(G)$.

- 1: Compute a numbering $(1, 2, \dots, n)$ of vertices of G satisfying the condition of Corollary 1.
 - 2: Compute $\text{low}(i)$ for all $i \in [1, n]$.
 - 3: Compute $\text{low}(j, i)$ for all $1 \leq j < i \leq n$.
 - 4: Compute $\text{maxlow}(i)$ for all $i \in [1, n]$.
 - 5: $\gamma_{[1,2]}^0(1) \leftarrow \infty$
 - 6: If $12 \in E$, then $\gamma_{[1,2]}^0(2) \leftarrow 1$, otherwise, $\gamma_{[1,2]}^0(2) \leftarrow \infty$
 - 7: $\gamma_{[1,2]}^0(1, 2) \leftarrow \infty$
 - 8: If $12 \in E$, then $\gamma_{[1,2]}^0(1, 2 : 1) \leftarrow 1$, otherwise, $\gamma_{[1,2]}^0(1, 2 : 1) \leftarrow \infty$
 - 9: $\gamma_{[1,2]}^1(1) \leftarrow 1$
 - 10: If $12 \in E$, then $\gamma_{[1,2]}^1(2) \leftarrow 1$, otherwise, $\gamma_{[1,2]}^1(2) \leftarrow 2$
 - 11: If $12 \in E$, then $\gamma_{[1,2]}^1(1, 2 : 2) \leftarrow 1$, otherwise, $\gamma_{[1,2]}^1(1, 2 : 2) \leftarrow \infty$
 - 12: $\gamma_{[1,2]}^1(1, 2 : 2, 1) \leftarrow \infty$
 - 13: $\gamma_{[1,2]}^{11}(1, 3 : 3, 2, 1) \leftarrow 3$
 - 14: **for** $i = 3$ to n **do**
 - 15: **for** $1 \leq j < i$ **do**
 - 16: Compute $\gamma_{[1,2]}^0(j, i)$ using Lemma 2.
 - 17: Compute $\gamma_{[1,2]}^1(j, i : i)$ using Lemma 6.
 - 18: **end for**
 - 19: **for** $1 \leq k \leq j < i$ **do**
 - 20: Compute $\gamma_{[1,2]}^0(k, i : j)$ using Lemma 3.
 - 21: Compute $\gamma_{[1,2]}^1(k, i : i, j)$ using Lemma 7.
 - 22: **end for**
 - 23: **for** $1 \leq l < k' < j < i$ **do**
 - 24: Compute $\gamma_{[1,2]}^1(l, i : i, j, k')$ using Lemma 8.
 - 25: Compute $\gamma_{[1,2]}^{11}(l, i : i, j, k')$ using Lemma 9.
 - 26: **end for**
 - 27: Compute $\gamma_{[1,2]}^0(i) = \gamma_{[1,2]}^0(i, i)$ using Lemma 2.
 - 28: Compute $\gamma_{[1,2]}^1(i)$ using Lemma 5.
 - 29: **end for**
 - 30: **return** $\min\{\gamma_{[1,2]}^0(n), \gamma_{[1,2]}^1(n)\}$
-

Theorem 1. Let $G = (V, E)$ be an interval graph of order n . Algorithm 1 computes $\gamma_{[1,2]}(G)$ in $O(n^4)$ time.

Proof. Let $(1, 2, \dots, n)$ be a numbering of vertices of G computed in Line 1. This numbering of vertices of $G = (V, E)$ can be computed in $O(|V| + |E|)$ time [12]. We need $O(n^3)$ time to compute $\text{low}(i)$ for all $i \in [1, n]$, $\text{low}(j, i)$ for all $1 \leq j < i \leq n$ and $\text{maxlow}(i)$ for all $i \in [1, n]$ in Lines 2-4. It is easy to compute the values $\gamma_{[1,2]}^0(1)$, $\gamma_{[1,2]}^0(2)$, $\gamma_{[1,2]}^0(1, 2)$, $\gamma_{[1,2]}^0(1, 2 : 1)$, $\gamma_{[1,2]}^1(1)$, $\gamma_{[1,2]}^1(2)$, $\gamma_{[1,2]}^1(1, 2 : 2)$, $\gamma_{[1,2]}^1(1, 2 : 2, 1)$ and $\gamma_{[1,2]}^{11}(1, 3 : 3, 2, 1)$ as seen in Lines 5-13 in $O(1)$ time. Let $i \in [3, n]$, $j \in [1, i]$, $k \in [1, j]$, $k' \in [2, j]$ and $l \in [1, k']$. By Lemmas 2, 3, 5, 6, 7, 8 and 9, compute $\gamma_{[1,2]}^0(i)$, $\gamma_{[1,2]}^0(j, i)$, $\gamma_{[1,2]}^0(k, i : j)$, $\gamma_{[1,2]}^1(i)$, $\gamma_{[1,2]}^1(j, i : i)$, $\gamma_{[1,2]}^1(k, i : i, j)$, $\gamma_{[1,2]}^1(k, i : i, j, k')$ and $\gamma_{[1,2]}^{11}(k, i : i, j, k')$ in Lines 15-29. We see that the running time of Lines 14-29 is $O(n^4)$. Hence, the running time of Algorithm 1 is $O(n^4)$. Clearly, $\gamma_{[1,2]}(G) = \min\{\gamma_{[1,2]}^0(n), \gamma_{[1,2]}^1(n)\}$. This completes the proof of the theorem. \square

3 $[1, 2]$ -domination in Circle Graphs

In this section, we prove hardness results for a $[1, 2]$ -domination problem in circle graphs. We start with the main result of this section.

Theorem 2. $[1, 2]$ -domination problem is NP-complete in circle graphs.

Proof. Clearly $[1, 2]$ -domination problem is in NP. To prove that the problem is NP-hard, We shall reduce (in polynomial time) the 3SAT problem to the problem of finding a $[1, 2]$ -dominating set. Let $X = \{x_1, x_2, \dots, x_n\}$ be the set of boolean variable and $C = \{C_1, C_2, \dots, C_m\}$ be the set of clauses in an arbitrary instance of 3SAT, like I . In the first part, we construct a circle graph $G(I)$ based on the instance I and in the second part, we choose an integer k and show $G(I)$ has $[1, 2]$ -dominating set of size k if and only if I is satisfiable. Let us describe the construction of $G(I)$ produced in reduction from 3SAT to $[1, 2]$ -domination problem. Without loss of generality, we assume that every variable appears once in each clause. We choose an arbitrary point of the circle as the origin. The circle graph $G(I)$ is defined as follows:

First Step: Constructed Gadget for Clause C_j .

- We divide the circle into m disjoint open intervals $[s_j, s'_j]$ for $1 \leq j \leq m$, corresponding each clause,




Fig. 10 Intervals corresponding each clause

- We create two chords t_j^l and f_j^l associated with each literal l , $1 \leq l \leq 3$, that appears in clause C_j . Including the chord t_j^l in $[1, 2]$ -dominating set will correspond to setting

the associated literal to true and including the f_j^l chord in $[1, 2]$ -dominating set will correspond to setting the associated literal to false,

- For each literal in clause C_j we create a pair of chords a_j^l ,
- For each literal in clause C_j we create a pair of chords p_j^l . The purpose of each such pair is to dominate them by including the chords t_j^l or f_j^l ,
- For each literal in clause C_j , the interval between the endpoint of t_j^l and the endpoint of the first chords of a_j^l is called the truth interval, and the interval between the right point of the second chord of p_j^l and the right point of f_j^l is called false interval for literal l of clause j . (See Figure 11)




Fig. 11 Part of constructed gadget for clause C_j

- For each clause C_j , we create three pairs of chords of types u and w . Based on the endpoints of a_j^l pairs in clause C_j , we have pairs $(u_{12}, w_{12}), (u_{13}, w_{13})$ and, (u_{23}, w_{23}) . Also, we have two guard chords, g_1 and g_2 , connected to two the claw graphs centered at c_1 and c_2 , respectively. The purpose of each will be discussed later.
 - The chord u_{13} incidence with only the pair a_j^1 and w_{13} incidence with only the pair a_j^3 ,
 - The left point of u_{12} is before pair a_j^1 and endpoint of it is before the pair a_j^3 . So, the chord u_{12} incidence with the pairs a_j^1 and a_j^2 . The chord w_{12} start before the endpoint u_{13} and after crossing the chord u_{13} , the pairs a_j^2 , and the chord w_{13} will be finished,
 - The left point of u_{23} is after pair a_j^1 and endpoint of it is after the pair a_j^3 . So, the chord u_{23} incidence with the pairs a_j^2 and a_j^3 . The chord w_{23} start before the endpoint u_{13} and also before the start point of w_{12} . After crossing the chord u_{13} , the pairs a_j^2 , and the chords w_{12}, w_{13} will be finished,
 - The guard chord g_1 , from the left endpoint is connected to a chord c_1 which three independent chords cross it. The purpose of these independent chords is to force that c_1 contains in any minimum $[1, 2]$ -dominating set. So, the guard chord g_1 is always dominated by c_1 once. The right endpoint of g_1 will be finished after the start point of u_{23} . So, g_1 will incidence with the chords u_{12}, u_{13} , and u_{23} ,
 - The left endpoint of g_2 will start before the left endpoint of u_{12} and finish after the right endpoint of w_{13} . So, g_2 will incidence with the chords u_{12}, u_{23} , and w_{13} . The guard chord g_2 , from the right endpoint, is connected to a chord c_2 which three independent chords cross it. Similar to chord g_1 , the purpose of these independent chords is to force that c_2 contains in any minimum $[1, 2]$ -dominating set. So, the guard chord g_2 is always dominated by c_2 once.

(see Fig. 12 for an illustration)




Fig. 12 Part of constructed gadget for clause C_j

The key idea of our proof is as follows. Suppose that C_j is satisfied by some assignment of truth values to a variable. Then we can construct a $[1, 2]$ -dominating set that contains at least one chord t_j^l corresponding to a true literal in C_j and a pair of u and w chords associated with C_j . The chord t_j^l dominates the pairs a_j^l while the other pairs a_j^l and (u, w) 's are dominated by an appropriate pair of (u, w) . To ensure that exactly one of the chords t_j^l and f_j^l associated with literal l in clause j appears in any $[1, 2]$ -dominating set, a pair of independent chords p_j^l associated with the literal l adjacent to both t_j^l and f_j^l and no other chords. In other words, since the independent chords p_j^l 's start from the left endpoint of f_j^l and the right endpoint t_j^l , to dominate them, t_j^l or f_j^l or both of chords in p_j^l should be select. Later we will argue that any minimum $[1, 2]$ -dominating set must contain exactly one of chords t_j^l or f_j^l otherwise both of p_j^l must be selected that leads to a larger dominating set.

The clause C_j is satisfied if and only if at least one of three literals involved in C_j is true. As we mentioned, this will correspond to at least one of the chords t_j^l appearing in the dominating set. So this chord will dominate t_j^l , f_j^l , and both pairs a_j^l and p_j^l . To dominate the remaining pairs a_j associated with other literals in C_j we use six chords of type u and w as follows. u_{13} and w_{13} cross the pairs a_j^1 and a_j^3 , respectively. If the second literal is true that leads t_j^2 be in $[1, 2]$ -dominating set, two pairs a_j^1 and a_j^3 remain undominated, so selecting u_{13} and w_{13} dominate them.

If the first literal is true the pairs a_j^2 and a_j^3 are not dominated. Although by choosing u_{23} both pairs a_j^2 and a_j^3 are dominated, the chords w_{12} and w_{23} remain undominated. As they cross each other, we choose one of them. Similarly, If the third literal is true the pairs a_j^1 and a_j^2 are not dominated. Although by choosing u_{12} both pairs a_j^1 and a_j^2 are dominated, the chords w_{12} and w_{23} remain undominated. As they cross each other, we choose one of them.

Second Step: Consistency Condition

In the above construction, we obtain m pairwise disjoint sections, each associated with one clause. In the second part of our construction, we use connection chords that

force a variable to have the same truth value throughout all clauses. It is known as the consistency condition. we create connection chords tt , ff , tf , and ft that connect the clauses as follows. So, the consistency of truth values of literals throughout all clauses will be preserved.

- If variable x_i appears as a positive or negative literal in two clauses C_j and C_k , we create a chord tf between two sections associated with clauses C_j and C_k such that the left endpoint of this chord is in the truth interval of literal x_i in clause C_j and its right endpoint is in the false interval of C_k . Also, we add a chord ft such that the left endpoint of this chord is in the false interval of literal x_i in the clause C_j and its right endpoint is in the truth interval of C_k .
- If variable x_i appears as a positive literal in clause C_j and as a negative literal clause C_k or vice versa, we create a chord tt between two sections associated with clauses C_j and C_k such that the left endpoint of this chord is in the truth interval of literal x_i in clause C_j and its right endpoint is in truth interval of C_k . Also, we add a chord ff such that the left endpoint of this chord is in the false interval of literal x_i in clause C_j and its right endpoint is in the false interval of C_k .
- The left endpoints of each connection chord intersect with a claw. In other words, three independent chords connect with another chord to a connection chord.

Figure 13 shows all chords associated with a two clause and four variables.




Fig. 13 Representation of chords between the clauses $(\bar{x}_1 \vee x_2 \vee x_3) \wedge (x_1 \vee x_2 \vee x_4)$

Let $k = 7m + 2t$ where m is the number of clauses and t is the total number of repetitions of variables in clauses. Theorem 2 directly follows from the following result.

Lemma 10. $G(I)$ has a $[1, 2]$ -dominating set D of cardinality at most $k = 7m + 2t$ if the instance I is satisfiable.

Proof. Given a satisfying truth assignment for I , we show how to construct a $[1, 2]$ -dominating set D of size $k = 7m + 2t$. Let us consider that the clause C_j contains three literals corresponding to variables x_i, x_k , and x_l where $i \leq k \leq l$. We assume the variable x_i appears in the first literal and x_k, x_l are the second and third literals

in C_j , respectively. Since I is satisfied, at least one of the literals x_i, x_k , and x_l must be true. Let's assume without loss of generality that the only first literal of C_j is true and the second and third literals are false according to the assignment. So we have included t_j^l, f_j^2 , and f_j^3 in D . This ensures that all the pairs p_1, p_2 , and p_3 have been dominated. Also, the pair a_j^1 is dominated, but the pairs a_j^2 and a_j^3 are not dominated. To dominate them and all pairs (u, w) in this clause, we put u_{23}, w_{23} to D . So to dominate this clause, we need at least five chords. Also the chords c_1 and c_2 must be put in D to dominate the three independent chords crossing them.

Generally, for each literal l in clause C_j that is true in the assignment of I , we include in D the chord(s) t_j^l where l is the literal number in clause C_j . Otherwise, we include in D the chord(s) f_j^l . The chords t_j^l dominate the pair a_j^l . So, to dominate the remaining a_j pairs and the pairs (u, w) in C_j , we should choose at least one of (u, w) pairs. Also the chords c_1 and c_2 must be put in D .

For each chord tt, tf, ft , and ff , the chord s connected to them put in dominating set so all these chords are dominated once. otherwise, we must choose three independent chords connected to s the leads s be dominated three times and contradiction. So for each chord tt, tf, ft , and ff , we need one chord in dominating set. According to the construction for each variable repeated in two clauses, we have two connection chords. If the total number of variable repartition is t , we include $2t$ chords in D . It is clear that D is a $[1, 2]$ -dominating set. □

Lemma 11. *The 3SAT instance I is satisfiable if $G(I)$ has a $[1, 2]$ -dominating set of cardinality at most $k = 7m + 2t$.*

Proof. Let D of size $k = 7m + 2t$ be the given $[1, 2]$ -dominating set. We show there is a truth assignment to the variables in X that satisfy I . Recall that associated with each literal l that appears in a clause C_j , there is a pair of chords p_j^l . Since no chord is adjacent to two different pairs p 's, D must contain, for each l and j , either t_j^l or f_j^l or both chords in p . If for each literal l in a clause j , D contains t_j^l or f_j^l (but not both) then exactly $3m$ type t or f chord in D suffice to dominate all pairs of chords p . Otherwise, if D contains any pairs of p more than $3m$ are necessary to dominate chords in pairs p . Also, all c_1 's and c_2 's, totally $2m$, chords are in D to dominate the three crossed chord to them.

There is no chord in $G(I)$ adjacent to (u, w) pairs associated with different clauses. Thus the set of chords in D which dominate the (u, w) pairs in different clauses are pairwise disjoint. Also, no chord in $G(I)$ is adjacent to all the three pairs (u, w) in a clause. Therefore D must contain at least two chords per a clause to dominate all (u, w) pairs in a clause. Combined with the argument in the previous paragraph, this means exactly $3m$ chords in D dominate p pairs and $2m$ chords dominate the (u, w) pairs. This further implies that for each variable x_i appearing in a clause C_j , either t_j^l or f_j^l belongs to D .

Suppose three literal l_1, l_2 , and l_3 in clause C_j , and assume none of t_j^1, t_j^2 , and t_j^3 contain in D (All f chords are selected). So, all pairs a_j^1, a_j^2 and a_j^3 are not dominated. There is no (u, w) pair that can dominate all three pairs a_j 's. So at least one t

chord must be selected to dominate one a_j pair, and the other remaining a_j 's will be dominated by the appropriate pair (u, w) .

Till now, we show that for each clause C_j , D contains at least one t chord and one pair (u, w) . Based on the construction of the gadget, each chord tt, tf, ft , and ff are attached to a claw chord, centered at s . So, all the chords s must be in domination and as we have $2t$ such chords, then the total number of our $[1, 2]$ -domination is equal to $|D| = 7m + 2t$. Now, we can construct a satisfying truth assignment for I as follows. If D contains a t chord associated with a literal involving x_i , then we set this literal (and the corresponding value to its variable) to true in I . Similarly, D contains a f chord associated with a literal involving x_i , then we set this literal to false in I . \square

The proof of theorem 2 concludes from Lemma 10 and 11. \square

Conflict of Interests: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

4 Conclusion

Applying a constraint that determines the maximum number of times a vertex outside the dominating set, like D , is dominated by the elements inside D , makes the problem easier or harder than the original domination problem. In this paper, first we have designed a polynomial algorithm for $[1, 2]$ -dominating problem in proper interval graphs. Although it seems $O(n^4)$ is huge, it is an affirmative answer to the open problem on the complexity class of $[1, 2]$ -Domination problem in non-proper interval graphs. Designing more efficient algorithm for this problem is an interesting open problem. Then, we have shown that the $[1, 2]$ -Dominating Set problem is NP -complete on circle graphs. It is thus natural to ask for which subclass of circle graphs, like distance-hereditary or k -polygon, the problem has polynomial-time algorithm. Meanwhile, it would be desirable to show the complexity of $[1, 2]$ -dominating problem on other classes of graphs marked with a question mark in Figure 3.

References

- [1] BISHNU, A., DUTTA, K., GHOSH, A., AND PAUL, S. $(1, j)$ -set problem in graphs. *Discrete Mathematics* 339, 10 (2016), 2515–2525.
- [2] BOOTH, K. S., AND LUEKER, G. S. Testing for the consecutive ones property, interval graphs, and graph planarity using pq-tree algorithms. *Journal of computer and system sciences* 13, 3 (1976), 335–379.
- [3] CHELLALI, M., HAYNES, T. W., HEDETNIEMI, S. T., AND MCRAE, A. $[1, 2]$ -sets in graphs. *Discrete Applied Mathematics* 161, 18 (2013), 2885–2893.

- [4] DEJTER, I. J. Quasiperfect domination in triangular lattices. *Discussiones Mathematicae Graph Theory*, 29 (2009), 179–198.
- [5] ETESAMI, O., GHAREGHANI, N., HABIB, M., HOOSHMANDASL, M., NASERASR, R., AND SHARIFANI, P. When an optimal dominating set with given constraints exists. *Theoretical Computer Science* 780 (2019), 54–65.
- [6] GAREY, M. R., AND JOHNSON, D. S. Computers and intractability. *freeman San Francisco*, (1979).
- [7] GOHARSHADY, A. K., HOOSHMANDASL, M. R., AND MEYBODI, M. A. $[1, 2]$ -sets and $[1, 2]$ -total sets in trees with algorithms. *Discrete Applied Mathematics* 198 (2016), 136–146.
- [8] GOLUBIC, M. C. Algorithmic graph theory and perfect graphs, acad. *Press, New York* (1980).
- [9] HAYNES, T. W., HEDETNIEMI, S., AND SLATER, P. *Fundamentals of domination in graphs*. CRC Press, (1998).
- [10] MEYBODI, M. A., FOMIN, F. V., MOUAWAD, A. E., AND PANOLAN, F. On the parameterized complexity of $[1, j]$ -domination problems. *Theoretical Computer Science* 804 (2020), 207–218.
- [11] POUREIDI, A. Algorithm and hardness results in double roman domination of graphs. *Theoretical Computer Science* 911 (2022), 70–79.
- [12] RAMALINGAM, G., AND RANGAN, C. P. A unified approach to domination problems on interval graphs. *Information Processing Letters* 27, 5 (1988), 271–274.
- [13] ROBERTS, F. S. Indifference graphs. proof techniques in graph theory. In *Proceedings of the Second Ann Arbor Graph Conference, Academic Press, New York* (1969).
- [14] SHARIFANI, P., HOOSHMANDASL, M., AND ALIKHANI, S. A linear-time algorithm to compute total $[1, 2]$ -domination number of block graphs. *AUT Journal of Mathematics and Computing* 1, 2 (2020), 263–270.
- [15] SHARIFANI, P., AND HOOSHMANDASL, M. R. A linear algorithm for computing $\gamma_{[1,2]}$ -set in generalized series-parallel graphs. *arXiv preprint arXiv:1707.06443* (2017).
- [16] SHARIFANI, P., HOOSHMANDASL, M. R., AND MEYBODI, M. A. An explicit construction of optimal dominating sets in grid. *AKCE International Journal of Graphs and Combinatorics*, 17 (3) (2020), 870–876.
- [17] TELLE, J. A. Complexity of domination-type problems in graphs. *Nord. J.*

Comput. 1, 1 (1994), 157–171.

- [18] TRIPATHI, V., KLOKS, T., PANDEY, A., PAUL, K., AND WANG, H.-L. Complexity of paired domination in at-free and planar graphs. In *Conference on Algorithms and Discrete Applied Mathematics* (2022), Springer, pp. 65–77.
- [19] YANG, X., AND WU, B. $[1, 2]$ -domination in graphs. *Discrete Applied Mathematics* 175 (2014), 79–86.