# Multipartite entanglement in the diagonal symmetric subspace

Jordi Romero-Pallejà, <sup>1</sup> Jennifer Ahiable, <sup>1</sup> Alessandro Romancino, <sup>2</sup> Carlo Marconi, <sup>3</sup> and Anna Sanpera<sup>1,4</sup>

<sup>1</sup>Física Teòrica: Informació i Fenòmens Quàntics. Departament de Física,
 Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain
 <sup>2</sup>Dipartimento di Fisica e Chimica "E. Segrè", Università degli Studi di Palermo, Via Archirafi 36, 90123 Palermo, Italy
 <sup>3</sup>Istituto Nazionale di Ottica (CNR-INO), Largo Enrico Fermi 6, 50125 Firenze, Italy
 <sup>4</sup>ICREA, Pg. Lluís Companys 23, 08010 Barcelona, Spain
 (Dated: March 11, 2024)

We investigate the entanglement properties in the symmetric subspace of N-partite d-dimensional systems (qudits). For diagonal symmetric states, we show that there is no bound entanglement for d=3,4 and N=3. Further, we present a constructive algorithm to map multipartite diagonal symmetric states of qudits onto bipartite symmetric states of larger local dimension. This technique greatly simplifies the analysis of multipartite states and allows to infer entanglement properties for any even  $N \ge 4$  due to the fact that the PPT conditions that arise from the bipartite symmetric states correspond to the same PPT conditions that appear in the multipartite diagonal symmetric states.

### I. INTRODUCTION

Symmetry and entanglement emerge as foundational concepts that profoundly shape our understanding of Nature. While symmetry reflects the invariance of physical systems under transformations, entanglement defines the role of genuine quantum correlations in composite systems. Here we unify these notions to reveal the role of symmetry in reducing the complexity of entanglement characterisation.

Over the years, entanglement has been exploited as a resource for many information tasks which would, otherwise, be unachievable (see, e.g., [1] and references therein, [2–4]). Consequently, the ability to access this resource in quantum states is a dire need for both theorists and experimentalists. Significant advances have been made in the characterisation and quantification of entanglement, despite the considerable challenges it presents. In fact, the separability problem, that is, determining whether a given quantum state is separable or not, is recognised as a difficult task, falling, in general, under the category of NP-hard problems [5]. Nevertheless, several methods have been developed to detect entanglement in specific quantum states, particularly in the case of bipartite systems. So far, the most powerful criterion for separability is based on the positivity of the partial transpose (PPT) of a quantum state [6]. The so called PPT-criterion becomes necessary and sufficient when the composite Hilbert space,  $\mathcal{H}_{AB}$ , has dimension dim( $\mathcal{H}_{AB}$ )  $\leq$  6 [7]. While states with a negative partial transposition are always entangled, the converse is not true: such states are known as PPT-entangled states (PPTES). A complete description of PPTES is still missing and further research is needed to uncover their structure, especially when dealing with multipartite systems, whose larger Hilbert space inevitably hinders the possibility of a simple characterisation.

Symmetric states, particularly in the bipartite scenario, have been previously addressed in [8–10]. In this work, we investigate multipartite diagonal symmetric states (DS), which correspond to mixtures of projectors onto symmetric pure states. While for bipartite DS states it is known under which conditions being PPT is necessary and sufficient to ensure separability [11–13], this is not the case in the multipartite sce-

nario. Here, we turn our attention to the latter, i.e.,  $\rho \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes N}))$ , where  $\mathcal{S}$  denotes the symmetric subspace of N parties of local dimension d.

Before proceeding further, let us present our main findings: first, we demonstrate that for DS states with N=3, PPT ensures separability for d=3,4, while for d>4, it is no longer the case. Second, we analyse multipartite DS states for a generic number of parties, N, and disclose the structure of their partial transpositions. Further, for N=2n (n>1), we map DS states in  $\mathcal{S}((\mathbb{C}^d)^{\otimes N})$  to bipartite symmetric states in  $\mathcal{S}(\mathbb{C}^m\otimes\mathbb{C}^m)$ , with m>d. With the aid of this technique, we infer the entanglement properties of DS states from their corresponding bipartite symmetric states, thus providing new insights on the structure of entanglement in multipartite systems.

This paper is organised as follows: in Section II, we introduce the main concepts and tools needed to address our problem. There, we also briefly review previous works on which our ideas are built. In Section III, we focus on entanglement in the multipartite scenario for arbitrary dimension. We discuss the consequences that emerge from imposing PPT conditions for N=3. Then, we move to the generic case of N parties and demonstrate that imposing PPT leads to a set of conditions that do not allow to ensure separability. In Section IV. we present an embedding that allows a significant simplification of the separability problem and illustrate our findings with an example. Remarkably, in Section V, we introduce a similar technique, based on DS states of qubits, which allows to the study symmetric bipartite states. Finally, we conclude summarising our results and presenting some open questions in the Conclusions.

## II. MATHEMATICAL PRELIMINARIES

## A. The symmetric subspace

Let  $\mathcal{H} = \mathcal{H}_1 \otimes \cdots \otimes \mathcal{H}_N$  be a finite dimensional N-partite Hilbert space, where  $\mathcal{H}_i = \mathbb{C}^d$  is the Hilbert space associated to the i-th party. The symmetric subspace,  $\mathcal{S}(\mathcal{H}) \subset \mathcal{H}$ , corre-

sponds to the convex set formed by the normalized pure states,  $|\Psi_S\rangle \in \mathcal{H}$ , that remain invariant under any permutation of the parties. In an abuse of language, we denote by symmetric quantum states,  $\rho_S \in \mathcal{B}(\mathcal{S}(\mathcal{H}))$ , the convex hull of projectors onto pure symmetric normalised states, i.e.,

$$\rho_S = \sum_k p_S^{(k)} |\Psi_S^{(k)}\rangle \langle \Psi_S^{(k)}| , \qquad (1)$$

with  $p_S^{(k)} \geq 0$ ,  $\sum_k p_S^{(k)} = 1$ . Thus, any  $\rho_S \in \mathcal{B}(\mathcal{S}(\mathcal{H}))$  is a positive semidefinite operator  $(\rho_S \succeq 0)$  with unit trace  $(\text{Tr}(\rho_S) = 1)$ , fulfilling the condition  $\Pi_S \rho_S \Pi_S = \Pi_S \rho_S = \rho_S \Pi_S = \rho_S$ , where  $\Pi_S$  is the projector onto the symmetric subspace. The dimension of this subspace is  $\binom{N+d-1}{d-1}$ , which is significantly smaller than the dimension of the Hilbert space  $\mathcal{H}$ , i.e.,  $\dim(\mathcal{H}) = d^N$ . Let us recall the corresponding definition of a separable state:

**Definition II.1.** Let  $\mathcal{H} = (\mathbb{C}^d)^{\otimes N}$  be a multipartite Hilbert space. A symmetric state  $\rho_S \in \mathcal{B}(\mathcal{S}(\mathcal{H}))$  is separable if it can be written as a convex combination of projectors onto symmetric product states, i.e.,

$$\rho_S = \sum_i p_i |e_i e_i \cdots e_i\rangle \langle e_i e_i \cdots e_i|. \tag{2}$$

A convenient basis for the symmetric subspace is given by the Dicke states which correspond to equally weighted superpositions of states with the same number of excitations. Such states are defined as follows:

**Definition II.2.** The Dicke state  $|D_{\mathbf{k}}\rangle$  corresponds to superpositions of  $k_0$  qudits in the state  $|0\rangle$ ,  $k_1$  qudits in the state  $|1\rangle$ , etc, and is expressed as

$$|D_{\mathbf{k}}\rangle = \mathcal{C}(N, \mathbf{k})^{-1/2} \sum_{\pi \in \mathcal{G}_N} \pi(|0\rangle^{\otimes k_0} \otimes \cdots \otimes |d-1\rangle^{\otimes k_{d-1}}), (3)$$

where  $G_N$  is the group of permutations of N elements,  $\pi$  is a permutation operator,  $\mathbf{k} = (k_0, k_1, \dots, k_{d-1})$  is a partition of N, i.e.,  $k_i \geq 0$ ,  $\sum_{i=0}^{d-1} k_i = N$ , and  $C(N, \mathbf{k})$  is a normalisation constant given by

$$C(N, \mathbf{k}) = \binom{N}{\mathbf{k}} = \frac{N!}{k_0! k_1! \cdots k_{d-1}!}.$$
 (4)

Thus, a symmetric quantum state can be compactly expressed in the Dicke basis as follows:

**Definition II.3.** Any symmetric state,  $\rho_S \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes N}))$ , can be written as

$$\rho_{S} = \sum_{\mathbf{k}\mathbf{k}'} \alpha_{\mathbf{k}}^{\mathbf{k}'} |D_{\mathbf{k}}\rangle\langle D_{\mathbf{k}'}| + \text{h.c.}$$
 (5)

with  $\alpha_{\mathbf{k}}^{\mathbf{k}'} \in \mathbb{C}$ , where  $\mathbf{k}, \mathbf{k}'$  are partitions of N.

Convex mixtures of projectors onto Dicke states are denoted as diagonal symmetric (DS), since they are diagonal in the Dicke basis. They form a convex subset of  $\mathcal S$  and are particularly relevant for our analysis.

**Definition II.4.** Any DS state,  $\rho_{DS} \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes N}))$ , is of the form

$$\rho_{DS} = \sum_{\mathbf{k}} p_{\mathbf{k}} |D_{\mathbf{k}}\rangle\langle D_{\mathbf{k}}|, \tag{6}$$

where **k** is a partition of N and  $p_k \ge 0$ ,  $\forall$  **k**,  $\sum_{\mathbf{k}} p_{\mathbf{k}} = 1$ .

Although in the following we will focus mainly on the multipartite scenario, we briefly recall some basic definitions regarding the symmetric bipartite space of two qudits, i.e.,  $\mathcal{S}(\mathbb{C}^d \otimes \mathbb{C}^d)$ . In this case, the Dicke states take the simpler expression,

$$|D_{ii}^{(d)}\rangle = |ii\rangle , \quad |D_{ij}^{(d)}\rangle = \frac{|ij\rangle + |ji\rangle}{\sqrt{2}} , \quad i \neq j ,$$
 (7)

where the superscript reminds that  $\{|i\rangle\}_{i=0}^{d-1}$  is an orthonormal basis of  $\mathbb{C}^d$ . Notice that the dimension of  $\mathcal{S}(\mathbb{C}^d \otimes \mathbb{C}^d)$  is now d(d+1)/2.

Equivalently, bipartite symmetric quantum states can be compactly expressed as follows:

**Definition II.5.** Any bipartite symmetric state,  $\rho_S \in \mathcal{B}(\mathcal{S}(\mathbb{C}^d \otimes \mathbb{C}^d))$ , can be written as

$$\rho_{S} = \sum_{\substack{0 \le i \le j < d \\ 0 \le k \le l < d}} \left( \rho_{ij}^{kl} | D_{ij}^{(d)} \rangle \langle D_{kl}^{(d)} | + \text{h.c.} \right) , \tag{8}$$

with  $\rho_{ij}^{kl} \in \mathbb{C}$ . Notice that, due to the symmetry of the Dicke states, it holds that  $\rho_{ij}^{kl} = \rho_{ii}^{kl} = \rho_{ii}^{lk} = \rho_{ii}^{lk} \forall i, j, k, l$ .

**Definition II.6.** Any DS state,  $\rho_{DS} \in \mathcal{B}(\mathcal{S}(\mathbb{C}^d \otimes \mathbb{C}^d))$ , is of the form

$$\rho_{DS} = \sum_{0 \le i \le j \le d} p_{ij} |D_{ij}^{(d)}\rangle\langle D_{ij}^{(d)}|, \tag{9}$$

with  $p_{ij} \geq 0$ ,  $\forall i, j \text{ and } \sum_{i,j} p_{i,j} = 1$ .

Due to their intrinsic structure, symmetric states possess a natural decomposition:

**Definition II.7.** Every symmetric state,  $\rho_S$ , can be decomposed as  $\rho_S = \rho_{DS} + \sigma_{CS}$ , where  $\rho_{DS}$  is a DS state and  $\sigma_{CS}$  is a traceless symmetric contribution which contains the coherences between Dicke states. In the bipartite case, this decomposition reads

$$\rho_{S} = \rho_{DS} + \sigma_{CS} \tag{10}$$

$$= \sum_{0 \le i \le j < d} p_{ij} |D_{ij}^{(d)}\rangle\langle D_{ij}^{(d)}| + \sum_{(i,j) \ne (k,l)} \left(\alpha_{ij}^{kl} |D_{ij}^{(d)}\rangle\langle D_{kl}^{(d)}| + \text{h.c.}\right),$$
with  $\alpha_{ii}^{kl} \in \mathbb{C}$  and  $\alpha_{ii}^{kl} = (\alpha_{bl}^{ij})^{*}$ .

## B. Entanglement in bipartite DS states

Previous works have shown that for bipartite DS states of low dimension (i.e., d = 3,4), separability is equivalent to the PPT criterion [12, 13]. Let us begin by observing that, for such states, the partial transpose has a special structure:

**Definition II.8.** The partial transpose of every bipartite DS state has the form

$$\rho_{DS}^{\Gamma} = M_d(\rho_{DS}) \bigoplus_{0 \le i \ne j < d} \frac{p_{ij}}{2} . \tag{11}$$

Here  $M_d(\rho_{DS})$  is a  $d \times d$  matrix with non-negative entries defined as

$$M_d(\rho_{DS}) = \begin{pmatrix} \bar{p}_{00} & \bar{p}_{01} & \cdots & \bar{p}_{0d-1} \\ \bar{p}_{01} & \bar{p}_{11} & \cdots & \bar{p}_{1d-1} \\ \vdots & \vdots & \ddots & \vdots \\ \bar{p}_{0d-1} & \bar{p}_{1d-1} & \cdots & \bar{p}_{d-1d-1} \end{pmatrix}$$
(12)

where  $\bar{p}_{ij} = \frac{p_{ij}}{N_{ij}}$  and  $N_{ij}$  is the normalisation constant associated to the Dicke state  $|D_{ij}\rangle$ . Notice that, due to symmetry, it is irrelevant w.r.t. which party the partial transposition is performed.

Clearly, positivity of  $\rho_{DS}^{\Gamma}$  is related to the one of the associated matrix  $M_d(\rho_{DS})$ . Before describing the known results, the following definitions are needed.

**Definition II.9.** A  $d \times d$  matrix, M, is said completely positive if and only if there exists a non-negative  $d \times k$  matrix, C, such that  $M = CC^T$ , for some  $k \ge 1$ . This set of matrices forms a convex cone, dubbed  $\mathcal{CP}_d$ .

**Definition II.10.** A  $d \times d$  matrix, M, is said to be doubly nonnegative if and only if it is positive semidefinite and its entries are non-negative. This set of matrices forms a convex cone, dubbed  $\mathcal{DNN}_d$ .

With the aid of the above definitions, one has the following Theorems:

**Theorem II.1** ([12, 13]). Let  $\rho_{DS} \in \mathcal{B}(\mathcal{S}(\mathbb{C}^d \otimes \mathbb{C}^d))$  be a DS state. Then,  $\rho_{DS}$  separable  $\iff M_d(\rho_{DS}) \in \mathcal{CP}_d$ .

**Theorem II.2** ([13]). Let  $\rho_{DS} \in \mathcal{B}(\mathcal{S}(\mathbb{C}^d \otimes \mathbb{C}^d))$  be a DS state. Then,  $\rho_{DS}^{\Gamma} \geq 0 \iff M_d(\rho_{DS}) \in \mathcal{DNN}_d$ .

Interestingly, the convex cones  $\mathcal{CP}_d$  and  $\mathcal{DNN}_d$  are related by the following Lemma:

**Lemma II.1** ([14]). The cone of doubly non-negative matrices and the cone of completely positive matrices coincide for  $d \le 4$ , i.e.,  $\mathcal{CP}_d = \mathcal{DNN}_d$ .

Hence, separability for bipartite DS states can be cast as:

**Theorem II.3** ([12, 13]). Let  $\rho_{DS} \in \mathcal{B}(\mathcal{S}(\mathbb{C}^d \otimes \mathbb{C}^d))$  be a DS state with  $d \leq 4$ . Then,

$$\rho_{DS}$$
 separable  $\iff \rho_{DS}$  PPT. (13)

Explicit examples of PPT-entangled DS states for  $d \ge 5$  have been provided in [13], along with the entanglement witnesses that are able to detect them [15].

### III. ENTANGLEMENT IN MULTIPARTITE DIAGONAL SYMMETRIC STATES

# **A.** Entanglement in *N*-partite **DS** states for N = 3

Let us start by showing that, when N=3, the results obtained for DS bipartite systems still hold true for any dimension d, i.e., for all  $\rho_{DS} \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes 3}))$ . To do so, we analyse the structure of the corresponding partial transpositions  $\rho_{DS}^{\Gamma_i}$ . Notice that, due to the symmetry, it is enough to examine only the partial transpositon w.r.t. a given partition, e.g., i=1, as all of them are identical.

**Lemma III.1.** Let  $\rho_{DS} \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes 3}))$ . Then, its partial transposition w.r.t any party leads to a  $d^2 \times d^2$  matrix  $M^{(3)}(\rho_{DS})$  of the form:

$$M^{(3)}(\rho_{DS}) = \bigoplus_{i=0}^{d-1} M_i^{(3)}(\rho_{DS}) , \qquad (14)$$

$$M_{i}^{(3)}(\rho_{DS}) = \begin{pmatrix} \bar{p}_{i,0,0} & \bar{p}_{i,0,1} & \cdots & \bar{p}_{i,0,d-1} \\ \bar{p}_{i,0,1} & \bar{p}_{i,1,1} & \cdots & \bar{p}_{i,1,d-1} \\ \vdots & \vdots & \ddots & \vdots \\ \bar{p}_{i,0,d-1} & \bar{p}_{i,1,d-1} & \cdots & \bar{p}_{i,d-1,d-1} \end{pmatrix}$$
(15)

with  $\bar{p}_{ijk} = \frac{p_{ijk}}{N_{ijk}}$  and  $N_{ijk}$  being the normalisation factor of the corresponding three-partite Dicke state, i.e.,

$$N_{ijk} = \begin{cases} 1 & for \quad i = j = k \\ 3 & for \quad i = j \neq k \\ 6 & for \quad i \neq j \neq k \end{cases}.$$

Since the partial transposition of a symmetric state does not belong, in general, to the symmetric subspace,  $\rho_{DS}^{\Gamma_i}$  has rank  $d^3$ . Nevertheless, DS states are endowed with an extra symmetry, which allows to strongly reduce the full rank of  $\rho_{DS}^{\Gamma_i}$  to an "effective rank" of  $d^2$ , corresponding to a matrix  $d^2 \times d^2$ . This can be seen after properly deleting the repeated rows and columns in the matrices and taking out all the  $1 \times 1$  block matrices corresponding to the coefficients  $p_{ijk} > 0$ . Moreover, as  $M^{(3)}(\rho_{DS})$  in Eq.(14) is a direct sum of d block matrices of size  $d \times d$ , the entanglement properties of  $\rho_{DS}$  are uniquely determined by the latter set of matrices. For such reason, we can rely on the results presented in Section II to derive the following theorems:

**Theorem III.1.** Let  $\rho_{DS} \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes 3}))$  be a DS with associated matrix  $M^{(3)}(\rho_{DS})$ . Then,

$$\rho_{DS} \text{ separable } \iff M^{(3)}(\rho_{DS}) \in \mathcal{CP}_{d^2}.$$

**Theorem III.2.** Let  $\rho_{DS} \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes 3}))$  be a DS with associated matrix  $M^{(3)}(\rho_{DS})$ . Then,

$$\rho_{DS} PPT \iff M^{(3)}(\rho_{DS}) \in \mathcal{DNN}_{d^2}$$
.

Making use of Lemma II.1 and taking into account the direct sum structure of Eq.(14), directly leads to:

**Theorem III.3.** Let  $\rho_{DS} \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes 3}))$  with  $d \leq 4$ . Then,  $\rho_{DS}$  separable  $\iff \rho_{DS}$  PPT.

An immediate consequence of Theorem III.3 is that there do not exist PPT-entangled DS states of three parties for  $d \leq 4$ . However, it is worth noting that, for d > 4, as it happens in the bipartite case, PPT is only necessary, but not sufficient, for separability (see Lemma II.1). In fact, when d > 4,  $M_i^{(d)}(\rho_{DS}) \in \mathcal{DNN}_d$  no longer implies that  $M_i^{(d)}(\rho_{DS}) \in \mathcal{CP}_d$ , strongly suggesting that there exist PPT entangled states of three parties for any dimension d > 4.

# **B.** Entanglement in *N*-partite **DS** states for $N \ge 4$

We now move to the case N=4. First, we prove that, although it is now necessary to consider PPT with respect to two partitions, namely 2:2 and 1:3, in the case of DS states, positivity w.r.t. the largest partition implies positivity w.r.t. the smaller. Although the separability problem for N=4 can be cast in a similar way to the case N=3, the dimension of the resulting matrices  $M_i^{(4)}$  prevents from ensuring that PPT implies separability for any d. Proving this last statement for an arbitrary number N of parties is not trivial. However, we provide strong evidence that this is the case.

**Lemma III.2.** Let  $\rho_{DS} \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes 4}))$ . Then, its partial transposition w.r.t the largest partition, i.e.,  $\rho_{DS}^{\Gamma_{2,2}}$ , leads to a  $d^4 \times d^4$  matrix  $M^{(4)}(\rho_{DS})$  of the form:

$$M^{(4)}(\rho_{DS}) = \begin{pmatrix} \bar{p}_{r_0,r_0} & \bar{p}_{r_0,r_1} & \dots & \bar{p}_{r_0,r_s} \\ \bar{p}_{r_0,r_1} & \bar{p}_{r_1,r_1} & \dots & \bar{p}_{r_1,r_s} \\ \vdots & \vdots & \ddots & \vdots \\ \bar{p}_{r_0,r_s} & \bar{p}_{r_1,r_s} & \dots & \bar{p}_{r_s,r_s} \end{pmatrix}$$

$$\stackrel{d-1}{\bigoplus_{\substack{i < j \\ i = 0}}} \begin{pmatrix} \bar{p}_{ij00} & \bar{p}_{ij01} & \dots & \bar{p}_{ij0d-1} \\ \bar{p}_{ij01} & \bar{p}_{ij11} & \dots & \bar{p}_{ij1d-1} \\ \vdots & \dots & \ddots & \vdots \\ \bar{p}_{ii0d-1} & \bar{p}_{ii1d-1} & \dots & \bar{p}_{iid-1d-1} \end{pmatrix},$$

$$(16)$$

where  $r_i$  denotes the i-th element of a list of dimension s = d(d+1)/2 corresponding to the ordered indices of the two-qudit Dicke states  $|D_{ij}\rangle$ .

As an example, when d=3, it is s=6 and  $r_i \in \{00,01,02,11,12,22\}$ . From the structure displayed in Eq.(16) stems the following observation:

**Lemma III.3.** Let  $\rho_{DS} \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes 4}))$ . Then,  $\rho_{DS}^{\Gamma_{2:2}} \succeq 0 \Longrightarrow \rho_{DS} PPT$ .

*Proof.* The lemma can be proved by looking at the general structure of the partial transposition with respect to any non-largest partition. When N=4 one simply has partition 1|3, whose corresponding partial transposed matrix,  $\rho_{DS}^{\Gamma_{1:3}}$ , takes the form:

$$\rho_{DS}^{\Gamma_{1:3}} = \bigoplus_{\substack{i \leq j \\ i,j=0}}^{d-1} \begin{pmatrix} \bar{p}_{ij00} & \bar{p}_{ij01} & \cdots & \bar{p}_{ij0d-1} \\ \bar{p}_{ij01} & \bar{p}_{ij11} & \cdots & \bar{p}_{ij1d-1} \\ \vdots & \ddots & \ddots & \vdots \\ \bar{p}_{ij0d-1} & \bar{p}_{ij1d-1} & \cdots & \bar{p}_{ijd-1d-1} \end{pmatrix}$$
(17)

Comparing Eq.(16) to Eq.(17), it can be observed that, choosing i < j, the matrices that arise in Eq.(17) are already included in  $M^{(4)}(\rho_{DS})$ . Taking i = j, the related matrices correspond to principal minors of the first matrix of Eq.(16), thus concluding the proof.

We are now ready to generalise the form of the matrix arising from the partial transpose with respect to the largest partition, to an arbitrary number of parties N:

**Lemma III.4.** Let  $\rho_{DS} \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes N}))$  be a DS state with N even. Then, its partial transpose w.r.t. the largest partition, i.e.,  $\rho_{DS}^{\Gamma_{N/2:N/2}}$ , leads to the following associated matrix  $M^{(N)}(\rho_{DS})$ :

$$M^{(N)}(\rho_{DS}) = \bigoplus_{i=0}^{N/2-1} \left( \bigoplus_{k=1}^{k=t} M_k^i \right) , \qquad (18)$$

$$M_{k}^{i} = \begin{pmatrix} \bar{p}_{r_{k},s_{0},s_{0}} & \bar{p}_{r_{k},s_{0},s_{1}} & \cdots & \bar{p}_{r_{k},s_{0},s_{l}} \\ \bar{p}_{r_{k},s_{0},s_{1}} & \bar{p}_{r_{k},s_{1},s_{1}} & \cdots & \bar{p}_{r_{k},s_{1},s_{l}} \\ \vdots & \vdots & \ddots & \vdots \\ \bar{p}_{r_{k},s_{0},s_{l}} & \bar{p}_{r_{k},s_{1},s_{l}} & \cdots & \bar{p}_{r_{k},s_{l},s_{l}} \end{pmatrix}$$
(19)

where  $M_k^i$  are square matrices of dimension  $\binom{\frac{N}{2}-i+d-1}{d-1}$ ,  $r_k, s_k$  denote the k-th element of the ordered lists containing the indices of the Dicke states that span  $\mathcal{S}((\mathbb{C}^d)^{\otimes 2i})$  with i equal indices (t different elements) and  $\mathcal{S}((\mathbb{C}^d)^{\otimes \frac{N}{2}-i})$ ), respectively. Here,  $s_l$  denotes the last element of the list.

**Lemma III.5.** Let  $\rho_{DS} \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes N}))$  be a DS state with N odd. Then, its partial transpose w.r.t. the largest partition, i.e.,  $\rho_{DS}^{\Gamma_{N-1}:\frac{N+1}{2}}$ , leads to the following associated matrix  $M^{(N)}(\rho_{DS})$ :

$$M^{(N)}(\rho_{DS}) = \bigoplus_{i=0}^{\frac{N-1}{2}-1} \left( \bigoplus_{k=1}^{k=t} \left( \bigoplus_{j=0}^{j=d-1} M_{jk}^i \right) \right) , \qquad (20)$$

$$M_{jk}^{i} = \begin{pmatrix} \bar{p}_{j,r_{k},s_{0},s_{0}} & \bar{p}_{j,r_{k},s_{0},s_{1}} & \cdots & \bar{p}_{j,r_{k},s_{0},s_{l}} \\ \bar{p}_{j,r_{k},s_{0},s_{1}} & \bar{p}_{j,r_{k},s_{1},s_{1}} & \cdots & \bar{p}_{j,r_{k},s_{1},s_{l}} \\ \vdots & \vdots & \ddots & \vdots \\ \bar{p}_{j,r_{k},s_{0},s_{l}} & \bar{p}_{j,r_{k},s_{1},s_{l}} & \cdots & \bar{p}_{j,r_{k},s_{l},s_{l}} \end{pmatrix}$$
(21)

where  $M^i_{jk}$  are square matrices of dimension  $\binom{N-1}{2}-i+d-1$ ,  $r_k, s_k$  denote the k-th element of the ordered lists containing the indices of the Dicke states that span  $\mathcal{S}((\mathbb{C}^d)^{\otimes 2i})$  with i equal indices (t different elements) and  $\mathcal{S}((\mathbb{C}^d)^{\otimes N-1}-i)$ , respectively. Here,  $s_l$  denotes the last element of the list.

Given the explicit structure of Eqs.(18)-(21), consisting of nested direct sums of matrices  $M^i$ , we reach the following conjecture:

**Conjecture III.1.** Let  $\rho_{DS} \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes N}))$  be an N-partite DS state. Then, PPT w.r.t. to its largest partition ensures PPT w.r.t any other partition, i.e.,

$$\rho_{DS}^{\Gamma_{N/2:N/2}} \succeq 0 \implies \rho_{DS} \ PPT \ for \ even \ N$$

$$\rho_{DS}^{\Gamma_{N-1}: \frac{N+1}{2}} \succeq 0 \implies \rho_{DS} \ PPT \ for \ odd \ N.$$

While it is not possible to explicitly prove that, in the generic case, PPT w.r.t. the largest partition also implies PPT w.r.t. any other partition, we provide strong analytical evidence that suggests that the above conjecture holds true.

**Theorem III.4.** Let  $\rho_{DS} \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes 4}))$  be a DS state with an associated matrix  $M^{(4)}(\rho_{DS})$  which arises from the partial transpose w.r.t the largest partition. Then,

$$\rho_{DS} \text{ separable} \iff M^{(4)}(\rho_{DS}) \in \mathcal{CP}$$
.

**Theorem III.5.** Let  $\rho_{DS} \in \mathcal{B}(\mathcal{S}((\mathbb{C}^d)^{\otimes 4}))$  be a DS state with an associated matrix  $M^{(4)}(\rho_{DS})$  which arises from the partial transpose w.r.t the largest partition. Then,

$$\rho_{DS} PPT \iff M^{(4)}(\rho_{DS}) \in \mathcal{DNN}$$
.

We conjecture that Theorems III.4-III.5 hold true for any N. This conjecture is further supported in the N even case by the embedding we present in the following section. Their proof is a straightforward generalisation of the bipartite case, as we have seen for N=3. However, since already for N=4 the associated matrices have size  $m \times m$ , with m>4, Lemma II.1 does not ensure that PPT is necessary and sufficient for separability, not even in the low-dimensional case, i.e., d=3,4.

# IV. MAPPING MULTIPARTITE DS-STATES TO BIPARTITE SYMMETRIC STATES

We now present a method to cast a multipartite DS state of N qudits, with N even, as a bipartite symmetric state of higher local dimension. In particular, a DS state acting on  $\mathcal{S}\left((\mathbb{C}^d)^{\otimes N}\right)$  is mapped to a symmetric state acting on  $\mathcal{S}\left(\mathcal{S}((\mathbb{C}^d)^{\otimes N/2})\otimes\mathcal{S}((\mathbb{C}^d)^{\otimes N/2})\right)$ . This mapping is based on the following steps:

- 1. Any Dicke states of *N* qudits can be expressed as a linear combination of tensor products of two Dicke states of *N*/2 qudits;
- 2. Since  $S((\mathbb{C}^d)^{\otimes N/2}) \cong \mathbb{C}^k$ , with  $k = \binom{N/2+d-1}{d-1}$ , each Dicke state of N/2 qudits is associated to a basis vector in  $\mathbb{C}^k$ ;
- 3. Any Dicke state of *N* qudits can now be expressed as a linear combination of *bipartite* Dicke states of local dimension *k*.
- 4. Hence, the initial state  $\rho_{DS}$  is transformed as  $\rho_{DS} \Longrightarrow \rho_S \in \mathcal{B}(\mathcal{S}(\mathbb{C}^k \otimes \mathbb{C}^k))$ .

It is worth stressing that this correspondence does not define an isomorphism between the two subspaces, since the first has dimension  $\binom{N+d-1}{d-1}$ , while the latter has dimension  $k(k+1)/2 = \binom{N/2+d-1}{d-1} \binom{N/2+d-1}{d-1} + 1/2$ , i.e., significantly greater. As a consequence, the resulting state through the mapping is generically a symmetric, although not DS, state.




Figure 1. Pictorial representation of the embedding of a multipartite DS state  $\rho_{DS}$  into a bipartite symmetric state  $\rho_S$  of systems with higher local dimension, i.e., k > d.

One may object that it might be more convenient to map a multipartite DS state into another symmetric state by means of an isomorphism, rather than considering an embedding into a higher dimensional subspace. However, it is necessary to take into account both the state and its partial transpositions with respect to different partitions. Let us clarify this point with an example. Since dim  $\mathcal{S}((\mathbb{C}^3)^{\otimes 4}) =$  $15 = \dim \mathcal{S}(\mathbb{C}^5 \otimes \mathbb{C}^5)$ , there exists an isomorphism between the two subspaces. Hence, one might be tempted to associate a state  $\rho_{DS} \in \mathcal{S}\left((\mathbb{C}^3)^{\otimes 4}\right)$  to a bipartite DS state  $\tilde{\rho}_{DS} \in$  $\mathcal{S}\left(\mathbb{C}^5\otimes\mathbb{C}^5\right)$ . However, since  $\mathrm{rank}(
ho_{DS}^{\Gamma_{2:2}})>\mathrm{rank}(\tilde{
ho}_{DS}^{\Gamma})$ , such a naive isomorphism is not able to encode the whole information of the state  $\rho_{DS}$ . Notice that this embedding is optimal in the sense that k corresponds to the smallest local dimension of the bipartite system such that  $rank(\rho_{DS}) < rank(\rho_{S})$ and  $\operatorname{rank}(\rho_{DS}^{\Gamma_{N/2:N/2}}) = \operatorname{rank}(\rho_S^{\Gamma})$ . This implies that, in order to check the conditions of  $\rho_{DS}$ , it is now sufficient to inspect the bipartite state  $\rho_S$  which results from the embedding. This is due to the fact that the PPT conditions on this bipartite symmetric state that results from the embedding, are identical to the PPT conditions arising in the multipartite DS case. This fact strongly supports that for even N, both the conjecture III.1 and the Theorems III.4-III.5 hold true. We expect the same behaviour for N odd even though this embedding cannot be applied in this case. The advantage of this approach becomes particularly evident when one compares the dimension of the partial transpositions, since  $\dim(\rho_S^\Gamma) = {N/2+d-1 \choose d-1}^2 < \dim(\rho_{DS}^{\Gamma_{N/2:N/2}}) = d^N$ , as it can be seen in Fig.2.

Let us illustrate the above embedding with an explicit example, where we map a multipartite DS state with N=4 and d=3 to a bipartite symmetric state with d=6, i.e.,  $\rho_{DS} \in \mathcal{B}\left(\mathcal{S}\left((\mathbb{C}^3)^{\otimes 4}\right)\right) \to \rho_S \in \mathcal{B}\left(\mathcal{S}\left(\mathbb{C}^6 \otimes \mathbb{C}^6\right)\right)$ . First, notice that the Dicke basis for four qutrits can be expressed as a linear combination of tensor products between Dicke states of




Figure 2. Scaling of the dimension (D) of  $(\rho_{DS})^{\Gamma_{N/2:N/2}}$  and  $\rho_S^{\Gamma}$  for an even number of parties (N). Here,  $\rho_{DS}$  is a DS state with N parties of dimension d (blue) and  $\rho_S$  is a bipartite symmetric state that results from the mapping of a DS state of local dimension  $d_E$  (orange).

two qutrits, i.e.,

$$\begin{split} |D_{iiij}^{(3)}\rangle &= |D_{ii}^{(3)}\rangle \, |D_{ii}^{(3)}\rangle \\ |D_{iiij}^{(3)}\rangle &= \frac{1}{\sqrt{2}} \left( |D_{ii}^{(3)}\rangle \, |D_{ij}^{(3)}\rangle + |D_{ij}^{(3)}\rangle \, |D_{ii}^{(3)}\rangle \right) \\ |D_{iiijj}^{(3)}\rangle &= \frac{1}{\sqrt{3}} \left( |D_{ii}^{(3)}\rangle \, |D_{jj}^{(3)}\rangle + |D_{jj}^{(3)}\rangle \, |D_{ii}^{(3)}\rangle \right) + \sqrt{\frac{2}{3}} \, |D_{ij}^{(3)}\rangle \, |D_{ij}^{(3)}\rangle \\ |D_{iijk}^{(3)}\rangle &= \frac{1}{\sqrt{6}} \left( |D_{ii}^{(3)}\rangle \, |D_{jk}^{(3)}\rangle + |D_{jk}^{(3)}\rangle \, |D_{ii}^{(3)}\rangle \right) + \\ \sqrt{\frac{1}{3}} \left( |D_{ij}^{(3)}\rangle \, |D_{ik}^{(3)}\rangle + |D_{ik}^{(3)}\rangle \, |D_{ij}^{(3)}\rangle \right) \end{split}$$

with  $i, j, k \in \{0, 1, 2\}$ ,  $i \neq j \neq k$  and where the superscript represents qutrits. Recalling that  $S(\mathbb{C}^3 \otimes \mathbb{C}^3) \cong \mathbb{C}^6$ , we have:

$$|D_{00}^{(3)}\rangle = |0\rangle, \quad |D_{01}^{(3)}\rangle = |1\rangle, \quad |D_{02}^{(3)}\rangle = |2\rangle, |D_{11}^{(3)}\rangle = |3\rangle, \quad |D_{12}^{(3)}\rangle = |4\rangle, \quad |D_{22}^{(3)}\rangle = |5\rangle.$$

As a consequence, a generic DS state in  $\mathcal{S}((\mathbb{C}^3)^{\otimes 4})$ , leads to a bipartite symmetric state  $\rho_S^{(6)} \in \mathcal{B}\left(\mathcal{S}\left(\mathbb{C}^6 \otimes \mathbb{C}^6\right)\right)$  of the form

$$\rho_S^{(6)} = \sum_{ij} p_{ij} |D_{ij}^{(6)}\rangle\langle D_{ij}^{(6)}| + \sum_{ijkl} \alpha_{ij}^{kl} |D_{ij}^{(6)}\rangle\langle D_{kl}^{(6)}| + \text{h.c.}$$
 (22)

Hence, the characterisation of PPT-entanglement is greatly simplified, since the size of the final state is now considerably smaller.

## V. MAPPING SYMMETRIC STATES OF QUDITS TO DS STATES OF QUBITS

In this Section we present a technique to recast multipartite qudit systems as DS states of qubits. Noticing that  $\mathcal{S}((\mathbb{C}^2)^{\otimes N}) \cong \mathbb{C}^{N+1}$ , one can set  $|D_i^N\rangle \equiv |i\rangle$ , where  $|D_i^N\rangle$  is the Dicke state of N qubits with i excitations, and  $|i\rangle \in \mathbb{C}^{N+1}$ .

Once the mapping has been performed, exploting the fact that PPT DS states of qubits are separable [16], we can deduce equivalent properties for the original qudit state. For the sake of clarity, in the following we restrict to bipartite states, although we stress that our approach can be generalised to any number of parties. Let us consider the symmetric space of two qutrits, i.e.,  $\mathcal{S}\left(\mathbb{C}^3\otimes\mathbb{C}^3\right)$ . Notice that  $\mathbb{C}^3\cong\mathcal{S}\left((\mathbb{C}^2)^{\otimes 2}\right)$ , since it is possible to write

$$|0\rangle = |D_0^2\rangle, \quad |1\rangle = |D_1^2\rangle, \quad |2\rangle = |D_2^2\rangle.$$
 (23)

Hence.

$$\mathcal{S}\left(\mathbb{C}^3\otimes\mathbb{C}^3\right) = \mathcal{S}\left(\mathcal{S}((\mathbb{C}^2)^{\otimes 2})\otimes\mathcal{S}((\mathbb{C}^2)^{\otimes 2})\right)\supset \mathcal{S}((\mathbb{C}^2)^{\otimes 4}))\;,$$

where the last inclusion implies that every Dicke state  $|D_i^4\rangle \in \mathcal{S}((\mathbb{C}^2)^{\otimes 4})$  can be expressed as a linear combination of the Dicke states  $|D_{ii}^{(3)}\rangle \in \mathcal{S}(\mathbb{C}^3\otimes\mathbb{C}^3)$ , i.e.,

$$|D_0^4\rangle = |D_{00}^{(3)}\rangle, \quad |D_1^4\rangle = |D_{01}^{(3)}\rangle,$$
 (24)

$$|D_2^4\rangle = \frac{1}{\sqrt{3}} |D_{02}^{(3)}\rangle + \sqrt{\frac{2}{3}} |D_{11}^{(3)}\rangle,$$
 (25)

$$|D_3^4\rangle = |D_{12}^{(3)}\rangle, \quad |D_4^4\rangle = |D_{22}^{(3)}\rangle.$$
 (26)

Inverting Eqs.(24)-(26) leads to

$$|D_{00}^{(3)}\rangle = |D_{0}^{4}\rangle, \quad |D_{01}^{(3)}\rangle = |D_{1}^{4}\rangle,$$
 (27)

$$|D_{02}^{(3)}\rangle = \sqrt{3} |D_2^4\rangle - \sqrt{2} |D_{11}^{(3)}\rangle,$$
 (28)

$$|D_{12}^{(3)}\rangle = |D_3^4\rangle, \quad |D_{22}^{(3)}\rangle = |D_4^4\rangle,$$
 (29)

from which it is evident that not every state in  $\mathcal{S}(\mathbb{C}^3 \otimes \mathbb{C}^3)$  can be mapped to a state in  $\mathcal{S}((\mathbb{C}^2)^{\otimes 4}))$ . However, imposing conditions on a state  $\rho_S \in \mathcal{S}(\mathbb{C}^3 \otimes \mathbb{C}^3)$  such that rank  $(\rho_S) = 5$ , allows to define an isomorphism to a four-qubit DS state  $\rho_{DS}^Q = \sum_k q_k |D_k^4\rangle\langle D_k^4|$ . In fact, consider the symmetric state  $\rho_S$  given by

$$\rho_S = \rho_{DS} + (\alpha_{02}^{11} | D_{11}^{(3)} \rangle \langle D_{02}^{(3)} | + \text{h.c.}).$$
 (30)

Imposing  $p_{11}=2p_{02}=\sqrt{2}\alpha_{02}^{11}$  and using Eqs.(27)-(29),  $\rho_S$  of Eq.(30) is mapped to a DS state  $\rho_{DS}^Q$  whose coefficients  $q_k$  satisfy

$$q_0 = p_{00}, \quad q_1 = p_{01},$$
 (31)

$$q_2 = 3p_{02}, \quad q_3 = p_{12}, \quad q_4 = p_{22}.$$
 (32)

As a consequence of this mapping we have the following result:

**Corollary V.1.** Let  $\rho_S \in \mathcal{S}(\mathbb{C}^3 \otimes \mathbb{C}^3)$  be a symmetric state of the form

$$\rho_S = \rho_{DS} + (\alpha_{02}^{11} | D_{11}^{(3)} \rangle \langle D_{02}^{(3)} | + \text{h.c.}), \qquad (33)$$

whose coefficients satisfy  $p_{11}=2p_{02}=\sqrt{2}\alpha_{02}^{11}$ . Then,  $\rho_S$  is separable iff it is PPT.

We conclude this section noting that the above result can be easily generalised to the case of other coherences of the same form, i.e.,  $\sigma_{CS} = \alpha_{jk}^{ii} |D_{ii}^{(3)}\rangle\langle D_{jk}^{(3)}| + \text{h.c.}$ , for  $i \neq j \neq k$ , by considering an alternative mapping.

## VI. CONCLUSIONS

We have investigated the entanglement properties of diagonal symmetric states of N-qudit systems. First, we have demonstrated that, for N = 3, PPT is necessary and sufficient for separability for d = 3,4. Second, we have considered the case  $N \ge 4$ , where the approach used is not enough to conclude whether PPT is necessary and sufficient for separability. Moreover, we have presented strong evidence that positivity w.r.t. the biggest partition ensures positivity w.r.t. any other partition. This conjecture is supported by embedding multipartite DS states into bipartite symmetric states with higher local dimension. The advantage of this approach is twofold: on the one hand, the entanglement properties of multipartite systems can be deduced from known results regarding separability of bipartite symmetric states; on the other, the dimension of the partial transpose after the embedding is greatly reduced, providing a valuable tool in the characterisation of the entanglement in the multipartite scenario. Further, we have provided some examples of separable bipartite symmetric states using a mapping onto diagonal symmetric qubits.

### VII. ACKNOWLEDGMENTS

J.R-P acknowledges financial support from Ministerio de Ciencia e Innovación of the Spanish Government

FPU22/01511. J.A. and A.S. acknowledges financial support from Ministerio de Ciencia e Innovación of the Spanish Government with funding from European Union NextGenerationEU (PRTR-C17.I1) and by Generalitat de Catalunya. C.M. acknowledges support from the European Union -NextGeneration EU, "Integrated infrastructure initiative in Photonic and Quantum Sciences" - I-PHOQS [IR0000016, ID D2B8D520, CUP B53C22001750006] . A.S. acknowledges financial support from the European Commission OuantERA grant ExTRaQT (Spanish MICIN project PCI2022-132965), by the Spanish MICIN (project PID2022-141283NB-I00) with the support of FEDER funds, and by the Ministry for Digital Transformation and of Civil Service of the Spanish Government through the QUANTUM ENIA project call -Quantum Spain project, and by the European Union through the Recovery, Transformation and Resilience Plan - NextGeneration EU within the framework of the Digital Spain 2026 Agenda.

- [1] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, "Quantum entanglement," *Reviews of modern physics*, vol. 81, no. 2, p. 865, 2009.
- [2] R. Jozsa and N. Linden, "On the role of entanglement in quantum-computational speed-up," *Proceedings of the Royal Society of London. Series A: Mathematical, Physical and En*gineering Sciences, vol. 459, no. 2036, pp. 2011–2032, 2003.
- [3] J. Yin, Y. Cao, Y.-H. Li, J.-G. Ren, S.-K. Liao, L. Zhang, W.-Q. Cai, W.-Y. Liu, B. Li, H. Dai, M. Li, Y.-M. Huang, L. Deng, L. Li, Q. Zhang, N.-L. Liu, Y.-A. Chen, C.-Y. Lu, R. Shu, C.-Z. Peng, J.-Y. Wang, and J.-W. Pan, "Satellite-to-ground entanglement-based quantum key distribution," *Phys. Rev. Lett.*, vol. 119, p. 200501, Nov 2017.
- [4] M. Epping, H. Kampermann, D. Bruß, et al., "Multi-partite entanglement can speed up quantum key distribution in networks," New Journal of Physics, vol. 19, no. 9, p. 093012, 2017.
- [5] L. Gurvits, "Classical deterministic complexity of edmonds' problem and quantum entanglement," in *Proceedings of the* thirty-fifth annual ACM symposium on Theory of computing, pp. 10–19, 2003.
- [6] A. Peres, "Separability criterion for density matrices," *Physical Review Letters*, vol. 77, no. 8, p. 1413, 1996.
- [7] M. Horodecki, P. Horodecki, and R. Horodecki, "Separability of mixed states: necessary and sufficient conditions," *Physics Letters A*, vol. 223, pp. 1–8, nov 1996.
- [8] G. Tóth and O. Gühne, "Entanglement and permutational symmetry," *Physical review letters*, vol. 102, no. 17, p. 170503, 2009.
- [9] L. Chen, D. Chu, L. Qian, and Y. Shen, "Separability of completely symmetric states in a multipartite system," *Phys. Rev. A*,

- vol. 99, p. 032312, Mar 2019.
- [10] K.-C. Ha, "Comment on "separability of completely symmetric states in a multipartite system"," *Phys. Rev. A*, vol. 104, p. 016401, Jul 2021.
- [11] E. Wolfe and S. Yelin, "Certifying separability in symmetric mixed states of n qubits, and superradiance," *Physical review letters*, vol. 112, no. 14, p. 140402, 2014.
- [12] N. Yu, "Separability of a mixture of dicke states," Phys. Rev. A, vol. 94, p. 060101, 12 2016.
- [13] J. Tura, A. Aloy, R. Quesada, M. Lewenstein, and A. Sanpera, "Separability of diagonal symmetric states: a quadratic conic optimization problem," *Quantum*, vol. 2, p. 45, 2018.
- [14] P. H. Diananda, "On non-negative forms in real variables some or all of which are non-negative," in *Mathematical Proceed*ings of the Cambridge Philosophical Society, vol. 58, pp. 17– 25, Cambridge University Press, 1962.
- [15] C. Marconi, A. Aloy, J. Tura, and A. Sanpera, "Entangled symmetric states and copositive matrices," *Quantum*, vol. 5, p. 561, Oct. 2021.
- [16] R. Quesada, S. Rana, and A. Sanpera, "Entanglement and non-locality in diagonal symmetric states of n qubits," *Physical Review A*, vol. 95, no. 4, p. 042128, 2017.