

A discrete formulation for three-dimensional winding number

Ken Shiozaki¹

¹Center for Gravitational Physics and Quantum Information,
Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

(Dated: March 13, 2024)

For a smooth map $g: X \rightarrow U(N)$, where X is a three-dimensional, oriented, and closed manifold, the winding number or the map's degree is defined by $W_3 = \frac{1}{24\pi^2} \int_X \text{Tr} [(g^{-1}dg)^3]$. We introduce a method to compute W_3 using a discrete approximation of X so that the result is manifestly quantized.

Introduction— Consider a three-dimensional closed and oriented manifold X . For a smooth map $g: X \rightarrow U(N)$, with $U(N)$ representing the group of $N \times N$ unitary matrices, the winding number, an integer value, is defined by the following expression:

$$W_3[g] = \frac{1}{2\pi} \int_X H \in \mathbb{Z}, \quad (1)$$

$$H = \frac{1}{12\pi} \text{Tr} [(g^{-1}dg)^3]. \quad (2)$$

The winding number $W_3[g]$ is pivotal in various branches of physics, including topological band theory, where it acts as the topological invariant for three-dimensional superconductors with time-reversal symmetry [1, 2], and in non-Abelian (lattice) gauge theory, appears in instanton number calculations [3, 4]. Often in these applications, the function g is defined only on a finite set of lattice points for numerical analysis. Therefore, an efficient numerical formulation with lattice approximation of manifolds is an important issue.

For the first Chern number ch_1 of a line bundle with

connection, a sort of two-dimensional counterpart of the winding number, a well-established discrete formulation with evident quantization exists [5, 6]. Furthermore, discrete line bundles over finite simplicial complexes have been explored, especially concerning applications in computer graphics [7]. This paper develops a method for evaluating $W_3[g]$ via a discretized approximation of X , ensuring the result remains manifestly quantized, provided the approximation of X is sufficiently refined relative to the spatial variation's length scale.

Formulation— Given that H is a closed three-form, it is locally exact, meaning that for a local patch, there exists a two-form B such that $H = dB$. To construct B explicitly, we introduce a gap condition for elements of $U(N)$. A matrix $g \in U(N)$ exhibits a θ -gap if none of its eigenvalues are $e^{i\theta}$ for a given real number $\theta \in [0, 2\pi)$ [8], as illustrated in Fig. 1 (a). Furthermore, we define $\log_\theta z$ for nonzero complex number $z \in \mathbb{C}^\times$ as

$$\log_\theta z = \log |z| + i \arg z, \quad \theta \leq \arg z < \theta + 2\pi. \quad (3)$$

For two distinct θ -gaps, θ_1 and θ_2 , the following relation holds:

$$\log_{\theta_1} z - \log_{\theta_2} z = \begin{cases} 2\pi i \times \text{sgn}(\theta_1 - \theta_2) & (\min(\theta_1, \theta_2) < \arg z < \max(\theta_1, \theta_2)), \\ 0 & (\text{otherwise}). \end{cases} \quad (4)$$

Consider $U \subset X$ as a three-dimensional subspace where $g(x)$ maintains a θ -gap for $x \in U$. Let $\gamma(x) = (u_1(x), \dots, u_N(x)) \in U(N)$ be a unitary matrix diagonalizing $g(x)$, i.e., $g(x) = \gamma(x)\Lambda(x)\gamma(x)^\dagger$ with $\Lambda(x) = \text{diag}(\lambda_1, \dots, \lambda_N)$, where $\lambda_n \in U(1)$ for $n = 1, \dots, N$.

The exact form B_θ is given by [9]

$$B_\theta = Q + R_\theta, \quad (5)$$

$$Q = \frac{1}{4\pi} \text{Tr}[\gamma^{-1}d\gamma\Lambda\gamma^{-1}d\gamma\Lambda^{-1}], \quad (6)$$

$$R_\theta = \frac{1}{2\pi} \text{Tr}[\log_\theta \Lambda(\gamma^{-1}d\gamma)^2]. \quad (7)$$

Note that while Q is independent of θ , but R_θ does. It is evident that X can be covered by $\{U_i\}_i$ such that in each patch U , $g(x)$ exhibits a specific θ -gap θ_i .




FIG. 1. (a) Illustration of a θ -gap. (b) Projection onto the eigenspace between two θ -gaps. (c) Smearing eigenvalues over a finite set of vertices. In each panel, the unit circle in the complex plane is depicted, with blue arcs representing the spectrum of the $U(N)$ -valued matrix $g(x)$ within a local region. Red lines mark the locations of θ -gaps.

The unitary matrix γ is not unique due to the transformation $\gamma \mapsto \gamma W$, where $W \in U(N)$ commutes with Λ , satisfying $W\Lambda W^{-1} = \Lambda$. This ambiguity, however, does not affect Q and R_θ . To illustrate, consider the N eigenvalues divided into groups of $|I|$ degenerate ones, each with eigenvalue λ_I so that $\Lambda = \bigoplus_I \lambda_I \mathbf{1}_{|I|}$. Introduce a block matrix notation $A_{IJ} = (u_i^\dagger du_j)_{i \in I, j \in J}$. The transformation matrix W is expressed as $W = \bigoplus_I W_I$ as well, with $W_I \in U(|I|)$, modifying A_{IJ} to $A_{IJ} = W_I^\dagger A_{IJ} W_J + \delta_{IJ} W_I^{-1} dW_I$. Consequently, Q and R_θ can be represented as:

$$Q = \frac{1}{4\pi} \sum_{I, J; I \neq J} \text{Tr}_I [A_{IJ} A_{JI}] \lambda_J \lambda_I^{-1}, \quad (8)$$

$$R_\theta = \frac{1}{2\pi} \sum_{I, J; I \neq J} \text{Tr}_I [A_{IJ} A_{JI}] \log_\theta \lambda_I, \quad (9)$$

where Tr_I denotes the trace over indices $i \in I$. In the summation $\sum_{I, J}$, terms with $I = J$ can be excluded due to $\text{Tr}_I [(A_{II})^2] = 0$. This demonstrates the invariance of Q and R_θ under the transformation $\gamma \mapsto \gamma W$.

Another noteworthy aspect is that the difference in B_θ between two θ -gaps is a total derivative. For $0 \leq \theta_1, \theta_2 < 2\pi$, and using (4), it follows that:

$$B_{\theta_1} - B_{\theta_2} = d\alpha_{\theta_1, \theta_2}, \quad (10)$$

where

$$\begin{aligned} \alpha_{\theta_1, \theta_2} &= -i \text{sgn}(\theta_1 - \theta_2) \text{Tr} [P_{\theta_1, \theta_2} \gamma^{-1} d\gamma] \\ &= -i \text{sgn}(\theta_1 - \theta_2) \\ &\quad \times \sum_{n; \min(\theta_1, \theta_2) < \arg \lambda_n < \max(\theta_1, \theta_2)} u_n^\dagger du_n, \end{aligned} \quad (11)$$

with

$$P_{\theta_1, \theta_2} = \sum_{n; \min(\theta_1, \theta_2) < \arg \lambda_n < \max(\theta_1, \theta_2)} u_n u_n^\dagger, \quad (12)$$

the orthogonal projection onto the eigenspace for eigenvalues that fulfill $\min(\theta_1, \theta_2) < \arg \lambda_n < \max(\theta_1, \theta_2)$. In cases where $\theta_1 = \theta_2$ or no eigenvectors meet the condition $\min(\theta_1, \theta_2) < \arg \lambda_n < \max(\theta_1, \theta_2)$, $\alpha_{\theta_1, \theta_2}$ is simply null.

Now, we express the winding number $W_3[g]$ as a sum of line integrals, utilizing a cubic decomposition L of the manifold X . (Any simplicial decomposition is equally valid.) Within each cube c of the lattice L , we select $\theta_c \in [0, 2\pi)$ such that g_x for $x \in c$ exhibits a θ -gap of θ_c . Thus, $W_3[g]$ can be reformulated as a sum of integrals over all plaquettes:

$$\begin{aligned} W_3[g] &= \frac{1}{2\pi} \sum_c \int_c dB_{\theta_c} \\ &= \frac{1}{2\pi} \sum_c \int_{\partial c} B_{\theta_c} \\ &= \frac{1}{2\pi} \sum_p \int_p (B_{\theta_p^-} - B_{\theta_p^+}). \end{aligned} \quad (13)$$

Here, \sum_p runs over all plaquettes p in the lattice L , with each p being oriented. The gap parameters θ_p^+ and θ_p^- correspond to the cubes adjacent to plaquette p , in directions parallel and antiparallel to p 's normal vector, respectively, as depicted in Fig. 2 (a). This formulation further simplifies to a sum of line integrals:

$$\begin{aligned} W_3[g] &= \frac{1}{2\pi} \sum_p \int_p d\alpha_{\theta_p^-, \theta_p^+} \\ &= \frac{1}{2\pi} \sum_p \oint_{\partial p} \alpha_{\theta_p^-, \theta_p^+}. \end{aligned} \quad (14)$$

The transition to the last expression is contingent upon $\alpha_{\theta_p^-, \theta_p^+}$ being smoothly defined across plaquette p . When $\alpha_{\theta_p^-, \theta_p^+}$ is solely determined along the loop ∂p bordering




FIG. 2. (a) A plaquette p within the cubic lattice, showing θ_p^+ and θ_p^- as the θ -gaps of cubes adjacent to p , aligned parallel and anti-parallel to p 's normal vector, respectively. The vertices v_0, v_1, v_2 , and v_3 are sequentially labeled around the perimeter of plaquette p . (b) An edge $v_a v_b$ of the cubic lattice, illustrating the θ -gaps of cubes adjacent to the edge $v_a v_b$.

plaquette p , a 2π ambiguity may arise from large gauge transformations $u_n \rightarrow u_n e^{i\chi_n}$, where $\oint_{\partial p} d\chi_n = 2\pi$, potentially altering $W_3[g]$ by an integer. However, if the cubic lattice L is sufficiently fine relative to g 's spatial variations, the integral $\oint_{\partial p} \alpha_{\theta_p^-, \theta_p^+}$ approximates $0 \bmod 2\pi$, permitting its interpretation as an \mathbb{R} -valued quantity devoid of the 2π ambiguity.

We claim that the winding number as expressed in (14) can be calculated solely using the diagonalizing matrices $\gamma(v)$ at the vertices v of lattice L . Diagonalizing $g(v)$ at vertices $v \in L$ yields the eigenvector and eigenvalue pairs $\{u_n(v), \lambda_n(v)\}_{n=1, \dots, N}$ for each vertex v . The ordering of eigenvectors $u_n(v)$ is such that the

angles of eigenvalues ascend, satisfying $0 \leq \lambda_1(v) \leq \dots \leq \lambda_N(v) < 2\pi$. (Note that eigenvalues λ near 0 may reorder significantly under minor perturbations, yet this does not contribute to the discrete formula below.) The gap parameter θ_c for each cube c is determined as follows: From the eight vertices of cube c , denoted as $v \in c$, we derive $8N$ eigenvalues $\{\lambda_n(v)\}_{v \in c, n=1, \dots, N}$. By smearing all eigenvalues $\lambda_n(v)$, we have a set of intervals:

$$I_c = \bigcup_{v \in c, n=1, \dots, N} \left\{ \arg(\lambda_n(v) e^{i\delta\phi}) \in [0, 2\pi) \right. \\ \left. - \frac{\beta}{2N} < \delta\phi < \frac{\beta}{2N} \right\}. \quad (15)$$

Here, $0 < \beta < 1$ is a constant smearing parameter ensuring that adjacent eigenvalues fall within the same smeared interval. For example, we can set $\beta = 1/2$. We select a θ from the set $[0, 2\pi) \setminus I_c$ to serve as θ_c for cube c . (Refer to Fig. 1 (c) for visualization.) With θ -gaps for all cubes in lattice L thus defined, the gap parameters θ_p^+ and θ_p^- for each plaquette p are specified. For each corner vertex v_0, v_1, v_2, v_3 of plaquette p , we define $\gamma_{\{\theta_p^+, \theta_p^-\}}(v_j)$ as a $N \times N_q$ matrix:

$$\gamma_{\{\theta_p^+, \theta_p^-\}}(v_a) = \left(u_{n_1}(v_a), \dots, u_{n_{N_q}}(v_a) \right), \quad (16)$$

comprising N_q eigenvectors, with eigenvalue angles satisfying $\min(\theta_p^+, \theta_p^-) < \arg \lambda_{n_1}(v_a) \leq \dots \leq \arg \lambda_{n_{N_q}}(v_a) < \max(\theta_p^+, \theta_p^-)$. The nonnegative integer N_q , indicating the count of eigenvalues between $e^{i\theta_p^+}$ and $e^{i\theta_p^-}$, should be in common for the four vertices v_0, v_1, v_2, v_3 of plaquette p , assuming the lattice L is fine enough. Then, the line integral $\oint_{\partial p} \alpha_{\theta_p^-, \theta_p^+}$ can be approximated as

$$\oint_{\partial p} \alpha_{\theta_p^-, \theta_p^+} \cong \text{sgn}(\theta_p^+ - \theta_p^-) \times \text{Arg det} \left[\gamma_{\{\theta_p^+, \theta_p^-\}}(v_0)^\dagger \gamma_{\{\theta_p^+, \theta_p^-\}}(v_3) \gamma_{\{\theta_p^+, \theta_p^-\}}(v_3)^\dagger \gamma_{\{\theta_p^+, \theta_p^-\}}(v_2) \right. \\ \left. \times \gamma_{\{\theta_p^+, \theta_p^-\}}(v_2)^\dagger \gamma_{\{\theta_p^+, \theta_p^-\}}(v_1) \gamma_{\{\theta_p^+, \theta_p^-\}}(v_1)^\dagger \gamma_{\{\theta_p^+, \theta_p^-\}}(v_0) \right] =: \Phi_p, \quad (17)$$

where Arg denotes the principal value $-\pi < \text{Arg } z < \pi$. Consequently, we obtain a discrete formula for the wind-

ing number:

$$W_3^{\text{dis}}[g] = \frac{1}{2\pi} \sum_p \Phi_p \in \mathbb{Z}, \quad (18)$$

which relies solely on the diagonalization of matrix $g(v)$ at vertices $v \in L$ of the cubic lattice L approximating the manifold X .

The discrete formula (18) is inherently quantized. To

demonstrate this, we express $e^{2\pi i W_3^{\text{dis}}[g]}$ as the product of edge contributions:

$$\begin{aligned}
e^{2\pi i W_3^{\text{dis}}[g]} &= \prod_p e^{i\Phi_p} \\
&= \prod_{v_a v_b \in \{\text{edges}\}} \exp \left[i \arg \left\{ \det [\gamma_{\{\theta_1, \theta_0\}}(v_a)^\dagger \gamma_{\{\theta_1, \theta_0\}}(v_b)]^{\text{sgn}(\theta_1 - \theta_0)} \det [\gamma_{\{\theta_2, \theta_1\}}(v_a)^\dagger \gamma_{\{\theta_2, \theta_1\}}(v_b)]^{\text{sgn}(\theta_2 - \theta_1)} \right. \right. \\
&\quad \left. \left. \times \det [\gamma_{\{\theta_3, \theta_2\}}(v_a)^\dagger \gamma_{\{\theta_3, \theta_2\}}(v_b)]^{\text{sgn}(\theta_3 - \theta_2)} \det [\gamma_{\{\theta_0, \theta_3\}}(v_a)^\dagger \gamma_{\{\theta_0, \theta_3\}}(v_b)]^{\text{sgn}(\theta_0 - \theta_3)} \right\} \right]. \tag{19}
\end{aligned}$$

Here, $v_a v_b$ denotes an individual edge, and $\theta_0, \dots, \theta_3$ are the gap parameters for cubes adjacent to the edge $v_a v_b$, ordered counterclockwise from the vector $\overrightarrow{v_b v_a}$ pointing out of the page. See Fig. 2 (b). Regardless of the relative magnitudes of $\theta_0, \theta_1, \theta_2$, and θ_3 , each edge's contribution in (19) cancels out exactly due to the property that for $\theta < \theta' < \theta''$,

$$\gamma_{\{\theta, \theta''\}}(v) = (\gamma_{\{\theta, \theta'\}}(v_a), \gamma_{\{\theta', \theta''\}}(v_a)). \tag{20}$$

Thus, $e^{2\pi i W_3^{\text{dis}}[g]} = 1$ is valid for any sufficiently fine discrete approximation of X .

Model calculation— Our formulation extends to computing the winding number for maps $g : X \rightarrow GL_N(\mathbb{C})$, where the target space consists of invertible matrices. Invertible matrices that cannot be diagonalized, known as "exceptional points," constitute a ring in the three-dimensional parameter space and are stable under minor perturbations. To circumvent these exceptional points, one can employ the singular value decomposition $g = U\Sigma V^\dagger$ to derive the unitary matrix UV^\dagger at each vertex within the discretized parameter space. We verified our formula (18) with the model $g(k_x, k_y, k_z) = t(\sin k_x \sigma_x + \sin k_y \sigma_y + \sin k_z \sigma_z) - i(m + \cos k_x + \cos k_y + \cos k_z) \mathbf{1}_2$ on the three-torus $(k_x, k_y, k_z) \in [-\pi, \pi]^{\times 3}$, employing a cubic lattice of $20 \times 20 \times 20$ mesh. Here, $\sigma_\mu \in \{\sigma_x, \sigma_y, \sigma_z\}$ denotes the Pauli matrices. We have checked that the winding number $W_3^{\text{dis}}[g]$

equals $-2\text{sgn}(t)$ for $|m| < 1$, $\text{sgn}(t)$ for $1 < |m| < 3$, and 0 for $|m| > 3$, which is consistent with the direct calculation of the analytic form $W_3[g]$.

Summary— In this work, we presented a formulation for calculating the three-dimensional winding number $W_3[g]$ for smooth maps $g : X \rightarrow U(N)$, utilizing a discrete approximation of the manifold X . Our approach allows for the computation of $W_3[g]$ exclusively through the diagonalization of matrices $g(v)$ at a finite number of vertices, ensuring the result is explicitly quantized to integer values. Discrete formulations that explicitly quantify topological invariants are currently limited in scope. Examples such as instanton numbers represented by the second Chern numbers, higher-dimensional winding numbers, and degrees of maps to more general symmetric spaces have yet to be explored. We look forward to future studies shedding more light on these topics.

ACKNOWLEDGMENTS

We were supported by JST CREST Grant No. JPMJCR19T2, and JSPS KAKENHI Grant No. 22H05118 and 23H01097.

[1] G. E. Volovik, *The universe in a helium droplet*, Vol. 117 (OUP Oxford, 2003).

[2] A. P. Schnyder, S. Ryu, A. Furusaki, and A. W. W. Ludwig,

- Classification of topological insulators and superconductors in three spatial dimensions, *Phys. Rev. B* **78**, 195125 (2008).
- [3] M. Nakahara, *Geometry, topology and physics* (CRC press, 2018).
- [4] M. Lüscher, Topology of lattice gauge fields, *Communications in Mathematical Physics* **85**, 39 (1982).
- [5] T. Fujiwara, H. Suzuki, and K. Wu, Topological Charge of Lattice Abelian Gauge Theory, *Progress of Theoretical Physics* **105**, 789 (2001), <https://academic.oup.com/ptp/article-pdf/105/5/789/5186102/105-5-789.pdf>.
- [6] T. Fukui, Y. Hatsugai, and H. Suzuki, Chern numbers in discretized brillouin zone: Efficient method of computing (spin) hall conductances, *Journal of the Physical Society of Japan* **74**, 1674 (2005), <https://doi.org/10.1143/JPSJ.74.1674>.
- [7] F. Knöppel and U. Pinkall, Complex line bundles over simplicial complexes and their applications, in *Advances in Discrete Differential Geometry*, edited by A. I. Bobenko (Springer Berlin Heidelberg, Berlin, Heidelberg, 2016) pp. 197–239.
- [8] D. Carpentier, P. Delplace, M. Fruchart, K. Gawędzki, and C. Tauber, Construction and properties of a topological index for periodically driven time-reversal invariant 2d crystals, *Nuclear Physics B* **896**, 779 (2015).
- [9] K. Gawędzki and N. Reis, WZW branes and gerbes, *Reviews in Mathematical Physics* **14**, 1281 (2002), <https://doi.org/10.1142/S0129055X02001557>.