

HIGHER-ORDER REVERSE ISOPERIMETRIC INEQUALITIES FOR LOG-CONCAVE FUNCTIONS

DYLAN LANGHARST, FRANCISCO MARÍN SOLA, AND JACOPO ULIVELLI

ABSTRACT. The Rogers-Shephard and Zhang’s projection inequalities are two reverse, affine isoperimetric inequalities relating the volumes of a convex body and its difference body and polar projection body, respectively. Following a classical work by Schneider, both inequalities have been extended to the so-called higher-order setting. In this work, we establish the higher-order analogues for these inequalities in the setting of log-concave functions.

1. INTRODUCTION

In the Brunn-Minkowski theory of convex bodies (i.e., convex, compact subsets of \mathbb{R}^n with non-empty interior, for $n \in \mathbb{N}$ fixed), two main affinely invariant inequalities concern the difference body and the polar projection body of a convex body K in \mathbb{R}^n . The starting points of this work are the two corresponding reverse isoperimetric inequalities. Namely, the Rogers-Shephard and Zhang’s projection inequalities, which we will briefly introduce now; see Yang’s survey [63] for an introduction to the theory of affine structures in convex geometry.

The Minkowski sum of two sets $A, B \subseteq \mathbb{R}^n$ is the set $A + B = \{a + b : a \in A, b \in B\}$. Then, for a convex body K , its difference body is defined as $DK := K + (-K)$. One has that

$$(1.1) \quad 2^n \leq \frac{\text{vol}_n(DK)}{\text{vol}_n(K)} \leq \binom{2n}{n},$$

where vol_n is the Lebesgue measure on \mathbb{R}^n . The left-hand side of (1.1), which follows from the Brunn-Minkowski inequality (see (2.3) later), has equality if and only if K is symmetric (we say K is symmetric about the origin if $K = -K$ and K is symmetric if a translate of K is symmetric about the origin). The right-hand side of (1.1) is known as the Rogers-Shephard inequality. Here equality holds if and only if K is an n -dimensional simplex [54]. We recall the convex hull of a set $A \subset \mathbb{R}^n$, denoted $\text{conv } A$, is the smallest convex set containing A ; an n -dimensional simplex is the convex hull of $(n + 1)$ affinely independent points.

2020 *Mathematics Subject Classification.* Primary 26B25, 52A38, 52A40; Secondary 52A20.

Key words and phrases. Rogers-Shephard inequality, Zhang’s projection inequality, log-concave functions, difference body, polar projection body.

Denoting $\text{vol}_{n-1}(P_{x^\perp}K)$ as the volume of the orthogonal projection of K onto the linear subspace orthogonal to $x \in \mathbb{R}^n \setminus \{o\}$, the polar projection body $\Pi^\circ K$ of a convex body K is the origin symmetric convex body corresponding to the norm

$$(1.2) \quad \|x\|_{\Pi^\circ K} = \frac{1}{2} \int_{\mathbb{S}^{n-1}} |\langle x, u \rangle| dS_K(u) = |x| \text{vol}_{n-1}(P_{x^\perp}K),$$

where S_K is the surface area measure of a convex body K (see Section 2.2). The following inequalities then hold, denoting the unit Euclidean ball in \mathbb{R}^n as B_2^n :

$$(1.3) \quad \frac{1}{n^n} \binom{2n}{n} \leq \text{vol}_n(K)^{n-1} \text{vol}_n(\Pi^\circ K) \leq \left(\frac{\text{vol}_n(B_2^n)}{\text{vol}_{n-1}(B_2^{n-1})} \right)^n.$$

The left-hand side is Zhang's projection inequality, where equality holds if and only if K is a n -dimensional simplex [64]. The right-hand side is Petty's projection inequality, with equality if and only if K is an ellipsoid [51].

We now introduce the first framework for the extensions we consider. We write $\bar{x} = (x_1, \dots, x_m)$, $x_i \in \mathbb{R}^n$, for vectors in $(\mathbb{R}^n)^m \simeq \mathbb{R}^{nm}$. For a convex body in \mathbb{R}^n , Schneider [58] introduced the difference body of order $m \in \mathbb{N}$,

$$(1.4) \quad D^m(K) := \left\{ \bar{x} = (x_1, \dots, x_m) \in \mathbb{R}^{nm} : K \bigcap_{i=1}^m (x_i + K) \neq \emptyset \right\} \subset \mathbb{R}^{nm},$$

and then established the following Rogers-Shephard inequality of order m

$$(1.5) \quad \frac{\text{vol}_{nm}(D^m(K))}{\text{vol}_n(K)^m} \leq \binom{n(m+1)}{n}.$$

Equality in (1.5) holds if and only if K is a n -dimensional simplex. Concerning a lower bound, Schneider showed for $n = 2$ and for arbitrary $m \in \mathbb{N}$ that the minimum is obtained for every symmetric convex body but, for $n \geq 3$ and $m \geq 2$, this is false. In the same work, Schneider conjectured that (1.5) is minimized for ellipsoids for such n and m .

The interest towards the construction (1.4) has been recently reignited by the work of Roysdon [57], where (1.5) was generalized to product measures with suitable concavity properties. This setting was further investigated by the first author and collaborators in [31], contextualizing (1.4) in the so-called higher-order setting. In [31], the following notion was introduced. For a convex body K in \mathbb{R}^n we denote by $\Pi^{\circ, m}K$ its higher-order polar projection body, which is characterized by the non-symmetric norm, or gauge

$$(1.6) \quad \|\bar{\theta}\|_{\Pi^{\circ, m}K} = \int_{\mathbb{S}^{n-1}} \max_{1 \leq i \leq m} \langle u, \theta_i \rangle - dS_K(u), \quad \bar{\theta} = (\theta_1, \dots, \theta_m) \in \mathbb{S}^{nm-1}.$$

Notice that $\Pi^{\circ, m}K$ is a subset of \mathbb{R}^{nm} . Then, the higher-order Zhang's projection inequality is given by

$$(1.7) \quad \frac{1}{n^{nm}} \binom{n(m+1)}{n} \leq \text{vol}_n(K)^{m(n-1)} \text{vol}_{nm}(\Pi^{\circ, m}K),$$

with equality if and only if K is a n -dimensional simplex. Similarly, the higher-order Petty's projection inequality is given by

$$\text{vol}_n(K)^{m(n-1)} \text{vol}_{nm}(\Pi^{\circ,m} K) \leq \text{vol}_n(B_2^n)^{m(n-1)} \text{vol}_{nm}(\Pi^{\circ,m} B_2^n),$$

with equality if and only if K is an ellipsoid. See Section 2 for the motivation of the definition of $\Pi^{\circ,m} K$.

Our next framework is the setting of log-concave functions (see (2.1) below). The works of Ball [11], Artstein-Avidan, Klartag, and Milman [10], and Klartag and Milman [35] instigated the geometrification of log-concave functions, where results and behaviours similar to those of convex bodies are shown to hold for log-concave functions. Several fundamental inequalities from convex geometry have received extensions to log-concave functions, such as the Santaló inequality (and its reverse inequality) [10, 26, 28, 27, 40, 39, 23, 12, 24], Grünbaum's inequality [46, 48, 50], Petty's projection inequality [65, 61, 42, 6, 31, 32], Zhang's projection inequality [2, 4, 38] and the Rogers-Shephard inequality [16, 3, 5, 9, 1].

In particular, Zhang's projection inequality (1.3) was generalized to integrable log-concave functions by Alonso-Gutiérrez, Bernués, and González Merino [2]. In this paper, we prove a higher-order generalization of Zhang's projection inequality (1.7) for integrable log-concave function; we save for Section 2.3 needed definitions concerning the setting of functions of bounded variation $BV(\mathbb{R}^n)$ and an overview of the rich theory developed for the established functional Zhang's projection inequality in [4, 2], which served as a guide for our work.

Given a non-identically zero function $f \in BV(\mathbb{R}^n)$ there exists a vector map $\sigma_f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and a non-negative Radon measure $|Df|$, known as the variation measure of f . Furthermore, there exists a unique convex body with center of mass at the origin $\langle f \rangle$, called the LYZ body of f , with the following property: for every 1-homogeneous function g on $\mathbb{R}^n \setminus \{o\}$

$$(1.8) \quad \int_{S^{n-1}} g(u) dS_{\langle f \rangle}(u) = \int_{\mathbb{R}^n} g(\sigma_f(x)) d|Df|(x).$$

In previous applications (compare [44, 61, 43, 18]) the LYZ Body of f denoted the origin symmetric convex body whose surface area measure is the even part of $S_{\langle f \rangle}$. In this case (1.8) holds only if g is even and 1-homogeneous. It is possible to construct $\langle f \rangle$ to be not necessarily origin symmetric (see Section 2.3 for further details), which is better suited for our aims. Indeed, for $f \in BV(\mathbb{R}^n)$, its polar projection LYZ body of order m is defined as the convex body containing the origin $\Pi^{\circ,m} \langle f \rangle \subset \mathbb{R}^{nm}$ satisfying

$$\|\bar{\theta}\|_{\Pi^{\circ,m} \langle f \rangle} = \int_{\mathbb{R}^n} \max_{1 \leq i \leq m} \langle \sigma_f(x), \theta_i \rangle_- d|Df|(x).$$

For a measurable function $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$, where $\mathbb{R}_+ = [0, +\infty)$, we denote the standard L^p norm of f as $\|f\|_{L^p(\mathbb{R}^n)} = \left(\int_{\mathbb{R}^n} f^p dx \right)^{\frac{1}{p}}$. Our first result is the following functional, higher-order Zhang's projection inequality.

Theorem 1.1. *Fix $n, m \in \mathbb{N}$. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be an integrable, log-concave function. Then,*

$$\frac{1}{(nm)!} \int_{\mathbb{R}^{nm}} \int_{\mathbb{R}^n} \min_{0 \leq i \leq m} \{f(y - x_i)\} dy d\bar{x} \leq \|f\|_{L^1(\mathbb{R}^n)}^{nm+1} \text{vol}_{nm}(\Pi^{\circ, m}(f)),$$

where equality holds if and only if $f(x) = \|f\|_{L^\infty(\mathbb{R}^n)} e^{-\|x-x'\|_{\Delta_n}}$ for an n -dimensional simplex Δ_n containing the origin and some $x' \in \mathbb{R}^n$.

Using (2.4) below and the fact that the maximum of gauges corresponds to the gauge of the Cartesian product, Theorem 1.1 indeed becomes the left-hand side of (1.3) when $f = e^{-\|\cdot\|_{\mathcal{K}}}$. In Section 3.2, we introduce the higher-order radial mean bodies, and in Theorem 3.2, we establish the associated sharp set-inclusions. This provides an alternative proof of Theorem 1.1 and generalizes on the established radial mean bodies (see Section 2.2).

Our second result is the following higher-order version of the Rogers-Shephard inequality (1.5), which extends the $m = 1$ case established by Alonso-Gutiérrez, González Merino, Jiménez, and Villa [3]. Given $m + 1$ functions g_0, \dots, g_m on \mathbb{R}^n , define $\bar{g}((x_0, \bar{x})) \in \mathbb{R}^{m+1}$ as

$$\bar{g}((x_0, \bar{x})) = (g_0(x_0), g_1(x_1), \dots, g_m(x_m)).$$

We next consider the two following types of convolutions:

$$(\bar{g})_{\oplus m}(\bar{x}) := \int_{\mathbb{R}^n} g_0(z) g_1(z - x_1) \cdots g_m(z - x_m) dz$$

and

$$(\bar{g})_{\star m}(\bar{x}) := \sup_{z \in \mathbb{R}^n} g_0(z) g_1(z - x_1) \cdots g_m(z - x_m).$$

With these definitions in hand, we follow the ideas from [3] and prove the following Rogers-Shephard-type theorems. The first is for $m + 1$ different log-concave functions.

Theorem 1.2. *Fix $n, m \in \mathbb{N}$. Let $g_0, g_1, \dots, g_m : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be integrable, log-concave functions. Then, setting $\bar{g} = (g_0, g_1, \dots, g_m)$:*

$$\|(\bar{g})_{\oplus m}\|_{L^\infty(\mathbb{R}^{nm})} \|(\bar{g})_{\star m}\|_{L^1(\mathbb{R}^{nm})} \leq \binom{n(m+1)}{n} \left(\prod_{i=0}^m \|g_i\|_{L^\infty(\mathbb{R}^n)} \|g_i\|_{L^1(\mathbb{R}^n)} \right).$$

When setting $g_0 = f$ and replacing, for $i = 1, \dots, m$, each g_i with $f(-\cdot)$ in Theorem 1.2, we can obtain a sharper Rogers-Shephard-type inequality using a slightly different approach.

Theorem 1.3. *Fix $n, m \in \mathbb{N}$. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be an integrable, log-concave function and set $\bar{f} := (f(\cdot), f(-\cdot), \dots, f(-\cdot))$. Then:*

$$(1.9) \quad \|(\bar{f})_{\star m}\|_{L^1(\mathbb{R}^{nm})} \leq \binom{n(m+1)}{n} \|f\|_{L^\infty(\mathbb{R}^n)}^m \|f\|_{L^{\frac{1}{m}}(\mathbb{R}^n)},$$

where equality holds if and only if $f/\|f\|_{L^\infty(\mathbb{R}^n)}$ is the characteristic function of an n -dimensional simplex.

In this work, Section 3 is dedicated to proving Theorem 1.1, while Section 4 is devoted to the proofs of Theorem 1.2 and Theorem 1.3. In Section 2, we recall some basic facts from convex geometry and the theory of log-concave functions.

2. PRELIMINARIES

2.1. Facts from Convex Geometry and log-concave functions. A function $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$ is log-concave if for every $\lambda \in [0, 1]$ and $x, y \in \mathbb{R}^n$, one has

$$(2.1) \quad f((1 - \lambda)x + \lambda y) \geq f(x)^{1-\lambda} f(y)^\lambda.$$

In this work, we will always assume that log-concave functions have full-dimensional support. A fundamental inequality for this class of functions is the Prékopa-Leindler inequality ([52, 53], see also [41, 14]). In its general statement, it reads as follows: If $\lambda \in (0, 1)$ and $h, f, g : \mathbb{R}^n \rightarrow \mathbb{R}_+$ are non-negative measurable functions such that, for any $x, y \in \mathbb{R}^n$,

$$h((1 - \lambda)x + \lambda y) \geq f(x)^{1-\lambda} g(y)^\lambda,$$

then

$$(2.2) \quad \int_{\mathbb{R}^n} h(z) dz \geq \left(\int_{\mathbb{R}^n} f(x) dx \right)^{1-\lambda} \left(\int_{\mathbb{R}^n} g(y) dy \right)^\lambda.$$

See Dubuc [21] for the equality cases.

We now turn to the geometric aspects. One cornerstone of the theory of convex bodies is the Brunn-Minkowski inequality. This result, which follows from, and which may be seen as the geometrical counterpart of, the Prékopa-Leindler inequality states that if K and L be convex bodies and $\lambda \in [0, 1]$, then

$$(2.3) \quad \text{vol}_n((1 - \lambda)K + \lambda L) \geq \text{vol}_n(K)^{1-\lambda} \text{vol}_n(L)^\lambda.$$

Using the homogeneity of the volume, one may easily check that this is equivalent to

$$\text{vol}_n((1 - \lambda)K + \lambda L)^{\frac{1}{n}} \geq (1 - \lambda) \text{vol}_n(K)^{\frac{1}{n}} + \lambda \text{vol}_n(L)^{\frac{1}{n}},$$

where equality holds for some $\lambda \in (0, 1)$ if and only if K and L are homothetic (i.e., $K = tL + x$, $t \in \mathbb{R}_+$, $x \in \mathbb{R}^n$). We refer to Gardner's survey [29] for more details on the Brunn-Minkowski inequality and its connections to the Prékopa-Leindler inequality.

Every convex body K is the unit ball of its Minkowski functional, or gauge: $\|x\|_K = \sup\{t > 0 : x \in tK\}$. Alternatively, K is characterized by its support function $h_K(u) = \max_{y \in K} \langle y, u \rangle$. Notice that if K contains the origin, its gauge and support function satisfy the relation $\|x\|_K = h_{K^\circ}(x)$, where $K^\circ = \{x \in \mathbb{R}^n : h_K(x) \leq 1\}$ is the polar body of K . This leads to the classical formula

$$(2.4) \quad \text{vol}_n(K) = \frac{1}{n!} \int_{\mathbb{R}^n} e^{-\|x\|_K} dx.$$

Let \mathcal{H}^{n-1} be the $(n-1)$ -dimensional Hausdorff measure, let ∂K denote the boundary of K , and $n_K : \partial K \rightarrow \mathbb{S}^{n-1}$ the (generalized) Gauss map. The latter associates a point on ∂K with the corresponding outer-unit normal vectors. Then, for a Borel set $E \subseteq \mathbb{S}^{n-1}$ and a convex body K in \mathbb{R}^n , the surface area measure of K is defined as

$$S_K(E) = \int_{n_K^{-1}(E)} d\mathcal{H}^{n-1}(y).$$

Using support functions, (1.2) can be written, for $\theta \in \mathbb{S}^{n-1}$, as

$$\|\theta\|_{\Pi^\circ K} = \frac{1}{2} \int_{\mathbb{S}^{n-1}} h_{[-\theta, \theta]}(u) dS_K(u),$$

where for $x, y \in \mathbb{R}^n$ we write $[x, y] = \{tx + (1-t)y : t \in [0, 1]\}$. Analogously, (1.6) can be written for $\bar{\theta} \in \mathbb{S}^{nm-1}$ as

$$\|\bar{\theta}\|_{\Pi^{\circ, m} K} = \int_{\mathbb{S}^{n-1}} h_{C_{-\bar{\theta}}}(u) dS_K(u).$$

Here, $C_{\bar{x}} = \text{conv}\{o, x_1, \dots, x_m\}$ for $(x_1, \dots, x_m) = \bar{x} \in \mathbb{R}^{nm}$. Using the fact that, for a convex body K ,

$$\int_{\mathbb{S}^{n-1}} \langle \theta, u \rangle_- dS_K(u) = \text{vol}_{n-1}(P_{\theta^\perp} K) = \frac{1}{2} \int_{\mathbb{S}^{n-1}} |\langle \theta, u \rangle| dS_K(u),$$

we indeed have that $\Pi^{\circ, 1} K = \Pi^\circ K$. The covariogram of a convex body K is given by $g_K(x) = \text{vol}_n(K \cap (K + x))$; see the recent survey by Bianchi [13] for a rich overview of this function. It is relevant here that the support of g_K is DK , and it was shown by Matheron [47] that, for $\theta \in \mathbb{S}^{n-1}$,

$$\left. \frac{d}{dr} g_K(r\theta) \right|_{r=0^+} = -\|\theta\|_{\Pi^\circ K}.$$

Fixing $m \in \mathbb{N}$, Schneider [58] introduced an extension of the covariogram function of a convex body K , namely

$$g_{K, m}(\bar{x}) = \text{vol} \left(K \bigcap_{i=1}^m (x_i + K) \right),$$

which is supported on $D^m(K)$. It follows from the Brunn-Minkowski inequality that $g_{K, m}(\bar{x})^{\frac{1}{m}}$ is concave for all $m \in \mathbb{N}$ (see e.g. [58, Lemma 5.1] for a proof). Moreover, it was established in [31] that

$$\left. \frac{d}{dr} g_{K, m}(r\bar{\theta}) \right|_{r=0^+} = -\|\bar{\theta}\|_{\Pi^{\circ, m} K},$$

which motivated the definition of $\Pi^{\circ, m} K$.

2.2. Radial Mean Bodies and the Mellin transform. The constant $\binom{2n}{n}$ appearing in both the Rogers-Shephard and Zhang's projection inequality is not a coincidence. Gardner and Zhang showed that $DK \subseteq n\text{vol}_n(K)\Pi^\circ K$ (with equality only for n -dimensional simplices). In fact, they showed more. To elaborate on this, we first recall that a set $L \subset \mathbb{R}^n$ is a star body if $[0, x] \subset L$ for all $x \in L$, if it is compact with non-empty interior, and if the function on \mathbb{S}^{n-1} given by $\theta \mapsto \rho_L(\theta) := \|\theta\|_L^{-1}$ is continuous when (-1) -homogeneously extended to $\mathbb{R}^n \setminus \{o\}$; ρ_L is called the radial function of L .

Given a non-negative function g on \mathbb{R}^n with finite p th moment, $p > 0$, its Ball body is the star body $K_p(g)$ given by the radial function

$$\rho_{K_p(g)}^p(\theta) = \frac{p}{\|g\|_{L^\infty(\mathbb{R}^n)}} \int_0^\infty g(r\theta)r^{p-1}dr.$$

If g is a function such that g^s is concave on its support for some $s > 0$, g is said to be s -concave; $s = 0$ corresponds to log-concavity. If $s > 0$, then every such s -concave function is also log-concave. It was shown that if g is log-concave, then $K_p(g)$ is a convex body (see [11] and [30] for a proof); if g is additionally even, then $K_p(g)$ is origin symmetric.

The Mellin transform of an integrable function $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is the analytic function

$$\mathcal{M}_\psi(p) = \begin{cases} \int_0^\infty t^{p-1}(\psi(t) - \psi(0))dt & \text{for } p \in (-1, 0), \\ \int_0^\infty t^{p-1}\psi(t)dt & \text{for } p > 0 \text{ such that } t^{p-1}\psi(t) \in L^1(\mathbb{R}). \end{cases}$$

In [65], Gardner and Zhang introduced the radial mean bodies $R_p K$ of a convex body K . They can be defined, for $p > -1, p \neq 0$, as the star bodies with radial function

$$\rho_{R_p K}^p(\theta) = \frac{p}{\text{vol}_n(K)} \mathcal{M}_{g_K(r\theta)}(p)$$

(see [37] for $p \in (-1, 0)$). Moreover, since g_K is log-concave and even, the bodies $R_p K$ are origin symmetric and convex for every $p \geq 0$ since they are Ball bodies of the covariogram. For $p \in (-1, 0)$, the convexity of $R_p K$ is an open problem. The main result by Gardner and Zhang is the following [30]: for $-1 < p \leq q < \infty$,

$$(2.5) \quad R_\infty K = DK \subseteq \binom{n+q}{q}^{\frac{1}{q}} R_q K \subseteq \binom{n+p}{p}^{\frac{1}{p}} R_p K \subseteq n\text{vol}_n(K)\Pi^\circ K,$$

where equality holds in each inclusion if and only if K is a n -dimensional simplex. The fact that $\text{vol}_n(R_n K) = \text{vol}_n(K)$ shows that the chain of inclusions (2.5) interpolates between (1.1) and (1.3). Higher-order extensions $R_p^m K$ were shown in [31], and we will generalize them for an alternative proof of Theorem 1.1.

One way to prove (2.5) and the higher-order analogue is the Mellin-Berwald inequality for s -concave measures by Fradelizi, Jiange, and Madiman [25]. We will need the $s = 0$ case for log-concave functions, established prior by Koldobsky, Pajor, and Yaskin [36]. Define the function

$$\psi_{\log}(t) = e^{-t} \chi_{(0, \infty)}(t).$$

Then, for all $p > -1$, one has $\mathcal{M}_{\psi_{\log}}(p) = \Gamma(p)$.

Proposition 2.1 (Mellin-Berwald inequality for log-concave functions). *Let $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be an integrable and log-concave function such that $\psi(0) > 0$. Then, for $p \in (p_0, 0) \cup (0, p_1)$, set*

$$\Omega_\psi(p) = \frac{\mathcal{M}_\psi(p)}{\mathcal{M}_{\psi_{\log}}(p)} = \Gamma(p)^{-1} \mathcal{M}_\psi(p),$$

where $\Omega_\psi(0) = \psi(0)$, p_0 and p_1 are defined implicitly by $p_0 = \inf\{p > -1 : \Omega_{f, \psi}(p) > 0\}$, and $p_1 = \sup\{p < \infty : \Omega_{f, \psi}(p) < \infty\}$. Moreover, set for $p \in (p_0, 0) \cup (0, p_1)$

$$G_\psi(p) = (\Omega_\psi(p))^{1/p} = (\Gamma(p)^{-1} \mathcal{M}_\psi(p))^{1/p}$$

and $G_\psi(0) = \exp(\log(\Omega_\psi)'(0))$. Then, $G_\psi(p)$ is non-increasing in p . It is a constant, say $G_\psi(p) \equiv \alpha > 0$, if and only if $\log \psi(t) = \log \psi(0) + \log \psi_{\log}(\frac{t}{\alpha})$, i.e. ψ is log-affine. Furthermore, if ψ is non-increasing, then $p_0 = -1$ and $p_1 = \infty$.

Notice that some of the conclusions of Proposition 2.1 are hidden in the proof and not stated explicitly; see [37] for more details.

Next, we will need the following lemma to analyze the behaviour of the radial mean bodies as $p \rightarrow -1$; see e.g. [33, Lemma 4] for a proof.

Lemma 2.1. *If $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a measurable function with $\lim_{t \rightarrow 0^+} \varphi(t) = \varphi(0)$ and such that $\int_0^\infty t^{-s_0} \varphi(t) dt < \infty$ for some $s_0 \in (0, 1)$, then*

$$\lim_{s \rightarrow 1^-} (1-s) \int_0^\infty t^{-s} \varphi(t) dt = \varphi(0).$$

2.3. Functional Setting. Recall the classical definition (see, e.g. [22]): a function f is said to be of bounded variation if there exists a vector valued measure Df such that, if $\sigma_f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is the Radon-Nykodim derivative of Df with respect to $|Df|$ (the variation measure of f) on \mathbb{R}^n , one has

$$\int_{\mathbb{R}^n} f(x) \operatorname{div}(\psi(x)) dx = - \int_{\mathbb{R}^n} \langle \sigma_f(x), \psi(x) \rangle d|Df|(x)$$

for every compactly supported smooth vector field $\psi : \mathbb{R}^n \rightarrow \mathbb{R}^n$. The variation measure for a log-concave function has been studied in a series of works by Colesanti and Fragalà [17], Cordero-Erausquin and Klartag [19], and Rotem [55, 56]. In particular, in [56] Rotem observed that, if f is log-concave,

$$(2.6) \quad d(Df) = -\nabla f dx + f n_{\operatorname{supp}(f)} d\mathcal{H}^{n-1}|_{\partial \operatorname{supp}(f)},$$

where $\text{supp}(f)$ denotes the support of f . We will not use this representation of Df explicitly, but this formula can, at any time, be inserted into the appropriate equations below. We remark that expressions in the form of the right-hand side of (2.6) appear naturally as first variations of the integral of suitable perturbations of log-concave functions (i.e., a functional concept of surface area), as investigated in [17, 19, 55, 56, 34, 60].

It was shown by Wang [61], extending the treatment on $W^{1,1}(\mathbb{R}^n)$ by Lutwak, Yang, and Zhang [45] that, given a non-identically zero function $f \in BV(\mathbb{R}^n)$, there exists a unique Borel measure μ_f on \mathbb{S}^{n-1} , not concentrated on any great hemisphere, such that

$$\int_{\mathbb{S}^{n-1}} g(u) d\mu_f(u) = \int_{\mathbb{R}^n} g(-\sigma_f(x)) d|Df|(x)$$

for every 1-homogeneous function g on $\mathbb{R}^n \setminus \{o\}$. As noticed, for example, in [31], the measure μ_f is centred. One can then apply Minkowski's existence theorem [59, Section 8.2] to show that there exists a unique convex body (the so-called LYZ body), denoted by $\langle f \rangle$, with the center of mass at the origin such that $\mu_f = \sigma_{-\langle f \rangle}$. This yields that, indeed, $\langle f \rangle$ satisfies (1.8).

The LYZ body appears naturally, for example, in the anisotropic Sobolev inequality (see [49, Appendix, Section 2.1] and [7, 20, 31] for further developments) in the form

$$\|f\|_{L^{\frac{n}{n-1}}(\mathbb{R}^n)} \leq \text{vol}_n(\langle f \rangle)^{\frac{n-1}{n}}.$$

Using this inequality together with the right-hand side of (1.3) yields the affine Sobolev inequality for $f \in BV(\mathbb{R}^n)$ established by Wang [61], elaborating on the case for smooth functions by Zhang [65]. This procedure works in the higher-order setting as well [31].

We now present some facts in the absolutely continuous case for the reader's convenience. Everything holds in wider generality for functions of bounded variation, as elaborated on below. If $f \in W^{1,1}(\mathbb{R}^n)$, then, for almost every $x \in \mathbb{R}^n$, $d|Df|(x) = |\nabla f(x)| dx$ and $\sigma_f = \nabla f / |\nabla f|$ when $\nabla f \neq o$, and o otherwise. Therefore, the change of variables formula for the LYZ body (1.8) becomes for such f

$$(2.7) \quad \int_{\mathbb{S}^{n-1}} g(u) dS_{\langle f \rangle}(u) = \int_{\mathbb{R}^n} g(\nabla f(x)) dx.$$

When using the change of variables formula (2.7) for $\Pi^\circ \langle f \rangle$ it is standard to set $g = h_{[-\theta, \theta]}$, but we instead set $g = \frac{1}{2} h_{[-\theta, \theta]}$. Therefore, the polar projection LYZ body of a log-concave function $f \in W^{1,1}(\mathbb{R}^n)$ (also denoted as $\Pi^* f$, for example in [2, 3, 6, 4]) is given by

$$\|x\|_{\Pi^\circ \langle f \rangle} = \frac{1}{2} \int_{\mathbb{R}^n} |\langle \nabla f(y), x \rangle| dy.$$

With this normalization, $\Pi^{\circ,1} \langle f \rangle = \Pi^\circ \langle f \rangle$ and $\Pi^\circ \langle e^{-\|x\|_K} \rangle = \frac{1}{(n-1)!} \Pi^\circ K$.

Recall that Klartag and Milman [35] defined the shadow of a log-concave function f as $P_{x^\perp} f(y) = \max_{s \in \mathbb{R}} f\left(y + s \frac{x}{|x|}\right)$. One then has for an absolutely continuous, log-concave $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$:

$$\|x\|_{\Pi^\circ \langle f \rangle} = \frac{1}{2} \int_{\mathbb{R}^n} |\langle \nabla f(y), x \rangle| dy = |x| \int_{x^\perp} P_{x^\perp} f(y) dy$$

by the co-area formula. To generalize this procedure, we will use the following version of the co-area formula for functions of bounded variation. It is implied by Proposition 3.15 and Theorem 3.40 in [8].

Lemma 2.2. *Let $f \in BV(\mathbb{R}^n)$ and h a real-valued, 1-homogeneous function on $\mathbb{R}^n \setminus \{0\}$. Then,*

$$(2.8) \quad \int_{\mathbb{R}^n} h(\sigma_f(y)) d|Df|(y) = \int_0^{\|f\|_{L^\infty(\mathbb{R}^n)}} \int_{\{f=t\}} h(\sigma_f(y)) d\mathcal{H}^{n-1}(y) dt.$$

We now recall the functional Zhang's projection inequality from [2]. First, we rewrite the inequality for convex bodies (the right-hand side of (1.3)) in a suitable form. Notice that

$$\text{vol}_n(K)^2 = \int_{\mathbb{R}^n} g_K(x) dx.$$

Then, rearranging (1.3) we obtain

$$\frac{1}{n^n} \binom{2n}{n} \int_{\mathbb{R}^n} g_K(x) dx \leq \text{vol}_n(K)^{n+1} \text{vol}_n(\Pi^\circ K).$$

To suit this formulation, the following functional covariogram was introduced in [2]:

$$g_f(x) = \int_{\mathbb{R}^n} \min \{f(y), f(y-x)\} dy.$$

This led, in the same work, to the functional form of Zhang's projection inequality. More precisely, they proved that if $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$ is an integrable log-concave function, then

$$\frac{1}{n!} \int_{\mathbb{R}^n} g_f(x) dx \leq \|f\|_{L^1(\mathbb{R}^n)}^{n+1} \text{vol}_n(\Pi^\circ \langle f \rangle).$$

Among the functions with $\|f\|_{L^\infty(\mathbb{R}^n)} = f(o)$, there is equality if and only if $f(x) = \|f\|_{L^\infty(\mathbb{R}^n)} e^{-\|x\|_{\Delta_n}}$ for any n -dimensional simplex Δ_n containing the origin.

As for the functional Rogers-Shephard inequality, one must first introduce an appropriate generalization of Minkowski addition. For two log-concave functions $f, g : \mathbb{R}^n \rightarrow \mathbb{R}_+$, their sup-convolution is

$$f \star g(z) = \sup_{x+y=z} f(x)g(y).$$

Setting $\bar{f}(x) = f(-x)$, Alonso-Gutiérrez, González Merino, Jiménez, and Villa [3, Theorem 2.2] showed, elaborating on a previous inequality by Colesanti [16], that

$$\int_{\mathbb{R}^n} f \star \bar{f}(z) dz \leq \binom{2n}{n} \|f\|_{L^\infty(\mathbb{R}^n)} \|f\|_{L^1(\mathbb{R}^n)},$$

with equality if and only if $f/\|f\|_{L^\infty(\mathbb{R}^n)}$ is the characteristic function of an n -dimensional simplex. In fact, they also established (see [3, Theorem 2.1]) a generalization when \bar{f} is replaced by any log-concave function g .

3. THE HIGHER-ORDER, FUNCTIONAL SETTING: ZHANG'S PROJECTION INEQUALITY

We now establish the functional, higher-order Zhang's projection inequality. We henceforth set $x_0 = o$. Then, for $\bar{x} = (x_1, \dots, x_m) \in \mathbb{R}^{nm}$, we set the higher-order covariogram of a function to be

$$g_{f,m}(\bar{x}) = \int_{\mathbb{R}^n} \min_{0 \leq i \leq m} \{f(y - x_i)\} dy.$$

Observe that $g_{f,m}(\bar{o}) = \|f\|_{L^1(\mathbb{R}^n)}$. We will prove the following theorem.

Theorem 1.1. *Fix $n, m \in \mathbb{N}$. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be an integrable, log-concave function. Then,*

$$\frac{1}{(nm)!} \int_{\mathbb{R}^{nm}} g_{f,m}(\bar{x}) d\bar{x} \leq \|f\|_{L^1(\mathbb{R}^n)}^{nm+1} \text{vol}_{nm}(\Pi^{\circ,m}\langle f \rangle).$$

There is equality if and only if $f(x) = \|f\|_{L^\infty(\mathbb{R}^n)} e^{-\|x-x'\|_{\Delta_n}}$ for an n -dimensional simplex Δ_n containing the origin and some $x' \in \mathbb{R}^n$.

3.1. The Polar Projection LYZ Body. It will be convenient to represent $\Pi^{\circ,m}\langle f \rangle$ in terms of $\Pi^{\circ,m}\{f \geq t\}$. Observe that, by definition,

$$(3.1) \quad \|\bar{x}\|_{\Pi^{\circ,m}\langle f \rangle} = \int_{\mathbb{S}^{n-1}} h_{C_{\bar{x}}}(u) d\sigma_{\langle f \rangle}(u),$$

where we recall that $C_{\bar{x}} = \text{conv}\{o, x_1, \dots, x_m\}$. In fact, using (3.1), one obtains $\|\bar{x}\|_{\Pi^{\circ,m}\langle e^{-\|x\|_K} \rangle} = (n-1)! \|x\|_{\Pi^{\circ,m}K}$, and so

$$\Pi^{\circ,m}\langle e^{-\|x\|_K} \rangle = \frac{1}{(n-1)!} \Pi^{\circ,m}K.$$

Next, we use (3.1), the change of variables formula for the LYZ body of a log-concave function f , the co-area formula (2.8), and the fact that $\{f = t\}$

is merely the boundary of $\{f \geq t\}$ with outer-unit normal (defined for \mathcal{H}^{n-1} -almost every y and up to a choice of sign) $n_{\{f \geq t\}}(y) = \sigma_f(y)$ to obtain

$$\begin{aligned}
\|\bar{x}\|_{\Pi^{\circ,m}\langle f \rangle} &= \int_{\mathbb{R}^n} h_{C_{-\bar{x}}}(\sigma_f(y)) d|Df|(y) \\
&= \int_0^{\|f\|_{L^\infty(\mathbb{R}^n)}} \int_{\{f=t\}} h_{C_{-\bar{x}}}(\sigma_f(y)) d\mathcal{H}^{n-1}(y) dt \\
(3.2) \quad &= \int_0^{\|f\|_{L^\infty(\mathbb{R}^n)}} \int_{\partial\{f \geq t\}} h_{C_{-\bar{x}}}(n_{\{f \geq t\}}(y)) d\mathcal{H}^{n-1}(y) dt \\
&= \int_0^{\|f\|_{L^\infty(\mathbb{R}^n)}} \|\bar{x}\|_{\Pi^{\circ,m}\{f \geq t\}} dt.
\end{aligned}$$

Remark 3.1. Notice that a Zhang's-projection-type inequality for integrals of $\text{vol}_{nm}(\Pi^{\circ,m}\{f \geq t\})$ is weaker than an inequality involving the LYZ body. Indeed, by Jensen's inequality and (3.2)

$$\begin{aligned}
\|f\|_{L^\infty(\mathbb{R}^n)}^{nm} \|\bar{x}\|_{\Pi^{\circ,m}\langle f \rangle}^{-nm} &= \left(\int_0^{\|f\|_{L^\infty(\mathbb{R}^n)}} \|\bar{x}\|_{\Pi^{\circ,m}\{f \geq t\}} \frac{dt}{\|f\|_{L^\infty(\mathbb{R}^n)}} \right)^{-nm} \\
&\leq \int_0^{\|f\|_{L^\infty(\mathbb{R}^n)}} \|\bar{x}\|_{\Pi^{\circ,m}\{f \geq t\}}^{-nm} \frac{dt}{\|f\|_{L^\infty(\mathbb{R}^n)}}.
\end{aligned}$$

Integrating over the sphere, we deduce

$$\|f\|_{L^\infty(\mathbb{R}^n)}^{nm+1} \text{vol}_{nm}(\Pi^{\circ,m}\langle f \rangle) \leq \int_0^{\|f\|_{L^\infty(\mathbb{R}^n)}} \text{vol}_{nm}(\Pi^{\circ,m}\{f \geq t\}) dt.$$

Before getting into the proof of Theorem 1.1, let us point out some crucial properties of the functional covariogram.

3.2. The Functional Covariogram. First, notice that there is a relation between the functional and geometric covariograms, mainly

$$\begin{aligned}
(3.3) \quad g_{f,m}(\bar{x}) &= \int_{\mathbb{R}^n} \min_{0 \leq i \leq m} \{f(y - x_i)\} dy = \int_{\mathbb{R}^n} \int_0^{\min_{0 \leq i \leq m} \{f(y - x_i)\}} dt dx \\
&= \int_0^\infty g_{\{f \geq t\},m}(\bar{x}) dt.
\end{aligned}$$

We next take the radial derivative of the functional covariogram. First, we need the following lemma. Recall that, for a function h on the sphere \mathbb{S}^{n-1} , the Wulff shape of h is given by $[h] = \{x \in \mathbb{R}^n : \max_{u \in \mathbb{S}^{n-1}} \langle x, u \rangle \leq h(u)\}$. If K is a convex body containing the origin, then $K = [h_K]$.

Lemma 3.1. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be an integrable, log-concave function, and let $\theta \in \mathbb{S}^{nm-1}$. Then, for $\varepsilon > 0$ suitably small,

- (a) For every $r \in [0, \varepsilon]$, $g_{\{f \geq t\},m}(r\bar{\theta})$ is integrable in $t \in [0, \|f\|_\infty]$,

(b) *The partial derivative*

$$\frac{d}{dr}g_{\{f \geq t\},m}(r\bar{\theta})$$

exists for every $r \in [0, \varepsilon]$ and $t \in [0, \|f\|_\infty)$,

(c) *There exists $c(t)$ integrable on $[0, \|f\|_\infty]$ such that*

$$g_{\{f \geq t\},m}(r\bar{\theta}) \leq c(t)$$

for every $r \in [0, \varepsilon]$ and $t \in [0, \|f\|_\infty]$.

Proof. Notice that (a) follows from (3.3) and the fact that f is integrable. Point (c) follows choosing $c(t) = \text{vol}_n(\{f \geq t\})$.

We now focus on point (b), which is slightly more delicate. Notice that, for a convex body K , we can write (cf. [31, Proposition 3.2])

$$g_{K,m}(r\bar{\theta}) = \text{vol}_n([h_K - rh_{C_{-\bar{\theta}}}]).$$

By the semigroup property of the Wulff shape (compare [62, Theorem 5.7]) and Aleksandrov's variational lemma [59, Lemma 7.5.3], we have for $\varepsilon > 0$ sufficiently small that

$$\frac{d}{dr}g_{\{f \geq t\},m}(r\bar{\theta})$$

is right-continuous in r on the interval $[0, \varepsilon]$. Since $g_{K,m}(r\bar{\theta})$ is continuous for $r \geq 0$ and has continuous right derivative on $[0, \varepsilon]$, it is differentiable on $[0, \varepsilon]$ (see, for example, [15, Theorem 1.3, page 40]). Notice that for every $t \in [0, \|f\|_\infty)$ the set $\{f \geq t\}$ has non-empty interior so that we can apply the previous procedure to every level set (except, at most, the one corresponding to the maximum). This concludes the proof of (b). \square

Theorem 3.1. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be an integrable, log-concave function. Then, for every $\bar{\theta} \in \mathbb{S}^{nm-1}$,*

$$\frac{d}{dr}g_{f,m}(r\bar{\theta})|_{r=0+} = -\|\bar{\theta}\|_{\Pi^{\circ,m}\langle f \rangle}.$$

Proof. Observe that (3.3) and (3.2) yield

$$\begin{aligned} \frac{d}{dr}g_{f,m}(r\bar{\theta})|_{r=0+} &= \int_0^\infty \frac{d}{dr}g_{\{f \geq t\},m}(r\bar{\theta})|_{r=0+} dt \\ &= - \int_0^\infty \|\bar{x}\|_{\Pi^{\circ,m}\{f \geq t\}} dt = -\|\bar{\theta}\|_{\Pi^{\circ,m}\langle f \rangle}, \end{aligned}$$

where we could exchange the signs of integral and derivative since, by Lemma 3.1, we can apply the Leibniz integral rule. \square

As a corollary, we obtain the following ‘‘tangent-line bound’’ for the functional covariogram.

Corollary 3.1. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be an integrable, log-concave function. Then, $g_{f,m}$ is log-concave and*

$$(3.4) \quad g_{f,m}(\bar{x}) \leq \|f\|_{L^1(\mathbb{R}^n)} e^{-\frac{\|\bar{x}\|_{\Pi^{\circ,m}\langle f \rangle}}{\|f\|_{L^1(\mathbb{R}^n)}}},$$

with equality if and only if $g_{f,m}(\bar{x})$ is log-affine.

Proof. First, observe that $g_{f,m}$ is log-concave. Indeed, the minimum of log-concave functions is log-concave (since the maximum of convex functions is convex). Then, from the Prékopa-Leindler (2.2) inequality applied in \mathbb{R}^{nm} , we obtain the log-concavity of $g_{f,m}$.

Next, we use $r \mapsto g_{f,m}(r\bar{\theta})$ is log-concave as well, to obtain from Theorem 3.1 that, for every $r > 0$ so that $r\bar{\theta}$ is in the support of $g_{f,m}$,

$$\begin{aligned} -\frac{\|\bar{\theta}\|_{\Pi^{\circ,m}\langle f \rangle}}{\|f\|_{L^1(\mathbb{R}^n)}} &= \frac{d}{dr} \log g_{f,m}(r\bar{\theta}) \Big|_{r=0^+} \geq \frac{\log g_{f,m}(r\bar{\theta}) - \log \|f\|_{L^1(\mathbb{R}^n)}}{r} \\ &= \frac{\log \left(\frac{g_{f,m}(r\bar{\theta})}{\|f\|_{L^1(\mathbb{R}^n)}} \right)}{r}. \end{aligned}$$

Consequently, we obtain

$$r\|\bar{\theta}\|_{\Pi^{\circ,m}\langle f \rangle} \leq -\|f\|_{L^1(\mathbb{R}^n)} \log \left(\frac{g_{f,m}(r\bar{\theta})}{\|f\|_{L^1(\mathbb{R}^n)}} \right).$$

This is precisely our claim after exponentiation. \square

In the following lemma, we show that equality occurs in Corollary 3.1 if and only if f is a multiple of $e^{-\|x-x'\|_{\Delta_n}}$ for some $x' \in \mathbb{R}^n$ and n -dimensional simplex Δ_n containing the origin.

Lemma 3.2. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be an integrable, log-concave function such that $g_{f,m}$ is log-affine. Then, one must have $f(x) = \|f\|_{L^\infty(\mathbb{R}^n)} e^{-\|x-x'\|_{\Delta_n}}$ for an n -dimensional simplex Δ_n containing the origin and some $x' \in \mathbb{R}^n$.*

Proof. Notice that one may verify via direct substitution that if $f(x) = \|f\|_{L^\infty(\mathbb{R}^n)} e^{-\|x-x'\|_{\Delta_n}}$ for an n -dimensional simplex Δ_n containing the origin and some $x' \in \mathbb{R}^n$, then $g_{f,m}$ is log-affine. It remains to show the converse direction.

We may suppose that $\|f\|_{L^\infty(\mathbb{R}^n)} = f(o)$. Indeed, for an arbitrary integrable, log-concave function f , take $x' \in \{x \in \mathbb{R}^n : f(x) = \|f\|_{L^\infty(\mathbb{R}^n)}\}$, and set $\tilde{f}(x) = f(x+x')$. Then, $\tilde{f}(x)$ has maximum at the origin and $g_{f,m}$ is log-affine if and only if $g_{\tilde{f},m}$ is log-affine. In fact, $g_{f,m} = g_{\tilde{f},m}$.

For $x \in \mathbb{R}^n$, let $\tilde{x} \in \mathbb{R}^{nm}$ be $\tilde{x} = (x, o, \dots, o)$. Then, $g_{f,m}(\tilde{x}) = g_{\tilde{f},m}(\tilde{x})$. Since the restriction of a log-affine function is still log-affine, we obtain that $g_{\tilde{f}}$ is log-affine. It is precisely the content of [3, Lemma 2.4.2] that, if $f(o) = 0$ and $g_{\tilde{f}}$ is log-affine, then $f(x) = \|f\|_{L^\infty(\mathbb{R}^n)} e^{-\|x\|_{\Delta_n}}$ for some n -dimensional simplex Δ_n containing the origin. \square

Proof of Theorem 1.1. Observe that, integrating both sides of (3.4) over \mathbb{R}^{nm} yields

$$\begin{aligned} \frac{1}{(nm)!} \int_{\mathbb{R}^{nm}} g_{f,m}(\bar{x}) d\bar{x} &\leq \|f\|_{L^1(\mathbb{R}^n)} \frac{1}{(nm)!} \int_{\mathbb{R}^{nm}} e^{-\frac{\|\bar{x}\|_{\Pi^{\circ,m}\langle f \rangle}}{\|f\|_{L^1(\mathbb{R}^n)}}} d\bar{x} \\ &= \|f\|_{L^1(\mathbb{R}^n)}^{nm+1} \frac{1}{(nm)!} \int_{\mathbb{R}^{nm}} e^{-\|\bar{x}\|_{\Pi^{\circ,m}\langle f \rangle}} d\bar{x} \\ &= \|f\|_{L^1(\mathbb{R}^n)}^{nm+1} \text{vol}_{nm}(\Pi^{\circ,m}\langle f \rangle). \end{aligned}$$

with equality if and only if $g_{f,m}$ is log-affine. Equality conditions then follow from Lemma 3.2. \square

3.3. The Approach of Radial Mean Bodies. We now introduce the functional analogue of the radial mean bodies. The higher-order, functional p th radial mean body of an integrable function f on \mathbb{R}^n for $-1 < p \leq q < \infty$ is the star body $R_p^m f$ on \mathbb{R}^{nm} given by the radial function: for $\bar{\theta} \in \mathbb{S}^{nm-1}$,

$$(3.5) \quad \rho_{R_p^m f}(\bar{\theta})^p = \begin{cases} \frac{p}{\|f\|_{L^1(\mathbb{R}^n)}} \int_0^\infty g_{f,m}(r\bar{\theta}) r^{p-1} dr, & p > 0, \\ \frac{p}{\|f\|_{L^1(\mathbb{R}^n)}} \int_0^\infty (g_{f,m}(r\bar{\theta}) - \|f\|_{L^1(\mathbb{R}^n)}) r^{p-1} dr, & p \in (-1, 0), \end{cases}$$

Notice that these are well defined since if the support of f is full dimensional, the support of $g_{f,m}$ is also full dimensional and contains the origin in its interior. Moreover, for $p \geq 0$, $R_p^m f$ is again a convex body (if $m = 1$, then it is always origin symmetric since g_f is always an even function, and if $m \geq 2$, $R_p^m f$ is origin symmetric if and only if f is even).

Notice that we can apply integration by parts to (3.5) to obtain for all $p \in (-1, 0) \cup (0, \infty)$

$$\rho_{R_p^m f}(\bar{\theta})^p = \int_0^\infty \left(-\frac{g_{f,m}(r\bar{\theta})}{\|f\|_{L^1(\mathbb{R}^n)}} \right)' r^p dr,$$

where the prime denotes derivative in r and we used Lebesgue's theorem to obtain that $g_{f,m}(r\bar{\theta})$ is differentiable almost everywhere on its support (in r) as it is monotonically decreasing in the variable r . For the next result, we define

$$p_{\max} = \sup \left\{ p < \infty : \left(\frac{1}{\Gamma(p+1)} \right)^{\frac{1}{p}} R_p^m f \neq \{\bar{o}\} \right\}.$$

Theorem 3.2. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be an integrable, log-concave function. Then, for $-1 < p < q < p_{\max}$,*

$$\left(\frac{1}{\Gamma(q+1)} \right)^{\frac{1}{q}} R_q^m f \subseteq \left(\frac{1}{\Gamma(p+1)} \right)^{\frac{1}{p}} R_p^m f \subseteq \|f\|_{L^1(\mathbb{R}^n)} \Pi^{\circ,m}\langle f \rangle,$$

with equality if and only if

$$g_{f,m}(\bar{x}) = \|f\|_{L^1(\mathbb{R}^n)} e^{-\frac{\|\bar{x}\|_{\Pi^{\circ,m}\langle f \rangle}}{\|f\|_{L^1(\mathbb{R}^n)}}}.$$

Equivalently, there is equality if and only if $f(x) = \|f\|_{L^\infty(\mathbb{R}^n)} e^{-\|x-x'\|_{\Delta_n}}$ for an n -dimensional simplex Δ_n containing the origin and some $x' \in \mathbb{R}^n$.

Proof. Notice that for a fixed $\bar{\theta}$ the function $r \mapsto \|f\|_{L^1(\mathbb{R}^n)}^{-1} g_{f,m}(r\bar{\theta})$ is non-increasing. Furthermore, we recall that $g_{f,m}(r\bar{\theta})$ is log-concave and thus, from Proposition 2.1, the function

$$p \mapsto \left(\frac{1}{\Gamma(p+1)} \right)^{\frac{1}{p}} \rho_{R_p^m} f(\bar{\theta})$$

is non-decreasing, and is constant if and only if for a fixed $\bar{\theta}$

$$(3.6) \quad g_{f,m}(r\bar{\theta}) = \|f\|_{L^1(\mathbb{R}^n)} e^{-cr}$$

for some $c > 0$, establishing our claim.

We now discuss the limiting behaviour in p (and determine c). Observe that

$$\lim_{p \rightarrow \infty} \left(\frac{1}{\Gamma(p+1)} \right)^{\frac{1}{p}} \rho_{R_p^m} f(\bar{\theta}) = 0$$

as $\Gamma(p+1)$ will be the dominating term. This yields

$$\lim_{p \rightarrow \infty} \left(\frac{1}{\Gamma(p+1)} \right)^{\frac{1}{p}} R_p^m f = \{\bar{o}\}.$$

To determine the behaviour as $p \rightarrow -1$, observe we can write

$$\frac{1}{\Gamma(p+1)} \rho_{R_p^m} f(\bar{\theta})^p = \frac{1}{\Gamma(p+2)} [(p+1) \rho_{R_p^m} f(\bar{\theta})^p].$$

We then obtain from Lemma 2.1 that

$$\begin{aligned} \lim_{p \rightarrow -1} \frac{1}{\Gamma(p+2)} [(p+1) \rho_{R_p^m} f(\bar{\theta})^p] &= -\frac{1}{\|f\|_{L^1(\mathbb{R}^n)}} \frac{d}{dr} g_{f,m}(r\bar{\theta}) \Big|_{r=0^+} \\ &= \frac{\|\bar{\theta}\|_{\Pi^{\circ,m}\langle f \rangle}}{\|f\|_{L^1(\mathbb{R}^n)}}. \end{aligned}$$

Now, again, suppose there is equality. In this case, when determining the behaviour as $p \rightarrow -1$, calculate the derivative of $g_{f,m}$ using (3.6) to conclude

$$c = \frac{\|\bar{\theta}\|_{\Pi^{\circ,m}\langle f \rangle}}{\|f\|_{L^1(\mathbb{R}^n)}}.$$

The equivalent form of the equality conditions follows from Lemma 3.2. \square

Alternative proof of Theorem 1.1. By integrating in polar coordinates, we obtain that

$$\begin{aligned} \frac{1}{\|f\|_{L^1(\mathbb{R}^n)}} \int_{\mathbb{R}^{nm}} g_{f,m}(\bar{x}) d\bar{x} &= \frac{1}{\|f\|_{L^1(\mathbb{R}^n)}} \int_{\mathbb{S}^{nm-1}} \int_0^\infty g_{f,m}(r\bar{\theta}) r^{nm-1} dr d\bar{\theta} \\ &= \frac{1}{nm} \int_{\mathbb{S}^{nm-1}} \rho_{R_{nm}^m} f(\bar{\theta})^{nm} d\bar{\theta} = \text{vol}_{nm}(R_{nm}^m f). \end{aligned}$$

Consequently, Theorem 1.1 follows from Theorem 3.2 by setting $p = nm$ and considering the volume of both sides. \square

Remark 3.2. *In the classical setting, $DK = \text{supp}(g_K)$, and the difference body appears in the radial mean body set inclusions. However, we see in Theorem 3.2 that $\text{supp}(g_{f,m})$ does not appear, and, in fact, for an arbitrary log-concave function, $\text{supp}(g_{f,m})$ might be \mathbb{R}^{nm} . Thus, in order to establish functional Rogers-Shephard inequalities, we must use different choices of “covariograms”.*

4. THE ROGERS-SHEPHARD INEQUALITY IN THE FUNCTIONAL, HIGHER-ORDER SETTING

In order to establish the functional, higher-order Rogers-Shephard inequality, some definitions are in order; these definitions generalize the $m = 1$ case in [3].

Recall the following from the introduction: given $m+1$ functions g_0, \dots, g_m on \mathbb{R}^n , consider the vector-valued function $\bar{g} : \mathbb{R}^{n(m+1)} \rightarrow \mathbb{R}^{m+1}$ as $\bar{g}((x_0, \bar{x})) = (g_0(x_0), g_1(x_1), \dots, g_m(x_m))$. Next, we define

$$\mathcal{A}_t^m(\bar{g})(\bar{x}) = \{z \in \text{supp } g_0 \cap \bigcap_{i=1}^m (x_i - \text{supp } g_i) \subset \mathbb{R}^n : \\ g_0(z)g_1(z - x_1) \cdots g_m(z - x_m) \geq t \prod_{i=0}^m \|g_i\|_{L^\infty(\mathbb{R}^n)}\}.$$

Then, the higher-order convolution body of \bar{g} is

$$\mathcal{C}_{\theta,t}^m(\bar{g}) = \{\bar{x} \in \mathbb{R}^{nm} : \mathcal{A}_t^m(\bar{g})(\bar{x}) \neq \emptyset, \text{vol}_n(\mathcal{A}_t^m(\bar{g})(\bar{x})) \geq \theta M_t\},$$

where $M_t = \max_{\bar{x}} \text{vol}_n(\mathcal{A}_t^m(\bar{g})(\bar{x}))$.

When for all $i = 0, \dots, m$ we have $g_i = f$ for some function f , we write $\mathcal{A}_t^m(f)(\bar{x}) = \mathcal{A}_t^m((f, \dots, f))(\bar{x})$. Then, we introduced the following two operations for these vector-valued maps, turning them into functions on \mathbb{R}^{nm} :

$$(\bar{g})_{\oplus_m}(\bar{x}) = \int_{\mathbb{R}^n} g_0(z)g_1(z - x_1) \cdots g_m(z - x_m) dz$$

and

$$(\bar{g})_{\star_m}(\bar{x}) = \sup_{z \in \mathbb{R}^n} g_0(z)g_1(z - x_1) \cdots g_m(z - x_m).$$

We now recall the Rogers-Shephard-type inequalities from the introduction, whose proofs will take the remainder of this section.

Theorem 1.2. *Fix $n, m \in \mathbb{N}$. Let $g_0, g_1, \dots, g_m : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be integrable, log-concave functions. Then, setting $\bar{g} = (g_0, g_1, \dots, g_m)$:*

$$\|(\bar{g})_{\oplus_m}\|_{L^\infty(\mathbb{R}^{nm})} \|(\bar{g})_{\star_m}\|_{L^1(\mathbb{R}^{nm})} \leq \binom{n(m+1)}{n} \left(\prod_{i=0}^m \|g_i\|_{L^\infty(\mathbb{R}^n)} \|g_i\|_{L^1(\mathbb{R}^n)} \right).$$

Theorem 1.3. *Fix $n, m \in \mathbb{N}$. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be an integrable, log-concave function and set $\bar{f}(\cdot) := (f(\cdot), f(-\cdot), \dots, f(-\cdot))$. Then:*

$$\|(\bar{f})_{\star_m}\|_{L^1(\mathbb{R}^{nm})} \leq \binom{n(m+1)}{n} \|f\|_{L^\infty(\mathbb{R}^n)}^m \|f\|_{L^{\frac{1}{m}}(\mathbb{R}^n)},$$

where equality holds, if and only if, $f/\|f\|_{L^\infty(\mathbb{R}^n)}$ is the characteristic function of an n -dimensional simplex.

We start by establishing some lemmas that will be used along the proofs. Notice that they are the analogues in our setting of Lemmas 4.1 and 4.3 in [3]. First, we show the following set inclusion.

Lemma 4.1. *Let $g_0, \dots, g_m : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be log-concave functions and set $\bar{g}((x_0, \bar{x})) = (g_0(x_0), \dots, g_m(x_m))$. Let $t \in [0, 1]$, and $\theta_1, \theta_2, \lambda_1, \lambda_2 \in [0, 1]$ such that $\lambda_1 + \lambda_2 \leq 1$. Then*

$$\lambda_1 \mathcal{C}_{\theta_1, t}^m(\bar{g}) + \lambda_2 \mathcal{C}_{\theta_2, t}^m(\bar{g}) \subseteq \mathcal{C}_{\theta, t}^m(\bar{g}),$$

with $1 - \theta^{1/n} = \lambda_1(1 - \theta_1^{1/n}) + \lambda_2(1 - \theta_2^{1/n})$.

Proof. Let $\bar{x} \in \mathcal{C}_{\theta_1, t}^m(\bar{g})$, $\bar{y} \in \mathcal{C}_{\theta_2, t}^m(\bar{g})$, $z_1 \in \mathcal{A}_t^m(\bar{g})(\bar{x})$ and $z_2 \in \mathcal{A}_t^m(\bar{g})(\bar{y})$. Let also $z_0 \in \mathcal{A}_t^m(\bar{g})(\bar{o})$, then we have that

$$\begin{aligned} \prod_{i=0}^m g_i(z_0)^m &\geq t \prod_{i=0}^m \|g_i\|_{L^\infty(\mathbb{R}^n)}, \\ g_0(z_1) \prod_{i=1}^m g_i(z_1 - x_i) &\geq t \prod_{i=0}^m \|g_i\|_{L^\infty(\mathbb{R}^n)}, \quad \text{and} \\ g_0(z_2) \prod_{i=1}^m g_i(z_2 - y_i) &\geq t \prod_{i=0}^m \|g_i\|_{L^\infty(\mathbb{R}^n)}. \end{aligned}$$

Moreover, since the g_i are log-concave functions, we deduce that

$$\begin{aligned} &g_0((1 - \lambda_1 - \lambda_2)z_0 + \lambda_1 z_1 + \lambda_2 z_2) \\ &\quad \times \prod_{i=1}^m g_i((1 - \lambda_1 - \lambda_2)z_0 - \lambda_1(z_1 - x_i) - \lambda_2(z_2 - y_i)) \\ &\geq \left(\prod_{i=0}^m g_i(z_0) \right)^{(1 - \lambda_1 - \lambda_2)} \left(g_0(z_1) \prod_{i=1}^m g_i(z_1 - x_i) \right)^{\lambda_1} \left(g_0(z_2) \prod_{i=1}^m g_i(z_2 - y_i) \right)^{\lambda_2} \\ &\geq t \prod_{i=0}^m \|g_i\|_{L^\infty(\mathbb{R}^n)}. \end{aligned}$$

Therefore,

$$\begin{aligned} \mathcal{A}_t^m(\bar{g})(\lambda_1 \bar{x} + \lambda_2 \bar{y}) &\supseteq (1 - \lambda_1 - \lambda_2) \mathcal{A}_t^m(\bar{g})(\bar{o}) \\ &\quad + \lambda_1 \mathcal{A}_t^m(\bar{g})(\bar{x}) + \lambda_2 \mathcal{A}_t^m(\bar{g})(\bar{y}). \end{aligned}$$

Using the Brunn-Minkowski inequality, we then obtain

$$\begin{aligned} \text{vol}_n(\mathcal{A}_t^m(\bar{g})(\lambda_1 \bar{x} + \lambda_2 \bar{y}))^{1/n} &\geq (1 - \lambda_1 - \lambda_2) \text{vol}_n(\mathcal{A}_t^m(\bar{g})(\bar{o}))^{1/n} \\ &\quad + \lambda_1 \text{vol}_n(\mathcal{A}_t^m(\bar{g})(\bar{x}))^{1/n} + \lambda_2 \text{vol}_n(\mathcal{A}_t^m(\bar{g})(\bar{y}))^{1/n}. \end{aligned}$$

Thus, taking into account that $\bar{x} \in C_{\theta_1, t}^m(\bar{g})$, $\bar{y} \in C_{\theta_2, t}^m(\bar{g})$, we deduce that

$$\begin{aligned} \text{vol}_n(\mathcal{A}_t^m(\bar{g})(\lambda_1\bar{x} + \lambda_2\bar{y}))^{1/n} &\geq (1 - \lambda_1 - \lambda_2)M_t^{1/n} + \lambda_1\theta_1M_t^{1/n} + \lambda_2\theta_2M_t^{1/n} \\ &= (1 - \lambda_1(1 - \theta_1^{1/n}) - \lambda_2(1 - \theta_2^{1/n}))M_t^{1/n}, \end{aligned}$$

concluding the proof. \square

Remark 4.1. Notice that from Lemma 4.1 trivially follows that, for every $0 \leq \theta_0 \leq \theta < 1$,

$$(4.1) \quad \frac{C_{\theta_0, t}^m(\bar{g})}{1 - \theta_0^{1/n}} \subseteq \frac{C_{\theta, t}^m(\bar{g})}{1 - \theta^{1/n}}.$$

We are now finally ready to prove the main results of this section. We start by providing the inequality for $m + 1$ functions.

Proof of Theorem 1.2. By the translation invariance of the Lebesgue measure and the homogeneity of the inequality, we may translate and dilate the functions g_i so that $M_t = \text{vol}_n(\mathcal{A}_t(\bar{g})(\bar{o}))$ for all t and $\|g_i\|_{L^\infty(\mathbb{R}^n)} = 1$ for all $i = 0, 1, \dots, m$. We write $C_{\theta, t}^m = C_{\theta, t}^m(\bar{g})$. Moreover, from (4.1) follows that $(1 - \theta^{1/n})C_{0, t}^m \subset C_{\theta, t}^m$ for all $\theta \in [0, 1)$. Thus, we have that

$$\int_0^1 (1 - \theta^{1/n})^{nm} \text{vol}_{nm}(C_{0, t}^m) d\theta \leq \int_0^1 \text{vol}_{nm}(C_{\theta, t}^m) d\theta.$$

Therefore,

$$\begin{aligned} \text{vol}_{nm}(C_{0, t}^m) &\leq \binom{n(m+1)}{n} \int_0^1 \text{vol}_{nm}(C_{\theta, t}^m) d\theta \\ &= \binom{n(m+1)}{n} \int_{\mathbb{R}^{nm}} \frac{\text{vol}_n(\mathcal{A}_t^m(\bar{g})(\bar{x}))}{M_t} d\bar{x}. \end{aligned}$$

Now, on the one hand, integrating with respect to the variable t , we deduce that

$$\begin{aligned} \int_0^1 M_t \text{vol}_{nm}(C_{0, t}^m) dt &\leq \binom{n(m+1)}{n} \int_0^1 \int_{\mathbb{R}^{nm}} \text{vol}_n(\mathcal{A}_t^m(\bar{g})(\bar{x})) d\bar{x} dt \\ &= \binom{n(m+1)}{n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^{nm}} g_0(z) \prod_{i=1}^m g_i(z - x_i) d\bar{x} dz \\ &= \binom{n(m+1)}{n} \left(\prod_{i=0}^m \int_{\mathbb{R}^n} g_i(x) dx \right). \end{aligned}$$

On the other hand,

$$\begin{aligned} \int_0^1 M_t \operatorname{vol}_{nm}(\mathcal{C}_{0,t}^m) dt &= \int_{\mathbb{R}^{nm}} \int_0^{(\bar{g})_{\star m}(\bar{x})} \max_{\bar{y}} \operatorname{vol}_n(\mathcal{A}_t^m(\bar{g})(\bar{y})) dt d\bar{x} \\ &\geq \max_{\bar{y}} \int_{\mathbb{R}^{nm}} \int_0^{(\bar{g})_{\star m}(\bar{x})} \operatorname{vol}_n(\mathcal{A}_t^m(\bar{g})(\bar{y})) dt d\bar{x} \\ &\geq \max_{\bar{y}} \int_{(\mathbb{R}^n)^{m+1}} \min \left\{ (\bar{g})_{\star m}(\bar{x}), g_0(z) \prod_{i=1}^m g_i(z - y_i) \right\} dz d\bar{x}, \end{aligned}$$

where we used the definition of $\mathcal{A}_t^m(\bar{g})(\bar{y})$. Our inequality continues:

$$\begin{aligned} \int_0^1 M_t \operatorname{vol}_{nm}(\mathcal{C}_{0,t}^m) dt &\geq \max_{\bar{y}} \int_{\mathbb{R}^n} g_0(z) \prod_{i=1}^m g_i(z - y_i) dz \int_{\mathbb{R}^{nm}} (\bar{g})_{\star m}(\bar{x}) d\bar{x} \\ &= \|(g)_{\oplus m}\|_{\infty} \int_{\mathbb{R}^{nm}} (\bar{g})_{\star m}(\bar{x}) d\bar{x}, \end{aligned}$$

therefore concluding the proof. \square

Next, we prove the inequality for a function and its reflection.

Proof of Theorem 1.3. Let $\bar{x} = (x_1, \dots, x_m) \in \mathbb{R}^{nm}$. Since f is log-concave, we have

$$\begin{aligned} &f\left(z - \frac{1}{m+1}(x_1 + \dots + x_m)\right) \\ &= f\left(\frac{1}{m+1}[z + (z - x_1) + \dots + (z - x_m)]\right) \geq \prod_{i=0}^m f(z - x_i)^{\frac{1}{m+1}}. \end{aligned}$$

Hence,

$$\begin{aligned} \mathcal{A}_t^m(f)(\bar{x}) &\subseteq \left\{ z \in \mathbb{R}^n : \frac{f(z - \frac{1}{m+1} \sum x_i)}{\|f\|_{L^\infty(\mathbb{R}^n)}} \geq t^{1/(m+1)} \right\} \\ &= \frac{1}{m+1} \sum_{i=1}^m x_i + \mathcal{A}_t^m(f)(\bar{o}). \end{aligned}$$

Moreover,

$$\begin{aligned} \mathcal{A}_t^m(f)(\bar{x}) &\supset \mathcal{A}_t^m(f)(\bar{o}) \cap_{i=1}^m (x_i + \mathcal{A}_t^m(f)(\bar{o})) \quad \text{and} \\ \mathcal{A}_t^m(f)(\bar{x}) &\subset \mathcal{A}_{t^{m+1}}^m(f)(\bar{o}) \cap_{i=1}^m (x_i + \mathcal{A}_{t^{m+1}}^m(f)(\bar{o})). \end{aligned}$$

Thus,

$$\mathcal{C}_{0,t}^m \subset D^m(\mathcal{A}_{t^{m+1}}^m(f)(\bar{o})).$$

Consequently, using the higher-order Rogers-Shephard inequality (1.5) yields

$$\operatorname{vol}_{nm}(\mathcal{C}_{0,t}^m) \leq \binom{n(m+1)}{n} \operatorname{vol}_n(\mathcal{A}_{t^{m+1}}^m(f)(\bar{o}))^m.$$

Now, by Minkowski's inequality for integrals, we infer

$$\begin{aligned} \int_0^1 \text{vol}_n(\mathcal{A}_{t^{m+1}}^m(f)(\bar{o}))^m dt &= \int_0^1 \left(\int_{\mathbb{R}^n} \chi_{\{f/\|f\|_{L^\infty(\mathbb{R}^n)} \geq t\}}(z) dz \right)^m dt \\ &\leq \left(\int_{\mathbb{R}^n} \left(\int_0^1 \chi_{\{f/\|f\|_{L^\infty(\mathbb{R}^n)} \geq t\}}(z) dt \right)^{1/m} dz \right)^m \\ &= \frac{1}{\|f\|_{L^\infty(\mathbb{R}^n)}} \left(\int_{\mathbb{R}^n} f(z)^{1/m} dz \right)^m = \frac{\|f\|_{L^{\frac{1}{m}}(\mathbb{R}^n)}}{\|f\|_{L^\infty(\mathbb{R}^n)}}. \end{aligned}$$

Moreover, taking into account that $\mathcal{A}_t^m(f)(\bar{x}) \neq \emptyset$ if and only if

$$t \leq (\bar{f})_{\star m}(\bar{x})/\|f\|_{L^\infty(\mathbb{R}^n)}^{m+1},$$

we obtain

$$\begin{aligned} \int_0^1 \text{vol}_{nm}(\mathcal{C}_{0,t}^m) dt &= \int_0^1 \int_{\mathbb{R}^{nm}} \chi_{\{\bar{x}: \mathcal{A}_t^m(f)(\bar{x}) \neq \emptyset\}} d\bar{x} dt \\ &= \int_{\mathbb{R}^{nm}} \int_0^1 \chi_{\{t: \mathcal{A}_t^m(f)(\bar{x}) \neq \emptyset\}} dt d\bar{x} \\ &= \int_{\mathbb{R}^{nm}} \int_0^{(\bar{f})_{\star m}(\bar{x})/\|f\|_{L^\infty(\mathbb{R}^n)}^{m+1}} dt d\bar{x} \\ &= \frac{1}{\|f\|_{L^\infty(\mathbb{R}^n)}^{m+1}} \int_{\mathbb{R}^{nm}} (\bar{f})_{\star m}(\bar{x}) d\bar{x}, \end{aligned}$$

as desired. \square

Equality case on Theorem 1.3. Fix $m \in \mathbb{N} \setminus \{1\}$, as the $m = 1$ case is already established ([3, Lemma 5.1]). On the one hand, if f is the characteristic function of an n -dimensional simplex, (1.9) becomes the higher-order Rogers-Shephard inequality (1.5) for an n -dimensional simplex, which is precisely the equality case.

On the other hand, equality in (1.9) implies equality in the following:

$$\begin{aligned} \int_0^1 \text{vol}_{nm}(\mathcal{C}_{0,t}^m) dt &\leq \int_0^1 \text{vol}\left(D^m(\mathcal{A}_{t^{m+1}}^m(f)(\bar{o}))\right) dt \\ &\leq \int_0^1 \binom{n(m+1)}{n} \text{vol}_n(\mathcal{A}_{t^{m+1}}^m(f)(\bar{o})) dt. \end{aligned}$$

Therefore, there is equality in

$$\text{vol}_{nm}(\mathcal{C}_{0,t}^m) \leq \text{vol}\left(D^m(\mathcal{A}_{t^{m+1}}^m(f)(\bar{o}))\right) \leq \binom{n(m+1)}{n} \text{vol}_n(\mathcal{A}_{t^{m+1}}^m(f)(\bar{o}))^m.$$

Using again the equality conditions in (1.5) one has that, for every $t \in (0, 1)$, $\mathcal{A}_{t^{m+1}}^m(f)(\bar{o})$ is an n -dimensional simplex. We next use the fact that equality implies equality in our use of Minkowski's integral inequality. We obtain that the function $(z, t) \mapsto \chi_{\{f/\|f\|_{L^\infty(\mathbb{R}^n)} \geq t\}}(z)$ equals $h(t)q(z)$ for some measurable functions $h : (0, 1] \mapsto \mathbb{R}_+$ and $q : \mathbb{R}^n \mapsto \mathbb{R}_+$ almost everywhere;

notice that since f is integrable, the set $\{f = \|f\|_{L^\infty(\mathbb{R}^n)}\}$ must be compact, and, since f is log-concave, the super-level sets of f are convex and f is continuous when restricted to the interior of its support. We deduce that the equality holds on $\mathbb{R}^n \times (0, 1]$, in which case we must have $h(t)$ is a constant (merely pick a $z_0 \in \{f = \|f\|_{L^\infty(\mathbb{R}^n)}\}$ to obtain $1 = h(t)q(z_0)$ for all $t \in (0, 1]$). Additionally this means, for a fixed z , the value of $\chi_{\{f/\|f\|_{L^\infty(\mathbb{R}^n)} \geq t\}}(z)$ is the same for all t . In terms of f , this means if z is such that $f(z) \geq t\|f\|_{L^\infty(\mathbb{R}^n)}$ for some $t \in (0, 1] \rightarrow f(z) \geq t\|f\|_{L^\infty(\mathbb{R}^n)}$ for all $t \in (0, 1]$. We deduce that $f(z) = \|f\|_{L^\infty(\mathbb{R}^n)}$ for all such z . Likewise, if z is such that $f(z) < t\|f\|_{L^\infty(\mathbb{R}^n)}$ for some $t \in (0, 1] \rightarrow f(z) < t\|f\|_{L^\infty(\mathbb{R}^n)}$ for all $t \in (0, 1]$, and so $f(z) = 0$. Therefore, f partitions \mathbb{R}^n into two sets: a convex body upon which $f(z) = \|f\|_{L^\infty(\mathbb{R}^n)}$, and the rest of \mathbb{R}^n . In particular, the latter is such that $f(z) = 0$, i.e. f is the characteristic function of a convex body. We now use that, for $t \in (0, 1)$, $\mathcal{A}_{t^{m+1}}^m(f)(\bar{o}) = \{f/\|f\|_{L^\infty(\mathbb{R}^n)} \geq t\} = \{f = \|f\|_{L^\infty(\mathbb{R}^n)}\}$ to deduce that this convex body is an n -dimensional simplex. \square

Acknowledgements: This work was completed while all three authors were in-residence at the Hausdorff Research Institute for Mathematics in Bonn, Germany, during Spring 2024 for the Dual Trimester Program: "Synergies between modern probability, geometric analysis and stochastic geometry". The three authors would like to express their sincere gratitude to the Hausdorff Institute, its dedicated staff, and the organizers of the trimester program for their invaluable support and hospitality.

Funding: The first named author was supported by a fellowship from the FSMP Post-doctoral program. The second named author is funded by the grant PID2021-124157NB-I00, financed by MCIN/AEI/10.13039/501100011033/ "ERDF A way of making Europe", and by Comunidad Autónoma de la Región de Murcia a través de la convocatoria de Ayudas a proyectos para el desarrollo de investigación científica y técnica por grupos competitivos, incluida en el Programa Regional de Fomento de la Investigación Científica y Técnica (Plan de Actuación 2022) de la Fundación Séneca-Agencia de Ciencia y Tecnología de la Región de Murcia, REF. 21899/PI/22. The third named author was supported by the Austrian Science Fund (FWF): 10.55776/P34446, and, in part, by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

REFERENCES

- [1] ALONSO-GUTIÉRREZ, D., ARTSTEIN-AVIDAN, S., GONZÁLEZ MERINO, B., JIMÉNEZ, C. H., AND VILLA, R. Rogers-Shephard and local Loomis-Whitney type inequalities. *Math. Ann.* 374, 3-4 (2019), 1719–1771.
- [2] ALONSO-GUTIÉRREZ, D., BERNUÉS, J., AND GONZÁLEZ MERINO, B. Zhang's inequality for log-concave functions. In *Geometric aspects of functional analysis. Vol. I*, vol. 2256 of *Lecture Notes in Math.* Springer, Cham, [2020] ©2020, pp. 29–48.

- [3] ALONSO-GUTIÉRREZ, D., GONZÁLEZ MERINO, B., JIMÉNEZ, C. H., AND VILLA, R. Rogers-Shephard inequality for log-concave functions. *J. Funct. Anal.* *271*, 11 (2016), 3269–3299.
- [4] ALONSO-GUTIÉRREZ, D., GONZÁLEZ MERINO, B., JIMÉNEZ, C. H., AND VILLA, R. John’s ellipsoid and the integral ratio of a log-concave function. *J. Geom. Anal.* *28*, 2 (2018), 1182–1201.
- [5] ALONSO-GUTIÉRREZ, D., HERNÁNDEZ CIFRE, M. A., ROYSDON, M., YEPES NICOLÁS, J., AND ZVAVITCH, A. On Rogers-Shephard type inequalities for general measures. *Int. Math. Res. Not. IMRN*, 10 (2021), 7224–7261.
- [6] ALVES DA SILVA, L., GONZÁLEZ MERINO, B., AND VILLA, R. Some remarks on petty projection of log-concave functions. *J. Geom. Anal.* *33*, 8 (2023), Paper No. 260, 14.
- [7] ALVINO, A., FERONE, V., TROMBETTI, G., AND LIONS, P.-L. Convex symmetrization and applications. *Ann. Inst. H. Poincaré C Anal. Non Linéaire* *14*, 2 (1997), 275–293.
- [8] AMBROSIO, L., FUSCO, N., AND PALLARA, D. *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2000.
- [9] ARTSTEIN-AVIDAN, S., EINHORN, K., FLORENTIN, D. I., AND OSTROVER, Y. On Godbersen’s conjecture. *Geom. Dedicata* *178* (2015), 337–350.
- [10] ARTSTEIN-AVIDAN, S., KLARTAG, B., AND MILMAN, V. The Santaló point of a function, and a functional form of the Santaló inequality. *Mathematika* *51*, 1-2 (2004), 33–48 (2005).
- [11] BALL, K. Logarithmically concave functions and sections of convex sets in \mathbf{R}^n . *Studia Math.* *88*, 1 (1988), 69–84.
- [12] BARTHE, F., BÖRÖCZKY, K. J., AND FRADELIZI, M. Stability of the functional forms of the Blaschke-Santaló inequality. *Monatsh. Math.* *173*, 2 (2014), 135–159.
- [13] BIANCHI, G. The covariogram problem. In *Harmonic Analysis and Convexity*, A. Koldobsky and A. Volberg, Eds., Adv. Anal. Geom. De Gruyter, 2023, pp. 37–82.
- [14] BRASCAMP, H. J., AND LIEB, E. H. On extensions of the Brunn-Minkowski and Prékopa-Leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation. *J. Functional Analysis* *22*, 4 (1976), 366–389.
- [15] BRUCKNER, A. *Differentiation of real functions*, second ed., vol. 5 of *CRM Monograph Series*. American Mathematical Society, Providence, RI, 1994.
- [16] COLESANTI, A. Functional inequalities related to the Rogers-Shephard inequality. *Mathematika* *53*, 1 (2006), 81–101.
- [17] COLESANTI, A., AND FRAGALÀ, I. The first variation of the total mass of log-concave functions and related inequalities. *Adv. Math.* *244* (2013), 708–749.
- [18] COLESANTI, A., LUDWIG, M., AND MUSSNIG, F. Minkowski valuations on convex functions. *Calc. Var. Partial Differential Equations* *56*, 6 (2017), Paper No. 162, 29.
- [19] CORDERO-ERAUSQUIN, D., AND KLARTAG, B. Moment measures. *J. Funct. Anal.* *268*, 12 (2015), 3834–3866.
- [20] CORDERO-ERAUSQUIN, D., NAZARET, B., AND VILLANI, C. A mass-transportation approach to sharp Sobolev and Gagliardo-Nirenberg inequalities. *Adv. Math.* *182*, 2 (2004), 307–332.
- [21] DUBUC, S. Critères de convexité et inégalités intégrales. *Ann. Inst. Fourier (Grenoble)* *27*, 1 (1977), x, 135–165.
- [22] EVANS, L. C. *Partial Differential Equations*, 2nd ed., vol. 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2010.
- [23] FRADELIZI, M., GORDON, Y., MEYER, M., AND REISNER, S. The case of equality for an inverse Santaló functional inequality. *Adv. Geom.* *10*, 4 (2010), 621–630.
- [24] FRADELIZI, M., GOZLAN, N., AND ZUGMEYER, S. Transport proofs of some functional inverse Santaló inequalities. In *High dimensional probability IX—the ethereal volume*, vol. 80 of *Progr. Probab.* Birkhäuser/Springer, Cham, [2023] ©2023, pp. 123–142.

- [25] FRADELIZI, M., LI, J., AND MADIMAN, M. Concentration of information content for convex measures. *Electron. J. Probab.* 25 (2020), Paper No. 20, 22.
- [26] FRADELIZI, M., AND MEYER, M. Some functional forms of Blaschke-Santaló inequality. *Math. Z.* 256, 2 (2007), 379–395.
- [27] FRADELIZI, M., AND MEYER, M. Increasing functions and inverse Santaló inequality for unconditional functions. *Positivity* 12, 3 (2008), 407–420.
- [28] FRADELIZI, M., AND MEYER, M. Some functional inverse Santaló inequalities. *Adv. Math.* 218, 5 (2008), 1430–1452.
- [29] GARDNER, R. J. The Brunn-Minkowski inequality. *Bull. Amer. Math. Soc. (N.S.)* 39, 3 (2002), 355–405.
- [30] GARDNER, R. J., AND ZHANG, G. Affine inequalities and radial mean bodies. *Amer. J. Math.* 120, 3 (1998), 505–528.
- [31] HADDAD, J., LANGHARST, D., PUTTERMAN, E., ROYSDON, M., AND YE, D. Affine isoperimetric inequalities for higher-order projection and centroid bodies. *arXiv:2304.07859* (2023).
- [32] HADDAD, J., LANGHARST, D., PUTTERMAN, E., ROYSDON, M., AND YE, D. General higher order lp isoperimetric and sobolev inequalities. *arXiv:2305.17468* (2023).
- [33] HADDAD, J. E., AND LUDWIG, M. Affine fractional sobolev and isoperimetric inequalities. *Preprint, arxiv: 2207.06375* (2022).
- [34] HUANG, Y., LIU, J., XI, D., AND ZHAO, Y. Dual curvature measures for log-concave functions. *J. Differential Geom.* (2024+).
- [35] KLARTAG, B., AND MILMAN, V. D. Geometry of log-concave functions and measures. *Geom. Dedicata* 112 (2005), 169–182.
- [36] KOLDOBSKY, A., PAJOR, A., AND YASKIN, V. Inequalities of the Kahane-Khinchin type and sections of L_p -balls. *Studia Math.* 184, 3 (2008), 217–231.
- [37] LANGHARST, D., AND PUTTERMAN, E. Weighted berwald’s inequality. *To appear in Amer. J. Math.* (2025).
- [38] LANGHARST, D., ROYSDON, M., AND ZVAVITCH, A. General measure extensions of projection bodies. *Proc. Lond. Math. Soc. (3)* 125, 5 (2022), 1083–1129.
- [39] LEHEC, J. A direct proof of the functional Santaló inequality. *C. R. Math. Acad. Sci. Paris* 347, 1-2 (2009), 55–58.
- [40] LEHEC, J. Partitions and functional Santaló inequalities. *Arch. Math. (Basel)* 92, 1 (2009), 89–94.
- [41] LEINDLER, L. On a certain converse of Hölder’s inequality. II. *Acta Sci. Math. (Szeged)* 33, 3-4 (1972), 217–223.
- [42] LIN, Y. The Petty projection inequality for sets of finite perimeter. *Calc. Var. Partial Differential Equations* 60, 5 (2021), Paper No. 196, 18.
- [43] LUDWIG, M. Valuations on function spaces. *Adv. Geom.* 11, 4 (2011), 745–756.
- [44] LUDWIG, M. Valuations on Sobolev spaces. *Amer. J. Math.* 134, 3 (2012), 827–842.
- [45] LUTWAK, E., YANG, D., AND ZHANG, G. Optimal Sobolev norms and the L^p Minkowski problem. *Int. Math. Res. Not.* (2006), Art. ID 62987, 21.
- [46] MARÍN SOLA, F. On general concavity extensions of Grünbaum type inequalities. *Bull. Braz. Math. Soc. (N.S.)* 55, 1 (2024), Paper No. 2, 17.
- [47] MATHERON, G. *Random Sets and Integral Geometry*. Wiley Series in Probability and Statistics. Wiley, Michigan, 1975.
- [48] MEYER, M., NAZAROV, F., RYABOGIN, D., AND YASKIN, V. Grünbaum-type inequality for log-concave functions. *Bulletin of the London Mathematical Society* 50 (2018), 745–752.
- [49] MILMAN, V. D., AND SCHECHTMAN, G. *Asymptotic theory of finite-dimensional normed spaces*, vol. 1200 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1986. With an appendix by M. Gromov.
- [50] MYROSHNYCHENKO, S., STEPHEN, M., AND ZHANG, N. Grünbaum’s inequality for sections. *J. Funct. Anal.* 275, 9 (2018), 2516–2537.

- [51] PETTY, C. M. Isoperimetric problems. In *Proceedings of the Conference on Convexity and Combinatorial Geometry (Univ. Oklahoma, Norman, Okla., 1971)*. Dept. Math., Univ. Oklahoma, Norman, Oklahoma, 1971, pp. 26–41.
- [52] PRÉKOPA, A. Logarithmic concave measures with application to stochastic programming. *Acta Sci. Math. (Szeged)* 32 (1971), 301–316.
- [53] PRÉKOPA, A. On logarithmic concave measures and functions. *Acta Sci. Math. (Szeged)* 34 (1973), 335–343.
- [54] ROGERS, C. A., AND SHEPHARD, G. C. The difference body of a convex body. *Arch. Math. (Basel)* 8 (1957), 220–233.
- [55] ROTEM, L. Surface area measures of log-concave functions. *J. Anal. Math.* 147, 1 (2022), 373–400.
- [56] ROTEM, L. The anisotropic total variation and surface area measures. In *Geometric aspects of functional analysis*, vol. 2327 of *Lecture Notes in Math.* Springer, Cham, [2023] ©2023, pp. 297–312.
- [57] ROYSDON, M. Rogers-Shephard type inequalities for sections. *J. Math. Anal. Appl.* 487, 1 (2020), 123958.
- [58] SCHNEIDER, R. Eine Verallgemeinerung des Differenzenkörpers. *Monatsh. Math.* 74 (1970), 258–272.
- [59] SCHNEIDER, R. *Convex Bodies: the Brunn-Minkowski Theory*, 2nd expanded ed., vol. 151 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, Cambridge, UK, 2014.
- [60] ULIVELLI, J. First variation of functional wulff shapes. *preprint* (2024), arXiv:2312.11172.
- [61] WANG, T. The affine Sobolev-Zhang inequality on $BV(\mathbb{R}^n)$. *Adv. Math.* 230, 4-6 (2012), 2457–2473.
- [62] WILLSON, S. J. A semigroup on the space of compact convex bodies. *SIAM J. Math. Anal.* 11, 3 (1980), 448–457.
- [63] YANG, D. Affine integral geometry from a differential viewpoint. In *Handbook of Geometric Analysis, No. 2, Advanced Lectures in Mathematics*. 2010, pp. 359–390.
- [64] ZHANG, G. Restricted chord projection and affine inequalities. *Geom. Dedicata* 39, 2 (1991), 213–222.
- [65] ZHANG, G. The affine Sobolev inequality. *J. Differential Geom.* 53, 1 (1999), 183–202.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU, SORBONNE UNIVERSITÉ, PARIS 75252, FRANCE

Email address: `dylan.langharst@imj-prg.fr`

Email address: `dlanghar@kent.edu`

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE MURCIA, CAMPUS DE ESPINARDO, 30100-MURCIA, SPAIN

Email address: `francisco.marin7@um.es`

INSTITUT FÜR DISKRETE MATHEMATIK UND GEOMETRIE, TU WIEN, WIEDNER HAUPTSTRASSE 8-10, E104-6, 1040 WIEN, AUSTRIA

Email address: `jacopo.ulivelli@tuwien.ac.at`