

HAMILTONICITY, PATH COVER, AND INDEPENDENCE NUMBER: AN FPT PERSPECTIVE*

Fedor V. Fomin[†]
fedor.fomin@uib.no

Petr A. Golovach[†]
petr.golovach@uib.no

Danil Sagunov[‡]
danilka.pro@gmail.com

Kirill Simonov[§]
kirillsimonov@gmail.com

Abstract

The connection between Hamiltonicity and the independence numbers of graphs has been a fundamental aspect of Graph Theory since the seminal works of the 1960s. This paper presents a novel algorithmic perspective on these classical problems. Our contributions are twofold.

First, we establish that a wide array of problems in undirected graphs, encompassing problems such as Hamiltonian Path and Cycle, Path Cover, Largest Linkage, and Topological Minor Containment are fixed-parameter tractable (FPT) parameterized by the independence number of a graph. To the best of our knowledge, these results mark the first instances of FPT problems for such parameterization.

Second, we extend the algorithmic scope of the Gallai-Milgram theorem. The original theorem by Gallai and Milgram, asserts that for a graph G with the independence number $\alpha(G)$, the vertex set of G can be covered by at most $\alpha(G)$ vertex-disjoint paths. We show that determining whether a graph can be covered by fewer than $\alpha(G) - k$ vertex-disjoint paths is FPT parameterized by k .

Notably, the independence number parameterization, which describes graph's density, departs from the typical flow of research in parameterized complexity, which focuses on parameters describing graph's sparsity, like treewidth or vertex cover.

1 Introduction

A significant portion of parameterized algorithms deals with optimization problems on sparse graphs when specific graph parameters characterize the sparsity and structure of the graph. Such parameters include treewidth, treedepth, or minimum vertex cover size. However, the parameterized complexity of many problems needs to be understood more when it comes to the parameters describing the graph's density, such as its independence number, that is, the size of the maximum independent set.

The initial step in exploring algorithmic properties of graphs with bounded independence numbers was taken by Ovetsky Fradkin and Seymour in their work [FS15]. They demonstrated that a generalization of the edge-disjoint path problem, which they call “fusion”, can be solved in polynomial time on digraphs with a constant independence number. The work of Ovetsky Fradkin and

*The research leading to these results has been supported by the Research Council of Norway via the project BWCA (grant no. 314528) and DFG Research Group ADYN via grant DFG 411362735.

[†]Department of Informatics, University of Bergen, Norway.

[‡]St. Petersburg State University, Russia.

[§]Hasso Plattner Institute, University of Potsdam, Germany.

Seymour, as well as the follow-up research [LLM⁺20, MSSZ23], indicates that the class of digraphs with bounded independence numbers exhibits sufficient structure for algorithmic exploitation.

In this paper, we leverage the structure of undirected graphs with bounded independence numbers to develop parameterized algorithms. Our approach is applicable to a wide range of combinatorial problems, including Hamiltonicity and path covers in undirected graphs. The interplay between Hamiltonicity and independence numbers is a fundamental theme in Graph Theory, stemming from the classic theorem of Gallai–Milgram [GM60]. Surprisingly, the computational complexity of the Graph Hamiltonicity problem remained unresolved, even for graphs with an independence number of $\alpha(G) = 3$, until the recent work by Jedlicková and Kratochvíl [JK23]. Jedlicková and Kratochvíl have demonstrated that not only Graph Hamiltonicity but also more general Hamiltonian- ℓ -Linkage problems (which involve connecting ℓ pairs of vertices with vertex-disjoint paths covering all the graph’s vertices) can be solved in polynomial time when the independence number $\alpha(G)$ of a graph is constant.

The algorithms proposed by Jedlicková and Kratochvíl for Graph Hamiltonicity is an XP algorithm for parameter $\alpha(G)$. This means that on an n -vertex graph G , the running times of these algorithms are the form of $n^{f(\alpha(G))}$ for some function f . To the best of our knowledge, no fundamental NP-hard problem has been found to be FPT parameterized by $\alpha(G)$. (The only trivial exception is coloring a graph G in a constant number of colors q because every yes-instance has at most $q \cdot \alpha(G)$ vertices.) This leads Jedlicková and Kratochvíl in [JK23] to an intriguing question: *What is the parameterized complexity of the Hamiltonian Path problem (and many other related problems) parameterized by $\alpha(G)$?*

We find the situation concerning parameterization by $\alpha(G)$ unusual: Most algorithmic findings in structural parameterization require that computing or approximating the corresponding graph structural parameter is in FPT. For example, deciding whether a graph contains a Hamiltonian cycle is FPT parameterized by the treewidth of a graph due to the following reasoning. First, for an input integer k , there are (either exact or approximation) algorithms that either correctly decide that the treewidth of the input graph is more than k or construct a tree decomposition of width $\mathcal{O}(k)$ in time $f(k) \cdot n^{\mathcal{O}(1)}$ [Bod96, Kor21, RS86]. Then, the tree decomposition is used to decide whether the graph is Hamiltonian in time $g(k) \cdot n^{\mathcal{O}(1)}$. In contrast, determining whether $\alpha(G) \geq k$ is W[1]-hard with parameter k [DF99]. Moreover, there is compelling evidence that the problem does not admit an FPT approximation [LRSW22, KK22]. To overcome the intractability of the independence number, we design algorithms with the following properties. For a given parameter k , the algorithm either solves an optimization problem at hand (Hamiltonicity, Path Cover, etc.) in FPT time with respect to k , or it correctly reports that the input graph does not fall within the appropriate domain, meaning it has an independence number greater than k . Such algorithms are what Raghavan and Spinrad referred to as *robust* [RS03].

Our contribution. To state formally our first main result, we need the following definitions.

Definition 1 (TM-model and TM-embedding). Let H, G be graphs, let M be a subgraph of G and $T \subseteq V(M) \subseteq V(G)$, such that all vertices of $M \setminus T$ have degree 2. We say that (M, T) is a *topological minor model* (TM-model) of H in G if H is isomorphic to a graph obtained from M by dissolving in M all vertices of $M \setminus T$. Here, *dissolving* a vertex of degree two means deleting it while making its neighbors adjacent. We further say that for the injective mapping $f : V(H) \rightarrow V(G)$ induced by the isomorphism above, (M, f) is a *topological minor embedding* (TM-embedding) of H in G . The *size* of a TM-model (M, T) and of a TM-embedding (M, f) is the number of vertices in M .

In other words, in a TM-embedding, the vertices of H are mapped to distinct vertices of G such that the edges of H could be mapped to internally vertex-disjoint paths of G connecting the

corresponding pairs of vertices of G . Moreover, the mapping from edges of H to paths in G can be “read off” from the corresponding model subgraph M .

We also consider a more general variant of TM-embedding. In this variant, every vertex of H is assigned a list of vertices of G where it could be mapped.

Definition 2 (List TM-embedding). For graphs H, G , and a mapping $L : V(H) \rightarrow 2^{V(G)}$, we say that TM-embedding $M, f : V(H) \rightarrow V(G)$ is a *list topological minor embedding* (list TM-embedding), if for each $h \in V(H)$, $f(h) \in L(h)$.

Our first result is the theorem establishing that computing the list TM-embedding of maximum size is FPT parameterized by the size of the maximum independent set in G and the size of the graph H .

Theorem 1. *There is an algorithm that, given graphs H, G , a list assignment $L : V(H) \rightarrow 2^{V(G)}$ and an integer parameter $k \geq 1$, in time $2^{|H|^{\mathcal{O}(k)}} \cdot 2^{k^{\mathcal{O}(k^2)}} \cdot |G|^{\mathcal{O}(1)}$*

- *either correctly reports that there is no list TM-embedding of H in G , or*
- *computes a TM-embedding of H in G of the maximum size, where the maximum is taken over all list TM-embeddings of H in G , or*
- *outputs an independent set in G of size k .*

List TM-embedding encompasses several vital problems in graph algorithms and Theorem 1 implies several corollaries.

- When graph H consists of a path P_2 on two vertices whose lists are all vertices of $V(G)$, then this is the problem of finding a longest path in G . If this path contains $|V(G)|$ vertices, this is a Hamiltonian path. For $H = C_3$, we get the problem of finding a longest cycle in G , the generalization of computing a Hamiltonian cycle in G . By Theorem 1, these tasks could be performed in time $f(\alpha(G)) \cdot |G|^{\mathcal{O}(1)}$. This positively answers the open question from [JK23].
- In PATH COVER, the task is to cover all vertices of G by at most p vertex-disjoint paths. By Gallai–Milgram’s theorem [GM60], if $p \geq \alpha(G)$ then a covering by at most p paths always exists and we can assume that $p < \alpha(G)$. We can solve the problem by taking H to be disjoint unions of at most p paths P_1 and P_2 assigning to each vertex of H the list containing all vertices of $V(G)$. Thus, Theorem 1 yields an algorithm of running time $f(\alpha(G)) \cdot |G|^{\mathcal{O}(1)}$ solving this problem. For CYCLE COVER, that is when the graph H consists of at most p disjoint copies of C_3 , we have the running time of $f(\alpha(G) + p) \cdot |G|^{\mathcal{O}(1)}$.
- Jedlicková and Kratochvíl in [JK23] defined the following generalization of PATH COVER. In the HAMILTONIAN- ℓ -LINKAGE problem, we are given ℓ pairs of vertices $(s_1, t_1), \dots, (s_\ell, t_\ell)$. The task is to connect these pairs by disjoint paths that altogether traverse all vertices of the graph. For $\ell = 1$ this is the problem of deciding whether a graph for selected vertices s and t contains a Hamiltonian s, t -path. HAMILTONIAN- ℓ -LINKAGE could be encoded as List TM-embedding with H consisting of ℓ disjoint copies of P_2 . Let the vertices of these P_2 ’s be $\{x_i, y_i\}$, $i \in \{1, \dots, \ell\}$. Then the list mapping L assigns $\{s_i\}$ to x_i and $\{t_i\}$ to y_i . By Theorem 1, HAMILTONIAN- ℓ -LINKAGE is solvable in time $f(\alpha(G) + \ell) \cdot |G|^{\mathcal{O}(1)}$, resolving another open question of Jedlicková and Kratochvíl. By playing with different graphs H in the List TM-embedding and Theorem 1, one easily obtains algorithms computing spanning subgraphs with various properties like a spanning tree with at most ℓ leaves.

- Another interesting scenario is the maximization variant of the T -CYCLE problem. In the LONGEST T -CYCLE problem, we are given a graph G and a set $T \subseteq V(G)$ of terminals. The task is to decide whether there is a cycle passing through all terminals [BHT12, Kaw08, Wah13]. We can reduce this problem to List TM-embedding by enumerating all $|T|!$ orderings in which terminal vertices occur in an optimal solution. For each guessed ordering, we apply Theorem 1 for H being a cycle on $|T|$ vertices such that each vertex of H is assigned to precisely one terminal vertex, and the cyclic ordering of H corresponds to the guess. This brings us to the $f(\alpha(G) + |T|) \cdot |G|^{\mathcal{O}(1)}$ algorithm computing a cycle of maximum length containing all T terminals. Similar arguments hold for a more general colored variant of the problem. In this variant, we are given a graph G , and some of the vertices of G are assigned colors from 1 to L . Then, the task is to find a longest cycle containing the maximum number of colors [FGK⁺23]. Note that an optimal cycle can contain many vertices of the same color. By guessing the colors participating in a solution cycle C and the ordering of their first appearance in some cyclic ordering of C , we construct a cycle H on at most L vertices. Every vertex of H is assigned the list of vertices of G corresponding to the guessed color. This brings us to the algorithm of running time $f(\alpha(G) + L) \cdot |G|^{\mathcal{O}(1)}$.

The Gallai–Milgram theorem from 1960 [GM60] states that the vertex set of every (directed or undirected) graph can be covered by at most $\alpha(G)$ vertex-disjoint paths. Known constructive proofs of the Gallai–Milgram theorem can be used to construct a graph path cover with a size of at most $\alpha(G)$. Our following theorem presents a significant generalization of this algorithmic result for undirected graphs: for an integer parameter k , deciding whether a graph G contains a path cover of size at most $\alpha(G) - k$ is FPT, parameterized by k .

One of the corollaries of Theorem 1 is that deciding whether a graph G contains a Hamiltonian path is FPT, parameterized by $\alpha(G)$. It is worth noting that Theorem 2 provides another non-trivial extension of this result as we solve this problem by taking $k = \alpha(G) - 1$ in the theorem.

Theorem 2. *There is an algorithm that, given an n -vertex graph G and an integer parameter $k \geq 1$, in time $2^{k^{\mathcal{O}(k^4)}} \cdot n^{\mathcal{O}(1)}$ outputs a path cover \mathcal{P} of G and, furthermore,*

- *either correctly reports that \mathcal{P} is a minimum-size path cover, or*
- *outputs together with \mathcal{P} an independent set of size $|\mathcal{P}| + k$ certifying that \mathcal{P} contains at most $\alpha(G) - k$ paths.*

Related Work. The exploration of long paths and cycles in graphs represents a well-established and actively pursued research direction within parameterized algorithms. Since the seminal introduction of the color-coding technique by Alon, Yuster, and Zwick [AYZ95], substantial progress in algorithmic methodologies has been achieved. Notably, a range of powerful algorithmic methods has been developed, including those by Bjorklund [Bjö14], Koutis [Kou08], Williams [Wil09], and Zehavi [Zeh16]. This progress has also contributed to our understanding of significantly more general and complex problems, such as topological minor embedding [GKMW11, FLP⁺20].

This progress also concerns structural parameterized complexity, especially in graphs of bounded treewidth [CNP⁺11, BCKN13, CKN13, FLPS16, BST23]. However, the success of treewidth parameterization does not extend to more general graph width parameters. For example, the problem of deciding graph Hamiltonicity is W[1]-hard when parameterized by the clique-width of a graph [FGLS10]. For a comprehensive overview and further details on the algorithms for finding long cycles and paths in graphs, the reader may refer to Chapters 10, 11, and 12 of the book by Cygan et al. [CFK⁺15].

The interplay of the independence number, connectivity, Hamiltonicity, and path cover is a popular theme in Graph Theory, with a historical foundation dating back to the works of Dilworth [Dil50], Gallai and Milgram [GM60], Nash-Williams [NW71], and Chvátal and Erdős [CE72]. An excellent introduction to this aspect of Graph Theory can be found in the survey by Jackson and Ordaz [JO90]. It is surprising that in graph algorithms, despite tremendous success in the study of Hamiltonicity, connectivity, and topological minors on graphs of bounded treewidth, which are sparse graphs, there has been almost no progress on graphs of bounded independence number. The notable exceptions are the works of Ovetsky Fradkin and Seymour [FS15] and Jedlicková and Kratochvíl [JK23]. The first work deals with the edge-disjoint path problem, and the second, most relevant to our first theorem, focuses on Graph Hamiltonicity and Hamiltonian- ℓ -Linkage problems. In the end of this section, we explain the most crucial differences between the approaches taken in our work and the work of Jedlicková and Kratochvíl. Both papers by Ovetsky Fradkin and Seymour and Jedlicková and Kratochvíl provide XP algorithms that run in polynomial time when the independence number of a graph is constant.

Theorem 1 could also be treated as a contribution to the rapidly developing area of algorithms on graphs, excluding some induced subgraphs like P_t or C_t [ACP⁺21, GLP⁺21]. (Every graph with an independence number of k excludes the graph kK_1 as an induced subgraph.) However, all results in this area concern the computation of the maximum independent set or, more generally, induced subgraphs of bounded treewidth. Hamiltonicity is a more complex problem; it becomes NP-complete already on P_5 -free graphs (a superclass of split graphs) [BLS99].

Theorem 2 contributes to another popular trend in parameterized algorithms that explores the algorithmic extensions of classic theorems from extremal graph theory. Examples of such works include the algorithmic extensions of Dirac’s theorem by Fomin, Golovach, Sagunov, and Simonov [FGSS20] or Hajnal–Szemerédi’s theorem by Gan, Han and Hu [GHH23].

1.1 Proof overview

Theorem 1: We explain here the main steps of the proof only for Hamiltonian- ℓ -linkages. An extension to arbitrary list TM-embeddings is more technical but builds on similar ideas.

Let us start by sketching the XP algorithm of Jedlicková and Kratochvíl [JK23]. Their algorithm solves HAMILTONIAN- ℓ -LINKAGE and runs in $n^{f(\alpha(G)+\ell)}$ time. Since Theorem 1 is a strict improvement of the XP algorithm, we highlight crucial differences that allow us to achieve fixed-parameter tractability for the more general problem. An essential step in the algorithm of Jedlicková and Kratochvíl is the following lemma: If a graph G is $\Omega(\alpha(G) \cdot \ell)$ -vertex-connected, then it is Hamiltonian- ℓ -linked, meaning that a path cover of size ℓ exists with any given endpoints of the paths.

The algorithm of [JK23] is recursive and requires $k \geq \alpha(G)$ to be given in the input. (This requirement is not a problem for an XP algorithm because k could be computed by a brute force in time $\mathcal{O}(n^{\alpha(G)})$.) When the input graph G is $\Omega(k \cdot \ell)$ -connected, the algorithm correctly reports that G is Hamiltonian- ℓ -linked. Otherwise, when the connectivity is smaller, a minimum vertex separator X of G has $\mathcal{O}(k \cdot \ell)$ vertices. Every connected component of $G - X$ has an independence number at most $\alpha(G) - 1$, and the algorithm proceeds recursively on each component. For that, the algorithm performs an exhaustive search of how the solution paths are split into parts by X . For every connected component of $G - X$ and each split, the algorithm also performs an exhaustive search of entry and exit points of the parts of the paths in the component. Since the number of parts is at most $\ell + 2|X|$, this constitutes $n^{\mathcal{O}(\ell+|X|)}$ possible choices. By treating each set of entry and exit points as new terminal vertices, the algorithm is called recursively on each component. By restricting the depth of the recursion by $\alpha(G)$, the recursion reaches the subproblem where the

graph has sufficiently high connectivity without significantly blowing up the number of paths.

Since the approach of Jedlicková and Kratochvíl exploits small separators separating highly connected components, the first natural direction to explore is whether the generic tools about cuts from the toolbox of parameterized algorithms could be leveraged to turn it into an FPT algorithm. The recursive understanding technique introduced by Chitnis et al. [CCH⁺16] is arguably the most powerful tool for designing FPT algorithms for such types of problems about cuts, as illustrated by the meta-theorem of Lokshtanov et al. [LRSZ18]. Unfortunately, recursive understanding is not of much help here due to the following reasons. To apply recursive understanding, one has to be able to deal with the primary case of so-called (p, q) -unbreakable graphs. However, for our problem, this primary case is not more straightforward than the case of general graphs.

This is why we employ a different method of decomposing the graph. To prevent the XP running time, our algorithm avoids recursive calls to itself. Instead, it encapsulates the recursion in a subroutine that in time $f(k, \ell) \cdot n^{\mathcal{O}(1)}$ either output an independent set of G of size k or find a vertex set S of size $k^{\mathcal{O}(k)} \cdot \ell$ with the following property: Each connected component of $G - S$ is $\Omega(k \cdot (\ell + |S|))$ -connected. Set S breaks solution paths into at most $\ell + 2|S|$ parts. Some parts belong to $G[S]$, while others belong to $G - S$. Since the size of S is in $k^{\mathcal{O}(k)} \cdot \ell$, the paths in $G[S]$ could be enumerated in FPT (in k and ℓ) time. However, guessing the entry and exit points of these parts, how it is done by Jedlicková and Kratochvíl, would be too time-consuming. Here is precisely the moment when the property of the decomposition comes into play. Each connected component of $G - S$ is highly connected and thus is Hamiltonian- $(\ell + 2|S|)$ -linked. Therefore, for any selection of the paths' endpoints, the required collection of vertex-disjoint paths spanning all component vertices always exists (unless empty). Therefore, it is sufficient for the algorithm to enumerate all possible "patterns" of paths' behavior between S and the components of $G - S$ without enumerating the entry-exit endpoints in $G - S$. The latter task reduces to computing a maximum matching in an auxiliary graph. This way, the whole enumeration can be done in FPT time.

Another complication comes from the fact that we do not require the $\alpha(G) < k$ assumption and instead aim to find a solution or an independent set of size k . For that, a constructive version of the high connectivity case is required. That is, given an $\Omega(k \cdot \ell)$ -connected graph and ℓ pairs of endpoints, we need to be able to construct a family of ℓ disjoint paths with the given endpoints spanning the graph or to find an independent set of size k . We employ the general result of Thomas and Wollan [TW05]: In a 10ℓ -connected graph, a collection of ℓ vertex-disjoint paths with the given endpoints always exists. However, the proof of Thomas and Wollan [TW05] is involved, and it is not apparent whether it could be made constructive. Instead, we employ the property that our graphs are of bounded independence number. We use a Ramsey-based argument, finding either an independent set of size k or a large clique, and in the latter case, we can find the desired ℓ paths by connecting the endpoints to the clique via Menger's theorem.

We remark here that the relation between connectivity and linkages in graphs is well-studied. In particular, a sequence of results, including the above-mentioned work of Thomas and Wollan [TW05], guarantees the existence of ℓ vertex-disjoint paths between given pairs of endpoints in a graph of connectivity $\Omega(\ell)$. However, there is a lack of simple and efficient algorithms that *compute* vertex-disjoint paths connecting the given ℓ pairs of endpoints, in an $\Omega(\ell)$ -connected, or even $\ell^{\Omega(1)}$ -connected graph. Our contributions in that direction can be of independent interest to the reader.

Finally, for Theorem 1, we additionally generalize the argument above from Hamiltonian- ℓ -linkages to arbitrary list TM-embeddings. Here, once more, the non-recursive structure of the algorithm proves helpful: With the same choice of the set S , we enumerate all possible behaviors of the target model on S and between S and the connected components of $G - S$. Again, we avoid

enumerating concrete terminals and connecting vertices in $G - S$, fixing only the general connection pattern to S . While considerably more technical, the enumeration can still be done in FPT time. Then, similarly to Hamiltonian- ℓ -linkages, it can be shown that one can always find a spanning list TM-model in a highly-connecting component (or an independent set of size k), with arbitrary locations of the terminals.

Theorem 2: To sketch our second main result, Theorem 2, we recall the proof of Gallai-Milgram theorem for undirected graphs. It is based on the following observation: if there are more than $\alpha(G)$ paths in a path cover of G , then there is an edge between two endpoints of distinct paths. Then these two paths can be joined into a single one and the size of the path cover is reduced by one. This proof gives a polynomial-time algorithm constructing a path cover of G consisting of at most $\alpha(G)$ paths. Note that the algorithm does not have to know or compute $\alpha(G)$ itself: it is enough to iteratively join a pair of paths with adjacent endpoints until it is no longer possible. The algorithm can start from a trivial path cover consisting of $|V(G)|$ empty paths.

The procedure above on its own does not provide any additional relation between the number of paths p in the obtained path cover \mathcal{P} and the independence number $\alpha(G)$ except for $p \leq \alpha(G)$. We propose a way to deal with this issue. First, we distinguish the paths in \mathcal{P} into *usual* and *special*. We call a path usual if its endpoints are distinct and there is no edge between them in G . Otherwise, we call a path *special*. We argue that having usual paths in \mathcal{P} is beneficial. For example, if all paths in \mathcal{P} are usual, then the endpoints of paths in \mathcal{P} form an independent set of size $2p$ as, otherwise, we would join two paths. In particular, this implies that $p \leq \alpha(G)/2$. Even having k usual paths in \mathcal{P} is enough since this yields an independent set of size $p + k$. Together with \mathcal{P} , this independent set is a feasible output for the algorithm of Theorem 2.

While special paths are unsuitable for constructing a large enough independent set, they are suitable for joining. A special path is either a path of length zero or one, or a cycle of length at least three (if considered together with the edge between the endpoints). Then, an edge between two arbitrary vertices of two special paths is as helpful as an edge between endpoints: it allows us to join the two paths and cover the vertices of their union by a single path. Our algorithm exploits this property of special paths and joins them exhaustively until there is no edge between special paths.

This property of special paths, in fact, gives us even more insight into the structure of G . Since special paths are independent of each other, and there are less than k usual paths in \mathcal{P} , taking just one vertex from each special path constitutes an independent set of size greater than $p - k$. Taking two non-adjacent vertices instead of one from $2k$ special paths gives an independent set of size at least $p + k$, solving the problem immediately. Hence, the remaining case is when all but at most $2k$ special paths have vertex sets inducing cliques in G .

However, this structural insight alone is only half of the solution. To proceed further, we need another and more sophisticated way of transforming the paths in \mathcal{P} . Instead of joining two paths as before, this transformation takes two special paths and one usual path, so there are two edges without common endpoints connecting the special paths to the usual path. The transformation produces three paths, and at least two of these paths are usual. Hence, this transformation reduces the number of special paths and does not increase \mathcal{P} .

The transformation above provides the remaining structural insight for the algorithm. When this transformation and all the transformations above cannot be applied, a vertex set S of size less than k separates all usual paths from at least $p - 4k$ special paths with vertex sets inducing cliques. In an auxiliary Lemma 4, we prove that the small size of S can be exploited and irrelevant cliques can be removed from G with respect to the size of an optimal path cover and the independence number. After applying this polynomial-time routine, at most $\mathcal{O}(k^2)$ special paths remain in \mathcal{P} .

The proof outline has come to the final case when $p = \mathcal{O}(k^2)$. In this case, with Theorem 1 in hand, we aim to find the optimal path cover of G as discussed previously in this section. The parameter choice in Theorem 1 is essential, and we choose it to be $p + k$. Thus, if the application of Theorem 1 does not give an optimal path cover, it gives an independent set of size $p + k$. Either way, the algorithm produces a solution feasible for Theorem 2.

We finish the outline of the proof of Theorem 2 by noting that the algorithm described above takes polynomial time, except for the application of Theorem 1.

2 Proof of Theorem 1: List TM-embedding parameterized by independence number

In this section, we prove Theorem 1. The proof consists of three main steps, deferred to their respective subsections. The first step is the proof of the stronger version of the theorem for highly connected graphs which we call Spanning Lemma. It shows that for any fixed injection $f : V(H) \rightarrow V(G)$ the desired TM-embedding respecting f or an independent set can be found; moreover, the found TM-embedding spans $V(G)$. The second step is the so-called Merging Lemma, that combines the solutions from several highly-connected components, provided that the size of the separator between them is small. Finally, to obtain the proof of Theorem 1 itself, we show an iterative procedure that either finds the desired separator, or an independent set.

2.1 Spanning Lemma

We first deal with highly-connected graphs. More precisely, we demonstrate that, given an injective mapping $f : V(H) \rightarrow V(G)$, we can either compute a TM-embedding of H in G respecting f and spanning $V(G)$, or find an independent set of size k in G ; this holds if the connectivity of G is sufficiently high. This step follows the approach of Jedlicková and Kratochvíl [JK23, Theorem 7], who proved that for every $k > 1$ and $\ell > 1$, if a graph G satisfies $\alpha(G) < k$ and G is $\max\{k\ell, 10\ell\}$ -connected, then G is Hamiltonian- ℓ -linked. Recall that a graph G is ℓ -linked if it has at least 2ℓ vertices and for every set of ℓ disjoint pairs (s_j, t_j) of distinct vertices, there exist ℓ vertex-disjoint paths in G such that the endpoints of the j -path are s_j and t_j , for each $j \in [\ell]$. A graph is called *Hamiltonian- ℓ -linked* if, in addition, the set of ℓ paths covers all vertices of G .

To adapt the approach of Jedlicková and Kratochvíl for our purposes, we need to address two issues. The first issue is minor; we need to slightly strengthen the connectivity requirement to simplify the case analysis and allow for possible coinciding endpoints of the paths (which is not allowed in [JK23]). The second issue is more serious—the proof of Jedlicková and Kratochvíl is not constructive, meaning that it cannot be used to construct the linkage. However, we need a procedure constructing the linkage for computing a TM-embedding of H . The proof of Jedlicková and Kratochvíl builds on the following result of Thomas and Wollan [TW05], which is also non-constructive.

Proposition 1 (Corollary 1.3, [TW05]). *If G is 10ℓ -connected then G is ℓ -linked.*

We do not know how to make the proof of Proposition 1 constructive in general. However, since the graphs of our interest have bounded independence number, we can use this property to construct linkages in highly connected graphs. The algorithm in the following lemma serves as a key starting subroutine in the proof of our result for highly-connected instances.

Lemma 1. *There is an algorithm that, given integers $k, \ell \geq 1$, a 10ℓ -connected graph G , and a family of vertex pairs $(s_1, t_1), \dots, (s_\ell, t_\ell)$ such that $s_i \neq t_i$ for each $i \in [\ell]$, in time $2^{(k+\ell)^{\mathcal{O}(k)}} + |G|^{\mathcal{O}(1)}$*

outputs either an independent set of size k or a family of ℓ internally vertex-disjoint $s_i t_i$ -paths P_i for $i \in [\ell]$, that is, the i -th path connects s_i and t_i and does not contain any other s_j or t_j as an internal vertex.

Proof. Let $T = \bigcup_{i=1}^{\ell} \{s_i, t_i\}$ be the set of vertices in the input pairs. We call the vertices in T *terminals*. For simplicity, our algorithm first ensures that $s_1, t_1, s_2, t_2, \dots, s_\ell, t_\ell$ are 2ℓ distinct vertices, i.e., $|T| = 2\ell$. For each vertex $v \in T$ that appears in $p > 1$ input pairs, the algorithm replaces v by p true twins, i.e., vertices that contain the neighborhood of v and are also adjacent to each other. Then, each repeating occurrence of v in the input pairs is replaced by a distinct copy. This transformation results in 2ℓ distinct terminal vertices. The requirement on 10ℓ -connectivity of the graph still holds, and the modification does not increase the maximum size of an independent set. Furthermore, given an independent set in the transformed graph, replacing each copy of a terminal vertex with the corresponding original vertex of G gives an independent set in the original graph. The same is true for paths avoiding terminal vertices: a path between copies is equivalent to the path between original vertices. Therefore, the transformed input is equivalent to the original input, and the respective solutions can be transformed in polynomial time. Henceforth, the algorithm works with the transformed input, meaning that all vertices in the input pairs are distinct.

Observe that by Proposition 1, a family of vertex-disjoint $s_i t_i$ -paths always exists. Hence, our task is to construct such paths. First, we show that if G contains a clique of size 2ℓ , then the paths can be constructed in polynomial time.

Claim 1. *Let C be a given clique of size 2ℓ in G . Then a family of internally vertex-disjoint paths P_1, P_2, \dots, P_ℓ , where for each $i \in [\ell]$, P_i connects s_i with t_i , can be constructed in polynomial time.*

Proof of Claim 1. Since G is 2ℓ -connected, by Menger's theorem [Men27, Die17], there are 2ℓ internally vertex-disjoint paths connecting vertices of $T = \bigcup_{i=1}^{\ell} \{s_i, t_i\}$ with distinct vertices of C . These paths can be constructed in the standard fashion via a network flow algorithm; in particular, using the Ford-Fulkerson algorithm [FF56] to find ℓ -flow in $\mathcal{O}(\ell \cdot |G|)$ running time. Then for each $i \in [\ell]$, we have two paths, one connecting s_i with c_{2i-1} and the other connecting t_i with c_{2i} for some $c_{2i-1}, c_{2i} \in C$. Note that it is possible that one or two of these paths are trivial, i.e., $s_i = c_{2i-1}$ or $t_i = c_{2i}$ can be true.

As C is a clique in G and $c_{2i-1} c_{2i} \in E(G)$, these paths can be concatenated, resulting in the $s_i t_i$ -path P_i . Since $c_1, c_2, \dots, c_{2\ell}$ are pairwise distinct, any two paths among P_1, P_2, \dots, P_ℓ do not share any common vertices. We obtain ℓ paths as required. \square

When G is of bounded size, we employ brute-force enumeration.

Claim 2. *If G has less than $(k+2\ell)^k$ vertices, then paths P_1, P_2, \dots, P_ℓ can be found in $\mathcal{O}(2^{(k+2\ell)2^k})$ time.*

Proof of Claim 2. It is sufficient to enumerate all possible subsets of edges of G contained in the desired family of paths. When a subset S of edges is fixed by the algorithm, it is sufficient to check that for each $i \in [\ell]$, there is a connected component in the graph with the vertex set $V(G)$ and the edge set S , that is a path between s_i and t_i . If S satisfies this for each $i \in [\ell]$, the algorithm stops and reports the corresponding components as P_1, P_2, \dots, P_ℓ . \square

It is left to explain how the algorithm constructs the clique C , given that G has at least $(k+2\ell)^k$ vertices. The essential tool here is the classical Ramsey's theorem [Ram30], which guarantees a large clique or independent set in a large enough graph. For completeness, we present the corresponding polynomial-time subroutine based on the Ramsey number bound given by Erdős and Szekeres in 1935 [ES09].

Claim 3. *There is a polynomial-time algorithm that, given a graph G and two integers $r, s > 0$ such that $|V(G)| \geq \binom{r+s-2}{r-1}$, outputs either an independent set of size r or a clique of size s in G .*

Proof of Claim 3. Note that G necessarily has either an independent set of size r or a clique of size s because the corresponding Ramsey number satisfies $R(r, s) \leq \binom{r+s-2}{r-1}$ [ES09]. The algorithm is recursive. Since $r, s > 0$, $|V(G)| > 1$ and G has at least one vertex, denote it by v . If $r = 1$ or $s = 1$, then algorithm outputs $\{v\}$ and stops. For $r, s \geq 2$, consider vertex sets $A = N_G(v)$ and $B = V(G) \setminus \{v\} \setminus A$. Since $|V(G)| = |A| + |B| + 1$ and $V(G) \geq \binom{r+s-2}{r-1}$ we have that either $|A| \geq \binom{r+s-3}{r-1}$ or $|B| \geq \binom{r+s-3}{r-2}$. If $|A| \geq \binom{r+s-3}{r-1}$ then $G[A]$ has either an independent set of size r or a clique of size $s-1$. In this case, the algorithm makes a recursive call to itself for the input $G[A]$, $r > 0$ and $s-1 > 0$. If its output is a clique of size $s-1$, the algorithm additionally appends v to it, resulting in a clique of size s . Thus, the algorithm obtains either an independent set of size r , or a clique of size s as required. The case of $|B| \geq \binom{r+s-3}{r-2}$ is symmetric, as v can be appended to any independent set in $G[B]$.

Because the values of $\binom{x+y-2}{y-1} = \binom{x+y-2}{x-1}$ for $x \leq r$ and $y \leq s$ can be computed in $\mathcal{O}(rs)$ time using the recursive formula for binomial coefficients, we obtain that the running time of the algorithm is $\mathcal{O}(n^2)$ where $n = |V(G)|$. The proof of the claim is complete. \square

Now, we are ready to complete the algorithm:

- If $|V(G)| \leq (k + 2\ell)^k$, the algorithm applies the subroutine of Claim 2 to find the solution, outputs it and stops.
- Otherwise, since $(k + 2\ell)^k \geq \binom{k+2\ell-2}{k-1}$, the algorithm applies the subroutine of Claim 3 with $r = k$ and $s = 2\ell$.
- If the output is an independent set of size k , the algorithm outputs it, and stops.
- Otherwise, the output is a clique of size 2ℓ in G . The algorithm uses the subroutine of Claim 1 to find the solution, then outputs it and stops.

The correctness of the algorithm and the upper bound on its running time follow from the claims. The proof of the lemma is complete. \square

We now proceed to present an algorithm that, for a highly connected graph G , a graph H , and a fixed mapping $f : V(H) \rightarrow V(G)$, either constructs a TM-embedding of H in G respecting f and spanning $V(G)$, or finds an independent set of size k in G .

Lemma 2 (Spanning Lemma). *Let H, G be graphs, such that H is non-empty. Let $f : V(H) \rightarrow V(G)$ be an injective mapping, and let k be an integer. Let G additionally be $(\max\{k + 2, 10\} \cdot h)$ -connected, where $h = |V(H)| + |E(H)|$. There is an algorithm with running time $2^{(h+k)\mathcal{O}(k)} + |G|^{\mathcal{O}(1)}$ that computes either a subgraph M of G so that (M, f) is a TM-embedding of H in G spanning $V(G)$, or an independent set of size k in G .*

Proof. First, the algorithm reduces to the case where H has no isolated vertices. Since each isolated vertex v in H has a fixed image $f(v)$ in G , the algorithm simply removes v from H and $f(v)$ from G . In the following steps of the algorithm, if a spanning TM-embedding of H is produced, the algorithm extends this embedding with the removed images of isolated vertices. Such an embedding remains spanning for G , while all described procedures are performed in polynomial time.

Note that removing an equal number of vertices from both G and H preserves the connectivity constraint in the statement. From now on, every vertex of H has at least one incident edge, while G is $(\max\{k + 2, 10\} \cdot |E(H)|)$ -connected.

In order to proceed, it is convenient to reformulate the problem in terms of disjoint paths: *Given ℓ vertex pairs $(s_1, t_1), (s_2, t_2), \dots, (s_\ell, t_\ell)$ of distinct vertices in G , find a family of internally-disjoint paths P_1, P_2, \dots, P_ℓ such that P_i connects s_i with t_i for each $i \in [\ell]$ and each vertex of G belongs to at least one path, and the union of the paths spans $V(G)$.*

To see that the reformulation is equivalent, put $\ell = |E(H)|$ and $(s_i, t_i) = (f(x_i), f(y_i))$, where x_i, y_i are the endpoints of the i -th edge of H . Then, given the paths P_i , it is straightforward to construct the subgraph M by taking the union of the paths; (M, f) then is the desired TM-embedding of H in G . The reverse is also possible: given a TM-embedding (M, f) of H in G spanning $V(G)$, the respective path can be found by traversing M from $f(x_i)$ to $f(y_i)$. From now on, we do not refer to TM-embeddings, but work with the equivalent disjoint-path statement above.

The first step of the algorithm is to find the initial family of paths P_1, P_2, \dots, P_ℓ , that satisfies all constraints except for spanning $V(G)$. That is, P_i is an $s_i t_i$ -path, and for each $i \neq j$, P_i and P_j have no common vertices except for possibly their endpoints. For this, the algorithm invokes the algorithm of Lemma 1 as a subroutine. The call is valid since G is 10ℓ -connected, therefore in time $2^{(k+\ell)\mathcal{O}(k)} + |G|^{\mathcal{O}(1)}$ the initial sequence P_1, P_2, \dots, P_ℓ is produced.

The only step left is to ensure that the spanning constraint is satisfied by the path sequence. The algorithm achieves this incrementally. That is, the algorithm enlarges one of the paths in the sequence without violating any other constraints. If we show how this can be done exhaustively and in polynomial time, it is clearly sufficient to find the desired TM-embedding.

We describe the enlargement process, akin to the method outlined in [JK23], with a key distinction. Unlike the scenario in [JK23], we lack assurance that G satisfies $\alpha(G) < k$, making the procedure potentially prone to failure. Therefore, we ensure that (a) the enlargement is completed within polynomial time and (b) if the enlargement fails, an independent set of size k is identified in G .

The procedure works as follows: If P_1, P_2, \dots, P_ℓ span all vertices of G , the algorithm achieves the desired outcome, reports the solution, and halts. Otherwise, let $S = \bigcup_{i=1}^{\ell} V(P_i)$ denote the span of the path sequence, with the assumption that $S \neq V(G)$. The algorithm selects any $x \in V(G) \setminus S$. By the $(k+2)\ell$ -connectivity of G , Menger's theorem guarantees the existence of $\min\{(k+2)\ell, |S|\}$ paths. Each path goes from x to S , and no two paths share any common vertex (including endpoints in S) but x . One can find such paths by utilizing the Ford-Fulkerson algorithm [FF56] to find a flow of size at most $(k+2)\ell$ in $\mathcal{O}(k\ell \cdot |G|)$ running time.

If $|S| \leq (k+2)\ell$, then two of these paths end in s_1 and v , where v is the neighbor of s_1 in P_1 . Denote these paths by Q_1 and Q_2 . These two paths are internally disjoint, both starting in x and ending in s_1 and v , respectively. The algorithm enlarges P_1 by replacing it with the path $(s_1 Q_1 x) \circ (x Q_2 v) \circ (v P_1 t_1)$, that is, x becomes embedded in P_1 between s_1 and v . Here and next, for a path P and vertices u, v on P we denote by uPv the subpath of P from u to v , and for two paths P_1 and P_2 sharing an endpoint, we denote by $P_1 \circ P_2$ their concatenation. This operation strictly increases the span of the path sequence.

It is left for the algorithm to process the case $(k+2)\ell < |S|$. There are precisely $(k+2)\ell$ paths that were found by the Ford-Fulkerson algorithm, so at least $k\ell$ paths end in internal vertices of P_1, P_2, \dots, P_ℓ . Hence, there is $i \in [\ell]$ such that at least k paths end in internal vertices of P_i . The algorithm takes arbitrary k of them. Denote the k selected paths by Q_1, Q_2, \dots, Q_k and their respective endpoints in P_i by v_1, v_2, \dots, v_k . For each $a \in [k]$, let u_a be the successor of v_a in P_i if one follows vertices on P_i from s_i to t_i .

If the set $\{u_1, u_2, \dots, u_k\}$ is an independent set in G , then the algorithm reports this independent set and stops. Otherwise, there is an edge $u_a u_b$ in G for some $a, b \in [k]$. Without loss of generality, v_a goes before v_b on P_i in the order from s_i to t_i . The algorithm then enlarges P_i by replacing it with $(s_i P_i v_a) \circ (v_a Q_a x) \circ (x Q_b v_b) \circ (v_b P_i u_a) \circ u_a u_b \circ (u_b P_i s_i)$. Again, x is embedded into P_i while

no original vertices of P_i are lost, increasing the total span.

The algorithm repeats the described procedure until paths P_1, P_2, \dots, P_ℓ span the whole vertex set of G , or an independent set of size k is encountered. This procedure is repeated at most $|V(G)|$ times, so the total running time of this part is polynomial in the size of the instance.

The correctness of the algorithm follows from the discussion. The proof of the lemma is complete. \square

2.2 Merging Lemma

With the highly-connected case at hand, we now move to the second key ingredient in the proof of Theorem 1. We show that if the graph is not necessarily highly-connected, but contains a small-sized vertex subset such that after its removal the remaining components are highly-connected, then we can solve the instance in FPT time. Here, the parameter is the size of the subset to remove plus the size of H plus k . In essence, the proof boils down to reconstructing the global solution from the solutions obtained on the highly-connected components via Lemma 2, therefore we dub the result Merging Lemma, stated next.

Lemma 3 (Merging Lemma). *Let G be a graph, let $S \subseteq V(G)$ be a vertex subset in G , and let C_1, \dots, C_t be the connected components of $G - S$. Let H be a graph, let k be a parameter, and $L : V(H) \rightarrow 2^{V(G)}$ be a list assignment. Assume that each component C_i is at least $\max\{k + 2, 10\} \cdot (3h + 3s)$ -connected, where $h = |V(H)| + |E(H)|$, $s = |S|$. There is an algorithm that either returns a maximum-size list TM-embedding of H in G respecting the list assignment L , or correctly reports that none exists, or returns an independent set in G of size k . The running time of the algorithm is $2^{(s+h+k)^{\mathcal{O}(k)}} \cdot |G|^{\mathcal{O}(1)}$.*

Proof. The main idea of the proof is to enumerate all possible interactions of the desired TM-model with S . We encapsulate the choice of the interaction in an object we call a *cut descriptor*, to be formally defined next. Intuitively, a cut descriptor captures the behavior of a TM-model of H in G in the subset S and its neighborhood: Given a TM-model (M, T) of H in G , its respective cut descriptor is obtained from M by dissolving all non-terminals outside of S , where vertices not in S are represented implicitly by the index of their component. The latter part is crucial to keep the number of choices the function of the parameter. As we will see later, such an implicit representation is sufficient since Lemma 2 always provides a spanning TM-model (or an independent set) and allows to fix an arbitrary injection of the terminals.

Formally, a cut descriptor is the tuple (W, M_W, ξ, f_W) , where W is a set of vertices (indexed implicitly from 1 to $|W|$), M_W is a graph with the vertex set $S' \cup W$ with $S' \subseteq S$, $\xi : W \rightarrow [t]$ is a mapping from W to indices of the components C_1, \dots, C_t , and $f_W : V(H) \rightarrow S' \cup W$ is an injective mapping from $V(H)$ to vertices of M_W . Denote $T_W = \text{Im } f_W$, and for each $i \in [t]$, denote $W_i = \{w \in W : \xi(w) \in C_i\}$. We require the following properties for (W, M_W, ξ, f_W) to be a cut descriptor:

- CD1** (M_W, f_W) is a TM-embedding of H in M_W ;
- CD2** every $w \in W$ is in $T_W \cup N_{M_W}(S)$;
- CD3** for each $w, w' \in W$ with $\xi(w) \neq \xi(w')$, w and w' are not adjacent in M_W ;
- CD4** $M_W[S]$ is a subgraph of G , and for each $s \in S, h \in V(H)$ with $f_W(h) = s$, $s \in L(h)$;
- CD5** there exists an injective assignment $\tau : W \rightarrow V(G)$ such that i) $\tau(w) \in C_{\xi(w)}$ ii) if there exists $h \in V(H)$ with $f_W(h) = w$, then $\tau(w) \in L(h)$ iii) $N_{M_W}(w) \cap S \subseteq N_G(\tau(w))$.

We first observe an upper bound on the number of extra vertices in a cut descriptor.

Claim 4. *If (W, M_W, ξ, f_W) is a cut descriptor, then $|W|$ is at most $h + 2s$.*

Proof of Claim 4. We call a vertex $w \in W$ a *terminal* if $w \in T_W$, and a *connector* otherwise. From **CD2**, every connector in w is adjacent to a vertex of S in M_W . Since f_W is injective, number of terminals in W is at most $|V(H)|$, therefore it remains to bound the number of connectors in W by $|E(H)| + 2s$. Let $w \in W$ be a connector that is adjacent to a terminal in S , the number of such connectors is at most $\sum_{s \in S \cap T_W} |N_{M_W}(s) \cap W| \leq |E(H)|$, since by **CD1** every $s \in S \cap T_W$ has degree $\deg_H(h)$, where h is the unique pre-image of s with respect to f_W , and no edge of H is counted twice. Finally, the number of connectors in W adjacent only to non-terminals of S is at most $2s$, since every non-terminal has degree two in M_W by **CD1**. \lrcorner

We also argue that if any list TM-embedding of H in G exists, then there exists also a “matching” cut descriptor. For a subgraph M of G and a subset $C \in V(G)$, we say that M *hits* C if $M[C]$ contains an edge, M *touches* C if $M[C]$ contains no edges but at least one vertex, and M *avoids* C if $V(M) \cap C = \emptyset$.

Claim 5. *Let (M, f) be a list TM-embedding of H in G . Then there exist a cut descriptor (W, M_W, ξ, f_W) such that $M_W[S] = M[S]$, and for each $i \in [t]$, M hits C_i if and only if M_W hits W_i , touches C_i if and only if M_W touches W_i and $|V(M) \cap C_i| = |W_i|$, and avoids C_i if and only if $W_i = \emptyset$.*

Proof of Claim 5. We construct the desired cut descriptor as follows. First, let W be the subset of vertices in G that are either terminals in $V(G) \setminus S$ or neighbors of S in the model M , i.e., $W = (\text{Im } f \setminus S) \cup N_M(S)$. We then dissolve all vertices in $V(M) \setminus (S \cup W)$, which are all non-terminals by definition of W , to obtain M_W from M . We set f_W as the respective mapping of the terminals obtained from f . Clearly, **CD1** is fulfilled for (M_W, T_W) , and **CD2** also holds immediately by construction. We set ξ so that $W_i = W \cap C_i$ for each $i \in [t]$. We now proceed to verify the remaining properties of a cut descriptor.

For **CD3**, consider $w, w' \in W$ with $\xi(w) \neq \xi(w')$. Let $w \in W_i, w' \in W_j$ in M_W , $i \neq j$, and $w \in C_i, w' \in C_j$ in G . Since S separates C_i and C_j in G , and M is a subgraph of G , w and w' are not adjacent in M . By construction of M_W , only non-terminals outside of S and W are dissolved; every such non-terminal lies in $C_{i'}$ together with its neighbors, for some $i' \in [t]$, and dissolving it only changes edges between vertices of $C_{i'}$. Therefore, w and w' are also not adjacent in M_W .

We now move to **CD4**, by construction $M_W[S] = M[S]$ and is therefore a subgraph of $G[S]$. Moreover, $f \equiv f_W$ on S ; let s be such that $f(h) = f_W(h) = s$, then $s \in L(h)$ since (M, f) is a list TM-embedding respecting L .

Finally, for **CD5**, the assignment τ is given directly by construction as an identity assignment on $W \subset V(G)$. For each $w \in W$, $\tau(w) \in C_{\xi(w)}$ by the construction of ξ . If $f_W(h) = w$, then $w = \tau(w) \in L(h)$ since $f(h) = w$ and (M, f) is a list TM-embedding respecting L . By construction of M_W , the edges between W and S are never changed between M and M_W , therefore $N_{M_W}(w) \cap S = N_M(w) \cap S$, and the latter is a subset of $N_G(w)$ since M is a subgraph of G .

For the last claim of the lemma, we argue again by construction: since the dissolving operation only affects edges inside C_i , for some i , and dissolving a vertex in C_i leaves C_i with at least one edge inside, the status of the component (hit/touched/avoided) is always preserved between M and M_W . Specifically, if an edge of M was present in C_i , then after the dissolving, an edge is still present in M_W in W_i . If no edge but some vertices of M are present in C_i then all these vertices are terminals or neighbors of S , and $|M \cap C_i| = |W_i|$, and if no vertices of M are in C_i , then also W_i is empty. \lrcorner

Now, the algorithm proceeds as follows. First, it branches over the choice of a cut descriptor. Then, for a fixed cut descriptor, the algorithm picks an arbitrary assignment fulfilling **CD5**. Since the assignment fixes the vertices of W in G , we run the algorithm of Lemma 2 on each C_i to find a TM-model connecting the vertices of W_i as prescribed by M_W . If any of the runs returns an independent set of size k , we output it and stop. Otherwise, we augment the model (M_W, T_W) to a model (M, T) of H in G via the returned model in each C_i . Out of all models of H in G obtained in the branches, the algorithm returns the one of maximum size. The above algorithm is shown in detail in Algorithm 1.

It remains to verify that Algorithm 1 is correct and to upper-bound its running time. The series of claims presented next verifies the correctness of the algorithm. While checking the conditions **CD1–CD5** for a fixed cut descriptor in Line 5 is generally straightforward, we provide more details for the condition **CD5** by constructing τ directly.

Claim 6. *In Line 5, the condition **CD5** can be verified, and a suitable assignment τ can be constructed via a call to a maximum matching algorithm on an auxiliary bipartite graph of polynomial size.*

Proof of Claim 6. We construct an auxiliary bipartite graph B where the two parts are W and $V(G)$, and $w \in W, v \in V(G)$ are adjacent if setting $\tau(w) = v$ does not violate any of the conditions i–iii) of **CD5**. That is, w and v are adjacent if and only if

- $v \notin S$ and $v \in C_{\xi(w)}$,
- if $w \in \text{Im } f_W$ with $f_W(h) = w$, then $v \in L(h)$,
- $N_{M_W} \cap S \subseteq N_G(v)$.

Clearly, a suitable assignment τ exists if and only if a matching in B that covers W exists. ┘

Then, we show that invoking Lemma 2 with the given arguments is possible.

Claim 7. *The call to Lemma 2 in Line 15 is valid.*

Proof of Claim 7. By the condition in Line 12, H_i is not empty. The mapping $\tau|_{T_i}$ is indeed an injective mapping from $T_i = V(H_i)$ to C_i , where the latter is guaranteed by **CD5**, i). It remains to verify that $G[C_i]$ is $\max\{k+2, 10\} \cdot (|V(H_i)| + |E(H_i)|)$ -connected. By the statement of the lemma, $G[C_i]$ is $\max\{k+2, 10\} \cdot (3h+3s)$ -connected, so it suffices to argue that $|V(H_i)| + |E(H_i)| \leq 3h+3s$.

We consider two kinds of edges in H_i separately. First, there are at most $2|E(H)|$ edges incident to vertices in $V(H_i) \cap T_W$, since for each $w \in T_w$, $\deg_{H_i}(w) \leq \deg_{M_W}(w)$, and $\sum_{w \in T_W} \deg_{M_W}(w) \leq 2|E(H)|$. Second, consider edges where both endpoints are connectors. By **CD1–CD2**, each connector has degree two in M_W and at least one neighbor in S . Therefore, two connectors adjacent in H_i have in total two distinct edges into S . The size of the cut between S and W in M_W is at most $2s + |E(H)|$, since the total degree of terminals in S towards W is at most $|E(H)|$, and every other vertex has degree two. Thus, the number of edges of the second type in H_i is at most $s + |E(H)|/2$. The total number of edges in H_i is then at most $\frac{3}{2}|E(H)| + s$.

For vertices, there are at most $|V(H)|$ vertices in H_i that are in T_W , and every other vertex is a connector in M_W . As above, every connector has at least one neighbor in S , and the cut size between S and W is at most $2s + |E(H)|$. Therefore, the total number of vertices in H_i is at most $|V(H)| + |E(H)| + 2s$, and $|V(H_i)| + |E(H_i)| \leq |V(H)| + \frac{5}{2}|E(H)| + 3s \leq 3h + 3s$. ┘

We are now able to show that we arrive at a TM-embedding of H_i in $G[C_i]$, either after invoking Lemma 2 such that the output is not an independent set or after taking the empty graph over $\tau(T_i)$.

Input: Graphs G and H , mapping $L : V(H) \rightarrow 2^{V(G)}$, integer k , vertex subset $S \subseteq V(G)$ splitting G into components C_1, \dots, C_t such that for each $i \in [t]$, $G[C_i]$ is $\max\{k+2, 10\}(3h+3s)$ -connected, where $h = |V(H)| + |E(H)|$, $s = |S|$.

Output: An independent set of size k in G , or a list TM-embedding of H in G respecting L , or the output that no such TM-embedding exists.

```

1  $\mathcal{M} \leftarrow \emptyset$ ;
2 for  $|W| \leftarrow 0$  to  $h+2s$  do
3    $W \leftarrow$  vertex set of size  $|W|$ ;
4   foreach  $M_W$  a graph on  $S \cup W$ ,  $\xi : W \rightarrow [t]$ ,  $f_W : V(H) \rightarrow S \cup W$  do
5     verify CD1–CD4, verify CD5 by a matching instance and pick arbitrary suitable  $\tau$ ;
6     if any of CD1–CD5 fail then
7       continue;
8     end
9      $M \leftarrow M_W$ ,  $f \leftarrow f_W$ ;
10    for  $i \leftarrow 1$  to  $t$  do
11       $T_i \leftarrow \{w \in W : \xi(w) = i\}$ ,  $H_i = M_W[T_i]$ ;
12      if  $H_i$  contains no edges then
13         $O \leftarrow$  empty graph over the vertex set  $\tau(T_i)$ ;
14      else
15         $O \leftarrow$  invoke Lemma 2 with  $H_i$ ,  $G[C_i]$ ,  $\tau|_{T_i}$ ,  $k$ ;
16      end
17      if  $O$  is an independent set of size  $k$  then
18        return  $O$ .
19      else
20         $M_i \leftarrow O$ ;  $M \leftarrow M$  where  $M[T_i]$  is replaced by  $M_i$ ;
21        foreach  $w \in (T_i \cap \text{Im } f_W)$  do
22           $f(f_W^{-1}(w)) \leftarrow \tau(w)$ ;
23        end
24      end
25    end
26     $\mathcal{M} \leftarrow \mathcal{M} \cup \{(M, f)\}$ ;
27  end
28 end
29 if  $\mathcal{M} \neq \emptyset$  then
30   return  $(M, f) \in \mathcal{M}$  with  $\max |M|$ .
31 else
32   return No solution.
33 end

```

Algorithm 1: Algorithm of Lemma 3.

Claim 8. *In Line 20, M_i is such that $(M_i, \tau|_{T_i})$ is a TM-embedding of H_i in $G[C_i]$.*

Proof of Claim 8. In case H_i is empty, in Line 13 the algorithm constructs the empty graph over $\tau(T_i)$, which is later assigned to M_i . Clearly, this is the desired TM-embedding.

If H_i is not empty, a call to Lemma 2 is made in Line 15. By Claim 7, the call is valid, and therefore O is either an independent set of size k in $G[C_i]$, or the subgraph M_i in the desired TM-embedding. Since the condition statement before Line 20 explicitly verifies that O is not an independent set of size k , in Line 20 the latter case holds. \square

In case for each $i \in [t]$ we obtain the desired TM-embedding of H_i in $G[C_i]$, we argue that the constructed (M, f) is indeed a valid list TM-embedding of H in G .

Claim 9. *At Line 26, (M, f) is a list TM-embedding of H in G respecting L .*

Proof of Claim 9. The algorithm starts with the TM-model M_W of H in M_W , and replaces $M_W[T_i]$ by M_i for each $i \in [t]$, where M_i is the TM-model of H_i in $G[C_i]$. The list condition for M is fulfilled immediately since f and f_W coincide, f_W respects L on S via **CD4**, and τ respects L via **CD5**, ii). We now argue that (M, f) is a TM-embedding of H in G .

First, we argue that M is a subgraph of G by considering three types of edges in M . Edges with two endpoints in S are not changed between M_W and M , and therefore are present in G since $M_W[S]$ is a subgraph of G by **CD4**. Consider an edge between $s \in S$ and $v \in C_i$, $i \in [t]$, in M . By construction of M , there exists an edge between s and $w \in W_i$ in M_W with $\tau(w) = v$. By **CD5**, $N_{M_W}(w) \cap S \subseteq N_G(\tau(w))$, meaning that s and $v = \tau(w)$ are adjacent in G . Finally, consider an edge of M with both endpoints outside S . By construction, the edge is a part of M_i , for some $i \in [t]$, and is present in G since M_i is a subgraph of $G[C_i]$.

Second, we claim that all vertices in $V(M) \setminus \text{Im } f$ have degree two, and each $v \in \text{Im } f$ with $v = f(h_v)$ has $\deg_M(v) = \deg_H(h_v)$. Based on the above, the vertices of S do not change the degree between M_W and M , and M_W is a TM-model of H . Consider a vertex $v \in V(M) \setminus \text{Im } f \setminus S$, $v \in C_i$ for some $i \in [t]$. If $v \notin T_i$, then v has degree two in M_i since M_i is a TM-model of H_i , and has the same neighborhood in M by construction of M . If $v \in T_i$, then its degree coincides with the degree of w in M_W with $\tau(w) = v$; the degree is also two since M_W is a TM-model of H and $w \notin \text{Im } f_W$. Finally, for a vertex $v \in \text{Im } f \setminus S$, $v \in C_i$ for some $i \in [t]$, $v = \tau(w)$ for some $w \in T_i \cap \text{Im } f_W$ and $\deg_{M_i}(v) = \deg_{H_i}(w)$ since M_i is a TM-model of H_i . Similarly, $N_{M_W}(w) \cap S = N_M(v) \cap S$, therefore $\deg_M(v) = \deg_{M_i}(v) + |N_M(v) \cap S| = \deg_{H_i}(w) + |N_{M_W}(w) \cap S| = \deg_H(h)$, where $v = f(h)$, $w = f_W(h)$, since (M_W, f_W) is a TM-model of H and all neighbors of w in M_W are either in S or in T_i by **CD5**.

We now argue that dissolving non-terminals in M results in a graph isomorphic to H via f . We call the graph obtained after dissolving by H' and identify $V(H')$ with a subset of $V(M)$. For $h_v, h_u \in V(H)$, we show that $f(h_v), f(h_u)$ are adjacent in H' . Denote by $v = f(h_v)$, $u = f(h_u)$. If $v \in S$, let $v' = v$, otherwise $v' \in W$ is such that $\tau(v') = v$. We define u' analogously to u . By the choice of v', u' , we have that $f_W(h_v) = v'$, $f_W(h_u) = u'$. Also, since (M_W, f_W) is a TM-embedding of H , v' and u' are connected by a path P' in M_W whose all internal vertices are non-terminals.

Based on P' , we construct a vu -path P in M whose all internal vertices are non-terminals. First, replace all vertices of W in P' with their images given by τ . Second, for each two consecutive vertices w_1, w_2 in P' , such that $w_1, w_2 \in W$, insert in P between $\tau(w_1), \tau(w_2)$ the $\tau(w_1)\tau(w_2)$ -path in M_i , whose all internal vertices are not in T_i . Such a path exists because $w_1, w_2 \in T_i$ for some $i \in [t]$ by **CD3**, and because M_i is a TM-model of $M_W[T_i]$ in $G[C_i]$. Since for any $s \in S$ and $w \in W$ that are neighbors in M_W , vertices s and $\tau(w)$ are also neighbors in G by **CD5**, iii), P is indeed a vu -path in G . By construction, all internal vertices are not in $\text{Im } f$. Hence v and u are adjacent

in H' . It remains to observe that $\deg_{H'}(v) = \deg_M(v) = \deg_H(h_v)$ for any v, h with $v = f(h_v)$. Therefore, H' is isomorphic to H via f , completing the proof. \lrcorner

Next, we show that trying each possible cut descriptor guarantees that the largest list TM-embedding is found.

Claim 10. *The TM-embedding (M, f) returned on Line 30 is of maximum size among all suitable list TM-embedding.*

Proof of Claim 10. Consider the list TM-embedding (M^*, f^*) of H in G of maximum size. By Claim 5, there exists a cut descriptor (W, M_W, ξ, f_W) with $M_W[S] = M^*[S]$, and M^* hits (touches/avoids) C_i if and only if M_W hits (touches/avoids) W_i for each $i \in [t]$. Consider the TM-embedding (M, f) constructed by the algorithm at Line 26 with the cut descriptor set to (W, M_W, ξ, f_W) . Note that since the algorithm reached Line 30, all calls to Lemma 2 resulted in a TM-model, and M is successfully constructed. By construction, $M[S] = M_W[S] = M^*[S]$. Observe also that if M_W hits W_i , then the TM-model M_i returned by Lemma 2 spans C_i , meaning $|M \cap C_i| \geq |M^* \cap C_i|$. If M^* touches C_i , then $|M^* \cap C_i| = |W_i|$, and M has at least as many vertices in C_i . If M^* avoids C_i , $|M^* \cap C_i| = 0$. Therefore, in each S , C_1, C_2, \dots, C_t , M has at least as many vertices as M^* , meaning $|M| \geq |M^*|$, proving the claim. \lrcorner

We also argue that if the algorithm reports that no suitable TM-model is found, then indeed none exists.

Claim 11. *If \mathcal{M} is empty by the end of the algorithm, then no list TM-model of H in G respecting L exists.*

Proof of Claim 11. Assume a list TM-model of H in G exists. Then by Claim 5, there exists also a cut descriptor. Consider the iteration of the algorithm with this cut descriptor. Since the algorithm reaches Line 32, all calls to Lemma 2 resulted in a suitable TM-embedding being found, therefore the algorithm necessarily adds a TM-embedding (M, f) to \mathcal{M} at Line 26. This contradicts the assumption that Line 32 is reached. \lrcorner

Finally, we estimate the running time. First note that all lines except Line 4 and Line 15 incur at most a polynomial factor. By Lemma 2, up to polynomial factors Line 15 takes time $2^{(h+s+k)\mathcal{O}(k)}$, for each invocation. This holds since $|V(H_i)| + |E(H_i)| \leq 3h + 3s$, see the proof of Claim 7. Moving to Line 4, let $w = h + 2s$ be the maximum size of W ; there are at most w^t choices for ξ , and at most $|V(H)|^{s+w}$ choices for f_W . We may assume $t \leq k$, since otherwise, an independent set of size k can be found trivially by taking an arbitrary vertex from each C_i . We now argue that with a fixed f_W , there are at most $(s+w)^{2(s+w)}$ choices for M_W that satisfy **CD1**. Indeed, terminals are fixed by f_W , therefore, it only remains to set the edges where at least one endpoint is a non-terminal. Since every non-terminal has degree two in M_W , there are at most $(s+w)^2$ choices for each non-terminal, and at most $(s+w)^{2(s+w)}$ choices in total. This part can also be upper-bounded by $(s+h)^{\mathcal{O}(s+h)} = 2^{(s+h)^{1+o(1)}}$. The overall running time is therefore at most $2^{(s+h+k)\mathcal{O}(k)} \cdot |G|^{\mathcal{O}(1)}$. \square

2.3 Proof of Theorem 1

We are now ready to show the main result of this section, which we restate for convenience.

Theorem 1. *There is an algorithm that, given graphs H, G , a list assignment $L : V(H) \rightarrow 2^{V(G)}$ and an integer parameter $k \geq 1$, in time $2^{|H|\mathcal{O}(k)} \cdot 2^{k\mathcal{O}(k^2)} \cdot |G|^{\mathcal{O}(1)}$*

- either correctly reports that there is no list TM-embedding of H in G , or
- computes a TM-embedding of H in G of the maximum size, where the maximum is taken over all list TM-embeddings of H in G , or
- outputs an independent set in G of size k .

Input: Graphs G and H , mapping $L : V(H) \rightarrow 2^{V(G)}$, integer k .
Output: An independent set of size k in G , or a list TM-embedding of H in G respecting L , or the output that no such TM-embedding exists.

```

1  $S \leftarrow \emptyset$ ;
2 for  $j \leftarrow 1$  to  $\infty$  do
3    $C_1, \dots, C_t \leftarrow$  connected components of  $G \setminus S$ ;
4   if  $t \geq k$  then
5      $I \leftarrow \{v_1, \dots, v_k\}$ , where  $v_i \in C_i$ ;
6     return  $I$ .
7   end
8    $\kappa_j \leftarrow 3 \cdot \max\{k + 2, 10\} \cdot (|V(H)| + |E(H)| + |S|)$ ;
9   if for each  $i \in [t]$ ,  $C_i$  is  $\kappa_j$ -connected then
10    break;
11  else
12    pick  $i$  s.t.  $C_i$  is not  $\kappa_j$ -connected;
13    find minimum separator  $S'$  in  $G[C_i]$ ;
14     $S \leftarrow S \cup S'$ ;
15  end
16 end
17  $O \leftarrow$  invoke Lemma 3 on  $G, H, L, k, S$ ;
18 return  $O$ .
```

Algorithm 2: Algorithm of Theorem 1.

Proof. We iteratively construct the vertex set S suitable for Lemma 3, then invoke Lemma 3. We start by setting $S = \emptyset$, then iteratively find a separator in one of the connected components of $G - S$, and add it to S . This happens until the connected components of $G - S$ are all κ -connected, where $\kappa = 3 \cdot \max\{k + 2, 10\} \cdot (|V(H)| + |E(H)| + |S|)$. If, at some point, there are at least k connected components, we return the independent set constructed by taking a vertex from each connected component. Otherwise, when the connectivity condition is fulfilled, we run Lemma 3 on the given input with the computed S , and return the output. See Algorithm 2 for a detailed algorithm description.

We now argue that the algorithm is correct. If the algorithm returns at Line 6, then the output is an independent set in G of size k . Otherwise, the algorithm eventually reaches Line 17, since at every iteration of the main loop, the size of S increases by at least one. It is left to verify that the call to Lemma 3 is valid. For that, we need that every connected component of $G - S$ is κ -connected, for $\kappa = 3 \cdot \max\{k + 2, 10\} \cdot (|V(H)| + |E(H)| + |S|)$. However, this is exactly the condition at Line 10, and this is the only option for the algorithm to break out of the main loop while reaching Line 6. Therefore, the algorithm's output is correct by Lemma 3.

It remains to upper-bound the running time of the algorithm. For that, it is crucial to upper-bound the size of S when the algorithm reaches Line 17. First, we observe that the main loop

performs at most k iterations before it stops either at Line 6 or at Line 10. Specifically, at the start of the j -th iteration, the number of connected components in $G - S$ is at least j , which implies that at iteration k , the loop necessarily stops at Line 6. This holds since each iteration increases the number of connected components by at least one, since a separator of one of the former components is added into S , while the remaining components are unchanged.

We now also observe that on the j -th iteration, the size of S increases by at most κ_j , since we add to S the minimum separator S' of some $G[C_i]$, and $G[C_i]$ is not κ_j -connected. Let s_j be the size of S at the start of the j -th iteration, and let $\alpha = 3 \cdot \max\{k + 2, 10\}$. We show the following bound.

Claim 12. *It holds that $s_j \leq (\alpha + 2)^j \cdot h$, where $h = |V(H)| + |E(H)|$.*

Proof of Claim 12. We show the claim by induction. The claim holds for $j = 1$ since $s_1 = 0$. Assume the claim holds for s_j , consider s_{j+1} :

$$\begin{aligned} s_{j+1} &\leq s_j + \kappa_j = s_j + \alpha \cdot (s_j + h) = (\alpha + 1) \cdot s_j + \alpha \cdot h \\ &\leq (\alpha + 1) \cdot (\alpha + 2)^j \cdot h + \alpha \cdot h \leq (\alpha + 2)^{j+1} \cdot h, \end{aligned}$$

where the last inequality holds since $\alpha \cdot h \leq (\alpha + 2)^j \cdot h$ for $j \geq 1$. \square

Since the number of iterations is at most k , the size of S by Line 17 is then at most $s_k = k^{\mathcal{O}(k)} \cdot h$. We now upper-bound the running time of Algorithm 1 by the running time bound of Lemma 3, since clearly up to Line 17 the algorithm takes polynomial time. By Lemma 3, the total running time is thus at most $2^{(hk^{\mathcal{O}(k)})^{\mathcal{O}(k)}} \cdot |G|^{\mathcal{O}(1)} = 2^{|H|^{\mathcal{O}(k)}} \cdot 2^{k^{\mathcal{O}(k^2)}} \cdot |G|^{\mathcal{O}(1)}$. \square

3 Proof of Theorem 2: Path cover below Gallai-Milgram

In this section, we prove Theorem 2. Recall that in the PATH COVER problem, we are given a graph G and integer $p \geq 1$, and the task is to decide whether it is possible to cover G with at most p vertex-disjoint paths. Throughout this section, by $\text{pc}(G)$, we denote the minimum number of vertex-disjoint paths required to cover G . A well-known theorem by Gallai and Milgram [GM60] guarantees that $\text{pc}(G) \leq \alpha(G)$. When we combine this theorem with Theorem 1, we obtain an algorithm solving PATH COVER in time $2^{\alpha(G)^{\mathcal{O}(\alpha(G)^2)}} \cdot |G|^{\mathcal{O}(1)}$. Our contribution in Theorem 2 makes a step further by establishing the tractability of PATH COVER for parameterization $k = \alpha(G) - p$, that is “below” the bound of Gallai-Milgram’s theorem. We restate Theorem 2 here for the reader’s convenience.

Theorem 2. *There is an algorithm that, given an n -vertex graph G and an integer parameter $k \geq 1$, in time $2^{k^{\mathcal{O}(k^4)}} \cdot n^{\mathcal{O}(1)}$ outputs a path cover \mathcal{P} of G and, furthermore,*

- *either correctly reports that \mathcal{P} is a minimum-size path cover, or*
- *outputs together with \mathcal{P} an independent set of size $|\mathcal{P}| + k$ certifying that \mathcal{P} contains at most $\alpha(G) - k$ paths.*

We require two auxiliary results to prove Theorem 2. The first result is dealing with small vertex separators of G . It provides a useful tool that for a small vertex separator S of G , reduces the number of connected components in $G - S$ and the value of $\text{pc}(G)$ simultaneously.

Lemma 4. *Let G be a graph, $S \subseteq V(G)$, and let $C_1, C_2, \dots, C_t \subseteq V(G) \setminus S$ be disjoint sets such that for each $i \in [t]$, C_i is a clique of G inducing a connected component of $G - S$. Then there is a polynomial time algorithm that, given G , S , and $\{C_i\}_{i \in [t]}$, outputs a set of at most $2|S|^2$ indices $X \subseteq [t]$ such that G has a minimum path cover containing a path with vertex set C_i for each $i \in [t] \setminus X$.*

Proof. The construction of X is based on the following observation. We say that a path P in a path cover \mathcal{P} of G is *degenerate* if $V(P) = C_i$ for some $i \in [t]$.

Claim 13. *In any minimum path cover \mathcal{P} of G , at most $2|S|$ of the sets C_1, C_2, \dots, C_t contain vertices of non-degenerate paths of \mathcal{P} .*

Proof of Claim 13. Notice that if \mathcal{P} has no path with an edge uv with $u \in S$ and $v \in C_i$ for some $i \in [t]$ then the clique C_i is covered by a degenerate path by the minimality of \mathcal{P} . Hence, if C_i contains a vertex of a non-degenerate path then \mathcal{P} has a path P with an edge with one endpoint in S and the other in C_i . Each vertex in S is incident to at most two edges in the paths from \mathcal{P} . Thus, at most $2|S|$ edges are in the path of \mathcal{P} with one endpoint in S and the other in one of C_1, C_2, \dots, C_t . Hence, at most $2|S|$ of the sets can contain vertices of non-degenerate paths. \square

To construct X , for each vertex $v \in S$, we *mark* at most $2|S|$ cliques from $\{C_i\}_{i \in [t]}$ that contain neighbors of v . If there are at most $2|S|$ such cliques, then we mark all of them. Otherwise, we mark arbitrary $2|S|$ cliques containing neighbors of v . Then X is defined as the set of indices of the marked cliques. It is straightforward that $|X| \leq 2|S|^2$ and X can be constructed in polynomial time. We claim that there is a minimum path cover \mathcal{P} of G such that for each $i \in [t] \setminus X$, there is a path $P \in \mathcal{P}$ satisfying $V(P) = C_i$. In other words, each unmarked clique is covered by a degenerate path.

Given a path cover \mathcal{P} of G , we say that an edge uv of a path $P \in \mathcal{P}$ is *bad* if $u \in S$ and v is a vertex of an unmarked clique. We show that there is a minimum path cover \mathcal{P} of G that does not contain paths with bad edges. The proof is by contradiction. We choose \mathcal{P} as a minimum path cover of G that minimizes the total number of bad edges in the paths. Suppose that there is $P \in \mathcal{P}$ that contain a bad edge uv with $u \in S$ and $v \in C_i$ for some $i \in [t]$. We have that C_i is unmarked and has a neighbor of u . Then, by the definition of the marking procedure, there are at least $2|S|$ marked cliques with neighbors of u . Because C_i contains a vertex of the non-degenerate path P , we obtain that there is a marked clique C_j for some $j \in [t]$ with a neighbor w of u such that C_j is covered by a degenerate path $P' \in \mathcal{P}$ by Claim 13. Because C_j is a clique, we can assume without loss of generality that w is an end-vertex of P' . Deleting the edge uv separates P into two paths Q and Q' having end-vertices v and u , respectively. We use the fact that $uw \in E(G)$ and construct the path \hat{Q} from Q' and P' by joining their end-vertices u and w by the edge uw . Notice that Q and \hat{Q} are vertex-disjoint and $V(Q) \cup V(\hat{Q}) = V(P) \cup V(P')$. This allows us to construct the path cover \mathcal{P}' by replacing P and P' in \mathcal{P} by Q and \hat{Q} . Because the number of paths in \mathcal{P}' is the same as in \mathcal{P} , \mathcal{P}' is a minimum path cover. However, the number of bad edges in the path of \mathcal{P}' is less than the number of bad edges in the paths of \mathcal{P} , contradicting the choice of \mathcal{P} . This proves that there is a minimum path cover \mathcal{P} of G that does not contain paths with bad edges.

To complete the proof, we remind that if a minimum path cover has no path with an edge uv with $u \in S$ and $v \in C_i$ for some $i \in [t]$ then the clique C_i is covered by a degenerate path. Thus, there is a minimum path cover \mathcal{P} of G such that for each $i \in [t] \setminus X$, there is a path $P \in \mathcal{P}$ satisfying $V(P) = C_i$. \square

Our following auxiliary result is a “lighter version” of Theorem 2. It deals with the particular case of $\text{pc}(G) < k^C$.

Lemma 5. *There is an algorithm with running time $2^{k^{\mathcal{O}(k^{2C})}} \cdot n^{\mathcal{O}(1)}$ that, given an n -vertex graph G , integer $k \geq 0$ and a path cover of G with at most $\mathcal{O}(k^C)$ paths for constant $C \geq 1$, outputs a path cover \mathcal{P} of G and*

- *either correctly reports that \mathcal{P} is a minimum-size path cover, or*
- *outputs together with \mathcal{P} an independent set of size $|\mathcal{P}| + k$ certifying that \mathcal{P} contains at most $\alpha(G) - k$ paths.*

Proof. We present an algorithm \mathcal{A} . The algorithm assumes that G contains at least one edge as, otherwise, $pc(G) = n$ and the minimum-size path cover consists of trivial paths.

Given G, k and a path cover \mathcal{P}' of G of size p' , the algorithm \mathcal{A} tries to find an optimal path cover of G . To achieve that, the algorithm iterates an integer i from 1 up to p' and an integer j from 0 to $i - 1$. On each iteration, the algorithm constructs a graph H_{ij} defined as the disjoint union of j copies of K_1 and $i - j$ copies of K_2 , that is, G contains j isolated vertices and $i - j \geq 1$ edges forming a matching. Notice that G admits a path cover with i paths if and only if H_{ij} has a spanning TM-model in G for some $j \in \{0, \dots, i - 1\}$ because G has at least one edge and, therefore, at least one path in a path cover of minimum size is nontrivial. The algorithm \mathcal{A} thus applies the algorithm of Theorem 1 as a subroutine to H_{ij}, G and trivial list assignment $L \equiv V(G)$, with integer parameter set to $p' + k = \mathcal{O}(k^C)$. The subroutine works in

$$2^{|H_{ij}|^{\mathcal{O}(p'+k)}} \cdot 2^{(p'+k)^{\mathcal{O}((p'+k)^2)}} \cdot n^{\mathcal{O}(1)} = 2^{k^{\mathcal{O}(k^{2C})}} \cdot n^{\mathcal{O}(1)}$$

running time.

If the outcome of the subroutine is an independent set I in G of size $p' + k$, then the algorithm outputs \mathcal{P}' and I as the solution. Since $|\mathcal{P}'| = p'$, $|I| \geq |\mathcal{P}'| + k$ is satisfied, so this solution satisfies the lemma statement.

If the subroutine decides that H_{ij} does not have TM-model in G or the maximum TM-model of H_{ij} in G is not spanning, then G cannot be covered with i paths and \mathcal{A} continues to the next iteration. If the subroutine gives a spanning TM-model of H_{ij} in G , then the minimum path cover size of G equals i and \mathcal{A} transforms this TM-model into a path cover of G with i paths. Then \mathcal{A} outputs this optimal path cover of G and stops.

Note that \mathcal{A} necessarily stops and produces a correct solution at some iteration, as it iterates i over $\{1, 2, \dots, p'\}$ while $1 \leq pc(G) \leq p'$. The proof of the lemma is complete. \square

We are ready to prove the main result of this section.

Proof of Theorem 2. We describe an algorithm that starts from a trivial path cover of G and then gradually transforms it. These transformations are indicated as *reduction rules* and are performed in polynomial time. We show that each such rule does not increase the number of paths in the path cover. Moreover, each reduction rule necessarily reduces either the total number of paths or the number of *special* (to be defined later) paths in the path cover.

A series of consecutive applications of the reduction rules always lead the algorithm to cases when the algorithm can produce a solution satisfying theorem statement in $f(k) \cdot n^{\mathcal{O}(1)}$ time and stop. We indicate such cases and the corresponding behavior of the algorithm as *solution subroutines*. The algorithm runtime thus always ends with a solution subroutine run.

In what follows, we present the *reduction rules* and *solution subroutines*. Each of the reduction rules reduces the number of paths in a path cover or reduces the number of some special paths. Solution subroutines output path covers and independent sets. The rules and subroutines are given in the order in which the algorithm checks their applicability. That is, the algorithm applies a

reduction rule or solution subroutine only if no reduction rule or solution subroutine before it can be applied.

When we reach the situation that the obtained path cover \mathcal{P} and graph G that are irreducible, we deduce that the instance has nice structural properties. In this case, the final solution subroutine calls the algorithms of Lemmas 4 and 5.

The algorithm starts with initializing the path cover \mathcal{P} with $|V(G)|$ paths of length 0. Suppose that at some stage of the algorithm, we have constructed a path cover with m paths for some $m \geq 1$. Let the current set of paths be $\mathcal{P} = \{P_1, P_2, \dots, P_m\}$. We denote the endpoints of P_i by s_i and t_i .

The first reduction rule is straightforward and comes from the proof of the Gallai-Milgram theorem for undirected graphs.

Reduction Rule 1. If there is an edge uv in G such that $u \in \{s_i, t_i\}$ and $v \in \{s_j, t_j\}$ and $i \neq j$, then join P_i and P_j into a single path via uv .

The correctness of Reduction Rule 1 is trivial, and it reduces the number of paths in \mathcal{P} by one. Clearly, the exhaustive application of Reduction Rule 1 to \mathcal{P} yields $m \leq \alpha(G)$, as $\{s_1, s_2, \dots, s_p\}$ induces an independent set in G . When Reduction Rule 1 cannot be applied, then the only edges in G allowed between the endpoints of paths in \mathcal{P} are the edges of form $s_i t_i$.

Special and usual paths. We call a path P_i in the path cover *special*, if either $s_i = t_i$, or P_i consists of only edge $s_i t_i$, or its endpoints, are connected by the edge $s_i t_i$. If a path P_i is not special, we call it *usual*. We claim that sufficiently many usual paths give a solution to the problem.

Claim 14. *If at least k paths are usual in the path cover $\mathcal{P} = P_1, \dots, P_m$ and Reduction Rule 1 is not applicable, then $m \leq \alpha(G) - k$. Moreover, an independent set I in G of size $m + k$ is computable in polynomial time.*

Proof of Claim 14. We show that G has an independent set I of size $m + k$. Initialize I with $\{s_1, s_2, \dots, s_p\}$. Since Reduction Rule 1 is not applicable, I is independent in G .

Let i_1, i_2, \dots, i_k be k distinct indices of usual paths. Since each P_{i_j} is usual, $t_{i_j} \neq s_{i_j}$ and s_{i_j} is not connected with t_{i_j} . Also, Reduction Rule 1 is not applicable, so there is no edge between t_{i_j} and any vertex in I , while $\{t_{i_1}, t_{i_2}, \dots, t_{i_k}\}$ is also an independent set in G .

Put $t_{i_1}, t_{i_2}, \dots, t_{i_k}$ in I . We obtain an independent set of size $m + k$ in G , consequently $\alpha(G) - k \geq m$. ┘

Claim 14 yields the corresponding solution subroutine for the algorithm.

Solution Subroutine 1. If there are at least k usual paths among $\mathcal{P} = P_1, P_2, \dots, P_m$, output \mathcal{P} and I as in Claim 14.

From now on, we assume that \mathcal{P} has less than k usual paths. The remaining reduction rules deal primarily with special paths. The following rule reduces the total number of paths given that G has an edge between *any* vertex of a special path and an endpoint of an arbitrary other path.

Reduction Rule 2. If P_i is a special path of length at least two and there is an edge in G connecting some $u \in V(P_i)$ with s_j (or t_j) for some $j \neq i$, then

1. Add edge $s_i t_i$ to P_i , so it becomes a cycle;
2. Remove edge uv from P_i , where v is any neighbor of u in P_i , so it becomes an uv -path;
3. Join P_i and P_j via us_j (or ut_j).

Reduction Rule 2 replaces P_i and P_j with a path covering $V(P_i) \cup V(P_j)$, so \mathcal{P} remains a path cover implying that the rule is correct. The following reduction rule deals with an edge between two vertices of distinct special paths. It is demonstrated in Figure 1a.

Reduction Rule 3. If P_i and P_j are special paths of length at least two for $i \neq j$, and there is an edge in G connecting $u_i \in V(P_i)$ and $u_j \in V(P_j)$, then

1. Add edge $s_j t_j$ to P_j , so it becomes a cycle;
2. Remove edge $u_j v_j$ from P_j , so it becomes an $u_j v_j$ -path.
3. Apply Reduction Rule 2 to P_i , P_j and $u_i u_j$.

Again, Reduction Rule 3 replaces P_i and P_j with a path covering $V(P_i) \cup V(P_j)$, thus reducing the number of paths in \mathcal{P} . We claim the following.

Claim 15. *If Reduction Rules 1, 2, and 3 are not applicable, then for each pair of distinct special paths P_i, P_j in \mathcal{P} , there is no edge between $V(P_i)$ and $V(P_j)$ in G .*

Proof of Claim 15. Assume that the reduction rules are not applicable, but there are paths $P_i, P_j \in \mathcal{P}$ with $i \neq j$ with an edge between $u_i \in V(P_i)$ and $u_j \in V(P_j)$. If both P_i and P_j consist of more than two vertices, then Reduction Rule 3 is applicable. Thus, without loss of generality, P_j consists of one or two vertices, so $V(P_j) = \{s_j, t_j\}$ and $u_j \in \{s_j, t_j\}$. If P_i consists of more than two vertices, then Reduction Rule 2 is applicable, which is a contradiction. Then, the length of P_i is less than three, implying that $u_i \in \{s_i, t_i\}$. In this case, Reduction Rule 1 is applicable, leading us to the final contradiction. \lrcorner

The claim is the basis for the next solution subroutine, which the algorithm applies next. We explain how it works by proving the following claim.

Claim 16. *If Solution Subroutine 1 and Reduction Rules 1, 2, and 3 are not applicable, and there are at least $2k$ special paths P_i such that $V(P_i)$ is not a clique in G , then $m < \alpha(G) - k$. Moreover, an independent set I in G of size more than $m + k$ is computable in polynomial time.*

Proof of Claim 16. We construct an independent set I consisting of more than $m + k$ vertices. Since Solution Subroutine 1 is not applicable, there are more than $m - k$ special paths. For each special path P_i such that $V(P_i)$ is not a clique, put two independent vertices of $V(P_i)$ in I . For each other special path, put its arbitrary vertex into I .

By Claim 15, I is an independent set in G . Since there are at least $2k$ non-cliques among vertex sets of special paths, $|I| > (m - k) + 2k = m + k$. \lrcorner

The solution subroutine based on Claim 16 is given below. The algorithm applies it when none of the reduction rules and solution subroutines above are applicable.

Solution Subroutine 2. If there are at least $2k$ special paths P_i such that $V(P_i)$ is not a clique in G , then output \mathcal{P} and an I as in Claim 16.

The algorithm then exploits the edges between special and usual paths in \mathcal{P} . The following reduction rule necessary reduces the number of special paths in \mathcal{P} while not increasing the total number of paths in \mathcal{P} .

Reduction Rule 4. If P_i and P_j are special paths for $i \neq j$, and there is a usual path P_ℓ , and there are two edges $u_i x_i, u_j x_j$ in G with $u_i \in V(P_i), u_j \in V(P_j), x_i, x_j \in V(P_\ell)$ and x_i goes before x_j in P_ℓ , then




Figure 1: Illustration of transformations performed in (a) Reduction Rule 3 and (b) Reduction Rule 4. Paths before the transformations are horizontal. Paths after the transformations are highlighted in bold and have pairwise-distinct colors. Dashed lines represent absence of an edge. Snake lines represent paths instead of single edges.

1. Make u_i endpoint of P_i and u_j endpoint of P_j similarly to the first two steps of Reduction Rule 2;
2. Break P_ℓ into three (or two) parts: the prefix part $s_\ell P_\ell x_i$, the suffix part $x_j P_\ell t_\ell$, and (if exists) the remaining middle part;
3. Join P_i with the prefix part via $u_i x_i$;
4. Join P_j with the suffix part via $u_j x_j$.

Figure 1b illustrates the result of application of Reduction Rule 4. The following claim addresses its correctness.

Claim 17. *If Reduction Rules 1, 2 are not applicable, then Reduction Rule 4 reduces the number of special paths in \mathcal{P} and does not increase the number of paths in \mathcal{P} .*

Proof of Claim 17. The second part of the claim is trivial, since Reduction Rule 4 produces three (or two, if $x_i x_j \in E(P_\ell)$) paths by transforming three distinct paths P_i, P_j, P_ℓ , so $|\mathcal{P}|$ cannot increase.

To prove the first part, we show that the resulting $v_i s_\ell$ -path and $v_j t_\ell$ -path are usual. We denote these two paths by Q_i and Q_j correspondingly. We give the proof only for Q_i , since the proof for Q_j is symmetric.

We first show that $|V(Q_i)| \geq 3$. Since $\{u_i, v_i, x_i, s_\ell\} \subset V(Q_i)$, it is enough to show that there are at least three distinct vertices among u_i, v_i, x_i, s_ℓ . The only two pairs of vertices that can coincide are u_i, v_i and x_i, s_ℓ , since they belong to disjoint paths P_i and P_ℓ correspondingly. Hence, if $|V(Q_i)| < 3$, then $u_i = v_i$ and $x_i = s_\ell$ both hold. But $u_i = v_i$ implies that P_i has length 0 and $s_i = t_i = u_i$. Then $u_i x_i = s_i s_\ell$, so $s_i s_\ell \in E(G)$, and Reduction Rule 1 is applicable for P_i and P_ℓ . This contradiction proves $|V(Q_i)| \geq 3$.

It is left to show that there is no edge between the endpoints of Q_i , v_i and s_ℓ . Indeed, if $v_i s_\ell \in E(G)$, then we have an edge between a special path P_i and an endpoint of P_ℓ , then Reduction Rule 2 is applicable. This contradiction proves the claim. \square

The list of reduction rules of the algorithm is exhausted. Starting from this point, the algorithm exploits the structure of the graph, assuming that no reduction rule is applicable. In what follows, we explain these structural properties. We also give the last solution subroutine of the algorithm.

Claim 18. *If none of the Reduction Rules 1-5 and Solution Subroutines 1,2 can be applied to \mathcal{P} , then there exists $S \subseteq V(G)$ with $|S| < k$ such that $V(P_i)$ is a connected component in $G - S$ for at least $m - 2k$ special paths P_i . Moreover, such a set S could be constructed in polynomial time.*

Proof of Claim 18. Since Claim 14 is not applicable, there are less than k usual paths in \mathcal{P} . We construct S by taking one or zero vertices from each usual path, so $|S| < k$ holds automatically. We now explain the choice of the vertex to put in S for each usual path.

Consider a usual path $P_i \in \mathcal{P}$. We call a vertex $u \in V(P_i)$ a *connector* if there is a special path P_j and a vertex $v \in V(P_j)$ with $uv \in E(G)$. If P_i has exactly one connector, then we put this connector in S . Otherwise, i.e. when P_i has zero or more than one connectors, we do not put any vertex of P_i in S .

We show that at least $m - 2k$ special paths that are isolated in $G - S$. First note that there are at least $m - k$ special paths in \mathcal{P} , since the number of usual paths is bounded.

Assume now that P_j is a special path that is not isolated in $G - S$. Then there is an edge uv with $u \in V(P_i)$, $v \in V(P_j)$, $i \neq j$ and $u \notin S$. By Claim 15, P_i can only be usual. Since u is a connector of P_i and $u \notin S$, P_i has two or more distinct connectors. As Reduction Rule 4 is not applicable, all these connectors can only connect P_i with P_j . Hence, each usual path P_i gives at most one non-isolated special path in $G - S$. Thus, at most k special paths in $G - S$ are not isolated, implying that at least $(m - k) - k = m - 2k$ special paths are isolated in $G - S$. \square

We are ready to present the final solution subroutine which combines all algorithmic results given previously in this section.

Solution Subroutine 3. Perform the following steps:

1. Obtain S according to Claim 18 and put $h := \max\{0, m - 4k\}$;
2. Order paths in \mathcal{P} so for each $i \in [h]$, the vertices $V(P_i)$ induce a complete connected component in $G - S$;
3. Apply the algorithm of Lemma 4 to G , S and $V(P_1), V(P_2), \dots, V(P_h)$ and obtain a set X of at most $2k^2$ indices, where each $i \in X$ satisfies $i \in [h]$;
4. Obtain G' and \mathcal{P}' by removing $V(P_j)$ from G and P_j from \mathcal{P} , for each j such that $j \in [h]$ and $j \notin X$;
5. Apply the algorithm of Lemma 5 as a subroutine to G' , \mathcal{P}' and $k' := 2k$ and obtain a path cover \mathcal{S}' of G' and (possibly) an independent set I' in G' ;
6. Obtain a path cover \mathcal{S} of G by adding all paths in $\mathcal{P} \setminus \mathcal{P}'$ to \mathcal{S}' ;
7. If I' was obtained, remove all vertices of S from I' and for each path $P_j \in \mathcal{P} \setminus \mathcal{P}'$, add one vertex of P_j to I' . Denote the resulting set by I ;
8. Output \mathcal{S} and (if obtained) I .

We conclude the proof with a claim certifying the correctness of the solution subroutine. In contrast with all previous rules and subroutines, Solution Subroutine 3 requires superpolynomial (but FPT in k) running time.

Claim 19. *If Reduction Rules 1-5 and Solution Subroutines 1,2 are not applicable, then Solution Subroutine 3 works in $2^{k^{\mathcal{O}(k^4)}} \cdot |G|^{\mathcal{O}(1)}$ time and produces a feasible solution.*

Proof of Claim 19. First note that at least $m - 4k$ complete connected components are guaranteed in $G - S$ by Claim 18 (S separates $m - 2k$ special paths) and inapplicability of Solution Subroutine 2 (at most $2k$ vertex sets of special paths are not cliques).

We now move on to establish the upper bound on the running time. All steps, except for applying Lemma 5, are carried out in polynomial time. We have

$$|\mathcal{P}'| = |\mathcal{P}| - h + |X| = m - h + |X| \leq 4k + 2k^2 = \mathcal{O}(k^2).$$

By applying Lemma 5 to G', k' and \mathcal{P}' , we obtain the running time $2^{k^{\mathcal{O}(k^4)}} \cdot |G|^{\mathcal{O}(1)}$.

Set \mathcal{S} is obtained from a path cover \mathcal{S}' of G' via adding paths covering $V(G) \setminus V(G')$. Hence it is also a path cover. On the other hand, I is (if obtained) an independent set in G because $I \subseteq V(G - S)$ and I is independent in $G - S$.

Now we prove that either \mathcal{S} is optimal or $|I| - |\mathcal{S}| \geq k$. First, let us consider the scenario where I is not produced. This can occur only when the algorithm of Lemma 5 does not produce I' . Then Lemma 5 guarantees that \mathcal{S}' is an optimal vertex cover of G' and $|\mathcal{S}'| = \text{pc}(G')$. By Lemma 4, removal of $V(P_j)$ for any $j \in [h] \setminus X$ decreases $\text{pc}(G)$ by one, so $\text{pc}(G') = \text{pc}(G) - (h - |X|)$. Since $|\mathcal{P} \setminus \mathcal{P}'| = h - |X|$, we have $|\mathcal{S}| = \text{pc}(G)$. It means that \mathcal{S} is an optimal path cover of G .

Finally, let us consider the situation where I is produced. In this case, the algorithm in Lemma 5 provides \mathcal{S}' and I' with $|I' - \mathcal{S}'| = k' = 2k$. We notice that

$$|I| \geq |I'| - |\mathcal{S}| + |\mathcal{P} \setminus \mathcal{P}'| > |I'| - k + (h - |X|).$$

As discussed in the previous paragraph, $|\mathcal{S}| - |\mathcal{S}'| = h - |X|$. We conclude that

$$|I| - |\mathcal{S}| = 2k + (|I| - |I'|) - (|\mathcal{S}| - |\mathcal{S}'|) > 2k + (-k + h - |X|) + (h - |X|) = k.$$

The feasibility of \mathcal{S} and I is proved. ┘

This completes the proof of Theorem 2. □

4 Conclusion

In conclusion, this paper shows that HAMILTONIAN PATH/CYCLE and PATH COVER are FPT parameterized by $\alpha(G)$. Furthermore, PATH COVER is FPT parameterized “below” the Gallai–Milgram bound.

We conclude with several open research questions. All our results concern undirected graphs. The complexity of all these problems on *directed* graphs of small independence number remains open. For example, we do not know whether HAMILTONIAN CYCLE is NP-complete on directed graphs with an independence number k for some fixed $k \geq 2$. But we cannot exclude that the problem may be in XP or even FPT parameterized by k . Parameterized complexity offers a powerful toolbox for obtaining lower bounds for structural parameterization [CNP⁺11, FGL⁺19, FGLS10, LMS18]. However, all these reductions use gadgets with large independence numbers. Obtaining algorithmic lower bounds for parameterization with small independence number appears more challenging.

A similar set of questions is about the analogue of Theorem 2 for directed graphs. The theorem of Gallai and Milgram holds for directed graphs. However, whether it could be extended algorithmically or not is not known. The “simplest” question here is whether there is a polynomial-time algorithm for computing a path cover of size $\alpha(G) - 1$?

References

- [ACP⁺21] Tara Abrishami, Maria Chudnovsky, Marcin Pilipczuk, Pawel Rzazewski, and Paul D. Seymour. Induced subgraphs of bounded treewidth and the container method. In *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1948–1964. SIAM, 2021.
- [AYZ95] Noga Alon, Raphael Yuster, and Uri Zwick. Color-coding. *J. ACM*, 42(4):844–856, 1995.
- [BCKN13] Hans L. Bodlaender, Marek Cygan, Stefan Kratsch, and Jesper Nederlof. Deterministic single exponential time algorithms for connectivity problems parameterized by treewidth. In *Proceedings of the 40th International Colloquium of Automata, Languages and Programming (ICALP)*, volume 7965 of *Lecture Notes in Comput. Sci.*, pages 196–207. Springer, 2013.
- [BHT12] Andreas Björklund, Thore Husfeldt, and Nina Taslamán. Shortest cycle through specified elements. In *Proceedings of the Twenty-Third Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1747–1753. SIAM, 2012.
- [Bjö14] Andreas Björklund. Determinant sums for undirected hamiltonicity. *SIAM J. Comput.*, 43(1):280–299, 2014.
- [BLS99] Andreas Brandstädt, Van Bang Le, and Jeremy P. Spinrad. *Graph Classes. A Survey*. SIAM Monographs on Discrete Mathematics and Applications. SIAM, Philadelphia, USA, 1999.
- [Bod96] Hans L. Bodlaender. A linear-time algorithm for finding tree-decompositions of small treewidth. *SIAM J. Computing*, 25(6):1305–1317, 1996.
- [BST23] Julien Baste, Ignasi Sau, and Dimitrios M. Thilikos. Hitting minors on bounded treewidth graphs. IV. an optimal algorithm. *SIAM J. Comput.*, 52(4):865–912, 2023.
- [CCH⁺16] Rajesh Chitnis, Marek Cygan, MohammadTaghi Hajiaghayi, Marcin Pilipczuk, and Michal Pilipczuk. Designing FPT algorithms for cut problems using randomized contractions. *SIAM J. Comput.*, 45(4):1171–1229, 2016.
- [CE72] V. Chvátal and P. Erdős. A note on Hamiltonian circuits. *Discrete Math.*, 2:111–113, 1972.
- [CFK⁺15] Marek Cygan, Fedor V. Fomin, Łukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michał Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015.
- [CKN13] Marek Cygan, Stefan Kratsch, and Jesper Nederlof. Fast hamiltonicity checking via bases of perfect matchings. In *Proceedings of the 45th Annual ACM Symposium on Theory of Computing (STOC)*, pages 301–310. ACM, 2013.
- [CNP⁺11] Marek Cygan, Jesper Nederlof, Marcin Pilipczuk, Michał Pilipczuk, Johan M. M. van Rooij, and Jakub Onufry Wojtaszczyk. Solving connectivity problems parameterized by treewidth in single exponential time. In *Proceedings of the 52nd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 150–159. IEEE, 2011.

- [DF99] Rodney G. Downey and Michael R. Fellows. *Parameterized complexity*. Springer-Verlag, New York, 1999.
- [Die17] Reinhard Diestel. *Graph theory*, volume 173 of *Graduate Texts in Mathematics*. Springer-Verlag, Berlin, 5th edition, 2017.
- [Dil50] R. P. Dilworth. A decomposition theorem for partially ordered sets. *Ann. of Math. (2)*, 51:161–166, 1950.
- [ES09] P. Erdős and G. Szekeres. A combinatorial problem in geometry. In *Classic Papers in Combinatorics*, pages 49–56. Birkhäuser Boston, 2009.
- [FF56] L R Ford, Jr and D R Fulkerson. Maximal flow through a network. *Canad. J. Math.*, 8(0):399–404, 1956.
- [FGK⁺23] Fedor V. Fomin, Petr A. Golovach, Tuukka Korhonen, Kirill Simonov, and Giannos Stamoulis. Fixed-parameter tractability of maximum colored path and beyond. In *Proceedings of the 2023 ACM-SIAM Symposium on Discrete Algorithms, (SODA)*, pages 3700–3712. SIAM, 2023.
- [FGL⁺19] Fedor V. Fomin, Petr A. Golovach, Daniel Lokshtanov, Saket Saurabh, and Meirav Zehavi. Clique-width III: hamiltonian cycle and the odd case of graph coloring. *ACM Trans. Algorithms*, 15(1):9:1–9:27, 2019.
- [FGLS10] Fedor V. Fomin, Petr A. Golovach, Daniel Lokshtanov, and Saket Saurabh. Intractability of clique-width parameterizations. *SIAM J. Comput.*, 39(5):1941–1956, 2010.
- [FGSS20] Fedor V. Fomin, Petr A. Golovach, Danil Sagunov, and Kirill Simonov. Algorithmic extensions of Dirac’s theorem. *CoRR*, abs/2011.03619, 2020.
- [FLP⁺20] Fedor V. Fomin, Daniel Lokshtanov, Fahad Panolan, Saket Saurabh, and Meirav Zehavi. Hitting topological minors is FPT. In *Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing (STOC)*, pages 1317–1326. ACM, 2020.
- [FLPS16] Fedor V. Fomin, Daniel Lokshtanov, Fahad Panolan, and Saket Saurabh. Efficient computation of representative families with applications in parameterized and exact algorithms. *J. ACM*, 63(4):29:1–29:60, 2016.
- [FS15] Alexandra Ovetsky Fradkin and Paul D. Seymour. Edge-disjoint paths in digraphs with bounded independence number. *J. Comb. Theory, Ser. B*, 110:19–46, 2015.
- [GHH23] Luyining Gan, Jie Han, and Jie Hu. An algorithmic version of the hajnal–szemerédi theorem, 2023.
- [GKMW11] Martin Grohe, Ken-ichi Kawarabayashi, Dániel Marx, and Paul Wollan. Finding topological subgraphs is fixed-parameter tractable. In *Proceedings of the 43rd ACM Symposium on Theory of Computing (STOC)*, pages 479–488, 2011.
- [GLP⁺21] Peter Gartland, Daniel Lokshtanov, Marcin Pilipczuk, Michal Pilipczuk, and Pawel Rzazewski. Finding large induced sparse subgraphs in c_t -free graphs in quasipolynomial time. In *Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing (STOC)*, pages 330–341. ACM, 2021.

- [GM60] T. Gallai and A. N. Milgram. Verallgemeinerung eines graphentheoretischen Satzes von Rédei. *Acta Sci. Math. (Szeged)*, 21:181–186, 1960.
- [JK23] Nikola Jedlicková and Jan Kratochvíl. Hamiltonian path and hamiltonian cycle are solvable in polynomial time in graphs of bounded independence number. *CoRR*, abs/2309.09228, 2023.
- [JO90] Bill Jackson and Oscar Ordaz. Chvátal and Erdős conditions for paths and cycles in graphs and digraphs. A survey. *Discrete Math.*, 84(3):241–254, 1990.
- [Kaw08] Ken-ichi Kawarabayashi. An improved algorithm for finding cycles through elements. In *Proceedings of the 13th International Conference on Integer Programming and Combinatorial Optimization (IPCO)*, volume 5035 of *Lecture Notes in Comput. Sci.*, pages 374–384. Springer, 2008.
- [KK22] Karthik C. S. and Subhash Khot. Almost polynomial factor inapproximability for parameterized k -clique. In *Proceedings of the 37th Computational Complexity Conference (CCC)*, volume 234 of *LIPICs*, pages 6:1–6:21. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2022.
- [Kor21] Tuukka Korhonen. A single-exponential time 2-approximation algorithm for treewidth. In *Proceedings of the 62nd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 184–192. IEEE, 2021.
- [Kou08] Ioannis Koutis. Faster algebraic algorithms for path and packing problems. In *Proceedings of the 35th International Colloquium on Automata, Languages and Programming (ICALP)*, volume 5125 of *Lecture Notes in Comput. Sci.*, pages 575–586. Springer, 2008.
- [LLM⁺20] William Lochet, Daniel Lokshtanov, Pranabendu Misra, Saket Saurabh, Roohani Sharma, and Meirav Zehavi. Fault tolerant subgraphs with applications in kernelization. In *Proceedings of the 11th Innovations in Theoretical Computer Science Conference (ITCS)*, volume 151 of *LIPICs*, pages 47:1–47:22. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- [LMS18] Daniel Lokshtanov, Dániel Marx, and Saket Saurabh. Known algorithms on graphs of bounded treewidth are probably optimal. *ACM Trans. Algorithms*, 14(2):13:1–13:30, 2018.
- [LRSW22] Bingkai Lin, Xuandi Ren, Yican Sun, and Xiuhan Wang. On lower bounds of approximating parameterized k -clique. In *Proceedings of the 49th International Colloquium on Automata, Languages, and Programming (ICALP)*, volume 229 of *LIPICs*, pages 90:1–90:18. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2022.
- [LRSZ18] Daniel Lokshtanov, M. S. Ramanujan, Saket Saurabh, and Meirav Zehavi. Reducing CMSO model checking to highly connected graphs. In Ioannis Chatzigiannakis, Christos Kaklamanis, Dániel Marx, and Donald Sannella, editors, *45th International Colloquium on Automata, Languages, and Programming, ICALP 2018, July 9-13, 2018, Prague, Czech Republic*, volume 107 of *LIPICs*, pages 135:1–135:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2018.

- [Men27] Karl Menger. Zur allgemeinen kurventheorie. *Fundamenta Mathematicae*, 10:96–115, 1927.
- [MSSZ23] Pranabendu Misra, Saket Saurabh, Roohani Sharma, and Meirav Zehavi. Sub-exponential time parameterized algorithms for graph layout problems on digraphs with bounded independence number. *Algorithmica*, 85(7):2065–2086, 2023.
- [NW71] C. St. J. A. Nash-Williams. Edge-disjoint Hamiltonian circuits in graphs with vertices of large valency. In *Studies in Pure Mathematics (Presented to Richard Rado)*, pages 157–183. Academic Press, London-New York, 1971.
- [Ram30] F. P. Ramsey. On a problem of formal logic. *Proceedings of the London Mathematical Society*, s2-30(1):264–286, 1930.
- [RS86] Neil Robertson and Paul D. Seymour. Graph minors. II. Algorithmic aspects of tree-width. *J. Algorithms*, 7(3):309–322, 1986.
- [RS03] Vijay Raghavan and Jeremy P. Spinrad. Robust algorithms for restricted domains. *J. Algorithms*, 48(1):160–172, 2003.
- [TW05] Robin Thomas and Paul Wollan. An improved linear edge bound for graph linkages. *European Journal of Combinatorics*, 26(3-4):309–324, April 2005.
- [Wah13] Magnus Wahlström. Abusing the Tutte matrix: An algebraic instance compression for the K -set-cycle problem. In *Proceedings of the 30th International Symposium on Theoretical Aspects of Computer Science (STACS)*, volume 20 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 341–352. Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2013.
- [Wil09] Ryan Williams. Finding paths of length k in $O^*(2^k)$ time. *Inf. Process. Lett.*, 109(6):315–318, 2009.
- [Zeh16] Meirav Zehavi. A randomized algorithm for long directed cycle. *Inf. Process. Lett.*, 116(6):419–422, 2016.