## Real-time observables in de Sitter thermodynamics

## Manvir Grewal<sup>a</sup> and Y.T. Albert Law<sup>b,c,d</sup>

<sup>a</sup> Center for Theoretical Physics, Columbia University, New York, NY 10027, USA
 <sup>b</sup> Center for the Fundamental Laws of Nature, Harvard University, Cambridge, MA 02138, USA
 <sup>c</sup> Black Hole Initiative, Harvard University, Cambridge, MA 02138, USA
 <sup>d</sup> Stanford Institute for Theoretical Physics, 382 Via Pueblo, Stanford, CA 94305, USA

manvir.grewal@columbia.edu , ytalaw@stanford.edu

#### Abstract

We study real-time finite-temperature correlators for free scalars of any mass in a  $dS_{d+1}$  static patch in any dimension. We show that whenever the inverse temperature is a rational multiple of the inverse de Sitter temperature, certain Matsubara poles of the symmetric Wightman function disappear. At the de Sitter temperature, we explicitly show how the Lorentzian thermal correlators can all be obtained by analytic continuations from the round  $S^{d+1}$ . We establish the precise relation between the Harish-Chandra character for SO(1, d+1) and the integrated spectral function, providing a novel dynamical perspective on the former and enabling generalizations. Furthermore, we study scalars with exceptional non-positive masses. We provide a physical picture for the distinctive structures of their characters. For the massless case, we perform a consistent static patch quantization, and find the unique  $S^{d+1}$  correlator that analytically continues to the correlators in the quantum theory.

## Contents

1	Introduction	2
2	Real-time static patch correlators	4
	2.1 Causal Green functions	6
	2.2 Analytic structures of worldline thermal correlators	8
3	Static patch correlators from $S^{d+1}$	11
	3.1 Worldline thermal correlators as resonance sums	11
	3.2 Analytic continuation from $S^{d+1}$	12
4	SO(1,d+1) Harish-Chandra character: a static patch viewpoint	13
	4.1 Scattering matrices and relative spectral density	14
	4.2 Relation with the spectral function	16
	4.3 Revisiting principal and complementary characters	19
5	Comments on the exceptional scalars	20
	5.1 Massless scalar	20
	5.2 General shift-symmetric scalars	26
	5.3 Harish-Chandra characters for the exceptional scalars	28
6	Outlook	29
A	Canonical quantization for free fields on the static patch	31
В	Different correlators in thermal field theory	32
$\mathbf{C}$	Spectral theory in 2D Rindler space	35

## 1 Introduction

Recent years have witnessed significant progress in understanding the semiclassical description of gravity and matter within a static patch of de Sitter (dS) space. For example, extensive calculations of the quantum-corrected dS horizon entropy, using Euclidean gravitational path integrals [1–11], significantly refine the original Gibbons-Hawking proposal [12], paving the way for constraining candidate microscopic models.<sup>1</sup> In parallel, considerable discussions have been devoted to properly understanding the dS horizon entropy in terms of the algebra of observables associated with a static patch [15, 16], stressing the role played by the observer [17].

Anticipating connecting these approaches and generalizations to interacting theories and beyond, we explore an object that played a key role in [1]: the Harish-Chandra character  $\chi(g)$ ,

<sup>&</sup>lt;sup>1</sup>In  $dS_3$ , the dS entropy up to the logarithmic correction is reproduced by a  $T\bar{T} + \Lambda_2$ -deformed CFT<sub>2</sub> [13, 14].

defined as a trace of a group element g over the representation space associated with UIRs of the isometry group SO(1, d+1) of  $dS_{d+1}$ . The character associated with the dS boost,  $g = e^{-i\hat{H}t}$ , admits an expansion in terms of quasinormal modes (QNMs) in the static patch:

$$\chi(t) = \sum_{z} D_z e^{-iz|t|} . {1.1}$$

Here z labels QNM frequencies and  $D_z$  their degeneracies. Interpreting the Fourier transform

$$\tilde{\rho}^{\mathrm{dS}}(\omega) \equiv \int_{-\infty}^{\infty} \frac{dt}{2\pi} e^{i\omega t} \chi(t) \tag{1.2}$$

as a spectral density, the thermal canonical partition function for free fields in a static patch can be rendered well-defined, and for the case of scalars and spinors agrees with the 1-loop determinant on  $S^{d+1}$ . In [18, 19], these results were extended to BTZ and Nariai black holes, and the precise meaning of (1.2) was clarified through a generalized Krein-Friedel-Lloyd formula [20],

$$\tilde{\rho}^{dS}(\omega) = \frac{1}{2\pi i} \partial_{\omega} \sum_{l} D_{l}^{d} \left( \log S_{l}(\omega) - \log \bar{S}(\omega) \right) , \qquad (1.3)$$

where  $S_l(\omega)$  and  $\bar{S}(\omega)$  are the scattering phases respectively for the scattering problems associated with the wave equation on the static patch and a reference Rindler-like space. l labels the spherical harmonics  $Y_l$  on  $S^{d-1}$  with degeneracy  $D_l^d$ . A short review of these ideas can be found in [21].

While the group-theoretic object (1.1) via the spectral density (1.2) offers insights into making sense of traces or partition functions, one might wonder if there is a physical picture of (1.1) directly in terms of observables within the static patch that is not intrinsically tied with the full dS isometries SO(1, d + 1). For one thing, a static patch is only invariant under the subgroup  $SO(1,1) \times SO(d)$ ; for another, the putative finite dimensionality of the microscopic Hilbert space [22] suggests that the description in terms of SO(1, d + 1) UIRs is likely an emergent one in the semi-classical limit [23, 24].

The thermal nature of the static patch [25] together with the fact that QNMs correspond to poles for the retarded propagator [17] naturally suggests that the relevant objects are real-time thermal correlators in the static patch. Hence, we start with a general discussion of real-time thermal correlators for a generic massive scalar in section 2. We will see that the spectral or commutator function

$$G^{C}(t, x, \Omega | 0, y, \Omega') \equiv \langle 0 | \left[ \hat{\phi}(t, x, \Omega), \hat{\phi}(0, y, \Omega') \right] | 0 \rangle , \qquad (1.4)$$

which is temperature independent for a free theory ( $|0\rangle$  is the vacuum defined with respect to the static patch Hamiltonian), plays a distinguished role. To probe the dependence on the inverse temperature  $\beta$ , we also study the symmetric Wightman function, and find that a subset of the Matsubara poles disappears whenever  $\beta$  is a rational multiple of the inverse dS temperature  $\beta_{\rm dS} = 2\pi\ell_{\rm dS}$  ( $\ell_{\rm dS}$  is the dS length). In section 3, we explicitly compute various Lorentzian correlators at  $\beta = \beta_{\rm dS}$  and show how they can all be obtained from an appropriate analytic continuation of the  $S^{d+1}$  correlator.

We begin in section 4 by reviewing the basic ideas for the formula (1.3), pointing out the distinction between resonance poles (poles of the retarded function) and scattering poles (poles of the scattering phase). Then, with an analysis inspired by the spectral theory on infinite-area hyperbolic surfaces [26], we establish the strikingly simple relation between the Harish-Chandra character (1.1) and the spectral function (1.4) for any scalars:

$$\chi(t) = 2i\frac{d}{dt}\tilde{\operatorname{tr}}\,\hat{G}^C(t) \,. \tag{1.5}$$

Here  $\hat{G}^{C}(t)$  is the spectral function understood as an integral operator with kernel (1.4). The trace  $\tilde{\mathbf{tr}}$  is implemented by an integration over the spatial slice, regulated in a suitable way. We stress that the right hand side of (1.5) is defined without reference to the dS isometries, and in principle can be extended to interacting theories.

Finally, in section 5, we study exceptional scalars labeled by a non-positive integer

$$\Delta = -k \;, \quad k = 0, 1, \cdots \;. \tag{1.6}$$

Despite the fact that the SO(1, d+1) UIRs corresponding to (1.6) generically appear in the multiparticle Hilbert spaces of massive fields [27, 28], their realizations in terms of local quantum field theories remain to be understood. For the k=0 (massless) case, we show that its static patch quantization works as in the massive case as long as we exclude the zero mode. We also find the unique  $S^{d+1}$  correlator that analytically continues to the various correlators in this quantum theory. For the  $k \geq 1$  (tachyonic) case, the causal correlators grow with time, indicating the need for a drastic modification in order to have a consistent quantization. Lastly, we give a physical picture for the peculiar structure of the Harish-Chandra characters associated with (1.6), which share similar features with (partially) massless fields [1, 29], which include the physically interesting examples of Maxwell or graviton.

Plan of the paper In section 2, we study Lorentzian thermal correlators in the static patch. In section 3, we work out the analytic continuation from  $S^{d+1}$ . In section 4 we establish the relation between the Harish-Chandra character and the spectral function. In section 5 we study scalars with exceptional non-positive masses. We conclude with some remarks in section 6. We review the canonical quantization for free massive scalars in static patch in appendix A, and some basic facts in thermal field theory in appendix B. In appendix C, we work out the spectral theory on 2D Rindler space.

# 2 Real-time static patch correlators

We consider a free scalar of mass  $m^2$ ,

$$S = -\frac{1}{2} \int \sqrt{-g} \left[ (\partial \phi)^2 + m^2 \phi^2 \right] , \qquad (2.1)$$

on a static patch in  $dS_{d+1}$  with  $d \geq 1$ , whose metric is most commonly expressed in the static form

$$ds^{2} = -\left(\ell^{2} - r^{2}\right)dt^{2} + \frac{dr^{2}}{1 - \frac{r^{2}}{\ell^{2}}} + r^{2}d\Omega^{2}, \qquad 0 \le r < \ell.$$
(2.2)

The observer is located at r=0, surrounded by a horizon at  $r=\ell$ , with temperature  $T_{\rm dS}=\frac{1}{2\pi\ell}$ . Throughout this work, we will extensively use the tortoise coordinate  $r=\ell \tanh x$  instead, with which the metric becomes

$$ds^{2} = \ell^{2} \frac{-dt^{2} + dx^{2} + \sinh^{2} x \, d\Omega^{2}}{\cosh^{2} x} \,, \qquad 0 \le x < \infty \,. \tag{2.3}$$

The observer is mapped to x=0 while the horizon is pushed to  $x\to\infty$ . From now on, we set the dS length  $\ell$  to 1. In this section, we focus on the case  $m^2>0$ , i.e. either principal- or complementary-series scalars [29]. We will study the case of exceptional scalars with  $m^2$  of some non-positive integer values in section 5.

After we rescale  $\phi$  and separate the angular dependence with spherical harmonics as<sup>2</sup>

$$\phi(t, x, \Omega) = \xi_l(t, x) Y_l(\Omega) = \tanh^{\frac{1-d}{2}} x \bar{\xi}(t, x) Y_l(\Omega) , \qquad (2.4)$$

the Klein-Gordon equation  $(-\nabla^2 + m^2) \phi = 0$  decomposes into an infinite set of 1D Schrödingerlike equations labeled by the SO(d) quantum number  $l \ge 0$ ,

$$\left(\partial_t^2 - \partial_x^2 + V_l(x)\right)\bar{\xi}_l(t, x) = 0, \qquad 0 \le x < \infty, \tag{2.5}$$

with a simple Pöschl-Teller-like potential

$$V_l(x) = \frac{(d+2l-3)(d+2l-1)}{4\sinh^2 x} - \frac{(2\Delta - d+1)(2\Delta - d-1)}{4\cosh^2 x}.$$
 (2.6)

Here  $\Delta$  is related to the mass through  $m^2 = \Delta(d - \Delta)$ . Since we will often go back and forth between quantities that are related by a  $\tanh^{\frac{1-d}{2}} x$  rescaling as in (2.4), we find it convenient to introduce the bar notation, which should not be confused with complex conjugation.

With the ansatz for the normal modes

$$\xi_l(t,x) = e^{-i\omega t} \,\psi_{\omega l}(x) = e^{-i\omega t} \tanh^{\frac{1-d}{2}} x \,\bar{\psi}_{\omega l}(x) \,, \qquad \omega \in \mathbb{R} \,, \tag{2.7}$$

a general solution to (2.5) is a linear combination

$$\psi_{\omega l}(x) = C_{\omega l}^{\text{n.}} \psi_{\omega l}^{\text{n.}}(x) + C_{\omega l}^{\text{n.n.}} \psi_{\omega l}^{\text{n.n.}}(x) . \qquad (2.8)$$

$$-\nabla_{S^{d-1}}^2 Y_l(\Omega) = l(l+d-2)Y_l(\Omega) , \qquad l \ge 0 ,$$

with degeneracy  $D_l^d = \frac{d+2l-2}{d-2} {d+l-3 \choose d-3}$ . For d=1, the angular dependence is trivial, and the range of x should be extended to  $-\infty < x < \infty$ . For d=2, the spherical harmonics are  $Y_l(\varphi) = \frac{e^{il\varphi}}{\sqrt{2\pi}}, l \in \mathbb{Z}$ . It is straightforward to modify the following discussions to the latter two cases.

<sup>&</sup>lt;sup>2</sup>For  $d \geq 3$ , the spherical harmonics  $Y_l(\Omega)$  on the unit round  $S^{d-1}$  satisfy

Here the (non-)normalizable mode<sup>3</sup>

$$\psi_{\omega l}^{\text{n.}}(x) = \psi_{\omega, 2-d-l}^{\text{n.n.}}(x) = \frac{\operatorname{sech}^{i\omega} x \tanh^{l} x}{\Gamma\left(\frac{d}{2} + l\right)} {}_{2}F_{1}\left(\frac{l + \Delta + i\omega}{2}, \frac{l + \bar{\Delta} + i\omega}{2}; \frac{d}{2} + l; \tanh^{2} x\right)$$
(2.9)

is (ir)regular at the location of the observer (x = 0):

$$\psi_{\omega l}^{\text{n.}}(x \approx 0) \propto x^{l}$$
,  $\psi_{\omega l}^{\text{n.n.}}(x \approx 0) \propto x^{2-d-l}$ . (2.10)

It is also useful to note that the solutions (2.9) are invariant under  $\omega \to -\omega$ :

$$\psi_{\omega l}^{\text{n.}}(x) = \psi_{-\omega l}^{\text{n.}}(x) , \qquad \psi_{\omega, 2-d-l}^{\text{n.n.}}(x) = \psi_{-\omega, 2-d-l}^{\text{n.n.}}(x) .$$
 (2.11)

Using the identity

$${}_{2}F_{1}(a,b;c;z) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} {}_{2}F_{1}(a,b;a+b-c+1;1-z) + \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} (1-z)^{c-a-b} {}_{2}F_{1}(c-a,c-b;c-a-b+1;1-z)$$
(2.12)

(valid for  $|\arg(1-z)| < \pi$ ), one finds that near the horizon  $(x \to \infty)$ 

$$\psi_{\omega l}^{\text{n.}}(x) = \psi_{\omega, 2-d-l}^{\text{n.n.}}(x) \stackrel{x \to \infty}{\approx} B_{\omega l} e^{i\omega x} + B_{-\omega l} e^{-i\omega x} , \qquad B_{\omega l} = \frac{\Gamma(i\omega)}{\Gamma\left(\frac{\Delta + l + i\omega}{2}\right) \Gamma\left(\frac{\bar{\Delta} + l + i\omega}{2}\right)} . \tag{2.13}$$

For a static patch observer not carrying any source, it is natural to set  $C_{\omega l}^{\text{n.n.}} = 0$  and discard the solution  $\psi_{\omega l}^{\text{n.n.}}(x)$  that blows up at x = 0. Viewing each normalizable mode  $\psi_{\omega l}^{\text{n.i.}}(x)$  as a harmonic oscillator (with the overall coefficient  $C_{\omega l}^{\text{n}}$  fixed by normalization with respect to the Klein-Gordon inner product), one can construct a Fock space following the usual canonical quantization procedure:  $\omega$  and l label single-particle eigenstates; multiparticle states are labeled by the associated occupation numbers. We review this construction in appendix A.

### 2.1 Causal Green functions

As a building block for other real-time finite-temperature correlators (reviewed in appendix B), we study the retarded two-point function, which for the free scalar theory coincides with the retarded Green functions associated with the Klein-Gordon equation,

$$\left(-\nabla_x^2 + m^2\right)G^R(t; x, \Omega|y, \Omega') = \frac{1}{\sqrt{-g}}\delta(t)\delta(x - y)\delta^{d-1}(\Omega, \Omega'), \qquad (2.14)$$

$$G^{R}(t; x, \Omega | y, \Omega') = 0, \qquad t < 0,$$
 (2.15)

where without loss of generality we have put one of the two points at t = 0. The advanced Green function  $G^A(t; x, \Omega | y, \Omega')$  is similarly defined but with the retarded condition (2.15) replaced by the

<sup>&</sup>lt;sup>3</sup>For even d, these expressions for  $\psi_{\omega l}^{\text{n.n.}}$  and  $\psi_{\omega l}^{\text{n.n.}}$  are not independent, and  $\psi_{\omega l}^{\text{n.n.}}$  needs to be replaced by a more complicated function. However,  $\psi_{\omega l}^{\text{n.n.}}$  are not required in our subsequent analysis and thus this issue can be neglected.

 $<sup>^4</sup>$ In d=1, both solutions (2.9) are normalizable, and should both be kept as physical excitations.

advanced condition that it is zero for t > 0 instead. It is straightforward to show that the retarded function for (2.5), i.e.

$$\left(\partial_t^2 - \partial_x^2 + V_l(x)\right) \bar{G}_l^R(t; x|y) = \delta(t)\delta(x - y) , \qquad (2.16)$$

is related to the full retarded function (2.14) by

$$G^{R}(t; x, \Omega | y, \Omega') = \tanh^{\frac{1-d}{2}} x \tanh^{\frac{1-d}{2}} y \sum_{l=0}^{\infty} \bar{G}_{l}^{R}(t; x | y) Y_{l}(\Omega) Y_{l}(\Omega') . \tag{2.17}$$

Similarly, all other correlators G have a rescaled version  $G_l$  for each fixed l. In writing sums like  $\sum_{l=0}^{\infty}$ , we implicitly sum over the degenerate spherical harmonics labeled by their magnetic quantum numbers as well, so for each l in the sum there are  $D_l^d$  terms.

We will be interested in the spectral properties of the single-particle Hamiltonian, so it is natural to work in the Fourier space:

$$\left(-\omega^2 - \partial_x^2 + V_l(x)\right) \bar{\mathcal{G}}_l^R(\omega; x|y) = \delta(x - y) . \tag{2.18}$$

Herein we use G and  $\mathcal{G}$  to denote correlators in the time (t) and frequency  $(\omega)$  domains respectively. Since the potential (2.6) exponentially decays as  $x \to \infty$ ,  $\bar{\mathcal{G}}_l^R(\omega; x|y)$  is meromorphic on the complex  $\omega$  plane [30]. The retarded condition (2.15) is implemented by requiring that in the inverse transform

$$\bar{G}_l^R(t;x|y) = \int_C d\omega \, e^{-i\omega t} \bar{\mathcal{G}}_l^R(\omega;x|y) , \qquad (2.19)$$

the contour C lies above all the poles of  $\bar{\mathcal{G}}_l^R(\omega;x|y)$ . Because of time-reversal symmetry, the advanced function is simply related to the retarded function by  $\bar{\mathcal{G}}_l^A(\omega;x|y) = \bar{\mathcal{G}}_l^R(-\omega;x|y)$ . The advanced condition is implemented by requiring that in the inverse transform the contour lies below all the poles of  $\bar{\mathcal{G}}_l^A(\omega;x|y)$ .

From the standard theory of Green functions for ordinary differential equations, we can immediately write down a formula for the retarded function [31]

$$2\pi \bar{\mathcal{G}}_{l}^{R}(\omega; x|y) = \frac{\bar{\psi}_{\omega l}^{\text{n.}}(x)\bar{\psi}_{\omega l}^{\text{Jost}}(y)}{W_{l}^{\text{Jost}}(\omega)}\theta(y-x) + \frac{\bar{\psi}_{\omega l}^{\text{n.}}(y)\bar{\psi}_{\omega l}^{\text{Jost}}(x)}{W_{l}^{\text{Jost}}(\omega)}\theta(x-y) . \tag{2.20}$$

Here  $\bar{\psi}_{\omega l}^{\rm n.}(x)=\tanh^{\frac{1-d}{2}}x\,\psi_{\omega l}^{\rm n.}(x)$  is the rescaled version of (2.9), and

$$\bar{\psi}_{\omega l}^{\text{Jost}}(x) = 2^{i\omega} \operatorname{sech}^{-i\omega} x \tanh^{\frac{d-1}{2}+l} x \,_{2}F_{1}\left(\frac{l+\Delta-i\omega}{2}, \frac{l+\bar{\Delta}-i\omega}{2}; 1-i\omega; \operatorname{sech}^{2} x\right)$$
(2.21)

is the so-called Jost solution that is purely outgoing

$$\bar{\psi}_{i,l}^{\text{Jost}}(x) \stackrel{x \to \infty}{\approx} e^{i\omega x}$$
 (2.22)

near the horizon. The Jost function, i.e. the Wronskian between  $\bar{\psi}_{\omega l}^{\rm n}$  and  $\bar{\psi}_{\omega l}^{\rm Jost}$ ,

$$W_l^{\text{Jost}}(\omega) \equiv \bar{\psi}_{\omega l}^{\text{Jost}}(x) \partial_x \bar{\psi}_{\omega l}^{\text{n.}}(x) - \partial_x \bar{\psi}_{\omega l}^{\text{Jost}}(x) \bar{\psi}_{\omega l}^{\text{n.}}(x) , \qquad (2.23)$$

appearing in (2.20), is independent of x and can be computed at the horizon:

$$W_l^{\text{Jost}}(\omega) = -2i\omega B_{-\omega l} = \frac{2\Gamma(1 - i\omega)}{\Gamma\left(\frac{\Delta + l - i\omega}{2}\right)\Gamma\left(\frac{\bar{\Delta} + l - i\omega}{2}\right)}.$$
 (2.24)

Note that the overall normalization (2.8) of  $\bar{\psi}_{\omega l}^{n}(x)$  does not affect that of (2.20). The locations of the zeros of (2.24), which we refer to as the **resonance** poles,

$$i\omega_{nl}^{\Delta} = \Delta + 2n + l$$
, or  $i\omega_{nl}^{\bar{\Delta}} = \bar{\Delta} + 2n + l$ ,  $0 \le n \in \mathbb{Z}$ , (2.25)

correspond to the poles of (2.20) on the complex  $\omega$ -plane. Similarly, the poles of  $\bar{\mathcal{G}}_l^A(\omega; x|y)$ , i.e. the **anti-resonance** poles, are given by the zeros of  $W_l^{\text{Jost}}(-\omega)$ .

The resonance poles (2.25), as poles of the retarded function, are also known as the quasinormal modes (QNMs) on a static patch. An alternative definition of QNMs characterizes them as being regular at the observer's location and purely outgoing at the horizon. However, these two definitions are not equivalent—a point we will elaborate on in Section 4.1. To clarify this distinction and avoid confusion, we will consistently use the term 'resonance poles.'

Before moving on, we also note that the *poles* of  $W_l^{\mathrm{Jost}}(\omega)$  and  $W_l^{\mathrm{Jost}}(-\omega)$ , corresponding to the zeros of  $\bar{\mathcal{G}}_l^R(\omega;x|y)$  and  $\bar{\mathcal{G}}_l^A(\omega;x|y)$  respectively, are located at

$$i\omega_k^{\mathrm{dS}} = k \in \mathbb{Z} \setminus \{0\} ,$$
 (2.26)

the (anti-)Matsubara frequencies associated with the dS temperature  $\beta_{dS} = 2\pi$ .

### 2.2 Analytic structures of worldline thermal correlators

Starting from the causal functions, we can build all other real-time thermal correlators, as reviewed in appendix B. Without loss of generality, we focus on the behaviors of the correlators near the observer's worldline, i.e. when  $x, y \approx 0$ . Expressing the answer in terms of a dS-invariant distance, the correlators at any two timelike separated points can be obtained.

#### 2.2.1 The spectral function

When  $x, y \approx 0$ , the full retarded function becomes

$$\mathcal{G}^{R}(\omega; x, \Omega | y, \Omega') \approx \sum_{l=0}^{\infty} \mathcal{G}_{l}^{R}(\omega) Y_{l}(\Omega) Y_{l}(\Omega') x^{l} y^{l} , \qquad (2.27)$$

where, according to a prescription motivated by the analogy with AdS/CFT [17],  $\mathcal{G}^R(\omega)$  is proportional to a ratio of the coefficients of the non-normalizable and normalizable modes such that the solution is purely outgoing at the horizon:

$$\mathcal{G}_l^R(\omega) \propto \frac{B_{-\omega,2-d-l}}{B_{-\omega l}}$$
 (2.28)

Alternatively, this can be obtained by taking  $x, y \to 0$  in (2.20), which fixes the overall  $\omega$ independent factor as well. The difference between the retarded and advanced functions gives
the so-called spectral or commutator function:

$$\mathcal{G}_{l}^{C}(\omega) = -i\left(\mathcal{G}_{l}^{R}(\omega) - \mathcal{G}_{l}^{A}(\omega)\right) \propto \sinh \pi \omega \left|\Gamma\left(\frac{\Delta + l - i\omega}{2}\right)\Gamma\left(\frac{\bar{\Delta} + l - i\omega}{2}\right)\right|^{2}.$$
 (2.29)

Other than the (anti-)resonance poles (see Figure 2.1) inherited from  $\mathcal{G}_l^{R/A}(\omega)$ , (2.29) has zeros at

$$i\omega_k^{\mathrm{dS}} = k \;, \qquad k \in \mathbb{Z} \;, \tag{2.30}$$

which are the (anti-)Matsubara frequencies associated with  $\beta_{\rm dS}=2\pi$ , including the  $\omega=0$  mode.




Figure 2.1:  $|\mathcal{G}_{l-1}^C(\omega)|$  for a massive scalar in  $dS_4$  in complex  $\omega$  plane.

The spectral function (2.29) plays a particularly important role because it encodes all the information for the single-particle Hamiltonian, and is directly related to the Harish-Chandra character. We will elaborate on this in section 4.2.

### 2.2.2 Thermal symmetric Wightman function

As reviewed in appendix B, the causal functions (and thus the spectral function) for a free theory at finite inverse temperature  $\beta$  are independent of  $\beta$ . To see temperature dependence, we must study other correlators, such as the thermal symmetric Wightman function:

$$G_{\beta}^{\text{Sym}}(t) \equiv \frac{1}{Z(\beta)} \operatorname{Tr} \left[ e^{-\beta \hat{H}} \left( \frac{\hat{\phi}(t)\hat{\phi}(0) + \hat{\phi}(0)\hat{\phi}(t)}{2} \right) \right] , \qquad Z(\beta) \equiv \operatorname{Tr} e^{-\beta \hat{H}} , \qquad (2.31)$$

where we have suppressed spatial dependence. This is related to the spectral function at a general inverse temperature  $\beta$  through

$$\mathcal{G}_{l}^{\mathrm{Sym}}(\beta,\omega) = \frac{1}{2} \coth \frac{\beta \omega}{2} \mathcal{G}_{l}^{C}(\omega) \propto \coth \frac{\beta \omega}{2} \sinh \pi \omega \left| \Gamma\left(\frac{\Delta + l - i\omega}{2}\right) \Gamma\left(\frac{\bar{\Delta} + l - i\omega}{2}\right) \right|^{2} . \quad (2.32)$$

The factor  $\coth \frac{\beta \omega}{2}$  reflects the fact that the particles obey bosonic statistics. From (2.32), we see that the analytic structure of (2.32) depends on whether  $\beta$  is a rational multiple of  $2\pi$ . Analogous dependence has been pointed out at the level of free energies in [32].

Generic  $\beta$  At a generic inverse temperature  $\beta$ , (2.32) has two kinds of poles. In addition to the resonance and anti-resonance poles inherited from the spectral function (2.29), (2.32) has poles at the Matsubara and anti-Matsubara frequencies associated with the inverse temperature  $\beta$ :

$$i\omega_k^{\text{Mat}} = \frac{2\pi k}{\beta} , \qquad k \in \mathbb{Z} \setminus \{0\} .$$
 (2.33)

An example is shown in figure 2.2a.

When  $\beta = 2\pi \mathbb{Q}$  At the inverse temperatures

$$\beta = \frac{2\pi p}{q} \,, \tag{2.34}$$

where  $p, q \in \mathbb{N}$  are co-prime integers, there are (partial) cancellations between the Matsubara poles (2.33) and the zeros (2.30). Specifically, the would-be Matsubara poles at

$$i\omega_k = kq$$
,  $k \in \mathbb{Z} \setminus \{0\}$ , (2.35)

are absent. An example is shown in figure 2.2b. For p = 1, the Matsubara poles (2.33) are completely absent.

When  $\beta = \beta_{dS} = 2\pi$  In this case of particular interest (see figure 2.2c), (2.32) becomes

$$\mathcal{G}_l^{\mathrm{Sym}}(2\pi,\omega) \propto \cosh \pi \omega \left| \Gamma\left(\frac{\Delta + l - i\omega}{2}\right) \Gamma\left(\frac{\bar{\Delta} + l - i\omega}{2}\right) \right|^2,$$
 (2.36)

where there is a one-to-one cancellation between the Matsubara poles (2.33) and the zeros (2.30).




Figure 2.2:  $\left|\mathcal{G}_{l=1}^{\mathrm{Sym}}(\beta,\omega)\right|$  for a principal-series scalar with  $\Delta=\frac{3}{2}+0.7i$  in  $dS_4$  in complex  $\omega$  plane.

#### 2.2.3 Two-sided function

Before ending this section, we point out that we have been assuming that the 2-sided correlator at the dS temperature  $\beta = 2\pi$ , related to the spectral function by

$$\mathcal{G}_{l}^{12}(\omega) = \frac{\mathcal{G}_{l}^{C}(\omega)}{\sinh \pi \omega} \propto \Gamma\left(\frac{\Delta + l + i\omega}{2}\right) \Gamma\left(\frac{\bar{\Delta} + l + i\omega}{2}\right) \Gamma\left(\frac{\Delta + l - i\omega}{2}\right) \Gamma\left(\frac{\bar{\Delta} + l - i\omega}{2}\right) , \quad (2.37)$$

can be written as an absolute value squared and is thus real and positive. This is not true for general complex values of  $\Delta$ . However, this is indeed the case for  $\Delta$  valued in the principal or complementary series, consistent with expectations from unitarity [33].

Another interesting comment is that using the Weierstrass form of the gamma function,

$$\Gamma(z) = \frac{e^{-\gamma z}}{z} \prod_{n=1}^{\infty} \left(1 + \frac{z}{n}\right)^{-1} e^{z/n} ,$$
 (2.38)

we can recast (2.37) into the product formula

$$\mathcal{G}_{l}^{12}(\omega) = \frac{\mathcal{G}_{l}^{12}(\omega = 0)}{\prod_{n=0}^{\infty} \left(1 - \left(\frac{\omega}{\omega_{nl}^{\Delta}}\right)^{2}\right) \left(1 - \left(\frac{\omega}{\omega_{nl}^{\Delta}}\right)^{2}\right)} . \tag{2.39}$$

A formula of this form was argued by [33] for holographic thermal correlators in AdS/CFT.

# 3 Static patch correlators from $S^{d+1}$

In this section, we explicitly compute the static patch correlators in the previous section and show how they can be obtained from analytically continued  $S^{d+1}$  correlators.

#### 3.1 Worldline thermal correlators as resonance sums

If one restricts the two points to be exactly on the worldline at x = y = 0, SO(d) invariance implies that only the l = 0 mode can contribute to the full worldline correlators:

$$G(t) = \frac{1}{\text{Vol}(S^{d-1})} \int_{-\infty}^{\infty} d\omega \, e^{-i\omega t} \mathcal{G}_{l=0}(\omega) . \tag{3.1}$$

Here G(t) can be any of the correlators discussed in the previous section. The factor  $\frac{1}{\operatorname{Vol}(S^{d-1})}$  comes from the l=0 spherical harmonics  $Y_{l=0}$  in the expansions of the full correlators in terms of the reduced ones, such as in (2.17). A general way to proceed is to close the contour on the lower (upper) half complex  $\omega$ -plane for t>0 (t<0), which then picks up contributions from poles of  $\mathcal{G}_{l=0}(\omega)$ . For example, the retarded function is a sum over resonances:

$$G^{R}(t) = \frac{\Gamma\left(\frac{d}{2} - \Delta\right)}{\Gamma\left(\frac{d}{2}\right)\Gamma\left(1 - \Delta\right)} \theta(t) \sum_{n=0}^{\infty} \frac{1}{n!} \frac{\left(\frac{d}{2}\right)_{n} (\Delta)_{n}}{\left(1 - \frac{d}{2} + \Delta\right)_{n}} e^{-(\Delta + 2n)t} + \left(\Delta \leftrightarrow \bar{\Delta}\right)$$

$$= \theta(t) \frac{\sin\left(\pi\Delta\right)\Gamma\left(\frac{d}{2} - \Delta\right)\Gamma\left(\Delta\right)}{\operatorname{Vol}(S^{d+1})\Gamma\left(\frac{d}{2} + 1\right)} e^{-\Delta t} {}_{2}F_{1}\left(\Delta, \frac{d}{2}, 1 - \frac{d}{2} + \Delta, e^{-2t}\right) + \left(\Delta \leftrightarrow \bar{\Delta}\right) . \tag{3.2}$$

We have used  $Vol(S^{d+1}) = \frac{2\pi}{d} Vol(S^{d-1})$ . Such a resonance/QNM sum was obtained in [34] for complementary-series scalars in  $dS_4$ , where a Fock space was built using the resonances normalized with respect to an 'R-norm'. Our approach bypasses the introduction of the R-norm and treats

both principal and complementary series uniformly. Similarly, the advanced function is a sum over anti-resonances. For completeness, we also write down the spectral function:

$$G^{C}(t) = \frac{-i\sin(\pi\Delta)\Gamma\left(\frac{d}{2} - \Delta\right)\Gamma\left(\Delta\right)}{\operatorname{Vol}(S^{d+1})\Gamma\left(\frac{d}{2} + 1\right)} e^{-\Delta|t|} {}_{2}F_{1}\left(\Delta, \frac{d}{2}, 1 - \frac{d}{2} + \Delta, e^{-2|t|}\right) + \left(\Delta \leftrightarrow \bar{\Delta}\right) . \tag{3.3}$$

As emphasized earlier, these functions are insensitive to  $\beta$ . For the  $\beta$ -sensitive functions such as the symmetric Wightman function  $G_{\beta}^{\text{Sym}}(t)$ , the contour could pick up Matsubara poles as discussed in the previous section. For generic  $\beta$ ,  $G_{\beta}^{\text{Sym}}(t)$  consists of two sums, one over the (anti-)Matsubrara poles and the other one over the (anti-)resonance poles, for which we have not found a closed-form formula. For  $\beta = 2\pi$ , the Matsubara sum is absent, and we have

$$G_{\beta=2\pi}^{\text{Sym}}(t) = \frac{\cos(\pi\Delta)\Gamma\left(\frac{d}{2} - \Delta\right)\Gamma(\Delta)}{2\text{Vol}(S^{d+1})\Gamma\left(\frac{d}{2} + 1\right)}e^{-\Delta|t|} {}_{2}F_{1}\left(\Delta, \frac{d}{2}, 1 - \frac{d}{2} + \Delta, e^{-2|t|}\right) + \left(\Delta \leftrightarrow \bar{\Delta}\right) . \tag{3.4}$$

We note that (3.2), (3.3), and (3.4) are valid for any dimensions  $d \ge 2$ .

## 3.2 Analytic continuation from $S^{d+1}$

We now show how these Lorentzian correlators can be obtained from suitable analytic continuation from the  $S^{d+1}$  Green function,

$$G_E(P(\mathbf{x}, \mathbf{y})) = \frac{\Gamma(\Delta)\Gamma(\bar{\Delta})}{\operatorname{Vol}(S^{d+1})\Gamma(d+1)} {}_{2}F_1\left(\Delta, \bar{\Delta}; \frac{d+1}{2}; \frac{1+P(\mathbf{x}, \mathbf{y})}{2}\right), \quad -1 < P < 1, \quad (3.5)$$

where  $\operatorname{Vol}(S^{d+1}) = \frac{2\pi^{\frac{d+2}{2}}}{\Gamma(\frac{d+2}{2})}$  is the volume for a unit round  $S^{d+1}$ , and  $P(\mathbf{x}, \mathbf{y}) \equiv \cos \sigma(\mathbf{x}, \mathbf{y})$  where  $\sigma(\mathbf{x}, \mathbf{y})$  the geodesic distance between  $\mathbf{x}, \mathbf{y} \in S^{d+1}$ . The Green function (3.5) satisfies the Euclidean equation of motion with a delta-function source,

$$\left(-\nabla_{\mathbf{x}}^2 + m^2\right) G_E(P(\mathbf{x}, \mathbf{y})) = \frac{1}{\sqrt{g}} \delta^{(d+2)}(\mathbf{x}, \mathbf{y}) , \qquad (3.6)$$

and can be represented as a path integral for the free scalar on  $S^{d+1}$ :

$$G_E(P(\mathbf{x}, \mathbf{y})) = \frac{\int \mathcal{D}\phi \,\phi(\mathbf{x})\phi(\mathbf{y}) \,e^{-S_E[\phi]}}{\int \mathcal{D}\phi \,e^{-S_E[\phi]}} \,, \qquad S_E = \frac{1}{2} \int_{S^{d+1}} \sqrt{g} \left[ (\partial\phi)^2 + m^2\phi^2 \right] \,. \tag{3.7}$$

Therefore, from the global dS perspective, (3.5) is the two point function for the free scalar in the Bunch-Davies vacuum where  $\mathbf{x}$  and  $\mathbf{y}$  are spacelike-separated.

From the static patch point of view, since  $S^{d+1}$  can be obtained by Euclideanizing the static patch time  $t \to -i\tau$  followed by the identification  $\tau \sim \tau + 2\pi$  (see figure 3.1), the same quantity (3.7) can be interpreted as a thermal correlator in a static patch at inverse temperature  $\beta_{\rm dS} = 2\pi$ , again with  ${\bf x}$  and  ${\bf y}$  spacelike-separated, i.e.

$$\frac{\int \mathcal{D}\phi \,\phi(\mathbf{x})\phi(\mathbf{y}) \,e^{-S_E[\phi]}}{\int \mathcal{D}\phi \,e^{-S_E[\phi]}} = \frac{1}{Z(\beta_{\mathrm{dS}})} \,\mathrm{Tr} \,\left(e^{-\beta_{\mathrm{dS}}\hat{H}}\hat{\phi}(\mathbf{x})\hat{\phi}(\mathbf{y})\right) \,, \quad Z(\beta) \equiv \mathrm{Tr} \,e^{-\beta\hat{H}} \,. \tag{3.8}$$

We recall that for spacelike separated points, the commutator-type functions are trivial, and all the Wightman-type functions are equivalent.




Figure 3.1: Left: Penrose diagram for a static patch. The yellow dot indicates the bifurcation surface at  $r = \ell$ . Right: Upon the Wick rotation  $t \to -i\tau$ ,  $\tau \sim \tau + 2\pi$ , the static patch becomes the sphere, with  $\tau$  parametrizing the thermal circle.

Analytic continuation to real time To continue to the region where  $\mathbf{x}$  and  $\mathbf{y}$  are timelike-separated, we need to Wick-rotate  $\sigma \to it$ , which takes P to  $P \in (1, \infty)$  where (3.5) has a branch cut. With an  $i\epsilon$  prescription  $P \to P \mp i\epsilon$  ( $\epsilon > 0$ ), we can safely perform the analytic continuation. After a linear transformation,

$${}_{2}F_{1}(a,b,c,z) = \frac{\Gamma(c)\Gamma(b-a)}{\Gamma(b)\Gamma(c-a)}(-z)^{-a} {}_{2}F_{1}\left(a,1-c+a;1-b+a;\frac{1}{z}\right) + \frac{\Gamma(c)\Gamma(a-b)}{\Gamma(a)\Gamma(c-b)}(-z)^{-b} {}_{2}F_{1}\left(b,1-c+b;1-a+b,\frac{1}{z}\right) ,$$
(3.9)

(valid for  $|\arg(-z)| < \pi$ ), and Kummer's quadratic transformation,

$$_{2}F_{1}(a,b,a-b+1,z) = (1+\sqrt{z})^{-2a} \, _{2}F_{1}\left(a,a-b+\frac{1}{2},2a-2b+1,\frac{4\sqrt{z}}{(1+\sqrt{z})^{2}}\right) ,$$
 (3.10)

we find $^5$ 

$$G_E(P \mp i\epsilon) = \frac{e^{\mp i\pi\Delta}\Gamma(\Delta)\Gamma(\frac{d}{2} - \Delta)}{2\text{Vol}(S^{d+1})\Gamma(\frac{d}{2} + 1)}e^{-\Delta|t|} {}_2F_1\left(\Delta, \frac{d}{2}, 1 - \frac{d}{2} + \Delta, e^{-2|t|}\right) + (\Delta \leftrightarrow \bar{\Delta}). \tag{3.11}$$

It is now clear how the Lorentzian  $dS_{d+1}$  correlators are obtained from those on  $S^{d+1}$ . In particular,

$$G^{C}(t) = G_{E}(P - i\epsilon) - G_{E}(P + i\epsilon) , \qquad G_{\beta = 2\pi}^{Sym}(t) = \frac{G_{E}(P - i\epsilon) + G_{E}(P + i\epsilon)}{2} . \qquad (3.12)$$

# 4 SO(1, d+1) Harish-Chandra character: a static patch viewpoint

Particles and fields in  $dS_{d+1}$  are naturally classified according to unitary irreducible representations (UIRs) of the de Sitter group SO(1, d+1) [35]. A SO(1, d+1) UIR  $\mathcal{V}_{[\Delta, \mathbf{s}]}$  is labeled by an SO(1, 1)

<sup>&</sup>lt;sup>5</sup>The  $i\epsilon$  prescription determines whether the phase of -z in the transformation (3.9) is  $+(\pi - \epsilon)$  or  $-(\pi - \epsilon)$ .

weight  $\Delta$  and an SO(d) weight  $\mathbf{s}$ , and can be realized as the single-particle Hilbert space in global quantization [29]. A useful object that encodes the information of a UIR is its Harish-Chandra character, defined as a trace over the representation space,

$$\chi(g) = \operatorname{tr}_{[\Delta,\mathbf{s}]} R(g) , \qquad g \in SO(1, d+1) , \qquad (4.1)$$

viewed as a distribution to be integrated against smooth test functions on SO(1,d+1). The classifications of SO(1,d+1) UIRs and their characters were studied in [36–38] (see [29] for a review). While it is not difficult to consider the character for a general SO(1,d+1) element, we will focus on the element  $g = e^{-i\hat{H}t}$  which generates the SO(1,1) subgroup.  $\hat{H}$  acts on the de Sitter geometry globally as a boost and on a static patch as the Hamiltonian. The character for a scalar with a mass squared  $m^2 = \Delta(d-\Delta) > 0$  reads explicitly

$$\chi(t) = \frac{e^{-\Delta t} + e^{-\bar{\Delta}t}}{|1 - e^{-t}|^d} \,. \tag{4.2}$$

Note that this satisfies the property  $\chi(-t) = \chi(t)$ .

There are two key related observations in [1]. First, the characters admit an expansion in terms of the resonance poles. For scalars, it is evident by expanding (4.2) in powers of  $e^{-|t|}$ :

$$\chi(t) = \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} D_l^d \left( e^{-i\omega_{nl}^{\Delta}|t|} + e^{-i\omega_{nl}^{\bar{\Delta}}|t|} \right) , \qquad (4.3)$$

where  $\omega_{nl}^{\Delta}$  and  $\omega_{nl}^{\Delta}$  are the resonance frequencies (2.25). Characters for more general UIRs also admit such an expansion [39]. The second observation is that the Fourier transform of the character,

$$\tilde{\rho}^{\mathrm{dS}}(\omega) \equiv \int_{-\infty}^{\infty} \frac{dt}{2\pi} e^{i\omega t} \chi(t) , \qquad (4.4)$$

defines a spectral density for the single-particle Hamiltonian. With (4.4), the Lorentzian calculation of the canonical thermal partition function for a scalar on the static patch can be rendered well-defined, and exactly reproduces the Euclidean path integrals on  $S^{d+1}$ .

#### 4.1 Scattering matrices and relative spectral density

In [18], the precise physical meaning of the spectral density (4.4) was clarified and generalized to other static spherically symmetric black holes. We now review the key ideas specializing to the case of a dS static patch, which will set the stage for our main result in section 4.2.

Resonance poles vs scattering poles For a fixed  $l \ge 0$ , the ratio between the outgoing and ingoing coefficients for the normalizable mode (2.13) defines a scattering phase,

$$S_l(\omega) \equiv \frac{B_{\omega l}}{B_{-\omega l}} \,, \tag{4.5}$$

which maps an incoming wave from the past horizon to an outgoing wave to the future horizon:




Figure 4.1: Left: The Penrose diagram for a static patch, with the black arrows indicating incoming and outgoing waves. Right: The scattering potential (2.6) for a principal-series scalar with  $\Delta = \frac{3}{2} + 0.1i$  and l = 3 in  $dS_4$ , which is hardly distinguishable from the Rindler potential (red) in (C.1) as  $x \to \infty$ . In these diagrams, the brown lines indicate a brick wall regulator [40] at x = R = 2.5.

The poles/zeros of (4.5), which we call the **scattering poles/zeros**, correspond to modes that are regular at the observer's location and purely outgoing/incoming at the horizon. Explicitly, (4.5) can be put into the product form

$$\mathcal{S}_{l}(\omega) = \mathcal{S}_{l}^{\mathrm{dS}}(\omega)\mathcal{S}^{\mathrm{Rin}}(2\pi, \omega) ,$$

$$\mathcal{S}_{l}^{\mathrm{dS}}(\omega) \equiv \frac{\Gamma\left(\frac{\Delta + l - i\omega}{2}\right)\Gamma\left(\frac{\bar{\Delta} + l - i\omega}{2}\right)}{\Gamma\left(\frac{\Delta + l + i\omega}{2}\right)\Gamma\left(\frac{\bar{\Delta} + l + i\omega}{2}\right)} , \qquad \mathcal{S}^{\mathrm{Rin}}(2\pi, \omega) = \frac{\Gamma(i\omega)}{\Gamma(-i\omega)} .$$
(4.6)

The poles (zeros) of  $\mathcal{S}_l^{\mathrm{dS}}(\omega)$  capture the (anti-)resonance poles (2.25). Meanwhile, as pointed out in [18] and reviewed in appendix C,  $\mathcal{S}^{\mathrm{Rin}}(2\pi,\omega)$  is exactly the scattering phase associated with the wave equation on a 2D Rindler space with  $\beta = \beta_{\mathrm{dS}} = 2\pi$ . The poles (zeros) of  $\mathcal{S}^{\mathrm{Rin}}(2\pi,\omega)$ ,

$$i\omega_k^{\text{Mat}} = \pm k \;, \qquad k \in \mathbb{N} \;, \tag{4.7}$$

correspond to the (anti-)Matsubara frequencies associated with the dS temperature.

We therefore have an interesting conclusion that (anti-)resonance poles do not coincide with scattering poles (zeros): there exist modes that satisfy the (anti-)QNM boundary conditions yet are not poles of the retarded (advanced) Green functions.

Scattering phases and spectral density While the density of states for a system with a continuous spectrum diverges, its *difference* relative to another system with a continuous spectrum is well-defined. The *relative* spectral density is captured by the (generalized) Krein-Friedel-Lloyd formula

$$\Delta \rho_l(\omega) = \frac{1}{2\pi i} \partial_\omega \left( \log S_l(\omega) - \log \bar{S}(\omega) \right) . \tag{4.8}$$

Here  $\bar{S}(\omega)$  is the scattering phase for the reference problem. An observation in [18] is that if we take the reference problem to be the one associated with the wave equation on the 2D Rindler

space with  $\beta = \beta_{dS} = 2\pi$ , i.e. we take  $\bar{S}(\omega) = S^{Rin}(2\pi, \omega)$ , the resulting spectral density

$$\Delta \rho_l^{\rm dS}(\omega) = \frac{1}{2\pi i} \partial_\omega \log \mathcal{S}_l^{\rm dS}(\omega) \tag{4.9}$$

coincides with the one defined by the Fourier-transform of the Harish-Chandra character (4.4) upon summing over  $l \ge 0$ , i.e.

$$\tilde{\rho}^{\mathrm{dS}}(\omega) = \sum_{l=0}^{\infty} D_l^d \Delta \rho_l^{\mathrm{dS}}(\omega) . \tag{4.10}$$

### 4.2 Relation with the spectral function

We are now ready to accomplish the main goal of this paper: establishing a direct connection between the Harish-Chandra character (4.2) and the correlators studied in detail in section 2. Specifically, the spectral function

$$G^{C}(t, x, \Omega | 0, y, \Omega') \equiv \langle 0 | \left[ \hat{\phi}(t, x, \Omega), \hat{\phi}(0, y, \Omega') \right] | 0 \rangle$$
(4.11)

plays a distinguished role. Our analysis starts by deriving a formula for the reduced version

$$\bar{\mathcal{G}}_{l}^{C}(\omega; x|y) = -i \left( \bar{\mathcal{G}}_{l}^{R}(\omega; x|y) - \bar{\mathcal{G}}_{l}^{A}(\omega; x|y) \right) , \qquad (4.12)$$

by noting that both  $\bar{\mathcal{G}}_l^R(\omega;x|y)$  and  $\bar{\mathcal{G}}_l^A(\omega;x|y)$  satisfy (2.18), implying

$$\bar{\mathcal{G}}_{l}^{R}(\omega;x|y) - \bar{\mathcal{G}}_{l}^{A}(\omega;x|y) = \int_{0}^{\infty} dz \left[ \bar{\mathcal{G}}_{l}^{A}(\omega;y|z) \partial_{z}^{2} \bar{\mathcal{G}}_{l}^{R}(\omega;x|z) - \bar{\mathcal{G}}_{l}^{R}(\omega;x|z) \partial_{z}^{2} \bar{\mathcal{G}}_{l}^{A}(\omega;y|z) \right] \\
= \left[ \bar{\mathcal{G}}_{l}^{A}(\omega;y|z) \partial_{z} \bar{\mathcal{G}}_{l}^{R}(\omega;x|z) - \bar{\mathcal{G}}_{l}^{R}(\omega;x|z) \partial_{z} \bar{\mathcal{G}}_{l}^{A}(\omega;y|z) \right]_{z=0}^{z=\infty} .$$
(4.13)

The last line can then be computed using the near-origin behaviors (2.10) and near-horizon asymptotics (2.22) of the Jost solution, resulting in

$$\overline{\mathcal{G}_{l}^{C}(\omega;x|y)} = \frac{\omega}{\pi} \frac{\psi_{\omega l}^{\mathrm{n.}}(x)\psi_{\omega l}^{\mathrm{n.}}(y)}{W_{l}^{\mathrm{Jost}}(\omega)W_{l}^{\mathrm{Jost}}(-\omega)}.$$
(4.14)

Notice that since the overall normalizations of  $\bar{\mathcal{G}}_l^R$  and  $\bar{\mathcal{G}}_l^A$  are uniquely fixed by (2.18), that of  $\bar{\mathcal{G}}_l^C$  is uniquely fixed as well.

The spectral trace Regarding (4.12) as the integration kernel associated with an integral operator  $\hat{\mathcal{G}}_{l}^{C}(\omega)$ , it is natural to consider its trace as an integral over the diagonal element

"tr" 
$$\hat{\mathcal{G}}_{l}^{C}(\omega) = \int_{0}^{\infty} dx \, \bar{\mathcal{G}}_{l}^{C}(\omega; x | x) = \frac{\omega}{\pi} \int_{0}^{\infty} dx \frac{\psi_{\omega l}^{\text{n.}}(x)\psi_{\omega l}^{\text{n.}}(x)}{W_{l}^{\text{Jost}}(\omega)W_{l}^{\text{Jost}}(-\omega)},$$
 (4.15)

which however is ill-defined because the integral diverges in the region  $x \to \infty$ . We cut off the integral at x = R (see figure 4.1) and study its large R asymptotics by the following trick. From (2.5), one can deduce

$$(a^{2} - \omega^{2})\psi_{\omega l}^{n}(x)\psi_{a l}^{n}(x) = \psi_{a l}^{n}(x)\partial_{x}^{2}\psi_{\omega l}^{n}(x) - \psi_{\omega l}^{n}(x)\partial_{x}^{2}\psi_{a l}^{n}(x).$$
(4.16)

Taking the derivative of both sides with respect to a and setting  $a = \omega$  implies the formula

$$\int_{0}^{R} dx \, \psi_{\omega l}^{\mathbf{n}.}(x) \psi_{\omega l}^{\mathbf{n}.}(x) = \frac{1}{2\omega} \int_{0}^{R} dx \, \left( \partial_{\omega} \psi_{\omega}^{\mathbf{n}.}(x) \partial_{x}^{2} \psi_{\omega l}^{\mathbf{n}.}(x) - \psi_{\omega l}^{\mathbf{n}.}(x) \partial_{\omega} \partial_{x}^{2} \psi_{\omega}^{\mathbf{n}.}(x) \right) \\
= \frac{1}{2\omega} \left[ \partial_{\omega} \psi_{\omega}^{\mathbf{n}.}(x) \partial_{x} \psi_{\omega l}^{\mathbf{n}.}(x) - \psi_{\omega l}^{\mathbf{n}.}(x) \partial_{\omega} \partial_{x} \psi_{\omega}^{\mathbf{n}.}(x) \right]_{x=0}^{x=R} .$$
(4.17)

Now, using (2.10) and (2.13) results in

$$\int_{0}^{R} dx \, \bar{\mathcal{G}}_{l}^{C}(\omega; x | x) = \frac{R}{2\pi\omega} + \frac{1}{4\pi i\omega} \partial_{\omega} \log \mathcal{S}_{l}(\omega) + O\left(\frac{1}{R}\right) . \tag{4.18}$$

Here  $S_l(\omega)$  is the scattering phase (4.5). Motivated by the reasoning in [18] leading to considering a relative spectral density (4.8), we consider an analogous object for a reference scattering problem. In particular, if we pick the reference problem to be the Rindler scattering problem (C.1) (with  $\beta = 2\pi$ ), we would have

$$\int_{-\infty}^{R} dx \, \bar{\mathcal{G}}_{l}^{\operatorname{Rin},C}(\omega;x|x) = \frac{R}{2\pi\omega} + \frac{1}{4\pi i\omega} \partial_{\omega} \log \mathcal{S}_{l}^{\operatorname{Rin}}(\omega) + O\left(\frac{1}{R}\right) . \tag{4.19}$$

Note that the lower limit for the x-integral is  $-\infty$ . Now, taking the difference between (4.18) and (4.19), we can safely send  $R \to \infty$ . Comparing the result with (4.9) then establishes the relation

$$\Delta \rho_l^{\mathrm{dS}}(\omega) = 2\omega \,\tilde{\mathrm{tr}} \,\hat{\mathcal{G}}_l^C(\omega) \,. \tag{4.20}$$

Here tr denotes the renormalized trace:

$$\tilde{\operatorname{tr}}\,\hat{\mathcal{G}}_{l}^{C}(\omega) \equiv \lim_{R \to \infty} \left( \int_{0}^{R} dx \,\bar{\mathcal{G}}_{l}^{C}(\omega; x | x) - \int_{-\infty}^{R} dx \,\bar{\mathcal{G}}_{l}^{\operatorname{Rin}, C}(\omega; x | x) \right) . \tag{4.21}$$

Taking the inverse Fourier transform of (4.20) and summing over l, we conclude that the Harish-Chandra character is proportional to the time derivative of the traced spectral function:

$$\chi(t) = 2i\frac{d}{dt}\tilde{\operatorname{tr}}\,\hat{G}^C(t) \,. \tag{4.22}$$

This is our central result. We stress that while the Harish-Chandra character on the left hand side of (4.22) is originally defined from the SO(1, d+1) representation theory standpoint, the right hand side is defined directly in terms of a physical observable in the static patch, which remains meaningful for interacting theories.

#### 4.2.1 A single-particle calculation of the character

Naively, one might interpret (4.2) as a trace over the single-particle Hilbert space in the static patch and wish to compute it in the basis  $|\omega l\rangle$  that diagonalizes the static patch Hamiltonian  $\hat{H}$  and SO(d) angular momentum **J**. This leads to a non-sensible result [1]

$$\chi(t) \stackrel{?}{=} \int d\omega \sum_{l} \langle \omega l | e^{-i\hat{H}t} | \omega l \rangle = 2\pi \sum_{l} \delta(0)\delta(t) . \tag{4.23}$$

As explained in [1, 29], one way to recover (4.2) as some single-particle trace is to compute it in a basis on which all SO(1, d+1) generators (in particular the SO(1, 1) generator) act regularly – one labeled by the  $global\ SO(d+1)$  angular momentum, for example. However, the static patch interpretation for these calculations is not completely transparent.

Our discussion leading to (4.22) suggests an elementary single-particle calculation for the character (4.2) from the static patch point of view. The starting point is the matrix element in the position basis:

$$\langle x, \Omega | e^{-i\hat{H}t} | y, \Omega' \rangle = \sum_{l=0}^{\infty} \int_{0}^{\infty} d\omega \, e^{-i\omega t} \, \langle x, \Omega | \omega, l \rangle \, \langle \omega, l | y, \Omega' \rangle . \tag{4.24}$$

Here the wave function

$$\langle x, \Omega | \omega, l \rangle = \sqrt{2\omega} |C_{\omega l}| \psi_{\omega l}^{\mathbf{n}}(x) Y_l(\Omega) , \qquad \omega > 0 , \qquad (4.25)$$

is  $\delta$ -function-normalized with respect to the canonical spatial norm:

$$\langle \omega', l' | \omega, l \rangle = \int_0^\infty dx \tanh^{d-1} x \int_{S^{d-1}} d\Omega \langle \omega', l' | x, \Omega \rangle \langle x, \Omega | \omega, l \rangle = \delta(\omega - \omega') \delta_{ll'}. \tag{4.26}$$

Note that (4.26) is the same as the normalization condition (A.5) for  $\omega, \omega' > 0$ , so the coefficient  $C_{\omega l}$  in (4.25) coincides (up to a constant phase) with the coefficient (2.8) for the normalizable mode.

A trace-like object is given by the integral of the diagonal element:

"tr" 
$$e^{-i\hat{H}t} = \int_0^\infty dx \int_{S^{d-1}} d\Omega \langle x, \Omega | e^{-i\hat{H}t} | x, \Omega \rangle$$
  

$$= \sum_{l=0}^\infty \int_0^\infty d\omega \, e^{-i\omega t} \int_0^\infty dx \, |\langle x, \Omega | \omega, l \rangle|^2 . \tag{4.27}$$

Again, this trace is ill-defined because the x-integral on the right-hand side diverges in the region  $x \to \infty$ . By now, we know very well how to proceed: cut off the integral at x = R, subtract off the analogous integral in a reference Rindler scattering problem, and send  $R \to \infty$  at the end. The calculation proceeds in the same way as the one leading to (4.20). The result is that

$$\Delta \rho_l^{\mathrm{dS}}(\omega) = \lim_{R \to \infty} \left( \int_0^R dx \, |\langle x, \Omega | \omega, l \rangle|^2 - \int_{-\infty}^R dx \, |\langle x, \Omega | \omega, l \rangle|_{\mathrm{Rin}}^2 \right) , \tag{4.28}$$

and therefore summing over l we have

$$\tilde{\operatorname{tr}} e^{-i\hat{H}t} = \sum_{l=0}^{\infty} D_l^d \int_0^{\infty} d\omega \, e^{-i\omega t} \Delta \rho_l^{\mathrm{dS}}(\omega) . \tag{4.29}$$

Here the renormalized trace  $\tilde{\text{tr}}$  is defined analogously as in (4.21). To finally make a connection with the character, we invert (4.4) to find

$$\chi(t) = \sum_{l=0}^{\infty} D_l^d \int_{-\infty}^{\infty} d\omega \, e^{-i\omega t} \Delta \rho_l^{\mathrm{dS}}(\omega) \,. \tag{4.30}$$

Notice that the integral over  $0 < \omega < \infty$  gives (4.29), while the integral over  $-\infty < \omega < 0$  gives the same quantity but with  $t \to -t$ . To summarize, the character is related to the elementary (renormalized) traces through

$$\chi(t) = \tilde{\operatorname{tr}}\left(e^{-i\hat{H}t} + e^{i\hat{H}t}\right). \tag{4.31}$$

## 4.3 Revisiting principal and complementary characters

In this section, we take a closer look at the analytic structure of the relative spectral density (4.4). By applying (2.38) in (4.9), one can separate the contributions from the resonance and anti-resonance poles as

$$\tilde{\rho}^{dS}(\omega) = \tilde{\rho}^{R}(\omega) + \tilde{\rho}^{A}(\omega) , \qquad (4.32)$$

where

$$\tilde{\rho}^{R}(\omega) = \tilde{\rho}^{A}(-\omega) = \sum_{l=0}^{\infty} \Delta \rho_{l}^{R}(\omega) = \frac{1}{2\pi i} \sum_{l=0}^{\infty} \sum_{n=0}^{\infty} D_{l}^{d} \left( \frac{1}{\omega_{nl}^{\Delta} - \omega} + \frac{1}{\omega_{nl}^{\bar{\Delta}} - \omega} \right) . \tag{4.33}$$

Here  $\omega_{nl}^{\Delta}$  and  $\omega_{nl}^{\bar{\Delta}}$  are the resonance frequencies (2.25).

Structure of the character To obtain the character, we perform the inverse Fourier transform

$$\chi(t) = \int_{-\infty}^{\infty} d\omega \, e^{-i\omega t} \tilde{\rho}^{dS}(\omega) . \qquad (4.34)$$

For both principal  $(\Delta = \frac{d}{2} + i\nu, \nu \in \mathbb{R})$  and complementary  $(0 < \Delta < d)$  series, the (anti-)resonance frequencies (2.25) all lie in the lower (upper) half of the complex  $\omega$ -plane.




Figure 4.2: Spectral densities for a principal scalar with  $\Delta = \frac{3}{2} + 0.7i$  in  $dS_4$  in complex  $\omega$  plane.

Therefore, for t > 0 (t < 0), closing the contour in the lower (upper) half plane, only the poles of  $\tilde{\rho}^R(\omega)$  ( $\tilde{\rho}^A(\omega)$ ) contribute, giving

$$\chi(t) = \theta(t) \int_{C_{-}} d\omega \, e^{-i\omega t} \tilde{\rho}^{R}(\omega) + \theta(-t) \int_{C_{+}} d\omega \, e^{-i\omega t} \tilde{\rho}^{A}(\omega)$$
(4.35)

where  $C_{-}$  and  $C_{+}$  are contours enclosing all the poles in the lower-half and upper-half planes respectively. Summing over these poles with the explicit expressions (2.25), we have

$$\chi(t) = \theta(t) \frac{e^{-\Delta t} + e^{-\bar{\Delta}t}}{(1 - e^{-t})^d} + \theta(-t) \frac{e^{\Delta t} + e^{\bar{\Delta}t}}{(1 - e^t)^d},$$
(4.36)

reproducing (4.2).

## 5 Comments on the exceptional scalars

In this section, we study free scalars with masses

$$m_k^2 = -k(k+d)$$
,  $k = 0, 1, 2, \cdots$ , (5.1)

corresponding to scaling dimensions of non-positive integers

$$\Delta = -k \ . \tag{5.2}$$

While k = 0 corresponds to a massless scalar, the scalars for any  $k \ge 1$  are all tachyonic. Classical theories of these scalars exhibit shift symmetries [41]. From the SO(1, d+1) representation theory point of view, (5.2) correspond to the so-called exceptional series I UIRs of [29], even though their realizations in terms of a local quantum field theory remain to be understood. Despite their tachyonic nature, they could be of physical interest because they generically appear in the multiparticle Hilbert spaces of massive fields [27, 28]. In 2D (d = 1), these UIRs coincide with the discrete series, whose Hilbert space realizations are explored in [42].

#### 5.1 Massless scalar

We start with the case of  $\Delta = 0$  [43, 44], for which the action is invariant under a constant shift

$$\phi \to \phi + c \ . \tag{5.3}$$

In sharp contrast with principal or complementary series, the resonance poles (supposedly) at

$$i\omega_{nl}^{\Delta=0} = l + 2n$$
 or  $i\omega_{nl}^{\bar{\Delta}=d} = d + l + 2n$ , (5.4)

largely coincide with the Matsubara frequencies (2.27). Also, there is a mode with l = n = 0 lying exactly at the origin. This drastically changes the analytic structure of the Jost function,

$$W_l^{\text{Jost}}(\omega) = \frac{2\Gamma(1 - i\omega)}{\Gamma(\frac{l - i\omega}{2})\Gamma(\frac{d + l - i\omega}{2})},$$
(5.5)

and therefore of the correlators as well. On one hand, except at l = n = 0, the first tower (5.4) are not zeros of (5.5), meaning they are not true resonance poles. On the other, the second tower are zeros of (5.5) in even d but not odd d. The occurrence of the would-be resonance poles (and the analogous phenomena for general  $k \ge 0$ ) is reminiscent of the pole-skipping phenomena [45–47].

One can also work out the worldline retarded function by an inverse transform,

$$G^{R}(t) = \frac{1}{\operatorname{Vol}(S^{d-1})} \int_{C^{+}} d\omega \, e^{-i\omega t} \mathcal{G}_{l=0}^{R}(\omega) , \qquad (5.6)$$

where the contour  $C^+$  lies above all the poles, including the one at  $\omega = 0$ , as illustrated in Figure 5.1. Performing the integral then leads to

$$G^{R}(t) = \frac{\theta(t)}{\text{Vol}(S^{d-1})} \left[ 1 - \cos\left(\frac{\pi d}{2}\right) \frac{4}{B\left(\frac{d}{2}, \frac{d}{2}\right)} \frac{e^{-dt}}{d} {}_{2}F_{1}\left(d, \frac{d}{2}, 1 + \frac{d}{2}, e^{-2t}\right) \right] . \tag{5.7}$$

One can check that this is equivalent to taking the limit  $\Delta \to 0$  in (3.2). Similarly, one can work out the advanced function, where the integration contour  $C^-$  must lie below the real line to avoid the pole at  $\omega = 0$ . With these, we can write down the spectral function:

$$G^{C}(t) = -i \left( G^{R}(t) - G^{A}(t) \right)$$

$$= \frac{-i}{\text{Vol}(S^{d-1})} \left[ 1 - \cos \left( \frac{\pi d}{2} \right) \frac{4}{B \left( \frac{d}{2}, \frac{d}{2} \right)} \frac{e^{-d|t|}}{d} {}_{2}F_{1} \left( d, \frac{d}{2}, 1 + \frac{d}{2}, e^{-2|t|} \right) \right] . \tag{5.8}$$

Notice the drastic difference between even and odd d: for the latter case (which includes the realistic d = 3 case), the second decaying term completely vanishes, leaving only the constant term.

Before we move on, we note that since  $\mathcal{G}^A(\omega; x|y) = \mathcal{G}^R(-\omega; x|y)$ , the (full) spectral function can be written as

$$G^{C}(t;x|y) = -i\left(G^{R}(t;x|y) - G^{A}(t;x|y)\right) = \frac{1}{\operatorname{Vol}(S^{d-1})} \int_{C^{+}} d\omega \left(e^{-i\omega t} - e^{i\omega t}\right) \mathcal{G}^{R}(\omega;x|y) . \quad (5.9)$$

Now, we deform the contour  $C^+ \to \mathbb{R}_\epsilon + C_\epsilon$  as follows:




Figure 5.1: The deformed contour consists of a semi-circle  $C_{\epsilon}$  of radius  $\epsilon > 0$  and  $\mathbb{R}_{\epsilon} = \mathbb{R} \setminus (-\epsilon, \epsilon)$ .

In the  $\epsilon \to 0$  limit, the contribution from the small semi-circle  $C_{\epsilon}$  is given by  $-\pi i$  times the residue at  $\omega = 0$ , which is zero. To conclude, we have

$$G^{C}(t;x|y) = \frac{1}{\operatorname{Vol}(S^{d-1})} \int_{-\infty}^{\infty} d\omega \, e^{-i\omega t} \mathcal{G}^{C}(\omega;x|y) , \qquad (5.10)$$

where the integral is understood in the sense of the Cauchy principal value

$$\int_{-\infty}^{\infty} d\omega \equiv \lim_{\epsilon \to 0^{+}} \int_{\mathbb{R} \setminus (-\epsilon, \epsilon)} d\omega . \tag{5.11}$$

#### 5.1.1 Canonical quantization

It is long known that the problem of quantizing a free massless scalar in de Sitter space is subtle. For example, the massive dS-invariant two-point function diverges in the massless limit, which is intimately tied with the fact that there is no dS-invariant Fock vacuum for a massless scalar [43]. However, there could exist vacua that preserve fewer symmetries, realized by quantizing the massless scalar in different patches [43, 44].

When we consider this problem in a static patch, we find that as long as we replace in (A.1)

$$\int_{-\infty}^{\infty} d\omega \to \int_{-\infty}^{\infty} d\omega \tag{5.12}$$

as defined in (5.11), the canonical quantization procedure in appendix A applies. In particular, we have a consistent *quantum* theory that reproduces correlators such as (5.7) and (5.8). Subsequently, we can also define the temperature-sensitive functions. As a demonstration, we compute the symmetric Wightman function at the dS temperature  $\beta = \beta_{dS} = 2\pi$ :

$$G_{\beta=2\pi}^{\text{Sym}}(t) = \frac{1}{\text{Vol}(S^{d-1})} \int_{-\infty}^{\infty} d\omega \, e^{-i\omega t} \mathcal{G}_{l=0}^{\text{Sym}}(2\pi, \omega) , \qquad (5.13)$$

where

$$\mathcal{G}_{l=0}^{\text{Sym}}(2\pi,\omega) = \frac{1}{8\pi^2} \frac{1}{\Gamma\left(\frac{d}{2}\right)^2} \cosh \pi\omega \left| \Gamma\left(\frac{-i\omega}{2}\right) \Gamma\left(\frac{d-i\omega}{2}\right) \right|^2 . \tag{5.14}$$

We will evaluate (5.13) explicitly and show that it can be obtained as an analytic continuation of an appropriate  $S^{d+1}$  correlator.

Clearly (5.14) has poles at the resonances (5.4) with l = 0,

$$i\omega_n^{\Delta=0} = 2n , \qquad i\omega_n^{\bar{\Delta}=d} = d + 2n , \qquad (5.15)$$

and the anti-resonances given by  $-i\omega_n^{\Delta=0}$  and  $-i\omega_n^{\bar{\Delta}=d}$ . In any dimension, the pole at  $\omega=\omega_{n=0}^{\Delta=0}=0$  is a double pole, where the integrand (5.13) has residue

$$-\pi i \operatorname{Res}_{\omega=0} e^{-i\omega t} \mathcal{G}_{l=0}^{\operatorname{Sym}}(2\pi, \omega) = -\frac{t}{2\pi} . \tag{5.16}$$

To proceed, we separate the cases of odd and even d.

When d is odd Except for the double pole at  $\omega = 0$ , both towers (5.15) are simple poles for the integrand (5.13), at which the residues are

$$-2\pi i \operatorname{Res}_{\omega = -i2n} e^{-i\omega t} \mathcal{G}_{l=0}^{\operatorname{Sym}}(2\pi, \omega) = \frac{(-1)^n}{2\pi\Gamma\left(\frac{d}{2}\right)^2} \Gamma\left(\frac{d}{2} - n\right) \Gamma\left(\frac{d}{2} + n\right) \frac{e^{-2nt}}{n} , \qquad n \ge 1 , \qquad (5.17)$$

and

$$-2\pi i \operatorname{Res}_{\omega = -i(d+2n)} e^{-i\omega t} \mathcal{G}_{l=0}^{\text{Sym}}(2\pi, \omega) = \frac{(-1)^{\frac{d-1}{2}}}{\Gamma(\frac{d}{2})^2} \frac{\Gamma(d+n)}{d+2n} \frac{e^{-(d+2n)t}}{n!} , \qquad n \ge 0 ,$$
 (5.18)

respectively. Therefore, for t > 0 (t < 0), closing the contour in the lower (upper) half plane, we sum over the residues and obtain

$$G_{\beta=2\pi}^{\text{Sym}}(t) = \frac{1}{\text{Vol}(S^{d-1})} \left[ -\frac{|t|}{2\pi} - \frac{d}{2\pi(d-2)} e^{-2|t|} \,_{3}F_{2}\left(1, 1, \frac{d}{2} + 1; 2, 2 - \frac{d}{2}; e^{-2|t|}\right) + \frac{(-)^{\frac{d-1}{2}}}{B\left(\frac{d}{2}, \frac{d}{2}\right)} \frac{e^{-d|t|}}{d} \,_{2}F_{1}\left(d, \frac{d}{2}, 1 + \frac{d}{2}, e^{-2|t|}\right) \right].$$

$$(5.19)$$

We note that the generalized hypergeometric function in the first line can be written as [48]

$$e^{-2|t|} {}_{3}F_{2}\left(1, 1, \frac{d}{2} + 1; 2, 2 - \frac{d}{2}; e^{-2|t|}\right) = -\frac{d-2}{d} \left[\frac{\partial}{\partial \Delta} {}_{2}F_{1}\left(\Delta, \frac{d}{2}; 1 - \frac{d}{2}; e^{-2t}\right)\Big|_{\Delta \to 0}\right]. \tag{5.20}$$

Therefore,

$$G_{\beta=2\pi}^{\text{Sym}}(t) = \frac{1}{\text{Vol}(S^{d-1})} \left[ -\frac{|t|}{2\pi} + \frac{1}{2\pi} \partial_{\Delta} {}_{2}F_{1} \left( \Delta, \frac{d}{2}; 1 - \frac{d}{2}; e^{-2|t|} \right) \Big|_{\Delta=0} + \frac{(-)^{\frac{d-1}{2}}}{B\left(\frac{d}{2}, \frac{d}{2}\right)} \frac{e^{-d|t|}}{d} {}_{2}F_{1} \left( d, \frac{d}{2}, 1 + \frac{d}{2}, e^{-2|t|} \right) \right].$$
 (5.21)

When d is even The integrand (5.13) has simple poles at the first tower of (5.15) for  $1 \le n \le \frac{d}{2} - 1$ , at which it has residues

$$-2\pi i \operatorname{Res}_{\omega=-i2n} e^{-i\omega t} \mathcal{G}_{l=0}^{\operatorname{Sym}}(2\pi,\omega) = \frac{(-1)^n}{2\pi\Gamma\left(\frac{d}{2}\right)^2} \Gamma\left(\frac{d}{2}-n\right) \Gamma\left(\frac{d}{2}+n\right) \frac{e^{-2nt}}{n} , \quad 1 \le n \le \frac{d}{2}-1 . \quad (5.22)$$

For all  $n \geq \frac{d}{2}$ , the integrand (5.13) has double poles instead, at which it has residues

$$-2\pi i \operatorname{Res}_{\omega=-i2n} e^{-i\omega t} \mathcal{G}_{l=0}^{\operatorname{Sym}}(2\pi,\omega)$$

$$= \frac{(-)^{d/2}}{2\pi\Gamma\left(\frac{d}{2}\right)^2} \frac{\Gamma\left(\frac{d}{2}+n\right)}{n} \left(\frac{1}{n}+2t+\psi\left(n-\frac{d}{2}+1\right)-\psi\left(n+\frac{d}{2}\right)\right) \frac{e^{-2nt}}{\Gamma\left(n-\frac{d}{2}+1\right)}, \quad n \ge \frac{d}{2}.$$
(5.23)

In this case, the second tower (5.15) completely overlaps with the  $n \ge \frac{d}{2}$  part of the first tower and hence does not lead to separate contributions.

Therefore, for t > 0 (t < 0), closing the contour in the lower (upper) half plane, we have

$$G_{\beta=2\pi}^{\text{Sym}}(t) = \frac{-1}{\text{Vol}(S^{d-1})} \left[ \frac{|t|}{2\pi} + 2\pi i \left( \sum_{n=0}^{\frac{d}{2}-1} + \sum_{n=\frac{d}{2}}^{\infty} \right) \underset{\omega=-i2n}{\text{Res}} e^{-i\omega|t|} \mathcal{G}_{l=0}^{\text{Sym}}(2\pi,\omega) \right] . \tag{5.24}$$

Shifting the second sum by  $n \to n + \frac{d}{2}$ , we have

$$G_{\beta=2\pi}^{\mathrm{Sym}}(t)$$

$$= \frac{1}{\text{Vol}(S^{d-1})} \left[ -\frac{|t|}{2\pi} + \frac{1}{B\left(\frac{d}{2}, \frac{d}{2}\right)} \sum_{n=0}^{\frac{d}{2}-1} \frac{(-1)^n}{2\pi\Gamma(d)} \Gamma\left(\frac{d}{2} - n\right) \Gamma\left(\frac{d}{2} + n\right) \frac{e^{-2n|t|}}{n} + \frac{(-)^{d/2}}{B\left(\frac{d}{2}, \frac{d}{2}\right)} \frac{e^{-d|t|}}{\pi} \sum_{n=0}^{\infty} \frac{1}{\Gamma(d)} \frac{\Gamma(d+n)}{d+2n} \left(\frac{2}{d+2n} + 2|t| + \psi(n+1) - \psi(n+d)\right) \frac{e^{-2n|t|}}{n!} \right]. \quad (5.25)$$

In the second equality, we have shifted the sum by  $n \to n + \frac{d}{2}$ 

#### 5.1.2 Euclidean correlator

A natural question is whether the real-time thermal correlators computed above correspond to some analytic continuation from a  $S^{d+1}$  correlator. This is not as trivial as one might have thought, because in contrast to the massive case, a massless scalar does not have a standard Green function satisfying (3.6); the Laplacian has a normalizable zero mode and is thus non-invertible. It is also reflected in the fact that the massive expression (3.5) diverges as  $\Delta \to 0$ . An object we can consider is a modified Green function that satisfies

$$-\nabla_{\mathbf{x}}^{2}\hat{G}_{E}(P(\mathbf{x},\mathbf{y})) = \frac{1}{\sqrt{g}}\delta^{(d+2)}(\mathbf{x},\mathbf{y}) - \frac{1}{\operatorname{Vol}(S^{d+1})}.$$
 (5.26)

Because of the constant term on the right hand side, it is an *inhomogeneous* equation even at non-coincident points. This constant term represents a non-local constraint of the problem, analogous to the Gauss law constraint that forbids putting a single charge on a sphere. A particular solution to (5.26) is given by [49]

$$\hat{\mathbf{G}}(\mathbf{x}, \mathbf{y}) \equiv \frac{d}{d\Delta} \left[ \frac{G_E^{\Delta}(P(\mathbf{x}, \mathbf{y}))}{\Gamma(\Delta)} \right] \Big|_{\Delta=0} , \qquad (5.27)$$

with  $G_E^{\Delta}(P(\mathbf{x}, \mathbf{y}))$  being the massive Green function (3.5). One can see that (5.27) is a solution to (5.26) by dividing (3.6) by  $\Gamma(\Delta)$  and evaluating its  $\Delta$ -derivative at  $\Delta = 0$ . Notice that (5.27) is not a unique solution to (5.26) since we can always add a constant C:

$$\hat{G}_E(P(\mathbf{x}, \mathbf{y})) = \hat{\mathbf{G}}(P(\mathbf{x}, \mathbf{y})) + C.$$
(5.28)

In the following we will demonstrate that upon choosing an appropriate constant C, the unique  $\hat{G}_E(P(\mathbf{x}, \mathbf{y}))$  would analytically continue to the real-time correlators as in the massive case.

The modified function (5.27) is convenient to work with since we can simply take limits in the previous massive results. We can apply the prescription (5.27) directly to the analytically continued expression (3.11), whose second term is finite for odd d and diverges for even d as  $\Delta \to 0$ .

When d is odd Applying (5.27) to (3.11) yields

$$\hat{\mathbf{G}}(P \mp i\epsilon) = \frac{1}{2\text{Vol}(S^{d+1})\Gamma\left(\frac{d}{2} + 1\right)} \partial_{\Delta} \left[ e^{\mp i\pi\Delta} \Gamma\left(\frac{d}{2} - \Delta\right) e^{-\Delta|t|} {}_{2}F_{1}\left(\Delta, \frac{d}{2}, 1 - \frac{d}{2} + \Delta, e^{-2|t|}\right) \right]_{\Delta = 0} - \frac{\Gamma(d)\Gamma(-\frac{d}{2})}{2\text{Vol}(S^{d+1})\Gamma\left(\frac{d}{2} + 1\right)} e^{-d|t|} {}_{2}F_{1}\left(d, \frac{d}{2}, 1 + \frac{d}{2}, e^{-2|t|}\right) .$$
(5.29)

Using the series representation for the hypergeometric function

$$_{2}F_{1}(a,b,c,z) = \sum_{n=0}^{\infty} \frac{(a)_{n}(b)_{n}}{(c)_{n}} \frac{z^{n}}{n!},$$
 (5.30)

the derivative on the first line can be computed to be

$$\partial_{\Delta} \left[ e^{\mp i\pi\Delta} e^{-\Delta|t|} \Gamma\left(\frac{d}{2} - \Delta\right) {}_{2}F_{1}\left(\Delta, \frac{d}{2}, 1 - \frac{d}{2} + \Delta, e^{-2|t|}\right) \right]_{\Delta=0}$$

$$= \Gamma\left(\frac{d}{2}\right) \left[ \mp i\pi - |t| + \partial_{\Delta} {}_{2}F_{1}\left(\Delta, \frac{d}{2}; 1 - \frac{d}{2}; e^{-2|t|}\right) \Big|_{\Delta=0} - \psi\left(\frac{d}{2}\right) \right]. \tag{5.31}$$

Here  $\psi(z) = \partial_z \log \Gamma(z)$  is the digamma function. Choosing

$$C = \frac{1}{2\pi \text{Vol}(S^{d-1})} \psi\left(\frac{d}{2}\right) \tag{5.32}$$

in (5.28), i.e.

$$\hat{G}_E(P(\mathbf{x}, \mathbf{y})) = \hat{\mathbf{G}}(P(\mathbf{x}, \mathbf{y})) + \frac{1}{2\pi \text{Vol}(S^{d-1})} \psi\left(\frac{d}{2}\right) \quad \text{(odd } d),$$
 (5.33)

we have the *unique* sphere correlator that analytically continues to the Lorentzian signature as in the massive case (3.12), i.e.

$$G^{C}(t) = \hat{G}_{E}(P - i\epsilon) - \hat{G}_{E}(P + i\epsilon) , \qquad G_{\beta=2\pi}^{Sym}(t) = \frac{\hat{G}_{E}(P - i\epsilon) + \hat{G}_{E}(P + i\epsilon)}{2} , \qquad (5.34)$$

where  $G^{C}(t)$  and  $G_{\beta=2\pi}^{\text{Sym}}(t)$  are defined in (5.8) and (5.19) respectively.

When d is even Applying (5.27) to (3.11) yields

$$\mathbf{G}(P \mp i\epsilon) = \frac{1}{2\text{Vol}(S^{d+1})\Gamma\left(\frac{d}{2} + 1\right)} \left(\partial_{\Delta} \left[e^{\mp i\pi\Delta}\Gamma\left(\frac{d}{2} - \Delta\right)e^{-\Delta|t|} {}_{2}F_{1}\left(\Delta, \frac{d}{2}, 1 - \frac{d}{2} + \Delta, e^{-2|t|}\right)\right]_{\Delta=0} + \partial_{\Delta} \left[e^{\mp i\pi\bar{\Delta}}\frac{\Gamma(\bar{\Delta})\Gamma\left(\frac{d}{2} - \bar{\Delta}\right)}{\Gamma(\Delta)}e^{-\bar{\Delta}|t|} {}_{2}F_{1}\left(\bar{\Delta}, \frac{d}{2}, 1 - \frac{d}{2} + \bar{\Delta}, e^{-2|t|}\right)\right]_{\Delta=0}\right).$$
(5.35)

The computation in this case is more subtle. For example, when using (5.30) to compute the first line, we note that in the  $\Delta \to 0$  limit, the term n=0 survives, and those with  $1 \le n \le \frac{d}{2} - 1$  go to zero, while all terms with  $n \ge \frac{d}{2}$  acquire a non-trivial limit:

$$\lim_{\Delta \to 0} {}_{2}F_{1}\left(\Delta, \frac{d}{2}, 1 - \frac{d}{2} + \Delta, e^{-2|t|}\right) = 1 + \frac{2(-)^{\frac{d}{2}+1}}{B\left(\frac{d}{2}, \frac{d}{2}\right)} e^{-d|t|} \sum_{n=0}^{\infty} \frac{1}{\Gamma(d)} \frac{\Gamma(d+n)}{d+2n} \frac{e^{-2n|t|}}{n!} . \tag{5.36}$$

After some computation with all limits carefully taken care of, we eventually obtain

$$\hat{\mathbf{G}}(P \mp i\epsilon)$$

$$= \frac{1}{\text{Vol}(S^{d-1})} \left[ \mp \frac{i}{2} - \frac{|t|}{2\pi} + \frac{1}{B\left(\frac{d}{2}, \frac{d}{2}\right)} \sum_{n=0}^{\frac{d}{2}-1} \frac{(-)^n}{2\pi\Gamma(d)} \Gamma\left(\frac{d}{2} + n\right) \Gamma\left(\frac{d}{2} - n\right) \frac{e^{-2n|t|}}{n} + \frac{(-)^{\frac{d}{2}}}{B\left(\frac{d}{2}, \frac{d}{2}\right)} \frac{e^{-d|t|}}{\pi} \sum_{n=0}^{\infty} \frac{1}{\Gamma(d)} \frac{\Gamma(d+n)}{d+2n} \left( \mp i2\pi + \frac{2}{d+2n} + 2|t| + \psi(n+1) - \psi(n+d) \right) \frac{e^{-2n|t|}}{n!} \right].$$
(5.37)

Choosing C=0 in (5.28), i.e.

$$\hat{G}_E(P(\mathbf{x}, \mathbf{y})) = \hat{\mathbf{G}}(P(\mathbf{x}, \mathbf{y})) \quad \text{(even } d),$$
 (5.38)

we have the *unique* sphere correlator that analytically continues to the Lorentzian signature as in the massive case (3.12), i.e.

$$G^{C}(t) = \hat{G}_{E}(P - i\epsilon) - \hat{G}_{E}(P + i\epsilon) , \qquad G^{\text{Sym}}_{\beta = 2\pi}(t) = \frac{\hat{G}_{E}(P - i\epsilon) + \hat{G}_{E}(P + i\epsilon)}{2} , \qquad (5.39)$$

where  $G^C(t)$  and  $G^{\mathrm{Sym}}_{\beta=2\pi}(t)$  are defined in (5.8) and (5.25) respectively.

To summarize, we have demonstrated that the prescription (5.12) leads to a consistent quantization for a massless scalar in the static patch, and performed a non-trivial check that the Lorentzian correlators can all be analytically continued from an appropriate  $S^{d+1}$  correlator. However, this is unlikely the full story. Specifically, the prescription (5.12) explicitly excludes the constant/soft mode  $\omega = 0$ , which could encode interesting physics.<sup>6</sup> For example, the constant mode seems closely related to the fact that extra data is required to fully determine the massless scalar 1-loop  $S^{d+1}$  partition functions [2], such as the target space geometry if the parent interacting theory is a non-linear sigma model.

### 5.2 General shift-symmetric scalars

For any  $\Delta = -k, k \ge 1$ , the scalars are tachyonic. See for example [49] and [52] for the study of such tachyonic scalars. Their free actions are invariant under the shift symmetries

$$\phi \to \phi + cf_k \,, \tag{5.40}$$

where  $f_k$  is a homogeneous polynomial in ambient space coordinates of degree k [41]. The k = 1 and k = 2 cases are the dS analogs for the Galileon and special Galileon theories in flat space [53], respectively. The former case can also be obtained by truncating linearized gravity to its conformal sector, and thus has been studied as a toy model for the latter [54].

Causal correlators Similar to the massless case, the resonance poles (supposedly) at

$$i\omega_{nl}^{\Delta=-k} = -k + l + 2n$$
 or  $i\omega_{nl}^{\bar{\Delta}=d+k} = d + k + l + 2n$  (5.41)

largely coincide with the Matsubara frequencies (2.27). In addition to the modes with  $\omega = 0$  (when l + 2n = k) lying exactly at the origin, there are a finite number of poles lying *above* the real line, namely those in the first tower with

$$l + 2n < k. (5.42)$$

The presence of these modes indicates a kind of superradiant instability, typical in tachyonic theories. Similar to the massless case, except when  $l + 2n \le k$ , the first tower (5.4) are not zeros of the Jost function,

$$W_l^{\text{Jost}}(\omega) = \frac{2\Gamma(1 - i\omega)}{\Gamma\left(\frac{-k + l - i\omega}{2}\right)\Gamma\left(\frac{d + k + l - i\omega}{2}\right)},$$
(5.43)

 $<sup>^6</sup>$ In the context of Minkowski space, the constant/soft modes play a crucial role in geometric soft theorems [50, 51].

while the second tower is zeros of (5.43) in even d but not odd d.

In computing the worldline retarded function,

$$G^{R}(t) = \frac{1}{\operatorname{Vol}(S^{d-1})} \int_{C} d\omega \, e^{-i\omega t} \mathcal{G}_{l=0}^{R}(\omega) , \qquad (5.44)$$

we must choose the contour lie above the highest pole  $\omega = ik$ . The result is equivalent to taking the limit  $\Delta \to -k$  in (3.2):

$$G^{R}(t) = \frac{\theta(t)}{\text{Vol}(S^{d-1})} \left[ \frac{1}{B(\frac{d}{2}, k)} \frac{e^{kt}}{k} {}_{2}F_{1}\left(-k, \frac{d}{2}, 1 - \frac{d}{2} - k, e^{-2t}\right) - \cos\left(\frac{\pi d}{2}\right) \frac{4}{B(\frac{d}{2}, \frac{d}{2} + k)} \frac{e^{-(d+k)t}}{d + 2k} {}_{2}F_{1}\left(d + k, \frac{d}{2}, 1 + \frac{d}{2} + k, e^{-2t}\right) \right].$$
 (5.45)

The advanced function can be similarly obtained. Taking the difference, we have the spectral function:

$$G^{C}(t) = \frac{-i}{\operatorname{Vol}(S^{d-1})} \left[ \frac{1}{B\left(\frac{d}{2}, k\right)} \frac{e^{k|t|}}{k} {}_{2}F_{1}\left(-k, \frac{d}{2}, 1 - \frac{d}{2} - k, e^{-2|t|}\right) - \cos\left(\frac{\pi d}{2}\right) \frac{4}{B\left(\frac{d}{2}, \frac{d}{2} + k\right)} \frac{e^{-(d+k)|t|}}{d + 2k} {}_{2}F_{1}\left(d + k, \frac{d}{2}, 1 + \frac{d}{2} + k, e^{-2|t|}\right) \right].$$
 (5.46)

Comments on quantization Notice that (5.45) or (5.46) implies that the full correlators grow indefinitely for two points at large timelike separations. It is not possible to reproduce these by a quantum free real scalar theory with usual properties, namely that we have a self-adjoint field operator  $\hat{\phi}(t)$  (suppressing the spatial dependence) that evolves in time unitarily as  $\hat{\phi}(t) = e^{i\hat{H}t}\hat{\phi}(0)e^{-i\hat{H}t}$ , generated by a self-adjoint Hamiltonian  $\hat{H}$  with respect to which we have a normalizable vacuum state  $|0\rangle$ . Otherwise, consider the smeared field operator

$$\hat{\phi}_f(T) = \int dt dx d\Omega \, \hat{\phi}(t+T, x, \Omega) f(t, x, \Omega)$$
(5.47)

for some smooth function  $f(t, x, \Omega)$  on the static patch, e.g. a Gaussian centered around t = 0 and x = 0. Then, the Wightman function for this smeared operator at general T > 0 must satisfy<sup>7</sup>

$$\langle 0|\phi_f(T)\phi_f(0)|0\rangle \leq \sqrt{\langle 0|\phi_f(t)\phi_f(t)|0\rangle}\sqrt{\langle 0|\phi_f(0)\phi_f(0)|0\rangle} = \langle 0|\phi_f(0)\phi_f(0)|0\rangle , \qquad (5.48)$$

i.e. it is bounded from above by its value at T=0. Since all other two-point functions are linear combinations of the Wightman function, their time evolutions are bounded as well. This clearly contradicts the exponentially growing behaviors of (5.45) or (5.46), since the smeared functions behave like the un-smeared ones for T much greater than the width of the function f.

This is consistent with the existing discussions in the literature: to have a self-consistent quantization, the shift symmetry (5.40) needs to be gauged in some way – coupling these scalars with gauge fields [42], for example, so that the modes responsible for the growing behaviors are puregauge.

<sup>&</sup>lt;sup>7</sup>The smearing (5.47) is to guarantee that  $\phi_f(0)|0\rangle$  has a finite norm.

## 5.3 Harish-Chandra characters for the exceptional scalars

In this section, we analyze the structures of the Harish-Chandra characters for the exceptional scalars along the lines of section 4.3. To that end, we stress that the analysis for the Klein-Gordon equation for the exceptional scalars are no different from the generic massive case, which implies that the discussion in section 4.1 remains valid. In particular, the S-matrices for the exceptional case are given by taking  $\Delta \to -k$  in (4.1):

$$S_{l}(\omega) = S_{l}^{\mathrm{dS}}(\omega) S^{\mathrm{Rin}}(2\pi, \omega) ,$$

$$S_{l}^{\mathrm{dS}}(\omega) \equiv \frac{\Gamma\left(\frac{-k+l-i\omega}{2}\right) \Gamma\left(\frac{d+k+l-i\omega}{2}\right)}{\Gamma\left(\frac{-k+l+i\omega}{2}\right) \Gamma\left(\frac{d+k+l+i\omega}{2}\right)} , \qquad S^{\mathrm{Rin}}(2\pi, \omega) = \frac{\Gamma(i\omega)}{\Gamma(-i\omega)} .$$
(5.49)

As discussed in section 5.1 and 5.2, a subset of (5.4) or (5.41) are not true resonance poles for the Klein-Gordon equation, since they are not zeros of the Jost function. However, they are poles of  $\mathcal{S}_l^{\mathrm{dS}}(\omega)$  (and therefore of  $\tilde{\rho}^R(\omega)$ ). Because of this, we will abuse the language in this section and call them the resonance poles. Similarly, we will call the zeros of  $\mathcal{S}_l^{\mathrm{dS}}(\omega)$  (poles of  $\tilde{\rho}^A(\omega)$ ) the anti-resonance poles.

Massless scalar As we saw in section 5.1, in the massless case the (anti-)resonance poles do not all lie in the lower (upper) half plane. In particular, there is a pole for  $\tilde{\rho}^R(\omega)$  and  $\tilde{\rho}^A(\omega)$  with exactly zero frequency  $\omega = 0$  (n = l = 0 in (5.4)). See figure 5.2. In the total relative spectral density (4.32), the contributions from these two poles cancel each other.

Therefore, closing the contour in the lower (upper) half plane for t > 0 (t < 0), the inverse transform (4.34) receives contributions from all (anti-)resonance frequencies except  $\omega = 0$ , yielding

$$\chi(t) = \frac{1 + e^{-dt}}{|1 - e^{-t}|^d} - 1.$$
 (5.50)

The subtraction by 1 accounts for the fact that the (anti-)resonance pole  $\omega = 0$  does not contribute.




Figure 5.2: Spectral densities for a massless scalar in  $dS_4$  in complex  $\omega$  plane.

General exceptional scalars In addition to a (anti-)resonance pole at  $\omega = 0$ , in the shift-symmetric case there are a finite number of resonance poles lying on the *upper*-half plane, namely

those with (5.42). By time reversal symmetry, we have the corresponding anti-resonance poles lying in the lower half plane. See figure 5.3.




Figure 5.3: Spectral densities for a exceptional scalar with  $\Delta = -3$  in  $dS_4$  in complex  $\omega$  plane.

While the pole at  $\omega = 0$  always gets canceled in the total relative spectral density (4.32), the poles (5.42) do not, and will contribute to the character upon the inverse Fourier transform (4.34). For t < 0, we close the contour in the lower half plane. This contour picks up all the poles of  $\Delta \rho^R(\omega)$  except  $\omega = 0$  or those satisfying (5.42); in addition, it picks up the poles of  $\Delta \rho^A(\omega)$  satisfying (5.42). Explicitly, we have

$$\int_{C_{-}} d\omega \, e^{-i\omega t} \tilde{\rho}^{R}(\omega) = \frac{e^{kt} + e^{-(d+k)t}}{(1 - e^{-t})^{d}} - 1 - \sum_{2n+l < k} D_{l}^{d} e^{(k-2n-l)t} ,$$

$$\int_{C_{-}} d\omega \, e^{-i\omega t} \tilde{\rho}^{A}(\omega) = \sum_{2n+l < k} D_{l}^{d} e^{-(k-2n-l)t} .$$
(5.51)

The t < 0 case works similarly. Summing these contributions results in

$$\chi(t) = \frac{e^{kt} + e^{-(d+k)t}}{|1 - e^{-t}|^d} - 1 - \sum_{2n+l < k} D_l^d \left( e^{(k-2n-l)|t|} - e^{-(k-2n-l)|t|} \right) . \tag{5.52}$$

This gives a nice physical picture for the "flipping" procedure in calculating the characters for the exceptional scalars [29]: there are a finite number of (anti-)resonance poles that lie in the upper (lower) half plane, and the subtraction by 1 accounts for the fact that the  $\omega = 0$  pole is never a pole of the total relative spectral density (4.32).

To end this section, we note that the Harish-Chandra characters associated with (partially) massless fields [55–63], which include Maxwell field or graviton that are of great physical interests, exhibit a similar flipping structure [1, 29]. It would be interesting to check if the physical mechanism we just revealed for such a structure in the exceptional scalar case applies to the partially massless fields as well.

### 6 Outlook

We conclude by outlining several avenues for future exploration.

**Spinning fields** Our discussion has primarily revolved around free scalars. A natural extension involves exploring fields with spins, whose  $S^{d+1}$  path integrals exhibit a distinct bulk-edge split [1] (extended to arbitrary static black holes in [19]). Indeed, elucidating the Lorentzian interpretation of this split originally motivated our work. The analysis is considerably more intricate and we will present the details in [64].

More general backgrounds The reader might have noticed the robustness of the analysis in section 4. In fact, it is easy to generalize it to free scalar theories on any (static) black holes, even in cases where the spectral functions have more complicated analytic structures such as branch cuts [30]. This is not too surprising, considering the parallel analyses for the partition functions that have been worked out in [18, 19]. This will be elaborated in [65].

Interacting theories and beyond Another natural next step is to generalize our considerations to interacting theories. In contrast to the free case, the spectral function in the interacting theory is no longer independent of the temperature, but it still plays a prominent role because all correlators can be written in terms of it by an integral representation [66, 67]. With the thermal correlators studied in this work, one can start performing perturbative calculations of various thermodynamic quantities. At the non-perturbative level, motivated by our key relation (4.22), one would naturally consider a character-like object

$$\chi(t) \equiv 2i \frac{d}{dt} \tilde{\text{tr}} \, \hat{G}^C(t) , \qquad (6.1)$$

where

$$G^{C}(t, x, \Omega | 0, y, \Omega') \equiv \left\langle \left[ \hat{O}(t, x, \Omega), \hat{O}(0, y, \Omega') \right] \right\rangle_{\beta = 2\pi}$$
(6.2)

is the spectral function for some scalar operator  $\hat{O}$  in the *interacting* theory at temperature  $\beta = 2\pi$ . Through a Källén–Lehmann-type representation [68, 69] for (6.2),<sup>8</sup> we can then write an analogous representation for (6.1)

$$\chi(t) = \int d\Delta \rho(\Delta) \chi_{\Delta}(t) . \tag{6.3}$$

It would be interesting to explore the usefulness of organizing the physical information in this way.

Our novel perspectives on the correlators might also be inspiring for the bottom-up approach to a microscopic model associated with a static patch. Conceptually, consider the role of an *integrated* correlator as a notable group-theoretic object. This could offer a "holistic" view that links together various proposals that posit microscopic degrees of freedom residing either on the stretched horizon [71, 72] or along the observer's worldline [15–17].

If the microscopic Hilbert space is indeed finite-dimensional, its algebra of observables would be type-I [73] for which notions such as traces and entropy are all well-defined. While one might not

<sup>&</sup>lt;sup>8</sup>A Källén–Lehmann representation is discussed in the context of static patch in [70].

expect all these notions would survive in the semi-classical limit because of the change in the algebra type [74], it is possible that data such as relative traces or relative entropy remains intact. This makes objects like (6.1) a natural candidate observable in a microscopic-macroscopic dictionary, encoding information of concepts like QNMs that emerge in the semi-classical limit [75, 76]. Another relevant comment is that the fact that (6.1) is invariant under  $T: t \to -t$  resonates with the idea that T should be gauged in quantum gravity [77, 78]. We leave these speculative ideas for future investigation.

## Acknowledgments

It is a great pleasure to thank Frederik Denef, Raghu Mahajan, Adel Rahman, Eva Silverstein and Douglas Stanford for stimulating conversations, and especially Dionysios Anninos, Klaas Parmentier, Zimo Sun and András Vasy for useful discussions and comments on the draft. AL was supported in part by the Croucher Foundation, the Black Hole Initiative at Harvard University, and the Stanford Science Fellowship. MG was supported in part by the U.S. Department of Energy grant de-sc0011941.

## A Canonical quantization for free fields on the static patch

We review the canonical quantization procedure for free scalars in a  $dS_{d+1}$  static patch, which has been discussed in d+1=4 [79], d+1=2 [80], and later extended to general d [32].

Following section 2, a complete set of solutions to the Klein-Gordon equation is given by

$$\phi(t,x,\Omega) = \sum_{l=0}^{\infty} \int_{-\infty}^{\infty} d\omega \,\phi_{\omega l}(t,x,\Omega) a_{\omega l} = \sum_{l=0}^{\infty} \int_{-\infty}^{\infty} d\omega \, e^{-i\omega t} \, C_{\omega l}^{\text{n.}} \psi_{\omega l}^{\text{n.}}(x) Y_l(\Omega) a_{\omega l} \,\,, \tag{A.1}$$

where  $\psi_{\omega l}^{\text{n.}}(x)$  is the normalizable solution (2.9). The coefficients  $C_{\omega l}^{\text{n.}}$  are fixed by requiring

$$(\phi_{\omega l}, \phi_{\omega' l'})_{KG} = \delta(\omega - \omega')\delta_{ll'}, \qquad \omega, \omega' > 0,$$
(A.2)

and

$$(\phi_{\omega l}, \phi_{\omega' l'})_{KG} = -\delta(\omega - \omega')\delta_{ll'}, \qquad \omega, \omega' < 0,$$
(A.3)

where

$$(\phi_1, \phi_2)_{KG} \equiv -i \int d^d x \sqrt{-g} g^{tt} \left( \phi_1 \partial_t \phi_2^* - \partial_t \phi_1 \phi_2^* \right) \tag{A.4}$$

is the Klein-Gordon inner product. Imposing these leads to the condition

$$2|\omega||C_{\omega l}^{\mathbf{n}}|^2 \int_0^\infty dx \, \bar{\psi}_{\omega l}^{\mathbf{n}}(x) \bar{\psi}_{\omega' l}^{\mathbf{n}}(x) = \delta(\omega - \omega') + \delta(\omega + \omega') , \qquad \omega, \omega' \in \mathbb{R} , \qquad (A.5)$$

where  $\bar{\psi}_{\omega l}^{n}(x)$  is the rescaled version of (2.9). To obtain this, we note that when  $\omega$  and  $\omega'$  have the same sign, (A.2) and (A.3) implies (A.5) without the second term on the right. To extend to the

case when  $\omega$  and  $\omega'$  have opposite signs, observe that the left hand side of (A.5) is invariant under  $\omega \to -\omega$  or  $\omega' \to -\omega'$ ; the term  $\delta(\omega + \omega')$  is required so that both sides have this property.

From (A.1), we can also write down the mode expansion for its conjugate momentum,

$$\pi \equiv \frac{\partial \mathcal{L}}{\partial (\partial_t \phi)} = \sqrt{-g} g^{tt} \partial_t \phi . \tag{A.6}$$

Upon quantization, (A.1) is promoted to a self-adjoint field operator  $\hat{\phi}$ , with operator-valued coefficients  $\hat{a}_{\omega l}$ . Since  $\psi_{\omega l}^{\mathbf{n}}(x)$  is real, the self-adjointness of  $\hat{\phi}$  implies

$$C_{\omega l} = C_{-\omega l}^* \quad \text{and} \quad \hat{a}_{-\omega l} = \hat{a}_{\omega l}^{\dagger} .$$
 (A.7)

The equal-time commutation relations are

$$\left[\hat{\phi}(t,x,\Omega), \hat{\pi}(t,y,\Omega')\right] = i\delta(x-y)\delta^{(d-1)}(\Omega,\Omega') ,$$

$$\left[\hat{\phi}(t,x,\Omega), \hat{\phi}(t,y,\Omega')\right] = \left[\hat{\pi}(t,x,\Omega), \hat{\pi}(t,y,\Omega')\right] = 0 ,$$
(A.8)

which translate into the commutation relation

$$[\hat{a}_{\omega l}, \hat{a}_{\omega' l'}] = \operatorname{sgn}(\omega)\delta(\omega + \omega')\delta_{ll'}. \tag{A.9}$$

The vacuum state is defined as one annihilated by  $\hat{a}_{\omega l}$  for any  $\omega > 0$ :

$$\hat{a}_{\omega l} |0\rangle = 0 , \qquad \forall \omega > 0 , \quad l \ge 0 .$$
 (A.10)

By construction, such a (Boulware) vacuum is  $SO(1,1) \times SO(d)$ -invariant. This static vacuum is distinct from the global Euclidean vacuum which is SO(1,d+1)-invariant. For example, the expectation value of the stress tensor for a conformally coupled scalar in the static vacuum diverges at the horizon [81]. Acting with creation operators  $\hat{a}_{\omega l}^{\dagger}$  ( $\omega > 0$ ) on  $|0\rangle$  then yields states labeled by occupation numbers  $n_{\omega l}$  for each  $(\omega, l)$  mode. On these, the actions of  $\hat{a}_{\omega l}$  and  $\hat{a}_{\omega l}^{\dagger}$  are

$$\hat{a}_{\omega l} | \cdots n_{\omega l} \cdots \rangle = \sqrt{n_{\omega l}} | \cdots n_{\omega l} - 1 \cdots \rangle , \quad \hat{a}_{\omega l}^{\dagger} | \cdots n_{\omega l} \cdots \rangle = \sqrt{n_{\omega l} + 1} | \cdots n_{\omega l} + 1 \cdots \rangle . \quad (A.11)$$

We end this appendix by recalling that the quantum retarded correlator in the static vacuum

$$i \langle 0 | \left[ \hat{\phi}(t, x, \Omega), \hat{\phi}(0, y, \Omega') \right] | 0 \rangle \theta(t)$$
 (A.12)

satisfies the defining equations (2.14) and (2.15) for the retarded Green function associated with the Klein-Gordon equation, which is the reason why the analysis in section 2.1 could be performed without reference to the quantum theory.

# B Different correlators in thermal field theory

In this appendix we review some basics for the real-time and imaginary-time formalisms in thermal field theory [66, 67]. When we put the theory into a canonical ensemble, we are interested in computing the thermal averages of operators. We will use the notation

$$\langle \cdots \rangle_{\beta} \equiv \frac{1}{Z(\beta)} \operatorname{Tr} \left( e^{-\beta \hat{H}} \cdots \right) , \qquad Z(\beta) \equiv \operatorname{Tr} e^{-\beta \hat{H}} ,$$
 (B.1)

to denote the thermal average. In the following, we focus on bosonic operators  $\hat{O}(t)$ , which can be elementary fields or composite operators. For notational simplicity, we suppress other spatial dependence and quantum numbers such as spins.

Real-time correlators It is convenient to define the Wightman functions

$$G_{\beta}^{>}(t-t') \equiv \langle \hat{O}(t)\hat{O}(t')\rangle_{\beta}, \qquad G_{\beta}^{<}(t-t') \equiv \langle \hat{O}(t')\hat{O}(t)\rangle_{\beta},$$
 (B.2)

where we have assumed time translation invariance so that the functions only depend on t - t'. Without loss of generality we will set t' = 0 from now on. Using

$$e^{-\beta \hat{H}}\hat{O}(t)e^{\beta \hat{H}} = \hat{O}(t+i\beta) \tag{B.3}$$

and the cyclicity of the trace, one can derive the Kubo-Martin-Schwinger (KMS) relation

$$G_{\beta}^{>}(t) = G_{\beta}^{<}(t + i\beta) , \qquad (B.4)$$

or in frequency space,

$$\mathcal{G}_{\beta}^{>}(\omega) = e^{\beta\omega}\mathcal{G}_{\beta}^{<}(\omega) .$$
 (B.5)

Starting from (B.2), one can construct all other correlators by taking (anti-)commutators and multiplying the step functions  $\theta(\pm t)$ . A simple one is to take the average of (B.2) to obtain the symmetric Wightman function or statistical correlator:

$$G_{\beta}^{\text{Sym}}(t) \equiv \frac{1}{2} \langle \hat{O}(t)\hat{O}(0) + \hat{O}(0)\hat{O}(t) \rangle_{\beta} . \tag{B.6}$$

It is sometimes useful to study the so-called 2-sided correlator:

$$G_{\beta}^{12}(t) = G_{\beta}^{>} \left( t - \frac{i\beta}{2} \right)$$
 (B.7)

In linear response theory, the retarded and advanced propagators,

$$G_{\beta}^{R}(t) \equiv i \left\langle \left[ \hat{O}(t), \hat{O}(0) \right] \right\rangle_{\beta} \theta(t) , \qquad G_{\beta}^{A}(t) \equiv -i \left\langle \left[ \hat{O}(t), \hat{O}(0) \right] \right\rangle_{\beta} \theta(-t) ,$$
 (B.8)

often arise. Closely related is the so-called spectral or commutator function,

$$G_{\beta}^{C}(t) \equiv \left\langle \left[ \hat{O}(t), \hat{O}(0) \right] \right\rangle_{\beta},$$
 (B.9)

which plays a distinguished role in section 4.

In frequency space, the Wightman functions are related to the spectral function by

$$\mathcal{G}_{\beta}^{>}(\omega) = \frac{1}{e^{\beta\omega} - 1} \mathcal{G}_{\beta}^{C}(\omega) , \qquad \mathcal{G}_{\beta}^{<}(\omega) = \frac{1}{1 - e^{-\beta\omega}} \mathcal{G}_{\beta}^{C}(\omega) , \qquad (B.10)$$

implying

$$\mathcal{G}_{\beta}^{\text{Sym}}(\omega) = \coth \frac{\beta \omega}{2} \mathcal{G}_{\beta}^{C}(\omega) .$$
 (B.11)

We also have

$$\mathcal{G}_{\beta}^{C}(\omega) = -i\left(\mathcal{G}_{\beta}^{R}(\omega) - \mathcal{G}_{\beta}^{A}(\omega)\right) , \tag{B.12}$$

and

$$\mathcal{G}_l^{12}(\omega) = \frac{\mathcal{G}_l^C(\omega)}{\sinh \pi \omega} \ . \tag{B.13}$$

**Euclidean correlators** In non-perturbative formulations, it is convenient to define a correlator in the imaginary time  $0 \le \tau \le \beta$ :

$$G_{\beta}^{E}(\tau) = \left\langle \hat{O}(-i\tau)\hat{O}(0)\right\rangle_{\beta}.$$
 (B.14)

The KMS condition is equivalent to the periodicity of the Euclidean correlator:

$$G_{\beta}^{E}(\tau) = G_{\beta}^{E}(\tau + \beta) . \tag{B.15}$$

We can expand this into a Fourier series,

$$G_{\beta}^{E}(\tau) = \frac{1}{\beta} \sum_{k \in \mathbb{Z}} e^{-i\omega_{k}^{\text{Mat}}\tau} \mathcal{G}_{\beta}^{E}(\omega_{k}) , \qquad (B.16)$$

where  $\omega_k^{\text{Mat}}$  are the so-called Matsubara frequencies

$$\omega_k^{\text{Mat}} = \frac{2\pi k}{\beta} , \qquad k \in \mathbb{Z} .$$
 (B.17)

Free scalars on the static patch For the discussion of this paper, we work out some formulas for the case of a free scalar in the static patch. Plugging (the quantized version of) (A.1) into the Wightman function (B.2) and using (A.11), one can show

$$G_{\beta}^{>}(t;x,\Omega|y,\Omega') = \sum_{l=0}^{\infty} \int_{0}^{\infty} d\omega \left[ n_{B}(\omega) e^{i\omega t} + (n_{B}(\omega) + 1) e^{-i\omega t} \right] |C_{\omega l}^{\mathbf{n}}|^{2} \psi_{\omega l}^{\mathbf{n}}(x) \psi_{\omega l}^{\mathbf{n}}(y) Y_{l}(\Omega) Y_{l}(\Omega') . \tag{B.18}$$

We have set  $t_x = t, t_y = 0$ . The factor

$$n_B(\omega) \equiv \frac{1}{e^{\beta\omega} - 1} \tag{B.19}$$

is the Bose-Einstein distribution. Similarly,  $G_{\beta}^{<}(t; x, \Omega|y, \Omega')$  is given by (B.18) but with  $t \to -t$ . Taking the difference, it is easy to see that the Bose-Einstein factor (B.19) drops out, implying that the spectral function, and thus the retarded and advanced functions, are independent of the temperature, and therefore exactly equal to that at zero temperature:

$$G_{\beta}^{C}(t; x, \Omega | y, \Omega') = G_{\beta = \infty}^{C}(t; x, \Omega | y, \Omega')$$

$$= \sum_{l=0}^{\infty} \int_{0}^{\infty} d\omega \left( e^{-i\omega t} - e^{i\omega t} \right) |C_{\omega l}^{\mathbf{n}}|^{2} \psi_{\omega l}^{\mathbf{n}}(x) \psi_{\omega l}^{\mathbf{n}}(y) Y_{l}(\Omega) Y_{l}(\Omega') . \tag{B.20}$$

Taking the average of (B.18) and  $G_{\beta}^{\leq}(t;x,\Omega|y,\Omega')$ , one can see that (B.11) is satisfied.

Before we end, we note that by comparing (4.14) with the mode expansion (B.20), we can read off the normalization such that the condition (A.5) is satisfied:

$$|C_{\omega l}^{\text{n.}}|^2 = C_{\omega l}^{\text{n.}} C_{-\omega l}^{\text{n.}} = \frac{|\omega|}{\pi W_l^{\text{Jost}}(\omega) W_l^{\text{Jost}}(-\omega)}.$$
(B.21)

## C Spectral theory in 2D Rindler space

In this section, we study the scattering problem

$$\left[ -\partial_x^2 + \left( \frac{4\pi}{\beta} \right)^2 e^{-\frac{4\pi}{\beta}x} \right] \psi(x) = \omega^2 \psi(x) , \qquad (C.1)$$

associated with the Klein-Gordon equation on the 2D Rindler space,

$$ds^2 = e^{-\frac{4\pi}{\beta}x} \left( -dt^2 + dx^2 \right) , \qquad -\infty < x < \infty , \qquad (C.2)$$

with  $x \to -\infty$  and  $x \to \infty$  corresponding to the asymptotic infinity and horizon respectively. The Klein-Gordon equation for a scalar on any non-extremal black hole reduces to this form near the horizon, with  $\beta$  set by the black hole temperature. The problem has been partially analyzed before in [18]. The normalizable solution to (C.1) is

$$\psi_{\omega}^{\text{n.}}(x) = K_{\frac{i\beta\omega}{2\pi}} \left( 2e^{-\frac{2\pi x}{\beta}} \right) , \qquad (C.3)$$

which exponentially decays as  $x \to -\infty$ :

$$K_{\frac{i\beta\omega}{2\pi}} \left( 2e^{-\frac{2\pi x}{\beta}} \right) \propto e^{\frac{x}{2} - e^{-x}}$$
 (C.4)

Near the Rindler horizon, it is a linear combination of plane waves:

$$\psi_{\omega}^{\text{n.}}(x) \stackrel{x \to \infty}{\approx} \frac{1}{2} \Gamma\left(\frac{i\beta\omega}{2\pi}\right) e^{i\omega x} + \frac{1}{2} \Gamma\left(-\frac{i\beta\omega}{2\pi}\right) e^{-i\omega x} .$$
 (C.5)

We also obtain the Jost solution

$$\psi_{\omega}^{\text{Jost}}(x) = \Gamma \left( 1 - \frac{i\beta\omega}{2\pi} \right) I_{-\frac{i\beta\omega}{2\pi}} \left( 2e^{-\frac{2\pi x}{\beta}} \right) , \qquad (C.6)$$

which is purely ingoing at the horizon:

$$\psi_{\omega}^{\text{Jost}}(x) \stackrel{x \to \infty}{\approx} e^{i\omega x}$$
 (C.7)

The ratio between the coefficients of the outgoing and incoming waves of the normalizable mode in (C.5) defines a unitary S-matrix:

$$S^{\text{Rin}}(\beta,\omega) = -\frac{\Gamma\left(\frac{i\beta\omega}{2\pi}\right)}{\Gamma\left(-\frac{i\beta\omega}{2\pi}\right)}.$$
 (C.8)

The retarded Green function for (C.1) is given by [31]

$$2\pi \mathcal{G}^{R}(\omega; x|y) = \frac{\psi_{\omega}^{\text{n.}}(x)\psi_{\omega}^{\text{Jost}}(y)}{W^{\text{Jost}}(\omega)}\theta(y-x) + \frac{\psi_{\omega}^{\text{n.}}(y)\psi_{\omega}^{\text{Jost}}(x)}{W^{\text{Jost}}(\omega)}\theta(x-y) , \qquad (C.9)$$

where the Jost function can be computed to be

$$W^{\text{Jost}}(\omega) = \frac{2\pi}{\beta} \Gamma \left( 1 - \frac{i\beta\omega}{2\pi} \right) . \tag{C.10}$$

The spectral function is given by the difference

$$\mathcal{G}^{C}(\omega; x|y) = -i\left(\mathcal{G}^{R}(\omega; x|y) - \mathcal{G}^{R}(-\omega; x|y)\right). \tag{C.11}$$

Using the fact that  $K_{i\lambda}(z) = K_{-i\lambda}(z), \lambda \in \mathbb{R}$  and the identity

$$I_{i\lambda}(z) - I_{-i\lambda}(z) = -\frac{2i\sinh(\pi\lambda)}{\pi} K_{i\lambda}(z) , \qquad (C.12)$$

we compute

$$\mathcal{G}^{C}(\omega; x|y) = \frac{\beta}{2\pi^{3}} \sinh \frac{\beta \omega}{2} \psi_{\omega}^{\mathbf{n}.}(x) \psi_{\omega}^{\mathbf{n}.}(y) . \tag{C.13}$$

## References

- [1] D. Anninos, F. Denef, Y. T. A. Law, and Z. Sun, "Quantum de Sitter horizon entropy from quasicanonical bulk, edge, sphere and topological string partition functions," <u>JHEP</u> **01** (2022) 088, arXiv:2009.12464 [hep-th].
- [2] Y. T. A. Law, "A compendium of sphere path integrals," <u>JHEP</u> **21** (2020) 213, arXiv:2012.06345 [hep-th].
- [3] J. R. David and J. Mukherjee, "Partition functions of p-forms from Harish-Chandra characters," JHEP **09** (2021) 094, arXiv:2105.03662 [hep-th].
- [4] D. Anninos and E. Harris, "Three-dimensional de Sitter horizon thermodynamics," <u>JHEP</u> 10 (2021) 091, arXiv:2106.13832 [hep-th].
- [5] D. Anninos, T. Bautista, and B. Mühlmann, "The two-sphere partition function in two-dimensional quantum gravity," JHEP **09** (2021) 116, arXiv:2106.01665 [hep-th].
- [6] D. Anninos and B. Mühlmann, "The semiclassical gravitational path integral and random matrices (toward a microscopic picture of a dS<sub>2</sub> universe)," <u>JHEP</u> 12 (2021) 206, arXiv:2111.05344 [hep-th].
- [7] M. Grewal and K. Parmentier, "Characters, quasinormal modes, and Schwinger pairs in  $dS_2$  with flux," arXiv:2112.07630 [hep-th].

- [8] B. Mühlmann, "The two-sphere partition function from timelike Liouville theory at three-loop order," JHEP **05** (2022) 057, arXiv:2202.04549 [hep-th].
- [9] N. Bobev, T. Hertog, J. Hong, J. Karlsson, and V. Reys, "Microscopics of de Sitter Entropy from Precision Holography," <u>Phys. Rev. X</u> 13 no. 4, (2023) 041056, arXiv:2211.05907 [hep-th].
- [10] A. Castro, I. Coman, J. R. Fliss, and C. Zukowski, "Keeping matter in the loop in dS<sub>3</sub> quantum gravity," JHEP **07** (2023) 120, arXiv:2302.12281 [hep-th].
- [11] A. Castro, I. Coman, J. R. Fliss, and C. Zukowski, "Coupling Fields to 3D Quantum Gravity via Chern-Simons Theory," <u>Phys. Rev. Lett.</u> 131 no. 17, (2023) 171602, arXiv:2304.02668 [hep-th].
- [12] G. W. Gibbons and S. W. Hawking, "Action Integrals and Partition Functions in Quantum Gravity," Phys. Rev. D 15 (1977) 2752–2756.
- [13] V. Shyam, " $T\overline{T} + \Lambda_2$  deformed CFT on the stretched dS<sub>3</sub> horizon," <u>JHEP</u> **04** (2022) 052, arXiv:2106.10227 [hep-th].
- [14] E. Coleman, E. A. Mazenc, V. Shyam, E. Silverstein, R. M. Soni, G. Torroba, and S. Yang, "De Sitter microstates from  $T\overline{T} + \Lambda_2$  and the Hawking-Page transition," <u>JHEP</u> **07** (2022) 140, arXiv:2110.14670 [hep-th].
- [15] V. Chandrasekaran, R. Longo, G. Penington, and E. Witten, "An algebra of observables for de Sitter space," JHEP 02 (2023) 082, arXiv:2206.10780 [hep-th].
- [16] E. Witten, "A Background Independent Algebra in Quantum Gravity," arXiv:2308.03663 [hep-th].
- [17] D. Anninos, S. A. Hartnoll, and D. M. Hofman, "Static Patch Solipsism: Conformal Symmetry of the de Sitter Worldline," <u>Class. Quant. Grav.</u> 29 (2012) 075002, arXiv:1109.4942 [hep-th].
- [18] Y. T. A. Law and K. Parmentier, "Black Hole Scattering and Partition Functions," arXiv:2207.07024 [hep-th].
- [19] M. Grewal, Y. T. A. Law, and K. Parmentier, "Black hole horizon edge partition functions," JHEP 06 (2023) 025, arXiv:2211.16644 [hep-th].
- [20] P. Cvitanovic, R. Artuso, R. Mainieri, G. Tanner, and G. Vattay, <u>Chaos: Classical and Quantum</u>. Niels Bohr Inst., Copenhagen, 2016. http://ChaosBook.org/.
- [21] Y. T. A. Law, "Characters, Quasinormal Modes, and Quantum de Sitter Thermodynamics," in CORFU2022: 22th Hellenic School and Workshops on Elementary Particle Physics and Gravity. 4, 2023. arXiv:2304.01471 [hep-th].

- [22] T. Banks, "Cosmological breaking of supersymmetry?," <u>Int. J. Mod. Phys. A</u> **16** (2001) 910–921, arXiv:hep-th/0007146.
- [23] E. Witten, "Quantum gravity in de Sitter space," in <u>Strings 2001: International Conference</u>. 6, 2001. arXiv:hep-th/0106109.
- [24] M. K. Parikh and E. P. Verlinde, "De Sitter holography with a finite number of states," JHEP **01** (2005) 054, arXiv:hep-th/0410227.
- [25] G. W. Gibbons and S. W. Hawking, "Cosmological event horizons, thermodynamics, and particle creation," Phys. Rev. D 15 (May, 1977) 2738–2751. https://link.aps.org/doi/10.1103/PhysRevD.15.2738.
- [26] D. Borthwick, Spectral Theory of Infinite-Area Hyperbolic Surfaces, vol. 318 of Progress in Mathematics. Springer International Publishing. http://link.springer.com/10.1007/978-3-319-33877-4.
- [27] V. K. Dobrev, G. Mack, V. B. Petkova, S. G. Petrova, and I. T. Todorov, <u>Harmonic Analysis on the n-Dimensional Lorentz Group and Its Application to Conformal Quantum Field Tool.</u> 63. 1977.
- [28] J. Repka, "Tensor products of unitary representations of sl2(r)," American Journal of Mathematics **100** no. 4, (1978) 747–774. http://www.jstor.org/stable/2373909.
- [29] Z. Sun, "A note on the representations of SO(1, d+1)," arXiv:2111.04591 [hep-th].
- [30] E. S. C. Ching, P. T. Leung, W. M. Suen, and K. Young, "Wave propagation in gravitational systems: Late time behavior," Phys. Rev. D 52 (1995) 2118–2132, arXiv:gr-qc/9507035.
- [31] D. Skinner, "Mathematical methods," <u>Green's functions for PDE's, Lecture notes,</u>

  Department of Applied Mathematics and Theoretical Physics, University of Cambridge,

  Source: http://www.damtp.cam.ac.uk/user/dbs26/1Bmethods.html;(Accessed: 26.10.

  2018) (2014).
- [32] E. T. Akhmedov and D. V. Diakonov, "Free energy and entropy in Rindler and de Sitter space-times," Phys. Rev. D 105 no. 10, (2022) 105003, arXiv:2112.14794 [hep-th].
- [33] M. Dodelson, C. Iossa, R. Karlsson, and A. Zhiboedov, "A thermal product formula," arXiv:2304.12339 [hep-th].
- [34] D. L. Jafferis, A. Lupsasca, V. Lysov, G. S. Ng, and A. Strominger, "Quasinormal quantization in de Sitter spacetime," JHEP 01 (2015) 004, arXiv:1305.5523 [hep-th].
- [35] T. Basile, X. Bekaert, and N. Boulanger, "Mixed-symmetry fields in de Sitter space: a group theoretical glance," JHEP 05 (2017) 081, arXiv:1612.08166 [hep-th].

- [36] T. Hirai, "The characters of irreducible representations of the Lorentz group of *n*-th order,"

  <u>Proceedings of the Japan Academy</u> **41** no. 7, (1965) 526 531.

  https://doi.org/10.3792/pja/1195522333.
- [37] T. Hirai, "On irreducible representations of the Lorentz group of *n*-th order," Proceedings of the Japan Academy 38 no. 6, (1962) 258 262. https://doi.org/10.3792/pja/1195523378.
- [38] T. Hirai, "On infinitesimal operators of irreducible representations of the Lorentz group of *n*-th order," Proceedings of the Japan Academy **38** no. 3, (1962) 83 87. https://doi.org/10.3792/pja/1195523460.
- [39] Z. Sun, "Higher spin de Sitter quasinormal modes," arXiv:2010.09684 [hep-th].
- [40] G. 't Hooft, "On the Quantum Structure of a Black Hole," <u>Nucl. Phys. B</u> **256** (1985) 727–745.
- [41] J. Bonifacio, K. Hinterbichler, A. Joyce, and R. A. Rosen, "Shift Symmetries in (Anti) de Sitter Space," JHEP **02** (2019) 178, arXiv:1812.08167 [hep-th].
- [42] D. Anninos, T. Anous, B. Pethybridge, and G. Şengör, "The discreet charm of the discrete series in dS<sub>2</sub>," J. Phys. A **57** no. 2, (2024) 025401, arXiv:2307.15832 [hep-th].
- [43] B. Allen, "Vacuum States in de Sitter Space," Phys. Rev. D 32 (1985) 3136.
- [44] B. Allen and A. Folacci, "The Massless Minimally Coupled Scalar Field in De Sitter Space," Phys. Rev. D 35 (1987) 3771.
- [45] S. Grozdanov, K. Schalm, and V. Scopelliti, "Black hole scrambling from hydrodynamics," Phys. Rev. Lett. **120** no. 23, (2018) 231601, arXiv:1710.00921 [hep-th].
- [46] M. Blake, H. Lee, and H. Liu, "A quantum hydrodynamical description for scrambling and many-body chaos," <u>JHEP</u> 10 (2018) 127, arXiv:1801.00010 [hep-th].
- [47] M. Blake, R. A. Davison, S. Grozdanov, and H. Liu, "Many-body chaos and energy dynamics in holography," JHEP 10 (2018) 035, arXiv:1809.01169 [hep-th].
- [48] W. Research, "Hypergeometric2f1: Symbolic differentiation." http://functions.wolfram.com/07.23.20.0049.02, 2024. Accessed: February 26, 2024.
- [49] H. Epstein and U. Moschella, "de Sitter tachyons and related topics," Commun. Math. Phys. 336 no. 1, (2015) 381–430, arXiv:1403.3319 [hep-th].
- [50] C. Cheung, A. Helset, and J. Parra-Martinez, "Geometric soft theorems," JHEP 04 (2022) 011, arXiv:2111.03045 [hep-th].

- [51] D. Kapec, Y. T. A. Law, and S. A. Narayanan, "Soft scalars and the geometry of the space of celestial conformal field theories," <u>Phys. Rev. D</u> 107 no. 4, (2023) 046024, arXiv:2205.10935 [hep-th].
- [52] J. Bros, H. Epstein, and U. Moschella, "Scalar tachyons in the de Sitter universe," Lett. Math. Phys. 93 (2010) 203–211, arXiv:1003.1396 [hep-th].
- [53] J. Bonifacio, K. Hinterbichler, A. Joyce, and D. Roest, "Exceptional scalar theories in de Sitter space," JHEP 04 (2022) 128, arXiv:2112.12151 [hep-th].
- [54] A. Folacci, "Zero modes, euclideanization, and quantization," Phys. Rev. D 46 (Sep. 1992) 2553–2559. https://link.aps.org/doi/10.1103/PhysRevD.46.2553.
- [55] A. Higuchi, "Forbidden Mass Range for Spin-2 Field Theory in De Sitter Space-time," <u>Nucl.</u> Phys. B **282** (1987) 397–436.
- [56] Y. M. Zinoviev, "On massive high spin particles in AdS," arXiv:hep-th/0108192.
- [57] S. Deser and R. I. Nepomechie, "Anomalous Propagation of Gauge Fields in Conformally Flat Spaces," Phys. Lett. B **132** (1983) 321–324.
- [58] S. Deser and R. I. Nepomechie, "Gauge invariance versus masslessness in de sitter spaces," <u>Annals of Physics</u> 154 no. 2, (1984) 396-420. <a href="https://www.sciencedirect.com/science/article/pii/0003491684901568">https://www.sciencedirect.com/science/article/pii/0003491684901568</a>.
- [59] L. Brink, R. R. Metsaev, and M. A. Vasiliev, "How massless are massless fields in AdS(d)," Nucl. Phys. B 586 (2000) 183–205, arXiv:hep-th/0005136.
- [60] S. Deser and A. Waldron, "Gauge invariances and phases of massive higher spins in (A)dS," Phys. Rev. Lett. 87 (2001) 031601, arXiv:hep-th/0102166.
- [61] S. Deser and A. Waldron, "Partial masslessness of higher spins in (A)dS," <u>Nucl. Phys. B</u> **607** (2001) 577–604, arXiv:hep-th/0103198.
- [62] S. Deser and A. Waldron, "Stability of massive cosmological gravitons," Phys. Lett. B 508 (2001) 347–353, arXiv:hep-th/0103255.
- [63] S. Deser and A. Waldron, "Null propagation of partially massless higher spins in (A)dS and cosmological constant speculations," <u>Phys. Lett. B</u> 513 (2001) 137–141, arXiv:hep-th/0105181.
- [64] M. Grewal and Y. T. A. Law, "Spinning thermal static patch correlators." In preparation.
- [65] Y. T. A. Law and A. Vasy In preparation.
- [66] M. Laine and A. Vuorinen, <u>Basics of Thermal Field Theory: A Tutorial on Perturbative Computations</u>, vol. 925 of <u>Lecture Notes in Physics</u>. Springer International Publishing. http://link.springer.com/10.1007/978-3-319-31933-9.

- [67] J. I. Kapusta and C. Gale, <u>Finite-Temperature Field Theory: Principles and Applications</u>. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 2 ed., 2006.
- [68] J. Bros, "Complexified de Sitter space: Analytic causal kernels and Kallen-Lehmann type representation," Nucl. Phys. B Proc. Suppl. 18 (1991) 22–28.
- [69] M. Loparco, J. Penedones, K. Salehi Vaziri, and Z. Sun, "The Källén-Lehmann representation in de Sitter spacetime," JHEP 12 (2023) 159, arXiv:2306.00090 [hep-th].
- [70] M. Mirbabayi and F. Riccardi, "Probing de Sitter from the horizon," JHEP 04 (2023) 053, arXiv:2211.11672 [hep-th].
- [71] L. Susskind, L. Thorlacius, and J. Uglum, "The Stretched horizon and black hole complementarity," Phys. Rev. D 48 (1993) 3743-3761, arXiv:hep-th/9306069.
- [72] E. Shaghoulian, "The central dogma and cosmological horizons," <u>JHEP</u> **01** (2022) 132, arXiv:2110.13210 [hep-th].
- [73] E. Witten, "APS Medal for Exceptional Achievement in Research: Invited article on entanglement properties of quantum field theory," Rev. Mod. Phys. 90 no. 4, (2018) 045003, arXiv:1803.04993 [hep-th].
- [74] E. Witten, "Why Does Quantum Field Theory In Curved Spacetime Make Sense? And What Happens To The Algebra of Observables In The Thermodynamic Limit?," arXiv:2112.11614 [hep-th].
- [75] V. Narovlansky and H. Verlinde, "Double-scaled SYK and de Sitter Holography," arXiv:2310.16994 [hep-th].
- [76] K. Parmentier, "Coherent spin states and emergent de Sitter quasinormal modes," arXiv:2312.08430 [hep-th].
- [77] D. Harlow and T. Numasawa, "Gauging spacetime inversions in quantum gravity," arXiv:2311.09978 [hep-th].
- [78] L. Susskind, "A Paradox and its Resolution Illustrate Principles of de Sitter Holography," arXiv:2304.00589 [hep-th].
- [79] A. Higuchi, "Quantization of Scalar and Vector Fields Inside the Cosmological Event Horizon and Its Application to Hawking Effect," <u>Class. Quant. Grav.</u> 4 (1987) 721.
- [80] E. T. Akhmedov, K. V. Bazarov, D. V. Diakonov, and U. Moschella, "Quantum fields in the static de Sitter universe," <u>Phys. Rev. D</u> 102 no. 8, (2020) 085003, arXiv:2005.13952 [hep-th].
- [81] N. D. Birrell and P. C. W. Davies, <u>Quantum Fields in Curved Space</u>. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 1982.